

C_p -Theory for Model Theorists

Clovis Hamel^{1,2} and Franklin D. Tall²

December 2, 2021

Abstract

We present applications of C_p -theory, the branch of general topology concerned with spaces of real-valued continuous functions, to model theory, mostly in the context of continuous logics. We include C_p -theoretic results and proofs in a self-contained way for model theorists who are not familiar with the techniques of this field. We further generalize some results of Casazza and Iovino, and of the authors, involving the definability of Banach spaces including isomorphic copies of c_0 or ℓ^p , after a problem posed by Odell and Gowers.

¹ The first author is an NSERC Vanier Scholar.

² Research supported by NSERC Grant A-7354.

2010 Mathematics Subject Classification. 03C45, 03C75, 03C95, 03C98, 54C35, 46B99
Key words and phrases. Tsirelson's space, Gowers' problem, explicitly definable Banach spaces, C_p -theory, model-theoretic stability, definability, double limit conditions, Grothendieck spaces.

1 Introduction

C_p -theory is a subfield of general topology. Given a topological space X , $C_p(X)$ is the collection of continuous real-valued functions on X , considered as a subspace of \mathbb{R}^X . C_p -theory is concerned with the interrelations between the topology on X and the topology on $C_p(X)$. First identified as a distinct field by A. V. Arhangel'skiĭ, it has grown to the point of having a monograph [2], a four-volume problem book [37], and many research papers devoted to it. Perhaps the most famous theorem is that of A. Grothendieck [17], proved long before Arhangel'skiĭ's work:

Theorem 1.1. *If X is countably compact and $A \subseteq C_p(X)$, then the closure of A in $C_p(X)$ is compact if and only if A is countably compact in $C_p(X)$.*

The usefulness of Grothendieck's work in model theory, especially with regard to stability was recognized by J. Iovino [21], I. Ben Yaacov [4] and A. Pillay [28], all in the context of logics satisfying the compactness theorem.

When Iovino described his use of Grothendieck's theorem in his solution with Casazza [7] of a problem of Gowers [16] to the second author, the latter realized it could be considerably generalized by using the methods and results of C_p -theory. This was done in the recent [18], with the goal of introducing model theory to C_p -theorists unfamiliar with mathematical logic in general, and with model theory in particular. Here we instead target a model-theoretic audience unfamiliar with C_p -theory and not very knowledgeable of general topology. Our goal is to present enough C_p -theory so that model theorists can use it, as well as to again give an example of its usefulness by further generalizing the results of [7] and [18].

It is worth mentioning that [7] contained errors in its model-theoretic component that have been reworked in [9]. We refer to definitions in [7] and the results from analysis that appear in it, which are correct. At the end of Section 3, we discuss the different approaches from [7] and [9], and how they have influenced our work.

We have omitted some C_p proofs which are very easy or not particularly germane to the main line of our exposition. On the other hand, we have included some C_p proofs which are long and not easy, because the techniques are central or instructive or not easy to find in the literature. We have put these in an Appendix. We will obtain

our undefinability results using special cases which are easy to prove, but thought we should include the general cases for those interested.

The main application of the results occurs with regard to Gowers' problem [16] which asks whether every infinite dimensional explicitly definable Banach space includes an isomorphic copy of either c_0 or ℓ^p . All the definitions are given below but a model theorist would probably recognize at once that this question intersects her field as it is mainly about definability. There are two fundamental facts that motivate Gowers' problem:

- (1) All classical Banach spaces include a copy of c_0 or ℓ^p .
- (2) B. Tsirelson [40] constructed a Banach space not including copies of c_0 or ℓ^p . Such a space is regarded as the "first truly nonclassical Banach space" [16].

An interesting characteristic is that the norm of Tsirelson's space appears on both sides of the equality in its definition; this has become what analysts call an *implicitly definable* norm, as opposed to *explicitly definable* norms which are simply defined by a formula. (Of course one has to prove that such a norm is well-defined; this can be found in the book [8] devoted to the Tsirelson space.) After Tsirelson, many other pathological Banach spaces were constructed using similar techniques. Of course, the first problem to face is to formalize the question, and E. Odell says in his book [26] that this is a logician's task. The natural logics to approach this problem are *continuous logics* as they provide ample room for approximation and the analyst's ε -play. [7] and [9] presented both a formulation and a solution of this problem in the context of compact continuous logics, i.e., continuous logics which satisfy the Compactness Theorem. We shall present here generalizations of some results obtained in [7] and [9] to a wide class of logics with the help of C_p -theory, in particular to non-compact, countable fragments of continuous $\mathcal{L}_{\omega_1, \omega}$.

We shall allow ourselves to repeat well-known C_p -theoretic results and proofs from the literature for a reasonably self-contained exposition so that this work can serve as an introduction for model theorists wishing to learn C_p -theory.

Perhaps our title - a reflection of [18] - is too ambitious, given the four-volume [37]. A more modest one would be " C_p -theory and Definability."

2 Preliminaries in C_p -theory

2.1 Some Basic Results

We shall assume basic knowledge of general topology such as can be found in any of the standard texts for a first graduate course, such as [41]. Nonetheless, in footnotes we shall remind the reader of definitions she may have forgotten. Unless otherwise stated, we assume that all the topological spaces we consider are infinite, completely regular, and Hausdorff.

Definition 2.1. Let X be a completely regular* Hausdorff topological space. $C_p(X)$ is the space of real-valued continuous functions on X , endowed with the subspace topology inherited from the product topology on \mathbb{R}^X . Thus, the basic open sets of $C_p(X)$ are just the basic open sets of \mathbb{R}^X intersected with $C_p(X)$, i.e.

$$W(g; x_0, \dots, x_n; \varepsilon) = \{f \in C_p(X) : (\forall i \leq n)(|f(x_i) - g(x_i)| < \varepsilon)\}.$$

Remark. It follows from the form of the basic open sets of \mathbb{R}^X and complete regularity that $C_p(X)$ is dense in \mathbb{R}^X (given any function $g \in \mathbb{R}^X$ and a basic open set in \mathbb{R}^X about g , say $U = U(g; x_0, \dots, x_n; \varepsilon) = \{f \in \mathbb{R}^X : (\forall i \leq n)(|f(x_i) - g(x_i)| < \varepsilon)\}$, we can find a continuous function $h \in C_p(X)$ such that $(\forall i \leq n)h(x_i) = f(x_i)$ and so $h \in U$). We shall occasionally deal with $C_p(X, [0, 1])$ and $C_p(X, 2)$, with the obvious meanings. Notice that each of these is a closed subspace of $C_p(X)$. Since the main properties we deal with are closed-hereditary, if we prove results for $C_p(X)$, they apply also to the other two cases. In particular, this means that our results can be applied to first order logic, in which $C_p(X, 2)$ is relevant.

Lemma 2.2. *Let X be any topological space, $Y \subseteq X$, and $\pi_Y : C_p(X) \rightarrow C_p(Y)$ the restriction map given by $\pi_Y(f) = f|_Y$, f restricted to Y . Then:*

(i) π_Y is continuous.

(ii) $\overline{\pi_Y(C_p(X))} = C_p(Y)$.

* A space X is *completely regular* if for each closed set F and each point x not in F there is a continuous real-valued $f : X \rightarrow \mathbb{R}$ such that $f(x) = 0$ and $f(F) = \{1\}$.

- (iii) If $Y \subseteq X$ is dense, then $\pi_Y : C_p(X) \rightarrow \pi_Y(C_p(X))$ is a continuous bijection.
- (iv) If $Y \subseteq X$ is closed, then $\pi_Y : C_p(X) \rightarrow \pi_Y(C_p(X))$ is an open map.

Proof.

- (i) Note that π_Y is continuous as the pre-image of a basic open set of $C_p(Y)$ is a basic open set of $C_p(X)$.
- (ii) Take $f \in C_p(Y)$ and an open neighbourhood

$$U = W(f; x_0, \dots, x_n; \varepsilon) \subseteq C_p(Y).$$

Since X is completely regular, we verify that there is a $g \in C_p(X)$ such that $(\forall i \leq n)(f(x_i) = g(x_i))$: for each $i \leq n$, let $g_i \in C_p(X)$ be such that $g_i(x_i) = 1$ and $g_i(x_j) = 0$ for each $i \neq j$. It suffices to take $g(x) = f(x_0)g_0(x) + \dots + f(x_n)g_n(x)$, which is continuous. Then $\pi_Y(g) \in U$.

- (iii) Injectivity follows from the following well-known fact: if $f, g \in C_p(X)$ and they agree on a dense subspace, then $f = g$.
- (iv) Take a basic open set $W(f; x_0, \dots, x_n; \varepsilon)$ of $C_p(X)$. Suppose $x_0, \dots, x_k \in Y$ and $x_{k+1}, \dots, x_n \in X \setminus Y$ for some $0 \leq k \leq n$. It suffices to show $\pi_Y[W(f; x_0, \dots, x_n; \varepsilon)] = W(\pi_Y(f); x_0, \dots, x_k; \varepsilon) \cap \pi_Y(C_p(X))$. The inclusion

$$\pi_Y[W(f; x_0, \dots, x_n; \varepsilon)] \subseteq W(\pi_Y(f); x_0, \dots, x_k; \varepsilon) \cap \pi_Y(C_p(X))$$

is immediate by the definition of $\pi_Y(f)$. To prove the other inclusion, suppose

$$g \in W(\pi_Y(f); x_0, \dots, x_k; \varepsilon) \cap \pi_Y(C_p(X))$$

with $\pi_Y(h) = g$. By complete regularity and the fact that Y is closed, there is a continuous function $l \in C_p(X)$ such that $l|_Y = 0$ and, for each $k+1 \leq i \leq n$, $l(x_i) = f(x_i) - h(x_i)$. Then $l+h \in W(f; x_0, \dots, x_n; \varepsilon)$ and $\pi_Y(l+h) = g$. \square

Recall that if $\varphi : X \rightarrow Y$ where X and Y are topological spaces, we say that φ is a *homeomorphism* if it is a continuous bijection with a continuous inverse.

A basic notion that will be relevant is the *dual map*: if $f : X \rightarrow Y$ is any function, the dual map of f is the function $\Phi_f : \mathbb{R}^Y \rightarrow \mathbb{R}^X$ given by $\Phi_f(g) = g \circ f$. The following result [2] collects the facts we shall need:

Lemma 2.3. *Let $f : X \rightarrow Y$ be any function. Then:*

- (i) Φ_f is continuous.
- (ii) If f is surjective, then Φ_f is a homeomorphism between \mathbb{R}^Y and $\Phi_f(\mathbb{R}^Y)$.
- (iii) If f is a continuous surjection, then $\Phi_f|_{C_p(Y)}$ is a homeomorphism between $C_p(Y)$ and $\Phi_f(C_p(Y)) \subseteq C_p(X)$.

Proof.

- (i) Suppose $\Phi_f(g) = h$. Consider a basic open neighbourhood $W(h; x_1, \dots, x_n; \varepsilon)$. Then

$$\Phi_f(W(g; f(x_1), \dots, f(x_n); \varepsilon)) \subseteq W(h; x_1, \dots, x_n; \varepsilon).$$

- (ii) Let $g_1, g_2 \in \mathbb{R}^Y$ be different functions. Take $y \in Y$ such that $g_1(y) \neq g_2(y)$. Since f is surjective, take $x \in f^{-1}(y)$. Then $\Phi_f(g_1)(x) \neq \Phi_f(g_2)(x)$. It remains to show that Φ_f^{-1} is continuous. Suppose $\Phi_f(g) = h$, and consider a basic open set $W(g; y_1, \dots, y_n; \varepsilon)$. By surjectivity, we can take $x_1, \dots, x_n \in X$ such that $f(x_1) = y_1, \dots, f(x_n) = y_n$. Then

$$\Phi_f^{-1}(W(h; x_1, \dots, x_n; \varepsilon) \cap \Phi_f(\mathbb{R}^Y)) \subseteq W(g; y_1, \dots, y_n; \varepsilon).$$

- (iii) Immediate from (ii) by noting that the range $\Phi_f|_{C_p(Y)}$ is included in $C_p(X)$.

□

The following results sample the interrelations between the topology of X and the one on $C_p(X)$. First, we introduce some topological cardinal functions.

Definition 2.4. Let (X, \mathcal{T}) be a topological space.

- (a) A *base* for X is a family \mathcal{B} of open sets such that $(\forall x \in X)(\forall U \in \mathcal{T})[(x \in U) \implies (\exists V \in \mathcal{B})(x \in V \subseteq U)]$. The *weight* $w(X)$ of X is the minimal cardinality of a base for X .
- (b) A *local base* of X at a point $x \in X$ is a family \mathcal{B}_x of open sets such that $x \in \bigcap \mathcal{B}_x$ and for any open set U containing x , there is some $V \in \mathcal{B}_x$ such that $x \in V \subseteq U$. The *character* of X at x , denoted $\chi(x, X)$, is the minimal cardinality of a local base of X at x . The *character* of X is defined by $\chi(X) = \sup\{\chi(x, X) : x \in X\}$.

Theorem 2.5. *Given a topological space X , the following equalities hold:*

$$|X| = \chi(C_p(X)) = w(C_p(X)).$$

Proof. It follows from the definitions (and our assumption that X is infinite) that

$$\chi(C_p(X)) \leq w(C_p(X)) \leq w(\mathbb{R}^X) \leq |X|,$$

so it suffices to prove $|X| \leq \chi(C_p(X))$. Suppose, on the contrary, that the strict inequality $\chi(C_p(X)) < |X|$ holds, and denote by $\mathbf{0}$ the function that is identically 0. Let \mathcal{B}_0 be a local base of $C_p(X)$ at $\mathbf{0}$ such that $|B_0| < |X|$. We may assume that \mathcal{B}_0 consists of only basic open sets of $C_p(X)$. Now consider the following set

$$Y = \{y \in X : (\exists n < \omega)(\exists x_0, \dots, x_n \in X)(\exists \varepsilon > 0) \\ (W(\mathbf{0}; x_0, \dots, x_n, y; \varepsilon) \in \mathcal{B}_0)\}.$$

By our assumption, $|Y| < |X|$. Take $y \in X \setminus Y$ and consider the open set $U = W(\mathbf{0}; y; 1)$, which is a neighbourhood of $\mathbf{0}$. For any $V = W(\mathbf{0}; x_0, \dots, x_n; \varepsilon) \in \mathcal{B}_0$, we have $(\forall i < n)(x_i \neq y)$. By complete regularity, there is a $g \in C_p(X)$ such that, for all $i < n$, $g(x_i) = 0$ and $g(y) = 1$. Then $g \in V \setminus U$, contradicting that \mathcal{B}_0 is a local base. \square

It follows from Theorem 2.5 that metrizability and first countability* are indistinguishable in $C_p(X)$, i.e. $C_p(X)$ is metrizable if and only if $C_p(X)$ is first countable if and only if X is countable.

On the way toward Grothendieck's theorem, we introduce some definitions:

Definition 2.6. Let X be a topological space.

- (a) A subset $A \subseteq X$ is *relatively compact* (in X) if \overline{A} is compact.
- (b) X is *countably compact* if every infinite subset Y of X has a *limit point*, i.e. a point x such that every open set about x contains infinitely many points of Y . If $A \subseteq B \subseteq X$, A is *countably compact in B* if every infinite subset of A has a limit point in B .

* A space X is first countable if $\chi(X) \leq \aleph_0$.

- (c) A set of real-valued functions $A \subseteq \mathbb{R}^X$ is *pointwise bounded* if for every $x \in X$ there is an $M_x > 0$ such that for all $f \in A$, $|f(x)| < M_x$.
- (d) The *Lindelöf number* $L(X)$ of X is the smallest infinite cardinal such that every open cover contains a subcover of cardinality $\leq L(X)$. X is *Lindelöf* if $L(X) = \aleph_0$.

Definition 2.6 (b) can be restated in terms that justify the choice of words for "countable compactness":

Proposition 2.7. *A T_1 space* X is countably compact if and only if every countable open cover of X includes a finite subcover.*

Proof. Suppose X is countably compact and suppose that there is a countable open cover $\{U_n : n < \omega\}$ that does not include a finite subcover. For each $n < \omega$, pick $x_n \in X \setminus (U_0 \cup \dots \cup U_{n-1})$, which is possible since $U_0 \cup \dots \cup U_{n-1} \neq X$. Then $\{x_n : n < \omega\}$ doesn't have a limit point[†], a contradiction.

Conversely, suppose every countable open cover of X includes a finite subcover and that there is an infinite subset $A \subseteq X$ with no limit points. Then A is closed and discrete and so all its subsets are also closed and discrete. Take a countable subset $B = \{b_n : n < \omega\} \subseteq A$ and let, for each $n < \omega$, $U_n = X \setminus \{b_k : k \geq n\}$. By T_1 , $\{U_n : n < \omega\}$ is an open cover of X which doesn't include a finite subcover, a contradiction. \square

The following standard result will be used in the proof of Grothendieck's theorem.

Proposition 2.8. *If X is a normal space and $A \subseteq X$ is countably compact in X , then \overline{A} is countably compact.*

Proof. Suppose that \overline{A} is not countably compact. Then there is a countably infinite closed discrete subset B of \overline{A} . Let $B = \{b_n\}_{n < \omega}$. It suffices to find disjoint open sets $\{U_n\}_{n < \omega}$, $b_n \in U_n$, for then we can use normality to obtain disjoint open sets U, W , $U \supseteq B$, $V \supseteq X \setminus \bigcup_{n < \omega} U_n$. Then the open sets $\{U \cap U_n\}_{n < \omega}$ form a discrete collection, i.e. each

* A space is T_1 if each point is a closed set.

† A point x is a limit point of a space Y if every open set about x intersects Y .

point in X is at most one of these. Take $a_n \in U \cap U_n \cap A$. Then $\{a_n : n < \omega\}$ has no limit points in X , contradicting A being countably compact in X . To find the required disjoint open sets, by normality find disjoint open sets V_n containing b_n and $V'_n \supseteq \{b_k : k \neq n\}$. Let $U_n = V_n \cap \bigcap \{V'_k : k < n\}$. \square

Proposition 2.8 is not true without some hypothesis such as normality. A standard example is the *tangent disk space* X . Consider the unit square, including the x -axis, Give the points above the x -axis their usual topology. A basic neighbourhood of a point on the axis is the point itself together with an open disk tangent to the axis at that point. Consider the set Y consisting of all points above the x -axis with both coordinates rational. Then Y is countably compact in X , but its closure is all of X , which includes the subset $[0, 1]$ of the closed discrete x -axis, so is not countably compact.

Now we proceed to prove Grothendieck's Theorem, which also relates properties of X to properties of $C_p(X)$. The proof given here is a variation on the one that appears in [39].

Theorem 2.9. *Let X be a countably compact topological space, and $A \subseteq C_p(X)$. Then A is relatively compact (in $C_p(X)$) if and only if it is countably compact in $C_p(X)$.*

Proof. If A is relatively compact in $C_p(X)$, then $\overline{A} \cap C_p(X)$ is compact and so A is countably compact in $C_p(X)$.

Conversely, suppose A is countably compact in $C_p(X)$; then it is pointwise bounded (otherwise there would be a sequence $\langle f_n : n < \omega \rangle$ in A and an $x \in X$ such that for each $n < \omega$, $|f_n(x)| > n$, and there can be no limit point of such a sequence). For each $x \in X$, let $M_x > 0$ witness that A is pointwise bounded. Then the closure \overline{A} in \mathbb{R}^X is compact since it is a closed subset of the compact $\prod_{x \in X} [-M_x, M_x]$. Then it suffices to show $\overline{A} \subseteq C_p(X)$.

Assume to the contrary that there is some $g \in \overline{A} \setminus C_p(X)$. Then since g is not continuous there are $\varepsilon > 0$ and $y \in X$ such that y is in the closure of $Y = X \setminus g^{-1}[(g(y) - \varepsilon, g(y) + \varepsilon)]$. We recursively build the following sequences: a sequence $\{U_n\}_{n < \omega}$ of open sets containing y ; a sequence of points $\{x_n\}_{n < \omega}$ in Y ; and a sequence of functions $\{f_n\}_{n < \omega}$ in A such that the following conditions are satisfied for each $n < \omega$:

- (i) $\overline{U_{n+1}} \subseteq U_n$.

- (ii) $(\forall x \in U_n)(|f_n(x) - f_n(y)| < \varepsilon/2^n)$.
- (iii) $x_n \in U_n \cap Y$.
- (iv) $(\forall i < n)(|f_{n+1}(x_i) - g(y)| > \varepsilon/2)$.
- (v) $|f_n(y) - g(y)| < \varepsilon/2^n$.

First, take $f_0 \in W(g; y; \varepsilon)$, U_0 the inverse image under f_0 of a ball of radius ε around $f_0(y)$ and $x_0 \in U_0 \cap Y$ (where $U_0 \cap Y$ is non-empty since y is a limit point of Y).

Suppose U_i , x_i , and f_i have been constructed for each $i \leq n$. Take

$$f_{n+1} \in W(g; x_1, \dots, x_n; \varepsilon/2) \cap W(g; y; \varepsilon/2^{n+1}) \cap A,$$

which is nonempty since g is a limit point of A . Since we have that $f_{n+1} \in W(g; y; \varepsilon/2^{n+1})$, it satisfies (v). Note that f_{n+1} also satisfies (iv) since for each $i < n$,

$$|f_{n+1}(x_i) - g(y)| \geq |g(x_i) - g(y)| - |f_{n+1}(x_i) - g(x_i)| > \varepsilon/2$$

by construction. Since f_{n+1} is continuous, we may take an open set U_{n+1} containing y satisfying (ii), and such that $\overline{U_{n+1}} \subseteq U_n$, which is (i); this is done by considering the inverse image under f_{n+1} of a ball of radius at most $\varepsilon/2^{n+1}$ and centred at $f_{n+1}(y)$, then taking its intersection with U_n and using regularity to shrink this to get $\overline{U_{n+1}} \subseteq U_n$. We may then pick some $x_{n+1} \in U_{n+1} \cap Y$ so that (iii) is satisfied.

Now X is countably compact, so let x_∞ be a limit point of $\{x_n\}_{n < \omega}$. Then $x_\infty \in \bigcap_{n < \omega} \overline{U_n}$, since $x_{n+1} \in U_{n+1} \subseteq \overline{U_{n+1}} \subseteq U_n$.

Let $S = \{x_n\}_{n < \omega} \cup \{x_\infty\}$ and consider the restriction map $\pi_S : C_p(X) \rightarrow C_p(S)$ given by $\pi_S(f) = f|_S$, f restricted to S , which is continuous by Lemma 2.2. Thus, $F = \pi_S(A)$ is a subset of \mathbb{R}^S , which is metrizable, since S is countable. By the continuity of π_S , F is countably compact in $C_p(S)$: a countably infinite subset of F is of the form $\{h_n|_S : n < \omega\}$ where each h_n belongs to A and so $\{h_n : n < \omega\}$ has a limit point $h \in C_p(X)$. Then $\pi_S(h) = h|_S \in C_p(S)$ is a limit point of the original set. Since $C_p(S)$ is a metrizable space (by Theorem 2.5), the closure \overline{F} of F in $C_p(S)$ is countably compact by Proposition 2.8. Countable compactness and compactness are equivalent for metrizable spaces (metrizable spaces are *paracompact* and paracompact countably compact spaces are compact, see [14]); thus \overline{F} is compact. In this particular case, there is also a simpler proof: S is countable, so \mathbb{R}^S has a countable base so is Lindelöf. Clearly, Lindelöf

countably compact spaces are compact. Now, let $h \in \overline{F} \subseteq C_p(S)$ be a limit point of $\{f_n|_S\}_{n < \omega}$. We verify that our construction implies $h(x_\infty) \notin \overline{\{h(x_n)\}_{n < \omega}}$.

First, notice that (ii) implies $h(x_\infty) = g(y)$. Assume for a contradiction that there is some $N < \omega$ such that $|h(x_N) - h(x_\infty)| < \varepsilon/4$, and take $M > N$ such that $f_M|_S \in W(h; x_N; \varepsilon/4)$. Thus,

$$|f_M(x_N) - g(y)| \leq |f_M(x_N) - h(x_N)| + |h(x_N) - h(x_\infty)| < \varepsilon/2,$$

which contradicts (iv). Thus there is some $r > 0$ such that

$$(\forall n < \omega)(|h(x_n) - h(x_\infty)| > r),$$

and so h is not continuous at x_∞ , contradicting $h \in C_p(S)$ and $x_\infty \in S$.

We conclude that $\overline{A} \subseteq C_p(X)$ and so A is relatively compact in $C_p(X)$. \square

Definition 2.10. A g -space is a topological space X satisfying the following property: for every $A \subseteq X$, if A is countably compact in X then A is relatively compact. We say that X is a *hereditary g -space* if every subspace of X is a g -space.

The Fréchet-Urysohn property arises naturally in the context of g -spaces.

Definition 2.11. Let X be a topological space. X is *Fréchet-Urysohn* if whenever $A \subseteq X$, every point p in \overline{A} is the limit of a sequence of points in A , i.e. every open set about p contains all but finitely many members of the sequence.

The following proposition from [3] will be useful for generalizing Grothendieck's Theorem:

Proposition 2.12. *A g -space X is a hereditary g -space if and only if every compact subspace of X is Fréchet-Urysohn.*

Proof. Let X be a g -space. First, assume that every compact subspace of X is Fréchet-Urysohn. Let $Y \subseteq X$, and let $A \subseteq Y$ be countably compact in Y , so it is also countably compact in X . Since X is a g -space, \overline{A} is compact. Then \overline{A} is Fréchet-Urysohn by our assumption. Since every element in \overline{A} is the limit of a sequence in A , we have $\overline{A} \cap (X \setminus Y) = \emptyset$. For if $x \in X \setminus Y$ were a limit of a sequence $\langle x_n : n < \omega \rangle$ from A , then $\{x_n : n < \omega\}$ would contradict A being

countably compact in Y . Thus $\overline{A} \subseteq Y$. This shows that \overline{A} is also compact in Y , showing that Y is a g -space as desired.

Conversely, suppose X is a hereditary g -space and assume for a contradiction that there is some $A \subseteq X$ such that \overline{A} is compact and there is an $x \in \overline{A}$ that is not the limit of a sequence in A . Then $Y = \overline{A} \setminus \{x\}$ is a g -space. We claim that A is countably compact in Y : take a countably infinite set $\{x_n : n < \omega\}$ in A . Any limit point of such a set must lie in \overline{A} . Then it is enough to check that x is not the unique limit point of that set. For the sake of contradiction, suppose that x is the unique limit point of that set. Now consider $\{x_n : n < \omega\}$ as the sequence $\langle x_n : n < \omega \rangle$. By assumption, no subsequence can converge to x . This means that there is a neighbourhood U containing x such that for any $N < \omega$ there are infinitely many $k \geq N$ such that $x_k \notin U$. These x_k 's constitute an infinite subset of A with no limit point in the compact \overline{A} , which is impossible. Then x cannot be the unique limit point of an infinite subset of A and so A is countably compact in Y , so it is relatively compact in the g -space Y . Since X is a hereditary g -space, A is compact, hence $A = \overline{A}$ is closed. But then $x \in A$, a contradiction. \square

The following definition is the starting point for generalizing Grothendieck's Theorem to a broader class of spaces:

Definition 2.13 ([3]). A topological space X is *Grothendieck* if $C_p(X)$ is a hereditary g -space. X is *weakly Grothendieck* if $C_p(X)$ is a g -space.

Any uncountable discrete space is an example of a space that is weakly Grothendieck but not Grothendieck [1].

The property of being a Grothendieck space is transferable from a dense subspace to the full space. This result appears in [3] which quotes Pryce [29]. In order to prove it, we need the following lemma, which is a variation on the well-known fact that a continuous bijection between compact spaces is necessarily a homeomorphism.

Lemma 2.14. *Let X be countably compact, Y Fréchet-Urysohn, and $f : X \rightarrow Y$ a continuous bijection. Then f is a homeomorphism.*

Proof. Suppose $f : X \rightarrow Y$ is a continuous bijection as above, $A \subseteq Y$, and $y \in \overline{A}$. Since Y is Fréchet-Urysohn we have a sequence $S = \{y_n\}_{n < \omega}$ in A converging to y . Then no other $x \neq f^{-1}(y)$ can be a limit point of $f^{-1}(S)$. However, by countable compactness, $f^{-1}(S)$ has a limit point, which must therefore be $f^{-1}(y)$. Then $f^{-1}(y) \in \overline{f^{-1}(A)}$,

so $f^{-1}(\overline{A}) \subseteq \overline{f^{-1}(A)}$ for any $A \subseteq Y$, hence f^{-1} is continuous and f is a homeomorphism. \square

Theorem 2.15. *Let Y be a dense subspace of X . If Y is Grothendieck, so is X .*

Proof. Consider the restriction map $\pi_Y : C_p(X) \rightarrow C_p(Y)$. By Lemma 2.2, π_Y is a continuous bijection of $C_p(X)$ onto a subspace Z of $C_p(Y)$. Since $C_p(Y)$ is a hereditary g -space, then so is Z .

If $A \subseteq C_p(X)$ is a subspace, and C is countably compact in A , then $\pi_Y(C) \subseteq \pi_Y(A)$ is countably compact in $\pi_Y(A)$ and thus relatively compact. By Proposition 2.12, $\overline{\pi_Y(C)}$ is Fréchet-Urysohn and we conclude by using Lemma 2.14. \square

Arhangel'skiĭ [3] proved the preservation of Grothendieck spaces under continuous images:

Theorem 2.16. *The continuous image of a Grothendieck space is Grothendieck.*

Proof. Suppose X is Grothendieck, Y a topological space, and $f : X \rightarrow Y$ a continuous surjection. By Lemma 2.3, $C_p(Y)$ is homeomorphic to a subspace of $C_p(X)$. Since $C_p(X)$ is a hereditary g -space, so is $C_p(Y)$. \square

Notice that the previous two results rely on $C_p(X)$ being a hereditary g -space, so the same proof does not work if we merely assume that the space is weakly Grothendieck.

2.2 Lindelöf Σ -spaces are Grothendieck

Definition 2.17. Let X be a topological space.

- (a) X is a *Lindelöf p -space* if and only if it can be perfectly mapped onto a space with a countable base. Recall that a map is *perfect* if it is continuous, closed, and the pre-images of points are compact.
- (b) X is a *Lindelöf Σ -space* if and only if it is the continuous image of a Lindelöf p -space.

These are not the original definitions but rather, equivalent ones. Perfect maps amount to just collapsing some compact sets to points. They are frequently used in general topology since they preserve many

properties, either in the direct image or the inverse image. The definitions of p -spaces and Σ -spaces are quite complicated. Notice that countable products of Lindelöf p -spaces—in particular, countable spaces—are also Lindelöf p -spaces: suppose $\{X_n : n < \omega\}$ is a family of Lindelöf p -spaces. For each $n < \omega$, fix a perfect surjection $f_n : X_n \rightarrow M_n$ where M_n is second countable*. Then the diagonal product $f : \prod_{n < \omega} X_n \rightarrow \prod_{n < \omega} M_n$ given by $f(x_0, x_1, \dots) = (f_0(x_0), f_1(x_1), \dots)$ is perfect (see [14], p. 185, for the details) and its range is a second countable space since second countability is hereditary and preserved by countable products. Thus, $\prod_{n < \omega} X_n$ is a Lindelöf p -space.

The class of spaces which are Grothendieck has been widely studied by Arhangel'skiĭ [3]. Following [3], we shall prove in the Appendix that

Theorem 2.18. *All Lindelöf Σ -spaces are Grothendieck.*

Lindelöf Σ -spaces constitute a nice class as they can be also described as follows: The class of Lindelöf Σ -spaces is the smallest class of spaces containing all compact spaces, all second countable spaces, and that is closed under finite products, closed subspaces, and continuous images ([38] contains a detailed exposition of these facts).

Examples of Lindelöf Σ -spaces include compact spaces, *Polish spaces*, and *K -analytic spaces* (see [38]). They coincide with the *K -countably-determined spaces* of [32]. We find it useful to think of Lindelöf Σ -spaces as a weakening of K -analytic spaces. Those latter spaces have found applications in Analysis as a generalization of *Polish* (separable completely metrizable) spaces. They can be characterized as continuous images of perfect pre-images of Polish spaces. Thus, Lindelöf Σ -spaces just drop “completeness” from K -analytic spaces. These more topological—as contrasted with descriptive set-theoretic—definitions of K -analytic and K -countably determined spaces were apparently not known to Rogers and Jayne. For further discussion see [35].

The proof that all Lindelöf Σ -spaces are Grothendieck is not easy. We have put it into the Appendix for those who want to learn more C_p -theory and are willing to invest some effort. It is put together from several important theorems of C_p -theory. It turns out, however, that for the particular application to undefinability of Banach spaces that we are interested in, we only need to prove that *countable* spaces are Grothendieck! The proof of this is quite easy so we give it in the main

* A space X is *second countable* if $w(X) \leq \aleph_0$.

text. The reason we can get away with this trivial case is that we only deal with countable languages, such as first order continuous logic and countable fragments of continuous $\mathcal{L}_{\omega_1, \omega}$.

Definition 2.19. The *tightness* $t(X)$ of X is the smallest infinite cardinal such that whenever $A \subseteq X$ and $x \in \overline{A}$, there exists a $B \subseteq A$ such that $|B| \leq t(X)$ and $x \in \overline{B}$. When $t(X) = \aleph_0$, we say that X is *countably tight*.

Countable tightness plays an important role in the study of Grothendieck spaces, as we shall see in the proof that Lindelöf Σ -spaces are Grothendieck. There, it is the countable tightness of $C_p(X)$ that is important, and this is obtained from the fact that countable products of Lindelöf Σ -spaces are Lindelöf. It is also important that

Theorem 2.20. *Countably tight spaces are weakly Grothendieck.*

Theorem 2.20 is stated in [3] as part of a much more general result. A different proof is in [36].

The second author [36] has recently shown that:

Theorem 2.21. *The Proper Forcing Axiom (PFA) implies that Lindelöf countably tight spaces are Grothendieck.*

The conclusion does not hold if $\mathbf{V} = \mathbf{L}$ (see [36]). Also in the realm of set theory is Arhangel'skiĭ's theorem which depends on work of Okunev [27]:

Theorem 2.22 ([3]). *\mathbf{MA}_{\aleph_1} implies countably tight spaces with finite powers Lindelöf are Grothendieck.*

As we said, all we will need in order to prove our definability result is:

Theorem 2.23. *Every countable space is Grothendieck.*

Proof. By Theorem 2.5, if X is countable, $C_p(X)$ has a countable base. It follows easily that all subspaces of $C_p(X)$ are Fréchet-Urysohn, so we need only show that a space with a countable base is a g -space. A space with a countable base is separable metrizable, so normal, so by Proposition 2.8, if A is countably compact in $C_p(X)$, then \overline{A} is countably compact. But $C_p(X)$ is Lindelöf, so \overline{A} is compact. \square

2.3 Grothendieck Spaces and Double Limit Conditions

The relationship of Grothendieck's Theorem with double limit conditions was explored by V. Pták [30]:

Theorem 2.24. *Let X be compact and $A \subseteq C_p(X)$ pointwise bounded. Then A is relatively compact in $C_p(X)$ if and only if the following double limit condition holds: whenever $\{x_n\}_{n < \omega}$ is a sequence in X and $\{f_m\}_{m < \omega}$ is a sequence in A , the double limits $\lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} f_m(x_n)$ and $\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} f_m(x_n)$ are equal whenever they exist.*

Pták's proof involves a combinatorial result on convex means which we omit, since we are only interested in the analogue of Pták's Theorem for *ultralimits* and without compactness assumptions; see Theorem 3.5. We shall frequently deal with ultralimits in what follows.

Definition 2.25. Let X be a topological space. Given an ultrafilter \mathcal{U} on a regular cardinal κ , and a κ -sequence $\{x_\alpha\}_{\alpha < \kappa}$ in X , we say that

$$\lim_{\alpha \rightarrow \mathcal{U}} x_\alpha = x$$

if and only if for every open neighbourhood U of x , $\{\alpha < \kappa : x_\alpha \in U\} \in \mathcal{U}$.

It follows from the fact that an ultrafilter cannot contain disjoint sets that if an ultralimit exists in a Hausdorff space, it is unique. The following result relates ultralimits and compactness:

Proposition 2.26. *A space X is compact if and only if every ultralimit in X exists.*

Proof. Suppose X is compact and assume for contradiction that there is an ultrafilter \mathcal{U} over κ such that $\lim_{\alpha \rightarrow \mathcal{U}} X_\alpha$ does not exist. Then for each $x \in X$ there is an open neighbourhood U_x of x such that $\{\alpha < \kappa : x_\alpha \in U_x\} \notin \mathcal{U}$. Since X is compact, the open cover $\{U_x : x \in X\}$ has a finite subcover $\{U_{x_0}, \dots, U_{x_{n-1}}\}$. Since $\bigcup_{i < n} U_{x_i} = X$, we must have $\{\alpha < \kappa : x_\alpha \in U_{x_i}\} \in \mathcal{U}$ for at least one $i < n$, a contradiction.

Conversely, suppose X is not compact. Let $\{U_\alpha\}_{\alpha < \kappa}$ be an open cover that does not have a finite subcover. Let κ be the minimal cardinal with that property. Then for each cardinal $\lambda < \kappa$, $\{U_\alpha : \alpha < \lambda\}$ is not a cover, so we can pick an $x_\lambda \notin \bigcup\{U_\alpha : \alpha < \lambda\}$. Then each $x \in X$ has a neighbourhood N_x that contains fewer than κ many

x_λ 's. Then $\{\{\alpha : x_\alpha \notin N_x\} : x \in X\}$ forms a filter on κ , which can be extended to an ultrafilter \mathcal{U} . Then \mathcal{U} has no ultralimit, for given any $x \in X$, $\{\alpha : x_\alpha \notin N_x\} \in \mathcal{U}$, so $\{\alpha : x_\alpha \in N_x\} \notin \mathcal{U}$. \square

In the following sections we shall present generalizations of the main results of [7] and [9] by working with Grothendieck spaces instead of with compact spaces. For this purpose we introduce some definitions which are different from the ones in [7] but identical with the added assumption of compactness.

Thorough introductions to continuous logic can be found in [11], [12] and [5]. In [18], the authors introduced in detail a setting for continuous logic, mostly following [7] and [12].

3 Stability, Definability, and Double (Ultra)limit Conditions

It is worth noting that there are different definitions of continuous logics in the literature. However, most authors differ only slightly from one another. We shall now introduce metric structures following Eagle [11] and [12] as this is the context we work in:

Definition 3.1.

- (a) If (M, d) and (N, ρ) are metric spaces and $f: M^n \rightarrow N$ is uniformly continuous, a *modulus of uniform continuity of f* is a function $\delta: (0, 1) \cap \mathbb{Q} \rightarrow (0, 1) \cap \mathbb{Q}$ such that whenever $\mathbf{a} = (a_1, \dots, a_n)$, $\mathbf{b} = (b_1, \dots, b_n) \in M^n$ and $\varepsilon \in (0, 1) \cap \mathbb{Q}$, $\sup\{d(a_i, b_i) : 1 \leq i \leq n\} < \delta(\varepsilon)$ implies $\rho(f(\mathbf{a}), f(\mathbf{b})) < \varepsilon$.
- (b) A *language* for metric structures is a set L which consists of constants, functions with an associated arity, and a modulus of uniform continuity; predicates with an associated arity and a modulus of uniform continuity; and a symbol d for a metric.
- (c) An *L -metric structure* \mathfrak{M} is a metric space (M, d^M) together with interpretations for each symbol in L : $c^{\mathfrak{M}} \in M$ for each constant $c \in L$; $f^{\mathfrak{M}}: M^n \rightarrow M$ a uniformly continuous function for each n -ary function symbol $f \in L$; $P^{\mathfrak{M}}: M^n \rightarrow [0, 1]$ a uniformly continuous function for each n -ary predicate symbol $P \in L$. Assume for the sake of notation simplicity that all metric structures have diameter 1.

- (d) The space of L -structures in a given language L is the family $\text{Str}(L)$ of all L -metric structures endowed with the topology generated by the following basic closed sets: $[\varphi] = \{\mathfrak{M} : \mathfrak{M} \models \varphi\}$ where φ is an L -sentence.

Remark. The continuous first-order formulas are constructed just as in the discrete setting except for the following addition: if $f : [0, 1]^n \rightarrow [0, 1]$ is a continuous function and $\varphi_0, \dots, \varphi_{n-1}$ are L -formulas, then $f(\varphi_0, \dots, \varphi_{n-1})$ is also an L -formula. It is customary to identify formulas φ of arity n with functions $\mathfrak{M}^n \rightarrow [0, 1]$. This allows an easy way to define the satisfaction relation, i.e. if φ is an L -formula, \mathfrak{M} an L -structure, and $a \in \mathfrak{M}^n$, then $\mathfrak{M} \models \varphi(a)$ if and only if $\varphi(a) = 1$. For example, under this identification, the conjunction, $\varphi \wedge \psi$, of two L -formulas φ and ψ is $\max(\varphi, \psi)$. In what follows, we shall introduce other continuous logics and study Gowers' problem in them; our main example is continuous $\mathcal{L}_{\omega_1, \omega}$ as introduced by Eagle [11], [12]. See [11], [12] and [18] for a detailed exposition.

A key technical concept introduced in [7] and used to talk about a pair of Banach space norms is the following:

Definition 3.2.

- (a) Let L be a language $L' \supseteq L$ is a *language for pairs of structures from L* , if L' includes two disjoint copies of L and there is a map $\text{Str}(L) \times \text{Str}(L) \rightarrow \text{Str}(L')$ which assigns to every pair of L -structures $(\mathfrak{M}, \mathfrak{N})$ an L' -structure $\langle \mathfrak{M}, \mathfrak{N} \rangle$.
- (b) Let L' be a language for pairs of structures from L , and X, Y function symbols from L . We say that a formula $\varphi(X, Y)$ is a *formula for pairs of structures from L* if

$$(\mathfrak{M}, \mathfrak{N}) \mapsto \varphi(X^{\mathfrak{M}}, Y^{\mathfrak{N}}) = \text{Val}(\varphi(X, Y), \langle \mathfrak{M}, \mathfrak{N} \rangle)$$

is separately continuous on $\text{Str}(L) \times \text{Str}(L)$. For simplicity, we write $\varphi(\mathfrak{M}, \mathfrak{N})$ instead of $\text{Val}(\varphi(X, Y), \langle \mathfrak{M}, \mathfrak{N} \rangle)$.

In this section we shall only be interested in φ -types when φ is a formula for pairs of structures:

Definition 3.3. Let L be a language, φ an L -formula for pairs of structures, and $\mathfrak{M} \in \text{Str}(L)$. The *left φ -type of \mathfrak{M}* is the function $\text{lt}_{\varphi, \mathfrak{M}} : \text{Str}(L) \rightarrow [0, 1]$ given by $\text{lt}_{\varphi, \mathfrak{M}}(\mathfrak{N}) = \varphi(\mathfrak{M}, \mathfrak{N})$.

The *space of left φ -types*, denoted S_φ^l is the closure of $\{\text{ltp}_{\varphi, \mathfrak{M}} : \mathfrak{M} \in \text{Str}(L)\}$ in $C_p(\text{Str}(L))$. If C is a subset of $\text{Str}(L)$, then $S_\varphi^l(C)$ is the closure of $\{\text{ltp}_{\varphi, \mathfrak{M}}|_C : \mathfrak{M} \in C\}$ in $C_p(C)$ and it is called the *space of left φ -types over C* .

The definitions for *right φ -types* are analogous.

Remark. Notice that the previous definition coincides with the one given in [7] for the compact case: if the logic is compact, that is to say $\text{Str}(L)$ is compact, then S_φ^l turns out to be the closure of $\{\text{ltp}_{\varphi, \mathfrak{M}} : \mathfrak{M} \in \text{Str}(L)\}$ in $[0, 1]^{\text{Str}(L)}$, which is then also compact. Since compact spaces are Grothendieck, we have that S_φ^l is always Fréchet-Urysohn when the logic is compact—see Proposition 2.12.

The notion of model-theoretic stability has been extensively studied in discrete model theory, whereas works like [7], [9] or [18] are devoted to understanding stability in the continuous setting. Among the multiple equivalent definitions of stability, we have found the one introduced in [21] involving a *double limit condition* is the most suitable for the continuous setting, even when it comes to non-compact logics:

Definition 3.4. Let L be a language for pairs of structures, φ a formula for pairs of structures, and $C \subseteq \text{Str}(L)$. We say that φ is *stable on C* if and only if whenever $\{\mathfrak{M}_i\}_{i < \omega}$ and $\{\mathfrak{N}_j\}_{j < \omega}$ are sequences in C , and \mathcal{U} and \mathcal{V} are ultrafilters on ω , we have

$$\lim_{i \rightarrow \mathcal{U}} \lim_{j \rightarrow \mathcal{V}} \varphi(\mathfrak{M}_i, \mathfrak{N}_j) = \lim_{j \rightarrow \mathcal{V}} \lim_{i \rightarrow \mathcal{U}} \varphi(\mathfrak{M}_i, \mathfrak{N}_j).$$

Remark. Notice that $\varphi(\mathfrak{M}_i, \mathfrak{N}_j)$ is always a number in $[0, 1]$ and so the ultralimits always exist by Proposition 2.26.

We now state the topological version of one of our main results, which varies the one that appears in [18] by replacing “weakly Grothendieck” by “Grothendieck” and “first countable” by “countably tight”:

Theorem 3.5. *Let X be Grothendieck and countably tight. A subset A of $C_p(X, [0, 1])$ is relatively compact in $C_p(X, [0, 1])$ if and only if it satisfies the following double limit condition: for every pair of sequences $\{f_n\}_{n < \omega} \subseteq A$ and $\{x_m\}_{m < \omega} \subseteq X$, and ultrafilters \mathcal{U} and \mathcal{V} on ω , the double limits*

$$\lim_{n \rightarrow \mathcal{U}} \lim_{m \rightarrow \mathcal{V}} f_n(x_m) = \lim_{m \rightarrow \mathcal{V}} \lim_{n \rightarrow \mathcal{U}} f_n(x_m)$$

agree whenever $\lim_{m \rightarrow \mathcal{V}} x_m$ exists.

Proof. Let \overline{A} denote the closure of A in $[0, 1]^X$. Suppose $\overline{A} \cap C_p(X)$ is not compact in $C_p(X)$. Then $\overline{A} \cap C_p(X)$ is closed but it is not countably compact in $C_p(X)$, since X is Grothendieck. Let $\{f_n\}_{n < \omega}$ be a sequence in A with no limit points in $\overline{A} \cap C_p(X)$. Since \overline{A} is a compact subset of $[0, 1]^X$, each ultralimit of the sequence exists, is a limit point, and is therefore discontinuous. Take a non-principal ultrafilter \mathcal{U} over ω and let $\lim_{n \rightarrow \mathcal{U}} f_n = g$, where g is discontinuous by assumption. Then there are $\varepsilon > 0$ and $y \in X$ such that $y \in \overline{Y}$, where

$$Y = X \setminus g^{-1}(g(y) - \varepsilon, g(y) + \varepsilon).$$

Since $t(X) = \aleph_0$, there is some $Z \subseteq Y$ with $|Z| = \aleph_0$ and $y \in \overline{Z}$. Suppose $Z = \{z_n\}_{n < \omega}$ and for each open neighbourhood U of y , let $M_U = \{n < \omega : z_n \in U\}$. Clearly, the family of all M_U is centred and so it can be extended to an ultrafilter \mathcal{V} on ω so that

$$\lim_{n \rightarrow \mathcal{U}} \lim_{m \rightarrow \mathcal{V}} f_n(x_m) = g(y)$$

since each f_n is continuous. On the other hand,

$$\lim_{m \rightarrow \mathcal{V}} \lim_{n \rightarrow \mathcal{U}} f_n(x_m) = \lim_{m \rightarrow \mathcal{V}} g(x_m)$$

exists by compactness of $[0, 1]$. However, by the choice of each x_m , we have $|g(y) - g(x_m)| > \varepsilon$ and so the ultralimits exist but are different, a contradiction.

Conversely, suppose $\overline{A} \cap C_p(X)$ is compact and $\lim_{m \rightarrow \mathcal{V}} x_m = y$. Then for any sequence $\{f_n\}_{n < \omega} \subseteq A$ and ultrafilter \mathcal{U} on ω , there is a continuous function $g = \lim_{n \rightarrow \mathcal{U}} f_n$. Thus,

$$\lim_{n \rightarrow \mathcal{U}} \lim_{m \rightarrow \mathcal{V}} f_n(x_m) = \lim_{n \rightarrow \mathcal{U}} f_n(y)g(y),$$

and

$$\lim_{m \rightarrow \mathcal{V}} \lim_{n \rightarrow \mathcal{U}} f_n(x_m) = \lim_{m \rightarrow \mathcal{V}} g(x_m) = g(y).$$

□

The main application we consider here deals with countable fragments of $\mathcal{L}_{\omega_1, \omega}$, for the corresponding space of types is Polish [25], so Grothendieck and countably tight.

The model-theoretic version of Theorem 3.5 will relate definability to a double limit condition just as in the discrete compact case. For this purpose, we introduce the notion of definability for φ -types, where φ is a formula for pairs of structures, following [7]:

Definition 3.6. Suppose L is a language for pairs of structures and φ is a formula for pairs of structures. Let $C \subseteq \text{Str}(L)$. A function $\tau : S_\varphi^r(C) \rightarrow [0, 1]$ is a *left global φ -type over C* if there is a sequence $\{\mathfrak{M}_\alpha\}_{\alpha < \kappa} \subseteq C$, and an ultrafilter \mathcal{U} on κ , such that for every type $t \in S_\varphi^r(C)$, say $t = \lim_{\beta \rightarrow \mathcal{V}} \text{rtp}_{\varphi, \mathfrak{M}_\beta}$, we have

$$\tau(t) = \lim_{\alpha \rightarrow \mathcal{U}} \lim_{\beta \rightarrow \mathcal{V}} \varphi(\mathfrak{M}_\alpha, \mathfrak{N}_\beta).$$

We say that τ is *explicitly definable* if it is continuous. If a left φ -type is given by $p = \lim_{\alpha \rightarrow \mathcal{U}} \text{ltp}_{\varphi, \mathfrak{M}_\alpha}$, we say that p is *explicitly definable* if the respective τ is continuous.

The previous definition is sound since in continuous logics the allowable formulas are continuous functions. Moreover, as with most concepts in continuous logic, such a definition reduces to the usual notion in the discrete setting. Also notice that, although our main application deals with Banach spaces, the context of the previous definition and the results in this section is that of the languages for pairs of structures rather than just Banach spaces. We proceed with the model-theoretic version of Theorem 3.5:

Theorem 3.7. *Let L be a language for pairs of structures, and φ a formula for pairs of structures. Suppose $C \subseteq \text{Str}(L)$ is such that $S_\varphi^l(C)$ is Grothendieck and countably tight. Then the following are equivalent:*

- (i) *Whenever a left type over C is given by $t = \lim_{i \rightarrow \mathcal{U}} \text{ltp}_{\varphi, \mathfrak{M}_i}$, and $\{\mathfrak{N}_j\}_{j < \omega} \subseteq C$ is a sequence in C , and \mathcal{V} is an ultrafilter on ω , then*

$$\lim_{i \rightarrow \mathcal{U}} \lim_{j \rightarrow \mathcal{V}} \varphi(\mathfrak{M}_i, \mathfrak{N}_j) = \lim_{j \rightarrow \mathcal{V}} \lim_{i \rightarrow \mathcal{U}} \varphi(\mathfrak{M}_i, \mathfrak{N}_j).$$

- (ii) *Whenever a left type over C is given by $t = \lim_{i \rightarrow \mathcal{U}} \text{ltp}_{\varphi, \mathfrak{M}_i}$, and $\{\mathfrak{N}_j\}_{j < \omega} \subseteq C$ is a sequence in C , then*

$$\sup_{i < j} \varphi(\mathfrak{M}_i, \mathfrak{N}_j) = \inf_{j < i} \varphi(\mathfrak{M}_i, \mathfrak{N}_j).$$

- (iii) *If τ is a left φ -type over C , then τ is explicitly definable.*

Proof. For a proof of the equivalence of (i) and (ii), see [21] or [18]. To see that (i) implies (iii), suppose $\langle \mathfrak{N}_\alpha : \alpha < \kappa \rangle$ is a κ -sequence in C such that

$$\tau(\lim_{i \rightarrow \mathcal{U}} \text{ltp}_{\varphi, \mathfrak{M}_i}) = \lim_{i \rightarrow \mathcal{U}} \lim_{\alpha \rightarrow \mathcal{V}} \varphi(\mathfrak{M}_i, \mathfrak{N}_\alpha)$$

defines a global φ -type, where \mathcal{V} is an ultrafilter on κ . Define $f_\alpha : S_\varphi^l(C) \rightarrow [0, 1]$ by

$$f_\alpha(\lim_{i \rightarrow \mathcal{U}} \text{ltp}_{\varphi, \mathfrak{M}_i}) = \lim_{i \rightarrow \mathcal{U}} \varphi(\mathfrak{M}_i, \mathfrak{N}_\alpha).$$

Then the set $A = \{f_\alpha : \alpha < \kappa\}$ is a subset $A \subseteq C_p(S_\varphi^l(C), [0, 1])$ and satisfies the double limit condition, so that $\overline{A} \cap C_p(S_\varphi^l(C), [0, 1])$ is compact and $\tau = \lim_{\alpha \rightarrow \mathcal{V}} f_\alpha$ is continuous. That (iii) implies (i) is immediate. \square

One often deals with the space of φ -types over a subspace C of $\text{Str}(L)$, $S_\varphi^l(C)$, which by Lemma 2.2 is a continuous image of S_φ^l . An important application of the C_p -theoretic machinery we have developed deals with the preservation of the Grothendieck property from S_φ^l to $S_\varphi^l(C)$ for any $C \subseteq \text{Str}(L)$:

Theorem 3.8. *Let L be a language for pairs of structures, and φ a formula for pairs of structures. If S_φ^l is Grothendieck and $C \subseteq \text{Str}(L)$, then $S_\varphi^l(C)$ is Grothendieck.*

Proof. Let $A = \{\text{ltp}_{\varphi, \mathfrak{M}} : \mathfrak{M} \in C\} \subseteq S_\varphi^l$. By Lemma 2.2, the restriction map $\pi_C : C_p(\text{Str}(L)) \rightarrow C_p(C)$ is continuous. Notice that $\pi_C(A)$ is Grothendieck by Theorem 2.16. By definition, $S_\varphi^l(C) = \overline{\pi_C(A)}$ and so it is Grothendieck by Theorem 2.15. \square

The following result is routine:

Theorem 3.9. *The continuous open image of a Fréchet-Urysohn space is Fréchet-Urysohn.*

Proof. Let X be Fréchet-Urysohn, and $f : X \rightarrow Y$ a continuous open surjection. Let $B \subseteq Y$ and take $y \in \overline{B}$. By continuity it is always true that $f^{-1}(\overline{B}) \supseteq \overline{f^{-1}(B)}$. To prove the reverse inclusion, take $x \in f^{-1}(\overline{B})$ and an open neighbourhood U of x . Since f is open, $f(U)$ is an open neighbourhood of $f(x) \in \overline{B}$ and so $f(U) \cap B \neq \emptyset$. Then $U \cap f^{-1}(B) \neq \emptyset$, and so $x \in \overline{f^{-1}(B)}$.

Next, suppose $f(x_0) = y$. Since X is Fréchet-Urysohn and $x_0 \in \overline{f^{-1}(B)}$, we can take a sequence in $f^{-1}(B)$ converging to x , say $\{z_n\}_{n < \omega}$. By continuity, $\lim_{n \rightarrow \infty} f(z_n) = f(x_0) = y$, where $f(z_n) \in B$ for each $n < \omega$. \square

Using Theorem 3.8, we get:

Corollary 3.10. *Let L be a language for pairs of structures, and φ a formula for pairs of structures. If S_φ^l is Fréchet-Urysohn and $C \subseteq \text{Str}(L)$ is closed, then $S_\varphi^l(C)$ includes a dense Fréchet-Urysohn subspace.*

Proof. Notice that $S_\varphi^l(C)$ is the closure of the image under the restriction map pi_C of S_φ^l . By Lemma 2.2, π_C is continuous and open. We conclude by Theorem 3.8 \square

Remark. The importance of compactness in Model Theory is well understood: the pioneers of the discipline built around this topological concept some of the fundamental pillars. Even though some famous open problems on compactness remain open, topologists have reached a deep understanding of the subject, and model theorists have also extensively explored the applications of compactness in first-order logic and its variations. In the last decades, the work of S. Shelah has brought to the foreground impressive model-theoretic results that rely on Combinatorics and Set Theory, making the use of these resources the fashionable approach in today's Model Theory.

Reasonable changes to first-order logic make the statement of the Compactness Theorem false — consider for instance $L_{\omega_1, \omega}$. This does not completely disable the combinatorial tools but definitely sets important limitations. Recent work [13] has shown that the intersection of Topology and Model Theory goes way beyond compactness. As opposed to the relation between compactness and first-order Model Theory, the connections between Topology and Model Theory beyond the Compactness Theorem have barely been explored. It is an interesting question whether it is reasonable to hope for a dictionary that translates model-theoretic terms into topological ones.

The authors have already shown how more advanced Topology can shine a light on modern model-theoretic results and questions on *metastability* and have shown how to convert topological spaces into abstract logics [13]. As mentioned before, this present work was inspired by the work of P. Casazza and J. Iovino [7] because of their use of Grothendieck's theorem in the context of compactness. Recently, E. Dueñez joined Casazza and Iovino, and [7] was reworked into [9].

There are clear advantages in the new approach of [9] but they come with a price tag for those who are interested in the underlying topology. Even though the results are essentially the same, [9] resorts to more advanced combinatorial results involving metastability

to yield their main proofs without the need of talking about languages for pairs of structures, which instead were omnipresent in [7]. In [9], the topology that we discuss here and that we use to produce generalizations hides away behind these combinatorial results which ultimately rely on compactness, limiting the applicability of their methods to other, non-compact logics.

The issue at hand is the following: given a language L , an L -formula $\varphi(x, y)$, an L -structure \mathcal{M} and a, b in the universe of \mathcal{M} , it is customary to associate $\varphi^{\mathcal{M}}(a, b)$ to be the truth value of $\varphi(a, b)$ in \mathcal{M} . The formulas are evaluated in elements of \mathcal{M} where usually \mathcal{M} carries little to no topology. This is the approach of [9]. On the other hand, working with a language for pairs of structures L turns the relevant formulas into continuous functions from $\text{Str}(L) \times \text{Str}(L)$ into $[0, 1]$. $\text{Str}(L)$, just as the space of types that appears further below, can carry quite interesting topologies; this is what we work with and what the original approach from [7] used. The final goal is to be able to combine two normed spaces into an enlarged structure with two norms where we can compare them — this is Definition 3.2.

Notice that an approach in which we start with a language with two function symbols $\|\cdot\|_1, \|\cdot\|_2$ for norms also reduces the maneuverability that the language for pairs of structures offer. One could certainly obtain continuous functions from $\text{Str}(L)$ to $[0, 1]$ by assigning to \mathcal{M} the truth value of $\varphi^{\mathcal{M}}(\|\cdot\|_1^{\mathcal{M}}, \|\cdot\|_2^{\mathcal{M}})$. This approach is an oversimplification since the applicability of the double limit condition becomes unclear once you work with only one variable. There is a similarity here with the use of bipartite formulas in classical model theory: $\varphi(\bar{x}, \bar{y})$, where \bar{x}, \bar{y} are tuples, instead of simply working with $\varphi(\bar{z})$ where \bar{z} is a longer tuple.

The authors realize that languages for pairs of structures can be cumbersome. However, they are a key component in the topological presentation of [7] and the corresponding generalization that appears in this work. As a result, we have kept the original approach of [7] here. This approach has its own limitations as it makes it more difficult to use similar techniques to talk about other types of languages, formulas or theories.

4 Applications and Examples Concerning the Undefinability of Pathological Banach Spaces

We first introduce some concepts from Analysis which we will deal with in what follows. We denote the Banach space of sequences of real numbers that are eventually 0 by c_{00} . The space of sequences of real numbers converging to 0 is denoted by c_0 . Finally, ℓ^p denotes the space of sequences $\{x_n\}_{n < \omega}$ of real numbers such that $\sum_{n < \omega} |x_n|^p < \infty$.

B. Tsirelson [40] constructed a Banach space which does not include a copy of any ℓ^p or c_0 . What is now called *Tsirelson's space* is due to T. Figiel and W. Johnson [15]: let $\{t_n\}_{n < \omega}$ be the canonical basis for c_{00} . If $x = \sum_{n < \omega} a_n x_n \in c_{00}$ and $E, F \in [\omega]^{<\omega}$, we denote $\sum_{n \in E} a_n x_n$ by Ex , and write $E \leq F$ if and only if $\max E \leq \min F$.

Recursively define

$$\|x\|_0 = \|x\|_{c_{00}},$$

$$\|x\|_{n+1} = \max \left\{ \|x\|_n, \frac{1}{2} \max_k \left\{ \sum_{i < k} \|E_i\|_n : \{k\} \leq E_1 < \dots < E_k \right\} \right\}.$$

Then,

$$\|x\|_n \leq \|x\|_{n+1} \leq \|x\|_{\ell^1}$$

and we denote $\|x\|_T = \lim_{n \rightarrow \infty} \|x\|_n$. Then,

$$\|x\|_T = \max \left\{ \|x\|_{c_{00}}, \frac{1}{2} \max \left\{ \sum_{i < k} \|E_i x\|_T : \{k\} \leq E_1 < \dots < E_k \right\} \right\}$$

and T is the norm completion of $\|\cdot\|_T$. The reader is referred to P. Casazza and T. Shura [8] for a detailed exposition of Tsirelson's space.

The context of the subsequent results is what is now known as *Gowers' problem*: does every infinite dimensional explicitly definable Banach space include an isomorphic copy of some ℓ^p or c_0 ? We found in [7] the inspiration for the framework and results presented in the current paper. The topological assumption used in [7] is countable compactness whereas ours is Grothendieck plus countable tightness.

There are mistakes in [7] (which we avoid here) which are corrected in the rewrite [9], which lead the authors to now only assert their results for compact logics. With the replacement of countable

compactness by compactness, [7] can be corrected, but with this assumption, the approach of [9] is more elegant.

Countable compactness does imply Grothendieck, but not countable tightness. In fact, even compactness does not imply countable tightness. In terms of results, we can trivially deduce their main definability results from ours because finitary continuous logic is a countable fragment (see below) of infinitary continuous logic. However the more general definability results for compact logics are obtained using a double limit condition on sequences obtained by Pták’s combinatorial approach.

To obtain our results, we needed to instead use a double limit condition involving ultralimits and assume countable tightness. In work in progress, we obtain a common generalization of “compactness” and “Grothendieck plus countable tightness” that yields the desired undefinability results.

Fix a language L for a pair of structures. Let C be the set of all L -structures which are normed spaces based on c_{00} (see definition below). We introduce the continuous logic formula for pairs of structures used in [7]: for norms $\|\cdot\|_1$ and $\|\cdot\|_2$, define

$$D(\|\cdot\|_1, \|\cdot\|_2) = \sup \left\{ \frac{\|x\|_1}{\|x\|_2} : \|x\|_{\ell^1} = 1 \right\}.$$

Then define

$$\varphi(\|\cdot\|_1, \|\cdot\|_2) = 1 - \frac{\log D(\|\cdot\|_1, \|\cdot\|_2)}{1 + \log D(\|\cdot\|_1, \|\cdot\|_2)}. \quad (\star)$$

It is not difficult to see that if $t = \lim_{i \rightarrow \mathcal{U}} \text{ltp}_{\varphi, \|\cdot\|_i}$ is realized by a structure $(\mathfrak{M}, \|\cdot\|_*)$, then

$$\lim_{i \rightarrow \mathcal{U}} \sup_{\|x\|_{\ell^1} = 1} \frac{\|x\|_i}{\|x\|} = \sup_{\|x\|_{\ell^1} = 1} \frac{\|x\|_*}{\|x\|} \quad (1)$$

for any structure $(c_{00}, \|x\|) \in C$. In particular, taking $\|\cdot\|_{\ell^1} = \|\cdot\|$ yields $\lim_{i \rightarrow \mathcal{U}} \|x\|_i = \|x\|_*$ for each $x \in c_{00}$. If in addition

$$\|\cdot\|_1 \leq \|\cdot\|_2 \leq \cdots \leq \|\cdot\|_n \leq \cdots$$

then $\lim_{i \rightarrow \infty} \|x\|_i = \|x\|_*$.

Conversely, if

$$\|\cdot\|_1 \leq \|\cdot\|_2 \leq \cdots \leq \|\cdot\|_n \leq \cdots$$

and also $\lim_{i \rightarrow \infty} \|x\|_i = \|x\|_*$ for every $x \in c_{00}$, then (1) holds for any $(c_{00}, \|\cdot\|) \in \mathcal{C}$ and any nonprincipal ultrafilter \mathcal{U} over ω .

This motivates the following definition from [7]:

Definition 4.1. Let \mathcal{C} be the set of all structures which are normed spaces based on c_{00} , and φ a formula for a pair of structures, and $\|\cdot\|_*$ a norm on c_{00} .

- (a) If $\{\|\cdot\|_i : i < \omega\}$ is a family of norms on c_{00} we say that $\{\text{tp}_{\varphi, \|\cdot\|_i} : i < \omega\}$ determines $\|\cdot\|_*$ uniquely if, for every $\mathcal{U} \in \beta\omega$, the type $t = \lim_{i \rightarrow \mathcal{U}} \text{tp}_{\varphi, \|\cdot\|_i}$ is realized, and $\|\cdot\|_*$ is its unique realization.
- (b) We say that $\|\cdot\|_*$ is uniquely determined by its φ -type over \mathcal{C} if there is a family of norms $\{\|\cdot\|_i : i < \omega\}$ on c_{00} such that $\{\text{tp}_{\varphi, \|\cdot\|_i} : i < \omega\}$ determines $\|\cdot\|_*$ uniquely.

Notice that if $\|\cdot\|_i$ denotes the i -th iterate of the Tsirelson norm, and $\|\cdot\|_T$ the Tsirelson norm, then $\lim_{i \rightarrow \infty} \|x\|_i = \|x\|_T$ for each $x \in c_{00}$, and so we have the following result from [7] that follows from the previous definition:

Proposition 4.2. Let L be a language for pairs of structures, \mathcal{C} the class of structures $(c_{00}, \|\cdot\|_{\ell^1}, \|\cdot\|)$ such that the norm completion of $(c_{00}, \|\cdot\|)$ is a Banach space including ℓ^p or c_{00} , and let $\varphi(X, Y)$ be the formula defined by (\star) above. Suppose $\{(c_{00}, \|\cdot\|_{\ell^1}, \|\cdot\|_i) : i < \omega\}$ is a family of structures in \mathcal{C} such that

$$\|\cdot\|_1 \leq \|\cdot\|_2 \leq \dots \leq \|\cdot\|_n \leq \dots$$

and the φ -type $t = \lim_{i \rightarrow \mathcal{U}} \text{tp}_{\varphi, \|\cdot\|_i}$ is realized by $(c_{00}, \|\cdot\|_{\ell^1}, \|\cdot\|_*)$ in $\text{Str}(L)$, then $\{\text{tp}_{\varphi, \|\cdot\|_i} : i < \omega\}$ uniquely determines $\|\cdot\|_*$ over \mathcal{C} . In particular, the Tsirelson norm is uniquely determined by its φ -type over \mathcal{C} .

The following result is a corollary in [7] and is purely analytical. The core analytical result is stated identically in [9].

Proposition 4.3. Let $\|\cdot\|_i$ be the i -th iterate in the construction of the Tsirelson norm. Then the following hold:

- (i) $\sup_{\|x\|_{\ell^1}=1} \frac{\|x\|_i}{\|x\|_j} \leq 1$ for $i < j$.
- (ii) $\sup_{\|x\|_{\ell^1}=1} \frac{\|x\|_i}{\|x\|_j} \geq j$ for $i > j$.

Thus, $\sup_{i < j} \varphi(\|\cdot\|_i, \|\cdot\|_j) \neq \inf_{j < i} \varphi(\|\cdot\|_i, \|\cdot\|_j)$.

Proposition 4.3 and Theorem 3.7 yield:

Theorem 4.4. *Let L be a language of pairs of structures, and \mathcal{C} the class of structures $(c_{00}, \|\cdot\|_{\ell^1}, \|\cdot\|)$ such that the norm completion of $(c_{00}, \|\cdot\|)$ is a Banach space including ℓ^p or c_0 , and let $\|\cdot\|_T$ be the Tsirelson norm. Let φ be the formula as in (\star) above. If the space $S_\varphi^l(\mathcal{C})$ of φ -types over \mathcal{C} is Grothendieck and countably tight, then $\|\cdot\|_T$ is uniquely determined by its φ -type over \mathcal{C} and that φ -type is not explicitly definable over \mathcal{C} .*

Remark. Notice that here we are dealing with languages for pairs of structures in which only first-order variables are considered. In such a context, definitions like Definition 4.1 are sound. An interesting logic with countably many formulas is second-order logic. Although it seems as if the topology is confined to subspaces of 2^ω , the model theory is not very well developed, even though Chang and Keisler [10] have the development of the model theory of second-order logic as one of the problems in their classic book. However, see [20]. It is unclear to the authors whether the results and even the definitions presented here apply to second-order logic, which has been called “set theory in sheep’s clothing” [31]. In particular, it would be interesting to know about the definability of Tsirelson space in second-order arithmetic, which is often conflated with Analysis.

We now prove a positive definability result which generalizes the one that appears in [7]. The proof is a variation of the original, in which their use of the Stone-Weierstrass Theorem is replaced by working with the double (ultra)limit condition. This is important because the Stone-Weierstrass Theorem is not nicely extendable beyond compact spaces.

Definition 4.5. A structure $(c_{00}, \|\cdot\|_{\ell^1}, \|\cdot\|, e_0, e_1, \dots)$ where $\|\cdot\|$ is an arbitrary norm and $\{e_n\}_{n < \omega}$ is the standard vector basis of c_{00} is called a *structure based on c_{00}* .

Theorem 4.6. *Let L be a language for pairs of structures, φ the formula defined by (\star) above, and \mathcal{C} a subclass of the class of structures based on c_{00} such that every closed subspace of a space in $\overline{\mathcal{C}}$ includes a copy of c_0 or ℓ^p . Assume that the space of φ -types over \mathcal{C} is Grothendieck and countably tight. If the φ -type of \mathfrak{M} is explicitly definable from \mathcal{C} , then \mathfrak{M} includes a copy of c_0 or ℓ^p .*

Proof. Suppose that the global φ -type τ of \mathfrak{M} is *approximated* by $\{\mathfrak{M}_i\}_{i < \omega} \subseteq C$ and $\mathcal{U} \in \beta\omega$, that is

$$\tau(\lim_{j \rightarrow \mathcal{V}} \text{rtp}_{j, \mathfrak{N}_j}) = \lim_{i \rightarrow \mathcal{U}} \lim_{j \rightarrow \mathcal{V}} \varphi(\mathfrak{M}_i, \mathfrak{N}_j).$$

We show that there is an $\mathfrak{N} \in \overline{C}$ such that $\varphi(\mathfrak{M}, \mathfrak{N}) > 0$. Otherwise, $\varphi(\mathfrak{M}, \mathfrak{N}) = 0$ for each $\mathfrak{N} \in \overline{C}$. Since τ is explicitly definable, we exchange the order of the ultralimits by taking $\tau(\lim_{j \rightarrow \mathcal{V}} \text{rtp}_{j, \mathfrak{N}_j}) = \lim_{j \rightarrow \mathcal{V}} \tau(\text{rtp}_{j, \mathfrak{N}_j})$ and then obtain

$$\lim_{i \rightarrow \mathcal{U}} \lim_{j \rightarrow \mathcal{V}} \varphi(\mathfrak{M}_i, \mathfrak{N}_j) = \lim_{j \rightarrow \mathcal{V}} \lim_{i \rightarrow \mathcal{U}} \varphi(\mathfrak{M}_i, \mathfrak{N}_j) = \lim_{j \rightarrow \mathcal{V}} \varphi(\mathfrak{M}, \mathfrak{N}_j) = 0.$$

But then consider $\varphi(\mathfrak{M}_i, \mathfrak{M}_j)$ and, by the symmetry of $D(\mathfrak{M}, \mathfrak{N})$, obtain $\varphi(\mathfrak{M}, \mathfrak{M}) = 0$, a contradiction. It is not hard to see that $\varphi(\mathfrak{M}, \mathfrak{N}) > 0$ implies that $D(\mathfrak{M}, \mathfrak{N}) < \infty$, which implies that $e_n^{\mathfrak{M}} \mapsto e_n^{\mathfrak{N}}$ determines an isomorphism (it is a linear bijection between the bases). This concludes the proof since \mathfrak{N} includes a copy of c_0 or ℓ^p . \square

Remark. Notice that regardless of the results proved above, a space can have a copy of c_0 or ℓ^p while failing to be explicitly definable.

Let L be a language for metric structures. The $\mathcal{L}_{\omega_1, \omega}(L)$ -formulas are defined recursively as follows:

- (a) All first-order L -formulas are $\mathcal{L}_{\omega_1, \omega}(L)$ -formulas.
- (b) If $\varphi_1, \dots, \varphi_n$ are $\mathcal{L}_{\omega_1, \omega}(L)$ -formulas, and $g : [0, 1]^n \rightarrow [0, 1]$ is continuous, then $g(\varphi_1, \dots, \varphi_n)$ is an $\mathcal{L}_{\omega_1, \omega}(L)$ -formula.
- (c) If $\{\varphi_n : n < \omega\}$ is a family of $\mathcal{L}_{\omega_1, \omega}(L)$ -formulas, then $\inf_n \varphi_n$ and $\sup_n \varphi_n$ are $\mathcal{L}_{\omega_1, \omega}(L)$ -formulas. These can also be denoted by $\bigwedge_n \varphi_n$ and $\bigvee_n \varphi_n$ respectively.
- (d) If φ is an $\mathcal{L}_{\omega_1, \omega}(L)$ -formula and x is a variable then $\inf_x \varphi$ and $\sup_x \varphi$ are $\mathcal{L}_{\omega_1, \omega}(L)$ -formulas.

An interesting feature of continuous $\mathcal{L}_{\omega_1, \omega}$, noted in [12], is that negation (\neg) becomes available in the classical sense: if L is a language and φ is an $\mathcal{L}_{\omega_1, \omega}(L)$ -formula, then we can define $\psi(x) = \bigvee_n \{\varphi(x) + \frac{1}{n}, 1\}$. Then $\mathfrak{M} \models \psi(a)$ if and only if there is an $n < \omega$ such that $\mathfrak{M} \models \max\{\varphi(a) + \frac{1}{n}, 1\}$; that is, $\max\{\varphi(a) + \frac{1}{n}, 1\} = 1$, which is the same as $\varphi(a) \leq 1 - \frac{1}{n}$, i.e. $\mathfrak{M} \not\models \varphi(a)$. Thus, $\mathfrak{M} \models \psi(x)$ if and only if $\mathfrak{M} \not\models \varphi(x)$, and so ψ corresponds to $\neg\varphi$.

It is sometimes useful to restrict one's attention to (countable) fragments of $\mathcal{L}_{\omega_1, \omega}(L)$, which are easier to work with than the full logic.

Let L be a language. A *fragment* \mathcal{F} of $\mathcal{L}_{\omega_1, \omega}(L)$ is a set of $\mathcal{L}_{\omega_1, \omega}(L)$ -formulas satisfying:

- (a) Every first-order formula in \mathcal{F} .
- (b) \mathcal{F} is closed under finitary conjunctions and disjunctions.
- (c) \mathcal{F} is closed under \inf_x and \sup_x .
- (d) \mathcal{F} is closed under subformulas.
- (e) \mathcal{F} is closed under substituting terms for free variables.

Notice that every subset of $\mathcal{L}_{\omega_1, \omega}(L)$ generates a fragment, and that every finite set of formulas generates a countable fragment. There are two arguments to support the idea of working with countable fragments of $\mathcal{L}_{\omega_1, \omega}(L)$. First, notice that a given proof involves only finitely many formulas which then generate a countable fragment of $\mathcal{L}_{\omega_1, \omega}(L)$. Notice that even if we adjoined to our proof system some sort of ω -rule, we would not have more than countably many formulas involved in a given proof. These formulas would still generate a countable fragment. Second (as noted independently by C. Eagle in a personal communication with the authors), if an object is definable in $\mathcal{L}_{\omega_1, \omega}(L)$, then it is definable in a countable fragment by the same argument. This remark does not hold for $\mathcal{L}_{\omega_1, \omega_1}$ because of (e) above.

As the work of Casazza and Iovino deals with continuous logics which are finitary in nature, it was natural to ask whether their result on the undefinability of Tsirelson's space could be proved for continuous $\mathcal{L}_{\omega_1, \omega}$, which is arguably a more natural language from the point of view of Banach space theorists.

In discrete model theory, countable fragments of $\mathcal{L}_{\omega_1, \omega}$ have been studied previously; for instance, as we mentioned previously, M. Morley [25] showed that the space of types of a countable fragment of $\mathcal{L}_{\omega_1, \omega}$ is Polish.

Remark. In the continuous case, we note that the space of types of a countable fragment \mathcal{F} of continuous $\mathcal{L}_{\omega_1, \omega}(L)$ can be seen as a subspace of $[0, 1]^{\mathcal{F}}$ which is metrizable and second countable, and so it is separable and first countable. Hence it is Grothendieck and countably tight and thus our results apply.

5 The NIP and the Bourgain-Fremlin-Talagrand Dichotomy

A series of papers by P. Simon [33], K. Khanaki [22, 23] and Khanaki and A. Pillay [24] has explored connections between the *Non Independence Property* (NIP) and the famous dichotomy of [6]:

Theorem 5.1. *Let X be a Polish space, and let $\{f_n\}_{n < \omega}$ be a sequence in $C_p(X)$ with an accumulation point $f \in \mathbb{R}^X$. Then one of the following holds:*

- (i) $\{f_n\}_{n < \omega}$ has a subsequence converging to f .
- (ii) The closure $\overline{\{f_n\}_{n < \omega}}$ in \mathbb{R}^X includes a copy of $\beta\omega$.

The informal conclusion is that such a sequence is either “good” (has a convergent subsequence) or is very complicated (its closure has cardinality $2^{2^{\aleph_0}}$ —the number of ultrafilters on ω). Although this dichotomy is widely attributed to [6], given that their work contains all the ideas for it, this more modern statement and a proof of it can be found in [39]. The model-theoretic translation of this result is presented in detail in [34].

At the suggestion of the editor, we have started looking at applications of C_p -theory to NIP. As opposed to the double limit conditions this paper deals with, the dichotomy above is not so easily generalized from the topological point of view. However, Haim Horowitz and the first author did realize that the dichotomy works for analytic spaces, i.e. continuous images of the Baire space ω^ω and, more generally, that if the dichotomy holds for a space X , then it holds for all continuous images of X . The proof goes as follows: suppose $f : X \rightarrow Y$ is continuous and that the dichotomy holds for X . Since the dichotomy is really a property of \mathbb{R}^X and by Lemma 2.3, \mathbb{R}^Y is homeomorphic to a subspace of \mathbb{R}^X through the dual map Φ_f , which also preserves continuity, then Y has the dichotomy.

Todorćević and Horowitz [19] have shown that there is no hope for it to hold for Lindelöf p -spaces as it fails for separable metrizable spaces: any maximal almost disjoint (MAD) family is a counterexample.

Definition 5.2. An *almost disjoint* family of subsets of ω is one in which any two members have finite intersection. The family is *MAD* if it is maximal, i.e. there is no bigger such family.

It is not hard to show there is such a MAD family, since by Zorn's Lemma, any almost disjoint family can be extended to a maximal one. Moreover, if $V = L$, a co-analytic MAD family can be found [19] to serve as a counterexample. Topologists like to consider almost disjoint families of rationals obtained by considering sequences of rationals converging to different irrationals.

Theorem 5.3 ([19]). *Let A be a MAD family, considered as a subspace of the Cantor set. Then the BFT dichotomy fails for A .*

Sketch of Proof. Consider the following family in $C_p(\mathcal{A})$: for each $n < \omega$, let $f_n : \mathcal{A} \rightarrow 2$ be defined by $f_n(x) = 1$ if and only if $n \in x$. Then the zero function $\mathbf{0}$ is in the closure $\overline{\{f_n\}_{n < \omega}}$. By MADness, there is no converging subsequence. One can also show that the closure $\overline{\{f_n\}_{n < \omega}}$ consists precisely of the f_n 's, the function $\mathbf{0}$, and the characteristic functions of each member of \mathcal{A} , i.e. the functions defined by $f_y(x) = 1$ if and only if $x = y$ for each $y \in \mathcal{A}$. \square

Angelic spaces were introduced by Fremlin and appeared in Pryce [29]. They are defined to be spaces X such that for every subset A of X , if A is countably compact in X , then A is relatively compact and Fréchet-Urysohn. Then we can restate Proposition 2.12 as: a space is a hereditary g -space if and only if it is angelic. In their seminal paper, Bourgain, Fremlin, and Talagrand proved that if X is Polish, then the first Baire class $B_1(X)$ (the class of pointwise limits of continuous real-valued functions on X) is angelic with the subspace topology inherited from the product topology on \mathbb{R}^X . In his lecture at the Moscow conference celebrating the 80th birthday of Prof. A. V. Arhangel'skiĭ, the second author suggested that topologists investigate $B_1(X)$ with the same vigour with which they have studied $C_p(X)$. This will likely provide new insights into NIP. We plan to do this ourselves.

Bourgain, Fremlin, and Talagrand prove their results for K -analytic spaces, which, as mentioned before, are continuous images of the perfect pre-images of separable completely metrizable spaces. Thus they are the subclass of Lindelöf Σ -spaces obtained by starting with separable completely metrizable spaces rather than arbitrary separable metrizable ones. Indeed, MAD families cannot be completely metrizable, or even analytic. This is a classical result. There is a “BFT-flavoured” proof in [19]. Thus a topological approach to NIP should start with K -analytic spaces. It is an interesting question whether one can go beyond that class. The second author is engaged in a project to

define K -projective sets in analogy to the projective sets in descriptive set theory [36], but it is not clear whether this is possible. If it is, does this enable NIP results to be extended to a larger class of theories?

Connections between NIP and compact spaces are explored in [34]. There is a brief survey of references for related topology in [18]. There is much more to be investigated, but that is beyond the scope of this paper.

6 Appendix: Proof that all Lindelöf Σ -spaces are Grothendieck

First we need the *Arhangel'skiĭ-Pytkeev Theorem* ([2]) and some definitions.

Theorem 6.1. $t(C_p(X)) = \sup\{L(X^{n+1}) : n < \omega\}$.

Proof. Suppose first that $L(X^n) \leq \kappa$ for all $n < \omega$. Let $A \subseteq C_p(X)$ and take $f \in \overline{A} \subseteq C_p(X)$. Now, for each $n < \omega$ and each $\xi = (x_0, \dots, x_n) \in X^{n+1}$, there is some $g_\xi \in A$ such that

$$(\forall i \leq n)(|g_\xi(x_i) - f(x_i)| < \frac{1}{n+1}).$$

Since both g_ξ and f are continuous, for each $i \leq n$ there is a neighbourhood $O(x_i)$ of x_i such that $(\forall y \in O(x_i))(|g_\xi(y) - f(y)| < \frac{1}{n+1})$.

Now define the neighbourhood $U_\xi = O(x_0) \times \dots \times O(x_n)$ and let $\mathcal{U}_{n+1} = \{U_\xi : \xi \in X^{n+1}\}$. Clearly, $\xi \in U_\xi$ and so \mathcal{U}_{n+1} covers X^{n+1} . Since $L(X^{n+1}) \leq \kappa$, we have some $\mathcal{U}_{n+1}^* \subseteq \mathcal{U}_{n+1}$, a cover of X^{n+1} , such that $|\mathcal{U}_{n+1}^*| \leq \kappa$, and we can define $B_{n+1} = \{g_\xi : U_\xi \in \mathcal{U}_{n+1}^*\}$, for each $n < \omega$. Now let $B = \bigcup_{n < \omega} B_{n+1}$. Clearly, $|B| \leq \kappa$ and $B \subseteq A$. We now claim that $f \in \overline{B}$. Indeed, let $y_0, \dots, y_n \in X$ and $\varepsilon > 0$, and consider the basic open neighbourhood $W(f; y_0, \dots, y_n; \varepsilon)$; we see that it has non-empty intersection with B . Without loss of generality, suppose $\frac{1}{n+1} < \varepsilon$. Since \mathcal{U}_{n+1}^* is a cover of X^{n+1} , let $\xi \in X^{n+1}$ be such that $(y_0, \dots, y_n) \in U_\xi$. Then $(\forall i \leq n) |g_\xi(y_i) - f(y_i)| < \frac{1}{n+1} < \varepsilon$.

The converse is more difficult to prove. We do not actually need it for our proof that Lindelöf Σ implies Grothendieck. Nonetheless, for the sake of completeness and exposure to C_p -theory, here it is. Suppose that $t(C_p(X)) \leq \kappa$ and fix $n < \omega$. Let \mathcal{V} be an open cover of X^{n+1} . Let \mathcal{E} be the family of all finite \mathcal{V} -small families, where a

family \mathcal{W} of open subsets of X is \mathcal{V} -small if for any $W_0, \dots, W_n \in \mathcal{W}$, there is some $V \in \mathcal{V}$ such that $W_0 \times \dots \times W_n \subseteq V$. For each $\mathcal{W} \in \mathcal{E}$, define

$$A_{\mathcal{W}} = \left\{ f \in C_p(X) : f \left(X \setminus \bigcup \mathcal{W} \right) = \{0\} \right\}$$

and let $A = \bigcup_{\mathcal{W} \in \mathcal{E}} A_{\mathcal{W}}$. We claim that $\overline{A} = C_p(X)$.

Indeed, let $f \in C_p(X)$ and let K be a finite subset of X . Take a finite family \mathcal{Q}_K of open sets in X with the following property: for any $(y_0, \dots, y_n) \in K^{n+1}$, there are $Q_0, \dots, Q_n \in \mathcal{Q}_K$ such that $y_i \in Q_i$ for each $i \leq n$ and $Q_0 \times \dots \times Q_n \subseteq G$ for some $G \in \mathcal{V}$. Clearly, $K \subseteq \bigcup \mathcal{Q}_K$. For each $x \in K$, define $W_x = \bigcap \{V \in \mathcal{Q}_K : x \in V\}$. Now, let $\mathcal{W}_K = \{W_x : x \in K\}$. Again, $K \subseteq \bigcup \mathcal{W}_K$. We now see that \mathcal{W}_K is \mathcal{V} -small: consider a set of the form $W_{x_0} \times \dots \times W_{x_n}$. By construction, there are $Q_0, \dots, Q_n \in \mathcal{Q}_K$ such that $(\forall i \leq n)(x_i \in Q_i)$ and $Q_0 \times \dots \times Q_n \subseteq G$ for some $G \in \mathcal{V}$. Since $W_{x_i} \subseteq Q_i$ for each $i \leq n$, we have $W_{x_0} \times \dots \times W_{x_n} \subseteq G$, proving the claim.

Now, by complete regularity, take a function $g \in C_p(X)$ such that $f|_K = g|_K$ and $g(X \setminus \bigcup \mathcal{W}) = \{0\}$. The point is that K is finite and $K \subseteq \bigcup \mathcal{W}_K$. So we can certainly get a continuous real-valued function h which is 1 on K and 0 on $X \setminus \bigcup \mathcal{W}$. Then we can add some continuous real-valued functions to h to get the desired values on K , since a finite sum of continuous functions is continuous. Clearly, $g \in A_{\mathcal{W}_K} \subseteq A$ and g lies in all basic neighbourhoods of f based on K (i.e. basic neighbourhoods of the form $W(f; x_0, \dots, x_n, \varepsilon)$ where $x_0, \dots, x_n \in K$ and $\varepsilon > 0$). Thus, $f \in \overline{A}$ and $\overline{A} = C_p(X)$.

If we take $\mathbf{1}$ to be the function identically 1 we have $\mathbf{1} \in \overline{A}$. Since $t(C_p(X)) \leq \kappa$, there is a $B \subseteq A$ such that $\mathbf{1} \in \overline{B}$ and $|B| \leq \kappa$. Then, there is a subfamily $\mathcal{E}_0 \subseteq \mathcal{E}$ for which $B \subseteq \bigcup_{\mathcal{W} \in \mathcal{E}_0} A_{\mathcal{W}}$ and $|\mathcal{E}_0| \leq \kappa$. Given any $\mathcal{W} \in \mathcal{E}_0$, we introduce the following notation: for each $V = (Q_0, \dots, Q_n) \in \mathcal{W}^{n+1}$, we fix $G_V \in \mathcal{V}$ such that $V \subseteq G_V$ and then define $\mathcal{V}_{\mathcal{W}} = \{G_V : V \in \mathcal{W}^{n+1}\}$. Since the family $\mathcal{V}_{\mathcal{W}}$ is finite and $|B| \leq \kappa$, then $\mathcal{V}^* = \bigcup_{\mathcal{W} \in \mathcal{E}_0} \mathcal{V}_{\mathcal{W}}$ always has cardinality $\leq \kappa$.

It remains to show that \mathcal{V}^* covers X^{n+1} . Take $(x_0, \dots, x_n) \in X^{n+1}$ and let $U = \{f_p(X) : (\forall i \leq n)(f(x_i) > 0)\}$. Notice that U is open in $C_p(X)$ and that $\mathbf{1} \in U$. Since $\mathbf{1} \in \overline{B}$ and $B \subseteq \bigcup_{\mathcal{W} \in \mathcal{E}_0} A_{\mathcal{W}}$, there is a $\mathcal{W}_0 \in \mathcal{E}_0$ such that $U \cap A_{\mathcal{W}_0} \neq \emptyset$. Now take $g \in U \cap A_{\mathcal{W}_0}$; then $g(x_i) > 0$ for each $i \leq n$ and $g(x) = 0$ for $x \in X \setminus \bigcup \mathcal{W}_0$. Thus, $x_i \in \bigcup \mathcal{W}_0$ for each $i \leq n$. Take $Q_i \in \mathcal{W}_0$ such that $x_i \in Q_i$, $i \leq n$. Then $(x_0, \dots, x_n) \in Q = Q_0 \times \dots \times Q_n \subseteq G_V \in \mathcal{V}_{\mathcal{W}_0} \subseteq \mathcal{V}^*$. \square

Our plan is to show Lindelöf p -spaces are weakly Grothendieck and

then use Proposition 2.12 to show they and (hence) Lindelöf Σ -spaces are Grothendieck. First, we introduce some important definitions.

Definition 6.2. A topological space is a *k-space* if subspaces are closed if and only if their intersection with each compact subspace is closed.

We will show

Proposition 6.3 ([3]). *Every k-space is weakly Grothendieck.*

Theorem 6.4. *Every Lindelöf p-space is a k-space.*

Actually, the “Lindelöf” is superfluous, but it is all we need and its inclusion makes the proof much easier. Metrizable spaces are easily seen to be *k*, since if a subspace W is not closed, there is a sequence converging from S to a point outside. That sequence together with its limit are a compact set with non-closed intersection with S . So all we have then to do is show:

Lemma 6.5. *A perfect pre-image of a k-space is a k-space.*

This is Theorem 3.7.25 in [14]. We shall sketch his proof, but it’s not C_p -theory, so the reader is welcome to skip it.

Proof. Given an arbitrary Hausdorff space X , one can put a finer topology on X by declaring closed all subsets of X that have closed intersections with each compact subspace of X . This new space is called kX and can be shown to be a *k-space*. Every continuous map f from X to a Hausdorff space Y induces a continuous map kf from kX to kY , by assigning to $x \in kX$ the image $f(x) \in kY$. It is not hard to show that kf is perfect since f is. Define a function $k_X : kX \rightarrow X$ by $k_X(x) = x$. Then $kf = f \circ k_X$ and it follows that k_X is perfect. But k_X is one-to-one, so it’s a homeomorphism, and so X is a *k-space*. \square

We shall also need:

Lemma 6.6 ([14, 3.7.25]). *A mapping from a k space X to a space Z is continuous if and only if for any compact K included in X , the restriction $f|K$ is continuous.*

Proof. Take any closed $B \subseteq Y$. For any compact $K \subseteq X$, we have $f^{-1}(B) \cap K = (f|K)^{-1}(B)$, which shows $f^{-1}(B)$ is closed, since X is a *k-space*. Thus f is continuous. \square

The following result is well-known:

Proposition 6.7. *The continuous image of a countably compact space is countably compact.*

Proof. Let X be countably compact and $f : X \rightarrow Y$ a continuous function. Suppose B is an infinite subset of $f(X)$, and let X' be obtained by picking one point from each $f^{-1}(\{b\})$, for each $b \in B$. Then X' is infinite and so has a limit point $x \in X$. Then $f(x)$ is a limit point of B , since if U is an open set about $f(x)$, the pre-image $f^{-1}(U)$ is an open set about x and so contains some $a \in X'$. Then U contains $f(a)$, which is in B . \square

Proof of Proposition 6.3. Let A be countably compact in $C_p(X)$, with X a k -space. Since $C_p(X)$ is canonically embedded in \mathbb{R}^X , A is also a subspace of \mathbb{R}^X . By continuity, by Proposition 6.7 the images of A under the natural projections of \mathbb{R}^X onto \mathbb{R} are also countably compact in \mathbb{R} . Since \mathbb{R} is metrizable, it is normal so the images of A have countably compact closures, but then by metrizability they have compact closures. Since A is included in the topological product of these projections, the closure of A in \mathbb{R}^X is a compact subspace F of \mathbb{R}^X . It remains to show that $F \subseteq C_p(X)$, i.e. each member of f is continuous.

The restriction map r mapping \mathbb{R}^X into \mathbb{R}^Y sends A to a dense subset of $r(F)$, since r is continuous and A is dense in F . Since r is continuous, $r(A)$ is countably compact in $C_p(Y)$ by Proposition 6.7. As a compact space, Y is Grothendieck and so the closure of $r(A)$ in $C_p(Y)$ is compact. But then that closure must equal $r(F)$, since, by continuity of r , $r(F)$ is the closure of $r(A)$ in \mathbb{R}^Y , so is the smallest closed set including $r(A)$ there. Thus, for each $f \in F$ and each compact $Y \subseteq X$, there is a g in the closure of A such that the restriction of f to Y coincides with the restriction of g to Y . Since X is a k -space, it follows that f is continuous. \square

Definition 6.8. Let X be a topological space.

- (a) The *density* $d(X)$ of X is the smallest cardinality of a dense subset of X .
- (b) A *network* of X is a family \mathcal{A} of subsets of X such that given any $x \in X$ and open set U containing x , there is an $A \in \mathcal{A}$ satisfying $x \in A \subseteq U$. The *network weight* $nw(X)$ of X is the smallest cardinality of a network of X .

- (c) The *i-weight* $iw(X)$ of X is the minimal weight of all the spaces onto which X can be continuously mapped.

We see that if X is compact, then $nw(X) = w(X)$ (see [2]): given any Hausdorff space X , we can construct a Hausdorff space Y such that $w(Y) \leq nw(X)$ and Y is a continuous one-to-one image of X . To see this, let \mathcal{A} be a network of X of size $nw(X)$. Consider all the pairs of elements $A_1, A_2 \in \mathcal{A}$ for which there are disjoint open subsets U_1 and U_2 such that $A_1 \subseteq U_1$ and $A_2 \subseteq U_2$. It suffices to define the topological space Y as the one with the same underlying set as X and with the topology generated by the all the open sets of X obtained as above. Thus the identity map is as desired. That $nw(X) = w(X)$ for every compact space X follows from the preceding discussion and the well-known fact that a continuous one-to-one map between a compact space and a Hausdorff space is a homeomorphism.

Some other relatively easy facts regarding these cardinals are the following: $iw(X) \leq nw(X)$, $d(X) \leq w(X)$, $nw(X) = nw(C_p(X))$ and $iw(X) = d(C_p(X))$ for every X . See [2], p. 26, for a detailed exposition. Now we proceed with two definitions that arise naturally when studying topological function spaces:

Definition 6.9. Let X be a topological space.

- (a) X is κ -*monolithic* if $nw(\overline{A}) \leq \kappa$ whenever $|A| \leq \kappa$. We say that X is *monolithic* if it is κ -monolithic for every κ .
- (b) X is κ -*stable* if for every continuous image Y of X we have: $iw(Y) \leq \kappa$ if and only if $nw(Y) \leq \kappa$. We say that X is *stable* if it is κ -stable for every κ .

Notice that being monolithic is inherited by all subspaces. Examples of monolithic spaces include all metric spaces and spaces with a countable network, whereas examples of stable spaces include all compact spaces and Lindelöf Σ -spaces (as we shall prove below). The following theorem shows that stability and monolithicity are intertwined notions for X and $C_p(X)$.

Theorem 6.10 (Arhangel'skiĭ). *X is κ -stable if and only if $C_p(X)$ is κ -monolithic.*

Proof. Assume $C_p(X)$ is κ -monolithic and let $f : X \rightarrow Y$ be a continuous surjection. Since always $iw(Y) \leq nw(Y)$, we also assume that $iw(Y) \leq \kappa$. By Lemma 2.2, $C_p(Y)$ is homeomorphic

to a subspace of $C_p(X)$ and so it's κ -monolithic. In particular, by looking at a dense subset of $C_p(Y)$ and using the definition of κ -monolithicity, $nw(C_p(Y)) \leq d(C_p(Y))$. Using that $iw(Y) = d(C_p(Y))$ and $nw(Y) = nw(C_p(Y))$, we obtain $nw(Y) \leq iw(Y)$ as desired. Conversely, assume that X is κ -stable and let $M \subseteq C_p(X)$ be such that $|M| < \kappa$. Define $f : X \rightarrow \mathbb{R}^M$ as the diagonal product of M , i.e. $f(x) = \langle g(x) : g \in M \rangle$, and let $Y = f(X)$. By our choice, $w(Y) \leq |M| \leq \kappa$ as in the proof of Theorem 2.5.

The rest of the proof is a bit dense. Although the theorem is important, the reader is unlikely to use the *real quotient topology*, so it is not unreasonable to take the rest on faith. But we shall proceed.

Let Y^* be the topological space with the same underlying set as Y but endowed with the *real quotient topology corresponding to f* , i.e. the topology on Y^* is the strongest completely regular topology for which f is continuous (see [2], p. 14). Then the identity $i : Y^* \rightarrow Y$ is a continuous bijection. By the definition of the i -weight, $iw(Y^*) \leq w(Y) \leq \kappa$.

Conversely, since f defines a continuous surjection from X onto Y^* , i.e. the real quotient map $f^* = i^{-1} \circ f$, and X is κ -stable, $nw(Y^*) \leq \kappa$ and $nw(C_p(Y^*)) = nw(Y^*) \leq \kappa$. Also, the space $C_p(Y^*)$ is homeomorphic to the closed subspace $F = \Phi_{f^*}[C_p(Y^*)] = \{g \circ f^* : g \in C_p(Y^*)\}$ of $C_p(X)$ since f^* is a real quotient map (see Section 0.4 of [2] for a detailed discussion of real quotient maps). Now let $g \in M$ and notice that $g = \pi_g \circ f = \pi_g \circ i \circ f^*$ where $\pi_g : \mathbb{R}^M \rightarrow \mathbb{R}$ is the projection $\pi_g(\langle h(x) : h \in M \rangle) = g(x)$. Also, $\pi_g \circ i : Y^* \rightarrow \mathbb{R}$ is continuous, so $g \in F$ and thus $M \subseteq F$. Thus $\overline{M} \subseteq F$ since F is closed, and we conclude by noticing that $nw(\overline{M}) \leq nw(F) = nw(C_p(Y^*)) \leq \kappa$. \square

Although we will not use this fact here, it is worth noting that the previous theorem is self-dual, i.e.: X is κ -monolithic if and only if $C_p(X)$ is κ -stable. See [2], p. 78, for a proof.

The following diagram-chasing lemma appears in [2], p. 80, and is used to prove that Lindelöf Σ -spaces are stable. We omit the proof here.

Lemma 6.11. *Let $f : X \rightarrow Y$, $g : X \rightarrow Z$ and $h : Z \rightarrow T$ be continuous surjections. Also assume that f is perfect and that h is a bijection. Then Z is a continuous image of a closed subspace of $Y \times T$.*

Theorem 6.12. *Lindelöf Σ -spaces are stable.*

Proof. Let Z be a Lindelöf Σ -space. Since Lindelöf Σ -spaces are continuous images of Lindelöf p -spaces, any continuous image of Z is also Lindelöf Σ so it suffices to see that $nw(Z) \leq iw(Z)$.

Suppose $iw(Z) = \kappa$ and let $h : Z \rightarrow T$ be a continuous bijection and suppose $w(T) \leq \kappa$. Again, by definition of Lindelöf Σ -spaces, there is a second countable space Y , a space X , a perfect surjective map $f : X \rightarrow Y$ and a continuous surjection $g : X \rightarrow Z$. By the previous lemma, Z is the continuous image of a closed subspace of $Y \times T$. Thus $nw(Z) \leq nw(Y \times T) \leq \kappa$ as desired. \square

Problem 1. *Is there a connection between C_p -theoretic stability (or monolithicity) and model-theoretic stability?*

Lemma 6.13. *A countably tight monolithic compact space is Fréchet-Urysohn.*

Proof. Let X be a countably tight monolithic compact space and suppose that $x \in \overline{A}$ for some $A \subseteq X$. Let $B \subseteq A$ be a countable subset such that $x \in \overline{B}$. Thus $w(\overline{B}) = nw(\overline{B}) = \aleph_0$ since \overline{B} is compact and monolithic. Then \overline{B} is second countable and so there is a sequence in B converging to x . \square

We now have all we need to prove Theorem 2.18 that says that all Lindelöf Σ spaces are Grothendieck:

Proof of Theorem 2.18. By Theorem 6.4, every Lindelöf p -space is a k -space. Then by Proposition 6.3 and Theorem 6.12, Lindelöf p -spaces are weakly Grothendieck and stable. By the Arhangel'skiĭ-Pytkeev Theorem, $C_p(X)$ for a Lindelöf p -space X is countably tight, since finite products of Lindelöf p -spaces are Lindelöf. Moreover, such $C_p(X)$ is monolithic by Theorem 6.10. We conclude that Lindelöf p -spaces are Grothendieck by Proposition 2.12 and Lemma 6.13. Finally, given that Lindelöf Σ spaces are continuous image of Lindelöf p -spaces, we conclude that Lindelöf Σ spaces are Grothendieck by Theorem 2.16. \square

Definition 6.14. A topological space X is σ -compact if it can be written as a countable union of compact subspaces. X is k -separable if it has a dense σ -compact subspace.

The following theorem collects well-known (to C_p -theorists) results on conditions that imply Grothendieck:

Theorem 6.15. *A topological space X is a Grothendieck space if it satisfies any of the following:*

- (i) X has a dense countably compact subspace.*
- (ii) X is k -separable.*
- (iii) X has a dense Lindelöf Σ -space.*

Proof.

- (i) Grothendieck's Theorem implies that countably compact spaces are weakly Grothendieck. Pryce [29] proved that they are indeed Grothendieck. The result follows from Theorem 2.15.
- (ii) Follows from (iii) (see [38]) and Theorem 2.15.
- (iii) This follows immediately from Theorems 2.15 and 2.18.

□

Problem 2. *Find a common generalization of these three conditions that implies Grothendieck.*

7 Acknowledgement

We have greatly appreciated the comments of the anonymous referee, who has prodded us to make this paper more accessible to model theorists.

References

- [1] A. Arhangel'skiĭ. Embedding in C_p -spaces. *Topology and its applications*, 85:9–33, 1998.
- [2] A. V. Arhangel'skiĭ. *Topological Function Spaces*, volume 78 of *Mathematics and its Applications (Soviet Series)*. Kluwer Academic Publishers Group, Dordrecht, 1992.
- [3] A. V. Arhangel'skiĭ. On a theorem of Grothendieck in C_p -theory. *Topology and its applications*, 80:21–41, 1997.
- [4] I. Ben Yaacov. Model theoretic stability and definability of types after A. Grothendieck. *The Bulletin of Symbolic Logic*, 20(4):491–496, 2014.
- [5] I. Ben Yaacov, A. Berenstein, C. W. Henson, and A. Usvyatsov. *Model Theory with Applications to Algebra and Analysis II*, volume 350 of *London Mathematical Society Lecture Note Series*, chapter 10: Model theory for metric structures, pages 315–427. Cambridge University Press, Cambridge, 2008.
- [6] J. Bourgain, D. H. Fremlin, and M. Talagrand. Pointwise compact sets of Baire-measurable functions. *American Journal of Mathematics*, 100:845–886, 1978.
- [7] P. Casazza and J. Iovino. On the undefinability of Tsirelson's space and its descendants. ArXiv preprint: 1710.05889.
- [8] P. Casazza and T. J. Shura. *Tsirelson's Space*. Lecture Notes in Mathematics. Springer-Verlag, 1989.
- [9] P. J. Casazza, E. Dueñez, and J. N. Iovino. The non-definability of the spaces of Tsirelson and Schlumprecht. Preprint, 2021.
- [10] C. C. Chang and H. J. Keisler. *Model theory*, volume 73 of *Studies in logic and the foundations of mathematics*. Amsterdam North-Holland Publishing Company, 1973.
- [11] C. J. Eagle. Omitting types for infinitary $[0, 1]$ -valued logic. *Annals of Pure and Applied Logic*, 165:913–932, 2014.
- [12] C. J. Eagle. *Topological Aspects of Real-Valued Logic*. PhD thesis, University of Toronto, 2015.
- [13] C. J. Eagle, C. Hamel, and F. D. Tall. Two applications of topology to model theory. *Ann. Pure Appl. Logic*, 172(5), 2021. Paper No. 102907.

- [14] R. Engelking. *General Topology*. Heldermann Verlag, Berlin, 1989.
- [15] T. Figiel and W. Johnson. A uniformly convex Banach space which contains no l_p . *Compositio Mathematica*, 29:179–190, 1974.
- [16] W. T. Gowers. Recent results in the theory of infinite-dimensional Banach spaces. In *Proceedings of the International Congress of Mathematicians Vol 1, 2 (Zürich, 1994)*, pages 933–942. Birkhäuser, Basel., 1995.
- [17] A. Grothendieck. Critères de compacité dans les espaces fonctionnels généraux. *American Journal of Mathematics*, 74:168–186, 1952.
- [18] C. Hamel and F. D. Tall. Model theory for C_p -theorists. *Topology and its Applications*, 281, 2020.
- [19] H. Horowitz and S. Todorćević. Compact sets of Baire class one functions and maximal almost disjoint families. Submitted. arXiv:1905.05898, 2019.
- [20] T. Hyttinen, K. Kangas, and J. Väänänen. On second-order characterizability. <https://arxiv.org/abs/1208.5167>, 2012.
- [21] J. Iovino. Stable models and reflexive Banach spaces. *Journal of Symbolic Logic.*, 64(4):1595–1600, 1999.
- [22] K. Khanaki. Stability, NIP, and NSOP; model theoretic properties of formulas via topological properties of function spaces. arXiv:1410.3339, 2014.
- [23] K. Khanaki. Correspondences between model theory and Banach space theory. arXiv:1512.08691, 2015.
- [24] K. Khanaki and A. Pillay. Remarks on NIP in a model. arXiv:1706.04674, 2017.
- [25] M. Morley. Applications of topology to $\mathcal{L}_{\omega_1, \omega}$. *Proceedings of Symposia in Pure Mathematics*, 25:233–240, 1974.
- [26] E. Odell. *Analysis and logic*, volume 262 of *London Mathematical Society Lecture Note Series*, chapter 3: On Subspaces, Asymptotic Structure, and Distortion of Banach Spaces; Connections with Logic, pages 189–267. Cambridge University Press, Cambridge, 2002.
- [27] O. G. Okunev. On Lindelöf sets of continuous functions. *Topology and its Applications*, 63:91–96, 1995.

- [28] A. Pillay. Stability and Grothendieck. *arXiv: 1703.04761*, 2017.
- [29] J. Pryce. A device of R. J. Whitley’s applied to pointwise compactness in spaces of continuous functions. *Proceedings of the London Mathematical Society*, 3-23:532–546, 1971.
- [30] V. Pták. A combinatorial lemma on the existence of convex means and its applications to weak compactness. In V. L. Klee, editor, *Convexity*, volume 7 of *Proceedings in Symposia in Pure Mathematics*, pages 437–450, 1963.
- [31] W. V. Quine. *Philosophy of Logic*. Harvard University Press, Cambridge, Mass, 2nd edition, 1986.
- [32] C. A. Rogers and J. E. Jayne. K -analytic sets. In C. A. Rogers, editor, *Analytic sets*, pages 2–175. Academic Press, London, 1980.
- [33] P. Simon. *A guide to NIP theories*. Lecture Notes in Logic. Cambridge University Press, Cambridge, 2015.
- [34] P. Simon. Rosenthal compacta and NIP formulas. *Fund. Math.*, 231(1):81–92, 2015.
- [35] F. D. Tall. Co-analytic spaces, K -analytic spaces, and definable versions of Menger’s conjecture. *Topology and its Applications*, 283, 2020.
- [36] F. D. Tall. Countable tightness and the Grothendieck property in C_p -theory. *Topology and its Applications*, 2021. To appear.
- [37] V. Tkachuk. *A C_p -theory problem book*. Problem Books in Mathematics. Vol. I-IV. Springer, 2011-2015.
- [38] V. V. Tkachuk. Lindelöf Σ : An omnipresent class. *Revista de la Real Academia de Ciencias Exactas Físicas y Naturales*, 104(2):221–244, 2010.
- [39] S. Todorćević. *Topics in Topology*. Lecture Notes in Mathematics. Springer-Verlag Berlin Heidelberg, 1997.
- [40] B. Tsirelson. It is impossible to imbed ℓ^p or c_0 into an arbitrary Banach space. *Funkcional Anal I Priložen*, 8(2):57–60, 1974.
- [41] S. Willard. *General Topology*. Dover, 2012.