

A RADON-NIKODÝM THEOREM FOR LOCAL COMPLETELY POSITIVE INVARIANT MULTILINEAR MAPS

ANINDYA GHATAK AND SANTHOSH KUMAR PAMULA

ABSTRACT. In this article, we introduce local completely positive k -linear maps between locally C^* -algebras and obtain Stinespring type representation by adopting the notion of “invariance” defined by J. Heo for k -linear maps between C^* -algebras. Also, we supply the minimality condition to make certain that minimal representation is unique up to unitary equivalence. As a consequence, we prove Radon-Nikodým theorem for unbounded operator-valued local completely positive invariant k -linear maps. The obtained Radon-Nikodým derivative is a positive contraction on some Hilbert space with several reducing subspaces.

1. INTRODUCTION

In 1987, Christensen and Sinclair introduced multilinear completely positive (in short CP) maps [4] between C^* -algebras and used this notion to develop theory on the cohomology group of the von Neuman algebras. These authors gave a representation for completely bounded symmetric multilinear maps in terms of $*$ -representations of the algebra and constructed suitable bridging operators between the Hilbert spaces by using the famous result of Wittstock’s matricial Hahn Banach theorem [24]. In the article [22], Sinclair and Smith proved that a von Neuman algebra is injective if and only if every bilinear completely bounded (CB) maps can be factorized in terms of bilinear CP-maps. Paulsen and Smith [21] extended this result to operator spaces by using Wittstock’s matricial Hahn-Banach theorem. Further, with the help of the representation of completely bounded symmetric multilinear maps, Effros et. al. showed in [7] that the completely bounded cohomology group from a C^* -algebra to an injective von Neumann algebra is zero.

To study the structure of multilinear CP-maps, Christensen established Stinespring type theorem for symmetric k -linear CP-maps (see [4, Section 3.] for details). Motivated by this work, Heo [15] came up with an additional property (called as **invariant**) on k -linear CP-maps as an analogue of invariant sesquilinear forms [1]. With this additional invariance property, Heo obtained a dilation for invariant k -linear CP-map to a multilinear representation which can be factorized as a product of m -number of commuting representations, where $m = \lceil \frac{k+1}{2} \rceil$. Further, the author developed a Radon-Nikodým theorem for completely bounded and completely positive invariant k -linear maps. Several authors studied invariant multilinear maps, for example, see [9, 11, 12, 13, 14, 19] and references therein. In this article, our main concern is to study the structure of k -linear local CP-maps between locally C^* -algebras. These results generalize the work of Heo [16].

We organize this manuscript into five sections. In the second section, we fix notations, recall some definitions and known results from the literature which are needed for the following sections. In the third section, firstly we construct a new quantized domain (contains the original quantized domain) associated to a given local completely positive invariant k -linear map defined from a locally C^* -algebra \mathcal{A} to a locally C^* -algebra of non-commutative continuous functions on a quantized domain. Next, we construct a commuting family of local contractive $*$ -homomorphisms from \mathcal{A} to a locally C^* -algebra

Date: December 31, 2021.

2010 Mathematics Subject Classification. 46L07, 46L05, 46G25, 47L40.

Key words and phrases. Locally C^* -algebra, Multilinear map, Local completely positive map, Stinespring’s theorem, Radon-Nikodým theorem.

of non-commutative continuous functions on the new quantized domain. Finally, we show that the product of these local contractive $*$ -homomorphisms is a desired dilation of the given invariant local completely positive k -linear map (see Theorem 3.2). This result extends the Anar Dosiev's Stinespring theorem [6, Theorem 5.1] to k -linear maps and a locally convex analogue of Heo's result [16].

In the fourth section, a suitable notion of minimality for our Stinespring type representation has been identified to ensure the uniqueness of minimal representation up to unitary equivalence (see Theorem 4.4). In the final section, we prove Radon-Nikodým theorem for multilinear local CP maps that characterizes the set of all maps dominated by a fixed local completely positive invariant k -linear map. Here, the obtained Radon-Nikodým derivative is a positive contraction on some Hilbert space with an upward filtered family of reducing sub spaces (see Theorem (5.5)).

2. PRELIMINARIES

A locally C^* -algebra \mathcal{A} is a $*$ -algebra which is complete in a locally convex topology determined by an upward filtered family $\{p_\alpha : \alpha \in \Lambda\}$ of C^* -seminorms. In this article, we denote \mathcal{A} and \mathcal{B} as the unital locally C^* -algebras with the upward filtered family of C^* -seminorms $\{p_\alpha : \alpha \in \Lambda\}$ and $\{q_\ell : \ell \in \Omega\}$ on \mathcal{A} and \mathcal{B} respectively. Locally C^* -algebras are the generalization of C^* -algebras, indeed it can be seen as the projective limit of the inverse system of C^* -algebras. For instance, let $I_\alpha = \{a \in \mathcal{A} : p_\alpha(a) = 0\}$. Then I_α is a closed two-sided ideal in \mathcal{A} and hence $\mathcal{A}_\alpha := \mathcal{A}/I_\alpha$ is a C^* -algebra with the C^* -norm $\|\cdot\|_\alpha$ given by

$$\|a_\alpha\|_\alpha := p_\alpha(a), \text{ for all } a_\alpha := a + I_\alpha \in \mathcal{A}_\alpha, \alpha \in \Lambda.$$

For every $\alpha, \beta \in \Lambda$ there exist a canonical projection $\pi_\alpha : \mathcal{A} \rightarrow \mathcal{A}_\alpha$ and a canonical $*$ -epimorphism $\pi_{\alpha, \beta} : \mathcal{A}_\beta \rightarrow \mathcal{A}_\alpha$ whenever $\alpha \leq \beta$ satisfying the relation $\pi_\alpha = \pi_{\alpha, \beta} \circ \pi_\beta$. The projective limit of an inverse system $(\{\mathcal{A}_\alpha\}_{\alpha \in \Lambda}; \{\pi_{\alpha, \beta}\}_{\alpha \leq \beta})$ of C^* -algebras is defined as

$$\varprojlim_{\alpha} \mathcal{A}_\alpha := \left\{ \{a_\alpha\}_{\alpha \in \Lambda} \in \prod_{\alpha \in \Lambda} \mathcal{A}_\alpha : \pi_{\alpha, \beta}(a_\beta) = a_\alpha \text{ whenever } \alpha \leq \beta, \alpha, \beta \in \Lambda \right\}.$$

Then \mathcal{A} is identified with $\varprojlim_{\alpha} \mathcal{A}_\alpha$ via the following map

$$(1) \quad a \rightarrow \{\pi_\alpha(a)\}_{\alpha \in \Lambda} \text{ for all } a \in \mathcal{A}.$$

From the identification given in Equation (1), it is natural to see the existence of local self-adjoints and local positive elements in local C^* -algebras. For further details, we refer the reader to [10].

Definition 2.1. [6] *An element $a \in \mathcal{A}$ is called*

- (1) *local self-adjoint if $a^* = a + x$, where $x \in \mathcal{A}$ such that $p_\alpha(x) = 0$ for some $\alpha \in \Lambda$;*
- (2) *local positive if $a = b^*b + y$, where $b, y \in \mathcal{A}$ such that $p_\beta(y) = 0$ for some $\beta \in \Lambda$.*

In other words, a is local self-adjoint (local positive) if and only if $\pi_\alpha(a)$ is self-adjoint (positive) in \mathcal{A}_α , for some $\alpha \in \Lambda$. For brevity, we write $a^* =_\alpha a$ if $\pi_\alpha(a)$ is self-adjoint in \mathcal{A}_α and $a \geq_\alpha 0$ if $\pi_\alpha(a)$ positive in \mathcal{A}_α . In 2008, Anar Dosi defined the notion of local completely positive maps between local operator systems, which is a locally convex analogues of operator systems. Initially, such maps between local operator spaces have been studied by Effros (see [8] for details).

Now we turn our discussion to multilinear maps between locally C^* -algebras. Let $\mathcal{A}^k = \mathcal{A} \times \dots \times \mathcal{A}$ denote the Cartesian product of k -copies of \mathcal{A} , where $k \in \mathbb{N}$. Then

$$\mathcal{A}^k := \{(a_1, \dots, a_k) : a_i \in \mathcal{A}, 1 \leq i \leq k\}$$

is a locally convex space with associated seminorm $\|\cdot\|_\alpha$ for each $\alpha \in \Lambda$ given by

$$\|(a_1, \dots, a_k)\|_\alpha = \max\{p_\alpha(a_i) : 1 \leq i \leq k\}, \text{ for all } (a_1, \dots, a_k) \in \mathcal{A}^k.$$

Definition 2.2. (α -symmetric) An element $(a_1, a_2, \dots, a_k) \in \mathcal{A}^k$ is said to be α -symmetric (for $\alpha \in \Lambda$) if $a_p^* =_\alpha a_{k+1-p}$ for all $p = 1, 2, \dots, m$ where $m = \lceil \frac{k+1}{2} \rceil$.

Now we define the notion of symmetric multilinear maps in local set up. A map $\varphi : \mathcal{A}^k \rightarrow \mathcal{B}$ is called k -linear map if

$$(2) \quad \varphi(a_1, \dots, a_i + \lambda b_i, \dots, a_k) = \varphi(a_1, \dots, a_i, \dots, a_k) + \lambda \varphi(a_1, \dots, b_i, \dots, a_k),$$

for each $a_i, b_i \in \mathcal{A}$, $\lambda \in \mathbb{C}$ and $1 \leq i \leq k$. The adjoint φ^* of a k -linear map φ is defined as,

$$(3) \quad \varphi^*(a_1, a_2, a_3, \dots, a_k) = \varphi(a_k^*, \dots, a_3^*, a_2^*, a_1^*)^*, \text{ for all } a_1, \dots, a_k \in \mathcal{A}.$$

A k -linear map is called *symmetric*, if $\varphi = \varphi^*$. In other words, φ is symmetric if and only if

$$\varphi(a_1, a_2, a_3, \dots, a_k)^* = \varphi(a_k^*, \dots, a_3^*, a_2^*, a_1^*), \text{ for all } a_1, \dots, a_k \in \mathcal{A}.$$

Since \mathcal{A} is a locally C^* -algebra, the set of all $n \times n$ matrices over \mathcal{A} denoted by $M_n(\mathcal{A})$ is a local C^* -algebra with an upward filtered family $\{p_\alpha^n : \alpha \in \Lambda\}$ of C^* -seminorms.

Definition 2.3. [16] For every $n \in \mathbb{N}$, the amplification map $\varphi_n : M_n(\mathcal{A})^k \rightarrow M_n(\mathcal{B})$ is defined by

$$\varphi_n(A_1, A_2, \dots, A_k) = \left[\sum_{r_1, r_2, \dots, r_{k-1}=1}^n \varphi(a_{1_{ir_1}}, a_{2_{r_1 r_2}}, \dots, a_{k_{r_{k-1}j}}) \right]_{i,j=1}^n,$$

where $A_t = [a_{t_{ij}}]_{i,j=1}^n \in M_n(\mathcal{A})$, for $t = 1, \dots, k$.

Now, we introduce local CP-maps and local CB-maps in this context (cf. [20] for CP-maps). Throughout this article, we write $m = \lceil \frac{k+1}{2} \rceil$, where $\lceil \cdot \rceil$ indicates the greatest integer function.

Definition 2.4. Let $\varphi : \mathcal{A}^k \rightarrow \mathcal{B}$ be a k -linear map. Then φ is said to be

(1) local contractive, if for each $\ell \in \Omega$, there exists an $\alpha_\ell \in \Lambda$ such that

$$q_\ell(\varphi(a_1, a_2, \dots, a_k)) \leq p_{\alpha_\ell}((a_1, a_2, \dots, a_k)), \text{ for all } (a_1, a_2, \dots, a_k) \in \mathcal{A}^k.$$

(2) local completely contractive, if the map φ_n is local contractive for each $n \in \mathbb{N}$ with an additional condition that given $\ell \in \Omega$, the existence of $\alpha_\ell \in \Lambda$ is independent of n .

Definition 2.5. Let $\varphi : \mathcal{A}^k \rightarrow \mathcal{B}$ be a k -linear map and let $m = \lceil \frac{k+1}{2} \rceil$. Then φ is said to be

(1) local positive, if for every $\ell \in \Omega$, there exists an $\alpha_\ell \in \Lambda$ such that

$$\begin{aligned} \varphi(a_1, a_2, \dots, a_k) &\geq_l 0, \text{ whenever } a_p^* =_{\alpha_\ell} a_{k+1-p} \text{ for all } p \in \{1, 2, 3, \dots, m\} \text{ and} \\ \varphi(a_1, a_2, \dots, a_k) &=_l 0 \text{ whenever } a_i =_{\alpha_\ell} 0 \text{ for some } 1 \leq i \leq k, \text{ and additionally} \\ a_m &\geq_{\alpha_\ell} 0 \text{ if } k \text{ is odd.} \end{aligned}$$

(2) local completely positive, if the map φ_n is local positive for each $n \in \mathbb{N}$ with an additional condition that given $\ell \in \Omega$, the existence of $\alpha_\ell \in \Lambda$ is independent of n .

Note that if $k = 1$, then Definitions 2.4, 2.5 coincide with the notion of local completely contractive and local completely positive linear maps defined between local C^* -algebras [6]. Furthermore, if \mathcal{A}, \mathcal{B} are C^* -algebras, then Definitions 2.4, 2.5 coincide with the notion of completely contractive and completely positive k -linear maps studied in [15, 16]. Now we introduce the notion of invariance from [16] to k -linear maps on locally C^* -algebras.

Definition 2.6. A k -linear map $\varphi : \mathcal{A}^k \rightarrow \mathcal{B}$ is said to be invariant if:

(1) for $k = 2m - 1$ (odd), we have

$$\begin{aligned} \varphi(a_1 c_1, \dots, a_{m-1} c_{m-1}, a_m, a_{m+1}, \dots, a_{2m-1}) \\ = \varphi(a_1, \dots, a_{m-1}, a_m, c_{m-1} a_{m+1}, \dots, c_1 a_{2m-1}), \end{aligned}$$

for all $a_1, a_2, \dots, a_{2m-1}, c_1, \dots, c_{m-1} \in \mathcal{A}$.
(2) for $k = 2m$ (even), we have

$$\begin{aligned} & \varphi(a_1 c_1, \dots, a_m c_m, a_{m+1}, \dots, a_{2m}) \\ & = \varphi(a_1, \dots, a_m, c_m a_{m+1}, \dots, c_1 a_{2m}), \end{aligned}$$

for all $a_1, a_2, \dots, a_{2m}, c_1, \dots, c_m \in \mathcal{A}$.

Now, we fix some notations to represent such maps.

- (1) $IC C_{loc}^s(\mathcal{A}^k, \mathcal{B})$: The space of all symmetric local completely contractive invariant k -linear maps from \mathcal{A}^k to \mathcal{B} .
- (2) $IC P_{loc}^s(\mathcal{A}^k, \mathcal{B})$: The set of all symmetric local completely positive invariant k -linear maps from \mathcal{A}^k to \mathcal{B} .
- (3) $IC P C C_{loc}^s(\mathcal{A}^k, \mathcal{B})$: The set of all symmetric local completely contractive and local completely positive invariant k -linear maps from \mathcal{A}^k to \mathcal{B} .

2.1. Quantized domain and Representation of local C^* -algebras. A quantized domain in a Hilbert space \mathcal{H} is represented by a triple $\{\mathcal{H}; \mathcal{E}; \mathcal{D}\}$, where $\mathcal{E} = \{\mathcal{H}_\ell : \ell \in \Omega\}$ is a directed upward filtered family of closed subspaces such that $\mathcal{D} = \bigcup_{\ell \in \Omega} \mathcal{H}_\ell$ is dense in \mathcal{H} . Like Hilbert space, quantized domain enjoys direct sum operations: That is, if $\mathcal{H}^n, \mathcal{H}_\ell^n, \mathcal{D}^n$ are the n -direct sum copies of $\mathcal{H}, \mathcal{H}_\ell, \mathcal{D}$ respectively and $\mathcal{E}^n = \{\mathcal{H}_\ell^n : \ell \in \Omega\}$ for $n \in \mathbb{N}$, then $\{\mathcal{H}^n; \mathcal{E}^n; \mathcal{D}^n\}$ represents quantized domain in the Hilbert space \mathcal{H}^n . Let $\{\mathcal{H}^{(i)}; \mathcal{E}^{(i)}; \mathcal{D}^{(i)}\}$ be a quantized domain in Hilbert space $\mathcal{H}^{(i)}$ and $\mathcal{E}^{(i)} = \{\mathcal{H}_\ell^{(i)} : \ell \in \Omega\}$ for $i = 1, 2$. Then the notation $\mathcal{E}^{(1)} \subseteq \mathcal{E}^{(2)}$ means that $\mathcal{H}_\ell^{(1)} \subseteq \mathcal{H}_\ell^{(2)}$ for all $\ell \in \Omega$.

Let \mathcal{H}, \mathcal{K} be two Hilbert spaces. A linear operator $T : \text{dom}(T) \subseteq \mathcal{H} \rightarrow \mathcal{K}$ is called *densely defined* if $\text{dom}(T)$ is dense in \mathcal{H} . Then adjoint of T is a linear map $T^\star : \text{dom}(T^\star) \rightarrow \mathcal{H}$ with domain $\text{dom}(T^\star)$ is given by

$$\text{dom}(T^\star) = \{\xi \in \mathcal{K} : \eta \mapsto \langle T\eta, \xi \rangle_{\mathcal{K}} \text{ is continuous, for every } \eta \in \text{dom}(T)\}.$$

It implies that

$$\langle T\eta, \xi \rangle_{\mathcal{K}} = \langle \eta, T^\star \xi \rangle_{\mathcal{H}}, \text{ for all } \eta \in \text{dom}(T) \text{ and } \xi \in \text{dom}(T^\star).$$

Let $\{\mathcal{H}; \mathcal{E}; \mathcal{D}\}$ be a quantized domain in the Hilbert space \mathcal{H} where $\mathcal{E} = \{\mathcal{H}_\ell : \ell \in \Omega\}$. Let P_ℓ be the projection of \mathcal{H} onto \mathcal{H}_ℓ for each $\ell \in \Omega$ and $\mathcal{L}(\mathcal{D})$ denote the set of all linear operators on \mathcal{D} . The algebra of all *non-commutative continuous functions on a quantized domain* is defined as

$$(4) \quad C_{\mathcal{E}}^*(\mathcal{D}) = \{T \in \mathcal{L}(\mathcal{D}) : TP_\ell = P_\ell TP_\ell, P_\ell T \subseteq TP_\ell, \ell \in \Omega\}.$$

It is easy to see that $C_{\mathcal{E}}^*(\mathcal{D})$ is a $*$ -algebra, where the $*$ -operation is defined by

$$T^* := T^\star|_{\mathcal{D}}, \text{ for all } T \in C_{\mathcal{E}}^*(\mathcal{D}).$$

Further, $C_{\mathcal{E}}^*(\mathcal{D})$ is a locally C^* -algebra determined by the family $\{\|\cdot\|_\ell : \ell \in \Omega\}$ of C^* -seminorms defined by

$$\|T\|_\ell := \|T|_{\mathcal{H}_\ell}\| \text{ for all } T \in C_{\mathcal{E}}^*(\mathcal{D}).$$

In 2010, Anar Dosiev obtained GNS type representation for locally C^* -algebras [6, Theorem 7.2]. The result states that every locally C^* -algebra \mathcal{A} is identified with a $*$ -subalgebra of $C_{\mathcal{E}}^*(\mathcal{D})$ for some quantized domain $\{\mathcal{H}; \mathcal{E}; \mathcal{D}\}$. Also, the author studied the Stinespring type representation for local completely contractive and local completely positive maps $\varphi : \mathcal{A} \rightarrow C_{\mathcal{E}}^*(\mathcal{D})$. It is regarded as a structure theorem for such maps in the sense that φ can be dilated to a unital local contractive $*$ -homomorphism [6]. Detailed explanation and motivation for studying these maps can be found in [5].

3. STINESPRING TYPE REPRESENTATION

In this section, we obtain Stinespring type representation for symmetric local completely contractive and local completely positive invariant k -linear maps. Our result is a locally convex analogue of Heo's work [16] as we deal with a multilinear local positive map between local C^* -algebras. In particular, if $k = 1$ then the result coincides with [6, Theorem 5.1].

Lemma 3.1. *Let $\varphi \in \text{ICPCC}_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$ and $m = \lfloor \frac{k+1}{2} \rfloor$. Then there exist a sesquilinear form on $\mathcal{A}^{\otimes m} \otimes \mathcal{D}$ and a quantized domain $\{\mathcal{H}^\varphi; \mathcal{E}^\varphi; \mathcal{D}^\varphi\}$ associated to φ .*

Proof. Let $\mathcal{A}^{\otimes m} \otimes \mathcal{D}$ be an algebraic tensor product of $\mathcal{A}^{\otimes m}$ with \mathcal{D} , where $\mathcal{A}^{\otimes m}$ denotes the m -fold tensor product of \mathcal{A} with itself. Now we define a map $\langle \cdot, \cdot \rangle_\varphi : (\mathcal{A}^{\otimes m} \otimes \mathcal{D}) \times (\mathcal{A}^{\otimes m} \otimes \mathcal{D}) \rightarrow \mathbb{C}$ by

$$\begin{aligned} & \left\langle \sum_{j=1}^n a_{1_j} \otimes \dots \otimes a_{m_j} \otimes g_j, \sum_{i=1}^t b_{1_i} \otimes \dots \otimes b_{m_i} \otimes h_i \right\rangle_\varphi \\ & := \begin{cases} \sum_{j,i=1}^{n,t} \langle \varphi(b_{m_i}^*, \dots, b_{2_i}^*, b_{1_i}^* a_{1_j}, a_{2_j}, \dots, a_{m_j}) g_j, h_i \rangle_{\mathcal{D}} & \text{if } k = 2m - 1 \\ \sum_{j,i=1}^{n,t} \langle \varphi(b_{m_i}^*, \dots, b_{2_i}^*, b_{1_i}^*, a_{1_j}, a_{2_j}, \dots, a_{m_j}) g_j, h_i \rangle_{\mathcal{D}} & \text{if } k = 2m, \end{cases} \end{aligned}$$

for all $a_{p_j}, b_{p_i} \in \mathcal{A}; g_j, h_i \in \mathcal{D}, j \in \{1, \dots, n\}, i \in \{1, \dots, t\}$ and $p \in \{1, \dots, m\}$. Since φ is k -linear, it follows that $\langle \cdot, \cdot \rangle_\varphi$ is linear in the first variable and anti-linear in the second variable. Firstly, we show that $\langle \cdot, \cdot \rangle_\varphi$ is a positive semi-definite map. Let $a_{p_j} \in \mathcal{A}, g_j \in \mathcal{D}, p \in \{1, \dots, m\}$ and $j \in \{1, \dots, n\}$. Since \mathcal{D} is the union space of upward filtered family \mathcal{E} , there is an $\ell \in \Omega$ such that $g_j \in \mathcal{H}_\ell$ for $1 \leq j \leq n$. Since $\varphi \in \text{ICPCC}_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$ and each \mathcal{H}_ℓ is invariant under $\varphi(b_1, b_2, \dots, b_k)$ for all $b_1, b_2, \dots, b_k \in \mathcal{A}$, we have

$$\begin{aligned} & \left\langle \sum_{j=1}^n a_{1_j} \otimes \dots \otimes a_{m_j} \otimes g_j, \sum_{j=1}^n a_{1_j} \otimes \dots \otimes a_{m_j} \otimes g_j \right\rangle_\varphi \\ & = \begin{cases} \sum_{j,i=1}^n \langle \varphi(a_{m_i}^*, \dots, a_{2_i}^*, a_{1_i}^* a_{1_j}, a_{2_j}, \dots, a_{m_j}) g_j, g_i \rangle_{\mathcal{H}_\ell} & \text{if } k = 2m - 1 \\ \sum_{j,i=1}^n \langle \varphi(a_{m_i}^*, \dots, a_{2_i}^*, a_{1_i}^*, a_{1_j}, a_{2_j}, \dots, a_{m_j}) g_j, g_i \rangle_{\mathcal{H}_\ell} & \text{if } k = 2m \end{cases} \\ & = \begin{cases} \left\langle \left[\varphi(a_{m_i}^*, \dots, a_{2_i}^*, a_{1_i}^* a_{1_j}, a_{2_j}, \dots, a_{m_j}) \right]_{i,j=1}^n \begin{bmatrix} g_1 \\ \vdots \\ g_n \end{bmatrix}, \begin{bmatrix} g_1 \\ \vdots \\ g_n \end{bmatrix} \right\rangle_{\mathcal{H}_\ell^n} & \text{if } k = 2m - 1 \\ \left\langle \left[\varphi(a_{m_i}^*, \dots, a_{2_i}^*, a_{1_i}^*, a_{1_j}, a_{2_j}, \dots, a_{m_j}) \right]_{i,j=1}^n \begin{bmatrix} g_1 \\ \vdots \\ g_n \end{bmatrix}, \begin{bmatrix} g_1 \\ \vdots \\ g_n \end{bmatrix} \right\rangle_{\mathcal{H}_\ell^n} & \text{if } k = 2m \end{cases} \\ & = \begin{cases} \langle \varphi_n(D_m^*, \dots, D_2^*, R_1^* R_1, D_2, \dots, D_m) \tilde{g}, \tilde{g} \rangle_{\mathcal{H}_\ell^n} & \text{if } k = 2m - 1 \\ \langle \varphi_n(D_m^*, \dots, D_2^*, R_1^*, R_1, D_2, \dots, D_m) \tilde{g}, \tilde{g} \rangle_{\mathcal{H}_\ell^n} & \text{if } k = 2m, \end{cases} \end{aligned}$$

where

$$(5) \quad \tilde{g} = \begin{bmatrix} g_1 \\ \vdots \\ g_n \end{bmatrix}, D_p = \begin{bmatrix} a_{p_1} & 0 & \dots & 0 \\ 0 & a_{p_2} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & a_{p_n} \end{bmatrix} \text{ and } R_1 = \begin{bmatrix} a_{1_1} & a_{1_2} \dots & a_{1_n} \\ 0 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 0 \end{bmatrix},$$

for $2 \leq p \leq m$. If $k = 2m - 1$, then

$$(D_m^*, \dots, D_2^*, R_1^* R_1, D_2, \dots, D_m)^* = (D_m^*, \dots, D_2^*, R_1^* R_1, D_2, \dots, D_m) \text{ and } R_1^* R_1 \geq_\alpha 0,$$

for every $\alpha \in \Lambda$. If $k = 2m$, then

$$(D_m^*, \dots, D_2^*, R_1^*, R_1, D_2, \dots, D_m)^* = (D_m^*, \dots, D_2^*, R_1^*, R_1, D_2, \dots, D_m).$$

Since φ is a local completely positive map, it follows that

$$\left\langle \sum_{j=1}^n a_{1_j} \otimes \dots \otimes a_{m_j} \otimes g_j, \sum_{j=1}^n a_{1_j} \otimes \dots \otimes a_{m_j} \otimes g_j \right\rangle_\varphi \geq 0.$$

Now by using the symmetric nature of φ , we show that $\langle \cdot, \cdot \rangle_\varphi$ satisfies Hermitian property. Let $a_{p_j}, b_{p_i} \in \mathcal{A}$, $g_j, h_i \in \mathcal{D}$ for $j \in \{1, \dots, n\}$, $i \in \{1, \dots, t\}$ and $p \in \{1, \dots, m\}$. Then

$$\begin{aligned} & \left\langle \sum_{j=1}^n a_{1_j} \otimes \dots \otimes a_{m_j} \otimes g_j, \sum_{i=1}^t b_{1_i} \otimes \dots \otimes b_{m_i} \otimes h_i \right\rangle_\varphi \\ &= \begin{cases} \sum_{j,i=1}^{n,t} \left\langle \varphi(b_{m_i}^*, \dots, b_{2_i}^*, b_{1_i}^* a_{1_j}, a_{2_j}, \dots, a_{m_j}) g_j, h_i \right\rangle_{\mathcal{H}_\ell} & \text{if } k = 2m - 1 \\ \sum_{j,i=1}^{n,t} \left\langle \varphi(b_{m_i}^*, \dots, b_{2_i}^*, b_{1_i}^*, a_{1_j}, a_{2_j}, \dots, a_{m_j}) g_j, h_i \right\rangle_{\mathcal{H}_\ell} & \text{if } k = 2m \end{cases} \\ &= \begin{cases} \sum_{j,i=1}^{n,t} \left\langle g_j, \varphi(b_{m_i}^*, \dots, b_{2_i}^*, b_{1_i}^* a_{1_j}, a_{2_j}, \dots, a_{m_j})^* h_i \right\rangle_{\mathcal{H}_\ell} & \text{if } k = 2m - 1 \\ \sum_{j,i=1}^{n,t} \left\langle g_j, \varphi(b_{m_i}^*, \dots, b_{2_i}^*, b_{1_i}^*, a_{1_j}, a_{2_j}, \dots, a_{m_j})^* h_i \right\rangle_{\mathcal{H}_\ell} & \text{if } k = 2m \end{cases} \\ &= \begin{cases} \sum_{j,i=1}^{n,t} \left\langle g_j, \varphi(a_{m_j}^*, \dots, a_{2_j}^*, a_{1_j}^* b_{1_i}, b_{2_i}, \dots, b_{m_i}) h_i \right\rangle_{\mathcal{H}_\ell} & \text{if } k = 2m - 1 \\ \sum_{j,i=1}^{n,t} \left\langle g_j, \varphi(a_{m_j}^*, \dots, a_{2_j}^*, a_{1_j}^*, b_{1_i}, b_{2_i}, \dots, b_{m_i}) h_i \right\rangle_{\mathcal{H}_\ell} & \text{if } k = 2m \end{cases} \\ &= \begin{cases} \sum_{j,i=1}^{n,t} \overline{\left\langle \varphi(a_{m_j}^*, \dots, a_{2_j}^*, a_{1_j}^* b_{1_i}, b_{2_i}, \dots, b_{m_i}) h_i, g_j \right\rangle_{\mathcal{H}_\ell}} & \text{if } k = 2m - 1 \\ \sum_{j,i=1}^{n,t} \overline{\left\langle \varphi(a_{m_j}^*, \dots, a_{2_j}^*, a_{1_j}^*, b_{1_i}, b_{2_i}, \dots, b_{m_i}) h_i, g_j \right\rangle_{\mathcal{H}_\ell}} & \text{if } k = 2m \end{cases} \\ &= \left\langle \sum_{i=1}^t b_{1_i} \otimes \dots \otimes b_{m_i} \otimes h_i, \sum_{j=1}^n a_{1_j} \otimes \dots \otimes a_{m_j} \otimes g_j \right\rangle_\varphi. \end{aligned}$$

Thus $\langle \cdot, \cdot \rangle_\varphi$ is a semi-inner product on $\mathcal{A}^{\otimes m} \otimes \mathcal{D}$. To prove the second part of the claim, let us define $\mathcal{N}_\varphi := \{\xi \in \mathcal{A}^{\otimes m} \otimes \mathcal{D} : \langle \xi, \xi \rangle_\varphi = 0\}$. Then by the Cauchy-Schwartz inequality of semi-inner product $\langle \cdot, \cdot \rangle_\varphi$ on $\mathcal{A}^{\otimes m} \otimes \mathcal{D}$, we have $\mathcal{N}_\varphi = \{\xi \in \mathcal{A}^{\otimes m} \otimes \mathcal{D} : \langle \xi, \eta \rangle_\varphi = 0, \text{ for every } \eta \in \mathcal{A}^{\otimes m} \otimes \mathcal{D}\}$. This shows

that \mathcal{N}_φ is a closed subspace of $\mathcal{A}^{\otimes m} \otimes \mathcal{D}$. The map $\langle \cdot, \cdot \rangle$ defined by

$$\langle \xi + \mathcal{N}_\varphi, \xi' + \mathcal{N}_\varphi \rangle := \langle \xi, \xi' \rangle_\varphi, \text{ for every } \xi, \xi' \in \mathcal{A}^{\otimes m} \otimes \mathcal{D}$$

is a sesquilinear form on $\mathcal{A}^{\otimes m} \otimes \mathcal{D}$. Let \mathcal{H}^φ be the Hilbert space obtained by the completion of $(\mathcal{A}^{\otimes m} \otimes \mathcal{D})/\mathcal{N}_\varphi$. Similarly, for each $\ell \in \Omega$ we define the Hilbert space \mathcal{H}_ℓ^φ as,

$$(6) \quad \mathcal{H}_\ell^\varphi = \overline{\text{span}}\{a_1 \otimes \dots \otimes a_m \otimes g + \mathcal{N}_\varphi : a_p \in \mathcal{A}, 1 \leq p \leq m, g \in \mathcal{H}_\ell\}.$$

Since \mathcal{E} is an upward filtered family, it follows from (6) that $\mathcal{E}^\varphi = \{\mathcal{H}_\ell^\varphi : \ell \in \Omega\}$ is an upward filtered family of closed subspaces of \mathcal{H}^φ . It remains to show that $\mathcal{D}^\varphi = \bigcup_{\ell \in \Omega} \mathcal{H}_\ell^\varphi$ is dense in \mathcal{H}^φ . Clearly,

$\overline{\mathcal{D}^\varphi} \subseteq \mathcal{H}^\varphi$. Suppose there is a non-zero $\eta \in (\overline{\mathcal{D}^\varphi})^\perp$, then there exists an $n_0 \in \mathbb{N}$ such that $\frac{1}{n_0} < \|\eta\|^2$.

Since $(\overline{\mathcal{A}^{\otimes m} \otimes \mathcal{D}})/\mathcal{N}_\varphi = \mathcal{H}^\varphi$, there exists a $\gamma \in \mathcal{A}^{\otimes m} \otimes \mathcal{D}/\mathcal{N}_\varphi$ of the form $\gamma = \sum_{j=1}^t a_{1j} \otimes \dots \otimes a_{mj} \otimes g_j + \mathcal{N}_\varphi$

such that $\|\eta - \gamma\|^2 < \frac{1}{n_0}$. Without loss of generality, for every $1 \leq j \leq t$ we have $g_j \in \mathcal{H}_\ell$ for some $\ell \in \Omega$. It follows that $\gamma \in \mathcal{H}_\ell^\varphi \subseteq \mathcal{D}^\varphi$ and hence $\langle \eta, \gamma \rangle = 0$. Therefore,

$$\|\eta\|^2 \leq \|\eta\|^2 + \|\gamma\|^2 = \|\eta - \gamma\|^2 < \frac{1}{n_0}.$$

This contradicts the fact that $\frac{1}{n_0} < \|\eta\|^2$. Thus \mathcal{D}^φ is dense in \mathcal{H}^φ . As a result, $\{\mathcal{H}^\varphi; \mathcal{E}^\varphi; \mathcal{D}^\varphi\}$ is a quantized domain associated to φ . \square

Next, we obtain Stinespring type representation for symmetric local completely contractive and local completely positive invariant k -linear maps. The original construction of the Stinespring theorem for CP-maps on C^* -algebras can be found in [23].

Theorem 3.2. *Let $\varphi \in ICPC C_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$. Then there is a quantized domain $\{\mathcal{H}^\varphi; \mathcal{E}^\varphi; \mathcal{D}^\varphi\}$, a contraction $V_\varphi \in \mathcal{B}(\mathcal{H}, \mathcal{H}^\varphi)$ and a commuting family $\{\pi_p^\varphi : 1 \leq p \leq m\}$ of unital local contractive $*$ -homomorphisms from \mathcal{A} to $C_{\mathcal{E}^\varphi}^*(\mathcal{D}^\varphi)$ such that $V_\varphi(\mathcal{E}) \subseteq \mathcal{E}^\varphi$ and*

(i) if $k = 2m - 1$ odd, then

$$\varphi(a_1, \dots, a_{m-1}, a_m, a_{m+1}, \dots, a_{2m-1}) \subseteq V_\varphi^* \pi_1^\varphi(a_m) \pi_2^\varphi(a_{m-1} a_{m+1}) \cdots \pi_m^\varphi(a_1 a_{2m-1}) V_\varphi,$$

for all $a_1, a_2, \dots, a_{2m-1} \in \mathcal{A}$.

(ii) if $k = 2m$ even, then

$$\varphi(a_1, \dots, a_{m-1}, a_m, a_{m+1}, \dots, a_{2m}) \subseteq V_\varphi^* \pi_1^\varphi(a_m a_{m+1}) \pi_2^\varphi(a_{m-1} a_{m+2}) \cdots \pi_m^\varphi(a_1 a_{2m}) V_\varphi,$$

for all $a_1, a_2, \dots, a_{2m} \in \mathcal{A}$.

Proof. By Lemma 3.1, there is a quantized domain $\{\mathcal{H}^\varphi; \mathcal{E}^\varphi; \mathcal{D}^\varphi\}$ associated to φ . Recall that \mathcal{H}^φ is obtained by the completion of the quotient space $(\mathcal{A}^{\otimes m} \otimes \mathcal{D})/\mathcal{N}_\varphi$. For $1 \leq p \leq m$ and $a \in \mathcal{A}$, we define a linear map $\pi_p^\varphi(a) : (\mathcal{A}^{\otimes m} \otimes \mathcal{D})/\mathcal{N}_\varphi \rightarrow (\mathcal{A}^{\otimes m} \otimes \mathcal{D})/\mathcal{N}_\varphi$ by

$$(7) \quad \begin{aligned} \pi_p^\varphi(a) \left(\sum_{j=1}^n a_{1j} \otimes \dots \otimes a_{pj} \otimes \dots \otimes a_{mj} \otimes g_j + \mathcal{N}_\varphi \right) \\ = \sum_{j=1}^n a_{1j} \otimes \dots \otimes a a_{pj} \otimes \dots \otimes a_{mj} \otimes g_j + \mathcal{N}_\varphi \end{aligned}$$

for all $a_{pj} \in \mathcal{A}$, $g_j \in \mathcal{D}$, $j \in \{1, \dots, n\}$. Now we show that π_p^φ is a well defined linear map. We can choose $\ell \in \Omega$ with $g_j \in \mathcal{H}_\ell$ for $j \in \{1, \dots, n\}$. Then

$$\begin{aligned}
& \left\| \pi_p^\varphi(a) \left(\sum_{j=1}^n a_{1_j} \otimes \dots \otimes a_{p_j} \otimes \dots \otimes a_{m_j} \otimes g_j + \mathcal{N}_\varphi \right) \right\|_{\mathcal{H}_\ell^\varphi}^2 \\
&= \left\| \sum_{i=1}^n a_{1_j} \otimes \dots \otimes a_{p_j} \otimes \dots \otimes a_{m_j} \otimes g_j + \mathcal{N}_\varphi \right\|_{\mathcal{H}_\ell^\varphi}^2 \\
&= \begin{cases} \sum_{i,j=1}^n \langle \varphi(a_{m_i}^*, \dots, a_{p_i}^* a^*, \dots, a_{2_i}^*, a_{1_i}^* a_{1_j}, \dots, a_{p_j}, \dots, a_{m_j}) g_j, g_i \rangle_{\mathcal{H}_\ell} & \text{if } k = 2m - 1 \\ \sum_{i,j=1}^n \langle \varphi(a_{m_i}^*, \dots, a_{p_i}^* a^*, \dots, a_{2_i}^*, a_{1_i}^*, a_{1_j}, \dots, a_{p_j}, \dots, a_{m_j}) g_j, g_i \rangle_{\mathcal{H}_\ell} & \text{if } k = 2m \end{cases} \\
(8) \quad &= \begin{cases} \langle \varphi_n(D_m^*, \dots, D_p^*(a^* \otimes I_n), \dots, R_1^* R_1, \dots, (a \otimes I_n) D_p, \dots, D_m) \tilde{g}, \tilde{g} \rangle_{\mathcal{H}_\ell^n} & \text{if } k = 2m - 1 \\ \langle \varphi_n(D_m^*, \dots, D_p^*(a^* \otimes I_n), \dots, R_1^*, R_1, \dots, (a \otimes I_n) D_p, \dots, D_m) \tilde{g}, \tilde{g} \rangle_{\mathcal{H}_\ell^n} & \text{if } k = 2m, \end{cases}
\end{aligned}$$

where \tilde{g}, R_1 and D_p for $2 \leq p \leq m$ are defined as in Equation (5). Since φ is a local completely positive map, for given $\ell \in \Omega$, there exists an $\alpha_\ell \in \Lambda$ such that $\varphi_n(B_1, B_2, \dots, B_k) \geq_\ell 0$ whenever $(B_1, B_2, \dots, B_k) =_{\alpha_\ell} (B_k^*, \dots, B_2^*, B_1^*)$ and $\varphi_n(B_1, B_2, \dots, B_k) =_\ell 0$ whenever $B_i =_{\alpha_\ell} 0$ for some $1 \leq i \leq k$, and additionally $B_m \geq_{\alpha_\ell} 0$ if $k = 2m - 1$ is odd. Let us consider the case $k = 2m - 1$. Notice that $\pi_{\alpha_\ell}(p_{\alpha_\ell}(a)^2 I - a^* a)$ is positive in $\mathcal{A}_{\alpha_\ell}$, thus there exists an element $b \in \mathcal{A}$ such that $\pi_{\alpha_\ell}(p_{\alpha_\ell}(a)^2 I - a^* a) = \pi_{\alpha_\ell}(b^* b)$. Therefore $p_{\alpha_\ell}(a)^2 I - a^* a = b^* b + c$ for some $c \in \mathcal{N}_{\alpha_\ell}$. As $c =_{\alpha_\ell} 0$, we have $(c \otimes I_n) D_p =_{\alpha_\ell} 0$. Applying these facts on φ_n , notice that $\varphi_n(D_m^*, \dots, D_p^*, \dots, D_2^*, R_1^* R_1, D_2, \dots, (c \otimes I_n) D_p, \dots, D_m) =_\ell 0$. Applying this argument as well as the invariant property of φ carefully, we obtain

$$\begin{aligned}
& \left\langle \varphi_n(D_m^*, \dots, D_p^*, \dots, D_2^*, R_1^* R_1, D_2, \dots, ((p_{\alpha_\ell}(a)^2 I - a^* a) \otimes I_n) D_p, \dots, D_m) \tilde{g}, \tilde{g} \right\rangle_{\mathcal{H}_\ell^n} \\
&= \left\langle \varphi_n(D_m^*, \dots, D_p^*, \dots, D_2^*, R_1^* R_1, D_2, \dots, (b^* b \otimes I_n) D_p, \dots, D_m) \tilde{g}, \tilde{g} \right\rangle_{\mathcal{H}_\ell^n} \\
&= \left\langle \varphi_n(D_m^*, \dots, D_p^*(b^* \otimes I_n), \dots, D_2^*, R_1^* R_1, D_2, \dots, (b \otimes I_n) D_p, D_m) \tilde{g}, \tilde{g} \right\rangle_{\mathcal{H}_\ell^n} \\
&\geq 0.
\end{aligned}$$

This implies that

$$\begin{aligned}
& \left\langle \varphi_n(D_m^*, \dots, D_p^*, \dots, D_2^*, R_1^* R_1, D_2, \dots, (a^* a \otimes I_n) D_p, \dots, D_m) \tilde{g}, \tilde{g} \right\rangle_{\mathcal{H}_\ell^n} \\
&\leq p_{\alpha_\ell}(a)^2 \left\langle \varphi_n(D_m^*, \dots, D_p^*, \dots, D_2^*, R_1^* R_1, D_2, \dots, D_p, \dots, D_m) \tilde{g}, \tilde{g} \right\rangle_{\mathcal{H}_\ell^n} \\
&= p_{\alpha_\ell}(a)^2 \sum_{i,j=1}^n \langle \varphi(a_{m_i}^*, \dots, a_{2_i}^*, a_{1_i}^* a_{1_j}, a_{2_j}, \dots, a_{m_j}) g_j, g_i \rangle_{\mathcal{H}_\ell} \\
&= p_{\alpha_\ell}(a)^2 \left\| \sum_{j=1}^n a_{1_j} \otimes a_{2_j} \otimes \dots \otimes a_{m_j} \otimes g_j + \mathcal{N}_\varphi \right\|_{\mathcal{H}_\ell^\varphi}^2
\end{aligned}$$

Similarly, the above inequality holds true for $k = 2m$. From Equation (8), we conclude that

$$(9) \quad \|\pi_p^\varphi(a)\|_{\mathcal{H}_\ell^\varphi} \leq p_{\alpha_\ell}(a); \text{ for every } a \in \mathcal{A}, \ell \in \Omega.$$

Equivalently, $\pi_p^\varphi : \mathcal{A} \rightarrow C_{\mathcal{E}^\varphi}^*(\mathcal{D}^\varphi)$ is a local contractive map. Next we show that $a \mapsto \pi_p^\varphi(a)$ is a unital *-homomorphism. Let $a_p, b_{p_i} \in \mathcal{A}$ and $g_j, h_i \in \mathcal{H}_\ell$ for $p \in \{1, \dots, m\}, j \in \{1, \dots, t\}, i \in \{1, \dots, n\}$. Then by using invariant property of φ , we get

$$\begin{aligned}
& \left\langle \pi_p^\varphi(a)^* \left(\sum_{j=1}^t a_{1_j} \otimes \dots \otimes a_{p_j} \otimes \dots \otimes a_{m_j} \otimes g_j + \mathcal{N}_\varphi \right), \sum_{i=1}^n b_{1_i} \otimes \dots \otimes b_{p_i} \otimes \dots \otimes b_{m_i} \otimes h_i + \mathcal{N}_\varphi \right\rangle_{\mathcal{H}_\ell^\varphi} \\
&= \left\langle \sum_{j=1}^t a_{1_j} \otimes \dots \otimes a_{p_j} \otimes \dots \otimes a_{m_j} \otimes g_j + \mathcal{N}_\varphi, \pi_p^\varphi(a) \left(\sum_{i=1}^n b_{1_i} \otimes \dots \otimes b_{p_i} \otimes \dots \otimes b_{m_i} \otimes h_i + \mathcal{N}_\varphi \right) \right\rangle_{\mathcal{H}_\ell^\varphi} \\
&= \begin{cases} \sum_{i,j=1}^{n,t} \left\langle \varphi(b_{m_i}^*, \dots, b_{p_i}^* a^*, \dots, b_{1_i}^* a_{1_j}, \dots, a_{p_j}, \dots, a_{m_j}) g_j, h_i \right\rangle_{\mathcal{H}_\ell} & \text{if } k = 2m - 1 \\ \sum_{i,j=1}^{n,t} \left\langle \varphi(b_{m_i}^*, \dots, b_{p_i}^* a^*, \dots, b_{1_i}^*, a_{1_j}, \dots, a_{p_j}, \dots, a_{m_j}) g_j, h_i \right\rangle_{\mathcal{H}_\ell} & \text{if } k = 2m \end{cases} \\
&= \begin{cases} \sum_{i,j=1}^{n,t} \left\langle \varphi(b_{m_i}^*, \dots, b_{p_i}^*, \dots, b_{1_i}^* a_{1_j}, \dots, a^* a_{p_j}, \dots, a_{m_j}) g_j, h_i \right\rangle_{\mathcal{H}_\ell} & \text{if } k = 2m - 1 \\ \sum_{i,j=1}^{n,t} \left\langle \varphi(b_{m_i}^*, \dots, b_{p_i}^*, \dots, b_{1_i}^*, a_{1_j}, \dots, a^* a_{p_j}, \dots, a_{m_j}) g_j, h_i \right\rangle_{\mathcal{H}_\ell} & \text{if } k = 2m \end{cases} \\
&= \left\langle \pi_p^\varphi(a^*) \left(\sum_{j=1}^t a_{1_j} \otimes \dots \otimes a_{p_j} \otimes \dots \otimes a_{m_j} \otimes g_j + \mathcal{N}_\varphi \right), \sum_{i=1}^n b_{1_i} \otimes \dots \otimes b_{p_i} \otimes \dots \otimes b_{m_i} \otimes h_i + \mathcal{N}_\varphi \right\rangle_{\mathcal{H}_\ell^\varphi}
\end{aligned}$$

It follows that $\pi_p^\varphi(a)^*|_{\mathcal{H}_\ell^\varphi} = \pi_p^\varphi(a^*)|_{\mathcal{H}_\ell^\varphi}$, for every $\ell \in \Omega$ and hence $\pi_p^\varphi(a)^* = \pi_p^\varphi(a^*)$ for all $a \in \mathcal{A}$. It implies that $\pi_p : \mathcal{A} \rightarrow C_{\mathcal{E}^\varphi}^*(\mathcal{D}^\varphi)$ is a $*$ -homomorphism and it is local contractive by Equation (9).

Now it remains to show that $\{\pi_p^\varphi\}_{p=1}^m$ is a commuting family. Let $1 \leq p < q \leq m$. Then

$$\begin{aligned}
& \pi_p^\varphi(a) \pi_q^\varphi(b) \left(\sum_{j=1}^t a_{1_j} \otimes \dots \otimes a_{m_j} \otimes g_j + \mathcal{N}_\varphi \right) \\
&= \sum_{j=1}^t a_{1_j} \otimes \dots \otimes a a_{p_j} \otimes \dots \otimes b a_{q_j} \otimes \dots \otimes a_{m_j} \otimes g_j + \mathcal{N}_\varphi \\
&= \pi_q^\varphi(b) \pi_p^\varphi(a) \left(\sum_{j=1}^t a_{1_j} \otimes \dots \otimes a_{m_j} \otimes g_j + \mathcal{N}_\varphi \right)
\end{aligned}$$

Hence by continuity, we have $\pi_p^\varphi(a) \pi_q^\varphi(b)|_{\mathcal{H}_\ell^\varphi} = \pi_q^\varphi(b) \pi_p^\varphi(a)|_{\mathcal{H}_\ell^\varphi}$ as both the maps agree on the dense subspace of \mathcal{H}_ℓ^φ . Since this is true for all $\ell \in \Omega$, we conclude that $\pi_p^\varphi(a) \pi_q^\varphi(b) = \pi_q^\varphi(b) \pi_p^\varphi(a)$.

Now we define a map $V_\varphi : \mathcal{H} \rightarrow \mathcal{H}^\varphi$ by $V_\varphi(g) = (1 \otimes \dots \otimes 1 \otimes g) + \mathcal{N}_\varphi$ for all $g \in \mathcal{H}$. Note that $V_\varphi(\mathcal{H}_\ell) \subseteq \mathcal{H}_\ell^\varphi$ for each $\ell \in \Omega$. That is, $V_\varphi(\mathcal{E}) \subseteq \mathcal{E}^\varphi$. Since φ is a local contractive map, for $g \in \mathcal{H}_\ell$, we see that V_φ is a contractive as follows:

$$\begin{aligned}
\|V_\varphi g\|_{\mathcal{H}_\ell^\varphi}^2 &= \left\langle (1 \otimes \dots \otimes 1 \otimes g) + \mathcal{N}_\varphi, (1 \otimes \dots \otimes 1 \otimes g) + \mathcal{N}_\varphi \right\rangle_{\mathcal{H}_\ell^\varphi} \\
&= \langle \varphi(1, \dots, 1)g, g \rangle_{\mathcal{H}_\ell} \\
&\leq \|\varphi(1, \dots, 1)g\|_{\mathcal{H}_\ell} \|g\|_{\mathcal{H}_\ell} \\
&\leq \|g\|_{\mathcal{H}_\ell}^2.
\end{aligned}$$

So V_φ is a contractive map on the dense subspace \mathcal{D}^φ and hence it is contractive on \mathcal{H}^φ . Thus $V_\varphi : \mathcal{H} \rightarrow \mathcal{H}^\varphi$ is a well defined contractive map. Moreover V_φ is an isometry when φ is unital. Suppose $k = 2m - 1$, $a_1, a_2, \dots, a_{2m-1} \in \mathcal{A}$ and $g, h \in \mathcal{H}_\ell$ for some $\ell \in \Omega$, we have

$$\left\langle V_\varphi^* \pi_1^\varphi(a_m) \pi_2^\varphi(a_{m-1} a_{m+1}) \cdots \pi_m^\varphi(a_1 a_{2m-1}) V_\varphi h, g \right\rangle_{\mathcal{H}_\ell}$$

$$\begin{aligned}
&= \left\langle \pi_1^\varphi(a_m) \pi_2^\varphi(a_{m-1} a_{m+1}) \cdots \pi_m^\varphi(a_1 a_{2m-1}) V_\varphi h, V_\varphi g \right\rangle_{\mathcal{H}_\ell} \\
&= \left\langle (a_m \otimes a_{m-1} a_{m+1} \otimes \cdots \otimes a_1 a_{2m-1} \otimes h) + \mathcal{N}_\varphi, (1 \otimes \cdots \otimes 1 \otimes g) + \mathcal{N}_\varphi \right\rangle_{\mathcal{H}_\ell^\varphi} \\
&= \left\langle \varphi(1, 1, \dots, a_m, a_{m-1} a_{m+1}, \dots, a_1 a_{2m-1}) h, g \right\rangle_{\mathcal{H}_\ell} \\
&= \left\langle \varphi(a_1, a_2, \dots, a_m, \dots, a_{2m-1}) h, g \right\rangle_{\mathcal{H}_\ell}
\end{aligned}$$

since φ is invariant map. Therefore,

$$\varphi(a_1, \dots, a_{m-1}, a_m, a_{m+1}, \dots, a_{2m-1}) \subseteq V_\varphi^* \pi_1^\varphi(a_m) \pi_2^\varphi(a_{m-1} a_{m+1}) \cdots \pi_m^\varphi(a_1 a_{2m-1}) V_\varphi,$$

for all $a_1, \dots, a_{2m-1} \in \mathcal{A}$. Similarly, the result holds for the case $k = 2m$. \square

Note 3.3. For a given $\varphi \in ICPC C_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$, we denote a Stinespring's triple associated to φ by $(\{\pi_p^\varphi\}_{p=1}^m, V_\varphi, \{\mathcal{H}^\varphi; \mathcal{E}^\varphi; \mathcal{D}^\varphi\})$, where the terms are defined as in Theorem 3.2.

4. MINIMALITY CONDITION

In this section, we identify a suitable notion of minimality for Stinespring's type representation so as to ensure the uniqueness up to unitary equivalence. In fact, W. Arveson was the first mathematician to construct a Radon-Nikodým derivative between two CP-maps [2] by using minimal Stinespring's representation. Here, we discuss the minimality condition of the Stinespring theorem for invariant multilinear local CP-maps and apply this to prove Radon-Nikodým theorem in the next section. Before that we fix some notations for simplicity: $[M] := \overline{\text{span}} M$, for any subset M of a Hilbert space;

$$\begin{aligned}
&\prod_{p=1}^m \pi_p^\varphi(a_{p_j}) := \pi_1^\varphi(a_{1_j}) \pi_2^\varphi(a_{2_j}) \cdots \pi_m^\varphi(a_{m_j}); \text{ and} \\
&\left(\prod_{p=1}^m \pi_p^\varphi(\mathcal{A}) \right) V_\varphi \mathcal{D} := \left\{ \prod_{p=1}^m \pi_p^\varphi(a_{p_j}) V_\varphi h_j : a_{1_j}, \dots, a_{m_j} \in \mathcal{A}, h_j \in \mathcal{D} \right\}.
\end{aligned}$$

The following lemma is a key observation to establish the minimality condition.

Lemma 4.1. Let $\varphi \in ICPC C_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$ and let $(\{\pi_p^\varphi\}_{p=1}^m, V_\varphi, \{\mathcal{H}^\varphi; \mathcal{E}^\varphi; \mathcal{D}^\varphi\})$ be a Stinespring triple of the map φ . Then

$$(10) \quad \bigcup_{\ell \in \Omega} \overline{\left[\left(\prod_{p=1}^m \pi_p^\varphi(\mathcal{A}) \right) V_\varphi \mathcal{H}_\ell \right]} = \overline{\left[\left(\prod_{p=1}^m \pi_p^\varphi(\mathcal{A}) \right) V_\varphi \mathcal{D} \right]}$$

Proof. The proof follows immediately from the fact that $\bigcup_{\ell \in \Omega} \mathcal{H}_\ell = \mathcal{D}$. \square

Proposition 4.2. Let $\varphi \in ICPC C_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$ and let $(\{\pi_p^\varphi\}_{p=1}^m, V_\varphi, \{\mathcal{H}^\varphi; \mathcal{E}^\varphi; \mathcal{D}^\varphi\})$ be a Stinespring triple of φ . Then one can construct another Stinespring triple $(\{\tilde{\pi}_p^\varphi\}_{p=1}^m, \tilde{V}_\varphi, \{\tilde{\mathcal{H}}^\varphi; \tilde{\mathcal{E}}^\varphi; \tilde{\mathcal{D}}^\varphi\})$ with the following additional properties

$$(11) \quad \tilde{\mathcal{E}}^\varphi \subseteq \mathcal{E}^\varphi \text{ and } \tilde{\mathcal{H}}_\ell^\varphi = \overline{\left[\left(\prod_{p=1}^m \tilde{\pi}_p^\varphi(\mathcal{A}) \right) \tilde{V}_\varphi \mathcal{H}_\ell \right]} \text{ for every } \ell \in \Omega.$$

Proof. First, we define a family of Hilbert spaces by $\tilde{\mathcal{H}}_\ell^\varphi := \overline{\left[\left(\prod_{p=1}^m \pi_p^\varphi(\mathcal{A}) \right) V_\varphi \mathcal{H}_\ell \right]}$ for each $\ell \in \Omega$ and

$\tilde{\mathcal{H}}^\varphi := \overline{\left[\left(\prod_{p=1}^m \pi_p^\varphi(\mathcal{A}) \right) V_\varphi \mathcal{D} \right]}$. Also take $\tilde{\mathcal{E}}^\varphi := \{\tilde{\mathcal{H}}_\ell^\varphi : \ell \in \Omega\}$ and $\tilde{\mathcal{D}}^\varphi := \bigcup_{\ell \in \Omega} \tilde{\mathcal{H}}_\ell^\varphi$. Then by Lemma 4.1, it

follows that $\{\widetilde{\mathcal{H}}^\varphi; \widetilde{\mathcal{E}}^\varphi; \widetilde{\mathcal{D}}^\varphi\}$ is a quantized domain. Since $\widetilde{\mathcal{H}}_\ell^\varphi \subseteq \mathcal{H}_\ell^\varphi$ for each $\ell \in \Omega$, we have $\widetilde{\mathcal{E}}^\varphi \subseteq \mathcal{E}^\varphi$. For each $1 \leq p \leq m$, we define a map $\widetilde{\pi}_p^\varphi : \mathcal{A} \rightarrow C_{\widetilde{\mathcal{E}}^\varphi}^*(\widetilde{\mathcal{D}}^\varphi)$ by

$$(12) \quad \widetilde{\pi}_p^\varphi(a) = \pi_p^\varphi(a)|_{\widetilde{\mathcal{E}}^\varphi}.$$

Suppose $\sum_{j=1}^n \prod_{p=1}^m \pi_p^\varphi(a_{p_j})V_\varphi h_j \in \text{Span}\{\prod_{p=1}^m \pi_p^\varphi(\mathcal{A})V_\varphi \mathcal{H}_\ell\}$, where $a_{p_j} \in \mathcal{A}$, and $h_j \in \mathcal{H}_\ell$ for $1 \leq j \leq n$. Then by using the fact that the family $\{\pi_p^\varphi\}_{p=1}^m$ is commutative, we see that

$$\pi_q^\varphi(a) \left(\sum_{j=1}^n \prod_{p=1}^m \pi_p^\varphi(a_{p_j})V_\varphi h_j \right) = \sum_{j=1}^n \pi_1^\varphi(a_{1_j}) \cdots \pi_q^\varphi(a_{q_j}) \cdots \pi_m^\varphi(a_{m_j})V_\varphi h_j \in \widetilde{\mathcal{H}}_\ell^\varphi$$

for each $1 \leq q \leq m$. It follows that

$$\pi_p^\varphi(a) (\text{Span}\{\prod_{p=1}^m \pi_p^\varphi(\mathcal{A})V_\varphi \mathcal{H}^\varphi\}) \subseteq \widetilde{\mathcal{H}}_\ell^\varphi \text{ for every } 1 \leq p \leq m.$$

Since $\pi_p^\varphi(a)|_{\widetilde{\mathcal{H}}_\ell^\varphi}$ is continuous, we have $\pi_p^\varphi(a)(\widetilde{\mathcal{H}}_\ell^\varphi) \subseteq \widetilde{\mathcal{H}}_\ell^\varphi$. Now from the definition of $\widetilde{\pi}_p^\varphi$ we conclude that $\widetilde{\pi}_p^\varphi(a)(\widetilde{\mathcal{H}}_\ell^\varphi) \subseteq \widetilde{\mathcal{H}}_\ell^\varphi$ for all $a \in \mathcal{A}$ and for each $\ell \in \Omega$. It follows that $\widetilde{\pi}_p^\varphi(a) \in C_{\widetilde{\mathcal{E}}^\varphi}^*(\widetilde{\mathcal{D}}^\varphi)$. Moreover, $\{\widetilde{\pi}_p^\varphi\}$ is a commuting family and hence $(\{\widetilde{\pi}_p^\varphi\}_{p=1}^m, \widetilde{V}_\varphi, \{\widetilde{\mathcal{H}}^\varphi; \widetilde{\mathcal{E}}^\varphi; \widetilde{\mathcal{D}}^\varphi\})$ is a Stinespring's triple for φ follows from Equation (12). \square

Definition 4.3. The Stinespring triple $(\{\pi_p^\varphi\}_{p=1}^m, V_\varphi, \{\mathcal{H}^\varphi; \mathcal{E}^\varphi; \mathcal{D}^\varphi\})$ of $\varphi \in \text{ICPCC}_{\text{loc}}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$ is called minimal, if

$$\mathcal{H}_\ell^\varphi = \left[\prod_{p=1}^m \pi_p^\varphi(\mathcal{A})V_\varphi \mathcal{H}_\ell \right], \text{ for each } \ell \in \Omega.$$

Now, our aim is to show that the minimality condition in the Definition 4.3 is appropriate in the sense that there is a unitary equivalence relation between any two minimal Stinespring representations.

Theorem 4.4. Let $(\{\widetilde{\pi}_p^\varphi\}_{p=1}^m, \widetilde{V}_\varphi, \{\widetilde{\mathcal{H}}^\varphi; \widetilde{\mathcal{E}}^\varphi; \widetilde{\mathcal{D}}^\varphi\})$ and $(\{\widehat{\pi}_p^\varphi\}_{p=1}^m, \widehat{V}_\varphi, \{\widehat{\mathcal{H}}^\varphi; \widehat{\mathcal{E}}^\varphi; \widehat{\mathcal{D}}^\varphi\})$ be two minimal Stinespring triples of the map $\varphi \in \text{ICPCC}_{\text{loc}}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$. Then there exists a unitary operator $U_\varphi : \widetilde{\mathcal{H}}^\varphi \rightarrow \widehat{\mathcal{H}}^\varphi$ such that

- (1) $U_\varphi \widetilde{V}_\varphi = \widehat{V}_\varphi$ and
- (2) $U_\varphi \prod_{p=1}^m \widetilde{\pi}_p^\varphi(a_p) = \prod_{p=1}^m \widehat{\pi}_p^\varphi(a_p)U_\varphi$ for every $a_p \in \mathcal{A}$, $p \in \{1, \dots, m\}$.

Proof. By minimality, we have $\left[\prod_{p=1}^m \widetilde{\pi}_p^\varphi(\mathcal{A})\widetilde{V}_\varphi \mathcal{H}_\ell \right] = \widetilde{\mathcal{H}}_\ell^\varphi$ and $\left[\prod_{p=1}^m \widehat{\pi}_p^\varphi(\mathcal{A})\widehat{V}_\varphi \mathcal{H}_\ell \right] = \widehat{\mathcal{H}}_\ell^\varphi$ for each $\ell \in \Omega$.

Then by Lemma 4.1, we get $\left[\prod_{p=1}^m \widetilde{\pi}_p^\varphi(\mathcal{A})\widetilde{V}_\varphi \mathcal{D} \right] = \widetilde{\mathcal{H}}^\varphi$ and $\left[\prod_{p=1}^m \widehat{\pi}_p^\varphi(\mathcal{A})\widehat{V}_\varphi \mathcal{D} \right] = \widehat{\mathcal{H}}^\varphi$. Let us define a

map $U_\varphi : \text{Span}\left\{ \prod_{p=1}^m \widetilde{\pi}_p^\varphi(\mathcal{A})\widetilde{V}_\varphi \mathcal{D} \right\} \rightarrow \text{Span}\left\{ \prod_{p=1}^m \widehat{\pi}_p^\varphi(\mathcal{A})\widehat{V}_\varphi \mathcal{D} \right\}$ by

$$U_\varphi \left(\sum_{j=1}^n \prod_{p=1}^m \widetilde{\pi}_p^\varphi(a_{p_j})\widetilde{V}_\varphi h_j \right) = \sum_{j=1}^n \prod_{p=1}^m \widehat{\pi}_p^\varphi(a_{p_j})\widehat{V}_\varphi h_j,$$

for all $a_{p_j} \in \mathcal{A}$, and $h_j \in \mathcal{D}$. Firstly, we show that U_φ is well defined. Suppose that $k = 2m - 1$ (the result follows for $k = 2m$) and without loss of generality there is an $\ell \in \Omega$ such that $h_j \in \mathcal{H}_\ell$ for $j \in \{1, \dots, n\}$.

It follows that

$$\begin{aligned}
\left\| U_\varphi \left(\sum_{j=1}^n \prod_{p=1}^m \tilde{\pi}_p^\varphi(a_{p_j}) \tilde{V}_\varphi h_j \right) \right\|_{\mathcal{H}_\ell^\varphi}^2 &= \left\| \sum_{j=1}^n \prod_{p=1}^m \hat{\pi}_p^\varphi(a_{p_j}) \hat{V}_\varphi h_j \right\|_{\mathcal{H}_\ell^\varphi}^2 \\
&= \sum_{i,j=1}^n \left\langle \prod_{p=1}^m \hat{\pi}_p^\varphi(a_{p_j}) \hat{V}_\varphi h_j, \prod_{p=1}^m \hat{\pi}_p^\varphi(a_{p_i}) \hat{V}_\varphi h_i \right\rangle_{\mathcal{H}_\ell^\varphi} \\
&= \sum_{i,j=1}^n \left\langle (\hat{V}_\varphi)^* \prod_{p=1}^m \hat{\pi}_p^\varphi(a_{p_i}^* a_{p_j}) \hat{V}_\varphi h_j, h_i \right\rangle_{\mathcal{H}_\ell} \\
&= \sum_{i,j=1}^n \left\langle \varphi(a_{m_i}^*, \dots, a_{p_i}^* a^*, \dots, a_{2_i}^*, a_{1_i}^* a_{1_j}, \dots, a a_{p_j}, \dots, a_{m_j}) h_j, h_i \right\rangle_{\mathcal{H}_\ell} \\
&= \sum_{i,j=1}^n \left\langle (\tilde{V}_\varphi)^* \prod_{p=1}^m \tilde{\pi}_p^\varphi(a_{p_i}^* a_{p_j}) \tilde{V}_\varphi h_j, h_i \right\rangle_{\mathcal{H}_\ell} \\
&= \left\| \sum_{j=1}^n \prod_{p=1}^m \tilde{\pi}_p^\varphi(a_{p_j}) \tilde{V}_\varphi h_j \right\|_{\tilde{\mathcal{H}}_\ell^\varphi}^2.
\end{aligned}$$

It implies that the linear map $U_\varphi: \text{Span} \left\{ \prod_{p=1}^m \tilde{\pi}_p^\varphi(\mathcal{A}) \tilde{V}_\varphi \mathcal{D} \right\} \rightarrow \text{Span} \left\{ \prod_{p=1}^m \hat{\pi}_p^\varphi(\mathcal{A}) \hat{V}_\varphi \mathcal{D} \right\}$ is a well defined and onto map. Moreover U_φ is an isometry. Since $\text{Span} \left\{ \prod_{p=1}^m \hat{\pi}_p^\varphi(\mathcal{A}) \hat{V}_\varphi \mathcal{D} \right\}$ is dense in $\widehat{\mathcal{H}}^\varphi$ and $\text{Span} \left\{ \prod_{p=1}^m \tilde{\pi}_p^\varphi(\mathcal{A}) \tilde{V}_\varphi \mathcal{D} \right\}$ is dense in $\widetilde{\mathcal{H}}^\varphi$, thus U_φ can be extended to a unitary from $\widetilde{\mathcal{H}}^\varphi$ to $\widehat{\mathcal{H}}^\varphi$. We denote this unitary map by U_φ itself. By construction of U_φ , it is easy to see that $U_\varphi(\widetilde{\mathcal{H}}_\ell^\varphi) \subseteq \widehat{\mathcal{H}}_\ell^\varphi$ for all $\ell \in \Omega$.

Proof of (1): Let $h \in \mathcal{D}$. Then $U_\varphi \tilde{V}_\varphi h = U_\varphi \left(\prod_{p=1}^m \tilde{\pi}_p^\varphi(1) \tilde{V}_\varphi h \right) = \prod_{p=1}^m \hat{\pi}_p^\varphi(1) \hat{V}_\varphi h = \hat{V}_\varphi h$. Since $U_\varphi \tilde{V}_\varphi, \hat{V}_\varphi$ are bounded linear operators which agree on \mathcal{D} , we conclude that $U_\varphi \tilde{V}_\varphi = \hat{V}_\varphi$.

Proof of (2): $\text{Dom} \left(U_\varphi \prod_{p=1}^m \tilde{\pi}_p^\varphi(a_p) \right) = \{ \xi \in \text{Dom} \left(\prod_{p=1}^m \tilde{\pi}_p^\varphi(a_p) \right) : \prod_{p=1}^m \tilde{\pi}_p^\varphi(a_p)(\xi) \in \text{Dom}(U_\varphi) \} = \widetilde{\mathcal{D}}^\varphi$, and $\text{Dom} \left(\prod_{p=1}^m \hat{\pi}_p^\varphi(a_p) U_\varphi \right) = \{ \xi \in \widetilde{\mathcal{H}}^\varphi : U_\varphi(\xi) \in \text{Dom} \left(\prod_{p=1}^m \hat{\pi}_p^\varphi(a_p) \right) \} = \{ \xi \in \widetilde{\mathcal{H}}^\varphi : U_\varphi(\xi) \in \widehat{\mathcal{D}}^\varphi \} = \widetilde{\mathcal{D}}^\varphi$.

Suppose ξ is of the form $\xi = \sum_{j=1}^n \prod_{p=1}^m \tilde{\pi}_p^\varphi(a_{p_j}) \tilde{V}_\varphi h_j$, where $a_{p_j} \in \mathcal{A}, h_j \in \mathcal{H}_\ell$. Then for all $b_p \in \mathcal{A}$, we have

$$\begin{aligned}
U_\varphi \prod_{p=1}^m \tilde{\pi}_p^\varphi(b_p) \xi &= U_\varphi \left(\sum_{j=1}^n \prod_{p=1}^m \tilde{\pi}_p^\varphi(b_p a_{p_j}) \tilde{V}_\varphi h_j \right) = \sum_{j=1}^n \prod_{p=1}^m \hat{\pi}_p^\varphi(b_p a_{p_j}) \hat{V}_\varphi h_j \\
&= \prod_{p=1}^m \hat{\pi}_p^\varphi(b_p) \left(\sum_{j=1}^n \prod_{p=1}^m \hat{\pi}_p^\varphi(a_{p_j}) \hat{V}_\varphi h_j \right) \\
&= \prod_{p=1}^m \hat{\pi}_p^\varphi(b_p) U_\varphi \xi.
\end{aligned}$$

This implies that $U_\varphi \prod_{p=1}^m \tilde{\pi}_p^\varphi(b_p)|_{\widetilde{\mathcal{H}}_\ell^\varphi} = \prod_{p=1}^m \hat{\pi}_p^\varphi(b_p) U_\varphi|_{\widetilde{\mathcal{H}}_\ell^\varphi}$ for all $\ell \in \Omega$. This completes the proof. \square

5. RADON-NIKODÝM THEOREM

In this section, we prove Radon-Nikodým type theorem for local completely positive invariant k -linear maps. In C^* -algebra theory, a Radon-Nikodým derivative is obtained between two CP-maps that are dominated by one with the other map. So, in this setup, it is necessary to define such relation “ \leq ” on $ICPCC_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$. The results proved in this section generalizes our earlier work with B. V. Rajarama Bhat [3] when $k = 1$.

Definition 5.1. Let $\varphi, \psi \in ICPCC_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$. If $\varphi - \psi \in ICPCC_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$, then ψ is said to be dominated by φ . In this case, we write $\psi \leq \varphi$.

Let $\varphi \in ICPCC_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$ and let $(\{\pi_p^\varphi\}_{p=1}^m, V_\varphi, \{\mathcal{H}^\varphi; \mathcal{E}^\varphi; \mathcal{D}^\varphi\})$ be the minimal Stinespring triple of φ . The commutant of the family $\{\pi_p^\varphi\}_{p=1}^m$ is defined by the subspace of $\mathcal{B}(\mathcal{H}^\varphi)$ as,

$$(13) \quad \bigcap_{p=1}^m \pi_p^\varphi(\mathcal{A})' = \{T \in \mathcal{B}(\mathcal{H}^\varphi) : T\pi_p^\varphi(a) \subseteq \pi_p^\varphi(a)T \text{ for all } a \in \mathcal{A}, 1 \leq p \leq m\}.$$

Lemma 5.2. Let $\varphi_1, \varphi_2 \in ICPCC_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$ and $m = \lfloor \frac{k+1}{2} \rfloor$. Let $(\{\pi_p^{\varphi_i}\}_{p=1}^m, V_{\varphi_i}, \{\mathcal{H}^{\varphi_i}; \mathcal{E}^{\varphi_i}; \mathcal{D}^{\varphi_i}\})$ be the minimal Stinespring's triple for φ_i with $i = 1, 2$. If $\varphi_1 \leq \varphi_2$, then there exists a unique contraction $T \in \mathcal{B}(\mathcal{H}^{\varphi_2}, \mathcal{H}^{\varphi_1})$ such that

- (i) $TV_{\varphi_2} = V_{\varphi_1}$.
- (ii) $T\left(\prod_{p=1}^m \pi_p^{\varphi_2}(a_p)\right) \subseteq \left(\prod_{p=1}^m \pi_p^{\varphi_1}(a_p)\right)T$, for all $a_p \in \mathcal{A}$ and $p = 1, 2, \dots, m$.

Proof. It is given that $(\varphi_2 - \varphi_1) \in ICPCC_{loc}(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$. Let us take D_p (for $2 \leq p \leq m$) and R_1 as in Equation (5). Suppose that $k = 2m - 1$, then $(D_m^*, \dots, D_2^*, R_1^*R_1, D_2, \dots, D_m) \in M_n(\mathcal{A})^{2m-1}$ is α -symmetric (see Definition 2.2) and $R_1^*R_1 \geq \alpha 0$, for every $\alpha \in \Lambda$. Moreover, for $a_{p_j} \in \mathcal{A}$, $h_j \in \mathcal{H}_\ell$ with $j \in \{1, \dots, n\}$ and $p \in \{1, \dots, m\}$, $n \in \mathbb{N}$ we see that

$$(14) \quad \begin{aligned} & \sum_{t,j=1}^n \left\langle \varphi_1(a_{m_t}^*, \dots, a_{2_t}^*, a_{1_t}^* a_{1_j}, a_{2_j}, \dots, a_{m_j}) h_j, h_t \right\rangle_{\mathcal{H}_\ell} \\ &= \left\langle (\varphi_1)_n(D_m^*, \dots, D_2^*, R_1^*R_1, D_2, \dots, D_m) \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix}, \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix} \right\rangle_{\mathcal{H}_\ell^n} \\ &\leq \left\langle (\varphi_2)_n(D_m^*, \dots, D_2^*, R_1^*R_1, D_2, \dots, D_m) \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix}, \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix} \right\rangle_{\mathcal{H}_\ell^n} \\ &= \sum_{t,j=1}^n \left\langle \varphi_2(a_{m_t}^*, \dots, a_{2_t}^*, a_{1_t}^* a_{1_j}, a_{2_j}, \dots, a_{m_j}) h_j, h_t \right\rangle_{\mathcal{H}_\ell}. \end{aligned}$$

The above observation is helpful in defining the desired contraction from \mathcal{H}^{φ_2} to \mathcal{H}^{φ_1} . Firstly, by the minimality condition from Definition 4.3, we have

$$\mathcal{H}_\ell^{\varphi_i} = \left[\prod_{p=1}^m \pi_p^{\varphi_i}(\mathcal{A}) V_{\varphi_i} \mathcal{H}_\ell \right], \text{ for every } \ell \in \Omega, i = 1, 2.$$

Thus by Lemma 4.1, it implies that $\mathcal{H}^{\varphi_i} = \left[\left(\prod_{p=1}^m \pi_p^{\varphi_i}(\mathcal{A}) \right) V_{\varphi_i} \mathcal{D} \right]$, for $i = 1, 2$. Let us define a map T on the dense subset of \mathcal{H}^{φ_2} as,

$$(15) \quad T \left(\sum_{j=1}^n \left(\prod_{p=1}^m \pi_p^{\varphi_2}(a_{p_j}) V_{\varphi_2} h_j \right) \right) = \sum_{j=1}^n \left(\prod_{p=1}^m \pi_p^{\varphi_1}(a_{p_j}) V_{\varphi_1} h_j \right),$$

for all $a_{p_j} \in \mathcal{A}$, $h_j \in \mathcal{D}$, $j = \{1, 2, \dots, n\}$ and $p = \{1, 2, \dots, m\}$. Now we show that T is a well defined bounded map. Since \mathcal{D} is the union space, there is an $\ell \in \Omega$ with $h_j \in \mathcal{H}_\ell$, for $j \in \{1, 2, \dots, n\}$. Then we have

$$\begin{aligned} \left\| \sum_{j=1}^n \left(\prod_{p=1}^m \pi_p^{\varphi_1}(a_{p_j}) V_{\varphi_1} h_j \right) \right\|_{\mathcal{H}_\ell^{\varphi_1}}^2 &= \left\langle \sum_{j=1}^n \left(\prod_{p=1}^m \pi_p^{\varphi_1}(a_{p_j}) V_{\varphi_1} h_j \right), \sum_{j=1}^n \left(\prod_{p=1}^m \pi_p^{\varphi_1}(a_{p_j}) V_{\varphi_1} h_j \right) \right\rangle_{\mathcal{H}_\ell^{\varphi_1}} \\ &= \sum_{t,j=1}^n \left\langle V_{\varphi_1}^* \pi_1^{\varphi_1}(a_{1_t}^*) \cdots \pi_m^{\varphi_1}(a_{m_t}^*) \pi_1^{\varphi_1}(a_{m_j}) \cdots \pi_1^{\varphi_1}(a_{1_j}) V_{\varphi_1} h_j, h_t \right\rangle_{\mathcal{H}_\ell} \\ &= \sum_{t,j=1}^n \left\langle V_{\varphi_1}^* \pi_1^{\varphi_1}(a_{1_t}^* a_{1_j}) \cdots \pi_2^{\varphi_1}(a_{2_t}^* a_{2_j}) \cdots \pi_m^{\varphi_1}(a_{m_t}^* a_{m_j}) V_{\varphi_1} h_j, h_t \right\rangle_{\mathcal{H}_\ell} \\ &= \sum_{t,j=1}^n \left\langle \varphi_1(a_{m_t}^*, \dots, a_{2_t}^*, a_{1_t}^* a_{1_j}, a_{2_j}, \dots, a_{m_j}) h_j, h_t \right\rangle_{\mathcal{H}_\ell}. \end{aligned}$$

Further by Equation (14), we get

$$\left\| T \left(\sum_{j=1}^n \left(\prod_{p=1}^m \pi_p^{\varphi_2}(a_{p_j}) V_{\varphi_2} h_j \right) \right) \right\| \leq \left\| \sum_{j=1}^n \left(\prod_{p=1}^m \pi_p^{\varphi_1}(a_{p_j}) V_{\varphi_1} h_j \right) \right\|.$$

This shows that T is a well defined contraction mapping from a dense subspace of \mathcal{H}^{φ_2} onto the dense subspace of \mathcal{H}^{φ_1} . Thus there is a unique contraction defined on \mathcal{H}^{φ_2} , again denote it by $T \in \mathcal{B}(\mathcal{H}^{\varphi_2}, \mathcal{H}^{\varphi_1})$. It is clear from the definition T in Equation (15) that $T(\mathcal{H}_\ell^{\varphi_2}) \subseteq \mathcal{H}_\ell^{\varphi_1}$ and $T|_{\mathcal{H}_\ell^{\varphi_2}} \in \mathcal{B}(\mathcal{H}_\ell^{\varphi_2}, \mathcal{H}_\ell^{\varphi_1})$, for every $\ell \in \Omega$.

Proof of (i): Let $g \in \mathcal{D}$. Then

$$TV_{\varphi_2} g = T \left(\prod_{p=1}^m \pi_p^{\varphi_2}(1) V_{\varphi_2} g \right) = \prod_{p=1}^m \pi_p^{\varphi_1}(1) V_{\varphi_1} g = V_{\varphi_1} g.$$

The result follows from the fact that \mathcal{D} is dense in \mathcal{H} .

Proof of (ii): Let $a_p, b_p \in \mathcal{A}$ with $p \in \{1, \dots, m\}$ and $h \in \mathcal{D}$. Then by using the fact that $\{\pi_p^{\varphi_i}\}_{p=1}^m$ is a commuting family of $*$ -homomorphisms from \mathcal{A} to $C_{\mathcal{G}^{\varphi_i}}^*(\mathcal{D}^{\varphi_i})$ for $i = 1, 2$ we have

$$\begin{aligned} T \left(\prod_{p=1}^m \pi_p^{\varphi_2}(a_p) \right) \left(\prod_{p=1}^m \pi_p^{\varphi_2}(b_p) V_{\varphi_2} h \right) &= T \left(\prod_{p=1}^m \pi_p^{\varphi_2}(a_p b_p) V_{\varphi_2} h \right) \\ &= \prod_{p=1}^m \pi_p^{\varphi_1}(a_p b_p) V_{\varphi_1} h \\ &= \prod_{p=1}^m \pi_p^{\varphi_1}(a_p) \left(\prod_{p=1}^m \pi_p^{\varphi_2}(b_p) V_{\varphi_1} h \right) \\ &= \prod_{p=1}^m \pi_p^{\varphi_1}(a_p) T \left(\prod_{p=1}^m \pi_p^{\varphi_2}(b_p) V_{\varphi_2} h \right). \end{aligned}$$

It follows that $T\left(\prod_{p=1}^m \pi_p^{\varphi_2}(a_p)\right) \subseteq \left(\prod_{p=1}^m \pi_p^{\varphi_1}(a_p)\right)T$, for all $a_p \in \mathcal{A}$ and $p \in \{1, \dots, m\}$. Similarly, if $k = 2m$, then $(D_m^*, \dots, D_2^*, R_1^*, R_1, D_2, \dots, D_m) \in \mathcal{A}^{2m}$ is α -symmetric, for every $\alpha \in \Lambda$ and hence the result follows same as in the odd case. \square

Lemma 5.3. Let $\varphi \in ICPC C_{loc}^s(\mathcal{A}^k, C_{\mathcal{E}}^*(\mathcal{D}))$ and $m = \lceil \frac{k+1}{2} \rceil$. For every $T \in \bigcap_{p=1}^m \pi_p^{\varphi}(\mathcal{A})' \cap C_{\mathcal{E}\varphi}^*(\mathcal{D}^{\varphi})$,

define the k -linear map $\varphi_T: \mathcal{A}^k \rightarrow C_{\mathcal{E}}(\mathcal{D})$ by

$$(16) \quad \varphi_T(a_1, \dots, a_k) = \begin{cases} V_{\varphi}^* T \pi_1^{\varphi}(a_m) \pi_2^{\varphi}(a_{m-1} a_{m+1}) \cdots \pi_m^{\varphi}(a_1 a_{2m-1}) V_{\varphi}, & \text{if } k = 2m - 1 \\ V_{\varphi}^* T \pi_1^{\varphi}(a_m a_{m+1}) \pi_2^{\varphi}(a_{m-1} a_{m+2}) \cdots \pi_m^{\varphi}(a_1 a_{2m}) V_{\varphi}, & \text{if } k = 2m, \end{cases}$$

for all $a_1, a_2, \dots, a_k \in \mathcal{A}$. Then the correspondence $T \mapsto \varphi_T$ is an injective linear map.

Proof. Let $T_1, T_2 \in \bigcap_{p=1}^m \pi_p^{\varphi}(\mathcal{A})' \cap C_{\mathcal{E}\varphi}^*(\mathcal{D}^{\varphi})$ and $\lambda \in \mathbb{C}$. Then

$$\begin{aligned} & \varphi_{(T_1 + \lambda T_2)}(a_1, \dots, a_k) \\ &= \begin{cases} V_{\varphi}^*(T_1 + \lambda T_2) \pi_1^{\varphi}(a_m) \pi_2^{\varphi}(a_{m-1} a_{m+1}) \cdots \pi_m^{\varphi}(a_1 a_{2m-1}) V_{\varphi}, & \text{if } k = 2m - 1 \\ V_{\varphi}^*(T_1 + \lambda T_2) \pi_1^{\varphi}(a_m a_{m+1}) \pi_2^{\varphi}(a_{m-1} a_{m+2}) \cdots \pi_m^{\varphi}(a_1 a_{2m}) V_{\varphi}, & \text{if } k = 2m \end{cases} \\ &= \varphi_{T_1}(a_1, \dots, a_k) + \lambda \varphi_{T_2}(a_1, \dots, a_k). \end{aligned}$$

Suppose that $\varphi_T = 0$ for some $T \in \bigcap_{p=1}^m \pi_p^{\varphi}(\mathcal{A})' \cap C_{\mathcal{E}\varphi}^*(\mathcal{D}^{\varphi})$. We claim that $T = 0$. Let us take $a_{p_j}, b_{p_i} \in \mathcal{A}$ and $h_j, g_i \in \mathcal{D}$, where $j \in \{1, \dots, n\}, i \in \{1, \dots, k\}$ and $p \in \{1, \dots, m\}$. Then there exists an $\ell \in \Omega$ such that each $g_j, h_i \in \mathcal{H}_{\ell}$ for all $j \in \{1, \dots, n\}, i \in \{1, \dots, k\}$. Then we compute that

$$\begin{aligned} & \left\langle T \left(\sum_{j=1}^n \prod_{p=1}^m \pi_p^{\varphi}(a_{p_j}) V_{\varphi} h_j \right), \left(\sum_{i=1}^t \prod_{p=1}^m \pi_p^{\varphi}(b_{p_i}) V_{\varphi} g_i \right) \right\rangle_{\mathcal{H}_{\ell}^{\varphi}} \\ &= \sum_{j=1}^n \sum_{i=1}^t \left\langle V_{\varphi}^* T \prod_{p=1}^m \pi_p^{\varphi}(b_{p_i}^* a_{p_j}) V_{\varphi} h_j, g_i \right\rangle_{\mathcal{H}_{\ell}} \\ &= \sum_{j=1}^n \sum_{i=1}^t \left\langle V_{\varphi}^* T \pi_1^{\varphi}(b_{1_i}^* a_{1_j}) \pi_2^{\varphi}(b_{2_i}^* a_{2_j}) \cdots \pi_m^{\varphi}(b_{m_i}^* a_{m_j}) V_{\varphi} h_j, g_i \right\rangle_{\mathcal{H}_{\ell}} \\ &= \begin{cases} \sum_{j=1}^n \sum_{i=1}^t \left\langle \varphi_T(b_{m_i}^*, \dots, b_{2_i}^*, b_{1_i}^* a_{1_j}, a_{2_j}, \dots, a_{m_j}) h_j, g_i \right\rangle & \text{if } k = 2m - 1 \\ \sum_{j=1}^n \sum_{i=1}^t \left\langle \varphi_T(b_{m_i}^*, \dots, b_{2_i}^*, b_{1_i}^*, a_{1_j}, a_{2_j}, \dots, a_{m_j}) h_j, g_i \right\rangle & \text{if } k = 2m \end{cases} \\ &= 0. \end{aligned}$$

Since $\mathcal{H}^{\varphi} = \left[\prod_{p=1}^m \pi_p^{\varphi}(\mathcal{A}) V_{\varphi} \mathcal{D} \right]$, we conclude that $T = 0$. \square

Proposition 5.4. Let $(\{\pi_p^\varphi\}_{p=1}^m, V_\varphi, \{\mathcal{H}^\varphi; \mathcal{E}^\varphi; \mathcal{D}^\varphi\})$ be the minimal Stinespring's triple for $\varphi \in ICPC C_{loc}^s(\mathcal{A}^k, C_{\mathcal{E}^\varphi}^*(\mathcal{D}^\varphi))$ and $T \in \bigcap_{p=1}^m \pi_p^\varphi(\mathcal{A})' \cap C_{\mathcal{E}^\varphi}^*(\mathcal{D}^\varphi)$. Let $\varphi_T: \mathcal{A}^k \rightarrow C_{\mathcal{E}^\varphi}^*(\mathcal{D})$ be a map defined in Equation (16). If $0 \leq T \leq I$, then the following assertions hold true.

- (i) $\varphi_T \in ICPC C_{loc}^s(\mathcal{A}^k, C_{\mathcal{E}^\varphi}^*(\mathcal{D}))$.
- (ii) $\varphi_T \leq \varphi$.

Proof. Proof of (i): For every $a_1, \dots, a_k \in \mathcal{A}$, we see that

$$\begin{aligned} & \varphi_T^*(a_1, \dots, a_k) \\ &= \varphi_T(a_k^*, \dots, a_1^*)^* \\ &= \begin{cases} V_\varphi^* \pi_m^\varphi(a_{2m-1}^* a_1^*)^* \cdots \pi_1^\varphi(a_m^*)^* T V_\varphi & \text{if } k = 2m - 1 \\ V_\varphi^* \pi_m^\varphi(a_{2m}^* a_1^*)^* \cdots \pi_1^\varphi(a_{m+1}^* a_m^*)^* T V_\varphi & \text{if } k = 2m \end{cases} \\ &= \begin{cases} V_\varphi^* T \pi_1^\varphi(a_m) \cdots \pi_m^\varphi(a_1 a_{2m-1}) V_\varphi, & \text{if } k = 2m - 1 \\ V_\varphi^* T \pi_1^\varphi(a_m a_{m+1}) \cdots \pi_m^\varphi(a_1 a_{2m}) V_\varphi, & \text{if } k = 2m \end{cases} \\ &= \varphi_T(a_1, \dots, a_k). \end{aligned}$$

Since $\{\pi_p^\varphi\}_{p=1}^m$ is a commuting family of $*$ -homomorphisms and $T \in \bigcap_{p=1}^m \pi_p(\mathcal{A})' \cap C_{\mathcal{E}^\varphi}^*(\mathcal{D})$ is positive.

This implies that $\varphi_T^* = \varphi_T$ and hence φ_T is symmetric.

Suppose $k = 2m - 1$ and let $a_1, a_2, \dots, a_{2m-1}, c_1, \dots, c_{m-1} \in \mathcal{A}$. Then

$$\begin{aligned} & \varphi_T(a_1 c_1, \dots, a_{m-1} c_{m-1}, a_m, a_{m+1}, \dots, a_{2m-1}) \\ &= V_\varphi^* T \pi_1^\varphi(a_m) \pi_2^\varphi(a_{m-1} c_{m-1} a_{m+1}) \cdots \pi_m^\varphi(a_1 c_1 a_{2m-1}) V_\varphi \\ &= \varphi_T(a_1, \dots, a_{m-1}, a_m, c_{m-1} a_{m+1}, \dots, c_1 a_{2m-1}). \end{aligned}$$

Similarly, if $k = 2m$, we have

$$\varphi_T(a_1 c_1, \dots, a_{m-1} c_{m-1}, a_m c_m, a_{m+1}, \dots, a_{2m}) = \varphi_T(a_1, \dots, a_{m-1}, a_m, c_m a_{m+1}, \dots, c_1 a_{2m}).$$

Therefore, φ_T is an invariant k -linear map. Now we show that φ_T is local completely contractive. Suppose that $k = 2m - 1$. Let $n \in \mathbb{N}$ and $A_t = [a_{t_{ij}}]_{i,j=1}^n \in M_n(\mathcal{A})^{2m-1}$ for $t = 1, \dots, (2m-1)$. We know that the amplification map $(\varphi_T)_n: M_n(\mathcal{A})^k \rightarrow C_{\mathcal{E}^\varphi}^*(\mathcal{D}^n)$ is defined by

$$\begin{aligned} & (\varphi_T)_n(A_1, A_2, \dots, A_{2m-1}) \\ &= \left[\sum_{r_1, r_2, \dots, r_{2(m-1)}=1}^n \varphi_T(a_{1_{ir_1}}, a_{2_{r_1 r_2}}, \dots, a_{2m-1_{r_{2(m-1)} j}}) \right]_{i,j=1}^n. \end{aligned}$$

For every $h_1, h_2, \dots, h_n \in \mathcal{D}$, due to the fact that \mathcal{D} is the union space of upper filtered family \mathcal{E} , there exists an $\ell \in \Omega$ such that $h_1, h_2, \dots, h_n \in \mathcal{H}_\ell$. Then we have

$$\begin{aligned} & \left\| (\varphi_T)_n(A_1, A_2, \dots, A_{2m-1}) \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix} \right\|_{\mathcal{H}_\ell^n}^2 \\ &= \sum_{i=1}^n \left\| \sum_{j=1}^n \sum_{r_1, r_2, \dots, r_{2(m-1)}=1}^n \varphi_T(a_{1_{ir_1}}, a_{2_{r_1 r_2}}, \dots, a_{2m-1_{r_{2(m-1)} j}}) h_j \right\|_{\mathcal{H}_\ell}^2 \end{aligned}$$

$$\begin{aligned}
&= \sum_{i=1}^n \left\| \sum_{j=1}^n \sum_{r_1, r_2, \dots, r_{2(m-1)}=1}^n V_\varphi^* T \pi_1^\varphi(a_{m_{r_{m-1} r_m}}) \cdots \pi_m^\varphi(a_{1_{ir_1} a_{2m-1} r_{2(m-1)j}}) V_\varphi h_j \right\|_{\mathcal{H}_\ell}^2 \\
&= \sum_{i=1}^n \left\| V_\varphi^* T \left(\sum_{j=1}^n \sum_{r_1, r_2, \dots, r_{2(m-1)}=1}^n \pi_1^\varphi(a_{m_{r_{m-1} r_m}}) \cdots \pi_m^\varphi(a_{1_{ir_1} a_{2m-1} r_{2(m-1)j}}) V_\varphi h_j \right) \right\|_{\mathcal{H}_\ell}^2 \\
&\leq \sum_{i=1}^n \left\| \sum_{j=1}^n \sum_{r_1, r_2, \dots, r_{2(m-1)}=1}^n \pi_1^\varphi(a_{m_{r_{m-1} r_m}}) \cdots \pi_m^\varphi(a_{1_{ir_1} a_{2m-1} r_{2(m-1)j}}) V_\varphi h_j \right\|_{\mathcal{H}_\ell^\varphi}^2 \\
&= \sum_{i=1}^n \left\| \sum_{j=1}^n \sum_{r_1, r_2, \dots, r_{2(m-1)}=1}^n \pi_m^\varphi(a_{1_{ir_1}}) \cdots \pi_1^\varphi(a_{m_{r_{m-1} r_m}}) \cdots \pi_m^\varphi(a_{2m-1} r_{2(m-1)j}) V_\varphi h_j \right\|_{\mathcal{H}_\ell^\varphi}^2 \\
&= \left\| \left[\pi_m^\varphi(a_{1_{ij}}) \right]_{i,j=1}^n \cdots \left[\pi_1^\varphi(a_{m_{ij}}) \right]_{i,j=1}^n \cdots \left[\pi_m^\varphi(a_{2m-1} r_{ij}) \right]_{i,j=1}^n \begin{bmatrix} V_\varphi h_1 \\ \vdots \\ V_\varphi h_n \end{bmatrix} \right\|_{\mathcal{H}_\ell^\varphi}^2 \\
&\leq \left\| \left[\pi_m^\varphi(a_{1_{ij}}) \right]_{i,j=1}^n \right\|_\ell^2 \cdots \left\| \left[\pi_1^\varphi(a_{m_{ij}}) \right]_{i,j=1}^n \right\|_\ell^2 \cdots \left\| \left[\pi_m^\varphi(a_{2m-1} r_{ij}) \right]_{i,j=1}^n \right\|_\ell^2 \left\| \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix} \right\|_{\mathcal{H}_\ell}^2
\end{aligned}$$

Since π_p^φ is a local completely contractive map for $p = 1, \dots, m$, there is an $\alpha_p \in \Lambda$ such that

$$\left\| \left[\pi_p^\varphi(b_{ij}) \right]_{i,j=1}^n \right\|_\ell \leq \left\| \left[(b_{ij}) \right]_{i,j=1}^n \right\|_{\alpha_p}, \text{ for } p = 1, \dots, m.$$

It follows from the above arguments that

$$\begin{aligned}
&\left\| (\varphi_T)_n(A_1, A_2, \dots, A_{2m-1}) \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix} \right\|_{\mathcal{H}_\ell^n} \\
(17) \quad &\leq \|A_1\|_{\alpha_m} \cdots \|A_m\|_{\alpha_1} \cdots \|A_{2m-1}\|_{\alpha_m} \left\| \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix} \right\|_{\mathcal{H}_\ell^n}.
\end{aligned}$$

As (Λ, \leq) is a directed set, we can choose $\beta \in \Lambda$ with $\beta \geq \alpha_p$ for each $p = 1, \dots, m$ and by Equation (17), we get

$$\begin{aligned}
&\left\| (\varphi_T)_n(A_1, A_2, \dots, A_{2m-1}) \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix} \right\|_{\mathcal{H}_\ell^n} \\
(18) \quad &\leq \|A_1\|_\beta \cdots \|A_m\|_\beta \cdots \|A_{2m-1}\|_\beta \left\| \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix} \right\|_{\mathcal{H}_\ell^n}.
\end{aligned}$$

It is clear from Equation (18) that for every $\ell \in \Omega$, there is a $\beta \in \Lambda$ such that

$$\left\| (\varphi_T)_n(A_1, A_2, \dots, A_{2m-1}) \right\|_\ell \leq 1, \text{ whenever } \|(A_1, A_2, \dots, A_{2m-1})\|_\beta \leq 1.$$

This implies that φ_T is local completely contractive. By following the similar arguments, the result holds true for $k = 2m$. We conclude that $\varphi_T \in ICC_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$.

Our next aim is to show that φ_T is a local completely positive map. Suppose that $k = 2m - 1$, $n \in \mathbb{N}$ and $A_t = [a_{t,ij}]_{i,j=1}^n \in M_n(\mathcal{A})$, for $t = 1, \dots, 2m - 1$. Then for every $\ell \in \Omega$ and $h_1, h_2, \dots, h_n \in \mathcal{H}_\ell$, we have

$$\begin{aligned}
& \left\langle (\varphi_T)_n(A_1, A_2, \dots, A_{2m-1}) \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix}, \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix} \right\rangle_{\mathcal{H}_\ell^n} \\
&= \sum_{i=1}^n \left\langle \sum_{j=1}^n \sum_{r_1, \dots, r_{2(m-1)}=1}^n \varphi_T(a_{1,ir_1}, \dots, a_{2m-1, r_{2(m-1)}j}) h_j, h_i \right\rangle_{\mathcal{H}_\ell} \\
&= \sum_{i=1}^n \left\langle \sum_{j=1}^n \sum_{r_1, \dots, r_{2(m-1)}=1}^n V_\varphi^* T \pi_1^\varphi(a_{m, r_{m-1}r_m}) \cdots \pi_m^\varphi(a_{1,ir_1} a_{2m-1, r_{2(m-1)}j}) V_\varphi h_j, h_i \right\rangle_{\mathcal{H}_\ell} \\
&= \sum_{i=1}^n \left\langle \sum_{j=1}^n \sum_{r_1, \dots, r_{2(m-1)}=1}^n \pi_m^\varphi(a_{1,ir_1}) \cdots \pi_1^\varphi(a_{m, r_{m-1}r_m}) \cdots \pi_m^\varphi(a_{2m-1, r_{2(m-1)}j}) \sqrt{T} V_\varphi h_j, \sqrt{T} V_\varphi h_i \right\rangle_{\mathcal{H}_\ell^\varphi} \\
&= \left\langle (\pi_1^\varphi)_n(A_m) \xi, \eta \right\rangle_{\mathcal{H}_\ell^\varphi \otimes n},
\end{aligned}$$

where

$$\xi = \prod_{p=2}^m (\pi_p^\varphi)_n(A_{m+p-1}) \begin{bmatrix} \sqrt{T} V_\varphi h_1 \\ \vdots \\ \sqrt{T} V_\varphi h_n \end{bmatrix} \text{ and } \eta = \prod_{p=2}^m (\pi_p^\varphi)_n(A_{m+1-p}^*) \begin{bmatrix} \sqrt{T} V_\varphi h_1 \\ \vdots \\ \sqrt{T} V_\varphi h_n \end{bmatrix}.$$

Since $(\pi_p^\varphi)_n$ is a local contractive map, there exists an $\alpha_p \in \Lambda$ for each $p = 2, \dots, m$ such that

$$(19) \quad \left\| (\pi_p^\varphi)_n(X) \right\|_\ell \leq \|X\|_{\alpha_p}.$$

We can choose $\alpha \geq \alpha_p$ for each p and by Equation (19), we get

$$(20) \quad \left\| (\pi_p^\varphi)_n(X) \right\|_\ell \leq \|X\|_\alpha, \text{ for every } p = 2, \dots, m.$$

Also by using the fact that $(\pi_1^\varphi)_n$ is a local positive map, there exist a $\beta \in \Lambda$ such that

$$(21) \quad (\pi_1^\varphi)_n(B) \geq_\ell 0, \text{ whenever } B \geq_\beta 0.$$

Since (Λ, \leq) is a directed set, we can choose $\gamma \in \Lambda$ with $\alpha, \beta \leq \gamma$. Suppose that

$$(22) \quad (A_1, \dots, A_m, \dots, A_{2m-1}) =_\gamma (A_{2m-1}^*, \dots, A_m^*, \dots, A_1^*)$$

and $A_m \geq_\gamma 0$, then due to the relation $\alpha \leq \gamma$, we have

$$(23) \quad A_{m+p-1} =_\alpha A_{m+1-p}^* \text{ and } A_m \geq_\alpha 0, \text{ for each } p = 2, \dots, m.$$

It follows from Equations (20), (23) that

$$\left\| (\pi_p^\varphi)_n(A_{m+p-1} - A_{m+1-p}^*) \right\|_\ell \leq \|A_{m+p-1} - A_{m+1-p}^*\|_\alpha = 0.$$

That is,

$$(\pi_p^\varphi)_n(A_{m+p-1})|_{\mathcal{H}_\ell^\varphi \otimes n} = (\pi_p^\varphi)_n(A_{m+1-p}^*)|_{\mathcal{H}_\ell^\varphi \otimes n}, \text{ for each } p = 2, \dots, m.$$

This implies that $\xi = \eta$ and hence by Equation (21), we conclude that

$$\left\langle (\pi_1^\varphi)_n(A_m) \xi, \xi \right\rangle_{\mathcal{H}_\ell^\varphi \otimes n} \geq 0, \text{ whenever } A_m \geq_\beta 0.$$

As $\gamma \geq \beta$, it is clear that $A_m \geq_\gamma 0$ whenever $A_m \geq_\beta 0$. Therefore, for each $\ell \in \Omega$, one can choose $\gamma \in \Lambda$ as described in Equation (22) such that

$$\left\langle (\varphi_T)_n(A_1, A_2, \dots, A_{2m-1}) \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix}, \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix} \right\rangle_{\mathcal{H}_\ell^n} \geq 0,$$

whenever $(A_1, \dots, A_m, \dots, A_{2m-1}) =_\gamma (A_{2m-1}^*, \dots, A_m^*, \dots, A_1^*)$ and $A_m \geq_\gamma 0$ and incase $A_j =_\gamma 0$ for some $1 \leq j \leq 2m-1$, then $(\varphi_T)_n =_\ell 0$ for all n . This shows that φ_T is a local completely positive map.

Suppose that $k = 2m$. Then

$$\left\langle (\varphi_T)_n(A_1, \dots, A_{2m}) \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix}, \begin{bmatrix} h_1 \\ \vdots \\ h_n \end{bmatrix} \right\rangle_{\mathcal{H}_\ell^n} = \left\langle (\pi_1^\varphi)_n(A_m A_{m+1}) \tilde{\xi}, \tilde{\eta} \right\rangle_{\mathcal{H}_\ell^{\varphi \otimes n}},$$

where

$$\tilde{\xi} = \left(\prod_{p=2}^m (\pi_p^\varphi)_n(A_{m+p}) \right) \begin{bmatrix} \sqrt{T} V_\varphi h_1 \\ \vdots \\ \sqrt{T} V_\varphi h_n \end{bmatrix} \text{ and } \tilde{\eta} = \left(\prod_{p=2}^m (\pi_p^\varphi)_n(A_{m+1-p}^*) \right) \begin{bmatrix} \sqrt{T} V_\varphi h_1 \\ \vdots \\ \sqrt{T} V_\varphi h_n \end{bmatrix}.$$

Following the similar arguments as in the case of $k = 2m-1$ and from Equations (20), (21), we conclude that there is a $\gamma \in \Lambda$ with

$$(A_1, \dots, A_m, \dots, A_{2m}) =_\gamma (A_{2m}^*, \dots, A_{m+1}^*, \dots, A_1^*),$$

then $\tilde{\xi} = \tilde{\eta}$ and

$$\left\langle (\pi_1^\varphi)_n(A_m^* A_m) \tilde{\xi}, \tilde{\xi} \right\rangle_{\mathcal{H}_\ell^{\varphi \otimes n}} \geq 0.$$

This implies that φ_T is a local completely positive.

Proof of (ii) : We claim that $(\varphi - \varphi_T) \in ICPC C_{loc}(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$. For every $a_1, a_2, \dots, a_k \in \mathcal{A}$, we have

$$\begin{aligned} & (\varphi - \varphi_T)(a_1, \dots, a_m, \dots, a_k) \\ &= \begin{cases} V_\varphi^*(I - T) \pi_1^\varphi(a_m) \pi_2^\varphi(a_{m-1} a_{m+1}) \cdots \pi_m^\varphi(a_1 a_{2m-1}) V_\varphi & \text{if } k = 2m-1 \\ V_\varphi^*(I - T) \pi_1^\varphi(a_m a_{m+1}) \pi_2^\varphi(a_{m-1} a_{m+2}) \cdots \pi_m^\varphi(a_1 a_{2m}) V_\varphi & \text{if } k = 2m \end{cases} \\ &= \varphi_{(I-T)}. \end{aligned}$$

Since $(I - T) \in \bigcap_{p=1}^m \pi_p^\varphi(\mathcal{A})' \cap C_{\mathcal{E}^\varphi}^*(\mathcal{D}^\varphi)$ and $0 \leq (I - T) \leq I$, it follows from the proof of (i) that $\varphi_{(I-T)} \in ICPC C_{loc}(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$. Equivalently, we obtain $\varphi_T \leq \varphi$. \square

The above results are a suitable recipe for Radon-Nikodým theorem for local completely positive invariant k -linear maps. This will provide a characterization of the set of all maps that are dominated by a fixed map from the class $ICPC C_{loc}(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$.

Theorem 5.5. *Let $\varphi, \psi \in ICPC C_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$ and $m = \lceil \frac{k+1}{2} \rceil$. Then $\psi \leq \varphi$ if and only if there is a unique $\Delta_\varphi(\psi) \in \bigcap_{p=1}^m \pi_p^\varphi(\mathcal{A})' \cap C_{\mathcal{E}^\varphi}^*(\mathcal{D}^\varphi)$ with $0 \leq \Delta_\varphi(\psi) \leq I$ such that $\psi = \varphi_{\Delta_\varphi(\psi)}$.*

Proof. Since $\psi \in ICPC C_{loc}^s(\mathcal{A}^k, C_\mathcal{E}^*(\mathcal{D}))$ is a local completely positive map, by Theorem 3.2 we have a minimal Stinespring's triple $(\{\pi_p^\psi\}_{p=1}^m, V_\psi, \{\mathcal{H}^\psi; \mathcal{E}^\psi; \mathcal{D}^\psi\})$ associated to ψ . Suppose that $\psi \leq \varphi$. Then

by Lemma 5.2, there is a unique contraction $T \in \mathcal{B}(\mathcal{H}^\varphi, \mathcal{H}^\psi)$ such that $TV_\varphi = V_\psi$ and

$$(24) \quad T\left(\prod_{p=1}^m \pi_p^\varphi(a_p)\right) \subseteq \left(\prod_{p=1}^m \pi_p^\psi(a_p)\right)T, \text{ for each } a_p \in \mathcal{A}.$$

Let us define $\Delta_\varphi(\psi) := T^*T$. Then clearly $\Delta_\varphi(\psi) \in \mathcal{B}(\mathcal{H}^\varphi)$ and $0 \leq \Delta_\varphi(\psi) \leq I$. Moreover, $\Delta_\varphi(\psi)|_{\mathcal{H}_\ell^\varphi} \in \mathcal{B}(\mathcal{H}_\ell^\varphi)$, for each $\ell \in \Omega$. In view of Equation (24), we compute that

$$\Delta_\varphi(\psi)\pi_p^\varphi(a) = T^*T\pi_p^\varphi(a) \prod_{t=1, t \neq p}^m \pi_t^\varphi(1) \subseteq T^*\pi_p^\psi(a) \prod_{t=1, t \neq p}^m \pi_t^\psi(1)T = \pi_p^\varphi(a)\Delta_\varphi(\psi),$$

for every $a \in \mathcal{A}$ and $p \in \{1, \dots, m\}$. As a result, $\Delta_\varphi(\psi) \in \bigcap_{p=1}^m \pi_p^\varphi(\mathcal{A})' \cap C_{\mathcal{E}^\varphi}^*(\mathcal{D}^\varphi)$. Let $a_1, \dots, a_k \in \mathcal{A}$ and $h, g \in \mathcal{D}$. Then there is an $\ell \in \Omega$ with $h, g \in \mathcal{H}_\ell$ and

$$\begin{aligned} & \left\langle \varphi_{\Delta_\varphi(\psi)}(a_1, \dots, a_k)h, g \right\rangle_{\mathcal{H}_\ell} \\ &= \begin{cases} \left\langle V_\varphi^* \Delta_\varphi(\psi) \pi_1^\varphi(a_m) \cdots \pi_m^\varphi(a_1 a_{2m-1}) V_\varphi h, g \right\rangle_{\mathcal{H}_\ell} & \text{if } k = 2m - 1 \\ \left\langle V_\varphi^* \Delta_\varphi(\psi) \pi_1^\varphi(a_m a_{m+1}) \cdots \pi_m^\varphi(a_1 a_{2m}) V_\varphi h, g \right\rangle_{\mathcal{H}_\ell} & \text{if } k = 2m \end{cases} \\ &= \begin{cases} \left\langle T \pi_1^\varphi(a_m) \cdots \pi_m^\varphi(a_1 a_{2m-1}) V_\varphi h, TV_\varphi g \right\rangle_{\mathcal{H}_\ell^\psi} & \text{if } k = 2m - 1 \\ \left\langle T \pi_1^\varphi(a_m a_{m+1}) \cdots \pi_m^\varphi(a_1 a_{2m}) V_\varphi h, TV_\varphi g \right\rangle_{\mathcal{H}_\ell^\psi} & \text{if } k = 2m \end{cases} \\ &= \begin{cases} \left\langle \pi_1^\psi(a_m) \cdots \pi_m^\psi(a_1 a_{2m-1}) V_\psi h, V_\psi g \right\rangle_{\mathcal{H}_\ell^\psi} & \text{if } k = 2m - 1 \\ \left\langle \pi_1^\psi(a_m a_{m+1}) \cdots \pi_m^\psi(a_1 a_{2m}) V_\psi h, V_\psi g \right\rangle_{\mathcal{H}_\ell^\psi} & \text{if } k = 2m \end{cases} \\ &= \begin{cases} \left\langle V_\psi^* \pi_1^\psi(a_m) \cdots \pi_m^\psi(a_1 a_{2m-1}) V_\psi h, g \right\rangle_{\mathcal{H}_\ell^\psi} & \text{if } k = 2m - 1 \\ \left\langle V_\psi^* \pi_1^\psi(a_m a_{m+1}) \cdots \pi_m^\psi(a_1 a_{2m}) V_\psi h, g \right\rangle_{\mathcal{H}_\ell^\psi} & \text{if } k = 2m \end{cases} \\ &= \left\langle \psi(a_1, \dots, a_k)h, g \right\rangle_{\mathcal{H}_\ell}. \end{aligned}$$

Therefore, $\psi = \varphi_{\Delta_\varphi(\psi)}$. Now conversely, suppose that $\psi = \varphi_{\Delta_\varphi(\psi)}$, where $\Delta_\varphi(\psi) \in \bigcap_{p=1}^m \pi_p^\varphi(\mathcal{A})' \cap C_{\mathcal{E}^\varphi}^*(\mathcal{D}^\varphi)$ with $0 \leq \Delta_\varphi(\psi) \leq I$. Then it follows from Proposition 5.4 that $\psi \in ICPC C_{loc}^s(\mathcal{A}^k, C_{\mathcal{E}^\varphi}^*(\mathcal{D}))$ and $\psi \leq \varphi$. This completes the proof. \square

Corollary 5.6. *Let $\varphi \in ICPC C_{loc}^s(\mathcal{A}^k, C_{\mathcal{E}^\varphi}^*(\mathcal{D}))$ and $m = \lceil \frac{k+1}{2} \rceil$. Then the map*

$$\eta: \left\{ T \in \bigcap_{p=1}^m \pi_p^\varphi(\mathcal{A})' \cap C_{\mathcal{E}^\varphi}^*(\mathcal{D}^\varphi) : 0 \leq T \leq I \right\} \longrightarrow [0, \varphi]$$

defined by $\eta(T) = \varphi_T$ is an affine order isomorphism onto $[0, \varphi]$.

Proof. We know that η is injective map by Lemma 5.3 and it is onto by Theorem 5.5. Suppose that $T_1, T_2 \in \bigcap_{p=1}^m \pi_p^\varphi(\mathcal{A})' \cap C_{\mathcal{E}^\varphi}^*(\mathcal{D}^\varphi)$ with $0 \leq T_1 \leq T_2 \leq I$ then $0 \leq (T_2 - T_1) \leq I$ and by Proposition 5.4, we have

$$\eta(T_2 - T_1) = \varphi_{(T_2 - T_1)} \in ICPC C_{loc}^s(\mathcal{A}^k, C_{\mathcal{E}}^*(\mathcal{D})).$$

This shows that $\eta(T_1) \leq \eta(T_2)$. Therefore, η is an order isomorphism onto $[0, \varphi]$. \square

Author Contributions. All authors contributed equally to this work.

Acknowledgements. The first named author would like to thank his doctoral supervisor Prof. Anil Kumar Karn for introducing the concept of local operator spaces and locally C^* -algebras. The second named author thanks NBHM (National Board for Higher Mathematics, India) for financial support with ref No. [0204/66/2017/R&D-II/15350] and BITS Pilani.

Both authors wish to express their sincere gratitude to Prof. B V Rajarama Bhat for mathematical discussions, support, encouragement and are grateful to Indian Statistical Institute, Bangalore for providing the necessary facilities to carry out this work. The authors thank anonymous reviewers for their careful reading and valuable comments.

REFERENCES

- [1] J.-P Antoine and A. Inoue, *Representability of invariant positive sesquilinear forms on partial $*$ -algebras*, Math. Proc. Cambridge Philos. Soc. **108** (1990), no. 2, 337–353.
- [2] W. Arveson, *A subalgebras C^* -algebras*, Acta Math. **123** (1969), 141–224.
- [3] B. V. Rajarama Bhat, Anindya Ghatak and Santhosh Kumar Pamula, *Stinespring’s theorem for unbounded operator valued local completely positive maps and its applications*, Indag. Math. (N.S.) **32** (2021), no. 2, 547–578.
- [4] E. Christensen, A. M. Sinclair, *Representations of completely bounded multilinear operators*, J. Funct. Anal. **72** (1987), no. 1, 151–181.
- [5] Dosi Anar, *Multinormed W^* -algebras and unbounded operators*, Proc. Amer. Math. Soc. **140** (2012), no. 12, 4187–4202.
- [6] A. Dosiev, *Local operator spaces, unbounded operators and multinormed C^* -algebras*, J. Funct. Anal. **255** (2008), 1724–1760.
- [7] E. Christensen, E. G. Effros and A. Sinclair, *Completely bounded multilinear maps and C^* -algebraic cohomology*, Invent. Math. **90** (1987), no. 2, 279–296.
- [8] E. G. Effros and C. Webster, *Operator analogues of locally convex spaces*, in *Operator algebras and applications (Samos, 1996)*, 163–207, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 495, Kluwer Acad. Publ., Dordrecht. MR1462680
- [9] Gheondea, A. S. Kavruk, *Absolute continuity for operator valued completely positive maps on C^* -algebras*, J. Math. Phys. **50** (2009), no. 2, 022102–29.
- [10] Gheondea, Aurelian, *Operator models for Hilbert locally C^* -modules*, Oper. Matrices **11** (2017), no. 3, 639–667.
- [11] Gudder, Stanley P, *A Radon-Nikodým theorem for $*$ -algebras*, Pacific J. Math. **80** (1979), no. 1, 141–149.
- [12] J. Heo and M. Joița, *A Stinespring type theorem for completely positive multilinear maps on Hilbert C^* -modules*, Linear and Multilinear Algebra **67** (2019), no. 1, 121–140.
- [13] J. Heo, *The structure of invariant α -CP multilinear maps and associated J -representations*, Linear and Multilinear Algebra **64** (2016), no. 7, 1295–1313.
- [14] J. Heo, Un Cig Ji, *Radon-Nikodým type theorem for α -completely positive maps*, J. Math. Phys. **51** (2010), no. 10, 103505, 10 pp
- [15] J. Heo, *Representations of invariant multilinear maps on Hilbert C^* -modules*, Israel J. Math. **118** (2000), 125–146.
- [16] J. Heo, *A Radon-Nikodým theorem for completely positive invariant multilinear maps and its applications*, Math. Proc. Cambridge Philos. Soc. **132** (2002), no. 1, 181–192.
- [17] M. Joița, *Corrigendum to “Comparison of completely positive maps on Hilbert C^* -modules”* [J. Math. Anal. Appl. 393 (2012) 644–650] [MR2921703], J. Math. Anal. Appl. **466** (2018).
- [18] M. Joița, *Comparison of completely positive maps on Hilbert C^* -modules*, J. Math. Anal. Appl. **393** (2012).
- [19] K. R. Parthasarathy, *Comparison of completely positive maps on a C^* -algebra and a Lebesgue decomposition theorem*, in *Athens Conference on Applied Probability and Time Series Analysis I, Lecture Series in Statistics 114* Springer-Verlag, Berlin, (1996), 34–54.
- [20] V. Paulsen, *Completely bounded maps and operator algebras*, Cambridge Studies in Advanced Mathematics, 78, Cambridge University Press, Cambridge, 2002.
- [21] V. I. Paulsen and R. R. Smith, *Multilinear maps and tensor norms on operator systems*, J. Funct. Anal. **73** (1987), no. 2, 258–276.

- [22] A. Sinclair, R. Smith, *Factorization of completely bounded maps bilinear operators and injectivity*, J. Funct. Anal. **157** (1998), 62–87.
- [23] W. F. Stinespring, *Positive functions on C^* -algebras*, Proc. Amer. Math. Soc. **6** (1955), 211–216.
- [24] G. Wittstock, *Extensions of completely bounded C^* -module homomorphisms*, in Proceedings of the Conference on Operator Algebras and Group Representations, Neptun 1980, Pitman, New York, 1983, pp. 238–250.

STATISTICS AND MATHEMATICS UNIT, INDIAN STATISTICAL INSTITUTE, R. V. COLLEGE POST, BANGALORE 560059, INDIA
Email address: `anindya.ghatak123@gmail.com`

DEPARTMENT OF MATHEMATICS, BIRLA INSTITUTE OF TECHNOLOGY AND SCIENCE (BITS PILANI), PILANI CAMPUS, RAJASTHAN 333031, INDIA.
Email address: `santhoshk.pamula@pilani.bits-pilani.ac.in`