

HIGH DIMENSIONAL LOGISTIC REGRESSION UNDER NETWORK DEPENDENCE

SOMABHA MUKHERJEE, SAGNIK HALDER, BHASWAR B. BHATTACHARYA,
AND GEORGE MICHAILIDIS

ABSTRACT. Logistic regression is one of the most fundamental methods for modeling the probability of a binary outcome based on a collection of covariates. However, the classical formulation of logistic regression relies on the independent sampling assumption, which is often violated when the outcomes interact through an underlying network structure, such as over a temporal/spatial domain or on a social network. This necessitates the development of models that can simultaneously handle both the network ‘peer-effect’ (arising from neighborhood interactions) and the effect of (possibly) high-dimensional covariates. In this paper, we develop a framework for incorporating such dependencies in a high-dimensional logistic regression model by introducing a quadratic interaction term, as in the Ising model, designed to capture the pairwise interactions from the underlying network. The resulting model can also be viewed as an Ising model, where the node-dependent external fields linearly encode the high-dimensional covariates. We propose a penalized maximum pseudo-likelihood method for estimating the network peer-effect and the effect of the covariates (the regression coefficients), which, in addition to handling the high-dimensionality of the parameters, conveniently avoids the computational intractability of the maximum likelihood approach. Consequently, our method is computationally efficient and, under various standard regularity conditions, we show that the corresponding estimate attains the classical high-dimensional rate of consistency. In particular, our results imply that even under network dependence it is possible to consistently estimate the model parameters at the same rate as in classical (independent) logistic regression, when the true parameter is sparse and the underlying network is not too dense. As a consequence of the general results, we derive the rates of consistency of our proposed estimator for various natural graph ensembles, such as bounded degree graphs, sparse Erdős-Rényi random graphs, and stochastic block models.

1. INTRODUCTION

Logistic regression [45, 55, 62] is one of the most popular methods for modeling the probability of a binary response based on multiple features/predictor variables. It is a mainstay of modern multivariate statistics that has found widespread applications in various fields, including machine learning, biological and medical sciences, economics, marketing and finance industries, and social sciences. For example, in machine learning it is regularly used for image classification and in the medical sciences it is often used to predict the risk of developing a particular disease based on the patients’ observed characteristics, among others. To describe the model formally, denote the vector of predictor variables (covariates) by $\mathbf{Z}_1, \dots, \mathbf{Z}_N \in \mathbb{R}^d$ and the independent response variables by $X_1, \dots, X_N \in \{-1, 1\}$. Then, the logistic regression model for the probability of positive outcome conditional on the covariates is given by

$$\mathbb{P}(X_i = 1 | \mathbf{Z}_i) = \frac{e^{\boldsymbol{\theta}^\top \mathbf{Z}_i}}{e^{\boldsymbol{\theta}^\top \mathbf{Z}_i} + e^{-\boldsymbol{\theta}^\top \mathbf{Z}_i}},$$

Key words and phrases. High-dimensional inference, Ising models, logistic regression, Markov random fields, network data, penalized regression, pseudo-likelihood, random graphs.

for $1 \leq i \leq N$, where $\boldsymbol{\theta} = (\theta_1, \theta_2, \dots, \theta_d)^\top \in \mathbb{R}^d$ is the vector of regression coefficients.¹ Using the independence of the response variables, the joint distribution of $\mathbf{X} := (X_1, \dots, X_N)$ given $\mathbf{Z} := (\mathbf{Z}_1, \dots, \mathbf{Z}_N) \in \mathbb{R}^{d \times N}$ can be written as:

$$\mathbb{P}(\mathbf{X}|\mathbf{Z}) = \prod_{i=1}^N \frac{e^{X_i \boldsymbol{\theta}^\top \mathbf{Z}_i}}{e^{\boldsymbol{\theta}^\top \mathbf{Z}_i} + e^{-\boldsymbol{\theta}^\top \mathbf{Z}_i}} = \frac{1}{\mathcal{Z}_N(\boldsymbol{\theta}, \mathbf{Z})} \exp \left\{ \sum_{i=1}^N X_i (\boldsymbol{\theta}^\top \mathbf{Z}_i) \right\}, \quad (1)$$

where $\mathcal{Z}_N(\boldsymbol{\theta}, \mathbf{Z}) = \prod_{i=1}^N \frac{1}{e^{\boldsymbol{\theta}^\top \mathbf{Z}_i} + e^{-\boldsymbol{\theta}^\top \mathbf{Z}_i}}$ is the normalizing constant. It is well-known from the classical theory of generalized linear models that the parameter $\boldsymbol{\theta}$ in (1) can be estimated at rate $O(1/\sqrt{N})$ for fixed dimensions, using the maximum likelihood (ML) method (see, for example, [52, 55, 79]).

The classical framework of logistic regression described above is, however, inadequate if the independence assumption on the response variables is violated, which is often the case if the observations are collected over a temporal or spatial domain or on a social network. The recent accumulation of dependent network data in modern statistics has, in particular, made it increasingly important to develop realistic and mathematically tractable methods for modeling high-dimensional distributions with underlying network dependencies (*network peer-effect*). Towards this, the Ising model, initially proposed in statistical physics to model ferromagnetism [46], has turned out to be a useful primitive for modeling such datasets, which arise naturally in spatial statistics, social networks, epidemic modeling, computer vision, neural networks, and computational biology, among others (see [4, 35, 40, 44, 57] and references therein). This can be viewed as a discrete exponential family with binary outcomes, where the sufficient statistic is a quadratic form designed to capture correlations arising from pairwise interactions. More formally, given an interaction matrix $A := ((a_{ij}))_{1 \leq i, j \leq N}$ and binary vector $\mathbf{X} = (X_1, X_2, \dots, X_N) \in \mathcal{C}_N = \{-1, 1\}^N$, the Ising model with parameters β and h encodes the dependence among the coordinates of \mathbf{X} as follows:

$$\mathbb{P}_{\beta, h}(\mathbf{X}) = \frac{1}{2^N \mathcal{Z}_N(\beta, h)} \exp \left\{ \beta \sum_{1 \leq i, j \leq N} a_{ij} X_i X_j + h \sum_{i=1}^N X_i \right\}, \quad (2)$$

where the *normalizing constant* $\mathcal{Z}_N(\beta, h)$ is determined by the condition $\sum_{\mathbf{X} \in \mathcal{C}_N} \mathbb{P}_{\beta, h}(\mathbf{X}) = 1$ (so that $\mathbb{P}_{\beta, h}$ is a probability measure). In statistical physics the parameters β and h are referred to as the *inverse temperature* and the *magnetic field*, respectively.

In this paper we are motivated by applications where in addition to peer effects, arising from an underlying network structure, there are other variables (covariates) associated with the nodes of the network, which affect the outcome of the response variables. For example, in the data collected by the National Longitudinal Study [43] students in grades 7–12 were asked to name up to 10 friends and answer many questions about age, gender, race, socio-economic background, personal and school life, and health, where it becomes imperative to model the peer-effect and the effect of the covariates simultaneously. Another example where high-dimensional covariates interact through an underlying network structure arises in spatial transcriptomics. This is a relatively new direction in biology made possible by technologies for massively parallelized measurement of cell transcriptomes/proteomes in situ that, unlike standard single cell sequencing methods, retains information regarding the spatial neighborhood of the cells. The spatial context of cells can be encoded into a Voronoi neighborhood graph, where the nodes are cells and edges link cells that are situated proximal to each other [34, 39, 64]. Each node has a high dimensional feature set, encoding

¹Note that we are parameterizing the outcomes as $\{-1, 1\}$ instead of the more standard $\{0, 1\}$ to integrate this within the framework of the Ising model (defined in (2)), where the $\{-1, 1\}$ notation is more common.

the measurements made for that cell, be it gene expression or protein expression, depending on the experimental protocol. Here, the goal is to understand how the spatial niche of a cell contribute to its phenotype. For other examples of network peer-effect in the presence of covariates, see [6, 24, 33, 38, 69, 75] and references therein.

In the aforementioned examples, it is natural to envisage a model that combines the logistic model in (1) (which encodes the node-specific covariates) and the Ising model (2) (for capturing the network dependency). Towards this, Daskalakis et al. [31] recently proposed the following model: Suppose for each node $1 \leq i \leq N$ of a network G_N on N vertices, one observes a d -dimensional covariate $\mathbf{Z}_i \in \mathbb{R}^d$. Then, the joint distribution of the binary outcomes $\mathbf{X} = (X_1, X_2, \dots, X_N)$, conditioned on the network G_N and the observed covariates $\mathbf{Z} := (\mathbf{Z}_1, \mathbf{Z}_2, \dots, \mathbf{Z}_N)$ is given by:

$$\mathbb{P}(\mathbf{X}|\mathbf{Z}) \propto \exp\left(\beta \mathbf{X}^\top \mathbf{A} \mathbf{X} + \sum_{i=1}^N X_i (\boldsymbol{\theta}^\top \mathbf{Z}_i)\right) \quad (3)$$

where $\mathbf{A} = ((a_{ij}))_{1 \leq i, j \leq N}$ is the (appropriately scaled) adjacency matrix of the network G_N , the parameter β is a measure of the strength of dependence (the network ‘peer effect’), and the parameter $\boldsymbol{\theta} \in \mathbb{R}^d$ measures the individual effects of the d -covariates. Note that as in the classical Ising model (2), the quadratic term captures the overall dependency in the network arising from pairwise interactions, while the linear term $\boldsymbol{\theta}^\top \mathbf{Z}_i$ encodes the strength of the covariates on the outcome at the i -th node, for $1 \leq i \leq N$, as in the logistic model (1). Moreover, when $\beta = 0$, which corresponds to the independent case, (3) reduces to (1), hence (3) can be viewed as a model for logistic regression with dependent observations.

The increasing relevance of models (2) and (3) for understanding covariate effects and nearest-neighbor interactions in network data, has made it imperative to develop computationally tractable methods for estimating the model parameters and understanding the statistical properties (rates of convergence) of the resulting estimates. A typical problem of interest is estimating the parameters of the model given a single sample of binary outcomes from an underlying network. For the classical Ising model as in (2), this problem has been extensively studied, beginning with the classical results on consistency and optimality of the maximum likelihood (ML) estimates for models where the underlying network is a spatial lattice [26, 37, 41, 65]. However, for general networks, parameter estimation using the ML method turns out to be notoriously hard due to the presence of an intractable normalizing constant in the likelihood. One approach to circumvent this issue that has turned out to be particularly useful, is the *maximum pseudolikelihood* (MPL) estimator [11, 12]. This provides a computationally efficient method for estimating the parameters of a Markov random field that avoids computing the partition function, by maximizing an approximation to the likelihood function (a ‘pseudo-likelihood’), based on conditional distributions. This approach was originally explored in the seminal paper of Chatterjee [21], where general conditions for \sqrt{N} -consistency of the MPL estimate for the model (2) were derived.² This result was subsequently extended to more general models in [7, 31, 32, 36, 60]. In particular, for model (3) Daskalakis et al. [32] showed that given a single sample of observations $(X_i, \mathbf{Z}_i)_{1 \leq i \leq N}$ from (3), the MPL estimate of the parameters $(\beta, \boldsymbol{\theta})$ is \sqrt{N} -consistent, when the dimension d is *fixed*, under various regularity assumptions on the underlying network and the parameters. This result has been subsequently extended to models with higher-order interactions in [32]. The high-dimensional analogue of this problem under an ℓ_1 norm constraint on the regression parameter has been studied very recently in [49].

²A sequence of estimators $\{\hat{\beta}_N\}_{N \geq 1}$ is said to be \sqrt{N} -consistent at β , if for every $\delta > 0$, there exists $M := M(\delta, \beta) > 0$ such that $\mathbb{P}(\sqrt{N}|\hat{\beta}_N(\mathbf{X}) - \beta| \leq M) > 1 - \delta$, for all N large enough.

In this paper, we consider the problem of parameter estimation in model (3) in the *high-dimensional regime* with sparsity constraints, that is, the number of covariates d grows with the size of the network N , but there are only a few non-zero regression coefficients. In other words, we assume that the parameter vector $\boldsymbol{\theta}$ is s -sparse, that is $\|\boldsymbol{\theta}\|_0 := \sum_{j=1}^d \mathbf{1}\{\theta_j \neq 0\} \leq s$, because, despite the fact that many covariates are available, we only expect a few of them to be relevant. Under this assumption, we want to estimate the parameters $(\beta, \boldsymbol{\theta})$ given a *single* sample of dependent observations $(X_i, \mathbf{Z}_i)_{1 \leq i \leq N}$ from model (3). One of the main reasons this problem is especially delicate, is that we only have access to a *single (dependent) sample* from the underlying model. Consequently, classical results from the M -estimation framework, which require the independence of multiple samples, are inapplicable. To deal with this dependence (which leads to the intractable normalizing constant as mentioned before) and the high-dimensionality of the parameter space, we propose a penalized maximum pseudo-likelihood approach for estimating the parameters. To this end, note that the conditional distribution of X_i given $(X_j)_{j \neq i}$ is:

$$\mathbb{P}(X_i = 1 | (X_j)_{j \neq i}, \mathbf{Z}) = \frac{e^{X_i \boldsymbol{\theta}^\top \mathbf{Z}_i + \beta X_i m_i(\mathbf{X})}}{e^{\boldsymbol{\theta}^\top \mathbf{Z}_i + \beta m_i(\mathbf{X})} + e^{-\boldsymbol{\theta}^\top \mathbf{Z}_i - \beta m_i(\mathbf{X})}}, \quad (4)$$

where, as before, $m_i(\mathbf{X}) = \sum_{j=1}^N a_{ij} X_j$ is the *local-effect* at node i , for $1 \leq i \leq N$. Therefore, by multiplying (4) over $1 \leq i \leq N$ and taking logarithms, we get the (negative) *log-pseudolikelihood* (LPL) function

$$\begin{aligned} L_N(\beta, \boldsymbol{\theta}) &= -\frac{1}{N} \sum_{i=1}^N \log \mathbb{P}(X_i = 1 | (X_j)_{j \neq i}, \mathbf{Z}) \\ &= -\frac{1}{N} \sum_{i=1}^N \{X_i(\boldsymbol{\theta}^\top \mathbf{Z}_i + \beta m_i(\mathbf{X})) - \log \cosh(\boldsymbol{\theta}^\top \mathbf{Z}_i + \beta m_i(\mathbf{X}))\} + \log 2. \end{aligned} \quad (5)$$

To encode the sparsity of the high-dimensional parameters we propose the *penalized maximum pseudo-likelihood* (PMPL) estimator of $(\beta, \boldsymbol{\theta}^\top)$, which, given a regularization (tuning) parameter λ , is defined as:

$$(\hat{\beta}, \hat{\boldsymbol{\theta}}^\top) := \arg \min_{(\beta, \boldsymbol{\theta})} \{L_N(\beta, \boldsymbol{\theta}) + \lambda \|(\beta, \boldsymbol{\theta}^\top)\|_1\}, \quad (6)$$

where $\|\boldsymbol{\theta}\|_1 := \sum_{j=1}^d |\theta_j|$. Under various regularity assumptions, we prove that if λ is chosen proportional to $\sqrt{\log d/n}$, then with high probability,

$$\|(\hat{\beta}, \hat{\boldsymbol{\theta}}^\top) - (\beta, \boldsymbol{\theta}^\top)\| \lesssim_s \sqrt{\log d/N},$$

whenever $d \rightarrow \infty$ such that $d = o(N)$ (Theorem 1). In particular, it follows from our results that for bounded degree networks, the MPL estimate attains the same rate (up to logarithmic factors) as in the independent logistic case (1) for models with bounded sparsity. We study the effect of dependence on estimating the regression parameters $\boldsymbol{\theta}$, where we obtain similar rates as above, even when d is much bigger than N , in particular $\log d = o(N)$ (Theorem 2). The proofs combine a conditioning technique introduced in [29] (which ensures that the dependency in the resulting conditional distribution is sufficiently weak) and general concentration inequalities for Ising models. As a consequence of our general result, we establish that the PMPL estimate is $O(1/\sqrt{N})$ -consistent (up to logarithmic factors) for the model (3) in various natural sparse graph ensembles, such as Erdős-Rényi random graphs and also inhomogeneous random graphs that include the popular in applications stochastic block model (Theorem 3 and Corollary 1).

1.1. Related Work. The asymptotic properties of penalized likelihood methods for logistic regression and generalized linear models in high dimensional settings have been extensively studied (see, for example, [16, 48, 70, 76, 78] and the references therein). These results allow d to be much bigger than N and provide rates of convergence for the high-dimensional parameters under various sparsity constraints. The performance of the ML estimate in the logistic regression model when the dimension d scales proportionally with N has also been studied in a series of recent papers [17, 72, 73]. However, as discussed earlier, ML estimation is both computationally and mathematically intractable in model (3), because of the dependency induced by the underlying network. That we are able to recover rates similar to those in the classical high-dimensional logistic regression, in spite of this underlying dependency, using the PMPL method, is one of the highlights of our results.

The problem of structure learning in Markov random fields is another related area of active research, where the goal is to estimate the underlying graph structure given access to *multiple* i.i.d. samples from an Ising model, or a more general graphical model. For more on these results refer to [3, 13, 14, 23, 42, 50, 67, 71, 80] and the references therein. In particular, [67] and [82] use a regularized pseudo-likelihood approach to learn the structure of the interaction \mathbf{A} given *multiple* i.i.d. samples from an Ising model. In a related direction, Daskalakis et al. [30] studied the related problems of identity and independence testing, and Neykov et al. [19, 63] considered problems in graph property testing, given access to multiple samples from an Ising model.

The aforementioned results, however, are in contrast with the present work, where the underlying graph structure is assumed to be *known* and the goal is to derive rates of estimation for the high-dimensional parameters given a *single* sample from the model. This is motivated by the applications mentioned above where it is often difficult, if not impossible, to generate many independent samples from the model within a reasonable amount of time. More closely related to the present work are the problems in hypothesis testing given a single sample from an Ising model [15, 58, 59]. For instance, in [58, 59] the authors considered the problem of global testing against sparse alternatives for the means of binary outcomes following an Ising model (with node-specific magnetic-field parameters).

1.2. Notation. The following notation will be used throughout the remainder of the paper. For a vector $\mathbf{a} := (a_1, \dots, a_s) \in \mathbb{R}^s$ and $0 < p < \infty$, $\|\mathbf{a}\|_p := (\sum_{i=1}^s |a_i|^p)^{\frac{1}{p}}$ denotes its p -th norm and $\|\mathbf{a}\|_\infty := \max_{1 \leq i \leq s} |a_i|$ its maximum norm, respectively. Moreover, $\|\mathbf{a}\|_0 := \sum_{i=1}^s \mathbf{1}\{a_i \neq 0\}$ denote the ‘zero-norm’ of \mathbf{a} , which counts the number of non-zero coordinates of \mathbf{a} .

For a matrix $\mathbf{M} := ((M_{ij}))_{1 \leq i \leq s, 1 \leq j \leq t} \in \mathbb{R}^{s \times t}$ we define the following norms:

- $\|\mathbf{M}\|_F := \sqrt{\sum_{i=1}^s \sum_{j=1}^t M_{ij}^2}$,
- $\|\mathbf{M}\|_\infty := \max_{1 \leq i \leq s} \sum_{j=1}^t |M_{ij}|$,
- $\|\mathbf{M}\|_1 := \max_{1 \leq j \leq t} \sum_{i=1}^s |M_{ij}|$,
- $\|\mathbf{M}\|_2 := \sigma_{\max}(\mathbf{M})$, where $\sigma_{\max}(\mathbf{M})$ denotes the largest singular value of \mathbf{M} .

Note that if \mathbf{M} is a square matrix, then $\|\mathbf{M}\|_2 = \max\{|\lambda_{\min}(\mathbf{M})|, |\lambda_{\max}(\mathbf{M})|\}$, where $\lambda_{\max}(\mathbf{M})$ and $\lambda_{\min}(\mathbf{M})$ denote the maximum and minimum eigenvalues of \mathbf{M} , respectively.

For positive sequences $\{a_n\}_{n \geq 1}$ and $\{b_n\}_{n \geq 1}$, $a_n = O(b_n)$ means $a_n \leq C_1 b_n$, and $a_n = \Omega(b_n)$ means $a_n \geq C_2 b_n$, for all n large enough and positive constants C_1, C_2 . Similarly, $a_n \lesssim b_n$ means $a_n = O(b_n)$, $a_n \gtrsim b_n$ means $a_n = \Omega(b_n)$. Moreover, subscripts in the above notations, for example \lesssim_\square , O_\square and Ω_\square denote that the hidden constants may depend on the subscripted parameters \square . Finally, we say that $a_n = \tilde{O}(b_n)$, if $a_n \leq C(\log n)^r b_n$ for all n large enough and some positive constants C, r .

1.3. Organization. The remainder of the paper is organized as follows. The main results are described in Section 2. The proofs of the main results are given in Sections 3,4 and 5. Proofs of various technical lemmas are given in the Appendix.

2. STATEMENTS OF THE MAIN RESULTS

In this section we present our main results. In Section 2.1 we discuss the rate of convergence of the PMPL estimator (6). The rate for estimating the regression parameters is presented in Section 2.2. We illustrate the applicability of our results on various graph ensembles in Section 2.3.

2.1. Consistency of the PMPL Estimate. We begin by stating the relevant assumptions:

Assumption 1. *The interaction matrix \mathbf{A} in (3) satisfies the following two conditions:*

$$\sup_{N \geq 1} \|\mathbf{A}\|_\infty < \infty \quad \text{and} \quad \liminf_{N \rightarrow \infty} \frac{1}{N} \|\mathbf{A}\|_F^2 > 0.$$

Assumption 2. *The design matrix $\mathbf{Z} := (\mathbf{Z}_1, \dots, \mathbf{Z}_N)^\top$ satisfies*

$$\liminf_{N \rightarrow \infty} \lambda_{\min} \left(\frac{1}{N} \mathbf{Z}^\top \mathbf{Z} \right) > 0.$$

Assumption 3. *The signal parameters $\boldsymbol{\theta}$ and the covariates $\{\mathbf{Z}_i\}_{1 \leq i \leq N}$ are uniformly bounded, that is, there exist positive constants Θ and M such that $\|\boldsymbol{\theta}\|_\infty < \Theta$ and $\|\mathbf{Z}_i\|_\infty < M$, for all $1 \leq i \leq N$.*

The above assumptions combined, aim to strike a balance between the signal from the peer-effects to that from the covariates, to ensure consistent estimation for all $\beta \geq 0$. In particular, a control on $\|\mathbf{A}\|_\infty$ is required to ensure that the peer effects coming from the quadratic dependence term in the probability mass function (3) do not overpower the effect of the signal $\boldsymbol{\theta}$ coming from the linear terms $\boldsymbol{\theta}^\top \mathbf{Z}_i$, thereby hindering joint recovery of the correlation term β and the signal term $\boldsymbol{\theta}$. At the same time, we also require the interaction matrix to be not too sparse, and its entries to be not too small, in order to ensure that the effect of the correlation parameter β is not nullified. This is guaranteed by the condition $\|\mathbf{A}\|_F^2 = \Omega(N)$. In fact, if this condition is violated, it may be impossible to estimate $(\beta, \boldsymbol{\theta})$ consistently. This phenomenon is observed in the Curie-Weiss Model (where the interaction matrix $a_{ij} = 1/N$, for $1 \leq i \neq j \leq N$) [27]. In this example, each entry of \mathbf{A} is $O(1/N)$, and hence, $\|\mathbf{A}\|_F^2 = O(1)$, and even when $d = 1$ (and $\mathbf{Z}_1 = \dots = \mathbf{Z}_N$) consistent estimation of the parameters β and θ is impossible (see Theorem 1.13 [36]).

The minimum eigenvalue condition in Assumption 2 is standard in the high-dimensional literature (see [54, 67]), and is crucial for establishing the strong-concavity of the log-pseudolikelihood function. In particular, this condition holds with high probability, if the covariates $\mathbf{Z}_1, \dots, \mathbf{Z}_N$ are i.i.d. realizations from a sub-Gaussian distribution on \mathbb{R}^d (see Theorem 2.1 in [32]).

Under the above assumptions we establish the rate of convergence of the PMPL estimate (6) given a single sample of observations from the model (3), when the parameter vector is sparse, that is, $\|(\beta, \boldsymbol{\theta}^\top)^\top\|_0 = s$. For notational convenience, we henceforth denote the $(d+1)$ -dimensional vector of parameters by $\boldsymbol{\gamma} := (\beta, \boldsymbol{\theta}^\top)^\top$ and the $(d+1)$ -dimensional vector of PMPL estimates obtained from (6) by $\hat{\boldsymbol{\gamma}} = (\hat{\beta}, \hat{\boldsymbol{\theta}}^\top)^\top$.

Theorem 1. *Suppose that Assumptions 1, 2, and 3 hold. Then, there exists a constant $\delta > 0$ such that by choosing $\lambda := \delta \sqrt{\log(d+1)/N}$ in the objective function in (16) we have,*

$$\|\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}\|_2 = O_s \left(\sqrt{\frac{\log d}{N}} \right), \quad (7)$$

with probability $1 - o(1)$, as $N \rightarrow \infty$ and $d \rightarrow \infty$ such that $d = o(N)$.

The proof of Theorem 1 is given in Section 3. Broadly speaking, it involves the following two steps:

- *Concentration of the gradient:* The first step in the proof of Theorem 1 is to show that the gradient of the logarithm of the pseudo-likelihood function L_N (recall (5)) is concentrated around zero in the ℓ_∞ norm. For this step, we use the conditioning trick introduced in [29], which reduces a general Ising model to an Ising model in the high-temperature regime, where exponential concentration inequalities for functionals of Ising models are available [20].
- *Strong-concavity of the pseudo-likelihood:* In the second step, we show that the logarithm of the pseudo-likelihood function is strongly concave with high probability. This entails showing that the lowest eigenvalue of the Hessian of L_N is bounded away from zero with high probability. For this step, we invoke the lower bounds on the variance of linear projections of \mathbf{X} developed in [29] and more delicate concentration arguments from [1].

Remark 1. Note that the rate in (7) suppresses the dependence on the sparsity parameter s in the order term. From the proof of Theorem 7 we will see that the dependence is, in general, exponential in s . However, if one replaces Assumption 3 with the stronger assumption that the ℓ_2 norms of the parameters and the covariates are *bounded*, that is, $\|\boldsymbol{\theta}\|_2 < \Theta$ and $\|\mathbf{Z}_i\|_\infty < M$, for all $1 \leq i \leq N$, then our proof can be easily modified to recover the standard $\sqrt{s \log d/n}$ rate. In fact, this stronger assumption has been used recently in [49] to derive rates of the pseudo-likelihood estimate under ℓ_1 norm constraints. However, since ℓ_2 -boundedness of the covariates and the parameters is quite restrictive in the high-dimensional setup, we have presented our results with the more realistic ℓ_∞ -bounded condition.

2.2. Estimation of the Regression Coefficients. In this section, we consider the problem of estimating the regression coefficients $\boldsymbol{\theta}$, for $\beta \geq 0$ fixed. The goal here is to understand how network dependence may affect our ability to estimate the high-dimensional regression coefficient under sparsity constraints. While there have been several recent attempts to understand the implications of dependence in high-dimensional inference, most of them have focused on Gaussian models. Going beyond Gaussian models, Mukherjee, Mukherjee, and Yuan [58] considered the problem of testing the global null (that is, $\boldsymbol{\theta} = \mathbf{0}$) against sparse alternatives in a special case of the model (3) (where $d = N$ and the design matrix $\mathbf{Z} = \mathbf{I}_N$ is the identity). Given this result, it is natural to wonder how estimation of the parameter $\boldsymbol{\theta}$ in model (3) depends on β and the network structure, when $\|\boldsymbol{\theta}\|_0 = s$. Towards this, we study the properties of the following PMPL estimator for the regression coefficients $\boldsymbol{\theta}$:

$$\hat{\boldsymbol{\theta}} := \arg \min_{\boldsymbol{\theta}} \{L_{\beta,N}(\boldsymbol{\theta}) + \lambda \|\boldsymbol{\theta}\|_1\}, \quad (8)$$

where λ is a regularization parameter and (recalling (5))

$$L_{\beta,N}(\boldsymbol{\theta}) = -\frac{1}{N} \sum_{i=1}^N \{X_i(\boldsymbol{\theta}^\top \mathbf{Z}_i + \beta m_i(\mathbf{X})) - \log \cosh(\boldsymbol{\theta}^\top \mathbf{Z}_i + \beta m_i(\mathbf{X}))\} + \log 2. \quad (9)$$

To handle the high-dimensional regime we assume that the design matrix \mathbf{Z} satisfies the following restricted eigenvalue condition:

Assumption 4. *There exists a constant $\kappa > 0$ such that*

$$\liminf_{N \rightarrow \infty} \inf_{\boldsymbol{\theta} \in \mathcal{C}(s)} \frac{\|\mathbf{Z}\boldsymbol{\theta}\|_2^2}{N\|\boldsymbol{\theta}\|_2^2} > \kappa,$$

where S denotes the support of the true parameter $\boldsymbol{\theta}$ and $\mathcal{C}(S) := \{\boldsymbol{\Delta} \in \mathbb{R}^d : \|\boldsymbol{\Delta}_{S^c}\|_1 \leq 3\|\boldsymbol{\Delta}_S\|_1\}$.

This type of condition is standard in the high-dimensional statistics literature (see [8, 56, 61, 66, 77] and the references therein), which is known to hold for many natural classes of design matrices. For example, if the design matrix has i.i.d. entries following a sub-Gaussian distribution, the restricted isometry property [18] holds with high probability, which in turn implies the restricted eigenvalue condition [8, 77]. More generally, design matrices with i.i.d. sub-Gaussian rows satisfy the restricted eigenvalue condition with high-probability [66, 68].

The following theorem shows that even under network dependence, one can consistently estimate the regression parameters of the model (3) at the same rate as the classical (independent) logistic regression model (1).

Theorem 2. *Fix $\beta \geq 0$. Suppose the interaction matrix \mathbf{A} in (3) satisfies $\sup_{N \geq 1} \|\mathbf{A}\|_\infty < \infty$, and Assumptions 1 and 4 hold. Then, there exists a constant $\delta > 0$ such that by choosing $\lambda := \delta \sqrt{\log d/N}$ in the objective function in (8) we have,*

$$\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}\|_2 = O_s \left(\sqrt{\frac{\log d}{N}} \right),$$

with probability $1 - o(1)$, as $N, d \rightarrow \infty$.

The proof of Theorem 2 is given in Section 4. While it follows the same outline as the proof of Theorem 1, contrary to Theorem 1, we can establish the strong concavity of the pseudo-likelihood even when d is much larger than N (up to $\log d = o(N)$), using the restricted eigenvalue condition on the design matrix. Consequently, we can establish the consistency of the PMPL estimate of the regression parameters in the entire high-dimensional regime. Moreover, as discussed in Remark 1, we can also recover the classical $\sqrt{s \log d/n}$ rate for estimating the regression parameters (as in the independent high-dimensional logistic regression model) by assuming a stronger ℓ_2 -boundedness condition.

2.3. Application to Various Network Structures. In this section, we apply Theorem 1 to establish the consistency of the PMPL estimator (16) for various natural network models. To this end, let $G_N = (V(G_N), E(G_N))$ be a sequence of graphs with $V(G_N) = [N] := \{1, 2, \dots, N\}$ and adjacency matrix $\mathcal{A}(G_N) = ((a_{ij}))_{1 \leq i, j \leq N}$. We denote by d_v the degree of the vertex $v \in V(G_N)$. To ensure that the model (3) has non-trivial scaling properties, one needs to choose the interaction matrix \mathbf{A} as the scaled adjacency matrix of G_N . In particular, define

$$\mathbf{A}_{G_N} := \frac{N}{2|E(G_N)|} \cdot \mathcal{A}(G_N) \tag{10}$$

Throughout this section we consider the model (3) with $\mathbf{A} = \mathbf{A}_{G_N}$ as above. We also assume that the number of non-isolated vertices in G_N (that is, the number of vertices in G_N with degree at least 1) is $\Omega(N)$. Note that this implies, $|E(G_N)| \gtrsim N$. Finally, we also assume that the sparsity $s = O(1)$ and consequently, absorb the dependence on s in the O -terms (recall Remark 1).

2.3.1. Bounded Degree Graphs. A sequence of graphs $\{G_N\}_{N \geq 1}$ is said to have bounded maximum degree if its maximum degree is uniformly bounded, that is, $\sup_{N \geq 1} d_{\max} < \infty$, where $d_{\max} := \max_{v \in V(G_N)} d_v$ is the maximum degree of G_N . Note that if G_N has bounded maximum degree and has $\Omega(N)$ non-isolated vertices, then $|E(G_N)| = \Theta(N)$.

Networks arising in certain applications, especially those with an underlying spatial or lattice structure generally have bounded degree. These include planar maps which encode neighborhood relations [5, 47], lattice models for capturing nearest-neighbor interactions between image pixels,

and demand-aware networks [2] among others. It is easy to check that the conditions in Assumption 1 are satisfied for bounded degree graphs. Towards this, note that, under the scaling in (10), the condition $\sup_{N \geq 1} \|\mathbf{A}\|_\infty < \infty$ is equivalent to $\|\mathbf{A}_{G_N}\|_\infty = \frac{N}{2|E(G_N)|} \max_{v \in V(G_n)} d_v$. Hence, under the scaling in (10), the assumption $\sup_{N \geq 1} \|\mathbf{A}_{G_N}\|_\infty < \infty$ is equivalent to

$$d_{\max} := \max_{v \in V(G_n)} d_v = O\left(\frac{|E(G)|}{N}\right), \quad (11)$$

that is, the maximum degree of G_N is of the same order as its average degree. Moreover, the condition $\liminf_{N \rightarrow \infty} \frac{1}{N} \|\mathbf{A}_{G_N}\|_F^2 > 0$ is equivalent to

$$\limsup_{N \rightarrow \infty} \frac{|E(G_N)|}{N} < \infty, \quad (12)$$

that is, the average degree of G_N is bounded. Therefore, (11) and (12) are together equivalent to the condition that G_N has bounded maximum degree. Hence, the PMPL estimate is $\sqrt{\log d/N}$ -consistent for any sequence of graphs of bounded maximum degree, whenever the assumptions in Theorem 1 hold.

2.3.2. Sparse Inhomogeneous Random Graphs. Although the discussion above shows that the graph G_N has to have bounded maximum degree for Assumption 1 to hold, our proofs can be easily adapted to establish similar rates of consistency of the PMPL estimate (up to polylog(N) factors), if the maximum degree G_N grows sufficiently slowly. This is summarized in the following result.

Theorem 3. *Suppose $\{G_N\}_{N \geq 1}$ is a sequence of graphs with $|E(G_N)| = O(N)$, $d_{\max} = \tilde{O}(1)$, and $\Omega(N)$ non-isolated vertices. Then for $\lambda = \tilde{O}\left(\sqrt{\log d/N}\right)$,*

$$\|\hat{\gamma} - \gamma\|_2 = \tilde{O}\left(\frac{1}{\sqrt{N}}\right)$$

with probability $1 - o(1)$ as $N, d \rightarrow \infty$ such that $d = o(N)$.

Theorem 3 is proved in Section 5.1. Note that Theorem 3 shows that the PMPL estimator is \sqrt{N} -consistent (up to polylog factors) whenever the maximum degree of G_N grows poly-logarithmically with the number of vertices N . In fact, from the proof of Theorem 3, it becomes evident that the PMPL estimator remains consistent even if the maximum degree grows at a slow polynomial rate (that is, $d_{\max} = O(N^\alpha)$ for $\alpha > 0$ sufficiently small), albeit with an additional polynomial (in N) factor in the rate of convergence.

Theorem 3 can be applied to obtain rates of convergence in sparse inhomogeneous random graph models.

Definition 1. [9] *Given a symmetric matrix $\mathbf{P}^{(N)} = ((p_{uv})) \in [0, 1]^{N \times N}$ with zeroes on the diagonal, the inhomogeneous random graph $\mathcal{G}(N, \mathbf{P}^{(N)})$ is the graph with vertex set $[N] := \{1, 2, \dots, N\}$ where the edge (u, v) is present with probability p_{uv} , independent of the other edges, for every $1 \leq u < v \leq N$.*

The class of inhomogeneous random graph models defined above includes several popular network models, such as the Chung-Lu model [25], the β -model [22], random dot product graphs [74], and stochastic block models [53]. Here, we will consider the sparse regime where

$$\max_{1 \leq u, v \leq N} p_{uv} = O\left(\frac{1}{N}\right). \quad (13)$$

In this regime, the expected degree remains bounded, although the maximum degree can diverge at rate $O(\log N)$ [10, 51]. We will also assume that there exists $\varepsilon \in (0, 1)$ and $\Omega(N)$ vertices $u \in G_N$, such that

$$\limsup_{N \rightarrow \infty} \prod_{v=1}^N (1 - p_{uv}) < \varepsilon. \quad (14)$$

This will ensure $\mathcal{G}(N, \mathbf{P}^{(N)})$ has $\Omega(N)$ non-isolated vertices. Under these assumptions we have the following result:

Corollary 1. *Suppose G_N is a realization of the inhomogeneous random graph $\mathcal{G}(N, \mathbf{P}^{(N)})$, where $\mathbf{P}^{(N)}$ satisfies the conditions in (13) and (14). Then for $\lambda = \tilde{O}\left(\sqrt{\log(d+1)/N}\right)$,*

$$\|\hat{\gamma} - \gamma\|_2 = \tilde{O}\left(\frac{1}{\sqrt{N}}\right),$$

with probability $1 - o(1)$ as $N, d \rightarrow \infty$ such that $d = o(N)$.

Corollary 1 is proved in Section 5.2. In the following example we illustrate how it can be applied to the sparse stochastic block models, in particular, sparse Erdős-Rényi random graphs.

Example 1 (Sparse stochastic block models). Fix $K \geq 1$, a vector of community proportions $\boldsymbol{\lambda} := (\lambda_1, \dots, \lambda_K) \in (0, 1)^K$, such that $\sum_{j=1}^K \lambda_j = 1$, and a symmetric probability matrix $\mathbf{B} := ((b_{ij}))_{1 \leq i, j \leq K}$, where $b_{ij} \in [0, 1]$, for all $1 \leq i, j \leq K$ and $b_{ij} > 0$ for some $1 \leq i, j \leq K$. The (sparse) stochastic block model with proportion vector $\boldsymbol{\lambda}$ and probability matrix $\boldsymbol{\Theta}$ is the inhomogeneous random graph $\mathcal{G}(N, \mathbf{P}^{(N)})$, with $\mathbf{P}^{(N)} = ((p_{uv}))_{1 \leq u, v \leq N}$ where

$$p_{uv} = \frac{b_{ij}}{N} \quad \text{for } (u, v) \in B_i \times B_j, \quad (15)$$

where $B_j := (N \sum_{i=1}^{j-1} \lambda_i, N \sum_{i=1}^j \lambda_i] \cap [N]$, for $j \in \{1, \dots, K\}$. In other words, the set of vertices is divided into K blocks (communities) B_1, B_2, \dots, B_K , such that the edge between vertices $u \in B_i$ and $v \in B_j$ occurs independently with probability b_{ij}/N . Clearly, in this case (13) holds. Next, to check that (14) holds, choose $1 \leq i, j \leq K$ such that $b_{ij} > 0$. Then for all $u \in B_i$,

$$\limsup_{N \rightarrow \infty} \prod_{v=1}^N (1 - p_{uv}) \leq \limsup_{N \rightarrow \infty} \prod_{v \in B_j} \left(1 - \frac{b_{ij}}{N}\right) = \exp(-\lambda_j b_{ij}) < 1,$$

which verifies (14), since $|B_j| = \Omega(N)$. Hence, by Corollary 1, the PMPL estimate (6) is $\tilde{O}(1/\sqrt{N})$ -consistent in this example. As a consequence, the PMPL estimate is $\tilde{O}(1/\sqrt{N})$ -consistent for sparse Erdős-Rényi random graphs $G(N, c/N)$, which corresponds to setting $b_{ij} = c$, for all $1 \leq i, j \leq K$ in (15).

3. PROOF OF THEOREM 1

Recall from (6) that our PMPL estimator is defined as:

$$(\hat{\beta}, \hat{\boldsymbol{\theta}}^\top) := \operatorname{argmin}_{(\beta, \boldsymbol{\theta}^\top) \in \mathbb{R}^{d+1}} L_N(\beta, \boldsymbol{\theta}) + \lambda \left\| (\beta, \boldsymbol{\theta}^\top) \right\|_1 \quad (16)$$

where $\lambda > 0$ is a tuning parameter and

$$L_N(\beta, \boldsymbol{\theta}) = \frac{1}{N} \sum_{i=1}^N \left[\log \cosh \left(\beta \sum_{j=1}^N A_{ij} X_j + \boldsymbol{\theta}^\top \mathbf{Z}_i \right) - X_i \left(\beta \sum_{j=1}^N A_{ij} X_j + \boldsymbol{\theta}^\top \mathbf{Z}_i \right) \right],$$

is as defined in (5) (where we have dropped the additive factor of $\log 2$). To begin, note that since Assumption 1 holds, by scaling the interaction matrix and the covariate vectors by $\|\mathbf{A}\|_\infty$ we can assume without loss of generality,

$$\sup_{N \geq 1} \|\mathbf{A}\|_\infty \leq 1, \quad (17)$$

The first step towards the proof of Theorem 1 is to establish the concentration of the pseudo-likelihood gradient vector $\nabla L_N(\hat{\gamma})$ in the ℓ_∞ norm. This is formalized in the following lemma which is proved in Section 3.1.

Lemma 1 (Concentration of the gradient). *For $\hat{\gamma}$ and any $\lambda > 0$,*

$$\|\nabla L_N(\hat{\gamma})\|_\infty \leq \lambda. \quad (18)$$

Moreover, there exists $\delta > 0$ such that with $\lambda := \delta \sqrt{\log(d+1)/N}$ the following holds:

$$\mathbb{P}\left(\|\nabla L_N(\gamma)\|_\infty > \frac{\lambda}{2}\right) = o(1), \quad (19)$$

where the $o(1)$ -term goes to infinity as $d \rightarrow \infty$.

The next lemma shows that the pseudo-likelihood function is strongly concave with high probability. The proof of this lemma is given in Section 3.2.

Lemma 2 (Strong concavity of pseudo-likelihood). *Suppose the assumptions of Theorem 1 hold. Then, there exists a constant $\kappa := \kappa(s, M, \beta, \Theta) > 0$, such that*

$$L_N(\hat{\gamma}) - L_N(\gamma) - \nabla L_N(\gamma)^\top (\hat{\gamma} - \gamma) \geq \kappa \|\hat{\gamma} - \gamma\|_2^2,$$

with probability $1 - o(1)$.

The proof of Theorem 1 can now be easily completed using the above lemmas. Towards this define:

$$S := \{1 \leq i \leq d+1 : \gamma_i \neq 0\}.$$

Moreover, for any vector $\mathbf{a} \in \mathbb{R}^p$ and any set $Q \subseteq \{1, \dots, p\}$, we denote by \mathbf{a}_Q the vector $(a_i)_{i \in Q}$. Now, for the constant κ as in Lemma 2, consider the event

$$\mathcal{E}_N := \left\{ \mathbf{X} \in \mathcal{C}_N : \|\nabla L_N(\gamma)\|_\infty \leq \frac{\lambda}{2} \text{ and } L_N(\hat{\gamma}) - L_N(\gamma) - \nabla L_N(\gamma)^\top (\hat{\gamma} - \gamma) \geq \kappa \|\hat{\gamma} - \gamma\|_2^2 \right\}.$$

Clearly, from Lemma 1 and Lemma 2, $\mathbb{P}(\mathcal{E}_N^c) = o(1)$.

Next, suppose $\mathbf{X} \in \mathcal{E}_N$. From the definition of $\hat{\gamma}$ it follows that

$$L_N(\hat{\gamma}) + \lambda \|\hat{\gamma}\|_1 \leq L_N(\gamma) + \lambda \|\gamma\|_1. \quad (20)$$

Hence,

$$\begin{aligned} \lambda(\|\gamma\|_1 - \|\hat{\gamma}\|_1) &\geq L_N(\hat{\gamma}) - L_N(\gamma) \\ &= \nabla L_N(\gamma)^\top (\hat{\gamma} - \gamma) + (L_N(\hat{\gamma}) - L_N(\gamma) - \nabla L_N(\gamma)^\top (\hat{\gamma} - \gamma)) \\ &\geq -\|\nabla L_N(\gamma)\|_\infty \|\hat{\gamma} - \gamma\|_1 + (L_N(\hat{\gamma}) - L_N(\gamma) - \nabla L_N(\gamma)^\top (\hat{\gamma} - \gamma)) \\ &\geq -\frac{\lambda \|\hat{\gamma} - \gamma\|_1}{2} + (L_N(\hat{\gamma}) - L_N(\gamma) - \nabla L_N(\gamma)^\top (\hat{\gamma} - \gamma)), \end{aligned} \quad (21)$$

where the last step uses $\|\nabla L_N(\gamma)\|_\infty \leq \frac{\lambda}{2}$, for $\mathbf{X} \in \mathcal{E}_N$. Next, note that

$$\begin{aligned} \|\hat{\gamma}\|_1 &= \|\gamma_S + (\hat{\gamma} - \gamma)_S\|_1 + \|(\hat{\gamma} - \gamma)_{S^c}\|_1 \geq \|\gamma_S\|_1 - \|(\hat{\gamma} - \gamma)_S\|_1 + \|(\hat{\gamma} - \gamma)_{S^c}\|_1 \\ &= \|\gamma\|_1 - \|(\hat{\gamma} - \gamma)_S\|_1 + \|(\hat{\gamma} - \gamma)_{S^c}\|_1. \end{aligned}$$

This implies,

$$\|(\hat{\gamma} - \gamma)_S\|_1 - \|(\hat{\gamma} - \gamma)_{S^c}\|_1 \geq \|\gamma\|_1 - \|\hat{\gamma}\|_1. \quad (22)$$

Combining (21) and (22) it follows that

$$\begin{aligned} \lambda \left(\|(\hat{\gamma} - \gamma)_S\|_1 - \|(\hat{\gamma} - \gamma)_{S^c}\|_1 + \frac{\|\hat{\gamma} - \gamma\|_1}{2} \right) &\geq L_N(\hat{\gamma}) - L_N(\gamma) - \nabla L_N(\gamma)^\top (\hat{\gamma} - \gamma) \\ &\geq \kappa \|\hat{\gamma} - \gamma\|_2^2, \end{aligned} \quad (23)$$

where the last inequality uses $L_N(\hat{\gamma}) - L_N(\gamma) - \nabla L_N(\gamma)^\top (\hat{\gamma} - \gamma) \geq \kappa \|\hat{\gamma} - \gamma\|_2^2$, for $\mathbf{X} \in \mathcal{E}_N$. Using $\|\hat{\gamma} - \gamma\|_1 = \|(\hat{\gamma} - \gamma)_S\|_1 + \|(\hat{\gamma} - \gamma)_{S^c}\|_1$ in the LHS of (23) now gives,

$$\begin{aligned} \|\hat{\gamma} - \gamma\|_2^2 &\leq \frac{\lambda}{\kappa} \left(\frac{3\|(\hat{\gamma} - \gamma)_S\|_1}{2} - \frac{\|(\hat{\gamma} - \gamma)_{S^c}\|_1}{2} \right) \lesssim_{\kappa} \lambda \|(\hat{\gamma} - \gamma)_S\|_1 \\ &\lesssim_{\kappa, s} \lambda \|(\hat{\gamma} - \gamma)_S\|_2 \\ &\leq \lambda \|\hat{\gamma} - \gamma\|_2. \end{aligned}$$

This implies, for $\mathbf{X} \in \mathcal{E}_N$,

$$\|\hat{\gamma} - \gamma\|_2 = O_{\kappa, s, \delta} \left(\sqrt{\frac{\log d}{N}} \right).$$

This completes the proof of Theorem 1, since $\mathbb{P}(\mathcal{E}_N) = 1 - o(1)$.

3.1. Proof of Lemma 1. First, we establish that $\|\nabla L_N(\hat{\gamma})\|_\infty \leq \lambda$. Fix $1 \leq i \leq d+1$ and define the univariate function:

$$f(x) := L_N(\hat{\gamma}_1, \dots, \hat{\gamma}_{i-1}, x, \hat{\gamma}_{i+1}, \dots, \hat{\gamma}_{d+1}).$$

Note that $f'(\hat{\gamma}_i) = \frac{\partial}{\partial \underline{\gamma}_i} \nabla L_N(\underline{\gamma})|_{\underline{\gamma}=\hat{\gamma}}$. Now, by the definition of $\hat{\gamma}$ we have, $f(\hat{\gamma}_i) + \lambda|\hat{\gamma}_i| \leq f(x) + \lambda|x|$, which implies,

$$f(x) - f(\hat{\gamma}_i) \geq \lambda(|\hat{\gamma}_i| - |x|).$$

Next, consider the following cases:

- $\hat{\gamma}_i > 0$: Then, for all $x > \hat{\gamma}_i$,

$$\frac{f(x) - f(\hat{\gamma}_i)}{x - \hat{\gamma}_i} \geq \lambda \frac{|\hat{\gamma}_i| - |x|}{x - \hat{\gamma}_i} = -\lambda.$$

Similarly, for all $0 < x < \hat{\gamma}_i$, $\frac{f(x) - f(\hat{\gamma}_i)}{x - \hat{\gamma}_i} \leq \lambda \frac{|\hat{\gamma}_i| - |x|}{x - \hat{\gamma}_i} = -\lambda$. This implies, $f'(\hat{\gamma}_i) = -\lambda$.

- $\hat{\gamma}_i < 0$: Then, for all $0 > x > \hat{\gamma}_i$,

$$\frac{f(x) - f(\hat{\gamma}_i)}{x - \hat{\gamma}_i} \geq \lambda \frac{|\hat{\gamma}_i| - |x|}{x - \hat{\gamma}_i} = \lambda.$$

Similarly, $x < \hat{\gamma}_i$, $\frac{f(x) - f(\hat{\gamma}_i)}{x - \hat{\gamma}_i} \leq \lambda \frac{|\hat{\gamma}_i| - |x|}{x - \hat{\gamma}_i} = \lambda$. Hence, in this case, $f'(\hat{\gamma}_i) = \lambda$.

- $\hat{\gamma}_i = 0$. In this case, for all $x > 0$,

$$\frac{f(x) - f(0)}{x} \geq -\lambda \frac{|x|}{x} = -\lambda$$

and for all $x < 0$, $\frac{f(x) - f(0)}{x} \leq -\lambda \frac{|x|}{x} = \lambda$. Since f' exists, this implies that $|f'(0)| \leq \lambda$.

Combining the above three cases, it follows that

$$\|\nabla L_N(\hat{\gamma})\|_\infty = \max_{j \in [d+1]} |f'(\hat{\gamma}_j)| \leq \lambda,$$

completing the proof of (18)

Next, we establish the concentration of $\|\nabla L_N(\gamma)\|_\infty$ as in (19). For this step, we require the following definitions. For $1 \leq i \leq N$, denote

$$m_i(\mathbf{X}) := \sum_{j=1}^N a_{ij} X_j.$$

Next, define functions $\phi_i : \mathcal{C}_N \rightarrow \mathbb{R}$, for $1 \leq i \leq N$, as follows:

$$\phi_i(\mathbf{x}) := -\frac{1}{N} \left\{ m_i(\mathbf{x}) \left(X_i - \tanh(\beta m_i(\mathbf{x}) + \boldsymbol{\theta}^\top \mathbf{Z}_i) \right) \right\}, \quad (24)$$

for $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathcal{C}_N$. Similarly, define functions $\phi_{i,s} : \mathcal{C}_N \rightarrow \mathbb{R}$, for $1 \leq i \leq N$ and $1 \leq s \leq d$, as follows:

$$\phi_{i,s}(\mathbf{x}) := -\frac{1}{N} \left\{ Z_{i,s} \left(x_i - \tanh(\beta m_i(\mathbf{x}) + \boldsymbol{\theta}^\top \mathbf{Z}_i) \right) \right\}. \quad (25)$$

Note that $\nabla L_N = (\frac{\partial L_N}{\partial \beta}, \frac{\partial L_N}{\partial \theta_1}, \dots, \frac{\partial L_N}{\partial \theta_d})^\top$ where

$$\frac{\partial L_N}{\partial \beta} = \sum_{i=1}^N \phi_i(\mathbf{X}) \quad \text{and} \quad \frac{\partial L_N}{\partial \theta_s} = \sum_{i=1}^N \phi_{i,s}(\mathbf{X}), \quad \text{for } 1 \leq s \leq d.$$

To establish the concentration of $\|\nabla L_N(\gamma)\|_\infty$, we use the conditioning trick from [29], which allows to reduce the model (3) to an Ising model in the Dobrushin regime (where the correlations are sufficiently weak and the model approximately behaves like a product measure), by conditioning on a subset of the nodes. To describe this, we need the following definition:

Definition 2. Suppose that $\boldsymbol{\sigma} \in \{-1, 1\}^N$ is a sample from the Ising model:

$$\mathbb{P}_{\beta, \mathbf{h}}(\boldsymbol{\sigma}) \propto \exp \left(\boldsymbol{\sigma}^\top \mathbf{D} \boldsymbol{\sigma} + \sum_{i=1}^N h_i \sigma_i \right), \quad (26)$$

where $\mathbf{h} = (h_1, h_2, \dots, h_n)^\top \in \mathbb{R}^n$ and \mathbf{D} is a symmetric matrix with zeros on the diagonal with $\sup_{N \geq 1} \|\mathbf{D}\|_\infty \leq R$. Moreover, suppose that with probability 1,

$$\min_{1 \leq i \leq N} \text{Var}(\sigma_i | \boldsymbol{\sigma}_{-i}) \geq \Upsilon,$$

for some $\Upsilon \geq 0$. Then, the model (26) is referred to as an (R, Υ) -Ising model.

Recently, Dagan et al. [29] developed a technique for reducing an (R, Υ) -Ising model to an (η, Υ) -Ising model, for $0 < \eta < R$, by conditioning on a subset of vertices. As a consequence, by choosing η one can ensure that the conditional model is in the Dobrushin high-temperature regime. Although the Ising model studied in [29] is different, the same proof extends to our model (3) as well. We formalize this in the following lemma, which is proved in Appendix A.2.

Lemma 3. Fix $R > 0$ and $\eta \in (0, R)$. Let $\mathbf{X} \in \{-1, 1\}^N$ be a sample from an (R, Υ) -Ising model. Then there exist subsets $I_1, \dots, I_\ell \subseteq [N]$ with $\ell \lesssim R^2 \log N / \eta^2$ such that:

(1) For all $1 \leq i \leq N$,

$$|\{j \in [\ell] : i \in I_j\}| = \lceil \eta \ell / 8R \rceil$$

(2) For all $1 \leq j \leq \ell$, the conditional distribution of \mathbf{X}_{I_j} given $\mathbf{X}_{I_j^c} := (X_u)_{u \in [N] \setminus I_j}$ is an (η, Υ) -Ising model.

Furthermore, for any non-negative vector $\mathbf{a} \in \mathbb{R}^N$, there exists $j \in \ell$ such that

$$\sum_{i \in I_j} a_i \geq \frac{\eta}{8R} \sum_{i=1}^N a_i. \quad (27)$$

We will apply the above result to our model (3). Towards this, set $\mathbf{D} = \beta \mathbf{A}$ and $h_i = \boldsymbol{\theta}^\top \mathbf{Z}_i$, for $1 \leq i \leq N$ in (26). Under this parametrization, (3) is an (R, Υ) -Ising model as shown below:

Lemma 4. *The model (3) is a (R, Υ) -Ising model with $R = \beta$ and any $\Upsilon = e^{-4\Theta Ms - 4\beta}$.*

Proof. Note that for every $j \in [N]$,

$$\text{Var}(X_j | \mathbf{X}_{[N] \setminus \{j\}}) = 4p(1-p),$$

where $p = \mathbb{P}(X_j = 1 | \mathbf{X}_{[N] \setminus \{j\}})$. Now, denoting the elements of the matrix \mathbf{D} as $(d_{ij})_{1 \leq i, j \leq N}$, note that

$$p = \frac{\exp\left(h_j + \sum_{v \in [N] \setminus \{j\}} d_{jv} X_v\right)}{2 \cosh\left(h_j + \sum_{v \in [N] \setminus \{j\}} d_{jv} X_v\right)}.$$

Then using the inequality $\frac{e^x}{2 \cosh(x)} \geq \frac{1}{2} e^{-2|x|}$ gives,

$$\min\{p, 1-p\} \geq \frac{1}{2} \exp\left(-2 \left| h_j + \sum_{v \in [N] \setminus \{j\}} d_{jv} X_v \right| \right). \quad (28)$$

Next, using

$$\left| h_j + \sum_{v \in [N] \setminus \{j\}} d_{jv} X_v \right| = \left| \boldsymbol{\theta}^\top \mathbf{Z}_j + \sum_{v \in [N] \setminus \{j\}} d_{jv} X_v \right| \leq |\boldsymbol{\theta}^\top \mathbf{Z}_j| + \|\mathbf{D}\|_\infty \leq \Theta Ms + \beta,$$

it follows from (28) that

$$\min\{p, 1-p\} \geq \frac{1}{2} e^{-2\Theta Ms - 2\beta}.$$

Hence, we have:

$$\text{Var}(X_j | \mathbf{X}_{[N] \setminus \{j\}}) \geq e^{-4\Theta Ms - 4\beta}.$$

This completes the proof of the lemma, since $\|\mathbf{D}\|_\infty \leq \beta$ (since $\|\mathbf{A}\|_\infty \leq 1$ by (17)). \square

By the above lemma, model (3) is an (R, Υ) -Ising model, with $R := \beta$ and $\Upsilon = e^{-4\Theta Ms - 4\beta}$.

Next, choose

$$\eta := \min\left\{\frac{1}{16}, \beta\right\},$$

and suppose I_1, \dots, I_ℓ be subsets of $[N]$ as in Lemma 3. Then, defining $\ell' := \lceil \eta \ell / R \rceil$ we get,

$$\left| \sum_{i=1}^N \phi_i(\mathbf{X}) \right| = \left| \frac{1}{\ell'} \sum_{r=1}^{\ell'} \sum_{i \in I_r} \phi_i(\mathbf{X}) \right| \leq \frac{1}{\ell'} \sum_{r=1}^{\ell'} \left| \sum_{i \in I_r} \phi_i(\mathbf{X}) \right| \leq \frac{\ell}{\ell'} \max_{r \in [\ell]} |Q_r(\mathbf{X})|, \quad (29)$$

where

$$Q_r(\mathbf{X}) := \sum_{i \in I_r} \phi_i(\mathbf{X}), \quad (30)$$

for $r \in [\ell]$. Similarly, it follows that, for $s \in [d]$,

$$\left| \sum_{i=1}^N \phi_{i,s}(\mathbf{X}) \right| = \left| \frac{1}{\ell'} \sum_{r=1}^{\ell} \sum_{i \in I_r} \phi_{i,s}(\mathbf{X}) \right| \leq \frac{1}{\ell'} \sum_{r=1}^{\ell} \left| \sum_{i \in I_r} \phi_{i,s}(\mathbf{X}) \right| \leq \frac{\ell}{\ell'} \max_{r \in [\ell]} |Q_{r,s}(\mathbf{X})|, \quad (31)$$

where

$$Q_{r,s}(\mathbf{X}) := \sum_{i \in I_r} \phi_{i,s}(\mathbf{X}). \quad (32)$$

The following lemma shows that the functions Q_r and $Q_{r,s}$ are Lipschitz in the Hamming metric. The proof is given in Appendix A.1.

Lemma 5. *For $r \in [\ell]$ and $s \in [d]$, let Q_r and $Q_{r,s}$ be as defined in (30) and (32), respectively. Then for any two vectors $\mathbf{X}, \mathbf{X}' \in \mathcal{C}_N$ differing in at most one coordinate, the following hold:*

(1) For $r \in [\ell]$,

$$|Q_r(\mathbf{X}) - Q_r(\mathbf{X}')| \leq \frac{2\beta + 6}{N}.$$

(2) Similarly, $r \in [\ell]$ and $s \in [d]$,

$$|Q_{r,s}(\mathbf{X}) - Q_{r,s}(\mathbf{X}')| \leq \frac{2M(1 + \beta)}{N}.$$

Using the above result together with Lemma 4, we can now establish the concentrations of $Q_r(\mathbf{X})$ and $Q_{r,s}(\mathbf{X})$, conditional on $\mathbf{X}_{I_r^c}$. To this end, recalling the definition of $\phi_i(\cdot)$ from (24) note that $\mathbb{E}(Q_r(\mathbf{X}) | \mathbf{X}_{I_r^c}) = 0$, for $r \in [\ell]$. Moreover, by Lemma 4, $\mathbf{X} | \mathbf{X}_{I_r^c}$ is an (η, Υ) -Ising model, where $\eta \leq \frac{1}{16}$. Therefore, since Q_r is $O_\beta(1/N)$ -Lipschitz (by Lemma 5), applying Theorem 4.3 and Lemma 4.4 in [20] gives, for every $t > 0$,

$$\mathbb{P}\left(|Q_r(\mathbf{X})| \geq t \mid \mathbf{X}_{I_r^c}\right) \leq 2e^{-O_\beta(Nt^2)}. \quad (33)$$

Similarly, recalling (25), it follows that for each $r \in [\ell]$ and $s \in [d]$, $\mathbb{E}(Q_{r,s}(\mathbf{X}) | \mathbf{X}_{I_r^c}) = 0$. Then, since $Q_{r,s}$ is $O_{\beta,M}(1/N)$ -Lipschitz (see Lemma 5), Theorem 4.3 and Lemma 4.4 in [20] gives, for every $t \geq 0$,

$$\mathbb{P}\left(|Q_{r,s}(\mathbf{X})| \geq t \mid \mathbf{X}_{I_r^c}\right) \leq 2e^{-O_{\beta,M}(Nt^2)}. \quad (34)$$

Hence, combining (29), (33), (31), (34), and Lemma 3 (which implies that $\ell = O(\log N)$) gives,

$$\mathbb{P}\left(\left|\sum_{i=1}^N \phi_i(\mathbf{X})\right| \geq t\right) \leq 2e^{-O_{\beta,M}(Nt^2)} \quad \text{and} \quad \mathbb{P}\left(\left|\sum_{i=1}^N \phi_{i,s}(\mathbf{X})\right| \geq t\right) \leq 2e^{-O_{\beta,M}(Nt^2)}, \quad (35)$$

for each $s \in [d]$. It thus follows from (35) and a union bound, that

$$\mathbb{P}(\|\nabla L_N(\gamma)\|_\infty \geq t) \leq 2(d+1)e^{-KNt^2}, \quad (36)$$

for some constant $K > 0$, depending only on β and M . Now, choosing $t = \frac{\lambda}{2} = \frac{1}{2}\delta\sqrt{\log(d+1)/N}$ in (36) above gives,

$$\mathbb{P}\left(\|\nabla L_N(\gamma)\|_\infty \geq \frac{\lambda}{2}\right) \leq 2(d+1)^{1-\frac{K\delta^2}{4}} = o(1),$$

whenever $\delta^2 < 4/K$. This completes the proof of Lemma 1.

3.2. Proof of Lemma 2. Define the following $(d+1) \times (d+1)$ dimensional matrix,

$$\mathbf{G} := \frac{1}{N} \begin{pmatrix} \mathbf{m}^\top \mathbf{m} & \mathbf{m}^\top \mathbf{Z} \\ \mathbf{Z}^\top \mathbf{m} & \mathbf{Z}^\top \mathbf{Z} \end{pmatrix}$$

The key step towards proving Lemma 2 is to show that the lowest eigenvalue of $\nabla^2 L_N$ is bounded away from 0 with high probability.

Lemma 6. *There exists a constant $C > 0$ (depending only on s, Θ and M), such that*

$$\mathbb{P}(\lambda_{\min}(\mathbf{G}) \geq C) \geq 1 - e^{-\Omega(N)}, \quad (37)$$

as $N, d \rightarrow \infty$, such that $d = o(N)$.

The proof of Lemma 6 is given in Section 3.2.1. We first show it can be used to complete the proof of Lemma 2. To this end, by a second order Taylor series expansion, we know that there exists $\alpha \in (0, 1)$ and $\underline{\gamma} = (\underline{\beta}, \underline{\boldsymbol{\theta}}^\top) = \alpha\gamma + (1-\alpha)\hat{\gamma}$ such that

$$\begin{aligned} L_N(\hat{\gamma}) - L_N(\gamma) - \nabla L_N(\gamma)^\top (\hat{\gamma} - \gamma) &= \frac{1}{2} (\hat{\gamma} - \gamma)^\top \nabla^2 L_N(\underline{\gamma}) (\hat{\gamma} - \gamma) \\ &= \frac{1}{2N} \sum_{i=1}^N \frac{(\hat{\gamma} - \gamma)^\top \mathbf{U}_i \mathbf{U}_i^\top (\hat{\gamma} - \gamma)}{\cosh^2(\underline{\beta} m_i(\mathbf{X}) + \underline{\boldsymbol{\theta}}^\top \mathbf{Z}_i)}, \end{aligned} \quad (38)$$

where $\mathbf{U}_i := (m_i(\mathbf{X}), \mathbf{Z}_i^\top)^\top$. Now, note that:

$$|\underline{\beta} m_i(\mathbf{X})| \leq |\underline{\beta}| |m_i(\mathbf{X})| \leq |\underline{\beta}| \|A\|_\infty \leq |\beta| + |\hat{\beta} - \beta| \leq |\beta| + \|\hat{\gamma} - \gamma\|_1 \quad (39)$$

and

$$\|\underline{\boldsymbol{\theta}}^\top \mathbf{Z}_i\| \leq \|\underline{\boldsymbol{\theta}}\|_1 \|\mathbf{Z}_i\|_\infty \leq M (\|\underline{\boldsymbol{\theta}} - \boldsymbol{\theta}\|_1 + \|\boldsymbol{\theta}\|_1) \leq M (\|\hat{\gamma} - \gamma\|_1 + s\Theta). \quad (40)$$

Since \cosh is an even function and increasing on the positive axis, we obtain

$$\frac{1}{2N} \sum_{i=1}^N \frac{(\hat{\gamma} - \gamma)^\top \mathbf{U}_i \mathbf{U}_i^\top (\hat{\gamma} - \gamma)}{\cosh^2(\underline{\beta} m_i(\mathbf{X}) + \underline{\boldsymbol{\theta}}^\top \mathbf{Z}_i)} \geq \frac{(\hat{\gamma} - \gamma)^\top \mathbf{G} \mathbf{v}}{2 \cosh^2(|\beta| + (M+1)(\|\hat{\gamma} - \gamma\|_1) + sM\Theta)}, \quad (41)$$

where $\mathbf{m} := (m_1(\mathbf{X}), \dots, m_N(\mathbf{X}))^\top$, $\mathbf{Z} = (\mathbf{Z}_1, \dots, \mathbf{Z}_N)^\top$, and

$$\mathbf{G} := \frac{1}{N} \begin{pmatrix} \mathbf{m}^\top \mathbf{m} & \mathbf{m}^\top \mathbf{Z} \\ \mathbf{Z}^\top \mathbf{m} & \mathbf{Z}^\top \mathbf{Z} \end{pmatrix}.$$

Combining (38) and (41) gives,

$$\begin{aligned} L_N(\hat{\gamma}) - L_N(\gamma) - \nabla L_N(\gamma)^\top (\hat{\gamma} - \gamma) &= \frac{1}{2} (\hat{\gamma} - \gamma)^\top \nabla^2 L_N(\underline{\gamma}) (\hat{\gamma} - \gamma) \\ &\geq \frac{(\hat{\gamma} - \gamma)^\top \mathbf{G} (\hat{\gamma} - \gamma)}{2 \cosh^2(|\beta| + (M+1)(\|\hat{\gamma} - \gamma\|_1) + sM\Theta)}. \end{aligned} \quad (42)$$

Next, we establish that $\|\hat{\gamma} - \gamma\|_1$ remains bounded with high probability whenever the conditions of Theorem 1 are satisfied.

Lemma 7. *Suppose (37) holds. Then, for $\lambda := \delta \sqrt{\log(d+1)/N}$ as in Lemma 1,*

$$\|\hat{\gamma} - \gamma\|_1 = O_s \left(\sqrt{\frac{\log d}{N}} \right).$$

with probability $1 - o(1)$, whenever $N, d \rightarrow \infty$ such that $\log d = o(N)$,

Proof. By the convexity of the function L_N it follows from (20) that

$$\begin{aligned}
\lambda(\|\gamma\|_1 - \|\hat{\gamma}\|_1) &\geq L_N(\hat{\gamma}) - L_N(\gamma) \\
&\geq \nabla L_N(\gamma)^\top (\hat{\gamma} - \gamma) \\
&\geq -\|\nabla L_N(\gamma)\|_\infty \|\hat{\gamma} - \gamma\|_1 \\
&\geq -\frac{\lambda\|\hat{\gamma} - \gamma\|_1}{2},
\end{aligned} \tag{43}$$

where the last step uses $\|\nabla L_N(\gamma)\|_\infty \leq \frac{\lambda}{2}$, for $\mathbf{X} \in \mathcal{E}_N$. Recall from (22) that $\|\hat{\gamma}\|_1 - \|\gamma\|_1 \geq \|(\hat{\gamma} - \gamma)_{S^c}\|_1 - \|(\hat{\gamma} - \gamma)_S\|_1$. Therefore, from (43), we have:

$$\|(\hat{\gamma} - \gamma)_{S^c}\|_1 - \|(\hat{\gamma} - \gamma)_S\|_1 \leq \|\hat{\gamma}\|_1 - \|\gamma\|_1 \leq \frac{\|\hat{\gamma} - \gamma\|_1}{2} = \frac{\|(\hat{\gamma} - \gamma)_S\|_1}{2} + \frac{\|(\hat{\gamma} - \gamma)_{S^c}\|_1}{2}.$$

This means, $\|(\hat{\gamma} - \gamma)_{S^c}\|_1 \leq 3\|(\hat{\gamma} - \gamma)_S\|_1$, and hence,

$$\|(\hat{\gamma} - \gamma)\|_1 = \|(\hat{\gamma} - \gamma)_S\|_1 + \|(\hat{\gamma} - \gamma)_{S^c}\|_1 \leq 4\|(\hat{\gamma} - \gamma)_S\|_1. \tag{44}$$

Next, for $t \in [0, 1]$, let $\gamma_t := t\hat{\gamma} + (1-t)\gamma$, and $g(t) := (\hat{\gamma} - \gamma)^\top \nabla L_N(\gamma_t)$. Then

$$|g(1) - g(0)| = |(\hat{\gamma} - \gamma)^\top (\nabla L_N(\hat{\gamma}) - \nabla L_N(\gamma))| \leq \|\hat{\gamma} - \gamma\|_1 \cdot \|\nabla L_N(\hat{\gamma}) - \nabla L_N(\gamma)\|_\infty. \tag{45}$$

Therefore,

$$\begin{aligned}
g'(t) &= (\hat{\gamma} - \gamma)^\top \nabla^2 L_N(\gamma_t) (\hat{\gamma} - \gamma) \\
&= \frac{1}{N} \sum_{i=1}^N \frac{(\hat{\gamma} - \gamma)^\top \mathbf{U}_i \mathbf{U}_i^\top (\hat{\gamma} - \gamma)}{\cosh^2(\beta_t m_i(\mathbf{X}) + \boldsymbol{\theta}_t^\top \mathbf{Z}_i)} \quad (\text{where } \mathbf{U}_i := (m_i(\mathbf{X}), \mathbf{Z}_i^\top)^\top) \\
&\geq \frac{(\hat{\gamma} - \gamma)^\top \mathbf{G} (\hat{\gamma} - \gamma)}{\cosh^2(|\beta| + (M+1)\|\gamma_t - \gamma\|_1 + sM\Theta)} \quad (\text{by (39) and (40)}) \\
&\geq \frac{\|\hat{\gamma} - \gamma\|_2^2 \lambda_{\min}(\mathbf{G})}{\cosh^2(|\beta| + (M+1)\|\gamma_t - \gamma\|_1 + sM\Theta)} \\
&\gtrsim \frac{\|\hat{\gamma} - \gamma\|_2^2}{\cosh^2(|\beta| + 4(M+1)\|(\hat{\gamma} - \gamma)_S\|_1 + Ms\Theta)},
\end{aligned}$$

where the last step uses (37) (which holds with probability $1 - o(1)$) and $\|\gamma_t - \gamma\|_1 = |t|\|(\hat{\gamma} - \gamma)\|_1 \leq 4\|(\hat{\gamma} - \gamma)_S\|_1$ (by (44)). Hence,

$$\begin{aligned}
|g(1) - g(0)| &\geq g(1) - g(0) = \int_0^1 g'(t) dt \\
&\geq \int_0^{\min\{1, \|(\hat{\gamma} - \gamma)_S\|_1^{-1}\}} g'(t) dt \\
&\gtrsim \|\hat{\gamma} - \gamma\|_2^2 \min\{1, \|(\hat{\gamma} - \gamma)_S\|_1^{-1}\}.
\end{aligned} \tag{46}$$

Combining (45) and (46) gives,

$$\begin{aligned}
\min\{\|(\hat{\gamma} - \gamma)_S\|_1, 1\} &\lesssim \frac{\|(\hat{\gamma} - \gamma)_S\|_1^2}{\|\hat{\gamma} - \gamma\|_2^2} \cdot \|\nabla L_N(\hat{\gamma}) - \nabla L_N(\gamma)\|_\infty \\
&\leq s \frac{\|(\hat{\gamma} - \gamma)_S\|_2^2}{\|\hat{\gamma} - \gamma\|_2^2} \|\nabla L_N(\hat{\gamma}) - \nabla L_N(\gamma)\|_\infty \\
&\leq s \|\nabla L_N(\hat{\gamma}) - \nabla L_N(\gamma)\|_\infty
\end{aligned} \tag{47}$$

Now, recall that, by Lemma 1, with probability $1 - o(1)$, $\|\nabla L_N(\boldsymbol{\gamma})\|_\infty \lesssim_\delta \sqrt{\log(d+1)/N}$ and $\|\nabla L_N(\hat{\boldsymbol{\gamma}})\|_\infty \lesssim \sqrt{\log(d+1)/N}$. Applying this in (47) shows that with probability $1 - o(1)$, $\min\{\|(\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})_S\|_1, 1\} = O_s(\sqrt{\log(d+1)/N})$. This implies, $\|(\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})_S\|_1 = O_s(\sqrt{\log(d+1)/N})$ with probability $1 - o(1)$, whenever $N, d \rightarrow \infty$ such that $\log d = o(N)$. Therefore, by (44), $\|\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}\|_1 = O_s(\sqrt{\log(d+1)/N})$ with probability $1 - o(1)$. \square

Using Lemma 6 and Lemma 7 in (42) it follows that, there exists κ (as the statement of Lemma 2) such that

$$\begin{aligned} L_N(\hat{\boldsymbol{\gamma}}) - L_N(\boldsymbol{\gamma}) - \nabla L_N(\boldsymbol{\gamma})^\top (\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}) &\geq \frac{(\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})^\top \mathbf{G} (\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})}{2 \cosh^2(|\beta| + (M+1)\|\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}\|_1 + sM\Theta)} \\ &\gtrsim_{\beta, M, \kappa} \|\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}\|_2^2 \end{aligned}$$

with probability $1 - o(1)$, as $N, d \rightarrow \infty$ such that $d = o(N)$. This completes the proof of Lemma 2.

3.2.1. *Proof of Lemma 6.* The first step towards proving Lemma 6 is to observe that:

$$\det(\mathbf{G} - \lambda \mathbf{I}) = \left(\frac{1}{N} \|\mathbf{F}\mathbf{m}\|_2^2 - \lambda \right) \cdot \det\left(\frac{1}{n} \mathbf{Z}^\top \mathbf{Z} - \lambda \mathbf{I} \right),$$

where $\mathbf{F} := \mathbf{I} - \mathbf{Z}(\mathbf{Z}^\top \mathbf{Z})^{-1} \mathbf{Z}^\top$. Hence,

$$\lambda_{\min}(\mathbf{G}) = \min \left\{ \lambda_{\min} \left(\frac{1}{N} \mathbf{Z}^\top \mathbf{Z} \right), \frac{1}{N} \|\mathbf{F}\mathbf{m}\|_2^2 \right\}.$$

In view of Assumption 2, to prove Lemma 6 it suffices to show that there exists a constant $C > 0$ (depending only on s, Θ and M), such that

$$\mathbb{P} \left(\frac{1}{N} \|\mathbf{F}\mathbf{m}\|_2^2 \geq C \right) = 1 - e^{-\Omega(N)}. \quad (48)$$

To this end, it suffices to prove the following conditional version of (48):

$$\mathbb{P} \left(\frac{1}{N} \|\mathbf{F}\mathbf{m}\|_2^2 \geq C \mid \mathbf{X}_{J^c} \right) = 1 - e^{-\Omega(N)} \quad (49)$$

where J is a suitably chosen subset of $[N]$ and $\mathbf{X}_{J^c} := (X_i)_{i \in J^c}$. To this end, note that (3) is a (R, Υ) -Ising model (Lemma 4). Now, applying Lemma 3 with

$$\eta := \min \left\{ \frac{1}{16}, \beta \|\mathbf{A}\|_\infty \right\},$$

gives subsets $I_1, \dots, I_\ell \subseteq [N]$ such that, for all $1 \leq j \leq \ell$, the conditional distribution of \mathbf{X}_{I_j} given $\mathbf{X}_{I_j^c} := (X_u)_{u \in [N] \setminus I_j}$ is an (η, Υ) -Ising model. Furthermore, for any vector \mathbf{a} , there exists $j \in [\ell]$ such that

$$\|\mathbf{a}_{I_j}\|_1 \geq \frac{\eta}{8\beta \|\mathbf{A}\|_\infty} \|\mathbf{a}\|_1. \quad (50)$$

The proof of (49) now proceeds in the following two steps: First, we show that there exists $j \in [\ell]$ such that the expectation of $N^{-1} \|\mathbf{F}\mathbf{m}\|_2^2$ conditioned on \mathbf{X}_{I_j} is $\Omega(1)$. Subsequently, we establish that conditioned on \mathbf{X}_{I_j} , $N^{-1} \|\mathbf{F}\mathbf{m}\|_2^2$ concentrates around its conditional expectation. These steps are verified in Lemmas 8 and 9, respectively.

Lemma 8. *Under the assumptions of Theorem 1, there exists $J \in \{I_1, I_2, \dots, I_\ell\}$ such that for all $N \geq 1$,*

$$\mathbb{E} \left(\frac{1}{N} \|\mathbf{F}\mathbf{m}\|_2^2 \middle| \mathbf{X}_{J^c} \right) \geq C,$$

where $C > 0$ is a constant depending only on Θ, M and s .

Proof. For any $(d+1) \times n$ dimensional matrix \mathbf{M} , we will denote the i -th row of \mathbf{M} by \mathbf{M}_i and the i -th largest singular value of \mathbf{M} by $\sigma_i(\mathbf{M})$, for $1 \leq i \leq d+1$. Also, for $J \subseteq [N]$ denote $(\mathbf{F}\mathbf{A})_{i,J} := ((\mathbf{F}\mathbf{A})_{i,j})_{j \in J}$. Note that for any $J \in \{I_1, I_2, \dots, I_\ell\} \subset [N]$, since $\mathbf{m} = \mathbf{A}\mathbf{X}$,

$$\begin{aligned} \mathbb{E} \left(\|\mathbf{F}\mathbf{m}\|_2^2 \middle| \mathbf{X}_{J^c} \right) &= \sum_{i=1}^N \mathbb{E} \left([(\mathbf{F}\mathbf{A})_i \mathbf{X}]^2 \middle| \mathbf{X}_{J^c} \right) \geq \sum_{i=1}^N \text{Var} \left((\mathbf{F}\mathbf{A})_i \mathbf{X} \middle| \mathbf{X}_{J^c} \right) \\ &= \sum_{i=1}^N \text{Var} \left((\mathbf{F}\mathbf{A})_{i,J} \mathbf{X}_J \middle| \mathbf{X}_{J^c} \right) \\ &\gtrsim \frac{\Upsilon^2}{\eta} \sum_{i=1}^N \|(\mathbf{F}\mathbf{A})_{i,J}\|_2^2, \end{aligned} \quad (51)$$

where the last step follows from Lemma 12. Now define a vector $\mathbf{a} = (a_1, a_2, \dots, a_N)^\top$, with $a_i = \|(\mathbf{F}\mathbf{A})_{\cdot,i}\|_2^2$, where $(\mathbf{F}\mathbf{A})_{\cdot,i}$ denotes the i -th column of the matrix $\mathbf{F}\mathbf{A}$. Then by (50) there exists $J \in \{I_1, I_2, \dots, I_\ell\} \subseteq [N]$ such that

$$\sum_{i=1}^N \|(\mathbf{F}\mathbf{A})_{i,J}\|_2^2 \geq \frac{\eta}{8\beta\|\mathbf{A}\|_\infty} \sum_{i=1}^N \|(\mathbf{F}\mathbf{A})_{\cdot,i}\|_2^2.$$

Therefore, by (51),

$$\mathbb{E} \left(\|\mathbf{F}\mathbf{m}\|_2^2 \middle| \mathbf{X}_{J^c} \right) \gtrsim \frac{\Upsilon^2}{\beta\|\mathbf{A}\|_\infty} \|\mathbf{F}\mathbf{A}\|_F^2 \gtrsim_{s,B,M} \|\mathbf{F}\mathbf{A}\|_F^2. \quad (52)$$

By Theorem 2 in [81], we get

$$\sum_{t=1}^N \sigma_t^2(\mathbf{F}\mathbf{A}) \geq \sum_{t=1}^N \sigma_t^2(\mathbf{F}) \sigma_{N-t+1}^2(\mathbf{A}) \quad (53)$$

Since \mathbf{F} is idempotent with trace $N-d$, it follows that $\sigma_1(\mathbf{F}) = \dots = \sigma_{N-d}(\mathbf{F}) = 1$ and $\sigma_{N-d+1}(\mathbf{F}) = \dots = \sigma_N(\mathbf{F}) = 0$. Hence, we have from (53),

$$\|\mathbf{F}\mathbf{A}\|_F^2 = \sum_{t=1}^N \sigma_t^2(\mathbf{F}\mathbf{A}) \geq \sum_{t=1}^{N-d} \sigma_{N-t+1}^2(\mathbf{A}) = \|\mathbf{A}\|_F^2 - \sum_{i=1}^d \sigma_i^2(\mathbf{A}). \quad (54)$$

where the last step uses $\sigma_i^2(\mathbf{A}) \leq 1$ for all $1 \leq i \leq N$, since $\|\mathbf{A}\|_2 \leq \|\mathbf{A}\|_\infty \leq 1$. Applying the bound in (54) to (52) gives,

$$\mathbb{E} \left(\|\mathbf{F}\mathbf{m}\|_2^2 \middle| \mathbf{X}_{J^c} \right) \gtrsim \Upsilon^2 (\|\mathbf{A}\|_F^2 - d). \quad (55)$$

Lemma 8 now follows from Assumption 1 and the hypothesis $d = o(N)$. \square

Next, we show that $\|\mathbf{F}\mathbf{m}\|_2^2$ concentrates around its conditional expectation $\mathbb{E}(\|\mathbf{F}\mathbf{m}\|_2^2 | \mathbf{X}_{J^c})$, for the set J as defined above.

Lemma 9. For any $t > 0$ and $J \in \{I_1, I_2, \dots, I_\ell\}$,

$$\begin{aligned} & \mathbb{P} \left(\|\mathbf{Fm}\|_2^2 < \mathbb{E}(\|\mathbf{Fm}\|_2^2 | \mathbf{X}_{J^c}) - t \mid \mathbf{X}_{J^c} \right) \\ & \leq 2 \exp \left(-C \cdot \min \left\{ \frac{t^2}{8N}, \frac{t}{2} \right\} \right) + 2 \exp \left(-\frac{t^2}{128N} \right), \end{aligned} \quad (56)$$

where C is a constant depending only on Θ, M and s .

Proof. Denote $\mathbf{W} := \mathbf{F}\mathbf{A}$ and let \mathbf{H} be the matrix obtained from $\mathbf{W}^\top \mathbf{W}$ by zeroing out all its diagonal elements. Moreover, throughout the proof we will denote $I := J^c$. Clearly,

$$\|\mathbf{Fm}\|_2^2 - \mathbb{E}(\|\mathbf{Fm}\|_2^2 | \mathbf{X}_I) = \mathbf{X}^\top \mathbf{H} \mathbf{X} - \mathbb{E} \left(\mathbf{X}^\top \mathbf{H} \mathbf{X} | \mathbf{X}_I \right).$$

By permuting the indices let us partition the vector \mathbf{X} as $(\mathbf{X}_I^\top, \mathbf{X}_J^\top)^\top$ and the matrix \mathbf{H} as:

$$\begin{pmatrix} \mathbf{H}_{I,I} & \mathbf{H}_{I,J} \\ \mathbf{H}_{I,J}^\top & \mathbf{H}_{J,J} \end{pmatrix}$$

where for two subsets A, B of $[N]$, we define $\mathbf{H}_{A,B} := ((H_{ij}))_{i \in A, j \in B}$. Note that

$$\begin{aligned} & \mathbf{X}^\top \mathbf{H} \mathbf{X} - \mathbb{E} \left(\mathbf{X}^\top \mathbf{H} \mathbf{X} | \mathbf{X}_I \right) \\ & = \mathbf{X}_J^\top \mathbf{H}_{J,J} \mathbf{X}_J - \mathbb{E} \left(\mathbf{X}_J^\top \mathbf{H}_{J,J} \mathbf{X}_J | \mathbf{X}_I \right) + 2\mathbf{X}_I^\top \mathbf{H}_{I,J} \mathbf{X}_J - 2\mathbb{E} \left(\mathbf{X}_I^\top \mathbf{H}_{I,J} \mathbf{X}_J | \mathbf{X}_I \right). \end{aligned} \quad (57)$$

Since $\mathbf{X} | \mathbf{X}_I$ is an (η, Υ) -Ising model, Example 2.5 in [1] implies (by taking the parameters α and ρ in [1] to be $\Theta Ms + 1/16$ and $7/8$, respectively),

$$\begin{aligned} & \mathbb{P} \left(\mathbf{X}_J^\top \mathbf{H}_{J,J} \mathbf{X}_J < \mathbb{E} \left(\mathbf{X}_J^\top \mathbf{H}_{J,J} \mathbf{X}_J | \mathbf{X}_I \right) - t \mid \mathbf{X}_I \right) \\ & \leq 2 \exp \left(-c \cdot \min \left\{ \frac{t^2}{\|\mathbf{H}_{J,J}\|_F^2 + \|\mathbb{E}(\mathbf{H}_{J,J} \mathbf{X}_J)\|_2^2}, \frac{t}{\|\mathbf{H}_{J,J}\|_2} \right\} \right), \end{aligned} \quad (58)$$

where c is a constant depending only on Θ, M and s . Clearly, one has $\|\mathbf{H}_{J,J}\|_F^2 \leq \|\mathbf{H}\|_F^2$ and, since the spectral norm of a matrix is always greater than or equal to the spectral norm of any of its submatrices, $\|\mathbf{H}_{J,J}\|_2 \leq \|\mathbf{H}\|_2$. Moreover, if

$$\mathbf{X}_0^\top := (\mathbf{0}^\top, \mathbf{X}_J^\top),$$

then $\mathbf{H}_{J,J} \mathbf{X}_J$ is a subvector of $\mathbf{H} \mathbf{X}_0$, and hence,

$$\|\mathbb{E}(\mathbf{H}_{J,J} \mathbf{X}_J)\|_2^2 \leq \|\mathbb{E}(\mathbf{H} \mathbf{X}_0)\|_2^2.$$

Combining all these, we have from (58)

$$\begin{aligned} & \mathbb{P} \left(\mathbf{X}_J^\top \mathbf{H}_{J,J} \mathbf{X}_J < \mathbb{E} \left(\mathbf{X}_J^\top \mathbf{H}_{J,J} \mathbf{X}_J | \mathbf{X}_I \right) - t \mid \mathbf{X}_I \right) \\ & \leq 2 \exp \left(-c \cdot \min \left\{ \frac{t^2}{\|\mathbf{H}\|_F^2 + \|\mathbb{E}(\mathbf{H} \mathbf{X}_0)\|_2^2}, \frac{t}{\|\mathbf{H}\|_2} \right\} \right) \end{aligned} \quad (59)$$

Next, note that, since $\mathbf{H} - \mathbf{W}^\top \mathbf{W}$ is a diagonal matrix,

$$\begin{aligned} \|\mathbf{H}\|_F^2 + \|\mathbb{E}(\mathbf{H} \mathbf{X}_0)\|_2^2 & \leq \|\mathbf{H}\|_F^2 + \left(\|\mathbb{E}(\mathbf{W}^\top \mathbf{W} \mathbf{X}_0)\|_2 + \|\mathbb{E}[(\mathbf{H} - \mathbf{W}^\top \mathbf{W}) \mathbf{X}_0]\|_2 \right)^2 \\ & \leq 2\|\mathbf{H}\|_F^2 + 2\mathbb{E}\|[(\mathbf{H} - \mathbf{W}^\top \mathbf{W}) \mathbf{X}_0]\|_2^2 + 2\|\mathbb{E}(\mathbf{W}^\top \mathbf{W} \mathbf{X}_0)\|_2^2 \end{aligned}$$

$$\begin{aligned}
&= 2 \sum_{1 \leq i \neq j \leq N} (\mathbf{W}^\top \mathbf{W})_{ij}^2 + 2 \sum_{i=1}^N (\mathbf{W}^\top \mathbf{W})_{ii}^2 + 2 \|\mathbb{E}(\mathbf{W}^\top \mathbf{W} \mathbf{X}_0)\|_2^2 \\
&= 2 \|\mathbf{W}^\top \mathbf{W}\|_F^2 + 2 \|\mathbb{E}(\mathbf{W}^\top \mathbf{W} \mathbf{X}_0)\|_2^2.
\end{aligned} \tag{60}$$

We also have $\|\mathbf{H}\|_2 \leq \|\mathbf{W}^\top \mathbf{W}\|_2$, since for any vector $\mathbf{a} \in \mathbb{R}^N$,

$$\mathbf{a}^\top \mathbf{H} \mathbf{a} = \mathbf{a}^\top \mathbf{W}^\top \mathbf{W} \mathbf{a} - \sum_{i=1}^N (\mathbf{W}^\top \mathbf{W})_{ii} a_i^2 \leq \mathbf{a}^\top \mathbf{W}^\top \mathbf{W} \mathbf{a}.$$

Hence, it follows from (59) and (60) that

$$\begin{aligned}
&\mathbb{P}\left(\mathbf{X}_J^\top \mathbf{H}_{J,J} \mathbf{X}_J < \mathbb{E}\left(\mathbf{X}_J^\top \mathbf{H}_{J,J} \mathbf{X}_J \mid \mathbf{X}_I\right) - t \mid \mathbf{X}_I\right) \\
&\leq 2 \exp\left(-c' \cdot \min\left\{\frac{t^2}{\|\mathbf{W}^\top \mathbf{W}\|_F^2 + \|\mathbb{E}(\mathbf{W}^\top \mathbf{W} \mathbf{X}_0)\|_2^2}, \frac{t}{\|\mathbf{W}^\top \mathbf{W}\|_2}\right\}\right)
\end{aligned} \tag{61}$$

where $c' := c/2$. Next, recall that for any two matrices \mathbf{U} and $(\hat{\gamma} - \gamma)$ such that $\mathbf{U}(\hat{\gamma} - \gamma)$ exists, we have $\|\mathbf{U}(\hat{\gamma} - \gamma)\|_F \leq \|\mathbf{U}\|_2 \|\hat{\gamma} - \gamma\|_F$. This implies,

$$\|\mathbf{W}^\top \mathbf{W}\|_F^2 \leq \|\mathbf{W}\|_2^2 \|\mathbf{W}\|_F^2, \tag{62}$$

Moreover, by the submultiplicativity of the matrix L^2 norm,

$$\|\mathbf{W}^\top \mathbf{W}\|_2 \leq \|\mathbf{W}^\top\|_2 \|\mathbf{W}\|_2 = \|\mathbf{W}\|_2^2 \tag{63}$$

and

$$\|\mathbb{E}(\mathbf{W}^\top \mathbf{W} \mathbf{X}_0)\|_2^2 = \|\mathbf{W}^\top \mathbb{E}(\mathbf{W} \mathbf{X}_0)\|_2^2 \leq \|\mathbf{W}\|_2^2 \cdot \|\mathbb{E}(\mathbf{W} \mathbf{X}_0)\|_2^2. \tag{64}$$

It follows from (61), (62), (63) and (64), that:

$$\begin{aligned}
&\mathbb{P}\left(\mathbf{X}_J^\top \mathbf{H}_{J,J} \mathbf{X}_J < \mathbb{E}\left(\mathbf{X}_J^\top \mathbf{H}_{J,J} \mathbf{X}_J \mid \mathbf{X}_I\right) - t \mid \mathbf{X}_I\right) \\
&\leq 2 \exp\left(-\frac{c'}{\|\mathbf{F} \mathbf{A}\|_2^2} \cdot \min\left\{\frac{t^2}{\|\mathbf{F} \mathbf{A}\|_F^2 + \|\mathbb{E}[\mathbf{W} \mathbf{X}_0]\|_2^2}, t\right\}\right).
\end{aligned} \tag{65}$$

Since $\|\mathbf{F} \mathbf{A}\|_2^2 \leq \|\mathbf{F}\|_2^2 \|\mathbf{A}\|_2^2 \leq 1$, $\|\mathbf{F} \mathbf{A}\|_F^2 \leq N \|\mathbf{F} \mathbf{A}\|_2^2 \leq N$, and $\|\mathbb{E}[\mathbf{W} \mathbf{X}_0]\|_2^2 \leq \mathbb{E}\|\mathbf{W} \mathbf{X}_0\|_2^2 \leq \|\mathbf{F} \mathbf{A}\|_2^2 \mathbb{E}\|\mathbf{X}_0\|_2^2 \leq N$, we can conclude from (65) that:

$$\mathbb{P}\left(\mathbf{X}_J^\top \mathbf{H}_{J,J} \mathbf{X}_J < \mathbb{E}\left(\mathbf{X}_J^\top \mathbf{H}_{J,J} \mathbf{X}_J \mid \mathbf{X}_I\right) - t \mid \mathbf{X}_I\right) \leq 2 \exp\left(-c' \cdot \min\left\{\frac{t^2}{2N}, t\right\}\right). \tag{66}$$

Next, let us define $\mathbf{y} := \mathbf{H}_{I,J}^\top \mathbf{X}_I$. Then,

$$\mathbf{X}_I^\top \mathbf{H}_{I,J} \mathbf{X}_J - \mathbb{E}\left(\mathbf{X}_I^\top \mathbf{H}_{I,J} \mathbf{X}_J \mid \mathbf{X}_I\right) = \mathbf{y}^\top \mathbf{X}_J - \mathbb{E}(\mathbf{y}^\top \mathbf{X}_J \mid \mathbf{X}_I).$$

By Lemma 4.4 in [20], Dobrushin's interdependence matrix for the model $\mathbf{X}_J \mid \mathbf{X}_I$ is given by $8\mathbf{D}$, where \mathbf{D} denotes the interaction matrix for the Ising model $\mathbf{X}_J \mid \mathbf{X}_I$. Since $\|8\mathbf{D}\|_2 \leq \frac{1}{2}$, by Theorem 4.3 in [20],

$$\mathbb{P}\left(\mathbf{y}^\top \mathbf{X}_J < \mathbb{E}(\mathbf{y}^\top \mathbf{X}_J \mid \mathbf{X}_I) - t \mid \mathbf{X}_I\right) \leq 2 \exp\left(-\frac{t^2}{8\|\mathbf{y}\|_2^2}\right). \tag{67}$$

Using the submultiplicativity of the matrix L^2 norm and the fact that the spectral norm of a matrix is always greater than or equal to the spectral norm of any of its submatrices gives,

$$\|\mathbf{y}\|_2^2 = \|\mathbf{H}_{I,J}^\top \mathbf{X}_I\|_2^2 \leq \|\mathbf{H}_{I,J}^\top\|_2^2 \cdot \|\mathbf{X}_I\|_2^2 \leq N \|\mathbf{H}\|_2^2 \leq N \|\mathbf{W}\|_2^4 = N \|\mathbf{F} \mathbf{A}\|_2^4 \leq N. \tag{68}$$

Combining (67) and (68),

$$\mathbb{P}\left(\mathbf{y}^\top \mathbf{X}_J < \mathbb{E}(\mathbf{y}^\top \mathbf{X}_J | \mathbf{X}_I) - t \mid \mathbf{X}_I\right) \leq 2 \exp\left(-\frac{t^2}{8N}\right). \quad (69)$$

Finally, combining (66), (69), and (57) gives,

$$\begin{aligned} & \mathbb{P}\left(\mathbf{X}^\top \mathbf{H} \mathbf{X} < \mathbb{E}\left(\mathbf{X}^\top \mathbf{H} \mathbf{X} | \mathbf{X}_I\right) - t \mid \mathbf{X}_I\right) \\ & \leq 2 \exp\left(-c' \cdot \min\left\{\frac{t^2}{8N}, \frac{t}{2}\right\}\right) + 2 \exp\left(-\frac{t^2}{128N}\right). \end{aligned}$$

This completes the proof of Lemma 9. \square

To complete the proof of (49), we choose J as in Lemma 8 and apply Lemma 9 with $t = \frac{1}{2}\mathbb{E}(\|\mathbf{F}\mathbf{m}\|_2^2 | \mathbf{X}_{J^c})$. This implies, there exists a constant C , depending only on Θ, M and s , such that

$$\mathbb{P}\left(\frac{1}{N}\|\mathbf{F}\mathbf{m}\|_2^2 \geq C \mid \mathbf{X}_{J^c}\right) = 1 - e^{-\Omega(N)}.$$

This proves (49) and completes the proof of Lemma 6. \square

4. PROOF OF THEOREM 2

Recall the definition of the log-pseudo-likelihood function $L_{\beta,N}(\cdot)$ from (9). As in Theorem 1, the proof of Theorem 2 entails showing the following: (1) concentration of the gradient of $L_{\beta,N}$ and (2) strong concavity of $L_{\beta,N}$. The concentration of the gradient $\nabla L_{\beta,N}$ follows by arguments identical to Lemma 1. Towards this, recall the definition of the functions $\phi_{i,s} : \mathcal{C}_N \rightarrow \mathbb{R}$, for $1 \leq i \leq N$ and $1 \leq s \leq d$, from (25). Note that $\nabla L_{\beta,N} = (\frac{\partial L_{\beta,N}}{\partial \theta_1}, \dots, \frac{\partial L_{\beta,N}}{\partial \theta_d})^\top$, where $\frac{\partial L_{\beta,N}}{\partial \theta_s} = \sum_{i=1}^N \phi_{i,s}(\mathbf{X})$ for $1 \leq s \leq d$. Then, by analogous arguments as in Lemma 1, it follows that there exists $\delta > 0$ such that with $\lambda := \delta \sqrt{\log d/N}$ the following holds:

$$\mathbb{P}\left(\|\nabla L_{\beta,N}(\boldsymbol{\theta})\|_\infty > \frac{\lambda}{2}\right) = o(1), \quad (70)$$

where the $o(1)$ -term goes to infinity as $d \rightarrow \infty$.

Next, recall the definition of $\mathcal{C}(S)$ from Assumption 4. The first step towards showing the strong concavity of $L_{\beta,N}$ is to establish $\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}$ belongs to the set $\mathcal{C}(S)$, whenever $\|\nabla L_{\beta,N}(\boldsymbol{\theta})\|_\infty \leq \frac{\lambda}{2}$.

Lemma 10. *The vector $\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}$ belongs to the set $\mathcal{C}(S)$, whenever $\|\nabla L_{\beta,N}(\boldsymbol{\theta})\|_\infty \leq \frac{\lambda}{2}$.*

Proof. By the convexity of the function $L_{\beta,N}$ it follows that

$$\begin{aligned} \lambda(\|\boldsymbol{\theta}\|_1 - \|\hat{\boldsymbol{\theta}}\|_1) & \geq L_{\beta,N}(\hat{\boldsymbol{\theta}}) - L_{\beta,N}(\boldsymbol{\theta}) \\ & \geq \nabla L_{\beta,N}(\boldsymbol{\theta})^\top (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}) \\ & \geq -\|\nabla L_{\beta,N}(\boldsymbol{\theta})\|_\infty \|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}\|_1 \\ & \geq -\frac{\lambda \|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}\|_1}{2}, \end{aligned} \quad (71)$$

where the last step uses $\|\nabla L_{\beta,N}(\boldsymbol{\theta})\|_\infty \leq \frac{\lambda}{2}$. Now, as in (22), note that $\|\hat{\boldsymbol{\theta}}\|_1 - \|\boldsymbol{\theta}\|_1 \geq \|(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})_{S^c}\|_1 - \|(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})_S\|_1$. Therefore, from (71), we have:

$$\|(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})_{S^c}\|_1 - \|(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})_S\|_1 \leq \|\hat{\boldsymbol{\theta}}\|_1 - \|\boldsymbol{\theta}\|_1 \leq \frac{\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}\|_1}{2} = \frac{\|(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})_S\|_1}{2} + \frac{\|(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})_{S^c}\|_1}{2}.$$

This implies that $\|(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})_{S^c}\|_1 \leq 3\|(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})_S\|_1$. \square

To show the strong convexity of $L_{\beta,N}$, by a second order Taylor series expansion of $L_{\beta,N}$, there exists $\alpha \in (0, 1)$ and $\underline{\boldsymbol{\theta}} = \alpha\boldsymbol{\theta} + (1 - \alpha)\hat{\boldsymbol{\theta}}$ such that

$$\begin{aligned} L_{\beta,N}(\hat{\boldsymbol{\theta}}) - L_{\beta,N}(\boldsymbol{\theta}) - \nabla L_{\beta,N}(\boldsymbol{\theta})^\top (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}) &= \frac{1}{2}(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})^\top \nabla^2 L_{\beta,N}(\underline{\boldsymbol{\theta}})(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}) \\ &= \frac{1}{2N} \sum_{i=1}^N \frac{(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})^\top \mathbf{Z}_i \mathbf{Z}_i^\top (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})}{\cosh^2(\beta m_i(\mathbf{X}) + \underline{\boldsymbol{\theta}}^\top \mathbf{Z}_i)}. \end{aligned} \quad (72)$$

Then, by (39) and (40), the RHS of (72) can be bounded by

$$\begin{aligned} L_{\beta,N}(\hat{\boldsymbol{\theta}}) - L_{\beta,N}(\boldsymbol{\theta}) - \nabla L_{\beta,N}(\boldsymbol{\theta})^\top (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}) &= \frac{1}{2}(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})^\top \nabla^2 L_{\beta,N}(\underline{\boldsymbol{\theta}})(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}) \\ &\gtrsim_{\beta,M,\kappa} \frac{1}{N} \cdot \frac{(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})^\top \mathbf{Z}^\top \mathbf{Z} (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})}{\cosh^2(|\beta| + M\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}\|_1 + sM\Theta)}. \end{aligned} \quad (73)$$

Now, if $\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}$ belongs to the set $\mathcal{C}(S)$ and Assumption 4 holds, then as in the proof of Lemma 7, it can be shown that, for $\lambda = O(\sqrt{\log d/N})$,

$$\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}\|_1 = O_s \left(\sqrt{\frac{\log d}{N}} \right).$$

with probability $1 - o(1)$, whenever $N, d \rightarrow \infty$ such that $\log d = o(N)$. This combined with (73) implies, whenever $\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}$ belongs to the set $\mathcal{C}(S)$,

$$L_{\beta,N}(\hat{\boldsymbol{\theta}}) - L_{\beta,N}(\boldsymbol{\theta}) - \nabla L_{\beta,N}(\boldsymbol{\theta})^\top (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}) \gtrsim_{\beta,M,\kappa} \left\| \hat{\boldsymbol{\theta}} - \boldsymbol{\theta} \right\|_2^2, \quad (74)$$

with probability $1 - o(1)$, as $N, d \rightarrow \infty$ such that $d = o(\log N)$. Since $\|\nabla L_{\beta,N}(\boldsymbol{\theta})\|_\infty \leq \frac{\lambda}{2}$ holds with probability $1 - o(1)$ by (70), applying Lemma 10 shows that $\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}$ belongs to the set $\mathcal{C}(S)$ with probability $1 - o(1)$, as well. This implies that (74) holds with probability $1 - o(1)$, whenever $N, d \rightarrow \infty$ such that $d = o(\log N)$. This establishes the required strong concavity of $L_{\beta,N}$.

The proof of Theorem 2 can now be completed as in Theorem 1, using the concentration of the gradient in (70) and the strong concavity of the pseudo-likelihood in (74). \square

5. PROOFS FROM SECTION 2.3

In this section, we prove the results stated in Section 2.3. We begin with the proof of Theorem 3 in Section 5.1. Corollary 1 is proved in Section 5.2.

5.1. Proof of Theorem 3. The proof of Theorem 5.1 follows along the same lines as in Theorem 1, so to avoid repetition we sketch the steps and highlight the relevant modifications. As in the proof of Theorem 5.1, the first step is to show the concentration of $\nabla L_N(\boldsymbol{\gamma})$. Towards this, following the proof of Lemma 5 shows that Q_r (as defined in (30)) is $O_\beta(\text{poly}(d_{\max})/N)$ -Lipschitz, for each $r \in [\ell]$. Therefore, by arguments as in Lemma 5,

$$\mathbb{P} \left(\left| \sum_{i=1}^N \phi_i(\mathbf{X}) \right| \geq t \right) \leq \mathbb{P} \left(\max_{r \in [\ell]} |Q_r(\mathbf{X})| \geq \frac{t\ell'}{\ell} \right) \lesssim e^{-O_{\beta,M}(Nt^2/\text{poly}(d_{\max}))}.$$

since $\ell = O(\log N)$ and $\ell'/\ell = \Theta(1)$. In this case, following the notations in the proof of Lemma 1, we have $\ell/\ell' = \Theta(d_{\max})$. Similarly, for $s \in [d]$,

$$\mathbb{P} \left(\left| \sum_{i=1}^N \phi_{i,s}(\mathbf{X}) \right| \geq t \right) \lesssim e^{-O_{\beta,M}(Nt^2/\text{poly}(d_{\max}))}.$$

Hence, choosing $\lambda := \delta \text{poly}(d_{\max}) \sqrt{\log(d+1)/N}$ and $t := \lambda/2$ gives, for some constant C depending only on β and M ,

$$\mathbb{P} \left(\left| \sum_{i=1}^N \phi_i(\mathbf{X}) \right| \geq \frac{\lambda}{2} \right) \leq (d+1)^{-C\delta^2/4} \quad \text{and} \quad \mathbb{P} \left(\left| \sum_{i=1}^N \phi_{i,j}(\mathbf{X}) \right| \geq \frac{\lambda}{2} \right) \leq (d+1)^{-C\delta^2/4},$$

for all $j \in [s]$. A final union bound over the $(d+1)$ coordinates now shows that

$$\mathbb{P} \left(\|\nabla L_N(\boldsymbol{\gamma})\|_{\infty} > \frac{\lambda}{2} \right) = o(1), \tag{75}$$

where $\lambda := \delta \text{poly}(d_{\max}) \sqrt{\log(d+1)/N}$ as above and the $o(1)$ -term goes to infinity as $d \rightarrow \infty$. This establishes the concentration of the gradient.

Next, we need to show the strong concavity of the Hessian. To this end, first note that since $|E(G_N)| = O(N)$ and the number of non-isolated vertices of G_N is $\Omega(N)$, there exist constants $L_1, L_2 > 0$, such that $|E(G_N)| \leq L_1 N$, and the number of non-isolated vertices of G_N is larger than $L_2 N$, for all N large enough. For $D \geq 1$ define,

$$V_N(D) := \{v \in V(G_N) : d_v \in [1, D]\}.$$

Note that

$$|V_N(D)| \geq N(1 - (2L_1/D)) - N(1 - L_2) = N(L_2 - (2L_1/D)),$$

since G_N has at least $N(1 - (2L_1/D))$ vertices with degree not exceeding D , among which at most $N(1 - L_2)$ are isolated. Hereafter, we choose $D := \lceil 4L_1/L_2 \rceil$, so that $|V_N(D)| \geq L_2 N/2$. Then, with notations as in (38) we have,

$$\begin{aligned} & L_N(\hat{\boldsymbol{\gamma}}) - L_N(\boldsymbol{\gamma}) - \nabla L_N(\boldsymbol{\gamma})^\top (\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}) \\ &= \frac{1}{2} (\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})^\top \nabla^2 L_N(\boldsymbol{\gamma}) (\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}) \\ &= \frac{1}{2N} \sum_{i=1}^N \frac{(\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})^\top \mathbf{U}_i \mathbf{U}_i^\top (\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})}{\cosh^2(\underline{\beta} m_i(\mathbf{X}) + \boldsymbol{\theta}^\top \mathbf{Z}_i)} \\ &\geq \frac{1}{2N} \sum_{i \in V_N(D)} \frac{(\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})^\top \mathbf{U}_i \mathbf{U}_i^\top (\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})}{\cosh^2(\underline{\beta} m_i(\mathbf{X}) + \boldsymbol{\theta}^\top \mathbf{Z}_i)} \\ &\geq \frac{1}{2N} \sum_{i \in V_N(D)} \frac{(\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})^\top \mathbf{U}_i \mathbf{U}_i^\top (\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})}{\cosh^2(|\beta|D + (D+M)\|\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}\|_1 + sM\Theta)}, \end{aligned} \tag{76}$$

where the last step uses the bound in (40) and the bound $|\underline{\beta} m_i(\mathbf{X})| \leq |\underline{\beta}| |m_i(\mathbf{X})| \leq |\underline{\beta}| D \leq |\beta| D + D \|\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}\|_1$. Next, using the bound $|V_N(D)| \geq L_2 N/2$ in (76) gives,

$$\begin{aligned} & L_N(\hat{\boldsymbol{\gamma}}) - L_N(\boldsymbol{\gamma}) - \nabla L_N(\boldsymbol{\gamma})^\top (\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}) \\ &\geq \frac{L_2}{4 \cosh^2(|\beta|D + (D+M)\|\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}\|_1 + sM\Theta)} \cdot \frac{1}{|V_N(D)|} \sum_{i \in V_N(D)} (\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma})^\top \mathbf{U}_i \mathbf{U}_i^\top (\hat{\boldsymbol{\gamma}} - \boldsymbol{\gamma}) \end{aligned}$$

$$= \frac{L_2}{4 \cosh^2(|\beta|D + (D + M)\|\hat{\gamma} - \gamma\|_1 + sM\Theta)} (\hat{\gamma} - \gamma)^\top \tilde{\mathbf{G}} (\hat{\gamma} - \gamma), \quad (77)$$

where

$$\tilde{\mathbf{G}} := \frac{1}{|V_N(D)|} \begin{pmatrix} \tilde{\mathbf{m}}^\top \tilde{\mathbf{m}} & \tilde{\mathbf{m}}^\top \tilde{\mathbf{Z}} \\ \tilde{\mathbf{Z}}^\top \tilde{\mathbf{m}} & \tilde{\mathbf{Z}}^\top \tilde{\mathbf{Z}} \end{pmatrix},$$

with $\tilde{\mathbf{m}} := (m_i(\mathbf{X}))_{i \in V_N(D)}^\top$ and $\tilde{\mathbf{Z}} = (\mathbf{Z}_i)_{i \in V_N(D)}^\top$. The calculation in (77) implies that in order to establish the strong concavity of the Hessian in the setting of Theorem 3, we need to show $\lambda_{\min}(\tilde{\mathbf{G}}) \gtrsim 1$ with probability going to 1, where $\lambda_{\min}(\tilde{\mathbf{G}})$ denotes the minimum eigenvalue of $\tilde{\mathbf{G}}$. For this step, we follow the steps of Lemma 6 with the matrix \mathbf{F} replaced by $\tilde{\mathbf{F}} := \mathbf{I} - \tilde{\mathbf{Z}}(\tilde{\mathbf{Z}}^\top \tilde{\mathbf{Z}})^{-1} \tilde{\mathbf{Z}}^\top$. Then, repeating the proof of (55) in Lemma 8, with \mathbf{A} replaced by $\tilde{\mathbf{A}} := \mathbf{A}|_{V_N(D) \times [N]}$, we can find a set $J \subseteq [N]$ such that

$$\mathbb{E} \left(\|\tilde{\mathbf{F}} \tilde{\mathbf{m}}\|_2^2 \middle| \mathbf{X}_{J^c} \right) \gtrsim \Upsilon^2 \left(\|\tilde{\mathbf{A}}\|_F^2 - d \cdot \text{polylog}(N) \right),$$

since $\|\tilde{\mathbf{A}}\|_2 \leq \|\mathbf{A}\|_\infty = O(\text{polylog}(N))$. This implies,

$$\mathbb{E} \left(\|\tilde{\mathbf{F}} \tilde{\mathbf{m}}\|_2^2 \middle| \mathbf{X}_{J^c} \right) \gtrsim N, \quad (78)$$

since $\|\tilde{\mathbf{A}}\|_F^2 \geq |V_N(D)| \gtrsim N$ (because every vertex in $V_N(D)$ has degree at least 1) and $d = o(N)$. The final step is to establish that $\|\tilde{\mathbf{F}} \tilde{\mathbf{m}}\|$ concentrates around $\|\tilde{\mathbf{F}} \tilde{\mathbf{m}}\|_2^2 \middle| \mathbf{X}_{J^c}$, conditional on \mathbf{X}_{J^c} . This follows by repeating the proof of Lemma 9, which introduces an extra $1/\text{polylog}(N)$ factor in each of the two exponential terms in the RHS of (56), since $\|\tilde{\mathbf{A}}\|_2 \leq \|\mathbf{A}\|_\infty = O(\text{polylog}(N))$. This, combined with (78), shows

$$\mathbb{P}(\lambda_{\min}(\tilde{\mathbf{G}}) \geq C) \geq 1 - e^{-\Omega(N/\text{polylog}(N))}, \quad (79)$$

for some constant $C > 0$. The proof of Theorem 3 can be now completed using (75) and (79), as in Theorem 1.

5.2. Proof of Corollary 1. To prove Corollary 1 we verify that the hypotheses of Theorem 3 are satisfied. Note that we can write

$$|E(G_N)| = \sum_{1 \leq u < v \leq N} B_{uv},$$

where $B_{uv} \sim \text{Ber}(p_{uv})$, and $\{B_{uv}\}_{1 \leq u < v \leq N}$ are independent. This implies, $\mathbb{E}|E(G_N)| = O(N)$ and $\text{Var}(|E(G_N)|) = O(N)$, since $\sup_{1 \leq i, j \leq N} p_{ij} = O(1/N)$ by (13). Hence, by Chebyshev's inequality,

$$|E(G_N)| \leq \mathbb{E}|E(G_N)| + N = O(N),$$

with probability $1 - o(1)$.

Next, we will show that $d_{\max} = \tilde{O}(1)$ holds with high probability. To this end, define $\eta_u = \sum_{v \neq u} p_{uv}$, for $1 \leq u \leq N$. Clearly, by assumption (13), $\max_{1 \leq u \leq N} \eta_u = O(1)$. Next, we establish that $\max_{1 \leq u \leq N} \eta_u = \Omega(1)$. Towards this, note that by assumption (14) there exists $\varepsilon \in (0, 1)$ and $u \in V(G_N)$ such that,

$$\limsup_{N \rightarrow \infty} \sum_{v=1}^N \log(1 - p_{uv}) < \log \varepsilon.$$

Next, using the inequality $\log(1-x) \geq -x/(1-x)$, for all $x < 1$, and taking N large enough such that $\sup_{u,v} p_{uv} < 1/2$, gives

$$\limsup_{N \rightarrow \infty} \left(-2 \sum_{v=1}^N p_{uv} \right) \leq \log \varepsilon \quad \implies \quad \liminf_{N \rightarrow \infty} \eta_u \geq \frac{1}{2} \log \left(\frac{1}{\varepsilon} \right).$$

This shows that $\max_{1 \leq u \leq N} \eta_u = \Omega(1)$. Therefore, by Proposition 1.11 in [10], $d_{\max} \leq O(\log N)$ with probability $1 - o(1)$.

Finally, we show that the number of non-isolated vertices of G_N is $\Omega(N)$ with high probability. To this end, for each $v \in V(G_N)$, define $Y_v := \mathbf{1}\{d_v = 0\}$. Then, $I_N := \sum_{v=1}^N Y_v$ is the total number of isolated vertices of G_N . Note that $Y_v \sim \text{Ber}(\prod_{u=1}^N (1 - p_{uv}))$. Therefore, by (14),

$$\mathbb{E}(I_N) = \sum_{u=1}^N \prod_{v=1}^N (1 - p_{uv}) \leq \alpha N,$$

for some $\alpha \in (0, 1)$ and all large N enough. Next, note that for any two distinct vertices $u, v \in V(G_N)$,

$$\text{Cov}(Y_u, Y_v) = p_{uv} (1 - p_{uv}) \prod_{w \notin \{u, v\}} [(1 - p_{uw})(1 - p_{vw})] \leq p_{uv} = O\left(\frac{1}{N}\right).$$

Similarly, it can be checked that $\text{Var}(Y_v) = O(1)$, for $1 \leq v \leq N$. This implies, $\text{Var}(I_N) = O(N)$. Hence, by Chebyshev's inequality,

$$\mathbb{P}\left(I_N \geq \left(\frac{1+\alpha}{2}\right) N\right) \leq \mathbb{P}\left(I_N \geq \mathbb{E}(I_N) + \left(\frac{1-\alpha}{2}\right) N\right) \leq \frac{\text{Var}(I_N)}{\left(\frac{1-\alpha}{2}\right)^2 N^2} = O\left(\frac{1}{N}\right).$$

This shows that the number of non-isolated vertices of G_N is at least $(1-\alpha)N/2$ with probability $1 - o(1)$. This completes the verification of the hypotheses of Theorem 3 and hence, Corollary 1 follows from Theorem 3.

Acknowledgment. B. B. Bhattacharya thanks Rajarshi Mukherjee and Nancy R. Zhang for many helpful discussions.

REFERENCES

- [1] R. Adamczak, M. Kotowski, B. Polaczyk and M. Strzelecki, A note on concentration for polynomials in the Ising model, *Electronic Journal of Probability*, Vol. 24 (42), 1-22, 2019.
- [2] C. Alvin, K. Mondal, S. Schmid, Demand-aware network designs of bounded degree, [arXiv:1705.06024](https://arxiv.org/abs/1705.06024), 2017.
- [3] A. Anandkumar, V. Y. F. Tan, F. Huang, and A. S. Willsky, High-dimensional structure estimation in Ising models: Local separation criterion, *Annals of Statistics*, Vol. 40 (3), 1346–1375 Vol. 2012.
- [4] S. Banerjee, B. P. Carlin, and A. E. Gelfand, Hierarchical modeling and analysis for spatial data, *Chapman and Hall/CRC*, 2014.
- [5] D. Batra, A. Gallagher, D. Parikh, and T. Chen, Beyond trees: MRF inference via outer-planar decomposition, *IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, 2496–2503, 2010.
- [6] M. Bertrand, E. F. P. Luttmer, and S. Mullainathan, Network effects and welfare cultures, *The Quarterly Journal of Economics*, Vol. 115 (3), 1019–1055, 2000.
- [7] B. B. Bhattacharya and S. Mukherjee, Inference in Ising models, *Bernoulli*, Vol. 24 (1), 493–525, 2018.
- [8] P. J. Bickel, Y. Ritov, and A. Tsybakov, Simultaneous analysis of Lasso and Dantzig selector, *Annals of Statistics*, Vol. 37 (4), 1705–1732, 2009.
- [9] B. Bollobas, S. Janson, and O. Riordan, The phase transition in inhomogeneous random graphs, *Random Structures and Algorithms*, Vol. 31 (1), 3–122, 2007.
- [10] F. Benaych-Georges, C. Bordenave and A. Knowles, Largest eigenvalues of sparse inhomogeneous Erdős-Rényi graphs, [arXiv:1704.02953](https://arxiv.org/abs/1704.02953), 2017.

- [11] J. Besag, Spatial interaction and the statistical analysis of lattice systems, *J. Roy. Stat. Soc. B*, Vol. 36, 192–236, 1974.
- [12] J. Besag, Statistical analysis of non-lattice data, *The Statistician*, Vol. 24 (3), 179–195, 1975.
- [13] G. Bresler, Efficiently learning Ising models on arbitrary graphs, *Proceedings Symposium on Theory of Computing (STOC)*, 771–782, 2015.
- [14] G. Bresler and M. Karzand, Learning a tree-structured Ising model in order to make predictions, *Annals of Statistics*, Vol. 48 (2), 713–737, 2020.
- [15] G. Bresler and D. Nagaraj, Optimal single sample tests for structured versus unstructured network data, *Conference On Learning Theory (COLT)*, 1657–1690, 2018.
- [16] F. Bunea, Honest variable selection in linear and logistic regression models via ℓ_1 and $\ell_1 + \ell_2$ penalization, *Electronic Journal of Statistics*, Vol. 2, 1153–1194, 2008.
- [17] E. J. Candés and P. Sur, The phase transition for the existence of the maximum likelihood estimate in high-dimensional logistic regression, *Annals of Statistics*, 48 (1), 27–42, 2020.
- [18] E. Candés and T. Tao, The Dantzig selector: Statistical estimation when p is much larger than n , *Annals of Statistics*, Vol. 35 (6), 2313–2351, 2007.
- [19] Yuan Cao, Matey Neykov and Han Liu, High temperature structure detection in ferromagnets, *Information and Inference: A Journal of the IMA*, to appear, 2021.
- [20] S. Chatterjee, Concentration inequalities with exchangeable pairs, [arXiv:math/0507526](https://arxiv.org/abs/math/0507526), 2016.
- [21] S. Chatterjee, Estimation in spin glasses: A first step, *Annals of Statistics*, Vol. 35 (5), 1931–1946, 2007.
- [22] S. Chatterjee, P. Diaconis, and A. Sly, Random graphs with a given degree sequence, *Annals of Applied Probability*, Vol. 21 (4), 1400–1435, 2011.
- [23] C. Chow and C. Liu, *Approximating discrete probability distributions with dependence trees*, *IEEE Transactions on Information Theory*, Vol. 14 (3), 462–467, 1968.
- [24] N. A. Christakis and J. H. Fowler, Social contagion theory: examining dynamic social networks and human behavior, *Statistics in medicine*, Vol. 32 (4), 556–577, 2013.
- [25] F. Chung and L. Lu, Connected components in random graphs with given expected degree sequences, *Annals of Combinatorics*, Vol. 6, 125–145, 2002.
- [26] F. Comets, On consistency of a class of estimators for exponential families of Markov random fields on the lattice, *Annals of Statistics*, Vol. 20 (1), 455–468, 1992.
- [27] F. Comets and B. Gidas, Asymptotics of maximum likelihood estimators for the Curie-Weiss model, *Annals of Statistics*, 19(2):557–578, 1991.
- [28] Y. Dagan, C. Daskalakis, N. Dikkala, and S. Jayanti, Learning from weakly dependent data under Dobrushin’s condition, *Conference on Learning Theory (COLT)*, 914–928, PMLR, 2019.
- [29] Y. Dagan, C. Daskalakis, N. Dikkala, and A. V. Kandiros, Learning Ising models from one or multiple samples, *Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing (STOC)*, 161–168, 2021.
- [30] C. Daskalakis, N. Dikkala, and G. Kamath, Testing Ising models, *IEEE Transactions on Information Theory*, Vol. 65 (11), 6829–6852, 2019.
- [31] C. Daskalakis, N. Dikkala, and I. Panageas, Logistic regression with peer-group effects via inference in higher-order Ising models, *Proceedings of the 23rd International Conference on Artificial Intelligence and Statistics (AISTATS)*, 3653–3663, 2020.
- [32] C. Daskalakis, N. Dikkala, and I. Panageas, Regression from dependent observations, *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing (STOC)*, 881–889, 2019.
- [33] E. Duflo and E. Saez, The role of information and social interactions in retirement plan decisions: Evidence from a randomized experiment, *The Quarterly Journal of Economics*, Vol.118(3), 815–842, 2003.
- [34] C.-H. L. Eng, M. Lawson, Q. Zhu, R. Dries, N. Koulena, Y. Takei, J. Yun et al., Transcriptome-scale super-resolved imaging in tissues by RNA seqFISH+, *Nature*, Vol. 568 (7751), 235–239, 2019.
- [35] S. Geman and C. Graffigne, Markov random field image models and their applications to computer vision, *Proceedings of the International Congress of Mathematicians*, 1496–1517, 1986.
- [36] P. Ghosal and S. Mukherjee, Joint estimation of parameters in Ising model, *Annals of Statistics*, Vol. 48 (2), 785–810, 2020.
- [37] B. Gidas, Consistency of maximum likelihood and pseudolikelihood estimators for Gibbs distributions, *In Stochastic Differential Systems, Stochastic Control Theory and Applications (W. Fleming and P.-L. Lions, eds.)*, 129–145, Springer, New York, 1988.
- [38] E. L. Glaeser, B. Sacerdote, and J. A. Scheinkman, Crime and social interactions, *The Quarterly Journal of Economics*, 111 (2):507–548, 1996.

- [39] Y. Goltsev, N. Samusik, J. Kennedy-Darling, S. Bhate, M. Hale, G. Vazquez, S. Black, and G. P. Nolan, Deep profiling of mouse splenic architecture with CODEX multiplexed imaging, *Cell*, Vol. 174 (4), 968–981, 2018.
- [40] P. J. Green and S. Richardson, Hidden markov models and disease mapping, *Journal of the American Statistical Association*, Vol. 97, 1055–1070, 2002.
- [41] X. Guyon and H. R. Künsch, Asymptotic comparison of estimators in the Ising Model, *Stochastic Models, Statistical Methods, and Algorithms in Image Analysis*, Lecture Notes in Statistics, Vol. 74, 177–198, 1992.
- [42] L. Hamilton, F. Koehler, and A. Moitra, Information theoretic properties of Markov Random Fields, and their algorithmic applications, *Advances in Neural Information Processing Systems (NIPS)*, 2463–2472, 2017.
- [43] K. M. Harris, National Longitudinal Study of Adolescent Health, et al. Waves i & ii, 1994-1996; wave iii, 2001-2002; wave iv, 2007-2009.
- [44] J. J. Hopfield, Neural networks and physical systems with emergent collective computational abilities, *Proceedings of the National Academy of Sciences USA*, 79:2554–2558, 1982.
- [45] D. W. Hosmer Jr, S. Lemeshow, and R. X Sturdivant, Applied logistic regression, *John Wiley & Sons*, Vol. 398, 2013.
- [46] E. Ising, Beitrag zur theorie der ferromagnetismus, *Zeitschrift für Physik*, Vol. 31, 253–258, 1925.
- [47] Jason K. Johnson, Diane Oyen, Michael Chertkov, and Praneeth Netrapalli, Learning planar Ising models, *Journal of Machine Learning Research* Vol. 17, 7539–7564, 2016.
- [48] S. Kakade, O. Shamir, K. Sindharan, and A. Tewari, Learning exponential families in high-dimensions: Strong convexity and sparsity, *Proceedings of the thirteenth international conference on artificial intelligence and statistics*, 381-388, 2010.
- [49] V. Kandiros, Y. Dagan, N. Dikkala, S. Goel, and C. Daskalakis, Statistical Estimation from Dependent Data, *International Conference on Machine Learning*, 5269–5278, PMLR, 2021.
- [50] A. R. Klivans and R. Meka, Learning Graphical Models Using Multiplicative Weights, *Foundations of Computer Science*, 343–354, 2017.
- [51] M. Krivelevich and B. Sudakov, The largest eigenvalue of sparse random graphs, *Combinatorics, Probability and Computing*, Vol. 12 (1), 61–72, 2003
- [52] E. L. Lehmann and J. P. Romano, *Testing statistical hypotheses*, Springer Science & Business Media, 2006.
- [53] J. Lei, A goodness-of-fit test for stochastic block models, *Annals of Statistics*, Vol. 44 (1), 401–424, 2016.
- [54] P. Loh, Statistical consistency and asymptotic normality for high-dimensional robust M -estimators, *Annals of Statistics*, Vol. 45 (2), 866-896, 2017.
- [55] P. McCullagh and J. A Nelder, *Generalized linear models*, Monograph on Statistics and Applied Probability, 1989
- [56] N. Meinshausen and B. Yu, Lasso-type recovery of sparse representations for high-dimensional data, *Annals of Statistics*, Vol. 37 (1), 246–270, 2009.
- [57] A. Montanari and Amin Saberi, The spread of innovations in social networks, *Proceedings of the National Academy of Sciences*, Vol. 107 (47), 20196–20201, 2010.
- [58] R. Mukherjee, S. Mukherjee and M. Yuan, Global testing against sparse alternatives under Ising models, *Annals of Statistics*, Vol. 46 (5), 2062–2093, 2018.
- [59] R. Mukherjee and G. Ray, On testing for parameters in Ising model, *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, to appear, 2021.
- [60] S. Mukherjee, J. Son, and B. B. Bhattacharya, Estimation in tensor Ising models, [arXiv:2008.12882](https://arxiv.org/abs/2008.12882), 2020.
- [61] S. N. Negahban, P. Ravikumar, M. J. Wainwright, and B. Yu, A unified framework for high-dimensional analysis of M -estimators with decomposable regularizers, *Statistical Science*, Vol. 27 (4), 538-557, 2012.
- [62] J. A. Nelder and R. W. M. Wedderburn, Generalized linear models, *Journal of the Royal Statistical Society, Series A (General)*, Vol. 135(3), 370–384, 1972.
- [63] M. Neykov and H. Liu, Property testing in high-dimensional Ising models, *Annals of Statistics*, Vol. 47 (5), 2472–2503, 2019.
- [64] J. M. Perkel, Starfish enterprise: finding RNA patterns in single cells, *Nature* 572, 549–551, 2019.
- [65] D. K. Pickard, Inference for discrete Markov Fields: the simplest nontrivial case, *Journal of the American Statistical Association*, Vol. 82 (397), 90–96, 1987.
- [66] G. Raskutti, M. J. Wainwright, and B. Yu, Minimax rates of estimation for high-dimensional linear regression over ℓ_q -balls, *IEEE Trans. Information Theory*, Vol. 57 (10), 6976–6994, 2011.
- [67] P. Ravikumar, M. J. Wainwright and J. Lafferty, High-dimensional Ising model selection using ℓ_1 -regularized logistic regression, *Annals of Statistics*, Vol. 38 (3), 1287–1319, 2010.

- [68] M. Rudelson and S. Zhou, Reconstruction from anisotropic random measurements, *Conference on Learning Theory (COLT)*, JMLR Workshop and Conference Proceedings, 2012.
- [69] B. Sacerdote, Peer effects with random assignment: Results for Dartmouth roommates, *The Quarterly Journal of Economics*, 116(2), 681–704, 2001.
- [70] F. Salehi, Ehsan Abbasi, and Babak Hassibi, The Impact of Regularization on High-dimensional Logistic Regression, [arXiv:1906.03761](https://arxiv.org/abs/1906.03761), 2019.
- [71] N. P. Santhanam and M. J. Wainwright, Information-theoretic limits of selecting binary graphical models in high dimensions, *IEEE Transactions on Information Theory*, Vol. 58 (7), 4117–4134, 2012.
- [72] P. Sur and E. J. Candés, A modern maximum-likelihood theory for high-dimensional logistic regression, *Proceedings of the National Academy of Sciences*, Vol. 116 (29), 14516–14525.
- [73] P. Sur, Y. Chen, and E. J. Candés, The likelihood ratio test in high-dimensional logistic regression is asymptotically a rescaled chi-square, *Probability Theory and Related Fields*, 1-72, 2017.
- [74] M. Tang, A. Athreya, D. L. Sussman, V. Lyzinski, and C. E. Priebe, A nonparametric two-sample hypothesis testing problem for random graphs, *Bernoulli*, Vol. 23, 1599–1630, 2017.
- [75] J. G. Trogon, J. Nonnemaker, and J. Pais. Peer effects in adolescent overweight, *Journal of Health Economics*, Vol. 27 (5), 1388–1399, 2008.
- [76] S. A. Van de Geer, High-dimensional generalized linear models and the lasso, *Annals of Statistics*, Vol. 36 (2), 614–645, 2008.
- [77] S. van de Geer and P. Bühlmann, On the conditions used to prove oracle results for the lasso, *Electronic Journal of Statistics*, Vol. 3, 1360–1392, 2009.
- [78] S.A. Van de Geer, Peter Bühlmann, Ya’acov Ritov, and Ruben Dezeure, On asymptotically optimal confidence regions and tests for high-dimensional models, *Annals of Statistics*, 42 (3), 1166–1202, 2014.
- [79] A. W. Van der Vaart, *Asymptotic Statistics*, Volume 3, Cambridge University Press, 2000.
- [80] M. Vuffray, S. Misra, A. Y. Likhov, and M. Chertkov, Interaction Screening: Efficient and Sample-Optimal Learning of Ising Models, *Neural Information Processing System (NeurIPS)*, 2595–2603, 2016.
- [81] B. Wang and B. Xi, Some inequalities for singular values of matrix products, *Linear algebra and its applications*, Vol. 264, 109–115, 1997.
- [82] L. Xue, H. Zou, and T. Cai, Non-concave penalized composite conditional likelihood estimation of sparse Ising models, *Annals of Statistics*, Vol. 40 (3), 1403–1429, 2012.

APPENDIX A. PROOFS OF TECHNICAL LEMMAS

In this section we collect the proofs of various technical lemmas. The section is organized as follows: In Appendix A.1 we prove Lemma 5. The proof of Lemma 3 is given in Appendix A.2. In Appendix A.3 we prove a variance lower bound for linear functions.

A.1. Proof of Lemma 5. To begin with, recall from (30) that

$$Q_r(\mathbf{X}) = -\frac{1}{N} \sum_{i \in I_r} m_i(\mathbf{X}) \left[X_i - \tanh(\beta m_i(\mathbf{X}) + \boldsymbol{\theta}^\top \mathbf{Z}_i) \right].$$

Hence, for any two $\mathbf{X}, \mathbf{X}' \in \{-1, 1\}^N$,

$$|Q_r(\mathbf{X}) - Q_r(\mathbf{X}')| = T_1 + T_2, \tag{80}$$

where

$$\begin{aligned} T_1 &= \frac{1}{N} \left| \sum_{i \in I_r} \{m_i(\mathbf{X})X_i - m_i(\mathbf{X}')X'_i\} \right| \\ T_2 &= \frac{1}{N} \left| \sum_{i \in I_r} \left\{ m_i(\mathbf{X}) \tanh(\beta m_i(\mathbf{X}) + \boldsymbol{\theta}^\top \mathbf{Z}_i) - m_i(\mathbf{X}') \tanh(\beta m_i(\mathbf{X}') + \boldsymbol{\theta}^\top \mathbf{Z}_i) \right\} \right|. \end{aligned} \tag{81}$$

Now, assume that \mathbf{X} and \mathbf{X}' differ only in the k -th coordinate, for some $k \in [N]$. Then

$$\begin{aligned}
T_1 &= \left| \sum_{i \in I_r} \sum_{j=1}^N a_{ij} (X_i X_j - X'_i X'_j) \right| \\
&\leq \left| \sum_{i \in I_r} a_{ik} (X_i X_k - X'_i X'_k) \right| + \left| \sum_{j=1}^N a_{kj} (X_k X_j - X'_k X'_j) \right| \\
&\leq 2 \sum_{i \in I_r} |a_{ik}| + 2 \sum_{j=1}^N |a_{kj}| \\
&\leq 4 \left| \sum_{i=1}^N a_{ik} \right| \leq 4 \|\mathbf{A}\|_1 \leq 4 \quad (\text{by (17)}).
\end{aligned} \tag{82}$$

Next, we proceed to bound T_2 . Towards this note that

$$T_2 \leq T_{21} + T_{22}, \tag{83}$$

where

$$\begin{aligned}
T_{21} &:= \left| \sum_{i \in I_r} m_i(\mathbf{X}) \left\{ \tanh(\beta m_i(\mathbf{X}) + \boldsymbol{\theta}^\top \mathbf{Z}_i) - \tanh(\beta m_i(\mathbf{X}') + \boldsymbol{\theta}^\top \mathbf{Z}_i) \right\} \right| \\
T_{22} &:= \left| \sum_{i \in I_r} a_{ik} (X_k - X'_k) \tanh(\beta m_i(\mathbf{X}') + \boldsymbol{\theta}^\top \mathbf{Z}_i) \right|,
\end{aligned}$$

since $m_i(\mathbf{X}) - m_i(\mathbf{X}') = a_{ik}(X_k - X'_k)$. Hence, using $|\tanh x - \tanh y| \leq |x - y|$ gives,

$$T_{21} \leq \beta \sum_{i \in I_r} |m_i(\mathbf{X})| |m_i(\mathbf{X}) - m_i(\mathbf{X}')| \leq 2\beta \sum_{i \in I_r} |m_i(\mathbf{X})| |a_{ik}| \leq 2\beta \|\mathbf{A}\|_\infty \|\mathbf{A}\|_1 \leq 1 \tag{84}$$

and using $\tanh x \leq 1$ gives,

$$T_{22} \leq 2 \sum_{i \in I_r} |a_{ir}| \leq 2 \|\mathbf{A}\|_1 \leq 2. \tag{85}$$

Combining (80), (81), (82), (83), (84), (85) the result in Lemma 5 (1) follows.

Next, we prove (2). To begin with, recall from (32) that

$$Q_{r,s}(\mathbf{X}) = -\frac{1}{N} \sum_{i \in I_r} \mathbf{Z}_{i,s} \left[X_i - \tanh(\beta m_i(\mathbf{X}) + \boldsymbol{\theta}^\top \mathbf{Z}_i) \right].$$

Hence, for any two $\mathbf{X}, \mathbf{X}' \in \{-1, 1\}^N$ differing in the k -th coordinate only,

$$\begin{aligned}
|Q_{r,s}(\mathbf{X}) - Q_{r,s}(\mathbf{X}')| &\leq \frac{1}{N} \left| \sum_{i \in I_r} \mathbf{Z}_{i,s} (X_i - X'_i) \right| \\
&\quad + \frac{1}{N} \left| \sum_{i \in I_r} \mathbf{Z}_{i,s} \left[\tanh(\beta m_i(\mathbf{X}) + \boldsymbol{\theta}^\top \mathbf{Z}_i) - \tanh(\beta m_i(\mathbf{X}') + \boldsymbol{\theta}^\top \mathbf{Z}_i) \right] \right| \\
&\leq \frac{2\|\mathbf{Z}\|_\infty}{N} + \frac{\beta}{N} \left| \sum_{i \in I_r} \mathbf{Z}_{i,s} (m_i(\mathbf{X}) - m_i(\mathbf{X}')) \right|
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{2M}{N} + \frac{\beta}{N} \sum_{i=1}^N |\mathbf{z}_{i,s} a_{ik} (X_k - X'_k)| \\
&\leq \frac{2M}{N} + \frac{2M\beta \|\mathbf{A}\|_1}{N} \leq \frac{2M(1+\beta)}{N},
\end{aligned}$$

as desired.

A.2. Proof of the Lemma 3. Suppose that \mathbf{X} comes from the model:

$$\mathbb{P}_{\beta, \mathbf{h}}(\mathbf{X}) \propto \exp\left(\sum_{i=1}^N h_i X_i + \mathbf{X}^\top \mathbf{D} \mathbf{X}\right), \quad (86)$$

which is assumed to be (R, Υ) -Ising. We will apply Lemma 17 in [29] on the matrix

$$\mathbf{D} := ((d_{ij}))_{1 \leq i, j \leq N} := \mathbf{A}/R,$$

where \mathbf{A} is the interaction matrix corresponding to the distribution of \mathbf{X} . Note that \mathbf{D} satisfies the hypotheses of Lemma 17 in [29]. For $\eta \in (0, R)$, define $\eta' := \eta/R$. This ensures that $\eta' \in (0, 1)$. By Lemma 17 in [29], there exist subsets $I_1, \dots, I_\ell \subseteq [N]$ with $\ell \lesssim R^2 \log N/\eta^2$, such that for all $1 \leq i \leq N$, $|\{j \in \ell : i \in I_j\}| = \lceil \eta\ell/8R \rceil$, and for all $j \in \ell$,

$$\|\mathbf{D}|_{I_j \times I_j}\|_\infty \leq \eta' \implies \|\mathbf{A}|_{I_j \times I_j}\|_\infty \leq \eta, \quad (87)$$

where for a matrix $\mathbf{M} = ((m_{ij})) \in \mathbb{R}^{s \times t}$ and for sets $S \subseteq \{1, \dots, s\}, T \subseteq \{1, \dots, t\}$, we define $\mathbf{M}|_{S \times T} := ((m_{ij}))_{i \in S, j \in T} \in \mathbb{R}^{|S| \times |T|}$.

Now, for $j \in [\ell]$,

$$\begin{aligned}
&\frac{\mathbb{P}(\mathbf{X}_{I_j} = \mathbf{y} | \mathbf{X}_{I_j^c} = \mathbf{x}_{-I_j})}{\mathbb{P}(\mathbf{X}_{I_j} = \mathbf{y}' | \mathbf{X}_{I_j^c} = \mathbf{x}_{-I_j})} \\
&= \frac{\exp\left(\mathbf{y}_{I_j}^\top \mathbf{D}|_{I_j \times I_j} \mathbf{y}_{I_j} + \sum_{u \in I_j, v \notin I_j} d_{uv} y_u x_v + \sum_{i \in I_j} h_i y_i\right)}{\exp\left(\mathbf{y}'_{I_j}^\top \mathbf{D}|_{I_j \times I_j} \mathbf{y}'_{I_j} + \sum_{u \in I_j, v \notin I_j} d_{uv} y'_u x_v + \sum_{i \in I_j} h_i y'_i\right)}.
\end{aligned} \quad (88)$$

Observe that the RHS of (88) is the probability mass function of an Ising model μ on $\{-1, 1\}^{|I_j|}$ with interaction matrix $\mathbf{D}|_{I_j \times I_j}$ and external magnetic field term at site i given by

$$h'_i = \sum_{v \notin I_j} D_{iv} x_v + h_i.$$

Recall from (87) that $\|\mathbf{A}|_{I_j \times I_j}\|_\infty \leq \eta$. The next step is to show that if $\mathbf{Y} \sim \mu$, then

$$\min_{1 \leq u \leq |I_j|} \text{Var}(Y_u | \mathbf{Y}_{-u}) \geq \Upsilon.$$

Towards this, note that for any $1 \leq u \leq |I_j|$,

$$\mathbb{P}(Y_u = 1 | \mathbf{Y}_{-u} = \mathbf{y}_{-u}) = \mathbb{P}(X_{I_j^u} = 1 | \mathbf{X}_{I_j \setminus \{I_j^u\}} = \mathbf{y}_{-u}, \mathbf{X}_{-I_j} = \mathbf{x}_{-I_j})$$

where I_j^u is the u -th smallest element of I_j . Hence,

$$\text{Var}(Y_u | \mathbf{Y}_{-u} = \mathbf{y}_{-u}) = \text{Var}(X_{I_j^u} | \mathbf{X}_{I_j \setminus \{I_j^u\}} = \mathbf{y}_{-u}, \mathbf{X}_{-I_j} = \mathbf{x}_{-I_j}) \geq \Upsilon,$$

since (86) is a (R, Υ) -Ising model. This implies that $X_{I_j} | \mathbf{X}_{I_j^c}$ is (η, Υ) -Ising model, for $j \in [\ell]$.

To prove (27), let us pick $j \in [\ell]$ uniformly at random, and note that for any vector \mathbf{a} ,

$$\mathbb{E} \left(\sum_{i \in I_j} a_i \right) = \sum_{i=1}^N a_i \mathbb{P}(I_j \ni i) = \sum_{i=1}^N a_i \frac{\lceil \eta \ell / 8R \rceil}{\ell} \geq \frac{\eta}{8R} \sum_{i=1}^N a_i.$$

This means that there is a fixed sample point j , such that

$$\sum_{i \in I_j} a_i \geq \frac{\eta}{8R} \sum_{i=1}^N a_i.$$

This completes the proof of Lemma 3.

A.3. Variance Lower Bound for Linear Functions. In this section we derive a variance lower bound for linear functions in (R, Υ) -Ising models. This follows the proof of Lemma 10 in [29] adapted to our setting. We begin with the following definition: For two probability measures μ and ν , the L^1 -Wasserstein distance is defined as:

$$W_1(\mu, \nu) := \min_{\pi \in \mathcal{C}_{\mu, \nu}} \mathbb{E}_{(\mathbf{U}, \mathbf{V}) \sim \pi} \|\mathbf{U} - \mathbf{V}\|_1,$$

where $\mathcal{C}_{\mu, \nu}$ denotes the set of all couplings of the probability measures μ and ν .

Lemma 11. *Let $\mathbf{X} \in \{-1, 1\}^N$ be a sample from an (R, Υ) -Ising model for some $R < 1/8$. Then, for each $i \in [N]$,*

$$W_1(\mathbb{P}_{\mathbf{X}_{-i}|X_i=1}, \mathbb{P}_{\mathbf{X}_{-i}|X_i=-1}) \leq \frac{16R}{1-8R}.$$

Proof. It follows from Lemma 4.9 in [28], that there exists a coupling π of the conditional measures $\mathbb{P}_{\mathbf{X}_{-i}|X_i=1}$ and $\mathbb{P}_{\mathbf{X}_{-i}|X_i=-1}$, such that:

$$\mathbb{E}_{(\mathbf{U}, \mathbf{V}) \sim \pi} [d_H(\mathbf{U}, \mathbf{V})] \leq \frac{\alpha}{1-\alpha}, \quad (89)$$

where d_H denotes the Hamming distance and α the Dobrushin coefficient. By Lemma 4.4 in [20], we know that Dobrushin's interdependence matrix is given by $8\mathbf{D}$. Hence, it follows from (89) that (Dobrushin's coefficient in Theorem 2.3 in [28] is given by $\alpha = 8\|\mathbf{D}\|_2$, see Theorem 4.3 in [20]),

$$W_1(\mathbb{P}_{\mathbf{X}_{-i}|X_i=1}, \mathbb{P}_{\mathbf{X}_{-i}|X_i=-1}) \leq 2 \mathbb{E}_{(\mathbf{U}, \mathbf{V}) \sim \pi} [d_H(\mathbf{U}, \mathbf{V})] \leq \frac{16\|\mathbf{D}\|_2}{1-8\|\mathbf{D}\|_2} \leq \frac{16R}{1-8R}.$$

which completes the proof of the lemma. \square

Using the above lemma, we now prove the desired variance lower bound:

Lemma 12. *Let \mathbf{X} be a sample from an (R, Υ) -Ising model for some $R > 0$. Then for any vector $\mathbf{a} \in \mathbb{R}^N$,*

$$\text{Var}(\mathbf{a}^\top \mathbf{X}) \gtrsim \frac{\|\mathbf{a}\|_2^2 \Upsilon^2}{R}.$$

Proof. First, consider the case $R \leq \Upsilon/32 \leq 1/32$. In this case, $1 - 8R \geq 3/4$, so

$$\frac{8R}{1-8R} \leq \frac{\Upsilon}{3} \quad (90)$$

Now, it follows from Lemma 11 that for every i ,

$$\sum_{j \in [N] \setminus \{i\}} |\mathbb{E}(X_j | X_i = 1) - \mathbb{E}(X_j | X_i = -1)| \leq \sum_{j \in [N] \setminus \{i\}} W_1(\mathbb{P}_{X_j|X_i=1}, \mathbb{P}_{X_j|X_i=-1})$$

$$\begin{aligned}
&\leq W_1 \left(\mathbb{P}_{\mathbf{X}_{-i}|X_i=1}, \mathbb{P}_{\mathbf{X}_{-i}|X_i=-1} \right) \\
&\leq \frac{16R}{1-8R}.
\end{aligned} \tag{91}$$

Next, note that:

$$\begin{aligned}
\text{Cov}(X_i, X_j) &= \mathbb{E}[(X_i - \mathbb{E}X_i)X_j] \\
&= \mathbb{E}[(X_i - \mathbb{E}X_i)\mathbb{E}(X_j|X_i)] \\
&= \mathbb{P}(X_i = 1)(1 - \mathbb{E}X_i)\mathbb{E}(X_j|X_i = 1) - \mathbb{P}(X_i = -1)(1 + \mathbb{E}X_i)\mathbb{E}(X_j|X_i = -1) \\
&= \frac{1 + \mathbb{E}X_i}{2}(1 - \mathbb{E}X_i)\mathbb{E}(X_j|X_i = 1) - \frac{1 - \mathbb{E}X_i}{2}(1 + \mathbb{E}X_i)\mathbb{E}(X_j|X_i = -1) \\
&= \frac{1}{2}[1 - (\mathbb{E}X_i)^2][\mathbb{E}(X_j|X_i = 1) - \mathbb{E}(X_j|X_i = -1)] \\
&\leq \frac{1}{2}[\mathbb{E}(X_j|X_i = 1) - \mathbb{E}(X_j|X_i = -1)]
\end{aligned} \tag{92}$$

Combining (90), (91) and (92), we have for every i ,

$$\sum_{j \in [N] \setminus \{i\}} |\text{Cov}(X_i, X_j)| \leq \frac{8R}{1-8R} \leq \frac{\Upsilon}{3}. \tag{93}$$

Hence, we have by (93),

$$\begin{aligned}
\text{Var}(\mathbf{a}^\top \mathbf{X}) &\geq \sum_{i=1}^N a_i^2 \text{Var}(X_i) - \sum_{i \neq j} |a_i a_j \text{Cov}(X_i, X_j)| \\
&\geq \sum_{i=1}^N a_i^2 \text{Var}(X_i) - \sum_{i \neq j} \frac{(a_i^2 + a_j^2) |\text{Cov}(X_i, X_j)|}{2} \\
&= \sum_{i=1}^N a_i^2 \left[\text{Var}(X_i) - \sum_{j \in [N] \setminus \{i\}} |\text{Cov}(X_i, X_j)| \right] \\
&\geq \sum_{i=1}^N a_i^2 \left(\Upsilon - \frac{\Upsilon}{3} \right) \\
&= \frac{2\Upsilon}{3} \|\mathbf{a}\|_2^2 \geq \frac{\|\mathbf{a}\|_2^2 \Upsilon^2}{R} \cdot \frac{2R}{3} \gtrsim \frac{\|\mathbf{a}\|_2^2 \Upsilon^2}{R}.
\end{aligned} \tag{94}$$

Now consider the case $R > \Upsilon/32$. By Lemma 3, we choose a subset I of $[N]$ such that conditioned on \mathbf{X}_{I^c} , \mathbf{X}_I is a $(\Upsilon/32, \Upsilon)$ - Ising model, and

$$\|\mathbf{a}_I\|_2^2 \geq \frac{\Upsilon}{256R} \|\mathbf{a}\|_2^2. \tag{95}$$

Hence, we have from (94) and (95),

$$\text{Var}(\mathbf{a}^\top \mathbf{X} | \mathbf{X}_{I^c}) = \text{Var}(\mathbf{a}_I^\top \mathbf{X}_I | \mathbf{X}_{I^c}) \geq \frac{2\Upsilon \|\mathbf{a}_I\|_2^2}{3} \geq \frac{\Upsilon^2 \|\mathbf{a}\|_2^2}{384R}. \tag{96}$$

Lemma 12 now follows from (96) on observing that $\text{Var}(\mathbf{a}^\top \mathbf{X}) \geq \mathbb{E}[\text{Var}(\mathbf{a}^\top \mathbf{X} | \mathbf{X}_{I^c})]$. \square

DEPARTMENT OF STATISTICS AND DATA SCIENCE, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE

Email address: somabha@nus.edu.sg

DEPARTMENT OF STATISTICS, UNIVERSITY OF FLORIDA, GAINESVILLE, USA

Email address: shalder@ufl.edu

DEPARTMENT OF STATISTICS AND DATA SCIENCE, UNIVERSITY OF PENNSYLVANIA, PHILADELPHIA, USA

Email address: bhaswar@wharton.upenn.edu

DEPARTMENT OF STATISTICS AND INFORMATICS INSTITUTE, UNIVERSITY OF FLORIDA, GAINESVILLE, USA

Email address: gmichail@ufl.edu