

# Hom-Leibniz bialgebras and BiHom-Leibniz dendriform algebras

ISMAIL LARAIEDH<sup>1,2</sup> AND SERGEI SILVESTROV<sup>3</sup>

<sup>1</sup>Departement of Mathematics, Faculty of Sciences,  
Sfax University, Box 1171, 3000 Sfax, Tunisia  
ismail.laraiedh@gmail.com

<sup>2</sup>Departement of Mathematics,  
College of Sciences and Humanities Al Quwaiyah,  
Shaqra University, Kingdom of Saudi Arabia  
ismail.laraiedh@su.edu.sa

<sup>3</sup>Division of Mathematics and Physics,  
School of Education, Culture and Communication,  
Mälardalen University, Box 883, 72123 Västerås, Sweden  
sergei.silvestrov@mdh.se

January 21, 2022

## Abstract

The notion of a Hom-Leibniz bialgebra is introduced and it is shown that matched pairs of Hom-Leibniz algebras, Manin triples of Hom-Leibniz algebras and Hom-Leibniz bialgebras are equivalent in a certain sense. The notion of Hom-Leibniz dendriform algebra is established, their bimodules and matched pairs are defined and their properties and theorems about their interplay and construction are obtained. Furthermore, the concept of BiHom-Leibniz dendriform algebras is introduced and discussed, their bimodules and matched pairs are constructed and properties are described. Finally, the connections between all these algebraic structures using  $\mathcal{O}$ -operators are shown.

## 1 Introduction

Hom-Lie algebras and quasi-Hom-Lie algebras were introduced first by Hartwig, Larson, and Silvestrov in 2003 in [29] devoted to a general method for construction of deformations and discretizations of Lie algebras of vector fields and deformations of Witt and Virasoro type algebras based on general twisted derivations ( $\sigma$ -derivations) obeying twisted Leibniz rule, and motivated also by the examples of  $q$ -deformed Jacobi identities in  $q$ -deformations of Witt and Visaroro algebras and in related  $q$ -deformed

---

2020 *Mathematics subject classification*: 17B61, 17D30

*Keywords*: Hom-Leibniz bialgebra, BiHom-Leibniz dendriform algebra, bimodule,  $\mathcal{O}$ -operator

algebras discovered in 1990'th in string theory, vertex models of conformal field theory, quantum field theory and quantum mechanics, and  $q$ -deformed differential calculi and  $q$ -deformed homological algebra [2, 16–19, 24, 30, 36, 46–48]. In 2005, Larsson and Silvestrov introduced quasi-Lie and quasi-Leibniz algebras [41] and color quasi-Lie and color quasi-Leibniz algebras [42] incorporating within the same framework Hom-Lie algebras and quasi-Hom-Lie algebras, the color Hom-Lie algebras and Hom-Lie superalgebras, the color quasi-Hom-Lie algebras and quasi-Hom-Lie superalgebras, as well as the quasi-Leibniz algebras, the color quasi-Leibniz algebras and super quasi-Leibniz algebras. The central extensions and cocycle conditions have been first considered for quasi-Hom-Lie algebras and Hom-Lie algebras in [29, 40] and for graded color quasi-Hom-Lie algebras in [76]. In quasi-Lie algebras, the skew-symmetry and the Jacobi identity are twisted in a special way by linear maps, and in general the quasi-Lie Jacobi identity contains six twisted triple bracket terms. In Hom-Lie algebras, the bilinear product satisfies the non-twisted skew-symmetry property as in Lie algebras, and the Hom-Lie algebras Jacobi identity has three terms twisted by a single linear map. Lie algebras are a special case of Hom-Lie algebras when the twisting linear map is the identity map. For other twisting linear maps however the Hom-Lie algebras are different and in many ways richer algebraic structures with classifications, deformations, representations, morphisms, derivations and homological structures in the fundamental ways dependent on joint properties of the twisting map and bilinear product which are in the intricate way interlinked by Hom-Jacobi identity. Hom-Lie admissible algebras have been considered first in [61], where the Hom-associative algebras and more general  $G$ -Hom-associative algebras including the Hom-Vinberg algebras (Hom-left symmetric algebras), Hom-pre-Lie algebras (Hom-right symmetric algebras), and some other new Hom-algebra structures have been introduced and shown to be Hom-Lie admissible, in the sense that the operation of commutator as new product in these Hom-algebras structures yields Hom-Lie algebras. Furthermore, in [61], flexible Hom-algebras have been introduced and connections to Hom-algebra generalizations of derivations and of adjoint derivations maps have been considered, investigations of the classification problems for Hom-Lie algebras have been initiated with constriction of families of the low-dimensional Hom-Lie algebras. Following the footsteps of [29, 40–43, 61], many Hom-algebra structures were developed and shown to link on the fundamental axiomatic level within common framework various deformed and non-deformed associative and non-associative algebraic structures, super-algebras and color algebras,  $n$ -ary algebraic structures, their formal deformations and quantum deformations and related non-commutative differential and homological algebra structures. In [1, 3, 4, 7–14, 21–23, 25, 31–34, 37–40, 43–45, 53–60, 62–74, 76–86] Hom-modules and Hom-bimodules, Hom-Lie admissible Hom-coalgebras, Hom-coalgebras, Hom-bialgebras, Hom-Hopf algebras,  $L$ -modules,  $L$ -comodules and Hom-Lie quasi-bialgebras,  $n$ -ary generalizations of BiHom-Lie algebras and BiHom-associative algebras and generalized derivations, Rota-Baxter operators, Hom-dendriform color algebras, Rota-Baxter bisystems and covariant bialgebras, Rota-Baxter cosystems, coquasi-triangular mixed bialgebras, coassociative Yang-Baxter pairs, coassociative Yang-Baxter equation and generalizations of Rota-Baxter systems and algebras, curved  $\mathcal{O}$ -operator systems and their connections with tridendriform systems and pre-Lie algebras, BiHom-algebras, BiHom-Frobenius algebras and double constructions, infinitesimal BiHom-bialgebras and Hom-dendriform  $D$ -bialgebras and Hom-omni-Lie algebras

and Hom-Gerstenhaber algebras and Hom-Lie algebroids.

Dendriform algebras, introduced by Loday as algebras with two operations which dichotomize the notion of associative algebras [51], are connected to K-theory, Hopf algebras, homotopy Gerstenhaber algebra, operads, homology, combinatorics, and quantum field theory where they occur in the theory of renormalization of Connes and Kreimer. Then, the tridendriform algebras were introduced by Loday and Ronco in their study of polytopes and Koszul duality [52]. Connections of dendriform algebras to Rota-Baxter algebras have been considered in [26]. Hom-tridendriform algebras, generalizing tridendriform algebras by twisting their axioms by arbitrary linear maps in a special ways, together with connections to Rota-Baxter Hom-algebras generalising Rota-Baxter algebras, have been introduced in [58]. Connections between Hom-bialgebras and Hom-coalgebras, Hom-dendriform and tridendriform systems and Rota-Baxter and Hom-Rota-Baxter Hom-algebra structures have been investigated in [53–56, 58, 65].

In this article, we introduce the notion of a Hom-Leibniz bialgebras and show that matched pairs of Hom-Leibniz algebras, Manin triples of Hom-Leibniz algebras and Hom-Leibniz bialgebras are equivalent in a certain sense. Next, we establish the notion of Hom-Leibniz dendriform algebra. Their bimodules and matched pairs are defined and the relevant properties and theorems are obtained. Furthermore, the concept of BiHom-Leibniz dendriform algebras is introduced and discussed. Their bimodules and matched pairs are also constructed, and related relevant properties are given. Finally, the connections between all these algebraic structures using  $\mathcal{O}$ -operators are shown.

The paper is organized as follows. In Section 2, we summarize the definition of Hom-Leibniz algebras and the concepts of bimodule and matched pairs and we give some relevant properties. In Section 3, we introduce the notions of a Manin triple of Hom-Leibniz algebras and a Hom-Leibniz bialgebra. We prove the equivalence between matched pairs of Hom-Leibniz algebras, Manin triples of Hom-Leibniz algebras and Hom-Leibniz bialgebras. In Section 4, we establish definition of Hom-Leibniz dendriform algebra. Similar to the connection from Hom-pre-Lie algebras to Hom-Lie algebras and from Hom-dendriform algebras to Hom-associative algebras, we show that a Hom-Leibniz dendriform algebra gives rise to a Hom-Leibniz algebra. Next, we develop some construction theorems. Their bimodule and matched pair are defined and their related relevant properties are also given. Section 5, is devoted to reminders of fundamental concepts for BiHom-Leibniz algebras. Some results of bimodules and matched pair are given. In Section 6, we introduce definition of BiHom-Leibniz dendriform algebra and we give some key of constructions. Their bimodule and matched pair are defined and their related relevant properties are also given. Also, we study the notion of  $\mathcal{O}$ -operator and we illustrate the relations existing between BiHom-Leibniz and BiHom-Leibniz dendriform algebras.

## 2 Preliminaries on Hom-Leibniz algebras

In this section definitions and basic facts on Hom-Leibniz algebras are reviewed.

Throughout this article all linear spaces are assumed to be over a field  $\mathbb{K}$ , and multilinear maps  $f: V_1 \times \cdots \times V_n \rightarrow W$  on finite direct products and linear maps  $F: V_1 \otimes \cdots \otimes V_n \rightarrow W$ , on finite tensor products of linear spaces are identified in the usual way via  $F(v_1 \otimes \cdots \otimes v_n) = f(v_1, \dots, v_n)$ .

Hom-modules are pairs  $(M, \alpha_M)$  consisting of a  $\mathbb{K}$ -module  $M$  over a field  $\mathbb{K}$  and a linear map  $\alpha_M : M \rightarrow M$ . A morphism  $f : (M, \alpha_M) \rightarrow (N, \alpha_N)$  of Hom-modules is a linear map  $f : M \rightarrow N$  such that  $f\alpha_M = \alpha_N f$ . Hom-algebras are triples  $(A, \mu, \alpha)$  where  $(A, \alpha)$  is a Hom-module and  $\mu : A \times A \rightarrow A$  is a bilinear map. If, moreover, the linear map  $\alpha : A \rightarrow A$  satisfies the multiplicativity,  $\alpha(\mu(x, y)) = \mu(\alpha(x), \alpha(y))$ , that is if  $\alpha$  is an algebra endomorphism of the algebra  $(A, \mu)$ , then the Hom-algebra  $(A, \mu(\cdot, \cdot), \alpha)$  is called multiplicative.

**Definition 2.1** ([41, 61]). Hom-Leibniz algebras are triples  $(A, [\cdot, \cdot], \alpha)$  consisting of a linear space  $A$  over a field  $\mathbb{K}$ , a bilinear map  $[\cdot, \cdot] : A \times A \rightarrow A$  and a linear map  $\alpha : A \rightarrow A$  satisfying for all  $x, y, z \in A$ ,

$$[\alpha(x), [y, z]] = [[x, y], \alpha(z)] + [\alpha(y), [x, z]]. \quad (\text{Hom-Leibniz identity}) \quad (2.1)$$

If, moreover, the linear map  $\alpha : A \rightarrow A$  is an algebra endomorphism, meaning that it satisfies also the multiplicativity property

$$\alpha([x, y]) = [\alpha(x), \alpha(y)], \quad (2.2)$$

then the Hom-Leibniz algebra is called multiplicative.

**Definition 2.2** ([29, 61]). Hom-Lie algebras are triples  $(A, [\cdot, \cdot], \alpha)$  consisting of a linear space  $A$  over a field  $\mathbb{K}$ , a bilinear map  $[\cdot, \cdot] : A \times A \rightarrow A$  and a linear map  $\alpha : A \rightarrow A$  satisfying for all  $x, y, z \in A$ ,

$$[x, y] = -[y, x] \quad (\text{Skew-symmetry identity}) \quad (2.3)$$

$$[\alpha(x), [y, z]] + [\alpha(y), [z, x]] + [\alpha(z), [x, y]] = 0. \quad (\text{Hom-Jacobi identity}) \quad (2.4)$$

When, moreover, the linear map  $\alpha : A \rightarrow A$  satisfies multiplicativity (2.2), that is when  $\alpha$  is an algebra endomorphism, the Hom-Lie algebra  $(A, [\cdot, \cdot], \alpha)$  is called multiplicative.

*Remark 2.3.* If  $\alpha = id_A$ , then  $(A, [\cdot, \cdot]) = (A, [\cdot, \cdot], id_A)$  in Definition 2.1 is a Leibniz algebra. So, Leibniz algebras are a special case of Hom-Leibniz algebras when the twisting linear map is the identity map, that is when the fundamental defining Leibniz identity is not deformed at all.

*Examples 2.4.* Some basic examples of Hom-Leibniz algebras are as follows.

1. Any Hom-Lie algebra is automatically a Hom-Leibniz algebra as in the presence of skew-symmetry and the Hom-Leibniz identity implies Hom-Jacobi identity.
2. Given a Leibniz algebra  $(A, [\cdot, \cdot])$  and a Leibniz algebra morphism  $\alpha : A \rightarrow A$ , one always get a Hom-Leibniz algebra  $(A, [\cdot, \cdot]_\alpha, \alpha)$ , where  $[x, y]_\alpha = [\alpha(x), \alpha(y)]$ .
3. Any Hom-module  $(M, \alpha_M)$ , together with the trivial zero bracket  $[\cdot, \cdot]$ , that is when  $[x, y] = 0$  for all  $x, y \in M$ , is a Hom-Leibniz algebra, called abelian or commutative Hom-Leibniz algebra.
4. The two-dimensional  $\mathbb{K}$ -linear space  $A$  with basis  $\{e_1, e_2\}$ , the bilinear operation given by  $[e_2, e_2] = e_1$  and zero elsewhere, and the endomorphism  $\alpha$  given by  $\alpha(e_1) = e_1$ ,  $\alpha(e_2) = e_1 + e_2$  define a Hom-Leibniz algebra. This Hom-Leibniz algebra is multiplicative, since

$$\alpha([e_j, e_k]) = \begin{cases} \alpha(0) = 0 = [\alpha(e_j), \alpha(e_k)], & (j, k) \neq (2, 2) \\ \alpha(e_1) = e_1 = [e_2, e_2] = [e_1 + e_2, e_1 + e_2] \\ \quad = [\alpha(e_2), \alpha(e_2)], & j = k = 2 \end{cases}$$

but it is not a Hom-Lie algebra since the skew-symmetry identity is not satisfied  $[e_2, e_2] = e_1 \neq -e_1 = -[e_2, e_2]$ .

**Definition 2.5** ([20]). Let  $(A, [\cdot, \cdot], \alpha)$  be a Hom-Leibniz algebra and let  $(V, \beta)$  be a Hom-module. Let  $l, r : A \rightarrow gl(V)$  be two linear maps. The quadruple  $(l, r, \beta, V)$  is called a bimodule of  $A$  if for all  $x, y \in A, v \in V$ :

$$l(\alpha(x))l(y)v = l([x, y])\beta(v) + l(\alpha(y))l(x)v, \quad (2.5)$$

$$l(\alpha(x))r(y)v = r(\alpha(y))l(x)v + r([x, y])\beta(v), \quad (2.6)$$

$$r([x, y])\beta(v) = r(\alpha(y))r(x)v + l(\alpha(x))r(y)v, \quad (2.7)$$

$$\beta(l(x)v) = l(\alpha(x))\beta(v), \quad (2.8)$$

$$\beta(r(x)v) = r(\alpha(x))\beta(v). \quad (2.9)$$

*Remark 2.6.* By using (2.6) and (2.7), observe that for all  $x, y \in A$  and  $v \in V$ ,

$$r(\alpha(y))l(x)v = -r(\alpha(y))r(x)v. \quad (2.10)$$

**Proposition 2.7.** Let  $(l, r, \beta, V)$  is a bimodule of a Hom-Leibniz algebra  $(A, [\cdot, \cdot], \alpha)$ . Then the direct sum of linear spaces,  $A \oplus V$ , is turned into a Hom-Leibniz algebra by defining multiplication in  $A \oplus V$  for all  $x_1, x_2 \in A, v_1, v_2 \in V$  by

$$\begin{aligned} [x_1 + v_1, x_2 + v_2]' &= [x_1, x_2] + (l(x_1)v_2 + r(x_2)v_1), \\ (\alpha \oplus \beta)(x_1 + v_1) &= \alpha(x_1) + \beta(v_1). \end{aligned} \quad (2.11)$$

*Proof.* For all  $v_1, v_2, v_3 \in V$  and  $x_1, x_2, x_3 \in A$ ,

$$\begin{aligned} & [(\alpha + \beta)(x_1 + v_1), [x_2 + v_2, x_3 + v_3]']' \\ & - [[x_1 + v_1, x_2 + v_2]', (\alpha + \beta)(x_3 + v_3)]' \\ & - [(\alpha + \beta)(x_2 + v_2), [x_1 + v_1, x_3 + v_3]']' \\ & = [\alpha(x_1) + \beta(v_1), [x_2, x_3] + l(x_2)v_3 + r(x_3)v_2]' \\ & - [[x_1, x_2] + l(x_1)v_2 + r(x_2)v_1, \alpha(x_3) + \beta(v_3)]' \\ & - [\alpha(x_2) + \beta(v_2), [x_1, x_3] + l(x_1)v_3 + r(x_3)v_1]' \\ & = [\alpha(x_1), [x_2, x_3]] + l(\alpha(x_1))l(x_2)v_3 + l(\alpha(x_1))r(x_3)v_2 + r([x_2, x_3])\beta(v_1) \\ & - [[x_1, x_2], \alpha(x_3)] - l([x_1, x_2])\beta(v_3) - r(\alpha(x_3))l(x_1)v_2 - r(\alpha(x_3))r(x_2)v_1 \\ & - [\alpha(x_2), [x_1, x_3]] - l(\alpha(x_2))l(x_1)v_3 - r(\alpha(x_3))l(x_1)v_2 - r(\alpha(x_3))r(x_2)v_1. \end{aligned}$$

Then, by (2.1) and (2.5)-(2.7),

$$\begin{aligned} & [(\alpha + \beta)(x_1 + v_1), [x_2 + v_2, x_3 + v_3]']' - [[x_1 + v_1, x_2 + v_2]', (\alpha + \beta)(x_3 + v_3)]' \\ & - [(\alpha + \beta)(x_2 + v_2), [x_1 + v_1, x_3 + v_3]']' = 0, \end{aligned}$$

which completes the proof.  $\square$

We denote such a Hom-Leibniz algebra  $(A \oplus V, [\cdot, \cdot]', \alpha + \beta)$ , or  $A \times_{l,r,\alpha,\beta} V$ .

*Example 2.8.* If  $(A, [\cdot, \cdot], \alpha)$  is a Hom-Leibniz algebra, then  $(L, 0, \alpha, A)$  and  $(L, R, \alpha, A)$  are bimodules of  $(A, [\cdot, \cdot], \alpha)$ , where  $L(a)b = [a, b]$  and  $R(a)b = [b, a]$  for all  $a, b \in A$ .

**Proposition 2.9.** *If  $(A, [\cdot, \cdot], \alpha)$  is a multiplicative Hom-Leibniz algebra, then the quadruple  $(\alpha \otimes L, \alpha \otimes R, \alpha \otimes \alpha, A \otimes A)$ , where for any  $x, a, b \in A$ ,*

$$\begin{aligned}(\alpha \otimes L)(x)(a \otimes b) &= (\alpha \otimes L(x))(a \otimes b) = \alpha(a) \otimes L(x)b = \alpha(a) \otimes [x, b], \\(\alpha \otimes R)(x)(a \otimes b) &= (\alpha \otimes R(x))(a \otimes b) = \alpha(a) \otimes R(x)b = \alpha(a) \otimes [b, x],\end{aligned}$$

*is a bimodule of  $A$ .*

*Proof.* By formulas for the maps and Hom-Leibniz identity (2.1), for all  $x, y, v_1, v_2 \in A$ :

$$\begin{aligned}(\alpha \otimes L)(\alpha(x))(\alpha \otimes L)(y)(v_1 \otimes v_2) &= (\alpha \otimes L(\alpha(x)))(\alpha(v_1)) \otimes L(y)v_2 \\&= v_1 \otimes L(\alpha(x))L(y)v_2 \\&= v_1 \otimes [\alpha(x), [y, v_2]], \\(\alpha \otimes L)([x, y])(\alpha \otimes \alpha)(v_1 \otimes v_2) &= (\alpha \otimes L)([x, y])(\alpha(v_1) \otimes \alpha(v_2)) \\&= v_1 \otimes L([x, y])\alpha(v_2) \\&= v_1 \otimes [[x, y], \alpha(v_2)], \\(\alpha \otimes L)(\alpha(y))(\alpha \otimes L)(x)(v_1 \otimes v_2) &= (\alpha \otimes L)(\alpha(y))(\alpha \otimes L(x))(v_1 \otimes v_2) \\&= (\alpha \otimes L)(\alpha(y))(\alpha(v_1) \otimes L(x)v_2) \\&= v_1 \otimes L(\alpha(y))L(x)v_2 \\&= v_1 \otimes [\alpha(y), [x, v_2]], \\(\alpha \otimes L)(\alpha(x))(\alpha \otimes L)(y)(v_1 \otimes v_2) &= v_1 \otimes [\alpha(x), [y, v_2]] \\&\stackrel{(2.1)}{=} v_1 \otimes ([[x, y], \alpha(v_2)] + [\alpha(y), [x, v_2]]) \\&= v_1 \otimes [[x, y], \alpha(v_2)] + v_1 \otimes [\alpha(y), [x, v_2]] \\&= (\alpha \otimes L)([x, y])(\alpha \otimes \alpha)(v_1 \otimes v_2) + (\alpha \otimes L)(\alpha(y))(\alpha \otimes L)(x)(v_1 \otimes v_2); \\(\alpha \otimes L)(\alpha(x))(\alpha \otimes R)(y)(v_1 \otimes v_2) &= (\alpha \otimes L(\alpha(x)))(\alpha \otimes R(y))(v_1 \otimes v_2) \\&= (\alpha \otimes L(\alpha(x)))(\alpha(v_1) \otimes R(y)v_2) \\&= v_1 \otimes [\alpha(x), [v_2, y]], \\(\alpha \otimes R)(\alpha(y))(\alpha \otimes L)(x)(v_1 \otimes v_2) &= (\alpha \otimes R(\alpha(y)))(\alpha \otimes L(x))(v_1 \otimes v_2) \\&= (\alpha \otimes R(\alpha(y)))(\alpha(v_1) \otimes L(x)v_2) \\&= v_1 \otimes [[x, v_2], \alpha(y)], \\(\alpha \otimes R)([x, y])(\alpha \otimes \alpha)(v_1 \otimes v_2) &= (\alpha \otimes R([x, y]))(\alpha(v_1) \otimes \alpha(v_2)) \\&= v_1 \otimes [\alpha(v_2), [x, y]], \\(\alpha \otimes L)(\alpha(x))(\alpha \otimes R)(y)(v_1 \otimes v_2) &= v_1 \otimes [\alpha(x), [v_2, y]] \\&\stackrel{(2.1)}{=} v_1 \otimes ([[x, v_2], \alpha(y)] + [\alpha(v_2), [x, y]]) \\&= v_1 \otimes [[x, v_2], \alpha(y)] + v_1 \otimes [\alpha(v_2), [x, y]] \\&= (\alpha \otimes R)(\alpha(y))(\alpha \otimes L)(x)(v_1 \otimes v_2) + (\alpha \otimes R)([x, y])(\alpha \otimes \alpha)(v_1 \otimes v_2); \\(\alpha \otimes R)([x, y])(\alpha \otimes \alpha)(v_1 \otimes v_2) &= (\alpha \otimes R([x, y]))(\alpha(v_1) \otimes \alpha(v_2)) \\&= v_1 \otimes [\alpha(v_2), [x, y]], \\(\alpha \otimes R)(\alpha(y))(\alpha \otimes R)(x)(v_1 \otimes v_2) &= (\alpha \otimes R(\alpha(y)))(\alpha \otimes R(x))(v_1 \otimes v_2) \\&= \alpha \otimes R(\alpha(y))(\alpha \otimes R(x))(v_1 \otimes v_2)\end{aligned}$$

$$\begin{aligned}
&= (\alpha \otimes R(\alpha(y)))(\alpha(v_1) \otimes R(x)v_2) \\
&= v_1 \otimes [[v_2, x], \alpha(y)], \\
(\alpha \otimes L)(\alpha(x))(\alpha \otimes R)(y)(v_1 \otimes v_2) &= (\alpha \otimes L(\alpha(x)))(\alpha \otimes R(y))(v_1 \otimes v_2) \\
&= (\alpha \otimes L(\alpha(x)))(\alpha(v_1) \otimes R(y)v_2) \\
&= v_1 \otimes [\alpha(x), [[v_2, y]]], \\
(\alpha \otimes R)([x, y])(\alpha \otimes \alpha)(v_1 \otimes v_2) &= v_1 \otimes [\alpha(v_2), [x, y]] \\
&\stackrel{(2.1)}{=} v_1 \otimes ([[v_2, x], \alpha(y)] + [\alpha(x), [v_2, y]]) \\
&= v_1 \otimes [[v_2, x], \alpha(y)] + v_1 \otimes [\alpha(x), [v_2, y]] \\
&= (\alpha \otimes R)(\alpha(y))(\alpha \otimes R)(x)(v_1 \otimes v_2) + (\alpha \otimes L)(\alpha(x))(\alpha \otimes R)(y)(v_1 \otimes v_2).
\end{aligned}$$

By formulas for the maps,

$$\begin{aligned}
(\alpha \otimes \alpha)(\alpha \otimes L)(x)(v_1 \otimes v_2) &= (\alpha \otimes \alpha)(\alpha \otimes L(x))(v_1 \otimes v_2) \\
&= (\alpha \otimes \alpha)(\alpha(v_1) \otimes L(x)v_2) \\
&= v_1 \otimes L(\alpha(x))\alpha(v_2), \\
(\alpha \otimes L)(\alpha(x))((\alpha \otimes \alpha)(v_1 \otimes v_2)) &= (\alpha \otimes L)(\alpha(x))(\alpha(v_1) \otimes \alpha(v_2)) \\
&= v_1 \otimes L(\alpha(x))\alpha(v_2).
\end{aligned}$$

Therefore,  $(\alpha \otimes \alpha)(\alpha \otimes L)(x)(v_1 \otimes v_2) = (\alpha \otimes L)(\alpha(x))((\alpha \otimes \alpha)(v_1 \otimes v_2))$ . Similarly,  $(\alpha \otimes \alpha)(\alpha \otimes R)(x)(v_1 \otimes v_2) = (\alpha \otimes R)(\alpha(x))((\alpha \otimes \alpha)(v_1 \otimes v_2))$ .  $\square$

*Remark 2.10.* The quadruple  $(L \otimes \alpha, R \otimes \alpha, \alpha \otimes \alpha, A \otimes A)$  is also a bimodule of  $A$ .

**Theorem 2.11** ([6]). *Let  $(A, [\cdot, \cdot], \alpha)$  and  $(B, \{\cdot, \cdot\}, \beta)$  be two Hom-Leibniz algebras. Suppose there are linear maps  $l_A, r_A : A \rightarrow gl(B)$  and  $l_B, r_B : B \rightarrow gl(A)$  such that the quadruple  $(l_A, r_A, \beta, B)$  is a bimodule of  $A$ , and  $(l_B, r_B, \alpha, A)$  is a bimodule of  $B$ , satisfying, for any  $x, y \in A, a, b \in B$ , the following conditions:*

$$\begin{aligned}
l_A(\alpha(x))(\{a, b\}) &= l_A(r_B(a)x)\beta(b) + r_A(r_B(b)x)\beta(a) \\
&\quad + \{l_A(x)a, \beta(b)\} + \{\beta(a), l_A(x)b\}, \quad (2.12)
\end{aligned}$$

$$\begin{aligned}
\{\beta(a), l_A(x)b\} + r_A(r_B(b)x)\beta(a) &= \{r_A(x)a, \beta(b)\} \\
&\quad + l_A(l_B(a)x)\beta(b) + l_A(\alpha(x))(\{a, b\}), \quad (2.13)
\end{aligned}$$

$$\begin{aligned}
\{\beta(a), r_A(x)b\} + r_A(l_B(b)x)\beta(a) &= r_A(\alpha(x))(\{a, b\}) \\
&\quad + \{\beta(b), r_A(x)a\} + r_A(l_B(a)x)\beta(b), \quad (2.14)
\end{aligned}$$

$$\begin{aligned}
l_B(\beta(a))([x, y]) &= l_B(r_A(x)a)\alpha(y) + r_B(r_A(y)a)\alpha(x) \\
&\quad + [l_B(a)x, \alpha(y)] + [\alpha(x), l_B(a)y], \quad (2.15)
\end{aligned}$$

$$\begin{aligned}
[\alpha(x), l_B(a)y] + r_B(r_A(y)a)\alpha(x) &= [r_B(a)x, \alpha(y)] \\
&\quad + l_B(l_A(x)a)\alpha(y) + l_B(\beta(a))([x, y]), \quad (2.16)
\end{aligned}$$

$$\begin{aligned}
[\alpha(x), r_B(a)y] + r_B(l_A(y)a)\alpha(x) &= r_B(\beta(a))([x, y]) \\
&\quad + [\alpha(y), r_B(a)x] + r_B(l_A(x)a)\alpha(y). \quad (2.17)
\end{aligned}$$

Then,  $(A, B, l_A, r_A, \beta, l_B, r_B, \alpha)$  is called a matched pair of Hom-Leibniz algebras. In this case, there is a Hom-Leibniz algebra structure on the direct sum  $A \oplus B$  of the underlying linear spaces of  $A$  and  $B$  given for all  $x, y \in A, a, b \in B$  by

$$\begin{aligned}
[x + a, y + b]' &= ([x, y] + l_B(a)y + r_B(b)x) + \{a, b\} + l_A(x)b + r_A(y)a, \\
(\alpha \oplus \beta)(x + a) &= \alpha(x) + \beta(a). \quad (2.18)
\end{aligned}$$

We denote this Hom-Leibniz algebra by  $(A \bowtie B, [\cdot, \cdot]', \alpha + \beta)$  or  $A \bowtie_{l_B, r_B, \alpha}^{l_A, r_A, \beta} B$ .

*Remark 2.12.* Condition 2.17, can be reformulated equivalently as

$$[l_B(a)x, \alpha(y)] + l_B(r_A(x)a)\alpha(y) + [r_B(a)x, \alpha(y)] + l_B(l_A(x)a)\alpha(y) = 0. \quad (2.19)$$

### 3 Manin triples and bialgebras of Hom-Leibniz algebras

In this section, we consider the multiplicative Hom-Leibniz algebra  $(A, [\cdot, \cdot], \alpha)$  such that  $\alpha$  is involutive, i.e,  $\alpha^2 = id_A$ .

**Definition 3.1.** Let  $V_1, V_2$  be two linear spaces. For a linear map  $\phi : V_1 \rightarrow V_2$ , we denote by  $\phi^* : V_2^* \rightarrow V_1^*$  the dual (linear) map given by

$$\langle v, \phi^*(u^*) \rangle = -\langle \phi(v), u^* \rangle \text{ for all } v \in V_1, u^* \in V_2^*.$$

**Lemma 3.2.** *Let  $(l, r, \beta, V)$  be a bimodule of a multiplicative involutive Hom-Leibniz algebra  $(A, [\cdot, \cdot], \alpha)$ . Then, for linear maps  $l^*, r^* : A \rightarrow gl(V^*)$  such that*

$$\forall x \in A, u^* \in V^*, v \in V : \langle l^*(x)u^*, v \rangle = -\langle l(x)v, u^* \rangle, \langle r^*(x)u^*, v \rangle = -\langle r(x)v, u^* \rangle,$$

- (i)  $(l^*, r^*, \beta^*, V^*)$  is a bimodule of  $(A, [\cdot, \cdot], \alpha)$ ;
- (ii)  $(l^*, -l^* - r^*, \beta^*, V^*)$  is a bimodule of  $(A, [\cdot, \cdot], \alpha)$ ;
- (iii)  $(l^*, 0, \beta^*, V^*)$  and  $(0, r^*, \beta^*, V^*)$  are bimodules of  $(A, [\cdot, \cdot], \alpha)$ .

*Proof.* (i) For any  $x, y \in A, u^* \in A^*$  and  $v \in V$ ,

$$\begin{aligned} \langle l^*(\alpha(x))l^*(y)u^*, v \rangle &= \langle l(y)l(\alpha(x))v, u^* \rangle \\ &= \langle l(\alpha^2(y))l(\alpha(x))v, u^* \rangle \\ \text{(by (2.5))} &= \langle l(\alpha^2(x))l(\alpha(y))v - l([\alpha(x), \alpha(y)])\beta(v), u^* \rangle \\ &= \langle l(x)l(\alpha(y))v - \beta(l([x, y])v), u^* \rangle \\ &= \langle l^*(\alpha(y))l^*(x)u^* + l^*([x, y])\beta^*(u^*), v \rangle, \\ \langle l^*(\alpha(x))r^*(y)u^*, v \rangle &= \langle r(y)l(\alpha(x))v, u^* \rangle \\ &= -\langle r(x)l(\alpha(y))v - r([\alpha(x), \alpha(y)])\beta(v), u^* \rangle \\ &= \langle r^*(\alpha(y))l^*(x)u^* + r^*([x, y])\beta^*(u^*), v \rangle, \\ \langle r^*([x, y])\beta^*(u^*), v \rangle &= -\langle r([\alpha(x), \alpha(y)])\beta(v)u^* \rangle \\ &= \langle r(y)l(\alpha(x))v - l(x)r(\alpha(y))v, u^* \rangle \\ &= \langle l^*(\alpha(x))r^*(y)u^* - r^*(\alpha(y))l^*(x)u^*, v \rangle. \end{aligned}$$

(ii) For any  $x, y \in A, u^* \in A^*$  and  $v \in V$ ,

$$\begin{aligned} \langle l^*(\alpha(x))l^*(y)u^*, v \rangle &= \langle l(y)l(\alpha(x))v, u^* \rangle \\ &= \langle l(\alpha^2(y))l(\alpha(x))v, u^* \rangle \\ \text{(by (2.5))} &= \langle l(\alpha^2(x))l(\alpha(y))v - l([\alpha(x), \alpha(y)])\beta(v), u^* \rangle \\ &= \langle l(x)l(\alpha(y))v - \beta(l([x, y])v), u^* \rangle \\ &= \langle l^*(\alpha(y))l^*(x)u^* + l^*([x, y])\beta^*(u^*), v \rangle, \\ \langle l^*(\alpha(x))(-l^* - r^*)(y)u^*, v \rangle &= -\langle (l + r)(y)l(\alpha(x))v, u^* \rangle \end{aligned}$$



$$\begin{aligned}
&= -\langle (l+r)(\alpha^2(y)l(\alpha(x))v, u^* \rangle \\
\text{(by (2.5))} &= \langle l(\alpha^2(x)l(\alpha(y))v - l([\alpha(x), \alpha(y)])\beta(v), u^* \rangle \\
&\quad - \langle l(\alpha^2(x)r(\alpha(y))v - r([\alpha(x), \alpha(y)])\beta(v), u^* \rangle \\
&= -\langle l(x)(l+r)(\alpha(y))v - (l+r)([\alpha(x), \alpha(y)])\beta(v), u^* \rangle \\
&= -\langle (l^* + r^*)(\alpha(y))l^*(x)u^* + (l^* + r^*)[x, y]\beta^*(u^*), v \rangle \\
&= \langle (-l^* - r^*)(\alpha(y))l^*(x)u^* + (-l^* - r^*)[x, y]\beta^*(u^*), v \rangle.
\end{aligned}$$

Similarly, one can show

$$(-l^* - r^*)([x, y])\beta^*(u^*) = (-l^* - r^*)(\alpha(y))(-l^* - r^*)(x)u^* + l(\alpha(x))(-l^* - r^*)(y)u^*.$$

(iii) It is proved analogously.  $\square$

**Definition 3.3.** Let  $(A, [\cdot, \cdot], \alpha)$  be a Hom-Leibniz algebra, and  $B : A \times A \rightarrow \mathbb{K}$  be a bilinear form on  $A$ . Then,

(i)  $B$  is said to be nondegenerate if

$$A^\perp = \{x \in A \mid B(y, x) = 0, \forall y \in A\} = \{0_A\}; \quad (3.1)$$

(ii)  $B$  is said to be skew-symmetric if for all  $x, y \in A$ ,

$$B(x, y) = -B(y, x); \quad (3.2)$$

(iii)  $B$  is said to be  $\alpha$ -invariant if for all  $x, y \in A$ ,

$$B([\alpha(x), \alpha(z)] + [\alpha(z), \alpha(x)], \alpha(y)) = B(\alpha(x), [\alpha(y), \alpha(z)]). \quad (3.3)$$

**Definition 3.4.** A quadratic Hom-Leibniz algebra is a Hom-Leibniz algebra equipped with a non-degenerate skew-symmetric invariant bilinear form.

Recall that a quadratic Hom-Lie algebra is a Hom-Lie algebra  $(\mathfrak{g}, [\cdot, \cdot], \alpha)$  equipped with a non-degenerate symmetric bilinear form  $B$  which is invariant in the sense that

$$B([\alpha(x), \alpha(y)], \alpha(z)) = B(\alpha(x), [\alpha(y), \alpha(z)]), \quad \forall x, y, z \in \mathfrak{g}.$$

**Definition 3.5.** Let  $(A, [\cdot, \cdot], \alpha)$  be a Hom-Leibniz algebras. A subalgebra  $H$  of  $A$  is called isotropic if  $H \subseteq H^\perp$ .

**Definition 3.6.** Manin triples for Hom-Leibniz algebras are defined as quadruples  $(A, A_1, A_2, \alpha)$  where

- (i)  $(A, \cdot, \alpha, B)$  is a quadratic Hom-Leibniz algebra;
- (ii) both  $(A_1, \cdot_1, \alpha_1)$  and  $(A_2, \cdot_2, \alpha_2)$  are isotropic Hom-Leibniz subalgebras of  $(A, \alpha)$  with  $\alpha = \alpha_1 + \alpha_2$ ;
- (iii)  $A = A_1 \oplus A_2$  as linear spaces.

Two Manin triples  $(A, A_1, A_2, \alpha)$  and  $(C, C_1, C_2, \beta)$  with the bilinear forms  $B_1$  and  $B_2$  respectively are isomorphic if there exists an isomorphism of Hom-Leibniz algebras  $f : A \rightarrow C$  such that for all  $x, y \in A$ ,

$$f \circ \alpha = \beta \circ f, \quad f(A_1) = C_1, \quad f(A_2) = C_2, \quad B_1(x, y) = B_2(f(x), f(y)). \quad (3.4)$$

*Example 3.7.* If  $(A, [\cdot, \cdot], \alpha)$  is a Hom-Leibniz algebra, then  $(A \times_{L^*, -L^* - R^*, \alpha, \alpha^*} A^*, A, A^*)$  is a Manin triple of Hom-Leibniz algebras, where the natural non-degenerate skew-symmetric bilinear form  $B$  on  $A \oplus A^*$  is given by

$$B(x + a^*, y + b^*) = \langle a^*, y \rangle - \langle b^*, x \rangle, \quad \forall x, y \in A, \quad a^*, b^* \in A^*, \quad (3.5)$$

which is called the standard Manin triple.

**Proposition 3.8.** *Every Manin triple is isomorphic to the standard Manin triple.*

For a Hom-Leibniz algebra  $(A^*, \{\cdot, \cdot\}, \alpha^*)$ , let  $\Delta : A \rightarrow \otimes^2 A$  be the dual map of  $\{\cdot, \cdot\} : \otimes^2 A^* \rightarrow A^*$ , that is

$$\langle \Delta x, a^* \otimes b^* \rangle = \langle x, \{a^*, b^*\} \rangle.$$

**Definition 3.9.** Let  $(A, [\cdot, \cdot], \alpha)$  and  $(A^*, \{\cdot, \cdot\}, \alpha^*)$  be a Hom-Leibniz algebras. Then  $(A, A^*)$  is called a Hom-Leibniz bialgebra if for all  $x, y \in A$ ,

$$(R_{[\cdot]}(x) \otimes id) \Delta \alpha(y) = \sigma((R_{[\cdot]}(\alpha(y)) \otimes id) \Delta x), \quad (3.6)$$

$$\begin{aligned} (\alpha \otimes id) \Delta[x, y] &= -(id \otimes R_{[\cdot]}(\alpha(y))) \Delta x - (id \otimes L_{[\cdot]}(\alpha(x))) \Delta y \\ &+ \left( L_{[\cdot]}(y) \otimes id + R_{[\cdot]}(y) \otimes id + \sigma(id \otimes L_{[\cdot]}(y) + id \otimes R_{[\cdot]}(y)) \right) \Delta \alpha(x) \\ &- (L_{[\cdot]}(x) \otimes id + R_{[\cdot]}(x) \otimes id) \Delta \alpha(y), \end{aligned} \quad (3.7)$$

where  $\sigma : A \otimes A \rightarrow A \otimes A$  is the exchange operator defined by  $\sigma(x \otimes y) = y \otimes x$ .

Now we are ready to give the main results of this section.

**Theorem 3.10.** *Let  $(A, [\cdot, \cdot], \alpha)$  be an involutive Hom-Leibniz algebra. Suppose that there is an involutive Hom-Leibniz algebra structure " $\{\cdot, \cdot\}$ " on its dual space  $A^*$ . Then, there is a standard Manin triple of  $(A, [\cdot, \cdot], \alpha)$  and  $(A^*, \{\cdot, \cdot\}, \alpha^*)$  if and only if  $(A, A^*, L_{[\cdot]}^*, -L_{[\cdot]}^* - R_{[\cdot]}^*, \alpha^*, L_{\{\cdot\}}^*, -L_{\{\cdot\}}^* - R_{\{\cdot\}}^*, \alpha)$  is a matched pair of involutive Hom-Leibniz algebras.*

*Proof.* Let us consider the four maps such that for all  $x, v, u \in A, x^*, v^*, u^* \in A^*$ ,

$$\begin{aligned} L_{[\cdot]}^* : A &\rightarrow gl(A^*), & \langle L_{[\cdot]}^*(x)u^*, v \rangle &= -\langle L_{[\cdot]}(x)v, u^* \rangle = -\langle [x, v], u^* \rangle, \\ R_{[\cdot]}^* : A &\rightarrow gl(A^*), & \langle R_{[\cdot]}^*(x)u^*, v \rangle &= -\langle R_{[\cdot]}(x)v, u^* \rangle = -\langle [v, x], u^* \rangle, \\ L_{\{\cdot\}}^* : A^* &\rightarrow gl(A), & \langle L_{\{\cdot\}}^*(x^*)u, v^* \rangle &= -\langle L_{\{\cdot\}}(x^*)v^*, u \rangle = -\langle \{x^*, v^*\}, u \rangle, \\ R_{\{\cdot\}}^* : A^* &\rightarrow gl(A), & \langle R_{\{\cdot\}}^*(x^*)u, v^* \rangle &= -\langle R_{\{\cdot\}}(x^*)v^*, u \rangle = -\langle \{v^*, x^*\}, u \rangle. \end{aligned}$$

If  $(A, A^*, L_{[\cdot]}^*, -L_{[\cdot]}^* - R_{[\cdot]}^*, \alpha^*, L_{\{\cdot\}}^*, -L_{\{\cdot\}}^* - R_{\{\cdot\}}^*, \alpha)$  is a matched pair of multiplicative Hom-Leibniz algebras, then  $(A \bowtie A^*, [\cdot, \cdot]', \alpha + \alpha^*)$  is a multiplicative Hom-Leibniz algebra with the product  $[\cdot, \cdot]'$  given by (2.18), and the bilinear form  $B(\cdot, \cdot)$  defined by (3.5) is  $(\alpha \oplus \alpha^*)$ -invariant, that is

$$\begin{aligned} B([\alpha(x) + \alpha^*(a^*), \alpha(z) + \alpha^*(c^*)]' + [\alpha(z) + \alpha^*(c^*), \alpha(x) + \alpha^*(a^*)]', \alpha(y) + \alpha^*(b^*)) \\ = B((\alpha(x) + \alpha^*(a^*), [\alpha(y) + \alpha^*(b^*), (\alpha(z) + \alpha^*(c^*))])' \end{aligned}$$

for all  $x, y \in A^*, a^*, b^* \in A^*$ , and

$$[(x + a^*), (y + b^*)]' = ([x, y] + l_{A^*}(a^*)y + r_{A^*}(b^*)x) + (\{a^*, b^*\} + l_A(x)b^* + r_A(y)a^*),$$

with  $l_A = L_{[\cdot]}^*, r_A = -L_{[\cdot]}^* - R_{[\cdot]}^*, l_{A^*} = L_{\{\cdot\}}^*, r_{A^*} = -L_{\{\cdot\}}^* - R_{\{\cdot\}}^*$ . Indeed,

$$\begin{aligned} & B(\alpha(x) + \alpha^*(a^*), [\alpha(y) + \alpha^*(b^*), \alpha(z) + \alpha^*(c^*)]') \\ &= B\left(\alpha(x) + \alpha^*(a^*), [\alpha(y), \alpha(z)] + l_{A^*}(\alpha^*(b^*))\alpha(z) + r_{A^*}(\alpha^*(c^*))\alpha(y) \right. \\ &\quad \left. + \{\alpha^*(b^*), \alpha^*(c^*)\} + l_A(\alpha(y))\alpha^*(c^*) + r_A(\alpha(z))\alpha^*(b^*)\right) \\ &= \langle \alpha^*(a^*), [\alpha(y), \alpha(z)] \rangle - \langle \{\alpha^*(b^*), \alpha^*(a^*)\}, \alpha(z) \rangle + \langle \{\alpha^*(c^*), \alpha^*(a^*)\}, \alpha(y) \rangle \\ &\quad + \langle \{\alpha^*(a^*), \alpha^*(c^*)\}, \alpha(y) \rangle - \langle \{\alpha^*(b^*), \alpha^*(c^*)\}, \alpha(x) \rangle + \langle [\alpha(y), \alpha(x)], \alpha^*(c^*) \rangle \\ &\quad - \langle [\alpha(z), \alpha(x)], \alpha^*(b^*) \rangle - \langle [\alpha(x), \alpha(z)], \alpha^*(b^*) \rangle, \end{aligned}$$

$$\begin{aligned} & B([\alpha(x) + \alpha^*(a^*), \alpha(z) + \alpha^*(c^*)]' + [\alpha(z) + \alpha^*(c^*), \alpha(x) + \alpha^*(a^*)]', \alpha(y) + \alpha^*(b^*)) \\ &= B\left([\alpha(x), \alpha(z)] + l_{A^*}(\alpha^*(a^*))\alpha(z) + r_{A^*}(\alpha^*(c^*))\alpha(y) + \{\alpha^*(a^*), \alpha^*(c^*)\} \right. \\ &\quad \left. + l_A(\alpha(x))\alpha^*(c^*) + r_A(\alpha(z))\alpha^*(a^*) + [\alpha(z), \alpha(x)] + l_{A^*}(\alpha^*(c^*))\alpha(x) \right. \\ &\quad \left. + r_{A^*}(\alpha^*(a^*))\alpha(z) + \{\alpha^*(c^*), \alpha^*(a^*)\} + l_A(\alpha(z))\alpha^*(a^*) + r_A(\alpha(y))\alpha^*(b^*)\right) \\ &= -\langle [\alpha(x), \alpha(z)], \alpha^*(b^*) \rangle + \langle \alpha(z), \{\alpha^*(a^*), \alpha^*(b^*)\} \rangle - \langle \alpha(x), \{\alpha^*(c^*), \alpha^*(b^*)\} \rangle \\ &\quad + \langle \alpha(y), \{\alpha^*(a^*), \alpha^*(c^*)\} \rangle + \langle [\alpha(y), \alpha(x)], \alpha^*(c^*) \rangle + \langle [\alpha(z), \alpha(y)], \alpha^*(a^*) \rangle \\ &\quad + \langle [\alpha(y), \alpha(z)], \alpha^*(a^*) \rangle - \langle [\alpha(z), \alpha(x)], \alpha^*(b^*) \rangle + \langle \alpha(x), \{\alpha^*(c^*), \alpha^*(b^*)\} \rangle \\ &\quad - \langle \{\alpha^*(b^*), \alpha^*(c^*)\}, \alpha(x) \rangle - \langle \alpha(z), \{\alpha^*(a^*), \alpha^*(b^*)\} \rangle - \langle \alpha(z), \{\alpha^*(b^*), \alpha^*(a^*)\} \rangle \\ &\quad + \langle \{\alpha^*(c^*), \alpha^*(a^*)\}, \alpha(y) \rangle - \langle [\alpha(z), \alpha(y)], \alpha^*(a^*) \rangle \\ &= \langle \alpha^*(a^*), [\alpha(y), \alpha(z)] \rangle - \langle \{\alpha^*(b^*), \alpha^*(a^*)\}, \alpha(z) \rangle + \langle \{\alpha^*(c^*), \alpha^*(a^*)\}, \alpha(y) \rangle \\ &\quad + \langle \{\alpha^*(a^*), \alpha^*(c^*)\}, \alpha(y) \rangle - \langle \{\alpha^*(b^*), \alpha^*(c^*)\}, \alpha(x) \rangle + \langle [\alpha(y), \alpha(x)], \alpha^*(c^*) \rangle \\ &\quad - \langle [\alpha(z), \alpha(x)], \alpha^*(b^*) \rangle - \langle [\alpha(x), \alpha(z)], \alpha^*(b^*) \rangle. \end{aligned}$$

Thus,  $B$  is well  $(\alpha \oplus \alpha^*)$ -invariant. Conversely, if  $(A \oplus A^*, A, A^*)$  is a Manin triple of Hom-Leibniz algebras. For all  $x \in A, a^*, b^* \in A^*$ ,

$$\begin{aligned} B(b^*, r_2(a^*)x) &= B(b^*, [x, a^*]') \\ &= B(\{b^*, a^*\} + \{a^*, b^*\}, x) \\ &= \langle R_{\{\cdot\}}(a^*)b^* + L_{\{\cdot\}}(a^*)b^*, x \rangle \\ &= -\langle b^*, (L_{\{\cdot\}}^* + R_{\{\cdot\}}^*)(x) \rangle, \end{aligned}$$

which implies that  $r_2 = -L_{\{\cdot\}}^* - R_{\{\cdot\}}^*$ . Now,

$$\begin{aligned} \langle L_1(x)a^*, y \rangle &= -B(y, [x, a^*]') \\ &= -B([y, a^*]' + [a^*, y]', x) \\ &= -B(a^*, [x, y]) \\ &= -\langle a^*, L_{[\cdot]}(x)y \rangle \\ &= \langle y, L_{[\cdot]}^*(x)a^* \rangle, \end{aligned}$$

which implies that  $L_1 = L_{[\cdot]}^*$ . Similarly,  $L_2 = L_{\{\cdot\}}^*$  and  $r_1 = -L_{[\cdot]}^* - R_{[\cdot]}^*$ . Thus,  $(A, A^*, L_{[\cdot]}^*, -L_{[\cdot]}^* - R_{[\cdot]}^*, \alpha^*, L_{\{\cdot\}}^*, -L_{\{\cdot\}}^* - R_{\{\cdot\}}^*, \alpha)$  is a matched pair.  $\square$

**Theorem 3.11.** *Let  $(A, [\cdot, \cdot], \alpha)$  be an involutive Hom-Leibniz algebra. Suppose that there is an involutive Hom-Leibniz algebra structure " $\{\cdot, \cdot\}$ " on its dual space  $(A^*, \alpha^*)$ . Then,  $(A, A^*, L_{[\cdot]}^*, -L_{[\cdot]}^* - R_{[\cdot]}^*, \alpha^*, L_{\{\cdot\}}^*, -L_{\{\cdot\}}^* - R_{\{\cdot\}}^*, \alpha)$  is a matched pair of involutive Hom-Leibniz algebras if and only if, for any  $x \in A$  and  $a^*, b^* \in A^*$ ,*

$$L_{\{\cdot\}}^*(R_{[\cdot]}^*(x)a^*)\alpha(y) + [R_{\{\cdot\}}^*a^*x, \alpha(y)] = 0, \quad (3.8)$$

$$\begin{aligned} & L_{\{\cdot\}}^*(\alpha^*(a^*))([x, y]) - [L_{\{\cdot\}}^*(a^*)x, \alpha(y)] - [\alpha(x), L_{\{\cdot\}}^*a^*y] \\ & + L_{\{\cdot\}}^*(L_{[\cdot]}^*(x)a^*)\alpha(y) + L_{\{\cdot\}}^*(R_{[\cdot]}^*(x)a^*)\alpha(y) - L_{\{\cdot\}}^*(L_{[\cdot]}^*(y)a^*)\alpha(x) \\ & - L_{\{\cdot\}}^*(R_{[\cdot]}^*(y)a^*)\alpha(x) - R_{\{\cdot\}}^*(L_{[\cdot]}^*(x)(a^*)\alpha(x) - R_{\{\cdot\}}^*(R_{[\cdot]}^*(x)a^*)\alpha(x) = 0. \end{aligned} \quad (3.9)$$

*Proof.* Obviously, (3.8) gives (2.16), and (3.9) reduces to (2.15) when

$$l_A = L_{[\cdot]}^*, \quad r_A = -L_{[\cdot]}^* - R_{[\cdot]}^*, \quad l_B = L_{\{\cdot\}}^*, \quad r_B = -L_{\{\cdot\}}^* - R_{\{\cdot\}}^*.$$

Now, we show that

$$(2.13) \iff (2.16);$$

$$\text{and } (2.12) \iff (2.14) \iff (2.15) \iff (2.17).$$

Suppose (3.8) and (3.9) are satisfied. We have for all  $x, y \in A, a^*, b^* \in A^*$ ,

$$\begin{aligned} \langle R_{[\cdot]}^*(x)a^*, y \rangle &= \langle L_{[\cdot]}^*(y)a^*, x \rangle = -\langle [y, x], a^* \rangle, \\ \langle R_{\{\cdot\}}^*(b^*)x, a^* \rangle &= \langle L_{\{\cdot\}}^*(a^*)x, b^* \rangle = -\langle \{a^*, b^*\}, x \rangle, \\ \alpha^*(R_{[\cdot]}^*(x)a^*) &= R_{[\cdot]}^*(\alpha(x))\alpha^*(a^*), \quad \alpha^*(L_{[\cdot]}^*(x)a^*) = L_{[\cdot]}^*(\alpha(x))\alpha^*(a^*), \\ \alpha(R_{\{\cdot\}}^*(a^*)x) &= R_{\{\cdot\}}^*(\alpha^*(a^*))\alpha(x), \quad \alpha(L_{\{\cdot\}}^*(a^*)x) = L_{\{\cdot\}}^*(\alpha^*(a^*))\alpha(x), \end{aligned}$$

Set  $\alpha(x) = z, \alpha(y) = t, \alpha^*(a^*) = c^*$  and  $\alpha^*(b^*) = d^*$ . Then (2.13)  $\iff$  (2.16) follows from

$$\begin{aligned} \langle L_{\{\cdot\}}^*(R_{[\cdot]}^*(x)a^*)\alpha(y), b^* \rangle &= -\langle \{(R_{[\cdot]}^*(x)a^*), b^*\}, \alpha(y) \rangle \\ &= \langle \alpha^*\{(R_{[\cdot]}^*(x)a^*), b^*\}, y \rangle \\ &= \langle \{R_{[\cdot]}^*(\alpha(x))\alpha^*(a^*), \alpha^*(b^*)\}, y \rangle \\ &= \langle \{R_{[\cdot]}^*(z)c^*, \alpha^*(b^*)\}, y \rangle; \\ \langle [R_{\{\cdot\}}^*(a^*)x, \alpha(y)], b^* \rangle &= -\langle \alpha(y), L_{[\cdot]}^*(R_{\{\cdot\}}^*(a^*)x)b^* \rangle \\ &= \langle y, \alpha^*(L_{[\cdot]}^*(R_{\{\cdot\}}^*(a^*)\alpha(x))b^*) \rangle \\ &= \langle y, L_{[\cdot]}^*(R_{\{\cdot\}}^*(c^*)z)\alpha^*(b^*) \rangle. \end{aligned}$$

The statement (2.14)  $\iff$  (2.17) follows from

$$\begin{aligned} \langle \{\alpha^*(a^*), (L_{[\cdot]}^*(x)b^*)\}, y \rangle &= -\langle \alpha^*(a^*), R_{\{\cdot\}}^*L_{[\cdot]}^*(x)b^*y \rangle \\ &= \langle a^*, \alpha[R_{\{\cdot\}}^*(L_{[\cdot]}^*(x)b^*)y] \rangle \\ &= \langle a^*, R_{\{\cdot\}}^*[\alpha^*(L_{[\cdot]}^*(x)b^*)\alpha(y)] \rangle \\ &= \langle a^*, R_{\{\cdot\}}^*(L_{[\cdot]}^*(\alpha(x))\alpha^*(b^*))\alpha(y) \rangle \\ &= \langle a^*, R_{\{\cdot\}}^*(L_{[\cdot]}^*(z)d^*)\alpha(y) \rangle; \\ \langle \{\alpha^*(a^*), (R_{[\cdot]}^*(x)b^*)\}, y \rangle &= -\langle \alpha^*(a^*), R_{\{\cdot\}}^*(R_{[\cdot]}^*(x)b^*)y \rangle \\ &= \langle a^*, \alpha[R_{\{\cdot\}}^*(R_{[\cdot]}^*(\alpha(x))\alpha^*(b^*))\alpha(y)] \rangle \\ &= \langle a^*, R_{\{\cdot\}}^*(R_{[\cdot]}^*(\alpha(x)\alpha^*(b^*))\alpha(y) \rangle \\ &= \langle a^*, R_{\{\cdot\}}^*(R_{[\cdot]}^*(z)d^*)\alpha(y) \rangle; \\ \langle L_{[\cdot]}^*(L_{\{\cdot\}}^*(b^*)x)\alpha^*(a^*), y \rangle &= -\langle [(L_{\{\cdot\}}^*(b^*)x), y], \alpha^*(a^*) \rangle \end{aligned}$$

$$\begin{aligned}
&= \langle a^*, [\alpha(L_{\{\cdot\}}^*(b^*)x), \alpha(y)] \rangle \\
&= \langle a^*, L_{\{\cdot\}}^*([\alpha^*(b^*)\alpha(x), \alpha(y)]) \rangle \\
&= \langle a^*, [L_{\{\cdot\}}^*(d^*)z, \alpha(y)] \rangle; \\
\langle R_{[\cdot]}^*(L_{\{\cdot\}}^*(a^*)x)\alpha^*(b^*), y \rangle &= -\langle [y, L_{\{\cdot\}}^*(a^*)x], \alpha^*(b^*) \rangle \\
&= \langle [\alpha(y), \alpha(L_{\{\cdot\}}^*(a^*)x)], b^* \rangle \\
&= \langle [\alpha(y), L_{\{\cdot\}}^*(\alpha^*(a^*))\alpha(x)], b^* \rangle \\
&= \langle [\alpha(y), L_{\{\cdot\}}^*(c^*)z], b^* \rangle; \\
\langle \{L^*(\alpha(x))(a^*, b^*)\}, y \rangle &= -\langle \{a^*, b^*\}, [\alpha(x), y] \rangle \\
&= \langle R_{\{\cdot\}}^*(b^*)([\alpha(x), y]), a^* \rangle \\
&= \langle R_{\{\cdot\}}^*(\alpha^*(d^*))([z, y]), a^* \rangle; \\
\langle \{R^*(\alpha(x))(a^*, b^*)\}, y \rangle &= -\langle \{a^*, b^*\}, [y, \alpha(x)] \rangle \\
&= -\langle L_{[\cdot]}^*(a^*)b^*, [y, z] \rangle \\
&= \langle b^*, L_{\{\cdot\}}^*(a^*)([y, z]) \rangle \\
&= \langle b^*, L_{\{\cdot\}}^*(\alpha^*(c^*))([y, z]) \rangle; \\
\langle \{\alpha^*(b^*), (L_{[\cdot]}^*(x)a^*)\}, y \rangle &= -\langle \alpha^*(b^*), R_{\{\cdot\}}^*(L_{[\cdot]}^*(x)a^*)y \rangle \\
&= \langle b^*, \alpha[R_{\{\cdot\}}^*(L_{[\cdot]}^*(x)a^*)y] \rangle \\
&= \langle b^*, R_{\{\cdot\}}^*[\alpha^*(L_{[\cdot]}^*(x)a^*)\alpha(y)] \rangle \\
&= \langle b^*, R_{\{\cdot\}}^*[L_{[\cdot]}^*(\alpha(x))\alpha^*(a^*)\alpha(y)] \rangle \\
&= \langle b^*, R_{\{\cdot\}}^*(L_{[\cdot]}^*(z)d^*)\alpha(y) \rangle; \\
\langle \{\alpha^*(b^*), (R_{[\cdot]}^*(x)a^*)\}, y \rangle &= -\langle R_{\{\cdot\}}^*(R_{\{\cdot\}}^*(x)a^*)y, \alpha^*(b^*) \rangle \\
&= -\langle \{\alpha^*(b^*), (R_{[\cdot]}^*(x)a^*)\}, y \rangle \\
&= \langle \alpha[R_{\{\cdot\}}^*(R_{\{\cdot\}}^*(x)a^*)y], b^* \rangle \\
&= \langle R_{\{\cdot\}}^*[R_{\{\cdot\}}^*(\alpha(x))\alpha^*(a^*)\alpha(y)], b^* \rangle \\
&= \langle R_{\{\cdot\}}^*(R_{[\cdot]}^*(z)c^*)\alpha(y), b^* \rangle; \\
\langle L_{[\cdot]}^*(L_{\{\cdot\}}^*(a^*)x)\alpha^*(b^*), y \rangle &= -\langle [(L_{\{\cdot\}}^*(a^*)x), y], \alpha^*(b^*) \rangle \\
&= \langle b^*, [\alpha(L_{\{\cdot\}}^*(a^*)x), \alpha(y)] \rangle \\
&= \langle b^*, [L_{\{\cdot\}}^*(\alpha^*(a^*))\alpha(x), \alpha(y)] \rangle \\
&= \langle b^*, [L_{\{\cdot\}}^*(c^*)z, \alpha(y)] \rangle; \\
\langle R_{[\cdot]}^*(L_{\{\cdot\}}^*(b^*)x)\alpha(a^*), y \rangle &= -\langle [y, L_{\{\cdot\}}^*(b^*)x], \alpha^*(a^*) \rangle \\
&= \langle L_{[\cdot]}^*(y)(\alpha^*(a^*)), L_{\{\cdot\}}^*(b^*)x \rangle \\
&= \langle L_{\{\cdot\}}^*(b^*)x, L_{[\cdot]}^*(y)(\alpha^*(a^*)) \rangle \\
&= -\langle \{b^*, (L_{[\cdot]}^*(y)(\alpha^*(a^*)))\}, x \rangle \\
&= \langle \{\alpha^*(c^*), (L_{[\cdot]}^*(y)(d^*))\}, x \rangle.
\end{aligned}$$

The statements (2.12)  $\iff$  (2.15) and (2.15)  $\iff$  (2.17) are proved analogously.  $\square$

**Theorem 3.12.** *Let  $(A, [\cdot, \cdot], \alpha)$  be an involutive Hom-Leibniz algebra. Suppose that there is an involutive Hom-Leibniz algebra structure  $\{\cdot, \cdot\}$  on its dual space  $(A^*, \alpha^*)$ . Then,  $(A, A^*)$  is a Hom-Leibniz bialgebra if and only if  $(A, A^*, L_{[\cdot]}^*, -L_{[\cdot]}^* - R_{[\cdot]}^*, \alpha^*, L_{\{\cdot\}}^*, -L_{\{\cdot\}}^* - R_{\{\cdot\}}^*, \alpha)$  is a matched pair of involutive Hom-Leibniz algebras.*

*Proof.* Let  $x, y \in A$  and  $a^*, b^* \in A^*$ . First, the left hand side of (3.8), is equal to

$$L_{\{\cdot\}}^*(R_{[\cdot]}^*(x)a^*)\alpha(y) + [R_{\{\cdot\}}^*(a^*)x, \alpha(y)].$$

Furthermore, we have

$$\begin{aligned}
&\langle L_{\{\cdot\}}^*(R_{[\cdot]}^*(x)a^*)\alpha(y) + [R_{\{\cdot\}}^*(a^*)x, \alpha(y)], b^* \rangle \\
&= \langle L_{\{\cdot\}}^*(R_{[\cdot]}^*(x)a^*)\alpha(y), b^* \rangle + \langle [R_{\{\cdot\}}^*(a^*)x, \alpha(y)], b^* \rangle
\end{aligned}$$

$$\begin{aligned}
&= \langle L_{\{\cdot\}}^*(R_{[\cdot]}^*(x)(a^*)\alpha(y), b^*) + \langle R_{[\cdot]}(\alpha(y))(R_{\{\cdot\}}^*(a^*)x), b^* \rangle \\
&= -\langle \alpha(y), \{R_{[\cdot]}^*(x)a^*, b^*\} \rangle + \langle x, \{R_{[\cdot]}^*(\alpha(y))b^*, a^*\} \rangle \\
&= -\langle \Delta\alpha(y), R_{[\cdot]}^*(x)a^* \otimes b^* \rangle + \langle \Delta x, R_{[\cdot]}^*(\alpha(y))b^* \otimes a^* \rangle \\
&= \langle (R_{[\cdot]}(x) \otimes id)\Delta\alpha(y), a^* \otimes b^* \rangle - \langle (R_{[\cdot]}(\alpha(y)) \otimes id)\Delta x, b^* \otimes a^* \rangle \\
&= \langle (R_{[\cdot]}(x) \otimes id)\Delta\alpha(y) - \sigma((R_{[\cdot]}(\alpha(y)) \otimes id)\Delta x), a^* \otimes b^* \rangle.
\end{aligned}$$

Thus, we conclude that (3.8) corresponds to (3.6).

Next, the left hand side of (3.9) is equal to

$$\begin{aligned}
&L_{\{\cdot\}}^*(\alpha^*(a^*))([x, y]) - [L_{\{\cdot\}}^*(a^*)x, \alpha(y)] - [\alpha(x), L_{\{\cdot\}}^*a^*y] \\
&+ L_{\{\cdot\}}^*(L_{[\cdot]}^*(x)a^*)\alpha(y) + L_{\{\cdot\}}^*(R_{[\cdot]}^*(x)a^*)\alpha(y) - L_{\{\cdot\}}^*(L_{[\cdot]}^*(y)a^*)\alpha(x) \\
&- L_{\{\cdot\}}^*(R_{[\cdot]}^*(y)a^*)\alpha(x) - R_{\{\cdot\}}^*(L_{[\cdot]}^*(x)a^*)\alpha(x) - R_{\{\cdot\}}^*(R_{[\cdot]}^*(x)a^*)\alpha(x).
\end{aligned}$$

Furthermore, we have

$$\begin{aligned}
&\langle L_{\{\cdot\}}^*(\alpha^*(a^*))([x, y]) - [L_{\{\cdot\}}^*(a^*)x, \alpha(y)] - [\alpha(x), L_{\{\cdot\}}^*a^*y] \\
&+ L_{\{\cdot\}}^*(L_{[\cdot]}^*(x)a^*)\alpha(y) + L_{\{\cdot\}}^*(R_{[\cdot]}^*(x)a^*)\alpha(y) - L_{\{\cdot\}}^*(L_{[\cdot]}^*(y)a^*)\alpha(x) \\
&- L_{\{\cdot\}}^*(R_{[\cdot]}^*(y)a^*)\alpha(x) - R_{\{\cdot\}}^*(L_{[\cdot]}^*(x)a^*)\alpha(x) - R_{\{\cdot\}}^*(R_{[\cdot]}^*(x)a^*)\alpha(x), b^* \rangle \\
&= \langle L_{\{\cdot\}}^*(\alpha^*(a^*))([x, y]), b^* \rangle - \langle [L_{\{\cdot\}}^*(a^*)x, \alpha(y)], b^* \rangle - \langle [\alpha(x), L_{\{\cdot\}}^*a^*y], b^* \rangle \\
&+ \langle L_{\{\cdot\}}^*(L_{[\cdot]}^*(x)a^*)\alpha(y), b^* \rangle + \langle L_{\{\cdot\}}^*(R_{[\cdot]}^*(x)a^*)\alpha(y), b^* \rangle - \langle L_{\{\cdot\}}^*(L_{[\cdot]}^*(y)a^*)\alpha(x), b^* \rangle \\
&- \langle L_{\{\cdot\}}^*(R_{[\cdot]}^*(y)a^*)\alpha(x), b^* \rangle - \langle R_{\{\cdot\}}^*(L_{[\cdot]}^*(x)a^*)\alpha(x), b^* \rangle - \langle R_{\{\cdot\}}^*(R_{[\cdot]}^*(x)a^*)\alpha(x), b^* \rangle \\
&= -\langle [x, y], \{\alpha^*(a^*), b^*\} \rangle - \langle x, \{a^*, R_{\{\cdot\}}^*(\alpha(y))b^*\} \rangle - \langle y, \{a^*, L_{[\cdot]}^*(\alpha(x))b^*\} \rangle \\
&- \langle \alpha(y), \{L_{[\cdot]}^*(x)a^*, b^*\} \rangle - \langle \alpha(y), \{R_{[\cdot]}^*(x)a^*, b^*\} \rangle + \langle \alpha(x), \{L_{[\cdot]}^*(y)a^*, b^*\} \rangle \\
&+ \langle \alpha(x), \{R_{[\cdot]}^*(y)a^*, b^*\} \rangle + \langle \alpha(x), \{a^*, L^*(y)a^*\} \rangle + \langle \alpha(x), \{b^*, R_{[\cdot]}^*(y)a^*\} \rangle \\
&= -\langle \Delta[x, y], \alpha^*(a^*) \otimes b^* \rangle - \langle \Delta x, a^* \otimes R_{[\cdot]}^*(\alpha(y))b^* \rangle - \langle \Delta y, a^* \otimes L_{[\cdot]}^*(\alpha(x))b^* \rangle \\
&- \langle \Delta\alpha y, L_{[\cdot]}^*(x)a^* \otimes b^* \rangle - \langle \Delta\alpha(y), R_{[\cdot]}^*(x)a^* \otimes b^* \rangle + \langle \Delta\alpha(x), L_{[\cdot]}^*(y)a^* \otimes b^* \rangle \\
&+ \langle \Delta\alpha(x), R_{[\cdot]}^*(y)a^* \otimes b^* \rangle + \langle \Delta\alpha(x), a^* \otimes L_{[\cdot]}^*(y)a^* \rangle + \langle \Delta\alpha(x), b^* \otimes R_{[\cdot]}^*(y)a^* \rangle \\
&= \langle (\alpha \otimes id)\Delta[x, y], a^* \otimes b^* \rangle + \langle (id \otimes R_{[\cdot]}(\alpha(y)))\Delta x, a^* \otimes b^* \rangle \\
&+ \langle (id \otimes L_{[\cdot]}(\alpha(x)))\Delta y, a^* \otimes b^* \rangle + \langle (L_{[\cdot]}(x) \otimes id)\Delta\alpha(y), a^* \otimes b^* \rangle \\
&+ \langle (R_{[\cdot]}(x) \otimes id)\Delta\alpha(y), a^* \otimes b^* \rangle - \langle (L_{[\cdot]}(y) \otimes id)\Delta\alpha(x), a^* \otimes b^* \rangle \\
&- \langle (R_{[\cdot]}(y) \otimes id)\Delta\alpha(x), a^* \otimes b^* \rangle - \langle (id \otimes L_{[\cdot]}(y))\Delta\alpha(x), b^* \otimes a^* \rangle \\
&- \langle (id \otimes R_{[\cdot]}(y))\Delta\alpha(x), b^* \otimes a^* \rangle \\
&= \langle (\alpha \otimes id)\Delta[x, y], a^* \otimes b^* \rangle + \langle (id \otimes R_{[\cdot]}(\alpha(y)))\Delta x, a^* \otimes b^* \rangle \\
&+ \langle (id \otimes L_{[\cdot]}(\alpha(x)))\Delta y, a^* \otimes b^* \rangle + \langle (L_{[\cdot]}(x) \otimes id)\Delta\alpha(y), a^* \otimes b^* \rangle \\
&+ \langle (R_{[\cdot]}(x) \otimes id)\Delta\alpha(y), a^* \otimes b^* \rangle - \langle (L_{[\cdot]}(y) \otimes id)\Delta\alpha(x), a^* \otimes b^* \rangle \\
&- \langle (R_{[\cdot]}(y) \otimes id)\Delta\alpha(x), a^* \otimes b^* \rangle - \langle \sigma((id \otimes L_{[\cdot]}(y))\Delta\alpha(x)), a^* \otimes b^* \rangle \\
&- \langle \sigma((id \otimes R_{[\cdot]}(y))\Delta\alpha(x)), a^* \otimes b^* \rangle \\
&= \langle (\alpha \otimes id)\Delta[x, y] + (id \otimes R_{[\cdot]}(\alpha(y)))\Delta x + (id \otimes L_{[\cdot]}(\alpha(x)))\Delta y \\
&+ (L_{[\cdot]}(x) \otimes id)\Delta\alpha(y) + (R_{[\cdot]}(x) \otimes id)\Delta\alpha(y)
\end{aligned}$$

$$\begin{aligned}
& - (L_{[\cdot]}(y) \otimes id)\Delta\alpha(x) - (R_{[\cdot]}(y) \otimes id)\Delta\alpha(x) \\
& - \sigma((id \otimes L_{[\cdot]}(y))\Delta\alpha(x)) - \sigma((id \otimes R_{[\cdot]}(y))\Delta\alpha(x)), a^* \otimes b^*.
\end{aligned}$$

Thus, we conclude that (3.9) corresponds to (3.7). This completes the proof.  $\square$

## 4 Hom-Leibniz dendriform algebras

**Definition 4.1.** Hom-Leibniz dendriform algebras are quadruples  $(A, \prec, \succ, \alpha)$  consisting of a linear space  $A$ , bilinear maps  $\prec, \succ : A \times A \rightarrow A$ , and linear maps  $\alpha : A \rightarrow A$  satisfying

$$([x, y]) \succ \alpha(z) = \alpha(x) \succ (y \succ z) - \alpha(y) \succ (x \succ z), \quad (4.1)$$

$$\alpha(x) \succ (y \prec z) = (x \succ y) \prec \alpha(z) + \alpha(y) \prec ([x, z]), \quad (4.2)$$

$$\alpha(x) \prec ([y, z]) = (x \prec y) \prec \alpha(z) + \alpha(y) \succ (x \prec z), \quad (4.3)$$

where

$$[x, y] = x \prec y + x \succ y. \quad (4.4)$$

A Hom-Leibniz dendriform algebra is called multiplicative if the linear map  $\alpha : A \rightarrow A$  is multiplicative with respect to  $\prec$  and  $\succ$ , that is, for all  $x, y \in A$ ,

$$\alpha(x \prec y) = \alpha(x) \prec \alpha(y), \quad \alpha(x \succ y) = \alpha(x) \succ \alpha(y).$$

**Definition 4.2.** Let  $(A, \prec, \succ, \alpha)$  and  $(A', \prec', \succ', \alpha')$  be two Hom-Leibniz dendriform algebras. A linear map  $f : A \rightarrow A'$  is a Hom-Leibniz dendriform algebra morphism if

$$\prec' \circ (f \otimes f) = f \circ \prec, \quad \succ' \circ (f \otimes f) = f \circ \succ \quad \text{and} \quad f \circ \alpha = \alpha' \circ f.$$

**Proposition 4.3.** Let  $(A, \prec, \succ, \alpha)$  be a Hom-Leibniz dendriform algebra. Then the binary operation  $[\cdot, \cdot] : A \otimes A \rightarrow A$ , given for all  $x, y \in A$  by  $[x, y] = x \prec y + x \succ y$ , defines a Hom-Leibniz algebra, which is called the sub-adjacent Hom-Leibniz algebra of  $(A, \prec, \succ, \alpha)$ , and  $(A, \prec, \succ, \alpha)$  is called a compatible Hom-Leibniz dendriform algebra structure on  $(A, [\cdot, \cdot], \alpha)$ .

*Proof.* For all  $x, y, z \in A$ ,

$$\begin{aligned}
& [\alpha(x), [y, z]] - [[x, y], \alpha(z)] - [\alpha(y), [x, z]] \\
& = [\alpha(x), y \prec z + y \succ z] - [x \prec y + x \succ y, \alpha(z)] - [\alpha(y), x \prec z + x \succ z] \\
& = \alpha(x) \prec (y \prec z + y \succ z) + \alpha(x) \succ (y \prec z + y \succ z) \\
& \quad - (x \prec y + x \succ y) \prec \alpha(z) - (x \prec y + x \succ y) \succ \alpha(z) \\
& \quad - \alpha(y) \prec (x \prec z + x \succ z) - \alpha(y) \succ (x \prec z + x \succ z) \\
& = \underbrace{\left( - [x, y] \succ \alpha(z) + \alpha(x) \succ (y \succ z) - \alpha(y) \succ (x \succ z) \right)}_{= 0 \text{ by (4.1)}} \\
& \quad + \underbrace{\left( \alpha(x) \succ (y \prec z) - (x \succ y) \prec \alpha(z) - \alpha(y) \succ ([x, z]) \right)}_{= 0 \text{ by (4.2)}} \\
& \quad + \underbrace{\left( \alpha(x) \prec ([y, z]) - (x \prec y) \prec \alpha(z) - \alpha(y) \succ (x \prec z) \right)}_{= 0 \text{ by (4.3)}} = 0. \quad \square
\end{aligned}$$

*Example 4.4.* Let  $V$  be a linear space. On the linear space  $\mathfrak{gl}(V) \oplus V$ , define two binary operations  $\prec, \succ: (\mathfrak{gl}(V) \oplus V) \otimes (\mathfrak{gl}(V) \oplus V) \rightarrow \mathfrak{gl}(V) \oplus V$  and a linear map  $\delta_\beta: \mathfrak{gl}(V) \oplus V \rightarrow \mathfrak{gl}(V) \oplus V$  by

$$\begin{aligned} (A + u) \prec (B + v) &= AB + Av, \\ (A + u) \succ (B + v) &= -BA, \\ \delta_\beta(A + u) &= Ad_\beta(A) + \beta(u), \quad \forall A, B \in \mathfrak{gl}(V), u, v \in V, \end{aligned}$$

with the adjoint action  $Ad_\beta(A) = \beta A \beta^{-1}$  for all  $\beta \in \mathfrak{gl}(V)$ . Then  $(\mathfrak{gl}(V) \oplus V, \prec, \succ, \delta_\beta)$  is a Hom-Leibniz dendriform algebra. Its sub-adjacent Hom-Leibniz algebra is exactly the one underlying a Hom-omni-Lie algebra [5, 74].

**Theorem 4.5.** *Let  $\mathcal{A} = (A, \prec, \succ)$  be a Leibniz dendriform algebra and  $\alpha: A \rightarrow A$  be a Leibniz dendriform algebras morphism of  $\mathcal{A}$ . With bilinear maps  $\prec_\alpha: A \times A \rightarrow A$  and  $\succ_\alpha: A \times A \rightarrow A$  defined for all  $x, y \in A$  by  $x \prec_\alpha y = \alpha(x \prec y)$  and  $x \succ_\alpha y = \alpha(x \succ y)$ , the quadruple  $\mathcal{A}_\alpha = (A_\alpha = A, \prec_\alpha, \succ_\alpha, \alpha)$  is a Hom-Leibniz dendriform algebra, called  $\alpha$ -twist (Yau twist) of  $\mathcal{A}$ . Moreover, assume that  $\mathcal{A}' = (A', \prec', \succ')$  is another Leibniz dendriform algebra, and  $\alpha': A' \rightarrow A'$  is a Leibniz dendriform algebras morphism. Let  $f: \mathcal{A} \rightarrow \mathcal{A}'$  be a Leibniz dendriform algebras morphism satisfying  $f \circ \alpha = \alpha' \circ f$ . Then,  $f: \mathcal{A}_\alpha \rightarrow \mathcal{A}'_{\alpha'}$  is a Hom-Leibniz dendriform algebras morphism.*

*Proof.* Being a Leibniz dendriform algebras morphism,  $\alpha: A \rightarrow A$  is a linear map which is multiplicative with respect to  $\prec$  and  $\succ$ , that is,

$$\forall x, y \in A: \quad \alpha(x \prec y) = \alpha(x) \prec \alpha(y), \quad \alpha(x \succ y) = \alpha(x) \succ \alpha(y).$$

The equalities (4.1), (4.2) and (4.3) in  $\mathcal{A}_\alpha$  are proved as follows:

$$\begin{aligned} ([x, y]_\alpha) \succ_\alpha \alpha(z) &= (x \prec_\alpha y + x \succ_\alpha y) \succ_\alpha \alpha(z) \\ &= (\alpha(x \prec y) + \alpha(x \succ y)) \succ_\alpha \alpha(z) \\ &= \alpha((\alpha(x \prec y) + \alpha(x \succ y)) \succ \alpha(z)) \\ \text{(multiplicativity)} &= (\alpha^2(x \prec y) + \alpha^2(x \succ y)) \succ \alpha^2(z) \\ \text{(multiplicativity)} &= (\alpha^2(x) \prec \alpha^2(y) + \alpha^2(x) \succ \alpha^2(y)) \succ \alpha^2(z) \\ &= [\alpha^2(x), \alpha^2(y)] \succ \alpha^2(z) \\ \text{(\mathcal{A} is Leibniz dendriform)} &= \alpha^2(x) \succ (\alpha^2(y) \succ \alpha^2(z)) - \alpha^2(y) \succ (\alpha^2(x) \succ \alpha^2(z)) \\ \text{(multiplicativity)} &= \alpha^2(x) \succ \alpha(\alpha(y) \succ \alpha(z)) - \alpha^2(y) \succ \alpha(\alpha(x) \succ \alpha(z)) \\ &= \alpha(x) \succ_\alpha (\alpha(y) \succ \alpha(z)) - \alpha(y) \succ_\alpha (\alpha(x) \succ \alpha(z)) \\ &= \alpha(x) \succ_\alpha (y \succ_\alpha z) - \alpha(y) \succ_\alpha (x \succ_\alpha z), \\ \alpha(x) \succ_\alpha (y \prec_\alpha z) &= \alpha(x \succ \alpha(y \prec z)) \\ \text{(multiplicativity)} &= \alpha^2(x \succ (y \prec z)) \\ &= \alpha^2((x \succ y) \prec z) + \alpha^2(x \succ (y \prec z)) - \alpha^2((x \succ y) \prec z) \\ &= \alpha^2((x \succ y) \prec z) + \alpha^2(x \succ (y \prec z)) - (x \succ y) \prec z \\ \text{(\mathcal{A} is Leibniz dendriform)} &= \alpha^2((x \succ y) \prec z) + \alpha^2(y \prec [x, z]) \\ \text{(multiplicativity)} &= \alpha(\alpha(x \succ y) \prec \alpha(z)) + \alpha(\alpha(y) \prec \alpha([x, z])) \end{aligned}$$



$$\begin{aligned}
&= \alpha(\alpha(x \succ y) \prec \alpha(z)) + \alpha(\alpha(y) \prec \alpha(x \prec z + x \succ z)) \\
&= \alpha(\alpha(x \succ y) \prec \alpha(z)) + \alpha(\alpha(y) \prec (\alpha(x \prec z) + \alpha(x \succ z))) \\
&= \alpha(\alpha(x \succ y) \prec \alpha(z)) + \alpha(\alpha(y) \prec (x \prec_{\alpha} z + x \succ_{\alpha} z)) \\
&= (x \succ_{\alpha} y) \prec_{\alpha} \alpha(z) + \alpha(y) \prec_{\alpha} ([x, z]_{\alpha}),
\end{aligned}$$

$$\begin{aligned}
\alpha(x) \prec_{\alpha} ([y, z]_{\alpha}) &= \alpha(x) \prec_{\alpha} (y \prec_{\alpha} z + y \succ_{\alpha} z) \\
&\text{(multiplicativity)} = \alpha^2(x \prec (y \prec z + y \succ z)) \\
&\text{(multiplicativity)} = \alpha^2(x \prec [y, z]) \\
\text{(\mathcal{A} is Leibniz dendriform)} &= \alpha^2((x \prec y) \prec z + y \succ (x \prec z)) \\
&= \alpha^2((x \prec y) \prec z) + \alpha^2(y \succ (x \prec z)) \\
&\text{(multiplicativity)} = \alpha(\alpha(x \prec y) \prec \alpha(z)) + \alpha(\alpha(y) \succ \alpha(x \prec z)) \\
&= (x \prec_{\alpha} y) \prec_{\alpha} \alpha(z) + \alpha(y) \succ_{\alpha} (x \prec_{\alpha} z).
\end{aligned}$$

The second assertion follows from

$$\begin{aligned}
f(x \prec_{\alpha} y) &= f(\alpha(x \prec y)) = \alpha'(f(x \prec' y)) = \alpha'(f(x) \prec' f(y)) = f(x) \prec'_{\alpha'} f(y), \\
f(x \succ_{\alpha} y) &= f(\alpha(x \succ y)) = \alpha'(f(x \succ' y)) = \alpha'(f(x) \succ' f(y)) = f(x) \succ'_{\alpha'} f(y). \quad \square
\end{aligned}$$

**Proposition 4.6.** *Let  $\mathcal{A} = (A, \prec, \succ, \alpha)$  be a Hom-Leibniz dendriform algebra and  $\alpha' : A \rightarrow A$  be a Hom-Leibniz dendriform algebras morphism of  $\mathcal{A}$ . With bilinear maps  $\prec_{\alpha'} : A \times A \rightarrow A$  and  $\succ_{\alpha'} : A \times A \rightarrow A$  defined for all  $x, y \in A$  by  $x \prec_{\alpha'} y = \alpha'(x \prec y)$  and  $x \succ_{\alpha'} y = \alpha'(x \succ y)$ , the quadruple  $\mathcal{A}_{\alpha'} = (A, \prec_{\alpha'}, \succ_{\alpha'}, \alpha \circ \alpha')$  is a Hom-Leibniz dendriform algebra.*

*Proof.* Being a Hom-Leibniz dendriform algebras morphism,  $\alpha' : A \rightarrow A$  is a linear map which is multiplicative with respect to  $\prec$  and  $\succ$ . Then (4.1), (4.2) and (4.3) in  $\mathcal{A}_{\alpha'}$  are proved for all  $x, y, z \in A$  as follows:

$$\begin{aligned}
([x, y]_{\alpha'}) \succ_{\alpha'} \alpha \alpha'(z) &= (x \prec_{\alpha'} y + x \succ_{\alpha'} y) \succ_{\alpha'} \alpha \alpha'(z) \\
&= (\alpha'(x \prec y) + \alpha'(x \succ y)) \succ_{\alpha'} \alpha \alpha'(z) \\
&= \alpha'((\alpha'(x \prec y) + \alpha'(x \succ y)) \succ \alpha \alpha'(z)) \\
&\text{(\alpha' morphism)} = (\alpha'^2(x \prec y) + \alpha'^2(x \succ y)) \succ \alpha' \alpha \alpha'(z) \\
&\text{(\alpha' morphism)} = (\alpha'^2(x) \prec \alpha'^2(y) + \alpha'^2(x) \succ \alpha'^2(y)) \succ \alpha' \alpha \alpha'(z) \\
&\text{(\alpha, \alpha' commute)} = (\alpha'^2(x) \prec \alpha'^2(y) + \alpha'^2(x) \succ \alpha'^2(y)) \succ \alpha \alpha'^2(z) \\
&= [\alpha'^2(x), \alpha'^2(y)] \succ \alpha \alpha'^2(z) \\
&\text{(\mathcal{A} Hom-Leib. dendr.)} = \alpha \alpha'^2(x) \succ (\alpha'^2(y) \succ \alpha'^2(z)) - \alpha \alpha'^2(y) \succ (\alpha'^2(x) \succ \alpha'^2(z)) \\
&\text{(\alpha' morphism)} = \alpha \alpha'^2(x) \succ \alpha'(\alpha'(y) \succ \alpha'(z)) - \alpha \alpha'^2(y) \succ \alpha'(\alpha'(x) \succ \alpha'(z)) \\
&\text{(\alpha, \alpha' commute)} = \alpha' \alpha \alpha'(x) \succ \alpha'(\alpha'(y) \succ \alpha'(z)) - \alpha' \alpha \alpha'(y) \succ \alpha'(\alpha'(x) \succ \alpha'(z)) \\
&= \alpha \alpha'(x) \succ_{\alpha'} (\alpha'(y) \succ \alpha'(z)) - \alpha \alpha'(y) \succ_{\alpha'} (\alpha'(x) \succ \alpha'(z)) \\
&= \alpha \alpha'(x) \succ_{\alpha'} (y \succ_{\alpha'} z) - \alpha \alpha'(y) \succ_{\alpha'} (x \succ_{\alpha'} z), \\
\alpha \alpha'(x) \succ_{\alpha'} (y \prec_{\alpha'} z) &= \alpha \alpha'(x) \succ_{\alpha'} (\alpha'(y \prec z)) \\
&= \alpha'(\alpha \alpha'(x) \succ \alpha'(y \prec z))
\end{aligned}$$

$$\begin{aligned}
& (\alpha' \text{ morphism}) = \alpha' \alpha \alpha' (x) \succ \alpha'^2 (y \prec z) \\
& (\alpha' \text{ morphism}) = \alpha' \alpha \alpha' (x) \succ (\alpha'^2 (y) \prec \alpha'^2 (z)) \\
& (\alpha, \alpha' \text{ commute}) = \alpha \alpha'^2 (x) \succ (\alpha'^2 (y) \prec \alpha'^2 (z)) \\
& (\mathcal{A} \text{ Hom-Leib. dendr.}) = (\alpha'^2 (x) \succ \alpha'^2 (y)) \prec \alpha \alpha'^2 (z) + \alpha \alpha'^2 (y) \prec ([\alpha'^2 (x), \alpha'^2 (z)]) \\
& \quad = (\alpha'^2 (x) \succ \alpha'^2 (y)) \prec \alpha \alpha'^2 (z) \\
& \quad \quad + \alpha \alpha'^2 (y) \prec (\alpha'^2 (x) \prec \alpha'^2 (z) + \alpha'^2 (x) \succ \alpha'^2 (z)) \\
& (\alpha, \alpha' \text{ commute}) = (\alpha'^2 (x) \succ \alpha'^2 (y)) \prec \alpha' \alpha \alpha' (z) \\
& \quad + \alpha' \alpha \alpha' (y) \prec (\alpha'^2 (x) \prec \alpha'^2 (z) + \alpha'^2 (x) \succ \alpha'^2 (z)) \\
& (\alpha' \text{ morphism}) = \alpha' (\alpha' (x) \succ \alpha' (y)) \prec \alpha' \alpha \alpha' (z) \\
& \quad + \alpha' \alpha \alpha' (y) \prec \alpha' (\alpha' (x) \prec \alpha' (z) + \alpha' (x) \succ \alpha' (z)) \\
& = (x \succ_{\alpha'} y) \prec_{\alpha'} \alpha \alpha' (z) + \alpha \alpha' (y) \prec_{\alpha'} (x \prec_{\alpha'} z + x \succ_{\alpha'} z) \\
& = (x \succ_{\alpha'} y) \prec_{\alpha'} \alpha \alpha' (z) + \alpha \alpha' (y) \prec_{\alpha'} ([x, z]_{\alpha'}), \\
\alpha \alpha' (x) \prec_{\alpha'} ([y, z]_{\alpha'}) & = \alpha \alpha' (x) \prec_{\alpha'} (y \prec_{\alpha'} z + y \succ_{\alpha'} z) \\
& = \alpha \alpha' (x) \prec_{\alpha'} (\alpha' (y \prec z) + \alpha' (y \succ z)) \\
& = \alpha' (\alpha \alpha' (x) \prec (\alpha' (y \prec z) + \alpha' (y \succ z))) \\
& (\alpha' \text{ morphism}) = \alpha' \alpha \alpha' (x) \prec (\alpha'^2 (y \prec z) + \alpha'^2 (y \succ z)) \\
& (\alpha' \text{ morphism}) = \alpha' \alpha \alpha' (x) \prec (\alpha'^2 (y) \prec \alpha'^2 (z) + \alpha'^2 (y) \succ \alpha'^2 (z)) \\
& \quad = \alpha' \alpha \alpha' (x) \prec ([\alpha'^2 (y), \alpha'^2 (z)]) \\
& (\alpha, \alpha' \text{ commute}) = \alpha \alpha'^2 (x) \prec ([\alpha'^2 (y), \alpha'^2 (z)]) \\
& (\mathcal{A} \text{ Hom-Leib. dendr.}) = (\alpha'^2 (x) \prec \alpha'^2 (y)) \prec \alpha \alpha'^2 (z) + \alpha \alpha'^2 (y) \succ (\alpha'^2 (x) \prec \alpha'^2 (z)) \\
& (\alpha, \alpha' \text{ commute}) = (\alpha'^2 (x) \prec \alpha'^2 (y)) \prec \alpha' \alpha \alpha' (z) + \alpha' \alpha \alpha' (y) \succ (\alpha'^2 (x) \prec \alpha'^2 (z)) \\
& (\alpha' \text{ morphism}) = \alpha' (\alpha' (x) \prec \alpha' (y)) \prec \alpha' \alpha \alpha' (z) + \alpha' \alpha \alpha' (y) \succ \alpha' (\alpha' (x) \prec \alpha' (z)) \\
& \quad = (\alpha' (x) \prec \alpha' (y)) \prec_{\alpha'} \alpha \alpha' (z) + \alpha \alpha' (y) \succ_{\alpha'} (\alpha' (x) \prec \alpha' (z)) \\
& \quad = (x \prec_{\alpha'} y) \prec_{\alpha'} \alpha \alpha' (z) + \alpha \alpha' (y) \succ_{\alpha'} (x \prec_{\alpha'} z). \quad \square
\end{aligned}$$

**Corollary 4.7.** *If  $\mathcal{A} = (A, \prec, \succ, \alpha)$  is a multiplicative Hom-Leibniz dendriform algebra, then for any positive integer  $n$ ,*

(i) *The  $n$ th derived Hom-Leibniz dendriform algebra of type 1 of  $\mathcal{A}$  is defined by*

$$\mathcal{A}_1^n = (A, \prec^{(n)} = \alpha^n \circ \prec, \succ^{(n)} = \alpha^n \circ \succ, \alpha^{n+1}).$$

(ii) *The  $n$ th derived Hom-Leibniz dendriform algebra of type 2 of  $\mathcal{A}$  is defined by*

$$\mathcal{A}_2^n = (A, \prec^{(2^n-1)} = \alpha^{2^n-1} \circ \prec, \succ^{(2^n-1)} = \alpha^{2^n-1} \circ \succ, \alpha^{2^n}).$$

*Proof.* Apply Proposition 4.6 with  $\alpha' = \alpha^n$  and  $\alpha' = \alpha^{2^n-1}$  respectively.  $\square$

*Example 4.8.* There is a three-dimensional multiplicative Hom-Leibniz dendriform algebras  $(A, \prec, \succ, \alpha)$  with multiplications tables for a basis  $\{e_1, e_2, e_3\}$ :

$\prec$	$e_1$	$e_2$	$e_3$
$e_1$	0	0	$-e_2$
$e_2$	0	0	0
$e_3$	$-e_2$	0	$2e_2$

$\succ$	$e_1$	$e_2$	$e_3$
$e_1$	0	0	0
$e_2$	0	0	0
$e_3$	$-e_2$	0	$2e_2$

$$\alpha(e_1) = -\frac{p}{2}e_1, \quad \alpha(e_2) = \frac{p^2}{2}e_2, \quad \alpha(e_3) = pe_3,$$

with a parameter  $p \in \mathbb{K} \setminus \{0\}$ .

Then there are Hom-Leibniz dendriform algebras  $\mathcal{A}_1^n$  and  $\mathcal{A}_2^n$  with multiplications tables respectively:

$\prec^{(n)}$	$e_1$	$e_2$	$e_3$
$e_1$	0	0	$-(-\frac{p^2}{2})^n e_2$
$e_2$	0	0	0
$e_3$	$-(-\frac{p^2}{2})^n e_2$	0	$2p^{2n}e_2$

$\succ^{(n)}$	$e_1$	$e_2$	$e_3$
$e_1$	0	0	0
$e_2$	0	0	0
$e_3$	$-(-\frac{p^2}{2})^n e_2$	0	$2p^{2n}e_2$

$$\alpha^{n+1}(e_1) = \left(-\frac{p}{2}\right)^{n+1} e_1, \quad \alpha^{n+1}(e_2) = \left(\frac{p^2}{2}\right)^{n+1} e_2, \quad \alpha^{n+1}(e_3) = p^{n+1}e_3,$$

$\prec^{(2^n-1)}$	$e_1$	$e_2$	$e_3$
$e_1$	0	0	$(\frac{p^2}{2})^{2^n-1} e_2$
$e_2$	0	0	0
$e_3$	$(\frac{p^2}{2})^{2^n-1} e_2$	0	$2(p^2)^{2^n-1} e_2$

$\succ^{(2^n-1)}$	$e_1$	$e_2$	$e_3$
$e_1$	0	0	0
$e_2$	0	0	0
$e_3$	$(\frac{p^2}{2})^{2^n-1} e_2$	0	$2(p^2)^{2^n-1} e_2$

$$\alpha^{2^n}(e_1) = \left(\frac{p}{2}\right)^{2^n} e_1, \quad \alpha^{2^n}(e_2) = \left(\frac{p^2}{2}\right)^{2^n} e_2, \quad \alpha^{2^n}(e_3) = p^{2^n}e_3.$$

Next we define the notion of bimodule of a Hom-Leibniz dendriform algebra.

**Definition 4.9.** Let  $\mathcal{A} = (A, \prec, \succ, \alpha)$  be a Hom-Leibniz dendriform algebra, and  $(V, \beta)$  be a Hom-module. Let  $l_\prec, r_\prec, l_\succ, r_\succ : A \rightarrow gl(V)$  be four linear maps. The sextuple  $(l_\prec, r_\prec, l_\succ, r_\succ, \beta, V)$  is called a bimodule of  $\mathcal{A}$  if for all  $x, y \in A, v \in V$ ,

$$l_\succ([x, y])\beta(v) = l_\succ(\alpha(x))l_\succ(y)v - l_\succ(\alpha(y))l_\succ(x)v, \quad (4.5)$$

$$r_\succ(\alpha(y))l(x)v = l_\succ(\alpha(x))r_\succ(y)v - r_\succ(x \succ y)\beta(v), \quad (4.6)$$

$$r_\succ(\alpha(y))r(x)v = r_\succ(x \succ y)\beta(v) - l_\succ(\alpha(x))r_\succ(y)v, \quad (4.7)$$

$$l_\succ(\alpha(x))l_\prec(y)v = l_\prec(x \succ y)\beta(v) + l_\prec(\alpha(y))l(x)v, \quad (4.8)$$

$$l_\prec(\alpha(x))r_\prec(y)v = r_\prec(\alpha(y))l_\prec(x)v + r_\prec([x, y])\beta(v), \quad (4.9)$$

$$r_\prec(x \prec y)\beta(v) = r_\prec(\alpha(y))r_\prec(x)v + l_\prec(\alpha(x))r(y)v, \quad (4.10)$$

$$l_\prec(\alpha(x))l(y)v = l_\prec(x \prec y)\beta(v) + l_\prec(\alpha(y))l_\prec(x)v, \quad (4.11)$$

$$l_\prec(\alpha(x))r(y)v = r_\prec(\alpha(y))l_\prec(x)v + r_\succ(x \prec y)\beta(v), \quad (4.12)$$

$$r_\prec([x, y])\beta(v) = r_\prec(\alpha(y))r_\prec(x)v + l_\succ(\alpha(x))r_\prec(y)v, \quad (4.13)$$

$$\beta(l_{\prec}(x)v) = l_{\prec}(\alpha(x))\beta(v) \quad (4.14)$$

$$\beta(r_{\prec}(x)v) = l_{\prec}(\alpha(x))\beta(v) \quad (4.15)$$

$$\beta(l_{\succ}(x)v) = l_{\succ}(\alpha(x))\beta(v) \quad (4.16)$$

$$\beta(r_{\succ}(x)v) = l_{\succ}(\alpha(x))\beta(v) \quad (4.17)$$

where  $[x, y] = x \prec y + x \succ y$ ,  $l = l_{\prec} + l_{\succ}$  and  $r = r_{\prec} + r_{\succ}$ .

**Proposition 4.10.** *Let  $(l_{\prec}, r_{\prec}, l_{\succ}, r_{\succ}, \beta, V)$  be a bimodule of a Hom-Leibniz-dendriform algebra  $(A, \prec, \succ, \alpha)$ . Then, the direct sum  $A \oplus V$  of linear spaces is a Hom-Leibniz dendriform algebra with multiplication given for all  $x_1, x_2 \in A, v_1, v_2 \in V$  by*

$$\begin{aligned} (x_1 + v_1) \prec' (x_2 + v_2) &= x_1 \prec x_2 + (l_{\prec}(x_1)v_2 + r_{\prec}(x_2)v_1), \\ (x_1 + v_1) \succ' (x_2 + v_2) &= x_1 \succ x_2 + (l_{\succ}(x_1)v_2 + r_{\succ}(x_2)v_1), \\ (\alpha \oplus \beta)(x_1 + v_1) &= \alpha(x_1) + \beta(v_1). \end{aligned}$$

*Proof.* We prove the axiom (4.1) in  $A \oplus V$  as the others relations are proved analogously. For any  $x_1, x_2, x_3 \in A$  and  $v_1, v_2, v_3 \in V$ ,

$$\begin{aligned} &([x_1 + v_1, x_2 + v_2]') \succ' (\alpha + \beta)(x_3 + v_3) \\ &\quad - (\alpha + \beta)(x_1 + v_1) \succ' ((x_2 + v_2) \succ' (x_3 + v_3)) \\ &\quad + (\alpha + \beta)(x_2 + v_2) \succ' ((x_1 + v_1) \succ' (x_3 + v_3)) \\ &= ([x_1, x_2]l(x_1)v_2 + r(x_2)v_1) \succ' (\alpha + \beta)(x_3 + v_3) \\ &\quad - (\alpha + \beta)(x_1 + v_1) \succ' (x_2 \succ x_3 + l_{\succ}(x_2)v_3 + r_{\succ}(x_3)v_2) \\ &\quad + (\alpha + \beta)(x_2 + v_2) \succ' (x_1 \succ x_3 + l_{\succ}(x_1)v_3 + r_{\succ}(x_3)v_1) \\ &= [x_1, x_2] \succ \alpha(x_3) + l_{\succ}([x_1, x_2])\beta(v_3) + r_{\succ}(\alpha(x_3))l(x_1)v_2 \\ &\quad + r_{\succ}(\alpha(x_3))r(x_2)v_1 - \alpha(x_1) \succ (x_2 \succ x_3) - l_{\succ}(\alpha(x_1))l_{\succ}(x_2)v_3 \\ &\quad - l_{\succ}(\alpha(x_1))r_{\succ}(x_3)v_2 - r_{\succ}(x_2 \succ x_3)\beta(v_1) + \alpha(x_2) \succ (x_1 \succ x_3) \\ &\quad + l_{\succ}(\alpha(x_2))l_{\succ}(x_1)v_3 + l_{\succ}(\alpha(x_2))r_{\succ}(x_3)v_1 + r_{\succ}(x_1 \succ x_3)\beta(v_2). \end{aligned}$$

By (4.5)-(4.7), and (4.1) in  $A$ ,

$$\begin{aligned} &([x_1 + v_1, x_2 + v_2]') \succ' (\alpha + \beta)(x_3 + v_3) - (\alpha + \beta)(x_1 + v_1) \succ' ((x_2 + v_2) \succ' (x_3 + v_3)) \\ &\quad + (\alpha + \beta)(x_2 + v_2) \succ' ((x_1 + v_1) \succ' (x_3 + v_3)) = 0. \quad \square \end{aligned}$$

We denote such a Hom-Leibniz dendriform algebra by  $(A \oplus V, \prec', \succ', \alpha + \beta)$ , or  $A \times_{l_{\prec}, r_{\prec}, l_{\succ}, r_{\succ}, \alpha, \beta} V$ .

*Example 4.11.* Let  $(A, \prec, \succ, \alpha)$  be a Hom-Leibniz dendriform algebra, and

$$\begin{aligned} \forall (x, y) \in A \times A : \\ L_{\prec}(x)y &= x \prec y, & R_{\prec}(x)y &= y \prec x, \\ L_{\succ}(x)y &= x \succ y, & R_{\succ}(x)y &= y \succ x. \end{aligned}$$

Then,  $(L_{\prec}, R_{\prec}, L_{\succ}, R_{\succ}, \alpha, A)$  is called a regular bimodule of  $(A, \prec, \succ, \alpha)$ .

**Proposition 4.12.** *Let  $(l_{\prec}, r_{\prec}, l_{\succ}, r_{\succ}, \beta, V)$  be a bimodule of a Hom-Leibniz-dendriform algebra  $(A, \prec, \succ, \alpha)$ . Let  $(A, [\cdot, \cdot], \alpha)$  be the associated Hom-Leibniz algebra. Then,*

- 1)  $(l_{\succ}, r_{\prec}, \beta, V)$  and  $(l_{\prec} + l_{\succ}, r_{\prec} + r_{\succ}, \beta, V)$  are bimodules of  $(A, [\cdot, \cdot], \alpha)$ .

- 2) For any bimodule  $(l, r, \beta, V)$  of  $(A, [\cdot, \cdot], \alpha)$ ,  $(0, l, r, 0, \beta, V)$  is a bimodule of  $(A, \prec, \succ, \alpha)$ .
- 3)  $(0, l_{\prec} + l_{\succ}, r_{\prec} + r_{\succ}, 0, \beta, V)$  and  $(0, l_{\succ}, r_{\prec}, 0, \beta, V)$  are bimodules of  $(A, \prec, \succ, \alpha)$ .
- 4) The Hom-Leibniz dendriform algebras  $A \times_{l_{\prec}, r_{\prec}, l_{\succ}, r_{\succ}, \alpha, \beta} V$  and  $A \times_{0, l_{\prec} + l_{\succ}, r_{\prec} + r_{\succ}, 0, \alpha, \beta} V$  have the same associated Hom-Leibniz algebra  $A \times_{l_{\prec} + l_{\succ}, r_{\prec} + r_{\succ}, \alpha, \beta} V$ .

*Proof.* 1) We prove that  $(l_{\succ}, r_{\prec}, \beta, V)$  and  $(l_{\prec} + l_{\succ}, r_{\prec} + r_{\succ}, \beta, V)$  respectively satisfies the axioms (2.5)- (2.7). For any  $x, y \in A$  and  $v \in V$ ,

$$l_{\succ}(\alpha(x))l_{\succ}(y)v - l_{\succ}(\alpha(y))l_{\succ}(x)v - l_{\succ}([x, y])\beta(v) = 0, \quad (\text{by (4.5)})$$

$$l_{\succ}(\alpha(x))r_{\prec}(y)v - r_{\prec}(\alpha(y))l_{\succ}(x)v - r_{\prec}([x, y])\beta(v) = 0, \quad (\text{by (4.9)})$$

$$r_{\prec}([x, y])\beta(v) - r_{\prec}(\alpha(y))r_{\prec}(x)v - l_{\succ}(\alpha(x))r_{\prec}(y)v = 0. \quad (\text{by (4.13)})$$

Then  $(l_{\succ}, r_{\prec}, \beta, V)$  is a bimodule of  $(A, [\cdot, \cdot], \alpha)$ .

$$\begin{aligned} & (l_{\prec} + l_{\succ})(\alpha(x))(l_{\prec} + l_{\succ})(y)v - (l_{\prec} + l_{\succ})([x, y])\beta(v) \\ & \quad - (l_{\prec} + l_{\succ})(\alpha(y))(l_{\prec} + l_{\succ})(x)v \\ &= \overbrace{l_{\prec}(\alpha(x))l_{\prec}(y)v + l_{\prec}(\alpha(x))l_{\succ}(y)v}^* + l_{\succ}(\alpha(x))l_{\prec}(y)v + l_{\succ}(\alpha(x))l_{\succ}(y)v \\ & \quad - \overbrace{l_{\prec}([x, y])\beta(v)}^{**} - l_{\succ}([x, y])\beta(v) \\ & \quad - \overbrace{l_{\prec}(\alpha(y))l_{\prec}(x)v - l_{\prec}(\alpha(y))l_{\succ}(x)v}^{***} - l_{\succ}(\alpha(y))l_{\prec}(x)v - l_{\succ}(\alpha(y))l_{\succ}(x)v \\ &= \underbrace{\left( \overbrace{l_{\prec}(\alpha(x))l_{\prec}(y)v}^* - \overbrace{l_{\prec}(x \prec y)\beta(v)}^{**} - l_{\succ}(\alpha(y))l_{\prec}(x)v \right)}_{= 0 \text{ by (4.11)}} \\ & \quad + \underbrace{\left( l_{\succ}(\alpha(x))l_{\prec}(y)v - \overbrace{l_{\prec}(x \succ y)\beta(v)}^{**} - \overbrace{l_{\prec}(\alpha(y))l_{\prec}(x)v}^{***} \right)}_{= 0 \text{ by (4.8)}} \\ & \quad + \underbrace{\left( l_{\succ}(\alpha(x))l_{\succ}(y)v - l_{\succ}([x, y])\beta(v) - l_{\succ}(\alpha(y))l_{\succ}(x)v \right)}_{= 0 \text{ by (4.5)}} = 0, \end{aligned}$$

$$\begin{aligned} & (l_{\prec} + l_{\succ})(\alpha(x))(r_{\prec} + r_{\succ})(y)v - (r_{\prec} + r_{\succ})(\alpha(y))(l_{\prec} + l_{\succ})(x)v \\ & \quad - (r_{\prec} + r_{\succ})([x, y])\beta(v) \\ &= \overbrace{l_{\prec}(\alpha(x))r_{\prec}(y)v + l_{\prec}(\alpha(x))r_{\succ}(y)v}^* + l_{\succ}(\alpha(x))r_{\prec}(y)v + l_{\succ}(\alpha(x))r_{\succ}(y)v \\ & \quad - r_{\prec}(\alpha(y))l_{\prec}(x)v - r_{\prec}(\alpha(y))l_{\succ}(x)v - \overbrace{r_{\succ}(\alpha(y))l_{\prec}(x)v - r_{\succ}(\alpha(y))l_{\succ}(x)v}^{***} \\ & \quad - r_{\prec}([x, y])\beta(v) - \overbrace{r_{\succ}([x, y])\beta(v)}^{**} \end{aligned}$$

$$\begin{aligned}
&= \underbrace{\left( \overbrace{l_{\prec}(\alpha(x))r(y)v - r_{\prec}(\alpha(y))l_{\prec}(x)v}^* - \overbrace{r_{\succ}(x \prec y)\beta(v)}^{**} \right)}_{=0 \text{ by (4.12)}} \\
&+ \underbrace{\left( \overbrace{l_{\succ}(\alpha(x))r_{\succ}(y)v - r_{\succ}(\alpha(y))l(x)v}^{***} - \overbrace{r_{\succ}(x \succ y)\beta(v)}^{**} \right)}_{=0 \text{ by (4.6)}} \\
&+ \underbrace{\left( l_{\succ}(\alpha(x))r_{\prec}(y)v - r_{\prec}(\alpha(y))l_{\succ}(x)v - r_{\prec}([x, y])\beta(v) \right)}_{=0 \text{ by (4.9)}} = 0, \\
&(r_{\prec} + r_{\succ})([x, y])\beta(v) - (r_{\prec} + r_{\succ})(\alpha(y))(r_{\prec} + r_{\succ})(x)v \\
&\quad - (l_{\prec} + l_{\succ})(\alpha(x))(r_{\prec} + r_{\succ})(y)v \\
&= r_{\prec}([x, y])\beta(v) + \overbrace{r_{\succ}([x, y])\beta(v)}^{**} \\
&\quad - r_{\prec}(\alpha(y))r_{\prec}(x)v - r_{\prec}(\alpha(y))r_{\succ}(x)v - \overbrace{r_{\succ}(\alpha(y))r_{\prec}(x)v - r_{\succ}(\alpha(y))r_{\succ}(x)v}^{***} \\
&\quad - \overbrace{l_{\prec}(\alpha(x))r_{\prec}(y)v - l_{\prec}(\alpha(x))r_{\succ}(y)v}^* - l_{\succ}(\alpha(x))r_{\prec}(y)v - l_{\succ}(\alpha(x))r_{\succ}(y)v \\
&= \underbrace{\left( r_{\prec}([x, y])\beta(v) - r_{\prec}(\alpha(y))r_{\prec}(x)v - l_{\succ}(\alpha(x))r_{\prec}(y)v \right)}_{=0 \text{ by (4.13)}} \\
&\quad + \underbrace{\left( \overbrace{r_{\succ}(x \prec y)\beta(v)}^{**} - r_{\prec}(\alpha(y))r_{\succ}(x)v - \overbrace{l_{\prec}(\alpha(x))r(y)v}^* \right)}_{=0 \text{ by (4.10)}} \\
&\quad + \underbrace{\left( \overbrace{r_{\succ}(x \succ y)\beta(v)}^{**} - \overbrace{r_{\succ}(\alpha(y))r(x)v}^{***} - l_{\succ}(\alpha(x))r_{\succ}(y)v \right)}_{=0 \text{ by (4.7)}} = 0.
\end{aligned}$$

Then  $(l_{\prec} + l_{\succ}, r_{\prec} + l_{\succ}, \beta, V)$  is a bimodule of  $(A, [\cdot, \cdot], \alpha)$ .

2) In this case, we prove only the axioms (4.5), (4.9) and (4.13), the others axioms satisfied automatically. For any  $x, y \in A$  and  $v \in V$ , we have

$$l([x, y])\beta(v) = l(\alpha(x))l(y)v - l(\alpha(y))l(x)v, \quad (\text{by (2.5)})$$

$$l(\alpha(x))r(y)v = r(\alpha(y))l(x)v + r([x, y])\beta(v), \quad (\text{by (2.6)})$$

$$r([x, y])\beta(v) = r(\alpha(y))r(x)v + l(\alpha(x))r(y)v. \quad (\text{by (2.7)})$$

Then  $(0, l, r, 0, \beta, V)$  is a bimodule of  $(A, \prec, \succ, \alpha)$ .

3) It follows directly by 1) and 2), after applying 2) with  $l = l_{\succ}$ ,  $r = r_{\prec}$  and  $l = l_{\prec} + l_{\succ}$ ,  $r = r_{\prec} + r_{\succ}$  respectively.

4) Let  $(A \oplus V, [\cdot, \cdot]'_1, \alpha + \beta)$  and  $(A \oplus V, [\cdot, \cdot]'_2, \alpha + \beta)$  the Hom-Leibniz algebras associated respectively to Hom-Leibniz dendriform algebras  $A \times_{l_{\prec}, r_{\prec}, l_{\succ}, r_{\succ}, \alpha, \beta} V$  and  $A \times_{0, l_{\prec} + l_{\succ}, r_{\prec} + r_{\succ}, 0, \alpha, \beta} V$ . For all  $x_1, x_2 \in A$ ,  $v_1, v_2 \in V$ ,

$$\begin{aligned}
[x_1 + v_1, x_2 + v_2]'_1 &= (x_1 + v_1) \prec'_1 (x_2 + v_2) + (x_1 + v_1) \succ'_1 (x_2 + v_2) \\
&= x_1 \prec x_2 + x_1 \succ x_2 + l_{\prec}(x_1)v_2 + l_{\succ}(x_1)v_2 + r_{\prec}(x_2)v_1 + r_{\succ}(x_2)v_1
\end{aligned}$$

$$\begin{aligned}
&= [x_1, x_2] + (l_{\prec} + l_{\succ})(x_1)v_2 + (r_{\prec} + r_{\succ})(x_2)v_1 \\
&= x_1 \prec x_2 + (l_{\prec} + l_{\succ})(x_1)v_2 + x_1 \succ x_2 + (r_{\prec} + r_{\succ})(x_2)v_1 \\
&= (x_1 + v_1) \prec'_2 (x_2 + v_2) + (x_1 + v_1) \succ'_2 (x_2 + v_2) \\
&= [x_1 + v_1, x_2 + v_2]'_2.
\end{aligned}$$

Hence,  $[\cdot, \cdot]'_1 = [\cdot, \cdot]'_2$ .  $\square$

**Proposition 4.13.** *If  $f : (A_1, \prec_1, \succ_1, \alpha) \rightarrow (A_2, \prec_2, \succ_2, \beta)$  is a Hom-Leibniz dendriform algebras morphism, then  $(l_{\prec_1}, r_{\prec_1}, l_{\succ_1}, r_{\succ_1}, \beta, A_2)$  becomes a bimodule of  $(A_1, \prec_1, \succ_1, \alpha)$  via  $f$ , that is, for all  $(x, y) \in A_1 \times A_2$ ,  $l_{\prec_1}(x)y = f(x) \prec_2 y$ ,  $r_{\prec_1}(x)y = y \succ_2 f(x)$ ,  $l_{\succ_1}(x)y = f(x) \succ_2 y$ ,  $r_{\succ_1}(x)y = y \succ_2 f(x)$ .*

*Proof.* We prove the axiom (4.5), as the others being proved similarly. For any  $x, y \in A_1$  and  $z \in A_2$ ,

$$\begin{aligned}
l_{\succ_1}([x, y]_1)\beta(z) &= f([x, y]_1) \succ_2 \beta(z) = [f(x), f(y)]_2 \succ_2 \beta(z) \\
&\text{(by (4.1))} = \beta(f(x)) \succ_2 (f(y) \succ_2 z) - \beta(f(y)) \succ_2 (f(x) \succ_2 z) \\
&= f(\alpha(x)) \succ_2 (f(y) \succ_2 z) - f(\alpha(y)) \succ_2 (f(x) \succ_2 z) \\
&= l_{\succ_1}(\alpha(x))(f(y) \succ_2 z) - l_{\succ_1}(\alpha(y))(f(x) \succ_2 z) \\
&= l_{\succ_1}(\alpha(x))l_{\succ_1}(y)z - l_{\succ_1}(\alpha(y))l_{\succ_1}(x)z.
\end{aligned}$$

$\square$

**Theorem 4.14.** *Let  $\mathcal{A} = (A, \prec, \succ, \alpha)$  be a Hom-Leibniz dendriform algebra, and let  $\mathcal{V} = (l_{\prec_1}, r_{\prec_1}, l_{\succ_1}, r_{\succ_1}, \beta, V)$  be a bimodule of  $\mathcal{A}$ . Let  $\alpha'$  be a Hom-Leibniz dendriform algebra morphism of  $\mathcal{A}$  (in particular,  $\alpha$  and  $\alpha'$  commute), and let  $\beta'$  be a linear map on  $V$  such that the maps  $\beta$  and  $\beta'$  commute. Furthermore, suppose that*

$$\begin{cases} \beta' \circ l_{\prec} = (l_{\prec} \circ \alpha')\beta', \\ \beta' \circ r_{\prec} = (r_{\prec} \circ \alpha')\beta', \\ \beta' \circ l_{\succ} = (l_{\succ} \circ \alpha')\beta', \\ \beta' \circ r_{\succ} = (r_{\succ} \circ \alpha')\beta', \end{cases}$$

$\mathcal{A}_{\alpha'}$  is the Hom-Leibniz dendriform algebra  $(A, \prec_{\alpha'}, \succ_{\alpha'}, \alpha\alpha')$ ,

$$\tilde{l}_{\prec} = (l_{\prec} \circ \alpha')\beta', \tilde{r}_{\prec} = (r_{\prec} \circ \alpha')\beta, \tilde{l}_{\succ} = (l_{\succ} \circ \alpha')\beta', \tilde{r}_{\succ} = (r_{\succ} \circ \alpha')\beta'$$

and  $\mathcal{V}_{\beta'} = (\tilde{l}_{\prec}, \tilde{r}_{\prec}, \tilde{l}_{\succ}, \tilde{r}_{\succ}, \beta\beta', V)$ , Then,  $\mathcal{V}_{\beta'}$  is a bimodule of  $\mathcal{A}_{\alpha'}$ .

*Proof.* We prove only the axiom (4.5), as the others being proved similarly. For any  $x, y \in A$  and  $v \in V$ ,

$$\begin{aligned}
\tilde{l}_{\succ}([x, y]_{\alpha'})\beta\beta'(v) &= \tilde{l}_{\succ}([\alpha'(x), \alpha'(y)])\beta\beta'(v) \\
&= l_{\succ}([\alpha'^2(x), \alpha'^2(y)])\beta'\beta\beta'(v) \\
&= l_{\succ}([\alpha'^2(x), \alpha'^2(y)])\beta\beta'^2(v) \\
&= l_{\succ}(\alpha\alpha'^2(x))l_{\succ}(\alpha'^2(y))\beta'^2(v) - l_{\succ}(\alpha\alpha'^2(y))l_{\succ}(\alpha'^2(x))\beta'^2(v) \\
&= \tilde{l}_{\succ}(\alpha\alpha'(x))\tilde{l}_{\succ}(y)v - \tilde{l}_{\succ}(\alpha\alpha'(y))\tilde{l}_{\succ}(x)v.
\end{aligned}$$

$\square$

**Corollary 4.15.** *Let  $\mathcal{A} = (A, \prec, \succ, \alpha)$  be a multiplicative Hom-Leibniz dendriform algebra, and  $(l_{\prec}, r_{\prec}, l_{\succ}, r_{\succ}, \beta, V)$  a bimodule of  $\mathcal{A}$ . Then,  $\mathcal{V}_{\beta^q}$  is a bimodule of  $\mathcal{A}_{\alpha^p}$  for any nonnegative integers  $p$  and  $q$ .*

**Theorem 4.16.** Let  $\mathcal{A} = (A, \prec_A, \succ_A, \alpha)$  and  $\mathcal{B} = (B, \prec_B, \succ_B, \beta)$  be two Hom-Leibniz-dendriform algebras. If there are linear maps  $l_{\prec_A}, r_{\prec_A}, l_{\succ_A}, r_{\succ_A} : A \rightarrow gl(B)$ , and  $l_{\prec_B}, r_{\prec_B}, l_{\succ_B}, r_{\succ_B} : B \rightarrow gl(A)$  such that  $(l_{\prec_A}, r_{\prec_A}, l_{\succ_A}, r_{\succ_A}, \beta, B)$  is a bimodule of  $\mathcal{A}$ , and  $(l_{\prec_B}, r_{\prec_B}, l_{\succ_B}, r_{\succ_B}, \alpha, A)$  is a bimodule of  $\mathcal{B}$ , satisfying the following relations for  $x, y \in A, a, b \in B$ , and  $l_A = l_{\prec_A} + l_{\succ_A}, r_A = r_{\prec_A} + r_{\succ_A}, l_B = l_{\prec_B} + l_{\succ_B}, r_B = r_{\prec_B} + r_{\succ_B}$  :

$$\begin{aligned} l_{\succ_A}(r_B(a)x)\beta(b) + l_A(x)a \succ_B \beta(b) &= l_{\succ_A}(\alpha'(x))(a \succ_B b) \\ &\quad - \beta(a) \succ_B l_{\succ_A}(x)b - r_{\succ_A}(r_{\succ_B}(b)x)\beta(a), \end{aligned} \quad (4.18)$$

$$\begin{aligned} r_A(x)a \succ_B \beta(b) + l_{\succ_A}(l_B(a)x)\beta(b) &= \beta(a) \succ_B (l_{\succ_A}(x)b) \\ &\quad + r_{\succ_A}(r_{\succ_B}(b)x)\beta(a) - l_{\succ_A}(\alpha(x))(a \succ_B b), \end{aligned} \quad (4.19)$$

$$\begin{aligned} r_{\succ_A}(\alpha(x))([a, b]) &= \beta(a) \succ_B (r_{\succ_A}(x)b) + r_{\succ_A}(l_{\succ_B}(b)x)\beta(a) \\ &\quad - \beta(b) \succ_B (r_{\succ_A}(x)a) - r_{\succ_A}(l_{\succ_B}(a)x)\beta(b), \end{aligned} \quad (4.20)$$

$$\begin{aligned} l_{\succ_A}(\alpha(x))(a \prec_B b) &= (l_{\succ_A}(x)a) \prec_B \beta(b) + l_{\prec_A}(r_{\succ_B}(a)x)\beta(b) \\ &\quad + \beta(a) \prec (l_A(x)b) + r_{\prec_A}(r_B(b)x)\beta(a), \end{aligned} \quad (4.21)$$

$$\begin{aligned} \beta(a) \succ_B (l_{\prec_A}(x)b) + r_{\succ_A}(r_{\prec_B}(b)x)\beta(a) &= (r_{\succ_A}(x)a) \prec_B \beta(b) \\ &\quad + l_{\prec_A}(l_{\succ_B}(a)x)\beta(b) + l_{\prec_A}(\alpha(x))([a, b]), \end{aligned} \quad (4.22)$$

$$\begin{aligned} \beta(a) \succ_B (r_{\prec_A}(x)b) + r_{\succ_A}(l_{\prec_B}(b)x)\beta(a) &= r_{\prec_A}(\alpha(x))(a \succ_B b) \\ &\quad + \beta(b) \prec_B (r_A(x)a) + r_{\prec_A}(l_B(a)x)\beta(b), \end{aligned} \quad (4.23)$$

$$\begin{aligned} l_{\prec}(\alpha(x))([a, b]) &= (l_{\prec_A}(x)a) \prec_B \beta(b) + l_{\prec_A}(r_{\prec_B}(a)x)\beta(b) \\ &\quad + \beta(a) \succ_B (l_{\prec_A}(x)b) + r_{\succ_A}(r_{\prec_B}(b)x)\beta(a), \end{aligned} \quad (4.24)$$

$$\begin{aligned} \beta(a) \prec_B (l_A(x)b) + r_{\prec_A}(r_B(b)x)\beta(a) &= (r_{\prec_A}(x)a) \prec_B \beta(b) \\ &\quad + l_{\prec_A}(l_{\prec_B}(a)x)\beta(b) + l_{\succ_A}(\alpha(x))(a \prec_B b), \end{aligned} \quad (4.25)$$

$$\begin{aligned} \beta(a) \prec_B (r_A(x)b) + r_{\prec_A}(l_B(b)x)\beta(a) &= r_{\prec_A}(\alpha(x))(a \prec_B b) \\ &\quad + \beta(b) \succ_B (r_{\prec_A}(x)a) + r_{\succ_A}(l_{\prec_B}(a)x)\beta(b), \end{aligned} \quad (4.26)$$

$$\begin{aligned} l_{\succ_B}(r_A(x)a)\alpha(y) + l_B(a)x \succ_A \alpha(y) &= l_{\succ_B}(\beta'(a))(x \succ_A y) \\ &\quad - \alpha(x) \succ_A l_{\succ_B}(a)y - r_{\succ_B}(r_{\succ_A}(y)a)\alpha(x) \end{aligned} \quad (4.27)$$

$$\begin{aligned} r_B(a)x \succ_A \beta(y) + l_{\succ_B}(l_A(x)a)\alpha(y) &= \alpha(y) \succ_A (l_{\succ_B}(a)y) \\ &\quad + r_{\succ_B}(r_{\succ_A}(y)a)\beta(x) - l_{\succ_B}(\beta(a))(x \succ_A y) \end{aligned} \quad (4.28)$$

$$\begin{aligned} r_{\succ_B}(\beta(a))([x, y]) &= \alpha(x) \succ_A (r_{\succ_B}(a)y) + r_{\succ_B}(l_{\succ_A}(y)a)\alpha(x) \\ &\quad - \alpha(y) \succ_A (r_{\succ_B}(a)x) - r_{\succ_B}(l_{\succ_A}(x)a)\alpha(y), \end{aligned} \quad (4.29)$$

$$\begin{aligned} l_{\succ_B}(\beta(a))(x \prec_A y) &= (l_{\succ_B}(a)x) \prec_A \alpha(y) + l_{\prec_B}(r_{\succ_A}(x)a)\alpha(y) \\ &\quad + \alpha(x) \prec (l_B(a)y) + r_{\prec_B}(r_A(y)a)\alpha(x), \end{aligned} \quad (4.30)$$

$$\begin{aligned} \alpha(x) \succ_A (l_{\prec_B}(a)y) + r_{\succ_B}(r_{\prec_A}(y)a)\alpha(x) &= (r_{\succ_B}(a)x) \prec_A \alpha(y) \\ &\quad + l_{\prec_B}(l_{\succ_A}(x)a)\alpha(y) + l_{\prec_B}(\beta(a))([x, y]), \end{aligned} \quad (4.31)$$

$$\begin{aligned} \alpha(x) \succ_A (r_{\prec_B}(a)y) + r_{\succ_B}(l_{\prec_A}(y)a)\alpha(x) &= r_{\prec_B}(\beta(a))(x \succ_A y) \\ &\quad + \alpha(y) \prec_A (r_B(a)x) + r_{\prec_B}(l_A(x)a)\alpha(y), \end{aligned} \quad (4.32)$$

$$\begin{aligned} l_{\prec}(\beta(a))([x, y]) &= (l_{\prec_B}(a)x) \prec_A \alpha(y) + l_{\prec_B}(r_{\prec_A}(x)a)\alpha(y) \\ &\quad + \beta(x) \succ_A (l_{\prec_B}(a)y) + r_{\succ_B}(r_{\prec_A}(y)a)\alpha(x), \end{aligned} \quad (4.33)$$

$$\begin{aligned} \alpha(x) \prec_A (l_B(a)y) + r_{\prec_B}(r_A(y)a)\alpha(x) &= (r_{\prec_B}(a)x) \prec_A \alpha(y) \\ &\quad + l_{\prec_B}(l_{\prec_A}(x)a)\alpha(y) + l_{\succ_B}(\beta(a))(x \prec_A y), \end{aligned} \quad (4.34)$$

$$\alpha(x) \prec_A (r_B(a)y) + r_{\prec_B}(l_A(y)a)\alpha(x) = r_{\prec_B}(\beta(a))(x \prec_A y)$$



$$+\alpha(y) \succ_A (r_{\prec_B}(a)x) + r_{\succ_B}(l_{\prec_A}(x)a)\alpha(y), \quad (4.35)$$

then, there is a Hom-Leibniz dendriform algebra structure on the direct sum  $A \oplus B$  of the underlying linear spaces of  $A$  and  $B$  given for any  $x, y \in A, a, b \in B$  by

$$\begin{aligned} (x+a) \prec (y+b) &= (x \prec_A y + r_{\prec_B}(b)x + l_{\prec_B}(a)y) + \\ &\quad (l_{\prec_A}(x)b + r_{\prec_A}(y)a + a \prec_B b), \\ (x+a) \succ (y+b) &= (x \succ_A y + r_{\succ_B}(b)x + l_{\succ_B}(a)y) + \\ &\quad (l_{\succ_A}(x)b + r_{\succ_A}(y)a + a \succ_B b), \\ (\alpha \oplus \beta)(x+a) &= \alpha(x) + \beta(a). \end{aligned}$$

*Proof.* It is obtained in a similar way as for Theorem 2.11.  $\square$

**Definition 4.17.** Let  $(A, \prec_A, \succ_A, \alpha)$  and  $(B, \prec_B, \succ_B, \beta)$  be two Hom-Leibniz-dendriform algebras. Suppose there are linear maps  $l_{\prec_A}, r_{\prec_A}, l_{\succ_A}, r_{\succ_A} : A \rightarrow gl(B)$  and  $l_{\prec_B}, r_{\prec_B}, l_{\succ_B}, r_{\succ_B} : B \rightarrow gl(A)$  such that  $(l_{\prec_A}, r_{\prec_A}, l_{\succ_A}, r_{\succ_A}, \beta)$  is a bimodule of  $A$ , and  $(l_{\prec_B}, r_{\prec_B}, l_{\succ_B}, r_{\succ_B}, \alpha)$  is a bimodule of  $B$ . If (4.18)-(4.35) are satisfied, then

$$(A, B, l_{\prec_A}, r_{\prec_A}, l_{\succ_A}, r_{\succ_A}, \beta, l_{\prec_B}, r_{\prec_B}, l_{\succ_B}, r_{\succ_B}, \alpha)$$

is called a matched pair of Hom-Leibniz dendriform algebras.

**Corollary 4.18.** *If  $(A, B, l_{\prec_A}, r_{\prec_A}, l_{\succ_A}, r_{\succ_A}, \beta, l_{\prec_B}, r_{\prec_B}, l_{\succ_B}, r_{\succ_B}, \alpha)$  is a matched pair of Hom-Leibniz dendriform algebras, then*

$$(A, B, l_{\prec_A} + l_{\succ_A}, r_{\prec_A} + r_{\succ_A}, \beta, l_{\prec_B} + l_{\succ_B}, r_{\prec_B} + r_{\succ_B}, \alpha)$$

is a matched pair of the associated Hom-Leibniz algebras  $(A, [\cdot, \cdot]_A, \alpha)$  and  $(B, [\cdot, \cdot]_B, \beta)$ .

*Proof.* Let  $(A, B, l_{\prec_A}, r_{\prec_A}, l_{\succ_A}, r_{\succ_A}, \beta, l_{\prec_B}, r_{\prec_B}, l_{\succ_B}, r_{\succ_B}, \alpha)$  be a matched pair of a Hom-Leibniz dendriform algebras  $(A, \prec_A, \succ_A, \alpha)$  and  $(B, \prec_B, \succ_B, \beta)$ . In view of Proposition 4.12, the linear maps  $l_{\prec_A} + l_{\succ_A}, r_{\prec_A} + r_{\succ_A} : A \rightarrow gl(B)$  and  $l_{\prec_B} + l_{\succ_B}, r_{\prec_B} + r_{\succ_B} : B \rightarrow gl(A)$  are bimodules of the underlying Hom-Leibniz algebras  $(A, *_A, \alpha)$  and  $(B, *_B, \beta)$ , respectively. Therefore, (2.12), (2.14) are equivalent to (4.18)-(4.26) and (2.15), (2.17) are equivalent to (4.27)-(4.35).  $\square$

**Definition 4.19.** Let  $(A, [\cdot, \cdot], \alpha)$  be a Hom-Leibniz algebra, and  $(l, r, \beta, V)$  be a bimodule. A linear map  $T : V \rightarrow A$  is called an  $\mathcal{O}$ -operator associated to  $(l, r, \beta, V)$ , if  $T$  satisfies

$$\begin{aligned} \alpha T &= T\beta, \\ [T(u), T(v)] &= T(l(T(u))v + r(T(v))u) \text{ for all } u, v \in V. \end{aligned}$$

*Example 4.20.* Let  $(A, [\cdot, \cdot], \alpha)$  be a multiplicative Hom-Leibniz algebra. Then, the identity map  $id$  is an  $\mathcal{O}$ -operator associated to the bimodule  $(L, 0, \alpha)$  or  $(0, R, \alpha)$ .

*Example 4.21.* Let  $\mathcal{A} = (A, [\cdot, \cdot], \alpha)$  be a multiplicative Hom-Leibniz algebra. A linear map  $K : A \rightarrow A$  is called a Rota-Baxter operator on  $\mathcal{A}$  of weight zero if  $K$  satisfies

$$\begin{aligned} \alpha \circ K &= K \circ \alpha, \\ [K(x), K(y)] &= K([K(x), y] + [x, K(y)]) \text{ for all } x, y \in A. \end{aligned}$$

In fact, a Rota-Baxter operator on  $\mathcal{A}$  is just an  $\mathcal{O}$ -operator associated to the regular bimodule  $(L, R, \alpha)$ .

**Theorem 4.22.** *Let  $(A, [\cdot, \cdot], \alpha)$  be a Hom-Leibniz algebra, and  $(l, r, \beta, V)$  be a bimodule. Let  $T : V \rightarrow A$  be an  $\mathcal{O}$ -operator associated to  $(l, r, \beta, V)$ . Then, there exists a Hom-Leibniz dendriform algebra structure on  $V$  given for all  $u, v \in V$  by*

$$u \prec v = r(T(v))u, \quad u \succ v = l(T(u))v.$$

*So, there is an associated Hom-Leibniz algebra structure on  $V$  given by the equation (4.4), and  $T$  is a homomorphism of Hom-Leibniz algebras. Moreover,  $T(V) = \{T(v) \mid v \in V\} \subset A$  is a Hom-Leibniz subalgebra of  $A$ , and there is an induced Hom-Leibniz dendriform algebra structure on  $T(V)$  given for all  $u, v \in V$  by*

$$T(u) \prec T(v) = T(u \prec v), \quad T(u) \succ T(v) = T(u \succ v). \quad (4.36)$$

*Its corresponding associated Hom-Leibniz algebra structure on  $T(V)$  given by the equation (4.4) is just the Hom-Leibniz subalgebra structure of  $A$ , and  $T$  is a homomorphism of Hom-Leibniz dendriform algebras.*

*Proof.* For any  $x, y, z \in V$ ,

$$\begin{aligned} ([x, y]) \succ \beta(z) &= \beta(x) \succ (y \succ z) - \beta(y) \succ (x \succ z) \\ &= (r(T(y))x + l(T(x)y) \succ \beta(z) - \beta(x) \succ (l(T(y))z) - \beta(y) \succ (l(T(x))z) \\ &= l(T(r(T(y))x)\beta(z) + l(T(l(T(x))y)\beta(z) - l(T(\beta(x))l(T(y))z - l(T(\beta(y))l(T(x))z) \\ &= l(T(r(T(y))x + l(T(x)y)\beta(z) - l(\alpha T(x))l(T(y))z - l(\alpha T(y))l(T(x))z) \\ &= l([T(x), T(y)])\beta(z) - l([T(x), T(y)])\beta(z) = 0. \end{aligned}$$

The two other axioms are proved similarly.  $\square$

**Corollary 4.23.** *Let  $T : V \rightarrow A$  be an  $\mathcal{O}$ -operator on a Hom-Leibniz algebra  $(A, [\cdot, \cdot], \alpha)$  with respect to a representation  $(l, r, \beta, V)$ . Let  $[u, v]_C = u \prec v + u \succ v$ , for all  $u, v \in V$ . Then,  $T$  is a morphism from the Hom-Leibniz algebra  $(V, [\cdot, \cdot]_C, \beta)$  to  $(A, [\cdot, \cdot], \alpha)$ .*

*Proof.* For all  $u, v \in V$ ,

$$T([u, v]_C) = T(u \prec v + u \succ v) = T(u) \prec T(v) + T(u) \succ T(v) = [T(u), T(v)]. \quad \square$$

**Corollary 4.24.** *Let  $(A, [\cdot, \cdot], \alpha)$  be a multiplicative Hom-Leibniz algebra. Then, there is a compatible multiplicative Hom-Leibniz dendriform algebra structure on  $A$  if and only if there exists an invertible  $\mathcal{O}$ -operator of  $(A, [\cdot, \cdot], \alpha)$ .*

*Proof.* If  $T$  is an invertible  $\mathcal{O}$ -operator associated to a bimodule  $(l, r, \beta, V)$ , then, the compatible multiplicative Hom-Leibniz dendriform algebra structure on  $A$  is given by

$$x \succ y = T(l(x)T^{-1}(y)), \quad x \prec y = T(r(y)T^{-1}(x)) \quad \text{for all } x, y \in A.$$

Conversely, let  $(A, \prec, \succ, \alpha)$  be a Hom-Leibniz dendriform algebra, and  $(A, [\cdot, \cdot], \alpha)$  be the associated multiplicative Hom-Leibniz algebra. Then, the identity map  $id$  is an  $\mathcal{O}$ -operator associated to the bimodule  $(L_\succ, R_\prec, \alpha)$  of  $(A, [\cdot, \cdot], \alpha)$ .  $\square$

**Proposition 4.25.** Let  $(l_{\prec}, r_{\prec}, l_{\succ}, r_{\succ}, \beta, V)$  be a bimodule of a Hom-Leibniz-dendriform algebra  $(A, \prec, \succ, \alpha)$ , and let  $(A, [\cdot, \cdot], \alpha)$  be the associated involutive Hom-Leibniz algebra. Let  $l_{\prec}^*, r_{\prec}^*, l_{\succ}^*, r_{\succ}^* : A \rightarrow gl(V^*)$  be the linear maps given by

$$\begin{aligned} \langle l_{\prec}^*(x)a^*, y \rangle &= -\langle l_{\prec}(x)y, a^* \rangle, & \langle r_{\prec}^*(x)a^*, y \rangle &= -\langle r_{\prec}(x)y, a^* \rangle, \\ \langle l_{\succ}^*(x)a^*, y \rangle &= -\langle l_{\succ}(x)y, a^* \rangle, & \langle r_{\succ}^*(x)a^*, y \rangle &= -\langle r_{\succ}(x)y, a^* \rangle. \end{aligned}$$

- 1)  $(0, l_{\prec}^* + l_{\succ}^*, r_{\prec}^* + r_{\succ}^*, 0, \beta^*, V^*)$ ,  $(0, l_{\prec}^*, -l_{\prec}^* - r_{\prec}^*, 0, \beta^*, V^*)$  and  $(0, l_{\succ}^*, r_{\succ}^*, 0, \beta^*, V^*)$  are bimodules of  $(A, [\cdot, \cdot], \alpha)$ ;
- 2)  $(l_{\prec}^* + l_{\succ}^*, r_{\prec}^* + r_{\succ}^*, \beta^*, V^*)$ ,  $(l_{\prec}^*, -l_{\prec}^* - r_{\prec}^*, \beta^*, V^*)$ , and  $(l_{\succ}^*, r_{\succ}^*, \beta^*, V^*)$  are bimodules of  $(A, [\cdot, \cdot], \alpha)$ .

*Example 4.26.* If  $(A, \prec, \succ, \alpha)$  is an involutive Hom-Leibniz dendriform algebra, then

$$(L_{\prec}, R_{\prec}, L_{\succ}, R_{\succ}, \alpha, A), \quad (0, L_{\succ}, R_{\prec}, 0, \alpha, A), \quad (0, L_{\prec} + L_{\succ}, R_{\prec} + R_{\succ}, 0, \alpha, A)$$

are bimodules of  $(A, \prec, \succ, \alpha)$ . On the other hand,

$$(0, L_{\prec}^*, -L_{\prec}^* - R_{\prec}^*, 0, \alpha^*, A^*), \quad (0, L_{\prec}^* + L_{\succ}^*, R_{\prec}^* + R_{\succ}^*, 0, \alpha^*, A^*),$$

are bimodules of  $(A, \prec, \succ, \alpha)$  too.

**Theorem 4.27.** Let  $(A, [\cdot, \cdot], \alpha)$  be an involutive Hom-Leibniz algebra, and let  $\omega$  be a non-degenerate skew-symmetric bilinear form on  $A$  such that the following identity (invariance condition) is satisfied for all  $x, y, z \in A$ ,

$$\omega([\alpha(x), \alpha(y)], \alpha(z)) + \omega([\alpha(y), \alpha(z)], \alpha(x)) + \omega([\alpha(z), \alpha(x)], \alpha(y)) = 0. \quad (4.37)$$

Then, there exists a compatible Hom-Leibniz dendriform algebra structure  $\prec, \succ$  on  $(A, \alpha)$  given by

$$\omega(x \prec y, z) = \omega(y, \{z, x\}), \quad \omega(x \succ y, z) = \omega(x, \{y, z\}) \text{ for all } x, y \in A. \quad (4.38)$$

*Proof.* Define a linear map  $T : A \rightarrow A^*$  by  $\langle T(x), y \rangle = \omega(x, y)$  for all  $x, y \in A$ . Then,  $T$  is invertible and  $T^{-1}$  is an  $\mathcal{O}$ -operator of the involutive Hom-Leibniz algebra  $(A, [\cdot, \cdot], \alpha)$  associated to the bimodule  $(L^*, -L^* - R^*, \alpha^*)$ . By Corollary 4.24, there is a compatible Hom-Leibniz dendriform algebra structure  $\prec, \succ$  on  $(A, [\cdot, \cdot])$  given by

$$x \prec y = T^{-1}L^*(x)T(y), \quad x \succ y = T^{-1}(-L^* - R^*)(y)T(x)$$

for all  $x, y \in A$ , which gives exactly (4.38).  $\square$

**Theorem 4.28.** Let  $(A, \prec_A, \succ_A, \alpha)$  be an involutive Hom-Leibniz dendriform algebra, and  $(A, [\cdot, \cdot], \alpha)$  be the associated involutive Hom-Leibniz algebra. Suppose there is an involutive Hom-Leibniz dendriform algebra structure " $\prec_{A^*}, \succ_{A^*}, \alpha^*$ " on its dual space  $A^*$ , and  $(A^*, \{\cdot, \cdot\}, \alpha^*)$  is the associated Hom-Leibniz algebra. Then, is a standard Manin triple of involutive Hom-Leibniz algebras associated to  $(A, [\cdot, \cdot], \alpha)$  and  $(A^*, \{\cdot, \cdot\}, \alpha^*)$  if and only if the octuple  $(A, A^*, L_{\succ_A}^*, -L_{\succ_A}^* - R_{\succ_A}^*, \alpha^*, L_{\succ_{A^*}}^*, -L_{\succ_{A^*}}^* - R_{\succ_{A^*}}^*, \alpha)$  is a matched pair of Hom-Leibniz algebras.

*Proof.* The conclusion can be obtained by a similar proof as in Theorem 3.10. Then, if  $(A, A^*, L_{\succ_A}^*, -L_{\succ_A}^* - R_{\prec_A}^*, \alpha^*, L_{\succ_{A^*}}^*, -L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*, \alpha)$  is a matched pair of the involutive Hom-Leibniz algebras  $(A, [\cdot, \cdot], \alpha)$  and  $(A^*, \{\cdot, \cdot\}, \alpha^*)$ , it is straightforward to show that the bilinear form (3.5) is an  $(\alpha_1 \oplus \alpha_1^*)$ -invariant on the Hom-Leibniz algebra  $A \bowtie_{L_{\succ_A}^*, -L_{\succ_A}^* - R_{\prec_A}^*, \alpha^*}^{L_{\succ_{A^*}}^*, -L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*, \alpha} A^*$  given by

$$(x + a^*) *_{A \oplus A^*} (y + b^*) = ([x, y] + L_{\succ_{A^*}}^*(a^*)y + (-L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*)(b^*)x) + (\{a^*, b^*\} + L_{\succ_A}^*(x)b^* + (-L_{\succ_A}^* - R_{\prec_A}^*)(y)a^*).$$

In fact, we have

$$\begin{aligned} & \omega(\alpha(x) + \alpha^*(a^*), \alpha(y) + \alpha^*(b^*))', \alpha(z) + \alpha^*(c^*) \\ & + \omega(\alpha(y) + \alpha^*(b^*), \alpha(z) + \alpha^*(c^*))', \alpha(x) + \alpha^*(a^*) \\ & + \omega(\alpha(z) + \alpha^*(c^*), \alpha(x) + \alpha^*(a^*))', \alpha(y) + \alpha^*(b^*) \\ = & -\langle [\alpha(x), \alpha(y)] + L_{\succ_{A^*}}^*(\alpha^*(a^*))\alpha(y) - (L_{\prec_{A^*}}^* + L_{\succ_{A^*}}^*)(\alpha^*(b^*))\alpha(x), \alpha^*(c^*) \rangle \\ & + \langle \{\alpha^*(a^*), \alpha^*(b^*)\} + L_{\succ_A}^*(\alpha(x))\alpha^*(b^*) - (L_{\prec_A}^* + R_{\succ_A}^*)(\alpha(y))\alpha^*(a^*), \alpha(z) \rangle \\ & - \langle [\alpha(y), \alpha(z)] + L_{\succ_{A^*}}^*(\alpha^*(a^*))\alpha(z) - (L_{\prec_{A^*}}^* + L_{\succ_{A^*}}^*)(\alpha^*(c^*))\alpha(y), \alpha^*(a^*) \rangle \\ & + \langle \{\alpha^*(b^*), \alpha^*(c^*)\} + L_{\succ_A}^*(\alpha(y))\alpha^*(c^*) - (L_{\prec_A}^* + R_{\succ_A}^*)(\alpha(z))\alpha^*(b^*), \alpha(x) \rangle \\ & - \langle [\alpha(x), \alpha(y)] + L_{\succ_{A^*}}^*(\alpha^*(a^*))\alpha(y) - (L_{\prec_{A^*}}^* + L_{\succ_{A^*}}^*)(\alpha^*(b^*))\alpha(x), \alpha^*(c^*) \rangle \\ & + \langle \{\alpha^*(a^*), \alpha^*(b^*)\} + L_{\succ_A}^*(\alpha(x))\alpha^*(b^*) - (L_{\prec_A}^* + R_{\succ_A}^*)(\alpha(y))\alpha^*(a^*), \alpha(z) \rangle \\ = & -\langle [\alpha(x), \alpha(y)], \alpha^*(c^*) \rangle + \langle \alpha(y), \alpha^*(a^*) \succ \alpha^*(c^*) - \langle \alpha(x), \alpha^*(b^*) \succ \alpha^*(c^*) \rangle \\ & - \langle \alpha(x), \alpha^*(c^*) \prec \alpha^*(b^*) \rangle + \langle \{\alpha^*(a^*), \alpha^*(b^*)\}, \alpha(z) \rangle - \langle \alpha(x) \succ \alpha(z), \alpha^*(b^*) \rangle \\ & + \langle \alpha(y) \succ \alpha(z), \alpha^*(a^*) \rangle + \langle \alpha(z) \prec \alpha(y), \alpha^*(a^*) \rangle - \langle [\alpha(y), \alpha(z)], \alpha^*(a^*) \rangle \\ & + \langle \alpha(z), \alpha^*(b^*) \succ \alpha^*(a^*) - \langle \alpha(y), \alpha^*(c^*) \succ \alpha^*(a^*) \rangle - \langle \alpha(y), \alpha^*(a^*) \prec \alpha^*(c^*) \rangle \\ & + \langle \{\alpha^*(b^*), \alpha^*(c^*)\}, \alpha(x) \rangle - \langle \alpha(y) \succ \alpha(x), \alpha^*(c^*) \rangle + \langle \alpha(z) \succ \alpha(x), \alpha^*(b^*) \rangle \\ & + \langle \alpha(x) \prec \alpha(z), \alpha^*(b^*) \rangle - \langle [\alpha(z), \alpha(x)], \alpha^*(b^*) \rangle + \langle \alpha(x), \alpha^*(c^*) \succ \alpha^*(b^*) \rangle \\ & - \langle \alpha(z), \alpha^*(a^*) \succ \alpha^*(b^*) \rangle - \langle \alpha(y), \alpha^*(b^*) \prec \alpha^*(a^*) \rangle + \langle \{\alpha^*(c^*), \alpha^*(a^*)\}, \alpha(y) \rangle \\ & - \langle \alpha(z) \succ \alpha(y), \alpha^*(a^*) \rangle + \langle \alpha(x) \succ \alpha(y), \alpha^*(c^*) \rangle + \langle \alpha(y) \prec \alpha(x), \alpha^*(c^*) \rangle = 0. \end{aligned}$$

Conversely, if there exists a standard Manin triple of involutive Hom-Leibniz algebras associated to  $(A, [\cdot, \cdot], \alpha)$  and  $(A^*, \{\cdot, \cdot\}, \alpha^*)$ , then, the octuple  $(A, A^*, L_{\succ_A}^*, -L_{\succ_A}^* - R_{\prec_A}^*, \alpha^*, L_{\succ_{A^*}}^*, -L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*, \alpha)$  is a matched pair of the involutive Hom-Leibniz algebras given by

$$\begin{aligned} L_{\succ_A}^*(\alpha(x))(\{a, b\}) &= L_{\succ_A}^*((-L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*)(a)x)\beta(b) \\ &+ (-L_{\succ_A}^* - R_{\prec_A}^*)((-L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*)(b)x)\beta(a) + \{L_{\succ_A}^*(x)a, \beta(b)\} \\ &+ \{\beta(a), L_{\succ_A}^*(x)b\}, \end{aligned} \tag{4.39}$$

$$\begin{aligned} & \{\beta(a), L_{\succ_A}^*(x)b\} + (-L_{\succ_A}^* - R_{\prec_A}^*)((-L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*)(b)x)\beta(a) \\ &= \{(-L_{\succ_A}^* - R_{\prec_A}^*)(x)a, \beta(b)\} + L_{\succ_A}^*(L_{\succ_{A^*}}^*(a)x)\beta(b) \\ &+ L_{\succ_A}^*(\alpha(x))([a, b]), \end{aligned} \tag{4.40}$$

$$\begin{aligned} & \{\beta(a), (-L_{\succ_A}^* - R_{\prec_A}^*)(x)b\} + (-L_{\succ_A}^* - R_{\prec_A}^*)(L_{\succ_{A^*}}^*(b)x)\beta(a) \\ &= (-L_{\succ_A}^* - R_{\prec_A}^*)(\alpha(x))([a, b]) + \{\beta(b), (-L_{\succ_A}^* - R_{\prec_A}^*)(x)a\} \\ &+ (-L_{\succ_A}^* - R_{\prec_A}^*)(L_{\succ_{A^*}}^*(a)x)\beta(b) \end{aligned} \tag{4.41}$$

$$\begin{aligned} L_{\succ_{A^*}}^*(\beta(a))(\{x, y\}) &= L_{\succ_{A^*}}^*((-L_{\succ_A}^* - R_{\prec_A}^*)(x)a)\alpha(y) \\ &+ (-L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*)((-L_{\succ_A}^* - R_{\prec_A}^*)(x)a)\alpha(x) + \{L_{\succ_{A^*}}^*(a)x, \alpha(y)\} \end{aligned}$$

$$+ \{\alpha(x), L_{\succ_{A^*}}^*(a)y\}, \quad (4.42)$$

$$\begin{aligned} & \{\alpha(x), L_{\succ_{A^*}}^*(a)y\} + (-L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*)((-L_{\succ_A}^* - R_{\prec_A}^*)(y)a)\alpha(x) \\ &= \{(-L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*)(a)x, \alpha(y)\} + L_{\succ_{A^*}}^*(L_{\succ_A}^*(x)a)\alpha(y) \\ &+ L_{\succ_{A^*}}^*(\beta(a))([x, y]), \end{aligned} \quad (4.43)$$

$$\begin{aligned} & \{\alpha(x), (-L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*)(a)y\} + (-L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*)(L_{\succ_{A^*}}^*(y)a)\alpha(x) \\ &= (-L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*)(\beta(a))([x, y]) + \{\alpha(y), (-L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*)(a)x\} \\ &+ (-L_{\succ_{A^*}}^* - R_{\prec_{A^*}}^*)(L_{\succ_A}^*(x)a)\alpha(y), \end{aligned} \quad (4.44)$$

which completes the proof.  $\square$

**Corollary 4.29.** *Let  $(A, \prec, \succ, \alpha)$  be an involutive Hom-Leibniz dendriform algebra, and the triple  $(L_{\succ}^*, -L_{\succ}^* - R_{\prec}^*, \alpha^*)$  be the bimodule of the associated involutive Hom-Leibniz algebra  $(A, [\cdot, \cdot], \alpha)$ . Then,  $(T(A) = A \times_{L_{\succ}^*, -L_{\succ}^* - R_{\prec}^*, \alpha, \alpha^*} A^*, \omega)$  is a standard Manin triple of involutive Hom-Leibniz algebras.*

## 5 BiHom-Leibniz algebra

**Definition 5.1.** A BiHom-Leibniz algebra is a quadruple  $(A, [\cdot, \cdot], \alpha, \beta)$  consisting of a linear space  $A$ , bilinear map  $[\cdot, \cdot] : A \times A \rightarrow A$ , linear maps  $\alpha, \beta : A \rightarrow A$  satisfying, for all  $x, y, z \in A$ ,

$$\alpha \circ \beta = \beta \circ \alpha, \quad (\text{commutativity}) \quad (5.1)$$

$$[\alpha\beta(x), [y, z]] = [[\beta(x), y], \beta(z)] + [\beta(y), [\alpha(x), z]]. \quad (\text{BiHom-Leibniz identity}) \quad (5.2)$$

If, in addition,  $\alpha$  and  $\beta$  satisfies the multiplicativity properties

$$\alpha([x, y]) = [\alpha(x), \alpha(y)], \quad \beta([x, y]) = [\beta(x), \beta(y)], \quad (5.3)$$

then  $(A, [\cdot, \cdot], \alpha, \beta)$  is said to be multiplicative.

**Definition 5.2.** A BiHom-module is a triple  $(M, \alpha, \beta)$ , where  $M$  is a  $\mathbb{K}$ -linear space, and  $\alpha, \beta : M \rightarrow M$  are two linear maps.

**Definition 5.3.** Let  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  be a BiHom-Leibniz algebra and let  $(V, \beta_1, \beta_2)$  be a BiHom-module. Let  $l, r : A \rightarrow gl(V)$  be two linear maps. The quintuple  $(l, r, \beta_1, \beta_2, V)$  is called a bimodule of  $A$  if for all  $x, y \in A, v \in V$ ,

$$l(\alpha_1\alpha_2(x))l(y)v = l([\alpha_2(x), y])\beta_2(v) + l(\alpha_2(y))l(\alpha_1(x))v, \quad (5.4)$$

$$l(\alpha_1\alpha_2(x))r(y)v = r(\alpha_2(y))l(\alpha_2(x))v + r([\alpha_1(x), y])\beta_2(v), \quad (5.5)$$

$$r([x, y])\beta_1\beta_2(v) = r(\alpha_2(y))r(x)\beta_2(v) + l(\alpha_2(x))r(y)\beta_1(v), \quad (5.6)$$

$$\beta_1(l(x)v) = l(\alpha_1(x))\beta_1(v), \quad (5.7)$$

$$\beta_1(r(x)v) = r(\alpha_1(x))\beta_1(v), \quad (5.8)$$

$$\beta_2(l(x)v) = l(\alpha_2(x))\beta_2(v), \quad (5.9)$$

$$\beta_2(r(x)v) = l(\alpha_2(x))\beta_2(v). \quad (5.10)$$

**Proposition 5.4.** *Let  $(l, r, \beta_1, \beta_2, V)$  be a bimodule of a BiHom-Leibniz algebra  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$ . Then, the direct sum  $A \oplus V$  of linear spaces is turned into a BiHom-Leibniz algebra by defining multiplication in  $A \oplus V$  for all  $x_1, x_2 \in A, v_1, v_2 \in V$  by*

$$\begin{aligned} [x_1 + v_1, x_2 + v_2]' &= [x_1, x_2] + (l(x_1)v_2 + r(x_2)v_1), \\ (\alpha_1 \oplus \beta_1)(x_1 + v_1) &= \alpha_1(x_1) + \beta_1(v_1), (\alpha_2 \oplus \beta_2)(x_1 + v_1) = \alpha_2(x_1) + \beta_2(v_1). \end{aligned}$$

*Proof.* For all  $v_1, v_2, v_3 \in V$  and  $x_1, x_2, x_3 \in A$ ,

$$\begin{aligned} & [(\alpha_1 + \beta_1)(\alpha_2 + \beta_2)(x_1 + v_1), [x_2 + v_2, x_3 + v_3]']' \\ & \quad - [[(\alpha_2 + \beta_2)(x_1 + v_1), x_2 + v_2]', (\alpha_2 + \beta_2)(x_3 + v_3)]' \\ & \quad - [(\alpha_2 + \beta_2)(x_2 + v_2), [(\alpha_1 + \beta_1)(x_1 + v_1), x_3 + v_3]']' \\ & = [\alpha_1\alpha_2(x_1) + \beta_1\beta_2(v_1), [x_2, x_3] + l(x_2)v_3 + r(x_3)v_2]' \\ & \quad - [[\alpha_2(x_1), x_2] + l((\alpha_2(x_1))v_2 + r(x_2)\beta_2(v_1), \alpha_2(x_3) + \beta_2(v_3)]' \\ & \quad - [\alpha_2(x_2) + \beta_2(v_2), [\alpha_1(x_1), x_3] + l(\alpha_1(x_1))v_3 + r(x_3)\beta_1(v_1)]' \\ & = [\alpha_1\alpha_2(x_1), [x_2, x_3]] + l(\alpha\alpha_2(x_1))l(x_2)v_3 + l(\alpha_1\alpha_2(x_1))r(x_3)v_2 \\ & \quad + r([x_2, x_3])\beta_1\beta_2(v_1) - [[\alpha_2(x_1), x_2], \alpha_2(x_3)] - l([\alpha_2(x_1), x_2])r(x_2)\beta_2(v_1) \\ & \quad - r(\alpha_2(x_3))r(x_2)\beta_2(v_1) - [\alpha_2(x_2), [\alpha_1(x_1), x_3]] - l(\alpha_2(x_2))l(\alpha_1(x_1))v_3 \\ & \quad - l(\alpha_2(x_2))r(x_3)\beta_1(v_1) - r([\alpha_1(x_1), x_3])\beta_2(v_2). \end{aligned}$$

Then, by (5.2) and (5.4)-(5.6),

$$\begin{aligned} & [(\alpha_1 + \beta_1)(\alpha_2 + \beta_2)(x_1 + v_1), [x_2 + v_2, x_3 + v_3]']' \\ & \quad - [[(\alpha_2 + \beta_2)(x_1 + v_1), x_2 + v_2]', (\alpha_2 + \beta_2)(x_3 + v_3)]' \\ & \quad - [(\alpha_2 + \beta_2)(x_2 + v_2), [(\alpha_1 + \beta_1)(x_1 + v_1), x_3 + v_3]']' = 0, \end{aligned}$$

which completes the proof.  $\square$

We denote such a BiHom-Leibniz algebra by  $(A \oplus V, [\cdot, \cdot]', \alpha_1 + \beta_1, \alpha_2 + \beta_2)$ , or  $A \times_{l,r,\alpha_1,\alpha_2,\beta_1,\beta_2} V$ .

*Example 5.5.* Let  $(A, [\cdot, \cdot], \alpha, \beta)$  be a multiplicative BiHom-Leibniz algebra. Then the quadruples  $(L, 0, \alpha, \beta)$  and  $(L, R, \alpha, \beta)$  are bimodules of  $(A, [\cdot, \cdot], \alpha, \beta)$ .

**Proposition 5.6.** *Let  $\mathcal{V} = (l, r, \beta_1, \beta_2, V)$  be bimodule of a multiplicative BiHom-Leibniz algebra  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$ . Then, the quintuples  $\mathcal{V}_1 = (l \circ \alpha_1^n, r \circ \alpha_1^n, \beta_1, \beta_2, V)$  and  $\mathcal{V}_2 = (l \circ \alpha_2^n, r \circ \alpha_2^n, \beta_1, \beta_2, V)$  are bimodules of  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$ , for any non-negative integer  $n$ , or if moreover  $\alpha_1$  and  $\alpha_2$  are invertible, then for any integer  $n$ .*

*Proof.* For all  $x, y \in A, v \in V$  and non-negative integer  $n$ , the defining equalities (5.4), (5.5) and (5.6) in  $\mathcal{V}_1$  are proved as follows:

$$\begin{aligned} (l \circ \alpha_1^n)(\alpha_1\alpha_2(x))(l \circ \alpha_1^n)(y)v &= l(\alpha_1^{n+1}\alpha_2(x))l(\alpha_1^n(y))v \\ &= l([\alpha_1^n\alpha_2(x), \alpha_1^n(y)]\beta_2(v) + l(\alpha_2\alpha_1^n(y))l(\alpha_1^{n+1}(x))v) \\ &= (l \circ \alpha_1^n)([\alpha_2(x), y])\beta_2(v) \\ & \quad + (l \circ \alpha_1^n)(\alpha_2(y))(l \circ \alpha_1^n)(\alpha_1(x)), \end{aligned}$$

$$(l \circ \alpha_1^n)(\alpha_1\alpha_2(x))(r \circ \alpha_1^n)(y)v = l(\alpha_1^{n+1}\alpha_2(x))r(\alpha_1^n(y))v$$

$$\begin{aligned}
&= r(\alpha_2 \alpha_1^n(y)) l(\alpha_1^n \alpha_2(x)) v + r([\alpha_1^{n+1}(x), \alpha_1^n(y)] \beta_2(v)) \\
&= (r \circ \alpha_1^n)(\alpha_2(y)) (l \circ \alpha_1^n)(\alpha_2(x)) v \\
&\quad + (r \circ \alpha_1^n)([\alpha_1(x), y]) \beta_2(v), \\
(r \circ \alpha_1^n)([x, y]) \beta_1 \beta_2(v) &= r([\alpha_1^n(x), \alpha_1^n(y)]) \beta_1 \beta_2(v) \\
&= r(\alpha_2 \alpha_1^n(y)) r(\alpha_1^n(x)) \beta_2(v) + l(\alpha_1^n \alpha_2(x)) r(\alpha_1^n(y)) \beta_2(v) \\
&= (r \circ \alpha_1^n)(\alpha_2(y)) (r \circ \alpha_1^n)(x) \beta_2(v) \\
&\quad + (l \circ \alpha_1^n)(\alpha_2(x)) (r \circ \alpha_1^n)(y) \beta_2(v).
\end{aligned}$$

Similarly it can be shown that for any non-negative integer  $n$ ,  $\mathcal{V}_2$  is a bimodule of  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$ . If moreover  $\alpha_1$  and  $\alpha_2$  are invertible, then in the same way it follows that  $\mathcal{V}_1 = (l \circ \alpha_1^n, r \circ \alpha_1^n, \beta_1, \beta_2, V)$  and  $\mathcal{V}_2 = (l \circ \alpha_2^n, r \circ \alpha_2^n, \beta_1, \beta_2, V)$  are bimodules of  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$ , for any integer  $n$ , since  $\alpha_1^{-1}$  and  $\alpha_2^{-1}$  are BiHom-Leibniz algebras morphisms, and  $\alpha_i^j \circ \alpha_k^l = \alpha_k^l \circ \alpha_i^j$ , for all  $(i, k) \in \{1, 2\}^2$  and  $(j, l) \in \{1, -1\}^2$ .  $\square$

*Example 5.7.* Let  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  be a multiplicative BiHom-Leibniz algebra, then,  $(L \circ \alpha_1^n, R \circ \alpha_1^n, \alpha_1, \alpha_2, A)$  and  $(L \circ \alpha_2^n, R \circ \alpha_2^n, \alpha_1, \alpha_2, A)$  are bimodules of  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  for any non-negative integer  $n$ , or for any integer  $n$  if  $\alpha_1$  and  $\alpha_2$  are invertible.

**Theorem 5.8.** *Let  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  and  $(B, \{\cdot, \cdot\}, \beta_1, \beta_2)$  be two biHom-Leibniz algebras. Suppose there exist linear maps  $l_A, r_A : A \rightarrow gl(B)$ , and  $l_B, r_B : B \rightarrow gl(A)$  such that  $(l_A, r_A, \beta_1, \beta_2, B)$  is a bimodule of  $A$ , and  $(l_B, r_B, \alpha_1, \alpha_2, A)$  is a bimodule of  $B$ , satisfying, for any  $x, y \in A, a, b \in B$  the following conditions:*

$$\begin{aligned}
l_A(\alpha_1 \alpha_2(x))(\{a, b\}) &= l_A(r_B(a) \alpha_2(x)) \beta_2(b) + r_A(r_B(b) \alpha_1(x)) \beta_2(a) \\
&\quad + \{l_A(\alpha_2(x)) a, \beta_2(b)\} + \{\beta_2(a), l_A(\alpha_1(x)) b\}, \tag{5.11}
\end{aligned}$$

$$\begin{aligned}
\{\beta_1 \beta_2(a), l_A(x) b\} + r_A(r_B(b) x) \beta_1 \beta_2(a) &= \{r_A(x) \beta_2(a), \beta_2(b)\} \\
&\quad + l_A(l_B(\beta_2(a)) x) \beta_2(b) + l_A(\alpha_2(x))(\{\beta_1(a), b\}), \tag{5.12}
\end{aligned}$$

$$\begin{aligned}
\{\beta_1 \beta_2(a), r_A(x) b\} + r_A(l_B(b) x) \beta_1 \beta_2(a) &= r_A(\alpha_2(x))(\{\beta_2(a), b\}) \\
&\quad + \{\beta_2(b), r_A(x) \beta_1(a)\} + r_A(l_B(\beta_1(a)) x) \beta_2(b), \tag{5.13}
\end{aligned}$$

$$\begin{aligned}
l_B(\beta_1 \beta_2(a))([x, y]) &= l_B(r_A(x) \beta_2(a)) \alpha_2(y) + r_B(r_A(y) \beta_1(a)) \alpha_2(x) \\
&\quad + [l_B(\beta_2(a)) x, \alpha_2(y)] + [\alpha_2(x), l_B(\beta_1(a)) y], \tag{5.14}
\end{aligned}$$

$$\begin{aligned}
[\alpha_1 \alpha_2(x), l_B(a) y] + r_B(r_A(y) a) \alpha_1 \alpha_2(x) &= [r_B(a) \alpha_2(x), \alpha_2(y)] \\
&\quad + l_B(l_A(\beta_2(x)) a) \alpha_2(y) + l_B(\beta_2(a))([\alpha_1(x), y]), \tag{5.15}
\end{aligned}$$

$$\begin{aligned}
[\alpha_1 \alpha_2(x), r_B(a) y] + r_B(l_A(y) a) \alpha_1 \alpha_2(x) &= r_B(\beta_2(a))([\alpha_2(x), y]) \\
&\quad + [\alpha_2(y), r_B(a) \alpha_1(x)] + r_B(l_A(\alpha_1(x)) a) \alpha_2(y). \tag{5.16}
\end{aligned}$$

Then, there is a BiHom-Leibniz algebra structure on the direct sum  $A \oplus B$  of the underlying linear spaces of  $A$  and  $B$  given for all  $x, y \in A, a, b \in B$  by

$$\begin{aligned}
[x + a, y + b]' &= ([x, y] + l_B(a) y + r_B(b) x) + (\{a, b\} + l_A(x) b + r_A(y) a), \\
(\alpha_1 \oplus \beta_1)(x + a) &= \alpha_1(x) + \beta_1(a), (\alpha_2 \oplus \beta_2)(x + a) = \alpha_2(x) + \beta_2(a). \tag{5.17}
\end{aligned}$$

We denote this BiHom-Leibniz algebra by  $(A \bowtie B, [\cdot, \cdot]', \alpha_1 + \beta_1, \alpha_2 + \beta_2)$  or  $A \bowtie_{l_B, r_B, \alpha_1, \alpha_2}^{l_A, r_A, \beta_1, \beta_2} B$ .

**Definition 5.9.** Let  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  and  $(B, \{\cdot, \cdot\}, \beta_1, \beta_2)$  be two BiHom-Leibniz algebras. Suppose there exist linear maps  $l_A, r_A : A \rightarrow gl(B)$ , and  $l_B, r_B : B \rightarrow gl(A)$  such that  $(l_A, r_A, \beta_1, \beta_2)$  is a bimodule of  $A$ , and  $(l_B, r_B, \alpha_1, \alpha_2)$  is a bimodule of  $B$ . Then,  $(A, B, l_A, r_A, \beta_1, \beta_2, l_B, r_B, \alpha_1, \alpha_2)$  is called a matched pair of BiHom-Leibniz algebras, if the conditions (5.11) - (5.16) are satisfied.

*Example 5.10.* We consider the multiplicative BiHom-Leibniz algebra  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$ . By Example 5.5, we have  $(L, 0, \alpha_1, \alpha_2, A)$  and  $(L, R, \alpha, \alpha_2, A)$  are bimodules of  $A$ . Then  $(A, A, L, 0, \alpha_1, \alpha_2, L, R, \alpha_1, \alpha_2)$  is a matched pair of BiHom-Leibniz algebras if and only if for all  $x, y, z \in A$ ,  $[[\alpha_2(x), y], \alpha_2(z)] = 0$

*Example 5.11.* Consider the three-dimensional BiHom-Leibniz algebras  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  and  $(A, \{\cdot, \cdot\}, \beta_1, \beta_2)$  with multiplications tables for a basis  $\{e_1, e_2, e_3\}$ :

$[\cdot, \cdot]$	$e_1$	$e_2$	$e_3$
$e_1$	$e_3$	$-e_3$	$0$
$e_2$	$-e_3$	$3e_3$	$0$
$e_3$	$0$	$0$	$0$

$$\begin{aligned} \alpha_1(e_1) &= -e_1, & \alpha_1(e_2) &= -e_2, & \alpha_1(e_3) &= e_3, \\ \alpha_2(e_1) &= -2e_1, & \alpha_2(e_2) &= -2e_2, & \alpha_2(e_3) &= 4e_3, \end{aligned}$$

$\{\cdot, \cdot\}$	$e_1$	$e_2$	$e_3$
$e_1$	$-e_3$	$0$	$0$
$e_2$	$0$	$2e_3$	$0$
$e_3$	$0$	$0$	$0$

$$\begin{aligned} \beta_1(e_1) &= \frac{1}{2}e_1, & \beta_1(e_2) &= \frac{1}{2}e_2, & \beta_1(e_3) &= \frac{1}{4}e_3, \\ \beta_2(e_1) &= \sqrt{2}e_1, & \beta_2(e_2) &= \sqrt{2}e_2, & \beta_2(e_3) &= 2e_3. \end{aligned}$$

It is clear that  $(L, R, \beta_1, \beta_2)$  is a bimodule of  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$ , and  $(\mathcal{L}, \mathcal{R}, \alpha_1, \alpha_2)$  is a bimodule of  $(A, \{\cdot, \cdot\}, \beta_1, \beta_2)$ , where

$$\begin{aligned} \forall (x, y) \in A \times A : \\ L(x)y &= [x, y], & R(x)y &= [y, x], \\ \mathcal{L}(x)y &= \{x, y\}, & \mathcal{R}(x)y &= \{y, x\}. \end{aligned}$$

It is easy to see that the conditions (5.11) - (5.16) are satisfied for  $l_A = L$ ,  $r_A = R$ ,  $l_B = \mathcal{L}$ ,  $r_B = \mathcal{R}$  and  $B = A$ . Then  $(A, A, L, R, \beta_1, \beta_2, \mathcal{L}, \mathcal{R}, \alpha_1, \alpha_2)$  is a matched pair of BiHom-Leibniz algebras.

## 6 BiHom-Leibniz dendriform algebras

**Definition 6.1.** A BiHom-Leibniz dendriform algebra is a quintuple  $(A, \prec, \succ, \alpha, \beta)$  consisting of a linear space  $A$  on which the operations  $\prec, \succ : A \otimes A \rightarrow A$  and  $\alpha, \beta : A \rightarrow A$  are linear maps satisfying

$$\alpha \circ \beta = \beta \circ \alpha, \tag{6.1}$$

$$([\beta(x), y]) \succ \beta(z) = \alpha\beta(x) \succ (y \succ z) - \beta(y) \succ (\alpha(x) \succ z), \tag{6.2}$$



$$\alpha\beta(x) \succ (y \prec z) = (\beta(x) \succ y) \prec \beta(z) + \beta(y) \prec ([\alpha(x), z]), \quad (6.3)$$

$$\alpha\beta(x) \prec ([y, z]) = (\beta(x) \prec y) \prec \beta(z) + \beta(y) \succ (\alpha(x) \prec z), \quad (6.4)$$

where  $[x, y] = x \prec y + x \succ y$ .

**Definition 6.2.** Let  $(A, \prec, \succ, \alpha, \beta)$  and  $(A', \prec', \succ', \alpha', \beta')$  be BiHom-Leibniz dendriform algebras. A linear map  $f : A \rightarrow A'$  is a BiHom-Leibniz dendriform algebra morphism if

$$\prec' \circ (f \otimes f) = f \circ \prec, \quad \succ' \circ (f \otimes f) = f \circ \succ, \quad f \circ \alpha = \alpha' \circ f \quad \text{and} \quad f \circ \beta = \beta' \circ f.$$

**Proposition 6.3.** Let  $(A, \prec, \succ, \alpha, \beta)$  be a BiHom-Leibniz dendriform algebra.

Then,  $(A, [\cdot, \cdot] = \prec + \succ, \alpha, \beta)$  is a BiHom-Leibniz algebra.

*Proof.* For all  $x, y, z \in A$ ,

$$\begin{aligned} & [\alpha\beta(x), [y, z]] - [[\beta(x), y], \beta(z)] - [\beta(y), [\alpha(x), z]] \\ &= [\alpha\beta(x), y \prec z + y \succ z] - [\beta(x) \prec y + \beta(x) \succ y, \beta(z)] \\ &\quad - [\beta(y), \alpha(x) \prec z + \alpha(x) \succ z] \\ &= [\alpha\beta(x), y \prec z + y \succ z] - [\beta(x) \prec y + \beta(x) \succ y, \beta(z)] \\ &\quad - [\beta(y), \alpha(x) \prec z + \alpha(x) \succ z] \\ &= \alpha\beta(x) \prec (y \prec z) + \alpha\beta(x) \succ (y \prec z) + \alpha\beta(x) \prec (y \succ z) \\ &\quad + \alpha\beta(x) \succ (y \succ z) - (\beta(x) \prec y) \prec \beta(z) - (\beta(x) \prec y) \succ \beta(z) \\ &\quad - (\beta(x) \succ y) \prec \beta(z) - (\beta(x) \succ y) \succ \beta(z) - \beta(y) \prec (\alpha(x) \prec z) \\ &\quad - \beta(y) \succ (\alpha(x) \prec z) - \beta(y) \prec (\alpha(x) \succ z) - \beta(y) \succ (\alpha(x) \succ z) \\ &= \underbrace{\left( \alpha\beta(x) \succ (y \succ z) - \beta(y) \succ (\alpha(x) \succ z) - (\beta(x) \prec y + \beta(x) \succ y)\beta(z) \right)}_{= 0 \text{ by (6.2)}} \\ &\quad + \underbrace{\left( \alpha\beta(x) \succ (y \prec z) - (\beta(x) \succ y) \prec \beta(z) - \beta(y) \prec (\alpha(x) \prec z + \alpha(x) \succ z) \right)}_{= 0 \text{ by (6.3)}} \\ &\quad + \underbrace{\left( \alpha\beta(x) \prec (y \prec z + y \succ z) - (\beta(x) \prec y) \prec \beta(z) - \beta(y) \succ (\alpha(x) \prec z) \right)}_{= 0 \text{ by (6.4)}} = 0. \quad \square \end{aligned}$$

We call  $(A, [\cdot, \cdot], \alpha, \beta)$  the BiHom-Leibniz algebra of  $(A, \prec, \succ, \alpha, \beta)$ , and the quintuple  $(A, \prec, \succ, \alpha, \beta)$  is called a compatible BiHom-Leibniz dendriform algebra structure on the BiHom-Leibniz algebra  $(A, [\cdot, \cdot], \alpha, \beta)$ .

The following theorem shows that any Leibniz dendriform algebra [73] together with two commuting morphisms yield BiHom-Leibniz dendriform algebra, and morphisms of Leibniz dendriform algebra naturally correspond to morphisms of BiHom-Leibniz dendriform algebra.

**Theorem 6.4.** Let  $\mathcal{A} = (A, \prec, \succ)$  be a Leibniz dendriform algebra and  $\alpha_1, \alpha_2 : A \rightarrow A$  be two morphisms of  $\mathcal{A}$  such that  $\alpha_1\alpha_2 = \alpha_2\alpha_1$ . Then,

- 1)  $\mathcal{A}_{\alpha_1, \alpha_2} := (A, \prec_{\alpha_1, \alpha_2} = \prec \circ (\alpha_1 \otimes \alpha_2), \succ_{\alpha_1, \alpha_2} = \succ \circ (\alpha_1 \otimes \alpha_2), \alpha_1, \alpha_2)$  is a BiHom-Leibniz dendriform algebra;

- 2) if  $(A', \prec', \succ')$  is another Leibniz dendriform algebra,  $\alpha'_1, \alpha'_2 : A' \rightarrow A'$  are two commuting Leibniz dendriform algebra morphisms, and  $f : A \rightarrow A'$  is a Leibniz dendriform algebra morphism satisfying  $f \circ \alpha_1 = \alpha'_1 \circ f$  and  $f \circ \alpha_2 = \alpha'_2 \circ f$ , then  $f : \mathcal{A}_{\alpha_1, \alpha_2} \rightarrow \mathcal{A}'_{\alpha'_1, \alpha'_2}$  is a BiHom-Leibniz dendriform algebra morphism.

*Proof.* We shall only prove relation (6.2) as the others are proved analogously.

For any  $x, y, z \in A$  :

$$\begin{aligned}
& ([\alpha_2(x), y]_{\alpha_1, \alpha_2}) \succ_{\alpha_1, \alpha_2} \alpha_2(z) \\
&= \alpha_1([\alpha_1\alpha_2(x), \alpha_2(y)]) \succ \alpha_2^2(z) = [\alpha_1^2\alpha_2(x), \alpha_1\alpha_2(y)] \succ \alpha_2^2(z) \\
&= \alpha_1^2\alpha_2(x) \succ (\alpha_1\alpha_2(y) \succ \alpha_2^2(z)) - \alpha_1\alpha_2(y) \succ (\alpha_1^2\alpha_2(x) \succ \alpha_2^2(z)) \\
&= \alpha_1^2\alpha_2(x) \succ \alpha_2(\alpha_1(y) \succ \alpha_2(z)) - \alpha_1\alpha_2(y) \succ \alpha_2(\alpha_1^2(x) \succ \alpha_2(z)) \\
&= \alpha_1\alpha_2(x) \succ_{\alpha_1, \alpha_2} (y \succ_{\alpha_1, \alpha_2} z) - \alpha_2(y) \succ_{\alpha_1, \alpha_2} (\alpha_1(x) \succ_{\alpha_1, \alpha_2} z).
\end{aligned}$$

The second assertion follows from

$$\begin{aligned}
f(x \prec_{\alpha_1, \alpha_2} y) &= f(\alpha_1(x) \prec \alpha_2(y)) = f(\alpha_1(x)) \prec' f(\alpha_2(y)) \\
&= \alpha'_1 f(x) \prec' \alpha'_2 f(y) = f(x) \prec'_{\alpha_1, \alpha_2} f(y).
\end{aligned}$$

Similarly,  $f(x \succ_{\alpha_1, \alpha_2} y) = f(x) \succ'_{\alpha'_1, \alpha'_2} f(y)$ . This completes the proof.  $\square$

**Proposition 6.5.** *Let  $\mathcal{A} = (A, \prec, \succ, \alpha_1, \alpha_2)$  be a BiHom-Leibniz dendriform algebra and  $\alpha'_1, \alpha'_2 : \mathcal{A} \rightarrow \mathcal{A}$  be a two BiHom-Leibniz dendriform algebra morphisms such that any two of the maps  $\alpha_1, \alpha_2, \alpha'_1, \alpha'_2$  commute. Then  $\mathcal{A}'_{\alpha'_1, \alpha'_2} = (A, \prec_{\alpha'_1, \alpha'_2}, \succ_{\alpha'_1, \alpha'_2}, \alpha_1 \circ \alpha'_1, \alpha_2 \circ \alpha'_2)$  is a BiHom-Leibniz dendriform algebra.*

*Proof.* We prove the axioms (6.2)-(6.4) in  $\mathcal{A}'_{\alpha'_1, \alpha'_2}$ . For any  $x, y, z \in A$ ,

$$\begin{aligned}
& ([\alpha_2\alpha'_2(x), y]_{\alpha'_1, \alpha'_2}) \succ_{\alpha'_1, \alpha'_2} \alpha_2\alpha'_2(z) \\
&= \alpha'_1[\alpha'_1\alpha'_2(x), \alpha'_2(y)] \succ \alpha_2\alpha'^2_2(z) \\
&= [\alpha'^2_1\alpha'_2\alpha_2(x), \alpha'_1\alpha'_2(y)] \succ \alpha_2\alpha'^2_2(z) \\
&= \alpha_1\alpha_2\alpha'^2_1\alpha'_2(x) \succ (\alpha'_1\alpha'_2 \succ \alpha'^2_2(z)) - \alpha_2\alpha'_1\alpha'_2(y) \succ (\alpha_1\alpha'_1\alpha'_2(x) \succ \alpha'^2_2(z)) \\
&= \alpha_1\alpha'^2_1\alpha_2\alpha'_2(x) \succ \alpha'_2(\alpha_1 \succ \alpha'_2(z)) - \alpha'_1\alpha_2\alpha'_2(y) \succ \alpha'_2(\alpha_1\alpha'^2_1(x) \succ \alpha'_2(z)) \\
&= \alpha_1\alpha'_1\alpha_2\alpha'_2(x) \succ_{\alpha'_1, \alpha'_2} (y \succ_{\alpha'_1, \alpha'_2} z) - \alpha_2\alpha'_2(y) \succ_{\alpha'_1, \alpha'_2} (\alpha_1\alpha'_1(x) \succ_{\alpha'_1, \alpha'_2} z), \\
& \alpha_1\alpha'_1\alpha_2\alpha'_2(x) \succ_{\alpha'_1, \alpha'_2} (y \prec_{\alpha'_1, \alpha'_2} z) \\
&= \alpha_1\alpha_2\alpha'^2_1\alpha'_2(x) \succ \alpha'_2(\alpha'_1(y) \prec \alpha'_2(z)) \\
&= \alpha_1\alpha_2\alpha'^2_1\alpha'_2(x) \succ (\alpha'_1\alpha'_2(y) \prec \alpha'^2_2(z)) \\
&= (\alpha'^2_1\alpha'_2(x) \succ \alpha'_1\alpha'_2(y)) \prec \alpha_2\alpha'^2_2(z) + \alpha_2\alpha'_1\alpha'_2(y) \succ ([\alpha_1\alpha'^2_1\alpha'_2(x), \alpha'_2(z)]) \\
&= (\alpha'_1\alpha_2\alpha'_2(x) \succ_{\alpha'_1, \alpha'_2} \alpha'_1(y)) \prec \alpha_2\alpha'^2_2(z) + \alpha_2\alpha'_1\alpha'_2(y) \prec ([\alpha_1\alpha'_1\alpha'_2(x), \alpha'_2(z)]_{\alpha'_1, \alpha'_2}) \\
&= (\alpha_2\alpha'_2(x) \succ_{\alpha'_1, \alpha'_2} y) \prec_{\alpha'_1, \alpha'_2} \alpha_2\alpha'_2(z) + \alpha_2\alpha'_2(y) \prec_{\alpha'_1, \alpha'_2} ([\alpha_1\alpha'_1(x), z]_{\alpha'_1, \alpha'_2}), \\
& \alpha_1\alpha'_1\alpha_2\alpha'_2(x) \prec_{\alpha'_1, \alpha'_2} ([y, z]_{\alpha'_1, \alpha'_2}) \\
&= \alpha_1\alpha'^2_1\alpha_2\alpha'_2(x) \prec [\alpha'_1\alpha'_2(y), \alpha'^2_2(z)]
\end{aligned}$$

$$\begin{aligned}
&= (\alpha_2 \alpha_1'^2 \alpha_2'(x) \prec \alpha_1' \alpha_2'(y)) \prec \alpha_2 \alpha_2'^2(z) + \alpha_2 \alpha_1' \alpha_2'(y) \succ (\alpha \alpha_1'^2 \alpha_2'(x) \prec \alpha_2'^2(z)) \\
&= (\alpha_2 \alpha_1' \alpha_2'(x) \prec_{\alpha_1', \alpha_2'} \alpha_1'(y)) \prec \alpha_2 \alpha_2'^2(z) + \alpha_2 \alpha_1' \alpha_2'(y) \succ (\alpha_1 \alpha_1' \alpha_2'(x) \succ_{\alpha_1', \alpha_2'} \alpha_2'(z)) \\
&= (\alpha_2 \alpha_2'(x) \prec_{\alpha_1', \alpha_2'} y) \prec_{\alpha_1', \alpha_2'} \alpha_2 \alpha_2'(z) + \alpha_2 \alpha_2'(y) \succ_{\alpha_1', \alpha_2'} (\alpha_1 \alpha_1'(x) \prec_{\alpha_1', \alpha_2'} z)
\end{aligned}$$

Then  $\mathcal{A}_{\alpha_1', \alpha_2'} = (A, \prec_{\alpha_1', \alpha_2'}, \succ_{\alpha_1', \alpha_2'}, \alpha_1 \circ \alpha_1', \alpha_2 \circ \alpha_2')$  is a BiHom-Leibniz dendriform algebra.  $\square$

*Example 6.6.* There are three-dimensional BiHom-Leibniz dendriform algebras  $(A, \prec, \succ, \alpha_1, \alpha_2)$  with multiplications tables for a basis  $\{e_1, e_2, e_3\}$ :

$\prec$	$e_1$	$e_2$	$e_3$	$\succ$	$e_1$	$e_2$	$e_3$
$e_1$	0	0	0	$e_1$	0	0	0
$e_2$	0	0	$-2e_1$	$e_2$	0	0	$-2e_1$
$e_3$	0	0	$3e_1$	$e_3$	0	$-2e_1$	$3e_1$

$$\begin{aligned}
\alpha_1(e_1) &= \lambda_1 e_1, & \alpha_1(e_2) &= 0, & \alpha_1(e_3) &= \lambda_2 e_3, \\
\alpha_2(e_1) &= 0, & \alpha_2(e_2) &= \mu_1 e_2, & \alpha_2(e_3) &= \mu_2 e_3,
\end{aligned}$$

where  $\lambda_1, \lambda_2, \mu_1$  and  $\mu_2$  are parameters in  $\mathbb{K} \setminus \{0\}$ . Using suitable algebra morphisms  $\alpha_1'$  and  $\alpha_2'$ , we obtain the BiHom-Leibniz dendriform algebras. The algebra morphisms  $\alpha_1', \alpha_2' : A \rightarrow A$  are given by

$$\begin{aligned}
\alpha_1'(e_1) &= \frac{p^2}{3} e_1, & \alpha_1'(e_2) &= -\frac{2p}{3} e_2, & \alpha_1'(e_3) &= p e_3, \\
\alpha_2'(e_1) &= \frac{q^2}{3} e_1, & \alpha_2'(e_2) &= -\frac{2q}{3} e_2, & \alpha_2'(e_3) &= q e_3,
\end{aligned}$$

where  $p$  and  $q$  are parameters in  $\mathbb{K} \setminus \{0\}$ . Then there is a BiHom-Leibniz dendriform algebra  $\mathcal{A}_{\alpha_1', \alpha_2'} = (A, \prec_{\alpha_1', \alpha_2'}, \succ_{\alpha_1', \alpha_2'}, \alpha_1 \circ \alpha_1', \alpha_2 \circ \alpha_2')$  with multiplications tables:

$\prec_{\alpha_1', \alpha_2'}$	$e_1$	$e_2$	$e_3$	$\succ_{\alpha_1', \alpha_2'}$	$e_1$	$e_2$	$e_3$
$e_1$	0	0	0	$e_1$	0	0	0
$e_2$	0	0	$\frac{4pq}{3} e_1$	$e_2$	0	0	$\frac{4pq}{3} e_1$
$e_3$	0	0	$3pq e_1$	$e_3$	0	$\frac{4pq}{3} e_1$	$3pq e_1$

$$\begin{aligned}
\alpha_1 \alpha_1'(e_1) &= \frac{\lambda_1 p^2}{3} e_1, & \alpha_1 \alpha_1'(e_2) &= 0, & \alpha_1 \alpha_1'(e_3) &= \lambda_2 p e_3, \\
\alpha_2 \alpha_2'(e_1) &= 0, & \alpha_2 \alpha_2'(e_2) &= -\frac{2\mu_1 q}{3} e_2, & \alpha_2 \alpha_2'(e_3) &= \mu_2 q e_3.
\end{aligned}$$

**Corollary 6.7.** *Let  $(A, \prec, \succ, \alpha_1, \alpha_2)$  be a multiplicative BiHom-Leibniz dendriform algebra. Let  $n$  be any positive integer.*

(i) *The  $n$ th derived BiHom-Leibniz dendriform algebra of type 1 of  $A$  is defined by*

$$\mathcal{A}_1^n = (A, \prec^{(n)} = \prec \circ (\alpha_1^n \otimes \alpha_2^n), \succ^{(n)} = \succ \circ (\alpha_1^n \otimes \alpha_2^n), \alpha_1^{n+1}, \alpha_2^{n+1}).$$

(ii) *The  $n$ th derived BiHom-Leibniz dendriform algebra of type 2 of  $A$  is defined by*

$$\mathcal{A}_2^n = (A, \prec^{(2^n-1)} = \prec \circ (\alpha_1^{2^n-1} \otimes \alpha_2^{2^n-1}), \succ^{(2^n-1)} = \succ \circ (\alpha_1^{2^n-1} \otimes \alpha_2^{2^n-1}), \alpha_1^{2^n}, \alpha_2^{2^n}).$$

*Proof.* Apply Proposition 6.5 with  $\alpha_1' = \alpha_1^n$ ,  $\alpha_2' = \alpha_2^n$  and  $\alpha_1' = \alpha_1^{2^n-1}$ ,  $\alpha_2' = \alpha_2^{2^n-1}$  respectively. We prove in more details, only the axiom (6.2) in both cases  $\mathcal{A}_1^n$  and  $\mathcal{A}_2^n$ , as others proved similarly.

(i) For any  $x, y, z \in A$ ,

$$\begin{aligned}
& ([\alpha_2^{n+1}(x), y]^{(n)}) \succ^{(n)} \alpha_2^{n+1}(z) \\
&= \alpha_1^n([\alpha_1^n \alpha_2^{n+1}(x), \alpha_2^n(y)]) \succ \alpha_2^{2n+1}(z) \\
&= [\alpha_1^{2n} \alpha_2^{n+1}(x), \alpha_1^n \alpha_2^n(y)] \succ \alpha_2^{2n+1}(z) \\
&= \alpha_1^{2n+1} \alpha_2^{n+1}(x) \succ (\alpha_1^n \alpha_2^n(y) \succ \alpha_2^{2n}(z)) - \alpha_1^n \alpha_2^{n+1}(y) \succ (\alpha_1^{2n+1} \alpha_2^n(x) \succ \alpha_2^{2n}(z)) \\
&= \alpha_1^{2n+1} \alpha_2^{n+1}(x) \succ (\alpha_2^n(y) \succ^{(n)} \alpha_2^n(z)) - \alpha_1^n \alpha_2^{n+1}(y) \succ (\alpha_1^{n+1} \alpha_2^n(x) \succ^{(n)} \alpha_2^n(z)) \\
&= \alpha_1^{n+1} \alpha_2^{n+1}(x) \succ^{(n)} (y \succ^{(n)} z) - \alpha_2^{n+1}(y) \succ^{(n)} (\alpha_1^{n+1}(x) \succ^{(n)} z).
\end{aligned}$$

(ii) For any  $x, y, z \in A$ ,

$$\begin{aligned}
& (\alpha_2^{2^n}(x), y]^{2^n-1}) \succ^{2^n-1} \alpha_2^{2^n}(z) \\
&= \alpha_1^{2^n-1}([\alpha_1^{2^n-1} \alpha_2^{2^n}(x), \alpha_2^{2^n-1}(y)]) \succ \alpha_2^{2^n-1} \alpha_2^{2^n}(z) \\
&= ([\alpha_1^{2^n-1} \alpha_1^{2^n-1} \alpha_2^{2^n}(x), \alpha_1^{2^n-1} \alpha_2^{2^n-1}(y)]) \succ \alpha_2^{2^n-1} \alpha_2^{2^n}(z) \\
&= \alpha_1^{2^n-1} \alpha_1^{2^n} \alpha_2^{2^n}(x) \succ (\alpha_2^{2^n-1}(y) \succ^{(2^n-1)} \alpha_2^{2^n-1}(z)) \\
&\quad - \alpha_1^{2^n-1} \alpha_2^{2^n}(y) \succ (\alpha_1^{2^n} \alpha_2^{2^n-1}(x) \succ^{(2^n-1)} \alpha_2^{2^n-1}(z)) \\
&= \alpha_1^{2^n-1} \alpha_1^{2^n} \alpha_2^{2^n}(x) \succ (\alpha_1^{2^n-1} \alpha_2^{2^n-1}(y) \succ \alpha_2^{2^n-1} \alpha_2^{2^n-1}(z)) \\
&\quad - \alpha_1^{2^n-1} \alpha_2^{2^n}(y) \succ (\alpha_1^{2^n} \alpha_1^{2^n-1} \alpha_2^{2^n-1}(x) \succ \alpha_2^{2^n-1} \alpha_2^{2^n-1}(z)) \\
&= \alpha_1^{2^n} \alpha_2^{2^n}(x) \succ^{(2^n-1)} (y \succ^{(2^n-1)} z) - \alpha_2^{2^n}(y) \succ^{(2^n-1)} (\alpha_1^{2^n}(x) \succ^{(2^n-1)} z). \quad \square
\end{aligned}$$

*Example 6.8.* There are three-dimensional multiplicative BiHom-Leibniz dendriform algebras  $(A, \prec, \succ, \alpha_1, \alpha_2)$  with multiplications tables for a basis  $\{e_1, e_2, e_3\}$ :

$\prec$	$e_1$	$e_2$	$e_3$	$\succ$	$e_1$	$e_2$	$e_3$
$e_1$	0	$-2e_3$	0	$e_1$	0	$-2e_3$	0
$e_2$	$-2e_3$	$3e_3$	0	$e_2$	$-2e_3$	$3e_3$	0
$e_3$	0	0	0	$e_3$	0	0	0

$$\begin{aligned}
\alpha_1(e_1) &= -\frac{2p}{3}e_1, & \alpha_1(e_2) &= pe_2, & \alpha_1(e_3) &= \frac{p^2}{3}e_3, \\
\alpha_2(e_1) &= -\frac{2q}{3}e_1, & \alpha_2(e_2) &= qe_2, & \alpha_2(e_3) &= \frac{q^2}{3}e_3,
\end{aligned}$$

where  $p$  and  $q$  are parameters in  $\mathbb{K} \setminus \{0\}$ . Then there are BiHom-Leibniz dendriform algebras  $\mathcal{A}_1^n$  and  $\mathcal{A}_2^n$  with multiplications tables respectively:

$\prec^{(n)}$	$e_1$	$e_2$	$e_3$
$e_1$	0	$-2(\frac{-2pq}{3})^n e_3$	0
$e_2$	$-2(\frac{-2pq}{3})^n e_3$	$3(pq)^n e_3$	0
$e_3$	0	0	0

$\succ^{(n)}$	$e_1$	$e_2$	$e_3$
$e_1$	0	$-2(\frac{-2pq}{3})^n e_3$	0
$e_2$	$-2(\frac{-2pq}{3})^n e_3$	$3(pq)^n e_3$	0
$e_3$	0	0	0

$$\begin{aligned}\alpha_1^{n+1}(e_1) &= \left(-\frac{2p}{3}\right)^{n+1}e_1, & \alpha_1^{n+1}(e_2) &= p^{n+1}e_2, & \alpha_1^{n+1}(e_3) &= \left(\frac{p^2}{3}\right)^{n+1}e_3, \\ \alpha_2^{n+1}(e_1) &= \left(-\frac{2q}{3}\right)^{n+1}e_1, & \alpha_2^{n+1}(e_2) &= q^{n+1}e_2, & \alpha_2^{n+1}(e_3) &= \left(\frac{q^2}{3}\right)^{n+1}e_3,\end{aligned}$$

$\prec^{(2^n-1)}$	$e_1$	$e_2$	$e_3$
$e_1$	0	$2\left(\frac{2pq}{3}\right)^{2^n-1}e_3$	0
$e_2$	$2\left(\frac{2pq}{3}\right)^{2^n-1}e_3$	$3(pq)^{2^n-1}e_3$	0
$e_3$	0	0	0

$\succ^{(2^n-1)}$	$e_1$	$e_2$	$e_3$
$e_1$	0	$2\left(\frac{2pq}{3}\right)^{2^n-1}e_3$	0
$e_2$	$2\left(\frac{2pq}{3}\right)^{2^n-1}e_3$	$3(pq)^{2^n-1}e_3$	0
$e_3$	0	0	0

$$\begin{aligned}\alpha_1^{2^n}(e_1) &= \left(\frac{2p}{3}\right)^{2^n}e_1, & \alpha_1^{2^n}(e_2) &= p^{2^n}e_2, & \alpha_1^{2^n}(e_3) &= \left(\frac{p^2}{3}\right)^{2^n}e_3, \\ \alpha_2^{2^n}(e_1) &= \left(\frac{2q}{3}\right)^{2^n}e_1, & \alpha_2^{2^n}(e_2) &= q^{2^n}e_2, & \alpha_2^{2^n}(e_3) &= \left(\frac{q^2}{3}\right)^{2^n}e_3.\end{aligned}$$

Next we define notion of bimodule of BiHomLeibniz dendriform algebra.

**Definition 6.9.** Let  $\mathcal{A} = (A, \prec, \succ, \alpha_1, \alpha_2)$  be a BiHom-Leibniz dendriform algebra, and  $V$  be a linear space. Let  $l_\prec, r_\prec, l_\succ, r_\succ : A \rightarrow gl(V)$ , and  $\beta_1, \beta_2 : V \rightarrow V$  be six linear maps. Then,  $(l_\prec, r_\prec, l_\succ, r_\succ, \beta_1, \beta_2, V)$  is called a bimodule of  $\mathcal{A}$  if for any  $x, y \in A, v \in V$  and  $[x, y] = x \prec y + x \succ y, l = l_\prec + l_\succ, r = r_\prec + r_\succ$ :

$$l_\succ([\alpha_2(x), y])\beta_2(v) = l_\succ(\alpha_1\alpha_2(x)l_\succ(y)v - l_\succ(\alpha_2(y))l_\succ((\alpha_1(x))v), \quad (6.5)$$

$$r_\succ(\alpha_2(y))l(\alpha_2(x))v = l_\succ(\alpha_1\alpha_2(x))r_\succ(y)v - r_\succ(\alpha_1(x) \succ y)\beta_2(v), \quad (6.6)$$

$$r_\succ(\alpha_2(y))r(x)\beta_2(v) = r_\succ(x \succ y)\beta_1\beta_2(v) - l_\succ(\alpha_2(x))r_\succ(y)\beta_1(v), \quad (6.7)$$

$$l_\succ(\alpha_1\alpha_2(x))l_\prec(y)v = l_\prec(\alpha_2(x) \succ y)\beta_2(v) + l_\prec(\alpha_2(y))l(\alpha_1(x))v, \quad (6.8)$$

$$l_\succ(\alpha_1\alpha_2(x))r_\prec(y)v = r_\prec(\alpha_2(y))l_\succ(\alpha_2(x))\beta_1(v) + r_\prec([\alpha_1(x), y])\beta_2(v), \quad (6.9)$$

$$r_\succ(x \prec y)\beta_1\beta_2(v) = r_\prec(\alpha_2(y))r_\succ(x)\beta_2(v) + l_\prec(\alpha_2(x))r(y)\beta_1(v), \quad (6.10)$$

$$l_\prec(\alpha_1\alpha_2(x))r(y)v = l_\prec(\alpha_2(x) \prec y)\beta_2(v) + l_\succ(\alpha_2(y))l_\prec(\alpha_1(x))v, \quad (6.11)$$

$$l_\prec(\alpha_1\alpha_2(x))r(y)v = r_\prec(\alpha_2(y))l_\prec(\alpha_2(x))v + r_\succ(\alpha_1(x) \prec y)\beta_2(v), \quad (6.12)$$

$$r_\prec([x, y])\beta_1\beta_2(v) = r_\prec(\alpha_2(y))r_\prec(x)\beta_2(v) + l_\succ(\alpha_2(x))r_\prec(y)\beta_1(v), \quad (6.13)$$

$$\beta_1(l_\prec(x)v) = l_\prec(\alpha_1(x))\beta_1(v), \quad \beta_1(r_\prec(x)v) = l_\prec(\alpha_1(x))\beta_1(v), \quad (6.14)$$

$$\beta_1(l_\succ(x)v) = l_\succ(\alpha_1(x))\beta_1(v), \quad \beta_1(r_\succ(x)v) = l_\succ(\alpha_1(x))\beta_1(v), \quad (6.15)$$

$$\beta_2(l_\prec(x)v) = l_\prec(\alpha_2(x))\beta_2(v), \quad \beta_2(r_\prec(x)v) = l_\prec(\alpha_2(x))\beta_2(v) \quad (6.16)$$

$$\beta_2(l_\succ(x)v) = l_\succ(\alpha_2(x))\beta_2(v), \quad \beta_2(r_\succ(x)v) = l_\succ(\alpha_2(x))\beta_2(v). \quad (6.17)$$

**Proposition 6.10.** Let  $(l_{\prec}, r_{\prec}, l_{\succ}, r_{\succ}, \beta_1, \beta_2, V)$  be a bimodule of a BiHom-Leibniz dendriform algebra  $(A, \prec, \succ, \alpha_1, \alpha_2)$ . On the direct sum  $A \oplus V$  of the linear spaces, there is a BiHom-Leibniz dendriform algebra structure given for  $x, y \in A, u, v \in V$  by

$$\begin{aligned}(x+u) \prec (y+v) &= x \prec y + l_{\prec}(x)v + r_{\prec}(y)u, \\ (x+u) \succ (y+v) &= x \succ y + l_{\succ}(x)v + r_{\succ}(y)u.\end{aligned}$$

We denote it by  $A \times_{l_{\prec}, r_{\prec}, l_{\succ}, r_{\succ}, \alpha_1, \alpha_2, \beta_1, \beta_2} V$ .

*Proof.* Let  $v_1, v_2, v_3 \in V$  and  $x_1, x_2, x_3 \in A$ . Setting and computing

$$\begin{aligned}& ([(\alpha_2 + \beta_2)(x_1 + v_1), x_2 + v_2]') \succ' (\alpha_2 + \beta_2)(x_3 + v_3) \\ & - (\alpha_1 + \beta_1)(\alpha_2 + \beta_2)(x_1 + v_1) \succ' ((x_2 + v_2) \succ' (x_3 + v_3)) \\ & + ((\alpha_2 + \beta_2)(x_2 + v_2) \succ' ((\alpha_1 + \beta_1)(x_1 + v_1) \succ x_3 + v_3)) \\ & = [\alpha_2(x_1), x_2] \succ \alpha_2(x_3) + l_{\succ}([\alpha_2(x_1), x_2])\beta_2(v_3) \\ & + r_{\succ}(\alpha_2(x_3))l_{\prec}(\alpha_2(x_1))v_2 + r_{\succ}(\alpha_2(x_3))r_{\prec}(x_2)\beta_2(v_3) \\ & + r_{\succ}(\alpha_2(x_3))l_{\succ}(\alpha_2(x_1))v_2 + r_{\succ}(\alpha_2(x_3))r_{\succ}(x_2)\beta_2(v_1) \\ & - \alpha_1\alpha_2(x_1) \succ (x_2 \succ x_3) - l_{\succ}(\alpha_1\alpha_2(x_1))l_{\succ}(x_2)v_3 \\ & - l_{\succ}(\alpha_1\alpha_2(x_1))r_{\succ}(x_3)v_2 - r_{\succ}(x_2 \succ x_3)\beta_1\beta_2(v_1) \\ & + \alpha_2(x_2) \succ (\alpha_1(x_1) \succ x_3) + l_{\succ}(\alpha_2(x_2))l_{\succ}(\alpha_1(x_1))v_3 \\ & + l_{\succ}(\alpha_2(x_2))r_{\succ}(x_3)\beta_1(v_1) + r_{\succ}(\alpha_1(x_1) \succ x_3)\beta_2(v_2).\end{aligned}$$

By (6.5)-(6.7), and (6.2) in  $A$ ,

$$\begin{aligned}& ([(\alpha_2 + \beta_2)(x_1 + v_1), x_2 + v_2]') \succ' (\alpha_2 + \beta_2)(x_3 + v_3) \\ & - (\alpha_1 + \beta_1)(\alpha_2 + \beta_2)(x_1 + v_1) \succ' ((x_2 + v_2) \succ' (x_3 + v_3)) \\ & + ((\alpha_2 + \beta_2)(x_2 + v_2) \succ' ((\alpha_1 + \beta_1)(x_1 + v_1) \succ x_3 + v_3)) = 0. \quad \square\end{aligned}$$

**Proposition 6.11.** Let  $(l_{\prec}, r_{\prec}, l_{\succ}, r_{\succ}, \beta_1, \beta_2, V)$  be a bimodule of a BiHom-Leibniz dendriform algebra  $(A, \prec, \succ, \alpha_1, \alpha_2)$ , and let  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  be the associated BiHom-Leibniz algebra. Then,

- 1)  $(l_{\succ}, r_{\prec}, \beta_2, \beta_1, V)$  and  $(l_{\prec} + l_{\succ}, r_{\prec} + r_{\succ}, \beta_1, \beta_2, V)$  are bimodules of  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$ ;
- 2) for any bimodule  $(l, r, \beta_1, \beta_2, V)$  of  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$ ,  $(0, l, r, 0, \beta_1, \beta_2, V)$  is a bimodule of  $(A, \prec, \succ, \alpha_1, \alpha_2)$ .
- 3)  $(0, l_{\prec} + l_{\succ}, r_{\prec} + r_{\succ}, 0, \beta_1, \beta_2, V)$  and  $(0, l_{\succ}, r_{\prec}, 0, \beta_1, \beta_2, V)$  are bimodules of  $(A, \prec, \succ, \alpha_1, \alpha_2)$ ;
- 4) the BiHom-Leibniz dendriform algebras

$$A \times_{l_{\prec}, r_{\prec}, l_{\succ}, r_{\succ}, \alpha_1, \alpha_2, \beta_1, \beta_2} V \text{ and } A \times_{0, l_{\prec} + l_{\succ}, r_{\prec} + r_{\succ}, 0, \alpha_1, \alpha_2, \beta_1, \beta_2} V$$

have the same associated BiHom-Leibniz algebra  $A \times_{l_{\prec} + l_{\succ}, r_{\prec} + r_{\succ}, \alpha_1, \alpha_2, \beta_1, \beta_2} V$ .

*Example 6.12.* Let  $(A, \prec, \succ, \alpha, \beta)$  be a BiHom-Leibniz dendriform algebra. Suppose that  $(A, [\cdot, \cdot], \alpha, \beta)$  is the associated BiHom-Leibniz algebra. Then,  $(L_{\prec} + L_{\succ}, R_{\prec} + R_{\succ}, \alpha, \beta, A)$  and  $(L_{\succ}, R_{\prec}, \alpha, \beta, A)$  are bimodule of  $(A, [\cdot, \cdot], \alpha, \beta)$ .

**Theorem 6.13.** *Let  $(A, \prec_A, \succ_A, \alpha_1, \alpha_2)$  and  $(B, \prec_B, \succ_B, \beta_1, \beta_2)$  be two BiHom-Leibniz dendriform algebras. If there are linear maps  $l_{\prec_A}, r_{\prec_A}, l_{\succ_A}, r_{\succ_A} : A \rightarrow gl(B)$ , and  $l_{\prec_B}, r_{\prec_B}, l_{\succ_B}, r_{\succ_B} : B \rightarrow gl(A)$  such that  $(l_{\prec_A}, r_{\prec_A}, l_{\succ_A}, r_{\succ_A}, \beta_1, \beta_2, B)$  is a bimodule of  $A$ , and  $(l_{\prec_B}, r_{\prec_B}, l_{\succ_B}, r_{\succ_B}, \alpha_1, \alpha_2, A)$  is a bimodule of  $B$ , and for any  $x, y \in A$ ,  $a, b \in B$  and  $l_A = l_{\prec_A} + l_{\succ_A}, r_A = r_{\prec_A} + r_{\succ_A}, l_B = l_{\prec_B} + l_{\succ_B}, r_B = r_{\prec_B} + r_{\succ_B}$ ,*

$$l_A(\alpha_2(x))a \succ_B \beta_2(b) + l_{\succ_A}(r_B(\alpha_2(x)))\beta_2(b) = l_{\succ_A}(\alpha_1\alpha_2(x))(a \succ b) - \beta_2(a) \succ_B (l_{\succ_A}(\alpha_1(x))b) - r_{\succ_A}(r_{\prec_B}(b)\alpha_1(x))\beta_2(a), \quad (6.18)$$

$$r_A(x)\beta_2(a) \succ_B \beta_2(b) + l_{\succ_A}(l_B(\beta_2(a))x)\beta_2(b) = \beta_1\beta_2(a) \succ_B (l_{\succ_A}(x)b) + r_A(r_B(b)x)\beta_1\beta_2(a) - l_{\succ_A}(\alpha_2(x))(\beta_1(a) \succ_B b), \quad (6.19)$$

$$r_{\succ_A}(\alpha_2(x))([\beta_2(a), b]) = \beta_1\beta_2(a) \succ_B (r_{\succ_A}(x)b) + r_A(r_B(b)x)\beta_1\beta_2(b) - \beta_2(a) \succ_B (r_{\succ_A}(x)\beta_1(b)) - r_A(l_{\succ_B}(\beta_1(b))x)\beta_2(a), \quad (6.20)$$

$$l_{\succ_A}(\alpha_1\alpha_2(x))(a \prec b) = (l_{\succ_A}(\alpha_2(x))a) \prec_B \beta_2(b) + l_{\prec_A}(r_{\prec_B}(a)\alpha_2(x))\beta_2(b) + \beta_2(a) \prec_B (l_A(\alpha_1(x))b) + r_{precA}(r_B(b)\alpha_1(x))\beta_2(a), \quad (6.21)$$

$$\beta_1\beta_2(a) \succ_B (l_{\prec_A}(x)b) + r_{\prec_A}(r_B(b)x)\beta_1\beta_2(a) = (r_{\prec_A}(x)\beta_2(a)) \prec_B \beta_2(b) + l_A(l_{\succ_B}(\beta_2(a))x)\beta_2(b) + l_{\prec_A}(\alpha_2(x))([\beta_1(a), b]), \quad (6.22)$$

$$\beta_1\beta_2(a) \succ_B (r_{\prec_A}(x)b) + r_A(l_B(b)x)\beta_1\beta_2(a) = r_{\prec_A}(\alpha_2(x))(\beta_2(a) \succ_B b) + r_{\prec_A}(l_B(\beta_1(b))x)\beta_2(a) + \beta_2(a) \prec_B (r_A(x)\beta_1(b)), \quad (6.23)$$

$$l_{\prec_A}(\alpha_1\alpha_2(x))([a, b]) = l_{\prec_A}(r_{\prec_B}(a)x)\beta_2(b) + (l_{\prec_A}(\alpha_2(x))a) \prec_B \beta_2(b) + \beta_2(a) \succ_B (l_{\prec_A}(\alpha_1(x))b) + r_A(r_B(b)\alpha_1(x))\beta_2(a), \quad (6.24)$$

$$\beta_1\beta_2(a) \prec_B (l_A(x)b) + r_{\prec_A}(r_B(b)x)\beta_1\beta_2(a) = (r_{\prec_A}(x)\beta_2(a)) \prec_B \beta_2(b) + l_{\prec_A}(l_{\prec_B}(\beta_2(a))x)\beta_2(b) + l_{\succ_A}(\alpha_2(x))(\beta_1(a) \prec b), \quad (6.25)$$

$$\beta_1\beta_2(a) \prec_B (r_A(x)b) + l_{\prec_A}(l_B(b)x)\beta_1\beta_2(a) = r_{\prec_A}(\alpha_2(x))(\beta_2(a) \prec b) + r_{\prec_A}(l_{\prec_B}(\beta_1(a))x)\beta_2(b) + \beta_2(b) \succ_B (r_{\prec_A}(x)\beta_1(a)), \quad (6.26)$$

$$l_B(\beta_2(a))x \succ_A \alpha_2(y) + l_{\succ_B}(r_A(\beta_2(a)))\alpha_2(y) = l_{\succ_B}(\beta_1\beta_2(a))(x \succ y) - \beta_2(x) \succ_A (l_{\succ_B}(\beta_1(a))y) - r_{\succ_B}(r_{\prec_A}(y)\beta_1(a))\alpha_2(x), \quad (6.27)$$

$$r_B(a)\alpha_2(x) \succ_A \alpha_2(y) + l_{\succ_B}(l_A(\alpha_2(x))a)\alpha_2(y) = \alpha_1\alpha_2(x) \succ_A (l_{\succ_B}(a)y) + r_B(r_A(y)a)\alpha_1\alpha_2(x) - l_{\succ_B}(\beta_2(a))(\alpha_1(x) \succ_A y), \quad (6.28)$$

$$r_{\succ_B}(\beta_2(a))([\alpha_2(x), y]) = \alpha_1\alpha_2(x) \succ_A (r_{\succ_B}(a)y) + r_B(r_A(y)a)\alpha_1\alpha_2(y) - \alpha_2(x) \succ_A (r_{\succ_B}(a)\alpha_1(y)) - r_B(l_{\succ_A}(\alpha_1(y))a)\alpha_2(x), \quad (6.29)$$

$$l_{\succ_B}(\beta_1\beta_2(a))(x \prec y) = (l_{\succ_B}(\beta_2(a))x) \prec_A \alpha_2(y) + l_{\prec_B}(r_{\prec_A}(x)\beta_2(a))\alpha_2(y) + \alpha_2(x) \prec_A (l_B(\beta_1(a))y) + r_{precB}(r_A(y)\beta_1(a))\alpha_2(x), \quad (6.30)$$

$$\alpha_1\alpha_2(x) \succ_A (l_{\prec_B}(a)y) + r_{\prec_B}(r_A(y)a)\alpha_1\alpha_2(x) = (r_{\prec_B}(a)\alpha_2(x)) \prec_A \alpha_2(y) + l_B(l_{\succ_A}(\alpha_2(x))a)\alpha_2(y) + l_{\prec_B}(\beta_2(a))([\alpha_1(x), y]), \quad (6.31)$$

$$\alpha_1\alpha_2(x) \succ_A (r_{\prec_B}(a)y) + r_B(l_A(y)a)\alpha_1\alpha_2(x) = r_{\prec_B}(\beta_2(a))(\alpha_2(x) \succ_A y) + r_{\prec_B}(l_A(\alpha_1(y)a)\alpha_2(x) + \alpha_2(x) \prec_A (r_B(a)\alpha_1(y))), \quad (6.32)$$

$$l_{\prec_B}(\beta_1\beta_2(a))([x, y]) = l_{\prec_B}(r_{\prec_A}(x)a)\alpha_2(y) + (l_{\prec_B}(\beta_2(a))x) \prec_A \alpha_2(y) + \alpha_2(x) \succ_A (l_{\prec_B}(\beta_1(a))y) + r_B(r_A(y)\beta_1(a))\alpha_2(x), \quad (6.33)$$

$$\alpha_1\alpha_2(x) \prec_A (l_B(a)y) + r_{\prec_B}(r_A(y)a)\alpha_1\alpha_2(x) = (r_{\prec_B}(a)\alpha_2(x)) \prec_A \alpha_2(y) + l_{\prec_B}(l_{\prec_A}(\alpha_2(x))a)\alpha_2(y) + l_{\succ_B}(\beta_2(a))(\alpha_1(x) \prec y), \quad (6.34)$$

$$\alpha_1\alpha_2(x) \prec_A (r_B(a)y) + l_{\prec_B}(l_A(y)a)\alpha_1\alpha_2(x) = r_{\prec_B}(\beta_2(a))(\alpha_2(x) \prec y)$$

$$+r_{>_B}(l_{<_A}(\alpha_1(x))a)\alpha_2(y) + \alpha_2(y) \succ_A (r_{>_B}(a)\alpha_1(x)), \quad (6.35)$$

then, there is a BiHom-Leibniz dendriform algebra structure on the direct sum  $A \oplus B$  of the underlying linear spaces of  $A$  and  $B$  given for any  $x, y \in A, a, b \in B$  by

$$\begin{aligned} (x+a) \prec (y+b) &= (x \prec_A y + r_{<_B}(b)x + l_{<_B}(a)y) \\ &\quad + (l_{<_A}(x)b + r_{<_A}(y)a + a \prec_B b), \\ (x+a) \succ (y+b) &= (x \succ_A y + r_{>_B}(b)x + l_{>_B}(a)y) \\ &\quad + (l_{>_A}(x)b + r_{>_A}(y)a + a \succ_B b). \end{aligned}$$

*Proof.* The proof is obtained in a similar way as for Theorem 2.11.  $\square$

Let  $A \bowtie_{\substack{l_{<_A}, r_{<_A}, l_{>_A}, r_{>_A}, \beta_1, \beta_2 \\ l_{<_B}, r_{<_B}, l_{>_B}, r_{>_B}, \alpha_1, \alpha_2}} B$  denote this BiHom-Leibniz dendriform algebra.

**Definition 6.14.** Let  $(A, \prec_A, \succ_A, \alpha_1, \alpha_2)$  and  $(B, \prec_B, \succ_B, \beta_1, \beta_2)$  be BiHom-Leibniz dendriform algebras. Suppose there exist linear maps  $l_{<_A}, r_{<_A}, l_{>_A}, r_{>_A} : A \rightarrow gl(B)$ , and  $l_{<_B}, r_{<_B}, l_{>_B}, r_{>_B} : B \rightarrow gl(A)$  such that  $(l_{<_A}, r_{<_A}, l_{>_A}, r_{>_A}, \beta_1, \beta_2)$  is a bimodule of  $A$ , and  $(l_{<_B}, r_{<_B}, l_{>_B}, r_{>_B}, \alpha_1, \alpha_2)$  is a bimodule of  $B$ . If (6.18) - (6.35) are satisfied,  $(A, B, l_{<_A}, r_{<_A}, l_{>_A}, r_{>_A}, \beta_1, \beta_2, l_{<_B}, r_{<_B}, l_{>_B}, r_{>_B}, \alpha_1, \alpha_2)$  is called a matched pair of BiHom-Leibniz dendriform algebras.

**Corollary 6.15.** *If  $(A, B, l_{<_A}, r_{<_A}, l_{>_A}, r_{>_A}, \beta_1, \beta_2, l_{<_B}, r_{<_B}, l_{>_B}, r_{>_B}, \alpha_1, \alpha_2)$  is an arbitrary matched pair of BiHom-Leibniz dendriform algebras, then  $(A, B, l_{<_A} + l_{>_A}, r_{<_A} + r_{>_A}, l_{<_B} + l_{>_B}, r_{<_B} + r_{>_B}, \alpha_1 + \beta_1, \alpha_2 + \beta_2)$  is a matched pair of the associated BiHom-Leibniz algebras  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  and  $(B, \{\cdot, \cdot\}, \beta_1, \beta_2)$ .*

*Proof.* Let  $(A, B, l_{<_A}, r_{<_A}, l_{>_A}, r_{>_A}, \beta_1, \beta_2, l_{<_B}, r_{<_B}, l_{>_B}, r_{>_B}, \alpha_1, \alpha_2)$  be some matched pair of BiHom-Leibniz dendriform algebras  $(A, \prec_A, \succ_A, \alpha_1, \alpha_2)$  and  $(B, \prec_B, \succ_B, \beta_1, \beta_2)$ . In view of Proposition 6.11, the linear maps  $l_{<_A} + l_{>_A}, r_{<_A} + r_{>_A} : A \rightarrow gl(B)$  and  $l_{<_B} + l_{>_B}, r_{<_B} + r_{>_B} : B \rightarrow gl(A)$  are bimodules of the underlying BiHom-Leibniz algebras  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  and  $(B, \{\cdot, \cdot\}, \beta_1, \beta_2)$ , respectively. Therefore, (5.11)-(5.13) are equivalent to (6.18)-(6.26) and (5.14)-(5.16) are equivalent to (6.27)-(6.35).  $\square$

**Definition 6.16.** Let  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  be a BiHom-Leibniz algebra, and  $(l, r, \beta_1, \beta_2, V)$  be a bimodule. A linear map  $T : V \rightarrow A$  is called an  $\mathcal{O}$ -operator associated to  $(l, r, \beta_1, \beta_2, V)$ , if  $T$  satisfies

$$\begin{aligned} \alpha_1 T &= T \beta_2, \quad \alpha_2 T = T \beta_1, \\ [T(u), T(v)] &= T(l(T(u))v + r(T(v))u) \text{ for all } u, v \in V. \end{aligned}$$

*Example 6.17.* Let  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  be a multiplicative BiHom-Leibniz algebra. Then, the identity map  $id$  is an  $\mathcal{O}$ -operator associated to the bimodule  $(L, 0, \alpha_1, \alpha_2)$ .

*Example 6.18.* Let  $\mathcal{A} = (A, [\cdot, \cdot], \alpha, \beta)$  be a multiplicative BiHom-Leibniz algebra. A linear map  $K : A \rightarrow A$  is called a Rota-Baxter operator on  $\mathcal{A}$  of weight zero if  $K$  satisfies

$$\begin{aligned} K \circ \alpha &= \alpha \circ K, \quad K \circ \beta = \beta \circ K, \\ [K(x), K(y)] &= K([K(x), y] + [x, K(y)]), \text{ for all } x, y \in A. \end{aligned}$$

A Rota-Baxter operator on  $\mathcal{A}$  is an  $\mathcal{O}$ -operator associated to the bimodule  $(L, R, \alpha, \beta)$ .



**Theorem 6.19.** *Let  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  be a BiHom-Leibniz algebra, and  $(l, r, \beta_1, \beta_2, V)$  be a bimodule. Let  $T : V \rightarrow A$  be an  $\mathcal{O}$ -operator associated to  $(l, r, \beta_1, \beta_2, V)$ . Then, there exists a BiHom-Leibniz dendriform algebra structure on  $V$  given by*

$$u \prec v = r(T(u))v, \quad u \succ v = l(T(v))u$$

for all  $u, v \in V$ . So, there is an associated BiHom-Leibniz algebra structure on  $V$  given by the equation (4.4), and  $T$  is a homomorphism of BiHom-Leibniz algebras. Moreover,  $T(V) = \{T(v) \mid v \in V\} \subset A$  is a BiHom-Leibniz subalgebra of  $A$ , and there is an induced BiHom-Leibniz dendriform algebra structure on  $T(V)$  given by

$$T(u) \succ T(v) = T(u \succ v), \quad T(u) \prec T(v) = T(u \prec v) \quad (6.36)$$

for all  $u, v \in V$ . Its corresponding associated BiHom-Leibniz algebra structure on  $T(V)$  given by (4.4) is just the BiHom-Leibniz subalgebra structure of  $A$ , and  $T$  is a homomorphism of BiHom-Leibniz dendriform algebras.

*Proof.* For any  $x, y, z \in V$ , we have

$$\begin{aligned} & ([\beta_2(x), y]) \succ \beta_2(z) - \beta_1\beta_2(x) \succ (y \succ z) + \beta_2(y) \succ (\beta_1(x) \succ z) \\ &= (r(T(y))\beta_2(x) + l(T((\beta_2(x))y)) \succ \beta_2(z) - \beta_1\beta_2(x) \succ (l(T(y))z) \\ &\quad + \beta_2(y) \succ (l(T(\beta_1(x))z) \\ &= l(T(r(T(y))\beta_2(x) + l(T(\beta_2(x))y))\beta_2(z) - l(T(\beta_1\beta_2(x)))l(T(y))z \\ &\quad + l(T(\beta_2(y))l(T(\beta_1(x))z) \\ &= l(T(r(T(y))\beta_2(x) + l(\alpha_2(T(x)))y)\beta_2(z) - l(\alpha_1\alpha_2(T(x)))l(T(y))z \\ &\quad + l(\alpha_2(T(y))l(\alpha_1(T(x))z) \\ &= l([T(\beta_2(x)), T(y)]\beta_2(z) - l([T(\beta_2(x)), T(y)]\beta_2(z) = 0. \end{aligned}$$

The two other axioms are checked in a similar way.  $\square$

*Example 6.20.* There is three-dimensional BiHom-Leibniz algebra  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  with multiplications tables for a basis  $\{e_1, e_2, e_3\}$ :

$[\cdot, \cdot]$	$e_1$	$e_2$	$e_3$
$e_1$	0	0	0
$e_2$	0	$e_1$	0
$e_3$	0	0	0

$$\begin{aligned} \alpha_1(e_1) &= e_1, & \alpha_1(e_2) &= -e_2, & \alpha_1(e_3) &= -2e_3, \\ \alpha_2(e_1) &= \frac{1}{4}e_1, & \alpha_2(e_2) &= \frac{1}{2}e_3, & \alpha_2(e_3) &= 2e_3. \end{aligned}$$

Let  $K$  the Rota-Baxter operator of weight zero defined, for the basis  $\{e_1, e_2, e_3\}$ , by

$$K(e_1) = \frac{\lambda_1}{2}e_1, \quad K(e_2) = \lambda_2e_1 + \lambda_1e_2 + \lambda_3e_3, \quad K(e_3) = \lambda_4e_1 + \lambda_5e_3,$$

where  $\lambda_i$  are parameters in  $\mathbb{K} \setminus \{0\}$ . Now, using the previous Theorem, there is a BiHom-Leibniz dendriform algebra structure on  $A$  with the multiplications " $\prec$ " and " $\succ$ " are given by  $x \prec y = [y, K(x)]$ ,  $x \succ y = [K(y), x]$  that is

$\prec$	$e_1$	$e_2$	$e_3$
$e_1$	0	0	0
$e_2$	0	$\lambda_1 e_1$	0
$e_3$	0	0	0

$\succ$	$e_1$	$e_2$	$e_3$
$e_1$	0	0	0
$e_2$	0	$\lambda_1 e_1$	0
$e_3$	0	0	0

$$\begin{aligned} \alpha_1(e_1) &= e_1, & \alpha_1(e_2) &= -e_2, & \alpha_1(e_3) &= -2e_3, \\ \alpha_2(e_1) &= \frac{1}{4}e_1, & \alpha_2(e_2) &= \frac{1}{2}e_3, & \alpha_2(e_3) &= 2e_3. \end{aligned}$$

**Corollary 6.21.** *Let  $T : V \rightarrow A$  be an  $\mathcal{O}$ -operator on a BiHom-Leibniz algebra  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$  with respect to a representation  $(l, r, \beta_1, \beta_2, V)$ . Let  $[u, v]_C = u \prec v + u \succ v$ , for all  $u, v \in V$ . Then,  $T$  is a morphism from the BiHom-Leibniz algebra  $(V, [\cdot, \cdot]_C, \beta_1, \beta_2)$  to  $(A, [\cdot, \cdot], \alpha_1, \alpha_2)$ .*

**Corollary 6.22.** *Let  $(A, [\cdot, \cdot], \alpha, \beta)$  be a multiplicative BiHom-Leibniz algebra. There is a compatible multiplicative BiHom-Leibniz dendriform algebra structure on  $A$  if and only if there exists an invertible  $\mathcal{O}$ -operator of  $(A, [\cdot, \cdot], \alpha, \beta)$ .*

*Proof.* In fact, if the homomorphism  $T$  is an invertible  $\mathcal{O}$ -operator associated to a bimodule  $(l, r, \alpha, \beta, V)$ , then the compatible multiplicative BiHom-Leibniz dendriform algebra structure on  $A$  is given for all  $x, y \in A$  by

$$x \prec y = T(l(x)T^{-1}(y)), \quad x \succ y = T(r(y)T^{-1}(x)).$$

Conversely, let  $(A, \prec, \succ, \alpha, \beta)$  be a multiplicative BiHom-Leibniz dendriform algebra, and  $(A, [\cdot, \cdot], \alpha, \beta)$  be its associated BiHom-Leibniz algebra. Then, the identity map  $id$  is an  $\mathcal{O}$ -operator associated to the bimodule  $(L_\succ, R_\prec, \alpha, \beta)$  of  $(A, [\cdot, \cdot], \alpha, \beta)$ .  $\square$

## References

- [1] Abdaoui, E., Mabrouk, S., Makhlouf, A.: Rota-Baxter operators on pre-Lie superalgebras, Bulletin of the Malaysian Math. Sci. Soc., 1-40 (2017)
- [2] Aizawa, N., Sato, H.:  $q$ -deformation of the Virasoro algebra with central extension, Phys. Lett. B **256**(2), 185-190 (1991) (Hiroshima Univ. preprint, HUPD-9012 (1990))
- [3] Ammar, F., Ejbehi, Z., Makhlouf, A.: Cohomology and deformations of Hom-algebras, J. Lie Theory **21**(4), 813-836 (2011). (arXiv:1005.0456 [math.RA])
- [4] Armakan A., Farhangdoost, M. R.: Geometric aspects of extensions of hom-Lie superalgebras, Int. J. Geom. Methods Mod. Phys. **14**(06), 1750085 (2017)
- [5] Armakan, A., Silvestrov, S.: Color Hom-Lie algebras, color Hom-Leibniz algebras and color omni-Hom-Lie algebras, arXiv:2010.06160 [math.RA], 17pp (2020)
- [6] Attan, S.: Representations and relative Rota-Baxter operators of Hom-Leibniz Poisson algebras, arXiv:2105.06310 [math.RA], 20pp (2021)
- [7] Attan, S., Laraiedh, I.: Construtions and bimodules of BiHom-alternative and BiHom-Jordan algebras, arXiv:2008.07020 [math.RA] (2020)
- [8] Bakayoko, I.: Laplacian of Hom-Lie quasi-bialgebras, International Journal of Algebra, **8**(15), 713-727 (2014)

- [9] Bakayoko, I.:  $L$ -modules,  $L$ -comodules and Hom-Lie quasi-bialgebras, African Diaspora Journal of Mathematics, **17**, 49-64 (2014)
- [10] Bakayoko, I., Banagoura, M.: Bimodules and Rota-Baxter relations. J. Appl. Mech. Eng. **4**(5) (2015)
- [11] Bakayoko, I., Silvestrov, S.: Multiplicative  $n$ -Hom-Lie color algebras, In: Silvestrov, S., Malyarenko, A., Rančić, M. (Eds.), Algebraic Structures and Applications, Springer Proceedings in Mathematics and Statistics **317**, Ch. 7, 159-187, Springer (2020). (arXiv:1912.10216[math.QA] (2019))
- [12] Bakayoko, I., Silvestrov, S.: Hom-left-symmetric color dialgebras, Hom-tridendriform color algebras and Yau's twisting generalizations, Afr. Mat. **32**, 941-958 (2021). (arXiv:1912.01441[math.RA])
- [13] Benayadi, S., Makhlouf, A.: Hom-Lie algebras with symmetric invariant nondegenerate bilinear forms, J. Geom. Phys. **76**, 38-60 (2014)
- [14] Ben Abdeljelil, A., Elhamdadi, M., Kaygorodov, I., Makhlouf, A.: Generalized Derivations of  $n$ -BiHom-Lie algebras, In: Silvestrov, S., Malyarenko, A., Rančić, M. (Eds.), Algebraic Structures and Applications, Springer Proceedings in Mathematics and Statistics **317**, Ch. 4, 81-97, Springer (2020). (arXiv:1901.09750[math.RA] (2019))
- [15] Ben Hassine, A., Chtioui, T., Mabrouk, S., Ncib, O.: Cohomology and linear deformation of BiHom-left-symmetric algebras, 19 pp, arXiv:1907.06979 [math.RA] (2019)
- [16] Chaichian, M., Ellinas, D., Popowicz, Z.: Quantum conformal algebra with central extension, Phys. Lett. B, **248**, 95-99 (1990)
- [17] Chaichian, M., Kulish, P., Lukierski, J.:  $q$ -Deformed Jacobi identity,  $q$ -oscillators and  $q$ -deformed infinite-dimensional algebras, Phys. Lett. B, **237**, 401-406 (1990)
- [18] Chaichian, M., Popowicz, Z., Prešnajder, P.:  $q$ -Virasoro algebra and its relation to the  $q$ -deformed KdV system, Phys. Lett. B, **249**, 63-65 (1990)
- [19] Chaichian, M., Isaev, A. P., Lukierski, J., Popowicz, Z.: Prešnajder, P:  $q$ -Deformations of Virasoro algebra and conformal dimensions, Phys. Lett. B, **262**(1), 32-38 (1991)
- [20] Cheng, Y., Su, Y.: (Co)homology and universal central extension of Hom-Leibniz algebras, Acta Math. Sin. (Engl. Ser.) **27**, 813-830 (2011)
- [21] Chtioui, T., Mabrouk, S., Makhlouf, A.: BiHom-alternative, BiHom-Malcev and BiHom-Jordan algebras." Rocky Mountain J. Math. **50**(1), 69-90 (2020)
- [22] Chtioui, T., Mabrouk, S., Makhlouf, A.: BiHom-pre-alternative algebras and BiHom-alternative quadri-algebras, Bull. Math. Soc. Sci. Math. Roumanie. **63** (111)(1), 3-21 (2020)
- [23] Dassoundo, M. L., Silvestrov, S.: Nearly associative and nearly Hom-associative algebras and bialgebras, arXiv:2101.12377 [math.RA] (2021)
- [24] Daskaloyannis, C.: Generalized deformed Virasoro algebras, Modern Phys. Lett. A, **7**(09), 809-816 (1992)
- [25] Graziani, G., Makhlouf, A., Menini, C., Panaite, F.: BiHom-Associative Algebras, BiHom-Lie Algebras and BiHom-Bialgebras, SIGMA **11**(086), 34 pp (2015)

- [26] Ebrahimi-Fard, K., Guo, L.: Rota-Baxter algebras and dendriform algebras, *J. Pure Appl. Algebra*, **212**(2), 320–339 (2008)
- [27] Graziani, G., Makhlouf, A., Menini, C., Panaite, F.: BiHom-Associative Algebras, BiHom-Lie Algebras and BiHom-Bialgebras, *SIGMA Symmetry Integrability Geom. Methods Appl.* 11, 086, 34 pp (2015)
- [28] Gürsey, F., Tze, C.-H.: On the role of division, Jordan and related algebras in particle physics, *World Scientific* (1996)
- [29] Hartwig, J. T., Larsson, D., Silvestrov, S. D.: Deformations of Lie algebras using  $\sigma$ -derivations, *J. Algebra*, **295**(2), 314-361 (2006) (Preprint in Mathematical Sciences 2003:32, LUTFMA-5036-2003, Centre for Mathematical Sciences, Department of Mathematics, Lund Institute of Technology, 52 pp. (2003))
- [30] Hu, N.:  $q$ -Witt algebras,  $q$ -Lie algebras,  $q$ -holomorph structure and representations, *Algebra Colloq.* **6**(1), 51-70 (1999)
- [31] Hassanzadeh, M., Shapiro, I., Sütlü, S.: Cyclic homology for Hom-associative algebras, *J. Geom. Phys.* **98**, 40-56 (2015)
- [32] Hounkonnou, M. N., Dassoundo, M. L.: Center-symmetric Algebras and Bialgebras: Relevant Properties and Consequences. In: Kielanowski P., Ali S., Bieliavsky P., Odziejewicz A., Schlichenmaier M., Voronov T. (eds) *Geometric Methods in Physics. Trends in Mathematics.* 2016, pp. 281-293. Birkhäuser, Cham (2016)
- [33] Hounkonnou, M. N., Houndedji, G. D., Silvestrov, S.: Double constructions of biHom-Frobenius algebras, 45pp, arXiv:2008.06645 [math.QA] (2020)
- [34] Hounkonnou, M. N., Dassoundo, M. L.: Hom-center-symmetric algebras and bialgebras. arXiv:1801.06539 (2018)
- [35] Jacobson, N.: Structure of alternative and Jordan bimodules, *Osaka J. Math.* **6**, 1-71 (1954)
- [36] Kassel, C.: Cyclic homology of differential operators, the Virasoro algebra and a  $q$ -analogue, *Comm. Math. Phys.* **146**(2), 343-356 (1992)
- [37] Kitouni, A., Makhlouf, A., Silvestrov, S.: On  $n$ -ary generalization of BiHom-Lie algebras and BiHom-associative algebras, In: Silvestrov, S., Malyarenko, A., RancicRančić, M. (Eds.), *Algebraic Structures and Applications, Springer Proceedings in Mathematics and Statistics* **317**, Springer, Ch. 5, 99-126 (2020)
- [38] Laraiedh, I.: Bimodules and matched pairs of noncommutative BiHom-(pre)-Poisson algebras, arXiv:2102.11364 [math.RA] (2021)
- [39] Larsson, D., Sigurdsson, G., Silvestrov, S. D.: Quasi-Lie deformations on the algebra  $\mathbb{F}[t]/(t^N)$ , *J. Gen. Lie Theory Appl.* **2**, 201-205 (2008)
- [40] Larsson, D., Silvestrov, S. D.: Quasi-Hom-Lie algebras, central extensions and 2-cocycle-like identities, *J. Algebra* **288**, 321-344 (2005) (Preprints in Mathematical Sciences 2004:3, LUTFMA-5038-2004, Centre for Mathematical Sciences, Department of Mathematics, Lund Institute of Technology, Lund University (2004))
- [41] Larsson, D., Silvestrov, S. D.: Quasi-Lie algebras, In: J. Fuchs, J. Mickelsson, G. Rozemnblioum, A. Stolin, A. Westerberg, *Noncommutative Geometry and Representation Theory in Mathematical Physics, Contemp. Math.* **391**, Amer. Math.

- Soc., Providence, RI, 241-248 (2005) (Preprints in Mathematical Sciences 2004:30, LUTFMA-5049-2004, Centre for Mathematical Sciences, Department of Mathematics, Lund Institute of Technology, Lund University (2004))
- [42] Larsson, D., Silvestrov, S. D.: Graded quasi-Lie algebras, Czechoslovak J. Phys. **55**(11), 1473-1478 (2005)
- [43] Larsson, D., Silvestrov, S. D.: Quasi-deformations of  $sl_2(\mathbb{F})$  using twisted derivations, Comm. Algebra, **35**(12), 4303-4318 (2007) (Preprint in Mathematical Sciences 2004:26, LUTFMA-5047-2004, Centre for Mathematical Sciences, Lund Institute of Technology, Lund University (2004). arXiv:math/0506172 [math.RA] (2005))
- [44] Larsson, D., Sigurdsson, G., Silvestrov, S. D.: Quasi-Lie deformations on the algebra  $\mathbb{F}[t]/(t^N)$ , J. Gen. Lie Theory Appl. **2**(3), 201-205 (2008)
- [45] Larsson, D., Silvestrov, S. D.: On generalized  $N$ -complexes coming from twisted derivations, In: Silvestrov, S., Paal, E., Abramov, V., Stolin, A. (Eds.), Generalized Lie Theory in Mathematics, Physics and Beyond, Springer-Verlag, Ch. 7, 81-88 (2009)
- [46] Liu, K. Q.: Quantum central extensions, C. R. Math. Rep. Acad. Sci. Canada **13**(4), 135-140 (1991)
- [47] Liu, K. Q.: Characterizations of the quantum Witt algebra, Lett. Math. Phys. **24**(4), 257-265 (1992)
- [48] Liu, K. Q.: The quantum Witt algebra and quantization of some modules over Witt algebra, PhD Thesis, Department of Mathematics, University of Alberta, Edmonton, Canada (1992)
- [49] Liu, L., Makhlouf, A., Menini, C., Panaite, F.: Rota-Baxter operators on BiHom-associative algebras and related structures, 27pp, arXiv:1703.07275 [math.RA] (2017)
- [50] Liu, Y., Chen, L., Ma, Y.: Hom-Nijenhuis operators and  $T^*$ -extensions of hom-Lie superalgebras, Linear Algebra Appl. **439**(7), 2131-2144 (2013)
- [51] Loday, J.-L.: Dialgebras, In: Loday, J.-L., Frabetti, A., Chapoton, F., Goichot, F., Dialgebras and related operads, Lecture Notes in Math. **1763**, Springer, Berlin, 7-66 (2001). (Prépublication de l'Inst de Recherche Math. Avancée (Strasbourg), **14**, 61 pp (1999))
- [52] Loday, J.-L., Ronco, M.: Trialgebras and families of polytopes, In: Goerss, P., Priddy, S., Homotopy Theory: relations with Algebraic Geometry, Group Cohomology, and Algebraic K-Theory, Contemp. Math. **346**, Amer. Math. Soc., Providence, RI, 369-673 (2004)
- [53] Ma, T., Makhlouf, A., Silvestrov, S.: Curved  $\mathcal{O}$ -operator systems, 17pp, arXiv:1710.05232 [math.RA] (2017)
- [54] Ma, T., Makhlouf, A., Silvestrov, S.: Rota-Baxter bisystems and covariant bialgebras, 30 pp, arXiv:1710.05161[math.RA] (2017)
- [55] Ma, T., Makhlouf, A., Silvestrov, S.: Rota-Baxter cosystems and coquasitriangular mixed bialgebras, J. Algebra Appl. **20**(04) 2150064 (2021)

- [56] Ma, T., Zheng, H.: Some results on Rota-Baxter monoidal Hom-algebras, *Results Math.* **72** (1-2), 145-170 (2017)
- [57] Mabrouk, S., Ncib, O., Silvestrov, S.: Generalized Derivations and Rota-Baxter Operators of  $n$ -ary Hom-Nambu Superalgebras, *Adv. Appl. Clifford Algebras*, **31**, 32, (2021). (arXiv:2003.01080[math.QA])
- [58] Makhlouf, A.: Hom-dendriform algebras and Rota-Baxter hom-algebras, In: C. Bai, L. Guo, J.-L. Loday, (eds.), *Operads and Universal Algebra*, Nankai Ser. Pure Appl. Math. Theoret. Phys., **9**, World Sci. Publ. 147-171 (2012)
- [59] Makhlouf, A.: Hom-alternative algebras and Hom-Jordan algebras, *Int. Elect. Journ. of Alg.*, **8**, 177-190 (2010). (arXiv:0909.0326 (2009))
- [60] Makhlouf, A.: Paradigm of nonassociative Hom-algebras and Hom-superalgebras, in: J. Carmona Tapia, A. Morales Campoy, A. M. Peralta Pereira, M. I. Ramirez lvarez (Eds.), *Proceedings of Jordan Structures in Algebra and Analysis Meeting*, Publishing house: Circulo Rojo, 145-177 (2009)
- [61] Makhlouf, A., Silvestrov, S. D.: Hom-algebra structures, *J. Gen. Lie Theory Appl.* **2**(2), 51-64 (2008) (Preprints in Mathematical Sciences 2006:10, LUTFMA-5074-2006, Centre for Mathematical Sciences, Department of Mathematics, Lund Institute of Technology, Lund University (2006))
- [62] Makhlouf, A., Silvestrov, S.: Hom-Lie admissible Hom-coalgebras and Hom-Hopf algebras, In: Silvestrov, S., Paal, E., Abramov, V., Stolin, A. (Eds.), *Generalized Lie Theory in Mathematics, Physics and Beyond*, Springer-Verlag, Berlin, Heidelberg, Ch. 17, 189-206 (2009) (Preprints in Mathematical Sciences, Lund University, Centre for Mathematical Sciences, Centrum Scientiarum Mathematicarum (2007:25) LUTFMA-5091-2007 and in arXiv:0709.2413 [math.RA] (2007))
- [63] Makhlouf, A., Silvestrov, S. D.: Hom-algebras and Hom-coalgebras, *J. Algebra Appl.* **9**(4), 553-589 (2010) (Preprints in Mathematical Sciences, Lund University, Centre for Mathematical Sciences, Centrum Scientiarum Mathematicarum, (2008:19) LUTFMA-5103-2008. arXiv:0811.0400 [math.RA] (2008))
- [64] Makhlouf, A., Silvestrov, S.: Notes on 1-parameter formal deformations of Hom-associative and Hom-Lie algebras, *Forum Math.* **22**(4), 715-739 (2010) (Preprints in Mathematical Sciences, Lund University, Centre for Mathematical Sciences, Centrum Scientiarum Mathematicarum, (2007:31) LUTFMA-5095-2007. arXiv:0712.3130v1 [math.RA] (2007))
- [65] Makhlouf, A., Yau, D.: Rota-Baxter Hom-Lie admissible algebras, *Comm. Alg.*, **23**(3), 1231-1257 (2014)
- [66] Mandal, A., Mishra, S. K.: On Hom-Gerstenhaber algebras and Hom-Lie algebroids, *J. Geom. Phys.* **133**, 287-302 (2018)
- [67] Mishra, S. K., Silvestrov, S.: A review on Hom-Gerstenhaber algebras and Hom-Lie algebroids, In: Silvestrov S., Malyarenko A., Rančić, M. (eds), *Algebraic Structures and Applications*, Springer Proceedings in Mathematics and Statistics, **317**, Springer, Ch. 11, 285-315 (2020)
- [68] Richard, L., Silvestrov, S. D.: Quasi-Lie structure of  $\sigma$ -derivations of  $\mathbb{C}[t^{\pm 1}]$ , *J. Algebra* **319**(3), 1285-1304 (2008) (arXiv:math/0608196[math.QA] (2006). Preprints

in mathematical sciences (2006:12), LUTFMA-5076-2006, Centre for Mathematical Sciences, Lund University (2006))

- [69] Richard, L., Silvestrov, S. D.: A note on quasi-Lie and Hom-Lie structures of  $\sigma$ -derivations of  $\mathbb{C}[z_1^{\pm 1}, \dots, z_n^{\pm 1}]$ , In: Silvestrov, S., Paal, E., Abramov, V., Stolin, A. (Eds.), *Generalized Lie Theory in Mathematics, Physics and Beyond*, Springer-Verlag, Ch. 22, 257-262, (2009)
- [70] Saadaou, N, Silvestrov, S.: On  $(\lambda, \mu, \gamma)$ -derivations of BiHom-Lie algebras, arXiv:2010.09148 [math.RA], (2020)
- [71] Sheng, Y.: Representations of Hom-Lie algebras, *Algebr. Represent. Theory* **15**, 1081-1098 (2012)
- [72] Sheng, Y., Bai, C.: A new approach to Hom-Lie bialgebras, *J. Algebra*, **399**, 232-250 (2014)
- [73] Sheng, Y., Tang, R.: Leibniz bialgebras, relative Rota-Baxter operators and the classical Leibniz Yang-Baxter equation, arXiv:1902.03033 [math.RA], 28pp (2017)
- [74] Sheng, Y., Xiong, Z.: On Hom-Lie algebras, *Linear and Multilinear Algebra*, **63**(12), 2379-2395 (2015)
- [75] Sigurdsson, G., Silvestrov, S. D.: Graded quasi-Lie algebras of Witt type, *Czechoslovak J. Phys.* **56**(10/11), 1287-1291 (2006)
- [76] Sigurdsson, G., Silvestrov, S.: Lie color and Hom-Lie algebras of Witt type and their central extensions, In: Silvestrov, S., Paal, E., Abramov, V., Stolin, A. (Eds.), *Generalized Lie Theory in Mathematics, Physics and Beyond*, Springer-Verlag, Berlin, Heidelberg, Ch. 21, 247-255 (2009)
- [77] Silvestrov, S.: Paradigm of quasi-Lie and quasi-Hom-Lie algebras and quasi-deformations, In "New techniques in Hopf algebras and graded ring theory", K. Vlaam. Acad. Belgie Wet. Kunsten (KVAB), Brussels, 165-177 (2007)
- [78] Silvestrov, S., Zargeh, C.: HNN-extension of involutive multiplicative Hom-Lie algebras, arXiv:2101.01319 [math.RA] (2021)
- [79] Sun, Q.: On Hom-Prealternative Bialgebras, *Algebr. Represent. Theor.* **19**, 657-677 (2016)
- [80] Wang, C., Zhang, Q., Wei, Z.: Hom-Leibniz superalgebras and hom-Leibniz Poisson superalgebras, *Hacettepe J. Math. and Statistics*, **44**(5), 1163-1179 (2015)
- [81] Yau D.: The Hom-Yang-Baxter equation, Hom-Lie algebras and quasi-triangular bialgebras, *J. Phys. A.: Math. Theor.* **42**(16), 165-202 (2009)
- [82] Yau, D.: Enveloping algebras of Hom-Lie algebras, *J. Gen. Lie Theory Appl.* **2**(2), 95-108 (2008). (arXiv:0709.0849 [math.RA] (2007))
- [83] Yau, D.: Module Hom-algebras, arXiv:0812.4695[math.RA] (2008)
- [84] Yau, D.: Hom-bialgebras and comodule Hom-algebras, *Int. Electron. J. Algebra* **8**, 45-64 (2010). (arXiv:0810.4866[math.RA] (2008))
- [85] Yau, D.: Hom-algebras and homology, *J. Lie Theory* **19**(2), 409-421 (2009)
- [86] Yau, D.: Hom-Malcev, Hom-alternative, and Hom-Jordan algebras, *Int. Electron. J. Algebra*, **11**, 177-217 (2012)