

A NOTE ON SET THEORETICAL SOLUTIONS OF THE YANG-BAXTER EQUATION WITH TRIVIAL RETRACTION

PABLO ZADUNAISKY

ABSTRACT. We show that every set theoretical solution to the YBE whose retraction is a flip linearizes to a twist of the flip solution by roots of unity. This generalizes a result of Gateva-Ivanova and Majid. To prove the result we use a new invariant associated to a solution, its Lie algebra. We show also that a solution retracts to a flip solution if and only if its Lie algebra is abelian.

1. Introduction

A set theoretical solution to the Yang-Baxter equation is a pair (X, r) where X is a set, and $r : X \times X \rightarrow X \times X$ is a map satisfying

$$r \times \text{Id}_X \circ \text{Id}_X \times r \circ r \times \text{Id}_X = \text{Id}_X \times r \circ r \times \text{Id}_X \circ \text{Id}_X \times r$$

Their study was suggested by Drinfeld in [Dri92], and begun in earnest with the seminal works of Gateva-Ivanova and van den Bergh [GIvdB98] and Etingof, Schedler and Soloviev [ESS99]. To study these solutions one usually constructs associated algebraic objects, such as groups, rings, Hopf algebras (all introduced in the cited papers) and more recently homology groups [LV17], braces [Rum07], and skew-braces [GV17], among many others. For an overview of the literature and open problems in the area and its connections to other areas of mathematics we refer the reader to [Ven19].

The space of all set theoretical solutions of the YBE is vast and one usually focuses on subfamilies to avoid drowning in it. A typical result will prove that the nice combinatorial properties of the solutions are reflected in nice algebraic properties of the associated objects. Common families to study are involutive solutions, or solutions of small size, etc. This paper is concerned with a subfamily of solutions which is close to trivial in the sense that its retraction is a flip solution (see section 2 for definitions). We first show that this is reflected in the structure of the linearization of the solution: it must be a flip solution twisted by roots of unity. This result was already observed by Gateva-Ivanova and Majid for involutive square-free solutions in [GIM11, Theorem 4.9]. We then introduce a new algebraic invariant, the Lie algebra associated to a solution, and show that a solution has trivial retraction if and only if its associated Lie algebra is abelian.

The author is a CONICET fellow and a postdoctoral researcher at Jacobs University. This work was supported in part by the projects UBACYT 20020170100613BA, PICT-2019-00099 and DFG PE 980/8-1.

Notation All vector spaces and tensor products are over \mathbb{C} .

2. Generalities on solutions

2.1. Quadratic sets. A *quadratic set* is a pair (X, r) where X is a set and $r : X \times X \rightarrow X \times X$ is a map; its cardinality is the cardinality of X . For a fixed quadratic set (X, r) and $x, y \in X$ we denote by $\mathcal{L}_x, \mathcal{R}_y$ the functions defined implicitly by

$$r(x, y) = (\mathcal{L}_x(y), \mathcal{R}_y(x)).$$

It is sometimes useful to see \mathcal{L}_x and \mathcal{R}_y as defining actions of X on itself, so we introduce the alternative notation

$$r(x, y) = (x \triangleright y, x \triangleleft y)$$

We say r is *non-degenerate* if the maps \mathcal{L}_x and \mathcal{R}_y are bijective for each $x \in X$. We say that it is *involutive* if $r^2 = \text{Id}_X$. We say it is *square-free* if $r(x, x) = (x, x)$ for all $x \in X$.

Set $r_1 = r \times \text{Id}_X$ and $r_2 = \text{Id}_X \times r$, which are functions of $X \times X \times X$ to itself. We say (X, r) is a solution to the set theoretical Yang-Baxter equation, or to be brief just a *solution*, if $r_1 \circ r_2 \circ r_1 = r_2 \circ r_1 \circ r_2$.

2.2. Examples. Suppose X is any finite set. If we choose arbitrary functions $f, g : X \rightarrow X$ then $r : X \times X \rightarrow X \times X$ given by $r(x, y) = (f(y), g(x))$ is a quadratic set. It is a solution if and only if f and g commute, it is non degenerate if and only if both f and g are bijective, and it is involutive if and only if $g = f^{-1}$. These solutions are known as *permutation solutions*. The *trivial* solution on X is the flip map, which is a permutation solution with $f = g = \text{Id}_X$.

We will use the following two examples in the sequel. Take $s : \llbracket 2 \rrbracket \times \llbracket 2 \rrbracket \rightarrow \llbracket 2 \rrbracket \times \llbracket 2 \rrbracket$ given by

$$\begin{array}{c|cc} s & 1 & 2 \\ \hline 1 & (2, 2) & (1, 2) \\ 2 & (2, 1) & (1, 1) \end{array}$$

This solution is involutive and non-degenerate but not square-free.

Now take $t : \llbracket 4 \rrbracket \times \llbracket 4 \rrbracket \rightarrow \llbracket 4 \rrbracket \times \llbracket 4 \rrbracket$ to be the permutation solution given by $f = (1, 2)(3, 4); g = (1, 2)$. Since g is not the inverse of f this solution is not involutive. However since it is a permutation solution, its retracion is trivial.

2.3. Associated vector space. From this point on (X, r) denotes a quadratic set. We denote by V the \mathbb{C} -span of X , and denote by $R : V \otimes V \rightarrow V \otimes V$ the map induced by r . We consider V to be a Hermitian space with the inner product uniquely determined by the fact that X is an orthogonal basis. Given a set $S \subset V$ we denote by p_S the orthogonal projection to the space generated by S .

2.4. Retractions. We define an equivalence relation in X denoted by \sim , where $x \sim y$ if and only if for each $z \in X$ we have $\mathcal{L}_x = \mathcal{L}_y$ and $\mathcal{R}_x = \mathcal{R}_y$. We denote by \bar{x}

the equivalence class of x and by a slight abuse of notation we write $\mathcal{L}_{\bar{x}} = \mathcal{L}_x$ and $\mathcal{R}_{\bar{x}} = \mathcal{R}_x$. We also denote by $\mathcal{L}_{\bar{x}}$ and $\mathcal{R}_{\bar{x}}$ the linear extension of these maps to V . We denote by \bar{X} the quotient X/\sim . The following lemma is standard. It was proved for involutive solutions (which are automatically non-degenerate) by Etingof, Schedler and Soloviev [ESS99, Subsection 3.2].

Lemma 2.1. *Suppose (X, r) is a quadratic set. Then (X, r) is a solution if and only if the following equalities hold in $\text{End}(V)$*

$$\begin{aligned}\mathcal{L}_{\bar{x}} \circ \mathcal{L}_{\bar{y}} &= \mathcal{L}_{\bar{x} \triangleright \bar{y}} \circ \mathcal{L}_{\bar{x} \triangleleft \bar{y}}; \\ \mathcal{R}_{\bar{z}} \circ \mathcal{R}_{\bar{y}} &= \mathcal{R}_{\bar{y} \triangleleft \bar{z}} \circ \mathcal{R}_{\bar{y} \triangleright \bar{z}}; \\ \mathcal{R}_{\overline{(x \triangleleft y) \triangleright z}} \circ \mathcal{L}_{\bar{x}} \circ p_{\bar{y}} &= \mathcal{L}_{\overline{x \triangleleft (y \triangleright z)}} \circ \mathcal{R}_{\bar{z}} \circ p_{\bar{y}}.\end{aligned}$$

If (X, r) is bijective then r induces a map $\bar{r} : \bar{X} \times \bar{X} \longrightarrow \bar{X} \times \bar{X}$ and (\bar{X}, \bar{r}) , which is also a solution.

Proof. For each $x, y, z \in X$ we have

$$\begin{aligned}r_1 r_2 r_1(x, y, z) &= (\mathcal{L}_{\bar{x} \triangleright \bar{y}}(\mathcal{L}_{\bar{x} \triangleleft \bar{y}}(z)), \mathcal{R}_{\overline{(x \triangleleft y) \triangleright z}}(\mathcal{L}_{\bar{x}}(y)), \mathcal{R}_{\bar{z}}(\mathcal{R}_{\bar{y}}(x))); \\ r_2 r_1 r_2(x, y, z) &= (\mathcal{L}_{\bar{x}}(\mathcal{L}_{\bar{y}}(z)), \mathcal{L}_{\overline{x \triangleleft (y \triangleright z)}}(\mathcal{R}_{\bar{z}}(y)), \mathcal{R}_{\bar{y} \triangleleft \bar{z}}(\mathcal{R}_{\bar{y} \triangleright \bar{z}}(x))).\end{aligned}$$

Since $y = p_{\bar{y}}(y)$ the equalities in the statement are equivalent to the fact that r is a solution to the YBE.

Now if $\bar{x} = \bar{x}'$ then $\bar{x} \triangleright \bar{y} = \bar{x}' \triangleright \bar{y}$. Thus

$$\mathcal{L}_{\bar{x} \triangleleft \bar{y}} = \mathcal{L}_{\bar{x}} \circ \mathcal{L}_{\bar{y}} \circ \mathcal{L}_{\bar{x} \triangleright \bar{y}}^{-1} = \mathcal{L}_{\bar{x}'} \circ \mathcal{L}_{\bar{y}} \circ \mathcal{L}_{\bar{x}' \triangleright \bar{y}}^{-1} = \mathcal{L}_{\bar{x}' \triangleleft \bar{y}}$$

and similarly $\mathcal{R}_{\bar{x} \triangleleft \bar{y}} = \mathcal{R}_{\bar{x}' \triangleleft \bar{y}}$. Thus $\bar{x} \triangleleft \bar{y}$ depends only on \bar{y} and \bar{x} . Similarly $\bar{y} \triangleright \bar{x} = \bar{y} \triangleright \bar{x}'$ and this result only depends on \bar{y} and \bar{x} . This says that the map $\bar{r} : \bar{X} \times \bar{X} \longrightarrow \bar{X} \times \bar{X}$ given by $\bar{r}(\bar{x}, \bar{y}) = (\bar{x} \triangleright \bar{y}, \bar{x} \triangleleft \bar{y})$ is well-defined. Since r is a solution, so is \bar{r} . \square

The solution $\text{Ret}(X, r) = (\bar{X}, \bar{r})$ is called the *retraction* of (X, r) . We say that a solution is retractable if $|\bar{X}| < |X|$. It may happen that the retraction of a retractable solution is itself retractable, so we set inductively $\text{Ret}^{n+1}(X, r) = \text{Ret}^n(\bar{X}, \bar{r})$. The *multipermutation level* of (X, r) , denoted $\text{mpl}(X, r)$, is the infimum of all $n \in \mathbb{N}$ such that $\text{Ret}^n(X, r)$ is a solution of cardinality 1.

Remark. *There is ambiguity in the notation $\mathcal{L}_{\bar{x}}$, as it could be a function from X to itself or from \bar{X} to itself. This issue will not arise in the sequel as we will always work at the level of X and its linearization V .*

2.5. Formulas. From now on we assume that (X, r) is a solution. To avoid overly complicated notation we denote by $[\bar{x}]$ the class of \bar{x} in $\text{Ret}^2(X, r)$, and by $[[\bar{x}]]$ the class of $[\bar{x}]$ in $\text{Ret}^3(X, r)$. Since $x \triangleright y$ depends only on the class of x it makes sense to write $\bar{x} \triangleright y$, and by a similar reasoning $\bar{x} \triangleright \bar{y} = [[\bar{x}]] \triangleright \bar{y}$, etc. We can thus rewrite the conditions

on the previous lemma as

$$\begin{aligned}\mathcal{L}_{\bar{x}} \circ \mathcal{L}_{\bar{y}} &= \mathcal{L}_{[\bar{x}] \triangleright \bar{y}} \circ \mathcal{L}_{\bar{x} \triangleleft [\bar{y}]}; \\ \mathcal{R}_{\bar{z}} \circ \mathcal{R}_{\bar{y}} &= \mathcal{R}_{\bar{y} \triangleleft [\bar{z}]} \circ \mathcal{R}_{[\bar{y}] \triangleright \bar{z}}; \\ \mathcal{R}_{([\bar{x}] \triangleleft [\bar{y}]) \triangleright \bar{z}} \circ \mathcal{L}_{\bar{x}} \circ p_{\bar{y}} &= \mathcal{L}_{\bar{x} \triangleleft ([\bar{y}] \triangleright [\bar{z}])} \circ \mathcal{R}_{\bar{z}} \circ p_{\bar{y}}.\end{aligned}$$

We also point out that

$$\mathcal{L}_{\bar{x}} \circ p_{\bar{y}} = p_{[\bar{x}] \triangleright \bar{y}} \circ \mathcal{L}_{\bar{x}}; \quad \mathcal{R}_{\bar{z}} \circ p_{\bar{y}} = p_{\bar{y} \triangleleft [\bar{z}]} \circ \mathcal{R}_{\bar{z}}; \quad p_{\bar{x}} \circ p_{\bar{y}} = p_{\bar{y}} \circ p_{\bar{x}} = \delta_{\bar{x}, \bar{y}} p_{\bar{x}}.$$

Finally, we can write R as

$$R \circ \tau = \sum_{\bar{x} \in \bar{X}} \sum_{\bar{y} \in \bar{X}} \mathcal{L}_{\bar{x}} \circ p_{\bar{y}} \otimes \mathcal{R}_{\bar{y}} \circ p_{\bar{x}}$$

where τ is the flip map given by $\tau(v \otimes w) = w \otimes v$. To see this, notice that given $v, w \in X$ we have

$$\begin{aligned}\left(\sum_{\bar{x} \in \bar{X}} \sum_{\bar{y} \in \bar{X}} \mathcal{L}_{\bar{x}} \circ p_{\bar{y}} \otimes \mathcal{R}_{\bar{y}} \circ p_{\bar{x}} \right) \tau(v \otimes w) \\ = \sum_{\bar{x} \in \bar{X}} \sum_{\bar{y} \in \bar{X}} \mathcal{L}_{\bar{x}} \circ p_{\bar{y}}(w) \otimes \mathcal{R}_{\bar{y}} \circ p_{\bar{x}}(v) \\ = \mathcal{L}_{\bar{v}}(w) \otimes \mathcal{R}_{\bar{w}}(v) = R(v \otimes w),\end{aligned}$$

so the desired equality holds over a basis of $V \otimes V$.

3. Multipermutation level 2

3.1. Formulas for multipermutation level 2. If $\text{mpl}(X, r) = 2$ then $[\bar{y}] = X$. Also there exist $f, g : \bar{X} \rightarrow \bar{X}$ such that $\bar{r}(\bar{x}, \bar{y}) = (f(\bar{y}), g(\bar{x}))$ so the formulas above simplify to

$$\begin{aligned}\mathcal{L}_{\bar{x}} \circ \mathcal{L}_{\bar{y}} &= \mathcal{L}_{f(\bar{y})} \circ \mathcal{L}_{g(\bar{x})}; \quad \mathcal{R}_{\bar{z}} \circ \mathcal{R}_{\bar{y}} = \mathcal{R}_{g(\bar{y})} \circ \mathcal{R}_{f(\bar{z})}; \quad \mathcal{R}_{f(\bar{z})} \circ \mathcal{L}_{\bar{x}} = \mathcal{L}_{g(\bar{x})} \circ \mathcal{R}_{\bar{z}}; \\ \mathcal{L}_{\bar{x}} \circ p_{\bar{y}} &= p_{f(\bar{y})} \circ \mathcal{L}_{\bar{x}} \quad \mathcal{R}_{\bar{x}} \circ p_{\bar{y}} = p_{g(\bar{y})} \circ \mathcal{R}_{\bar{x}}\end{aligned}$$

In particular, if $\text{Ret}(X, r)$ is the trivial solution, or equivalently if $f = g = \text{Id}_{\bar{X}}$, then the maps $\mathcal{L}_{\bar{x}}, \mathcal{R}_{\bar{y}}, p_{\bar{z}}$ commute for all $x, y, z \in X$.

3.2. The following result is a generalization of [GIM11, Theorem 4.9] to non-involutive solutions. Also, we replace the hypothesis that r is square-free with the weaker hypothesis that the first retraction is a trivial solution.

Proposition 3.1. *Suppose $\text{Ret}(X, r)$ is trivial. Then there exist a basis $\{v_1, \dots, v_n\}$ of V and a set of roots of unity $\mathbf{q} = (q_{i,j})_{1 \leq i, j \leq n} \subset \mathbb{C}$ such that $R(v_i \otimes v_j) = q_{i,j} v_j \otimes v_i$.*

Proof. Since $p_{\bar{z}}$ is a projection operator it is diagonalizable, and its eigenvalues are either 1 or 0. Since $\mathcal{L}_{\bar{x}}$ and $\mathcal{R}_{\bar{y}}$ are induced by bijections of the set X , they are also diagonalizable and their eigenvalues are roots of unity. As mentioned above the fact that $\text{Ret}(X, r)$ is trivial means that the maps $\mathcal{L}_{\bar{x}}, \mathcal{R}_{\bar{y}}$ commute, and they commute with $p_{\bar{z}}$, for all $\bar{x}, \bar{y}, \bar{z} \in \bar{X}$. Also since X is the disjoint union of the classes \bar{z} the projectors $p_{\bar{z}}$ are orthogonal and add up to the identity, so they also commute with each other.

By a standard result of linear algebra the fact that these linear operators are diagonalizable and commute implies that they are *simultaneously* diagonalizable, i.e. there exists a basis of V consisting of eigenvectors of all these maps, say $\{v_1, \dots, v_n\}$. Write $\lambda_{i,\bar{x}}$, resp. $\mu_{i,\bar{y}}$, for the eigenvalue of v_i with respect to $\mathcal{L}_{\bar{x}}$, resp. $\mathcal{R}_{\bar{y}}$. Using the fact that $R = \left(\sum_{\bar{x}, \bar{y} \in \bar{X}} \mathcal{L}_{\bar{x}} \circ p_{\bar{y}} \otimes \mathcal{R}_{\bar{y}} \circ p_{\bar{x}} \right) \circ \tau$ and that for each i there is exactly one $\bar{z}(i)$ such that $p_{\bar{z}(i)}(v_i) \neq 0$ we get

$$R(v_i \otimes v_j) = \lambda_{j,\bar{z}(i)} \mu_{i,\bar{z}(j)} v_j \otimes v_i$$

Since $\lambda_{i,j}$ and $\mu_{i,j}$ are roots of unity, so is their product and we are done. \square

3.3. The algebra $Q(X, r)$ associated to a solution is the quotient of the tensor algebra $T(V)$ by the ideal generated by the image of the map $\text{Id}_{V \otimes V} - R$. This is a quadratic algebra whose properties have been studied intensively, see for example [GIJ03, GI04, GI12]. Proposition 3.1 arose from attempts to study this algebras. It follows from it that if $\text{Ret}(X, r)$ is a trivial solution then $Q(X, r)$ is a twisted polynomial algebra, and furthermore it is integral if and only if r is involutive. A similar result holds for the Hopf algebra associated to the solution (see [ESS99, section 2.9]). These results appear in the MsC thesis of Agustín Muñoz [Muñ20].

3.4. Examples. Recall the involutive, non square-free solution $([[2]], s)$ from 2.2. Taking $v_1 = x_1 + x_2$ and $v_2 = v_1 - v_2$ the corresponding linear solution is given by

R	v_1	v_2
v_1	$v_1 \otimes v_1$	$-v_2 \otimes v_1$
v_2	$-v_1 \otimes v_2$	$v_2 \otimes v_2$

The algebra $Q(X, r)$ is then the polynomial algebra in anticommuting variables.

Now consider solution $([[4]], t)$, which is a non-involutive permutation solution. Using the notation from the proposition we have eigenvectors $v_1 = x_1 + x_2, v_2 = x_1 - x_2, v_3 = x_3 + x_4, v_4 = x_3 - x_4$. The linearized solution is given by $R(v_i \otimes v_j) = q_{i,j} v_j \otimes v_i$ where

$$\mathbf{q} = \begin{pmatrix} 1 & -1 & 1 & -1 \\ -1 & 1 & -1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & 1 & -1 \end{pmatrix}.$$

Thus for example $v_4^2 = 0$ in $Q(X, r)$. If r is not involutive then there will always be elements with square zero in $Q(X, r)$.

4. A Lie algebra associated to a solution

Throughout this section (X, r) denotes a finite solution. We associate to (X, r) a Lie algebra contained in $\mathfrak{gl}(V)$ and show that it characterizes solutions with trivial retraction.

Definition 4.1. The Lie algebra associated to the solution (X, r) , denoted by $\mathfrak{g}(X, r)$, is the Lie subalgebra of $\mathfrak{gl}(V)$ generated by the operators $\{\mathcal{L}_{\bar{x}} \circ p_{\bar{y}}, \mathcal{R}_{\bar{x}} \circ p_{\bar{y}} \mid \bar{x}, \bar{y} \in \bar{X}\}$.

4.1. From the formulas in 2.5 we deduce the commutation formulas

$$\begin{aligned} [\mathcal{L}_{\bar{x}} \circ p_{\bar{y}}, \mathcal{L}_{\bar{z}} \circ p_{\bar{w}}] &= p_{[\bar{x}] \triangleright \bar{y}} \circ p_{[[\bar{x}] \triangleright (\bar{z}] \triangleright \bar{w}]} \circ \mathcal{L}_{\bar{x}} \circ \mathcal{L}_{\bar{z}} - p_{[\bar{z}] \triangleright \bar{w}} \circ p_{[[\bar{z}] \triangleright ([\bar{x}] \triangleright \bar{y})]} \circ \mathcal{L}_{\bar{z}} \circ \mathcal{L}_{\bar{x}}, \\ [\mathcal{R}_{\bar{x}} \circ p_{\bar{y}}, \mathcal{R}_{\bar{z}} \circ p_{\bar{w}}] &= p_{\bar{y} \triangleleft [\bar{x}]} \circ p_{(\bar{w} \triangleleft [\bar{z}]) \triangleleft [[\bar{x}]]} \circ \mathcal{R}_{\bar{x}} \circ \mathcal{R}_{\bar{z}} - p_{\bar{w} \triangleleft [\bar{z}]} \circ p_{(\bar{y} \triangleleft [\bar{x}]) \triangleleft [[\bar{z}]]} \circ \mathcal{R}_{\bar{z}} \circ \mathcal{R}_{\bar{x}}, \\ [\mathcal{L}_{\bar{x}} \circ p_{\bar{y}}, \mathcal{R}_{\bar{z}} \circ p_{\bar{w}}] &= p_{[\bar{x}] \triangleright \bar{y}} \circ p_{[[\bar{x}] \triangleright (\bar{w} \triangleleft [\bar{z}])]} \circ \mathcal{L}_{\bar{x}} \circ \mathcal{R}_{\bar{z}} - p_{\bar{w} \triangleleft [\bar{z}]} \circ p_{([\bar{x}] \triangleright \bar{y}) \triangleleft [[\bar{z}]]} \circ \mathcal{R}_{\bar{z}} \circ \mathcal{L}_{\bar{x}}. \end{aligned}$$

It also follows that the linear solution R is a solution to the classical YBE in $\mathfrak{g}(X, r)$. The crucial fact in the proof of Proposition 3.1 is that the operators $\{\mathcal{L}_{\bar{x}} \circ p_{\bar{y}}, \mathcal{R}_{\bar{z}} \circ p_{\bar{w}}\}$ commute when $\text{Ret}(X, r)$ is a flip solution, so in this case R is a solution to the YBE in an abelian Lie algebra.

Theorem 4.2. Let (X, r) be a finite non-degenerate solution and let $\mathfrak{g} = \mathfrak{g}(X, r)$. Then the following are equivalent.

- (a) The solution $\text{Ret}(X, r)$ is trivial.
- (b) The span of the set $\{\mathcal{L}_{\bar{x}}, \mathcal{R}_{\bar{x}}, p_{\bar{x}}\}$ is an abelian Lie subalgebra of $\mathfrak{gl}(V)$.
- (c) The algebra $\mathfrak{g}(X, r)$ is abelian.

Proof. The implication (a) \Rightarrow (b) was shown in the proof of Proposition 3.1, and is also easy to deduce from the commutation formulas displayed above. The implication (b) \Rightarrow (c) is immediate.

Now suppose $\mathfrak{g}(X, r)$ is abelian. Choose arbitrary elements $x, y, z \in X$ and choose $w \in X$ such that $y = w \triangleleft z$. Since $[\mathcal{L}_{\bar{x}} \circ p_{\bar{y}}, \mathcal{R}_{\bar{z}} \circ p_{\bar{w}}] = 0$ we deduce from the formulas in 2.5 that

$$p_{[\bar{x}] \triangleright \bar{y}} \circ \mathcal{L}_{\bar{x}} \circ \mathcal{R}_{\bar{z}} = p_{\bar{y}} \circ p_{([\bar{x}] \triangleright \bar{y}) \triangleleft [[\bar{z}]]} \circ \mathcal{R}_{\bar{z}} \circ \mathcal{L}_{\bar{x}}.$$

Since $\mathcal{L}_{\bar{x}}$ and $\mathcal{R}_{\bar{z}}$ are automorphisms of V , the morphism on the left hand side of the equality is nonzero and its image is the subspace of V spanned by $[\bar{x}] \triangleright \bar{y}$. Thus the map on the right hand side is also nonzero, which implies that $\bar{y} = ([\bar{x}] \triangleright \bar{y}) \triangleleft [[\bar{z}]]$, and its image is the subspace spanned by \bar{y} , which implies that $\bar{y} = [\bar{x}] \triangleright \bar{y}$. Both inequalities together imply that \bar{y} is a fixed point for the action of $[\bar{x}]$ on the left and for the action of $[\bar{z}]$ on the right. Since x, y, z were chosen arbitrarily, this means that $\mathcal{L}_{[\bar{x}]} = \mathcal{R}_{[\bar{z}]} = \text{Id}_{\bar{V}}$ and hence $\text{Ret}(X, r)$ is a trivial solution. \square

We would like to have a more intrinsic definition of $\mathfrak{g}(X, r)$ but so far this is the best description we have. Computer experiments show that for all $|X| \leq 8$ and r involutive the derived algebra of $\mathfrak{g}(X, r)$ is semisimple of type A . Thus the multipermutation level is not directly related to the solubility or nilpotency of $\mathfrak{g}(X, r)$, as one might suppose from the statement. We intend to return to this matter in future work.

Acknowledgements I thank Leandro Vendramin and Tatiana Gateva-Ivanova for reading and commenting on an early version of this article. The computer experiments

mentioned above were ran by Agustín Muñoz as part of his MsC. thesis, using Vendramin's GAP package YBE.

References

- [Dri92] V. G. Drinfel'd, *On some unsolved problems in quantum group theory*, Quantum groups (Leningrad, 1990), Lecture Notes in Math., vol. 1510, Springer, Berlin, 1992, pp. 1–8.
- [ESS99] P. Etingof, T. Schedler, and A. Soloviev, *Set-theoretical solutions to the quantum Yang-Baxter equation*, Duke Math. J. **100** (1999), no. 2, 169–209.
- [GI12] T. Gateva-Ivanova, *Quadratic algebras, Yang-Baxter equation, and Artin-Schelter regularity*, Adv. Math. **230** (2012), no. 4–6.
- [GI04] ———, *Binomial skew polynomial rings, Artin-Schelter regularity, and binomial solutions of the Yang-Baxter equation*, Serdica Math. J. **30** (2004), no. 2–3, 431–470.
- [GIJ03] T. Gateva-Ivanova, E. Jespers, and J. Okniński, *Quadratic algebras of skew type and the underlying monoids*, J. Algebra **270** (2003), no. 2, 635–659.
- [GIM11] T. Gateva-Ivanova and S. Majid, *Quantum spaces associated to multipermutation solutions of level two*, Algebr. Represent. Theory **14** (2011), no. 2, 341–376.
- [GIvdB98] T. Gateva-Ivanova and M. van den Bergh, *Semigroups of I-type*, J. Algebra **206** (1998), no. 1, 97–112.
- [GV17] L. Guarnieri and L. Vendramin, *Skew braces and the Yang-Baxter equation*, Math. Comp. **86** (2017), no. 307, 2519–2534.
- [LV17] V. Lebed and L. Vendramin, *Homology of left non-degenerate set-theoretic solutions to the Yang-Baxter equation*, Adv. Math. **304** (2017), 1219–1261.
- [Muñ20] A. Muñoz, *Álgebras cuantizadas y álgebras de Lie asociadas a soluciones conjuntistas de la ecuación de Yang-Baxter (in spanish)*, 2020. MsC thesis, available at <http://cms.dm.uba.ar/academico/carreras/licenciatura/tesis/2020/AMunoz.pdf>.
- [Rum07] W. Rump, *Braces, radical rings, and the quantum Yang-Baxter equation*, J. Algebra **307** (2007), no. 1, 153–170.
- [Ven19] L. Vendramin, *Problems on skew left braces*, Adv. Group Theory Appl. **7** (2019), 15–37.

JACOBS UNIVERSITY, BREMEN AND CAECE UNIVERSITY, BUENOS AIRES

Email address: pzadunaisky@gmail.com