

LEARNING FROM NON-IRREDUCIBLE MARKOV CHAINS

NIKOLA SANDRIĆ AND STJEPAN ŠEBEK

ABSTRACT. Most of the existing literature on supervised learning problems focuses on the case when the training data set is drawn from an i.i.d. sample. However, many practical supervised learning problems are characterized by temporal dependence and strong correlation between the marginals of the data-generating process, suggesting that the i.i.d. assumption is not always justified. This problem has been already considered in the context of Markov chains satisfying the Doeblin condition. This condition, among other things, implies that the chain is not singular in its behavior, i.e. it is irreducible. In this article, we focus on the case when the training data set is drawn from a not necessarily irreducible Markov chain. Under the assumption that the chain is uniformly ergodic with respect to the L^1 -Wasserstein distance, and certain regularity assumptions on the hypothesis class and the state space of the chain, we first obtain a uniform convergence result for the corresponding sample error, and then we conclude learnability of the approximate sample error minimization algorithm and find its generalization bounds. At the end, a relative uniform convergence result for the sample error is also discussed.

1. INTRODUCTION

Let (X, \mathcal{X}) and (Y, \mathcal{Y}) be measurable spaces, let $Z \subseteq X \times Y$ be a complete separable metric space with bounded metric ρ , and let $\text{pr}_X : Z \rightarrow X$ and $\text{pr}_Y : Z \rightarrow Y$ be the projection mappings. We endow (Z, ρ) with its Borel σ -algebra $\mathfrak{B}(Z)$. Assume that $\text{pr}_X^{-1}(\text{pr}_X(Z) \cap \mathcal{X}) \subseteq \mathfrak{B}(Z)$ and $\text{pr}_Y^{-1}(\text{pr}_Y(Z) \cap \mathcal{Y}) \subseteq \mathfrak{B}(Z)$. This, in particular, implies that pr_X and pr_Y are measurable.

Further, let $(\Omega, \mathcal{F}, \mathbb{P}_z, \{Z_n\}_{n \geq 0}, \{\mathcal{F}_n\}_{n \geq 0})_{z \in Z}$, denoted by $\{Z_n\}_{n \geq 0}$ in the sequel, be a temporally-homogeneous Markov chain on $(Z, \mathfrak{B}(Z))$, in the sense of [MT09]. The process $\{Z_n\}_{n \geq 0} = \{(\text{pr}_X(Z_n), \text{pr}_Y(Z_n))\}_{n \geq 0}$ is viewed as a sequence of input data (the first component) together with their outputs (the second component).

The main aim of this article is to learn a function (from a given hypothesis class $\mathcal{H} \subseteq Y^X$) that, given a training set of data $z_0, \dots, z_{n-1} \in Z$ drawn from the first n samples of $\{Z_n\}_{n \geq 0}$, best approximates the relationship between the input and output observable. We remark that even if there is an exact functional relationship, this function is in general not known to us and it does not have to belong to the class \mathcal{H} .

This type of problem has already been considered in the literature, see the literature review part below. However, in all these works a crucial assumption is that the process $\{Z_n\}_{n \geq 0}$ is not singular in its behavior, i.e. it is irreducible. In this article, we study the learning problem in the case when $\{Z_n\}_{n \geq 0}$ does not necessarily possess this property. We follow the standard statistical learning theory scheme: we first obtain a uniform convergence result for the corresponding sample error, and then we conclude learnability of the approximate sample error minimization (ASEM) algorithm and find its generalization bounds.

Main results. We start with the description of the model. Assume the following:

- (A1):** $Y \subseteq \mathbb{R}$, $\mathcal{Y} = \mathfrak{B}(\mathbb{R}) \cap Y$ (the standard relative Euclidean σ -algebra), and $\mathcal{H} \subseteq Y^X$ is a totally bounded metric space, i.e. \mathcal{H} admits a metric $\rho_{\mathcal{H}}$ such that for each $\epsilon > 0$

2010 *Mathematics Subject Classification.* 68W40, 68T10, 60J05.

Key words and phrases. Approximate sample error minimization algorithm, Generalization bounds, Markov chain, Wasserstein distance.

there is finite $\mathcal{H}_\epsilon \subseteq \mathcal{H}$ with the property that for every $h \in \mathcal{H}$ there is $h_\epsilon \in \mathcal{H}_\epsilon$ with $\rho_{\mathcal{H}}(h, h_\epsilon) < \epsilon$.

For $\epsilon > 0$ let

$$\mathcal{N}(\epsilon, \mathcal{H}, \rho_{\mathcal{H}}) := \min\{|\mathcal{H}_\epsilon| : \mathcal{H}_\epsilon \text{ is an } \epsilon\text{-covering of } \mathcal{H} \text{ in the sense of (A1)}\}$$

be the ϵ -covering number of \mathcal{H} . Typical examples of classes of functions satisfying (A1) are the following:

- (i) Let $X = [0, T]$, let $Y = [-V/2, V/2]$ and let $\mathcal{H} \subset Y^X$ be a class of functions with total variation at most V , for some $T, V > 0$. Further, let $\rho_{\mathcal{H}}$ be the corresponding L^1 -metric (with respect to the Lebesgue measure). Then, in [BKP97, Theorem 1] it has been shown that

$$\mathcal{N}(\epsilon, \mathcal{H}, \rho_{\mathcal{H}}) \leq 2^{13VT/\epsilon}.$$

- (ii) Let $X \subset \mathbb{R}^d$ be compact and let $\mathcal{H} \subset C^\gamma(X, \mathbb{R})$ be bounded (with respect to the corresponding Hölder metric) for some $0 < \gamma \leq 1$. Then, for $\rho_{\mathcal{H}}$ being the Hölder or sup-metric, in [Zho03, Theorem 4] it has been shown that there is $C > 0$ such that

$$(1.1) \quad \mathcal{N}(\epsilon, \mathcal{H}, \rho_{\mathcal{H}}) \leq e^{C\epsilon^{-2d/\gamma}}.$$

Further, let $\ell : Y \times Y \rightarrow [0, \infty)$ be measurable (the loss function), and for each $h \in \mathcal{H}$ define $\ell_h : Z \rightarrow [0, \infty)$ by

$$\ell_h(z) := \ell(h(\text{pr}_X(z)), \text{pr}_Y(z)).$$

We next assume:

- (A2):** there are $L, \bar{L} \geq 0$ such that $|\ell_{h_1}(z_1) - \ell_{h_2}(z_2)| \leq L\rho(z_1, z_2) + \bar{L}\rho_{\mathcal{H}}(h_1, h_2)$ for all $z_1, z_2 \in Z$ and $h_1, h_2 \in \mathcal{H}$.

Examples of classes of functions for which (A2) holds are the following:

- (i) Let X and \mathcal{H} be as in (ii) above, with $\gamma = 1$ (the class of Lipschitz continuous functions). Assume also that Y is bounded. Then, (A2) holds with $\ell(y, \bar{y}) = |y - \bar{y}|^2$, and ρ and $\rho_{\mathcal{H}}$ being the Euclidean and sup metric, respectively.
- (ii) Let X and Y be as in (i), and let $\mathcal{H} \subseteq \{h \in Y^X : h \text{ is Lipschitz continuous and } h(x_0) = y_0\}$ for some $x_0 \in X$ and $y_0 \in Y$. Then, (A2) holds with $\ell(y, \bar{y}) = |y - \bar{y}|^2$, and ρ and $\rho_{\mathcal{H}}$ being the Euclidean and Lipschitz metric, respectively.

Denote by $\mathcal{P}(z, d\bar{z})$ the (one-step) transition kernel of $\{Z_n\}_{n \geq 0}$, and let $\mathcal{P}_1(Z)$ be the class of all probability measures on Z having finite first moment. Recall that the L^1 -Wasserstein distance on $\mathcal{P}_1(Z)$ is defined by

$$\mathcal{W}(\mu_1, \mu_2) := \inf_{\Pi \in \mathcal{C}(\mu_1, \mu_2)} \int_{Z \times Z} \rho(z, \bar{z}) \Pi(dz, d\bar{z}),$$

where $\mathcal{C}(\mu_1, \mu_2)$ is the family of couplings of $\mu_1(dz)$ and $\mu_2(dz)$, i.e. $\Pi \in \mathcal{C}(\mu_1, \mu_2)$ if, and only if, $\Pi(dz, d\bar{z})$ is a probability measure on $Z \times Z$ having $\mu_1(dz)$ and $\mu_2(dz)$ as its marginals. We finally assume:

- (A3):** Y is bounded and there is $\eta \in (0, 1)$ such that

$$\mathcal{W}(\mathcal{P}(z_1, d\bar{z}), \mathcal{P}(z_2, d\bar{z})) \leq (1 - \eta) \rho(z_1, z_2).$$

An example satisfying (A3) is as follows. Let $\{\chi_n\}_{n \geq 1}$ be a sequence of i.i.d. \mathbb{R} -valued random variables defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, satisfying $\mathbb{P}(\chi_n = 0) = \mathbb{P}(\chi_n = 1/2) = 1/2$. Define

$$X_{n+1} := \frac{1}{2}X_n + \chi_{n+1}$$

with $X_0 = x \in [0, 1]$. Clearly, $\{X_n\}_{n \geq 0}$ is a Markov chain on $[0, 1]$ with transition kernel $\mathcal{P}_X(x, d\bar{x}) = \mathbb{P}(\chi_1 + x/2 \in d\bar{x})$. Further, let $f : [0, 1] \rightarrow \mathbb{R}$ be Lipschitz continuous with Lipschitz constant $\text{Lip}(f) < \sqrt{3}$ (hence, $f(x)$ is of bounded variation), and set $Z_n := (X_n, f(X_n))$ for $n \geq 0$. Then, $\{Z_n\}_{n \geq 0}$ is a Markov chain on $Z = \{(x, f(x)) : x \in [0, 1]\}$ with transition kernel $\mathcal{P}(z, d\bar{z}) = \mathbb{P}(\chi_1 + \text{pr}_X(z)/2 \in \text{dpr}_X(\bar{z}))$, see Lemma 3.1. For ρ we take the standard Euclidean metric. Clearly, $\{Z_n\}_{n \geq 0}$ is not irreducible. Namely, for $z \in Z$ such that $\text{pr}_X(z) \in \mathbb{Q}$ it holds that $\mathcal{P}^n(z, Z \cap (\mathbb{Q}^c \times \mathbb{R})) = 0$ for all $n \geq 1$, and analogously for $z \in Z$ such that $\text{pr}_X(z) \in \mathbb{Q}^c$ it holds that $\mathcal{P}^n(z, Z \cap (\mathbb{Q} \times \mathbb{R})) = 0$ for all $n \geq 1$. Further, let $\text{Length}(dz)$ be the arc-length measure on Z . It is then easy to see that $\pi(dz) := \text{Length}(dz)/\text{Length}(Z)$ satisfies $\int_Z \mathcal{P}(z, d\bar{z})\pi(dz) = \pi(d\bar{z})$, i.e. $\pi(dz)$ is an invariant probability measure for $\{Z_n\}_{n \geq 0}$. Here, $\text{Length}(Z)$ is the length of Z . Observe that $\pi(d\bar{z})$ and $\mathcal{P}(z, \bar{z})$ are mutually singular, implying that the relation in eq. (1.4) cannot hold. Finally, we show the relation in (A3). By Kantorovich-Rubinstein theorem we have that

$$\mathcal{W}(\mu_1, \mu_2) = \sup_{\{g : \text{Lip}(g) \leq 1\}} |\mu_1(g) - \mu_2(g)|,$$

where the supremum is taken over all Lipschitz continuous functions $g : Z \rightarrow \mathbb{R}$ with Lipschitz constant $\text{Lip}(g) \leq 1$ and, for a probability measure $\mu(dz)$ on Z and a measurable function $f : Z \rightarrow \mathbb{R}$, the symbol $\mu(f)$ stands for $\int_Z f(z)\mu(dz)$, whenever the integral is well defined. Thus, for any $z_1 = (x_1, f(x_1))$, $z_2 = (x_2, f(x_2)) \in Z$ it holds that

$$\begin{aligned} & \mathcal{W}(\mathcal{P}(z_1, \bar{z}), \mathcal{P}(z_2, \bar{z})) \\ &= \sup_{\{g : \text{Lip}(g) \leq 1\}} \left| \int_Z g(\bar{z}) \mathbb{P}(\chi_1 + \text{pr}_X(z_1)/2 \in \text{dpr}_X(\bar{z})) - \int_Z g(\bar{z}) \mathbb{P}(\chi_1 + \text{pr}_X(z_2)/2 \in \text{dpr}_X(\bar{z})) \right| \\ &= \sup_{\{g : \text{Lip}(g) \leq 1\}} \left| \frac{1}{2} (g(x_1/2, f(x_1/2)) - g(x_2/2, f(x_2/2))) \right. \\ & \quad \left. + \frac{1}{2} (g((x_1+1)/2, f((x_1+1)/2)) - g((x_2+1)/2, f((x_2+1)/2))) \right| \\ &\leq \frac{1}{2} \rho((x_1/2, f(x_1/2)), (x_2/2, f(x_2/2))) \\ & \quad + \frac{1}{2} \rho(((x_1+1)/2, f((x_1+1)/2)), ((x_2+1)/2, f((x_2+1)/2))) \\ &\leq \frac{\sqrt{1 + \text{Lip}(f)^2}}{2} |x_1 - x_2| \\ &\leq \frac{\sqrt{1 + \text{Lip}(f)^2}}{2} \rho(z_1, z_2), \end{aligned}$$

which proves (A3) with $\eta = 1 - \sqrt{1 + \text{Lip}(f)^2}/2$ (recall that $\text{Lip}(f) < \sqrt{3}$).

According to [But14, Theorem 2.1] (by taking $V \equiv 1$, $\varphi(t) = t$ and $K \geq 1$) the relation in (A3) implies that

- (i) $\{Z_n\}_{n \geq 0}$ admits a unique invariant probability measure $\pi(dz)$;
- (ii) there are $C_1, C_2 > 0$ such that

$$(1.2) \quad \mathcal{W}(\mathcal{P}^n(z, d\bar{z}), \pi(d\bar{z})) \leq C_1 e^{-C_2 n}.$$

Before stating the main results of this article, we introduce some notation we need. For $n \in \mathbb{N}$, $z \in Z$ and $h \in \mathcal{H}$, let $\hat{\text{er}}_n(h) := \frac{1}{n} \sum_{i=0}^{n-1} \ell_h(Z_i)$, $\text{er}_\pi(h) := \pi(\ell_h)$, $\ell_h^\pi(z) := \ell_h(z) - \text{er}_\pi(h)$ and $\text{opt}_\pi(\mathcal{H}) := \inf_{h \in \mathcal{H}} \text{er}_\pi(h)$. Observe that, according to (A2) and boundedness of ρ , $\text{er}_\pi(h)$ and $\ell_h^\pi(z)$ are well defined. Next, for $\varepsilon > 0$ the ε -ASEM algorithm for \mathcal{H} is defined as a mapping

$\mathcal{A}^\varepsilon : \bigcup_{n=1}^{\infty} \mathcal{Z}^n \rightarrow \mathcal{H}$ satisfying

$$\frac{1}{n} \sum_{i=0}^{n-1} \ell_{\mathcal{A}^\varepsilon(z_0, \dots, z_{n-1})}(z_i) < \inf_{h \in \mathcal{H}} \frac{1}{n} \sum_{i=0}^{n-1} \ell_h(z_i) + \varepsilon.$$

Theorem 1.1. *Assume (A1)-(A3), and fix $\varepsilon > 0$ and $\delta \in (0, 1)$. Then, for any initial distribution μ of $\{Z_n\}_{n \geq 0}$,*

$$\mathbb{P}_\mu \left(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_0, \dots, Z_{n-1})) - \text{opt}_\pi(\mathcal{H})| < 5\varepsilon \right) \geq 1 - \delta, \quad \forall n \geq n_1(\varepsilon, \delta),$$

where

$$n_1(\varepsilon, \delta) := \max \left\{ \frac{16C_1L}{\varepsilon(1 - e^{-C_2})}, \frac{128C_1^2L^2 \ln \frac{\mathcal{N}(\varepsilon/4\bar{L}, \mathcal{H}, d_{\mathcal{H}})}{\delta}}{\varepsilon^2(1 - e^{-C_2})^2} \right\}.$$

The main step in the proof of Theorem 1.1 is the following sample error uniform convergence result

$$\mathbb{P}_z \left(\sup_{h \in \mathcal{H}} |\hat{\text{er}}_n(h) - \text{er}_\pi(h)| > \varepsilon \right) \leq \mathcal{N}(\varepsilon/4\bar{L}, \mathcal{H}, d_{\mathcal{H}}) e^{-\left(\frac{\varepsilon n(1 - e^{-C_2})}{4C_1L} - 2\right)^2 / (2n)}, \quad \forall n \geq \frac{8C_1L}{\varepsilon(1 - e^{-C_2})},$$

(see eq. (2.4)). Notice that in the case when $\mathcal{N}(\varepsilon, \mathcal{H}, d_{\mathcal{H}}) = e^{C\varepsilon^{-2d/\gamma}}$ (see eq. (1.1)) the sample size (in the above relation or Theorem 1.1) is of order $\varepsilon^{-(2+2d/\gamma)}$. On the other hand, one expects that the sample size n needed to ensure that $(1 + \alpha)\hat{\text{er}}_n(h) + \varepsilon \geq \text{er}_\pi(h)$ for some $\alpha, \varepsilon > 0$ is of much smaller order. This problem was studied in [AB99, Subsection 5.5]. Crucial step in the proof of this result is to study the uniform relative error

$$\sup_{h \in \mathcal{H}} \frac{|\text{er}_\pi(h) - \hat{\text{er}}_n(h)|}{\sqrt{\text{er}_\pi(h)}},$$

instead of the uniform standard error

$$\sup_{h \in \mathcal{H}} |\text{er}_\pi(h) - \hat{\text{er}}_n(h)|.$$

Studying this quantity is additionally motivated by the observation that a uniform convergence bound fails to capture the phenomenon that for those functions $h \in \mathcal{H}$ for which the true error $\text{er}_\pi(h)$ is small, the deviation $\text{er}_\pi(h) - \hat{\text{er}}_n(h)$ is also small with large probability. However, the relative deviation $(\text{er}_\pi(h) - \hat{\text{er}}_n(h)) / \sqrt{\text{er}_\pi(h)}$ is more appropriate for capturing this phenomenon (see [ZZX09, Section 4]). With relative error we penalize the difference between the sample error and the true error much more when the true error is very small in the first place. Notice that if there is an exact functional relationship between the input and the output observable, and if this function is in the class \mathcal{H} , then the logic behind the relative error tells us that every function (except the true one) is infinitely worse than the true one. Hence, studying relative error makes sense only when the true error is bounded away from zero. Even though this assumption does seem a bit unnatural, in many practical applications there will not even be a true function in the background, but some distribution, and even if there will be a functional relationship between the input and the output observable, this relationship can often be pretty complex and hence not contained in the class \mathcal{H} .

Theorem 1.2. *Assume (A1)-(A3) and assume additionally that there exist constants $m, M > 0$ such that*

$$(1.3) \quad m \leq \inf_{h \in \mathcal{H}} \text{er}_\pi(h) \leq \sup_{h \in \mathcal{H}} \text{er}_\pi(h) \leq M.$$

Fix $\varepsilon > 0$ and $\delta \in (0, 1)$. Then, for any initial distribution μ of $\{Z_n\}_{n \geq 0}$,

$$\mathbb{P}_\mu \left(\sup_{h \in \mathcal{H}} \frac{|\hat{e}r_n(h) - e r_\pi(h)|}{\sqrt{e r_\pi(h)}} \geq \varepsilon \right) \leq \delta, \quad \forall n \geq n_2(\varepsilon, \delta),$$

where

$$n_2(\varepsilon, \delta) := \max \left\{ \frac{2C_1 L}{\varepsilon(\sqrt{m} \wedge 1)(1 - e^{-C_2})}, \frac{1}{\xi_1 \varepsilon^2} \left(\xi_2 \varepsilon + \ln \left(\frac{4\mathcal{N}(\varepsilon/\bar{L}, \mathcal{H}, d_{\mathcal{H}})}{\delta} \right) \right) \right\}$$

and

$$\xi_1 = \frac{49m^4(1 - e^{-C_2})^2}{72M(M + 6m)^2 C_1^2 L^2}, \quad \xi_2 = \frac{7m^2(1 - e^{-C_2})}{6\sqrt{M}(M + 6m)C_1 L}.$$

As a consequence of Theorem 1.2 we conclude the following.

Corollary 1.3. Assume (A1)-(A3) and eq. (1.3). Fix $\alpha, \varepsilon > 0$ and $\delta \in (0, 1)$. Then, for any initial distribution μ of $\{Z_n\}_{n \geq 0}$,

$$\mathbb{P}_\mu \left(\exists h \in \mathcal{H} : e r_\pi(h) > (1 + \alpha)\hat{e}r_n(h) + \varepsilon \right) \leq \delta, \quad \forall n \geq n_3(\varepsilon, \delta),$$

where

$$n_3(\varepsilon, \delta) := \max \left\{ \frac{2C_1 L \sqrt{1 + 1/\alpha}}{\sqrt{\varepsilon}(\sqrt{m} \wedge 1)(1 - e^{-C_2})}, \frac{\alpha + 1}{\alpha \xi_1} \cdot \frac{1}{\varepsilon} \left(\frac{\xi_2}{\sqrt{1 + 1/\alpha}} \cdot \varepsilon^{1/2} + \ln \left(\frac{4\mathcal{N} \left(\frac{\varepsilon^{1/2}}{\bar{L}\sqrt{1+1/\alpha}}, \mathcal{H}, d_{\mathcal{H}} \right)}{\delta} \right) \right) \right\},$$

and constants ξ_1 and ξ_2 are as in Theorem 1.2.

Remark 1.4. Notice that for $\mathcal{N}(\varepsilon, \mathcal{H}, d_{\mathcal{H}}) = e^{C\varepsilon^{-2d/\gamma}}$, we need to have n of order $\varepsilon^{-(1+d/\gamma)}$ (which is much smaller than $\varepsilon^{-(2+2d/\gamma)}$).

1.1. Literature review. Our work contributes to the understanding of statistical properties of supervised learning problems. Most of the existing literature focuses on the case when the training data set is drawn from an i.i.d. sample, see the classical monographs [AB99] and [SSBD14]. However, many practical supervised learning problems (e.g. speech recognition, object tracking, market prediction, etc.) are characterized by temporal dependence and strong correlation between the marginals of the data-generating process, suggesting that the i.i.d. assumption is not always justified. This problem has been first addressed in [Yu94], [Mei00] and [Vid03] in the context of stationary β -mixing sequences, and in [MM96] in the case of stationary m -dependent α -mixing sequences. In [SHS09] the authors relax the stationarity assumption and require that the data-generating process satisfies a certain law of large numbers only. This, for example, includes (not necessarily stationary) α -mixing sequences and Markov chains satisfying the Doeblin condition. Recall that the Doeblin condition implies irreducibility, aperiodicity and uniform exponential ergodicity of the chain (see [MT09, Theorem 16.0.2]). The learning problem of the later model was further investigated in a series of articles [Gam03], [ZZX09], [ZLX12], [ZLX⁺13] and [ZXX14]. We remark that a crucial step in all these works is the Hoeffding's inequality for uniformly ergodic Markov chains, obtained in [GO02]. Generalization of these results to irreducible aperiodic \mathcal{V} -ergodic Markov chains has been obtained in [ZXX14]. Actually the proof of the main results of this article (Theorems 1 and 2) is taken from [Vid03, Theorem 3.5]. We also remark that these results do not give generalization bounds in the full sense. Namely, a crucial assumption in the article is that the data-generating process starts from its invariant measure, which is unrealistic. A way to partly fix this gap is presented below.

Recall that the chain $\{Z_n\}_{n \geq 0}$ is said to be

- (i) irreducible if there exists a σ -finite measure $\varphi(dz)$ on Z such that whenever $\varphi(B) > 0$ we have $\sum_{n=0}^{\infty} \mathcal{P}^n(z, B) > 0$ for all $z \in Z$;
- (ii) aperiodic if there does not exist a partition $\{B_1, \dots, B_k\} \subseteq \mathfrak{B}(Z)$ with $k \geq 2$ of Z such that $\mathcal{P}(z, B_{i+1}) = 1$ for all $z \in B_i$ and all $1 \leq i \leq k-1$, and $\mathcal{P}(z, B_1) = 1$ for all $z \in B_k$.
- (iii) \mathcal{V} -ergodic if it admits an invariant probability measure $\pi(dz)$ and there are $\mathcal{V} : Z \rightarrow [1, \infty)$ with $\int_Z \mathcal{V}(z)\pi(dz) < \infty$, $C > 0$ and $\kappa \in (0, 1)$, such that

$$(1.4) \quad \|\mathcal{P}^n(z, d\bar{z}) - \pi(d\bar{z})\|_{\mathcal{V}} \leq C\mathcal{V}(z)\kappa^n$$

for all $z \in Z$ and $n \geq 0$.

Here, for a signed measure $\mu(dz)$ on Z ,

$$\|\mu(dz)\|_{\mathcal{V}} := \sup_{|f| \leq \mathcal{V}} |\mu(f)|.$$

Further, the β -mixing (or complete regularity, or the Kolmogorov) coefficient of the chain $\{Z_n\}_{n \geq 0}$ with initial distribution $\mu(dz)$ is defined as

$$\beta^{\mu}(n) := \sup_{m \geq 0} \mathbb{E}_{\mu} \left[\sup_{B \in \sigma\{Z_k : k \geq m+n\}} |\mathbb{P}_{\mu}(B | \mathcal{F}_m) - \mathbb{P}_{\mu}(B)| \right].$$

If $\pi(dz)$ is an invariant probability measure of $\{Z_n\}_{n \geq 0}$, then, by employing the Markov property and stationarity,

$$\beta^{\pi}(n) = \int_Z \|\mathcal{P}^n(z, d\bar{z}) - \pi(d\bar{z})\|_{\text{TV}} \pi(dz).$$

In particular, if $\{Z_n\}_{n \geq 0}$ is φ -irreducible, aperiodic and \mathcal{V} -ergodic with invariant probability measure $\pi(dz)$, then

$$\beta^{\pi}(n) \leq \int_Z \|\mathcal{P}^n(z, d\bar{z}) - \pi(d\bar{z})\|_{\mathcal{V}} \pi(dz) \leq C\pi(\mathcal{V})\kappa^n$$

for all $n \geq 0$. Under this assumptions, (A1) and

$$\overline{\text{(A2)}}: B := \sup_{z \in Z, h \in \mathcal{H}} \ell_h(z) < \infty \text{ and } |\ell_{h_1}(z) - \ell_{h_2}(z)| \leq \bar{L}\rho_{\mathcal{H}}(h_1, h_2) \text{ for some } \bar{L} > 0, \text{ and} \\ \text{all } z \in Z \text{ and } h_1, h_2 \in \mathcal{H}$$

in [ZXC12, Theorem 2] (as we have already commented, the proof of this result has been taken from [Vid03, Theorem 3.5]) it has been shown that for fixed $\varepsilon \in (0, 3B)$,

$$\mathbb{P}_{\pi} \left(\sup_{h \in \mathcal{H}} |\hat{r}_n(h) - \text{er}_{\pi}(h)| > \varepsilon \right) \leq 2(1 + C\pi(\mathcal{V})e^{-2})\mathcal{N}(\varepsilon/4\bar{L}, \mathcal{H}, \rho_{\mathcal{H}})e^{-\frac{\sqrt{n}\sqrt{\ln(\kappa^{-1})}\varepsilon^2}{16\sqrt{2}B^2}}$$

for all

$$n \geq n_0(\varepsilon) := \max \left\{ \frac{16 + 8\sqrt{4 + \ln(\kappa^{-1})^2} + 8\ln(\kappa^{-1})^2}{\ln(\kappa^{-1})^2}, \frac{\ln(\kappa^{-1})}{8} \right\}.$$

From this it follows (see the proof of Theorem 1.1) that for fixed $\varepsilon > 0$ and $\delta \in (0, 1)$, and an ε -ASEM algorithm $\mathcal{A}^{\varepsilon}$ for \mathcal{H} , we have that

$$(1.5) \quad \mathbb{P}_{\pi} \left(|\text{er}_{\pi}(\mathcal{A}^{\varepsilon}(Z_0, \dots, Z_{n-1})) - \text{opt}_{\pi}(\mathcal{H})| < 5\varepsilon \right) \geq 1 - \delta \quad \forall n \geq n_0(\varepsilon, \delta),$$

where

$$n_0(\varepsilon, \delta) := \max \left\{ n_0(\varepsilon), \frac{512B^4}{\ln(\kappa^{-1})\varepsilon^4} \ln \left(\frac{2(1 + C\pi(\mathcal{V})e^{-2})\mathcal{N}(\varepsilon/4\bar{L}, \mathcal{H}, \rho_{\mathcal{H}})}{\delta} \right)^2 \right\}.$$

However, as we have already commented, this result is restrictive and impractical since it assumes that the process is initially distributed according to the invariant distribution $\pi(dz)$ which

is in general unknown to us. It is more natural, and in the spirit of generalization bounds, that the relation in eq. (1.5) holds for every initial distribution of $\{Z_n\}_{n \geq 0}$. This result can be slightly generalized by employing (i) Hölder inequality, or (ii) ergodicity property in eq. (1.4).

- (i) Let $\mu(dz)$ be any initial distribution of $\{Z_n\}_{n \geq 0}$ such that $\mu \ll \pi$. Assume also that there is $p \in (1, \infty]$ such that $\frac{d\mu}{d\pi} \in L^p(\pi)$. Then,

$$\begin{aligned} & \mathbb{P}_\mu \left(\sup_{h \in \mathcal{H}} |\hat{e}r_n(h) - e r_\pi(h)| > \varepsilon \right) \\ &= \int_{\mathcal{Z}} \mathbb{P}_z \left(\sup_{h \in \mathcal{H}} |\hat{e}r_n(h) - e r_\pi(h)| > \varepsilon \right) \mu(dz) \\ &= \int_{\mathcal{Z}} \mathbb{P}_z \left(\sup_{h \in \mathcal{H}} |\hat{e}r_n(h) - e r_\pi(h)| > \varepsilon \right) \frac{d\mu}{d\pi}(z) \pi(dz) \\ &\leq \left\| \frac{d\mu}{d\pi} \right\|_p \left(\int_{\mathcal{Z}} \mathbb{P}_z \left(\sup_{h \in \mathcal{H}} |\hat{e}r_n(h) - e r_\pi(h)| > \varepsilon \right)^q \pi(dz) \right)^{1/q} \\ &\leq \left\| \frac{d\mu}{d\pi} \right\|_p \mathbb{P}_\pi \left(\sup_{h \in \mathcal{H}} |\hat{e}r_n(h) - e r_\pi(h)| > \varepsilon \right)^{1/q}, \end{aligned}$$

where $q \in [1, \infty)$ is such that $1/p + 1/q = 1$. Thus, for fixed $\varepsilon > 0$ and $\delta \in (0, 1)$, and an ε -ASEM algorithm \mathcal{A}^ε for \mathcal{H} , we have that

$$\mathbb{P}_\mu \left(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_0, \dots, Z_{n-1})) - \text{opt}_\pi(\mathcal{H})| < 5\varepsilon \right) \geq 1 - \delta$$

for all

$$n \geq \max \left\{ n_0(\varepsilon, \delta), \frac{512B^4 q^2}{\ln(\kappa^{-1})\varepsilon^4} \ln \left(\frac{\left((2(1 + C\pi(\mathcal{V})\varepsilon^{-2})\mathcal{N}(\varepsilon/4\bar{L}, \mathcal{H}, \rho_{\mathcal{H}})) \right)^{1/q} \left\| \frac{d\mu}{d\pi} \right\|_p \right)^2}{\delta} \right\}.$$

- (ii) Let now $\mu(dz)$ be any initial distribution of $\{Z_n\}_{n \geq 0}$ such that $\mu(\mathcal{V}) < \infty$. Fix $\varepsilon > 0$ and $\delta \in (0, 1)$, and let \mathcal{A}^ε be an ε -ASEM algorithm for \mathcal{H} . Then, for any $n \geq n_0(\varepsilon, \delta)$ the relation in eq. (1.5) implies

$$\begin{aligned} & \mathbb{P}_\mu \left(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_n, \dots, Z_{2n-1})) - \text{opt}_\pi(\mathcal{H})| \geq 5\varepsilon \right) \\ &\leq \mathbb{P}_\pi \left(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_0, \dots, Z_{n-1})) - \text{opt}_\pi(\mathcal{H})| \geq 5\varepsilon \right) \\ &\quad + \left| \mathbb{P}_\mu \left(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_n, \dots, Z_{2n-1})) - \text{opt}_\pi(\mathcal{H})| \geq 5\varepsilon \right) \right. \\ &\quad \left. - \mathbb{P}_\pi \left(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_0, \dots, Z_{n-1})) - \text{opt}_\pi(\mathcal{H})| \geq 5\varepsilon \right) \right| \\ &< \delta + \left| \mathbb{E}_\mu \left[\mathbb{P}_{Z_n} \left(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_0, \dots, Z_{n-1})) - \text{opt}_\pi(\mathcal{H})| \geq 5\varepsilon \right) \right] \right. \\ &\quad \left. - \mathbb{E}_\pi \left[\mathbb{P}_{Z_n} \left(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_0, \dots, Z_{n-1})) - \text{opt}_\pi(\mathcal{H})| \geq 5\varepsilon \right) \right] \right| \\ &\leq \delta + \left| \int_{\mathcal{Z}} \mathbb{P}_z \left(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_0, \dots, Z_{n-1})) - \text{opt}_\pi(\mathcal{H})| \geq 5\varepsilon \right) (\mu \mathcal{P}^n(dz) - \pi(dz)) \right| \\ &\leq \delta + \|\mu \mathcal{P}^n(dz) - \pi(dz)\|_{\text{TV}} \\ &\leq \delta + C\mu(\mathcal{V})\kappa^n. \end{aligned}$$

In addition to the above assumptions assume that $\{Z_n\}_{n \geq 0}$ is also reversible, i.e.

$$\int_A \mathcal{P}(z, B) \pi(dz) = \int_B \mathcal{P}(z, A) \pi(dz) \quad \forall A, B \in \mathfrak{B}(\mathcal{Z}).$$

Then, according to [KM12, Lemma 2.2] and [RR97, Theorem 2.1], it admits a spectral gap in $L^2(\pi)$, i.e.

$$\lambda := \sup_{f \in L^2(\pi), \|f\|_2=1} \|\mathcal{P}f - \pi(f)\|_2 < 1.$$

Now, from [Mia14, Theorem 3.3] it follows that

$$\mathbb{P}_\pi(|\hat{\text{er}}_n(h) - \text{er}_\pi(h)| > \varepsilon) \leq e^{-2\frac{1-\lambda}{1+\lambda}\varepsilon^2 n} \quad \forall n \geq 0.$$

From this, analogously as in the proof of Theorem 1.1, we conclude that

$$\mathbb{P}_\pi\left(\sup_{h \in \mathcal{H}} |\hat{\text{er}}_n(h) - \text{er}_\pi(h)| > \varepsilon\right) \leq \mathcal{N}(\varepsilon/4\bar{L}, \mathcal{H}, \rho_{\mathcal{H}}) e^{-\frac{1-\lambda}{2(1+\lambda)}\varepsilon^2 n} \quad \forall n \geq 0.$$

Let $\mu(dz)$ be any initial distribution of $\{Z_n\}_{n \geq 0}$ such that $\mu \ll \pi$ and $\frac{d\mu}{d\pi} \in L^p(\pi)$ for some $p \in (1, \infty]$. Then,

$$\mathbb{P}_\mu\left(\sup_{h \in \mathcal{H}} |\hat{\text{er}}_n(h) - \text{er}_\pi(h)| > \varepsilon\right) \leq \left\| \frac{d\mu}{d\pi} \right\|_p \mathbb{P}_\pi\left(\sup_{h \in \mathcal{H}} |\hat{\text{er}}_n(h) - \text{er}_\pi(h)| > \varepsilon\right)^{1/q},$$

where again $q \in [1, \infty)$ is such that $1/p + 1/q = 1$. From this it follows that for $\varepsilon > 0$, $\delta \in (0, 1)$ and an ε -ASEM algorithm \mathcal{A}^ε for \mathcal{H} , we have that

$$\mathbb{P}_\mu(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_0, \dots, Z_{n-1})) - \text{opt}_\pi(\mathcal{H})| < 5\varepsilon) \geq 1 - \delta$$

for all

$$n \geq \frac{2q(1+\lambda)}{(1-\lambda)} \log \left(\frac{\left\| \frac{d\mu}{d\pi} \right\|_p \mathcal{N}(\varepsilon/4\bar{L}, \mathcal{H}, \rho_{\mathcal{H}})^{1/q}}{\delta} \right).$$

Additionally, if $p = 2$ from [KM12, Proposition 1.5] we conclude

$$\begin{aligned} & \mathbb{P}_\mu(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_n, \dots, Z_{2n-1})) - \text{opt}_\pi(\mathcal{H})| \geq 5\varepsilon) \\ & \leq \mathbb{P}_\pi(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_0, \dots, Z_{n-1})) - \text{opt}_\pi(\mathcal{H})| \geq 5\varepsilon) \\ & \quad + \left| \mathbb{P}_\mu(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_n, \dots, Z_{2n-1})) - \text{opt}_\pi(\mathcal{H})| \geq 5\varepsilon) \right. \\ & \quad \left. - \mathbb{P}_\pi(|\text{er}_\pi(\mathcal{A}^\varepsilon(Z_0, \dots, Z_{n-1})) - \text{opt}_\mu(\mathcal{H})| \geq 5\varepsilon) \right| \\ & \leq \delta + \|\mu \mathcal{P}^n(dz) - \pi(dz)\|_{\text{TV}} \\ & \leq \delta + \frac{1}{2} \left(\int_{\mathcal{Z}} \left(1 - \frac{d\mu}{d\pi}(z)\right)^2 \pi(dz) \right)^{1/2} (1-\lambda)^n \end{aligned}$$

for all

$$n \geq \frac{2(1+\lambda)}{(1-\lambda)} \log \left(\frac{\mathcal{N}(\varepsilon/4\bar{L}, \mathcal{H}, \rho_{\mathcal{H}})}{\delta} \right).$$

2. PROOF OF THE MAIN RESULTS

In this section, we prove Theorems 1.1 and 1.2.

Proof of Theorem 1.1. Fix $h \in \mathcal{H}$ and $z_0 \in Z$. Then, (A2) and the boundedness of the metric ρ on Z , imply that $\|\ell_h^\pi\|_\infty \leq 2\|\ell_h\|_\infty \leq 2 \sup_{z_1, z_2 \in Z} \rho(z_1, z_2) + 2|\ell_h(z_0)| < \infty$, $\pi(\ell_h^\pi) = 0$, ℓ_h and ℓ_h^π are Lipschitz continuous, and $\text{Lip}(\ell_h^\pi) = \text{Lip}(\ell_h)$. Further, by employing Kantorovich-Rubinstein theorem, for any $z \in Z$, from eq. (1.2) it follows that

$$\mathbb{E}_z[\ell_h^\pi(Z_n)] \leq \text{Lip}(\ell_h^\pi) \mathcal{W}(\mathcal{P}^n(z, d\bar{z}), \pi(d\bar{z})) \leq C_1 \text{Lip}(\ell_h) e^{-C_2 n} \leq C_1 L e^{-C_2 n},$$

where in the last line we used assumption (A2), more precisely, we used that $|\ell_h(z_1) - \ell_h(z_2)| \leq L\rho(z_1, z_2)$. Thus, the function

$$(2.1) \quad g(z) := \sum_{n \geq 0} \mathbb{E}_z[\ell_h^\pi(Z_n)]$$

is well defined. Furthermore, the function $g(z)$ satisfies

$$(2.2) \quad \|g\|_\infty \leq C_1 L / (1 - e^{-C_2})$$

and solves the equation

$$(2.3) \quad g(z) - \mathbb{E}_z[g(Z_1)] = \ell_h^\pi(z).$$

Set now $W_i := \ell_h(Z_i)$ and $S_n := \sum_{i=0}^{n-1} W_i$ for $i \geq 0$ and $n \geq 1$. Thus,

$$S_n - n\pi(\ell_h) = n\hat{\text{er}}_n(h) - n\text{er}_\pi(h).$$

Fix now arbitrary $\epsilon > 0$. Analogously as in the proof of [GO02, Theorem 2] we see that

$$\begin{aligned} \mathbb{P}_z(|\hat{\text{er}}_n(h) - \text{er}_\pi(h)| > \epsilon) &= \mathbb{P}_z(|S_n - n\pi(\ell_h)| > n\epsilon) \\ &\leq e^{-(\epsilon n / \|g\|_\infty - 2)^2 / (2n)} \\ &\leq e^{-\left(\frac{\epsilon n(1 - e^{-C_2})}{2C_1 L} - 2\right)^2 / (2n)} \quad \forall n \geq \frac{4C_1 L}{\epsilon(1 - e^{-C_2})}. \end{aligned}$$

Further, let $\mathcal{C}_\epsilon = \{h_1, \dots, h_{\mathcal{N}(\epsilon, \mathcal{H}, d_{\mathcal{H}})}\}$ be the ϵ -cover of \mathcal{H} . Thus, for any $h \in \mathcal{H}$ there is $h_i \in \mathcal{C}_\epsilon$ such that $d_{\mathcal{H}}(h, h_i) < \epsilon$. We now have,

$$\begin{aligned} \mathbb{P}_z\left(\sup_{h \in \mathcal{H}} |\hat{\text{er}}_n(h) - \text{er}_\pi(h)| > 4\bar{L}\epsilon\right) &\leq \mathbb{P}_z\left(\bigcup_{i=1}^{\mathcal{N}(\epsilon, \mathcal{H}, d_{\mathcal{H}})} \left\{ \sup_{h \in \{\bar{h} : d_{\mathcal{H}}(\bar{h}, h_i) < \epsilon\}} |\hat{\text{er}}_n(h) - \text{er}_\pi(h)| > 4\bar{L}\epsilon \right\}\right) \\ &\leq \sum_{i=1}^{\mathcal{N}(\epsilon, \mathcal{H}, d_{\mathcal{H}})} \mathbb{P}_z\left(\sup_{h \in \{\bar{h} : d_{\mathcal{H}}(\bar{h}, h_i) < \epsilon\}} |\hat{\text{er}}_n(h) - \text{er}_\pi(h)| > 4\bar{L}\epsilon\right). \end{aligned}$$

For $h \in \{\bar{h} : d_{\mathcal{H}}(\bar{h}, h_i) < \epsilon\}$, from (A2), it follows that

$$|\hat{\text{er}}_n(h) - \text{er}_\pi(h) - \hat{\text{er}}_n(h_i) + \text{er}_\pi(h_i)| \leq 2\bar{L}\rho_{\mathcal{H}}(h, h_i) < 2\bar{L}\epsilon.$$

Thus,

$$\begin{aligned} \mathbb{P}_z\left(\sup_{h \in \{\bar{h} : d_{\mathcal{H}}(\bar{h}, h_i) < \epsilon\}} |\hat{\text{er}}_n(h) - \text{er}_\pi(h)| > 4\bar{L}\epsilon\right) &\leq \mathbb{P}_z(|\hat{\text{er}}_n(h_i) - \text{er}_\pi(h_i)| > 2\bar{L}\epsilon) \\ &\leq e^{-\left(\frac{\epsilon \bar{L} n(1 - e^{-C_2})}{C_1 L} - 2\right)^2 / (2n)} \quad \forall n \geq \frac{2C_1 L}{\bar{L}\epsilon(1 - e^{-C_2})}, \end{aligned}$$

which implies

$$(2.4) \quad \mathbb{P}_z\left(\sup_{h \in \mathcal{H}} |\hat{\text{er}}_n(h) - \text{er}_\pi(h)| > 4\bar{L}\epsilon\right) \leq \mathcal{N}(\epsilon, \mathcal{H}, d_{\mathcal{H}}) e^{-\left(\frac{\epsilon \bar{L} n(1 - e^{-C_2})}{C_1 L} - 2\right)^2 / (2n)}$$

for all $n \geq 2C_1 L / \bar{L}\epsilon(1 - e^{-C_2})$.

Finally, set $\varepsilon := 4\bar{L}\varepsilon$, let \mathcal{A}^ε be the ε -ASEM algorithm for \mathcal{H} , and fix $\delta \in (0, 1)$. Then,

$$\mathbb{P}_z \left(\sup_{h \in \mathcal{H}} |\hat{\text{er}}_n(h) - \text{er}_\pi(h)| \leq \varepsilon \right) \geq 1 - \delta \quad \forall n \geq n_1(\varepsilon, \delta).$$

Consequently, for all $n \geq n_0(\varepsilon, \delta)$ it follows that with probability at least $1 - \delta$,

$$\begin{aligned} \text{er}_\pi(\mathcal{A}^\varepsilon(Z_0, \dots, Z_{n-1})) &\leq \text{er}_\pi(h_{(Z_0, \dots, Z_{n-1})}) + \varepsilon/4 \\ &\leq \hat{\text{er}}_n(h_{(Z_0, \dots, Z_{n-1})}) + 5\varepsilon/4 \\ &\leq \hat{\text{er}}_n(\mathcal{A}^\varepsilon(Z_0, \dots, Z_{n-1})) + 6\varepsilon/4 \\ &\leq \inf_{h \in \mathcal{H}} \hat{\text{er}}_n(h) + 10\varepsilon/4 \\ &\leq \hat{\text{er}}_n(\bar{h}) + 5\varepsilon/2 \\ &\leq \hat{\text{er}}_n(h_i) + 11\varepsilon/4 \\ &\leq \text{er}_\pi(h_i) + 15\varepsilon/4 \\ &\leq \text{er}_\pi(\bar{h}) + 4\varepsilon \\ &< \text{opt}_\pi(\mathcal{H}) + 5\varepsilon, \end{aligned}$$

where $h_{(Z_0, \dots, Z_{n-1})} \in \mathcal{C}_{\varepsilon/4\bar{L}}$, $\bar{h} \in \mathcal{H}$ is such that $\text{er}_\pi(\bar{h}) < \text{opt}_\pi(\mathcal{H}) + \varepsilon$, and $h_i \in \mathcal{C}_{\varepsilon/4\bar{L}}$ such that $d_{\mathcal{H}}(\bar{h}, h_i) < \varepsilon/4\bar{L}$. \square

Proof of Theorem 1.2. We again follow the proof of [GO02, Theorem 2]. For $i \geq 0$ and $z \in \mathcal{Z}$, set

$$D_i := g(Z_i) - \mathbb{E}_z[g(Z_i) \mid Z_0, \dots, Z_{i-1}] = g(Z_i) - \mathbb{E}_{Z_{i-1}}[g(Z_i)].$$

Recall that the function $g(z)$ is given by eq. (2.1). Since $\mathbb{E}[D_i] = 0$ for all $i \geq 0$, $\{D_i\}_{i \geq 0}$ is a martingale difference sequence. Using Markov property and eq. (2.3) we get

$$S_n - n\pi(\ell_h) = \sum_{i=1}^n D_i + g(Z_0) - g(Z_n) \leq \sum_{i=1}^n D_i + 2\|g\|_\infty.$$

From this it follows that

$$\begin{aligned} \mathbb{E}_z \left[e^{\theta(S_n - n\pi(\ell_h))} \right] &\leq e^{2\theta\|g\|_\infty} \mathbb{E}_z \left[e^{\theta \sum_{i=1}^n D_i} \right] \\ &= e^{2\theta\|g\|_\infty} \mathbb{E}_z \left[e^{\theta \sum_{i=1}^{n-1} D_i} \mathbb{E} \left[e^{\theta D_n} \mid Z_0, \dots, Z_{n-1} \right] \right]. \end{aligned}$$

Since $|D_n| \leq 2\|g\|_\infty$, applying [DGL96, Lemma 8.1] with expectation conditional to Z_0, \dots, Z_{n-1} , $a = -2\|g\|_\infty$ and $b = 2\|g\|_\infty$ gives us

$$\mathbb{E}_z \left[e^{\theta D_n} \mid Z_0, \dots, Z_{n-1} \right] \leq e^{2\theta^2\|g\|_\infty^2}.$$

Iterating this procedure results in

$$\mathbb{E}_z \left[e^{\theta \sum_{i=1}^n D_i} \right] \leq e^{2n\theta^2\|g\|_\infty^2}.$$

Hence, we have

$$\mathbb{E}_z \left[e^{\theta(S_n - n\pi(\ell_h))} \right] \leq e^{2\theta\|g\|_\infty + 2n\theta^2\|g\|_\infty^2}.$$

Using Markov's inequality, we get

$$\mathbb{P}_z \left(S_n - n\pi(\ell_h) \geq n\varepsilon \right) \leq \frac{\mathbb{E}_z \left[e^{\theta(S_n - n\pi(\ell_h))} \right]}{e^{\theta n\varepsilon}} \leq e^{2n\|g\|_\infty^2\theta^2 + (2\|g\|_\infty - n\varepsilon)\theta}.$$

Notice that if we define D_i as

$$D_i := \mathbb{E}_z \left[g(Z_i) \mid Z_0, \dots, Z_{i-1} \right] - g(Z_i)$$

and repeat the same steps, we would get

$$\mathbb{P}_z \left(n \pi(\ell_h) - S_n \geq n\varepsilon \right) \leq e^{2n\|g\|_\infty^2 \theta^2 + (2\|g\|_\infty - n\varepsilon)\theta}.$$

Therefore

$$\begin{aligned} \mathbb{P}_z \left(|S_n - n \pi(\ell_h)| \geq n\varepsilon \right) &= \mathbb{P}_z \left(S_n - n \pi(\ell_h) \geq n\varepsilon \right) + \mathbb{P}_z \left(n \pi(\ell_h) - S_n \geq n\varepsilon \right) \\ &\leq 2e^{2n\|g\|_\infty^2 \theta^2 + (2\|g\|_\infty - n\varepsilon)\theta}. \end{aligned}$$

This relation holds for all $\theta > 0$. The best bound is obtained for θ that minimizes the function $\theta \mapsto 2n\|g\|_\infty^2 \theta^2 + (2\|g\|_\infty - n\varepsilon)\theta$. It is easy to see that the value of θ that minimizes this function is

$$\theta_n = \frac{n\varepsilon - 2\|g\|_\infty}{4n\|g\|_\infty^2}.$$

Hence,

$$\mathbb{P}_z \left(|S_n - n \pi(\ell_h)| \geq n\varepsilon \right) \leq 2 \exp \left(-\frac{(n\varepsilon - 2\|g\|_\infty)^2}{8n\|g\|_\infty^2} \right).$$

Using eq. (2.2), for every $n \geq \frac{2C_1 L}{\varepsilon(1-e^{-C_2})}$ we get

$$(n\varepsilon - 2\|g\|_\infty)^2 \geq \left(n\varepsilon - \frac{2C_1 L}{1-e^{-C_2}} \right)^2.$$

This, together with $8n\|g\|_\infty^2 \leq 8nC_1^2 L^2 (1-e^{-C_2})^{-2}$ gives

$$-\frac{(n\varepsilon - 2\|g\|_\infty)^2}{8n\|g\|_\infty^2} \leq -\frac{(n\varepsilon - 2C_1 L(1-e^{-C_2})^{-1})^2}{8nC_1^2 L^2 (1-e^{-C_2})^{-2}} = -\frac{(n\varepsilon(1-e^{-C_2}) - 2C_1 L)^2}{8nC_1^2 L^2}.$$

Therefore,

$$\mathbb{P}_z \left(|S_n - n \pi(\ell_h)| \geq n\varepsilon \right) \leq 2 \exp \left(-\frac{(n\varepsilon(1-e^{-C_2}) - 2C_1 L)^2}{8nC_1^2 L^2} \right).$$

Combining this with relations $\hat{e}_n(h) - e_{r_\pi}(h) = \frac{1}{n}(S_n - n \pi(\ell_h))$ and $e_{r_\pi}(h) \geq m$ (recall that for every $h \in \mathcal{H}$ it holds that $e_{r_\pi}(h) \geq m$), we get that for every $n \geq \frac{2C_1 L}{\varepsilon \sqrt{m(1-e^{-C_2})}}$ it holds

$$\begin{aligned} \mathbb{P}_z \left(\frac{|\hat{e}_n(h) - e_{r_\pi}(h)|}{\sqrt{e_{r_\pi}(h)}} \geq \varepsilon \right) &\leq \mathbb{P}_z \left(|S_n - n \pi(\ell_h)| \geq n\varepsilon \sqrt{m} \right) \\ &\leq 2 \exp \left(-\frac{(n\varepsilon \sqrt{m}(1-e^{-C_2}) - 2C_1 L)^2}{8nC_1^2 L^2} \right). \end{aligned}$$

Let $\varepsilon > 0$ be arbitrary and denote with $\mathcal{C}_{\varepsilon/\bar{L}} = \{h_1, h_2, \dots, h_{\mathcal{N}(\varepsilon/\bar{L}, \mathcal{H}, d_{\mathcal{H}})}\}$ the ε/\bar{L} -cover of \mathcal{H} . By subadditivity, we have

$$\mathbb{P}_z \left(\sup_{h \in \mathcal{H}} \frac{|\hat{e}_n(h) - e_{r_\pi}(h)|}{\sqrt{e_{r_\pi}(h)}} \geq \varepsilon \right) \leq \sum_{i=1}^{\mathcal{N}(\varepsilon/\bar{L}, \mathcal{H}, d_{\mathcal{H}})} \mathbb{P}_z \left(\sup_{h \in \mathcal{H}_i} \frac{|\hat{e}_n(h) - e_{r_\pi}(h)|}{\sqrt{e_{r_\pi}(h)}} \geq \varepsilon \right),$$

where $\mathcal{H}_i := \{\bar{h} : d_{\mathcal{H}}(\bar{h}, h_i) < \varepsilon/\bar{L}\}$. Our goal now is to provide an upper bound for the summands on the right hand side. Since it can happen that $\hat{e}_n(h) \geq e_{r_\pi}(h)$ and $\hat{e}_n(h) \leq e_{r_\pi}(h)$, we will inspect expressions $e_{r_\pi}(h) - \hat{e}_n(h)$ and $\hat{e}_n(h) - e_{r_\pi}(h)$ separately.

Part 1: $e_{r_\pi}(h) - \hat{e}_n(h)$. For $\eta \in (0, 1)$, define

$$\phi(h) := (1 - \eta)e_{r_\pi}(h) - \hat{e}_n(h).$$

For any $h \in \mathcal{H}_i$ we have that

$$\begin{aligned} \phi(h) - \phi(h_i) &= (1 - \eta)(\text{er}_\pi(h) - \text{er}_\pi(h_i)) + (\hat{\text{er}}_n(h_i) - \hat{\text{er}}_n(h)) \\ &\leq (1 - \eta) \int_{\mathcal{Z}} |\ell_h(z) - \ell_{h_i}(z)| \pi(\text{d}z) + \frac{1}{n} \sum_{j=0}^{n-1} |\ell_{h_i}(z_j) - \ell_h(z_j)| \\ &\leq (2 - \eta)\varepsilon, \end{aligned}$$

where we used assumption (A2), more precisely, we used that

$$|\ell_h(z) - \ell_{h_i}(z)| \leq \bar{L} \text{d}_{\mathcal{H}}(h, h_i) \leq \bar{L} \cdot \frac{\varepsilon}{\bar{L}} = \varepsilon.$$

Hence, if $\sup_{h \in \mathcal{H}_i} \phi(h) \geq 2\varepsilon(2 - \eta)$, then $\phi(h_i) \geq \varepsilon(2 - \eta)$, i.e. for $i \in \{1, 2, \dots, \mathcal{N}(\varepsilon/\bar{L}, \mathcal{H}, \text{d}_{\mathcal{H}})\}$ we have

$$\underbrace{\mathbb{P}_z(\phi(h_i) \geq \varepsilon(2 - \eta))}_{=: I_1} \geq \underbrace{\mathbb{P}_z\left(\sup_{h \in \mathcal{H}_i} \phi(h) \geq 2\varepsilon(2 - \eta)\right)}_{=: I_2}.$$

Recall that for every $h \in \mathcal{H}$ it holds $m \leq \text{er}_\pi(h) \leq M$. Setting $\eta := \varepsilon/\text{er}_\pi(h_i)$ and taking $0 < \varepsilon < 2m/3$, we get

$$\begin{aligned} I_1 &= \mathbb{P}_z\left((1 - \eta)\text{er}_\pi(h_i) - \hat{\text{er}}_n(h_i) \geq \varepsilon(2 - \eta)\right) \\ &= \mathbb{P}_z\left(\text{er}_\pi(h_i) - \hat{\text{er}}_n(h_i) \geq \varepsilon + \varepsilon\left(2 - \frac{\varepsilon}{\text{er}_\pi(h_i)}\right)\right) \\ &= \mathbb{P}_z\left(\frac{\text{er}_\pi(h_i) - \hat{\text{er}}_n(h_i)}{\sqrt{\text{er}_\pi(h_i)}} \geq \frac{\varepsilon}{\sqrt{\text{er}_\pi(h_i)}}\left(3 - \frac{\varepsilon}{\text{er}_\pi(h_i)}\right)\right) \\ &\leq \mathbb{P}_z\left(\frac{\text{er}_\pi(h_i) - \hat{\text{er}}_n(h_i)}{\sqrt{\text{er}_\pi(h_i)}} \geq \frac{\varepsilon}{\sqrt{M}}\left(3 - \frac{\varepsilon}{m}\right)\right) \\ &\leq \mathbb{P}_z\left(\frac{\text{er}_\pi(h_i) - \hat{\text{er}}_n(h_i)}{\sqrt{\text{er}_\pi(h_i)}} \geq \frac{7\varepsilon}{3\sqrt{M}}\right). \end{aligned}$$

In a similar fashion one can obtain

$$I_2 \geq \mathbb{P}_z\left(\sup_{h \in \mathcal{H}_i} \frac{\text{er}_\pi(h) - \hat{\text{er}}_n(h)}{\sqrt{\text{er}_\pi(h)}} \geq \frac{M + 4m}{m^{3/2}}\varepsilon\right).$$

Since $I_1 \geq I_2$, we have

$$\mathbb{P}_z\left(\frac{\text{er}_\pi(h_i) - \hat{\text{er}}_n(h_i)}{\sqrt{\text{er}_\pi(h_i)}} \geq \frac{7\varepsilon}{3\sqrt{M}}\right) \geq \mathbb{P}_z\left(\sup_{h \in \mathcal{H}_i} \frac{\text{er}_\pi(h) - \hat{\text{er}}_n(h)}{\sqrt{\text{er}_\pi(h)}} \geq \frac{M + 4m}{m^{3/2}}\varepsilon\right).$$

Part 2: $\hat{\text{er}}_n(h) - \text{er}_\pi(h)$. For $\eta \in (0, 1)$, define

$$\tilde{\phi}(h) := \hat{\text{er}}_n(h) - (1 + \eta)\text{er}_\pi(h).$$

Analogously as in *Part 1*, we obtain

$$\mathbb{P}_z\left(\sup_{h \in \mathcal{H}_i} \frac{\hat{\text{er}}_n(h) - \text{er}_\pi(h)}{\sqrt{\text{er}_\pi(h)}} \geq \frac{M + 4m}{m^{3/2}}\varepsilon\right) \leq \mathbb{P}_z\left(\frac{\hat{\text{er}}_n(h_i) - \text{er}_\pi(h_i)}{\sqrt{\text{er}_\pi(h_i)}} \geq \frac{3\varepsilon}{\sqrt{M}}\right).$$

Since $(M + 6m)/m^{3/2} > (M + 4m)/m^{3/2}$ and $3/\sqrt{M} > 7/(3\sqrt{M})$, we get

$$\begin{aligned}
& \mathbb{P}_z \left(\sup_{h \in \mathcal{H}_i} \frac{|\hat{e}_n(h) - e_{r_\pi}(h)|}{\sqrt{e_{r_\pi}(h)}} \geq \frac{M + 6m}{m^{3/2}} \varepsilon \right) \\
& \leq \mathbb{P}_z \left(\sup_{h \in \mathcal{H}_i} \frac{e_{r_\pi}(h) - \hat{e}_n(h)}{\sqrt{e_{r_\pi}(h)}} \geq \frac{M + 6m}{m^{3/2}} \varepsilon \right) + \mathbb{P}_z \left(\sup_{h \in \mathcal{H}_i} \frac{\hat{e}_n(h) - e_{r_\pi}(h)}{\sqrt{e_{r_\pi}(h)}} \geq \frac{M + 6m}{m^{3/2}} \varepsilon \right) \\
& \leq \mathbb{P}_z \left(\frac{|e_{r_\pi}(h_i) - \hat{e}_n(h_i)|}{\sqrt{e_{r_\pi}(h_i)}} \geq \frac{7\varepsilon}{3\sqrt{M}} \right) + \mathbb{P}_z \left(\frac{|\hat{e}_n(h_i) - e_{r_\pi}(h_i)|}{\sqrt{e_{r_\pi}(h_i)}} \geq \frac{7\varepsilon}{3\sqrt{M}} \right) \\
& \leq 2\mathbb{P}_z \left(\frac{|e_{r_\pi}(h_i) - \hat{e}_n(h_i)|}{\sqrt{e_{r_\pi}(h_i)}} \geq \frac{7\varepsilon}{3\sqrt{M}} \right).
\end{aligned}$$

Setting $\varepsilon' := m^{3/2}\varepsilon/(M + 6m)$, we have

$$\begin{aligned}
& \mathbb{P}_z \left(\sup_{h \in \mathcal{H}} \frac{|\hat{e}_n(h) - e_{r_\pi}(h)|}{\sqrt{e_{r_\pi}(h)}} \geq \varepsilon \right) \\
& \leq \sum_{i=1}^{\mathcal{N}(\varepsilon/\bar{L}, \mathcal{H}, d_{\mathcal{H}})} \mathbb{P}_z \left(\sup_{h \in \mathcal{H}_i} \frac{|\hat{e}_n(h) - e_{r_\pi}(h)|}{\sqrt{e_{r_\pi}(h)}} \geq \varepsilon \right) \\
& = \sum_{i=1}^{\mathcal{N}(\varepsilon/\bar{L}, \mathcal{H}, d_{\mathcal{H}})} \mathbb{P}_z \left(\sup_{h \in \mathcal{H}_i} \frac{|\hat{e}_n(h) - e_{r_\pi}(h)|}{\sqrt{e_{r_\pi}(h)}} \geq \frac{M + 6m}{m^{3/2}} \varepsilon' \right) \\
(2.5) \quad & \leq 2 \sum_{i=1}^{\mathcal{N}(\varepsilon/\bar{L}, \mathcal{H}, d_{\mathcal{H}})} \mathbb{P}_z \left(\frac{|\hat{e}_n(h_i) - e_{r_\pi}(h_i)|}{\sqrt{e_{r_\pi}(h_i)}} \geq \frac{7\varepsilon'}{3\sqrt{M}} \right) \\
& \leq 4 \sum_{i=1}^{\mathcal{N}(\varepsilon/\bar{L}, \mathcal{H}, d_{\mathcal{H}})} \exp \left(-\frac{(7\varepsilon' n \sqrt{m}(1 - e^{-C_2}) - 2C_1 L)^2}{72Mn C_1^2 L^2} \right) \\
& \leq 4\mathcal{N}(\varepsilon/\bar{L}, \mathcal{H}, d_{\mathcal{H}}) e^{-\xi_1 \varepsilon^2 n + \xi_2 \varepsilon},
\end{aligned}$$

where

$$\xi_1 = \frac{49m^4(1 - e^{-C_2})^2}{72M(M + 6m)^2 C_1^2 L^2} \quad \text{and} \quad \xi_2 = \frac{7m^2(1 - e^{-C_2})}{6\sqrt{M}(M + 6m)C_1 L}.$$

Hence, to have

$$\mathbb{P}_z \left(\sup_{h \in \mathcal{H}} \frac{|\hat{e}_n(h) - e_{r_\pi}(h)|}{\sqrt{e_{r_\pi}(h)}} \geq \varepsilon \right) \leq \delta,$$

it suffices to have

$$4\mathcal{N}(\varepsilon/\bar{L}, \mathcal{H}, d_{\mathcal{H}}) e^{-\xi_1 \varepsilon^2 n + \xi_2 \varepsilon} \leq \delta,$$

which is equivalent to

$$n \geq \frac{1}{\xi_1 \varepsilon^2} \left(\xi_2 \varepsilon + \log \left(\frac{4\mathcal{N}(\varepsilon/\bar{L}, \mathcal{H}, d_{\mathcal{H}})}{\delta} \right) \right),$$

and this is exactly what we wanted to prove. \square

Proof of Corollary 1.3. Define

$$\tilde{\varepsilon} := \sqrt{\frac{\varepsilon}{1 + \frac{1}{\alpha}}},$$

where $\varepsilon > 0$ is the one from the statement of this result. Using

$$n \geq \frac{2C_1 L \sqrt{1 + 1/\alpha}}{\sqrt{\varepsilon}(\sqrt{m} \wedge 1)(1 - e^{-C_2})} = \frac{2C_1 L}{\tilde{\varepsilon}(\sqrt{m} \wedge 1)(1 - e^{-C_2})},$$

together with eq. (2.5) we get

$$\mathbb{P}_z \left(\sup_{h \in \mathcal{H}} \frac{|\hat{e}_n(h) - e_{\pi}(h)|}{\sqrt{e_{\pi}(h)}} \geq \tilde{\varepsilon} \right) \leq 4\mathcal{N}(\tilde{\varepsilon}/\bar{L}, \mathcal{H}, d_{\mathcal{H}}) e^{-\xi_1 \tilde{\varepsilon}^2 n + \xi_2 \tilde{\varepsilon}}.$$

Hence, we obviously have

$$(2.6) \quad \mathbb{P}_z \left(\exists h \in \mathcal{H} : \frac{e_{\pi}(h) - \hat{e}_n(h)}{\sqrt{e_{\pi}(h)}} \geq \tilde{\varepsilon} \right) \leq 4\mathcal{N}(\tilde{\varepsilon}/\bar{L}, \mathcal{H}, d_{\mathcal{H}}) e^{-\xi_1 \tilde{\varepsilon}^2 n + \xi_2 \tilde{\varepsilon}}.$$

We now follow the proof of [AB99, Theorem 5.8]. Suppose that

$$(2.7) \quad e_{\pi}(h) - \hat{e}_n(h) \leq \tilde{\varepsilon} \sqrt{e_{\pi}(h)}.$$

Then, for $\alpha > 0$ from the statement of this result (arbitrary, but fixed) we have two cases.

Case 1: If $e_{\pi}(h) < (1 + 1/\alpha)^2 \tilde{\varepsilon}^2$ then eq. (2.7) implies

$$e_{\pi}(h) \leq \hat{e}_n(h) + \tilde{\varepsilon}^2 \left(1 + \frac{1}{\alpha}\right).$$

Case 2: If $e_{\pi}(h) \geq (1 + 1/\alpha)^2 \tilde{\varepsilon}^2$ then $\tilde{\varepsilon} \leq \sqrt{e_{\pi}(h)}/(1 + 1/\alpha)$. Hence, from eq. (2.7) it follows that

$$e_{\pi}(h) \leq \hat{e}_n(h) + \frac{\alpha}{\alpha + 1} \cdot e_{\pi}(h),$$

which is equivalent to

$$e_{\pi}(h) \leq (1 + \alpha) \hat{e}_n(h).$$

Therefore, we have just showed that eq. (2.7) implies

$$e_{\pi}(h) \leq (1 + \alpha) \hat{e}_n(h) + \tilde{\varepsilon}^2 \left(1 + \frac{1}{\alpha}\right) = (1 + \alpha) \hat{e}_n(h) + \varepsilon.$$

Combining this with eq. (2.6), we get

$$\begin{aligned} \mathbb{P}_z \left(\exists h \in \mathcal{H} : e_{\pi}(h) > (1 + \alpha) \hat{e}_n(h) + \varepsilon \right) &\leq \mathbb{P}_z \left(\exists h \in \mathcal{H} : \frac{e_{\pi}(h) - \hat{e}_n(h)}{\sqrt{e_{\pi}(h)}} > \tilde{\varepsilon} \right) \\ &\leq 4\mathcal{N}(\tilde{\varepsilon}/\bar{L}, \mathcal{H}, d_{\mathcal{H}}) e^{-\xi_1 \tilde{\varepsilon}^2 n + \xi_2 \tilde{\varepsilon}} \\ &= 4\mathcal{N} \left(\frac{\varepsilon^{1/2}}{\bar{L} \sqrt{1 + 1/\alpha}}, \mathcal{H}, d_{\mathcal{H}} \right) \exp \left(-\xi_1 \cdot \frac{\varepsilon}{1 + \frac{1}{\alpha}} \cdot n + \xi_2 \cdot \sqrt{\frac{\varepsilon}{1 + \frac{1}{\alpha}}} \right), \end{aligned}$$

and this is clearly less than or equal to δ for

$$n \geq \frac{\alpha + 1}{\alpha \xi_1} \cdot \frac{1}{\varepsilon} \left(\frac{\xi_2}{\sqrt{1 + 1/\alpha}} \cdot \varepsilon^{1/2} + \ln \left(\frac{4\mathcal{N} \left(\frac{\varepsilon^{1/2}}{\bar{L} \sqrt{1 + 1/\alpha}}, \mathcal{H}, d_{\mathcal{H}} \right)}{\delta} \right) \right),$$

which is exactly what we wanted to prove. \square

3. APPENDIX

In this section, we prove the auxiliary result used in the construction of the example satisfying condition (A3).

Lemma 3.1. *Let $\{X_n\}_{n \geq 0}$ be a Markov chain on X with transition kernel $\mathcal{P}_X(x, d\bar{x})$, and let $f : X \rightarrow f(X) \subseteq Y$ be measurable. The family of random variables $\{f(X_n)\}_{n \geq 0}$ is a Markov chain on Y if $\{f^{-1}(\{y\})\}_{y \in f(X)}$ forms a family of atoms for $\{X_n\}_{n \geq 0}$ on $f^{-1}(f(X))$, i.e. for any $y \in f(X)$ and $x_1, x_2 \in f^{-1}(\{y\})$,*

$$\mathcal{P}_X(x_1, f^{-1}(B)) = \mathcal{P}_X(x_2, f^{-1}(B)).$$

In that case, transition kernel of $\{f(X_n)\}_{n \geq 0}$ is given by

$$\mathcal{P}(y, B) = \mathcal{P}_X(x, f^{-1}(B))$$

for $y \in f(X)$ and $x \in f^{-1}(\{y\})$.

Proof. By assumption, $\{X_n\}_{n \geq 0}$ is defined on $(\Omega, \mathcal{F}, \mathbb{P}_x)_{x \in X}$. Now, on (Ω, \mathcal{F}) define a family of probability measures $\{\bar{\mathbb{P}}_y\}_{y \in f(X)}$ by $\bar{\mathbb{P}}_y(f(X_n) \in B) := \mathbb{P}_x(X_n \in f^{-1}(B))$ for $n \geq 0$, $y \in f(X)$ and $x \in f^{-1}(\{y\})$. By assumption, $\{\bar{\mathbb{P}}_y\}_{y \in f(X)}$ is well defined, and the assertion now follows directly. \square

ACKNOWLEDGEMENTS

Financial support through *Alexander von Humboldt Foundation* (No. HRV 1151902 HFST-E) and *Croatian Science Foundation* under project 8958 (for N. Sandrić), and the *Croatian Science Foundation* under project 4197 (for S. Šebek) is gratefully acknowledged.

REFERENCES

- [AB99] M. Anthony and P. L. Bartlett. *Neural network learning: theoretical foundations*. Cambridge University Press, Cambridge, 1999.
- [BKP97] P. L. Bartlett, S. R. Kulkarni, and S. E. Posner. Covering numbers for real-valued function classes. *IEEE Trans. Inform. Theory*, 43(5):1721–1724, 1997.
- [But14] O. Butkovsky. Subgeometric rates of convergence of Markov processes in the Wasserstein metric. *Ann. Appl. Probab.*, 24(2):526–552, 2014.
- [DGL96] L. Devroye, L. Györfi, and G. Lugosi. *A probabilistic theory of pattern recognition*, volume 31 of *Applications of Mathematics (New York)*. Springer-Verlag, New York, 1996.
- [Gam03] D. Gamarnik. Extension of the PAC framework to finite and countable Markov chains. *IEEE Trans. Inform. Theory*, 49(1):338–345, 2003.
- [GO02] P. W. Glynn and D. Ormoneit. Hoeffding’s inequality for uniformly ergodic Markov chains. *Statist. Probab. Lett.*, 56(2):143–146, 2002.
- [KM12] I. Kontoyiannis and S. P. Meyn. Geometric ergodicity and the spectral gap of non-reversible Markov chains. *Probab. Theory Related Fields*, 154(1-2):327–339, 2012.
- [Mei00] R. Meir. Nonparametric time series prediction through adaptive model selection. *Machine Learning*, 39(1):5–34, 2000.
- [Mia14] B. Miasojedow. Hoeffding’s inequalities for geometrically ergodic Markov chains on general state space. *Statist. Probab. Lett.*, 87:115–120, 2014.
- [MM96] D. S. Modha and E. Masry. Minimum complexity regression estimation with weakly dependent observations. *IEEE Trans. Inform. Theory*, 42(6, part 2):2133–2145, 1996.
- [MT09] S. Meyn and R. L. Tweedie. *Markov chains and stochastic stability*. Cambridge University Press, Cambridge, second edition, 2009.
- [RR97] G. O. Roberts and J. S. Rosenthal. Geometric ergodicity and hybrid Markov chains. *Electron. Comm. Probab.*, 2:13–25, 1997.
- [SHS09] I. Steinwart, D. Hush, and C. Scovel. Learning from dependent observations. *J. Multivariate Anal.*, 100(1):175–194, 2009.
- [SSBD14] S. Shalev-Shwartz and S. Ben-David. *Understanding Machine Learning - From Theory to Algorithms*. Cambridge University Press, 2014.

- [Vid03] M. Vidyasagar. *Learning and generalization*. Springer-Verlag London, Ltd., London, 2003.
- [Yu94] B. Yu. Rates of convergence for empirical processes of stationary mixing sequences. *Ann. Probab.*, 22(1):94–116, 1994.
- [Zho03] D. X. Zhou. Capacity of reproducing kernel spaces in learning theory. *IEEE Trans. Inform. Theory*, 49(7):1743–1752, 2003.
- [ZLX12] B. Zou, L. Li, and Z. Xu. Generalization performance of least-square regularized regression algorithm with Markov chain samples. *J. Math. Anal. Appl.*, 388(1):333–343, 2012.
- [ZLX⁺13] B. Zou, L. Li, Z. Xu, T. Luo, and Y. Y. Tang. Generalization performance of Fisher linear discriminant based on Markov sampling. *IEEE Trans. Neural. Netw. Learn. Syst.*, 24(2):288–300, 2013.
- [ZXC12] B. Zou, Z. Xu, and X. Chang. Generalization bounds of ERM algorithm with V -geometrically ergodic Markov chains. *Adv. Comput. Math.*, 36(1):99–114, 2012.
- [ZXX14] B. Zou, Z. Xu, and J. Xu. Generalization bounds of ERM algorithm with Markov chain samples. *Acta Math. Appl. Sin. Engl. Ser.*, 30(1):223–238, 2014.
- [ZZX09] B. Zou, H. Zhang, and Z. Xu. Learning from uniformly ergodic Markov chains. *J. Complexity*, 25(2):188–200, 2009.

(Nikola Sandrić) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ZAGREB, ZAGREB, CROATIA
Email address: nikola.sandric@math.hr

(Stjepan Šebek) DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF ELECTRICAL ENGINEERING AND COMPUTING, UNIVERSITY OF ZAGREB, ZAGREB, CROATIA
Email address: stjepan.sebek@fer.hr