

# DYNAMIC PRECISE AND IMPRECISE PROBABILITY KINEMATICS

MICHELE CAPRIO AND RUOBIN GONG

**ABSTRACT.** We introduce a new method for updating subjective beliefs based on Jeffrey’s rule of conditioning, called dynamic (precise) probability kinematics (DPK). We also give its generalization in order to work with sets of probabilities, called dynamic imprecise probability kinematics (DIPK). Updating a set of probabilities may be computationally costly. To this end, we provide bounds for the lower probability associated with the updated probability set, characterizing the set completely. The behavior of the updated sets of probabilities are studied, including contraction, dilation, and sure loss. We discuss the application of DPK and DIPK to survey sampling studies in which coarse and imprecise observations are anticipated.

## 1. INTRODUCTION

In uncertainty reasoning, probability distributions are the single most widely used mathematical object to encode subjective belief and uncertain evidence. There are situations in which sets of probabilities, rather than a single precise probability, better captures the imprecision occurring when an agent’s initial beliefs are elicited and translated. The structural imprecision in a set of probabilities allows for the analysis to be robust against coarsely specified, or potentially misspecified model evidence.

In this paper, we present a new way to update subjective beliefs based on Jeffrey’s rule of conditioning in a dynamic setting. The method can be applied to beliefs in the form of precise probabilities, which we call dynamic probability kinematics (DPK), and can be generalized to work with sets of probabilities, which we call dynamic imprecise probability kinematics (DIPK). The contribution of the DPK and DIPK frameworks proposed in this paper hinges on the premise that Jeffrey’s rule presents a compelling alternative to Bayes’ rule as a reasonable way to update initial beliefs. In this Section, we motivate DPK and DIPK through justifying the importance of its constituent elements. We begin with the Jeffrey’s updating rule.

**1.1. Why Jeffrey’s updating?** The most famous and widely used technique to describe updating of beliefs is Bayesian updating. It prescribes the scholar to form an initial opinion on the plausibility of the event  $A$  of interest, where  $A$  is a subset of the state space  $\Omega$ , and to express it by specifying a probability measure  $P$ , so that  $P(A)$  can be quantified. Once some data  $E$  is collected, the Bayesian updating mechanism revises the initial opinion by applying

---

2010 *Mathematics Subject Classification.* Primary: 62A01; Secondary: 62A01, 60A99.

*Key words and phrases.* Subjective probability; Jeffrey’s updating; Sets of probabilities; Lower probabilities; Contraction; Dilation; Sure loss.

the Bayes' rule

$$P^*(A) \equiv P(A | E) = \frac{P(A \cap E)}{P(E)} = \frac{P(E | A)P(A)}{P(E)} \propto P(E | A)P(A),$$

provided that  $P(E) \neq 0$ . In [8, 9, 10], Richard Jeffrey makes a compelling case of the fact that Bayes' rule is not the only reasonable way of updating. For example, its use presupposes that both  $P(E)$  and  $P(A \cap E)$  have been quantified before event  $E$  takes place: this can be a very challenging task, for example when  $E$  is not anticipated. The following example illustrates a situation in which Bayes' rule is not directly applicable to compute the updated probability of an event, but Jeffrey's rule can be applied.

**Example 1.1.** [5, Section 1.1] Three trials of a new surgical procedure are to be conducted at a hospital. Let 1 denote a successful outcome, and 0 an unsuccessful one. The state space has the form  $\Omega = \{000, 001, 010, 011, 100, 101, 110, 111\}$ . A colleague informs us that another hospital performed this type of procedure 100 times, registering 80 successful outcomes. This information is relevant and should influence our opinion about the outcome of the three trials, but it cannot be put in direct terms about events in the original  $\Omega$ , thus the Bayes' rule is not directly applicable.

Since the description contains no information about the order of the three trials, our initial opinion  $P$  assumes that they are exchangeable. That is, consider the partition  $\{E_0, E_1, E_2, E_3\}$  of  $\Omega$  where  $E_j$  is the set of all outcomes with exactly  $j$  successes, exchangeability implies that we assign equal probabilities to atomic events within each partition. In other words,  $P(\{001\}) = P(\{100\}) = P(\{010\})$  and  $P(\{110\}) = P(\{101\}) = P(\{011\})$ .

The success rate at the other hospital informs our opinion over the partition  $\{E_j\}$  only, and nothing more. In relation to our old opinion  $P$ , our updated opinion  $P^*$  satisfies  $P(A | E_j) = P^*(A | E_j)$  for all  $A \subset \Omega$  and all  $j \in \{0, \dots, 3\}$ . Upon specifying a new subjective assessment of the  $P^*(E_j)$ 's, the updated probability measure  $P^*$  can be fully reassessed by the relation

$$P^*(A) = \sum_{j=0}^3 P^*(A | E_j)P^*(E_j) = \sum_{j=0}^3 P(A | E_j)P^*(E_j).$$

It is within our liberty to reassess the  $P^*(E_j)$ 's. We may, for example, regard the three trials as a random subsample of size three from those of the other hospital. This would equate  $P^*(E_j)$  to the probability of obtaining  $j$  successes from a Hypergeometric(100, 80, 3) distribution.  $\triangle$

The rule  $P^*(A) = \sum_{E_j \in \mathcal{E}} P(A | E_j)P^*(E_j)$  is known as Jeffrey's rule of conditioning. It is valid when there is a partition  $\mathcal{E}$  of the sample space  $\Omega$  such that

$$P^*(A | E_j) = P(A | E_j), \quad \forall A \subset \Omega, \forall E_j \in \mathcal{E}. \quad (1)$$

It has the practical advantage of reducing the assessment of  $P^*$  to the simpler task of assessing  $P^*(E_j)$ , for all  $E_j \in \mathcal{E}$ . In the above example, instead of a full reassessment of probabilities on  $\Omega$ , the agent only needs to deliberate new assessment of the four probabilities  $P^*(E_0)$  through  $P^*(E_3)$  based on the given information.

Jeffrey's rule of conditioning is also a generalization of Bayesian conditioning. If we consider partition  $\{E, E^c\}$ , for some  $E \subset \Omega$ , then if  $P^*(E) = 1$ , we have that  $P^*(A) = P(A |$

$E)P^*(E) + P(A | E^c)P^*(E^c) = P(A | E)$ , which is Bayes' rule. In addition, as studied in [5, Section 2], if we are given the couple  $\{P, P^*\}$  of probability measures, we can always reconstruct a partition  $\{E_j\}$  for which  $\{P, P^*\}$  could have arisen via Jeffrey's updating rule, unlike Bayesian conditionalization.

**1.2. Precise and imprecise probabilities.** In this work, we consider the dynamic updating of beliefs encoded both in the form of precise and imprecise probabilities. Precise probabilities are widely employed as the central vocabulary of many modes of uncertainty reasoning, nearly exclusively so in statistical inference, for example. Yet, there are many reasons to work with imprecise probabilities, the most important of which are summarized in [18, Section 1.1.4]. In this paper, we work with lower and upper probabilities resulting from specifying sets of probability measures. In the subjective probability literature, the agent's initial beliefs about an event  $A \subset \Omega$  are usually encapsulated in a (prior) probability measure, that is then refined once new information in the form of data become available. As Walley points out in [18, Section 1.1.4.(e)], though, due to bounded rationality, the agent may be unable to assess probabilities precisely in practice, even if that is possible in principle. This is due to the lack of time or of computational ability, or because it is extremely difficult to analyze a complex body of evidence. This is also confirmed by Berger in [2] and in the de Finetti lecture delivered at ISBA 2021. Berger pointed out how people that are not "trained properly", when asked to report a prior variance, produce a value that is way too small; the folklore says by a factor of 3. People simply think that they know more than they actually do.

The agent may only be able to evaluate upper and lower bounds of a precise probability. Formally, this can be translated in the agent specifying a set of probabilities. Generally speaking, the farther apart (e.g. in the total variation distance) the "boundary elements" of the set (i.e. its infimum and supremum), the higher the agent's uncertainty.

Another more practical reason for the agent to specify a set of probabilities at the beginning of the procedure is to conduct a robust analysis. This is inspected in [18, Section 1.1.4.(j)]: the conclusions of a statistical analysis are robust when a realistically wide class of probability models all lead to essentially the same conclusions. To check the robustness of an analysis using a precise probability model, the agent would need to define a "realistically wide class" of models by varying those assumptions of the precise model that are arbitrary or unreliable. In effect, this replaces the precise probability model by an imprecise one. Conclusions drawn from the imprecise model are automatically robust, because they do not rely on arbitrary or doubtful assumptions. A way for the agent to replace the precise probability model with an imprecise one is to start the analysis by specifying a set of (prior) probabilities.

Robustness is the main reason we work with sets of probabilities in this paper. As Section 6 will explain, dynamic probability kinematics may be prior sensitive. To conduct a robust analysis we need to begin with a set of prior probabilities. This leads to the introduction of dynamic imprecise probability kinematics.

As Section 6 will discuss, the lower probabilities associated with the updated sets characterize these latter. That is why in Section 8 we study the behavior of the updated sets (contraction, dilation, sure loss) by giving sufficient conditions involving lower (and upper) probabilities. So despite DIPK can be considered as an imprecise probability model, we

do not consider DIPK lower probabilities as ill-known precise probabilities, as they simply germinate from the agent specifying a set of precise probabilities, the wider the higher their initial uncertainty.

**1.3. Structure of the paper.** The paper is organized as follows. In Section 2, we discuss the connection between our work and the related literature. In Section 3, we introduce dynamic probability kinematics (DPK). In Section 4, we provide the mechanical version of DPK that we use throughout the rest of the work. It is mechanical in the sense that it does not require the agent to subjectively specify the probability to attach to the elements of the partition, but it does so mechanically. In Section 5, we explain how to subsequently update prior  $P$  as more and more data become available. In Section 6, we introduce dynamic imprecise probability kinematics (DIPK). In Section 7, we give bounds for the upper and lower probabilities associated with the updated probability set. This allows to avoid the exhaustive procedure to update lower and upper probabilities, which involves updating the whole probability set. In Section 8, we study the behavior of updated sets of probabilities, namely contraction, dilation, and sure loss. Section 9 discusses an application of DPK and DIPK to survey sampling studies. Section 10 concludes our work. Appendix A contains the proofs of our results.

## 2. RELATED LITERATURE

In this Section, we discuss the relation between our work and other papers dealing with Jeffrey's updating in the context of imprecise probability models. Jeffrey's updating has been generalized to be put to use in the context of Dempster-Shafer theory, evidence theory, neighborhood models theory, and credal sets theory. DIPK belongs to this last category.

In [15], Shafer discusses Jeffrey's updating from a philosophical perspective, and is the first to consider its application to the context of Dempster-Shafer theory, for which belief functions – functions representing the degree of belief of the agent on a given event – and Dempster's rule of combination play a central operational role. In [7] and [17] the authors further study the generalization of Jeffrey's updating for belief functions defined on a finite sample space. In [7], the authors point out how Shafer's approach is different from the normative Bayesian approach and is not a straight generalization of Jeffrey's rule, so they propose rules of conditioning for which Jeffrey's rule is a direct consequence of a special case. In [17], the author generalizes the results in [7]. He shows that several forms of Jeffrey's updating rule can be defined so that they correspond to the geometrical rule of conditioning and to Dempster's rule of conditioning, respectively.

In [12], the authors provide a generalization of both Jeffrey's rule and Dempster conditioning to propose an effective revision rule in the field of evidence theory. This is very interesting since when one source of evidence is less reliable than another, the idea is to let prior knowledge of an agent be altered only by some of the input information. The change problem is thus intrinsically asymmetric. To this extent, their model takes into account inconsistency between prior and input information.

In [16], the author discusses the application of Jeffrey's rule to neighborhood models theory. In this field, uncertainty is captured by neighborhoods of a classical probability measure  $P$ , presented in the form of interval probabilities  $[L, U]$ . This means that  $P(A) \in [L(A), U(A)]$ ,

for all  $A \subset \Omega$ , where  $\Omega$  is the state space of interest. The author shows that a neighborhood  $[L, U]$  of a probability measure  $P$  whose lower envelope  $L$  is convex or bi-elastic with respect to the base probability measure – see [16, Definitions 3 and 4] – is closed with respect to Jeffrey’s rule of conditioning. This means that Jeffrey’s posterior for  $Q \in [L, U]$  still belongs to the interval.

In [13], the authors generalize Jeffrey’s rule to credal set theory. In this field, a set of probability measures (called a credal set) is used to capture either the agent’s initial uncertainty, or inconsistency/imprecision in the process of collecting data. The authors introduce imaginary kinematics, see [13, Definition 7]. They combine Jeffrey’s rule with Lewis’ imaging [11] for credal sets to be able to update beliefs when possibly inconsistent probabilistic evidence is gathered. The main differences between our work and [13] are two. In [13], the authors require  $\Omega$  to be a space of atoms, while we do not put any restriction on the nature of  $\Omega$ . Also, while we use sets of probability measures in order to capture the impossibility of the agent to specify a single prior probability, and in order to conduct a robust analysis, in [13] the authors use credal sets to tackle the problem of the agent gathering inconsistent data. In the future, we plan to generalize DIPK to account for imprecise data collection, as we point out in Section 10.

It is worth noting that in [3] the authors provide an ergodic theory for the limit of a sequence of successive DIPK updates of a set representing the initial beliefs of an agent. As a consequence, they formulate a strong law of large numbers.

### 3. A NEW WAY TO UPDATE SUBJECTIVE BELIEFS

In this Section, we describe a new way to update subjective beliefs based on Jeffrey’s rule of conditioning (see e.g. [5, 8, 9, 10]), which we call dynamic probability kinematics (DPK). Let  $\Omega$  be the state space of interest. Suppose that  $P$  is a probability measure on  $(\Omega, \mathcal{F})$ , and we want to compute its Jeffrey’s update. We collect data, i.e. we observe  $x_1, \dots, x_n$ . These latter are images of random variables, that is,

$$\exists E_1, \dots, E_n \in \mathcal{F} : \forall \omega \in E_j, X_j(\omega) = x_j, \forall j \in \{1, \dots, n\}.$$

Let us denote by  $\mathcal{X}$  the range of  $X$ , that is, the image under  $X$  of the domain of  $X$ ; notationally, this means that  $\mathcal{X} = \text{Im}_X(\text{Dom}(X))$ . We drop the index since we assume  $X_1, \dots, X_n$  to be identically distributed. We also assume that  $\mathcal{X}$  is finite or countable.

Notice that  $E_j$  can be written as

$$E_j := \{\omega \in \Omega : X_j(\omega) = x_j\}.$$

$E_1, \dots, E_n$  induce a partition

$$\mathcal{E} = \left\{ E_1, \dots, E_n, E_{n+1} := \left( \bigcup_{j=1}^n E_j \right)^c \right\}, \tag{2}$$

where  $E_{n+1}$  is the remainder set to the union of  $E_1, \dots, E_n$ .

Note that the purpose of the observed data points, or  $x_j$ ’s, is to inform the structure of the partition  $\mathcal{E}$ . Each unique value expressed by  $x_j$  conveys the information for a unique element  $E_j$  of the partition. Therefore for our purpose, it suffices that we keep track of only the unique values in the collection of observations  $x_1, \dots, x_n$ , rather than treating it as a sample

from an underlying population, as is typically the case in Bayesian or frequentist statistical inference. Furthermore, the assumption that  $X_1, \dots, X_n$  are identically distributed is needed because otherwise we may have  $E_k, E_s \in \mathcal{E}$  such that  $E_k \cap E_s \neq \emptyset$ .<sup>1</sup>

As an update to  $P$ , we propose the following

$$P_{\mathcal{E}} : \mathcal{F} \rightarrow [0, 1], \quad A \mapsto P_{\mathcal{E}}(A) := \sum_{E_j \in \mathcal{E}} P(A | E_j) P_{\mathcal{E}}(E_j) \quad (3)$$

such that  $P_{\mathcal{E}}(E_j) \geq 0, \forall j \in \{1, \dots, n\}$ , and  $\sum_{E_j \in \mathcal{E}} P_{\mathcal{E}}(E_j) = 1$ .

We have the following.

**Proposition 3.1.**  *$P_{\mathcal{E}}$  is a probability measure, and it is a Jeffrey's posterior for  $P$ .*

In general, Jeffrey's rule of conditioning – as presented in [5, Equation 1.1] – is given by  $P^*(A) = \sum_j P(A | E_j) P^*(E_j)$ , where  $P^*$  is Jeffrey's posterior for  $P$ . It is valid when Jeffrey's condition is met, that is, when there is a given partition  $\{E_j\}$  of the state space  $\Omega$  such that  $P(A | E_j) = P^*(A | E_j)$  is true for all  $A \in \mathcal{F}$  and all  $j$ . Specifically, this condition is met by  $P_{\mathcal{E}}$ . Since  $P_{\mathcal{E}}$  is a probability measure by Proposition 3.1, it is true that, for all  $A \in \mathcal{F}$ ,  $P_{\mathcal{E}}(A) = \sum_{E_j \in \mathcal{E}} P_{\mathcal{E}}(A | E_j) P_{\mathcal{E}}(E_j)$ . But given our definition for  $P_{\mathcal{E}}$ , we also have that  $P_{\mathcal{E}}(A) = \sum_{E_j \in \mathcal{E}} P(A | E_j) P_{\mathcal{E}}(E_j)$ . This implies that there is a partition  $\mathcal{E}$  for which  $P(A | E_j) = P_{\mathcal{E}}(A | E_j)$  is true for all  $A \in \mathcal{F}$  and all  $E_j \in \mathcal{E}$ .

In the next Section, we show how computing  $P_{\mathcal{E}}(A)$  can be a mechanical procedure that does not require subjective assessment.

#### 4. A MECHANICAL PROCEDURE TO COMPUTE $P_{\mathcal{E}}$

In this Section, we show how computing  $P_{\mathcal{E}}(A)$  can be performed as a mechanical procedure, if the analyst is unwilling or unable to make full subjective probabilistic assessment for the elements of  $\mathcal{E}$ . Recall that  $\mathcal{X}$  is the range of  $X$ , that is,  $\mathcal{X} = \text{Im}_X(\text{Dom}(X))$ . Every  $X_j$  has its probability mass function  $P_{X_j}(\{x\})$  that tells us what is the probability that  $X_j = x$ , for all  $x \in \mathcal{X}$ . Since we assumed  $X_1, \dots, X_n$  to be identically distributed, we have that  $P_{X_1} = P_{X_2} = \dots = P_{X_n}$ . Let us denote it by  $P_X$ . Probability mass function  $P_X$  is the pushforward probability measure for a probability measure  $Q$  defined on  $(\Omega, \mathcal{F})$ . That is, we have that  $P_X(\{x\}) = Q(X^{-1}(x))$ , for all  $x \in \mathcal{X}$ . Let us now focus on the data  $x_1, \dots, x_n$  that we collect. Notice that

$$X_j^{-1}(x_j) := \{\omega \in \Omega : X_j(\omega) = x_j\} = E_j, \quad \forall j \in \{1, \dots, n\}.$$

So, if we let  $P_{\mathcal{E}}(E_j) = P_{X_j}(\{x_j\}) = Q(X_j^{-1}(x_j))$  and  $P_{\mathcal{E}}((\cup_{j=1}^n E_j)^c) = 1 - \sum_{j=1}^n P_{\mathcal{E}}(E_j)$ , we have that the probabilities assigned by  $P_{\mathcal{E}}$  to the elements of  $\mathcal{E}$  are assigned mechanically. Jeffrey's updating procedure, then, becomes easier (the analyst does not have to subjectively specify the probability assigned to all the elements of  $\mathcal{E}$ , a task that can be mentally and mathematically challenging). Let us give an example.

<sup>1</sup>To be specific, suppose that  $X_1 \sim F$  and  $X_2 \sim G$ , where  $F$  and  $G$  are two different distributions. Then, there may exist  $\omega \in \Omega$  such that  $X_1(\omega) = a$ ,  $X_2(\omega) = b$ , and  $a \neq b$ . Call then  $E_1 := \{\omega \in \Omega : X_1(\omega) = a\}$  and  $E_2 := \{\omega \in \Omega : X_2(\omega) = b \neq a\}$ . In this case,  $E_1 \cap E_2 \neq \emptyset$ . So  $E_1$  and  $E_2$  do not induce a partition, since the elements of a partition must not intersect with each other.

**Example 4.1.** Suppose  $X_1, X_2, X_3 \sim \text{Bin}(n, p)$  iid, and let  $n = 10$  and  $p = 0.8$ . Then, suppose that we observe  $x_1 = 3$ ,  $x_2 = 5$ , and  $x_3 = 7$ . Then,  $P_{X_1}(\{x_1\}) = \binom{10}{3}0.8^30.2^{10-3} \approx 0.00079$ ,  $P_{X_2}(\{x_2\}) = \binom{10}{5}0.8^50.2^{10-5} \approx 0.02642$ ,  $P_{X_3}(\{x_3\}) = \binom{10}{7}0.8^70.2^{10-7} \approx 0.20133$ . Now, we let  $P_{\mathcal{E}}(E_1) = P_{X_1}(\{x_1\})$ ,  $P_{\mathcal{E}}(E_2) = P_{X_2}(\{x_2\})$ ,  $P_{\mathcal{E}}(E_3) = P_{X_3}(\{x_3\})$ , and  $P_{\mathcal{E}}((\cup_{j=1}^3 E_j)^c) = 1 - \sum_{j=1}^3 P_{\mathcal{E}}(E_j) \approx 0.77146$ . Given these, computing  $P_{\mathcal{E}}(A)$  becomes an easier task: we just need to compute  $P(A | E_j)$ , for all  $E_j \in \mathcal{E}$ , which is another mechanical task that does not require subjective assessments of probabilities.

In the remainder of this document, we are going to use the mechanical procedure we just described to assign updated probabilities to the elements of  $\mathcal{E}$ . We have the following useful lemma.

**Lemma 1.** *Let  $E \in \mathcal{E}$ .  $P_{\mathcal{E}}(E) = 0$  if and only if  $E = \emptyset$ .*

**Remark 1.** There is a subtlety in moving from the prior to the posterior. Requiring that  $P_{\mathcal{E}}(E_j) = P_{X_j}(\{x_j\}) = Q(X_j^{-1}(x_j))$  and  $P_{\mathcal{E}}((\cup_{j=1}^n E_j)^c) = 1 - \sum_{j=1}^n P_{\mathcal{E}}(E_j)$ , means that the restriction  $P_{\mathcal{E}}|_{\sigma(\mathcal{E})}$  of  $P_{\mathcal{E}}$  agrees with the restriction  $Q|_{\sigma(\mathcal{E})}$  of  $Q$  on the sigma algebra  $\sigma(\mathcal{E})$  generated by the elements of  $\mathcal{E}$ .  $P_{\mathcal{E}}|_{\sigma(\mathcal{E})}$  is then extended to  $(\Omega, \mathcal{F})$  through  $P(\cdot | E_j)$ , for all  $j$ , via

$$P_{\mathcal{E}}(A) = \sum_{E_j \in \mathcal{E}} P(A | E_j) P_{\mathcal{E}}(E_j).$$

## 5. SUBSEQUENT UPDATES

Once we observe new data points  $x_{n+1}, \dots, x_m$ , we update  $P_{\mathcal{E}} \equiv P_{\mathcal{E}_1}$  to  $P_{\mathcal{E}_2}$  via the same mechanical procedure depicted in Section 4. With this, we mean the following. We now have observed data  $x_1, \dots, x_n, x_{n+1}, \dots, x_m$ . Then, we compute  $P_{\mathcal{E}_2}(A) = \sum_{E_j \in \mathcal{E}_2} P_{\mathcal{E}_1}(A | E_j) P_{\mathcal{E}_2}(E_j)$ , where  $\mathcal{E}_2 = \{E_j\}_{j=1}^{m+1}$ . In particular,  $E_j = \{\omega \in \Omega : X(\omega) = x_j\}$ ,  $j \in \{1, \dots, m\}$ , and  $E_{m+1} = (\cup_{j=1}^m E_j)^c$ . Also, we assign probability  $P_{\mathcal{E}_2}(E_j) = P_X(\{x_j\})$ , for  $j \in \{1, \dots, m\}$ , and  $P_{\mathcal{E}_2}(E_{m+1}) = 1 - \sum_{j=1}^m P_{\mathcal{E}_2}(E_j)$ . Clearly, Propositions 3.1 and Lemma 1 are true also for  $P_{\mathcal{E}_2}$ .

If we keep on collecting observations, eventually our sequence of Jeffrey's updates converges.

**Proposition 5.1.** *The limit  $\lim_{n \rightarrow \infty} P_{\mathcal{E}_n}$  exists and is well defined.*

Notice that, for any  $n$ , partition  $\mathcal{E}_n$  is a refinement of partition  $\mathcal{E}_{n-1}$ . To see this, suppose  $\mathcal{E}_{n-1}$  has  $\ell + 1$  many elements, that is,  $\mathcal{E}_{n-1} = \{E_1^{\mathcal{E}_{n-1}}, \dots, E_{\ell}^{\mathcal{E}_{n-1}}, E_{\ell+1}^{\mathcal{E}_{n-1}}\}$ . By (2), this means that  $E_{\ell+1}^{\mathcal{E}_{n-1}} = (\cup_{j=1}^{\ell} E_j^{\mathcal{E}_{n-1}})^c$ . Now suppose that in the next updating step we only observe one new element  $x$  of  $\mathcal{X}$ . Then we have that  $\mathcal{E}_n$  has  $\ell + 2$  many elements. In particular,  $E_j^{\mathcal{E}_{n-1}} = E_j^{\mathcal{E}_n}$ , for all  $j \in \{1, \dots, \ell\}$ , and  $E_{\ell+1}^{\mathcal{E}_{n-1}} = E_{\ell+1}^{\mathcal{E}_n} \cup E_{\ell+2}^{\mathcal{E}_n}$ . Of course, if we observe more elements, we further refine  $E_{\ell+1}^{\mathcal{E}_{n-1}}$ . Call now  $\tilde{\mathcal{E}}$  the partition resulting from our updating process having possibly countably many elements, and such that  $\emptyset \in \tilde{\mathcal{E}}$ . Then, we have the following.

**Proposition 5.2.**  *$\tilde{\mathcal{E}}$  cannot be refined as a result of the updating process described in Sections 3 and 4.*

Since the main driver of the update from  $P_{\mathcal{E}_{n-1}}$  to  $P_{\mathcal{E}_n}$  is the refinement of partition  $\mathcal{E}_{n-1}$  to partition  $\mathcal{E}_n$ , we denote by  $P_{\mathcal{E}}$  the limit we found in Proposition 5.1.

Note that the larger the cardinality of the range of the random variables that drive the updating process, the more steps are needed to reach convergence. Consider two random variables  $X$  and  $Y$ , call  $\mathcal{X} := \text{Im}_X(\text{Dom}(X))$  and  $\mathcal{Y} := \text{Im}_Y(\text{Dom}(Y))$ , and suppose that  $\#\mathcal{X} < \#\mathcal{Y} < \infty$ , where  $\#$  denotes the cardinality operator. Call  $k$  the steps that  $(P_{\mathcal{E}_n})$  takes to converge when the convergence is driven by the elements of  $\mathcal{X}$ , and  $\ell$  the steps that  $(P_{\mathcal{E}_n})$  takes to converge when the convergence is driven by the elements of  $\mathcal{Y}$ .

**Proposition 5.3.** *If at every step of the updating procedure we observe the same number of new elements from  $\mathcal{X}$  and  $\mathcal{Y}$ , then  $k < \ell$ .*

Note that the requirement of observing the same number of new elements from  $\mathcal{X}$  and  $\mathcal{Y}$  “levels the playing field” in Proposition 5.3. Had we not required it to hold, we could have had a situation in which in the first step of our updating procedure, we would have observed all the elements in  $\mathcal{Y}$ , and, say, only one element of  $\mathcal{X}$ . The validity of the proposition is evident in the extreme case that  $\#\mathcal{X}$  is finite, yet  $\#\mathcal{Y}$  is infinite. In that case, it is clear that  $k < \ell$ , whether or not we observe the same number of new elements at every step.

**Remark 2.** Notice that Lemma 1 holds for all  $P_{\mathcal{E}_n}$ ,  $n \in \mathbb{N}$ , not just for  $P_{\mathcal{E}_1}$  and  $P_{\mathcal{E}_2}$ . We also have the following. If  $\emptyset$  belongs to the partition we are considering, computing the conditional probability of any  $A \subset \Omega$  given  $\emptyset$  is a problem, in that it gives rise to an indeterminate form. Indeed, for a generic probability measure  $P$ , we have that  $P(A \mid \emptyset) = \frac{P(A \cap \emptyset)}{P(\emptyset)} = \frac{0}{0}$ , an indeterminate form. We solve this indeterminacy by requiring that for dynamic probability kinematics this ratio is equal to 0. This because the information conveyed by the data collected on the event  $A$  is already encapsulated in all the other elements of the partition.

**Remark 3.** Dynamic probability kinematics, being based on Jeffrey’s updating, is not commutative. With this we mean the following. Consider an initial probability  $P$  and compute its dynamic probability kinematics update  $P_{\mathcal{E}_1}$  based on partition  $\mathcal{E}_1$ ; then compute the DPK update of  $P_{\mathcal{E}_1}$  based on partition  $\mathcal{E}_2$ , and call this update  $P_{\mathcal{E}_1\mathcal{E}_2}$ . If we proceed in the opposite direction, that is, if we first update  $P$  to  $P_{\mathcal{E}_2}$ , and then update this latter to  $P_{\mathcal{E}_2\mathcal{E}_1}$ , we have that, in general,  $P_{\mathcal{E}_1\mathcal{E}_2} \neq P_{\mathcal{E}_2\mathcal{E}_1}$ .

This comes from [5, Theorem 3.2], where the authors show that, given two generic partitions  $\mathcal{E}$  and  $\mathcal{F}$  of  $\Omega$ ,  $P_{\mathcal{E}\mathcal{F}} = P_{\mathcal{F}\mathcal{E}}$  if and only if  $\mathcal{E}$  and  $\mathcal{F}$  are Jeffrey independent. This happens when  $P_{\mathcal{E}}(F_j) = P(F_j)$  and  $P_{\mathcal{F}}(E_i) = P(E_i)$ , for all  $i$  and all  $j$ , where the  $F_j$ ’s are the elements of  $\mathcal{F}$ , and the  $E_i$ ’s are the elements of  $\mathcal{E}$ .

It is immediate to see that this necessary and sufficient condition does not hold for DPK. Indeed, consider the following toy example. Let  $\Omega = [0, 1]$ , and  $\mathcal{X} = \{1, 2, 3, 4\}$ . Let  $P_X(\{1\}) = \frac{1}{6}$ ,  $P_X(\{2\}) = \frac{1}{3}$ ,  $P_X(\{3\}) = \frac{1}{8}$ ,  $P_X(\{4\}) = \frac{3}{8}$ . Let also the initial probability measure  $P$  be the Lebesgue measure on  $[0, 1]$ . Suppose then that at time  $n = 1$  we observe  $x_1 = 1$ , and that the partition induced by this data point is given by  $\mathcal{E}_1 = \{E_1^{\mathcal{E}_1} = [0, \frac{1}{8}), E_2^{\mathcal{E}_1} = [\frac{1}{8}, 1]\}$ . At time  $n = 2$ , we observe  $x_2 = 2$ ,  $x_3 = 3$ ,  $x_4 = 4$ , and the partition induced by these data points is  $\mathcal{E}_2 \equiv \tilde{\mathcal{E}} = \{E_1^{\mathcal{E}_2} = [0, \frac{1}{8}), E_2^{\mathcal{E}_2} = [\frac{1}{8}, \frac{1}{4}), E_3^{\mathcal{E}_2} = [\frac{1}{4}, \frac{3}{4}), E_4^{\mathcal{E}_2} = [\frac{3}{4}, 1], E_5^{\mathcal{E}_2} = \emptyset\}$ . To verify whether  $\mathcal{E}_1$  and  $\mathcal{E}_2$  are Jeffrey independent, we need to check, for example, whether

$P_{\mathcal{E}_1}(E_2^{\mathcal{E}_2}) = P(E_2^{\mathcal{E}_2})$ . Now,  $P(E_2^{\mathcal{E}_2}) = \frac{1}{4} - \frac{1}{8} = \frac{1}{8}$ . Instead,

$$\begin{aligned} P_{\mathcal{E}_1}(E_2^{\mathcal{E}_2}) &= \sum_{E_j^{\mathcal{E}_1} \in \mathcal{E}_1} P(E_2^{\mathcal{E}_2} \mid E_j^{\mathcal{E}_1}) P_{\mathcal{E}_1}(E_j^{\mathcal{E}_1}) \\ &= \frac{P(E_2^{\mathcal{E}_2} \cap E_2^{\mathcal{E}_1})}{P(E_2^{\mathcal{E}_1})} P_{\mathcal{E}_1}(E_2^{\mathcal{E}_1}) \\ &= \frac{P(E_2^{\mathcal{E}_2})}{P(E_2^{\mathcal{E}_1})} P_{\mathcal{E}_1}(E_2^{\mathcal{E}_1}) \\ &= \frac{1/8 \cdot 5}{7/8 \cdot 6} = \frac{5}{42} \neq \frac{1}{8}. \end{aligned}$$

This simple example shows that DPK is not commutative. Should this worry the agent that intends to use DPK to update their beliefs? The answer is no; it is actually something desirable. Since in DPK the partitions become finer as more data is collected, if  $P_{\mathcal{E}_1 \dots \mathcal{E}_{n-1} \mathcal{E}_n}$  were to be equal to  $P_{\mathcal{E}_1 \dots \mathcal{E}_n \mathcal{E}_{n-1}}$ , that would mean that losing granularity in the partition of the state space would yield no loss of information on the event  $A \subset \Omega$  of interest. This is undesirable: the more we know about the composition of  $\Omega$ , the better we want our assessment to be on the plausibility of event  $A$ . As Diaconis and Zabell point out in [5, Section 4.2, Remark 2], “noncommutativity is not a real problem for successive Jeffrey updating”.

Before concluding this Remark, we point out how, despite the order in which new data is collected matters for the updating process, the limit probability  $P_{\tilde{\mathcal{E}}}$  is the same regardless of the order in which data is collected. Let us give an example.

Continuing the toy example from above, let  $A = [\frac{1}{4}, \frac{3}{4}]$  be our set of interest. Suppose at time  $n = 1$  we observe  $x_1 = 1$ . We have that  $P_{\mathcal{E}_1}(A) = \sum_{E_j^{\mathcal{E}_1} \in \mathcal{E}_1} P(A \mid E_j^{\mathcal{E}_1}) P_{\mathcal{E}_1}(E_j^{\mathcal{E}_1}) = \frac{P(A)}{P(E_2^{\mathcal{E}_1})} P_{\mathcal{E}_1}(E_2^{\mathcal{E}_1}) = \frac{1/2 \cdot 5}{7/8 \cdot 6} = \frac{10}{21}$ . Then, suppose at time  $n = 2$  we observe  $x_2 = 2, x_3 = 3, x_4 = 4$ . Then,  $P_{\mathcal{E}_2}(A) = \sum_{E_j^{\mathcal{E}_2} \in \mathcal{E}_2} P_{\mathcal{E}_1}(A \mid E_j^{\mathcal{E}_2}) P_{\mathcal{E}_2}(E_j^{\mathcal{E}_2}) = P_{\mathcal{E}_2}(E_3^{\mathcal{E}_2}) = \frac{1}{8}$ . Suppose instead that at time  $n = 1$  we observe  $x_1 = 3$ . Then, the induced partition is  $\mathcal{E}'_1 = \{E_1^{\mathcal{E}'_1} = [\frac{1}{4}, \frac{3}{4}], E_2^{\mathcal{E}'_1} = [0, \frac{1}{4}] \cup [\frac{3}{4}, 1]\}$ . We have that  $P_{\mathcal{E}'_1}(A) = \sum_{E_j^{\mathcal{E}'_1} \in \mathcal{E}'_1} P(A \mid E_j^{\mathcal{E}'_1}) P_{\mathcal{E}'_1}(E_j^{\mathcal{E}'_1}) = P_{\mathcal{E}'_1}(E_1^{\mathcal{E}'_1}) = \frac{1}{8}$ . Suppose then that at time  $n = 2$  we observe  $x_2 = 1, x_3 = 2, x_4 = 4$ . We have that the induced partition is  $\mathcal{E}'_2 = \{E_1^{\mathcal{E}'_2} = [\frac{1}{4}, \frac{3}{4}], E_2^{\mathcal{E}'_2} = [0, \frac{1}{8}), E_3^{\mathcal{E}'_2} = [\frac{1}{8}, \frac{1}{4}), E_4^{\mathcal{E}'_2} = [\frac{3}{4}, 1], E_5^{\mathcal{E}'_2} = \emptyset\}$ . As we can see,  $\mathcal{E}'_2 = \mathcal{E}_2$ . Then,  $P_{\mathcal{E}'_2}(A) = \sum_{E_j^{\mathcal{E}'_2} \in \mathcal{E}'_2} P_{\mathcal{E}'_1}(A \mid E_j^{\mathcal{E}'_2}) P_{\mathcal{E}'_2}(E_j^{\mathcal{E}'_2}) = P_{\mathcal{E}'_2}(E_1^{\mathcal{E}'_2}) = \frac{1}{8}$ . We see how  $P_{\mathcal{E}_1}(A) \neq P_{\mathcal{E}'_1}(A)$ , but  $P_{\mathcal{E}_2}(A) = P_{\mathcal{E}'_2}(A) = P_{\tilde{\mathcal{E}}}(A)$ . This because despite the order in which the observation are collected is different, in the limit the information conveyed by the data around the plausibility of event  $A$  is the same, no matter the order in which data is gathered. The following is the formal result.

Suppose we collect the elements of  $\mathcal{X}$  in a different order in two different experiments. We call  $(\mathcal{E}_n)$  and  $(\mathcal{E}'_n)$  the sequences of finer partitions in the first and second experiments, respectively. Call  $\tilde{\mathcal{E}}$  and  $\tilde{\mathcal{E}}'$  the limit partitions for the first and second experiments, respectively. That is, they are the finest partitions compatible with the elements of  $\mathcal{X}$  according to DPK.

**Proposition 5.4.** *Call  $P_{\tilde{\mathcal{E}}} := \lim_{n \rightarrow \infty} P_{\mathcal{E}_n}$  and  $P_{\tilde{\mathcal{E}'}} := \lim_{n \rightarrow \infty} P_{\mathcal{E}'_n}$ . Then,  $P_{\tilde{\mathcal{E}}} = P_{\tilde{\mathcal{E}'}}$ .*

## 6. WORKING WITH SETS OF PROBABILITIES

In this Section, we generalize dynamic probability kinematics to dynamic imprecise probability kinematics. To do so, we first need to introduce the concepts of lower probability, upper probability, and core of a lower probability.

Consider a generic set of probabilities  $\Pi$  on a measurable space  $(\Omega, \mathcal{F})$ . The lower probability  $\underline{P}$  associated with  $\Pi$  is defined as

$$\underline{P}(A) := \inf_{P \in \Pi} P(A), \quad \forall A \in \mathcal{F}.$$

The upper probability  $\overline{P}$  associated with  $\Pi$  is defined as the conjugate to  $\underline{P}$ , that is,

$$\overline{P}(A) := 1 - \underline{P}(A^c) = \sup_{P' \in \Pi} P'(A), \quad \forall A \in \mathcal{F}.$$

Let us denote by  $\Delta(\Omega, \mathcal{F})$  the set of all probability measures on  $(\Omega, \mathcal{F})$ . Lower probability  $\underline{P}$  completely characterizes the convex set

$$\begin{aligned} \text{core}(\underline{P}) &:= \{P \in \Delta(\Omega, \mathcal{F}) : P(A) \geq \underline{P}(A), \forall A \in \mathcal{F}\} \\ &= \{P \in \Delta(\Omega, \mathcal{F}) : \overline{P}(A) \geq P(A) \geq \underline{P}(A), \forall A \in \mathcal{F}\}, \end{aligned}$$

where the second equality is a characterization (see e.g. [4, Page 3389]). By complete characterization, we mean that it is sufficient to know  $\underline{P}$  to be able to completely specify  $\text{core}(\underline{P})$ . To emphasize this aspect, some authors say that  $\underline{P}$  is *compatible* with  $\text{core}(\underline{P})$  (see e.g. [6]).

To generalize DPK to DIPK, we first prescribe the agent to specify a finite set of probabilities, then to compute the lower probability associated with it. The core of such lower probability represents the agent's initial beliefs. To update their beliefs, the agent computes the DPK update of the elements of the finite set, obtaining an updated finite set. Their updated beliefs are represented by the core of the lower probability associated with the updated finite set. We work with sets of probabilities because DPK is a prior sensitive procedure, so probability sets allow for a robust analysis. In addition, we require the agent's beliefs to be represented by the core for two main reasons. The first, mathematical, one is to ensure that the belief set can be completely characterized by the lower probability. The second, philosophical, one is to further hedge against ambiguity. Indeed, it is immediate to see that the finite set is contained in the core, so requiring this latter to represent the agent's beliefs corresponds to the agent adopting a "doubly robust" approach: not only is the agent incapable of specifying a single prior probability measure, but also specifying a finite set would be too precise. From an information theoretic perspective, performing an analysis starting with a finite set or with the core of a lower probability are two different problems. In particular, the second entails a higher initial degree of ignorance of the agent. We are aware of this distinction, and for DIPK we choose to work with the core to achieve a more robust outcome.

We start our analysis by specifying a set of prior probability measures on  $\Omega$ . We call it  $\mathcal{P} \equiv \mathcal{P}_{\mathcal{E}_0} \subset \Delta(\Omega, \mathcal{F})$ , and we require that it is finite. This makes it easy to specify. We then consider  $\underline{P}_{\mathcal{E}_0}$ , the lower probability associated with  $\mathcal{P}_{\mathcal{E}_0}$ . The set representing the agent's

initial beliefs, then, is given by  $\mathcal{P}_{\mathcal{E}_0}^{\text{co}} = \text{core}(\underline{P}_{\mathcal{E}_0})$ , where superscript co denotes the fact that  $\mathcal{P}_{\mathcal{E}_0}^{\text{co}}$  is convex. The importance of this property is explained in Remark 4.

We then update every element in  $P_{\mathcal{E}_0} \in \mathcal{P}_{\mathcal{E}_0}$  using Jeffrey's procedure explained in Section 4 (DPK), and we obtain

$$\mathcal{P}_{\mathcal{E}_1} := \left\{ P_{\mathcal{E}_1} \in \Delta(\Omega, \mathcal{F}) : P_{\mathcal{E}_1}(A) = \sum_{E_j \in \mathcal{E}_1} P_{\mathcal{E}_0}(A | E_j) P_{\mathcal{E}_1}(E_j), \forall A \in \mathcal{F}, P_{\mathcal{E}_0} \in \mathcal{P}_{\mathcal{E}_0} \right\}.$$

After that, we compute  $\underline{P}_{\mathcal{E}_1}$ , and so  $\mathcal{P}_{\mathcal{E}_1}^{\text{co}} = \text{core}(\underline{P}_{\mathcal{E}_1})$ . Repeating this procedure, we build two sequences,  $(\mathcal{P}_{\mathcal{E}_n})$  and  $(\mathcal{P}_{\mathcal{E}_n}^{\text{co}})$ . Notice that for any  $n \in \mathbb{N}$ , the lower and upper probabilities associated with  $\mathcal{P}_{\mathcal{E}_n}$  are equal to the lower and upper probabilities associated with  $\mathcal{P}_{\mathcal{E}_n}^{\text{co}}$ , respectively, by construction. Call

$$\mathcal{P}_{\bar{\mathcal{E}}} := \left\{ P_{\bar{\mathcal{E}}} \in \Delta(\Omega, \mathcal{F}) : P_{\bar{\mathcal{E}}} = \lim_{n \rightarrow \infty} P_{\mathcal{E}_n}, P_{\mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n} \right\}.$$

That is,  $\mathcal{P}_{\bar{\mathcal{E}}}$  is the set of limits (in the total variation metric) of the sequences  $(P_{\mathcal{E}_n})$  whose elements  $P_{\mathcal{E}_n}$  are elements of set  $\mathcal{P}_{\mathcal{E}_n}$  representing the agent's updated beliefs. Notice that  $\mathcal{P}_{\bar{\mathcal{E}}}$  is finite and has the same cardinality as  $\mathcal{P}_{\mathcal{E}_n}$ , for every  $n$ . We are sure it is not empty by Propositions 5.1 and 5.2. Then, by construction, we have that

$$d_H(\mathcal{P}_{\mathcal{E}_n}, \mathcal{P}_{\bar{\mathcal{E}}}) = \max \left( \sup_{P \in \mathcal{P}_{\mathcal{E}_n}} d_{TV}(P, \mathcal{P}_{\bar{\mathcal{E}}}), \sup_{P' \in \mathcal{P}_{\bar{\mathcal{E}}}} d_{TV}(\mathcal{P}_{\mathcal{E}_n}, P') \right) \xrightarrow[n \rightarrow \infty]{} 0, \quad (4)$$

where  $d_H$  denotes the Hausdorff metric, and, in general,  $d_{TV}(P, \mathcal{Q}) := \inf_{Q \in \mathcal{Q}} d_{TV}(P, Q)$ , for all  $P \in \Delta(\Omega, \mathcal{F})$  and all  $\mathcal{Q} \subset \Delta(\Omega, \mathcal{F})$ . Such a convergence is true also for  $\mathcal{P}_{\mathcal{E}_n}^{\text{co}}$  and  $\mathcal{P}_{\bar{\mathcal{E}}}^{\text{co}}$ .

**Proposition 6.1.** *The following is true*

$$d_H(\mathcal{P}_{\mathcal{E}_n}^{\text{co}}, \mathcal{P}_{\bar{\mathcal{E}}}^{\text{co}}) \xrightarrow[n \rightarrow \infty]{} 0.$$

Before concluding this Section, let us highlight a characteristic of our updating method involving sets of probabilities. Fix any  $n$ , and suppose  $\mathcal{E}_n$  has  $\ell + 1$  components. Pick two probability measures  $P_{1, \mathcal{E}_n}, P_{2, \mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n}$ . By the procedure described in Section 5, we have that  $P_{1, \mathcal{E}_n}(A) = \sum_{E_j \in \mathcal{E}_n} P_{1, \mathcal{E}_{n-1}}(A | E_j) P_{1, \mathcal{E}_n}(E_j)$  and  $P_{2, \mathcal{E}_n}(A) = \sum_{E_j \in \mathcal{E}_n} P_{2, \mathcal{E}_{n-1}}(A | E_j) P_{2, \mathcal{E}_n}(E_j)$ , for all  $A \in \mathcal{F}$ . By the mechanical procedure described in Section 4, then, we have that  $P_{1, \mathcal{E}_n}(E_j) = P_{2, \mathcal{E}_n}(E_j) = Q(E_j) = P_X(\{x_j\})$ , for  $j \in \{1, \dots, \ell\}$ , and  $P_{1, \mathcal{E}_n}(E_{\ell+1}) = P_{2, \mathcal{E}_n}(E_{\ell+1}) = 1 - \sum_{j=1}^{\ell} P_{1, \mathcal{E}_n}(E_j) = 1 - \sum_{j=1}^{\ell} P_{2, \mathcal{E}_n}(E_j)$ . As we can see then,  $P_{1, \mathcal{E}_n}(A)$  and  $P_{2, \mathcal{E}_n}(A)$  only differ by the conditional probability the priors  $P_{1, \mathcal{E}_{n-1}}$  and  $P_{2, \mathcal{E}_{n-1}}$  assign to  $A$  given  $E_j$ , for all  $E_j \in \mathcal{E}_n$ . Of course this is true for every element of  $\mathcal{P}_{\mathcal{E}_n}$ , for all  $n$ . This means that our procedure is sensitive to the prior: this shows how, to perform a robust analysis using our method, it is crucial is to work with sets of probabilities.

**Remark 4.** Consider  $\mathcal{P}_{\mathcal{E}_n}$ , and take two of its components  $P_{1, \mathcal{E}_n}, P_{2, \mathcal{E}_n}$ . We have that  $\alpha P_{1, \mathcal{E}_n}(A) + (1 - \alpha) P_{2, \mathcal{E}_n}(A) \in [\underline{P}_{\mathcal{E}_n}(A), \overline{P}_{\mathcal{E}_n}(A)]$ , for some  $\alpha \in [0, 1]$  and all  $A \in \mathcal{F}$ , but  $\alpha P_{1, \mathcal{E}_n} + (1 - \alpha) P_{2, \mathcal{E}_n} \notin \mathcal{P}_{\mathcal{E}_n}$ . To this extent, upper and lower probabilities  $\overline{P}_{\mathcal{E}_n}$  and  $\underline{P}_{\mathcal{E}_n}$  are not able to capture "holes" and "dents" in the set  $\mathcal{P}_{\mathcal{E}_n}$ . That is why we need the sequence of convex sets  $(\mathcal{P}_{\mathcal{E}_n}^{\text{co}})$  to represent the agent's belief updating procedure.

**Remark 5.** For generic sets  $\mathcal{P}, \mathcal{Q} \subset \Delta(\Omega, \mathcal{F})$ , we have that

$$\sup_{P \in \mathcal{P}} d_{TV}(P, \mathcal{Q}) = \sup_{P \in \mathcal{P}} \inf_{Q \in \mathcal{Q}} d_{TV}(P, Q) = d_{TV}(\overline{\mathcal{P}}, \underline{\mathcal{Q}}),$$

where  $\overline{\mathcal{P}}$  is the upper probability associated with  $\mathcal{P}$  and  $\underline{\mathcal{Q}}$  is the lower probability associated with  $\mathcal{Q}$ . The total variation distance is well defined (that is, it is indeed a metric) also when computed between upper and lower probabilities,  $d_{TV}(\overline{\mathcal{P}}, \underline{\mathcal{Q}}) := \sup_{A \in \mathcal{F}} |\overline{\mathcal{P}}(A) - \underline{\mathcal{Q}}(A)|$ . In light of this, the interpretation we can give to equation (4) is the following: as we collect more and more data points, the distance between the “boundary elements”  $\underline{P}_{\mathcal{E}_n}$  and  $\overline{P}_{\mathcal{E}_n}$  of  $\mathcal{P}_{\mathcal{E}_n}$  and the “boundary elements”  $\underline{P}_{\mathcal{E}}$  and  $\overline{P}_{\mathcal{E}}$  of  $\mathcal{P}_{\mathcal{E}}$  becomes progressively smaller.

## 7. PROCEDURES TO OBTAIN AND BOUND UPPER AND LOWER PROBABILITIES

As we have seen in Section 6, upper and lower probabilities associated with  $\mathcal{P}_{\mathcal{E}_n}^{\text{co}}$  encode all the information contained in the set. In a sense, they summarize all the information in the set. It is natural, then, that we focus our attention on  $\underline{P}_{\mathcal{E}_n}$  and  $\overline{P}_{\mathcal{E}_n}$ .

Let us start from the first round of belief updating. We specify  $\mathcal{P}_{\mathcal{E}_0} = \{P_{1, \mathcal{E}_0}, \dots, P_{k, \mathcal{E}_0}\} \subset \Delta(\Omega, \mathcal{F})$ . Then, we compute  $\underline{P}_{\mathcal{E}_0}$ . The set representing the agent’s initial beliefs is  $\mathcal{P}_{\mathcal{E}_0}^{\text{co}} = \text{core}(\underline{P}_{\mathcal{E}_0})$ . The exhaustive procedure to update  $\underline{P}_{\mathcal{E}_0}$  is given by the following.

- (1) We collect data  $x_1, \dots, x_n$ ;
- (2) We update all the elements in  $\mathcal{P}_{\mathcal{E}_0}$  according to the mechanical procedure in Section 4, so to obtain  $\mathcal{P}_{\mathcal{E}_1}$ ;
- (3) We calculate  $\underline{P}_{\mathcal{E}_1}(A) = \inf_{P \in \mathcal{P}_{\mathcal{E}_1}} P(A)$ , for all  $A \in \mathcal{F}$ .

$\underline{P}_{\mathcal{E}_1}$  and its conjugate  $\overline{P}_{\mathcal{E}_1}$  are the updates to  $\underline{P}_{\mathcal{E}_0}$  and  $\overline{P}_{\mathcal{E}_0}$ , respectively.

This procedure holds for updating every  $\underline{P}_{\mathcal{E}_n}$  to  $\underline{P}_{\mathcal{E}_{n+1}}$  and every  $\overline{P}_{\mathcal{E}_n}$  to  $\overline{P}_{\mathcal{E}_{n+1}}$ , not just for  $n = 0$ .

In practice, step (2) appears to be the bottleneck of the algorithm: it may be computationally intensive to compute the DPK update of all the elements in  $\mathcal{P}_{\mathcal{E}_n}$ . To overcome this shortcoming, we give a lower bound for the updated lower probability and an upper bound for the updated upper probability that can be calculated without updating all of the elements in  $\mathcal{P}_{\mathcal{E}_n}$ .

By (3), we know that for every  $P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}$  and every  $A \in \mathcal{F}$ ,

$$P_{\mathcal{E}_{n+1}}(A) = \sum_{E_j \in \mathcal{E}_{n+1}} P_{\mathcal{E}_n}(A | E_j) P_{\mathcal{E}_{n+1}}(E_j). \quad (5)$$

Also, by the intermediate step described in Remark 1, we know that, assuming the number of components of  $\mathcal{E}_{n+1}$  is  $\ell + 1$ , for all  $P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}$ ,

$$P_{\mathcal{E}_{n+1}}(E_j) = Q(E_j) = P_X(\{x_j\}), \forall j \in \{1, \dots, \ell\}, \quad P_{\mathcal{E}_{n+1}}(E_{\ell+1}) = Q(E_{\ell+1}) = 1 - \sum_{j=1}^{\ell} Q(E_j). \quad (6)$$

So, for all  $P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}$  and all  $A \in \mathcal{F}$ , we can rewrite (5) as

$$P_{\mathcal{E}_{n+1}}(A) = \sum_{E_j \in \mathcal{E}_{n+1}} P_{\mathcal{E}_n}(A | E_j) Q(E_j). \quad (7)$$

Notice that the fact that (6) holds for all  $P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}$  implies that, for all  $E_j \in \mathcal{E}_{n+1}$ ,

$$\underline{P}_{\mathcal{E}_{n+1}}(E_j) = Q(E_j) = \overline{P}_{\mathcal{E}_{n+1}}(E_j). \quad (8)$$

Both equalities in (8) hold because we implicitly assume that  $P_X$  is known. That is, the collected data are precise (i.e. with no missingness) and from a known generative model (i.e. no inference required). If this assumption were to be relaxed, these equalities need not hold.

For any  $A \in \mathcal{F}$ , and any  $E_j \in \mathcal{E}_n$ , define

$$\underline{P}_{\mathcal{E}_n}^B(A | E_j) := \inf_{P \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}} P(A | E_j) = \inf_{P \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}} \frac{P(A \cap E_j)}{P(E_j)}$$

and

$$\overline{P}_{\mathcal{E}_n}^B(A | E_j) := \sup_{P \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}} P(A | E_j) = \sup_{P \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}} \frac{P(A \cap E_j)}{P(E_j)}.$$

These are called the generalized Bayes' conditional lower and upper probabilities [19], respectively. Then, we have the following.

**Proposition 7.1.** *For any  $A \in \mathcal{F}$  and any  $n \in \mathbb{N}_0$ ,*

$$\underline{P}_{\mathcal{E}_{n+1}}(A) \geq \sum_{E_j \in \mathcal{E}_{n+1}} \underline{P}_{\mathcal{E}_n}^B(A | E_j) Q(E_j) \quad (9)$$

and

$$\overline{P}_{\mathcal{E}_{n+1}}(A) \leq \sum_{E_j \in \mathcal{E}_{n+1}} \overline{P}_{\mathcal{E}_n}^B(A | E_j) Q(E_j). \quad (10)$$

**Corollary 7.1.1.** *For all  $n \in \mathbb{N}_0$ , all  $P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}^{\text{co}}$ , and all  $A \in \mathcal{F}$ ,*

$$P_{\mathcal{E}_{n+1}}(A) \in \left[ \sum_{E_j \in \mathcal{E}_{n+1}} \underline{P}_{\mathcal{E}_n}^B(A | E_j) Q(E_j), \sum_{E_j \in \mathcal{E}_{n+1}} \overline{P}_{\mathcal{E}_n}^B(A | E_j) Q(E_j) \right].$$

The bounds we presented allow us to circumvent the bottleneck presented by step (2) in the exhaustive procedure. This because it is easier to compute  $\underline{P}_{\mathcal{E}_n}^B(A | E_j)$  and  $\overline{P}_{\mathcal{E}_n}^B(A | E_j)$ , for all  $A \in \mathcal{F}$  and all  $E_j \in \mathcal{E}_{n+1}$  rather than first computing the DIPK update of  $\mathcal{P}_{\mathcal{E}_n}$  to  $\mathcal{P}_{\mathcal{E}_{n+1}}$ , and then calculating  $\underline{P}_{\mathcal{E}_{n+1}}$  and  $\overline{P}_{\mathcal{E}_{n+1}}$ .

There are two other ways to define lower and upper conditional probabilities. The first one, called geometric update, is such that for all  $n \in \mathbb{N}_0$ , all  $A \in \mathcal{F}$ , and all  $E_j \in \mathcal{E}_0$ ,

$$\underline{P}_{\mathcal{E}_n}^G(A | E_j) := \frac{\inf_{P \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}} P(A \cap E_j)}{\inf_{P \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}} P(E_j)} = \frac{\underline{P}_{\mathcal{E}_n}(A \cap E_j)}{\underline{P}_{\mathcal{E}_n}(E_j)} \quad \text{and} \quad \overline{P}_{\mathcal{E}_n}^G(A | E_j) = \frac{\overline{P}_{\mathcal{E}_n}(A \cap E_j)}{\overline{P}_{\mathcal{E}_n}(E_j)}.$$

The other one, called Dempster's rule of conditioning, is the natural dual to the geometric procedure. It differs from this latter from the operational point of view (see [6, Section 2]), but since mathematically they are the same, we are not going to cover Dempster's rule in the present work.

An interpretation of how generalized Bayes' and geometric rules come about when a generic partition  $\{E_j\}$  of  $\Omega$  is available is the following. Let  $\sqcup$  denote the union of disjoint sets, and

$\underline{P}$  a generic lower probability. We know that lower probabilities are superadditive, so since given any  $A \in \mathcal{F}$  we have that  $A = \sqcup_j (A \cap E_j)$ , it follows that

$$\underline{P}(A) \geq \sum_j \underline{P}(A \cap E_j). \quad (11)$$

Now,  $\underline{P}(A \cap E_j)$  can be interpreted as the lowest possible probability attached to event  $A \cap E_j$ , in which case we retrieve generalized Bayes' rule. It can also be rewritten as  $\frac{\underline{P}(A \cap E_j)}{\underline{P}(E_j)} \underline{P}(E_j)$ ; in this latter case, we retrieve the geometric rule. It is worth noting that, for any lower probability  $\underline{P}$ , by [6, Lemma 5.3] we have that

$$\underline{P}^B(A | B) \leq \underline{P}^G(A | B) \leq \overline{P}^G(A | B) \leq \overline{P}^B(A | B). \quad (12)$$

for all  $A, B \in \mathcal{F}$ .

**7.1. No free lunch.** As we have seen in Proposition 7.1, generalized Bayes' comes naturally from our updating procedure. This because, as shown in Section 1.1, Jeffrey's rule is a generalization of Bayesian conditioning. Given the inequalities in (12), we can sharpen the bounds we found using generalized Bayes' rule by using the geometric rule, but this comes at the cost of an assumption. As the proverb goes, there is no free lunch. We begin with a lemma, followed by the result.

**Lemma 2.** *For any  $n \in \mathbb{N}_0$ , the following are true.*

$$\sum_{E_j \in \mathcal{E}_{n+1}} \underline{P}_{\mathcal{E}_n}^G(A | E_j) Q(E_j) \geq \sum_{E_j \in \mathcal{E}_{n+1}} \underline{P}_{\mathcal{E}_n}^B(A | E_j) Q(E_j)$$

and

$$\sum_{E_j \in \mathcal{E}_{n+1}} \overline{P}_{\mathcal{E}_n}^G(A | E_j) Q(E_j) \leq \sum_{E_j \in \mathcal{E}_{n+1}} \overline{P}_{\mathcal{E}_n}^B(A | E_j) Q(E_j).$$

**Proposition 7.2.** *In the event that  $\frac{\underline{P}_{\mathcal{E}_{n+1}}(A \cap E_j)}{\underline{P}_{\mathcal{E}_{n+1}}(E_j)} = \frac{\underline{P}_{\mathcal{E}_n}(A \cap E_j)}{\underline{P}_{\mathcal{E}_n}(E_j)}$  for all  $n \in \mathbb{N}_0$ , all  $A \in \mathcal{F}$ , and all  $E_j \in \mathcal{E}_1$ , the following hold*

$$\underline{P}_{\mathcal{E}_{n+1}}(A) \geq \sum_{E_j \in \mathcal{E}_{n+1}} \underline{P}_{\mathcal{E}_n}^G(A | E_j) Q(E_j) \quad (13)$$

and

$$\overline{P}_{\mathcal{E}_{n+1}}(A) \leq \sum_{E_j \in \mathcal{E}_{n+1}} \overline{P}_{\mathcal{E}_n}^G(A | E_j) Q(E_j). \quad (14)$$

Condition  $\frac{\underline{P}_{\mathcal{E}_{n+1}}(A \cap E_j)}{\underline{P}_{\mathcal{E}_{n+1}}(E_j)} = \frac{\underline{P}_{\mathcal{E}_n}(A \cap E_j)}{\underline{P}_{\mathcal{E}_n}(E_j)}$  in Proposition 7.2 requires that the relative lower probability assigned to  $A \cap E_j$  with respect to the one assigned to  $E_j$ , for all  $A \in \mathcal{F}$  and all  $E_j$  in the updated partition, has to stay constant when we perform the DIPK update of a probability set. In a sense, this condition is an assumption governs the dynamic of the lower

probability assignment. Since Lemma 2 implies that

$$\begin{aligned} & \left[ \sum_{E_j \in \mathcal{E}_{n+1}} \underline{P}_{\mathcal{E}_n}^G(A | E_j)Q(E_j), \sum_{E_j \in \mathcal{E}_{n+1}} \overline{P}_{\mathcal{E}_n}^G(A | E_j)Q(E_j) \right] \\ & \subset \left[ \sum_{E_j \in \mathcal{E}_{n+1}} \underline{P}_{\mathcal{E}_n}^B(A | E_j)Q(E_j), \sum_{E_j \in \mathcal{E}_{n+1}} \overline{P}_{\mathcal{E}_n}^B(A | E_j)Q(E_j) \right], \end{aligned}$$

the extra cost of the assumption in Proposition 7.2 helps retrieve tighter bounds for  $\underline{P}_{\mathcal{E}_{n+1}}(A)$  and  $\overline{P}_{\mathcal{E}_{n+1}}(A)$ , and also obtain a tighter interval around  $P_{\mathcal{E}_{n+1}}(A)$ , for all  $A \in \mathcal{F}$  and all  $n \in \mathbb{N}_0$ .

## 8. BEHAVIOR OF UPDATED SETS OF PROBABILITIES

In the imprecise probabilities literature, three concepts are crucial regarding the behavior of updated sets of probabilities. They are contraction, dilation, and sure loss. In this Section, we adapt the definition of these phenomena to our updating procedure, and we give sufficient conditions for them to take place.

Fix some  $n \in \mathbb{N}$ . We say that  $\mathcal{P}_{\mathcal{E}_n}^{\text{co}}$  *contracts with respect to*  $\mathcal{P}_{\mathcal{E}_{n-1}}^{\text{co}}$  for some  $A \in \mathcal{F}$  if  $\underline{P}_{\mathcal{E}_n}(A) \geq \underline{P}_{\mathcal{E}_{n-1}}(A)$  and  $\overline{P}_{\mathcal{E}_n}(A) \leq \overline{P}_{\mathcal{E}_{n-1}}(A)$ . It *strictly contracts* if the inequalities are strict. In addition, we say that sequence  $(\mathcal{P}_{\mathcal{E}_n}^{\text{co}})$  *contracts for some*  $A \in \mathcal{F}$  if for any  $n \in \mathbb{N}$ , we have that  $\underline{P}_{\mathcal{E}_n}(A) \geq \underline{P}_{\mathcal{E}_{n-1}}(A)$ ,  $\overline{P}_{\mathcal{E}_n}(A) \leq \overline{P}_{\mathcal{E}_{n-1}}(A)$ , and the inequalities are strict for some  $n$ .

*Dilation* is defined analogously, by inverting the inequality signs.

Finally, we say that  $\mathcal{P}_{\mathcal{E}_n}^{\text{co}}$  exhibits *sure loss with respect to*  $\mathcal{P}_{\mathcal{E}_{n-1}}^{\text{co}}$  for some  $A \in \mathcal{F}$  if  $\underline{P}_{\mathcal{E}_n}(A) > \overline{P}_{\mathcal{E}_{n-1}}(A)$  or  $\overline{P}_{\mathcal{E}_n}(A) < \underline{P}_{\mathcal{E}_{n-1}}(A)$ .

Despite being similar to the classical definitions of contraction, dilation, and sure loss – see e.g. [6, Section 3] – our definitions differ slightly to take into account our updating procedure. For this reason, we find results that in a classical imprecise probabilities setting may be impossible. For example, as shown in [6, Section 5.1], in a classical imprecise probabilities setting, generalized Bayes' rule cannot contract nor induce sure loss. Instead, we have the following.

**Proposition 8.1.** *For any  $n \in \mathbb{N}$ , sufficient conditions for  $\mathcal{P}_{\mathcal{E}_n}^{\text{co}}$  to contract with respect to  $\mathcal{P}_{\mathcal{E}_{n-1}}^{\text{co}}$  for some  $A \in \mathcal{F}$  are the following*

$$\sum_{E_j \in \mathcal{E}_n} \underline{P}_{\mathcal{E}_{n-1}}^B(A | E_j)Q(E_j) \geq \underline{P}_{\mathcal{E}_{n-1}}(A) \quad \text{and} \quad \sum_{E_j \in \mathcal{E}_n} \overline{P}_{\mathcal{E}_{n-1}}^B(A | E_j)Q(E_j) \leq \overline{P}_{\mathcal{E}_{n-1}}(A).$$

Notice that we obtain strict contraction if the inequalities are strict. We have the same results if we use geometric lower conditional probabilities instead of the generalized Bayes' ones, provided that the assumption in Proposition 7.2 holds.

We also have the following.

**Proposition 8.2.** *For any  $n \in \mathbb{N}$ , sufficient conditions for  $\mathcal{P}_{\mathcal{E}_n}^{\text{co}}$  to exhibit sure loss with respect to  $\mathcal{P}_{\mathcal{E}_{n-1}}^{\text{co}}$  for some  $A \in \mathcal{F}$  are the following*

$$\sum_{E_j \in \mathcal{E}_n} \underline{P}_{\mathcal{E}_{n-1}}^B(A | E_j)Q(E_j) > \overline{P}_{\mathcal{E}_{n-1}}(A) \quad \text{or} \quad \sum_{E_j \in \mathcal{E}_n} \overline{P}_{\mathcal{E}_{n-1}}^B(A | E_j)Q(E_j) < \underline{P}_{\mathcal{E}_{n-1}}(A).$$

Again, we obtain the same conditions if we use geometric lower conditional probabilities instead of the generalized Bayes' ones, provided that the assumption in Proposition 7.2 holds.

Giving a sufficient condition for dilation without directly computing  $\underline{P}_{\mathcal{E}_n}(A)$  and  $\overline{P}_{\mathcal{E}_n}(A)$  is less straightforward. This because, unlike contraction and sure loss for which the upper and lower probabilities from DIPK updating are conservative bounds to the exhaustive methods, the same bounds cannot be used to sufficiently establish dilation. We have the following.

**Proposition 8.3.** *For any  $n \in \mathbb{N}$  and some  $A \in \mathcal{F}$ , if there exist  $P_{s,\mathcal{E}_n}, P_{k,\mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}$  such that  $\underline{P}_{\mathcal{E}_{n-1}}(A) \geq P_{s,\mathcal{E}_n}(A)$  and  $\overline{P}_{\mathcal{E}_{n-1}}(A) \leq P_{k,\mathcal{E}_n}(A)$ , then  $\mathcal{P}_{\mathcal{E}_n}^{\text{co}}$  dilates with respect to  $\mathcal{P}_{\mathcal{E}_{n-1}}^{\text{co}}$  for  $A$ .*

We obtain strict dilation if the inequalities in Proposition 8.3 are strict. As we can see, we do not need to directly compute  $\underline{P}_{\mathcal{E}_n}(A)$  and  $\overline{P}_{\mathcal{E}_n}(A)$ . We only need to find  $P_{s,\mathcal{E}_{n-1}}, P_{k,\mathcal{E}_{n-1}} \in \mathcal{P}_{\mathcal{E}_{n-1}}^{\text{co}}$  such that their updates satisfy the assumptions in Proposition 8.3.

We can give a result, similar to Proposition 8.3 that provides sufficient conditions for  $\mathcal{P}_{\mathcal{E}_n}^{\text{co}}$  to contract with respect to  $\mathcal{P}_{\mathcal{E}_{n-1}}^{\text{co}}$  for some  $A \in \mathcal{F}$ . This is interesting because, contrary to what we have in Proposition 8.1, we do not use the notions of lower and upper conditional probabilities. It is less useful from a computational perspective, though, since it requires the computation of both  $\underline{P}_{\mathcal{E}_n}(A)$  and  $\overline{P}_{\mathcal{E}_n}(A)$ .

**Proposition 8.4.** *For any  $n \in \mathbb{N}$  and some  $A \in \mathcal{F}$ , if there exist  $P_{s,\mathcal{E}_{n-1}}, P_{k,\mathcal{E}_{n-1}} \in \mathcal{P}_{\mathcal{E}_{n-1}}^{\text{co}}$  such that  $\underline{P}_{\mathcal{E}_n}(A) \geq P_{k,\mathcal{E}_{n-1}}(A)$  and  $\overline{P}_{\mathcal{E}_n}(A) \leq P_{s,\mathcal{E}_{n-1}}(A)$ , then  $\mathcal{P}_{\mathcal{E}_n}^{\text{co}}$  contracts with respect to  $\mathcal{P}_{\mathcal{E}_{n-1}}^{\text{co}}$  for  $A$ .*

We obtain strict contraction if the inequalities in Proposition 8.4 are strict.

Notice that we cannot give a result similar to Propositions 8.3 and 8.4 for sure loss because we cannot require any assumption on any  $P_{\mathcal{E}_{n-1}} \in \mathcal{P}_{\mathcal{E}_{n-1}}^{\text{co}}$ ,  $P_{\mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}$  to make them “sit in between”  $\underline{P}_{\mathcal{E}_n}(A)$  and  $\overline{P}_{\mathcal{E}_{n-1}}(A)$ , or  $\overline{P}_{\mathcal{E}_n}(A)$  and  $\underline{P}_{\mathcal{E}_{n-1}}(A)$ .

## 9. AN APPLICATION TO SURVEY SAMPLING STUDIES

Survey questions typically standardize their answers into a finite number of bins. For example, a simple questionnaire, consisting of just one question about age, can have the following choices:

- (1) under 18;
- (2) 18 to 64;
- (3) 65 and above,

for which  $\mathcal{X}_1 = \{1, 2, 3\}$ , and the corresponding partition is  $\check{\mathcal{E}}_1 = \{[0, 18), [18, 65), [65, \infty)\}$ . A second simple questionnaire, either newly administered or as a follow up to the previous, can refine this into

- (1) under 18;

- (2) 18 to 20;
- (3) 21 to 44;
- (4) 45 to 64;
- (5) 65 and above,

for which  $\mathcal{X}_2 = \{1, 2, 3, 4, 5\}$ , and the corresponding partition is  $\check{\mathcal{E}}_2 = \{[0, 18), [18, 21), [21, 45), [45, 65), [65, \infty)\}$ .  $\check{\mathcal{E}}_2$  is a refinement of  $\check{\mathcal{E}}_1$ , although  $\mathcal{X}_2$  does not have a particular bearing to  $\mathcal{X}_1$  since the nestedness of options is not preserved (it can be, if needed). A third questionnaire can have two questions, add to the five-category age question a second question about race, with answers

- (1) White (W);
- (2) Black (B);
- (3) Asian and pacific islander (A);
- (4) Other or two or more (O),

for which  $\mathcal{X}_3 = \mathcal{X}_2 \times \{1, 2, 3, 4\}$ , and the corresponding partition is  $\check{\mathcal{E}}_3 = \check{\mathcal{E}}_2 \times \{W, B, A, O\}$ .

How do we connect these progressively more refined surveys to our model? Let  $\mathcal{X} = \mathcal{X}_3$ , the finest partition of concern as defined the three questionnaires taken together. A random batch of subjects were interviewed on their age and race, and we obtained the following unique reports  $x_1 = (1, 1), x_2 = (1, 2), x_3 = (1, 3), x_4 = (1, 4), x_5 = (5, 1), x_6 = (5, 2), x_7 = (5, 3), x_8 = (5, 4)$ , invoking the corresponding partition

$$\begin{aligned} \mathcal{E}_1 &= \{E_1 = ([0, 18), W), E_2 = ([0, 18), B), E_3 = ([0, 18), A), E_4 = ([0, 18), O), \\ &E_5 = ([65, \infty), W), E_6 = ([65, \infty), B), E_7 = ([65, \infty), A), E_8 = ([65, \infty), O), \\ &E_9 = \Omega \setminus \cup_{j=1}^8 E_j\}. \end{aligned}$$

Then, we have that  $P_{\mathcal{E}_1}(E_j) = Q(E_j) = P_X(\{x_j\})$ , for  $j \in \{1, \dots, 8\}$ , and  $P_{\mathcal{E}_1}(E_9) = Q(E_9) = 1 - \sum_{j=1}^8 Q(E_j)$ . The above observations allow to compute  $P_{\mathcal{E}_1}(A)$  mechanically for every  $A \in \mathcal{F}$ .

We first attempt to make sense of these observations in light of the first survey questionnaire, consisting of just three age categories. The outcomes in relation to the first survey are obtained by letting

$$\check{\mathcal{E}}_1 = \{\check{E}_1 = \cup_{j=1}^4 E_j, \check{E}_2 = \cup_{j=5}^8 E_j, \check{E}_3 = E_9\}.$$

Putting  $P_{\check{\mathcal{E}}_1}(\check{E}_1) = \sum_{j=1}^4 Q(E_j)$ ,  $P_{\check{\mathcal{E}}_1}(\check{E}_2) = \sum_{j=5}^8 Q(E_j)$ ,  $P_{\check{\mathcal{E}}_1}(\check{E}_3) = P_{\mathcal{E}_1}(E_9) = Q(E_9)$  allows to compute  $P_{\check{\mathcal{E}}_1}(A)$  mechanically for every  $A \in \mathcal{F}$ .

We proceed to interview a second batch of randomly chosen subjects, who provided 12 more unique reports  $x_9 = (2, 1), x_{10} = (2, 2), x_{11} = (2, 3), x_{12} = (2, 4), x_{13} = (3, 1), x_{14} = (3, 2), x_{15} = (3, 3), x_{16} = (3, 4), x_{17} = (4, 1), x_{18} = (4, 2), x_{19} = (4, 3), x_{20} = (4, 4)$ . Taken

together, the two batches of reports invoke the partition

$$\begin{aligned} \mathcal{E}_2 = \{ & E_1 = ([0, 18), W), E_2 = ([0, 18), B), E_3 = ([0, 18), A), E_4 = ([0, 18), O), \\ & E_5 = ([65, \infty), W), E_6 = ([65, \infty), B), E_7 = ([65, \infty), A), E_8 = ([65, \infty), O), \\ & E_9 = ([18, 21), W), E_{10} = ([18, 21), B), E_{11} = ([18, 21), A), E_{12} = ([18, 21), O), \\ & E_{13} = ([21, 45), W), E_{14} = ([21, 45), B), E_{15} = ([21, 45), A), E_{16} = ([21, 45), O), \\ & E_{17} = ([45, 65), W), E_{18} = ([45, 65), B), E_{19} = ([45, 65), A), E_{20} = ([45, 65), O), \\ & E_{21} = \emptyset\}. \end{aligned}$$

Thus we have that  $P_{\mathcal{E}_2}(E_j) = Q(E_j) = P_X(\{x_j\})$ , for  $j \in \{1, \dots, 20\}$ , and  $P_{\mathcal{E}_2}(E_{21}) = 0$ . This allows to compute  $P_{\mathcal{E}_2}(A)$  mechanically for every  $A \in \mathcal{F}$ .

With this, we would like to update our opinion by making sense of the reports in light of the second survey, consisting five age categories. Let

$$\begin{aligned} \check{\mathcal{E}}_2 = \{ & \check{E}_1 = \cup_{j=1}^4 E_j, \check{E}_2 = \cup_{j=5}^8 E_j, \check{E}_3 = \cup_{j=9}^{12} E_j, \\ & \check{E}_4 = \cup_{j=13}^{16} E_j, \check{E}_5 = \cup_{j=17}^{20} E_j, \check{E}_6 = E_{21} = \emptyset\}, \end{aligned}$$

and put  $P_{\check{\mathcal{E}}_2}(\check{E}_1) = \sum_{j=1}^4 Q(E_j)$ ,  $P_{\check{\mathcal{E}}_2}(\check{E}_2) = \sum_{j=5}^8 Q(E_j)$ ,  $P_{\check{\mathcal{E}}_2}(\check{E}_3) = \sum_{j=9}^{12} Q(E_j)$ ,  $P_{\check{\mathcal{E}}_2}(\check{E}_4) = \sum_{j=13}^{16} Q(E_j)$ ,  $P_{\check{\mathcal{E}}_2}(\check{E}_5) = \sum_{j=17}^{20} Q(E_j)$ ,  $P_{\check{\mathcal{E}}_2}(\check{E}_6) = 0$ . This allows us to compute  $P_{\check{\mathcal{E}}_2}(A)$  mechanically for every  $A \in \mathcal{F}$ .

We now further update our opinion relative to the combination of the third survey, consisting of  $5 \times 4 = 20$  unique age-race categories. Since we have already seen all the elements  $x$  in  $\mathcal{X}$ , there is no need to observe new data points, but we adopt the finer partition  $\check{\mathcal{E}}_3 = \mathcal{E}_2$  and directly apply our method.

This example is set up in a way that the three surveys offer successively finer partition of the state space. To derive result for the first two surveys calls for an extra ‘‘coarsening step’’, by gathering the more informative observations into the coarser partition in which they belong. This slight variation allows for the use of dynamic probability kinematics when dealing with the update of initial opinions given by the results of a survey sample analysis where increasingly finer questions, or two or more ‘‘dimesions’’ are surveyed. Notice that  $d_{TV}(P_{\check{\mathcal{E}}_n}, P_{\mathcal{E}_n}) \rightarrow 0$  as  $n \rightarrow \infty$ , and so  $(P_{\check{\mathcal{E}}_n})$  converges in the total variation metric to  $P_{\check{\mathcal{E}}}$ .

The generalization of this variation to our method to sets of probabilities is straightforward. We consider a finite set  $\mathcal{P}$  of probability measures, compute the lower probability associated with it, and set  $\mathcal{P}^{\text{co}}$  equal to the core of such lower probability.  $\mathcal{P}^{\text{co}}$  represents the agent’s initial opinions. To update it, we update all of the elements  $P \in \mathcal{P}$  to  $P_{\check{\mathcal{E}}_1}$ , so to obtain set  $\mathcal{P}_{\check{\mathcal{E}}_1}$ . We then compute its lower probability  $\underline{P}_{\check{\mathcal{E}}_1}$ , and set  $\mathcal{P}_{\check{\mathcal{E}}_1}^{\text{co}} = \text{core}(\underline{P}_{\check{\mathcal{E}}_1})$ . After that, we proceed along the lines of what we did in Section 6. It is immediate to see that  $d_H(\mathcal{P}_{\check{\mathcal{E}}_n}, \mathcal{P}_{\check{\mathcal{E}}}) \rightarrow 0$  as  $n \rightarrow \infty$ , and that  $d_H(\mathcal{P}_{\check{\mathcal{E}}_n}^{\text{co}}, \mathcal{P}_{\check{\mathcal{E}}}^{\text{co}}) \rightarrow 0$  as  $n \rightarrow \infty$ . Equivalent results to those in Sections 7 and 8 can be found for the variation to our method by replacing  $(\mathcal{P}_{\check{\mathcal{E}}_n}^{\text{co}})$  with  $(\mathcal{P}_{\check{\mathcal{E}}_n}^{\text{co}})$ .

## 10. CONCLUSION

In this paper, we presented the dynamic probability kinematics (DPK) and dynamic imprecise probability kinematics (DIPK), a pair of methods to dynamically update subjective

beliefs stated in terms of precise and imprecise probabilities based on Jeffrey's rule of conditioning. In the case of DIPK, we provided bounds for the upper and lower probabilities associated with the updated sets, and studied their set-specific behavior including contraction, dilation, and sure loss. An example of survey sampling studies is used to illustrate the application of the proposal.

In the future, we plan to deal with imprecise or inconsistent data collection, in the spirit of [13]. This happens when  $P_X$  that characterizes the distribution of the random variables driving the updating process is not known or is incompletely specified. A way to account for this could be to specify a vacuous prior for  $P_X$ , i.e. a prior belonging to the maximal set of probabilities defined on  $\mathcal{X}$ . We also plan to further generalize DPK and DIPK by allowing the range  $\mathcal{X}$  of the random variables driving the updating process to be uncountable.

#### ACKNOWLEDGEMENTS

The authors thank Sayan Murkherjee for inspiring this project and helpful discussions, Teddy Seidenfeld for a fruitful discussion about the commutativity of dynamic probability kinematics, and Xiao-Li Meng for an insightful dialogue on information theory. Research of R. Gong is supported in part by the National Science Foundation (DMS-1916002). M. Caprio too would like to acknowledge partial funding by the National Science Foundation (CCF-1934964).

#### APPENDIX A. PROOFS

*Proof of Proposition 3.1.* We begin by showing that  $P_{\mathcal{E}}$  is a probability measure. We verify the Kolmogorovian axioms for a probability measure. First, we have that  $P_{\mathcal{E}}(A) \geq 0$ , for all  $A \in \mathcal{F}$ . This comes by its definition, since it is defined as the summation of products of nonnegative quantities. Second, we have that  $P_{\mathcal{E}}(\Omega) = 1$ . This comes from the following

$$P_{\mathcal{E}}(\Omega) = \sum_{E_j \in \mathcal{E}} P(\Omega | E_j) P_{\mathcal{E}}(E_j) = \sum_{E_j \in \mathcal{E}} P_{\mathcal{E}}(E_j) = 1.$$

Finally, we have that if  $\{A_i\}_{i \in I}$  is a countable, pairwise disjoint collection of events, then  $P_{\mathcal{E}}(\cup_{i \in I} A_i) = \sum_{i \in I} P_{\mathcal{E}}(A_i)$ . This because

$$\begin{aligned} P_{\mathcal{E}}(\cup_{i \in I} A_i) &= \sum_{E_j \in \mathcal{E}} P(\cup_{i \in I} A_i | E_j) P_{\mathcal{E}}(E_j) \\ &= \sum_{E_j \in \mathcal{E}} \frac{P([\cup_{i \in I} A_i] \cap E_j)}{P(E_j)} P_{\mathcal{E}}(E_j) \\ &= \sum_{E_j \in \mathcal{E}} \frac{P(\cup_{i \in I} [A_i \cap E_j])}{P(E_j)} P_{\mathcal{E}}(E_j) \\ &= \sum_{E_j \in \mathcal{E}} \frac{\sum_{i \in I} P(A_i \cap E_j)}{P(E_j)} P_{\mathcal{E}}(E_j) \\ &= \sum_{i \in I} \sum_{E_j \in \mathcal{E}} \frac{P(A_i \cap E_j)}{P(E_j)} P_{\mathcal{E}}(E_j) = \sum_{i \in I} P_{\mathcal{E}}(A_i). \end{aligned}$$

We now show that  $P_{\mathcal{E}}$  is a Jeffrey's posterior for  $P$ . We use [5, Theorem 2.1]: it states that  $P^*$  is a Jeffrey's posterior for  $P$  if and only if there exists a constant  $B \geq 1$  such that  $P^*(\{\omega\}) \leq BP(\{\omega\})$ , for all  $\omega \in \Omega$ . Fix any  $\omega \in \Omega$ . We have that  $P_{\mathcal{E}}(\{\omega\}) = \sum_{E_j \in \mathcal{E}} P(\{\omega\} | E_j) P_{\mathcal{E}}(E_j)$ . Call  $E_{\omega}$  the element in  $\mathcal{E}$  such that  $\{\omega\} \cap E_{\omega} \neq \emptyset$ . Then, we have that

$$P_{\mathcal{E}}(\{\omega\}) = \sum_{E_j \in \mathcal{E}} P(\{\omega\} | E_j) P_{\mathcal{E}}(E_j) = \frac{P_{\mathcal{E}}(E_{\omega})}{P(E_{\omega})} P(\{\omega\}). \quad (15)$$

Now, let  $B_{\omega} := \lceil \frac{P_{\mathcal{E}}(E_{\omega})}{P(E_{\omega})} + 1 \rceil$ . We have that  $P_{\mathcal{E}}(\{\omega\}) < B_{\omega} P(\{\omega\})$ . Consider then the well-ordered collection  $\{B_{\omega}\}_{\omega \in \Omega}$ . If we let  $B := \sup_{B'_{\omega} \in \{B_{\omega}\}} B'_{\omega}$ , we conclude that  $P_{\mathcal{E}}(\{\omega\}) < BP(\{\omega\})$ , for all  $\omega \in \Omega$ .  $\square$

Let us discuss about the ratio  $\frac{P_{\mathcal{E}}(E_{\omega})}{P(E_{\omega})}$  in (15). In the analysis in Section 3, we made a tacit assumption, that is,  $P(E_j) = 0 \iff E_j = \emptyset$ ,  $E_j \in \mathcal{E}$ . This is needed for the ratio  $\frac{P_{\mathcal{E}}(E_{\omega})}{P(E_{\omega})}$  to be well defined for all  $\omega \in \Omega$ . Notice that if there is an  $E_j \in \mathcal{E}$  such that  $E_j = \emptyset$ , this means that no  $\omega$  belongs to  $E_j$ , so the ratio  $\frac{P_{\mathcal{E}}(E_{\omega})}{P(E_{\omega})}$  is well defined for all  $\omega \in \Omega$ .

*Proof of Lemma 1. If.* Let  $E = \emptyset$ . Then,  $P_{\mathcal{E}}(E) = 0$ .

**Only if.** Let  $P_{\mathcal{E}}(E) = 0$ . Notice that, since  $\mathcal{X} = \text{Im}_X(\text{Dom}(X))$ , we have that  $P_X(\{x\}) > 0$ , for all  $x \in \mathcal{X}$ . By the mechanical procedure in Section 4, we have that  $P_{\mathcal{E}}(E_j) = P_X(\{x_j\})$ , for all  $j \in \{1, \dots, n\}$ . Then, because  $P_X(\{x\}) > 0$ , for all  $x \in \mathcal{X}$ , we get that  $P_{\mathcal{E}}(E_j) > 0$ , for all  $j \in \{1, \dots, n\}$ . This leaves us with the fact that the only  $E \in \mathcal{E}$  that can have probability 0 is  $E_{n+1} = (\cup_{j=1}^n E_j)^c$ . Then, we have that  $0 = P_{\mathcal{E}}(E_{n+1}) = 1 - \sum_{j=1}^n P_{\mathcal{E}}(E_j)$ , which implies that  $\sum_{j=1}^n P_{\mathcal{E}}(E_j) = 1$ . This in turn implies that  $\sum_{j=1}^n P_X(\{x_j\}) = 1$ . But this latter holds if and only if  $\cup_{j=1}^n x_j = \mathcal{X}$ . This implies that  $\cup_{j=1}^n E_j = \Omega$ , and so  $E_{n+1} = \Omega^c = \emptyset$ .  $\square$

*Proof of Proposition 5.1.* We have two cases.

**Case 1** Let  $\mathcal{X}$  be finite, that is,  $\cup_{j=1}^{\ell} x_j = \mathcal{X}$ , for some  $\ell$ . Then, suppose at time  $T \in \mathbb{N}$  we observe all  $x_j$ 's in  $\mathcal{X}$ . Then,  $\mathcal{E}_T = \{E_1, \dots, E_{\ell}, E_{\ell+1} = \emptyset\}$  and  $\mathcal{E}_t = \mathcal{E}_T$ , for all  $t \geq T$ . In turn, this implies that  $P_{\mathcal{E}_T} = P_{\mathcal{E}_t}$ , for all  $t \geq T$ . This concludes the proof for the finite  $\mathcal{X}$  case.

**Case 2** Let now  $\mathcal{X}$  be countable. We first show that  $(\Delta(\Omega, \mathcal{F}), d_{TV})$  is a complete space, where  $\Delta(\Omega, \mathcal{F})$  is the space of probability measures on  $(\Omega, \mathcal{F})$ , and  $d_{TV}$  is the total variation metric induced by the total variation norm. Let us call  $\mathcal{M}(\Omega, \mathcal{F})$  the linear space of all bounded (signed) measures on  $(\Omega, \mathcal{F})$ . As shown in [1, 14], this is a Banach space with respect to the total variation norm

$$\|\mu\|_{TV} := \sup_{A \in \mathcal{F}} (|\mu(A)| + |\mu(A^c)|).$$

Hence,  $(\mathcal{M}(\Omega, \mathcal{F}), \|\cdot\|_{TV})$  is complete. Define now  $f : \mathcal{M}(\Omega, \mathcal{F}) \rightarrow \mathbb{R}$ ,  $\mu \mapsto f(\mu) := \mu(\Omega)$ . This map is continuous,  $|f(\mu) - f(\nu)| \leq \|\mu - \nu\|_{TV}$ . Hence,  $\mathcal{M}_1(\Omega, \mathcal{F}) := \{\mu \in \mathcal{M}(\Omega, \mathcal{F}) : f(\mu) = 1\}$  is a closed subspace of  $\mathcal{M}(\Omega, \mathcal{F})$ . The space of all non-negative measures  $\mathcal{M}_+(\Omega, \mathcal{F}) := \{\mu \in \mathcal{M}(\Omega, \mathcal{F}) : \mu \geq 0\}$  can be characterized as  $\mathcal{M}_+(\Omega, \mathcal{F}) = \{\mu \in \mathcal{M}(\Omega, \mathcal{F}) : \|\mu\|_{TV} - f(\mu) = 0\}$ , and so it is a closed subspace of  $\mathcal{M}(\Omega, \mathcal{F})$  as well. The space of probability measures  $\Delta(\Omega, \mathcal{F}) = \mathcal{M}_1(\Omega, \mathcal{F}) \cap \mathcal{M}_+(\Omega, \mathcal{F})$  is then a closed subspace of a complete space, hence

complete itself. Notice also that the total variation distance  $d_{TV}$  between two probability measures  $\mu, \nu \in \Delta(\Omega, \mathcal{F})$  is the metric induced by the total variation norm,

$$d_{TV}(\mu, \nu) := \|\mu - \nu\|_{TV} = 2 \sup_{A \in \mathcal{F}} |\mu(A) - \nu(A)|.$$

Factor 2 can be dropped without loss of generality, and  $(\Delta(\Omega, \mathcal{F}), d_{TV})$  is still a complete space.

Now, to show that  $\lim_{n \rightarrow \infty} P_{\mathcal{E}_n}$  exists and is well defined, we need to show that  $(P_{\mathcal{E}_n})_n$  is a Cauchy sequence. That is, we need to show that  $d_{TV}(P_{\mathcal{E}_{n-1}}, P_{\mathcal{E}_n})$  goes to 0 as  $n \rightarrow \infty$ . Let us assume, without loss of generality, that we collect only one data point at a time. Then, the number of elements of partition  $\mathcal{E}_n$  are  $n + 1$ , for all  $n \in \mathbb{N}$ . Let us denote by  $E_j^{\mathcal{E}_n}$  the elements of partition  $\mathcal{E}_n$ .  $E_1^{\mathcal{E}_n}, \dots, E_n^{\mathcal{E}_n}$  are the elements induced by all the observations  $x_1, \dots, x_n$  available at time  $n$ , and  $E_{n+1}^{\mathcal{E}_n} := (\cup_{j=1}^n E_j^{\mathcal{E}_n})^c$ .

Notice now that we can write

$$P_{\mathcal{E}_n}(A) = \sum_{E_j^{\mathcal{E}_n}} P_{\mathcal{E}_{n-1}}(A | E_j^{\mathcal{E}_n}) P_{\mathcal{E}_n}(E_j^{\mathcal{E}_n}),$$

for all  $A \in \mathcal{F}$ , by (3), and

$$P_{\mathcal{E}_{n-1}}(A) = \sum_{E_j^{\mathcal{E}_n}} P_{\mathcal{E}_{n-1}}(A | E_j^{\mathcal{E}_n}) P_{\mathcal{E}_{n-1}}(E_j^{\mathcal{E}_n}),$$

for all  $A \in \mathcal{F}$ , by the fact that  $P_{\mathcal{E}_{n-1}}$  is a probability measure. Then, we have that

$$\begin{aligned} d_{TV}(P_{\mathcal{E}_{n-1}}, P_{\mathcal{E}_n}) &:= \sup_{A \in \mathcal{F}} |P_{\mathcal{E}_{n-1}}(A) - P_{\mathcal{E}_n}(A)| \\ &= \sup_{A \in \mathcal{F}} \left| \sum_{E_j^{\mathcal{E}_n}} P_{\mathcal{E}_{n-1}}(A | E_j^{\mathcal{E}_n}) [P_{\mathcal{E}_{n-1}}(E_j^{\mathcal{E}_n}) - P_{\mathcal{E}_n}(E_j^{\mathcal{E}_n})] \right| \\ &\leq \sup_{A \in \mathcal{F}} \sum_{E_j^{\mathcal{E}_n}} P_{\mathcal{E}_{n-1}}(A | E_j^{\mathcal{E}_n}) |P_{\mathcal{E}_{n-1}}(E_j^{\mathcal{E}_n}) - P_{\mathcal{E}_n}(E_j^{\mathcal{E}_n})| \\ &\leq \sum_{E_j^{\mathcal{E}_n}} |P_{\mathcal{E}_{n-1}}(E_j^{\mathcal{E}_n}) - P_{\mathcal{E}_n}(E_j^{\mathcal{E}_n})| \end{aligned} \quad (16)$$

$$= |P_{\mathcal{E}_{n-1}}(E_n^{\mathcal{E}_n}) - P_{\mathcal{E}_n}(E_n^{\mathcal{E}_n})| + |P_{\mathcal{E}_{n-1}}(E_{n+1}^{\mathcal{E}_n}) - P_{\mathcal{E}_n}(E_{n+1}^{\mathcal{E}_n})|. \quad (17)$$

Equation (16) is true because  $P_{\mathcal{E}_{n-1}}(A | E_j^{\mathcal{E}_n}) \in [0, 1]$ , for all  $j$ , for all  $A \in \mathcal{F}$ . Equation (17) is true because, given our assumption that we only observe one new data point at a time, we have that  $E_j^{\mathcal{E}_{n-1}} = E_j^{\mathcal{E}_n}$ , for all  $j \in \{1, \dots, n-1\}$ , and

$$(\cup_{j=1}^{n-1} E_j^{\mathcal{E}_{n-1}})^c \equiv E_n^{\mathcal{E}_{n-1}} = E_n^{\mathcal{E}_n} \cup E_{n+1}^{\mathcal{E}_n} \equiv E_n^{\mathcal{E}_n} \cup (\cup_{j=1}^n E_j^{\mathcal{E}_n})^c.$$

In turn, given the update procedure described in Section 4, this implies that the only elements of  $\mathcal{E}_n$  that  $P_{\mathcal{E}_{n-1}}$  and  $P_{\mathcal{E}_n}$  do not agree on – that is, the only elements of  $\mathcal{E}_n$  that are not assigned the same probability by both  $P_{\mathcal{E}_{n-1}}$  and  $P_{\mathcal{E}_n}$  – are  $E_n^{\mathcal{E}_n}$  and  $E_{n+1}^{\mathcal{E}_n}$ .

Now, we first notice that  $|P_{\mathcal{E}_{n-1}}(E_{n+1}^{\mathcal{E}_n}) - P_{\mathcal{E}_n}(E_{n+1}^{\mathcal{E}_n})|$  goes to 0 as  $n \rightarrow \infty$ . This because  $(\cup_{j=1}^n x_j)^c \rightarrow \mathcal{X}^c = \emptyset$  as  $n \rightarrow \infty$ , so  $P_X((\cup_{j=1}^n x_j)^c) \rightarrow P_X(\emptyset) = 0$  as  $n \rightarrow \infty$ . In turn, this implies that  $P_{\mathcal{E}_n}(E_{n+1}^{\mathcal{E}_n}) \rightarrow 0$  as  $n \rightarrow \infty$ .

Let us now focus on  $|P_{\mathcal{E}_{n-1}}(E_n^{\mathcal{E}_n}) - P_{\mathcal{E}_n}(E_n^{\mathcal{E}_n})|$ . Fix any  $\epsilon > 0$ , and consider  $N$  such that  $P_{\mathcal{E}_{N-1}}(E_N^{\mathcal{E}_{N-1}}) \leq \epsilon$ .

Notice that  $E_N^{\mathcal{E}_{N-1}} = E_N^{\mathcal{E}_N} \cup E_{N+1}^{\mathcal{E}_N}$ , so  $P_{\mathcal{E}_{N-1}}(E_N^{\mathcal{E}_N}) < \epsilon$  by the monotonicity of probability measures.<sup>2</sup> Notice also that

$$\epsilon \geq P_{\mathcal{E}_{N-1}}(E_N^{\mathcal{E}_{N-1}}) = P_{\mathcal{E}_N}(E_N^{\mathcal{E}_N}) + P_{\mathcal{E}_N}(E_{N+1}^{\mathcal{E}_N}) \quad (18)$$

This is true because  $P_{\mathcal{E}_{N-1}}(E_j^{\mathcal{E}_{N-1}}) = P_{\mathcal{E}_N}(E_j^{\mathcal{E}_N})$  for all  $j \in \{1, \dots, N-1\}$  by construction: once we observe  $x_j$ , we do not change the probability we assign to  $\{\omega \in \Omega : X(\omega) = x_j\}$  in future iterations. From (18) we have that  $P_{\mathcal{E}_N}(E_N^{\mathcal{E}_N}) < \epsilon$ . But then we can conclude that  $|P_{\mathcal{E}_{N-1}}(E_N^{\mathcal{E}_N}) - P_{\mathcal{E}_N}(E_N^{\mathcal{E}_N})| < \epsilon$ , and it will stay smaller than  $\epsilon$ , for all  $n \geq N$ . This shows that  $|P_{\mathcal{E}_{n-1}}(E_n^{\mathcal{E}_n}) - P_{\mathcal{E}_n}(E_n^{\mathcal{E}_n})|$  goes to 0 as  $n \rightarrow \infty$ , and hence concludes the proof.  $\square$

*Proof of Proposition 5.2.* If  $\emptyset \in \tilde{\mathcal{E}}$ , this means that  $\cup_{E_j \in \tilde{\mathcal{E}} \setminus \{\emptyset\}} E_j = \Omega$ , which implies that  $\cup_j x_j = \mathcal{X}$ , where  $\{x_j\}_j$  is the collection of data points we gathered. Then, since we observed all the elements of  $\mathcal{X}$ , and given the procedure in Sections 3 and 4 to refine the partition, it is immediate to see that  $\tilde{\mathcal{E}}$  cannot be further refined.  $\square$

*Proof of Proposition 5.3.* Notice that  $k, \ell < \infty$  because we assumed  $\#\mathcal{X}, \#\mathcal{Y} < \infty$ . Then,  $k < \ell$  follows immediately from our assumption and  $\#\mathcal{X} < \#\mathcal{Y}$ .  $\square$

*Proof of Proposition 5.4.* We first point out that  $\tilde{\mathcal{E}} = \tilde{\mathcal{E}}'$ . This because, no matter the order in which we collect data points  $x_j \in \mathcal{X}$ , in the limit we collect all of them. So if  $\tilde{\mathcal{E}}$  is finer than  $\tilde{\mathcal{E}}'$ , this means that there exists an  $\omega$  that is mapped by  $X$  into two different values, a contradiction. If instead  $\tilde{\mathcal{E}}$  is coarser than  $\tilde{\mathcal{E}}'$ , this means that  $\tilde{\mathcal{E}}$  can be further refined, which contradicts Proposition 5.2.

Then, the claim follows by the uniqueness of the limit of a sequence.  $\square$

*Proof of Proposition 6.1.* Recall that  $d_H(\mathcal{P}_{\mathcal{E}_n}, \mathcal{P}_{\tilde{\mathcal{E}}})$  goes to 0 as  $n \rightarrow \infty$  by construction, since every element in the finite limit set  $\mathcal{P}_{\tilde{\mathcal{E}}}$  is a limit of an element in set  $\mathcal{P}_{\mathcal{E}_n}$ . Then, after realizing that  $d_H(\mathcal{P}_{\mathcal{E}_n}, \mathcal{P}_{\tilde{\mathcal{E}}}) = d_H(\mathcal{P}_{\mathcal{E}_n}^{\text{co}}, \mathcal{P}_{\tilde{\mathcal{E}}}^{\text{co}})$ , the claim follows. This realization comes from the fact that, by construction, the ‘‘boundaries’’ of  $\mathcal{P}_{\mathcal{E}_n}$  and those of  $\mathcal{P}_{\mathcal{E}_n}^{\text{co}}$  coincide, for all  $n$ . That is, upper and lower probabilities associated with  $\mathcal{P}_{\mathcal{E}_n}$  are equal to those associated with  $\mathcal{P}_{\mathcal{E}_n}^{\text{co}}$ , for all  $n$ . This is also true – again by construction – for the limit sets  $\mathcal{P}_{\tilde{\mathcal{E}}}$  and  $\mathcal{P}_{\tilde{\mathcal{E}}}^{\text{co}}$ . Then, since the Hausdorff metric measures the distance between the ‘‘boundary elements’’  $\underline{\mathcal{P}}_{\mathcal{E}_n}$  and  $\overline{\mathcal{P}}_{\mathcal{E}_n}$ , and the ‘‘boundary elements’’  $\underline{\mathcal{P}}_{\tilde{\mathcal{E}}}$  and  $\overline{\mathcal{P}}_{\tilde{\mathcal{E}}}$ , we immediately see how  $d_H(\mathcal{P}_{\mathcal{E}_n}, \mathcal{P}_{\tilde{\mathcal{E}}}) = d_H(\mathcal{P}_{\mathcal{E}_n}^{\text{co}}, \mathcal{P}_{\tilde{\mathcal{E}}}^{\text{co}})$ .  $\square$

---

<sup>2</sup>It cannot be equal to  $\epsilon$  because that would imply that  $E_{N+1}^{\mathcal{E}_N} = \emptyset$ , which would in turn imply that  $\mathcal{X}$  is finite, and this would contradict the assumption we made for **Case 2**.

*Proof of Proposition 7.1.* Fix any  $A \in \mathcal{F}$  and any  $n \in \mathbb{N}_0$ . Notice that

$$\underline{P}_{\mathcal{E}_{n+1}}(A) := \inf_{\substack{P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}^{\text{co}} \\ P_{\mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}}} P_{\mathcal{E}_{n+1}}(A) = \inf_{\substack{P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}^{\text{co}} \\ P_{\mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}}} P_{\mathcal{E}_{n+1}}(A).$$

Then, we have that

$$\begin{aligned} \underline{P}_{\mathcal{E}_{n+1}}(A) &= \inf_{\substack{P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}^{\text{co}} \\ P_{\mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}}} \sum_{E_j \in \mathcal{E}_{n+1}} P_{\mathcal{E}_n}(A | E_j) P_{\mathcal{E}_{n+1}}(E_j) \\ &\geq \sum_{E_j \in \mathcal{E}_{n+1}} \inf_{\substack{P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}^{\text{co}} \\ P_{\mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}}} [P_{\mathcal{E}_n}(A | E_j) P_{\mathcal{E}_{n+1}}(E_j)] \end{aligned} \quad (19)$$

$$\geq \sum_{E_j \in \mathcal{E}_{n+1}} \inf_{\substack{P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}^{\text{co}} \\ P_{\mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}}} P_{\mathcal{E}_n}(A | E_j) \inf_{\substack{P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}^{\text{co}} \\ P_{\mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}}} P_{\mathcal{E}_{n+1}}(E_j) \quad (20)$$

$$= \sum_{E_j \in \mathcal{E}_{n+1}} \left[ \inf_{\substack{P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}^{\text{co}} \\ P_{\mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}}} P_{\mathcal{E}_n}(A | E_j) \right] \underline{P}_{\mathcal{E}_{n+1}}(E_j) \quad (21)$$

$$= \sum_{E_j \in \mathcal{E}_{n+1}} \left[ \inf_{\substack{P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}^{\text{co}} \\ P_{\mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}}} P_{\mathcal{E}_n}(A | E_j) \right] Q(E_j) \quad (22)$$

$$= \sum_{E_j \in \mathcal{E}_{n+1}} \left[ \inf_{P_{\mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}} P_{\mathcal{E}_n}(A | E_j) \right] Q(E_j)$$

$$= \sum_{E_j \in \mathcal{E}_{n+1}} \underline{P}_{\mathcal{E}_n}^B(A | E_j) Q(E_j).$$

The inequality in (19) comes from the well known fact that the sum of the infima is at most equal to the infimum of the sum. The inequality in (20) comes from the fact that for differentiable functions, the product of the infima is at most equal to the infimum of the product. Equation (21) comes from the fact that

$$\inf_{\substack{P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}^{\text{co}} \\ P_{\mathcal{E}_n} \in \mathcal{P}_{\mathcal{E}_n}^{\text{co}}}} P_{\mathcal{E}_{n+1}}(E_j) = \inf_{P_{\mathcal{E}_{n+1}} \in \mathcal{P}_{\mathcal{E}_{n+1}}^{\text{co}}} P_{\mathcal{E}_{n+1}}(E_j) = \underline{P}_{\mathcal{E}_{n+1}}(E_j).$$

Finally, equation (22) comes from (8). A similar argument – together with the facts that the supremum of the sum is at most equal to the sum of the suprema, and that for differentiable functions, the supremum of the product is at most equal to the product of the suprema – gives us the stated upper bound for  $\overline{P}_{\mathcal{E}_{n+1}}(A)$ .  $\square$

*Proof of Corollary 7.1.1.* Immediate from Proposition 7.1 and the definitions of upper and lower probabilities.  $\square$

*Proof of Lemma 2.* Immediate from the inequalities in (12).  $\square$

*Proof of Proposition 7.2.* Pick any  $A \in \mathcal{F}$  and any  $n \in \mathbb{N}$ . Then, the following holds

$$\underline{P}_{\mathcal{E}_{n+1}}(A) \geq \sum_{E_j \in \mathcal{E}_{n+1}} \underline{P}_{\mathcal{E}_{n+1}}(A \cap E_j) \quad (23)$$

$$\begin{aligned} &= \sum_{E_j \in \mathcal{E}_{n+1}} \frac{\underline{P}_{\mathcal{E}_{n+1}}(A \cap E_j)}{\underline{P}_{\mathcal{E}_{n+1}}(E_j)} \underline{P}_{\mathcal{E}_{n+1}}(E_j) \\ &= \sum_{E_j \in \mathcal{E}_{n+1}} \frac{\underline{P}_{\mathcal{E}_n}(A \cap E_j)}{\underline{P}_{\mathcal{E}_n}(E_j)} \underline{P}_{\mathcal{E}_{n+1}}(E_j) \end{aligned} \quad (24)$$

$$= \sum_{E_j \in \mathcal{E}_{n+1}} \underline{P}_{\mathcal{E}_n}^G(A | E_j) \underline{P}_{\mathcal{E}_{n+1}}(E_j) \quad (25)$$

$$= \sum_{E_j \in \mathcal{E}_{n+1}} \underline{P}_{\mathcal{E}_n}^G(A | E_j) Q(E_j). \quad (26)$$

The inequality in (23) comes from equation (11). Equation (24) comes from our assumption. Equation (25) comes from the definition of geometric rule's lower conditional probability. Equation (26) comes from (8).

A similar argument gives us the stated upper bound for  $\overline{P}_{\mathcal{E}_{n+1}}(A)$ .  $\square$

*Proof of Proposition 8.1.* By Proposition 7.1, we have that  $\underline{P}_{\mathcal{E}_n}(A) \geq \sum_{E_j \in \mathcal{E}_n} \underline{P}_{\mathcal{E}_{n-1}}^B(A | E_j) Q(E_j)$ , so if

$$\sum_{E_j \in \mathcal{E}_n} \underline{P}_{\mathcal{E}_{n-1}}^B(A | E_j) Q(E_j) \geq \underline{P}_{\mathcal{E}_{n-1}}(A),$$

then  $\underline{P}_{\mathcal{E}_n}(A) \geq \underline{P}_{\mathcal{E}_{n-1}}(A)$ . A similar reasoning gives us that  $\sum_{E_j \in \mathcal{E}_n} \overline{P}_{\mathcal{E}_{n-1}}^B(A | E_j) Q(E_j) \leq \overline{P}_{\mathcal{E}_{n-1}}(A)$  implies  $\overline{P}_{\mathcal{E}_n}(A) \leq \overline{P}_{\mathcal{E}_{n-1}}(A)$ . In turn, we obtain the desired contraction.  $\square$

*Proof of Proposition 8.2.* By Proposition 7.1, we have that  $\underline{P}_{\mathcal{E}_n}(A) \geq \sum_{E_j \in \mathcal{E}_n} \underline{P}_{\mathcal{E}_{n-1}}^B(A | E_j) Q(E_j)$ , so if

$$\sum_{E_j \in \mathcal{E}_n} \underline{P}_{\mathcal{E}_{n-1}}^B(A | E_j) Q(E_j) > \overline{P}_{\mathcal{E}_{n-1}}(A),$$

then  $\underline{P}_{\mathcal{E}_n}(A) > \overline{P}_{\mathcal{E}_{n-1}}(A)$ . A similar reasoning gives us that  $\sum_{E_j \in \mathcal{E}_n} \overline{P}_{\mathcal{E}_{n-1}}^B(A | E_j) Q(E_j) < \underline{P}_{\mathcal{E}_{n-1}}(A)$  implies  $\overline{P}_{\mathcal{E}_n}(A) < \underline{P}_{\mathcal{E}_{n-1}}(A)$ . In turn, we obtain the desired sure loss.  $\square$

*Proof of Proposition 8.3.* Fix any  $n \in \mathbb{N}$  and consider some  $A \in \mathcal{F}$ . By the definitions of lower and upper probabilities, we have that  $P_{s, \mathcal{E}_n}(A'), P_{k, \mathcal{E}_n}(A') \in [\underline{P}_{\mathcal{E}_n}(A'), \overline{P}_{\mathcal{E}_n}(A')]$ , for all  $A' \in \mathcal{F}$ . Then, if our hypotheses hold, we have that, for the set  $A$  we have chosen,

$$\underline{P}_{\mathcal{E}_{n-1}}(A) \geq P_{s, \mathcal{E}_n}(A) \geq \underline{P}_{\mathcal{E}_n}(A)$$

and

$$\overline{P}_{\mathcal{E}_{n-1}}(A) \leq P_{k, \mathcal{E}_n}(A) \leq \overline{P}_{\mathcal{E}_n}(A).$$

This concludes the proof.  $\square$

*Proof of Proposition 8.4.* Fix any  $n \in \mathbb{N}$  and consider some  $A \in \mathcal{F}$ . By the definitions of lower and upper probabilities, we have that  $P_{s,\varepsilon_{n-1}}(A'), P_{k,\varepsilon_{n-1}}(A') \in [\underline{P}_{\varepsilon_{n-1}}(A'), \overline{P}_{\varepsilon_{n-1}}(A')]$ , for all  $A' \in \mathcal{F}$ . Then, if our hypotheses hold, we have that, for the set  $A$  we have chosen,

$$\underline{P}_{\varepsilon_n}(A) \geq P_{k,\varepsilon_{n-1}}(A) \geq \underline{P}_{\varepsilon_{n-1}}(A)$$

and

$$\overline{P}_{\varepsilon_n}(A) \leq P_{s,\varepsilon_{n-1}}(A) \leq \overline{P}_{\varepsilon_{n-1}}(A).$$

This concludes the proof. □

## REFERENCES

- [1] John Armstrong. The Banach space of totally finite signed measures. *Available online*, 2010.
- [2] James O. Berger. The robust Bayesian viewpoint. In Joseph B. Kadane, editor, *Robustness of Bayesian Analyses*. Amsterdam : North-Holland, 1984.
- [3] Michele Caprio and Sayan Mukherjee. Ergodic theorems in dynamic imprecise probability kinematics. *Available on ResearchGate*, 2021.
- [4] Simone Cerreia-Vioglio, Fabio Maccheroni, and Massimo Marinacci. Ergodic Theorems for Lower Probabilities. *Proceedings of the American Mathematical Society*, 144:3381–3396, 2015.
- [5] Persi Diaconis and Sandy L. Zabell. Updating subjective probability. *Journal of the American Statistical Association*, 77(380):822–830, 1982.
- [6] Ruobin Gong and Xiao-Li Meng. Judicious judgment meets unsettling updating: dilation, sure loss, and Simpson’s paradox. *Statistical Science*, 36(2):169–190, 2021.
- [7] Hidetomo Ichihashi and Hideo Tanaka. Jeffrey-like rules of conditioning for the dempster-shafer theory of evidence. *International Journal of Approximate Reasoning*, 3(2):143–156, 1989.
- [8] Richard C. Jeffrey. *Contributions to the Theory of Inductive Probability*. PhD Thesis, Princeton University, Dept. of Philosophy, 1957.
- [9] Richard C. Jeffrey. *The Logic of Decision*. Chicago : University of Chicago Press, 1965.
- [10] Richard C. Jeffrey. Probable Knowledge. In Imre Lakatos, editor, *The Problem of Inductive Logic*, volume 51 of *Studies in Logic and the Foundations of Mathematics*, pages 166 – 190. Elsevier, 1968.
- [11] David Lewis. Probabilities of conditionals and conditional probabilities. *The Philosophical Review*, 85(3):297–315, 1976.
- [12] Jianbing Ma, Weiru Lu, Didier Dubois, and Henri Prade. Bridging Jeffrey’s rule, AGM revision and Dempster conditioning in the theory of evidence. *International Journal on Artificial Intelligence Tools*, 20(4):691–720, 2011.
- [13] Sabina Marchetti and Alessandro Antonucci. Imaginary kinematics. In Amir Globerson and Ricardo Silva, editors, *UAI 2018: Proceedings of the Thirty-Fourth Conference on Uncertainty in Artificial Intelligence*, pages 104–113, Monterey, California, USA, 2018. AUAI Press.
- [14] Walter Rudin. *Real and complex analysis*. McGraw-Hill, New York, 3rd edition, 1987.
- [15] Glenn Shafer. Jeffrey’s Rule of Conditioning. *Philosophy of Science*, 48(3):337–362, 1981.
- [16] Damjan Škulj. Jeffrey’s conditioning rule in neighbourhood models. *International Journal of Approximate Reasoning*, 42(3):192–211, 2006.
- [17] Philippe Smets. Jeffrey’s rule of conditioning generalized to belief functions. In *Proceedings of the Ninth international conference on Uncertainty in Artificial Intelligence*, pages 500–505, 1993.
- [18] Peter Walley. *Statistical reasoning with imprecise probabilities*, volume 42 of *Monographs on Statistics and Applied Probability*. London : Chapman and Hall, 1991.
- [19] Larry A. Wasserman and Joseph B. Kadane. Bayes’ theorem for Choquet capacities. *The Annals of Statistics*, 18(3):1328–1339, 1990.

DEPARTMENT OF STATISTICAL SCIENCE, DUKE UNIVERSITY, 214 OLD CHEMISTRY, DURHAM, NC  
27708-0251

*Email address:* `michele.caprio@duke.edu`

*URL:* <https://mc6034.wixsite.com/caprio>

DEPARTMENT OF STATISTICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854

*Email address:* `ruobin.gong@rutgers.edu`

*URL:* <https://ruobingong.github.io/>