

Logical Foundations of Quantitative Equality (long version)

Francesco Dagnino*, Fabio Pasquali†

Abstract

Quantitative reasoning provides a flexible approach capable to deal with the uncertainty and imprecision that affects modern software systems due to their complexity. Its distinguishing feature is the use of *distances*, instead of equivalence relations, so that one can measure how much two objects are similar, rather than just saying whether they are equivalent or not. In this paper we aim at providing a solid logical ground to quantitative reasoning, using the categorical language of Lawvere’s hyperdoctrines. The key idea is to see distances as equality predicates in Linear Logic. Adding equality to Linear Logic, however, is not as trivial as it might appear. The linear version of the usual rules for equality, asserting that it is reflexive and substitutive, has the consequence that equality collapses to an equivalence relation, as it can be used an arbitrary number of times. To overcome this issue, we consider the extension of Linear Logic with *graded modalities* and use them to write a resource sensitive substitution rule that keeps equality quantitative. We introduce a deductive calculus for (Graded) Linear Logic with quantitative equality and the notion of Lipschitz doctrine to give semantics to it. Main examples are based on metric spaces and Lipschitz maps and on quantitative realisability.

1 Introduction

The increasing complexity and heterogeneity of software systems makes it very difficult to describe and reason about their behaviour, hence having formal guarantees on such systems is more and more challenging. In this complex and error-prone environment, fully reliable and precise information

*DIBRIS - Università di Genova, email: francesco.dagnino@dibris.unige.it

†DIMA - Università di Genova, email: pasquali@dima.unige.it

on the system is basically not feasible, hence, we have to accept a certain degree of noise and use formal tools capable to deal with it.

Quantitative reasoning provides a promising approach to cope with this situation. In very general terms, the basic idea is that one is interested in measuring *how much* a property holds, rather than just saying whether it holds or not. For instance, we could measure how far an element is from having a given property, using non-negative real numbers, or we could measure the probability that an element has such a property. This approach allows a much finer analysis and, indeed, it is increasingly used to study different aspects of programs, such as differential privacy [40, 6, 45, 7], denotational semantics [5, 16], quantitative algebras [34, 35], and program/behavioural metrics [17, 43, 11, 21, 15, 14].

At the core of quantitative reasoning is the concept of *distance*, which is the quantitative counterpart of an equivalence relation. A distance measures how much two objects are similar, rather than just saying whether they are equivalent or not. Furthermore, many typical properties of equivalence relations can be extended to distances. The most important is the *substitutivity* property, stating that we can always replace an object by an equivalent one. This is particularly important when reasoning about programs, as substitutivity enables a form of *compositionality*: if we have a program t , which contains a component u , and we know that u is equivalent to another component u' , then, replacing u by u' , we obtain a new program t' which is equivalent to t , that is, the substitution does not affect the behaviour of the original program. Moving to the quantitative setting, this property becomes a *metric preservation* condition, which can be expressed as *Lipschitz continuity*, meaning that the distance after the substitution is proportional to the distance between the substituted objects. In other words, we want the substitution not to affect too much the context. That is, when substituting in t the component u by u' , the behaviour of the resulting program t' should not be too far from the original one.

Linear Logic [22], treating formulas as consumable resources, enables a quantitative interpretation of them, thus providing a natural setting for quantitative reasoning. However, in order to allow the formal comparison of terms, such a logic needs to be enriched by a *quantitative equality* predicate, that is, a substitutive binary predicate which can be interpreted as a distance. In this paper, we develop syntax and categorical semantics of a minimal linear logic with quantitative equality, thus obtaining a logical framework with solid foundations for quantitative reasoning.

A first naive attempt to introduce such an equality in Linear Logic could be to take the linear version of equality rules in First Order Logic. They

force equality to be reflexive and substitutive, that is, equality has to satisfy the following entailments:

$$\top \vdash t = t \quad \phi[t/x] \ \& \ t = u \vdash \phi[u/x]$$

One can take the linear version of these properties by replacing $\&$ by \otimes and \top by $\mathbf{1}$. That is, one can consider Linear Logic extended by a binary predicate $=$ that satisfies

$$\mathbf{1} \vdash t = t \quad \phi[t/x] \ \otimes \ t = u \vdash \phi[u/x]$$

However smooth, this approach raises the unexpected issue that the equality predicate turns out to be non-quantitative, in the sense that it can be used an arbitrary number of times. Indeed, as already observed by Dozen [18], the following entailments are derivable from the two properties above:

$$t = u \vdash \mathbf{1} \quad t = u \vdash (t = u) \ \otimes \ (t = u)$$

In other words, weakening and contraction, which in general are banned from Linear Logic, are instead admissible for the equality predicates. This is undesirable in a linear setting, where the fundamental purpose is to control the use of formulas, by avoiding uncontrolled weakening and contraction. But it is even worse for our goal: the distance interpreting equality trivialises collapsing to an equivalence relation. For instance, for a distance $d : X \times X \rightarrow [0, \infty]$ interpreting equality, the entailment $x = y \vdash x = y \ \otimes \ x = y$ becomes $d(x, y) \geq d(x, y) + d(x, y)$, which forces $d(x, y)$ to be either 0 or ∞ , hence, de facto, an equivalence relation.

In this paper, we propose a novel approach to equality in the linear setting, able to keep equality quantitative, developing both a deductive calculus and its categorical semantics. As a consequence, equality can be interpreted as a non-trivial distance and the resulting logic provides a formal framework for quantitative reasoning.

The starting observation is that the usual substitutive property is not resource sensitive, as it does not take into account resources needed to perform the substitution: the equality $t = u$ has to be substitutive *for any* formula ϕ , while the “cost” of substituting t by u may be different for different formulas (for instance, it may depend on the number of occurrences of t in the formula). Hence, we need to rephrase the substitutive property to explicitly consider such resources. To this end, we work in (fragments of) Linear Logic enriched by *graded modalities* [8]. Roughly, one adds to the calculus a family of unary connectives $!_r$, where r is taken from a structure of resources, which make explicit “how much” a formula can be used.

Using graded modalities, the resource sensitive variant of the substitutive property becomes

$$\phi[t/x] \otimes !_r(t = u) \vdash \phi[u/x]$$

where r depends on ϕ (in a way that will be described in the paper) and encodes the amount of resources needed to derive the substitution. In this way, the equality predicate cannot be duplicated, hence it admits a quantitative interpretation. Furthermore, from this axiom we can derive the entailment $!_r u = v \vdash t[u/x] = t[v/x]$, where r depends on t , which can be seen as a syntactic form of Lipschitz continuity of the term t .

The main mathematical tool that we will use throughout the paper is that of *Lawvere's doctrines*, which provide an elegant categorical/algebraic understanding of logic. The key feature of this approach, that goes back to Lawvere [28], is that theories are written as functors, named doctrines, whose domain category models contexts and terms, and the functor maps each object to a poset that models formulas on that object ordered by logical entailment. In this framework, logical phenomena can be explained algebraically, abstracting away from details and focusing only on minimal essential features. For instance, the non-linear nature of the standard equality in the linear setting has a neat algebraic explanation: such an equality is a left adjoint, as pioneered by Lawvere [28, 29], and, as we will show, predicates defined by left adjoints are always replicable, even in the linear setting. This shows that a quantitative equality cannot be a left adjoint, however our notion of equality shares other fundamental structural properties with the standard one described by Emmenegger et al. [19].

The main contributions of the paper are the following:

- we introduce R -graded doctrines, that is, doctrines modelling the $(\otimes, \mathbf{1})$ -fragment of Linear Logic enriched by R -graded modalities, where R is an ordered semiring of resources;
- we introduce R -Lipschitz doctrines, that is, R -graded doctrines with quantitative equality;
- we present a deductive calculus for quantitative equality, based on that by Breuvert and Pagani [8], describing its sound and complete categorical semantics in R -Lipschitz doctrines;
- we prove that R -Lipschitz doctrines arise as coalgebras for a 2-comonad on the 2-category of R -graded doctrines, defining a universal construction which yields a R -Lipschitz doctrine from an R -graded one, and we use this construction to generate semantics for the calculus;

- we relate our quantitative equality to the usual non-quantitative one;
- we extend the results above to richer fragments of Linear Logic.

Outline Section 2 recalls known facts on equality in non-linear doctrines. These will be rephrased in the linear setting in Section 3, where we also discuss the non-quantitative nature of standard equality defined by left adjoints. Main contributions are in Section 4. In Section 4.1 we introduce R -graded doctrines and R -Lipschitz doctrines. We present a minimal calculus for quantitative equality in Section 4.2 and its categorical semantics in R -Lipschitz doctrines in Section 4.3. In Section 4.4 we show the coalgebraic nature of R -Lipschitz doctrines and the universal construction that generates an R -Lipschitz doctrine out of any R -graded doctrine. Section 5 relates quantitative equality with equality as left adjoint. In Section 6 we extend previous results to richer fragments of Linear Logic up to full LL. Section 7 compares our approach with quantitative equational theories by Mardare et al. [34, 35], providing many examples of theories in our calculus. We conclude in Section 8 discussing related work and future developments. Appendix A collects the proofs omitted in the paper.

2 Background on categorical logic

Denote by \mathcal{Pos} the category of posets and monotone functions. A left adjoint to a monotone function $g : K \rightarrow H$ is a monotone function $f : H \rightarrow K$ such that for every x in K and y in H , both $y \leq gf(y)$ and $fg(x) \leq x$ hold. Equivalently $y \leq g(x)$ if and only if $f(y) \leq x$.

A *doctrine* is a pair $\langle \mathcal{C}, P \rangle$ where \mathcal{C} is a category with finite products and $P : \mathcal{C}^{\text{op}} \rightarrow \mathcal{Pos}$ is a functor. Traditionally the category \mathcal{C} is named the *base* of the doctrine. For X an object in \mathcal{C} the poset $P(X)$ is called *fibre over X* . For $f : X \rightarrow Y$ an arrow in \mathcal{C} , the monotone function $P_f : P(Y) \rightarrow P(X)$ is called *reindexing along f* . We will often refer to a doctrine $\langle \mathcal{C}, P \rangle$ using only the functor $P : \mathcal{C}^{\text{op}} \rightarrow \mathcal{Pos}$.

A doctrine $P : \mathcal{C}^{\text{op}} \rightarrow \mathcal{Pos}$ is *primary* if all fibres have finite meets preserved by reindexing. A primary doctrine $P : \mathcal{C}^{\text{op}} \rightarrow \mathcal{Pos}$ is a *first order doctrine* if P factors through the category of Heyting algebras and their homomorphisms and there are left and right adjoints along any projection $\pi : A \times B \rightarrow B$ in \mathcal{C} and these adjoints are natural in B .

The naturality of left and right adjoints is often called Beck-Chevalley condition, but we do not use this terminology here to avoid confusion with the Beck-Chevalley condition of equality predicates (cf. Definition 2.2).

There is a large variety of examples of primary and first order doctrines [25]. Here we consider the following ones to exemplify definitions throughout the paper.

- Examples 2.1.** 1. A many-sorted signature L consists of a collection of sorts σ, τ , a collection of function symbols f, g and a collection of predicate symbols p, q . Consider a theory \mathcal{T} in the $(\&, \top)$ -fragment of First Order Logic over a many-sorted signature L . The category of contexts \mathbf{Ctx}_L has contexts $\vec{\sigma} = \langle x_1 : \sigma_1, \dots, x_n : \sigma_n \rangle$ as objects and lists of terms as arrows, i.e. an arrow from $\vec{\sigma}$ to $\langle y_1 : \tau_1, \dots, y_k : \tau_k \rangle$ is a list of terms $\langle t_1, \dots, t_k \rangle$ such that each t_i has sort τ_i in the context $\vec{\sigma}$. As detailed by Maietti and Rosolini [32, Example 2.2] the category \mathbf{Ctx}_L has finite products and is the base of the *syntactic doctrine* $\mathbf{Prp}_{\mathcal{T}}$ mapping each context $\vec{\sigma}$ to the Lindenbaum-Tarski algebra of well-formed formulas over L in the $(\&, \top)$ -fragment with free variables in $\vec{\sigma}$. More precisely, the elements of the fibre over $\vec{\sigma}$ are equivalence classes $[\phi]$ of formulas modulo logical equivalence in the theory \mathcal{T} , ordered by entailment in \mathcal{T} , that is, $[\phi] \leq [\psi]$ iff $\phi \vdash_{\vec{\sigma}} \psi$ is provable in \mathcal{T} . The doctrine $\mathbf{Prp}_{\mathcal{T}}$ is primary where conjunctions give finite meets. If instead of the $(\&, \top)$ -fragment we consider full First Order Logic, then $\mathbf{Prp}_{\mathcal{T}}$ is a first order doctrine, where conjunctions, disjunctions and implications give the corresponding operations of the Heyting algebras, while left and right adjoints to projections are given by existential and universal quantification, respectively.
2. Suppose H is a poset, the functor $\mathcal{P}_H : \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Pos}$ sends a set A to H^A , the set of functions from A to H ordered pointwise. For a function $f : X \rightarrow A$ and α in $\mathcal{P}_H(A)$ the function $\mathcal{P}_H(f)(\alpha)$ is the composite αf in $\mathcal{P}_H(X)$. The doctrine \mathcal{P}_H is primary when H is a meet-semilattice and is first order when H is a complete Heyting algebra. For $H = \{0, 1\}$ the doctrine \mathcal{P}_H is the contravariant powerset functor that will be denoted simply by \mathcal{P} .
3. Suppose $A = (|A|, \cdot, K, S)$ is a (partial) combinatory algebra. The doctrine $\mathcal{R}_A : \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Pos}$ maps each set X to the poset reflection of the preorder $\mathcal{P}(|A|)^X$ where for $\alpha, \beta : X \rightarrow \mathcal{P}(|A|)$, $\alpha \leq \beta$ whenever there is a in $|A|$ such that, for every x in X and b in $\alpha(x)$, the application $a.b$ is defined and belongs to $\beta(x)$. One says that a realises $\alpha \leq \beta$. The action of \mathcal{R}_A on functions is given by pre-composition. \mathcal{R}_A is first order [44].

Doctrines are the objects of the 2-category \mathbf{Dtn} whose 1-arrows are pairs $\langle F, f \rangle : \langle \mathcal{C}, P \rangle \rightarrow \langle \mathcal{D}, Q \rangle$ where the first component $F : \mathcal{C} \rightarrow \mathcal{D}$ is a finite product preserving functor and the second component $f : P \rightarrow QF^{\text{op}}$ is a natural transformation; the 2-arrows $\theta : \langle F, f \rangle \Rightarrow \langle F', f' \rangle$ are such that $\theta : F \rightarrow F'$ is a natural transformation where $f_X \leq_{FX} Q_{\theta_X} \circ f'_X$ for each object X in \mathcal{C} . The composition of 1-arrows $\langle G, g \rangle : \langle \mathcal{B}, M \rangle \rightarrow \langle \mathcal{C}, P \rangle$ and $\langle F, f \rangle : \langle \mathcal{C}, P \rangle \rightarrow \langle \mathcal{D}, Q \rangle$ is (essentially) pairwise $\langle FG, (fG^{\text{op}}) \circ g \rangle : \langle \mathcal{B}, M \rangle \rightarrow \langle \mathcal{D}, Q \rangle$. Composition of 2-arrows $\theta : \langle F, f \rangle \Rightarrow \langle F', f' \rangle$ and $\zeta : \langle F', f' \rangle \Rightarrow \langle F'', f'' \rangle$ is the 2-arrow $(\zeta_X \circ \theta_X)_{X \in \mathcal{C}_0} : \langle F, f \rangle \Rightarrow \langle F'', f'' \rangle$, which is well-defined since, for any object X in \mathcal{C} ,

$$f_X \leq_{FX} Q_{\theta_X} \circ f'_X \leq_{FX} Q_{\theta_X} \circ Q_{\zeta_X} \circ f''_X \leq_{FX} Q_{((\zeta \circ \theta)_X)} \circ f''_X$$

Primary doctrines are the objects of the 2-full 2-subcategory \mathbf{PD} of \mathbf{Dtn} , where a 1-arrow from P to Q in \mathbf{PD} is a 1-arrow $\langle F, f \rangle$ in \mathbf{Dtn} such that each component of f preserves finite meets. First order doctrines are the objects of the 2-full 2-subcategory \mathbf{FOD} of \mathbf{PD} , where a 1-arrow in \mathbf{FOD} is a 1-arrow in \mathbf{PD} that preserves the Heyting algebra structure as well as left and right adjoints. Hence we have the following inclusions

$$\mathbf{FOD} \hookrightarrow \mathbf{PD} \hookrightarrow \mathbf{Dtn}$$

The 2-categorical structure of doctrines has a relevant logical meaning: objects correspond to theories, 1-arrows to models/translations between theories and 2-arrows to homomorphisms of models/translations. Under this interpretation, note that the hom-category $\mathbf{Dtn}(P, Q)$ is the category of models of P into Q . For example, given a theory \mathcal{T} in First Order Logic over a signature L , a categorical semantics of \mathcal{T} in a first order doctrine P is an object of $\mathbf{FOD}(Prp_{\mathcal{T}}, P)$.

A categorical description of equality in terms of left adjoints goes back to the work of Lawvere [29]. More recently, Maietti and Rosolini [31] identify primary doctrines as the essential framework where equality can be studied. Primary doctrines with equality are called elementary by Maietti and Rosolini [31] and can be defined as follows.

Definition 2.2. A primary doctrine $P : \mathcal{C}^{\text{op}} \rightarrow \mathcal{P}os$ is *elementary* if for all X and A in \mathcal{C} the map $P_{\text{id}_X \times \Delta_A} : P(X \times A \times A) \rightarrow P(X \times A)$ has a left adjoint $\mathcal{E}_{\text{id}_X \times \Delta_A} : P(X \times A) \rightarrow P(X \times A \times A)$ such that

1. $P_{f \times \text{id}_A \times \text{id}_A} \mathcal{E}_{\text{id}_X \times \Delta_A} = \mathcal{E}_{\text{id}_Y \times \Delta_A} P_{f \times \text{id}_A}$, for every arrow $f : Y \rightarrow X$,
and

2. $\mathcal{E}_{\text{id}_X \times \Delta_A}(\alpha) \wedge \beta = \mathcal{E}_{\text{id}_X \times \Delta_A}(\alpha \wedge P_{\text{id}_X \times \Delta_A}(\beta))$, for every $\alpha \in P(X \times A)$ and $\beta \in P(X \times A \times A)$.

A first order elementary doctrine is a *hyperdoctrine*.

This definition of hyperdoctrine coincides with the one in [39]. Condition 1) is often called Beck-Chevalley condition, while condition 2) is often called Frobenius reciprocity.

Examples 2.3. 1. The syntactic doctrine $Prp_{\mathcal{T}}$ associated to a theory \mathcal{T} in the $(\&, \top, =)$ -fragment of First Order Logic as in Examples 2.1(1), is elementary. If the full First Order Logic with equality is considered, $Prp_{\mathcal{T}}$ is a hyperdoctrine.

2. The first order doctrine \mathcal{P}_H of Example 2.1(2) is elementary where for a set X the function $\mathcal{E}_{\Delta_X}(\top_X) : X \times X \rightarrow H$ maps (x, x') to \top if $x = x'$ and to \perp otherwise.
3. The first order doctrine \mathcal{R}_A of Example 2.1(3) is elementary where $\mathcal{E}_{\Delta_X}(\top_X)$ maps (x, x') to $|A|$ if $x = x'$ and to \emptyset otherwise [44].

Elementary doctrines are the objects of the 2-category **ED** where a 1-arrow $\langle F, f \rangle : P \rightarrow Q$ is a 1-arrow in **PD** such that f commutes with left adjoints along arrows of the form $\text{id}_X \times \Delta_A$, while 2-arrows are those of **PD**. The 2-full 2-subcategory of **ED** on first order doctrines, *i.e.* on hyperdoctrines, is **HD**.

As studied in [19] equality is coalgebraic in the sense that the obvious inclusion of **ED** into **PD** has a right 2-adjoint and **ED** is isomorphic to the 2-category of coalgebras of the induced 2-comonad. This 2-adjunction lifts to full first order doctrines as depicted below

$$\begin{array}{ccc}
 \mathbf{ED} & \begin{array}{c} \longleftarrow \\ \top \\ \longrightarrow \end{array} & \mathbf{PD} \\
 \uparrow & & \uparrow \\
 \mathbf{HD} & \begin{array}{c} \longleftarrow \\ \top \\ \longrightarrow \end{array} & \mathbf{FOD}
 \end{array} \tag{1}$$

A straightforward linearisation of all these results is presented in the next section.

3 Equality in primary linear doctrines

Primary linear doctrines are the linear version of primary doctrines: they model the $(\otimes, \mathbf{1})$ -fragment of Linear Logic instead of the $(\&, \top)$ -fragment of

First Order Logic. They capture the minimal fragment of linear hyperdoctrines, by Seely [41], needed to study equality in the linear setting.

Definition 3.1. A *primary linear doctrine* is a triple $\langle P, *, \kappa \rangle$ where $P : \mathcal{C}^{\text{op}} \rightarrow \mathcal{Pos}$ is a doctrine and $* : P \times P \rightarrow P$ and $\kappa : 1_{\mathcal{C}} \rightarrow P$ are natural transformations such that, for all objects X in \mathcal{C} , $\langle PX, *_X, \kappa_X \rangle$ is a commutative monoid.

Spelling out the definition, a primary linear doctrine is determined by a doctrine $P : \mathcal{C}^{\text{op}} \rightarrow \mathcal{Pos}$ and, for each object X in \mathcal{C} , a monotone function $*_X : P(X) \times P(X) \rightarrow P(X)$ and an element $\kappa_X \in P(X)$ such that $*_X$ is associative and commutative and κ_X is the neutral element for $*_X$ and, for all arrows $f : X \rightarrow Y$ in \mathcal{C} and all α, β in $P(Y)$, we have $P_f(\alpha) *_X P_f(\beta) = P_f(\alpha *_Y \beta)$ and $\kappa_X = P_f(\kappa_Y)$. We shall often omit subscripts from $*$ and κ when these are obvious.

Examples 3.2. 1. Consider a theory \mathcal{T} in the $(\otimes, \mathbf{1})$ -fragment of Linear Logic over a signature L as in [22]. The syntactic doctrine $Prp_{\mathcal{T}}$ built as in Examples 2.1(1) is such that $\langle Prp_{\mathcal{T}}, \otimes, \mathbf{1} \rangle$ is primary linear.

2. Suppose $\langle H, *, \kappa \rangle$ is a monoidal poset. The doctrine $\mathcal{P}_H : \mathbf{Set}^{\text{op}} \rightarrow \mathcal{Pos}$ as in Examples 2.1(2) is part of a primary linear doctrine $\langle \mathcal{P}_H, *, \kappa \rangle$ where for α, β in $\mathcal{P}_H(A) = H^A$ the tensor $\alpha *_A \beta$ is the function that maps a to $\alpha(a) *_A \beta(a)$. Hence the unit is the constant function mapping a to κ .

3. Suppose $A = (|A|, \cdot, B, C, I)$ is a BCI algebra (see [3]). Let \mathcal{R}_A be as in Examples 2.1(3). Each $\mathcal{R}_A(X)$ is a poset as I and B combinators realise reflexivity and transitivity of \leq , respectively. Given $\alpha, \beta : X \rightarrow \mathcal{P}(|A|)$ in $\mathcal{R}_A(X)$, define $\alpha * \beta$ as the function that maps x to $\{\mathbf{P}ab \in |A| \mid a \in \alpha(x) \text{ and } b \in \beta(x)\}$, where \mathbf{P} is the BCI pairing (i.e. $\mathbf{P}ab = \lambda x.xab$ see also [24]), and κ_X as the function that maps x to $\{I\}$. Then $\langle \mathcal{R}_A, *, \kappa \rangle$ is a primary linear doctrine.

The following definition generalises to the framework of primary linear doctrines the definition of elementary doctrines.

Definition 3.3. An *elementary primary linear doctrine* is a primary linear doctrine $\langle P, *, \kappa \rangle$ such that, for every X and every A in the base of P , the map $P_{\text{id}_X \times \Delta_A} : P(X \times A \times A) \rightarrow P(X \times A)$ has a left adjoint $\mathcal{E}_{\text{id}_X \times \Delta_A} : P(X \times A) \rightarrow P(X \times A \times A)$ satisfying the Beck-Chevalley condition and the linear Frobenius Reciprocity, i.e. $\mathcal{E}_{\text{id}_X \times \Delta_A}(\alpha) *_\beta = \mathcal{E}_{\text{id}_X \times \Delta_A}(\alpha *_\beta)$, for all α in $P(X \times A)$ and β in $P(X \times A \times A)$.

The element $\mathcal{E}_{\Delta_A}(\kappa_A)$ in $P(A \times A)$ will be called *equality predicate* over A . When $*$ is a meet operation, Definition 3.3 reduces to Definition 2.2 so every elementary doctrine is an elementary primary linear doctrine. More examples will follow after the next characterisation, that rephrases in the linear setting a similar result from [19].

Proposition 3.4. *Let $\langle P, *, \kappa \rangle$ be a primary linear doctrine. The following are equivalent:*

(i) $\langle P, *, \kappa \rangle$ is elementary

(ii) for every object A there is δ_A in $P(A \times A)$ such that for every object X and α in $P(A)$ the following hold

$$(a) \quad \kappa \leq P_{\Delta_A}(\delta_A)$$

$$(b) \quad \delta_{A \times X} = P_{\langle \pi_1, \pi_3 \rangle}(\delta_A) * P_{\langle \pi_2, \pi_4 \rangle}(\delta_X)$$

$$(c) \quad P_{\pi_1}(\alpha) * \delta_A \leq P_{\pi_2}(\alpha)$$

(iii) for every object A there is δ_A in $P(A \times A)$ such that for every object X and α in $P(X \times A)$ the following hold

$$(a) \quad \kappa \leq P_{\Delta_A}(\delta_A)$$

$$(b) \quad P_{\langle \pi_1, \pi_2 \rangle}(\alpha) * P_{\langle \pi_2, \pi_3 \rangle}(\delta_A) \leq P_{\langle \pi_1, \pi_3 \rangle}(\alpha)$$

Examples 3.5. 1. Let the $(\otimes, \mathbf{1}, =)$ -fragment of Linear Logic be the $(\otimes, \mathbf{1})$ -fragment extended with the rules

$$[\text{R}] \frac{}{\vdash t = t} \qquad [\text{SUBST}] \frac{\Gamma \vdash \phi[t/x] \quad \Delta \vdash t = u}{\Gamma, \Delta \vdash \phi[u/x]}$$

This is the calculus for equality in non-modal Linear Logic considered in [18]. Given a theory \mathcal{T} in this fragment, the syntactic primary linear doctrine $\langle \text{Prp}\mathcal{T}, \otimes, \mathbf{1} \rangle$ is elementary by virtue of Proposition 3.4(iii).

2. If H is a monoidal poset with an absorbing bottom element \mathbf{o} then $\langle \mathcal{P}_H, *, \kappa \rangle$ is elementary where $\delta_X : X \times X \rightarrow H$ sends (x, x') to κ when $x = x'$ and to \mathbf{o} otherwise. It is easy to check that δ_X satisfies conditions in Proposition 3.4(ii).
3. For a BCI algebra A , the primary linear doctrine \mathcal{R}_A defined in Example 3.2(3) is elementary in virtue of Proposition 3.4(ii), where $\delta_X : X \times X \rightarrow \mathcal{P}(|A|)$ maps (x, x') to $|A|$ when $x = x'$ and to \emptyset otherwise.

Equality predicates in elementary doctrines are proved to be equivalence relations. The linear counterpart of an equivalence relation is a distance.

Definition 3.6. Let $\langle P, *, \kappa \rangle$ be a primary linear doctrine and A an object in the base of P . A P -distance on A is an element $\rho \in P(A \times A)$ satisfying the following properties:

1. $\kappa \leq P_{\Delta_A}(\rho)$ (reflexivity)
2. $\rho \leq P_{\text{sw}}(\rho)$ (symmetry)
3. $P_{\langle \pi_1, \pi_2 \rangle}(\rho) * P_{\langle \pi_2, \pi_3 \rangle}(\rho) \leq P_{\langle \pi_1, \pi_3 \rangle}(\rho)$ (transitivity)

where sw is the isomorphism $\langle \pi_2, \pi_1 \rangle : A \times A \rightarrow A \times A$.

Proposition 3.7. *Equality predicates in an elementary primary linear doctrine $\langle P, *, \kappa \rangle$ are P -distances.*

Primary linear doctrines are the objects of the 2-category **LD** where a 1-arrow $\langle F, f \rangle : \langle P, *^P, \kappa^P \rangle \rightarrow \langle Q, *^Q, \kappa^Q \rangle$ is a 1-arrow $\langle F, f \rangle : P \rightarrow Q$ in **Dtn** such that $f : P \rightarrow Q^{F^{\text{op}}}$ is a natural monoid homomorphism, that is, for every A in the base of P , the map $f_A : P(A) \rightarrow Q(FA)$ satisfies $f_A(\alpha) *^Q f_A(\beta) = f_A(\alpha *^P \beta)$, for all α, β in $P(A)$, and $\kappa_{FA}^Q = f_A(\kappa_A^P)$. Finally, 2-arrows are those of **Dtn**.

3.1 Equality as left adjoint is not quantitative

Given a primary linear doctrine $\langle P, *, \kappa \rangle$ an element α in $P(X)$ is *affine* if $\alpha \leq \kappa$, while α is *replicable* if $\alpha \leq \alpha * \alpha$. On affine and replicable elements the monoidal operation $*$ behaves as a meet operation, hence, they can be deleted and duplicated in a proof. As mentioned in the introduction, equality as a left adjoint is affine and replicable as shown by the following propositions.

Proposition 3.8. *Let $\langle P, *, \kappa \rangle$ be a primary linear doctrine and $f : X \rightarrow Y$ an arrow in the base of P such that P_f has a left adjoint \mathcal{A}_f . The element $\mathcal{A}_f(\kappa_X)$ is affine and replicable.*

Proof. $\mathcal{A}_f(\kappa_X) = \mathcal{A}_f P_f(\kappa_Y) \leq \kappa_Y$ proves affineness. Replicability follows from $\kappa_X = \kappa_X * \kappa_X \leq P_f(\mathcal{A}_f \kappa_X) * P_f(\mathcal{A}_f \kappa_X) = P_f(\mathcal{A}_f(\kappa_X) * \mathcal{A}_f(\kappa_X))$ and the adjunction between \mathcal{A}_f and P_f . \square

Recall that equality predicates have the form $\mathcal{A}_{\Delta_A}(\kappa_A)$, hence the following is an immediate corollary of Proposition 3.8.

Corollary 3.9. *Let $\langle P, *, \kappa \rangle$ be an elementary primary linear doctrine. Equality predicates are affine and replicable.*

Corollary 3.9 instantiated to the syntactic doctrine over the $(\otimes, \mathbf{1}, =)$ -fragment of Linear Logic (see Examples 3.5(1)) ensures that both $t = u \vdash \mathbf{1}$ and $t = u \vdash (t = u) \otimes (t = u)$ are derivable.

Remark 3.10. Suppose $\langle P, *, \kappa \rangle$ is an elementary primary linear doctrine. For every $f : X \rightarrow A$ in \mathcal{C} it holds $\kappa_X = P_f(\kappa_A) \leq P_f(P_{\Delta_A}(\delta_A)) = P_{\Delta_X}(P_{f \times f}(\delta_A))$. Since equality predicates are left adjoints, this implies $\delta_X \leq P_{f \times f}(\delta_A)$. Syntactically, it means that, for every term t , the entailment $x = y \vdash t(x) = t(y)$ is derivable. Aiming at taking resources into account, this entailment is less innocent as it might appear, as it suffices alone to prove replicability of equality: $P_{\Delta_A \times \Delta_A}(\delta_{A \times A}) = P_{\Delta_A \times \Delta_A}(P_{\langle \pi_1, \pi_3 \rangle} \delta_A * P_{\langle \pi_2, \pi_4 \rangle}(\delta_A)) = \delta_A * \delta_A$, whence from $\delta_A \leq P_{\Delta_A \times \Delta_A}(\delta_{A \times A})$ replicability follows.

In the next section we introduce quantitative equality working in the more general setting of graded doctrines, namely, primary linear doctrines endowed with graded modalities.

4 Quantitative equality via graded modalities

Graded modalities indexed by a semiring R are used in many linear calculi and type theories to enable resource sensitive reasoning [40, 9, 20, 37, 1, 14, 36]. In this section, we introduce graded modalities in the framework of primary linear doctrines and use them to formulate a quantitative variant of elementary primary linear doctrines. We will call them Lipschitz doctrines, motivating the terminology after Corollary 4.8. Then, we introduce a calculus for quantitative equality and its categorical semantics, proved to be sound. We will show that, like in the non-linear case [19], Lipschitz doctrines arise as coalgebras for a 2-comonad on graded doctrines, providing a universal construction to generate Lipschitz doctrines from graded ones.

4.1 Graded doctrines and Lipschitz doctrines

A *resource semiring* is a tuple $R = \langle |R|, \preceq, +, \cdot, 0, 1 \rangle$ such that $\langle |R|, +, \cdot, 0, 1 \rangle$ is a semiring and \preceq is a partial order on $|R|$ making $+$ and \cdot monotone. More explicitly, we have that

- $\langle |R|, +, 0 \rangle$ is a commutative monoid,

- $\langle |R|, \cdot, 1 \rangle$ is a monoid, and
- for all $r, s, q \in |R|$ the following hold:

$$\begin{aligned} r \cdot (s + q) &= r \cdot s + r \cdot q & (s + q) \cdot r &= s \cdot r + q \cdot r \\ r \cdot 0 &= 0 & 0 \cdot r &= 0 \end{aligned}$$

- $+$ and \cdot are monotone w.r.t. \preceq .

Let $R = \langle |R|, \preceq, +, \cdot, 0, 1 \rangle$ be a resource semiring. To define R -graded modalities on a primary linear doctrine, we adapt the approach to modalities on doctrines by Dagnino and Rosolini [13], following the definition of graded linear exponential comonads on a symmetric monoidal category [9, 20, 26].

Given a resource semiring R and a doctrine $P : \mathcal{C}^{\text{op}} \rightarrow \mathcal{Pos}$, we denote by $R \times P$ the doctrine on \mathcal{C} mapping every object A to $R \times P(A)$ and an arrow $f : X \rightarrow A$ to $\text{id}_R \times P_f : R \times P(A) \rightarrow R \times P(X)$.

Definition 4.1. Let $\langle P, *, \kappa \rangle$ be a primary linear doctrine. A R -graded (linear exponential) modality on $\langle P, *, \kappa \rangle$ is a natural transformation $\flat : R^{\text{op}} \times P \rightarrow P$ satisfying the following axioms for all objects X in \mathcal{C} , α, β in $P(X)$ and r, s in $|R|$:

1. $\kappa \leq \flat(r, \kappa)$ (lax-monoidality-1)
2. $\flat(r, \alpha) * \flat(r, \beta) \leq \flat(r, \alpha * \beta)$ (lax-monoidality-2)
3. $\flat(0, \alpha) \leq \kappa$ (weakening)
4. $\flat(r + s, \alpha) \leq \flat(r, \alpha) * \flat(s, \alpha)$ (contraction)
5. $\flat(1, \alpha) \leq \alpha$ (counit)
6. $\flat(r \cdot s, \alpha) \leq \flat(r, \flat(s, \alpha))$ (comultiplication)

An R -graded (primary linear) doctrine is a primary linear doctrine together with an R -graded modality on it.

In the following, we will write $\flat_r \alpha$ for $\flat(r, \alpha)$; note that, for all $r \in |R|$, \flat_r is a natural transformation from P to itself.

The naturality condition implies additional properties of a graded modality: it is stable under reindexing, covariant in the second argument and contravariant in the first one. In particular, for all $r, s \in |R|$, if $r \preceq s$ then $\flat_s \leq \flat_r$. Basically, the graded modality behaves as a graded linear exponential comonad on each fibre.

Depending on the choice of the resource semiring, we can define several kinds of modalities as sketched below.

Examples 4.2. 1. The singleton set $\{\infty\}$ with the trivial order and the trivial operations is a resource semiring. An $\{\infty\}$ -graded modality on a primary linear doctrine P satisfies the following inequalities in each fibre: $b_\infty\alpha \leq \kappa$, $b_\infty\alpha \leq b_\infty\alpha * b_\infty\alpha$, $b_\infty\alpha \leq \alpha$ and $b_\infty\alpha \leq b_\infty b_\infty\alpha$. ensuring that b_∞ is a linear exponential comonad on P , hence it models the usual bang modality of Linear Logic [22]. Syntactic doctrines associated to the $(\otimes, \mathbf{1}, !)$ -fragment of Linear Logic are $\{\infty\}$ -graded.

2. Let \mathbb{N} be the semiring of natural numbers $\langle \mathbb{N}, +, \cdot, 0, 1 \rangle$ and \mathbb{N}_∞ its extension by a new point ∞ with $\infty + x = x + \infty = \infty \cdot x = x \cdot \infty = \infty$ for all $x \in \mathbb{N}_\infty$. Denote by \mathbb{N}^\equiv and \mathbb{N}_∞^\equiv the semirings \mathbb{N} and \mathbb{N}_∞ ordered by the equality relation. If b is an \mathbb{N}^\equiv -graded modality on P , since $n = 1 + \dots + 1$ (n times), contraction and counit imply

$$b_n\alpha \leq \underbrace{b_1\alpha * \dots * b_1\alpha}_n \leq \underbrace{\alpha * \dots * \alpha}_n$$

Therefore, $b_n\alpha$ provides exactly n copies of α , that is, we have to use α exactly n times. Syntactically, this corresponds to the *exact usage* modality $!_n$ of Bounded Linear Logic [23] and, indeed, syntactic doctrines built out from it are \mathbb{N}^\equiv -graded.

If we consider \mathbb{N}_∞^\equiv , we get the additional modality b_∞ which behaves as in Item 1, modelling arbitrary usage.

3. Every resource semiring R gives rise to the *interval semiring* $\text{Int}(R) = \langle |\text{Int}(R)|, \sqsubseteq, +, \cdot, \langle 0, 0 \rangle, \langle 1, 1 \rangle \rangle$ where $|\text{Int}(R)|$ is the set of pairs $\langle r, s \rangle \in |R| \times |R|$ such that $r \preceq s$, $\langle r, s \rangle \sqsubseteq \langle r', s' \rangle$ iff $r' \preceq r$ and $s \preceq s'$, while the other operations are defined pointwise. Let \mathbb{N}_∞^\leq be the semiring \mathbb{N}_∞ in Item 2 ordered by the usual order on extended natural numbers. Any $\text{Int}(\mathbb{N}_\infty^\leq)$ -graded modality on a primary linear doctrine satisfies, for all $\langle n, m \rangle \in |\text{Int}(\mathbb{N}_\infty^\leq)|$ and $k \in n..m$ and $k \neq \infty$,

$$b_{\langle n, m \rangle}\alpha \leq b_{\langle k, k \rangle}\alpha \leq \underbrace{b_{\langle 1, 1 \rangle}\alpha * \dots * b_{\langle 1, 1 \rangle}\alpha}_k \leq \underbrace{\alpha * \dots * \alpha}_k$$

by contravariance, contraction and counit, as $\langle k, k \rangle \sqsubseteq \langle n, m \rangle$ and $\langle k, k \rangle = \langle 1, 1 \rangle + \dots + \langle 1, 1 \rangle$ (k times). Then the modality $b_{\langle n, m \rangle}$ models bounded usage as $b_{\langle n, m \rangle}\alpha$ provides a number of copies of α between n and m . Note that modalities of shape $b_{\langle n, \infty \rangle}$ allow unbounded usage.

Doctrines endowed by this family of modalities encode several substructural logics [37, 14], for instance, $b_{\langle 1, 1 \rangle}$ encodes linear usage, $b_{\langle 0, 1 \rangle}$

encodes affine usage, $\flat_{\langle 0, \infty \rangle}$ encodes intuitionistic usage and $\flat_{\langle 1, \infty \rangle}$ encodes relevant usage.

4. Consider the resource semiring of non-negative real numbers $\mathbb{R}_{\geq 0} = \langle [0, \infty), \leq, +, \cdot, 0, 1 \rangle$ and take the primary linear doctrine $\langle \mathcal{P}_{[0, \infty]}, +, 0 \rangle$ as in Example 3.2(2) where $[0, \infty]$ is the Lawvere quantale [30]. For every $\alpha : X \rightarrow [0, \infty]$ in $\mathcal{P}_{[0, \infty]}(X)$ and $r \in [0, \infty)$, the assignment $\alpha \mapsto \mathbf{m}_r \alpha$, where $\mathbf{m}_r \alpha(x) = r \cdot \alpha(x)$, determines an $\mathbb{R}_{\geq 0}$ -graded modality on $\mathcal{P}_{[0, \infty]}$, with $r \cdot \infty = \infty$ when $r \neq 0$ and it is 0 otherwise.
5. We recall from Atkey [4] that an R -Linear Combinatory Algebra (R -LCA) is a BCI algebra A with a semiring R and a function $!_r : |A| \rightarrow |A|$ for r in $|R|$ and elements $K, W_{r,s}, D, d_{r,s}, F_r$ of $|A|$ satisfying

$$\begin{aligned} K.x.!_0y &= x & d_{r,s}.!_{r,s}x &= !_r!_s x \\ W_{r,s}.x.!_{r+s}y &= x.!_r y.!_s y & F_r.!_r x.!_r y &= !_r(x.y) \\ D.!_1x &= x \end{aligned}$$

LCAs [3] are generalised by R -LCAs, as every LCA is a $\{\infty\}$ -LCA where $\{\infty\}$ is the singleton semiring of Item 1. When R is ordered (i.e. it is a resource semiring), we define an ordered R -LCA to be a R -LCA together with a combinator $O_{r,s}$ for all r, s in $|R|$ with $r \preceq s$, satisfying

$$O_{r,s}.!_s x = !_r x$$

Every R -LCA is an ordered R -LCA once we endow R with the trivial order ($r \preceq s$ when $r = s$), in this case $O_{r,s} = I$. Any affine LCA $(A, \cdot, !)$, such as those described by Abramsky [2], determines an ordered $\mathbb{N}_{\infty}^{\leq}$ -LCA as follows: $!_0 x = I$, $!_n x$ is given by the linear tupling of n copies of x and $!_{\infty} x = !x$ (here O combinators are obtained by K , which in affine LCAs satisfies $K.x.y = x$).

Recall from Example 3.2(3) primary linear doctrines of the form \mathcal{R}_A . If A is an ordered R -LCA, for $\alpha : X \rightarrow \mathcal{P}(|A|)$ in $\mathcal{R}_A(X)$ and r in $|R|$, define $\flat_r \alpha : X \rightarrow \mathcal{P}(|A|)$ as the function mapping x to $\{!_r a \mid a \in \alpha(x)\}$; then, $\langle \mathcal{R}_A, \flat \rangle$ is an R -graded doctrine.

R -graded doctrines are the objects of the 2-category \mathbf{LD}_R where a 1-arrow from $\langle P, \flat \rangle$ to $\langle Q, \sharp \rangle$ is a 1-arrow $\langle F, f \rangle : P \rightarrow Q$ in \mathbf{LD} preserving modalities, that is, such that, for all $r \in |R|$ and object X in the base of P , $\sharp_r F_X \circ f_X = f_X \circ \flat_r X$ holds. A 2-arrow from $\langle F, f \rangle$ to $\langle G, g \rangle$ is a 2-arrow $\theta : \langle F, f \rangle \Rightarrow \langle G, g \rangle$ in \mathbf{LD} . Compositions and identities are those of \mathbf{LD} .

The following proposition shows that (non-linear) primary doctrines are in particular R -graded, i.e. there is an inclusion $\mathbf{PD} \hookrightarrow \mathbf{LD}_R$.

Proposition 4.3. *Let $\langle P, *, \kappa \rangle$ be a primary linear doctrine and $\flat_r : P \rightarrow P$ be the identity for all $r \in |R|$. Then, P is a primary doctrine iff \flat is an R -graded modality on it.*

Proof. Let X be an object in the base category of P and α an element in $P(X)$. The left-to-right implication is easy to check: the non-trivial axiom to verify is weakening, indeed, we have $\flat_0 \alpha = \alpha \leq \kappa$, as κ is the top element of $P(X)$. For the converse we have to show that κ is the top element of $P(X)$ and $\alpha \leq \alpha * \alpha$ for all $\alpha \in P(X)$. By weakening $\alpha = \flat_0 \alpha \leq \kappa$ and contraction gives $\alpha = \flat_{r+r} \alpha \leq \flat_r \alpha * \flat_r \alpha = \alpha * \alpha$. \square

We can now define the notion of quantitative equality for R -graded doctrines.

Definition 4.4. An R -Lipschitz doctrine is a triple $\langle P, \flat, \mathbf{d} \rangle$ where $\langle P, \flat \rangle$ is an R -graded doctrine and, for each object A in the base, \mathbf{d}_A is a P -distance on A such that, for all objects A and X and α in $P(X \times A)$

(a) there is an r in $|R|$ such that

$$P_{\langle \pi_1, \pi_2 \rangle}(\alpha) * \flat_r P_{\langle \pi_2, \pi_3 \rangle}(\mathbf{d}_A) \leq P_{\langle \pi_1, \pi_3 \rangle}(\alpha);$$

(b) $\mathbf{d}_{A \times X} = P_{\langle \pi_1, \pi_3 \rangle}(\mathbf{d}_A) * P_{\langle \pi_2, \pi_4 \rangle}(\mathbf{d}_X)$;

(c) $\mathbf{d}_{A \times X} \leq P_{\langle \pi_1, \pi_3 \rangle}(\mathbf{d}_A)$ and $\mathbf{d}_{A \times X} \leq P_{\langle \pi_2, \pi_4 \rangle}(\mathbf{d}_X)$;

(d) $\mathbf{d}_1 = \kappa$.

The key difference between elementary and R -Lipschitz doctrines is the substitutive property, which, taking advantage of graded modalities, in the latter is stated in a resource sensitive way. Indeed, to prove a substitution, we need to have *enough equality resources*.

Since symmetry and transitivity are no longer derivable from substitutivity, for Lipschitz doctrines we assumed \mathbf{d}_A to be a P -distance (and not only a reflexive relation), as these are natural properties for equality. Moreover, the proof of Proposition 3.7, that ensures that equality in elementary primary linear doctrines is a P -distance, does not use replicability of δ , suggesting that to be a P -distance is a truly linear behaviour.

The product $A \times B$ mimics the concatenation of the context A with the context B , hence axioms (b), (c) and (d) encode the *independence of*

contexts. This property allows us to use independently the equalities in a product, for instance, if $r = r_1 + \dots + r_n$, then we can derive $b_r \mathbf{d}_{X_1 \times \dots \times X_n} \leq b_{r_1} P_{\langle \pi_1, \pi_{n+1} \rangle}(\mathbf{d}_{X_1}) * \dots * b_{r_n} P_{\langle \pi_n, \pi_{2n} \rangle}(\mathbf{d}_{X_n})$ and use separately each equality.

Axiom (a), describing substitution, can be equivalently rephrased as follows.

Proposition 4.5. *Let $\langle P, \flat \rangle$ be an R -graded doctrine and \mathbf{d} a family of P -distances. The following are equivalent:*

1. $\langle P, \flat, \mathbf{d} \rangle$ is R -Lipschitz;
2. $\langle P, \flat, \mathbf{d} \rangle$ satisfies axioms of Definition 4.4 where (a) is replaced by the following:

(a') for all objects A and α in $P(A)$, there is $r \in |R|$ such that

$$P_{\pi_1}(\alpha) * b_r \mathbf{d}_A \leq P_{\pi_2}(\alpha).$$

Proof. 1) \Rightarrow 2) take as X the terminal object of \mathcal{C} to get (a'). 2) \Rightarrow 1) Take α in $P(X \times A)$, then $P_{\langle \pi_1, \pi_2 \rangle}(\alpha) * b_r \mathbf{d}_{X \times A} \leq P_{\langle \pi_3, \pi_4 \rangle}(\alpha)$. By (b) one has $P_{\langle \pi_1, \pi_2 \rangle}(\alpha) * b_r P_{\langle \pi_2, \pi_4 \rangle} \mathbf{d}_A \leq P_{\langle \pi_3, \pi_4 \rangle}(\alpha)$. Reindexing along $\langle \pi_1, \pi_2, \pi_1, \pi_3 \rangle : X \times A \times A \rightarrow X \times A \times X \times A$ completes the proof. \square

Axioms (c) and (d) of Definition 4.4 are equivalent to affineness of equality.

Proposition 4.6. *Let $\langle P, \flat \rangle$ be an R -graded doctrine and \mathbf{d} be a family of P -distances. The following are equivalent:*

1. $\langle P, \flat, \mathbf{d} \rangle$ is R -Lipschitz;
2. \mathbf{d} satisfies the following axioms:

(a'') axioms (a) and (b) of Definition 4.4 hold,

(b'') \mathbf{d}_A is affine.

Proof. 1) \Rightarrow 2) follows from $\mathbf{d}_{A \times 1} = P_{\langle \pi_1, \pi_3 \rangle}(\mathbf{d}_A)$ and $\mathbf{d}_{A \times 1} \leq \mathbf{d}_1 = \kappa$. 2) \Rightarrow 1) is immediate. \square

The quantitative nature of this notion of equality can be noticed also from its relationship with reindexing, described by the following proposition.

Proposition 4.7. *Let $\langle P, \flat, \mathbf{d} \rangle$ be an R -Lipschitz doctrine. For any arrow $f : A_1 \times \dots \times A_n \rightarrow B$ in the base category of P , there are $r_1, \dots, r_n \in |R|$ such that*

$$\flat_{r_1} P_{\langle \pi_1, \pi_{n+1} \rangle}(\mathbf{d}_{A_1}) * \dots * \flat_{r_n} P_{\langle \pi_n, \pi_{2n} \rangle}(\mathbf{d}_{A_n}) \leq P_{f \times f}(\mathbf{d}_B)$$

Proof. For all $i \in 1..n$, let $\vec{\pi}_i = \langle \pi_i, \dots, \pi_n \rangle$. By Item (a) of Definition 4.4 we know that, for all $i \in 1..n$, there is $r_i \in |R|$ such that

$$P_{\langle \pi_1, \dots, \pi_{i-1}, \vec{\pi}_i, \pi_{n+1}, \dots, \pi_{n+i-1}, \vec{\pi}_i \rangle}(P_{f \times f}(\mathbf{d}_B)) * \flat_{r_i}(P_{\langle \pi_1, \pi_{n+i} \rangle}(\mathbf{d}_{A_i})) \leq P_{\langle \pi_1, \dots, \pi_i, \vec{\pi}_{i+1}, \pi_{n+1}, \dots, \pi_{n+i}, \vec{\pi}_{i+1} \rangle}(P_{f \times f}(\mathbf{d}_B))$$

holds in $A_1 \times \dots \times A_n \times A_1 \times \dots \times A_n$. Combining these inequalities, using monotonicity of $*$ and transitivity of \leq , we get

$$P_{\langle \pi_1, \dots, \pi_n, \pi_1, \dots, \pi_n \rangle}(P_{f \times f}(\mathbf{d}_B)) * \flat_{r_1}(P_{\langle \pi_1, \pi_{n+1} \rangle}(\mathbf{d}_{A_1})) * \dots * \flat_{r_n}(P_{\langle \pi_n, \pi_{2n} \rangle}(\mathbf{d}_{A_n})) \leq P_{f \times f}(\mathbf{d}_B)$$

Note that $P_{\langle \pi_1, \dots, \pi_n, \pi_1, \dots, \pi_n \rangle}(P_{f \times f}(\mathbf{d}_B)) = P_{f \circ \langle \pi_1, \dots, \pi_n \rangle}(P_{\Delta_B}(\mathbf{d}_B))$. Then, the thesis follows because $\kappa \leq P_{f \circ \langle \pi_1, \dots, \pi_n \rangle}(P_{\Delta_B}(\mathbf{d}_B))$ holds by reflexivity of \mathbf{d}_B . \square

Corollary 4.8. *Let $\langle P, \flat, \mathbf{d} \rangle$ be an R -Lipschitz doctrine. For any arrow $f : A \rightarrow B$ in the base category of P , there exists $r \in |R|$ such that $\flat_r \mathbf{d}_A \leq P_{f \times f}(\mathbf{d}_B)$.*

Since \mathbf{d} is a family of distances, the property described by Corollary 4.8 resembles the Lipschitz condition on maps between metric spaces and motivates the name in Definition 4.4.

Finally, also R -Lipschitz doctrines can be organised into a 2-category \mathbf{LLD}_R , whose objects are R -Lipschitz doctrines, a 1-arrow from $\langle P, \flat, \mathbf{d} \rangle$ to $\langle Q, \flat', \mathbf{d}' \rangle$ is 1-arrow $\langle F, f \rangle : \langle P, \flat \rangle \rightarrow \langle Q, \flat' \rangle$ preserving the distance, that is, such that, for each object A in the base category of P , it holds $\mathbf{d}'_{FA} = f_{A \times A}(\mathbf{d}_A)$, and a 2-arrow from $\langle F, f \rangle$ to $\langle G, g \rangle$ is a 2-arrow $\theta : \langle F, f \rangle \Rightarrow \langle G, g \rangle$ in \mathbf{LD}_R .

4.2 A deductive calculus for quantitative equality

In the examples presented so far, we saw that theories \mathcal{T} over a signature L generate the syntactic doctrines $Prp_{\mathcal{T}}$. In this section we describe a calculus whose associated syntactic doctrine is R -Lipschitz.

Assume $R = \langle |R|, \preceq, +, \cdot, 0, 1 \rangle$ is a resource semiring and L a first order signature. A judgment has the form $\Gamma \vdash_{\vec{\sigma}} \phi$ where $\Gamma = \phi_1, \dots, \phi_k$ is a finite multiset of formulas, ϕ is a formula and all formulas $\phi_1, \dots, \phi_k, \phi$ are in the

$$\begin{array}{c}
\text{[W]} \frac{\Gamma \vdash \phi}{\Gamma, !_0\psi \vdash \phi} \qquad \text{[C]} \frac{\Gamma, !_r\psi, !_s\psi \vdash \phi}{\Gamma, !_{r+s}\psi \vdash \phi} \qquad \text{[DER]} \frac{\Gamma, \psi \vdash \phi}{\Gamma, !_1\psi \vdash \phi} \\
\text{[PRO]} \frac{!_{s_1}\psi_1, \dots, !_{s_n}\psi_n \vdash \phi}{!_{r \cdot s_1}\psi_1, \dots, !_{r \cdot s_n}\psi_n \vdash !_r\phi} \qquad \text{[DECR]} \frac{\Gamma \vdash !_s\phi}{\Gamma \vdash !_r\phi} \quad r \preceq s
\end{array}$$

Figure 1: Rules for graded bang modality

context $\vec{\sigma} = \langle x_1 : \sigma_1, \dots, x_n : \sigma_n \rangle$. We will often omit context $\vec{\sigma}$ to improve readability.

First of all, we introduce PLL_R , a calculus extending the $(\otimes, \mathbf{1})$ -fragment of Linear Logic by a family of modalities $!_r$, for $r \in |R|$. The front ‘P’ is for primary as a syntactic doctrine associated with PLL_R is a primary linear doctrine (endowed by an R -graded modality).

Rules of PLL_R are a subset of the calculus presented by Breuvert and Pagani [8], in Fig. 1 we report rules for modalities, while structural rules and rules for \otimes and $\mathbf{1}$ are standard.

The first four rules are the graded variant of the standard rules for the bang modality of Linear Logic. Rules [W] and [C] encode graded structural rules: by weakening we can add formulas marked as not used (labelled by 0), while contraction keeps track of the usage by addition. Rule [DER] tells that hypotheses with grade 1 can be treated as linear hypotheses, while rule [PRO] introduces $!_r$, scaling by r the grades of the hypotheses. Rule [DECR] ensures that one can decrease the usage of a formula following (contravariantly) the order of the semiring.

We now extend PLL_R by quantitative equality. Intuitively, we need a way to assign to each formula ϕ and each variable x the cost, in terms of resources, of substituting x in ϕ . To this end, we first have to enrich the notions of signature to take into account these resources.

An R -graded signature is a first order signature L where symbols have an R -graded arity, that is, an assignment $|-|$ as the following:

$$|f| = \langle r_1, \sigma_1 \rangle, \dots, \langle r_n, \sigma_n \rangle \rightarrow \tau \qquad |p| = \langle r_1, \sigma_1 \rangle, \dots, \langle r_n, \sigma_n \rangle$$

where r_1, \dots, r_n are elements of R , $\sigma_1, \dots, \sigma_n, \tau$ are sorts and f and p are a function and a predicate symbol, respectively. Intuitively, the sort σ_i says which sort the i -th argument belongs to, while r_i says how much the cost of substituting a variable in the i -th position is amplified. We write $|f|_i$

(resp. $|p|_i$) in place of r_i in the assignments above. Note that a graded arity determines a standard arity just by forgetting resources.

Denote by V and Trm the sets of variables and terms inductively constructed from symbols in L in the usual way (using the standard arity obtained from the graded one by forgetting resources). The resources $|f|_i$ determines a function $\mathbf{gr} : Trm \times V \rightarrow |R|$ as follows:

- $\mathbf{gr}(z, x) = 0$ if $x \neq z$
- $\mathbf{gr}(x, x) = 1$
- $\mathbf{gr}(f(t_1, \dots, t_n), x) = |f|_1 \cdot \mathbf{gr}(t_1, x) + \dots + |f|_n \cdot \mathbf{gr}(t_n, x)$

Intuitively, $\mathbf{gr}(t, x)$ represents the cost of substituting the variable x inside the term t . Note that $\mathbf{gr}(t, x)$ depends on the number of occurrences of x in t . For instance, if $f_0, f_1(x), f_2(x, x), \dots, f_n(x, x, \dots, x)$ are terms where f_i is a function symbol with i arguments and $|f_i|_i = 1$ for all $i \in 0..n$, then $\mathbf{gr}(f_i(x, \dots, x), x) = 1 + \dots + 1$ (i times). In particular, for constants, i.e. function symbols f with no arguments, we always have $\mathbf{gr}(f, x) = 0$.

Denote by Wff the set of well formed formulas constructed from symbols in L , the equality symbol $=$ and using \otimes , $\mathbf{1}$ and $!_r$ as connectives. We extend the function $\mathbf{gr} : Trm \times V \rightarrow |R|$ to a function $\mathbf{gr} : Wff \times V \rightarrow |R|$ mapping a formula ϕ and a variable x to the amount of resources one needs to substitute x in ϕ .

- $\mathbf{gr}(p(t_1, \dots, t_n), x) = |p|_1 \cdot \mathbf{gr}(t_1, x) + \dots + |p|_n \cdot \mathbf{gr}(t_n, x)$,
- $\mathbf{gr}(t =_\tau u, x) = \mathbf{gr}(t, x) + \mathbf{gr}(u, x)$,
- $\mathbf{gr}(\phi \otimes \psi, x) = \mathbf{gr}(\phi, x) + \mathbf{gr}(\psi, x)$,
- $\mathbf{gr}(\mathbf{1}, x) = 0$,
- $\mathbf{gr}(!_r \phi, x) = r \cdot \mathbf{gr}(\phi, x)$.

Similarly to terms, the cost for substituting the variable x in a formula depends on the number of occurrences of x in ϕ . For instance, we have $\mathbf{gr}(p(x) \otimes p(y), x) = |p|_1$, while $\mathbf{gr}(p(x) \otimes p(x), x) = |p|_1 + |p|_1$. Again, for a predicate symbol p with no arguments, we have $\mathbf{gr}(p, x) = 0$, independently from the value of $|p|$.

The calculus PLL_R extended by rules in Fig. 2 will be called $LPLL_R$, where the first ‘L’ is for Lipschitz.

$$\begin{array}{c}
\text{[R]} \frac{}{\vdash t \equiv_{\tau} t} \quad \text{[S]} \frac{\Gamma \vdash t \equiv_{\tau} u}{\Gamma \vdash u \equiv_{\tau} t} \quad \text{[T]} \frac{\Gamma \vdash t \equiv_{\tau} u \quad \Delta \vdash u \equiv_{\tau} v}{\Gamma, \Delta \vdash t \equiv_{\tau} v} \\
\text{[W-EQ]} \frac{\Gamma \vdash \phi}{\Gamma, t \equiv_{\tau} u \vdash \phi} \quad \text{[SUBST]} \frac{\Gamma \vdash \phi[t/x] \quad \Delta \vdash !_{\mathbf{gr}(\phi, x)} t \equiv_{\tau} u}{\Gamma, \Delta \vdash \phi[u/x]}
\end{array}$$

Figure 2: Rules for graded equality.

Rules are pretty standard: we have rules for reflexivity, symmetry and transitivity of \equiv_{σ} and a rule for weakening. Note the substitution rule: a substitution is derivable only if *enough equality resources* are available and these are determined by the function \mathbf{gr} .

Remark 4.9. For a function symbol f with arity $|f| = \langle |f|_1, \sigma_1 \rangle \dots \langle |f|_n, \sigma_n \rangle \rightarrow \tau$, one can prove the entailment

$$!_{|f|_1}(x_1 \equiv_{\sigma_1} y_1), \dots, !_{|f|_n}(x_n \equiv_{\sigma_n} y_n) \vdash f(x_1, \dots, x_n) \equiv_{\tau} f(y_1, \dots, y_n)$$

Intuitively, interpreting the equality as a distance, this means that the application of the function f amplifies the distance between x_i and y_i by a factor $|f|_i$. Similarly, if p is a predicate symbol of arity $|p| = \langle |p|_1, \sigma_1 \rangle \dots \langle |p|_n, \sigma_n \rangle$, one can prove

$$p(x_1, \dots, x_n), !_{|p|_1}(x_1 \equiv_{\sigma_1} y_1), \dots, !_{|p|_n}(x_n \equiv_{\sigma_n} y_n) \vdash p(y_1, \dots, y_n)$$

Intuitively, this means that we need to amplify equality between x_i and y_i by a factor $|p|_i$ to derive $p(y_1, \dots, y_n)$ from $p(x_1, \dots, x_n)$.

Let \mathcal{T} be a theory in LPLL_R over the R -graded signature L , then the syntactic doctrine $\text{Prp}_{\mathcal{T}}$ is an R -Lipschitz doctrine with R -graded modality given by $!$ and a family of distances \mathbf{d}^{\equiv} inductively defined by

- $\mathbf{d}_{\langle \rangle}^{\equiv} = \mathbf{1}$;
- $\mathbf{d}_{\langle \vec{\sigma}, x_{n+1} : \sigma_{n+1} \rangle}^{\equiv} = \mathbf{d}_{\vec{\sigma}}^{\equiv} \otimes (x_{n+1} \equiv_{\sigma_{n+1}} x'_{n+1})$.

Basically, checking that $\text{Prp}_{\mathcal{T}}$ is R -Lipschitz amounts to prove that rules in Fig. 1 imply axioms in Definition 4.1, $!_r$, \otimes , $\mathbf{1}$ and \vdash replace b_r , $*$, κ and \leq , and rules in Fig. 2 imply conditions in Definition 4.4. Note that the rule [W-EQ] is needed to ensure context independence (see Proposition 4.6).

4.3 Semantics in R -Lipschitz doctrines

The interpretation of a theory in a doctrine is standard [39, 25]: it maps contexts and terms to objects and arrows of the base and formulas to elements of the fibres, respecting entailments in the theory. The interpretation of LPLL_R in an R -Lipschitz doctrine essentially follows this way, except that we have to make the function gr agree with the structure of the doctrine. To this end, we define a notion of interpretation that takes into account resources in the graded arity of the signature.

Let L be an R -graded signature. An R -graded interpretation of L into a R -Lipschitz doctrine $\langle P, \flat, \mathbf{d} \rangle$ assigns to every sort σ an object $\llbracket \sigma \rrbracket$ in the base of P , to every function symbol f of arity $\langle |f|_1, \sigma_1 \rangle \dots \langle |f|_n, \sigma_n \rangle \rightarrow \tau$ an arrow $\llbracket f \rrbracket : \llbracket \sigma_1 \rrbracket \times \dots \times \llbracket \sigma_n \rrbracket \rightarrow \llbracket \tau \rrbracket$ in the base of P such that

$$\prod_{i=1}^n \flat_{|f|_i} P_{\langle \pi_i, \pi_{n+i} \rangle}(\mathbf{d}_{\llbracket \sigma_i \rrbracket}) \leq P_{\llbracket f \rrbracket \times \llbracket f \rrbracket}(\mathbf{d}_{\llbracket \tau \rrbracket}) \quad (2)$$

and to every predicate symbol p of arity $\langle |p|_1, \sigma_1 \rangle \dots \langle |p|_n, \sigma_n \rangle$ an element $\llbracket p \rrbracket$ in $P(\llbracket \sigma_1 \rrbracket \times \dots \times \llbracket \sigma_n \rrbracket)$ such that

$$P_{\langle \pi_1, \dots, \pi_n \rangle}(\llbracket p \rrbracket) * \prod_{i=1}^n \flat_{|p|_i} P_{\langle \pi_i, \pi_{n+i} \rangle}(\mathbf{d}_{\llbracket \sigma_i \rrbracket}) \leq P_{\langle \pi_{n+1}, \dots, \pi_{2n} \rangle}(\llbracket p \rrbracket) \quad (3)$$

Intuitively, we can pick as interpretation of a function symbol f an arrow $\llbracket f \rrbracket$ for which the cost of substituting the i -th argument is exactly $|f|_i$, and similarly for predicate symbols. In other words, the graded arity of f determines valid Lipschitz constants for its interpretation $\llbracket f \rrbracket$.

An R -graded interpretation of L induces an interpretation of contexts and terms in the base of P and an interpretation of formulas in the fibres of P . More precisely, a context $\vec{\sigma} = \langle x_1 : \sigma_1, \dots, x_n : \sigma_n \rangle$ is interpreted by an object $\llbracket \vec{\sigma} \rrbracket_{\text{Trm}} = \llbracket \sigma_1 \rrbracket \times \dots \times \llbracket \sigma_n \rrbracket$, a term t of sort τ in $\vec{\sigma}$ by an arrow $\llbracket t \rrbracket_{\text{Trm}} : \llbracket \vec{\sigma} \rrbracket_{\text{Trm}} \rightarrow \llbracket \tau \rrbracket$ and a formula ϕ in $\vec{\sigma}$ by an element $\llbracket \phi \rrbracket_{\text{Wff}} \in P(\llbracket \vec{\sigma} \rrbracket_{\text{Trm}})$, inductively defined as follows:

$$\begin{aligned} \llbracket x_i \rrbracket_{\text{Trm}} &= \pi_i & \llbracket \mathbf{1} \rrbracket_{\text{Wff}} &= \kappa \\ \llbracket f(t_1, \dots, t_n) \rrbracket_{\text{Trm}} &= \llbracket f \rrbracket \circ \langle \llbracket t_1 \rrbracket_{\text{Trm}}, \dots, \llbracket t_n \rrbracket_{\text{Trm}} \rangle & \llbracket \phi \otimes \psi \rrbracket_{\text{Wff}} &= \llbracket \phi \rrbracket_{\text{Wff}} * \llbracket \psi \rrbracket_{\text{Wff}} \\ \llbracket p(t_1, \dots, t_n) \rrbracket_{\text{Wff}} &= P_{\langle \llbracket t_1 \rrbracket_{\text{Trm}}, \dots, \llbracket t_n \rrbracket_{\text{Trm}} \rangle}(\llbracket p \rrbracket) & \llbracket !_r \phi \rrbracket_{\text{Wff}} &= \flat_r \llbracket \phi \rrbracket_{\text{Wff}} \\ \llbracket t \equiv_{\tau} u \rrbracket_{\text{Wff}} &= P_{\langle \llbracket t \rrbracket_{\text{Trm}}, \llbracket u \rrbracket_{\text{Trm}} \rangle}(\mathbf{d}_{\llbracket \tau \rrbracket}) \end{aligned}$$

An R -graded interpretation of a theory \mathcal{T} of LPLL_R over the signature L , is an R -graded interpretation $\llbracket - \rrbracket$ of L such that $\llbracket - \rrbracket_{\text{Wff}}$ respects entail-

ments in \mathcal{T} . The following theorem states soundness for the semantics of LPLL_R .

Theorem 4.10. *Let $\llbracket - \rrbracket$ be an R -graded interpretation of a theory \mathcal{T} in LPLL_R over L into an R -Lipschitz $\langle P, \flat, \mathbf{d} \rangle$. Then, $\langle \llbracket - \rrbracket_{\text{Trm}}, \llbracket - \rrbracket_{\text{Wff}} \rangle : \langle \text{Prp}_{\mathcal{T}}, !, \mathbf{d}^{\perp} \rangle \rightarrow \langle P, \flat, \mathbf{d} \rangle$ is a 1-arrow in \mathbf{LLD}_R .*

The proof is carried out by the usual induction on rules [39, 25], relying on the following lemma (where we omitted subscripts Trm and Wff from $\llbracket - \rrbracket$):

Lemma 4.11. *For every R -graded interpretation $\llbracket - \rrbracket$ of L into $\langle P, \flat, \mathbf{d} \rangle$ the following hold:*

1. for any term $t : \tau$ in the context $\vec{\sigma} = \langle x_1 : \sigma_1, \dots, x_n : \sigma_n \rangle$

$$\prod_{i=1}^n \flat_{\text{gr}(t, x_i)} P_{\langle \pi_i, \pi_{n+i} \rangle}(\mathbf{d}_{\llbracket \sigma_i \rrbracket}) \leq P_{\llbracket t \rrbracket \times \llbracket t \rrbracket}(\mathbf{d}_{\llbracket \tau \rrbracket})$$

2. for any formula ϕ in the context $\vec{\sigma} = \langle x_1 : \sigma_1, \dots, x_n : \sigma_n \rangle$

$$P_{\text{pr}_1}(\llbracket \phi \rrbracket) * \prod_{i=1}^n \flat_{\text{gr}(\phi, x_i)} P_{\langle \pi_i, \pi_{n+i} \rangle}(\mathbf{d}_{\llbracket \sigma_i \rrbracket}) \leq P_{\text{pr}_2}(\llbracket \phi \rrbracket)$$

where $\text{pr}_1 = \langle \pi_1, \dots, \pi_n \rangle$ and $\text{pr}_2 = \langle \pi_{n+1}, \dots, \pi_{2n} \rangle$.

The following proposition states completeness of the semantics of LPLL_R :

Proposition 4.12. *Let \mathcal{T} be a theory in LPLL_R over a signature L and ϕ, ψ formulas over L . If, for all R -Lipschitz doctrines $\langle P, \flat, \mathbf{d} \rangle$ and for all interpretations $\llbracket - \rrbracket$ of \mathcal{T} in $\langle P, \flat, \mathbf{d} \rangle$, we have $\llbracket \phi \rrbracket_{\text{Wff}} \leq \llbracket \psi \rrbracket_{\text{Wff}}$, then $\phi \vdash \psi$ is provable in \mathcal{T} .*

The proof of this result is straightforward, as we can take the trivial interpretation in the syntactic doctrine $\text{Prp}_{\mathcal{T}}$, which is R -Lipschitz. Hence, R -Lipschitz doctrines provide a sound and complete categorical semantics of LPLL_R .

Remark 4.13. One could prove a stronger completeness result, showing an *internal language theorem*. This consists in proving the equivalence between the 2-category of R -Lipschitz doctrines and a suitable 2-category of LPLL_R theories. As it is well-known, the key lemma of this kind of results shows

that from any doctrine one can extract a theory, where, loosely speaking, sorts and function symbols are objects and arrows in the base, while predicate symbols and axioms are elements and entailments in the fibres. One could do the same thing for R -Lipschitz doctrines, where the difficult part is to determine grades in the R -graded arities. This is achieved thanks to Proposition 4.7 with the use of the axiom of choice. A rigorous proof is left for future work.

So far we did not list any example of Lipschitz doctrines. The next subsection describes a construction producing Lipschitz doctrines from R -graded ones. This gives semantics for LPLL_R .

4.4 R -Lipschitz doctrines as coalgebras

The construction is given by a 2-functor $\text{R}_{\text{Lip}} : \mathbf{LD}_R \rightarrow \mathbf{LLD}_R$. This turns out to be right 2-adjoint to the obvious forgetful 2-functor $\text{U}_{\text{Lip}} : \mathbf{LLD}_R \rightarrow \mathbf{LD}_R$ and R -Lipschitz doctrines are coalgebras of the 2-comonad induced by the 2-adjunction. This shows that quantitative equality has a coalgebraic nature exactly as the standard equality in First Order Logic, as proved by Emmenegger et al. [19].

To construct a right 2-adjoint to U_{Lip} , we build an R -Lipschitz doctrine $D_{\langle P, b \rangle} : \mathcal{L}_{\langle P, b \rangle}^{\text{op}} \rightarrow \mathcal{Pos}$ out of any R -graded doctrine $\langle P, b \rangle$. The category $\mathcal{L}_{\langle P, b \rangle}$ is as follows:

- objects are pairs $\langle A, \rho \rangle$ where A is an object in \mathcal{C} and ρ is an affine P -distance on A ,
- an arrow from $\langle A, \rho \rangle$ to $\langle B, \sigma \rangle$ is an arrow $f : A \rightarrow B$ in \mathcal{C} such that $b_r \rho \leq P_{f \times f}(\sigma)$, for some $r \in |R|$,
- composition and identities are those of \mathcal{C} .

Given $f : \langle A, \rho \rangle \rightarrow \langle B, \sigma \rangle$ we say that r tracks f if r is such that $b_r \rho \leq P_{f \times f}(\sigma)$. So $f : A \rightarrow B$ underlies an arrow $f : \langle A, \rho \rangle \rightarrow \langle B, \sigma \rangle$ if there is r that tracks f .

It is easy to check that $\mathcal{L}_{\langle P, b \rangle}$ is a category: identities are well-defined as for every $\langle A, \rho \rangle$ the identity id_A is tracked by 1 thanks to the counit axiom of b , while given $f : \langle A, \rho \rangle \rightarrow \langle B, \sigma \rangle$ and $g : \langle B, \sigma \rangle \rightarrow \langle C, \tau \rangle$ if r tracks f and s tracks g , then $s \cdot r$ tracks gf as the comultiplication axiom and naturality of b_s lead to

$$b_{s \cdot r} \rho \leq b_s b_r \rho \leq b_s P_{f \times f}(\sigma) = P_{f \times f}(b_s \sigma) \leq P_{f \times f}(P_{g \times g}(\tau))$$

Before proving that $\mathcal{L}_{\langle P, \flat \rangle}$ has finite products, we introduce a useful notation: given elements $\rho \in P(A \times A)$ and $\sigma \in P(B \times B)$, we denote by $\rho \boxtimes \sigma$ the element $P_{\langle \pi_1, \pi_3 \rangle}(\rho) * P_{\langle \pi_2, \pi_4 \rangle}(\sigma)$ in $P(A \times B \times A \times B)$, which has the following straightforward property:

Proposition 4.14. *Let $\langle P, *, \kappa \rangle$ be a primary linear doctrine. If ρ and σ are P -distances, $\rho \boxtimes \sigma$ is a P -distance. If ρ and σ are affine, $\rho \boxtimes \sigma$ is affine.*

Proposition 4.15. *The category $\mathcal{L}_{\langle P, \flat \rangle}$ has finite products.*

Proof. (Sketched) The terminal object in $\mathcal{L}_{\langle P, \flat \rangle}$ is $\langle 1, \kappa_{1 \times 1} \rangle$, where 1 is the terminal object in \mathcal{C} , and 0 tracks the unique arrow $A \rightarrow 1$. The product of $\langle A, \rho \rangle$ and $\langle B, \sigma \rangle$ is $\langle A \times B, \rho \boxtimes \sigma \rangle$. If π is a projection, then 1 tracks π . If r tracks f and s tracks g then $r + s$ tracks $\langle f, g \rangle$. \square

Given an affine P -distance ρ on an object A in \mathcal{C} , an element α in $P(A)$ and r in $|R|$, we write r tracks α when $P_{\pi_1}(\alpha) * \flat_r \rho \leq P_{\pi_2}(\alpha)$. The suborder of $P(A)$ of R -graded descent data is:

$$\mathcal{Des}_\rho(A) = \{\alpha \in P(A) \mid r \text{ tracks } \alpha \text{ for some } r \in |R|\}$$

Consider $f : \langle A, \rho \rangle \rightarrow \langle B, \sigma \rangle$ in $\mathcal{L}_{\langle P, \flat \rangle}$ and $\beta \in \mathcal{Des}_\sigma(B)$, then, r tracks f and s tracks β for some $r, s \in |R|$. By comultiplication and naturality of \flat_s , we get $\flat_{s \cdot r} \rho \leq \flat_s \flat_r \rho \leq \flat_s P_{f \times f}(\sigma) = P_{f \times f}(\flat_s \sigma)$. Thus, we have $P_{\pi_1}(P_f(\beta)) * \flat_{s \cdot r} \rho \leq P_{f \times f}(P_{\pi_1}(\beta) * \flat_s \sigma) \leq P_{f \times f}(P_{\pi_2}(\beta)) = P_{\pi_2}(P_f(\beta))$, showing that $s \cdot r$ tracks $P_f(\beta)$. In other words P_f applies $\mathcal{Des}_\sigma(B)$ to $\mathcal{Des}_\rho(A)$ and the assignments

$$D_{\langle P, \flat \rangle} \langle A, \rho \rangle := \mathcal{Des}_\rho(A) \quad D_{\langle P, \flat \rangle}(f) := P_f$$

determine a functor $D_{\langle P, \flat \rangle} : \mathcal{L}_{\langle P, \flat \rangle}^{\text{op}} \rightarrow \mathcal{Pos}$.

Proposition 4.16. *$D_{\langle P, \flat \rangle}$ is a R -Lipschitz doctrine.*

Proof. (Sketched) Take $\langle A, \rho \rangle$ in $\mathcal{L}_{\langle P, \flat \rangle}$. By weakening of \flat it holds that 0 tracks κ and if r tracks α and s tracks β , then $r + s$ tracks $\alpha * \beta$. So $\mathcal{Des}_\rho(A)$ is closed under the monoidal operations of $P(A)$. If r tracks α , then $s \cdot r$ tracks $\flat_s \alpha$, whence $\langle D_{\langle P, \flat \rangle}, \flat \rangle$ is R -graded. Let $\mathbf{d}_{\langle A, \rho \rangle} = \rho$, Proposition 4.5 proves that $\langle D_{\langle P, \flat \rangle}, \flat \rangle$ is R -Lipschitz. \square

The construction described above, mapping an R -graded doctrine into an R -Lipschitz one, extends to a 2-functor $R_{\text{Lip}} : \mathbf{LD}_R \rightarrow \mathbf{LLD}_R$ as follows.

Let $\langle F, f \rangle : \langle P, \flat \rangle \rightarrow \langle Q, \flat' \rangle$ be a 1-arrow in \mathbf{LD}_R and $h : \langle A, \rho \rangle \rightarrow \langle B, \sigma \rangle$ an arrow in $\mathcal{L}_{\langle P, \flat \rangle}$, the following assignments

$$\overline{F}\langle A, \rho \rangle := \langle FA, f_{A \times A}(\rho) \rangle \quad \overline{F}h := Fh$$

define a functor $\overline{F} : \mathcal{L}_{\langle P, \flat \rangle} \rightarrow \mathcal{L}_{\langle Q, \flat' \rangle}$. Given a 2-arrow $\theta : \langle F, f \rangle \Rightarrow \langle G, g \rangle$ in \mathbf{LD}_R define

$$\mathbf{R}_{\text{Lip}}\langle F, f \rangle := \langle \overline{F}, f \rangle \quad \mathbf{R}_{\text{Lip}}(\theta) := \theta$$

Proposition 4.17. *With the assignments above, $\mathbf{R}_{\text{Lip}} : \mathbf{LD}_R \rightarrow \mathbf{LLD}_R$ is a 2-functor.*

Note that, for each R -graded doctrine $\langle P, \flat \rangle$, there is an obvious forgetful 1-arrow $\langle U^P, u^P \rangle : \langle D_{\langle P, \flat \rangle}, \flat \rangle \rightarrow \langle P, \flat \rangle$ in \mathbf{LD}_R , where U^P forgets the distance and u^P is the inclusion. Furthermore, if $\langle P, \flat, \mathbf{d} \rangle$ is R -Lipschitz, there is a 1-arrow $\langle E^P, e^P \rangle : \langle P, \flat \rangle \rightarrow \langle D_{\langle P, \flat \rangle}, \flat \rangle$ in \mathbf{LLD}_R , where E^P maps $f : A \rightarrow B$ to $f : \langle A, \mathbf{d}_A \rangle \rightarrow \langle B, \mathbf{d}_B \rangle$ and e^P is the identity on fibres. Note that E^P is well-defined on objects as $\langle P, \flat, \mathbf{d} \rangle$ is R -Lipschitz and on arrows by Corollary 4.8; moreover, it preserves finite products by Items (b) and (d) of Definition 4.4, hence $\langle E^P, e^P \rangle$ is well-defined.

Theorem 4.18. *The 2-functor $\mathbf{U}_{\text{Lip}} : \mathbf{LLD}_R \rightarrow \mathbf{LD}_R$ is 2-comonadic with right 2-adjoint \mathbf{R}_{Lip} .*

Proof. (Sketched) The counit is $\langle U^P, u^P \rangle : \langle D_{\langle P, \flat \rangle}, \flat \rangle \rightarrow \langle P, \flat \rangle$. Take $\langle F, f \rangle : \langle Q, \flat' \rangle \rightarrow \langle P, \flat \rangle$ where $\langle Q, \flat' \rangle$ is R -Lipschitz. The universal arrow is $\mathbf{R}_{\text{Lip}}\langle F, f \rangle \circ \langle E^Q, e^Q \rangle : \langle Q, \flat' \rangle \rightarrow \langle D_{\langle P, \flat \rangle}, \flat \rangle$. \square

The restriction of \mathbf{R}_{Lip} to \mathbf{PD} gives the 2-adjunctions between \mathbf{PD} and \mathbf{ED} discussed in [19] as depicted by the commutative diagram

$$\begin{array}{ccc} \mathbf{LD}_R & \xrightarrow{\mathbf{R}_{\text{Lip}}} & \mathbf{LLD}_R \\ \uparrow & \xleftarrow{\top} & \uparrow \\ \mathbf{PD} & \xrightarrow{\top} & \mathbf{ED} \end{array}$$

The functor $\mathbf{R}_{\text{Lip}} : \mathbf{LD}_R \rightarrow \mathbf{LLD}_R$ can be used to construct R -Lipschitz doctrines from R -graded ones and obtain semantics for \mathbf{LPLL}_R . The application of \mathbf{R}_{Lip} to $\langle P, \flat \rangle$ gives $\langle D_{\langle P, \flat \rangle}, \flat \rangle$ that we call *Lipschitz completion* of $\langle P, \flat \rangle$.

We conclude the section with two concrete examples.

Consider the $\mathbb{R}_{\geq 0}$ -graded doctrine $\mathcal{P}_{[0, \infty]}$ presented in Examples 4.2(4) with $\mathbb{R}_{\geq 0}$ -graded modality \mathbf{m}_r , for $r \in [0, \infty)$, mapping $\alpha : X \rightarrow [0, \infty]$ to the function $\mathbf{m}_r \alpha = x \mapsto r \cdot \alpha(x)$. Here, a $\mathcal{P}_{[0, \infty]}$ -distance ρ on a set X is exactly a pseudometric $\rho : X \times X \rightarrow [0, \infty]$ in the common sense (i.e. a metric except that the identity of indiscernibles is dropped). The $\mathbb{R}_{\geq 0}$ -Lipschitz doctrine generated by \mathbf{R}_{Lip} is $D_{\langle \mathcal{P}_{[0, \infty]}, \mathbf{m} \rangle} : \mathcal{L}_{\langle \mathcal{P}_{[0, \infty]}, \mathbf{m} \rangle}^{\text{op}} \rightarrow \mathcal{Pos}$, whose base category $\mathcal{L}_{\langle \mathcal{P}_{[0, \infty]}, \mathbf{m} \rangle}$ is the category of pseudometric spaces and Lipschitz functions (cf. Corollary 4.8).

As described in Section 4.3, we can soundly interpret $\text{LPLL}_{\mathbb{R}_{\geq 0}}$ (over an $\mathbb{R}_{\geq 0}$ -graded signature L) in $D_{\langle \mathcal{P}_{[0, \infty]}, \mathbf{m} \rangle}$, by mapping sorts to pseudometric spaces, a function symbol f of arity $\langle |f|_1, \sigma_1 \rangle \dots \langle |f|_n, \sigma_n \rangle \rightarrow \tau$ to a Lipschitz function $\llbracket f \rrbracket : \llbracket \sigma_1 \rrbracket \times \dots \times \llbracket \sigma_n \rrbracket \rightarrow \llbracket \tau \rrbracket$, which will have as a Lipschitz constant the sum $|f|_1 + \dots + |f|_n$ and a predicate symbol p of arity $\langle |p|_1, \sigma_1 \rangle \dots \langle |p|_n, \sigma_n \rangle$ to a function $\llbracket p \rrbracket : \llbracket \sigma_1 \rrbracket \times \dots \times \llbracket \sigma_n \rrbracket \rightarrow [0, \infty]$, where $\llbracket \sigma \rrbracket$ is the underling set of the pseudometric space $\llbracket \sigma \rrbracket$, such that $\llbracket p \rrbracket(x_1, \dots, x_n) + |p|_1 \mathbf{d}_{\llbracket \sigma_1 \rrbracket}(x_1, y_1) + \dots + |p|_n \mathbf{d}_{\llbracket \sigma_n \rrbracket}(x_n, y_n) \geq \llbracket p \rrbracket(y_1, \dots, y_n)$.

Realisability semantics are built in a similar way. Consider an ordered R -LCA A (see Example 4.2(5)) and the associated R -graded doctrine $\langle \mathcal{R}_A, \flat \rangle$. By Lipschitz completion, we get the R -Lipschitz doctrine $D_{\langle \mathcal{R}_A, \flat \rangle} : \mathcal{L}_{\langle \mathcal{R}_A, \flat \rangle}^{\text{op}} \rightarrow \mathcal{Pos}$. An interpretation of an R -graded signature in such a doctrine is described below. The interpretation of a sort σ is an object $\llbracket \sigma \rrbracket$ in $\mathcal{L}_{\langle \mathcal{R}_A, \flat \rangle}$, which is a pair $\langle X, \rho \rangle$ such that there are four elements in $|A|$ realizing reflexivity, symmetry, transitivity and affineness of ρ . The interpretation of a function symbol f is an arrow $\llbracket f \rrbracket$ of $\mathcal{L}_{\langle \mathcal{R}_A, \flat \rangle}$, respecting the arity, such that there is an element in $|A|$ realising the condition in (2). Similarly, the interpretation of a predicate symbol p is an element in the correct fibre of $D_{\langle \mathcal{R}_A, \flat \rangle}$ such that there is in $|A|$ a realizer of the condition (3).

5 Relating notions of equality

In this section we establish a relation between the quantitative equality introduced in Section 4 and the traditional Lawvere's notion of equality as a left adjoint.

Since R -graded doctrines are in particular primary linear doctrines, one can always consider those that are elementary.

Definition 5.1. An R -graded doctrine $\langle P, \flat \rangle$ is *elementary* if P is elementary as primary linear doctrine (cf. Definition 3.3).

We have already seen in previous sections that the standard equality

predicate is replicable (cf. Corollary 3.9). In the R -graded setting, this non-linearity is even more evident as detailed by the next proposition.

Given an R -graded doctrine $\langle P, \flat \rangle$ an element α in $P(A)$ is \flat -intuitionistic if $\alpha \leq \flat_r \alpha$ for all $r \in |R|$. Notice that, by lax monoidality of \flat , we have that κ is \flat -intuitionistic and if α and β are \flat -intuitionistic, so is $\alpha * \beta$. Proposition 3.8 and Corollary 3.9 are generalised as follows.

Proposition 5.2. *Let $\langle P, \flat \rangle$ be an R -graded doctrine and $f : X \rightarrow Y$ an arrow in the base of P such that P_f has a left adjoint \mathcal{A}_f . The element $\mathcal{A}_f(\kappa_X)$ is \flat -intuitionistic.*

Proof. We have $\kappa_X \leq \flat_r \kappa_X \leq \flat_r P_f(\mathcal{A}_f(\kappa_X)) = P_f(\flat_r \mathcal{A}_f(\kappa_X))$, using naturality and lax monoidality of \flat_r . Then, the thesis follows as \mathcal{A}_f is left adjoint of P_f . \square

Corollary 5.3. *Let $\langle P, \flat \rangle$ be an elementary R -graded doctrine. Equality predicates are \flat -intuitionistic.*

Proof. Immediate after Proposition 5.2 as equality predicate is of the form $\mathcal{A}_{\Delta_A}(\kappa_A)$. \square

Elementary R -graded doctrines are the objects of the 2-category \mathbf{ELD}_R where a 1-arrow from $\langle P, \flat \rangle$ to $\langle Q, \sharp \rangle$ is a 1-arrow $\langle F, f \rangle : \langle P, \flat \rangle \rightarrow \langle Q, \sharp \rangle$ in \mathbf{LD}_R such that $\mathcal{A}_{\text{id}_{FX} \times \Delta_{FA}} \circ f_{X \times A} = f_{X \times A \times A} \circ \mathcal{A}_{\text{id}_X \times \Delta_A}$, for all objects X and A in the base of P . A 2-arrow from $\langle F, f \rangle$ to $\langle G, g \rangle$ is a 2-arrow $\theta : \langle F, f \rangle \Rightarrow \langle G, g \rangle$ in \mathbf{LD}_R . Compositions and identities are those of \mathbf{LD}_R .

Elementary R -graded doctrines are in particular R -Lipschitz, so there is an inclusion $\mathbf{I}_{\text{EL}} : \mathbf{ELD}_R \hookrightarrow \mathbf{LLD}_R$.

If $\langle P, \flat, \mathbf{d} \rangle$ is an R -Lipschitz doctrine, denote by P_{\flat} the restriction of P to \mathcal{C}_{\flat} , the full subcategory of \mathcal{C} on those objects A such that \mathbf{d}_A is \flat -intuitionistic.

Proposition 5.4. *$\langle P_{\flat}, \flat \rangle$ is an elementary R -graded doctrine.*

Proof. For α in $P(A)$, there is r such that $P_{\pi_1}(\alpha) * \flat_r \mathbf{d}_A = P_{\pi_1}(\alpha) * \mathbf{d}_A \leq P_{\pi_2}(\alpha)$. Then apply Proposition 3.4. \square

The construction of $\langle P_{\flat}, \flat \rangle$ from $\langle P, \flat \rangle$ extends also on arrows \mathbf{LLD}_R giving a 2-functor $\mathbf{R}_{\text{EL}} : \mathbf{LLD}_R \rightarrow \mathbf{ELD}_R$.

Theorem 5.5. *The 2-functor $\mathbf{I}_{\text{EL}} : \mathbf{ELD}_R \rightarrow \mathbf{LLD}_R$ is 2-comonadic with right 2-adjoint \mathbf{R}_{EL} .*

Composing Theorem 5.5 and Theorem 4.18 we get a 2-comonadic 2-adjunction between \mathbf{LD}_R and \mathbf{ELD}_R , which, by Proposition 4.3, generalises the 2-adjunction in the intuitionistic case as below:

$$\begin{array}{ccccc}
 & & \xrightarrow{\text{R}_{\text{Lip}}} & & \xrightarrow{\text{R}_{\text{EL}}} \\
 \mathbf{LD}_R & \xleftarrow{\text{T}} & \mathbf{LLD}_R & \xleftarrow{\text{T}} & \mathbf{ELD}_R \\
 \uparrow & & & \searrow & \uparrow \\
 \mathbf{PD} & \xleftarrow{\text{T}} & & & \mathbf{ED}
 \end{array}$$

6 Richer fragments of Linear Logic

Formulas in the calculus LPLL_R that we presented in Section 4.2 are written in the fragment $(!_r, \otimes, \mathbf{1}, =)$. In this section we extend LPLL_R to larger fragments of Linear Logic.

Each new connective comes with usual rules, hence we do not report them. On the other hand, we have to extend the definition of the function gr to new connectives, in order to determine resources they require to derive substitutions.

Since constants do not depend on variables, there is nothing to substitute, hence the cost to derive a substitution in a constant is zero. Thus we set

$$\text{gr}(\mathbf{0}, x) = \text{gr}(\top, x) = 0$$

A substitution in a formula built out of binary multiplicative connectives costs the sum of the resources that the two subformulas need to derive the substitution, because both of them can be subsequently used. Thus we set

$$\text{gr}(\phi \multimap \psi, x) = \text{gr}(\phi, x) + \text{gr}(\psi, x)$$

A substitution in a formula built out of quantifiers costs the same resources as the subformula, only if the substituted and the quantified variable differs, otherwise the cost is zero. Thus we set

$$\text{gr}(\forall z.\phi, x) = \text{gr}(\exists z.\phi, x) = \begin{cases} \text{gr}(\phi, x) & \text{if } x \neq z \\ 0 & \text{if } x = z \end{cases}$$

This is because, since z is bounded in $\forall z.\phi$ and $\exists z.\phi$, substitutions propagate to subformulas only if the substituted variable is different from z , otherwise $\forall z.\phi$ and $\exists z.\phi$ behave as constants.

The extension of LPLL_R by additive connectives requires additional structure on R .

A straightforward sufficient condition is that R has binary suprema denoted by \vee . Then, the cost of a substitution in a formula built out of binary additive connectives is the supremum of the resources that the two subformulas need to derive a substitution, because only one of them can be subsequently used. Thus we set

$$\mathbf{gr}(\phi \& \psi, x) = \mathbf{gr}(\phi \oplus \psi, x) = \mathbf{gr}(\phi, x) \vee \mathbf{gr}(\psi, x)$$

When all the connectives above are considered we abbreviate the resulting calculus by \mathbf{ILL}_R .

A way to obtain classical calculi is by adding a constant \perp which is dualising, namely, satisfying the entailment $(\phi \multimap \perp) \multimap \perp \vdash \phi$. Being a constant \perp is such that $\mathbf{gr}(\perp, x) = 0$. The classical version of \mathbf{ILL}_R is denoted by \mathbf{LL}_R .

It is straightforward to define R -graded doctrines that correspond to the various fragments of (classical or intuitionistic) Linear Logic. Here we consider only the following.

Definition 6.1. A *multiplicative (linear) doctrine* is a primary linear doctrine $\langle P, *, \kappa \rangle$ such that

- for every A and every α, β, γ in $P(A)$ there is $\alpha \multimap \beta$ in $P(A)$ such that $\gamma \leq \alpha \multimap \beta$ if and only if $\gamma * \alpha \leq \beta$
- for every $f : X \rightarrow A$ and every α, β in $P(A)$, it holds $P_f(\alpha \multimap \beta) = P_f(\alpha) \multimap P_f(\beta)$;

A *multiplicative-additive (linear) doctrine* is a multiplicative linear doctrine $\langle P, *, \kappa \rangle$ such that $P : \mathcal{C}^{\text{op}} \rightarrow \mathcal{Pos}$ factors through the category of lattices.

A multiplicative linear doctrine P has *quantifiers* if for every projection $\pi : A \times B \rightarrow B$ the map P_π has a left adjoint \mathcal{E}_π and a right adjoint \mathcal{V}_π and these are natural in B .

A multiplicative linear doctrine is *classical* if for each object A there is \perp_A in $P(A)$ such that $(\alpha \multimap \perp_A) \multimap \perp_A = \alpha$, which is preserved by reindexing.

A *first order linear doctrine* is a multiplicative-additive linear doctrine with quantifiers.

The syntactic doctrine associated with the $(\otimes, \mathbf{1}, \multimap)$ -fragment of \mathbf{ILL} is a multiplicative linear doctrine; the one associated with the $(\otimes, \mathbf{1}, \multimap, \oplus, \mathbf{0}, \&, \top)$ -fragment of \mathbf{ILL} is a multiplicative-additive linear doctrine; the one associated with full first order \mathbf{ILL} is a first order linear doctrine. Similarly, syntactic doctrines associated with classical variants of the above fragments provides classical versions of the corresponding doctrines.

Proposition 6.2. *Suppose $\langle P, \flat \rangle$ is an R -graded doctrine and consider its Lipschitz completion $D_{\langle P, \flat \rangle} : \mathcal{L}_{\langle P, \flat \rangle}^{\text{op}} \rightarrow \mathcal{Pos}$.*

- *If P is multiplicative, so is $D_{\langle P, \flat \rangle}$;*
- *if P is multiplicative and classical, so is $D_{\langle P, \flat \rangle}$;*
- *if P is multiplicative and has quantifier, so is $D_{\langle P, \flat \rangle}$.*

Proof. • Suppose P is multiplicative. Take $\langle A, \rho \rangle$ in $\mathcal{L}_{\langle P, \flat \rangle}$ and α, β in $\mathcal{Des}_\rho(A)$, hence there are r, s in $|R|$ such that $P_{\pi_1}(\alpha) * \flat_r \rho \leq P_{\pi_2}(\alpha)$ and $P_{\pi_1}(\beta) * \flat_s \rho \leq P_{\pi_2}(\beta)$. Then, using contraction of \flat and symmetry of ρ , we have $P_{\pi_2}(\alpha) * P_{\pi_1}(\alpha \multimap \beta) * \flat_{r+s} \rho \leq P_{\pi_2}(\alpha) * P_{\pi_1}(\alpha \multimap \beta) * \flat_r \rho * \flat_s \rho \leq P_{\pi_1}(\alpha) * P_{\pi_1}(\alpha \multimap \beta) * \flat_s \rho \leq P_{\pi_1}(\beta) * \flat_s \rho \leq P_{\pi_2}(\beta)$, showing that $P_{\pi_1}(\alpha \multimap \beta) * \flat_{r+s} \rho \leq P_{\pi_2}(\alpha \multimap \beta)$. This makes $\alpha \multimap \beta$ an element of \mathcal{Des}_ρ .

- Suppose P is multiplicative and classical and take $\langle A, \rho \rangle$ in $\mathcal{L}_{\langle P, \flat \rangle}$. By weakening of \flat , it holds $P_{\pi_1}(\perp_A) * \flat_0 \rho \leq P_{\pi_1}(\perp_A) * \kappa = P_{\pi_1}(\perp_A) = \perp_{A \times A} = P_{\pi_2}(\perp_A)$ so \perp_A is in $\mathcal{Des}_\rho(A)$.
- Suppose P is multiplicative with quantifiers and take γ in $\mathcal{Des}_{\rho \boxtimes \sigma}(A \times B)$. By naturality $P_{\pi_1}(\mathcal{A}_{\pi_1} \alpha) * \flat_r \sigma = \mathcal{A}_{\langle \pi_1, \pi_2 \rangle}(P_{\langle \pi_1, \pi_3 \rangle}(\alpha)) * \flat_r \sigma \leq \mathcal{A}_{\langle \pi_1, \pi_2 \rangle}(P_{\langle \pi_1, \pi_3 \rangle}(\alpha) * \flat_r P_{\langle \pi_1, \pi_2 \rangle}(\sigma)) \leq \mathcal{A}_{\langle \pi_1, \pi_2 \rangle} P_{\langle \pi_2, \pi_3 \rangle} \alpha = P_{\pi_2} \mathcal{A}_{\pi_1} \alpha$ for some r in R . So descent data are closed under left adjoints along projections. Similarly, for right adjoints we have $P_{\pi_1}(V_{\pi_1} \alpha) * \flat_r \sigma = V_{\langle \pi_1, \pi_2 \rangle}(P_{\langle \pi_1, \pi_3 \rangle}(\alpha)) * \flat_r \sigma \leq V_{\langle \pi_1, \pi_2 \rangle}(P_{\langle \pi_1, \pi_3 \rangle}(\alpha) * \flat_r P_{\langle \pi_1, \pi_2 \rangle}(\sigma)) \leq V_{\langle \pi_1, \pi_2 \rangle} P_{\langle \pi_2, \pi_3 \rangle} \alpha = P_{\pi_2} V_{\pi_1} \alpha$ for some r in R . \square

Proposition 6.3. *Let $\langle P, \flat \rangle$ be an R -graded doctrine where R has finite suprema and consider its Lipschitz completion $D_{\langle P, \flat \rangle} : \mathcal{L}_{\langle P, \flat \rangle}^{\text{op}} \rightarrow \mathcal{Pos}$.*

- *if P is multiplicative-additive, so is $D_{\langle P, \flat \rangle}$;*
- *if P is first order linear, so is $D_{\langle P, \flat \rangle}$.*


Proof. • Suppose P is multiplicative-additive. Take $\langle A, \rho \rangle$ in $\mathcal{L}_{\langle P, \flat \rangle}$ and α, β in $\mathcal{Des}_\rho(A)$, hence there are r, s in $|R|$ such that $P_{\pi_1}(\alpha) * \flat_r \rho \leq P_{\pi_2}(\alpha)$ and $P_{\pi_1}(\beta) * \flat_s \rho \leq P_{\pi_2}(\beta)$. Then, $P_{\pi_1}(\alpha \wedge \beta) * \flat_{r \vee s} \rho \leq (P_{\pi_1}(\alpha) * \flat_{r \vee s} \rho) \wedge (P_{\pi_1}(\beta) * \flat_{r \vee s} \rho)$. Since $\flat_{r \vee s} \rho \leq \flat_r \rho$ and $\flat_{r \vee s} \rho \leq \flat_s \rho$, it follows $P_{\pi_1}(\alpha \wedge \beta) * \flat_{r \vee s} \rho \leq (P_{\pi_1}(\alpha) * \flat_r \rho) \wedge (P_{\pi_1}(\beta) * \flat_s \rho) \leq P_{\pi_2}(\alpha) \wedge P_{\pi_2}(\beta)$, which proved $\mathcal{Des}_\rho(A)$ is closed under binary meets. By a similar argument proves the closure under binary joins. Top and bottom elements belong to $\mathcal{Des}_\rho(A)$ as one can prove using the same arguments as for \perp .

- It follows by previous item and Proposition 6.2. □

Corollary 6.4. *Let $\langle P, b \rangle$ be an R -graded doctrine where R has finite suprema. If P is classical and first order linear, so is the Lipschitz completion $D_{\langle P, b \rangle} : \mathcal{L}_{\langle P, b \rangle}^{\text{op}} \rightarrow \mathcal{Pos}$.*

Proof. Immediate by Proposition 6.2 and Proposition 6.3. □

Denote by \mathbf{FLD}_R the 2-full 2-subcategory of \mathbf{LD}_R on first order R -graded doctrines (i.e. R -graded doctrines $\langle P, b \rangle$ where P is first order linear) and 1-arrows that preserve the first order linear structure. As first order doctrines which are elementary are called hyperdoctrines, a first order R -graded doctrine which is also R -Lipschitz will be named R -quantitative hyperdoctrine. The 2-full 2-subcategory of \mathbf{LLD}_R on R -quantitative hyperdoctrines is \mathbf{QHD}_R . Proposition 6.3 shows that (if R has finite suprema) the 2-adjunction between \mathbf{LD}_R and \mathbf{LLD}_R restricts to a 2-adjunction between \mathbf{FLD}_R and \mathbf{QHD}_R . Thus, the diagram (1) in Section 2 can be refined to



Any R -quantitative hyperdoctrine (i.e. any object in \mathbf{QHD}_R) provides a domain for interpreting the \mathbf{ILL}_R calculus with quantitative equality, it suffices to extend Lemma 4.11 to all the other connectives given the definition of \mathbf{gr} at the beginning of this section. So in particular, when $R = \mathbb{R}_{\geq 0}$ as in Examples 4.2(4), the $\mathbb{R}_{\geq 0}$ -graded doctrine $\langle \mathcal{P}_{[0, \infty]}, +, 0 \rangle$ is a first order $\mathbb{R}_{\geq 0}$ -graded doctrine. Its Lipschitz completion gives a $\mathbb{R}_{\geq 0}$ -quantitative hyperdoctrine. Similarly for realisability examples. The $\mathbb{N}_{\infty}^{\leq}$ -graded doctrine $\langle \mathcal{R}_A, b \rangle$ built over an affine LCA algebra in Example 4.2(5) is first order linear. Whence its Lipschitz completion gives a $\mathbb{N}_{\infty}^{\leq}$ -quantitative hyperdoctrine.

7 Examples from quantitative equational theories and beyond

Mardare et al. [34, 35] introduced the notion of *quantitative equational theory* (QET) as a formal tool to describe and reason about quantitative algebras, that is, algebras whose carrier is a metric space and whose operations are non-expansive maps. In this section, we show how the calculus LPLL_R and its semantics can be used to reason about quantitative algebras, comparing it with QET and obtaining at the same time a range of examples of LPLL_R theories. Mardare et al. [34, 35] study QETs in great detail using the language of monads and proving many completeness results. Here we just focus on examples illustrating how LPLL_R works, leaving for future work a formal and systematic comparison between LPLL_R and QET.

We first recall the key features of a QET. Syntactically, terms are built out from a single-sorted signature Ω of possibly infinitary function symbols, we write $f : I \in \Omega$, where I is a set, when f is a function symbol in Ω whose arity is I . In order to deal with distances, the key idea is to explicitly handle quantities, by working with labelled equations $t \vDash_\epsilon u$, where ϵ is a non-negative rational number. Such an equation states that the distance between t and u is bounded by ϵ . A sequent has shape $\Gamma \vdash t \vDash_\epsilon u$, where Γ is a possibly infinite set of equations, and the calculus is defined by the following rules, where $f : I \in \Omega$ and σ a substitution.

$$\begin{array}{ll}
\text{(Ref)} & \vdash t \vDash_0 t \\
\text{(Symm)} & t \vDash_\epsilon u \vdash u \vDash_\epsilon t \\
\text{(Triang)} & \{t \vDash_\epsilon u, u \vDash_{\epsilon'} v\} \vdash t \vDash_{\epsilon+\epsilon'} v \\
\text{(Arch)} & \{t \vDash_{\epsilon'} u \mid \epsilon' > \epsilon\} \vdash t \vDash_\epsilon u \\
\text{(NExp)} & \{t_i \vDash_\epsilon u_i \mid i \in I\} \vdash f((t_i)_{i \in I}) \vDash_\epsilon f((u_i)_{i \in I}) \\
\text{(Subst)} & \Gamma \vdash t \vDash_\epsilon u \text{ implies } \sigma(\Gamma) \vdash \sigma(t) \vDash_\epsilon \sigma(u) \\
\text{(Assumpt)} & \text{if } t \vDash_\epsilon u \in \Gamma, \text{ then } \Gamma \vdash t \vDash_\epsilon u \\
\text{(Cut)} & \text{if } \Gamma \vdash \psi \text{ for all } \psi \in \Theta \text{ and } \Theta \vdash t \vDash_\epsilon u, \\
& \text{then } \Gamma \vdash t \vDash_\epsilon u
\end{array}$$

A first crucial difference between a QET and LPLL_R is that the underlying logic of the former is the usual non-linear one, as contraction and weakening are admissible: the hypotheses Γ form a set, the rule **(Cut)** is in additive form and the rule **(Assumpt)** disregards some hypotheses. Therefore, to reason quantitatively, one needs to explicitly manage quantities (as labels of equations), while in LPLL_R this does not happen because, being the underlying logic linear, it has a direct quantitative interpretation of equations. Working with explicit quantities makes it necessary to add specific rules to manage them (such as **(Arch)**) and to allow sequents with infinitely many hypotheses. Note also that infinitary sequents are needed to deal with infinitary function symbols, which are allowed in QETs, while LPLL_R only deals with finitary ones to keep sequents finitary as well.

Another important difference is that in QETs function symbols are forced to be non-expansive by the rule **(NExp)**, while in LPLL_R they can be arbitrary Lipschitz functions. Since non-expansive maps are Lipschitz with constant equal to 1, LPLL_R can treat them, but, as we will see, it can potentially go beyond this limitation, dealing with quantitative algebras where operations are arbitrary Lipschitz functions.

Remaining rules are instead similar: rules **(Refl)**, **(Symm)** and **(Triang)** above formalise the axioms of a distance and correspond to rules [R], [s] and [T] of LPLL_R , while rules **(Subst)** and **(Cut)** correspond to monotonicity of substitution and the cut rule of LPLL_R .

We now focus on examples. We start by rephrasing the example of QETs presented by Mardare et al. [34] as LPLL_R theories where R is the semiring $\mathbb{R}_{\geq 0}$ of non-negative real numbers, showing that the intended semantics of each QETs gives rise to a sound interpretation of the corresponding LPLL_R theory in the doctrine of metric spaces with Lipschitz functions. We will use the following shorter notation for functions symbols in a single-sorted graded signature: if f is a function symbol with $|f| = \langle r_1, \star \rangle \dots \langle r_n, \star \rangle \rightarrow \star$, we write $f : \langle r_1, \dots, r_n \rangle$.

We first consider the theory \mathbf{QS}_0 of quantitative semilattices with zero. To write \mathbf{QS}_0 as a QET, one takes a signature whose function symbols are $+ : 2$ and $0 : 0$, while, to write \mathbf{QS}_0 as an LPLL_R theory, one takes a $\mathbb{R}_{\geq 0}$ -graded signature with one sort, no predicate symbols and whose function symbols are $+ : \langle 1, 1 \rangle$ and $0 : \langle \rangle$, meaning that $+$ is non-expansive in both arguments. Below we list on the left the axioms of \mathbf{QS}_0 given by Mardare et al. [34] and on the right the corresponding ones in LPLL_R .

QET	LPLL _R
(S0) $\vdash x + 0 =_0 x$	$\vdash x + 0 = x$
(S1) $\vdash x + x =_0 x$	$\vdash x + x = x$
(S2) $\vdash x + y =_0 y + x$	$\vdash x + y = y + x$
(S3) $\vdash (x + y) + z =_0 x + (y + z)$	$\vdash (x + y) + z = x + (y + z)$
(S4) $\{x =_{\epsilon} x', y =_{\epsilon'} y'\} \vdash x + y =_{\epsilon \vee \epsilon'} x' + y'$	

The axiom **(S0)** says that the distance between $x + 0$ and x is less than 0. This information in LPLL_R is given forcing $\mathbf{1} \vdash x + 0 = x$, thanks to the fact that $\mathbf{1}$ is interpreted as the function constantly equal to 0. Axioms **(S1)**, **(S2)** and **(S3)** are translated similarly. Axiom **(S4)** in the QET refines the non-expansiveness of $+$ and it is needed to properly handle labels of equations. Since in LPLL_R there are no labels, this axiom is not needed, as non-expansiveness of $+$ follows from rules of LPLL_R and the graded arity in the considered signature. Note that the axioms in LPLL_R are the

usual ones of the theory of semilattices: $+$ is associative, commutative and idempotent and 0 is the neutral element.

A semantics for \mathbf{QS}_0 as QET is given by Mardare et al. [34] as follows. Let $\langle M, d \rangle$ be a metric space and consider the metric space $\langle C_d, H_d \rangle$ where C_d the set of compact subsets of M with respect to d and $H_d(A, B) = \max\{\sup_{a \in A} d(a, B), \sup_{b \in B} d(b, A)\}$, with $d(m, X) = \inf_{x \in X} d(m, x)$ for $m \in M$ and $X \subseteq M$. Then, 0 is interpreted as the empty set and $+$ as the function of type $\langle C_d, H_d \rangle \times \langle C_d, H_d \rangle \rightarrow \langle C_d, H_d \rangle$ mapping two compact sets to their union. This assignment gives rise to a semantics also of \mathbf{QS}_0 seen as an LPLL_R theory in the $\mathbb{R}_{\geq 0}$ -Lipschitz doctrine $D_{\langle \mathcal{P}_{[0, \infty]}, \mathbf{m} \rangle} : \mathcal{L}_{\langle \mathcal{P}_{[0, \infty]}, \mathbf{m} \rangle}^{\text{op}} \rightarrow \mathbf{Pos}$. We interpret a context with n variables as the product of n copies of $\langle C_d, H_d \rangle$ and a term with n free variables as a Lipschitz function from $\langle C_d, H_d \rangle^n$ to $\langle C_d, H_d \rangle$.

A difference between the semantics in the two approaches is the way one defines products of metric spaces, hence the domain of the functions interpreting terms: for QET a product has as distance the supremum of the distances of the factors, while in $\mathcal{L}_{\langle \mathcal{P}_{[0, \infty]}, \mathbf{m} \rangle}$ the distance of a product is the sum of the distances of the factors.

Other examples discussed by Mardare et al. [34] deals with distances between probability distributions. The considered theories are defined over a signature whose function symbols are $+_e : 2$ for all $e \in [0, 1]$, which can be thought of as probabilistic choice operators with probability e , that is, intuitively, $x +_e y$ is x with probability e and y with probability $1 - e$. The QET \mathbf{IBA}_p of p -Interpolative Barycentric Algebras, with $p \geq 1$, can be rephrased in LPLL_R using a single-sorted $\mathbb{R}_{\geq 0}$ -graded signature with no predicate symbols and whose function symbols are $+_e : \langle e^{1/p}, (1 - e)^{1/p} \rangle$ for all $e \in [0, 1]$, meaning that $+_e$ scales the distance of the first argument by a factor $e^{1/p}$ and that of the second argument by a factor $(1 - e)^{1/p}$. We report the axioms below.

QET	LPLL _R
(B1) $\vdash x +_1 y =_0 x$	$\vdash x +_1 y = x$
(B2) $\vdash x +_e x =_0 x$	$\vdash x +_e x = x$
(SC) $\vdash x +_e y =_0 y +_{1-e} x$	$\vdash x +_e y = y +_{1-e} x$
(SA) $\vdash (x +_e y) +_{e'} z =_0 x +_{ee'} (y +_{\frac{e'-ee'}{1-ee'}} z)$	$\vdash (x +_e y) +_{e'} z = x +_{ee'} (y +_{\frac{e'-ee'}{1-ee'}} z)$
(IB_p) $\{x =_{\epsilon_1} y, x' =_{\epsilon_2} y'\} \vdash x +_e y =_{\delta} x' +_e y'$ $(\delta \geq (e\epsilon_1^p + (1-e)\epsilon_2^p)^{1/p})$	

The axioms are translated as for the theory \mathbf{QS}_0 . The last one has no counterpart in LPLL_R as it essentially states that $+_e$ is Lipschitz with

coefficients $e^{1/p}$ and $(1 - e)^{1/p}$,¹ and this in LPLL_R follows from the the fact that $+_e$ has gaded arity $\langle e^{1/p}, (1 - e)^{1/p} \rangle$ and so we can derive the entailment $!_{e^{1/p}}x = y, !_{(1-e)^{1/p}}x' = y' \vdash x +_e y = x' +_e y'$.

Mardare et al. [34] define a semantics of this theory taking as domain of the interpretation the set $\Delta[M]$ of Borel probability measures over a metric space $\langle M, d \rangle$ with the p -Wasserstein distance W_d^p defined by

$$W_d^p(\mu, \nu) = \sup \left\{ \left| \int f^p d\mu - \int f^p d\nu \right|^{1/p} \mid f : M \rightarrow [0, \infty) \text{ is non-expansive} \right\}$$

and interpreting $+_e$ as the convex combination of two distributions, that is, the function mapping a pair of distributions $\langle \mu, \nu \rangle$ to $e\mu + (1 - e)\nu$. This choice, induces an interpretation of the corresponding LPLL_R theory in $D_{\langle \mathcal{P}_{[0, \infty)}, \mathbf{m} \rangle} : \mathcal{L}_{\langle \mathcal{P}_{[0, \infty)}, \mathbf{m} \rangle}^{\text{op}} \rightarrow \mathcal{Pos}$, as it is easy to see that the convex combination operation has Lipschitz constants $e^{1/p}$ and $(1 - e)^{1/p}$ with respect to W_d^p .

A specialised version the of theory \mathbf{IBA}_p is the theory \mathbf{BA} of Left-Invariant Barynetric Algebras, which is obtained by replacing the axioms (\mathbf{IB}_P) by the axiom (\mathbf{LI}) stating $\vdash x +_e z =_{\epsilon} y +_e z$ where $\epsilon \geq e$. This axiom, together with the others, basically requires that Lipschitz constants of $+_e$ are e and $1 - e$, but it also forces the distance to be smaller than 1. To rephrase this theory in LPLL_R , we need to consider a single-sorted $\mathbb{R}_{\geq 0}$ -graded signature where function symbols are $+_e : \langle e, 1 - e \rangle$, for all $e \in [0, 1]$, and having a constant predicate symbol $u : \langle \rangle$ modelling the real number 1. This matches the intuition that in our setting real numbers are “truth values”, hence they can be represented syntactically by constant predicate symbols. Finally, we have to add to the theory above the axiom $u \vdash x = y$, stating that the distance is smaller than 1. Note that, using the rules of LPLL_R , we can prove $!_{\epsilon}u \vdash x = x +_e z = y +_e z$, for all $\epsilon \geq e$, which corresponds exactly to the axiom (\mathbf{LI}) .

Following again Mardare et al. [34], a semantics of the theory \mathbf{BA} in the Lipschitz doctrine $D_{\langle \mathcal{P}_{[0, \infty)}, \mathbf{m} \rangle}$ can be constructed by taking as interpretation of the sort the set $\Delta[S, \Sigma]$ of probability measures over a measurable space $\langle S, \Sigma \rangle$, with the total variation distance $T(\mu, \nu) = \sup_{E \in \Sigma} |\mu(E) - \nu(E)|$, and interpreting $+_e$ again as the convex combination of distributions.

Mardare et al. [34, 35] want function symbols to be interpreted by non-expansive maps and in all the above examples this is possible because the

¹To check this we rely on the inequality $(a + b)^{1/p} \leq a^{1/p} + b^{1/p}$, which holds for non-negative real numbers a, b .

Lipschitz constants of these function symbols are smaller than 1. We now consider an example where function symbols are not non-expansive: the theory \mathbb{R} -**MVS** of metric vector spaces over the field \mathbb{R} of real numbers. The single-sorted $\mathbb{R}_{\geq 0}$ -graded signature has no predicate symbols and consists of function symbols $+$: $\langle 1, 1 \rangle$, $-$: $\langle 1 \rangle$ and 0 : $\langle \rangle$ for the group structure and λ_a : $\langle |a| \rangle$ for scalar multiplication by a , for all $a \in \mathbb{R}$. Note that the scalar multiplication λ_a is required to scale the distance by a factor $|a|$, hence in general is not non-expansive; indeed, the entailment $!_{|a|}x = y \vdash \lambda_a(x) = \lambda_a(y)$ is provable. Below we report the axioms:

$$\begin{array}{ll}
\text{(G1)} \quad \vdash (x + y) + z = x + (y + z) & \text{(L1)} \quad \vdash \lambda_a(x + y) = \lambda_a(x) + \lambda_a(y) \\
\text{(G2)} \quad \vdash x + y = y + x & \text{(L2)} \quad \vdash \lambda_{a+b}(x) = \lambda_a(x) + \lambda_b(x) \\
\text{(G3)} \quad \vdash x + 0 = x & \text{(L3)} \quad \vdash \lambda_{ab}(x) = \lambda_a(\lambda_b(x)) \\
\text{(G4)} \quad \vdash x + (-x) = 0 & \text{(L4)} \quad \vdash \lambda_1(x) = x
\end{array}$$

These axioms are actually the standard ones for vector spaces, the only difference is that equality is interpreted as a distance. For instance, a semantics for this theory can be given by the n -dimensional real vector space \mathbb{R}^n with the Euclidean distance, as it is easy to see that scalar multiplication by a is a Lipschitz function with constant $|a|$. More generally, if V is a real vector space with a norm $\|-\|$, the distance $d(u, v) = \|u - v\|$ makes the group operations non-expansive and the scalar multiplication by a a Lipschitz function with constant $|a|$ as $\|au - av\| = |a| \cdot \|u - v\|$ holds.

This example shows that LPLL_R provide theories for quantitative algebras where operations are arbitrary Lipschitz functions which are not necessarily non-expansive.

8 Related and future works

A work on equality in non-modal Linear Logic is by Dozen [18], where rules for equality are those in Example 3.5(1). The author proves that these rules lead to affineness and replicability of equality using a syntactic approach. In the same year, a work in full Linear Logic appears by Coniglio and Miraglia [12], where equality predicate is intuitionistic in the sense that $x = y \vdash !(x = y)$ holds, hence it is again affine and replicable. To recover linearity, Dozen [18] restricts the substitution rule in such a way that, given an equality $t = u$ and a formula ϕ , one can replace exactly one occurrence of t in ϕ . This approach is a specific instance of our calculus, when the semiring is $\mathbb{N}^=$ and all symbols have grade equal to 1.

In Section 7 we have compared by examples our approach with Quantitative Equational Theories by Mardare et al. [34, 35]. In future work, we

plan to deepen this comparison. Notably, Mardare et al. [34, 35] prove that every QET \mathbf{U} determines a monad on the category of metric spaces and non-expansive functions, and semantics for \mathbf{U} are exactly the algebras for such a monad. This is a strong result that attests that \mathbf{U} axiomatise the algebras of the monad induced by itself. We could state a similar result in our setting, showing that an equational theory in LPLL_R induces a monad on categories of the form $\mathcal{L}_{\langle P, \flat \rangle}$ and proving that the category of models for such a theory is equivalent to the category of algebras for the induced monad. This would show that equational theories in LPLL_R axiomatise quantitative algebras where operations are Lipschitz maps.

Another potential application of Lipschitz doctrines is in the definition of denotational models for various forms of what we could call *graded λ -calculi*, see, *e.g.*, works by Petricek et al. [38], Abel and Bernardy [1]. The key feature of these calculi is that variables in typing contexts are annotated by grades from a semiring, describing how the term uses that variable. Accordingly, arrow types are labelled by a grade describing how the function uses its parameter. These data can be interpreted in (the base category of) a Lipschitz doctrine: types are objects, contexts are just the product of their types and terms are arrows whose Lipschitz constants agree with grades reported in the context. Moreover, to interpret the labelled arrow type, it is possible, exploiting the structure of a Lipschitz doctrine, to define a notion of *graded exponential*, which captures all Lipschitz arrows between two objects having some fixed Lipschitz constant.

Last but not least, it would be interesting to move the ideas discussed here to a proof-relevant setting, studying connections with type theories. It is known that every type theory gives rise to a doctrine $Ty : \mathbf{Ctx}^{\text{op}} \rightarrow \mathbf{Pos}$ where \mathbf{Ctx}_T denotes the category of contexts and Ty maps a context Γ to the (poset reflection of the) preorder whose elements are types depending on Γ and where $A \leq B$ over Γ if there is a term $\Gamma \vdash p : A \rightarrow B$. The action of Ty on arrows is given by substitution. Linear dependent type theories, where linear and intuitionistic types are kept separated in contexts, originate from the work of Cervesato and Pfenning [10] and they give rise to a doctrine $Ty : \mathbf{Ctx}^{\text{op}} \rightarrow \mathbf{Pos}$ as before, where \mathbf{Ctx} contains intuitionistic context and $A \leq B$ if there is a term $p : A \multimap B$. Studies of Id-Types in linear type theories are available in the literature, but to the best of our knowledge, they treat Id-Types as intuitionistic types (see Krishnaswami et al. [27] and Vákár [42]). There are also dependent type theories using grading, *e.g.*, by Atkey [4] and Moon et al. [36], but they do not consider Id-Types. Therefore, a direction that we will explore is to use the machinery introduced in this paper, possibly extended to indexed categories and fibrations, to

study quantitative Id-Types in linear type theories.

Doctrines (equivalent to those) of the form $Ty : \mathbf{Ctx}^{\text{op}} \rightarrow \mathbf{Pos}$ are presented by Maietti and Rosolini [32, 33], where the authors observed that the construction $\mathbf{PD} \rightarrow \mathbf{ED}$, when applied to Ty , can be used to generate setoids models à la Bishop. Similarly, if we move to the linear setting, we suspect that $\mathbf{RLip} : \mathbf{LD}_R \rightarrow \mathbf{LLD}_R$ can be applied to linear type theories (rearranged as doctrines of the form Ty) to give categories of setoids having a linear and quantitative dress.

References

- [1] Andreas Abel and Jean-Philippe Bernardy. A unified view of modalities in type systems. *Proceedings of ACM on Programming Languages*, 4 (ICFP):90:1–90:28, 2020. doi: 10.1145/3408972.
- [2] Samson Abramsky. A structural approach to reversible computation, 2011. URL <http://arxiv.org/abs/1111.7154>.
- [3] Samson Abramsky, Esfandiar Haghverdi, and Philip J. Scott. Geometry of interaction and linear combinatory algebras. *Mathematical Structures in Computer Science*, 12(5):625–665, 2002. doi: 10.1017/S0960129502003730.
- [4] Robert Atkey. Syntax and semantics of quantitative type theory. In Anuj Dawar and Erich Grädel, editors, *Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2018*, pages 56–65. ACM, 2018. doi: 10.1145/3209108.3209189.
- [5] Christel Baier and Mila E. Majster-Cederbaum. Denotational semantics in the CPO and metric approach. *Theoretical Computer Science*, 135 (2):171–220, 1994. doi: 10.1016/0304-3975(94)00046-8.
- [6] Gilles Barthe, Boris Köpf, Federico Olmedo, and Santiago Zanella Béguelin. Probabilistic relational reasoning for differential privacy. *ACM Transactions on Programming Languages and Systems*, 35(3):9:1–9:49, 2013. doi: 10.1145/2492061.
- [7] Gilles Barthe, Marco Gaboardi, Justin Hsu, and Benjamin C. Pierce. Programming language techniques for differential privacy. *ACM SIGLOG News*, 3(1):34–53, 2016.

- [8] Flavien Breuvert and Michele Pagani. Modelling coeffects in the relational semantics of linear logic. In Stephan Kreutzer, editor, *24th EACSL Annual Conference on Computer Science Logic, CSL 2015*, volume 41 of *LIPICs*, pages 567–581. Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2015. doi: 10.4230/LIPICs.CSL.2015.567.
- [9] Aloïs Brunel, Marco Gaboardi, Damiano Mazza, and Steve Zdancewic. A core quantitative coeffect calculus. In Zhong Shao, editor, *Programming Languages and Systems - 23rd European Symposium on Programming, ESOP 2014*, volume 8410 of *Lecture Notes in Computer Science*, pages 351–370. Springer, 2014. doi: 10.1007/978-3-642-54833-8_19.
- [10] Iliano Cervesato and Frank Pfenning. A linear logical framework. *Inform. and Comput.*, 179(1):19–75, 2002. doi: 10.1006/inco.2001.2951.
- [11] Konstantinos Chatzikokolakis, Daniel Gebler, Catuscia Palamidessi, and Lili Xu. Generalized bisimulation metrics. In Paolo Baldan and Daniele Gorla, editors, *Concurrency Theory - 25th International Conference, CONCUR 2014*, volume 8704 of *Lecture Notes in Computer Science*, pages 32–46. Springer, 2014. doi: 10.1007/978-3-662-44584-6_4.
- [12] Marcelo E. Coniglio and Francisco Miraglia. Equality in linear logic. *Logique et Analyse*, 39(153/154):113–151, 1996. ISSN 00245836, 22955836. URL <http://www.jstor.org/stable/44084560>.
- [13] Francesco Dagnino and Giuseppe Rosolini. Doctrines, modalities and comonads. *Mathematical Structures in Computer Science*, page 1–30, 2021. doi: 10.1017/S0960129521000207.
- [14] Ugo Dal Lago and Francesco Gavazzo. Modal reasoning = metric reasoning, via lawvere, 2021. URL <https://arxiv.org/abs/2103.03871>.
- [15] Ugo Dal Lago, Francesco Gavazzo, and Akira Yoshimizu. Differential logical relations, part I: the simply-typed case. In Christel Baier, Ioannis Chatzigiannakis, Paola Flocchini, and Stefano Leonardi, editors, *46th International Colloquium on Automata, Languages, and Programming, ICALP 2019*, volume 132 of *LIPICs*, pages 111:1–111:14. Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2019. doi: 10.4230/LIPICs.ICALP.2019.111.

- [16] Arthur Azevedo de Amorim, Marco Gaboardi, Justin Hsu, Shin-ya Katsumata, and Ikram Cherigui. A semantic account of metric preservation. In Giuseppe Castagna and Andrew D. Gordon, editors, *Proceedings of the 44th Annual ACM Symposium on Principles of Programming Languages, POPL'17*, pages 545–556. ACM, 2017.
- [17] Josée Desharnais, Vineet Gupta, Radha Jagadeesan, and Prakash Panangaden. Metrics for labelled markov processes. *Theoretical Computer Science*, 318(3):323–354, 2004. doi: 10.1016/j.tcs.2003.09.013.
- [18] Kosta Dozen. Equality in substructural logics. In W. Hodges, M. Hyland, C. Steinhorn, and J. K. Truss, editors, *Logic: from foundations to applications: European Logic Colloquium*, 1996.
- [19] Jacopo Emmenegger, Fabio Pasquali, and Giuseppe Rosolini. Elementary doctrines as coalgebras. *Journal of Pure and Applied Algebra*, 224(12):106445, 2020. doi: <https://doi.org/10.1016/j.jpaa.2020.106445>.
- [20] Marco Gaboardi, Shin-ya Katsumata, Dominic A. Orchard, Flavien Breuvert, and Tarmo Uustalu. Combining effects and coeffects via grading. In Jacques Garrigue, Gabriele Keller, and Eijiro Sumii, editors, *Proceedings of the 21st ACM International Conference on Functional Programming, ICFP 2016*, pages 476–489. ACM, 2016. doi: 10.1145/2951913.2951939.
- [21] Francesco Gavazzo. Quantitative behavioural reasoning for higher-order effectful programs: Applicative distances. In Anuj Dawar and Erich Grädel, editors, *Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2018*, pages 452–461. ACM, 2018. doi: 10.1145/3209108.3209149.
- [22] Jean-Yves Girard. Linear logic. *Theoretical Computer Science*, 50: 1–102, 1987. doi: 10.1016/0304-3975(87)90045-4.
- [23] Jean-Yves Girard, Andre Scedrov, and Philip J. Scott. Bounded linear logic: A modular approach to polynomial-time computability. *Theoretical Computer Science*, 97(1):1–66, 1992. doi: 10.1016/0304-3975(92)90386-T.
- [24] Naohiko Hoshino. Linear realizability. In Jacques Duparc and Thomas A. Henzinger, editors, *Computer Science Logic*, pages 420–434, Berlin, Heidelberg, 2007. Springer Berlin Heidelberg. ISBN 978-3-540-74915-8.

- [25] Bart P. F. Jacobs. *Categorical Logic and Type Theory*, volume 141 of *Studies in logic and the foundations of mathematics*. North-Holland, 2001. ISBN 978-0-444-50853-9.
- [26] Shin-ya Katsumata. A double category theoretic analysis of graded linear exponential comonads. In Christel Baier and Ugo Dal Lago, editors, *Foundations of Software Science and Computation Structures - 21st International Conference, FOSSACS 2018*, volume 10803 of *Lecture Notes in Computer Science*, pages 110–127. Springer, 2018. doi: 10.1007/978-3-319-89366-2_6.
- [27] Neelakantan R. Krishnaswami, Pierre Pradic, and Nick Benton. Integrating linear and dependent types. In Sriram K. Rajamani and David Walker, editors, *Proceedings of the 42nd Annual ACM Symposium on Principles of Programming Languages, POPL’15*, pages 17–30. ACM, 2015. doi: 10.1145/2676726.2676969.
- [28] F. William Lawvere. Adjointness in foundations. *Dialectica*, 23:281–296, 1969.
- [29] F. William Lawvere. Equality in hyperdoctrines and comprehension schema as an adjoint functor. In A. Heller, editor, *Proc. New York Symposium on Application of Categorical Algebra*, pages 1–14. Amer.Math.Soc., 1970.
- [30] F. William Lawvere. Metric spaces, generalized logic, and closed categories. *Rend. Sem. Mat. Fis. Milano*, 43:135–166, 1973.
- [31] Maria Emilia Maietti and Giuseppe Rosolini. Elementary quotient completion. *Theory Appl. Categ.*, 27(17):445–463, 2013.
- [32] Maria Emilia Maietti and Giuseppe Rosolini. Quotient completion for the foundation of constructive mathematics. *Log. Univers.*, 7(3):371–402, 2013. doi: 10.1007/s11787-013-0080-2.
- [33] Maria Emilia Maietti and Giuseppe Rosolini. Unifying exact completions. *Appl. Categ. Structures*, 23(1):43–52, 2015. doi: 10.1007/s10485-013-9360-5.
- [34] Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. Quantitative algebraic reasoning. In Martin Grohe, Eric Koskinen, and Natarajan Shankar, editors, *Proceedings of the 31st Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2016*, pages 700–709. ACM, 2016. doi: 10.1145/2933575.2934518.

- [35] Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. On the axiomatizability of quantitative algebras. In *Proceedings of the 32nd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2017*, pages 1–12. IEEE Computer Society, 2017. doi: 10.1109/LICS.2017.8005102.
- [36] Benjamin Moon, Harley Eades III, and Dominic Orchard. Graded modal dependent type theory. In Nobuko Yoshida, editor, *Programming Languages and Systems - 30th European Symposium on Programming, ESOP 2021*, volume 12648 of *Lecture Notes in Computer Science*, pages 462–490. Springer, 2021. doi: 10.1007/978-3-030-72019-3_17.
- [37] Dominic Orchard, Vilem-Benjamin Liepelt, and Harley Eades III. Quantitative program reasoning with graded modal types. *Proceedings of ACM on Programming Languages*, 3(ICFP):110:1–110:30, 2019. doi: 10.1145/3341714.
- [38] Tomas Petricek, Dominic A. Orchard, and Alan Mycroft. Coeffects: a calculus of context-dependent computation. In Johan Jeuring and Manuel M. T. Chakravarty, editors, *Proceedings of the 19th ACM International Conference on Functional programming, ICFP 2014*, pages 123–135. ACM, 2014. doi: 10.1145/2628136.2628160.
- [39] Andrew M. Pitts. Categorical logic. In *Handbook of logic in computer science, Vol. 5*, volume 5 of *Handbook of Logic in Computer Science*, pages 39–128. Oxford Univ. Press, New York, 2000.
- [40] Jason Reed and Benjamin C. Pierce. Distance makes the types grow stronger: a calculus for differential privacy. In Paul Hudak and Stephanie Weirich, editors, *Proceedings of the 15th ACM International Conference on Functional programming, ICFP 2010*, pages 157–168. ACM, 2010. doi: 10.1145/1863543.1863568.
- [41] Robert Seely. Linear logic, *-autonomous categories and cofree coalgebras. *Contemporary Mathematics*, 92, 1989.
- [42] Matthijs Vákár. A categorical semantics for linear logical frameworks. In Andrew M. Pitts, editor, *Foundations of Software Science and Computation Structures - 18th International Conference, FoSSaCS 2015*, volume 9034 of *Lecture Notes in Computer Science*, pages 102–116. Springer, 2015. doi: 10.1007/978-3-662-46678-0_7.

- [43] Franck van Breugel and James Worrell. A behavioural pseudometric for probabilistic transition systems. *Theoretical Computer Science*, 331 (1):115–142, 2005. doi: 10.1016/j.tcs.2004.09.035.
- [44] Jap van Oosten. *Realizability: An Introduction to its Categorical Side*, volume 152 of *Studies in Logic and the Foundations of Mathematics*. North Holland Publishing Company, 2008.
- [45] Lili Xu, Konstantinos Chatzikokolakis, and Huimin Lin. Metrics for differential privacy in concurrent systems. In Erika Ábrahám and Catuscia Palamidessi, editors, *Formal Techniques for Distributed Objects, Components, and Systems - 34th IFIP International Conference, FORTE 2014*, volume 8461 of *Lecture Notes in Computer Science*, pages 199–215. Springer, 2014. doi: 10.1007/978-3-662-43613-4_13.

A Proofs

A.1 Proofs of Section 3

Proposition 3.4. The proof of (i) \Rightarrow (ii) is a direct rephrasing of part of Proposition 2.5 in [19] where replicability and affineness of δ is not used. To show (ii) \Rightarrow (iii) take inequality (ii)-(c) and by replace $\delta_{X \times A}$ with its expression in (ii)-(b). Reindexing both side of the inequality along $\langle \pi_1, \pi_2, \pi_1, \pi_3 \rangle : X \times A \times A \rightarrow X \times A \times X \times A$ gives (iii)-(b). To show (iii) \Rightarrow (i) take β in $P(X \times A \times A)$ and suppose $P_{\langle \pi_1, \pi_2 \rangle}(\alpha) * P_{\langle \pi_2, \pi_3 \rangle}(\delta_A) \leq \beta$, reindexing along $\text{id}_X \times \Delta_A$ gives $\alpha \leq P_{\text{id}_X \times \Delta_A}(\beta)$. Conversely if $\alpha \leq P_{\text{id}_X \times \Delta_A}(\beta)$, then $P_{\langle \pi_1, \pi_2 \rangle}(\alpha) * P_{\langle \pi_2, \pi_3 \rangle}(\delta_A) \leq P_{\langle \pi_1, \pi_2, \pi_2 \rangle}(\alpha) * P_{\langle \pi_2, \pi_3 \rangle}(\delta_A)$. By (iii)-(b), seeing β as an element in $(X \times A) \times A$ one gets $P_{\langle \pi_1, \pi_2 \rangle}(\alpha) * P_{\langle \pi_2, \pi_3 \rangle}(\delta_A) \leq \beta$, establishing the assignments that maps α to $P_{\langle \pi_1, \pi_2 \rangle}(\alpha) * P_{\langle \pi_2, \pi_3 \rangle}(\delta_A)$ as a left adjoint to $P_{\langle \text{id}_X \times \Delta_A \rangle}$. The Beck-Chevalley condition and the Frobenius reciprocity are immediate. \square

Proposition 3.7. Equality predicate are reflexive. Taking A as X and δ_A as α in Proposition 3.4 (iii)-(b) one gets transitivity. While taking A as X and $P_{\text{sw}}(\delta_A)$ as α one has $P_{\langle \pi_1, \pi_2 \rangle} P_{\text{sw}}(\delta_A) * P_{\langle \pi_2, \pi_3 \rangle}(\delta_A) \leq P_{\langle \pi_1, \pi_3 \rangle} P_{\text{sw}}(\delta_A)$. Reindexing along $\Delta_A \times \text{id}_A : A \times A \rightarrow A \times A \times A$ gives $P_{\pi_1} P_{\Delta_A}(\delta_A) * \delta_A \leq P_{\text{sw}}(\delta_A)$. Symmetry follows as δ_A is reflexive. \square

A.2 Proofs of Section 4.3

Lemma 4.11. 1) is proved by induction on $t : \sigma$. If $t : \sigma$ is a variable in $\vec{\sigma}$, i.e. $t : \sigma$ is $x_p : \sigma_p$ for $1 \leq p \leq n$, then $\llbracket t \rrbracket$ is the projection $\pi_p : \llbracket \sigma_1 \rrbracket \times \dots \times \llbracket \sigma_n \rrbracket \rightarrow \llbracket \sigma_p \rrbracket$ and $\text{gr}(t, x_j) = 1$ if $p = j$ and is 0 if $p \neq j$, so $\flat_{\text{gr}(t, x_j)} \mathbf{d}_{\llbracket \sigma_j \rrbracket} \leq \kappa$ (for $j \neq p$ by weakening of \flat) and $\flat_{\text{gr}(t, x_p)} \mathbf{d}_{\llbracket \sigma_p \rrbracket} \leq \mathbf{d}_{\llbracket \sigma_p \rrbracket}$ (by counit of \flat), so

$$\prod_{i=1}^n P_{\langle \pi_i, \pi_{n+i} \rangle} (\flat_{\text{gr}(t, x_i)} \mathbf{d}_{\llbracket \sigma_i \rrbracket}) \leq P_{\langle \pi_p, \pi_{n+p} \rangle} (\flat_{\text{gr}(t, x_p)} \mathbf{d}_{\llbracket \sigma_i \rrbracket})$$

whence the claim as $\langle \pi_p, \pi_{n+p} \rangle = \pi_p \times \pi_p = \llbracket t \rrbracket \times \llbracket t \rrbracket$.

Suppose $t : \tau$ is of the form $f(t_1, \dots, t_m)$ where f is a function symbol of arity $\tau_1, \dots, \tau_m \rightarrow \tau$, and each t_k is a term of type τ_k in the context $\sigma_1, \dots, \sigma_n$, satisfying the inductive hypothesis, i.e.

$$\prod_{i=1}^n P_{\langle \pi_i, \pi_{n+i} \rangle} (\flat_{\text{gr}(t_k, x_i)} \mathbf{d}_{\llbracket \sigma_i \rrbracket}) \leq P_{\llbracket t_k \rrbracket \times \llbracket t_k \rrbracket} (\mathbf{d}_{\llbracket \tau_k \rrbracket})$$

Multiply both sides by $\flat_{|f|_k}$ and use comultiplication of \flat to get

$$\prod_{i=1}^n P_{\langle \pi_i, \pi_{n+i} \rangle} (\flat_{|f|_k} \text{gr}(t_k, x_i) \mathbf{d}_{\llbracket \sigma_i \rrbracket}) \leq P_{\llbracket t_k \rrbracket \times \llbracket t_k \rrbracket} (\flat_{|f|_k} \mathbf{d}_{\llbracket \tau_k \rrbracket})$$

From $\llbracket t_k \rrbracket \times \llbracket t_k \rrbracket = \langle \pi_k, \pi_{m+k} \rangle \circ (\langle \llbracket t_1 \rrbracket, \dots, \llbracket t_m \rrbracket \rrbracket \times \langle \llbracket t_1 \rrbracket, \dots, \llbracket t_m \rrbracket \rrbracket)$ one has

$$P_{\llbracket t_k \rrbracket \times \llbracket t_k \rrbracket} (\flat_{|f|_k} \mathbf{d}_{\llbracket \tau_k \rrbracket}) = P_{\langle \llbracket t_1 \rrbracket, \dots, \llbracket t_m \rrbracket \rrbracket \times \langle \llbracket t_1 \rrbracket, \dots, \llbracket t_m \rrbracket \rrbracket} P_{\langle \pi_k, \pi_{m+k} \rangle} (\flat_{|f|_k} \mathbf{d}_{\llbracket \tau_k \rrbracket})$$

The way in which the interpretation of an R -graded signature is defined ensures the satisfaction of inequality (2) for f .

$$\prod_{k=1}^m P_{\langle \pi_k, \pi_{m+k} \rangle} (\flat_{|f|_k} \mathbf{d}_{\llbracket \tau_k \rrbracket}) \leq P_{\llbracket f \rrbracket \times \llbracket f \rrbracket} (\mathbf{d}_{\llbracket \tau \rrbracket})$$

Since $\llbracket t \rrbracket = \llbracket f(t_0, \dots, t_m) \rrbracket = \llbracket f \rrbracket \circ \langle \llbracket t_1 \rrbracket, \dots, \llbracket t_m \rrbracket \rrbracket$, evaluating both side of the inequality along $\langle \llbracket t_1 \rrbracket, \dots, \llbracket t_m \rrbracket \rrbracket \times \langle \llbracket t_1 \rrbracket, \dots, \llbracket t_m \rrbracket \rrbracket$ one has

$$\prod_{k=1}^m P_{\llbracket t_k \rrbracket \times \llbracket t_k \rrbracket} (\flat_{|f|_k} \mathbf{d}_{\llbracket \tau_k \rrbracket}) \leq P_{\llbracket t \rrbracket \times \llbracket t \rrbracket} (\mathbf{d}_{\llbracket \tau \rrbracket})$$

whence

$$\prod_{k=1}^m \prod_{i=1}^n P_{\langle \pi_i, \pi_{n+i} \rangle} (\flat_{|f|_k} \text{gr}(t_k, x_i) \mathbf{d}_{\llbracket \sigma_i \rrbracket}) \leq P_{\llbracket t \rrbracket \times \llbracket t \rrbracket} (\mathbf{d}_{\llbracket \tau \rrbracket})$$

finally recall that $\text{gr}(t, x_i) = |f|_1 \cdot \text{gr}(t_1, x_i) + \dots + |f|_m \cdot \text{gr}(t_m, x_i)$. Contraction of \flat completes the proof.

2) If ϕ is $p(t_1, \dots, t_m)$ where p predicate symbol of L of arity τ_1, \dots, τ_m and t_i is a term of L of arity $\sigma_1, \dots, \sigma_n \rightarrow \tau_i$, then $\text{gr}(p, x_i) = |p|_1 \cdot \text{gr}(t_1, x_i) + \dots + |p|_n \cdot \text{gr}(t_n, x_i)$. Condition displayed in (3) together with contraction of \flat gives

$$P_{\text{pr}_1}(\llbracket p \rrbracket) * \prod_{i=1}^n \flat_{\text{gr}(\phi, x_i)} P_{\langle \pi_i, \pi_{n+i} \rangle}(\mathbf{d}_{\llbracket \sigma_i \rrbracket}) \leq P_{\text{pr}_2}(\llbracket p \rrbracket)$$

where $\text{pr}_1 = \langle \pi_1, \dots, \pi_m \rangle$ and $\text{pr}_2 = \langle \pi_{m+1}, \dots, \pi_{2m} \rangle$.

The reindexing of both sides along $\langle \llbracket t_1 \rrbracket, \dots, \llbracket t_m \rrbracket \rangle$ yields that the right hand side, that becomes $P_{\langle \pi_{n+1}, \dots, \pi_{2n} \rangle}(\llbracket \phi \rrbracket)$, is greater or equal to the left hand side, that becomes

$$P_{\langle \pi_1, \dots, \pi_n \rangle}(\llbracket \phi \rrbracket) * \prod_{k=1}^m P_{\langle \pi_k, \pi_{n+k} \rangle}(\flat_{\text{gr}(p, x_k)} P_{\llbracket t_k \rrbracket \times \llbracket t_k \rrbracket} \mathbf{d}_{\llbracket \tau_k \rrbracket})$$

By point 1) of this proposition, the displayed formula is greater than or equal to

$$P_{\langle \pi_1, \dots, \pi_n \rangle}(\llbracket \phi \rrbracket) * \prod_{k=1}^m \prod_{i=1}^n P_{\langle \pi_1, \pi_{n+1} \rangle}(\flat_{\text{gr}(p, x_k) \cdot \text{gr}(t_k, x_i)} \mathbf{d}_{\llbracket \sigma_i \rrbracket})$$

recall that $\text{gr}(\phi, x_i) = |p|_1 \cdot \text{gr}(t_1, x_i) + \dots + |p|_m \cdot \text{gr}(t_m, x_i)$, so contraction of \flat leads to the claim.

If ϕ is $t \equiv_{\tau} u$ where t, u are terms of arity $\sigma_1, \dots, \sigma_n \rightarrow \tau$, then $\llbracket \phi \rrbracket = P_{\langle \llbracket t \rrbracket, \llbracket u \rrbracket \rangle}(\mathbf{d}_{\llbracket \tau \rrbracket})$ and $\text{gr}(\phi, x_i) = \text{gr}(t, x_i) + \text{gr}(u, x_i)$, so by of \flat and conditions on t and u it holds that

$$\begin{aligned} & P_{\langle \pi_1, \dots, \pi_n \rangle} P_{\langle \llbracket t \rrbracket, \llbracket u \rrbracket \rangle}(\mathbf{d}_{\llbracket \tau \rrbracket}) * \prod_{i=1}^n P_{\langle \pi_i, \pi_{n+i} \rangle}(\flat_{\text{gr}(\phi, x_i)} \mathbf{d}_{\llbracket \sigma_i \rrbracket}) \\ & \leq P_{\langle \pi_1, \dots, \pi_n \rangle} P_{\langle \llbracket t \rrbracket, \llbracket u \rrbracket \rangle}(\mathbf{d}_{\llbracket \tau \rrbracket}) * P_{\llbracket t \rrbracket \times \llbracket t \rrbracket}(\mathbf{d}_{\llbracket \tau \rrbracket}) * P_{\llbracket u \rrbracket \times \llbracket u \rrbracket}(\mathbf{d}_{\llbracket \tau \rrbracket}) \\ & \leq P_{\langle \pi_{n+1}, \dots, \pi_{2n} \rangle} P_{\langle \llbracket t \rrbracket, \llbracket u \rrbracket \rangle}(\mathbf{d}_{\llbracket \tau \rrbracket}) \end{aligned}$$

by transitivity of $\mathbf{d}_{\llbracket \tau \rrbracket}$.

If ϕ is $\psi \otimes \theta$ where both ψ and θ are well formed formulas in the context $\vec{\sigma}$ satisfying the inductive hypothesis, then $\llbracket \phi \rrbracket = \llbracket \psi \rrbracket * \llbracket \theta \rrbracket$ and $\text{gr}(\phi, x_i) =$

$\text{gr}(\psi, x_i) + \text{gr}(\theta, x_i)$. By contraction of \flat , it follows

$$\begin{aligned}
& P_{\langle \pi_1, \dots, \pi_n \rangle}(\llbracket \psi \rrbracket * \llbracket \theta \rrbracket) * \prod_{i=1}^n P_{\langle \pi_i, \pi_{n+i} \rangle}(\flat_{\text{gr}(\psi, x_i) + \text{gr}(\theta, x_i)} \mathbf{d}_{\llbracket \sigma_i \rrbracket}) \\
& \leq P_{\langle \pi_1, \dots, \pi_n \rangle}(\llbracket \psi \rrbracket * \llbracket \theta \rrbracket) * \prod_{i=1}^n P_{\langle \pi_i, \pi_{n+i} \rangle}(\flat_{\text{gr}(\psi, x_i)} \mathbf{d}_{\llbracket \sigma_i \rrbracket} * \flat_{\text{gr}(\theta, x_i)} \mathbf{d}_{\llbracket \sigma_i \rrbracket}) \\
& = P_{\langle \pi_1, \dots, \pi_n \rangle}(\llbracket \psi \rrbracket) * \prod_{i=1}^n P_{\langle \pi_i, \pi_{n+i} \rangle}(\flat_{\text{gr}(\psi, x_i)} \mathbf{d}_{\llbracket \sigma_i \rrbracket}) * P_{\langle \pi_1, \dots, \pi_n \rangle}(\llbracket \theta \rrbracket) * \prod_{i=1}^n P_{\langle \pi_i, \pi_{n+i} \rangle}(\flat_{\text{gr}(\theta, x_i)} \mathbf{d}_{\llbracket \sigma_i \rrbracket})
\end{aligned}$$

By hypothesis on ψ the first formulas in the square brackets is less than or equal to $P_{\langle \pi_{n+1}, \dots, \pi_{2n} \rangle}(\llbracket \psi \rrbracket)$ and the same for θ , whence the claim.

If ϕ is $\mathbf{1}$ then $\text{gr}(\phi, x_i) = 0$. Then $\llbracket \mathbf{1} \rrbracket = \kappa$ and the claim is trivial by weakening fo \flat .

If ϕ is $!_r \psi$ for some well formed formula ψ in the context $\vec{\sigma}$, then $\llbracket \phi \rrbracket = \flat_r \llbracket \psi \rrbracket$ and $\text{gr}(\phi, x_i) = r \cdot \text{gr}(\psi, x_i)$. By the hypothesis on ψ it is

$$P_{\langle \pi_1, \dots, \pi_n \rangle}(\llbracket \psi \rrbracket) * \prod_{i=1}^n P_{\langle \pi_i, \pi_{n+i} \rangle}(\flat_{\text{gr}(\psi, x_i)} \mathbf{d}_{\llbracket \sigma_i \rrbracket}) \leq P_{\langle \pi_{n+1}, \dots, \pi_{2n} \rangle}(\llbracket \psi \rrbracket)$$

It suffices to multiply both sides by \flat_r and use comultiplication of \flat to prove the claim. \square

A.3 Proofs of Section 4.4

Proposition 4.15. The terminal object in $\mathcal{L}_{\langle P, \flat \rangle}$ is $\langle 1, \kappa_{1 \times 1} \rangle$, where 1 is the terminal object in \mathcal{C} . Indeed, for any object $\langle A, \rho \rangle$ in $\mathcal{L}_{\langle P, \flat \rangle}$, the unique map $t_A : A \rightarrow 1$ in \mathcal{C} , induces a map from $\langle A, \rho \rangle$ to $\langle 1, \kappa_{1 \times 1} \rangle$ since by weakening it holds $\flat_0 \rho \leq \kappa_{A \times A} = P_{t_A \times t_A}(\kappa_{1 \times 1})$. Given $\langle A, \rho \rangle$ and $\langle B, \sigma \rangle$ in $\mathcal{L}_{\langle P, \flat \rangle}$, the pair $\langle A \times B, \rho \boxtimes \sigma \rangle$ is an object in $\mathcal{L}_{\langle P, \flat \rangle}$ by Proposition 4.14, as both ρ and σ are affine P -distances.

Since both ρ and σ are affine, projections $\pi_A : A \times B \rightarrow A$ and $\pi_B : A \times B \rightarrow B$ induce maps from $\langle A \times B, \rho \boxtimes \sigma \rangle$ to $\langle A, \rho \rangle$ and $\langle B, \sigma \rangle$, respectively; indeed, using affineness and counit, we have $\flat_1(\rho \boxtimes \sigma) \leq \flat_1 P_{\langle \pi_1, \pi_3 \rangle}(\rho) \leq P_{\langle \pi_1, \pi_3 \rangle}(\rho) = P_{\pi_A \times \pi_A}(\rho)$ and similarly for π_B . Finally, to check the universal property, let $\langle C, \tau \rangle$ be an object in $\mathcal{L}_{\langle P, \flat \rangle}$ and $f : \langle C, \tau \rangle \rightarrow \langle A, \rho \rangle$ and $g : \langle C, \tau \rangle \rightarrow \langle B, \sigma \rangle$ arrows in $\mathcal{L}_{\langle P, \flat \rangle}$. It suffices to show that the arrow $\langle f, \flat \rangle : C \rightarrow A \times B$ in \mathcal{C} induced by f and g underlines an arrow from $\langle C, \tau \rangle$ to $\langle A \times B, \rho \boxtimes \sigma \rangle$ in $\mathcal{L}_{\langle P, \flat \rangle}$. By hypothesis, we know that there are $r, s \in |R|$

such that $b_r\tau \leq P_{f \times f}(\rho)$ and $b_s\tau \leq P_{g \times g}(\sigma)$, hence, using contraction, we get

$$b_{r+s}\tau \leq b_r\tau * b_s\tau \leq P_{f \times f}(\rho) * P_{g \times g}(\sigma)$$

whence the claim. \square

Proposition 4.16. To show that that $D_{\langle P, b \rangle}$ is R -graded, amounts to prove the following three points, for all $\langle A, \rho \rangle$ in $\mathcal{L}_{\langle P, b \rangle}$.

1. We have $\kappa \in \mathcal{Des}_\rho$, as $\kappa * b_0\rho \leq \kappa$, by weakening.
2. Let $\alpha, \beta \in \mathcal{Des}_\rho$, hence we have $P_{\pi_1}(\alpha) * b_r\rho \leq P_{\pi_2}(\alpha)$ and $P_{\pi_1}(\beta) * b_s\rho \leq P_{\pi_2}(\beta)$, for some $r, s \in |R|$. Then, by contraction and commutativity of $*$, we get

$$\begin{aligned} P_{\pi_1}(\alpha * \beta) * b_{r+s}\rho &\leq P_{\pi_1}(\alpha) * b_r\rho * P_{\pi_1}(\beta) * b_s\rho \\ &\leq P_{\pi_2}(\alpha) * P_{\pi_2}(\beta) = P_{\pi_2}(\alpha * \beta) \end{aligned}$$

3. Let $\alpha \in \mathcal{Des}_\rho$ and $r \in |R|$, hence we have $P_{\pi_1}(\alpha) * b_s\rho \leq P_{\pi_2}(\alpha)$, for some $s \in |R|$. Then, by comultiplication, lax-monoidality and naturality of b_r , we get

$$\begin{aligned} P_{\pi_1}(b_r\alpha) * b_{r \cdot s}\rho &\leq b_r P_{\pi_1}(\alpha) * b_r b_s\rho \\ &\leq b_r(P_{\pi_1}(\alpha) * b_s\rho) \\ &\leq b_r P_{\pi_2}(\alpha) = P_{\pi_2}(b_r\alpha) \end{aligned}$$

In the end, to show that $D_{\langle P, b \rangle}$ is R -Lipschitz, we set $d_{\langle A, \rho \rangle} := \rho$, for each object $\langle A, \rho \rangle$ in $\mathcal{L}_{\langle P, b \rangle}$. First of all, note that $d_{\langle A, \rho \rangle} \in D_{\langle P, b \rangle}(\langle A, \rho \rangle \times \langle A, \rho \rangle) = D_{\langle P, b \rangle}\langle A \times A, \rho \boxtimes \rho \rangle$ since, by symmetry and transitivity of ρ and counit, we have $P_{\langle \pi_1, \pi_2 \rangle}(\rho) * b_1(\rho \boxtimes \rho) \leq P_{\langle \pi_3, \pi_4 \rangle}(\rho)$. Then, the thesis follows by Proposition 4.6. \square

Proposition 4.17. To prove the thesis, we just have to show that R_{Lip} is well-define, then algebraic identities follow immediately. Let $\langle F, f \rangle : \langle P, b \rangle \rightarrow \langle Q, b' \rangle$ be a 1-arrow in \mathbf{LD}_R . The fact that $\overline{F} : \mathcal{L}_{\langle P, b \rangle} \rightarrow \mathcal{L}_{\langle Q, b' \rangle}$ is a well-defined product-preserving functor follows from the following two facts.

- Let ρ be an affine P -distance on A , then $f_{A \times A}(\rho)$ is an affine Q -dittance on FA . This is immediate as f is a natural transformation preserving the monoidal structure.

- Let $h : \langle A, \rho \rangle \rightarrow \langle B, \sigma \rangle$ be an arrow in $\mathcal{L}_{\langle P, b \rangle}$, then $Fh : \langle FA, f_{A \times A}(\rho) \rangle \rightarrow \langle FB, f_{B \times B}(\sigma) \rangle$ is an arrow in $\mathcal{L}_{\langle Q, b' \rangle}$. By definition of arrows in $\mathcal{L}_{\langle P, b \rangle}$, we know that $b_r \rho \leq P_h(\sigma)$ holds for some $r \in |R|$. Since $\langle F, f \rangle$ is a 1-arrow in \mathbf{LD}_R , we know that $b'_{rFA \times FA} \circ f_{A \times A} \leq f_{A \times A} \circ b_{rA \times A}$, hence, applying $f_{A \times A}$ we get

$$b'_r f_{A \times A}(\rho) \leq f_{A \times A}(b_r \rho) \leq f_{A \times A}(P_h(\sigma)) = Q_{Fh}(f_{B \times B}(\sigma))$$

To check that f_A applies $D_{\langle P, b \rangle} \langle A, \rho \rangle = \mathcal{Des}_\rho$ into $D_{\langle Q, b' \rangle} \langle FA, f_{A \times A}(\rho) \rangle = \mathcal{Des}_{f_{A \times A}(\rho)}$, it is enough to note that, since, for all $\alpha \in \mathcal{Des}_\rho$, we have $P_{\pi_1}(\alpha) *^P b_r \rho \leq P_{\pi_2}(\alpha)$, for some $r \in |R|$, we

$$\begin{aligned} Q_{F\pi_1}(f_A(\alpha)) *^Q b'_r f_{A \times A}(\rho) &\leq Q_{f\pi_1}(f_A(\alpha)) *^Q f_{A \times A}(b_r \rho) \\ &= f_{A \times A}(P_{\pi_1}(\alpha) *^P b_r \rho) \\ &\leq f_{A \times A} P_{\pi_2}(\alpha) \\ &= Q_{F\pi_2}(f_A(\alpha)) \end{aligned}$$

Since the R -graded structure of $D_{\langle P, b \rangle}$ and $D_{\langle Q, b' \rangle}$ are, respectively, that of $\langle P, b \rangle$ and $\langle Q, b' \rangle$, f preserves the structure by hypothesis, hence $\langle \overline{F}, f \rangle$ is a well-defined 1-arrow in \mathbf{LLD}_R .

Let $\theta : \langle F, f \rangle \Rightarrow \langle G, g \rangle$ be a 2-arrow in \mathbf{LD}_R . The fact that $\mathbf{R}_{\text{Lip}}(\theta) = \theta$ is a well-defined 2-arrow in \mathbf{LLD}_R follows from the following observation: let $\langle A, \rho \rangle$ be an object in $\mathcal{L}_{\langle P, b \rangle}$, then $\theta_A : \langle FA, f_{A \times A}(\rho) \rangle \rightarrow \langle GA, g_{A \times A}(\rho) \rangle$ is an arrow in $\mathcal{L}_{\langle Q, b' \rangle}$. This is immediate as, by definition of 2-arrow and counit, we get $b'_1 f_{A \times A}(\rho) \leq f_{A \times A}(\rho) \leq Q_{\theta_{A \times A}}(g_{A \times A}(\rho)) = Q_{\theta_A \times \theta_A}(g_{A \times A}(\rho))$. \square

Theorem 4.18. First we prove that $\mathbf{U}_{\text{Lip}} \dashv \mathbf{R}_{\text{Lip}}$. Consider the following 2-natural transformations:

$$\epsilon_{\langle P, b \rangle} := \langle U^P, u^P \rangle \quad \eta_{\langle Q, \sharp \rangle} := \langle E^Q, e^Q \rangle$$

for each object $\langle P, b \rangle$ in \mathbf{LD}_R and $\langle Q, \sharp, e \rangle$ in \mathbf{LLD}_R . We have that $\epsilon : \mathbf{U}_{\text{Lip}} \circ \mathbf{R}_{\text{Lip}} \xrightarrow{\cdot} \text{Id}_{\mathbf{LD}_R}$ is a 2-natural transformation because, for any 1-arrow $\langle F, f \rangle : \langle P, b \rangle \rightarrow \langle P', b' \rangle$ in \mathbf{LD}_R , it holds $U^{P'} \circ \overline{F} = F \circ U^P$, by definition of \mathbf{R}_{Lip} . On the other hand, $\eta : \text{Id}_{\mathbf{LLD}_R} \xrightarrow{\cdot} \mathbf{R}_{\text{Lip}} \circ \mathbf{U}_{\text{Lip}}$ is a 2-natural transformation because each 1-arrow $\langle F, f \rangle : \langle Q, \sharp, d \rangle \rightarrow \langle Q', \sharp', d' \rangle$ in \mathbf{LLD}_R preserves distances, hence $E^{Q'} F A = \langle F A, d'_{FA} \rangle = \langle F A, f_{A \times A}(d_A) \rangle = \overline{F} E^Q A$, for any object A in the base category of Q . The two triangular identities hold as the following diagrams (in \mathbf{LD}_R) commute for any R -graded doctrine

$\langle P, \flat \rangle$ and R -Lipschitz doctrine $\langle Q, \sharp, \mathbf{e} \rangle$:

$$\begin{array}{ccc}
\langle D_{\langle P, \flat \rangle}, \flat \rangle & \xrightarrow{\langle E^{D_{\langle P, \flat \rangle}}, e^{D_{\langle P, \flat \rangle}} \rangle} & \langle D_{\langle D_{\langle P, \flat \rangle}, \flat \rangle}, \flat \rangle \\
& \searrow \langle \text{Id}, \text{id} \rangle & \downarrow \langle U^{D_{\langle P, \flat \rangle}}, u^{D_{\langle P, \flat \rangle}} \rangle \\
& & \langle D_{\langle P, \flat \rangle}, \flat \rangle
\end{array}$$

$$\begin{array}{ccc}
\langle Q, \sharp \rangle & \xrightarrow{\langle E^Q, e^Q \rangle} & \langle D_{\langle Q, \sharp \rangle}, \sharp \rangle \\
& \searrow \langle \text{Id}, \text{id} \rangle & \downarrow U^Q, u^Q \\
& & \langle Q, \sharp \rangle
\end{array}$$

Let \mathbf{T} be the 2-comonad on \mathbf{LD}_R induced by the adjunction. In order to prove that \mathbf{LLD}_R is isomorphic to the 2-category $\mathbf{LLD}_R^{\mathbf{T}}$ of coalgebras for \mathbf{T} , note that there is a comparison 2-functor $\mathbf{K} : \mathbf{LLD}_R \rightarrow \mathbf{LD}_R^{\mathbf{T}}$ mapping an R -Lipschitz doctrine $\langle Q, \sharp, \mathbf{e} \rangle$ to the coalgebra $\langle E^Q, e^Q \rangle : \langle Q, \sharp \rangle \rightarrow \langle D_{\langle Q, \sharp \rangle}, \sharp \rangle$ and being the identity on arrows. On the other hand, given a coalgebra $\langle F, f \rangle : \langle P, \flat \rangle \rightarrow \langle D_{\langle P, \flat \rangle}, \flat \rangle$ on an R -graded doctrine $\langle P, \flat \rangle$, since $\langle U^P, u^P \rangle \circ \langle F, f \rangle = \langle \text{Id}, \text{id} \rangle$, we have $f = \text{id}$ and $FA = \langle A, \rho \rangle$ for any object A in the base category of P and $P(A) = \mathcal{D}es_{\rho}(A)$. Hence $\langle P, \flat, \mathbf{d} \rangle$ is R -Lipschitz with $\mathbf{d}_A = \rho$. On arrows the 2-functor is the identity, because being a coalgebra morphism is exactly the same as preserving distances. Such a 2-functor is the inverse of \mathbf{K} . \square

A.4 Proofs of Section 5

Theorem 5.5. The composition $\mathbf{R}_{\text{EL}} \circ \mathbf{I}_{\text{EL}}$ is the identity on \mathbf{ELD}_R , because equality predicates are \flat -intuitionistic by Corollary 5.3, hence the unit of the 2-adjunction is the identity. On the other hand, for any R -Lipschitz doctrine $\langle P, \flat, \mathbf{d} \rangle$, there is an obvious inclusion 1-arrow $\langle I^P, \iota^P \rangle : \langle P_!, \flat, \mathbf{d} \rangle \rightarrow \langle P, \flat, \mathbf{d} \rangle$, which is the counit of the 2-adjunction. The fact that the unit and the counit satisfy triangular identities is immediate, hence we have the 2-adjunction $\mathbf{I}_{\text{EL}} \dashv \mathbf{R}_{\text{EL}}$.

Let \mathbf{T} be the 2-comonad $\mathbf{I}_{\text{EL}} \circ \mathbf{R}_{\text{EL}}$. It is easy to check that the comparison 2-functor from \mathbf{ELD}_R to the 2-category of \mathbf{T} -coalgebras is full and faithful as \mathbf{I}_{EL} is an inclusion. Hence, to conclude, we just have to prove that the underlying R -graded doctrine of every \mathbf{T} -coalgebra $\langle F, f \rangle : \langle P, \flat, \mathbf{d} \rangle \rightarrow \langle P_!, \flat, \mathbf{d} \rangle$ is elementary. This follows because the triangle of the definition of

T-coalgebra imposes that $\langle F, f \rangle$ is the identity, hence $P = P_1$, which proves the thesis. \square