

METABELIAN LIE AND PERM ALGEBRAS

F. A. MASHUROV AND B. K. SARTAYEV

ABSTRACT. It is well-known that any Lie algebra can be embedded into an associative algebra. We prove that any metabelian Lie algebra can be embedded into an algebra in the subvariety of associative algebras defined by the identity $abc - acb = 0$. In addition, we give a technical method to universal enveloping perm algebra for a given metabelian Lie algebra.

1. INTRODUCTION

Determining a set of identities for an adjoint class of a variety through commutator or anti-commutator products is a popular question in algebra. This question has been solved or largely answered for some varieties, but it is still open for others. It is well known that a variety of associative algebras are a classic illustration of this approach. Poincaré-Birkhoff-Witt's (PBW) theorem provides us the entire collection of identities for commutator algebras of associative algebras. In other words, anti-commutativity and the Jacobi identities result in any identity satisfied by the commutator product in any associative algebra. Moreover, it follows that any Lie algebra can be embedded into an associative algebra. However, anti-commutator algebras of associative algebras have an entirely different story.

A variety of metabelian Lie algebras is defined by the additional identity

$$[[a, b], [c, d]] = 0.$$

The metabelian variety of Lie algebras has received a lot of attention and considered in [6], [10], [11], [15], [16], [17]. A basis of free metabelian Lie algebras was given in [1].

In this paper we show that any metabelian Lie algebra can be embedded into an algebra in the subvariety of associative algebras defined by the right-commutative identity:

$$abc - acb = 0. \tag{1}$$

An associative algebra with the right-commutative identity (1) is called a *perm* algebra, see [5], [20]. In [13] perm algebras were considered with derivation.

2010 *Mathematics Subject Classification.* 17B01, 17B35, 13P10.

Key words and phrases. Perm algebras, Lie and Jordan elements, polynomial identities, Gröbner bases.

The authors were supported by grant AP08052405 of MES RK. The authors are grateful to Professor A. Dzhumadil'dayev and Professor P. Kolesnikov for their essential comments.

Let $A^{(-)} = (A, [\cdot, \cdot])$ be a subalgebra of A under commutator i.e. multiplication is defined by the Lie bracket $[a, b] = ab - ba$. An algebra $A^{(+)} = (A, \{\cdot, \cdot\})$ can be defined in the similar way i.e. with the anticommutator product $\{a, b\} = ab + ba$.

Let $\mathcal{P}erm$ be a variety of perm algebras. Define $\mathcal{P}erm^{(-)}$ and $\mathcal{P}erm^{(+)}$ as a class of algebras of the form $P^{(-)}$ and $P^{(+)}$, respectively, where $P \in \mathcal{P}erm$. An algebra B is called *special* concerning Lie (*Jordan*) brackets if there exists a perm algebra P such that B is a subalgebra of $P^{(-)}$ (*respectively* $P^{(+)}$), otherwise it is called *exceptional*. The algebra P is called a perm envelope for B . We show that any algebra of $\mathcal{P}erm^{(-)}$ is a metabelian Lie algebra. We prove that any identity satisfied by the commutator product in all perm algebras is a consequence of the anticommutativity, Jacobi, and metabelian identities. In the case of anti-commutator product, we show that there are two identities in degree 4.

Let $X = \{x_1, x_2, \dots\}$ be a set and $P(X)$ be a free perm algebra generated by X . A polynomial in $P(X)$ is called Lie (*Jordan*) element of $P(X)$ if it can be expressed by elements of X in terms of commutators (*anticommutators*). For associative algebras, there is a well-known Dynkin-Specht-Wever criterion for Lie elements [19], and for the Jordan elements, the question remains open. Also, metabelian Lie algebras can be obtained from bicommutative algebras by commutator. Criteria for Lie and Jordan elements in a free bicommutative algebra were found recently in [9]. P.M. Cohn gave a criterion for the Jordan elements generated by three elements [4]. In addition, he gave a criterion of homomorphic images to be special and showed that the homomorphic image of the free special Jordan algebra generated by three elements is exceptional.

In this study, we present conditions for Lie elements and describe all Jordan elements in a free perm algebra. The major results of the paper are derived by using Cohn's criterion for speciality of algebras and the obtained criteria. If an algebra B can be embedded into A , then it is natural to find the base of the universal envelope algebra of A . For this reason, we use the technique of Gröbner-Shirshov bases for perm algebras. We show a method to find the basis of the universal enveloping perm algebra for a given metabelian Lie algebra.

In the last section, we consider a commutative algebra with two identities of degree four, which is obtained from $P^{(+)}$ in terms of operads. We use computer algebra methods to show that all special identities up to degree 10 of $P^{(+)}$ follow from identities of degree 4. In addition, the operad in question satisfies Glennie's identity.

2. MAIN RESULTS

We consider all algebras over a field K of characteristic 0.

2.1. Theorem. *Every identity satisfied by the commutator in every free perm algebra is a consequence of anti-commutativity, the Jacobi and the metabelian identities:*

$$[a, b] = -[b, a], \quad (2)$$

$$[[a, b], c] + [[b, c], a] + [[c, a], b] = 0, \quad (3)$$

$$[[a, b], [c, d]] = 0, \quad (4)$$

where $a, b, c, d \in P(X)$.

For $x_1, \dots, x_n \in P(X)$ denote by $x_1x_2 \cdots x_n$ a left-bracketed element $(\cdots(x_1x_2) \cdots x_{n-1})x_n$. The Dynkin map $D : P(X) \rightarrow P(X)$ is a linear map, defined on base elements of $P(X)$ as follows

$$x_{i_1}x_{i_2} \cdots x_{i_n} \mapsto [[\cdots[x_{i_1}, x_{i_2}], \cdots], x_{i_n}],$$

where $i_2 \leq \dots \leq i_n$.

We define *head* of $f \in P(X)$ as a polynomial where the first generator of each monomial in f is not x_1 . For example, if $f = x_1x_2x_3x_4 + x_2x_1x_3x_4 - x_3x_1x_2x_4 + 2x_4x_1x_2x_3$, then

$$\text{head}(f) = x_2x_1x_3x_4 - x_3x_1x_2x_4 + 2x_4x_1x_2x_3.$$

Let $L(X)$ be a free special metabelian Lie algebra generated by a set X .

2.2. Theorem. *Let $f \in P(X)$. Then f is a Lie element if and only if $D(\text{head}(f)) = f$.*

2.3. Theorem. *Every homomorphic image of $L(X)$ is special.*

Consequently, from Theorem 2.1 and Theorem 2.3 we have:

2.4. Corollary. *Every metabelian Lie algebra has a perm enveloping algebra.*

Now we consider algebras from $\mathcal{P}\text{erm}^{(+)}$.

2.5. Theorem. *Every perm algebra under anticommutator satisfies the following identities:*

$$\{a, b\} = \{b, a\}, \quad (5)$$

$$\{\{a, b\}, \{c, d\}\} = \{\{a, d\}, \{b, c\}\}, \quad (6)$$

$$2\langle \{a, b\}, c, d \rangle = \langle \{a, b\}, d, c \rangle + \langle \{a, c\}, b, d \rangle + \langle \{b, c\}, a, d \rangle, \quad (7)$$

for $a, b, c, d \in P(X)$ and where $\langle a, b, c \rangle = \{\{a, b\}, c\} - \{a, \{b, c\}\}$.

2.6. Theorem. *Every element with degree more than 2 in a free perm algebra is a Jordan element.*

Let $J(X)$ be a free special algebra on X under anticommutator product, i.e. subalgebra of $P(X)^{(+)} = (P(X), \{, \})$ generated by X . The next theorem is an analogue of Cohn's theorem on exceptional homomorphic image of the special Jordan algebra in three generators [4]. In our case, we have an exceptional homomorphic image in two generators.

2.7. Theorem. *An algebra $J(\{x, y\})$ has an exceptional homomorphic image.*

2.8. Corollary. *The class of all $P(X)^{(+)}$ algebras is not variety.*

Using computer algebra method we show that there is no a new special identity up to degree ten.

2.9. Theorem. *Every identity of $J(X)$ of degree ≤ 10 follows from the commutativity, (6) and (7) identities.*

3. PROOF OF THEOREMS

In this section we give proofs of main results.

3.1. Lemma. *For $n \geq 2$, we have*

$$[\cdots [[x_1, x_2], x_3], \cdots, x_n] = [x_1, x_2]x_3 \cdots x_n. \quad (8)$$

Proof. We prove it by induction on n . The base of induction $n = 2$. Suppose that the equality (8) holds for $n - 1$.

Then we have

$$\begin{aligned} & [\cdots [[x_1, x_2], x_3], \cdots, x_n] = \\ & [\cdots [[x_1, x_2], x_3], \cdots, x_{n-1}]x_n - x_n[\cdots [[x_1, x_2], x_3], \cdots, x_{n-1}] \end{aligned}$$

It is easy to see that

$$x_n[\cdots [[x_1, x_2], x_3], \cdots, x_{n-1}] = 0 \quad (9)$$

by (1) and by the inductive hypothesis, we obtain

$$[\cdots [[x_1, x_2], x_3], \cdots, x_n] = [x_1, x_2]x_3 \cdots x_n.$$

□

An ideal I of the free perm algebra $P(X)$ is called *T-ideal* if $\xi(I) \subseteq I$ for all endomorphisms ξ of $P(X)$. Denote by $T(u)$ the *T-ideal* in $P(X)$ generated by $u \in P(X)$.

3.2. Proposition. *An algebra $(P, [\cdot, \cdot])$ satisfies the metabelian identity.*

Proof. It is easy to see that $[[a, b], [c, d]] \in T(a[b, c])$. Since the identity (1) holds in $P(X)$, hence (4) is an identity in $(P, [\cdot, \cdot])$.

□

3.1. Proof of Theorem 2.2.

Proof. Let $f \in P(X)$ and suppose that $D(\text{head}(f)) = f$. Since the image of Dynkin map D is the subspace of $SL(X)$, f is Lie element.

Now, suppose that $f \in L(X)$ and

$$f = \sum \lambda [\cdots [[x_1, x_{i_1}], x_{i_2}], \cdots, x_{i_n}],$$

for $\lambda \in K$ and $i_1 > 1$. By Lemma 3.1, for monomial $\lambda [\cdots [[x_1, x_{i_1}], x_{i_2}], \cdots, x_{i_n}]$ we have

$$\text{head}(\lambda [\cdots [[x_1, x_{i_1}], x_{i_2}], \cdots, x_{i_n}]) = -\lambda x_{i_1} x_1 x_{i_3} \cdots x_{i_n}.$$

If we apply Dynkin map for the monomial of $P(X)$ on the right hand of the equality, then

$$D(-\lambda x_{i_1} x_1 x_{i_3} \cdots x_{i_n}) = -\lambda D(x_{i_1} x_1 x_{i_3} \cdots x_{i_n}) =$$

(and by anti-commutativity)

$$\lambda[[\cdots [[x_1, x_{i_1}], x_{i_2}] \cdots], x_{i_n}].$$

Applying the linear mapping D to other basis elements in $head(f)$, we obtain $D(head(f)) = f$. \square

Let us recall a free base of metabelian Lie algebra on a set X of degree n [1]:

$$[[\cdots [[x_{i_1}, x_1], x_{i_2}], \cdots], x_{i_{n-1}}], \quad (10)$$

where $i_1 > 1$ and $1 \leq i_2 \leq \cdots \leq i_{n-1} \leq n$.

3.2. Proof of Theorem 2.1.

Proof. Since the number basis monomials of the space of the *head* of $P(X)$ is equal to the dimension of the free metabelian Lie algebra and they are independent in $P(X)$, it follows that any identity consequence of (2), (3) and (4). \square

3.3. Corollary. *The free metabelian Lie algebra is special.*

3.3. Proof of Theorem 2.3. Now we consider the algebras $SL(X)$ and $P(X)$ with ideals α and $\{\alpha\}$, respectively, where $\{\alpha\}$ generated by the set α .

Proof. To prove the factor-algebra $SL(X)/\alpha$ is special for any α , by Cohn's criterion [4], we need to show that $SL(X) \cap \{\alpha\} \subseteq \alpha$.

Assume that g_i ($i \in I$) are generators of the ideal α . Let w be a non-zero element of $SL(X) \cap \{\alpha\}$. Then

$$s = \sum \lambda_k s_k, \text{ where } \lambda_k \in K,$$

is a linear combination of base monomials in elements of X and g_i , such that each monomial s_k is linear in at least one generator of α . We write the sum as follows

$$\sum \lambda_k s_k = \sum \mu_l u_l + \sum \mu_m v_m, \text{ where } \mu_l, \mu_m \in K,$$

such that every u_l depends on a generator of α whose degree is more than one and every v_m depends on a generator of α whose degree is equal to one.

Assume that u_l is linear by some $g_i \in \alpha$ and for some l . Since $g_i \notin X$, we have $g_i \in [P(X), P(X)]$ and consequently, by (9) $x_k g_i = 0$. Then

$$u_l = g_i x_{i_1} \cdots x_{i_n}$$

where $x_{i_1}, \dots, x_{i_n} \in X$. One can write

$$\begin{aligned} g_i x_{i_1} \cdots x_{i_n} &= (g_i x_{i_1}) x_{i_2} \cdots x_{i_n} - (x_{i_1} g_i) x_{i_2} \cdots x_{i_n} = \\ &= [g_i, x_{i_1}] x_{i_2} \cdots x_{i_n} = [[\cdots [g_i, x_{i_1}], x_{i_2}] \cdots], x_{i_n}. \end{aligned}$$

Hence $u_l \in \alpha$ for all l . Then $\sum \mu_l u_l \in \alpha$. It implies that $\sum \mu_m v_m \in SL(X)$ and consequently $\sum \mu_m v_m \in \alpha$. Therefore $s \in \alpha$. \square

3.4. Proof of Theorem 2.5.

Proof. The identities (6) and (7) can be easily verified using the following expansions of the elements

$$\begin{aligned} \{\{a, b\}, \{c, d\}\} &= 2abcd + 2bacd + 2cabd + 2dabc, \\ \langle \{a, b\}, c, d \rangle &= -abcd - bacd + 4dabc. \end{aligned} \quad (11) \quad \square$$

3.5. Proof of Theorem 2.6.

Proof. Let us write a congruence relation $a \equiv 0$ for $a \in P(X)^{(\+)}$. Then we have to show that

$$x_1 x_2 \cdots x_n \equiv 0. \quad (12)$$

We use induction on n . If $n = 3$, then

$$x_1 x_2 x_3 = -\frac{1}{4} \{\{x_1, x_2\}, x_3\} + \frac{3}{4} \{\{x_2, x_3\}, x_1\} - \frac{1}{4} \{\{x_1, x_3\}, x_2\}.$$

Assume that (12) holds for elements with degree less than n . Then

$$x_1 x_2 \cdots x_{n-1} \equiv 0.$$

Set $x_{n-1} = \{x_{n-1}, x_n\}$. Then by (1), we obtain

$$0 \equiv x_1 x_2 \cdots x_{n-2} \{x_{n-1}, x_n\} = 2x_1 x_2 \cdots x_n.$$

Hence

$$x_1 x_2 \cdots x_n \equiv 0. \quad \square$$

3.6. Proof of Theorem 2.7.

Proof. Let us consider an ideal I of the free algebra $J(\{x, y\})$ generated by one element $e_1 = \{x, y\}$ and $\{I\}$ is an ideal of $P(\{x, y\})$ generated by the e_1 .

By the criterion of speciality of the homomorphic image in [4], we have to show that the homomorphic image of $J(\{x, y\})$ is exceptional. That is, the equality $\{I\} \cap J(\{x, y\}) = I$ does not hold.

We consider $u = \frac{1}{2} x e_1 = xxy$. By the Theorem 2.6, the element u belongs to $J(\{x, y\})$. Therefore $u \in \{I\} \cap J(\{x, y\})$. However, u does not belong to I , by the reason that there is no $\lambda \in K$ such that $u = \lambda \{e_1, x\}$. Consequently, $J(\{x, y\})/I$ is exceptional. \square

4. EXAMPLES OF UNIVERSAL ENVELOPING ALGEBRA FOR GIVEN METABELIAN LIE ALGEBRA

In this section, we consider some examples of metabelian Lie algebras and find the basis of universal enveloping algebras for them. In fact, the described method works for arbitrary metabelian Lie algebra. We use the technique of Gröbner-Shirshov bases for perm algebras that can be derived from [12]. More explicitly, in [12] given the method of finding the Gröbner-Shirshov bases for *di-Var* algebras using the theory of Gröbner-Shirshov bases of *Var* algebras.

In our case, the free perm algebra is a free di-commutative algebra, and the theory of Gröbner bases for commutative algebras developed in [3]. So, we have

$$\text{Perm}\langle X \rangle = \text{di-Com}\langle X \rangle \simeq V \subseteq \text{Com}\langle X \cup \dot{X} \rangle,$$

where V is a subalgebra of $\text{Com}\langle X \cup \dot{X} \rangle$ generated by \dot{X} and operations of V are \vdash and \dashv such that $a \vdash b = \dot{a}\dot{b}$ and $a \dashv b = \dot{a}b$.

Let us consider two-dimensional metabelian Lie algebra A with multiplication table

$$[e_1, e_2] = -[e_2, e_1] = e_2.$$

Now, we find the Gröbner-Shirshov base of the free perm algebra with relation

$$f = e_2e_1 - e_1e_2 + e_2$$

or in terms of the free di-commutative algebra, a relation f can be written as

$$f = e_2 \dashv e_1 - e_1 \dashv e_2 + e_2.$$

According to the general scheme, which is given in [12], to work in commutative algebra we should transform the relation f in terms of commutative algebra into relations

$$f_1 = \dot{e}_2e_1 - \dot{e}_1e_2 + \dot{e}_2 \quad \text{and} \quad f_2 = e_2.$$

It is clear that f_1 and f_2 form a Gröbner base in a free commutative algebra $\text{Com}\langle e_1, e_2, \dot{e}_1, \dot{e}_2 \rangle$ and basis monomials of $U(A)$ are

$$e_2 \quad \text{and} \quad (\cdots (e_1 \dashv e_1) \cdots) \dashv e_1$$

or in terms of the perm algebra e_2, e_1^n and $A \hookrightarrow U(A)$.

Let us consider three dimensional Heisenberg algebra \mathcal{H} i.e. algebra with the following multiplication table:

$$[e_1, e_2] = -[e_2, e_1] = e_3.$$

An algebra \mathcal{H} is a metabelian Lie. Now, we consider the free perm algebra with relations

$$f_1 := e_2e_1 - e_1e_2 + e_3, f_2 := e_3e_1 - e_1e_3, f_3 := e_3e_2 - e_2e_3 \quad (13)$$

As in previous case, we rewrite (13) in terms of $\text{Com}\langle X \cup \dot{X} \rangle$:

$$f_1 = \dot{e}_2e_1 - \dot{e}_1e_2 + \dot{e}_3, f_2 = \dot{e}_3e_1 - \dot{e}_1e_3, f_3 = \dot{e}_3e_2 - \dot{e}_2e_3, f_4 = e_3.$$

So we have

$$f_1 = \dot{e}_2 e_1 - \dot{e}_1 e_2 + \dot{e}_3, \quad f_2 = \dot{e}_3 e_1, \quad f_3 = \dot{e}_3 e_2, \quad f_4 = e_3. \quad (14)$$

It is an easy task to find the Gröbner base for a given set of relations. That is (14) with additional relation $f_4 = \dot{e}_3^2$. So basis monomials of $U(\mathcal{H})$ are

$$e_3, \quad (\cdots (e_2 \dashv e_2) \cdots) \dashv e_2$$

and

$$(\cdots (((\cdots (e_1 \dashv e_1) \cdots) \dashv e_1) \dashv e_2) \cdots) \dashv e_2.$$

In terms of perm algebra, basis monomials of $U(\mathcal{H})$ are e_3 , e_2^n , $e_1^k e_2^l$ and $\mathcal{H} \hookrightarrow U(\mathcal{H})$.

4.1. Remark. Note that the pair of varieties $(\mathcal{P}erm, \mathcal{ML}ie)$ is not a PBW-pair in the sense of [18]. Indeed, the Gel'fand–Kirillov dimension of $U(\mathcal{H})$ equals two, but the universal enveloping of the 3-dimensional abelian Lie algebra is just the polynomial algebra in 3 variables, so its Gel'fand–Kirillov dimension is 3.

5. ON THE GRÖBNER BASIS OF \mathcal{J} -OPERAD

Let us denote by \mathcal{J} a commutative symmetric operad defined by identities (6), (7) and $\mathcal{J}(n)$ with $n \geq 1$ is a linear space in the category $\text{Vec}_{\mathbb{k}}$ of linear spaces over a field \mathbb{k} , equipped with linear composition maps

$$\gamma_{n_1, \dots, n_m}^m : \mathcal{J}(m) \otimes \mathcal{J}(n_1) \otimes \cdots \otimes \mathcal{J}(n_m) \rightarrow \mathcal{J}(n_1 + \cdots + n_m)$$

for all integers $m, n_1, \dots, n_m \geq 1$, each $\mathcal{J}(n)$ is a module over the symmetric group S_n (for more details, see [2]).

Shuffle operads give a method for the computation of Gröbner bases and normal forms in an operad defined by generators and relations [8]. As shown in [2] every symmetric operad may be considered as a shuffle one provided that the operation set is extended.

Let us give a definition of shuffle operad. A shuffle operad \mathcal{J} is a collection of spaces $\mathcal{J}(n)$, $n \geq 1$, equipped with a collection of compositions

$$\gamma_\pi : \mathcal{J}(m) \otimes \mathcal{J}(n_1) \otimes \cdots \otimes \mathcal{J}(n_m) \rightarrow \mathcal{J}(n_1 + \cdots + n_m)$$

where π is a *shuffle partition* of the set $\{1, \dots, n\}$, $n = m_1 + \cdots + m_m$, into m disjoint subsets I_j , $j = 1, \dots, m$, such that $\min I_1 < \min I_2 < \cdots < \min I_m$.

Following [2], we may convert the symmetric operad \mathcal{J} into a shuffle operad \mathcal{J}^f , a homomorphic image of the shuffle tree operad $\mathcal{T}_{\text{III}}(\mathcal{X})$ for an appropriate language \mathcal{X} .

5.1. Example. Let us consider an operad Lie with operation μ . Applying a functor f , we rewrite monomials of operad Lie as elements of operad $\mathcal{T}_{\text{III}}(\{x\})$ as follows:

$$\mu(x_1, x_2) = x(1 \ 2), \quad \mu(x_2, x_1) = -\mu(x_1, x_2) = -x(1 \ 2).$$

Converting the Jacoby identity, we obtain the following relation:

$$x(x(1 \ 2) \ 3) - x(1 \ x(2 \ 3)) - x(x(1 \ 3) \ 2)$$

and it is easy to see that the relation defined by the Jacoby identity forms the Gröbner basis.

5.2. Example. Let us consider another symmetric operad defined by the following identity:

$$\begin{aligned} (x_i x_j) x_k &= (x_j x_i) x_k + x_k (x_i x_j) - x_k (x_j x_i) - (x_j x_k) x_i + (x_k x_j) x_i + x_i (x_j x_k) \\ &\quad - x_i (x_k x_j) - (x_k x_i) x_j + (x_i x_k) x_j + x_j (x_k x_i) - x_j (x_i x_k). \end{aligned} \quad (15)$$

That is defining identity of Lie-admissible operad and denote the operation of such operad by ρ . We convert the identity (15) to elements of $\mathcal{T}_{\text{III}}(\{x, y\})$ as follows:

$$\rho(x_1, x_2) = x(1\ 2), \quad \rho(x_2, x_1) = y(1\ 2).$$

We obtain

$$\begin{aligned} &x(x(1\ 2)\ 3) - x(y(1\ 2)\ 3) - y(x(1\ 2)\ 3) + y(y(1\ 2)\ 3) + y(1\ x(2\ 3)) - y(1\ y(2\ 3)) \\ &- x(1\ x(2\ 3)) + x(1\ y(2\ 3)) + x(y(1\ 3)\ 2) - x(x(1\ 3)\ 2) - y(y(1\ 3)\ 2) + y(x(1\ 3)\ 2). \end{aligned} \quad (16)$$

Direct calculations show that the relation (16) forms a Gröbner basis. Another example of applying a functor f on Gelfand-Dorfman-operad is given in [14] and the computation was made up to degree 6. There appear nontrivial compositions in each degree and the explicit form of Gröbner basis is unknown.

Analogically, let us rewrite defining identities of the operad \mathcal{J} as elements of $\mathcal{T}_{\text{III}}(\{x\})$. Commutativity gives us that we may convert defining identities of operad \mathcal{J} without additional operations by the reason that we define operation γ as

$$\gamma(x_1, x_2) = x(1\ 2) \quad \text{and} \quad \gamma(x_2, x_1) = \gamma(x_1, x_2) = x(1\ 2).$$

Converting and simplifying (6), we get

$$\begin{aligned} &x(x(1\ 4)\ x(2\ 3)) - x(x(1\ x(3\ 4))\ 2) - x(1\ x(x(2\ 3)\ 4)) \\ &\quad - x(1\ x(x(2\ 4)\ 3)) + 2x(1\ x(2\ x(3\ 4))), \\ &x(x(1\ 3)\ x(2\ 4)) - x(x(1\ x(3\ 4))\ 2) - x(1\ x(x(2\ 3)\ 4)) \\ &\quad - x(1\ x(x(2\ 4)\ 3)) + 2x(1\ x(2\ x(3\ 4))), \\ &x(x(1\ 2)\ x(3\ 4)) - x(x(1\ x(3\ 4))\ 2) - x(1\ x(x(2\ 3)\ 4)) \\ &\quad - x(1\ x(x(2\ 4)\ 3)) + 2x(1\ x(2\ x(3\ 4))). \end{aligned}$$

Converting and simplifying (7), we get

$$\begin{aligned} &x(x(1\ x(2\ 4))\ 3) - x(x(1\ x(3\ 4))\ 2) - 3x(1\ x(x(2\ 4)\ 3)) + 3x(1\ x(2\ x(3\ 4))), \\ &x(x(x(1\ 4)\ 2)\ 3) - 3/2x(x(1\ x(3\ 4))\ 2) - 3x(1\ x(x(2\ 4)\ 3)) + 7/2x(1\ x(2\ x(3\ 4))), \\ &x(x(x(1\ 3)\ 2)\ 4) - 3/2x(x(1\ x(3\ 4))\ 2) - 3x(1\ x(x(2\ 3)\ 4)) + 7/2x(1\ x(2\ x(3\ 4))), \\ &x(x(x(1\ 3)\ 4)\ 2) - 3/2x(x(1\ x(3\ 4))\ 2) - x(1\ x(x(2\ 3)\ 4)) + 3/2x(1\ x(2\ x(3\ 4))), \\ &x(x(x(1\ 4)\ 3)\ 2) - 3/2x(x(1\ x(3\ 4))\ 2) - x(1\ x(x(2\ 4)\ 3)) + 3/2x(1\ x(2\ x(3\ 4))), \end{aligned}$$

$$\begin{aligned}
& x(x(1 x(2 3)) 4) - x(x(1 x(3 4)) 2) - 3x(1 x(x(2 3) 4)) + 3x(1 x(2 x(3 4))), \\
& \quad x(x(x(1 2) 4) 3) - 3/2x(x(1 x(3 4)) 2) - x(1 x(x(2 3) 4)) \\
& \quad \quad - 3x(1 x(x(2 4) 3)) + 9/2x(1 x(2 x(3 4))), \\
& \quad x(x(x(1 2) 3) 4) - 3/2x(x(1 x(3 4)) 2) - 3x(1 x(x(2 3) 4)) \\
& \quad \quad - x(1 x(x(2 4) 3)) + 9/2x(1 x(2 x(3 4))).
\end{aligned}$$

The obtained elements generate the ideal of defining relations for the operad \mathcal{J}^f . Calculating the Gröbner base of the operad \mathcal{J}^f by means of the package [7], we get the following result:

n	1	2	3	4	5	6	7	8	9	10
$\dim(\mathcal{J}(n))$	1	1	3	4	5	6	7	8	9	10

In particular, we have $\dim J(n) = \dim P^{(+)}(n)$ for $n \leq 10$ which proofs **Theorem 2.9**.

It is well-known that a free Jordan algebra does not satisfy the Glennie identity (that is identity of degree 8) but for subvariety \mathcal{J} of variety of Jordan algebras we have the following result:

5.3. Corollary. *A free algebra $\mathcal{J}(X)$ satisfies the Glennie identity.*

Computing the Gröbner basis of operad \mathcal{J} , we obtain nontrivial compositions in each degree up to degree 10. It is an interesting problem to find the Gröbner basis of operad \mathcal{J} .

REFERENCES

- [1] Bahturin Yu. A., *Identities in Lie algebras I*, Vest. Moskov. Univ. Ser. I. Mat. Mekh., **28**, (1973), 12–18.
- [2] Bremner M.R., Dotsenko V. *Algebraic Operads An Algorithmic Companion*, (2016).
- [3] Buchberger B., *An algorithmical criteria for the solvability of algebraic systems of equations*, Aequationes Mathematicae **4** (1970), 374–383.
- [4] Cohn P.M., *On homomorphic images of special Jordan algebras*, Canadian Journal of Mathematics **6** (1954), 253–264.
- [5] Chapoton F., *Un endofoncteur de la catégorie des opérades*, In: Dialgebras and Related Operads. Lecture Notes in Mathematics, **1763**, Springer, Berlin, Heidelberg, 2001.
- [6] Dangovski R., Drensky V., Findik Ş., *Weitzenböck derivations of free metabelian Lie algebras*, Linear Algebra and its Applications, **439** (2013), 3279–3296.
- [7] Dotsenko V., Heijltjes W., *Gröbner bases for operads*, <http://irma.math.unistra.fr/~dotsenko/Operads.html> (2019).
- [8] Dotsenko V., Khoroshkin A., *Gröbner bases for operads*, Duke Math. J. 2010. V. 153, N 2. P. 363–396.
- [9] Dzhumadil'daev A.S., and Ismailov N.A., *Polynomial identities of bicommutative algebras, Lie and Jordan elements*, Communication in algebra, **46** (2018), 5241–5251.
- [10] Gauger M. A., *Duality Theories for Metabelian Lie Algebras*, Transactions of the American Mathematical Society, **187** (1974), 89–102.
- [11] Scheuneman J., *Two-step nilpotent Lie algebras*, Journal of Algebra, **7** (1967), 152–159.
- [12] Kolesnikov P.S., *Gröbner—Shirshov bases for replicated algebras*, Algebra Colloquium **24** (2017), 563–576.

- [13] Kolesnikov P.S., Sartayev B.K., *On the embedding of left-symmetric algebras into differential Perm algebras*, arXiv:2106.00367.
- [14] Kolesnikov P.S., Sartayev B.K., *On the special identities of Gelfand–Dorfman algebras*, <https://arxiv.org/abs/2105.13815>.
- [15] Kurlin V. *The Baker–Campbell–Hausdorff Formula in the Free Metabelian Lie Algebra*, Journal of Lie Theory, **17** (2007), 525–538.
- [16] Leger G., Luks E., *On a duality for metabelian Lie algebras*, Journal of Algebra **21** (1972), 266–270.
- [17] Romankov V., *On the automorphism group of a free metabelian Lie algebra*, International Journal of Algebra and Computation **18**(2) (2008), 209–226.
- [18] Mikhalev A. A., Shestakov I. P., *PBW-pairs of varieties of linear algebras*, Commun. in Algebra **42** (2014) 667–687.
- [19] Specht W., *Die lineare Beziehungen zwischen höheren Kommutatoren*, Mathematische Zeitschrift volume **51** (1948), 367–376.
- [20] Zinbiel G.W., *Encyclopedia of types of algebras*, Proceedings of the International Conference in Nankai Series in Pure Applied Mathematics and Theoretical Physics **9** (2012), 217–298.

INSTITUTE OF MATHEMATICS AND MATHEMATICAL MODELING, ALMATY, KAZAKHSTAN, SULEYMAN DEMIREL UNIVERSITY, KASKELEN, KAZAKHSTAN
Email address: `f.mashurov@gmail.com`

SOBOLEV INSTITUTE OF MATHEMATICS, NOVOSIBIRSK, RUSSIA AND SULEYMAN DEMIREL UNIVERSITY, KASKELEN, KAZAKHSTAN
Email address: `baurjai@gmail.com`