

Maskin Meets Abreu and Matsushima*

Yi-Chun Chen[†] Takashi Kunimoto[‡] Yifei Sun[§] Siyang Xiong[¶]

October 14, 2021

Abstract

The theory of full implementation has been criticized for using integer/modulo games which admit no equilibrium (Jackson (1992)). To address the critique, we revisit the classical Nash implementation problem due to Maskin (1999) but allow for the use of lotteries and monetary transfers as in Abreu and Matsushima (1992, 1994). We unify the two well-established but somewhat orthogonal approaches in full implementation theory. We show that Maskin monotonicity is a necessary and sufficient condition for (exact) mixed-strategy Nash implementation by a finite mechanism. In contrast to previous papers, our approach possesses the following features: finite mechanisms (with no integer or modulo game) are used; mixed strategies are handled explicitly; neither undesirable outcomes nor transfers occur in equilibrium; the size of transfers can be made arbitrarily small; and our mechanism is robust to information perturbations. Finally, our result can be extended to infinite/continuous settings and ordinal settings.

*We owe special thanks to a coeditor and four anonymous referees for their insightful comments as well as to Kim-Sau Chung, Eddie Dekel, and Phil Reny for long discussions which led to major improvements of the paper. We also thank Soumen Banerjee, Olivier Bochet, Matthew Jackson, Hitoshi Matsushima, Stephen Morris, Daisuke Nakajima, Hamid Sabourian, Roberto Serrano, Balázs Szentes, Xiangqian Yang, and participants at various seminars and conference presentations for helpful comments. Part of this paper was written while the four authors were visiting Academia Sinica, and we would like to thank the institution for its hospitality and support.

[†]Department of Economics and Risk Management Institute, National University of Singapore. Email: ecsycc@nus.edu.sg

[‡]School of Economics, Singapore Management University. Email: tkunimoto@smu.edu.sg

[§]School of International Trade and Economics, University of International Business and Economics. Email: sunyifei@uibe.edu.cn

[¶]Department of Economics, University of California, Riverside. Email: siyang.xiong@ucr.edu

1 Introduction

Implementation theory can be seen as reverse engineering game theory. Suppose that a society has decided on a social choice rule – a recipe for choosing the socially-optimal alternatives on the basis of individuals’ preferences over alternatives. The individuals’ preferences vary across states and the realized state is common knowledge among the agents but unknown to a social planner/mechanism designer. To (fully) implement the social choice rule, the designer chooses a mechanism so that at each state, the equilibrium outcomes of the mechanism coincide with the outcomes designated by the social choice rule.

We study Nash implementation by a finite mechanism where agents report only their preferences and preference profiles. We focus on the monotonicity condition (hereafter, Maskin monotonicity) which Maskin shows is necessary and “almost sufficient” for Nash implementation. We aim to implement *social choice functions* (henceforth, SCFs) that are Maskin-monotonic in mixed-strategy Nash equilibria without making use of the *integer game* or the *modulo game* which prevails in the full implementation literature.

In the integer game, each agent announces some integer and the person who announces the highest integer gets to name his favorite outcome. When the agents’ favorite outcomes differ, an integer game has no pure-strategy Nash equilibria. The questionable feature is also shared by modulo games. The modulo game is considered a finite version of the integer game in which agents announce integers from a finite set. The agent who makes the integer announcement matching the modulo of the sum of the integers gets to name the allocation. In order to “knock out” undesirable equilibria in general environments, most constructive proofs in the literature, following Maskin (1999), have either taken advantage of the fact that the integer/modulo game has no solution in pure strategies and/or restricted attention to pure-strategy Nash equilibria.

Instead of invoking integer/modulo games, we study Nash implementation in a restricted domain where the designer can invoke both lotteries and (off-the-equilibrium) transfers in designing the implementing mechanism. We study a finite environment in which a finite mechanism is to be anticipated.¹ Finite mechanisms are also bounded in the sense of Jackson (1992) and have no aforementioned questionable features. Indeed, Jackson (1992,

¹More precisely, the implementing mechanism which we construct is finite as long as both the set of agents and the set of states are finite. In Section 5.4, we consider infinite environments in which we construct an infinite yet “well-behaved” implementing mechanism to achieve the same goal.

Example 4) shows that when no domain restriction on the environment is imposed, some Maskin-monotonic SCF is not implementable in mixed-strategy Nash equilibria by any finite mechanism. It raises the question as to whether *every* Maskin-monotonic SCF is mixed-strategy Nash implementable with domain restrictions imposed by lotteries and transfers.²

Our main result (Theorem 1) shows that when the designer can make use of lotteries and transfers off the equilibrium, Maskin monotonicity is indeed a necessary and sufficient condition for mixed-strategy Nash implementation by a finite mechanism. In our finite mechanism, each agent is asked to report only his preference and a preference profile. That is, we replace the integer announcement in Maskin’s mechanism with an announcement of each agent’s own preference. The preference announcement plays the same role as an integer in knocking out unwanted equilibria, albeit in a different manner. Following the idea of [Abreu and Matsushima \(1994\)](#), we design the mechanism so that whenever an “unwanted equilibrium” occurs, the agents’ reports must be “truthful,” namely, they announce their own preference and preference profile as prescribed under the true state. That in turn implies, through cross-checking the (truthful) preferences and the preference profiles reported by the agents, that the unwanted equilibrium could not have happened. In our finite mechanism, a pure-strategy (truth-telling) equilibrium exists, and all mixed-strategy equilibria achieve the desirable social outcome in each state.

We also provide several extensions of our main results. First, we show that our implementation is robust to information perturbations. Second, we extend Theorem 1 to cover social choice correspondences (i.e., multi-valued social choice rules), studied in [Maskin \(1999\)](#) as well as in many subsequent papers. Formally, we show that when there are at least three agents, every Maskin-monotonic social choice correspondence is mixed-strategy Nash implementable (Theorem 2). Moreover, as long as the social choice correspondence is finite-valued, our implementing mechanism remains finite. Third, we show that if there are at least three agents and the SCF satisfies Maskin monotonicity in the restricted domain without any transfer, then it is implementable in mixed-strategy Nash equilibria by a finite mechanism in which the size of transfers remains zero on the equilibrium and can be made arbitrarily small off the equilibrium (Theorem 3). Fourth, we consider an infinite setting in which the state space is a compact set, and the utility functions and the SCF are all continuous. In this

²Another direction one can take is to characterize, without making any domain restriction, the subclass of Maskin-monotonic SCFs which can be implemented in mixed-strategy Nash equilibria in a finite mechanism. For that goal, our exercise serves as a clarification of whether in certain environments, the class of implementable SCFs is as permissive as it can be.

setting, we show that Maskin monotonicity is a necessary and sufficient condition for mixed-strategy Nash implementation by a mechanism with a compact message space, a continuous outcome function, and a continuous transfer rule (Theorem 4). The extension covers many applications and verifies that our finite settings approximates settings with a continuum of states. To our knowledge, such an extension to an infinite setting has not appeared in the literature. Moreover, a compact and continuous mechanism allows each player to find a best response to every (pure or mixed) strategy profiles of the other players. Indeed, in discussing the notion of “well-behaved” mechanisms, [Abreu and Matsushima \(1992, footnote 8\)](#) also regard compactness and continuity as “plausible necessary desiderata”.

Finally, the extension to an infinite setting yields another interesting extension. Specifically, in proving Theorem 1, we have assumed that each agent is an expected utility maximizer with a fixed cardinal utility function over pure alternatives. This raises the question as to whether our result is an artifact of the fixed finite set of cardinalizations. To answer the question, we study the concept of *ordinal* Nash implementation proposed by [Mezzetti and Renou \(2012a\)](#). The notion requires that the implementing mechanism achieve mixed-strategy Nash implementation for *every* cardinal representation of preferences over lotteries. By making use of our implementing mechanism in the infinite setting, we show that ordinal almost monotonicity, as defined in [Sanver \(2006\)](#), is a necessary and sufficient condition for ordinal Nash implementation (Theorem 5). This extension also verifies that our implementation result does not suffer from the critique raised by [Jackson, Palfrey, and Srivastava \(1994, p. 490\)](#) to the dependence of the result of [Abreu and Matsushima \(1994\)](#) on a finite set of cardinalizations.

The rest of the paper is organized as follows. In Section 2, we position our paper in the literature. In Section 3, we present the basic setup and definitions. Section 4 proves our main result. We discuss the extensions of our main result in Section 5 and conclude in Section 6. The appendix contains all proofs, which are omitted from the main text.

2 Related Literature

[Maskin \(1999\)](#) proposes the notion of Maskin monotonicity and implements a Maskin-monotonic social choice correspondence by constructing an infinite mechanism with integer games. While integer games are useful in achieving positive results in general settings, the

hope has been that for more specific environments, more realistic mechanisms, or mechanisms without the “questionable features,” may suffice. The research program has been proposed by [Jackson \(1992\)](#). One such class of specific environments is the one with lotteries and transfers which our paper, as well as the partial implementation literature, focuses on.

In environments with lotteries and transfers, [Abreu and Matsushima \(1992, 1994\)](#) obtain permissive full implementation results using finite mechanisms without the aforementioned questionable features. Like our implementing mechanisms, their mechanisms also make use of only payoff relevant messages, such as preferences or preference profiles. However, [Abreu and Matsushima \(1992, 1994\)](#) do not investigate Nash implementation but rather appeal to a different notion of implementation: virtual implementation (in [Abreu and Matsushima \(1992\)](#)) or exact implementation under iterated weak dominance (in [Abreu and Matsushima \(1994\)](#)).³

Virtual implementation means that the planner contents herself with implementing the SCF with arbitrarily high probability. In contrast, by studying *exact* Nash implementation in the specific setting, we unify the two well-established but somewhat orthogonal approaches to implementation theory which are due to [Maskin \(1999\)](#) and to [Abreu and Matsushima \(1992, 1994\)](#). Our exercise is directly comparable to [Maskin \(1999\)](#) and highlights the pivotal trade-off between domain restrictions and the feature of implementing mechanisms. We consider it to be one step in advancing the research program proposed by [Jackson \(1992\)](#).

An alternative approach to handling mixed-strategy equilibria is to resort to refinements such as undominated Nash equilibria or subgame-perfect equilibria. With such refinements, essentially every SCF, whether Maskin-monotonic or not, is implementable in a complete-information environment; see, for example, [Moore and Repullo \(1988\)](#) and [Abreu and Matsushima \(1994\)](#). However, according to [Chung and Ely \(2003\)](#) and [Aghion, Fudenberg, Holden, Kunimoto, and Tercieux \(2012\)](#), if we were to achieve exact implementation in these refinements which are also robust to a *small amount of incomplete information*, then Maskin monotonicity would be restored as a necessary condition. Those permissive implementation results, which are driven by the lack of the closed-graph property of the refinements, cast doubt on the success of taking care of non-Maskin-monotonic SCFs by resorting to equilibrium refinements. In contrast, our [Theorem 1](#) establishes exact and *ro-*

³Iterated weak dominance in [Abreu and Matsushima \(1994\)](#) also yields the unique undominated Nash equilibrium outcome. For undominated Nash implementation by “well-behaved” mechanisms, see also [Jackson, Palfrey, and Srivastava \(1994\)](#) and [Sjöström \(1994\)](#).

best implementation in mixed-strategy Nash equilibria to the maximal extent that every Maskin-monotonic SCF is implementable (Proposition 3).⁴

Ollár and Penta (2017) study a full implementation problem using transfers both on and off the equilibrium. Specifically, Theorem 2 of Ollár and Penta (2017) provides a sufficient condition restricting agents’ beliefs via moment conditions under which their notion of robust full implementation is possible in a direct mechanism. Their notion of robustness is a “global notion” which accommodates arbitrary information structures consistent with a fixed payoff environment.⁵ In contrast, our paper follows the classical implementation literature in dealing with the specific belief restriction of complete information, and our notion of robustness (in Section 5.1) accommodates only perturbations around the complete-information benchmark. With the specific belief restriction, we prove that Maskin monotonicity is both necessary and sufficient for implementation in mixed-strategy Nash equilibria in a finite, indirect mechanism with only off-the-equilibrium transfers.

3 Preliminaries

3.1 Environment

Consider a finite set of agents $\mathcal{I} = \{1, 2, \dots, I\}$ with $I \geq 2$; a finite set of possible states Θ ; and a set of pure alternatives A . We consider an environment with lotteries and transfers. Specifically, we work with the space of allocations/outcomes $X \equiv \Delta(A) \times \mathbb{R}^I$ where $\Delta(A)$ denotes the set of lotteries on A that have a countable support, and \mathbb{R}^I denotes the set of transfers to the agents. We identify $a \in A$ with a degenerate lottery in $\Delta(A)$.

For each $x = (\ell, (t_i)_{i \in \mathcal{I}}) \in X$, agent i receives the utility $\tilde{u}_i(x, \theta) = v_i(\ell, \theta) + t_i$ for some bounded expected utility function $v_i(\cdot, \theta)$ over $\Delta(A)$. That is, we work with an environment with a transferable utility (TU) on agents’ preferences, which Maskin (1999) does not impose. We abuse notation to identify $\Delta(A)$ with a subset of X , i.e., each $\ell \in \Delta(A)$ is identified with the allocation $(\ell, 0, \dots, 0)$ in X .

We focus on a *complete-information* environment in which a true state θ is common

⁴Harsanyi (1973) shows that a mixed-strategy Nash equilibrium outcome may occur as the limit of a sequence of pure-strategy Bayesian Nash equilibria for “nearby games” in which players are uncertain about the exact profile of preferences. Hence, ignoring mixed-strategy equilibria would be particularly problematic if we were to achieve implementation which is robust to information perturbations.

⁵See Ollár and Penta (2021) for a further extension of the approach of Ollár and Penta (2017).

knowledge among the agents but unknown to a mechanism designer. The designer's objective is specified by a *social choice function* $f : \Theta \rightarrow X$, namely, if the state is θ , the designer would like to implement the social outcome $f(\theta)$.

3.2 Mechanism and Solution

Let $\mathcal{M} = ((M_i, \tau_i)_{i \in \mathcal{I}}, g)$ be a finite mechanism where M_i is a nonempty finite *set of messages* available to agent i ; $g : M \rightarrow X$ (where $M \equiv \times_{i=1}^I M_i$) is the *outcome function*; and $\tau_i : M \rightarrow \mathbb{R}$ is the *transfer rule* which specifies the payment to agent i . At each state $\theta \in \Theta$, the environment and the mechanism together constitute a *game with complete information* which we denote by $\Gamma(\mathcal{M}, \theta)$. Note that the restriction of M_i to a finite set rules out the use of integer games à la Maskin (1999).

Let $\sigma_i \in \Delta(M_i)$ be a mixed *strategy* of agent i in the game $\Gamma(\mathcal{M}, \theta)$. A strategy profile $\sigma = (\sigma_1, \dots, \sigma_I) \in \times_{i \in \mathcal{I}} \Delta(M_i)$ is said to be a mixed-strategy *Nash equilibrium* of the game $\Gamma(\mathcal{M}, \theta)$ if, for all agents $i \in \mathcal{I}$ and all messages $m_i \in \text{supp}(\sigma_i)$ and $m'_i \in M_i$, we have

$$\begin{aligned} & \sum_{m_{-i} \in M_{-i}} \prod_{j \neq i} \sigma_j(m_j) [\tilde{u}_i(g(m_i, m_{-i}), \theta) + \tau_i(m_i, m_{-i})] \\ & \geq \sum_{m_{-i} \in M_{-i}} \prod_{j \neq i} \sigma_j(m_j) [\tilde{u}_i(g(m'_i, m_{-i}), \theta) + \tau_i(m'_i, m_{-i})]. \end{aligned}$$

A pure-strategy Nash equilibrium is a mixed-strategy Nash equilibrium σ such that each agent i 's mixed-strategy σ_i assigns probability one to some $m_i \in M_i$. For any message profile $m \in M$, let $\sigma(m) \equiv \prod_{j \in \mathcal{I}} \sigma_j(m_j)$.

Let $NE(\Gamma(\mathcal{M}, \theta))$ denote the set of mixed-strategy Nash equilibria of the game $\Gamma(\mathcal{M}, \theta)$. We also denote by $\text{supp}(NE(\Gamma(\mathcal{M}, \theta)))$ the set of message profiles that can be played with positive probability under some mixed-strategy Nash equilibrium $\sigma \in NE(\Gamma(\mathcal{M}, \theta))$, i.e.,

$$\text{supp}(NE(\Gamma(\mathcal{M}, \theta))) = \{m \in M : \text{there exists } \sigma \in NE(\Gamma(\mathcal{M}, \theta)) \text{ such that } \sigma(m) > 0\}.$$

We now define our notion of Nash implementation.

Definition 1 *An SCF f is implementable in mixed-strategy Nash equilibria by a finite mechanism if there exists a finite mechanism $\mathcal{M} = ((M_i, \tau_i)_{i \in \mathcal{I}}, g)$ such that for every state $\theta \in \Theta$, (i) there exists a pure-strategy Nash equilibrium in the game $\Gamma(\mathcal{M}, \theta)$;*

and (ii) $m \in \text{supp} (NE(\Gamma(\mathcal{M}, \theta))) \Rightarrow g(m) = f(\theta)$ and $\tau_i(m) = 0$ for every $i \in \mathcal{I}$.

Our definition is adapted from mixed-strategy Nash implementation in Maskin (1999) to (1) require that the implementing mechanism be finite; and (2) accommodate our quasi-linear environments with transfers. In particular, Condition (ii) requires that transfers be imposed only off the equilibrium. Mezzetti and Renou (2012a) propose another definition of Nash implementation that keeps Condition (ii) but weakens Condition (i) in requiring only the existence of a mixed-strategy Nash equilibrium, which, by Nash's theorem, is guaranteed in a finite mechanism. In their sufficiency result, however, Mezzetti and Renou (2012a) construct an infinite mechanism with integer games.

3.3 Maskin Monotonicity

We now restate the definition of Maskin monotonicity that Maskin (1999) proposes for Nash implementation.

Definition 2 *An SCF f satisfies **Maskin monotonicity** if, for every pair of states $\tilde{\theta}$ and θ with $f(\tilde{\theta}) \neq f(\theta)$, some agent $i \in \mathcal{I}$ and some allocation $x \in X$ exist such that*

$$\tilde{u}_i(x, \tilde{\theta}) \leq \tilde{u}_i(f(\tilde{\theta}), \tilde{\theta}) \text{ and } \tilde{u}_i(x, \theta) > \tilde{u}_i(f(\tilde{\theta}), \theta). \quad (1)$$

To illustrate how the idea of Maskin monotonicity is applied, suppose that the SCF f is implemented in Nash equilibria by a mechanism. When $\tilde{\theta}$ is the true state, there exists a pure-strategy Nash equilibrium $m \in M$ in $\Gamma(\mathcal{M}, \tilde{\theta})$ which induces $f(\tilde{\theta})$. If $f(\tilde{\theta}) \neq f(\theta)$ and θ is the true state, then m cannot be a Nash equilibrium, i.e., there exists some agent i who has a profitable deviation. Suppose that the deviation induces outcome x , i.e., agent i strictly prefers x to $f(\tilde{\theta})$ at state θ . Since m is a Nash equilibrium at state $\tilde{\theta}$, such a deviation cannot be profitable in state $\tilde{\theta}$; that is, agent i weakly prefers $f(\tilde{\theta})$ to x at state $\tilde{\theta}$. In other words, x belongs to agent i 's lower contour set at $f(\tilde{\theta})$ of state $\tilde{\theta}$, whereas it belongs to the strict upper-contour set at $f(\tilde{\theta})$ of state θ . Therefore, Maskin monotonicity is a necessary condition for Nash implementation; in fact, it is a necessary condition even for Nash implementation that restricts attention to pure-strategy equilibria (i.e., to require that condition (ii) of Definition 1 hold only for pure-strategy Nash equilibria).

4 Main Result

In this section, we present our main result, which shows that Maskin monotonicity is necessary and sufficient for mixed-strategy Nash implementation. We formally state the result as follows.

Theorem 1 *An SCF f is implementable in mixed-strategy Nash equilibria by a finite mechanism if and only if it satisfies Maskin monotonicity.*

In the rest of this section, we will establish Theorem 1 and discuss the issues regarding the theorem. Section 4.1 details how our implementing mechanism is constructed. In Section 4.2, we prove Theorem 1 by making use of the implementing mechanism constructed in Section 4.1. Section 4.3 illustrates two special cases in which our implementing mechanism can be made into a direct mechanism where each agent reports a state. In Section 4.4, we discuss the necessity of domain restrictions in establishing Theorem 1.

4.1 The Mechanism

We construct a mechanism $\mathcal{M} = ((M_i, \tau_i)_{i \in \mathcal{I}}, g)$ which will be used to prove Theorem 1. The mechanism shares a number of features of the implementing mechanisms in Maskin (1999) and in Abreu and Matsushima (1992, 1994), which we summarize at the end of the subsection. The construction involves two major building blocks that we call the *best challenge scheme* and *dictator lotteries*, respectively. After introducing these building blocks, we will define the message space, allocation rule, and transfer rule of our implementing mechanism.

For each agent i , as a preliminary step, we define

$$\Theta_i \equiv \{v_i(\cdot, \theta) : \theta \in \Theta\}.$$

That is, Θ_i is the set of expected utility functions of agent i induced by some state θ . Denote by $\theta_i \in \Theta_i$ the expected utility function of agent i obtained at state $\theta \in \Theta$, namely, that $\theta_i = v_i(\cdot, \theta)$. We call θ_i the type of player i at state θ . We denote by $u_i(\cdot, \theta_i)$ the quasilinear utility function which corresponds to type θ_i , namely, that for each $x = (\ell, (t_i)_{i \in \mathcal{I}}) \in X$, we have $u_i(x, \theta_i) \equiv v_i(\ell, \theta) + t_i$.

Remark 1 Note that every state θ uniquely determines a type profile $(\theta_i)_{i \in \mathcal{I}}$, while an arbitrary type profile may not identify a state in Θ . For example, suppose we have two states, $\{\alpha, \beta\}$, and two agents who have the same expected utility function at each state such that $\alpha_A = \alpha_B \neq \beta_A = \beta_B$. In this example, $\{(\alpha_A, \alpha_B), (\alpha_A, \beta_B), (\beta_A, \alpha_B), (\beta_A, \beta_B)\}$ are four type profiles, and yet neither the type profile (α_A, β_B) nor (β_A, α_B) corresponds to a state.

4.1.1 Best Challenge Scheme

For $(x, \theta_i) \in X \times \Theta_i$, we use $\mathcal{L}_i(x, \theta_i)$ to denote the lower-contour set at allocation x in X for type θ_i , i.e.,

$$\mathcal{L}_i(x, \theta_i) = \{x' \in X : u_i(x, \theta_i) \geq u_i(x', \theta_i)\}.$$

We use $\mathcal{SU}_i(x, \theta_i)$ to denote the strict upper-contour set of $x \in X$ for type θ_i , i.e.,

$$\mathcal{SU}_i(x, \theta_i) = \{x' \in X : u_i(x', \theta_i) > u_i(x, \theta_i)\}.$$

Hence, according to Definition 2, an SCF f satisfies Maskin monotonicity if and only if for every pair of states $\tilde{\theta}$ and θ in Θ ,

$$f(\tilde{\theta}) \neq f(\theta) \Rightarrow \exists i \in \mathcal{I} \text{ s.t. } \mathcal{L}_i(f(\tilde{\theta}), \tilde{\theta}_i) \cap \mathcal{SU}_i(f(\tilde{\theta}), \tilde{\theta}_i) \neq \emptyset. \quad (2)$$

Agent i in (2) is called a “whistle-blower” or a “test agent,” and an allocation in $\mathcal{L}_i(f(\tilde{\theta}), \tilde{\theta}_i) \cap \mathcal{SU}_i(f(\tilde{\theta}), \tilde{\theta}_i)$ is called a “test allocation” for agent i and the ordered pair of states $(\tilde{\theta}, \theta)$. We now define a notion called *the best challenge scheme*, which plays a crucial role in proving Theorem 1. We say that a mapping $x : \Theta \times \Theta_i \rightarrow X$ is a *challenge scheme* for an SCF f if and only if, for each pair of state $\tilde{\theta} \in \Theta$ and type $\theta_i \in \Theta_i$,

$$\begin{cases} x(\tilde{\theta}, \theta_i) \in \mathcal{L}_i(f(\tilde{\theta}), \tilde{\theta}_i) \cap \mathcal{SU}_i(f(\tilde{\theta}), \tilde{\theta}_i), & \text{if } \mathcal{L}_i(f(\tilde{\theta}), \tilde{\theta}_i) \cap \mathcal{SU}_i(f(\tilde{\theta}), \tilde{\theta}_i) \neq \emptyset; \\ x(\tilde{\theta}, \theta_i) = f(\tilde{\theta}), & \text{if } \mathcal{L}_i(f(\tilde{\theta}), \tilde{\theta}_i) \cap \mathcal{SU}_i(f(\tilde{\theta}), \tilde{\theta}_i) = \emptyset. \end{cases}$$

We may think of state $\tilde{\theta}$ as an announcement made by one or more other agents that agent i of type θ_i could “challenge” (as a whistle-blower). The following lemma shows that there is a challenge scheme in which each whistle-blower i facing state announcement $\tilde{\theta}$ finds it weakly best to challenge $\tilde{\theta}$ by simply reporting his true type θ_i .

Lemma 1 *There is a challenge scheme $x(\cdot, \cdot)$ for an SCF f such that for every state $\tilde{\theta}$ and type θ_i ,*

$$u_i(x(\tilde{\theta}, \theta_i), \theta_i) \geq u_i(x(\tilde{\theta}, \theta'_i), \theta_i), \forall \theta'_i \in \Theta_i. \quad (3)$$

We relegate its formal proof to Appendix A.1.⁶ In defining the implementing mechanism, we shall invoke a challenge scheme which satisfies (3). We call such a challenge scheme *the best challenge scheme*. In words, under the best challenge scheme, for any state $\tilde{\theta}$, agent i of type θ_i weakly prefers the allocation $x(\tilde{\theta}, \theta_i)$ to any other $x(\tilde{\theta}, \theta'_i)$ induced by announcing $\theta'_i \neq \theta_i$.

4.1.2 Dictator Lotteries

Let $\tilde{X} \equiv A \cup \bigcup_{i \in \mathcal{I}, \theta_i \in \Theta_i, \tilde{\theta} \in \Theta} x(\tilde{\theta}, \theta_i)$. We can then conclude that \tilde{X} is a finite set over which all agents' utilities are bounded, because $v_i(\cdot, \theta)$ is bounded, Θ is finite, and we pre-specify $x(\tilde{\theta}, \theta_i)$ for each $i \in \mathcal{I}$, type $\theta_i \in \Theta_i$, and state $\tilde{\theta} \in \Theta$. Hence, we can choose $\eta' > 0$ as an upper bound on the monetary value of a change of allocation in \tilde{X} , that is,

$$\eta' > \sup_{i \in \mathcal{I}, \theta_i \in \Theta_i, x, x' \in \tilde{X}} |u_i(x, \theta_i) - u_i(x', \theta_i)|. \quad (4)$$

We now state a result which ensures the existence of what we call *dictator lotteries* for agent i . In particular, a collection of lotteries is called dictator lotteries of agent i if it satisfies Conditions (5) and (6) stated in Lemma 2. Condition (5) shows that within dictator lotteries, each agent has a strict incentive to reveal his true type, whereas Condition (6) says that these dictator lotteries are strictly less preferred than any allocations in \tilde{X} .

Lemma 2 *For each agent $i \in \mathcal{I}$, there exists a collection of lotteries $\{y_i(\theta_i)\}_{\theta_i \in \Theta_i}$ such that for all types $\theta_i, \theta'_i \in \Theta_i$ with $\theta_i \neq \theta'_i$, we have*

$$u_i(y_i(\theta_i), \theta_i) > u_i(y_i(\theta'_i), \theta_i); \quad (5)$$

moreover, for each $j \in \mathcal{I}$ and type $\theta'_j \in \Theta_j$, we also have that, for every $x \in \tilde{X}$,

$$u_i(y_j(\theta'_j), \theta_i) < u_i(x, \theta_i). \quad (6)$$

⁶We owe special thanks to Phil Reny for suggesting the lemma which simplifies the implementing mechanism adopted in an earlier version of our paper.

Since two distinct types in Θ_i induce different expected utility functions over $\Delta(A)$, it follows from (Abreu and Matsushima, 1992, Lemma) that we can prove the existence of lotteries $\{y'_i(\cdot)\} \subset \Delta(A)$ that satisfy Condition (5). To satisfy Condition (6), we simply add a penalty of η' to each outcome of the lotteries $\{y'_i(\theta_i)\}_{\theta_i \in \Theta_i}$. More precisely, for each $\theta_i \in \Theta_i$, we set

$$y_i(\theta_i) = (y'_i(\theta_i), -\eta', \dots, -\eta') \in X.$$

4.1.3 Message Space

A generic message of agent i is described as follows:

$$m_i = (m_i^1, m_i^2) \in M_i = M_i^1 \times M_i^2 = \Theta_i \times [\times_{j=1}^I \Theta_j].$$

That is, agent i is asked to make (1) a report of his own type (which we denote by m_i^1); and (2) a report of a type profile (which we denote by m_i^2). To simplify the notation, we write $m_{i,j}^2 = \tilde{\theta}_j$ if agent i reports in m_i^2 that agent j is of type $\tilde{\theta}_j$. Recall that agents have complete information about the true state. If the true state is θ , we say that agent i sends a truthful first report if $m_i^1 = \theta_i$ and a truthful second report if $m_i^2 = (\theta_j)_{j \in \mathcal{I}}$. Recall the difficulties in identifying the state from a type profile which we explain Remark 1. Here each agent is asked to report a type profile in M_i^2 instead of a state.

It is useful to compare the message space of our mechanism with that of the implementing mechanism in Maskin (1999). In Maskin's mechanism (see Maskin (1999, p. 31)), each agent is asked to report a preference profile and an integer, as well as an allocation. The allocation need not be specified in the case of SCFs, since there is no ambiguity about the socially desirable outcome assigned to each state. In contrast, we ask each agent to report a preference/type profile and a *type*. The type component of the message space plays the role of an integer in Maskin's mechanism in knocking out unwanted equilibria, albeit in a different manner. As the integer game admits no equilibrium when there is disagreement over most preferred outcomes, it is used to assure that undesirable message profiles do not form an equilibrium. However, the logic of their argument no longer works when the goal is to achieve implementation in mixed-strategy Nash equilibria by a finite mechanism. Indeed, there is no a priori obvious way to rule out *any* message profile because any of them might be played with positive probability in a mixed-strategy Nash equilibrium.

By making use of the type component m_i^1 in the message space, we appeal to the

approach of Abreu and Matsushima (1992, 1994)) to resolve the issue. More precisely, we design the mechanism so that when an unwanted message profile is triggered in equilibrium, the type report m_i^1 must coincide with agent i 's preference under the true state. Through the cross-checking of the preferences and preference profiles reported by the agents (in a similar manner to Abreu and Matsushima (1992, 1994)), it further implies that the unwanted message profile could not have happened. Unlike Abreu and Matsushima (1992, 1994), however, to ensure that m_i^1 is truthful, we must guarantee that the designer's twin goals of allowing for whistle-blowing/challenges (as in Maskin (1999)) and eliciting the truth (from the dictator lotteries, as in Abreu and Matsushima (1992, 1994)) can be aligned perfectly. It is achieved through Lemmas 1 and 2: since truth-telling is weakly optimal for the former and strictly optimal for the latter, we can make the truth-telling in m_i^1 strictly optimal by taking a convex combination of the best challenge scheme and dictator lotteries. Hence, Maskin meets Abreu and Matsushima. We formalize the idea in Section 4.1.4.

4.1.4 Allocation Rule

For each message profile $m \in M$, the allocation is determined as follows:

$$g(m) = \frac{1}{I(I-1)} \sum_{i \in \mathcal{I}} \sum_{j \neq i} \left[e_{i,j}(m_i, m_j) \left(\frac{1}{2} \sum_{k=i,j} y_k(m_k^1) \right) \oplus (1 - e_{i,j}(m_i, m_j)) x(m_i^2, m_j^1) \right],$$

where $\{y_k(\cdot)\}$ are the dictator lotteries for agent k obtained from Lemma 2, and $\alpha x \oplus (1 - \alpha) x'$ denotes the outcome which corresponds to the compound lottery that outcome x occurs with probability α , and outcome x' occurs with probability $1 - \alpha$;⁷ moreover, we define

$$e_{i,j}(m_i, m_j) = \begin{cases} 0, & \text{if } m_i^2 \in \Theta, m_i^2 = m_j^2, \text{ and } x(m_i^2, m_j^1) = f(m_i^2); \\ \varepsilon, & \text{if } m_i^2 \in \Theta, \text{ and } [m_i^2 \neq m_j^2 \text{ or } x(m_i^2, m_j^1) \neq f(m_i^2)]; \\ 1, & \text{if } m_i^2 \notin \Theta. \end{cases}$$

In what follows, we say that the second reports of agent i and agent j are *consistent* if $m_i^2 = m_j^2$ and the common type profile identifies a state in Θ ; moreover, we say that agent

⁷More precisely, if $x = (\ell, (t_i)_{i \in \mathcal{I}})$ and $x' = (\ell', (t'_i)_{i \in \mathcal{I}})$ are two outcomes in X , we identify $\alpha x \oplus (1 - \alpha) x'$ with the outcome $(\alpha \ell \oplus (1 - \alpha) \ell', (\alpha t_i + (1 - \alpha) t'_i)_{i \in \mathcal{I}})$. For simplicity, we also write the compound lottery $\frac{1}{2} y_i(m_i^1) \oplus \frac{1}{2} y_j(m_j^1)$ as $\frac{1}{2} \sum_{k=i,j} y_k(m_k^1)$.

j does not challenge agent i if $x(m_i^2, m_j^1) = f(m_i^2)$.⁸

In words, the designer first chooses an ordered pair of distinct agents (i, j) with equal probability. The outcome function distinguishes three cases: (1) if the second reports of agent i and agent j are consistent *and* agent j does not challenge agent i , then we implement $f(m_i^2)$; (2) if agent i reports a type profile which does not identify a state in Θ , then we implement the dictator lottery $\frac{1}{2} \sum_{k=i,j} y_k(m_k^1)$; (3) otherwise, we implement the compound lottery:

$$C_{i,j}^\varepsilon(m_i, m_j) \equiv \varepsilon \left(\frac{1}{2} \sum_{k=i,j} y_k(m_k^1) \right) \oplus (1 - \varepsilon) x(m_i^2, m_j^1).$$

That is, $C_{i,j}^\varepsilon(m_i, m_j)$ is an $(\varepsilon, 1 - \varepsilon)$ -combination of (i) the two dictator lotteries— $y_i(m_i^1)$ and $y_j(m_j^1)$ —which occur with equal probability; and (ii) the allocation specified by the best challenge scheme $x(m_i^2, m_j^1)$.

By (4), we can choose $\varepsilon > 0$ sufficiently small, and $\eta > 0$ sufficiently large⁹ such that firstly we have

$$\eta > \sup_{i \in \mathcal{I}, \theta_i \in \Theta_i, m, m' \in M} |u_i(g(m), \theta_i) - u_i(g(m'), \theta_i)|; \quad (7)$$

secondly it does not disturb the “effectiveness” of agent j ’s challenge; put formally,

$$\begin{aligned} x(m_i^2, m_j^1) &\neq f(m_i^2) \Rightarrow \\ u_j(C_{i,j}^\varepsilon(m_i, m_j), m_{i,j}^2) &< u_j(f(m_i^2), m_{i,j}^2) \text{ and } u_j(C_{i,j}^\varepsilon(m_i, m_j), m_j^1) > u_j(f(m_i^2), m_j^1). \end{aligned} \quad (8)$$

It means that whenever agent j challenges agent i , the lottery $C_{i,j}^\varepsilon(m_i, m_j)$ is strictly worse than $f(m_i^2)$ for agent j when agent i tells the truth about agent j ’s preference in m_i^2 ; moreover, the lottery $C_{i,j}^\varepsilon(m_i, m_j)$ is strictly better than $f(m_i^2)$ for agent j when agent j tells the truth in m_j^1 , a fact which implies that agent i tells a lie about agent j ’s preference.

⁸Observe that we make the first report of both agents i and j effective (through affecting the compound lottery $\frac{1}{2} \sum_{k=i,j} y_k(m_k^1)$), regardless of whether pair (i, j) or pair (j, i) is picked. The construction will be used in proving Claim 1, which, in turn, is used to prove Claim 4.

⁹Instead of using η' defined in (4), we choose η because the mechanism may produce a strictly larger finite set of alternatives than those contained in \tilde{X} . For instance, allocations from the dictator lotteries may occur from the mechanism but are not contained in \tilde{X} .

4.1.5 Transfer Rule

We now define the transfer rule. For every message profile $m \in M$ and every agent $i \in \mathcal{I}$, we specify the transfer received by agent i as follows:

$$\tau_i(m) = \sum_{j \neq i} [\tau_{i,j}^1(m_i, m_j) + \tau_{i,j}^2(m_i, m_j)],$$

where for each agent $j \neq i$, we define

$$\tau_{i,j}^1(m_i, m_j) = \begin{cases} 0, & \text{if } m_{i,j}^2 = m_{j,j}^2; \\ -\eta, & \text{if } m_{i,j}^2 \neq m_{j,j}^2 \text{ and } m_{i,j}^2 \neq m_j^1; \\ \eta, & \text{if } m_{i,j}^2 \neq m_{j,j}^2 \text{ and } m_{i,j}^2 = m_j^1. \end{cases} \quad (9)$$

$$\tau_{i,j}^2(m_i, m_j) = \begin{cases} 0, & \text{if } m_{i,i}^2 = m_{j,i}^2; \\ -\eta, & \text{if } m_{i,i}^2 \neq m_{j,i}^2. \end{cases} \quad (10)$$

Recall that $\eta > 0$ is chosen to be larger than the maximal utility difference from the outcome function $g(\cdot)$; see (7).

In words, for each pair of agents (i, j) , if their second reports on agent j 's type coincide ($m_{i,j}^2 = m_{j,j}^2$), then no transfer will be made; if their second reports on agent j 's type differ ($m_{i,j}^2 \neq m_{j,j}^2$), then we consider the following two subcases: (i) if agent i 's second report about agent j 's type matches agent j 's first report ($m_{i,j}^2 = m_j^1$), then agent j pays η to agent i ; (ii) if agent i 's second report about agent j 's type does not match agent j 's first report ($m_{i,j}^2 \neq m_j^1$), then both agents pay η to the designer. Note that the first report m_i^1 does not affect the transfer to agent i .

4.2 Proof of Theorem 1

As we argue in Section 3.3, Maskin monotonicity is a necessary condition for Nash implementation. We therefore focus on the “if” part of the proof. Fix an arbitrary true state θ throughout the proof. Recall that θ_i stands for agent i 's type at state θ and $(\theta_i)_{i \in \mathcal{I}}$ denotes the true type profile.

We argue that the truth-telling message profile m (i.e., $m_i = (\theta_i, \theta)$ for each agent i) constitutes a pure-strategy Nash equilibrium. Since m is truthful, for all agents i and j , we have $e_{i,j}(m_i, m_j) = 0$ and $\tau_i(m) = 0$ (consistency and no challenge). Consider a possible

deviation \tilde{m}_i of agent i from m . First, if $\tilde{m}_{i,j}^2 = \theta'_j \neq \theta_j$ for some $j \in \mathcal{I}$, then the message profile (\tilde{m}_i, m_{-i}) induces the penalty of η from rule $\tau_{i,j}^1(\cdot)$ if $j \neq i$, and rule $\tau_{i,j}^2(\cdot)$ if $j = i$. As a result, \tilde{m}_i is strictly worse against m_{-i} than m_{-i} .

Second, if $\tilde{m}_i^1 \neq \theta_i$ and $\tilde{m}_i^2 = \theta$, (\tilde{m}_i, m_{-i}) leads either to $x(\theta, \tilde{m}_i^1) = f(\theta)$ and thereby the same payoff, or to $x(\theta, \tilde{m}_i^1) \neq f(\theta)$. In the latter case, the message profile (\tilde{m}_i, m_{-i}) results in the outcome $C_{i,j}^\varepsilon(\tilde{m}_i, m_j)$ which, by (8), is strictly worse than $f(\theta)$ induced by m . Furthermore, deviating from m_i to \tilde{m}_i does not affect the transfer of agent i . Therefore, the truth-telling message profile m constitutes a pure-strategy Nash equilibrium.

We next show that for every Nash equilibrium σ of the game $\Gamma(\mathcal{M}, \theta)$ and every message profile m reported with positive probability under σ , we must achieve the socially desirable outcome, i.e., $g(m) = f(\theta)$ and $\tau_i(m) = 0$ for every agent i . The proof is divided into three steps.

Step 1: Contagion of truth. If agent j announces his type truthfully in his first report with probability one, then everyone must also report agent j 's type truthfully in their second report;

Step 2: Consistency. Every agent reports the same state $\tilde{\theta}$ in the second report;

Step 3: No challenge. No agent challenges the common reported state $\tilde{\theta}$, i.e., $x(\tilde{\theta}, m_j^1) = f(\tilde{\theta})$ for every agent $j \in \mathcal{I}$.

Consistency implies that $\tau_i(m) = 0$ for every agent $i \in \mathcal{I}$, whereas no challenge together with Maskin monotonicity of the SCF f implies that $g(m) = f(\tilde{\theta}) = f(\theta)$. It completes the proof of Theorem 1. We now proceed to establish these three steps. In the rest of the proof, we fix σ as an arbitrary mixed-strategy Nash equilibrium of the game $\Gamma(\mathcal{M}, \theta)$.

As a consequence of Lemmas 2 and 1, the mechanism has the following crucial property which we will make use of in establishing the implementation.

Claim 1 *Let σ be a Nash equilibrium of the game $\Gamma(\mathcal{M}, \theta)$. If $m_i^1 \neq \theta_i$ for some $m_i \in \text{supp}(\sigma_i)$, then for every agent $j \neq i$, we have $e_{i,j}(m_i, m_j) = e_{j,i}(m_j, m_i) = 0$ with σ_j -probability one.*

The claim essentially follows from Lemmas 1 and 2. Indeed, the two lemmas together imply that agents must have a strict incentive to tell the truth in their first report, as long as switching from a lie to truth affects the allocation with positive probability. A detailed verification of the claim, however, is tedious, as it involves checking different cases of the value of functions $e_{i,j}(\cdot)$ and $e_{j,i}(\cdot)$. We relegate its formal proof to Appendix A.4.

Step 1: Contagion of Truth

Claim 2 *The following two statements hold:*

- (a) *If agent j sends a truthful first report with σ_j -probability one, then every agent $i \neq j$ must report agent j 's type truthfully in his second report with σ_i -probability one.*
- (b) *If every agent $i \neq j$ reports the same type $\tilde{\theta}_j$ of agent j in his second report with σ_i -probability one, then agent j must also report the type $\tilde{\theta}_j$ in his second report with σ_j -probability one.*

Proof. We first prove (a). Suppose instead that there exist some agent $i \in \mathcal{I}$ and some message m_i played with σ_i -positive probability such that m_i misreports agent j 's type in the second report, i.e., $m_{i,j}^2 \neq \theta_j$. Let \tilde{m}_i be a message that differs from m_i only in reporting j 's type truthfully $\tilde{m}_{i,j}^2 = \theta_j$. Such a change affects only $\tau_{i,j}^1(\cdot)$. For every m_{-i} played with σ_{-i} -positive probability, we consider the following two cases.

Case 1: $m_{j,j}^2 = \theta_j$

Since agent j sends a truthful first report with σ_j -probability one, due to the construction of $\tau_{i,j}^1(\cdot)$, we have $\tau_{i,j}^1(m_i, m_{-i}) = -\eta$ whereas $\tau_{i,j}^1(\tilde{m}_i, m_{-i}) = 0$.

Case 2: $m_{j,j}^2 \neq \theta_j$

Since agent j sends a truthful type in the first report with σ_j -probability one, according to the construction of $\tau_{i,j}^1(\cdot)$, we have $\tau_{i,j}^1(m_i, m_{-i})$ is either 0 or $-\eta$ whereas $\tau_{i,j}^1(\tilde{m}_i, m_{-i}) = \eta$.

Thus, in terms of transfers, the gain from reporting \tilde{m}_i rather than m_i is at least η , which is larger than the maximal utility loss from the outcome function $g(\cdot)$ by (7). Hence, \tilde{m}_i is a profitable deviation from m_i against σ_{-i} . As it contradicts the hypothesis that $m_i \in \text{supp}(\sigma_i)$, we have established (a).

We now prove (b). Suppose, on the contrary, that there exists some message m_j played with σ_j -positive probability such that $m_{j,j}^2 \neq \tilde{\theta}_j$. Let \tilde{m}_j be a message that is identical to m_j except that $\tilde{m}_{j,j}^2 = \tilde{\theta}_j$. Such a change affects only $\tau_{j,i}^2(\cdot)$. According to the construction of $\tau_{j,i}^2(\cdot)$ and since every agent $i \neq j$ reports $\tilde{\theta}_j$ in the second report with σ_i -probability one, agent j saves the penalty of $(I-1)\eta$ from reporting \tilde{m}_j instead of m_j . Again, since η is greater than the maximal utility difference by (7), we conclude that \tilde{m}_j is a profitable deviation from m_j against σ_{-i} . It contradicts the hypothesis that $m_j \in \text{supp}(\sigma_j)$. Hence, we prove (b). ■

Step 2: Consistency

Claim 3 shows that in equilibrium, all agents must announce the same state $\tilde{\theta}$ with probability one.

Claim 3 *There exists a state $\tilde{\theta} \in \Theta$ such that every agent announces $\tilde{\theta}$ in their second report with probability one.*

Proof. We consider the following two cases:

Case 1: *Everyone tells the truth in the first report with probability one, i.e., $m_i^1 = \theta_i$ with σ_i -probability one for every agent $i \in \mathcal{I}$.*

It follows directly from Claim 2 that $m_i^2 = \theta$ with σ_i -probability one for every agent $i \in \mathcal{I}$.

Case 2: *There exists agent i who tells a lie in the first report with σ_i -positive probability.*

That is, there exists $m_i \in \text{supp}(\sigma_i)$ such that $m_i^1 \neq \theta_i$. By Claim 1, (m_i, m_{-i}) is consistent with σ_{-i} -probability one. In particular, there exists $\tilde{\theta} \in \Theta$ such that every agent $j \neq i$ must report

$$m_j^2 = m_i^2 = \tilde{\theta} \text{ with } \sigma_j\text{-probability one.} \quad (11)$$

Hence, by Claim 2(b), for every $\tilde{m}_i \in \text{supp}(\sigma_i)$, we have

$$\tilde{m}_{i,i}^2 = m_{i,i}^2 = \tilde{\theta}. \quad (12)$$

We now prove that for every $\tilde{m}_i \in \text{supp}(\sigma_i)$, we have $\tilde{m}_i^2 = m_i^2 = \tilde{\theta}$, which would complete the proof. We prove it by contradiction, i.e., suppose there exists $\tilde{m}_i \in \text{supp}(\sigma_i)$ such that

$$\tilde{m}_i^2 \neq m_i^2. \quad (13)$$

Furthermore, (11) and (13) imply that for every agent $j \neq i$, $e_{j,i}(m_j, \tilde{m}_i) = \varepsilon$ with σ_j -probability one. Hence, by Claim 1, every agent $j \neq i$ must tell the truth in the first report, i.e., $m_j^1 = \theta_j$ with σ_j -probability one. As a result, Claim 2(a) implies for every agent $j \neq i$

$$\tilde{m}_{i,j}^2 = m_{i,j}^2 = \theta_j \text{ with } \sigma_i\text{-probability one.} \quad (14)$$

Finally, (12) and (14) imply $\tilde{m}_i^2 = m_i^2$, contradicting (13). ■

Step 3: No Challenge

By Claim 3, there exists a common state $\tilde{\theta} \in \Theta$ with σ_i -probability one for every agent $i \in \mathcal{I}$. We now show in Claim 4 that no one challenges the common state $\tilde{\theta}$.

Claim 4 *No agent challenges with positive probability the common state $\tilde{\theta}$ announced in the second report.*

Proof. Suppose by way of contradiction that $x(\tilde{\theta}, m_i^1) \neq f(\tilde{\theta})$ for some message $m_i \in \text{supp}(\sigma_i)$. By Claim 3, we have $x(m_j^2, m_i^1) \neq f(m_j^2)$ for every message $m_j \in \text{supp}(\sigma_j)$ and every agent $j \neq i$. It implies that $e_{j,i}(m_j, m_i) = \varepsilon$ with σ_j -probability one for every $j \neq i$ and $m_j \in \text{supp}(\sigma_j)$. By Claim 1, we have $m_j^1 = \theta_j$ with σ_j -probability one and $m_i^1 = \theta_i$. Thus, we obtain $x(\tilde{\theta}, \theta_i) \neq f(\tilde{\theta})$. By the construction of the best challenge scheme, we also have $x(\tilde{\theta}, \theta_i) \in \mathcal{L}_i(f(\tilde{\theta}), \tilde{\theta}_i) \cap \mathcal{SU}_i(f(\tilde{\theta}), \theta_i)$. Then, by (8), every message \tilde{m}_i with $x(\tilde{\theta}, \tilde{m}_i^1) = f(\tilde{\theta})$ cannot be a best response against σ_{-i} . Indeed, since $x(\tilde{\theta}, \theta_i) \in \mathcal{SU}_i(f(\tilde{\theta}), \theta_i)$, it is a profitable deviation to replace \tilde{m}_i^1 by θ_i . Hence, $x(\tilde{\theta}, \tilde{m}_i) \neq f(\tilde{\theta})$ and $e_{j,i}(m_j, \tilde{m}_i) = \varepsilon$ for every $\tilde{m}_i \in \text{supp}(\sigma_i)$. Once again, by Claim 1, we have $\tilde{m}_i^1 = \theta_i$ with σ_i -probability one. Therefore, every agent's first report is truthful with probability one. By Claim 2, we conclude that $\tilde{\theta} = \theta$. Since $x(\tilde{\theta}, \theta_i) \neq f(\tilde{\theta})$, it follows that $x(\tilde{\theta}, \theta_i)$ belongs to the empty intersection $\mathcal{L}_i(f(\theta), \theta_i) \cap \mathcal{SU}_i(f(\theta), \theta_i)$, which is impossible. ■

4.3 Implementation in a Direct Mechanism

In this section we present two special cases in which our implementing mechanism can be made into a direct mechanism. Both cases require three or more agents. A direct (revelation) mechanism is a mechanism $((M_i), g, (\tau_i))_{i \in \mathcal{I}}$ in which (i) agents are asked to report the state (i.e., $M_i = \Theta$ for every agent i), and (ii) a unanimous report leads to the socially desirable outcome with no transfers (i.e., $g(\theta, \dots, \theta) = f(\theta)$, and $\tau_i(\theta) = 0$, for every $i \in \mathcal{I}$ and $\theta \in \Theta$). Our notion of direct mechanism is adopted in, for example, [Dutta and Sen \(1991\)](#) and ([Osborne and Rubinstein, 1994](#), Definition 179.2) both of which ask each agent to report a state.

Although direct mechanisms invoke a simpler message space than the augmented mechanisms used in the full implementation literature, the literature on partial implementation has attempted to construct mechanisms that are simpler or easier to implement than direct mechanisms, allowing lotteries and transfers. See, for example, [Dasgupta and Maskin \(2000\)](#)

and [Perry and Reny \(2002\)](#). While our result complements these papers, our main focus is to study full implementation in mixed-strategy Nash equilibrium without making use of integer or modulo games.

The first case shows that every Maskin-monotonic SCF is fully implementable in pure-strategy Nash equilibria in a direct mechanism. Pure-strategy Nash implementation means that we only require that each pure-strategy Nash equilibrium achieve desirable outcomes, i.e., condition (ii) of [Definition 1](#) holds only for pure-strategy Nash equilibria. Indeed, one might expect that by penalizing disagreement with transfers, the designer can easily obtain a unanimous state announcement without using integer/modulo games. Once there is a unanimous state announcement in equilibrium, Maskin monotonicity will ensure implementation, as it does in [Maskin \(1999\)](#). The following proposition formalizes the idea; see [Appendix A.2](#) for a proof.

Proposition 1 *Suppose that there are at least three agents and the SCF f satisfies Maskin monotonicity. Then, f is implementable in pure-strategy Nash equilibria by a direct mechanism.*

The idea of “penalizing disagreement” becomes problematic once we consider mixed-strategy equilibria. Indeed, the direct mechanism which we construct in proving [Proposition 1](#) is reminiscent of modulo games, which, as is well known, admit unwanted mixed-strategy equilibria. Thus, it should come at no surprise that the direct mechanism also admits unwanted mixed-strategy equilibria.

The second case establishes mixed-strategy Nash implementation in direct mechanisms by considering a state space of a “product form,” i.e., there is a one-to-one correspondence between Θ and $\times_{i=1}^I \Theta_i$. We state the following result and relegate its proof to [Appendix A.3](#):

Proposition 2 *Suppose that there are at least three agents, there is a one-to-one correspondence between Θ and $\times_{i=1}^I \Theta_i$, and the SCF f satisfies Maskin monotonicity. Then, f is implementable in mixed-strategy Nash equilibria by a direct mechanism.*

[Proposition 2](#) represents an extreme case in which mixed-strategy Nash implementation can be achieved in a direct mechanism. Product state space naturally arises in a Bayesian setup with a full-support common prior. While such a full-support prior is precluded by the

complete-information assumption, it is consistent with “almost complete information” which we will introduce in Section 5.1.

4.4 Implementation without Off-the-Equilibrium Transfers

The following example illustrates the fact of that without any domain restriction such as quasilinear preferences with transfers, some Maskin-monotonic SCF cannot be implemented by mixed-strategy Nash equilibria in finite mechanisms.

Example 1 (Example 4 of Jackson (1992)) *Consider the environment with two agents 1 and 2. Suppose that there are four alternatives a, b, c , and d and two states θ and θ' . Suppose that agent 1 has the state-independent preference $a \succ_1 b \succ_1 c \sim_1 d$, and agent 2 has the preference $a \succ_2^\theta b \succ_2^\theta d \succ_2^\theta c$ at state θ and preference $b \succ_2^{\theta'} a \succ_2^{\theta'} c \sim_2^{\theta'} d$ at state θ' . Consider the SCF f such that $f(\theta) = a$ and $f(\theta') = c$.*

With no restrictions on agents’ preferences, Jackson (1992) shows that for every finite mechanism which implements f in pure-strategy Nash equilibria, there must also exist a “bad” mixed-strategy Nash equilibrium such that at state θ' the equilibrium outcome differs from c with positive probability.¹⁰ Since f satisfies Maskin monotonicity, the example shows that without imposing any domain restrictions on the environment, it is impossible to implement any Maskin-monotonic SCF in mixed-strategy equilibria by a finite mechanism. However, regardless of the cardinal representation of the preferences in Jackson’s example, the SCF f can actually be implemented in mixed-strategy equilibria with arbitrarily small transfers off the equilibrium; for more discussion, see Section 5.3 and in particular, footnote 15.

¹⁰We briefly recap the argument here. Let \mathcal{M} be a finite mechanism which implements the SCF f in pure-strategy Nash equilibria. Consider a mechanism which restricts the message space of \mathcal{M} such that, against any message of agent i , the opponent agent j can choose a message that induces either outcome a or b . The restricted set of messages is nonempty since the equilibrium message profile at state θ leads to outcome a . It follows that at state θ' , the game induced by the restricted mechanism must have a mixed-strategy Nash equilibrium. Moreover, the equilibrium outcome must be a or b with positive probability; otherwise, agent 2 can deviate to induce outcome a or b with positive probability. Since c and d are ranked lowest by both agents at state θ' , the mixed-strategy equilibrium must remain an equilibrium at state θ' in the game induced by \mathcal{M} ; moreover, the equilibrium fails to achieve $f(\theta') = c$.

5 Extensions

We now establish several extensions of our main result (Theorem 1). In Section 5.1, we show that the implementation result is robust to information perturbations. That is, we establish that our implementation result remains valid in any incomplete-information environment that is close to our complete-information benchmark. In Section 5.2, we extend our result to the case of social choice correspondences (henceforth, SCCs). Section 5.3 clarifies how the designer can modify the implementing mechanism to make the size of transfers arbitrarily small. For the sake of clarity, we will not discuss any combination of multiple extensions. For instance, we will study the case of SCCs only in Section 5.2 but focus entirely on SCFs in the rest of the paper.

The extensions involve more technical details. Thus, we assume, in this section, that the set A (of pure alternatives) is finite and relegate all the proofs to the appendix.

5.1 Robustness to Information Perturbations

Chung and Ely (2003) and Aghion, Fudenberg, Holden, Kunimoto, and Tercieux (2012) consider a designer who not only wants all equilibria of her mechanism to yield a desirable outcome under complete information, but is also concerned about the possibility that agents may entertain small doubts about the true state. They argue that such a designer should insist on implementing the SCF in the closure of a solution concept as the amount of incomplete information about the state vanishes. Chung and Ely (2003) adopt undominated Nash equilibrium and Aghion, Fudenberg, Holden, Kunimoto, and Tercieux (2012) adopt subgame-perfect equilibrium as a solution concept in studying the robustness issue.

To allow for information perturbations, suppose that the agents do not observe the state directly but are informed of the state via signals. The set of agent i 's signals is denoted as S_i , which is identified with Θ , i.e., $S_i \equiv \Theta$.¹¹ A signal profile is an element $s = (s_1, \dots, s_I) \in S \equiv \times_{i \in \mathcal{I}} S_i$. When the realized signal profile is s , agent i observes only his own signal s_i . Let s_i^θ denote the signal which corresponds to state θ , and we write $s^\theta = (s_i^\theta)_{i \in \mathcal{I}}$. State and signals are drawn from some prior distribution over $\Theta \times S$. In particular, complete information can be modelled as a prior μ such that $\mu(\theta, s) = 0$ whenever

¹¹We adopt the formulation from Chung and Ely (2003) and Aghion, Fudenberg, Holden, Kunimoto, and Tercieux (2012). Our result holds for any alternative formulation under which the (Bayesian) Nash equilibrium correspondence has a closed graph.

$s \neq s^\theta$. Such a μ will be called a *complete-information prior*. We assume that for each agent i , the marginal distribution on i 's signals places a strictly positive weight on each of i 's signals, that is, $\text{marg}_{S_i} \mu(s_i) > 0$ for every $s_i \in S_i$, so that the posterior belief given every signal is well defined. For every prior ν , we also write $\nu(\cdot|s_i)$ for the conditional distribution of ν on signal s_i .

The distance between two priors is measured by the uniform metric. That is, for every two priors μ and ν , we have $d(\mu, \nu) \equiv \max_{\theta, s} |\mu(\theta, s) - \nu(\theta, s)|$. Write $\nu^\varepsilon \rightarrow \mu$ if $d(\nu^\varepsilon, \mu) \rightarrow 0$ as $\varepsilon \rightarrow 0$. A prior ν together with a mechanism $\mathcal{M} = ((M_i, \tau_i)_{i \in \mathcal{I}}, g)$ induces an incomplete-information game, which we denote by $\Gamma(\mathcal{M}, \nu)$. A (mixed-)strategy of agent i is now a mapping $\sigma_i : S_i \rightarrow \Delta(M_i)$.

The designer may resort to a solution concept \mathcal{E} for the game $\Gamma(\mathcal{M}, \nu)$ (such as Bayesian Nash equilibrium) which induces a set of mappings from $\Theta \times S$ to X , which we call *acts*, following [Chung and Ely \(2003\)](#). For instance, each Bayesian Nash equilibrium σ induces the act α_σ with $\alpha_\sigma(\theta, s) \equiv \sigma(s) \circ (g, (\tau_i)_{i \in \mathcal{I}})^{-1}$, where we abuse the notation to identify the finite-support distribution $\sigma(s) \circ (g, (\tau_i)_{i \in \mathcal{I}})^{-1}$ on X with an allocation in X . We denote the set of acts induced by the solution concept \mathcal{E} as $\mathcal{E}(\mathcal{M}, \nu)$. We endow X with a topology with respect to which the utility function u_i is continuous on X .¹² We now define $\overline{\mathcal{E}}$ -implementation.

Definition 3 *An SCF f is $\overline{\mathcal{E}}$ -implementable under the complete-information prior μ if there exists a mechanism $\mathcal{M} = ((M_i), g, (\tau_i))_{i \in \mathcal{I}}$ such that for every $(\theta, s) \in \text{supp}(\mu)$ and every sequence of priors $\{\nu^n\}$ converging to μ , the following two requirements hold: (i) there is a sequence of acts $\{\alpha_n\}$ with $\alpha_n \in \mathcal{E}(\mathcal{M}, \nu_n)$ such that $\alpha_n(\theta, s) \rightarrow f(\theta)$; and (ii) for every sequence of acts $\{\alpha_n\}$ with $\alpha_n \in \mathcal{E}(\mathcal{M}, \nu_n)$, we have $\alpha_n(\theta, s) \rightarrow f(\theta)$.*

[Chung and Ely \(2003\)](#) and [Aghion, Fudenberg, Holden, Kunimoto, and Tercieux \(2012\)](#) show that Maskin monotonicity is a necessary condition for \overline{UNE} -implementation and \overline{SPE} -implementation, respectively.¹³ The result of [Chung and Ely \(2003\)](#) implies that implementation of a non-Maskin-monotonic SCF in undominated Nash equilibria such as the result in [Abreu and Matsushima \(1994\)](#) is necessarily vulnerable to information perturbations. More-

¹²For instance, it is the case if A is a (Hausdorff) topological space, $v_i(a, \theta)$ is bounded and continuous in a , and $\Delta(A)$ is endowed with the weak*-topology. Then, $X \equiv \Delta(A) \times \mathbb{R}^I$, endowed with the product topology, is also a Hausdorff topological space.

¹³[Aghion, Fudenberg, Holden, Kunimoto, and Tercieux \(2012\)](#) adopt sequential equilibrium as the solution concept for the incomplete-information game $\Gamma(\mathcal{M}, \nu)$.

over, both [Chung and Ely \(2003, Theorem 2\)](#) and [Aghion, Fudenberg, Holden, Kunimoto, and Tercieux \(2012\)](#) establish the sufficiency result by using an infinite mechanism with an integer game and restricting attention to pure-strategy equilibria. It raises the question as to whether their robustness test may be too demanding when it is applied to finite mechanisms where mixed-strategy equilibria have to be taken seriously, since the implementing mechanism of [Jackson, Palfrey, and Srivastava \(1994\)](#), that of [Abreu and Matsushima \(1994\)](#), or the simple mechanism in Section 5 of [Moore and Repullo \(1988\)](#) are considered examples of such finite mechanisms.

The canonical mechanism which we propose in the proof of [Theorem 1](#) is indeed finite, and we show that that finite mechanism implements every Maskin-monotonic SCF in mixed-strategy Nash equilibria. Since the solution concept of Bayesian Nash equilibrium, viewed as a correspondence on priors, has a closed graph, that finite mechanism also achieves \overline{NE} -implementation. We now obtain the following result as a corollary of [Theorem 1](#) in our setup with lotteries and transfers.

Proposition 3 *Let \mathcal{E} be a solution concept such that $\emptyset \neq \mathcal{E}(\mathcal{M}, \mu) \subseteq NE(\mathcal{M}, \mu)$ for each finite mechanism \mathcal{M} and a complete-information prior μ . Then, every Maskin-monotonic SCF f is $\overline{\mathcal{E}}$ -implementable.*

The condition $\emptyset \neq \mathcal{E}(\mathcal{M}, \mu) \subseteq NE(\mathcal{M}, \mu)$ is satisfied for virtually every refinement of Nash equilibrium, because we allow for mixed-strategy equilibria and $\Gamma(\mathcal{M}, \mu)$ is a finite game.

5.2 Social Choice Correspondences

A large portion of the implementation literature strives to deal with social choice correspondences (hereafter, SCCs), i.e., multivalued social choice rules. In this section, we extend our Nash implementation result to cover the case of SCCs. We suppose that the designer's objective is specified by an SCC $F : \Theta \rightrightarrows X$; and for simplicity, we assume that $F(\theta)$ is a finite set for each state $\theta \in \Theta$. It includes the special case where the co-domain of F is A . Following [Maskin \(1999\)](#), we first define the notion of Nash implementation for SCCs.

Definition 4 *An SCC F is **implementable in mixed-strategy Nash equilibria by a finite mechanism** if there exists a finite mechanism $\mathcal{M} = ((M_i, \tau_i)_{i \in \mathcal{I}}, g)$ such that for*

every state $\theta \in \Theta$, the following two conditions are satisfied: (i) for every $x \in F(\theta)$, there exists a pure-strategy Nash equilibrium m in the game $\Gamma(\mathcal{M}, \theta)$ such that $g(m) = x$ and $\tau_i(m) = 0$ for every agent $i \in \mathcal{I}$; and (ii) for every $m \in \text{supp}(NE(\Gamma(\mathcal{M}, \theta)))$, we have $\text{supp}(g(m)) \subseteq F(\theta)$ and $\tau_i(m) = 0$ for every agent $i \in \mathcal{I}$.

Second, we state the definition of Maskin monotonicity for SCCs.

Definition 5 *An SCC F satisfies **Maskin monotonicity** if, for each pair of states $\tilde{\theta}$ and θ and $x \in F(\tilde{\theta}) \setminus F(\theta)$, some agent $i \in \mathcal{I}$ and some allocation $x \in X$ exist such that*

$$\tilde{u}_i(x, \tilde{\theta}) \leq \tilde{u}_i(f(\tilde{\theta}), \tilde{\theta}) \text{ and } \tilde{u}_i(x, \theta) > \tilde{u}_i(f(\tilde{\theta}), \theta).$$

We now state our Nash implementation result for SCCs and relegate the proof to Appendix 2.¹⁴

Theorem 2 *Suppose there are at least three agents. An SCC F is implementable in mixed-strategy Nash equilibria by a finite mechanism if and only if it satisfies Maskin monotonicity.*

Compared with Theorem 1 for SCFs, Theorem 2 needs to overcome additional difficulties. In the case of SCFs, when the agents' second reports are consistent at a common state $\tilde{\theta}$, they will be associated with a single outcome $f(\tilde{\theta})$. Hence, if agent i 's second report is challenged, then *every* second report which is played with positive probability by any agent must also be challenged in equilibrium. Together with Claim 1, it implies that every agent must tell the truth in their first and second report, which leads to a contradiction in the proof of Claim 4.

In the case of SCCs, each allocation $x \in F(\theta)$ has to be implemented as the outcome of some pure-strategy equilibrium. Hence, each agent must also report an allocation to be implemented. It also follows that a challenge scheme for an SCC must be defined for a type θ_i to challenge a pair $(\tilde{\theta}, x)$ with $x \in F(\tilde{\theta})$. As a result, even when the agents' second reports are consistent at state $\tilde{\theta}$ (which still holds by Claim 3), they might still be randomizing between two allocations x and x' in $F(\tilde{\theta})$ such that $(\tilde{\theta}, x)$ is challenged and yet $(\tilde{\theta}, x')$ is not. Hence, we cannot follow a similar argument as in Claim 1 to derive a contradiction. Instead,

¹⁴When there are only two agents, we can still show that every Maskin-monotonic SCC F is *weakly* implementable in Nash equilibria, that is, there exists a mechanism which has a pure-strategy Nash equilibrium and satisfies requirement (ii) in Definition 4.

we build on the implementing mechanism in Section 4.1 and show that agent i will not report $(\tilde{\theta}, x)$ which can be challenged either by (i) agent $j \neq i$ or by (ii) agent i himself. We ensure Case (i) by imposing a large penalty on agent i conditional on agent j 's challenging $(\tilde{\theta}, x)$, whereas we deal with Case (ii) by allowing agent i to challenge himself without penalty.

Remark. Mezzetti and Renou (2012b) also consider deterministic SCCs in a separable environment studied in Jackson, Palfrey, and Srivastava (1994). Mezzetti and Renou (2012b) identify a condition (which they call top- D inclusiveness) under which an SCC is implementable in mixed-strategy Nash equilibria in finite mechanisms if and only if it satisfies *set-monotonicity* (proposed by Mezzetti and Renou (2012a)). There are several differences between our Theorem 2 and their result. First, Mezzetti and Renou (2012b) require only the existence of mixed-strategy equilibria but we follow Maskin (1999) in requiring the existence of pure-strategy equilibria. Second, Mezzetti and Renou (2012b) consider an ordinal setup, while we consider a cardinal setup. These two features of Mezzetti and Renou (2012b) are the reason why they use set-monotonicity as a relevant necessary condition for characterizing their ordinal Nash implementation. We also study ordinal Nash implementation for the case of SCFs in Section 5.5. Third, our quasilinear environments with transfers are more restrictive than the separable environments considered by Mezzetti and Renou (2012b). Finally, Mezzetti and Renou (2012b) need “top D -inclusiveness” as an additional condition, which requires that there exists at least one agent for whom the SCC contains the agent’s best outcome within the range of the SCC for every state of the world, whereas we impose no conditions beyond Maskin monotonicity for the SCF.

5.3 Small Transfers

One potential drawback of the mechanism we propose for Theorem 1 is that the size of transfers may be large. To tackle the problem, we use the technique introduced by Abreu and Matsushima (1994) to show that if the SCF satisfies Maskin monotonicity in the restricted domain without any transfer, then it is Nash-implementable with arbitrarily small transfers.

We first propose a notion of Nash implementation with bounded transfers off the equilibrium and still no transfers on the equilibrium.

Definition 6 *An SCF $f : \Theta \rightarrow \Delta(A)$ is implementable in mixed-strategy Nash equilibria by a finite mechanism **with transfers bounded by $\bar{\tau}$** if there exists a finite mechanism*

$\mathcal{M} = ((M_i, \tau_i)_{i \in \mathcal{I}}, g)$ such that for every state $\theta \in \Theta$ and $m \in M$, (i) there exists a pure-strategy Nash equilibrium in the game $\Gamma(\mathcal{M}, \theta)$; (ii) for each m in $\text{supp}(NE(\Gamma(\mathcal{M}, \theta)))$, we have $g(m) = f(\theta)$ and $\tau_i(m) = 0$ for every agent $i \in \mathcal{I}$; and (iii) $|\tau_i(m)| \leq \bar{\tau}$ for every $m \in M$ and every agent $i \in \mathcal{I}$.

Next, we propose a notion of Nash implementation in which there are no transfers on the equilibrium and only arbitrarily small transfers off the equilibrium.

Definition 7 *An SCF f is implementable in mixed-strategy Nash equilibria by a finite mechanism **with arbitrarily small transfers** if, for every $\bar{\tau} > 0$, the SCF f is implementable in Nash equilibria by a finite mechanism with transfers bounded by $\bar{\tau}$.*

We say that an SCF f satisfies Maskin monotonicity in the restricted domain $\Delta(A)$ if $f(\tilde{\theta}) \neq f(\theta)$ implies that there are an agent i and some lottery $x(\tilde{\theta}, \theta_i)$ in $\Delta(A)$ such that $x(\tilde{\theta}, \theta_i)$ belongs to $\mathcal{L}_i(f(\tilde{\theta}), \tilde{\theta}_i) \cap \mathcal{SU}_i(f(\tilde{\theta}), \theta_i)$. Here, for $(\ell, \theta_i) \in \Delta(A) \times \Theta_i$, we use $\mathcal{L}_i(\ell, \theta_i)$ to denote the lower-contour set at allocation ℓ in $\Delta(A)$ for type θ_i , i.e.,

$$\mathcal{L}_i(\ell, \theta_i) = \{\ell' \in \Delta(A) : v_i(\ell, \theta) \geq v_i(\ell', \theta)\}.$$

In a similar fashion, \mathcal{SU}_i is defined. Clearly, Maskin monotonicity in the restricted domain $\Delta(A)$ is stronger than Maskin monotonicity in the domain X , as the former requires that the test allocation be a lottery over alternatives without transfer. In Appendix A.6, we assume there are at least three agents, and prove the following result.¹⁵

Theorem 3 *Suppose there are at least three agents. An SCF $f_A : \Theta \rightarrow \Delta(A)$ is implementable in mixed-strategy Nash equilibria by a finite mechanism with arbitrarily small transfers if f_A satisfies Maskin monotonicity in the restricted domain.*

5.4 Infinite State Space

One significant assumption we have made in this paper is that the state space is finite. In Appendix A.7, we extend Theorem 1 to an infinite state space in which the agents' utility

¹⁵In the case with only two agents, Theorem 3 still holds if there exists an alternative $w \in A$ which is the worst alternative for any agent at any state. In that case, we can simply modify the “voting rule” ϕ in the proof of Theorem 3 to be $\phi(m^h) = f(\tilde{\theta})$ if both agents announce a common type profile which identifies a state $\tilde{\theta}$ in m^h ; and $\phi(m^h) = w$ otherwise. In particular, $w = c$ in Example 4 of Jackson (1992) and thus the SCF can be implemented with arbitrarily small transfers. Moreover, the conclusion holds regardless of the utility representation of the agents' preferences. However, note that we assume that agents have quasilinear utilities while Jackson's example does not make such an assumption.

functions are continuous. A similar extension was raised as an open question for virtual implementation in [Abreu and Matsushima \(1992\)](#) (see their Section 5) and it has not been answered to our knowledge.

In [Appendix A.7](#), we construct an extension of the implementing mechanism for mixed-strategy Nash implementation which accommodates an infinite state space. We state the result as follows:

Theorem 4 *Suppose that Θ is a Polish space. Then, an SCF f satisfies Maskin monotonicity if and only if there exists a mechanism which implements f in mixed-strategy Nash equilibria. Moreover, if Θ is compact and both the utility function $\{v_i(a, \cdot)\}_{a \in A}$ and the SCF are continuous functions on Θ , then the implementing mechanism has a compact message space together with a continuous outcome function and continuous transfer rules.*

One notable feature of this extension is that as long as the setting is the one with a compact state space, the continuous SCF, and continuous utility functions, the resulting implementing mechanism will also be compact and continuous. This feature ensures that best responses are always well defined in our mechanism; hence, it differentiates our construction from the traditional one invoking the use of integer games.

The proof of [Theorem 4](#) needs to overcome two difficulties. First, in a finite state space, the transfer rules $\tau_{i,j}^1$ and $\tau_{i,j}^2$ which we define in [\(9\)](#) and [\(10\)](#) impose either a large penalty and/or a large reward as long as the designer sees a discrepancy in the agents' announcements. With a continuum of states/types, however, such a drastic change in transfer scale is precluded by the continuity requirement. Hence, our first challenge is to suitably define $\tau_{i,j}^1$ and $\tau_{i,j}^2$ so that they vary continuously yet still incentivize them to tell the truth.

Second, in an infinite setting, we know of no way to construct a challenge scheme which pre-selects a test allocation in a continuous manner. As a result, we cannot have the agents report their type, let alone the true type, to cast a challenge to state $\tilde{\theta}$. Instead, we will restore the continuity of the outcome function by asking them to report a test allocation x directly. Despite this change, we will establish a counterpart of [Condition \(8\)](#) as [Lemmas 5](#) and [6](#) in [Appendix A.7](#).

5.5 The Ordinal Approach

We have assumed that the agents are expected utility maximizers. This leaves open the issue as to whether, and to what extent, our implementation result depends on the designer's knowledge about the cardinalization of the agents' preferences over lotteries. To address the issue, we discuss how our result can accommodate an ordinal setting.

First, we introduce the notion of *ordinal Nash implementation*. The notion requires that the mixed-strategy Nash implementation be obtained for *any* cardinal representation of the ordinal preferences over the finite set of pure alternatives A . Formally, we follow the approach proposed by [Mezzetti and Renou \(2012a\)](#).

Suppose that at state θ , the agents commonly know only that their ordinal rankings over the set of pure alternatives A . We write the induced ordinal preference profile at state θ by $(\succeq_i^\theta)_{i \in \mathcal{I}}$. We also assume no redundancy, i.e., $\theta \neq \theta'$ implies $\succeq_i^\theta \neq \succeq_i^{\theta'}$ for some agent i . It is taken for granted that distinct pair of \succeq_i^θ and $\succeq_i^{\theta'}$ induce different preference orderings over A , and also that no players are indifferent over all elements of A . For each $i \in \mathcal{I}$, let $v_i : A \times \Theta \rightarrow \mathbb{R}$ be a *cardinal representation* of $(\succeq_i^\theta)_{\theta \in \Theta}$ over A , i.e., for each pair of alternatives $a, a' \in A$, agent $i \in \mathcal{I}$, and state $\theta \in \Theta$, we have

$$v_i(a, \theta) \geq v_i(a', \theta) \Leftrightarrow a \succeq_i^\theta a'.$$

We assume that each function v_i takes a value in $[0, 1]$. In addition, each cardinal representation v_i induces an expected utility function on $\Delta(A)$ which, by abuse of notation, we also denote by $v_i : \Delta(A) \times \Theta \rightarrow \mathbb{R}$. We denote by V_i^θ the set of all cardinal representations $v_i(\cdot, \theta)$ of \succeq_i^θ . Following [Mezzetti and Renou \(2012a\)](#), we focus our discussion on the case of a deterministic SCF, i.e., $f : \Theta \rightarrow A$. We say that an SCF f is *ordinally Nash implementable* if it is implementable in mixed-strategy Nash equilibria independently of the cardinal representation. We formalize this idea in the following definition.

Definition 8 *An SCF f is ordinally Nash implementable if there exists a mechanism \mathcal{M} such that, for every $\theta \in \Theta$ and every profile of cardinal representations $v = (v_i)_{i \in \mathcal{I}}$ of $(\succeq_i^\theta)_{i \in \mathcal{I}, \theta \in \Theta}$, the following two conditions are satisfied: (i) there exists a pure-strategy Nash equilibrium m in the game $\Gamma(\mathcal{M}, \theta, v)$ such that $g(m) = f(\theta)$ and $\tau_i(m) = 0$ for every $i \in \mathcal{I}$; and (ii) for every $m \in \text{supp}(NE(\Gamma(\mathcal{M}, \theta, v)))$, we have $\text{supp}(g(m)) = f(\theta)$ and $\tau_i(m) = 0$ for every $i \in \mathcal{I}$.*

In Appendix A.8, we introduce the notion of *ordinal almost monotonicity* proposed by Sanver (2006). It roughly says that whenever the SCF designates different outcomes at states θ and θ' , there must be an agent with a *deterministic* test allocation which displays a suitable preference reversal with respect to the socially desirable outcome at the two states. We obtain the following result:

Theorem 5 *An SCF f is ordinally Nash implementable if and only if it satisfies ordinal almost monotonicity.*

It is worth emphasizing that Theorem 4 becomes the key to proving this result. We show that ordinal almost monotonicity of an SCF implies Maskin monotonicity of the SCF consistently extended from Θ (now the set of ordinal preference profiles) to the set of cardinalizations. Since the set of all cardinalizations extended from Θ becomes an infinite state space, the extension of ordinal almost monotonicity renders Theorem 4 applicable.

Instead of relying on ordinal almost monotonicity, Mezzetti and Renou (2012a) propose a notion called *set-monotonicity* for SCCs. They show that the notion of set-monotonicity is weaker than Maskin monotonicity and is necessary and “almost sufficient” in their notion of implementation in mixed-strategy Nash equilibria. There are three further differences between the results of Mezzetti and Renou (2012a) and ours. First, Mezzetti and Renou (2012a) require only the existence of mixed-strategy equilibria but we follow Maskin (1999) in requiring the existence of a pure-strategy equilibrium. This difference makes our ordinal implementation notion more demanding than that of Mezzetti and Renou (2012a). Second, we use monetary transfers, while Mezzetti and Renou (2012a) do not. Although ordinal almost monotonicity is weaker than set-monotonicity, by our use of transfers, we obtain our stronger notion of ordinal mixed-strategy Nash implementation à la Maskin (1999) for the case of SCFs by means of ordinal almost monotonicity. Finally, Mezzetti and Renou (2012a) also study the case of SCCs which we omit here.

6 Concluding Remarks

Despite its tremendous success, implementation theory has also been criticized on various fronts. A major critique is that the mechanisms used to achieve full implementation are not “natural”. To address the critique, Jackson (1992) proposes to characterize the class of

SCFs which can be fully implemented in “natural” mechanisms, even at the cost of imposing domain restrictions.

We consider our results as an important step in advancing Jackson’s research program. Specifically, we propose to recast an implementation problem by requiring that the implementing mechanism be finite/well-behaved and have no unwanted mixed-strategy equilibria. Such requirements are to be anticipated, when the setting of interest is indeed finite/well-behaved to start with. We prove a first set of benchmark results on mixed-strategy Nash implementation by considering environments with lotteries and transfers. We also show that our results are robust to information perturbations and amenable to handling prominent extensions such as SCCs, small transfers, infinite settings, and ordinal settings.

There have been some past attempts at tackling implementation in Bayesian (incomplete-information) environments, such as [Mookherjee and Reichelstein \(1990\)](#) and [d’Aspremont, Crémer, and Gérard-Varet \(2003\)](#). Both papers are considered as previous attempts at full Bayesian implementation by well-behaved mechanisms in incomplete-information environments with transfers. However, they do not consider mixed-strategy equilibria and we are aware of no results on mixed-strategy Bayesian implementation achieved by well-behaved mechanisms. The novelty of our paper is to take care of mixed-strategy equilibria by a well-behaved mechanism. We take complete-information environments with a restricted preference domain as a natural starting point and leave for future research the important yet more challenging extensions to incomplete-information environments.¹⁶

Our implementation results are obtained by imposing transfers off the equilibrium. This feature is intimately related to the burgeoning literature on repeated implementation, such as [Lee and Sabourian \(2011\)](#) and [Mezzetti and Renou \(2017\)](#), in which continuation values can serve as transfers in our construction.¹⁷ It is an important future direction to investigate the formal connection.

A Appendix

In this Appendix, we provide the proofs omitted from the main body of the paper.

¹⁶Allowing for integer games, [Kunimoto \(2019\)](#) and [Serrano and Vohra \(2010\)](#) characterize mixed-strategy Bayesian implementation in more general environments.

¹⁷We thank Hamid Sabourian for drawing our attention to this point.

A.1 Proof of Lemma 1

First, we elaborate the proof of Lemma 1 here.

Proof. Consider a challenge scheme $\bar{x}(\cdot, \cdot)$. First, we show that we can modify $\bar{x}(\cdot, \cdot)$ into a new challenge scheme $x(\cdot, \cdot)$ such that

$$x(\tilde{\theta}, \theta_i) \neq f(\tilde{\theta}) \text{ and } x(\tilde{\theta}, \theta'_i) \neq f(\tilde{\theta}) \Rightarrow u_i(x(\tilde{\theta}, \theta_i), \theta_i) \geq u_i(x(\tilde{\theta}, \theta'_i), \theta_i). \quad (15)$$

To construct $x(\cdot, \cdot)$, for each player i , we distinguish two cases: (a) if $\bar{x}(\tilde{\theta}, \theta_i) = f(\tilde{\theta})$ for all $\theta_i \in \Theta_i$, then set $x(\tilde{\theta}, \theta_i) = \bar{x}(\tilde{\theta}, \theta_i) = f(\tilde{\theta})$; (b) if $\bar{x}(\tilde{\theta}, \theta_i) \neq f(\tilde{\theta})$ for some $\theta_i \in \Theta_i$, then define $x(\tilde{\theta}, \theta_i)$ as the most preferred allocation of type θ_i in the finite set

$$X(\tilde{\theta}) = \left\{ \bar{x}(\tilde{\theta}, \theta'_i) : \theta'_i \in \Theta_i \text{ and } \bar{x}(\tilde{\theta}, \theta'_i) \neq f(\tilde{\theta}) \right\}.$$

Since $\bar{x}(\tilde{\theta}, \theta'_i) \in \mathcal{L}_i(f(\tilde{\theta}), \tilde{\theta}_i)$, we have $u_i(x(\tilde{\theta}, \theta_i), \tilde{\theta}_i) \leq u_i(f(\tilde{\theta}), \tilde{\theta}_i)$; moreover, since $x(\tilde{\theta}, \theta_i)$ as the most preferred allocation of type θ_i in $X(\tilde{\theta})$ and $\bar{x}(\tilde{\theta}, \theta_i) \in \mathcal{SU}_i(f(\tilde{\theta}), \theta_i)$, it follows that $u_i(x(\tilde{\theta}, \theta_i), \theta_i) > u_i(f(\tilde{\theta}), \theta_i)$. In other words, $x(\cdot, \cdot)$ remains a challenge scheme. Moreover, $x(\cdot, \cdot)$ satisfies (15) by construction.

Next, for each state $\tilde{\theta}$ and type θ_i , we show that $x(\cdot, \cdot)$ satisfies (3). We proceed by considering the following two cases. First, suppose that $x(\tilde{\theta}, \theta_i) \neq f(\tilde{\theta})$. Then, by (15), it suffices to consider type θ'_i with $x(\tilde{\theta}, \theta'_i) = f(\tilde{\theta})$. Since $x(\tilde{\theta}, \theta'_i) = f(\tilde{\theta})$ and $x(\tilde{\theta}, \theta_i) \neq f(\tilde{\theta})$, then it follows from $x(\tilde{\theta}, \theta_i) \in \mathcal{SU}_i(f(\tilde{\theta}), \theta_i)$ that $u_i(x(\tilde{\theta}, \theta_i), \theta_i) > u_i(x(\tilde{\theta}, \theta'_i), \theta_i)$. Hence, (3) holds. Second, suppose that $x(\tilde{\theta}, \theta_i) = f(\tilde{\theta})$. Then, it suffices to consider type θ'_i with $x(\tilde{\theta}, \theta'_i) \neq f(\tilde{\theta})$. Since $x(\tilde{\theta}, \theta_i) = f(\tilde{\theta})$, we have $\mathcal{L}_i(f(\tilde{\theta}), \tilde{\theta}_i) \cap \mathcal{SU}_i(f(\tilde{\theta}), \theta_i) = \emptyset$. Moreover, $x(\tilde{\theta}, \theta'_i) \neq f(\tilde{\theta})$ implies that $x(\tilde{\theta}, \theta'_i) \in \mathcal{L}_i(f(\tilde{\theta}), \tilde{\theta}_i)$. Hence, we must have $x(\tilde{\theta}, \theta'_i) \notin \mathcal{SU}_i(f(\tilde{\theta}), \theta_i)$. That is, $u_i(x(\tilde{\theta}, \theta_i), \theta_i) \geq u_i(x(\tilde{\theta}, \theta'_i), \theta_i)$, i.e., (3) holds. ■

A.2 Proof of Proposition 1

To facilitate the comparison with Maskin (1999), we assume that there are three or more agents and define the following direct mechanism, denoted by \mathcal{M}^D , according to three rules:

Rule 1. If there exists state $\tilde{\theta}$ such that every agent announces $\tilde{\theta}$, then implement the outcome $f(\tilde{\theta})$.

Rule 2. If there exists state $\tilde{\theta}$ such that everyone except agent i announces $\tilde{\theta}$ and agent i

announces $\tilde{\theta}'$, then implement a test allocation $x(\tilde{\theta}, \tilde{\theta}'_i)$ for agent i and the ordered pair of states $(\tilde{\theta}, \tilde{\theta}')$; and if there is no such test allocation, implement $f(\tilde{\theta})$. Moreover, charge agent $i + 1 \pmod{I}$ a large penalty 2η , where the scale η dominates any difference in utility from allocation.

Rule 3. Otherwise, implement $f(m_1)$. Moreover, charge each agent i a penalty of η if i reports a state which is not reported by the unique majority (i.e., $\{m_i\} \neq \arg \max_{\tilde{\theta}} |\{j \in \mathcal{I} : m_j = \tilde{\theta}\}|$).¹⁸

Now let the true state be θ .

It follows from Rule 2 that since θ is the true state, $x(\theta, \tilde{\theta}'_i) \neq f(\theta)$ implies that $x(\theta, \tilde{\theta}'_i) \in \mathcal{L}_i(f(\theta), \theta_i)$. Hence, everyone reporting the true state constitutes a pure-strategy Nash equilibrium.

Now fix an arbitrary pure-strategy Nash equilibrium m . First, we claim that m cannot trigger Rule 2. Suppose that Rule 2 is triggered, and let agent i be the odd man out. Then, agent $i + 1$ finds it strictly profitable to deviate to announce m_i . After such a deviation, since $I \geq 3$, either Rule 3 is triggered or it remains in Rule 2, but agent i is no longer the odd man out. Thus, agent $i + 1$ saves at least η (from paying 2η to paying η or 0). Such a deviation may also change the allocation selected by the outcome function $g(\cdot)$, which induces utility change less than η . Hence, agent $i + 1$ strictly prefers deviating to announce m_i , which contradicts the hypothesis that m is a Nash equilibrium.

Second, we claim that m cannot trigger Rule 3 either. Suppose that Rule 3 is triggered. Pick an arbitrary state reported by some (not necessarily unique) majority of agents, i.e., $\hat{\theta} \in \arg \max_{\tilde{\theta}} |\{j \in I : m_j = \tilde{\theta}\}|$. Let $\mathcal{I}_{\hat{\theta}}$ be the set of agents who report $\hat{\theta}$. Clearly, $\mathcal{I}_{\hat{\theta}} \subsetneq \mathcal{I}$, because Rule 3 (rather than Rule 1) is triggered. Then, we can find an agent $i^* \in \mathcal{I}_{\hat{\theta}}$ such that agent $i^* + 1 \pmod{I}$ is not in $\mathcal{I}_{\hat{\theta}}$. Since agent $i^* + 1$ does not belong to the unique majority, he must pay η under m . Then, agent $i^* + 1$ will strictly prefer deviating to announce $m_{i^*} = \hat{\theta}$. After such a deviation, either Rule 3 is triggered, and agent $i^* + 1$ falls in the unique majority who reports $\hat{\theta}$; or Rule 2 is triggered, but agent i^* cannot be the odd man out. Thus, agent $i^* + 1$ saves η (from paying η to paying 0) and η' is larger than the maximal utility change induced by different allocations in $g(\cdot)$. The existence of profitable deviation of agent $i^* + 1$ contradicts the hypothesis that m is a Nash equilibrium.

Hence, we conclude that m must trigger Rule 1. It follows that $f(\tilde{\theta}) = f(\theta)$. Otherwise,

¹⁸Note that Rule 3 penalizes every agent by η , if each of them reports a different state.

by Maskin monotonicity, a whistle blower can deviate to trigger Rule 2.

A.3 Proof of Proposition 2

The proof is based on modifying the implementing mechanism and the proof of Theorem 1. We only provide a sketch here. Set $M_i = M_i^1 \times M_i^2$ where $M_i^1 = \Theta_i$ and $M_i^2 = \times_{j \neq i} \Theta_j$. Since $I \geq 3$, the type of each agent is reported by at least two agents in their second report. For each message profile $m = (m_i)_{i=1}^I$, denote by $\tilde{\Theta}(m)$ the set of state induced from the agents' second report, namely that $\tilde{\theta} \in \tilde{\Theta}(m)$ iff for every $i \in \mathcal{I}$, we have $\tilde{\theta}_i = m_{j,i}^2$ for some agent $j \in \mathcal{I}$ (possibly $j = i$). Then, we modify the outcome function:

$$g(m) = \frac{1}{I |\tilde{\Theta}(m)|} \sum_{i \in \mathcal{I}} \sum_{\tilde{\theta} \in \tilde{\Theta}(m)} \left[e(m) \frac{1}{I} \sum_{j \in \mathcal{I}} y_j(m_j^1) \oplus (1 - e(m)) x(\tilde{\theta}, m_i^1) \right]$$

where $e(m) = 0$ if (i) $\tilde{\Theta}(m)$ contains a unique state (consistency); and (ii) $x(\tilde{\theta}, m_i^1) = f(\tilde{\theta})$ for every agent i and every $\tilde{\theta} \in \tilde{\Theta}(m)$ (no challenge); otherwise, $e(m) = \varepsilon$.¹⁹ For the transfer rule, we define

$$\hat{\tau}_{i,j}^1(m_i, m_{-i}) = \begin{cases} 0 & \text{if } m_{i,j}^2 = m_{k,j}^2 \text{ for all } k \in \mathcal{I} \setminus \{i, j\}; \\ -\eta & \text{if } m_{i,j}^2 \neq m_{k,j}^2 \text{ for some } k \in \mathcal{I} \setminus \{i, j\} \text{ and } m_{i,j}^2 \neq m_j^1; \\ \eta & \text{if } m_{i,j}^2 \neq m_{k,j}^2 \text{ for some } k \in \mathcal{I} \setminus \{i, j\} \text{ and } m_{i,j}^2 = m_j^1. \end{cases}$$

Set $\tau_i(m) = \sum_{j \neq i} \hat{\tau}_{i,j}^1(m)$. As the agents no longer report their own type in the second report, we do not need to define $\tau_{i,j}^2(\cdot)$.

The proof of implementation follows the same steps as the proof of Theorem 1 and we only highlight the difference. First, for the contagion of truth argument, we can only establish Claim 2(a) because in the modified mechanism, the agents no longer report their own type in the second report so that we do not have rule $\tau_{i,j}^2(\cdot)$. For the consistency argument, it turns out that Claim 2(a) suffices. Specifically, consider an arbitrary message $m_i \in \text{supp}(\sigma_i)$ such that $m_i^1 \neq \theta_i$. The same argument as in the proof of Claim 3 implies that (m_i, m_{-i}) is consistent for every $m_{-i} \in \text{supp}(\sigma_{-i})$. To show that (\tilde{m}_i, m_{-i}) is consistent for any other $\tilde{m}_i \in \text{supp}(\sigma_i)$, we make use of the assumption that we have three or more agents. In particular, since (m_i, m_{-i}) is consistent for every $m_{-i} \in \text{supp}(\sigma_{-i})$, if (\tilde{m}_i, m_{-i}) is

¹⁹Here we do not have the case with $e(m) = 1$ since $\Theta = \times_{i=1}^I \Theta_i$ implies that $\tilde{\Theta}(m) \subseteq \Theta$.

inconsistent, it must be $\tilde{m}_{i,k}^2 \neq m_{j,k}^2$ for some $j \neq i$, $k \neq i$, and $k \neq j$. By Claim 1 agent k must report his true type with probability one. Then, it follows from Claim 2(a) that $\tilde{m}_{i,k}^2 = m_{j,k}^2$ with probability one and we have reached a contradiction. The argument for no challenge remains the same.

A.4 Proof of Claim 1

Suppose that $m_i^1 \neq \theta_i$ for some $m_i \in \text{supp}(\sigma_i)$. Consider a message \tilde{m}_i which differs from m_i only in sending a truthful first report, i.e., $\tilde{m}_i^1 = \theta_i$ and $\tilde{m}_i^2 = m_i^2$. We prove the claim by showing that \tilde{m}_i is not a strictly better response than m_i against m_j only when $e_{i,j}(m_i, m_j) = e_{j,i}(m_j, m_i) = 0$. Recall that the first report of agent i has no effect on his own transfer.

We consider first the case that the designer uses agent j 's report to check agent i 's report. In that situation, the first report of agent i has no effect on the function $e_{i,j}(\cdot, m_j)$ for every m_j . Hence, we have $e_{i,j}(\tilde{m}_i, m_j) = e_{i,j}(m_i, m_j)$. Moreover, if $m_i^2 \notin \Theta$, then $e_{i,j}(\tilde{m}_i, m_j) = e_{i,j}(m_i, m_j) = 1$; thus, by Lemma 2, \tilde{m}_i is a strictly better response than m_i against m_j . Hence, we may assume $m_i^2 \in \Theta$ and consider the following two cases:

Case 1.1. $e_{i,j}(\tilde{m}_i, m_j) = e_{i,j}(m_i, m_j) = \varepsilon$.

It follows from Lemmas 2 and 1 that

$$u_i(C_{i,j}^\varepsilon(\tilde{m}_i, m_j), \theta_i) - u_i(C_{i,j}^\varepsilon(m_i, m_j), \theta_i) > 0.$$

Hence, \tilde{m}_i is a strictly better response than m_i against m_j .

Case 1.2. $e_{i,j}(\tilde{m}_i, m_j) = e_{i,j}(m_i, m_j) = 0$.

Since $m_i^2 = \tilde{m}_i^2$, both (m_i, m_j) and (\tilde{m}_i, m_j) lead to the same outcome $x(m_i^2, m_j^1) = x(\tilde{m}_i^2, m_j^1) = f(m_i^2)$.

Next, suppose that the designer uses agent i 's report to check agent j 's report. Again, if $m_j^2 \notin \Theta$, then $e_{j,i}(m_j, \tilde{m}_i) = e_{j,i}(m_j, m_i) = 1$; thus, by Lemma 2, \tilde{m}_i is a strictly better response than m_i against m_j . Hence, we may assume $m_j^2 \in \Theta$ and consider the following four cases:

Case 2.1. $e_{j,i}(m_j, m_i) = \varepsilon$ and $e_{j,i}(m_j, \tilde{m}_i) = 0$.

It follows from (6) and Lemma 1 that

$$u_i(f(m_j^2), \theta_i) - u_i(C_{j,i}^\varepsilon(m_j, m_i), \theta_i) > 0,$$

where $f(m_j^2)$ is the outcome induced by (m_j, \tilde{m}_i) .

Case 2.2. $e_{j,i}(m_j, m_i) = 0$ and $e_{j,i}(m_j, \tilde{m}_i) = \varepsilon$.

Since $e_{j,i}(m_j, m_i) = 0$, we have $m_i^2 = \tilde{m}_i^2 = m_j^2$. Hence, $e_{j,i}(m_j, \tilde{m}_i) = \varepsilon$ implies that $x(m_j^2, \tilde{m}_i^1) = x(m_j^2, \theta_i) \neq f(m_j^2)$. Thus, it follows from (8) that

$$u_i(C_{j,i}^\varepsilon(m_j, \tilde{m}_i), \theta_i) - u_i(f(m_j^2), \theta_i) > 0,$$

where $f(m_j^2)$ is the outcome induced by (m_j, m_i) .

Case 2.3. $e_{j,i}(m_j, m_i) = e_{j,i}(\tilde{m}_j, m_i) = \varepsilon$.

It follows from Lemmas 2 and 1 that

$$u_i(C_{j,i}^\varepsilon(m_j, \tilde{m}_i), \theta_i) - u_i(C_{j,i}^\varepsilon(m_j, m_i), \theta_i) > 0.$$

Case 2.4. $e_{j,i}(m_j, m_i) = e_{j,i}(\tilde{m}_j, m_i) = 0$.

Both (m_j, m_i) and (m_j, \tilde{m}_i) lead to the same outcome $x(m_j^2, \tilde{m}_i^1) = x(m_j^2, m_i^1) = f(m_j^2)$.

In sum, as long as $e_{i,j}(m_i, m_j) = \varepsilon$ or $e_{j,i}(m_j, m_i) = \varepsilon$ (Case 1.1 and Cases 2.1-2.3), \tilde{m}_i is a strictly better response than m_i against m_j . Hence, in order for \tilde{m}_i not to be a profitable deviation, we must have $e_{i,j}(m_i, m_j) = e_{j,i}(m_j, m_i) = 0$.

A.5 Proof of Theorem 2

We first extend the notion of a *challenge scheme* to the case of SCCs. Fix agent i of type θ_i . For each state $\tilde{\theta} \in \Theta$, and $x \in F(\tilde{\theta})$, if $\mathcal{L}_i(x, \tilde{\theta}_i) \cap \mathcal{SU}_i(x, \theta_i) \neq \emptyset$, we select some $x(\tilde{\theta}, x, \theta_i) \in \mathcal{L}_i(x, \tilde{\theta}_i) \cap \mathcal{SU}_i(x, \theta_i)$; otherwise, we set $x(\tilde{\theta}, x, \theta_i) = x$. In the sequel, we define $F(\Theta) \equiv \bigcup_{\theta \in \Theta} F(\theta)$. Observe that $F(\Theta)$ is a finite set since each $F(\theta)$ is assumed to be finite.

As in the case of SCFs, the following lemma shows that there is a challenge scheme under which truth-telling induces the best allocation. In addition, here we choose the challenge scheme such that for every agent i , type θ_i , and state $\tilde{\theta}$ under which the challenge is effective (i.e. $x(\tilde{\theta}, x, \theta_i) \neq x$), no type $\theta_i'' \in \Theta_i$ is indifferent between $x(\tilde{\theta}, x, \theta_i)$ and an

allocation x' in $F(\Theta)$. The property can be achieved by adding a small transfer to $x(\tilde{\theta}, x, \theta_i)$ with $x(\tilde{\theta}, x, \theta_i) \neq x$, thanks to the finiteness of $F(\Theta)$ and Θ_i .

Lemma 3 *There is a challenge scheme $\{x(\tilde{\theta}, x, \theta_i)\}$ for an SCC F such that for every state $\tilde{\theta}$, every $x \in F(\tilde{\theta})$ and type θ_i ,*

$$u_i(x(\tilde{\theta}, x, \theta_i), \theta_i) \geq u_i(x(\tilde{\theta}, x, \theta'_i), \theta_i), \forall \theta'_i \in \Theta_i; \quad (16)$$

moreover, whenever, $x(\tilde{\theta}, x, \theta_i) \neq x$, we have

$$u_i(x(\tilde{\theta}, x, \theta_i), \theta''_i) \neq u_i(x', \theta''_i), \forall \theta''_i \in \Theta_i, \forall x' \in F(\Theta). \quad (17)$$

Proof. We first prove (17) by constructing a challenge scheme $\{x(\tilde{\theta}, x, \theta_i)\}$. Fix agent i of type θ_i . For each state $\tilde{\theta} \in \Theta$ and $x \in F(\tilde{\theta})$, if $\mathcal{L}_i(x, \tilde{\theta}_i) \cap \mathcal{SU}_i(x, \theta_i) = \emptyset$, we let $x(\tilde{\theta}, x, \theta_i) = x$. We define

$$\mathcal{S}(i, x, \tilde{\theta}, \theta) = \left\{ x'' \in X : u_i(x'', \tilde{\theta}_i) < u_i(x, \tilde{\theta}_i) \text{ and } u_i(x'', \theta_i) > u_i(x, \theta_i) \right\}.$$

Thus, $\mathcal{S}(i, x, \tilde{\theta}, \theta)$ is a nonempty open set. Now consider

$$\begin{aligned} & \mathcal{S}^*(i, x, \tilde{\theta}, \theta) \\ \equiv & \mathcal{S}(i, x, \tilde{\theta}, \theta) \setminus \{x'' \in X : u_i(x'', \theta''_i) = u_i(x', \theta''_i) \text{ for some } \theta''_i \in \Theta_i \text{ and some } x' \in F(\Theta)\}. \end{aligned}$$

Thanks to the finiteness of $F(\Theta)$ and Θ_i , $\mathcal{S}^*(i, x, \tilde{\theta}, \theta)$ is still a nonempty open set when we delete a finite set of elements. Now we choose an element $x(\tilde{\theta}, x, \theta_i) \in \mathcal{S}^*(i, x, \tilde{\theta}, \theta)$. Hence, we obtain (17). The proof of (16) is the same as the proof of Lemma 1 applied to the challenge scheme $\{x(\tilde{\theta}, x, \theta_i)\}$. ■

Next, we propose a mechanism $\mathcal{M} = ((M_i), g, (\tau_i))_{i \in \mathcal{I}}$ which will be used to prove the if-part of Theorem 2. We define the message space, allocation rule, and transfer rule below. First, a generic message of agent i is described as follows:

$$\begin{aligned} m_i &= (m_i^1, m_i^2, m_i^3) \in M_i = M_i^1 \times M_i^2 \times M_i^3 = \Theta_i \times \left[\times_{j=1}^I \Theta_j \right] \times F(\Theta) \text{ s.t.} \\ & m_i^2 \in \Theta \Rightarrow m_i^3 \in F(m_i^2). \end{aligned}$$

That is, agent i is asked to make (1) an announcements of agent i 's own type (which we denote by m_i^1); (2) an announcement of a type profile (which we denote by m_i^2); (3) an allocation which must belong to $F(m_i^2)$ if m_i^2 is a state. As we do in the case of SCFs, we write $m_{i,j}^2 = \tilde{\theta}_j$ if agent i 's report in m_i^2 that agent j 's type is $\tilde{\theta}_j$.

First, the allocation rule is defined as follows. For each $m \in M$, we apply exactly one of the following two rules:

Rule 1: If there exist $\tilde{\theta} \in \Theta$ and $x \in F(\tilde{\theta})$ such that every agent reports state $\tilde{\theta}$ in his second report and $I-1$ agents report lottery $m_i^3(\tilde{\theta}) = x$ according to their third announcement, then

$$g(m) = \frac{1}{I^2} \sum_{i \in \mathcal{I}} \sum_{j \in \mathcal{I}} \left[e_{i,j}(m_i, m_j) \frac{1}{I} \sum_{k \in \mathcal{I}} y_k(m_k^1) \oplus (1 - e_{i,j}(m_i, m_j)) x(\tilde{\theta}, x, m_j^1) \right]$$

where $\{y_k(\cdot)\}$ are the dictator lotteries for agent k as defined in Lemma 2 and

$$e_{i,j}(m_i, m_j) = \begin{cases} 0, & \text{if } m_i^2 \in \Theta, m_i^2 = m_j^2, \text{ and } x(m_i^2, m_i^3, m_j^1) = m_i^3; \\ \varepsilon, & \text{if } m_i^2 \in \Theta, \text{ and } [m_i^2 \neq m_j^2 \text{ or } x(m_i^2, m_i^3, m_j^1) \neq m_i^3]; \\ 1, & \text{if } m_i^2 \notin \Theta. \end{cases}$$

Rule 2: Otherwise,

$$g(m) = \frac{1}{I^2} \sum_{i \in \mathcal{I}} \sum_{j \in \mathcal{I}} \left[e_{i,j}(m_i, m_j) \frac{1}{I} \sum_{k \in \mathcal{I}} y_k(m_k^1) \oplus (1 - e_{i,j}(m_i, m_j)) x(m_i^2, m_i^3, m_j^1) \right].$$

Next, we define

$$C_{i,j}^\varepsilon(m) \equiv \varepsilon \times \frac{1}{I} \sum_{k \in \mathcal{I}} y_k(m_k^1) \oplus (1 - \varepsilon) \times x(m_i^2, m_i^3, m_j^1).$$

For every message profile m and agent j , we can choose $\varepsilon > 0$ sufficiently small such that (i) it does not disturb the ‘‘effectiveness’’ of agent j 's challenge, i.e.,

$$\begin{aligned}
x(m_i^2, m_i^3, m_j^1) &\neq m_j^3 \Rightarrow \\
u_j(C_{i,j}^\varepsilon(m), m_{i,j}^2) &< u_j(m_i^3, m_{i,j}^2) \text{ and } u_j(C_{i,j}^\varepsilon(m), m_j^1) > u_j(m_i^3, m_j^1); \tag{18}
\end{aligned}$$

moreover, (ii) an “effective self-challenge” of agent j induces a generic outcome such that at each state, no agent is indifferent between the resulting outcome and any outcome in $F(\Theta)$, i.e.,

$$\begin{aligned}
&x(m_j^2, m_j^3, m_j^1) \neq m_j^3 \Rightarrow \\
\frac{1}{I}u_j(C_{j,j}^\varepsilon(m), \theta_j) + \left(1 - \frac{1}{I}\right)u_j(m_j^3, \theta_j) &\neq \frac{1}{I}u_j(x, \theta_j) + \left(1 - \frac{1}{I}\right)u_j(x, \theta_j) \tag{19} \\
&\text{for any } \theta \text{ and any } x \in F(\Theta).
\end{aligned}$$

Observe that (ii) is possible because inequality (17) holds in Lemma 3; moreover, by (6) in Lemma 2, $u_j(C_{j,j}^\varepsilon(m), \theta_j)$ is a strictly decreasing function in ε .

Second, the transfer rule from agent i is specified as follows:

$$\tau_i(m) = \sum_{j \neq i} [\bar{\tau}_{i,j}^1(m) + \bar{\tau}_{i,j}^2(m) + \tau_{i,j}^3(m)]$$

where $\bar{\tau}_{i,j}^1(m)$ double the scale of $\tau_{i,j}^1(m)$ and $\tau_{i,j}^2(m)$ defined in Section 4.1.5 (i.e., $\bar{\tau}_{i,j}^1(m) = 2\tau_{i,j}^1(m)$ and $\bar{\tau}_{i,j}^2(m) = 2\tau_{i,j}^2(m)$); moreover, we introduce one more transfer rule:

$$\tau_{i,j}^3(m) = \begin{cases} 0, & \text{if } x(m_i^2, m_i^3, m_j^1) = m_j^3; \\ -\eta, & \text{if } x(m_i^2, m_i^3, m_j^1) \neq m_j^3. \end{cases}$$

That is, agent i is asked to pay η if his reported outcome m_i^3 is challenged by agent $j \neq i$. Note that we still require that η be greater than the payoff difference by (7) in Section 4.1.5.

To prove Theorem 2, we first observe that we have a stronger statement than Claim 1 since each agent i 's dictator lotteries are triggered whenever there is a pair of agents (j, k) such that $e_{j,k}(m_j, m_k) = \varepsilon$ where $j, k \in \mathcal{I}$. The proof of the additional claim is identical to the proof of Case 1.1 in Claim 1.

Claim 5 *Let σ be a Nash equilibrium of the game $\Gamma(\mathcal{M}, \theta)$. If $m_i^1 \neq \theta_i$ for some $m_i \in \text{supp}(\sigma_i)$, then for each agent $j \neq i$, we have $e_{i,j}(m_i, m_j) = e_{j,i}(m_j, m_i) = 0$ with σ_j -probability one.*

In addition, for any each agents $j, k \in \mathcal{I} \setminus \{i\}$, $m_j \in \text{supp}(\sigma_j)$, and $m_k \in \text{supp}(\sigma_k)$, we have $e_{j,k}(m_j, m_k) = e_{k,j}(m_k, m_j) = 0$.

Then, Claims 2 and 3 hold with exactly the same proof. Again, we denote the true state by θ and (by Claim 3) the common state announced in the agents' second report by $\tilde{\theta}$. In the following, we establish Claim 6 as the counterpart of Claim 4 in Theorem 1 in the modified mechanism above.

Claim 6 *No one challenges an allocation announced in the third report of the other agents, i.e., for every pair of agents i and j where $i \neq j$, $m_i \in \text{supp}(\sigma_i)$, and $m_j \in \text{supp}(\sigma_j)$, we have $x(\tilde{\theta}, m_i^3, m_j^1) = m_i^3$.*

Proof. For each $x \in F(\Theta)$, we define the following set of agents:

$$\mathcal{J}(x) \equiv \left\{ j \in \mathcal{I} : \mathcal{L}_j(x, \tilde{\theta}_j) \cap \mathcal{SU}_j(x, \theta_j) = \emptyset \right\}.$$

First, if $j \in \mathcal{J}(m_i^3)$, then $x(\tilde{\theta}, m_i^3, m_j^1) \neq m_i^3$ implies that $x(\tilde{\theta}, m_i^3, m_j^1)$ is weakly worse than m_i^3 under the true type of agent j . In addition, the dictator lotteries are triggered when $x(\tilde{\theta}, m_i^3, m_j^1) \neq m_i^3$. By (6) in Lemma 2, the dictator lotteries deliver worse allocations than every alternative in $F(\Theta)$, we know that agent j can profitably deviate from m_j to $\tilde{m}_j = (\theta_j, m_j^2, m_j^3)$. Hence, to establish the claim, it suffices to prove that $\mathcal{J}(m_i^3) = \mathcal{I}$ for each message $m_i \in \text{supp}(\sigma_i)$ and each agent $i \in \mathcal{I}$.

Suppose to the contrary that for some agent i and some message $m_i \in \text{supp}(\sigma_i)$, we have agent $j \notin \mathcal{J}(m_i^3)$ and $j \neq i$. That is, we have

$$\mathcal{L}_j(m_i^3, \tilde{\theta}_j) \cap \mathcal{SU}_j(m_i^3, \theta_j) \neq \emptyset. \quad (20)$$

First, we claim that agent j will challenge m_i^3 with probability one, i.e., $x(\tilde{\theta}, m_i^3, m_j^1) \neq m_i^3$ for every $m_j \in \text{supp}(\sigma_j)$. Observe that if $x(\tilde{\theta}, m_i^3, m_j^1) = m_i^3$ for some m_j^1 , agent j can deviate to announce $\tilde{m}_j = (\theta_j, m_j^2, m_j^3)$. The deviation is profitable, since $x(\tilde{\theta}, m_i^3, \theta_j) \in \mathcal{SU}_j(f(\tilde{\theta}), \theta_j)$.

Second, we claim that if agent j will challenge m_i^3 with probability one, then by playing m_i agent i suffers from the penalty η according to $\tau_{i,j}^3$. We derive a contradiction by showing that agent i can profitably deviate in each of the following two cases. Firstly, if there is some $x \in F(\tilde{\theta})$ such that $x(\tilde{\theta}, x, m_k^1) = x$ with σ_k -probability one for agent $k \neq i$, then agent i can profitably deviate to \tilde{m}_i which is identical to m_i except that $\tilde{m}_i^3 = x$. By doing so agent i

avoids paying the penalty η for being challenged by agent $k \neq i$. By (7), it is a profitable deviation.

Secondly, suppose that for each $x \in F(\tilde{\theta})$, there is some agent $k_x \neq i$ such that $x(\tilde{\theta}, x, m_{k_x}^1) \neq x$ for some $m_{k_x} \in \text{supp}(\sigma_{k_x})$. That is, for every $m_k \in \text{supp}(\sigma_k)$ with $m_k^3 = x$, there exists some agent $k_x \neq i$ with a message $m_{k_x} \in \text{supp}(\sigma_{k_x})$ such that $e_{k, k_x}(m_k, m_{k_x}) = \varepsilon$. Then, by Claim 5, we know that $m_k^1 = \theta_k$ with σ_k -probability one. By Claim 2, we conclude that $\tilde{\theta} = \theta$. That, together with (20), implies that $\mathcal{L}_j(m_i^3, \theta_j) \cap \mathcal{SU}_j(m_i^3, \theta_j) \neq \emptyset$, which is impossible. ■

Claim 7 *No one challenges an allocation announced in his own third report, i.e., for every agent i , $m_i \in \text{supp}(\sigma_i)$, we have $x(\tilde{\theta}, m_i^3, m_i^1) = m_i^3$.*

Proof. Note that all agents report a common state $\tilde{\theta}$ by Claim 3. Suppose to the contrary that there exist agent i and some message $\tilde{m}_i \in \text{supp}(\sigma_i)$ such that $x(\tilde{\theta}, m_i^3, \tilde{m}_i^1) \neq \tilde{m}_i^3$. It follows that

$$\mathcal{L}_i(\tilde{m}_i^3, \tilde{\theta}_j) \cap \mathcal{SU}_i(\tilde{m}_i^3, \theta_i) \neq \emptyset. \quad (21)$$

By Claim 5, for every $\tilde{m}_i \in \text{supp}(\sigma_i)$ such that $x(\tilde{\theta}, m_i^3, \tilde{m}_i^1) \neq \tilde{m}_i^3$, we have $\tilde{m}_i^1 = \theta_i$, and $m_k^1 = \theta_k$ for each $m_k \in \text{supp}(\sigma_k)$ and agent $k \neq i$. We further conclude that for every $\tilde{m}_i \in \text{supp}(\sigma_i)$, we have $x(\tilde{\theta}, m_i^3, \tilde{m}_i^1) \neq \tilde{m}_i^3$. Suppose on the contrary that there exists some $m_i \in \text{supp}(\sigma_i)$ such that $x(\tilde{\theta}, m_i^3, m_i^1) = m_i^3$. For every $m_{-i} \in \text{supp}(\sigma_{-i})$, we consider four different situations: (i) when (j, i) with $j \neq i$ is chosen, by Claim 6, both (\tilde{m}_i, m_{-i}) and (m_i, m_{-i}) induce the outcome m_j^3 ; (ii) when (j, j) with $j \neq i$ is chosen, since we have $m_j^1 = \theta_j$ with σ_j -probability one, it follows from Claim 2 that $m_{j,j}^2 = \theta_j$ with σ_j -probability one. Hence, we know that agent j does not challenge himself. Thus, (\tilde{m}_i, m_{-i}) and (m_i, m_{-i}) both induce the outcome m_j^3 ; (iii) when (i, j) with $j \neq i$ is chosen, by Claim 6 again, the outcome induced by (\tilde{m}_i, m_{-i}) is \tilde{m}_i^3 and the outcome induced by (m_i, m_{-i}) is m_i^3 ; and (iv) when (i, i) is chosen, (\tilde{m}_i, m_{-i}) induces the outcome $C_{i,i}^\varepsilon(\tilde{m}_i, m_{-i})$ and (m_i, m_{-i}) induces m_i^3 . To summarize, against every $m_{-i} \in \text{supp}(\sigma_{-i})$ the payoff difference between m_i and \tilde{m}_i , is as

follows,

$$\begin{aligned}
& \frac{1}{I^2} u_i(C_{i,i}^\varepsilon(\tilde{m}_i, m_{-i}), \theta_i) + \frac{1}{I^2} \sum_{j \neq i} u_i(\tilde{m}_i^3, \theta_i) - \left[\frac{1}{I^2} u_i(m_i^3, \theta_i) + \frac{1}{I^2} \sum_{j \neq i} u_i(m_i^3, \theta_i) \right] \\
&= \frac{1}{I^2} u_i(C_{i,i}^\varepsilon(\tilde{m}_i, m_{-i}), \theta_i) + \frac{I-1}{I^2} u_i(\tilde{m}_i^3, \theta_i) - \left[\frac{1}{I^2} u_i(x, \theta_i) + \frac{I-1}{I^2} u_i(x, \theta_i) \right] \\
&\neq 0
\end{aligned}$$

where the last non-equality follows from (19), i.e., agent i is never indifferent between the outcomes induced by m_i and \tilde{m}_i . It contradicts the fact that $m_i \in \text{supp}(\sigma_i)$. Hence, for every $\tilde{m}_i \in \text{supp}(\sigma_i)$, we have $\tilde{m}_i^1 = \theta_i$. Hence, $m_j^1 = \theta_j$ for each $m_j \in \text{supp}(\sigma_k)$ and agent $j \in \mathcal{I}$. By Claim 2, we conclude that $\tilde{\theta} = \theta$. It is a contradiction to (21). ■

It only remains to prove the existence of pure-strategy Nash equilibrium.

Claim 8 *For every $\theta \in \Theta$ and $x \in F(\theta)$, there exists a pure-strategy Nash equilibrium $m \in M$ of the game $\Gamma(\mathcal{M}, \theta)$ such that $g(m) = x$ and $\tau_i(m) = 0$ for every $i \in \mathcal{I}$.*

Proof. Fix an arbitrary allocation $x \in F(\theta)$. We argue that truth-telling (i.e., $m_i = (\theta_i, \theta, x)$ for each i) constitutes a pure-strategy equilibrium of the game $\Gamma(\mathcal{M}, \theta)$. Note that reporting \tilde{m}_i with $\tilde{m}_i^1 = \theta_i$, $\tilde{m}_i^2 = \theta$, and $\tilde{m}_i^3 \neq x$ instead of m_i affects neither the allocation nor the transfer. The argument for proving that either $\tilde{m}_i^1 \neq \theta_i$ or $\tilde{m}_i^2 \neq \theta$ cannot be a profitable unilateral deviation for every agent i is identical to the proof of Theorem 1. ■

A.6 Proof of Theorem 3

Recall that in the mechanism which we use to prove Theorem 1, agent i 's generic message is $m_i = (m_i^1, m_i^2) \in \Theta_i \times [\times_{j=1}^I \Theta_j]$. We expand m_i^2 into H copies of $[\times_{j=1}^I \Theta_j]$ and define

$$m_i = (m_i^1, m_i^2, \dots, m_i^{H+1}) \in \Theta_i \times \underbrace{[\times_{j=1}^I \Theta_j] \times \dots \times [\times_{j=1}^I \Theta_j]}_{H \text{ terms}}$$

where H is a positive integer to be chosen later. For each message profile $m \in M$, the allocation is defined as follows:

$$g(m) = \frac{1}{I(I-1)} \sum_{i \in \mathcal{I}} \sum_{j \neq i} \left[e_{i,j}(m_i, m_j) \frac{1}{2} \sum_{k=i,j} y_k(m_k^1) \oplus \frac{1 - e_{i,j}(m_i, m_j)}{H} \left[x(m_i^2, m_j^{H+2}) \oplus \sum_{h=3}^{H+1} \phi(m^h) \right] \right]$$

where $\{y_k(\cdot)\}$ are the dictator lotteries²⁰ for agent k defined in Lemma 2, $\phi(\cdot)$ is an outcome function such that

$$\phi(m^h) = \begin{cases} f(\tilde{\theta}), & \text{if } m_i^h = \tilde{\theta} \in \Theta \text{ for at least } I - 1 \text{ agents;} \\ b, & \text{otherwise, where } b \text{ is an arbitrary outcome in } A, \end{cases}$$

and

$$e_{i,j}(m_i, m_j) = \begin{cases} 0, & \text{if } m_i^2 \in \Theta, m_i^2 = m_j^2 = m_i^h = m_j^h \text{ and } x(m_i^2, m_j^{H+2}) = f(m_i^2), \forall h \in \{3, \dots, H+1\}; \\ 1, & \text{if } m_i^2 \notin \Theta; \\ \varepsilon, & \text{otherwise.} \end{cases}$$

We now define the transfer rule. For every message profile $m \in M$ and agent $i \in \mathcal{I}$, we specify the transfer to agent i as follows:

$$\tau_i(m) = \sum_{j \neq i} [\tau_{i,j}^{1,2}(m) + \tau_{i,j}^{2,2}(m)] + \sum_{h=3}^{H+1} \tau_i^h(m) + d_i(m^2, \dots, m^{H+1})$$

where $\gamma, \kappa, \xi > 0$ (their size are determined later)

$$\begin{aligned} \tau_{i,j}^{1,2}(m) &= \begin{cases} 0, & \text{if } m_{i,j}^2 = m_{j,j}^2; \\ -\gamma & \text{if } m_{i,j}^2 \neq m_{j,j}^2 \text{ and } m_{i,j}^2 \neq m_j^1; \\ \gamma & \text{if } m_{i,j}^2 \neq m_{j,j}^2 \text{ and } m_{i,j}^2 = m_j^1. \end{cases} \\ \tau_{i,j}^{2,2}(m) &= \begin{cases} 0, & \text{if } m_{i,i}^2 = m_{j,i}^2; \\ -\gamma, & \text{if } m_{i,i}^2 \neq m_{j,i}^2; \end{cases} \end{aligned}$$

moreover, for every $h \in \{3, \dots, H+1\}$,

$$\tau_i^h(m) = \begin{cases} -\kappa, & \text{if there exists } \tilde{\theta} \text{ such that } m_i^h \neq \tilde{\theta} \text{ but } m_j^h = \tilde{\theta} \text{ for all } j \neq i; \\ 0, & \text{otherwise,} \end{cases}$$

²⁰Although the dictator lotteries may contain transfers, we do not take into account the scale of transfers in it. To dispense with the transfers in the dictator lotteries, we can use an arbitrarily small amount of money to make the best challenge schemes and social choice function generic, i.e., any two resulting outcomes are distinct from each other.

and

$$d_i(m^2, \dots, m^{H+1}) = \begin{cases} -\xi, & \text{if there exists } h \in \{3, \dots, H+1\} \text{ such that } m_i^h \neq m_i^2 \text{ and } m_j^{h'} = m_j^2, \\ & \text{for all } h' \in \{2, \dots, h-1\} \text{ and all } j \neq i; \\ 0, & \text{otherwise.} \end{cases}$$

Finally, we choose positive numbers γ, ξ, H, κ , and ε such that

$$\begin{aligned} \bar{\tau} &> \gamma + (H-1)\kappa + \xi \\ \gamma &> \xi + \varepsilon\eta \\ \kappa &> \varepsilon\eta \\ \xi &> \frac{1}{H}\eta + \kappa. \end{aligned}$$

More precisely, we first fix $\bar{\tau}$ and choose $\gamma < \frac{1}{3}\bar{\tau}$ and $\xi < \min\{\frac{1}{3}\bar{\tau}, \gamma\}$. Second, we choose H large enough so that $\xi > \frac{1}{H}\eta$. Third, we choose κ small enough such that $(H-1)\kappa < \frac{1}{3}\bar{\tau}$ and $\xi > \frac{1}{H}\eta + \kappa$. Fourth, we choose ε small enough such that $\gamma > \xi + \varepsilon\eta$ and $\kappa > \varepsilon\eta$. We can now prove Theorem 3 following the three steps as in the proof of Theorem 1.

A.6.1 Contagion of Truth

First, note that Claims 1 and 2 hold. The proof of Claim 2 applies with only one minor difference: Here m_i^2 may affect agent i 's payoff through $d_i(\cdot)$. However, a similar argument follows, since we have $\gamma > \xi + \varepsilon\eta$.²¹ Let θ denote the true state.

Claim 9 *If every agent j reports the truth in his first report σ_j -probability one, then every agent j reports the truth in his 2nd, ..., (H + 1)th report. That is, $m_j^h = \theta$ for $h = 2, \dots, H+1$.*

By Claims 1 and 2, every agent j reports the state truthfully in his 2nd report. Then, we can follow verbatim the argument on p. 12 of [Abreu and Matsushima \(1994\)](#) which shows that every agent j reports the state truthfully in his h th report for every $h = 2, \dots, H+1$.

A.6.2 Consistency

Claim 10 *There exists a state $\tilde{\theta}$ such that every agent announces $\tilde{\theta}$ in the second report all the way to the last/(H + 1)th report with probability one.*

²¹It corresponds to Property (b) in [Abreu and Matsushima \(1994\)](#).

Proof. We prove consistency by considering the two cases as in the proof of Claim 3. The proof for the first case remains the same. For the second case, suppose that one agent, say i , tells a lie in the first report. As agent i believes that all the other agents report the same state $\tilde{\theta}$ in their second all the way to the last report. By the same argument in the second case in the proof of Claim 3, we can show that agent i announces $\tilde{\theta}$ in the second report with probability one. In addition, for every $h = 2, \dots, H + 2$, as agent i believes that all the other agent report the same state $\tilde{\theta}$, by the rule $\phi(m^h)$ and $\tau_i^h(m^h)$, we know $m_i^h = \tilde{\theta}$. ■

A.6.3 No Challenge

Claim 11 *No agent challenges with positive probability the common state $\tilde{\theta}$ announced in the second report.*

Proof. The argument is the same as the proof of Claim 4. ■

A.7 Proof of Theorem 4

Recall that we assume that set of alternatives A is a finite set, and the state space Θ is a Polish (i.e., separable and complete metric) space. Recall the definition of Θ_i introduced in Section 4.1. For every $\ell \in \Delta(A)$, we write $v_i(\ell, \theta) = \ell \cdot \bar{v}_i(\theta_i)$ where $\bar{v}_i(\theta_i) \in [0, 1]^{|A|}$ is a vector of utilities over A induced by $v_i(\cdot, \theta)$. Assume that agents' utilities remain bounded; if f specifies some transfers in outcome, the transfers are also bounded in money and $\eta > 0$ still satisfies condition (4). Let $\bar{X} \equiv \Delta(A) \times [-2\eta, 2\eta]^I$ and identify \bar{X} with a compact subset of $\mathbb{R}^{I+|A|}$ (endowed with the Euclidean topology). Let d denote the metric on Θ , d_i the metric on Θ_i , and $\rho : \bar{X} \times \bar{X} \rightarrow \mathbb{R}$ a metric on the outcome space. We endow $\times_{j=1}^I \Theta_j$ with the product topology and $\times_{j=1}^I \Theta_j$ and \bar{X} with the Borel σ -algebra. We say that the setting is *compact and continuous* if Θ is compact and $(v_i(a, \cdot))_{a \in A, i \in \mathcal{I}}$ and the SCFs are all continuous functions on Θ .

We introduce the following version of challenge scheme. First, for $(x, \theta_i) \in X \times \Theta_i$, we use $\mathcal{SL}_i(x, \theta_i)$ to denote the strict lower-contour set at allocation x for type θ_i , i.e.,

$$\mathcal{SL}_i(x, \theta_i) = \{x' \in X : u_i(x, \theta_i) > u_i(x', \theta_i)\}.$$

For agent i of type θ_i , allocation $x \in \bar{X}$, and state $\tilde{\theta} \in \Theta$, we construct the following

compound lottery:

$$\ell(x, \tilde{\theta}) = \frac{\rho(x, f(\tilde{\theta}))}{1 + \rho(x, f(\tilde{\theta}))} x \oplus \frac{1}{1 + \rho(x, f(\tilde{\theta}))} f(\tilde{\theta}).$$

Define

$$x(\tilde{\theta}, x) = \begin{cases} \ell(x, \tilde{\theta}), & \text{if } x \in \mathcal{SL}_i(f(\tilde{\theta}), \tilde{\theta}_i); \\ f(\tilde{\theta}), & \text{otherwise.} \end{cases}$$

As we mentioned in the main text, in the infinite setting we know of no way to construct a challenge scheme by pre-selecting a test allocation (depending on type θ_i) in a continuous manner. As a result, we cannot have the agents report their type (let alone the true type) to cast a challenge to state $\tilde{\theta}$. Instead, we will restore the continuity of the outcome function by asking them to report an allocation x directly (see Section A.7.1). The definition of strict lower-contour sets can be found in Section A.5.

Claim 12 $x(\tilde{\theta}, x)$ is a continuous function on $\bar{X} \times \Theta$.

Proof. Since $\rho(\cdot, \cdot)$ is continuous, we have that $\ell(\cdot, \cdot)$ is continuous. We show that

$$x(\tilde{\theta}[n], x[n]) \rightarrow x(\tilde{\theta}, x)$$

in each of the following two cases.

Case 1. $x \in \mathcal{SL}_i(f(\tilde{\theta}), \tilde{\theta}_i)$.

In this case, $x(\tilde{\theta}, x) = \ell(x, f(\tilde{\theta}))$. Since f and u_i are both continuous, it follows that $x[n] \in \mathcal{SL}_i(f(\tilde{\theta}[n]), \tilde{\theta}_i[n])$ for each n large enough. Thus, $x(\tilde{\theta}[n], x[n]) = \ell(x[n], \tilde{\theta}[n])$. Hence, $x(\tilde{\theta}[n], x[n]) \rightarrow \ell(x, f(\tilde{\theta}))$ as $(x[n], \tilde{\theta}[n]) \rightarrow (x, \tilde{\theta})$.

Case 2. $x \notin \mathcal{SL}_i(f(\tilde{\theta}), \tilde{\theta}_i)$.

In this case, $x(\tilde{\theta}, x) = f(\tilde{\theta})$. If there is some integer \bar{n} such that $x[n] \notin \mathcal{SL}_i(f(\tilde{\theta}[n]), \tilde{\theta}_i[n])$ for every $n \geq \bar{n}$, then $x(\tilde{\theta}[n], x[n]) = f(\tilde{\theta}[n])$. Since f is continuous and $\tilde{\theta}[n] \rightarrow \tilde{\theta}$, it follows that $x(\tilde{\theta}[n], x[n]) \rightarrow f(\tilde{\theta})$. Now suppose that there is a subsequence of $x[n], \tilde{\theta}[n]$, say itself, such that $x[n] \in \mathcal{SL}_i(f(\tilde{\theta}[n]), \tilde{\theta}_i[n])$ for every n . Then, we have $x(\tilde{\theta}[n], x[n]) = \ell(x[n], f(\tilde{\theta}[n]))$. Since ρ is jointly continuous, we must have $\rho(x[n], f(\tilde{\theta}[n])) \rightarrow \rho(x, f(\tilde{\theta}))$. Since $x \notin \mathcal{SL}_i(f(\tilde{\theta}), \tilde{\theta}_i)$, it follows that $\rho(x, f(\tilde{\theta})) = 0$. By construction of $\ell(x[n], f(\tilde{\theta}[n]))$, we must have $\ell(x[n], f(\tilde{\theta}[n])) \rightarrow f(\tilde{\theta})$. Hence, $x(\tilde{\theta}[n], x[n]) \rightarrow f(\tilde{\theta})$. ■

Lemma 4 For each $i \in \mathcal{I}$, there exists a continuous function $y_i : \Theta_i \rightarrow X$ such that for all types θ_i and θ'_i of agent i with $\theta_i \neq \theta'_i$, we have

$$u_i(y_i(\theta_i), \theta_i) > u_i(y_i(\theta'_i), \theta_i); \quad (22)$$

and for each type θ'_j of agent $j \in \mathcal{I}$, we also have that for every $x \in f(\Theta)$

$$u_i(y_j(\theta'_j), \theta_i) < u_i(x, \theta_i). \quad (23)$$

Moreover, $y_i(\cdot)$ is continuous on Θ_i .

Proof. We construct the dictator lotteries in the infinite state space. We first construct $\ell_i(\theta_i) \in \Delta(A)$ for each $\theta_i \in \Theta_i$ and let

$$y_i(\theta_i) = (\ell_i(\theta_i), -2\eta, \dots, -2\eta).$$

Hence, we obtain $u_i(a, \theta_i) > u_i(y_k(\theta'_k), \theta_i)$ for all type θ_i and type θ'_k of agents i and k . Let ℓ^* be the uniform lottery over A , i.e., $\ell^*[a] = 1/|A|$. Pick $r < 1/|A|$. Consider the maximization problem as follows:

$$\begin{aligned} & \max_{\ell \in \Delta(A)} \ell \cdot \bar{v}_i(\theta_i) \\ \text{s.t. } & \|\ell - \ell^*\| \leq r \end{aligned}$$

The Kuhn-Tucker condition for $\ell_i(\theta_i)$ to be the solution is

$$\bar{v}_i(\theta_i) - 2\lambda_i(\theta_i)(\ell_i(\theta_i) - \ell^*) = 0$$

We know that $\bar{v}_i(\theta_i)$ is not a zero vector. Hence, $\lambda_i(\theta_i) > 0$ and $\ell_i(\theta_i)$ is equal to the normalization of $\frac{1}{2} \left(\frac{\bar{v}_i(\theta_i)}{\lambda_i(\theta_i)} + \ell^* \right)$ as a lottery. For every $\theta_i \neq \theta'_i$, since $\bar{v}_i(\theta_i)$ is not an affine transform of $\bar{v}_i(\theta'_i)$, it follows that $\ell_i(\theta_i) \neq \ell_i(\theta'_i)$. Moreover, by Theorem of the maximum, $\ell_i(\cdot)$ is a continuous function on Θ_i . ■

A.7.1 The Mechanism

A generic message of agent i is described as follows:

$$m_i = (m_i^1, m_i^2, m_i^3) \in M_i = M_i^1 \times M_i^2 \times M_i^3 = \Theta_i \times [\times_{j=1}^I \Theta_j] \times \bar{X}.$$

That is, agent i is asked to make (1) one announcement of agent i 's type (i.e., m_i^1); and (2) one announcement of a type profile (i.e., m_i^2); and (3) one announcement of an allocation (i.e., m_i^3). As in the main text, we write $m_{i,j}^2 = \tilde{\theta}_j$ if agent i reports in m_i^2 that agent j 's type is $\tilde{\theta}_j$.

A.7.1.1 Allocation Rule

For each message profile $m \in M$, the allocation is defined as follows:

$$g(m) = \frac{1}{I(I-1)} \sum_{i \in \mathcal{I}} \sum_{j \neq i} \left[e_{i,j}(m_i, m_j) \frac{1}{2} \sum_{k=i,j} y_k(m_k^1) \oplus (1 - e_{i,j}(m_i, m_j)) x(m_i^2, m_j^3) \right]$$

where $y_k(\theta_k) = (l_k(\theta_k), t_1(\theta_k), \dots, t_I(\theta_k))$ is the dictator lottery for agent k with type θ_k defined in Lemma 4 and we define

$$e_{i,j}(m_i, m_j) \equiv \min \left\{ \max \left\{ \tilde{d}(m_i^2, m_j^2), d(m_i^2, m_j^2), \rho(m_j^3, f(m_i^2))^3 \right\}, 1 \right\},$$

where²²

$$\tilde{d}(m_i^2, m_j^2) = \inf_{\theta \in \Theta} d(m_i^2, \theta) + \inf_{\theta \in \Theta} d(m_j^2, \theta). \quad (24)$$

For each message profile (m_i, m_j) of agents i and j , let

$$C_{i,j}(m_i, m_j) \equiv e_{i,j}(m_i, m_j) \frac{1}{2} \sum_{k=i,j} y_k(m_k^1) \oplus (1 - e_{i,j}(m_i, m_j)) x(m_i^2, m_j^3).$$

In sum, with probability $\frac{1}{I(I-1)}$ an ordered pair (i, j) is chosen, then $C_{i,j}(m_i, m_j)$ is implemented.

Claim 13 *The outcome function g is continuous.*

²²In comparison with the $e_{i,j}(\cdot)$ defined in the proof of Theorem 1, here the terms of \tilde{d} and d correspond to the consistency check and the term ρ corresponds to the no challenge check.

Proof. It follows from Claim 12 and Lemma 4. ■

A.7.1.2 Transfer Rule

We now define the transfer rule. For every message profile $m \in M$ and agent $i \in \mathcal{I}$, we specify the transfer to agent i as

$$\tau_i(m) = \sum_{j \neq i} [\tau_{i,j}^1(m) + \tau_{i,j}^2(m)],$$

where $\tau_{i,j}^1$ and $\tau_{i,j}^2$ will be defined as follows: Given a message profile m and agent j , let $\tilde{m}_i^m = (m_i^1, (m_j^1, m_{i,-j}^2), m_i^3)$ (which replaces $m_{i,j}^2$ in m_i by m_j^1), $\hat{m}_i^m = (m_i^1, (m_{j,j}^2, m_{i,-j}^2), m_i^3)$ (which replaces $m_{i,j}^2$ in m_i by $m_{j,j}^2$), and $\bar{m}_i^m = (m_i^1, (m_{j,i}^2, m_{i,-i}^2), m_i^3)$ (which replaces $m_{i,i}^2$ by $m_{j,i}^2$). We define $\tau_{i,j}^1$ as follows:

$$\begin{aligned} \tau_{i,j}^1(m) &= - \sup_{\theta'_i} |u_i(g(m), \theta'_i) - u_i(g(\tilde{m}_i^m, m_{-i}), \theta'_i)| \\ &\quad + \sup_{\theta'_i} |u_i(g(\hat{m}_i^m, m_{-i}), \theta'_i) - u_i(g(\tilde{m}_i^m, m_{-i}), \theta'_i)| \\ &\quad + d_j(m_{j,j}^2, m_j^1) - d_j(m_{i,j}^2, m_j^1). \end{aligned} \tag{25}$$

Observe that $\tau_{i,j}^1$ satisfies two important properties:

1. neither $u_i(g(\hat{m}_i^m, m_{-i}), \theta'_i) - u_i(g(\tilde{m}_i^m, m_{-i}), \theta'_i)$ nor $d_j(m_{j,j}^2, m_j^1)$ depends on agent i 's choice of $m_{i,j}^2$, since $m_{i,j}^2$ has been replaced by agent j 's announcements in both \hat{m}_i^m and \tilde{m}_i^m ;
2. $\tau_{i,j}^1(m) = 0$ if $m_{i,j}^2 = m_{j,j}^2$.

We next define $\tau_{i,j}^2$ as follows:

$$\tau_{i,j}^2(m) = - \sup_{\theta'_i} |u_i(g(m), \theta'_i) - u_i(g(\bar{m}_i^m, m_{-i}), \theta'_i)| - d_i(m_{i,i}^2, m_{j,i}^2) \tag{26}$$

We say that a function $\alpha(\cdot)$ between two metric spaces S and Y , both endowed with the Borel σ -algebra, is *analytic* if its pre-image of every open set on Y is an analytic set. Since every analytic set is universally measurable, an analytic function is “almost” a measurable function (see pp. 498-499 of [Stinchcombe and White \(1992\)](#)). We show below that the

mechanism which we are about to construct has an analytic outcome function and transfer rule. Hence, whenever we fix a mixed-strategy Nash equilibrium which is a Borel probability measure on M , we can work with the σ -completion of the Borel σ -algebra on M to make all the expected payoffs well defined.²³

Claim 14 *The transfer rule $\tau_i : M \rightarrow \mathbb{R}$ is an analytic function. Moreover, if the setting is compact and continuous, then $\tau_i(\cdot)$ is a continuous function.*

Proof. It follows from Theorem 2.17 of [Stinchcombe and White \(1992\)](#) that $\tau_i(\cdot)$ is an analytic function. Suppose that the setting is compact and continuous. Then, by Claim 13, g is also continuous on M . Moreover, by the theorem of maximum, $\tau_{ij}^1(\cdot)$ and $\tau_{ij}^2(\cdot)$ are continuous on M . Hence $\tau_{ij}^1(\cdot)$ and $\tau_{ij}^2(\cdot)$ are both continuous. ■

With the claims above, the implementing mechanism $\mathcal{M} = ((M_i), g, (\tau_i))_{i \in \mathcal{I}}$ has been defined; moreover, when the setting is compact and continuous, \mathcal{M} is a mechanism with compact sets of message and continuous outcome function and transfer rule. To show that implementation is achieved by the constructed mechanism, we only emphasize arguments different from those in a finite state space. Before we provide the main argument, we establish two lemmas which play an important role in the proof of Theorem 4.

Throughout the proof, we denote by θ the true state. First, we show that it is strictly worse for every agent to challenge the truth.

Lemma 5 *Let (m_i, m_j) be a message profile of agents i and j with $x(m_i^2, m_j^3) \neq f(m_i^2)$. Then, $u_j(C_{i,j}(m_i, m_j), m_{i,j}^2) < u_j(f(m_i^2), m_{i,j}^2)$.*

Proof. Since $x(m_i^2, m_j^3) \neq f(m_i^2)$, we have $u_j(x(m_i^2, m_j^3), m_{i,j}^2) < u_j(f(m_i^2), m_{i,j}^2)$. Moreover, by (23) for every $x \in f(\Theta)$ we have $u_i(x, \theta_i) > u_i(y_k(\theta'_k), \theta_i)$ for each type θ_i and type θ'_k of agents i and k , we conclude that

$$u_j(C_{i,j}(m_i, m_j), m_{i,j}^2) < u_j(f(m_i^2), m_{i,j}^2). \quad (27)$$

This completes the proof. ■

Second, whenever $\mathcal{SL}_j(f(\tilde{\theta}), \tilde{\theta}_i) \cap \mathcal{SU}_j(f(\tilde{\theta}), \theta_i) \neq \emptyset$, we show that it is strictly better for agent j to challenge.

²³As will be clear in the argument later, the outcome function and transfer rule which we construct is a value function of some optimization problem. Such value functions are not necessarily continuous when Θ is not compact (which is the case when we apply Theorem 4 to prove Theorem 5 later).

Lemma 6 Let (m_i, m_j) be a message profile of agents i and j with $m_i^2 = m_j^2 = \tilde{\theta}$ with $\tilde{\theta} \in \Theta$ and $\mathcal{SL}_j(f(\tilde{\theta}), \tilde{\theta}_j) \cap \mathcal{SU}_j(f(\tilde{\theta}), m_j^1) \neq \emptyset$. Then, we can choose $m_j^3 \in \bar{X}$ such that $u_j(C_{i,j}(m_i, m_j), m_j^1) > u_j(f(m_i^2), m_j^1)$.

Proof. First, we fix an arbitrary $x \in \mathcal{SL}_j(f(\tilde{\theta}), \tilde{\theta}_j) \cap \mathcal{SU}_j(f(\tilde{\theta}), m_j^1)$. Let

$$m_j^3 = \alpha x \oplus (1 - \alpha) f(\tilde{\theta}),$$

where $\alpha \in (0, 1)$. Note that $m_j^3 \in \mathcal{SL}_j(f(\tilde{\theta}), \tilde{\theta}_j) \cap \mathcal{SU}_j(f(\tilde{\theta}), m_j^1)$ for every $\alpha \in (0, 1)$. As $\alpha \rightarrow 0$, we have

$$\rho(m_j^3, f(\tilde{\theta})) \rightarrow 0 \text{ and } u_j(m_j^3, m_j^1) \rightarrow u_j(f(\tilde{\theta}), m_j^1).$$

Observe that $\rho(m_j^3, f(\tilde{\theta})) = \alpha \|x - f(\tilde{\theta})\|$ (recall that \bar{X} is a compact subset of $\mathbb{R}^{I+|A|}$). Hence, we can choose $\alpha > 0$ small enough such that

$$\rho(m_j^3, f(\tilde{\theta})) < 1; \tag{28}$$

$$\rho(m_j^3, f(\tilde{\theta}))^3 (-3\eta) + \alpha \rho(m_j^3, f(\tilde{\theta})) (u_j(x, m_j^1) - u_j(f(\tilde{\theta}), m_j^1)) > 0. \tag{29}$$

Now, we have

$$\begin{aligned} & u_j(C_{i,j}(m_i, m_j), m_j^1) - u_j(f(\tilde{\theta}), m_j^1) \\ &= e_{i,j}(m_i, m_j) u_j\left(\frac{1}{2} \sum_{k=i,j} y_k(m_k^1), m_j^1\right) + (1 - e_{i,j}(m_i, m_j)) u_j(x(m_i^2, m_j^3), m_j^1) - u_j(f(\tilde{\theta}), m_j^1) \\ &= \rho(m_j^3, f(\tilde{\theta}))^3 u_j\left(\frac{1}{2} \sum_{k=i,j} y_k(m_k^1), m_j^1\right) + (1 - \rho(m_j^3, f(\tilde{\theta}))^3) u_j(\ell(m_j^3, f(\tilde{\theta})), m_j^1) - u_j(f(\tilde{\theta}), m_j^1) \\ &= \rho(m_j^3, f(\tilde{\theta}))^3 \left[u_j\left(\frac{1}{2} \sum_{k=i,j} y_k(m_k^1), m_j^1\right) - u_j(f(\tilde{\theta}), m_j^1) \right] \\ &\quad + (1 - \rho(m_j^3, f(\tilde{\theta}))^3) \left[u_j\left(\frac{\rho(m_j^3, f(\tilde{\theta}))}{1 + \rho(m_j^3, f(\tilde{\theta}))} m_j^3 \oplus \frac{1}{1 + \rho(m_j^3, f(\tilde{\theta}))} f(\tilde{\theta}), m_j^1\right) - u_j(f(\tilde{\theta}), m_j^1) \right] \\ &> \rho(m_j^3, f(\tilde{\theta}))^3 (-3\eta) + (1 - \rho(m_j^3, f(\tilde{\theta}))^3) \left[\frac{\rho(m_j^3, f(\tilde{\theta}))}{1 + \rho(m_j^3, f(\tilde{\theta}))} (u_j(m_j^3, m_j^1) - u_j(f(\tilde{\theta}), m_j^1)) \right] \\ &= \rho(m_j^3, f(\tilde{\theta}))^3 (-3\eta) + (1 - \rho(m_j^3, f(\tilde{\theta}))^3) \left[\frac{\alpha \rho(m_j^3, f(\tilde{\theta}))}{1 + \rho(m_j^3, f(\tilde{\theta}))} (u_j(x, m_j^1) - u_j(f(\tilde{\theta}), m_j^1)) \right] \\ &> \rho(m_j^3, f(\tilde{\theta}))^3 (-3\eta) + \frac{1}{2} \alpha \rho(m_j^3, f(\tilde{\theta})) (u_j(x, m_j^1) - u_j(f(\tilde{\theta}), m_j^1)) \end{aligned}$$

where the second equality follows because $m_i^2 = m_j^2 = \tilde{\theta}$ with $\tilde{\theta} \in \Theta$ and $\rho(m_j^3, f(\tilde{\theta}))^3 < 1$ (by (28)); the third equality follows from the definition of $\ell(m_j^3, f(\tilde{\theta}))$; the fourth equality follows from the linearity of u_j in allocation; the first inequality follows because

$$u_j \left(\frac{1}{2} \sum_{k=i,j} y_k(m_k^1), m_j^1 \right) - u_j(f(\tilde{\theta}), m_j^1) > -3\eta;$$

the last inequality follows from (28). Hence, it follows from (29) that

$$u_j(C_{i,j}(m_i, m_j), m_j^1) - u_j(f(\tilde{\theta}), m_j^1) > 0.$$

This completes the proof. ■

A.7.2 Existence of Good Equilibrium

Consider an arbitrary true state θ . The proof consists of two parts. In the first part, we argue that truth-telling m where $m_i = (\theta_i, \theta, x)$ for each $i \in \mathcal{I}$ constitutes a pure-strategy equilibrium, where $x(\theta, x) = f(\theta)$. Under this message profile m , $e_{i,j}(m_i, m_j) = 0$. Firstly, reporting \tilde{m}_i with either $\tilde{m}_{i,i}^2 \neq \theta_i$ or $\tilde{m}_{i,j}^2 \neq \theta_j$ suffers the penalty of $\tau_{i,j}^2(m)$ or $\tau_{i,j}^1(m)$ and hence cannot be a profitable deviation by Claim 15. Secondly, reporting \tilde{m}_i with $\tilde{m}_i^2 = \theta$ and $\tilde{m}_i^3 = x' \neq x$ either leads to $x(\theta, x') = f(\theta)$ and results in no change in payoff or $x(\theta, x') \neq f(\theta)$ which is strictly worse than $f(\theta)$. By Lemma 5, this is not a profitable deviation. Finally, reporting \tilde{m}_i with $\tilde{m}_i^2 = \theta$, $\tilde{m}_i^3 = \theta_i$, and $\tilde{m}_i^1 \neq \theta_i$ does not affect the allocation or transfer, since we still have $\tau_i(m) = 0$ and $e_{j,k}(m_j, m_k) = 0$ for every j and k .

In the second part, we show that for every Nash equilibrium σ of the game $\Gamma(\mathcal{M}, \theta)$ and every $m \in \text{supp}(\sigma)$, we have that $g(m) = f(\theta)$ and $\tau_i(m) = 0$ for every $i \in \mathcal{I}$. The proof of the second part is divided into three steps: (Step 1) *contagion of truth*: if agent j announces his type truthfully in his first report, then every agent must also report agent j 's type truthfully in their second report; (Step 2) *consistency*: every agent reports the same state $\tilde{\theta}$ in the second report; and (Step 3) *no challenge*: no agent challenges the common reported state $\tilde{\theta}$, i.e., $x(\tilde{\theta}, m_j^3) = f(\tilde{\theta})$ for every $j \in \mathcal{I}$. Then, consistency implies that $\tau_i(m) = 0$ for every $i \in \mathcal{I}$, whereas no challenge is invoked so that monotonicity of f together with Lemma 6 implies that $g(m) = f(\tilde{\theta}) = f(\theta)$.

As we do not make use of the notion of best challenge scheme in the infinite setting, the

proof of Claim 1 is more straightforward. Here m_i^1 only affects agent i 's own payoff through controlling the dictator lottery y_i ; in particular, both $e_{i,j}(m_i, m_j)$ and $e_{j,i}(m_j, m_i)$ are not determined by m_i^1 or m_j^1 . Hence, if $e_{i,j}(m_i, m_j) = \varepsilon$ or $e_{j,i}(m_j, m_i) = \varepsilon$, then $m_i^1 = \theta_i$ by (22). We now establish Steps 1-3.

A.7.3 Contagion of Truth

Claim 15 *We establish two results:*

- (a) *If agent j sends a truthful first report with σ_j -probability one, then every agent $i \neq j$ must report agent j 's type truthfully in his second report with σ_i -probability one.*
- (b) *If every agent $i \neq j$ reports a type $\tilde{\theta}_j$ of agent j in his second report with σ_i -probability one, then agent j must also report $\tilde{\theta}_j$ in his second report with σ_j -probability one.*

Proof. First, we prove part (a). Suppose instead that there exists some message m_i played with σ_i -positive probability such that agent i misreports agent j 's type in his second report, i.e., $m_{i,j}^2 \neq \theta_j$. Let \tilde{m}_i be a message which differs from m_i only in reporting j 's type truthfully $\tilde{m}_{i,j}^2 = \theta_j$. Then, for each m_{-i} played with σ_{-i} -positive probability, since agent j reports his type truthfully ($m_j^1 = \theta_j$), we have $\tilde{m}_i = \tilde{m}_i^m$ where $m = (m_i, m_{-i})$. Recall the definition of \tilde{m}_i^m in Section A.7.1.2. Hence, we reach a contradiction if we show \tilde{m}_i is a profitable deviation, i.e., for each m_{-i} played with σ_{-i} -positive probability, we have

$$u_i(g(\tilde{m}_i^m, m_{-i}), \theta_i) + \tau_i(\tilde{m}_i^m, m_{-i}) > u_i(g(m), \theta_i) + \tau_i(m).$$

First, observe that

$$\begin{aligned} \tau_i(\tilde{m}_i^m, m_{-i}) - \tau_i(m) &= \tau_{i,j}^1(\tilde{m}_i^m, m_{-i}) - \tau_{i,j}^1(m) \\ &= \sup_{\theta'_i} |u_i(g(m), \theta'_i) - u_i(g(\tilde{m}_i^m, m_{-i}), \theta'_i)| + d_j(m_{i,j}^2, m_j^1) \end{aligned}$$

where the first equality follows because \tilde{m}_i^m only differs from m_i in agent i 's second report

of agent j 's type. Thus we have

$$\begin{aligned}
& [u_i(g(\tilde{m}_i^m, m_{-i}), \theta_i) + \tau_i(\tilde{m}_i^m, m_{-i})] - [u_i(g(m), \theta_i) + \tau_i(m)] \\
= & u_i(g(\tilde{m}_i^m, m_{-i}), \theta_i) - u_i(g(m), \theta_i) \\
& + \sup_{\theta'_i} |u_i(g(m), \theta'_i) - u_i(g(\tilde{m}_i^m, m_{-i}), \theta'_i)| + d_j(m_{i,j}^2, m_j^1) \\
> & 0.
\end{aligned}$$

where the last inequality follows because $m_{i,j}^2 \neq \theta_j = m_j^1$.

Second, we prove part (b). Suppose, on the contrary, that there exists some message m_j played with σ_j -positive probability such that agent j misreports agent i 's type in his second report, i.e., $m_{j,j}^2 \neq \tilde{\theta}_j$. Let \bar{m}_j be a message that is identical to m_j except that $\bar{m}_{j,j}^2 = \tilde{\theta}_j$. Then, for each m_{-j} played with σ_{-j} -positive probability, since every agent $i \neq j$ reports $\tilde{\theta}_j$ in agent i 's second report, we have $\bar{m}_j = \bar{m}_j^m$ where $m = (m_j, m_{-j})$. Recall the definition of \bar{m}_i^m in Section A.7.1.2. Hence, we reach a contradiction if we show that \bar{m}_j is a profitable deviation, i.e., for each m_{-j} played with σ_{-j} -positive probability, we have

$$u_j(g(\bar{m}_j^m, m_{-j}), \theta_j) + \tau_j(\bar{m}_j^m, m_{-j}) > u_j(g(m), \theta_j) + \tau_j(m).$$

Notice that $\tau_{j,i}^1(\bar{m}_j^m, m_{-j}) = \tau_{j,i}^1(m)$.

$$\begin{aligned}
\tau_j(\bar{m}_j^m, m_{-j}) - \tau_j(m) &= \sum_{i \neq j} \{ \tau_{j,i}^2(\bar{m}_j^m, m_{-j}) - \tau_{j,i}^2(m) \} \\
&= \sum_{i \neq j} \left\{ \sup_{\theta'_j} |u_j(g(m), \theta'_j) - u_j(g(\bar{m}_j^m, m_{-j}), \theta'_j)| \right. \\
&\quad \left. + d_j(m_{j,j}^2, m_{i,j}^2) \right\}.
\end{aligned}$$

Thus, we have

$$\begin{aligned}
& [u_j(g(\bar{m}_j^m, m_{-j}), \theta_j) + \tau_j(\bar{m}_j^m, m_{-j})] - [u_j(g(m), \theta_j) + \tau_j(m)] \\
= & u_j(g(\bar{m}_j^m, m_{-j}), \theta_j) - u_j(g(m), \theta_j) \\
& + \sum_{i \neq j} \left\{ \sup_{\theta'_j} |u_j(g(m), \theta'_j) - u_j(g(\bar{m}_j^m, m_{-j}), \theta'_j)| \right. \\
& \quad \left. + d_j(m_{j,j}^2, m_{i,j}^2) \right\} \\
> & 0.
\end{aligned}$$

where the last inequality follows because $m_{j,j}^2 \neq \tilde{\theta}_j = m_{i,j}^2$. This completes the proof. ■

A.7.4 Consistency

The argument for consistency follows verbatim the proof of Claim 3 in the proof of Theorem 1.

A.7.5 No Challenge

Claim 16 *No agent challenges with positive probability the common state $\tilde{\theta}$ announced in the second report.*

Proof. Provided that a consistent report of $\tilde{\theta}$ is achieved, it suffices to show that $\mathcal{SL}_j(f(\tilde{\theta}), \tilde{\theta}_j) \cap \mathcal{SU}_j(f(\tilde{\theta}), \theta_j) = \emptyset$ for every agent $j \in \mathcal{I}$. Suppose to the contrary that $\mathcal{SL}_j(f(\tilde{\theta}), \tilde{\theta}_j) \cap \mathcal{SU}_j(f(\tilde{\theta}), \theta_j) \neq \emptyset$ for some agent j . Then, we first show that $x(\tilde{\theta}, m_j^3) \neq f(\tilde{\theta})$ for every $m_j \in \text{supp}(\sigma_j)$. By Lemma 6, there exists $x \in \mathcal{SL}_j(f(\tilde{\theta}), \tilde{\theta}_j) \cap \mathcal{SU}_j(f(\tilde{\theta}), \theta_j)$. If $x(\tilde{\theta}, m_j^3) = f(\tilde{\theta})$, then $\tilde{m}_j = (m_j^1, m_j^2, x)$ is a strictly profitable deviation from announcing m_j . This deviation results in a better allocation $x(\tilde{\theta}, x) \in \mathcal{SU}_j(f(\tilde{\theta}), \theta_j)$. Hence, we have $x(\tilde{\theta}, m_j^3) \neq f(\tilde{\theta})$ for every $m_j \in \text{supp}(\sigma_j)$. It follows that the dictator lottery is triggered with positive probability. Thus, by (22), each agent i has a strict incentive to announce the true type in his first report (i.e., $m_i^1 = \theta_i$) with σ_i -probability one. By Claim 15, we conclude that $\tilde{\theta} = \theta$ and hence $\mathcal{SL}_j(f(\theta), \theta_j) \cap \mathcal{SU}_j(f(\theta), \theta_j) \neq \emptyset$, which is impossible. ■

A.8 Proof of Theorem 5

First, we obtain a stronger version of Lemma 2, namely, there is a set of dictator lotteries which can be used to elicit the agents' true types, regardless of their cardinal representation. This follows from the same proof of Lemma in Abreu and Matsushima (1992); moreover, the dictator lottery constructed remains valid (in the sense of (30)) as long as the preferences exhibit monotonicity with respect to first-order stochastic dominance. Recall that for each $x = (\ell, (t_i)_{i \in \mathcal{I}}) \in X$, $u_i(x, \theta_i) \equiv v_i(\ell, \theta) + t_i$.

Lemma 7 *For each $i \in \mathcal{I}$, there exists a function $y_i : \Theta_i \rightarrow X$ such that for every $\theta_i, \theta'_i \in \Theta_i$ with $\theta_i \neq \theta'_i$ and every cardinal representation $v_i(\cdot)$ of $(\succeq_i^\theta)_{\theta \in \Theta}$,*

$$u_i(y_i(\theta_i), \theta_i) > u_i(y_i(\theta'_i), \theta_i); \tag{30}$$

moreover, for each type θ'_j of agent $j \in \mathcal{I}$, we also have that for every $x \in f(\Theta)$

$$u_i(y_j(\theta'_j), \theta_i) < u_i(x, \theta_i). \quad (31)$$

To satisfy condition (31), we simply add a penalty of η to each outcome of the lotteries $\{y'_i(\theta_i)\}$ where η is chosen in the same fashion as in (4) given each cardinal representation $v_i(\cdot)$. Moreover, since each $v_i(\cdot)$ takes values in $[0, 1]$, we also save the notation and take η to be independent of $v_i(\cdot)$.

We first introduce the following definitions of contour sets under ordinal preferences. For $(a, \theta_i) \in A \times \Theta_i$, under ordinal preference \succeq_i^θ , we write the upper-contour set, the lower-contour set, the strict upper-contour set, and the strict lower-contour set as follows:

$$\begin{aligned} U_i(a, \theta_i) &= \{a' \in A : a' \succeq_i^\theta a\}; \\ L_i(a, \theta_i) &= \{a' \in A : a \succeq_i^\theta a'\}; \\ SU_i(a, \theta_i) &= \{a' \in A : a' \succ_i^\theta a\}; \\ SL_i(a, \theta_i) &= \{a' \in A : a \succ_i^\theta a'\}; \end{aligned}$$

where \succ_i^θ denotes the strict preference induced by \succeq_i^θ . We now introduce the notion of ordinal almost monotonicity proposed by [Sanver \(2006\)](#) as the key condition for [Theorem 5](#).

Definition 9 *An SCF f satisfies **ordinal almost monotonicity** if, for every pair of states θ and $\tilde{\theta}$, with $f(\tilde{\theta}) \neq f(\theta)$, there is some agent $i \in \mathcal{I}$ such that either*

$$L_i(f(\tilde{\theta}), \tilde{\theta}_i) \cap SU_i(f(\tilde{\theta}), \theta_i) \neq \emptyset,$$

or

$$SL_i(f(\tilde{\theta}), \tilde{\theta}_i) \cap U_i(f(\tilde{\theta}), \theta_i) \neq \emptyset.$$

A.8.1 Proof of the Only-If Part

In the proof, we make use of Claim C from [Mezzetti and Renou \(2012a\)](#) which is reproduced as follows:

Claim 17 *Suppose that $L_i(f(\theta), \theta_i) \subseteq L_i(f(\theta), \tilde{\theta}_i)$ and $SL_i(f(\theta), \theta_i) \subseteq SL_i(f(\theta), \tilde{\theta}_i)$. Then, given every cardinal representation $v_i(\cdot, \theta)$ of \succ_i^θ , there exists a cardinal representation*

$v_i(\cdot, \tilde{\theta})$ of $\succsim_i^{\tilde{\theta}}$ such that $v_i(a, \tilde{\theta}) \leq v_i(a, \theta)$ for all $a \in A$ and $v_i(f(\theta), \tilde{\theta}) = v_i(f(\theta), \theta)$

Suppose f is ordinally Nash implementable in a mechanism $\mathcal{M} = ((M_i), g, (\tau_i))_{i \in \mathcal{I}}$ but not almost monotonic. That is, there exists $\theta, \tilde{\theta} \in \Theta$ such that for each agent $i \in \mathcal{I}$, we have $L_i(f(\theta), \theta_i) \subseteq L_i(f(\theta), \tilde{\theta}_i)$ and $SL_i(f(\theta), \theta_i) \subseteq SL_i(f(\theta), \tilde{\theta}_i)$, but $f(\theta) \neq f(\tilde{\theta})$. Since \mathcal{M} ordinally Nash implements f , we know that for every cardinal representation $(v_i)_{i \in \mathcal{I}}$, there exists a pure-strategy Nash equilibrium m^* in the game $\Gamma(\mathcal{M}, \theta, v)$ such that $g(m^*) = f(\theta)$. Since $f(\theta) \neq f(\tilde{\theta})$, the message profile m^* cannot be a Nash equilibrium at state $\tilde{\theta}$ for every cardinal representation v_i . Then, there exist agent i and message $m_i \in M_i$ such that

$$\begin{aligned} v_i(g(m_i^*, m_{-i}^*), \theta) + \tau_i(m_i^*, m_{-i}^*) &\geq v_i(g(m_i, m_{-i}^*), \theta) + \tau_i(m_i, m_{-i}^*); \\ v_i(g(m_i^*, m_{-i}^*), \tilde{\theta}) + \tau_i(m_i^*, m_{-i}^*) &< v_i(g(m_i, m_{-i}^*), \tilde{\theta}) + \tau_i(m_i, m_{-i}^*). \end{aligned}$$

Summing up the two inequalities, we obtain that for every cardinal representation v_i

$$v_i(g(m_i^*, m_{-i}^*), \theta) - v_i(g(m_i^*, m_{-i}^*), \tilde{\theta}) > v_i(g(m_i, m_{-i}^*), \theta) - v_i(g(m_i, m_{-i}^*), \tilde{\theta}). \quad (32)$$

Note that $g(m_i^*, m_{-i}^*) = f(\theta)$. By Claim 17, however, we can construct a cardinal utility representation $v_i(\cdot, \tilde{\theta})$ of $\succsim_i^{\tilde{\theta}}$ such that $v_i(a, \tilde{\theta}) \leq v_i(a, \theta)$ for all $a \in A$ and $v_i(f(\theta), \tilde{\theta}) = v_i(f(\theta), \theta)$. Therefore, the left-hand side of (32) is zero, while the right-hand side is non-negative. This is a contradiction.

A.8.2 Proof of the If Part

Let f be an SCF which is ordinally almost monotonic on Θ . Recall that V_i^θ denotes the set of all cardinal representations $v_i(\cdot, \theta)$ of \succeq_i^θ . Define $V^\theta = \times_{i \in \mathcal{I}} V_i^\theta$ with a generic element v^θ . By no redundancy of Θ , $\theta \neq \theta'$ implies that $\succeq_i^\theta \neq \succeq_i^{\theta'}$ for some agent $i \in \mathcal{I}$. Hence, $\{V^\theta : \theta \in \Theta\}$ forms a partition of $\Theta^* \equiv \bigcup_{\theta \in \Theta} V^\theta$ which is the set of all cardinal utility profiles of agent i induced by Θ . Observe that Θ^* is a Polish space.²⁴ For the sake of notational simplicity,

²⁴Since any product or disjoint union of a countable family of Polish spaces remains a Polish space (see Proposition A.1(b) in p. 550 of Treves (2016)), it suffices to argue that V_i^θ is a Polish space. Let $V = [0, 1]^{|A|}$ be the set of possible cardinalizations. We may write $V_i^\theta = \bigcap_{a \in A} V_{i,a}^\theta$ where for each $a \in A$, we set

$$V_{i,a}^\theta \equiv \bigcap_{\{b \in A : a \succ_i^\theta b\}} \{v \in V : v(a) > v(b)\} \bigcap \bigcap_{\{b \in A : a \sim_i^\theta b\}} \{v \in V : v(a) = v(b)\}.$$

we write θ_i^* as a generic element in Θ_i^* and $\theta^* = (\theta_i^*)_{i \in \mathcal{I}}$. Let $f^* : \Theta^* \rightarrow A$ be the SCF on Θ^* induced by $f : \Theta \rightarrow A$ such that $f^*(\theta^*) = f(\theta)$ if and only if $\theta^* \in V^\theta$.

We prove the if-part by establishing two claims: First, we show that f^* is Maskin-monotonic in Claim 18. Hence, Theorem 4 implies that f^* is implementable in Nash equilibria on Θ^* . Second, it follows from Claim 19 that f is ordinally Nash implementable on Θ .

Claim 18 *If f is ordinally almost monotonic on Θ , then f^* is strictly Maskin-monotonic on Θ^* .*

Proof. Let θ^* and $\tilde{\theta}^*$ be in Θ^* such that $f^*(\theta^*) \neq f^*(\tilde{\theta}^*)$. Since $f^*(\theta^*) = f(\theta)$ if and only if $\theta^* \in V^\theta$, we must have two states θ and $\tilde{\theta} \in \Theta$ such that $\theta^* \in V^\theta$ and $\tilde{\theta}^* \in V^{\tilde{\theta}}$, and $f(\theta) \neq f(\tilde{\theta})$. Since f satisfies ordinal almost monotonicity, there exists agent $i \in \mathcal{I}$ and outcomes $a, a' \in A$ such that either $a \in L_i(f(\tilde{\theta}), \tilde{\theta}_i) \cap SU_i(f(\tilde{\theta}), \theta_i)$ or $a' \in SL_i(f(\tilde{\theta}), \tilde{\theta}_i) \cap U_i(f(\tilde{\theta}), \theta_i)$. Then, either choose $t_i < 0$ such that $(a, (t_i, \mathbf{0})) \in \mathcal{SL}_i(f^*(\tilde{\theta}^*), \tilde{\theta}_i^*) \cap \mathcal{SU}_i(f^*(\tilde{\theta}^*), \theta_i^*)$ or $t_i > 0$ such that $(a', (t_i, \mathbf{0})) \in \mathcal{SL}_i(f^*(\tilde{\theta}^*), \tilde{\theta}_i^*) \cap \mathcal{SU}_i(f^*(\tilde{\theta}^*), \theta_i^*)$ where $\mathbf{0} \in \mathbb{R}^{I-1}$ means that every agent $j \neq i$ incurs no transfer. Hence, f^* is strictly Maskin-monotonic on Θ^* . ■

Claim 19 *If f^* is implementable in Nash equilibria, then f is ordinally Nash implementable.*

Proof. Suppose the SCF f^* is implementable in Nash equilibria on Θ^* . Then, there exists a mechanism $\mathcal{M} = ((M_i), g, (\tau_i))_{i \in \mathcal{I}}$ such that for every state $\theta^* \in \Theta^*$ and message profile $m \in M$, (i) there exists a pure-strategy Nash equilibrium in the game $\Gamma(\mathcal{M}, \theta^*)$; and (ii) $m \in \text{supp}(NE(\Gamma(\mathcal{M}, \theta^*))) \Rightarrow g(m) = f^*(\theta^*)$ and $\tau_i(m) = 0$ for every $i \in \mathcal{I}$. Thus, for every state $\theta^* \in V^\theta$, we must have (i) there exists a pure-strategy Nash equilibrium in the game $\Gamma(\mathcal{M}, \theta, v^{\theta^*})$; and (ii) $m \in \text{supp}(NE(\Gamma(\mathcal{M}, \theta, v^{\theta^*}))) \Rightarrow g(m) = f(\theta)$ and $\tau_i(m) = 0$ for every $i \in \mathcal{I}$. Hence, f is ordinally Nash implementable on Θ . ■

It follows that V_i^θ is a finite intersection of open subsets and closed subsets of the Polish space V and hence remains a Polish space (see Proposition A.1(a)(c)(e) in p. 550 of Treves (2016)).

References

- ABREU, D., AND H. MATSUSHIMA (1992): “Virtual Implementation in Iteratively Undominated Strategies: Complete Information,” *Econometrica*, 60, 993–1008.
- (1994): “Exact Implementation,” *Journal of Economic Theory*, 64, 1–19.
- AGHION, P., D. FUDENBERG, R. HOLDEN, T. KUNIMOTO, AND O. TERCIEUX (2012): “Subgame-Perfect Implementation under Information Perturbations,” *The Quarterly Journal of Economics*, 127(4), 1843–1881.
- CHUNG, K.-S., AND J. C. ELY (2003): “Implementation with Near-Complete Information,” *Econometrica*, 71(3), 857–871.
- DASGUPTA, P., AND E. MASKIN (2000): “Efficient Auctions,” *The Quarterly Journal of Economics*, 115(2), 341–388.
- D’ASPROMONT, C., J. CRÉMER, AND L.-A. GÉRARD-VARET (2003): “Correlation, Independence, and Bayesian Incentives,” *Social Choice and Welfare*, 21(2), 281–310.
- DUTTA, B., AND A. SEN (1991): “A Necessary and Sufficient Condition for Two-Person Nash implementation,” *The Review of Economic Studies*, 58(1), 121–128.
- HARSANYI, J. C. (1973): “Games with Randomly Disturbed Payoffs: A New Rationale for Mixed-Strategy Equilibrium Points,” *International Journal of Game Theory*, 2(1), 1–23.
- JACKSON, M. O. (1992): “Implementation in Undominated Strategies: A Look at Bounded Mechanisms,” *The Review of Economic Studies*, 59(4), 757–775.
- JACKSON, M. O., T. R. PALFREY, AND S. SRIVASTAVA (1994): “Undominated Nash Implementation in Bounded Mechanisms,” *Games and Economic Behavior*, 6(3), 474–501.
- KUNIMOTO, T. (2019): “Mixed Bayesian Implementation in General Environments,” *Journal of Mathematical Economics*, 82, 247–263.
- LEE, J., AND H. SABOURIAN (2011): “Efficient Repeated Implementation,” *Econometrica*, 79(6), 1967–1994.

- MASKIN, E. (1999): “Nash Equilibrium and Welfare Optimality,” *Review of Economic Studies*, 66, 23–38.
- MEZZETTI, C., AND L. RENO (2012a): “Implementation in Mixed Nash Equilibrium,” *Journal of Economic Theory*, 147(6), 2357–2375.
- (2012b): “Mixed Nash Implementation with Finite Mechanisms,” *mimeo*.
- (2017): “Repeated Nash Implementation,” *Theoretical Economics*, 12(1), 249–285.
- MOOKHERJEE, D., AND S. REICHELSTEIN (1990): “Implementation via Augmented Revelation Mechanisms,” *The Review of Economic Studies*, 57(3), 453–475.
- MOORE, J., AND R. REPULLO (1988): “Subgame Perfect Implementation,” *Econometrica*, 56, 1191–1220.
- OLLÁR, M., AND A. PENTA (2017): “Full Implementation and Belief Restrictions,” *American Economic Review*, 107(8), 2243–77.
- (2021): “A network solution to robust implementation: The case of identical but unknown distributions,” .
- OSBORNE, M. J., AND A. RUBINSTEIN (1994): *A Course in Game Theory*. MIT press.
- PERRY, M., AND P. J. RENY (2002): “An Efficient Auction,” *Econometrica*, 70(3), 1199–1212.
- SANVER, M. R. (2006): “Nash Implementing Non-Monotonic Social Choice Rules by Awards,” *Economic Theory*, 28(2), 453–460.
- SERRANO, R., AND R. VOHRA (2010): “Multiplicity of Mixed Equilibria in Mechanisms: A Unified Approach to Exact and Approximate Implementation,” *Journal of Mathematical Economics*, 46(5), 775–785.
- SJÖSTROM, T. (1994): “Implementation in Undominated Nash Equilibria without Integer Games,” *Games and Economic Behavior*, 6(3), 502–511.
- STINCHCOMBE, M. B., AND H. WHITE (1992): “Some Measurability Results for Extrema of Random Functions over Random Sets,” *The Review of Economic Studies*, 59(3), 495–514.

TREVES, F. (2016): *Topological Vector Spaces, Distributions and Kernels: Pure and Applied Mathematics*, vol. 25. Elsevier.