

PSEUDO-ANOSOV ARE EXPONENTIALLY GENERIC IN MAPPING CLASS GROUPS

INHYEOK CHOI

ABSTRACT. Given a finite generating set S , let us endow the mapping class group of a closed hyperbolic surface with the word metric for S . We discuss the following question: does the proportion of non-pseudo-Anosov mapping classes in the ball of radius R decrease to 0 as R increases? We show that any finite subset S' of the mapping class group is contained in a finite generating set S such that this proportion decreases exponentially. Our strategy applies to weakly hyperbolic groups and does not refer to the automatic structure of the group.

Keywords. Mapping class group, pseudo-Anosov map, random walk

MSC classes: 20F67, 30F60, 57K20, 57M60, 60G50

1. INTRODUCTION

Let Σ be a closed hyperbolic surface. We denote by $\text{Mod}(\Sigma)$, $\mathcal{T}(\Sigma)$ and $\mathcal{C}(\Sigma)$ the mapping class group, the Teichmüller space and the curve complex of Σ , respectively. When X is a Gromov hyperbolic space or $\mathcal{T}(\Sigma)$ and $g \in \text{Isom}(X)$, we denote by $\tau_X(g)$ the (asymptotic) translation length of g . For a group G generated by a finite set S , we denote by $B_S(n)$ the ball of radius n with respect to the word metric for S . We also denote by $\partial B_S(n)$ the corresponding sphere of radius n . Our main result is as follows.

Theorem A (Translation length grows linearly). *Let X be either a Gromov hyperbolic space or $\mathcal{T}(\Sigma)$. Let also G be a finitely generated non-elementary subgroup of $\text{Isom}(X)$ and $S' \subseteq G$ be a finite subset. Then there exist $L, K > 0$ and a finite generating set $S \supseteq S'$ of G such that*

$$\frac{\#\{g \in B_S(n) : \tau_X(g) \leq Ln\}}{\#B_S(n)} \leq Ke^{-n/K}$$

holds for each n .

Non-pseudo-Anosov mapping classes have translation length zero in $\mathcal{C}(\Sigma)$. As a result, we affirmatively answer the following version of a folklore conjecture, at least for infinitely many generating sets S .

Corollary 1.1 (Genericity of pAs, cf. [Far06, Conjecture 3.15]). *Let G be a finitely generated non-elementary subgroup of $\text{Mod}(\Sigma)$. Then there exists a finite generating set $S \subseteq G$ such that the proportion of non-pseudo-Anosov mapping classes in the ball $B_S(n)$ decays exponentially as $n \rightarrow \infty$.*

Date: November 3, 2021.

Note that $\text{Mod}(\Sigma)$ can act on both $\mathcal{T}(\Sigma)$ and $\mathcal{C}(\Sigma)$. Comparing the translation lengths of mapping classes on these two spaces is an interesting question. Thanks to the linear growth in Theorem A, we can deduce:

Corollary 1.2. *Let G be a finitely generated non-elementary subgroup of $\text{Mod}(\Sigma)$ and $S' \subseteq G$ be a finite subset. Then there exist $L, K > 0$ and a finite generating set $S \supseteq S'$ of G such that the following holds. For*

$$P := \left\{ g \in G : \frac{1}{L} \leq \frac{\tau_{\mathcal{T}(\Sigma)}(g)}{\tau_{\mathcal{C}(\Sigma)}(g)} \leq L \right\},$$

and each n , we have

$$\frac{\#B_S(n) \cap P^c}{\#B_S(n)} \leq K e^{-n/K}.$$

1.1. History and related problems. We remark that Theorem A may not be optimal for Gromov hyperbolic spaces. Gekhtman, Taylor and Tiozzo have proved the genericity of loxodromics in hyperbolic groups acting on separable Gromov hyperbolic spaces, in terms of the word metric for *any* finite generating set. This was generalized to relatively hyperbolic groups, RAAGs and RACGs with *particular* finite generating sets (see [GTT18], [GTT20b], [GTT20a]). Gekhtman-Taylor-Tiozzo's example [GTT20b, Example 1] shows that the genericity of loxodromics is not achieved for all weakly hyperbolic groups for all finite generating sets.

In the setting of relatively hyperbolic groups (and many more), Yang also established that loxodromics are exponentially generic when the action is proper and cocompact [Yan20]. Hence, at least for hyperbolic groups acting on Gromov hyperbolic spaces properly and cocompactly, Theorem A is weaker than previous results in the sense that the finite generating set cannot be arbitrary. It is however stronger in the sense that

- (1) it does not require the action of G to be proper and cocompact,
- (2) it discusses the exponential genericity with respect to a linearly growing threshold, not a static threshold.

In fact, combining our strategy with the theory of Gekhtman, Taylor and Tiozzo yields the following.

Proposition 1.3. *Let X, G be as in Theorem A and S be a finite generating set of G . Suppose moreover that G itself is a hyperbolic group. Then there exist $\lambda > 0$ such that the following hold. Below, ν denotes the Patterson-Sullivan measure with respect to S .*

- (1) *For any $x \in X$ and ν -a.e. $\eta \in \partial G$, if $(g_n)_{n \geq 0}$ is a geodesic in G converging to η , then*

$$\lim_{n \rightarrow \infty} \frac{d_X(x, g_n x)}{n} = \lambda.$$

(2) For any $\epsilon > 0$, there exists $K > 0$ such that

$$\frac{\#\{g \in \partial B_S(n) : \tau_X(g) \notin [\lambda - \epsilon, \lambda + \epsilon]\}}{\#\partial B_S(n)} \leq Ke^{-n/K}$$

holds for each n .

Note that this implies the following. For any hyperbolic subgroup G of $\text{Mod}(\Sigma)$ and any finite generating set S of G , let $\lambda_{\mathcal{T}}$, $\lambda_{\mathcal{C}}$ be the escape rate of G on $\mathcal{T}(\Sigma)$ and $\mathcal{C}(\Sigma)$, respectively, in terms of the Patterson-Sullivan measure for S . Then for any $\epsilon > 0$, there exists K such that

$$\frac{\#\{g \in B_S(n) : \frac{\lambda_{\mathcal{T}}}{\lambda_{\mathcal{C}}} - \epsilon \leq \frac{\tau_{\mathcal{T}(\Sigma)}(g)}{\tau_{\mathcal{C}(\Sigma)}(g)} \leq \frac{\lambda_{\mathcal{T}}}{\lambda_{\mathcal{C}}} + \epsilon\}}{\#B_S(n)} \leq Ke^{-n/K}$$

holds for each n . For the sake of completeness, we sketch the proof of Proposition 1.3 in Appendix B.

Meanwhile, Theorem A is new for $\text{Mod}(\Sigma)$. The progress so far was that the proportion of pseudo-Anosov elements in the word metric ball stays bounded away from zero [CW18]. See [Yan20] and [EST20] for counting pseudo-Anosovs in other viewpoints. We can further ask:

Question 1.4. *Are pseudo-Anosovs exponentially generic with respect to any finite generating set? For example, are they exponentially generic with respect to Humphries' generators? If not, are they generic at least?*

Question 1.5. *Does Proposition 1.3 hold for $G = \text{Mod}(\Sigma)$ and at least one S ?*

Question 1.5 is intimately related to the (geodesic) automaticity of $\text{Mod}(\Sigma)$.

Let us finally mention a problem investigated by I. Kapovich. Let μ be a discrete measure on a group G . We define the *non-backtracking random walk* generated by μ as follows. The first alphabet g_1 is chosen from G with the law of μ ; for each $n \geq 2$, g_n is chosen from $G \setminus \{g_{n-1}^{-1}\}$ with the law

$$\mathbb{P}(g_n = g) = \frac{1}{\mu(G \setminus \{g_{n-1}^{-1}\})} \mu(g).$$

In this setting, Gekhtman-Taylor-Tiozzo proved that $\mathbb{P}(\omega_n \text{ is loxodromic})$ tends to 1 as $n \rightarrow \infty$ [GTT20b, Theorem 2.8]. With an adequate modification, our argument yields the following.

Proposition 1.6. *Let X be as in Theorem A, and μ be a non-elementary discrete measure μ on $\text{Isom}(X)$. Consider the non-backtracking random walk ω generated by μ . Then there exists $L, K > 0$ such that*

$$\mathbb{P}\{\tau_X(\omega_n) \leq Ln\} \leq Ke^{-n/K}$$

holds for each n .

1.2. Strategy for Theorem A. One method for the counting problem is to utilize (geodesic) automatic structures of the group. Lacking such structures, we instead consider the random walk ω on G generated by the uniform measure on S . Then Gouëzel's and Baik-Choi-Kim's theories imply that the unwanted probability decays exponentially ([Gou21], [BCK21]). There are at least two more theories that provide this exponential decay. One is Maher's theory [Mah12] for random walks with bounded support, which led to Maher-Tiozzo's more general theory [MT18]. Another one is Boulanger-Mathieu-Sert-Sisto's theory [BMSS20] of large deviation principles for random walks with finite exponential moment.

Unavoidably, random walks cannot count lattice points in a one-to-one manner. If S is nicely populated by the self-convolution of a Schottky set S_0 , however, then we have a one-to-one correspondence between some portion of lattice points and (non-backtracking) paths of alphabets in S_0 . This leads to the estimate $\#B_S(e, n) \sim \#(\text{paths from the random walk}) \cdot r^n$ with $r \sim 1$. We arrive at the desired estimate by forcing the exponential decay of probability to be much faster than the decay of r^n .

We remark that the notions and statements in Section 2, 3 and 4 are mostly established in [Gou21], [BCK21] and [Cho21]. Nonetheless, to make the exposition self-contained, we present all details except the proofs of Fact 2.5, 2.6, 2.8, 2.9 and A.1. See [Cho21] for their proofs.

Acknowledgments. The author truly thanks Hyungryul Baik, Dongryul M. Kim, Jing Tao, Samuel Taylor and Wenyan Yang for helpful discussion.

The author was partially supported by Samsung Science & Technology Foundation grant No. SSTF-BA1702-01.

2. WITNESSING AND SCHOTTKY SETS

In this section, we gather relevant facts about Gromov hyperbolic spaces and Teichmüller spaces. All facts in this section are proved in [Cho21].

Given a metric space (X, d) and a triple $x, y, z \in X$, we define the *Gromov product* of y, z with respect to x by

$$(y, z)_x = \frac{1}{2}[d(x, y) + d(x, z) - d(y, z)].$$

Throughout the article, we will frequently use the following property: for $x, y, z \in X$ and $g \in \text{Isom}(X)$, we have

$$(gy, gz)_{gx} = (y, z)_x.$$

X is said to be *Gromov hyperbolic* if there exists a constant $\delta > 0$ such that every quadruple $x, y, z, w \in X$ satisfies the following inequality called the *Gromov inequality*:

$$(2.1) \quad (x, y)_w \geq \min\{(x, z)_w, (y, z)_w\} - \delta.$$

From now on, we permanently fix $\delta > 0$.

In $\mathcal{T}(\Sigma)$, we say that a surface $x \in \mathcal{T}(\Sigma)$ is ϵ -thin if it has a curve with the extremal length less than ϵ ; if not, it is said to be ϵ -thick. For $x, y \in \mathcal{T}(\Sigma)$, $[x, y]$ denotes the Teichmüller geodesic from x to y ; it is said to be ϵ -thick if it is composed of ϵ -thick points.

In δ -hyperbolic spaces, we regard every point as ϵ -thick for any $\epsilon > 0$. Here $[x, y]$ denotes the pair of points (x, y) , which is considered ϵ -thick for any $\epsilon > 0$. In either space, $[x, y]$ are called *segments* and their lengths are defined by $d(x, y)$.

Definition 2.1 (Witnessing in δ -hyperbolic spaces). *Let $x, y, \{x_i\}_{i=1}^n, \{y_i\}_{i=1}^n$ be points in a δ -hyperbolic space X , and $D > 0$. We say that $[x, y]$ is D -witnessed by $([x_1, y_1], \dots, [x_n, y_n])$ if:*

- (1) $(x_{i-1}, x_{i+1})_{x_i} < D$ for $i = 1, \dots, n$, where $x = x_0$ and $y = x_{n+1}$;
- (2) $(y_{i-1}, y_{i+1})_{y_i} < D$ for $i = 1, \dots, n$, where $x = y_0$ and $y = y_{n+1}$, and
- (3) $(y_{i-1}, y_i)_{x_i}, (x_i, x_{i+1})_{y_i} < D$ for $i = 1, \dots, n$.

Definition 2.2 (Witnessing in $\mathcal{T}(\Sigma)$). *Let $x, y, \{x_i\}_{i=1}^n, \{y_i\}_{i=1}^n$ be points in $X = \mathcal{T}(\Sigma)$, and $D > 0$. We say that $[x, y]$ is D -witnessed by $([x_1, y_1], \dots, [x_n, y_n])$ if the geodesic $[x, y]$ contains subsegments $[x'_i, y'_i]$ such that*

- (1) x'_{i+1} is not closer to x than y'_i for $i = 1, \dots, n - 1$, and
- (2) $[x_i, y_i]$ and $[x'_i, y'_i]$ D -fellow travel.

From now on, we permanently fix X , a δ -hyperbolic space or $\mathcal{T}(\Sigma)$.

Definition 2.3. *Let x, y and $\{x_i, y_i, z_i\}_{i=1}^N$ be points in X . We say that $[x, y]$ is D -marked with $([x_i, y_i])_{i=1}^N, ([z_i, x_i])_{i=1}^N$ if the following hold:*

- (1) $(y_i, z_i)_{x_i} < D$ for each $i = 1, \dots, N$;
- (2) $[x_{i-1}, x_i]$ is D -witnessed by $([x_{i-1}, y_{i-1}], [z_i, x_i])$ for each $i = 1, \dots, N + 1$, where we set $x_0 = y_0 = x$ and $x_{N+1} = z_{N+1} = y$;

In this case, we also say that $[x_1, x_n]$ is fully D -marked with $([x_i, y_i])_{i=1}^{N-1}, ([z_i, x_i])_{i=2}^N$.




FIGURE 1. Schematics for $[x, y]$ being D -marked with $([x_i, y_i])_{i=1}^4, ([z_i, x_i])_{i=1}^4$.

The following observation is immediate.

Lemma 2.4. *Let $\{x_i\}_{i=0}^N, \{y_{i-1}, z_i\}_{i=1}^N$ be points in X . Suppose that for each $i = 1, \dots, N$, $[x_{i-1}, x_i]$ is fully D -marked with sequences of segments $(\gamma_{i,j})_{j=1}^{n_{i-1}}, (\eta_{i,j})_{j=2}^{n_i}$, where $\gamma_{i,1} = [x_{i-1}, y_{i-1}]$ and $\eta_{i,n_i} = [z_i, x_i]$. Suppose also*

that $(y_i, z_i)_{x_i} \leq D$ for $i = 1, \dots, N-1$. Then $[x_0, x_N]$ is fully D -marked with

$$\begin{aligned} & (\gamma_{1,1}, \dots, \gamma_{1,n_1-1}, \gamma_{2,1}, \dots, \gamma_{2,n_2-1}, \dots, \gamma_{N,n_N-1}), \\ & (\eta_{1,2}, \dots, \eta_{1,n_1}, \eta_{2,2}, \dots, \eta_{2,n_2}, \dots, \eta_{N,n_N}). \end{aligned}$$

Our aim is to prove that if $[x, y]$ is D -marked with sufficiently thick and long segments, then $[x, y]$ is witnessed by those segments. In order to prove this, we need the following consequences of Rafi's fellow-travelling and thin triangle theorems [Raf14].

Fact 2.5 ([Cho21, Lemma 3.9]). *For each $D, \epsilon > 0$, there exist $E, L > D$ that satisfy the following property. Let x_1 be an ϵ -thick point and $\{x_i\}_{i=2}^3, \{y_i\}_{i=1}^5$ be points in X such that:*

- (1) $[y_1, y_2], [y_3, y_4], [y_4, y_5]$ are ϵ -thick and longer than L ,
- (2) $(y_3, y_5)_{y_4} \leq D$,
- (3) $[x_1, x_2]$ is D -witnessed by $([y_1, y_2], [y_3, y_4])$, and
- (4) $[x_2, x_3]$ is E -witnessed by $[y_4, y_5]$,

then $[x_1, x_3]$ is E -witnessed by $[y_1, y_2]$.

Fact 2.6 ([Cho21, Lemma 3.10]). *For each $E, \epsilon > 0$, there exists $F, L > E$ that satisfies the following property. Let $\{x_i\}_{i=1}^3, \{y_i\}_{i=1}^3$ be points in X such that:*

- (1) $[y_1, y_2], [y_2, y_3]$ are ϵ -thick and longer than L ,
- (2) $(y_1, y_3)_{y_2} \leq E$,
- (3) $[x_1, x_2]$ is E -witnessed by $[y_1, y_2]$, and
- (4) $[x_2, x_3]$ is E -witnessed by $[y_2, y_3]$,

then $[x_1, x_3]$ is F -witnessed by $[y_1, y_2]$ and $[y_2, y_3]$. In particular, $|(x_1, x_3)_{x_2} - d(x_2, y_2)| < F$.

Combining Fact 2.5 and 2.6 yields the following.

Corollary 2.7. *Let $D, \epsilon > 0$ and*

- $E = E(\epsilon, D)$, $L_1 = L(\epsilon, D)$ as in Fact 2.5, and
- $F = F(\epsilon, E)$, $L_2 = L(\epsilon, E)$ as in Fact 2.6.

Suppose that $x, y, \{x_i, y_i, z_i\}_{i=1}^N$ are points in X such that:

- (1) $[x, y]$ is D -marked with $([x_i, y_i])_{i=1}^N, ([z_i, x_i])_{i=1}^N$, and
- (2) $[x_i, y_i], [z_i, x_i]$ are ϵ -thick and longer than $\max(L_1, L_2)$ for $i = 1, \dots, N$.

Then we have that:

- (1) $[x, y]$ is F -witnessed by $[x_i, y_i]$ and $[z_i, x_i]$ for each $i = 1, \dots, N$,
- (2) $(x, y)_{x_i}, (x, y)_{y_i}, (x, y)_{z_i}$ are smaller than F for each $i = 1, \dots, N$.

Proof. It is assumed that $[x_N, y]$ is E -witnessed by $[x_N, y_N]$. Moreover, by assumption, $[x_{k-1}, x_k]$ is D -witnessed by $([x_{k-1}, y_{k-1}], [z_k, x_k])$ where $[x_{k-1}, y_{k-1}], [z_k, x_k]$ are ϵ -thick and longer than L_1 . Note also that $(y_k, z_k)_{x_k} \leq D$. Hence, if $[x_k, y]$ is E -witnessed by $[x_k, y_k]$, then $[x_{k-1}, y]$ is E -witnessed by $[x_{k-1}, y_{k-1}]$ by Fact 2.5. Thus, inductively, we deduce that $[x_i, y]$ is E -witnessed by $[x_i, y_i]$ for each i . Similarly, $[x, x_i]$ is E -witnessed by $[z_i, x_i]$.




FIGURE 2. Schematics for Fact 2.5 and 2.6.

Now Fact 2.6 asserts that $[x, y]$ is F -witnessed by $[x_i, y_i]$ and $[z_i, y_i]$, which also implies the second item. \square

We finally need two facts that guarantee witnessing by a pair of segments.

Fact 2.8 ([Cho21, Lemma 3.6]). *For each $C, \epsilon > 0$, there exists $D > C$ that satisfies the following. If $x, y, z, x' \in X$ satisfy that*

- (1) $[x, y], [z, x']$ are ϵ -thick;
- (2) $(x, z)_y, (y, x')_z < C$, and
- (3) $d(y, z) \geq d(x, y), d(z, x'), 3D$,

then $[x, x']$ is D -witnessed by $([x, y], [z, x'])$.

Fact 2.9 ([Cho21, Lemma 3.7]). *For each $C, \epsilon > 0$, there exists $D > C$ that satisfies the following condition. If $x, y, x', z \in X$ satisfy that*

- (1) $[x, y], [x', z]$ are ϵ -thick and
- (2) $(x, z)_y, (x, x')_z < C$,

then $[x, x']$ is D -witnessed by $([x, y], [z, x'])$.

From now on, we permanently fix a base point $o \in X$ and a finitely generated subgroup G of $\text{Isom}(X)$ that contains independent loxodromics a, b . For $S \subset G$ and $i \in \mathbb{Z}$, we employ the notation $S^{(i)} := \{s^i : s \in S\}$.

Let us introduce the notion of Schottky set that originated from [Gou21].

Definition 2.10 (Schottky set). *Let $K, K', \epsilon > 0$. A finite set S of isometries of X is said to be (K, K', ϵ) -Schottky if the following hold:*

- (1) for all $x, y \in X$, $|\{s \in S : (x, s^i y)_o \geq K \text{ for some } i > 0\}| \leq 2$;
- (2) for all $x, y \in X$, $|\{s \in S : (x, s^i y)_o \geq K \text{ for some } i < 0\}| \leq 2$;




FIGURE 3. Schematics for Fact 2.8, 2.9.

- (3) for all $s \in S$ and $i \neq 0$, $0 \leq iK' - d(o, s^i o) < iK'/2000$; and
- (4) for all $s \in S$ and $i \in \mathbb{Z}$, the geodesic $[o, s^i o]$ is ϵ -thick.
- (5) for all $x \in X$, $|\{s \in S : (x, s^i o)_o \geq K \text{ for some } i > 0\}| \leq 1$;
- (6) for all $x \in X$, $|\{s \in S : (x, s^i o)_o \geq K \text{ for some } i < 0\}| \leq 1$;
- (7) for all $s_1, s_2 \in S$ and $i, j > 0$, $(s_1^i o, s_2^{-j} o)_o < K$.

Remark 2.11. Gou zel's original definition of Schottky set in [Gou21] and Choi's definition in [Cho21] do not require (5), (6) and (7); however, as remarked there, Schottky sets constructed in [Cho21] automatically satisfy (5), (6) and (7).

We now claim the existence of certain Schottky sets.

Proposition 2.12 (cf. [Cho21, Proposition 4.2]). *There exists $\epsilon > 0$ such that the following holds. For any $n, n' > 0$, there exist $K(n) > 0$ and $K'(n, n') > n'$ such that G has a (K, K', ϵ) -Schottky subset S with cardinality at least n .*

Proof. The proof of [Cho21, Proposition 4.2] asserts the existence of $\epsilon > 0$, F, N_0 such that the following holds for all $N > N_0$:

- (a) for any $0 \leq m \leq n$ and $\phi_i \in \{a, b, a^{-1}, b^{-1}\}$ such that $\phi_i \notin \phi_{i+1}^{-1}$, $[o, \phi_1^{2N} \cdots \phi_n^{2N} o]$ is ϵ -thick and $(o, \phi_1^{2N} \cdots \phi_n^{2N} o)_{\phi_1^{2N} \cdots \phi_m^{2N} o} \leq F$;
- (b) for all n and k , the set

$$S_{n,k,N} := \{(\phi_1^{2N} \cdots \phi_n^{2N})^{2k} : \phi_i \in \{a, b\}\}$$

satisfies Property (1), (2) and (4) of Schottky sets for

$$K(n, N) := \max\{d(o, \phi_1^{2N} \cdots \phi_n^{2N} o) : \phi_i \in \{a, b\}\}.$$

In fact, $S_{n,k,N}$ also satisfies Property (5) and (6).

The proof of this fact is given in Appendix A. Assuming this, we now take large enough N such that $d(o, a^{2N} o), d(o, b^{2N} o) > 10000F$ and fix $K(n) := K(n, N)$. Let \mathcal{S} be the collection of concatenations of n copies of a^{2N} and n copies of b^{2N} . For $s \in \mathcal{S}$ and k , the property (a) above implies

$$0 \leq [2nk d(o, a^{2N} o) + 2nk d(o, b^{2N} o)] - d(o, s^k o) \leq 8nkF \leq \frac{1}{2500} \cdot 2nk [d(o, a^{2N} o), d(o, b^{2N} o)].$$

We finally fix k such that $K'(n, n') := 2nk[d(o, a^{2N}o) + d(o, b^{2N}o)]$ is larger than n' . Then $\mathcal{S}^{(k)}$ is (K, K', ϵ) -Schottky and has cardinality $\binom{2n}{n} \geq n$. \square

From now on, we permanently fix the choice $\epsilon > 0$ from Proposition 2.12. Now for each $C > 0$, we fix:

- $D = D(C, \epsilon)$ that works in Fact 2.8 and 2.9,
- $E = E(D, \epsilon)$, $L_1 = L(D, \epsilon)$ as in Fact 2.5,
- $F = F(E, \epsilon)$, $L_2 = L(D, \epsilon)$ as in Fact 2.6

and $L = \max(L_1, L_2)$. Note that D , F and L ultimately depend on the values of C and ϵ ; we will write them as $D(C, \epsilon)$, $F(C, \epsilon)$ and $L(C, \epsilon)$.

Lemma 2.13. *Let $K > 0$, $K' > 2L(K, \epsilon) + 5000F(K, \epsilon)$ and S_1 be a (K, K', ϵ) -Schottky set. Then S_1 and $S_1^{(-1)}$ are disjoint.*

Moreover, if a sequence $\{s_i\}_{i=1}^N$ of elements of $S_1 \cup S_1^{(-1)}$ satisfy $s_i \neq s_{i+1}^{-1}$ for $i = 1, \dots, N-1$, then $s_1 \cdots s_N \neq id$.

Proof. For each $s \in S_1^{(\pm 1)}$, we claim that $\{s' \in S_1 \cup S_1^{(-1)} : (so, s'o)_o > K\} = \{s\}$. Indeed, $(so, so)_o = d(o, so) > K$ and Properties (5), (6), (7) of Schottky sets imply $\{s' \in S_1 \cup S_1^{(-1)} : (so, s'o)_o > K\} = \{s' \in S_1^{(\pm 1)} : (so, s'o)_o > K\} = \{s\}$. In particular, $s' \in S_1^{(\mp 1)}$ cannot belong to this set; this settles the first claim.

We now let $x_i = s_1 \cdots s_i o$ for $i = 0, \dots, N$. By the above claim, we realize that $(x_i, x_{i+2})_{x_i} = (s_{i+1}^{-1}o, s_{i+2}o)_o < K < D(K, \epsilon)$ for $i = 0, \dots, N-2$. Then Lemma 2.9 implies that $[x_i, x_2]$ is $D(K, \epsilon)$ -witnessed by $([x_i, x_{i+1}], [x_{i+1}, x_2])$. Note also that $[x_i, x_{i+1}]$ is trivially $D(K, \epsilon)$ -witnessed by $([x_i, x_{i+1}], [x_{i+1}, x_{i+1}])$, and by $([x_i, x_i], [x_i, x_{i+1}])$. Combining these observations, we deduce that $[x_0, x_N]$ is $D(K, \epsilon)$ -marked with $([x_{2i-1}, x_{2i}])_{i=1}^{\lfloor N/2 \rfloor}$, $([x_{2i-2}, x_{2i-1}])_{i=1}^{\lfloor N/2 \rfloor}$. Since $[x_i, x_{i+1}]$ are ϵ -thick and longer than $0.999K' > L(K, \epsilon)$, Corollary 2.7 implies that $(x_0, x_{i+1})_{x_i} < F(K, \epsilon)$ for each i and $d(x_0, x_N) \geq (0.999K' - F(K, \epsilon))N \geq 0.9K'N$. Hence, $s_1 \cdots s_N \neq id$. \square

Corollary 2.14. *Let S_1 be as in Lemma 2.13. Then the correspondence $a \mapsto a^2$ from $S_1 \cup S_1^{(-1)}$ to $S_1^{(2)} \cup S_1^{(-2)}$ is 1-1.*

Definition 2.15. *We say that a finite set S is nicely populated by S_0 if $S_0 \subseteq S$ and $\#S_0 \geq 0.99 \cdot \#S + 400$.*

Lemma 2.16. *Given a finite set $S' \subseteq G$, there exist a (K, K', ϵ) -Schottky subset S_1 of G such that $K' > 2L(K, \epsilon) + 5000F(K, \epsilon)$, and a finite symmetric generating set S of G such that $S' \cup \{e\} \subseteq S$ and S is nicely populated by $S_1^{(2)} \cup S_1^{(-2)}$.*

Proof. We first enlarge S' into a finite symmetric generating set S'' containing e . Let $n = 100\#S'' + 40000$ and take $K = K(n) > 0$ from Proposition 2.12. We then take $F = F(K, \epsilon)$ and $L = L(K, \epsilon)$ as described before. Next, we take $K' > 2L + 5000F$ and a (K, K', ϵ) -Schottky subset S_1 of G with cardinality at least n , from Proposition 2.12. Thanks to Corollary 2.14, we

also have $\#(S_1^{(2)} \cup S_1^{(-2)}) = 2 \cdot \#S_1 \geq n$. Hence, the union of S'' , $S_1^{(2)}$, and $S_1^{(-2)}$ satisfy the desired property. \square

From now on, we fix constants $K > 0$, $K' > 2L(K, \epsilon) + 5000F(K, \epsilon)$, a (K, K', ϵ) -Schottky set S_1 with $K' > 2L(K, \epsilon) + 5000F(K, \epsilon)$, and a finite symmetric generating set $S \ni e$ of G that is nicely populated by $S_1^{(2)} \cup S_1^{(-2)}$. For $g \in G$ and $s \in S_1 \cup S_1^2$, we call $[go, gso]$ a Schottky segment. One should keep in mind that Schottky segments are ϵ -thick and longer than $0.999K'$. Finally, we will denote $S_1 \cup S_1^{(-1)}$ by S_0 .

3. PIVOTING IN A RANDOM WALK

We will make use of the random walk on G generated by the uniform measure μ_S on S , which is constructed as follows. We consider the *step space* $(G^{\mathbb{Z}}, \mu_S^{\mathbb{Z}})$, the product space of G equipped with the product measure of μ_S . Each step path (g_n) induces a sample path (ω_n) by

$$\omega_n = \begin{cases} g_1 \cdots g_n & n > 0 \\ e & n = 0 \\ g_0^{-1} \cdots g_{n+1}^{-1} & n < 0, \end{cases}$$

which constitutes a random walk with transition probability μ_S .

Given $\mathbf{g} = (g_1, \dots, g_n) \in G^n$, we define

$$\Theta(\mathbf{g}) = \{\vartheta(1) < \dots < \vartheta(N)\} := \left\{ 1 \leq i \leq n/2 : g_{2i-1}, g_{2i} \in S_1^{(2)} \cup S_1^{(-2)} \right\}.$$

For each $1 \leq i \leq N$, let a_i, b_i be the elements of $S_0 = S_1 \cup S_1^{-1}$ such that

$$a_i^2 = g_{2\vartheta(i)-1}, \quad b_i^2 = g_{2\vartheta(i)}.$$

We also define $w_i := g_{2\vartheta(i)+1} \cdots g_{2\vartheta(i+1)-2}$ for $1 \leq i \leq N-1$ and $w_0 := g_1 \cdots g_{2\vartheta(1)-2}$, $w_m := g_{2\vartheta(N)+1} \cdots g_n$.

We set $w_{0,0} = w_{0,2}^+ := e$ and define

$$w_{i,2}^- = \omega_{2\vartheta(i)-2}, \quad w_{i,1}^- = \omega_{2\vartheta(i)-2} a_i, \quad w_{i,0} = \omega_{2\vartheta(i)-1}, \quad w_{i,2}^+ = \omega_{2\vartheta(i)}$$

for $1 \leq i \leq N$. This amounts to saying


$$w_{i,2}^- = w_{i-1,2}^+ w_{i-1}, \quad w_{i,1}^- = w_{i,2}^- a_i, \quad w_{i,0} = w_{i,2}^- a_i^2, \quad w_{i,2}^+ = w_{i,2}^- a_i^2 b_i^2.$$

We also set $w_{N+1,2}^- = \omega_n$. Finally, let $y_{i,t}^\pm := w_{i,t}^\pm o$ and $y_{i,0} := w_{i,0} o$.

We now inductively define sets $P_k(\mathbf{g}) \subseteq \{1, \dots, k\}$ and the moving point z_k for $k = 0, \dots, N$. First take $P_0 := \emptyset$ and $z_0 := o$. Now given (P_{k-1}, z_{k-1}) , (P_k, z_k) is determined as follows.

- (A) If $a_k \neq b_k^{-1}$, $(z_{k-1}, y)_{y_{k,2}^-} < K$ for $y \in \{y_{k,1}^-, y_{k,0}\}$, and $(y_{k,0}, y_{k+1,2}^-)_{y_{k,2}^+} < K$, then we set $P_k := P_{k-1} \cup \{k\}$ and $z_k := y_{k,0}$.
- (B) If not, we search for sequences $\{i(1) < \dots < i(M)\} \subseteq P_{k-1}$ ($M \geq 2$) such that:

- (i) $[y_{i(1),0}, y_{i(M),1}^-]$ is fully $D(K, \epsilon)$ -marked with
- $$(\gamma_i)_{i=1}^{M-1} := \left([y_{i(1),0}, y_{i(1),2}^+], [y_{i(2),1}^-, y_{i(2),0}], \dots, [y_{i(M-1),1}^-, y_{i(M-1),0}] \right),$$
- $$(\eta_i)_{i=2}^M := \left([y_{i(2),2}^-, y_{i(2),1}^-], [y_{i(3),2}^-, y_{i(3),1}^-], \dots, [y_{i(M),2}^-, y_{i(M),1}^-] \right),$$
- (ii) $(y_{i(M),1}^-, y_{k+1,2}^-)_{y_{i(M),0}} < K$.
 If exists, let $\{i(1) < \dots < i(M)\}$ be such sequence with maximal $i(1)$; we set $P_k := P_{k-1} \cap \{1, \dots, i(1)\}$ and $z_n := y_{i(M),1}^-$. If such sequence does not exist, then we set $P_k := \emptyset$ and $z_k := o$.¹


 FIGURE 4. Loci $y_{i,k}^{\pm}$ inside a trajectory.

Note the following facts:

- (1) $P_k(\mathbf{g})$ is measurable with respect to the choice of g_i .
- (2) $i \in P_m$ only if $P_i = P_{i-1} \cup \{i\}$ and i survives during steps $i+1, \dots, m$.
- (3) If $i \in P_m$ and $i \in P_n$, then $\{1, \dots, i\} \cap P_m = \{1, \dots, i\} \cap P_n$.

Lemma 3.1. *Let $l < m$ be consecutive elements in P_k and $y \in \{y_{m,0}, y_{m,1}^-\}$. Then there exist $\{l = i(1) < \dots < i(M) = m\} \subseteq P_k$ such that $[y_{l,0}, y]$ is fully $D(K, \epsilon)$ -marked with*

$$(\gamma_i)_{i=1}^{M-1} := \left([y_{i(1),0}, y_{i(1),2}^+], [y_{i(2),1}^-, y_{i(2),0}], \dots, [y_{i(M-1),1}^-, y_{i(M-1),0}] \right),$$

$$(\eta_i)_{i=2}^M := \left([y_{i(2),2}^-, y_{i(2),1}^-], [y_{i(3),2}^-, y_{i(3),1}^-], \dots, [y_{i(M),2}^-, y] \right).$$

Proof. $l, m \in P_n$ implies that $l \in P_l$ and $l, m \in P_m$. In particular, l (m , resp.) is newly chosen at step l (m , resp.) by fulfilling Criterion (A). Hence, $(y_{l,0}, y_{l+1,2}^-)_{y_{l,2}^+} < K$ and $z_l = y_{l,0}$. Moreover, $P_m = P_{m-1} \cup \{m\}$ and $l = \max P_{m-1}$.

¹When there are several sequences that realize maximal $i(1)$, we choose the maximum in the lexicographic order on the length of sequences and $i(2), i(3), \dots$

If $l = m - 1$ and m was newly chosen at step $m = l + 1$, we have $(y_{l,0}, y)_{y_{m,2}^-} = (z_l, y)_{y_{m,2}^-} < K$ also. Then Lemma 2.9 implies that $[y_{l,0}, y]$ is fully $D(K, \epsilon)$ -marked with $[y_{l,0}, y_{l,2}^+]$ and $[y_{m,2}^-, y]$.

If $l < m - 1$, then $l = \max P_{m-1}$ has survived at step $m - 1$ by fulfilling Criterion (B); there exist $l = i(1) < \dots < i(M)$ in P_{m-2} such that $[y_{l,0}, y_{m,2}^-]$ is fully $D(K, \epsilon)$ -marked with

$$\left([y_{l,0}, y_{l,2}^+], [y_{i(2),1}^-, y_{i(2),0}^-], \dots, [y_{i(M-1),1}^-, y_{i(M-1),0}^-] \right), \left([y_{i(2),2}^-, y_{i(2),1}^-], \dots, [y_{i(M),2}^-, y_{i(M),1}^-] \right),$$

and $(y_{i(M),1}^-, y_{m,2}^-)_{y_{i(M),0}} < K$. We also have $z_{m-1} = y_{i(M),1}^-$.

Since m is newly chosen based on Criterion (A), we have $(y_{i(M),1}^-, y)_{y_{m,2}^-} = (z_{m-1}, y)_{y_{m,2}^-} < K$. Then Lemma 2.9 implies that $[y_{i(M),1}^-, y]$ is also $D(K, \epsilon)$ -witnessed by $([y_{i(M),1}^-, y_{i(M),0}^-], [y_{m,2}^-, y])$. Finally, we observe that $(y_{i(M),0}, y_{i(M),2}^-)_{y_{i(M),1}^-} = (a_{i(M)} o, a_{i(M)}^{-1} o)_o < K < D(K, \epsilon)$. Combining these implies that $[y_{l,0}, y]$ is fully $D(K, \epsilon)$ -marked with

$$\begin{aligned} & \left([y_{l,0}, y_{l,2}^+], [y_{i(2),1}^-, y_{i(2),0}^-], \dots, [y_{i(M),1}^-, y_{i(M),0}^-] \right), \\ & \left([y_{i(2),2}^-, y_{i(2),1}^-], \dots, [y_{i(M),2}^-, y_{i(M),1}^-], [y_{m,2}^-, y] \right). \quad \square \end{aligned}$$

Lemma 3.2. *Let $P_N(\mathbf{g}) = \{i(1) < \dots < i(m)\}$. Then there exist Schottky segments $(\gamma_l)_{l=1}^M$, $(\eta_l)_{l=1}^M$ and $1 \leq l(1) < \dots < l(m) \leq M$ such that:*

- (1) $[o, \omega_n o]$ is $D(K, \epsilon)$ -marked with (γ_l) , (η_l) , and
- (2) $\gamma_{l(t)} = [y_{i(t),0}, y_{i(t),2}^+]$ and $\eta_{l(t)} = [y_{i(t),2}^-, y_{i(t),0}]$.

Proof. We will apply Lemma 2.4. Since we have established Lemma 3.1, it suffices to prove the following items:

- (1) $[o, y_{i(1),0}]$ is fully $D(K, \epsilon)$ -marked with $[o, o]$, $[y_{i(1),2}^-, y_{i(1),0}]$;
- (2) $[y_{i(m),0}, \omega_n o]$ is fully $D(K, \epsilon)$ -marked with $(\gamma'_j)_{j=1}^M$, $(\eta'_j)_{j=2}^{M+1}$, where $\gamma'_1 = [y_{i(m),0}, y_{i(m),2}^+]$ and $(\gamma'_j)_{j=1}^M$, $(\eta'_j)_{j=2}^M$ are Schottky;
- (3) $(y_{i(t),2}^-, y_{i(t),2}^+)_{y_{i(t),0}} < D(K, \epsilon)$ for each $t = 1, \dots, m$.

First, $i(1) = \min P_N$ implies that $P_{i(1)-1} = \emptyset$, $z_{i(1)-1} = o$ and that $i(1)$ is newly chosen at step $i(1)$. Hence, we have $(o, y_{i(1),0})_{y_{i(1),2}^-} = (z_{i(1)-1}, y_{i(1),0})_{y_{i(1),2}^-} < K$ and Lemma 2.9 implies the first item.

Next, we observe how $i(m)$ survived in P_N . If $i(m) = N$, then it was newly chosen at step N ; $(y_{i(m),0}, y_{N+1,2}^-)_{y_{i(m),2}^+} < K$ holds and Lemma 2.9 implies that $[y_{i(m),0}, \omega_n o]$ is fully $D(K, \epsilon)$ -marked with $[y_{i(m),0}, y_{i(m),2}^+]$, $[\omega_n o, \omega_n o]$. If $i(m) \neq N$, then it has survived at step N by fulfilling Criterion (B). Thus, there exist $i \in P_{N-1}$ such that $(y_{i,1}^-, y_{N+1,2}^-)_{y_{i,0}} < K$, and Schottky segments $(\gamma'_j)_{j=1}^{M'-1}$, $(\eta'_j)_{j=2}^{M'}$ such that $[y_{i(m),0}, y_{i,1}^-]$ is fully $D(K, \epsilon)$ -witnessed by (γ'_j) , (η'_j) , and $\gamma'_1 = [y_{i(m),0}, y_{i(m),2}^+]$, $\eta'_{M'} = [y_{i,2}^-, y_{i,1}^-]$. By Lemma 2.9, we realize

that $[y_{i,1}^-, y_{N+1,2}^-] = [y_{i,1}^-, \omega_n o]$ is $D(K, \epsilon)$ -witnessed by $([y_{i,1}^-, y_{i,0}], [\omega_n o, \omega_n o])$. Since $(y_{i,2}^-, y_{i,0})_{y_{i,1}^-} = (a_i^{-1} o, a_i o)_o < K < D(K, \epsilon)$, we deduce that $[y_{i(m),0}, \omega_n o]$ is fully $D(K, \epsilon)$ -marked with

$$(\gamma'_1, \dots, \gamma'_{M'-1}, [y_{i,1}^-, y_{i,0}]), (\eta'_2, \dots, \eta'_{M'}, [\omega_n o, \omega_n o]).$$

This settles the second item.

Finally, for each $t = 1, \dots, m$, $i(t) \in P_N(\mathbf{g})$ implies that $i(t)$ was newly chosen at step $i(t)$ by fulfilling Criterion (A); hence, $a_{i(t)}^{-1} \neq b_{i(t)}$ and we deduce $(y_{i(t),2}^-, y_{i(t),0}^+)_{y_{i(t),0}} = (a_{i(t)}^{-2} o, b_{i(t)}^2 o)_o < K$ from Properties (5), (6), (7) of Schottky sets. \square

The same proof also yields the following lemma:

Lemma 3.3. *Let $k < k'$ be elements of $P_N(\mathbf{g})$ and $y \in \{y_{k',0}, y_{k',1}^-\}$. Then there exist Schottky segments $(\gamma_l)_{l=1}^{M-1}$, $(\eta_l)_{l=2}^M$ such that:*

- (1) $[y_{k,0}, y]$ is fully $D(K, \epsilon)$ -marked with (γ_l) , (η_l) , and
- (2) $\gamma_1 = [y_{k,0}, y_{k,2}^+]$ and $\eta_M = [y_{k',2}^-, y]$.

Having established properties of the set $P_N(\mathbf{g})$, our next goal is to estimate the size of $P_N(\mathbf{g})$. We first fix N , $\Theta = \{\vartheta(1) < \dots < \vartheta(N)\}$ and the choices $g_i \in S$ for $i \notin \{2j-1, 2j : j \in \Theta\}$ such that $(g_{2j-1}, g_{2j}) \notin \left(S_1^{(2)} \cup S_1^{(-2)}\right)^2$ for $j \notin \Theta$. Conditioned on these choices, we draw $(g_{2\theta(1)-1}, g_{2\theta(1)}, \dots, g_{2\theta(N)-1}, g_{2\theta(N)})$ from $\left(S_1^{(2)} \cup S_1^{(-2)}\right)^{2N}$, with the product measure of the uniform measure on $S_1 \cup S_1^{-1}$. In particular, we are fixing $\Theta(\mathbf{g}) = \Theta$ and w_i 's, and regard $P_k(\mathbf{g})$ as an RV of $s = (a_1, b_1, \dots, a_k, b_k)$; here, a_i, b_i are independently drawn from S_0 with the uniform measure.

Lemma 3.4. *For $0 \leq k < N$ and $s \in S_0^{2k}$, we have*

$$\mathbb{P}\left(\#P_{k+1}(s, a_{k+1}, b_{k+1}) = \#P_k(s) + 1\right) \geq 9/10.$$

Proof. Recall Criterion (A) for $\#P_{k+1} = \#P_k + 1$. Note that the condition

$$(3.1) \quad (y_{k,0}, y_{k+1,2}^-)_{y_{k,2}^+} = (b_k^{-2} o, w_k o)_o < K$$

depends only on b_k and not on other a_i 's and b_i 's. This holds for at least $(\#S_1 - 1)$ choices of b_k in S_1 and $(\#S_1 - 1)$ choices in $S_1^{(-1)}$.

We fix one such choice $b_k \in S_1^{(\pm)}$, and also $a_1, b_1, \dots, a_{k-1}, b_{k-1}$ that determine $w_{k,2}^-$ and z_{k-1} . Then

$$(3.2) \quad \begin{aligned} (z_{k-1}, y_{k,1}^-)_{y_{k,2}^-} &= ((w_{k,2}^-)^{-1} z_{k-1}, a_k o)_o < K, \\ (z_{k-1}, y_{k,0})_{y_{k,2}^-} &= ((w_{k,2}^-)^{-1} z_{k-1}, a_k^2 o)_o < K, \\ a_k &\neq b_k^{-1} \end{aligned}$$

hold for at least $(\#S_1 - 1)$ choices of a_k in $S_1^{(\pm 1)}$ and $(\#S_1 - 2)$ choices in $S_1^{(\mp 1)}$. Since Condition 3.1 and Condition 3.2 together constitute Criterion (A), we obtain

$$\mathbb{P}(\#P_{k+1} = \#P_k + 1) \geq \frac{2\#S_1 - 2}{2\#S_1} \cdot \frac{2\#S_1 - 3}{2\#S_1} \geq 0.9. \quad \square$$

Given $a_1, b_1, \dots, a_{k-1}, b_{k-1}$ and b_k , we define the set \tilde{S}'_k of elements a_k in S_0 that satisfy Condition 3.2. In the proof above, we have observed that $\# [S_0 \setminus \tilde{S}'_k] \leq 3$.

Lemma 3.5. *Let $i \in P_k(s)$ for a choice $s = (a_1, b_1, \dots, a_N, b_N)$, and \bar{s} be obtained from s by replacing a_i with $\bar{a}_i \in \tilde{S}'_i(a_1, b_1, \dots, a_{i-1}, b_{i-1}, b_i)$. Then $P_l(s) = P_l(\bar{s})$ and $\tilde{S}'_l(s) = \tilde{S}'_l(\bar{s})$ for any $1 \leq l \leq k$.*

Proof. Since $a_1, b_1, \dots, a_{i-1}, b_{i-1}$ are intact, $P_l(s) = P_l(\bar{s})$ and $\tilde{S}'_l(s) = \tilde{S}'_l(\bar{s})$ hold for $l = 0, \dots, i-1$. At step i , b_i satisfies Condition 3.1 (since $i \in P_k(s)$) and \bar{a}_i satisfies Condition 3.2; hence, $i \in P_k(\bar{s})$ and $P_i(s) = P_i(\bar{s})$. We also have $\tilde{S}'_i(s) = \tilde{S}'_i(\bar{s})$. At this step, however, we have $\bar{z}_i = \bar{y}_{i,0} = gy_{i,0} = gz_i$ where $g := w_{i,2}^- \bar{a}_i^2 (w_{i,2}^- a_i^2)^{-1}$. More generally, we have

$$(3.3) \quad \begin{aligned} w_{l,t}^- &= gw_{l,t}^- & (t \in \{1, 2\}, l > i), \\ w_{l,0} &= gw_{l,0} & (l \geq i), \\ w_{l,2}^+ &= gw_{l,2}^+ & (l \geq i). \end{aligned}$$

We now claim the following for $i < l \leq k$:

- (1) If s fulfills Criterion (A) at step l , then so does \bar{s} .
- (2) If not and $\{i(1) < \dots < i(M)\} \subseteq P_{l-1}(s)$ is the maximal sequence for s in Criterion (B) at step l , then it is also the maximal one for \bar{s} at step l .
- (3) In both cases, we have $P_l(s) = P_l(\bar{s})$ and $\bar{z}_l = gz_l$.

Assuming the third item for $l-1$: $P_{l-1}(s) = P_{l-1}(\bar{s})$ and $\bar{z}_{l-1} = gz_{l-1}$, Equality 3.3 implies the first item. In this case we also deduce $P_l(s) = P_{l-1}(s) \cup \{l\} = P_{l-1}(\bar{s}) \cup \{l\} = P_l(\bar{s})$ and $\bar{z}_l = \bar{y}_{l,0} = gy_{l,0} = gz_l$, the third item for l .

Furthermore, Equality 3.3 also implies that a sequence $\{i(1) < \dots < i(M)\}$ in $P_{l-1}(s) \cap \{i, \dots, l-1\} = P_{l-1}(\bar{s}) \cap \{i, \dots, l-1\}$ works for s in Criterion (B) if and only if it works for \bar{s} . Note that $i \in P_l(s)$ since $i \in P_k(s)$ and $l \leq k$; hence, such sequences exist and the maximal sequence is chosen among them. Therefore, the maximal sequence $\{i(1) < \dots < i(M)\}$ for s is also maximal for \bar{s} . We then deduce $P_l(s) = P_{l-1}(s) \cap \{1, \dots, i(1)\} = P_{l-1}(\bar{s}) \cap \{1, \dots, i(1)\} = P_l(\bar{s})$ and $\bar{z}_l = \bar{y}_{i(M),1}^- = gy_{i(M),1}^- = gz_l$, the third item for l .

Since we have $\bar{z}_i = gz_i$, induction shows that $P_l(s) = P_l(\bar{s})$ for each $i < l \leq k$. Moreover, Equality 3.3 and $\bar{z}_l = gz_l$ implies that $\tilde{S}'_l(s) = \tilde{S}'_l(\bar{s})$. \square

Given $1 \leq k \leq N$ and a partial choice $s = (a_1, b_1, \dots, a_k, b_k)$, we say that $\bar{s} = (\bar{a}_1, \bar{b}_1, \dots, \bar{a}_k, \bar{b}_k)$ is *pivoted from* s if $b_j = \bar{b}_j$ for all $1 \leq j \leq k$, $\bar{a}_i \in \tilde{S}'_i(s)$ for $i \in P_k(s)$, and $a_j = \bar{a}_j$ for all other $j \notin P_k(s)$. Then Lemma 3.5 asserts that being pivoted from each other is an equivalence relation. For each $s \in S_0^{2k}$, let $\mathcal{E}_k(s)$ be the equivalence class of s .

Lemma 3.6. *For $0 \leq k < N$, $j \geq 0$ and $s \in S_0^{2k}$, we have*

$$\mathbb{P} \left(\#P_{k+1}(\tilde{s}, a_{k+1}, b_{k+1}) < \#P_k(s) - j \mid \tilde{s} \in \mathcal{E}_k(s), (a_{k+1}, b_{k+1}) \in S_0^2 \right) \leq 1/10^{j+1}.$$

Proof. Let us fix $s = (a_1, b_1, \dots, a_k, b_k) \in S_0^{2k}$ and

$$\mathcal{A} := \left\{ (a_{k+1}, b_{k+1}) \in S_0^2 : \#P_{k+1}(s, a_{k+1}, b_{k+1}) = \#P_k(s) + 1 \right\}.$$

Then Lemma 3.4 implies that $\mathbb{P}(\mathcal{A} \mid S_0^2) \geq 0.9$. Moreover, for $(a_{k+1}, b_{k+1}) \in \mathcal{A}$ and $\tilde{s} \in \mathcal{E}_k(s)$, $(\tilde{s}, a_{k+1}, b_{k+1})$ is pivoted from (s, a_{k+1}, b_{k+1}) also since $P_k(s) \subseteq P_{k+1}(s, a_{k+1}, b_{k+1})$. Lemma 3.5 then implies that $P_{k+1}(\tilde{s}) = P_{k+1}(s) = P_k(s) \cup \{k+1\} = P_k(\tilde{s}) \cup \{k+1\}$, and we have

$$\mathbb{P} \left(\#P_{k+1}(\tilde{s}, a_{k+1}, b_{k+1}) < \#P_k(\tilde{s}) \mid \tilde{s} \in \mathcal{E}_k(s), (a_{k+1}, b_{k+1}) \in S_0^2 \right) \leq 1 - \mathbb{P}(\mathcal{A}) \leq 1/10.$$

This settles the case $j = 0$.

Now let $j = 1$. The event under discussion becomes void when $\#P_k(s) \leq 2$. Excluding such cases, let $l < m$ be the last 2 elements of $P_k(s)$. For $\tilde{s} \in \mathcal{E}_k(s)$ and each $A \subseteq \tilde{S}'_m(s)$ we define

$$E(\tilde{s}, A) := \left\{ \bar{s} = (\bar{a}_i, \bar{b}_i)_{i=1}^k : \bar{b}_i = \tilde{b}_i \text{ for all } i, \bar{a}_i = \tilde{a}_i \text{ for } i \neq m, \bar{a}_m \in A \right\}.$$

Then $\{E(\tilde{s}, \tilde{S}'_m(s)) : \tilde{s} \in \mathcal{E}_k(s)\}$ partitions $\mathcal{E}_k(s)$ by Lemma 3.5.

We now fix $(a_{k+1}, b_{k+1}) \in S_0^2$ and $\tilde{s} \in \mathcal{E}_k(s)$, and let $A' \subseteq \tilde{S}'_m(s)$ be the collection of elements \bar{a}_m that satisfies

$$(3.4) \quad (\bar{a}_m^{-1} o, (\tilde{w}_{m,0})^{-1} \tilde{w}_{k,2}^{-1} a_{k+1}^2 b_{k+1}^2 w_{k+1} o)_o = (\bar{a}_m^{-1} o, \tilde{w}_m^2 w_m \cdots \tilde{a}_k^2 \tilde{b}_k^2 w_k a_{k+1}^2 b_{k+1}^2 w_{k+1} o)_o < K.$$

By Property (5), (6) of Schottky sets, we have $\# \left[\tilde{S}'_m(s) \setminus A' \right] \leq 2$.

We now claim that $\#P_{k+1}(\bar{s}, a_{k+1}, b_{k+1}) \geq \#P_k(s) - 1$ for $\bar{s} \in E(\tilde{s}, A')$. First, since $l < m$ are consecutive elements in $P_k(\bar{s})$, Lemma 3.1 gives a sequence $\{l = i(1) < \dots < i(M) = m\} \subseteq P_k$ such that $[\bar{y}_{l,0}, \bar{y}_{m,1}]$ is fully $D(K, \epsilon)$ -marked with

$$\begin{aligned} (\gamma_i)_{i=1}^{M-1} &:= \left([\bar{y}_{i(1),0}, \bar{y}_{i(1),2}^+], [\bar{y}_{i(2),1}, \bar{y}_{i(2),0}], \dots, [\bar{y}_{i(M-1),1}, \bar{y}_{i(M-1),0}] \right), \\ (\eta_i)_{i=2}^M &:= \left([\bar{y}_{i(2),2}, \bar{y}_{i(2),1}], [\bar{y}_{i(3),2}, \bar{y}_{i(3),1}], \dots, [\bar{y}_{i(M),2}, \bar{y}_{i(M),1}] \right). \end{aligned}$$

Moreover, Condition 3.4 implies that

$$(\bar{y}_{m,1}, \bar{y}_{k+2,2})_{\bar{y}_{m,0}} = (\bar{a}_m^{-1} o, (\tilde{w}_{m,0})^{-1} \tilde{w}_{k,2}^{-1} a_{k+1}^2 b_{k+1}^2 w_{k+1} o)_o < K.$$

In summary, $\{l = i(1) < \dots < i(M)\} \subseteq P_k(\bar{s})$ works for \bar{s} in Criterion (B) at step $k+1$, which implies $P_{k+1}(\bar{s}) \supseteq P_k(\bar{s}) \cap \{1, \dots, l\}$, hence the claim.

As a result, we deduce

$$\begin{aligned} & \mathbb{P} \left(\#P_{k+1}(\bar{s}, a_{k+1}, b_{k+1}) < \#P_k(s) - 1 \mid \bar{s}, (a_{k+1}, b_{k+1}), \bar{s} \in E(\tilde{s}, \tilde{S}'_m) \right) \\ & \leq \frac{\# \left[\tilde{S}'_m(s) \setminus A' \right]}{\#\tilde{S}'_m(s)} \leq \frac{2}{\#S_0 - 3} \leq 0.1. \end{aligned}$$

Since $E(\tilde{s}, \tilde{S}'_m)$'s for $\tilde{s} \in \mathcal{E}_k(s)$ partition $\mathcal{E}_k(s)$, we deduce

$$\mathbb{P} \left(\#P_{k+1}(\tilde{s}, a_{k+1}, b_{k+1}) < \#P_k(s) - 1 \mid \tilde{s} \in \mathcal{E}_k(s), (a_{k+1}, b_{k+1}) \right) \leq 0.1.$$

Moreover, the above probability vanishes when $(a_{k+1}, b_{k+1}) \in \mathcal{A}$. Since $\mathbb{P}(\mathcal{A} \mid S_0^2) \geq 0.9$, we deduce

$$(3.5) \quad \mathbb{P} \left(\#P_{k+1}(\tilde{s}, a_{k+1}, b_{k+1}) < \#P_k(s) - 1 \mid \tilde{s} \in \mathcal{E}_k(s), (a_{k+1}, b_{k+1}) \in S_0^2 \right) \leq 0.01.$$

For $j = 2$, we similarly assume $\#P_k(s) \geq 3$ and let $l' < l < m$ be the last 3 elements. We define \mathcal{A}_1 to be the set of $(\bar{a}_m, a_{k+1}, b_{k+1})$ in $\tilde{S}'_m(s) \times S_0^2$ such that

$$\#P_{k+1} \left(\underbrace{a_1, b_1, \dots, \bar{a}_m, b_m, \dots, a_k, b_k}_{\text{obtained from } s \text{ by replacing } a_m \text{ with } \bar{a}_m}, a_{k+1}, b_{k+1} \right) \geq \#P_k(s) - 1,$$

or equivalently,

$$P_k(s) \cap \{1, \dots, l\} \subseteq P_{k+1}(a_1, b_1, \dots, \bar{a}_m, b_m, \dots, a_k, b_k, a_{k+1}, b_{k+1}).$$

Now, if $\tilde{s} = (\tilde{a}_1, \tilde{b}_1, \dots, \tilde{a}_k, \tilde{b}_k) \in \mathcal{E}_k(s)$ is such that $(\tilde{a}_m, a_{k+1}, b_{k+1}) \in \mathcal{A}_1$, then $(\tilde{s}, a_{k+1}, b_{k+1})$ is pivoted from $(a_1, b_1, \dots, \bar{a}_m, b_m, \dots, a_{k+1}, b_{k+1})$ since they only differ at a_i 's for $i \in P_k(s) \cap \{1, \dots, l\} \subseteq P_{k+1}(a_1, b_1, \dots, \tilde{a}_m, b_m, \dots, a_{k+1}, b_{k+1})$. Lemma 3.5 then implies that $P_{k+1}(\tilde{s}, a_{k+1}, b_{k+1})$ also contains $P_k(s) \cap \{1, \dots, l\}$. This fact and Inequality 3.5 implies that

$$\begin{aligned} & \mathbb{P}(\mathcal{A}_1 \mid \tilde{S}'_m(s) \times S_0^2) \\ & = \mathbb{P} \left((\tilde{a}_m, a_{k+1}, b_{k+1}) \in \mathcal{A}_1 \mid \tilde{s} \in \mathcal{E}_k(s), (a_{k+1}, b_{k+1}) \in S_0^2 \right) \\ & \geq \mathbb{P} \left(\#P_{k+1}(\tilde{s}, a_{k+1}, b_{k+1}) \geq \#P_k(s) - 1 \mid \tilde{s} \in \mathcal{E}_k(s), (a_{k+1}, b_{k+1}) \in S_0^2 \right) \geq 0.99. \end{aligned}$$

We now define for $\tilde{s} \in \mathcal{E}_k(s)$ and each $A \subseteq \tilde{S}'_l(s)$

$$E_1(\tilde{s}, A) := \left\{ \bar{s} = (\bar{a}_i, \bar{b}_i)_{i=1}^k : \bar{b}_i = \tilde{b}_i \text{ for all } i, \bar{a}_i = \tilde{a}_i \text{ for } i \neq l, a_l \in A \right\}.$$

Then $\{E(\tilde{s}, \tilde{S}'_l(s)) : \tilde{s} \in \mathcal{E}_k(s)\}$ partitions $\mathcal{E}_k(s)$ by Lemma 3.5.

Now fixing $(a_{k+1}, b_{k+1}) \in S_0^2$ and $\tilde{s} \in \mathcal{E}_k(s)$, let $A'_1 \subseteq \tilde{S}'_l(s)$ be the collection of elements $\bar{a}_l \in \tilde{S}'_l(s)$ that satisfies

$$(3.6) \quad (\bar{a}_l^{-1} o, (\tilde{w}_{l,0})^{-1} \tilde{w}_{k,2}^- a_{k+1}^2 b_{k+1}^2 w_{k+1} o)_o = (\bar{a}_l^{-1} o, \tilde{b}_l^2 w_l \cdots \tilde{a}_k^2 \tilde{b}_k^2 w_k a_{k+1}^2 b_{k+1}^2 w_{k+1} o)_o < K.$$

By Property (5), (6) of Schottky sets, we have $\# \left[\tilde{S}'_m(s) \setminus A'_1 \right] \leq 2$.

We now claim that $\#P_{k+1}(\bar{s}, a_{k+1}, b_{k+1}) \geq \#P_k(s) - 2$ for $\bar{s} \in E_1(\tilde{s}, A'_1)$. First, since $l' < l$ are consecutive elements in $P_k(\bar{s})$, Lemma 3.1 gives a sequence $\{l' = i(1) < \dots < i(M) = l\} \subseteq P_k$ such that $[\bar{y}_{l',0}, \bar{y}_{l',1}]$ is fully $D(K, \epsilon)$ -marked with

$$\begin{aligned} (\gamma_i)_{i=1}^{M-1} &:= \left([\bar{y}_{i(1),0}, \bar{y}_{i(1),2}^+], [\bar{y}_{i(2),1}, \bar{y}_{i(2),0}], \dots, [\bar{y}_{i(M-1),1}, \bar{y}_{i(M-1),0}] \right), \\ (\eta_i)_{i=2}^M &:= \left([\bar{y}_{i(2),2}, \bar{y}_{i(2),1}], [\bar{y}_{i(3),2}, \bar{y}_{i(3),1}], \dots, [\bar{y}_{i(M),2}, \bar{y}_{i(M),1}] \right). \end{aligned}$$

Moreover, Condition 3.6 implies that

$$(\bar{y}_{l,1}, \bar{y}_{k+2,2})_{\bar{y}_{l,0}} = (\bar{a}_l^{-1}o, (\tilde{w}_{l,0})^{-1}\tilde{w}_{k,2}a_{k+1}^2b_{k+1}^2w_{k+1}o)_o < K.$$

In summary, $\{l' = i(1) < \dots < i(M)\} \subseteq P_k(\bar{s})$ works for \bar{s} in Criterion (B) at step $k+1$, which implies that $P_{k+1}(\bar{s}) \supseteq P_k(\bar{s}, a_{k+1}, b_{k+1}) \cap \{1, \dots, l'\}$, hence the claim.

As a result, we deduce

$$\mathbb{P} \left(\#P_{k+1}(\bar{s}, a_{k+1}, b_{k+1}) < \#P_k(s) - 2 \mid \tilde{s}, (a_{k+1}, b_{k+1}), \bar{s} \in E_1(\tilde{s}, \tilde{S}'_l) \right) \leq 0.1.$$

Here, for \tilde{s} and (a_{k+1}, b_{k+1}) such that $(\tilde{a}_m, a_{k+1}, b_{k+1}) \in \mathcal{A}_1$, the above probability vanishes. Since

$$\begin{aligned} &\mathbb{P} \left[\bigcup \{ E_1(\tilde{s}, \tilde{S}'_l) \times (a_{k+1}, b_{k+1}) : (\tilde{a}_m, a_{k+1}, b_{k+1}) \notin \mathcal{A}_1 \} \mid \mathcal{E}_k(s) \times S_0^2 \right] \\ &= \mathbb{P} \left[(\tilde{a}_m, a_{k+1}, b_{k+1}) \notin \mathcal{A}_1 \mid \tilde{S}'_m(s) \times S_0^2 \right] \leq 0.01, \end{aligned}$$

we sum up the conditional probabilities to obtain

$$(3.7) \quad \mathbb{P} \left(\#P_{k+1}(\bar{s}, a_{k+1}, b_{k+1}) < \#P_k(s) - 2 \mid \tilde{s} \in \mathcal{E}_k(s) \right) \leq 0.001.$$

We repeat this procedure to cover all $j < \#P_k(s)$. The case $j \geq \#P_k(s)$ is void. \square

Corollary 3.7. *Conditioned on paths $\mathbf{g} \in G^n$ such that $\#\Theta(\mathbf{g}) = N$, $\#P_N(\mathbf{g})$ is greater in distribution than the sum of N i.i.d. X_i , whose distribution is given by*

$$(3.8) \quad \mathbb{P}(X_i = j) = \begin{cases} 9/10 & \text{if } j = 1, \\ 9/10^{-j+1} & \text{if } j < 0, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Lemma 3.4 and Lemma 3.6 imply the following: for $0 \leq k < N$ and any i ,

$$(3.9) \quad \mathbb{P} \left(\#P_{n+1}(\mathbf{g}) \geq i + j \mid \#P_k(\mathbf{g}) = i \right) \geq \begin{cases} 1 - 1/10 & \text{if } j = 1, \\ 1 - 1/10^{-j+1} & \text{if } j < 0, \\ 0 & \text{otherwise.} \end{cases}$$

Hence, there exists a nonnegative RV U_k such that $\#P_{n+1} - U_k$ and $\#P_k + X'$ have the same distribution, where X' is an i.i.d. copy of X_{k+1} that is independent from $\#P_k$.

For each $1 \leq k \leq N$, we claim that $\mathbb{P}(\#P_k(\mathbf{g}) \geq i) \geq \mathbb{P}(X_1 + \dots + X_k \geq i)$ for each i . For $k = 1$, we have $\#P_{k-1}(\mathbf{g}) = 0$ always and the claim follows from Inequality 3.9. Given the claim for $k - 1$, we have

$$\begin{aligned} \mathbb{P}(\#P_{n+1} \geq i) &\geq \mathbb{P}(\#P_k + X' \geq i) = \sum_j \mathbb{P}(\#P_k \geq j) \mathbb{P}(X' = i - j) \\ &\geq \sum_j \mathbb{P}(X_1 + \dots + X_k \geq j) \mathbb{P}(X_{k+1} = i - j) \\ &= \mathbb{P}(X_1 + \dots + X_k + X_{k-1} \geq i). \quad \square \end{aligned}$$

Given $\mathbf{g} \in G^n$ with $\#\Theta(\mathbf{g}) = N$ and $P_N(\mathbf{g}) = \{i(1) < \dots < i(m)\}$, we finally define the l -th pivotal time $\rho(l)$ of \mathbf{g} by $2\vartheta(i(l)) - 1$ and the set of pivotal times $P_n^*(\mathbf{g})$ by

$$P_n^*(\mathbf{g}) := \{\rho(1), \dots, \rho(m)\} = \{2\vartheta(i) - 1 : i \in P_N(\mathbf{g})\}.$$

Moreover, we define $\tilde{S}_l(\mathbf{g}) := \tilde{S}'_{i(l)}(\mathbf{g})$ for $l = 1, \dots, m$. $\bar{\mathbf{g}} \in G^n$ is said to be *pivoted from* \mathbf{g} if $g_j = \bar{g}_j$ unless $j = \rho(l)$ for some $l = 1, \dots, m$, in which case we require $\bar{g}_j \in \tilde{S}_l(\mathbf{g})$. As before, being pivoted from each other is an equivalence relation in G^n .

Note that

$$w_{i(l),2}^- = \omega_{\rho(l)-1}, \quad w_{i(l),0} = \omega_{\rho(l)}, \quad w_{i(l),2}^+ = \omega_{\rho(l)+1}.$$

Lemma 3.2 and Corollary 2.7 imply that

$$(3.10) \quad (\omega_{\rho(i)} o, \omega_{\rho(k)} o)_{\omega_{\rho(j)} o} < F(K, \epsilon) < K'/4000$$

for $0 \leq i \leq j \leq k \leq m + 1$, where we set $\rho(0) := 0$ and $\rho(m + 1) := n$. Moreover, for any $0 \leq i < j \leq k \leq m$, we have

$$(3.11) \quad \begin{aligned} 1.999K' &\leq d(\omega_{\rho(k)-1} o, \omega_{\rho(k)} o) = d(o, a_{i(k)}^2 o) \leq 2K', \\ (\omega_{\rho(i)} o, \omega_{\rho(k)} o)_{\omega_{\rho(j)-1} o} &< F(K, \epsilon) < K'/4000. \end{aligned}$$

4. PIVOTING AND TRANSLATION LENGTHS

We will now define another equivalence relation on paths with sufficiently many pivots. For $\mathbf{g} \in G^n$ with m pivotal times, where $n/5 \leq m \leq n/2$, we define

$$\begin{aligned} D_f(\mathbf{g}) &= \sum_{l=1}^{\lfloor n/12 \rfloor} [d(\omega_{\rho(l-1)} o, \omega_{\rho(l)-1} o) + 2K'], \\ D_b(\mathbf{g}) &= \sum_{l=m-\lfloor n/12 \rfloor+1}^m [d(\omega_{\rho(l-1)} o, \omega_{\rho(l)-1} o) + 2K'] + d(\omega_{\rho(m)} o, \omega_n o), \\ D_t(\mathbf{g}) &= \sum_{l=1}^m [d(\omega_{\rho(l-1)} o, \omega_{\rho(l)-1} o) + 2K'] + d(\omega_{\rho(m)} o, \omega_n o). \end{aligned}$$

Note the inequality

$$(4.1) \quad \begin{aligned} D_f(\mathbf{g}) &\geq \sum_{l=1}^{\lfloor n/12 \rfloor} [d(\omega_{\rho(l-1)} o, \omega_{\rho(l-1)} o) + d(\omega_{\rho(l)-1} o, \omega_{\rho(l)} o)] \\ &\geq d(o, \omega_{\rho(i)-1}), d(o, \omega_{\rho(i)}) \end{aligned}$$

for $i = 1, \dots, \lfloor n/12 \rfloor$. Similarly, D_t dominates $d(o, \omega_n o)$. Moreover, due to Inequality 3.10 and 3.11, we have $|d(o, \omega_n o) - D_t| < 2F(K, \epsilon)n \leq K'n/1000$. We also observe that at least one of D_f, D_b is smaller than $D_t/2 - K'n/20$; indeed, $D_t - D_f - D_b$ is the sum of at least $n/30$ terms of the form $d(\omega_{\rho(l-1)} o, \omega_{\rho(l)-1} o) + 2K' \geq 3K'$.

If $D_f \leq D_b$, then we allow pivoting at the first $\lfloor n/12 \rfloor$ pivotal times. Otherwise, we allow pivoting at the last $\lfloor n/12 \rfloor$ pivotal times. Since D_f, D_b, D_t , and the set of pivotal times are invariant under pivoting, this rule partitions $\{\mathbf{g} \in G^n : \#P_n^*(\mathbf{g}) \geq n/5\}$ into equivalence classes $\mathcal{F}(\mathbf{g})$'s.

We are now ready to prove the core lemma for Theorem A.

Lemma 4.1. *Let $n > 25$ and suppose that $\mathbf{g} \in G^n$ satisfies*

$$\#P_n^*(\mathbf{g}) \geq n/5, \quad D_f(\mathbf{g}) \leq D_b(\mathbf{g}).$$

Let also $1 \leq k < k' \leq \lfloor n/12 \rfloor$ and $\tilde{g}_l \in \tilde{S}_l(\mathbf{g})$ for $l = 1, \dots, k-1, k'+1, \dots, \lfloor n/12 \rfloor$.

Then there exist $A \subseteq \tilde{S}_k(\mathbf{g})$ and $A' \subseteq \tilde{S}_{k'}(\mathbf{g})$, each of cardinality at most 2, such that the following holds: for any $\bar{\mathbf{g}} \in \mathcal{F}(\mathbf{g})$ such that $\bar{g}_{\rho(l)} = \tilde{g}_l$ for $l = 1, \dots, k-1, k'+1, \dots, \lfloor n/12 \rfloor$ and $\bar{g}_{\rho(k)} \notin A^{(2)}, \bar{g}_{\rho(k')} \notin A'^{(2)}$, we have $\tau(\bar{\omega}_n) \geq K'n/12$.

Proof. By the assumption, $\bar{\mathbf{g}} \in \mathcal{F}(\mathbf{g})$ and \mathbf{g} can differ only at step $\rho(1), \dots, \rho(\lfloor n/12 \rfloor)$. Hence, for any $\bar{\mathbf{g}} \in \mathcal{F}(\mathbf{g})$ such that $\bar{g}_{\rho(l)} = \tilde{g}_l$ for $l = 1, \dots, k-1, k'+1, \dots, \lfloor n/12 \rfloor$, the isometry

$$v := (\bar{\omega}_{\rho(k')})^{-1} \bar{\omega}_n \bar{\omega}_{\rho(k)-1} = \bar{g}_{\rho(k')+1} \cdots \bar{g}_n \cdot \bar{g}_1 \cdots \bar{g}_{\rho(k)-1}$$

is uniform. We define

$$\begin{aligned} A &:= \{g \in \tilde{S}_k(\mathbf{g}) : (v^{-1}o, g^2o)_o \geq K\}, \\ A' &:= \{g \in \tilde{S}_{k'}(\mathbf{g}) : (g^{-1}o, vo)_o \geq K\}. \end{aligned}$$

Since $\tilde{S}_k(\mathbf{g}) \subseteq S_0 = S_1 \cup S_1^{(-1)}$, Property (5), (6) of Schottky sets imply that $\#A \leq 2$. Similarly we have $\#A' \leq 2$.

Let us now fix $h \in \tilde{S}_k(\mathbf{g}) \setminus A$, $h' \in \tilde{S}_{k'}(\mathbf{g}) \setminus A'$ and consider $\bar{\mathbf{g}} \in \mathcal{F}(\mathbf{g})$ such that $\bar{g}_{\rho(l)} = \tilde{g}_l$ for $l = 1, \dots, k-1, k'+1, \dots, \lfloor n/12 \rfloor$ and $\bar{g}_{\rho(k)} = h^2$, $\bar{g}_{\rho(k')} = h'^2$.

Since $\rho(k), \rho(k') \in P_n^*(\mathbf{g}) = P_n^*(\bar{\mathbf{g}})$ and $h'^2 = \bar{g}_{\rho(k')}$, Lemma 3.3 gives Schottky segments $(\gamma_l)_{l=1}^{M-1}, (\eta_l)_{l=2}^M$ such that $[\bar{\omega}_{\rho(k)} o, \bar{\omega}_{\rho(k)-1} h' o]$ is fully $D(K, \epsilon)$ -marked with $(\gamma_l), (\eta_l)$ and $\gamma_1 = [\bar{\omega}_{\rho(k)} o, \bar{\omega}_{\rho(k)+1} o], \eta_M = [\bar{\omega}_{\rho(k')-1} o, \bar{\omega}_{\rho(k')-1} h' o]$.

Next, Inequality 4.1 implies that

$$\begin{aligned} d(o, vo) &\geq d(o, \bar{\omega}_n o) - d(o, \bar{\omega}_{\rho(k')} o) - d(o, \bar{\omega}_{\rho(k)-1} o) \\ &\geq D_t - K'n/1000 - 2D_f \\ &\geq K'n/12 \geq 2K' + 3D(K, \epsilon). \end{aligned}$$

Since we also have $(h'^{-1}o, vo)_o \leq K$ and $(o, vh^2o)_{vo} \leq K$, Lemma 2.8 implies that $[h'^{-1}o, vh^2o]$ is $D(K, \epsilon)$ -witnessed by $([h'^{-1}o, o], [vo, vh^2o])$. By applying isometry $\bar{\omega}_n^{i-1}\bar{\omega}_{\rho(k')}$, we deduce that $[\bar{\omega}_n^{i-1}\bar{\omega}_{\rho(k')-1}h'o, \bar{\omega}_n^i\bar{\omega}_{\rho(k)}o]$ is $D(K, \epsilon)$ -witnessed by Schottky segments

$$[\bar{\omega}_n^{i-1}\bar{\omega}_{\rho(k')-1}h'o, \bar{\omega}_n^{i-1}\bar{\omega}_{\rho(k')}o], \quad [\bar{\omega}_n^i\bar{\omega}_{\rho(k)-1}o, \bar{\omega}_n^i\bar{\omega}_{\rho(k)}o].$$

We now claim that $[\bar{\omega}_{\rho(k)}o, \bar{\omega}_n^i\bar{\omega}_{\rho(k)}o]$ is fully $D(K, \epsilon)$ -witnessed by

$$\begin{aligned} &(\gamma_1, \dots, \gamma_{M-1}, [\bar{\omega}_{\rho(k')-1}h'o, \bar{\omega}_{\rho(k')}o], \quad \bar{\omega}_n\gamma_1, \dots, \bar{\omega}_n\gamma_{M-1}, [\bar{\omega}_n\bar{\omega}_{\rho(k')-1}h'o, \bar{\omega}_n\bar{\omega}_{\rho(k')}o], \dots), \\ &(\eta_2, \dots, \eta_M, [\bar{\omega}_n\bar{\omega}_{\rho(k)-1}o, \bar{\omega}_n\bar{\omega}_{\rho(k)}o], \quad \bar{\omega}_n\eta_2, \dots, \bar{\omega}_n\eta_M, [\bar{\omega}_n^2\bar{\omega}_{\rho(k)-1}o, \bar{\omega}_n^2\bar{\omega}_{\rho(k)}o], \dots). \end{aligned}$$

To apply Lemma 2.4, it remains to check

$$\begin{aligned} (\bar{\omega}_{\rho(k')-1}o, \bar{\omega}_{\rho(k')}o)_{\bar{\omega}_{\rho(k')-1}h'o} &= (h'^{-1}o, h'o)_o < D(K, \epsilon), \\ (\bar{\omega}_{\rho(k)-1}o, \bar{\omega}_{\rho(k)+1}o)_{\bar{\omega}_{\rho(k)}o} &= (h^{-2}o, g_{\rho(k)+1}o)_o < D(K, \epsilon). \end{aligned}$$

The first item follows from Property (7) of Schottky sets, and the second item follows from $h \in \tilde{S}_k(\mathbf{g})$; hence the claim. In particular, Corollary 2.7 implies $(\bar{\omega}_{\rho(k)}o, \bar{\omega}_n^i\bar{\omega}_{\rho(k)}o)_{\bar{\omega}_n^{i-1}\bar{\omega}_{\rho(k)}o} < F(K, \epsilon)$ for each $i \geq 1$ and

$$\begin{aligned} \frac{1}{i}d(\bar{\omega}_{\rho(k)}o, \bar{\omega}_n^i\bar{\omega}_{\rho(k)}o) &\geq d(\bar{\omega}_{\rho(k)}o, \bar{\omega}_n\bar{\omega}_{\rho(k)}o) - F(K, \epsilon) \\ &\geq [(o, \omega_n o) - d(o, \bar{\omega}_{\rho(k)}o) - d(\bar{\omega}_n o, \bar{\omega}_n\bar{\omega}_{\rho(k)}o)] - K'n/1000 \\ &\geq D_t - 2D_f - K'n/500 \geq K'n/12. \end{aligned}$$

By sending $i \rightarrow \infty$, we deduce that $\tau(\bar{\omega}_n) \geq K'n/12$. \square

5. PROOF OF THEOREM A

Let us now prove Theorem A.

Proof. Given a finite set $S' \subseteq G$, we take a (K, K', ϵ) -Schottky subset S_1 of G such that $K' > 2L(K\epsilon) + 5000F(K, \epsilon)$ and a finite symmetric generating set $S \supseteq S'$ such that $e \in S$ and S is nicely populated by $S_1^{(2)} \cup S_1^{(-2)}$, by using Lemma 2.16.

As before, we consider the random walk on G generated by the uniform measure μ_S on S . We first claim

$$\mathbb{P}(\#P_n^* \leq n/5) \leq \frac{n}{3} \cdot 0.9^n + 0.9886^n.$$

for large n . The first term of the RHS is for the event of trajectories \mathbf{g} with $\#\Theta(\mathbf{g}) \leq n/3$, whose probability is at most

$$\sum_{i=0}^{\lfloor n/3 \rfloor} \binom{\lfloor n/2 \rfloor}{i} \cdot 0.99^i \cdot 0.01^{\lfloor n/2 \rfloor - i} \leq \sum_{i=0}^{\lfloor n/3 \rfloor} \binom{\lfloor n/2 \rfloor}{\lfloor n/3 \rfloor} \cdot 0.99^{\lfloor n/3 \rfloor} \cdot 0.01^{n/6-1} \leq \frac{n}{3} \cdot 0.9^n.$$

Here the final inequality is deduced from the fact that

$$\binom{3(m+1)}{2(m+1)} \cdot 0.01^{m+1} = \binom{3m}{2m} \cdot \frac{(3m+3)(3m+2)(3m+1)}{(m+1)(2m+1)(2m+2)} \cdot 0.01^{m+1} \leq \binom{3m}{2m} \cdot 0.01^m \cdot 0.07$$

for sufficiently large m , and that $0.07^{1/6} < 0.9$.

The second term is an estimation for the sum of m i.i.d. RVs X_i of the distribution in Equation 3.8. We have

$$\mathbb{P} \left(\sum_{i=1}^m X_i < n/5 \right) \cdot 1.4^{-n/5} \leq \mathbb{E} \left[1.4^{-\sum_{i=1}^m X_i} \right] = \prod_{i=1}^m \mathbb{E} \left[1.4^{-X_i} \right] = \left(\frac{1188}{1505} \right)^m$$

and the desired estimate follows for $m \geq n/3$.

We now consider an equivalence class $\mathcal{F}(\mathbf{g})$ of $\mathbf{g} \in G^n$ such that $\#P_n(\mathbf{g}) \geq n/5$ and $D_f(\mathbf{g}) \leq D_b(\mathbf{g})$. For $h_i \in \tilde{S}_i(\mathbf{g})$ and $k = 1, \dots, \lfloor n/24 \rfloor$, Lemma 4.1 gives the sets

$$\begin{aligned} A_k(\mathbf{g}, \{h_i, h_{\lfloor n/12 \rfloor - i}\}_{i=1}^{k-1}) &\subseteq \tilde{S}_{i(k)}(\mathbf{g}), \\ A'_k(\mathbf{g}, \{h_i, h_{\lfloor n/12 \rfloor - i}\}_{i=1}^{k-1}) &\subseteq \tilde{S}_{i(\lfloor n/12 \rfloor - k)}(\mathbf{g}) \end{aligned}$$

with cardinality at most 2, such that the following holds: $\bar{g} \in \mathcal{F}(\mathbf{g})$ satisfies $\tau(\bar{\omega}_n) < K'n/12$ only if $\bar{g}_{\rho(k)} \in A_k(\mathbf{g}, \{\bar{g}_{\rho(i)}, \bar{g}_{\rho(\lfloor n/12 \rfloor - i)}\}_{i=1}^{k-1})$ and $\bar{g}_{\rho(\lfloor n/12 \rfloor - k)} \in A'_k(\mathbf{g}, \{\bar{g}_{\rho(i)}, \bar{g}_{\rho(\lfloor n/12 \rfloor - i)}\}_{i=1}^{k-1})$ for each $k = 1, \dots, \lfloor n/24 \rfloor$. This implies that

$$\begin{aligned} &\mathbb{P} \left(\tau(\bar{\omega}_n) \leq K'n/12 \mid \bar{\mathbf{g}} \in \mathcal{F}(\mathbf{g}) \right) \\ &\leq \prod_{l=1}^{\lfloor n/24 \rfloor} \frac{(\#\tilde{S}_l)(\#\tilde{S}_{\lfloor n/12 \rfloor - l}) - (\#\tilde{S}_l - 2)(\#\tilde{S}_{\lfloor n/12 \rfloor - l} - 2)}{(\#\tilde{S}_l)(\#\tilde{S}_{\lfloor n/20 \rfloor - l})} \\ &\leq \prod_{l=1}^{\lfloor n/24 \rfloor} \left[\frac{2}{\#\tilde{S}_l} + \frac{2}{\#\tilde{S}_{\lfloor n/12 \rfloor - l}} \right] \\ &\leq \prod_{l=1}^{\lfloor n/24 \rfloor} \left[\frac{2}{0.99\#S} + \frac{2}{0.99\#S} \right] = (0.2475\#S)^{-\lfloor n/24 \rfloor}. \end{aligned}$$

A similar argument leads to the same conclusion for $\mathcal{F}(\mathbf{g})$ where $D_f(\mathbf{g}) > D_b(\mathbf{g})$. Since these $\mathcal{F}(\mathbf{g})$'s partition $\{\#P_n \geq n/5\}$, we conclude that the number of n -step trajectories ω such that $\tau(\omega_n) \leq K'n/12$ is bounded by

$$(\#S)^n \cdot [0.91^n + 0.9886^n + (0.2475\#S)^{-n/24}] \leq 0.999^n \cdot (0.99\#S)^n$$

for sufficiently large n . Since any mapping class in $B_S(n)$ is obtained from an n -step trajectory, we conclude that the number of mapping classes in $B_S(n)$ with translation length less than $K'n/12$ is bounded by $0.999^n \cdot (0.99\#S)^n$.

Meanwhile, the set

$$S_{Schottky} = \{(a_1, \dots, a_n) : a_i \in S_1 \cup S_1^{-1}, a_i \neq a_{i+1}^{-1}\}$$

is composed of at least $(\#S_0 - 1)^n \geq (0.99\#S)^n$ sequences. We claim that if $(a_1, \dots, a_n), (b_1, \dots, b_n)$ are distinct sequences in $S_{Schottky}$, then $a_1^2 \cdots a_n^2, b_1^2 \cdots b_n^2$ are distinct elements in $B_S(n)$. Indeed, the sequence

$$(a_n^{-1}, a_n^{-1} \cdots, a_1^{-1}, a_1^{-1}, b_1, b_1, \dots, b_n, b_n)$$

will not completely cancel out and their product will not become an identity by Lemma 2.13. Hence, we have at least $(0.99\#S)^n$ distinct elements in $B_S(n)$. We thus finally have

$$\frac{\#\{g \in B_S(n) : \tau_X(g) \leq K'n/12\}}{\#B_S(n)} \leq 0.999^n$$

for large n , finishing the proof. \square

APPENDIX A. REMARK ON LEMMA 2.12

In this section, we prove our claim during the proof of Lemma 2.12. We first recall the following lemma:

Fact A.1 ([Cho21, Lemma 3.12]). *For each $F, \epsilon > 0$, there exists $H, L > F$ that satisfies the following condition. If x, y, z, p_1, p_2 in X satisfy that:*

- (1) $[p_1, p_2]$ is ϵ -thick and longer than L ,
- (2) $[x, y]$ is F -witnessed by $[p_1, p_2]$, and
- (3) $(x, z)_y \geq d(p_1, y) - F$,

then $[z, y]$ is H -witnessed by $[p_1, p_2]$.

Recall that we have fixed $o \in X$ and independent loxodromics $a, b \in G$. By Lemma 4.3 and 4.4 of [Cho21], there exists $\epsilon_0, C_0 > 0$ such that the following hold:

- (1) $[o, a^i o], [o, b^i o]$ are ϵ_0 -thick for $i \in \mathbb{Z}$, and
- (2) $(\phi^i o, \psi^j o)_o < C_0$ for $i, j > 0$ and $\phi, \psi \in \{a, b, a^{-1}, b^{-1}\}$ such that $\phi \neq \psi$.

We then define:

- $D_0 = D(C = C_0, \epsilon_0)$ as in Lemma 2.9;
- $E_0 = E(D = D_0, \epsilon_0)$, $L_0 = L(D = D_0, \epsilon_0)$ as in Fact 2.5;
- $F_0 = F(E = E_0, \epsilon_0)$, $L_1 = L(E = E_0, \epsilon_0)$ as in Fact 2.6;
- $H_0 = H(F = F_0, \epsilon_0)$, $L_2 = L(F = F_0, \epsilon_0)$ as in Fact A.1;
- $H_1 = H(F = H_0, \epsilon_0)$, $L_3 = L(F = H_0, \epsilon_0)$ as in Fact A.1;
- $F_1 = 2F_0 + H_1 + \delta + 1$;
- $F_2 = F(E = H_0, \epsilon_0)$ and $L_4 = L(E = H_0, \epsilon_0)$ in Fact 2.6;
- $L_5 = \max(L_0, L_1, L_2, L_3, L_4, 2F_0 + F_1 + 2F_2)$.

There exists N_0 such that $d(o, \phi^N o) > L_5$ for all $\phi \in \{a, b, a^{-1}, b^{-1}\}$ and $N > N_0$. Let us fix $N > N_0$. We now consider a sequence $\{\phi_i\}$ in $\{a, b, a^{-1}, b^{-1}\}$ such that $\phi_i \neq \phi_{i+1}^{-1}$. Then we observe the following:

- (1) $(\phi_i^{-N} o, \phi_{i+1}^N o)_o < C_0$ by the assumption $\phi_i \neq \phi_{i+1}^{-1}$.
- (2) $[o, \phi_i^N \phi_{i+1}^N o]$ is D_0 -witnessed by $[o, \phi_i^N o]$ and $[\phi_i^N o, \phi_{i+1}^N \phi_i^N o]$ by Lemma 2.9.
- (3) $[o, \phi_i^N o]$ are ϵ -thick and longer than L_5 .

Then as in Corollary 2.7, Lemma 2.5 and Lemma 2.6 imply that $[o, \phi_1^{2N} \cdots \phi_n^{2N} o]$ is F_0 -witnessed by ϵ_0 -thick segments

$$[o, \phi_1^N o], [\phi_1^N o, \phi_1^N \phi_2^N o], \dots, [\phi_1^N \cdots \phi_{n-1}^N o, \phi_1^N \cdots \phi_n^N o].$$

Consequently, $[o, \phi_1^N \cdots \phi_n^N o]$ is ϵ -thick for $\epsilon = \epsilon_0 e^{-8F_0}$. It is also clear that

$$(o, \phi_1^N \cdots \phi_n^N o)_{\phi_1^N \cdots \phi_m^N o} < F_0 \quad (0 \leq m \leq n)$$

and

$$(A.1) \quad d(o, \phi_1^N \cdots \phi_n^N o) \geq d(o, \phi_1^N \cdots \phi_{n-1}^N o) + L_1 - 2F_0 \geq d(o, \phi_1^N \cdots \phi_{n-1}^N o) + F_1.$$

We now define

$$\begin{aligned} S_{n,N} &:= \{g_1, \dots, g_{2^n}\} = \{\phi_1^{2N} \cdots \phi_n^{2N} : \phi_i \in \{a, b\}\}, \\ V(g_i^\pm) &:= \{x \in X : (x, g_i^{\pm 2} o)_o \geq d(o, g_i^{\pm 1} o) - F_1\}, \\ V'(g_i^\pm) &:= \{x \in X : (x, g_i^{\pm 2} o)_o \geq d(o, g_i^{\pm 1} o)\}. \end{aligned}$$

Our first claim is that $V(g_1^+), \dots, V(g_{2^n}^+), V(g_1^-), \dots, V(g_{2^n}^-)$ are all disjoint. To show this, let $(\phi_i)_{i=1}^n, (\psi_i)_{i=1}^n$ be distinct sequences in $\{a, b\}^n \cup \{a^{-1}, b^{-1}\}^n$ and $\Phi = \phi_1^{2N} \cdots \phi_n^{2N}, \Psi = \psi_1^{2N} \cdots \psi_n^{2N}$. Let $t = \min\{1 \leq i \leq 10 : \phi_i \neq \psi_i\}$ and $w = \phi_1^{2N} \cdots \phi_{t-1}^{2N}$. If some x belongs to $V(\Phi) \cap V(\Psi)$, then

$$(x, \Phi^2 o)_o \geq d(o, \Phi o) - F_1 \geq d(o, \phi_1^{2N} \cdots \phi_{t-1}^{2N} \phi_t^N o) = d(o, w \phi_t^N o)$$

by Inequality A.1. Similarly, we have $(x, \Psi^2 o)_o \geq d(o, w \psi_t^N o)$. Since $[o, \Phi^2 o]$ is F_0 -witnessed by $[wo, w \phi_t^N o]$, Fact A.1 asserts that $[o, x]$ is H_0 -witnessed by $[wo, w \phi_t^N o]$. Similarly, $[o, x]$ is H_0 -witnessed by $[wo, w \psi_t^N o]$. Since $(\phi_t o, \psi_t o)_o < C_0 < H_0$, Lemma 2.6 implies that $[x, x]$ is F_2 -witnessed by $[wo, w \psi_t^N o]$, whose length is at least $L_5 > 2F_2$: such x does not exist.

The next claim is that if $x \notin V(g_i^-)$, then $g_i^2 x \in V'(g_i)$. Indeed, we know that $(o, g_i^2 o)_{g_i o} \leq F_0 \leq F_1/2$ and

$$\begin{aligned} (g_i^2 x, g_i^2 o)_o &= (x, o)_{g_i^{-2} o} = d(o, g_i^{-2} o) - (x, g_i^{-2} o)_o \\ &\geq d(o, g_i^2 o) - d(o, g_i o) + F_1 \geq d(o, g_i o). \end{aligned}$$

Since $V'(g_i) \subseteq V(g_i)$ and $V(g_i) \cap V(g_i^-) = \emptyset$, we can iterate this to deduce $g_i^{2k} x \in V'(g_i)$ for $k > 0$. Similarly, if $x \notin V(g_i)$, then $g_i^{-2k} x \in V'(g_i^-)$ for $k > 0$.

Now let $x, y \in X$. Since $\{V(g_i^+), V(g_i^-)\}$ are disjoint, $y \in V(g_i^-)$ for at most one $g_i \in S_{n,N}$ and $x \in V(g_j^+)$ for at most one $g_j \in S_{n,N}$. Suppose $s = \phi_1^{2N} \cdots \phi_n^{2N} \in S_{n,N}$ is neither of them, and let $k > 0$. We then have

$$(x, s^2 o)_o < d(o, so) - F_1, \quad (s^{2k} y, s^2 o)_o \geq d(o, so).$$

As $[o, s^2 o]$ is F_0 -witnessed by $[s\phi_n^{-N} o, so]$, Fact ?? implies that $[o, s^{2k} y]$ is H_0 -witnessed by $[s\phi_n^{-N} o, so]$. Now if we suppose that $(x, s^{2k} y)_o > d(o, so)$, then $[o, x]$ is also H_1 -witnessed by $[s\phi_n^{-N} o, so]$, again by Fact ?. Meanwhile:

- if X is a δ -hyperbolic space, we deduce that

$$\begin{aligned} (x, s^2 o)_o &\geq \min\{(x, so)_o, (so, s^2 o)_o\} - \delta \\ &= d(o, so) - \max\{(x, o)_{so}, (s^2 o, o)_{so}\} - \delta \\ &\geq d(o, so) - (G_1 + F_0 + \delta) \geq d(o, so) - F_1. \end{aligned}$$

- if $X = \mathcal{T}(\Sigma)$, then we deduce that

$$\begin{aligned} (x, s^2 o)_o &\geq d(o, so) - d(so, [o, x]) - d(so, [o, s^2 o]) \\ &\geq d(o, so) - (G_1 + F_0) \geq d(o, so) - F_1. \end{aligned}$$

In either case we obtain a contradiction. Hence, $(x, s^{2k} y)_o \leq d(o, so)$.

Similarly, if $s \neq g_i$ such that $y \in V(g_i^+)$ and $s \neq g_j$ such that $x \in V(g_j^-)$, then $(x, s^{-2k} y)_o \leq d(o, s^{-1} o)$ for all $k > 0$. Finally, note that $y = o$ cannot belong to any of $V(g_j^\pm)$ since $d(o, g_j^{\pm 1} o) \geq L_5 > F_1$ for any $g_j \in S_{n,N}$. This settles the desired claim.

APPENDIX B. SKETCH OF THE PROOF OF PROPOSITION 1.3

We borrow definitions and notations in [GTT18].

In [GTT18], the authors consider the automatic structure of G , a directed graph Γ that records exactly one geodesic between e and g for each $g \in G$. Hence, the vertex set of its universal cover $\tilde{\Gamma}$ and G has 1-1 correspondence. Let LG be the set of vertices with large growth. For $g \in G$ and $0 < \epsilon < 1$, we denote by \hat{g}_ϵ the element along the path from e to g at distance ϵn from e . Then for any $0 < \epsilon < 1$, the ratio

$$(B.1) \quad \frac{\#\{g \in \partial B_S(n) : \hat{g}_\epsilon \notin LN\}}{\#\partial B_S(n)}$$

decays exponentially (cf. [GTT18, Proposition 2.5]).

The authors then construct a Markov chain on Γ whose n -step distribution is denoted by \mathbb{P}^n . There exists $c > 1$ such that for any $A \subseteq \tilde{\Gamma}$, the proportion of $A \cap LG$ in $\partial B_S(n)$ is at least $(1/c) \mathbb{P}^n(A)$ and at most $c \mathbb{P}^n(A)$.

We now denote by \mathcal{R} the set of recurrent vertices. For $v \in \mathcal{R}$, the loop semigroup Γ_v associated to v is nonelementary, i.e., there exist independent loxodromics $a_v, b_v \in \Gamma_v$ [GTT18, Corollary 6.11]. Let us now condition on

the paths growing from v . Let $n(k, v, \omega)$ be the k -th return time to v and $T_v = \mathbb{E}_v n(1, v, \omega)$. Then for each $\epsilon > 0$,

$$(B.2) \quad \mathbb{P}_v \left\{ \left| \frac{n(k, v, \omega)}{k} - T_v \right| > \epsilon \right\}$$

is exponentially decaying as $k \rightarrow \infty$ (cf. [GTT18, Lemma 6.13]). For each n , we also define the last return time $\tilde{n}(\omega) = \max(\{n(k, v, \omega) : k \in \mathbb{N}\} \cap \{1, \dots, n\})$ to v . Then for each $\epsilon > 0$,

$$(B.3) \quad \mathbb{P}_v^n \left\{ \frac{n - \tilde{n}(\omega)}{n} > \epsilon \right\}$$

decays exponentially.

We now strengthen Theorem 6.14 of [GTT18]. For each $\epsilon > 0$,

$$(B.4) \quad \mathbb{P}_v \left\{ \left| \frac{d(\omega_{n(k,v)} o, o)}{k} - l_v \right| > \epsilon \right\}$$

decays exponentially as $k \rightarrow \infty$, since μ_v actually has finite exponential moment. The proof for deviations from above can be found in [BMSS20]. For deviations from below, [BMSS20] and [Gou21] deals with the case that X is Gromov hyperbolic. When X is the Teichmüller space, one can use Choi's modification of Gouëzel's construction in [Cho21]. Now together with the control on quantities B.2 and B.3, we obtain the following: for any $\epsilon > 0$,

$$(B.5) \quad \mathbb{P}_v^n \left\{ \left| \frac{d(\omega_n o, o)}{n} - \frac{l_v}{T_v} \right| > \epsilon \right\}$$

decays exponentially. Now the proof of [GTT18, Theorem 6.14] shows that l_v/T_v is uniform for all $v \in \mathcal{R}$, which we denote by λ . Since the arrival time at \mathcal{R} (beginning at e) also has finite exponential moment, we conclude that

$$(B.6) \quad \mathbb{P}_e^n \left\{ \left| \frac{d(\omega_n o, o)}{n} - \lambda \right| > \epsilon \right\}$$

decays exponentially. By combining this with the decay of B.1, we deduce that

$$\frac{\#\{g \in \partial B_S(n) : \left| \frac{d(o, go)}{n} - \lambda \right| > \epsilon\}}{\#\partial B_S(n)}$$

decays exponentially (cf. [GTT18, Theorem 7.3]).

We now need to discuss translation lengths instead of displacements. For each recurrent component C of Γ , we pick $v = v_C \in C$ and take a Schottky set as a subset of $\{w = g_1 \cdots g_n : g_i = a_v \text{ or } b_v\}$. We now consider the loop random walk generated by μ_v : recall the decomposition of the random walk into usual steps and ‘‘Schottky steps’’ for the pivot construction in [Gou21] and [Cho21].

Until step $T_v n/4$ in the loop random walk, we have Kn slots for Schottky steps for some $K > 0$ outside an event of exponentially decaying probability. Moreover, $T_v n/4$ steps in the loop random walk occur before step $n/2$ in the Markov process based at v_C , outside an event of exponentially

decaying probability (quantity B.2). Finally, the Markov process beginning from e arrives at $\{v_C : C \text{ recurrent}\}$ within step $n/2$ outside an event of exponentially decaying probability.

In summary, giving up an event of exponentially decaying probability, a random path in the Markov process has at least Kn slots for Schottky loops based at some v_C . By pivoting the choice of Schottky loops at these slots, we can guarantee at least $K'n$ eventual pivots until step n for some $K' > 0$, outside an event of exponentially decaying probability.

Given these results, it now suffices to focus on the elements g such that:

- (1) $\check{g}_{1-\epsilon} \in LG$ and $d(o, \check{g}_{1-\epsilon}o) \geq (1 - 2\epsilon)\lambda n$,
- (2) $d(o, \check{g}_\epsilon o) \leq 2\epsilon\lambda n$, and
- (3) the subpath $[e, \check{g}_\epsilon]$ possesses at least $K'\epsilon n$ pivots for $[e, g]$.

We then consider the equivalence class by pivoting at the first $K'\epsilon n$ pivots. By early pivoting, one can show that only few elements inside the equivalence class satisfy $\tau_X(g) \leq (1 - 4\epsilon - MK')n$, for some suitable $M > 0$. By modulating ϵ and K' , we establish the desired result.

REFERENCES

- [BCK21] Hyungryl Baik, Inhyeok Choi, and Dongryul M Kim. Linear growth of translation lengths of random isometries on Gromov hyperbolic spaces and Teichmüller spaces. *arXiv preprint arXiv:2103.13616*, 2021.
- [BMSS20] Adrien Boulanger, Pierre Mathieu, Çağrı Sert, and Alessandro Sisto. Large deviations for random walks on hyperbolic spaces. *arXiv preprint arXiv:2008.02709*, 2020.
- [Cho21] Inhyeok Choi. Central limit theorem and geodesic tracking on hyperbolic spaces and Teichmüller spaces. *arXiv preprint arXiv:2106.13017*, 2021.
- [CW18] María Cumplido and Bert Wiest. A positive proportion of elements of mapping class groups is pseudo-Anosov. *Bulletin of the London mathematical society*, 50(3):390–394, 2018.
- [EST20] Viveka Erlandsson, Juan Souto, and Jing Tao. Genericity of pseudo-Anosov mapping classes, when seen as mapping classes. *L'Enseignement Mathématique*, 66(3/4):419–439, 2020.
- [Far06] Benson Farb. Some problems on mapping class groups and moduli space. In Benson Farb, editor, *Problems on mapping class groups and related topics*, volume 74 of *Proceedings of Symposia in Pure Mathematics*, pages 11–46. American Mathematical Society, 2006.
- [Gou21] Sébastien Gouëzel. Exponential bounds for random walks on hyperbolic spaces without moment conditions. *arXiv preprint arXiv:2102.01408*, 2021.
- [GTT18] Ilya Gekhtman, Samuel J. Taylor, and Giulio Tiozzo. Counting loxodromics for hyperbolic actions. *Journal of Topology*, 11(2):379–419, 2018.
- [GTT20a] Ilya Gekhtman, Samuel J. Taylor, and Giulio Tiozzo. Central limit theorem for counting measures in coarse negative curvature. *arXiv preprint arXiv:2004.13084*, 2020.
- [GTT20b] Ilya Gekhtman, Samuel J. Taylor, and Giulio Tiozzo. Counting loxodromics in graph products and relatively hyperbolic groups. *Israel Journal of Mathematics*, 237:311–371, 2020.
- [Mah12] Joseph Maher. Exponential decay in the mapping class group. *Journal of the London Mathematical Society*, 86(2):366–386, 2012.

- [MT18] Joseph Maher and Giulio Tiozzo. Random walks on weakly hyperbolic groups. *Journal für die reine und angewandte Mathematik (Crelle's Journal)*, 2018(742):187–239, 2018.
- [Raf14] Kasra Rafi. Hyperbolicity in Teichmüller space. *Geometry & Topology*, 18(5):3025–3053, 2014.
- [Yan20] Wen-yuan Yang. Genericity of contracting elements in groups. *Mathematische Annalen*, 376:823–861, 2020.

DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, 291 DAEHAK-RO YUSEONG-GU,
DAEJEON, 34141, SOUTH KOREA

Email address: inhyeokchoi@kaist.ac.kr