

Long-time shadow limit for reaction-diffusion-ODE systems

Chris Kowall^a, Anna Marciniak-Czochra^b, Andro Mikelić^{c,1}

^a*Institute of Applied Mathematics and IWR, Heidelberg University, Im Neuenheimer Feld 205, 69120 Heidelberg, Germany*

^b*Institute of Applied Mathematics, IWR and BIOQUANT, Heidelberg University, Im Neuenheimer Feld 205, 69120 Heidelberg, Germany*

^c*Univ Lyon, Université Claude Bernard Lyon 1, CNRS UMR 5208, Institut Camille Jordan, 43 blvd. du 11 novembre 1918, F-69622 Villeurbanne cedex, France*

Abstract

Shadow systems are an approximation of reaction-diffusion-type problems obtained in the infinite diffusion coefficient limit. They allow reducing complexity of the system and hence facilitate its analysis. The quality of approximation can be considered in three time regimes: (i) short-time intervals taking account for the initial time layer, (ii) long-time intervals scaling with the diffusion coefficient and tending to infinity for diffusion tending to infinity, and (iii) asymptotic state for times up to $T = \infty$. In this paper we focus on uniform error estimates in the long-time case. Using linearization at a time-dependent shadow solution, we derive sufficient conditions for control of the errors. The employed methods are cut-off techniques and L^p -estimates combined with stability conditions for the linearized shadow system. Additionally, we show that the global-in-time extension of the uniform error estimates may fail without stronger assumptions on the model linearization. The approach is presented on example of reaction-diffusion equations coupled to ordinary differential equations (ODEs) and is also applied to a classical reaction-diffusion system. The results are illustrated by examples showing necessity and applicability of the established conditions.

Keywords: Shadow limit, reaction-diffusion equation, model reduction, uniform long-time approximation

1. Introduction

Reaction-diffusion systems with zero flux boundary conditions and a large diffusion coefficient can be studied by means of their model reduction as diffusion tends to infinity. The resulting shadow limit facilitates analysis of the spatio-temporal evolution of solutions to reaction-diffusion systems. Originally considered for a system of two coupled reaction-diffusion equations, starting from [15, 27, 41, 49], the shadow limit has attracted a considerable interest also in the case of reaction-diffusion-ODE systems [3, 22, 32, 37, 39]. Many ecological or biological applications lead to models coupling reaction-diffusion equations with ordinary differential equations, e.g., [25, 28, 36, 38]. The non-diffusing components may lead to a lower spatial regularity of model solutions and even loss of stability of regular stationary solutions [19, 30]. In this paper, we focus on a rigorous proof of a large diffusion limit for such models, pursuing the work [32] and extending the analysis to nonlinear models on long-time intervals. Our results provide understanding of the long-term dynamics of the reaction-diffusion-ODE system from results obtained for its shadow limit.

We consider reaction-diffusion-ODE systems, but as shown in Section 4, the presented theory also applies to the classical models. We assume that $\Omega \subset \mathbb{R}^n$ is a bounded domain with Lipschitz boundary $\partial\Omega \in C^{0,1}$

Email addresses: kowall@math.uni-heidelberg.de (Chris Kowall), anna.marciniak@iwr.uni-heidelberg.de (Anna Marciniak-Czochra)

¹Dedicated to the Memory of Professor Andro Mikelić, deceased on November 28, 2020.

and consider the following quasilinear system

$$\begin{aligned} \frac{\partial \mathbf{u}_\varepsilon}{\partial t} &= \mathbf{f}(\mathbf{u}_\varepsilon, v_\varepsilon, x, t) & \text{in } \Omega_T := \Omega \times (0, T), & \quad \mathbf{u}_\varepsilon(\cdot, 0) = \mathbf{u}^0 & \text{in } \Omega, \\ \frac{\partial v_\varepsilon}{\partial t} - \frac{1}{\varepsilon} \Delta v_\varepsilon &= g(\mathbf{u}_\varepsilon, v_\varepsilon, x, t) & \text{in } \Omega_T, & \quad v_\varepsilon(\cdot, 0) = v^0 & \text{in } \Omega, \\ \frac{\partial v_\varepsilon}{\partial \nu} &= 0 & \text{on } \partial\Omega \times (0, T), & \end{aligned} \quad (1.1)$$

where $\varepsilon > 0$ is a small parameter. This problem consists of a non-diffusing, vector-valued component denoted by \mathbf{u}_ε and a diffusing, scalar component v_ε satisfying zero-flux boundary conditions.

The shadow limit reduction of system (1.1) yields the following system of integro-differential equations

$$\begin{aligned} \frac{\partial \mathbf{u}}{\partial t} &= \mathbf{f}(\mathbf{u}, v, x, t) & \text{in } \Omega_T, & \quad \mathbf{u}(\cdot, 0) = \mathbf{u}^0 & \text{in } \Omega, \\ \frac{dv}{dt} &= \langle g(\mathbf{u}(\cdot, t), v(t), \cdot, t) \rangle_\Omega & \text{in } (0, T), & \quad v(0) = \langle v^0 \rangle_\Omega, \end{aligned} \quad (1.2)$$

where we abbreviate the spatial mean value for $z \in L^1(\Omega)$ by

$$\langle z \rangle_\Omega = \frac{1}{|\Omega|} \int_\Omega z(x) \, dx.$$

For reaction-diffusion-ODE models, the shadow limit has been studied in [3] and independently in [32, 39], showing an error estimate in terms of the large diffusion parameter $1/\varepsilon$. In both approaches, the obtained estimates depend exponentially on the length of the time interval, see [32] for details. This means that for long time intervals the error estimate deteriorates significantly and no conclusion on the long-time behavior of solutions to system (1.1) can be drawn from the behavior of its shadow limit reduction (1.2).

The current paper is devoted to analysis of the uniform shadow-limit approximation for long-time intervals, i.e., time intervals $(0, T)$ of the length $T \sim \varepsilon^{-\ell}$ for some $\ell > 0$. We provide error estimates in terms of powers of the inverse ε of the large diffusion parameter, which are uniform with respect to space and time. Departing from the analysis of a linear reaction-diffusion-ODE system with space-independent coefficients, considered in [32], we approach the nonlinear problem using linearization of the system of errors at the shadow solution. This step requires extending the analysis of [32] to a system with space-dependent coefficients and formulating sufficient stability conditions.

The presented analysis involves several technical challenges arising from low regularity of model parameters, initial data or the considered domain. We consider mild solutions given by an implicit integral equation of the reaction-diffusion-type system and its shadow limit. The spatial mean values in the integro-differential system (1.2) implies that the shadow system is a singular limit of the partly diffusive system (1.1). Hence, in the error estimates, we involve a correction term ψ_ε . This term includes the initial layer which originates from different initial values v^0 and $\langle v^0 \rangle_\Omega$. Moreover, there is a discrepancy between nonlinearities of both systems, which we also include in this correction term ψ_ε to streamline the analysis. When comparing both solutions, estimates for the error $\mathbf{u}_\varepsilon - \mathbf{u}$ and $v_\varepsilon - v - \psi_\varepsilon$ are established with respect to the $L^\infty(\Omega_T)$ norm where the correction term ψ_ε is negligible for larger times. The choice of $L^\infty(\Omega_T)$ is suitable for bounded, discontinuous solutions of a wide range of nonlinear problems which can be solved using the method of Rothe [46]. At the same time, boundary regularity of Ω can be relaxed to Lipschitz regularity which is necessary for validity of Sobolev embeddings, existence of the heat semigroup and a proper notion of a boundary condition.

The system of errors is linearized at a globally defined, uniformly bounded shadow limit. To control the growth of the nonlinear parts, we apply a cut-off technique, similar to [39]. Since the cut-off does not affect the nonlinear part in a small neighborhood of $\mathbf{0}$, the truncation allows obtaining uniform estimates

of the truncated errors. Showing invariance of those neighborhoods under certain stability assumptions, for fixed times T and sufficiently large diffusion coefficients $D \geq D(T)$, provides estimates for the original errors. To control the truncated errors, we formulate stability conditions on a linearized shadow system. If the evolutionary system is uniformly stable with respect to $L^\infty(\Omega)^m \times \mathbb{R}$, we obtain error estimates with explicit dependence on the time interval length T . Stability implies estimates that are valid on Ω_T for $T \sim \varepsilon^{-\ell}$ and some $0 < \ell < 1$. Unfortunately, as examples in [32] show, accuracy of the approximation for transient states does not imply a valuable asymptotic approximation as $T \rightarrow \infty$. However, assuming uniform exponential stability of the linearized shadow system yields global error estimates on $\Omega \times \mathbb{R}_{\geq 0}$. The latter stability condition can also be considered in $L^p(\Omega)^m \times \mathbb{R}$ for sufficiently large $p < \infty$. In such case, parabolic L^p estimates are employed to conclude on L^∞ bounds for the diffusive component. An additional uniform stability of the ODE subsystem in L^∞ as in [32] yields an estimate for the non-diffusive components.

It is usually difficult to check directly the obtained stability conditions at the shadow limit solutions. However, in the stationary case, stability properties of the linearization can be deduced from the spectrum of the corresponding linear operator. This is a consequence of the well-known spectral mapping theorem for analytic semigroups. The spectrum of the linearized shadow operator is characterized for bounded stationary solutions. For analysis of the spectrum, which does not need to be purely discrete, we apply a spectral decomposition for block operator matrices [2]. This is based on properties of bounded multiplication operators, which are induced by the ODE subsystem on $L^p(\Omega)^m$ for $1 \leq p \leq \infty$. Moreover, the semigroup framework in this work allows applying the uniform shadow limit approximation to classical reaction-diffusion systems as well. In this context, we also study dissipativity conditions for the linearized evolution system which imply the stability conditions.

The paper is organized as follows. We introduce the notion of a mild solution for both problems (1.1) and (1.2), and provide preliminary results in Section 2. In Section 3, we prove uniform error estimates for long-time intervals. The main results are given in Theorem 3.6 and Theorem 3.11. Section 4 is devoted to classical reaction-diffusion systems, where also dissipativity conditions are considered. In Section 5, a linear model and a predator-prey system exemplify the results. Basic properties concerning the heat equation and the spectral analysis of a linear shadow operator is deferred to the Appendix.

2. Preliminary results

2.1. Reaction-diffusion-ODE problem

Let us recall short-time well-posedness of the reaction-diffusion-ODE system (1.1) for given nonlinearities

$$\mathbf{f} : \mathbb{R}^m \times \mathbb{R} \times \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}^m \quad \text{and} \quad g : \mathbb{R}^m \times \mathbb{R} \times \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}.$$

Existence of a solution $\mathbf{u}_\varepsilon : \Omega_T \rightarrow \mathbb{R}^m$ and $v_\varepsilon : \Omega_T \rightarrow \mathbb{R}$ for a short time follows from standard theory [34, 45, 46]. Since we want to impose only low regularity, we recollect some definitions and the main idea of the proof of [46, Part II, Theorem 1]. Therefore, we rewrite system (1.1) as a system of $m + 1$ reaction-diffusion-type equations:

$$\frac{\partial \Psi}{\partial t} - \mathbf{D}_\varepsilon \Delta \Psi = \mathbf{h}(\Psi, x, t) \quad \text{in } \Omega_T, \quad \Psi(\cdot, 0) = (\mathbf{u}^0, v^0) \quad \text{in } \Omega, \quad \frac{\partial \Psi_{m+1}}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times (0, T) \quad (2.1)$$

Here, $\Psi = (\mathbf{u}_\varepsilon, v_\varepsilon)$, $\mathbf{h} = (\mathbf{f}, g)$, and $\mathbf{D}_\varepsilon = \text{diag}(0, \dots, 0, \varepsilon^{-1}) \in \mathbb{R}_{\geq 0}^{(m+1) \times (m+1)}$ is a diagonal matrix. If we consider this system for the decoupled case, i.e., for $\mathbf{h} = \mathbf{0}$, the solution v_ε induces an analytic semigroup $(S_\Delta(\tau))_{\tau \in \mathbb{R}_{\geq 0}}$ on $L^p(\Omega)$ for each $p \in (1, \infty)$ by $v_\varepsilon(\cdot, t) = S_\Delta(t/\varepsilon)v^0$ for initial values $v^0 \in L^p(\Omega)$, see Proposition A.1. This semigroup then can be restricted to $L^\infty(\Omega)$ independently of p and we obtain a formal semigroup on $L^\infty(\Omega)$ which is not strongly continuous [46, Part I, Lemma 2]. However, due to the maximum principle, the semigroup $(S_\Delta(\tau))_{\tau \in \mathbb{R}_{\geq 0}}$ is contractive on $L^\infty(\Omega)$, i.e.,

$$\|S_\Delta(\tau)z\|_{L^\infty(\Omega)} \leq \|z\|_{L^\infty(\Omega)} \quad \forall z \in L^\infty(\Omega), \tau \in \mathbb{R}_{\geq 0}.$$

Using this semigroup approach, we define similarly to Rothe, [46, Part II, Definition 2]:

Definition 2.1. Let $0 < T \leq \infty$. A mild solution of problem (2.1) on the interval $[0, T]$ for initial values $\Psi^0 = (\mathbf{u}^0, v^0) \in L^\infty(\Omega)^{m+1}$ is a measurable function $\Psi : \Omega \times [0, T] \rightarrow \mathbb{R}^{m+1}$ satisfying for all $t \in (0, T)$

(i) $\Psi(\cdot, t) \in L^\infty(\Omega)^{m+1}$ and $\sup_{s \in (0, t)} \|\Psi(\cdot, s)\|_{L^\infty(\Omega)^{m+1}} < \infty$,

(ii) the integral representation

$$\Psi(\cdot, t) = \mathbf{S}(t/\varepsilon)\Psi^0 + \int_0^t \mathbf{S}((t-s)/\varepsilon)\mathbf{h}(\Psi(\cdot, s), \cdot, s) ds, \quad (2.2)$$

where the integral is an absolute converging Bochner integral in $L^\infty(\Omega)^{m+1}$ and the semigroup $(\mathbf{S}(\tau))_{\tau \in \mathbb{R}_{\geq 0}}$ on $L^\infty(\Omega)^{m+1}$ has components $S_k(\tau) = I$ for $k = 1, \dots, m$ and $S_{m+1}(\tau) = S_\Delta(\tau)$.

We follow the proof of [46, Part II, Theorem 1] and its assumptions to obtain a mild solution for problem (2.1). Hence, we assume a local Lipschitz condition for \mathbf{h} and bounded initial data.

Assumption 1 (Existence and uniqueness). For every fixed $(\mathbf{u}, v) \in \mathbb{R}^{m+1}$ $\mathbf{h} = (\mathbf{f}, g)$ is a measurable function in the variables $(x, t) \in \bar{\Omega} \times \mathbb{R}_{\geq 0}$. Moreover, for every bounded set $B \subset \mathbb{R}^{m+1} \times \bar{\Omega} \times \mathbb{R}_{\geq 0}$ there exists a constant $L(B) > 0$ such that for all $(\mathbf{u}, v, x, t), (\mathbf{y}, z, x, t) \in B$

$$\begin{aligned} |\mathbf{h}(\mathbf{u}, v, x, t)| &\leq L(B), \\ |\mathbf{h}(\mathbf{u}, v, x, t) - \mathbf{h}(\mathbf{y}, z, x, t)| &\leq L(B) (|\mathbf{u} - \mathbf{y}| + |v - z|). \end{aligned}$$

Assumption 2 (Bounded initial data). Initial values $(\mathbf{u}^0, v^0) \in L^\infty(\Omega)^{m+1}$.

Although [46] uses higher regularity of the boundary $\partial\Omega$, the proof is analog, applying a Picard iteration on $L^\infty(\Omega)^{m+1}$ to the implicit integral representation (2.2) and using the semigroup $(S(\tau))_{\tau \in \mathbb{R}_{\geq 0}}$ for $\partial\Omega \in C^{0,1}$.

Theorem 2.2. Let Assumption 1–2 hold. Then for each $\varepsilon > 0$ there exists a maximal time interval $[0, T_{\max})$ for $T_{\max} = T_{\max}(\varepsilon) > 0$, such that problem (1.1) has a unique mild solution Ψ on $[0, T_{\max})$ satisfying $\Psi = (\mathbf{u}_\varepsilon, v_\varepsilon) \in C([0, T]; L^\infty(\Omega)^m \times L^\infty(\Omega_T))$ for each $T < T_{\max}$. Furthermore, the solutions $(\mathbf{u}_\varepsilon, v_\varepsilon)_{\varepsilon > 0}$ locally exist on the same time interval, i.e., $\inf_{\varepsilon \in (0, 1]} T_{\max}(\varepsilon) > 0$.

Remark 2.3. The integral representation for \mathbf{u}_ε yields higher regularity for the ODE component in time. If we choose more regular initial data, nonlinearities and a more regular boundary, we obtain Hölder continuous solutions that satisfy the equations in the classical sense [46, Part II, Theorem 1].

2.2. Shadow problem

We consider the shadow system (1.2), i.e.,

$$\begin{aligned} \frac{\partial \mathbf{u}}{\partial t} &= \mathbf{f}(\mathbf{u}, v, x, t) & \text{in } \Omega_T, & \quad \mathbf{u}(\cdot, 0) = \mathbf{u}^0 & \text{in } \Omega, \\ \frac{dv}{dt} &= \langle g(\mathbf{u}(\cdot, t), v(t), \cdot, t) \rangle_\Omega & \text{in } (0, T), & \quad v(0) = \langle v^0 \rangle_\Omega. \end{aligned}$$

Applying a Picard iteration yields a unique solution with the integral representation

$$\begin{aligned} \mathbf{u}(\cdot, t) &= \mathbf{u}^0 + \int_0^t \mathbf{f}(\mathbf{u}(\cdot, s), v(s), \cdot, s) ds & \text{in } \Omega, \\ v(t) &= \langle v^0 \rangle_\Omega + \int_0^t \langle g(\mathbf{u}(\cdot, s), v(s), \cdot, s) \rangle_\Omega ds. \end{aligned} \quad (2.3)$$

By Assumption 1–2, [39, Theorem 1] yields a unique local-in-time solution $(\mathbf{u}, v) \in C([0, T]; L^\infty(\Omega)^m \times \mathbb{R})$ for some $T > 0$. Since we are looking for long-time behavior of the model, we additionally assume

Assumption 3 (Bounded shadow limit). The solution (\mathbf{u}, v) of the shadow limit (1.2) is globally defined and uniformly bounded, i.e., $\mathbf{u} \in L^\infty(\Omega \times \mathbb{R}_{\geq 0})^m$ and $v \in L^\infty(\mathbb{R}_{\geq 0})$.

Clearly, using equation (2.3), Assumption 3 implies $(\mathbf{u}, v) \in C(\mathbb{R}_{\geq 0}; L^\infty(\Omega)^m \times \mathbb{R})$ for the shadow limit.

2.3. Mean value correction

This problem is linked to the fact that in the shadow limit equation (1.2) only the mean value of g resp. v^0 appears. We will use the mean value correction to compare both solutions, the diffusive solution and its shadow limit. In [39], it is distinguished between initial time layer and a mean value correction for g , but – since we are interested in long-time behavior – we combine them. In the remainder, we use the mean value correction ψ_ε which solves

$$\begin{aligned} \frac{\partial \psi_\varepsilon}{\partial t} - \frac{1}{\varepsilon} \Delta \psi_\varepsilon &= g(\mathbf{u}, v, x, t) - \langle g(\mathbf{u}, v, \cdot, t) \rangle_\Omega \quad \text{in } \Omega \times (0, \infty), & \frac{\partial \psi_\varepsilon}{\partial \nu} &= 0 \quad \text{on } \partial\Omega \times (0, \infty), \\ \psi_\varepsilon(\cdot, 0) &= v^0 - \langle v^0 \rangle_\Omega \quad \text{in } \Omega, \end{aligned} \quad (2.4)$$

where (\mathbf{u}, v) is the given shadow limit. By Assumption 3, the right-hand side is bounded at least for finite times, and the well-known Duhamel formula reads

$$\psi_\varepsilon(\cdot, t) = S_\Delta(t/\varepsilon)(v^0 - \langle v^0 \rangle_\Omega) + \int_0^t S_\Delta((t-s)/\varepsilon) \left\{ g(\mathbf{u}(\cdot, s), v(s), \cdot, s) - \langle g(\mathbf{u}(\cdot, s), v(s), \cdot, s) \rangle_\Omega \right\} ds.$$

The mean value correction ψ_ε fulfills $\langle \psi_\varepsilon(\cdot, t) \rangle_\Omega = 0$ for all $t \in \mathbb{R}_{\geq 0}$ such that the exponential decay estimate (A.3) implies the estimate

$$\begin{aligned} \|\psi_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} &\leq \bar{C} e^{-\lambda_1 t/\varepsilon} \|v^0 - \langle v^0 \rangle_\Omega\|_{L^\infty(\Omega)} \\ &\quad + \bar{C} \int_0^t e^{-\lambda_1(t-s)/\varepsilon} \|g(\mathbf{u}(\cdot, s), v(s), \cdot, s) - \langle g(\mathbf{u}(\cdot, s), v(s), \cdot, s) \rangle_\Omega\|_{L^\infty(\Omega)} ds. \end{aligned}$$

Thus, it is natural for long-time behaviour to assume

Assumption 4 (Decay of mean value correction). *The mean value correction ψ_ε satisfies the estimate*

$$\|\psi_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)} \leq C_{v^0} e^{-\lambda_1 t/\varepsilon} + C_g \varepsilon \quad \forall \varepsilon > 0, t \in \mathbb{R}_{\geq 0} \quad (2.5)$$

for some constants $C_{v^0}, C_g > 0$ that do not depend on t or ε , but on bounds of v^0 resp. g .

Clearly, if $g - \langle g \rangle_\Omega$ is bounded in the time variable $t \in \mathbb{R}_{\geq 0}$ on bounded subsets of $\mathbb{R}^{m+1} \times \bar{\Omega}$, then Assumption 4 is satisfied. Especially, for nonlinearities g that do not depend explicitly on time, Assumption 4 holds true if Assumption 1–3 is satisfied.

3. Error estimates

We introduce the error functions

$$\mathbf{U}_\varepsilon = \mathbf{u}_\varepsilon - \mathbf{u} \quad \text{and} \quad V_\varepsilon = v_\varepsilon - v - \psi_\varepsilon, \quad (3.1)$$

where we took into account the mean value correction ψ_ε . Our goal is to estimate the error functions in terms of the small parameter ε . In this section, we consider the semilinear problem (1.1) and follow the idea of truncation as in the case of short-time intervals [39]. To assure boundedness of the model variables in the subsequent analysis, we localize the problem by introducing certain cut-off functions. This is similar as in the classical center manifold theory in [5, 17] or [51].

Our strategy of proof is depicted in Figure 1, starting in the left upper corner. Let $(\alpha_\varepsilon, \beta_\varepsilon)$ be the solution to the truncated problem (3.8). Using suitable estimates, we will show that the truncated solution $(\alpha_\varepsilon, \beta_\varepsilon)$ is located within the small neighborhood of $\mathbf{0}$ with radius ε^{δ_0} for sufficiently large diffusion $1/\varepsilon$. However, in this small neighborhood of the origin, the cut-off does not affect the nonlinearities if one restricts to the trajectory of the solution $(\alpha_\varepsilon, \beta_\varepsilon)$. As solutions to problem (3.2) are unique, this implies $(\alpha_\varepsilon, \beta_\varepsilon) = (\mathbf{U}_\varepsilon, V_\varepsilon)$, and we obtain error estimates for the original error as well. In order to estimate the truncated solution $(\alpha_\varepsilon, \beta_\varepsilon)$,




Figure 1: Strategy of proof of error estimates.

we decompose β_ε into its spatial mean $b_\varepsilon = \langle \beta_\varepsilon \rangle_\Omega$ and the residual $W_\varepsilon = \beta_\varepsilon - \langle \beta_\varepsilon \rangle_\Omega$ with $\langle W_\varepsilon \rangle_\Omega = 0$ to exploit asymptotic properties of the heat semigroup [15]. As already figured out in the linear case in [32], an additional stability condition is necessary to obtain long-time estimates. It turns out in Theorem 3.6 below that assuming stability conditions 6–7 for a linearized shadow system yields $L^\infty(\Omega_T)$ estimates for $(\alpha_\varepsilon, \beta_\varepsilon)$ which are valid on long-time ranges $(0, T)$. Optimality of the made assumptions in this section is exemplified for the linear case in Model 5.1.

3.1. Truncation of errors

Let us start from the system

$$\begin{aligned}
\frac{\partial \mathbf{U}_\varepsilon}{\partial t} &= \mathbf{f}(\mathbf{u}_\varepsilon, v_\varepsilon, x, t) - \mathbf{f}(\mathbf{u}, v, x, t) \quad \text{in } \Omega_T, & \mathbf{U}_\varepsilon(\cdot, 0) &= \mathbf{0} \quad \text{in } \Omega, \\
\frac{\partial V_\varepsilon}{\partial t} - \frac{1}{\varepsilon} \Delta V_\varepsilon &= g(\mathbf{u}_\varepsilon, v_\varepsilon, x, t) - g(\mathbf{u}, v, x, t) \quad \text{in } \Omega_T, & V_\varepsilon(\cdot, 0) &= 0 \quad \text{in } \Omega, \\
\frac{\partial V_\varepsilon}{\partial \nu} &= 0 \quad \text{on } \partial\Omega \times (0, T)
\end{aligned} \tag{3.2}$$

for the errors $\mathbf{U}_\varepsilon, V_\varepsilon$. Using Taylor’s expansion, we write for $\mathbf{h} = (\mathbf{f}, g)$

$$\mathbf{h}(\mathbf{u}_\varepsilon, v_\varepsilon, x, t) - \mathbf{h}(\mathbf{u}, v, x, t) = \nabla_{\mathbf{u}} \mathbf{h}(\mathbf{u}, v, x, t) \mathbf{U}_\varepsilon + \partial_v \mathbf{h}(\mathbf{u}, v, x, t) (V_\varepsilon + \psi_\varepsilon) + \mathbf{H}(\mathbf{U}_\varepsilon, V_\varepsilon + \psi_\varepsilon, x, t), \tag{3.3}$$

where the remainder $\mathbf{H} = (\mathbf{F}, G)$ is given due to the mean value theorem by

$$\mathbf{H}(\mathbf{y}, z + \psi_\varepsilon, x, t) = \int_0^1 \nabla \mathbf{h}(\mathbf{u} + \vartheta \mathbf{y}, v + \vartheta(z + \psi_\varepsilon), x, t) - \nabla \mathbf{h}(\mathbf{u}, v, x, t) \, d\vartheta \cdot \begin{pmatrix} \mathbf{y} \\ z + \psi_\varepsilon \end{pmatrix}.$$

For this to be valid and for the following procedure let us assume, in addition to Assumption 1–4, a differentiability assumption.

Assumption 5 (Regularity for linearization). *Let \mathbf{f}, g be continuously differentiable with respect to the variables $(\mathbf{u}, v) \in \mathbb{R}^{m+1}$ and let their derivatives $\nabla_{\mathbf{u}} f_i, \nabla_v f_i, i = 1, \dots, m$, and $\nabla_{\mathbf{u}} g, \partial_v g$ satisfy Assumption 1. Let the linearized parts $\nabla_{(\mathbf{u}, v)} \mathbf{f}, \nabla_{(\mathbf{u}, v)} g$, evaluated at the shadow limit, be uniformly bounded in $(x, t) \in \bar{\Omega} \times \mathbb{R}_{\geq 0}$.*

For instance, such a regularity is satisfied for an autonomous right-hand side (\mathbf{f}, g) which is of class C^2 with respect to the unknown variables (\mathbf{u}, v) . If, in addition, we have a uniformly bounded shadow limit, then Assumption 5 is fulfilled.

Following the idea of truncation, we construct a suitable cut-off for the possibly unbounded right-hand side $\mathbf{H} = (\mathbf{F}, G)$. We first modify the \mathbf{y} -component using a cut-off function $\Theta \in C^{0,1}(\mathbb{R})$ defined by

$$\Theta(z) = \begin{cases} \operatorname{sgn}(z) \cdot \varepsilon^{\delta_0} & \text{for } |z| > \varepsilon^{\delta_0}, \\ z & \text{for } |z| \leq \varepsilon^{\delta_0} \end{cases} \quad (3.4)$$

for $\delta_0 \in (0, 1/2]$. The function Θ satisfies $|\Theta(z)| \leq \min\{\varepsilon^{\delta_0}, |z|\}$. For simplicity, in the vector-valued case, we write $\Theta(\mathbf{y}) := (\Theta(y_1), \dots, \Theta(y_m))$.

Remark 3.1. Notice that the choice of δ_0 is restricted to the interval $(0, 1/2]$. This restriction is linked to the long-time error estimates in Proposition 3.5 in which the convergence rate is weaker than in case of finite time analysis in [39].

We further consider the function $\overline{\mathbf{H}}_\varepsilon = (\overline{\mathbf{F}}_\varepsilon, \overline{G}_\varepsilon)$ given by $\overline{\mathbf{H}}_\varepsilon(\mathbf{y}, z, x, t) := \mathbf{H}(\Theta(\mathbf{y}), z + \psi_\varepsilon, x, t)$. To control the z -component of the truncation, we use a symmetric cut-off function $\rho \in C_c^\infty(\mathbb{R}; [0, 1])$ defined by

$$\rho(z) = \begin{cases} 1 & \text{for } |z| \leq L, \\ 0 & \text{for } |z| \geq 2L. \end{cases} \quad (3.5)$$

Here, $L := C_{v_0} + 2$, where $C_{v_0} > 0$ is the same constant from the time decay estimate (2.5) of ψ_ε . Such a choice of ρ is possible by mollifying the characteristic function $\mathbb{1}_{(-r, r)}$ where $L < r < 2L$ [1, Theorem 2.29]. We define a further truncation by

$$\mathbf{H}_\varepsilon(\mathbf{y}, z, x, t) := \rho\left(\frac{\varepsilon^{-\delta_0}|\bar{z}|}{L}\right) \overline{\mathbf{H}}_\varepsilon(\mathbf{y}, z, x, t) \left(1 - \rho\left(\frac{2\lambda_1 t}{-\varepsilon \log \varepsilon}\right)\right) + \rho\left(\frac{|\bar{z}|}{L}\right) \overline{\mathbf{H}}_\varepsilon(\mathbf{y}, z, x, t) \rho\left(\frac{2\lambda_1 t}{-\varepsilon \log \varepsilon}\right) \quad (3.6)$$

for $\bar{z} := z + \psi_\varepsilon$. A similar modification is done in [39, Lemma 2]. Properties of the truncated function $\mathbf{H}_\varepsilon = (\mathbf{F}_\varepsilon, G_\varepsilon)$ are given next.

Lemma 3.2. *Let Assumption 1–5 be satisfied. For each $\delta_0 \in (0, 1/2]$ there is a constant $C > 0$, independent of $\varepsilon \leq 1$ but which depends on L defined in (3.5), such that for $\bar{z} = z + \psi_\varepsilon$ we have*

$$|\mathbf{F}_\varepsilon(\mathbf{y}, z, x, t)|, |G_\varepsilon(\mathbf{y}, z, x, t)| \leq C \left(\varepsilon^{2\delta_0} + \mathbb{1}_{\{t \leq -\varepsilon \log \varepsilon / \lambda_1\}}(t) \cdot \min\{1, |\bar{z}|\} \right) \quad (3.7)$$

for all $(\mathbf{y}, z) \in \mathbb{R}^{m+1}$, $t \in \mathbb{R}_{\geq 0}$ and for a.e. $x \in \Omega$. Here, $\mathbb{1}_{\{t \leq c\}}$ is the characteristic function on the time interval $[0, c]$.

Proof. The first term in definition (3.6) can be estimated as $\overline{\mathbf{H}}_\varepsilon$. However, only $\varepsilon^{-\delta_0}|\bar{z}| \leq 2L$ has to be considered due to $\rho \in C_c^\infty(\mathbb{R})$ defined in (3.5). By Assumption 3–5, we infer a constant $C > 0$ such that

$$|\overline{\mathbf{H}}_\varepsilon(\mathbf{y}, z, \cdot, \cdot)| \leq C |(\Theta(\mathbf{y}), \bar{z})|^2.$$

This constant C depends on the time-independent bounds on \mathbf{u}, v in Assumption 3, on the one for ψ_ε in Assumption 4, and on Lipschitz bounds in Assumption 5 for the derivatives, but neither on diffusion nor on time t . For the second term in definition (3.6), we use the estimate

$$\left| \rho\left(\frac{|\bar{z}|}{L}\right) \overline{\mathbf{H}}_\varepsilon(\mathbf{y}, z, x, t) \rho\left(\frac{2\lambda_1 t}{-\varepsilon \log \varepsilon}\right) \right| \leq C (\varepsilon^{2\delta_0} + |\bar{z}|^2) \rho\left(\frac{|\bar{z}|}{L}\right) \mathbb{1}_{\{t \leq -\varepsilon \log \varepsilon / \lambda_1\}}(t)$$

which results from the estimate for $\overline{\mathbf{H}}_\varepsilon$. The right-hand side is at most non-zero if $|\bar{z}| \leq 2L$ and the quadratic term is (linearly) bounded. \square

In the following, we study the localized problem using the truncation \mathbf{H}_ε associated with the error system (3.2). A substitution of \mathbf{H} in equation (3.3) yields

$$\begin{aligned} \frac{\partial \alpha_\varepsilon}{\partial t} &= \nabla_{\mathbf{u}} \mathbf{f} \cdot \alpha_\varepsilon + \nabla_v \mathbf{f} \cdot (\beta_\varepsilon + \psi_\varepsilon) + \mathbf{F}_\varepsilon(\alpha_\varepsilon, \beta_\varepsilon, x, t) \quad \text{in } \Omega_T, \\ \frac{\partial \beta_\varepsilon}{\partial t} - \frac{1}{\varepsilon} \Delta \beta_\varepsilon &= \nabla_{\mathbf{u}} \mathbf{g} \cdot \alpha_\varepsilon + \nabla_v \mathbf{g} \cdot (\beta_\varepsilon + \psi_\varepsilon) + G_\varepsilon(\alpha_\varepsilon, \beta_\varepsilon, x, t) \quad \text{in } \Omega_T, \\ \alpha_\varepsilon(\cdot, 0) &= \mathbf{0}, \quad \beta_\varepsilon(\cdot, 0) = 0 \quad \text{in } \Omega, \quad \frac{\partial \beta_\varepsilon}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times (0, T). \end{aligned} \quad (3.8)$$

Herein, we abbreviated the Jacobians $\nabla_{\mathbf{u}}\mathbf{f}, \nabla_v\mathbf{f}, \nabla_{\mathbf{u}}g, \nabla_vg$ which are evaluated at the shadow solution (\mathbf{u}, v) and depend in general on space and time.

The focal idea using cut-offs is to find estimates for the solution $(\alpha_\varepsilon, \beta_\varepsilon)$ of the truncated problem (3.8) to show that its solution actually is located for sufficiently large diffusions $1/\varepsilon$ within a small neighborhood of $\mathbf{0}$, with radius ε^δ for some $\delta > 0$, where the cut-offs have no effect if $\delta \geq \delta_0$.

Proposition 3.3. *Let Assumption 1–5 hold and let $\varepsilon \leq 1$. Then there exists a unique mild solution $(\alpha_\varepsilon, \beta_\varepsilon)$ of the truncated problem (3.8) which exists globally-in-time. Furthermore, for all finite times $T > 0$, we have $(\alpha_\varepsilon, \beta_\varepsilon) \in L^\infty(\Omega_T)^{m+1}$ and $\beta_\varepsilon \in L^\infty(0, T; H^1(\Omega))$ is a weak solution with weak derivative $\partial_t\beta_\varepsilon \in L^2(\Omega_T)$.*

Proof. To apply Rothe's method as in Theorem 2.2, we write system (3.8) as $m + 1$ differential equations

$$\frac{\partial \Psi_\varepsilon}{\partial t} - \mathbf{D}_\varepsilon \Delta \Psi_\varepsilon = \mathbf{h}_\varepsilon(\Psi_\varepsilon, x, t) \quad \text{in } \Omega_T, \quad \Psi_\varepsilon(\cdot, 0) = \mathbf{0} \quad \text{in } \Omega.$$

The function β_ε is endowed with zero Neumann boundary conditions and \mathbf{h}_ε is given by

$$\mathbf{h}_\varepsilon(\Psi_\varepsilon, x, t) = \mathbf{J}(x, t) \cdot \begin{pmatrix} \alpha_\varepsilon \\ \beta_\varepsilon + \psi_\varepsilon(x, t) \end{pmatrix} + \begin{pmatrix} \mathbf{F}_\varepsilon(\alpha_\varepsilon, \beta_\varepsilon, x, t) \\ G_\varepsilon(\alpha_\varepsilon, \beta_\varepsilon, x, t) \end{pmatrix}.$$

Here and in the sequel, we use the notation \mathbf{J} for the Jacobian

$$\mathbf{J}(x, t) := \begin{pmatrix} \mathbf{A}(x, t) & \mathbf{B}(x, t) \\ \mathbf{C}(x, t) & D(x, t) \end{pmatrix} := \begin{pmatrix} \nabla_{\mathbf{u}}\mathbf{f}(\mathbf{u}(x, t), v(t), x, t) & \nabla_v\mathbf{f}(\mathbf{u}(x, t), v(t), x, t) \\ \nabla_{\mathbf{u}}g(\mathbf{u}(x, t), v(t), x, t) & \nabla_vg(\mathbf{u}(x, t), v(t), x, t) \end{pmatrix} \quad (3.9)$$

evaluated at the shadow solution (\mathbf{u}, v) . Local Lipschitz continuity of \mathbf{h}_ε in the variable Ψ_ε on bounded sets in $\overline{\Omega} \times \mathbb{R}_{\geq 0}$ carries over from \mathbf{h} since \mathbf{F}_ε and G_ε are locally Lipschitz in the sense of Assumption 1, see definition (3.6). Following the proof of Proposition 2.2, there exists a local-in-time mild solution in the sense of Definition 2.1. We obtain the integral representation

$$\Psi_\varepsilon(\cdot, t) = \int_0^t \mathbf{S}(t - \tau) \mathbf{h}_\varepsilon(\Psi_\varepsilon(\cdot, \tau), \cdot, \tau) \, d\tau.$$

The function \mathbf{h}_ε is linearly bounded in the variable Ψ_ε due to Lemma 3.2. By the same reasoning as in [46, Part II, Theorem 1], this implies that no blow-up is possible and we obtain a unique mild solution with $\Psi_\varepsilon \in L^\infty(\Omega_T)^{m+k}$ for all $T < \infty$.

In order to improve regularity for the diffusing component, we apply parabolic L^2 theory performed in Proposition A.2. Recall for this that β_ε solves equation (A.6) with $R \in L^\infty(\Omega_T)$. \square

To obtain estimates for very long time intervals of order $\varepsilon^{-\ell}$ for some $\ell > 0$, we need a more detailed estimation of the quantities which are involved. We decompose β_ε into its spatial mean $b_\varepsilon = \langle \beta_\varepsilon \rangle_\Omega$ and the residual $W_\varepsilon = \beta_\varepsilon - \langle \beta_\varepsilon \rangle_\Omega$ with $\langle W_\varepsilon \rangle_\Omega = 0$ to exploit asymptotic properties of the heat semigroup. The differential equation of β_ε from problem (3.8) may be replaced by

$$\begin{aligned} \frac{\partial W_\varepsilon}{\partial t} - \frac{1}{\varepsilon} \Delta W_\varepsilon &= \nabla_{\mathbf{u}}g \cdot \alpha_\varepsilon - \langle \nabla_{\mathbf{u}}g \cdot \alpha_\varepsilon \rangle_\Omega + \nabla_vg \cdot b_\varepsilon - \langle \nabla_vg \cdot b_\varepsilon \rangle_\Omega \\ &\quad + \nabla_vg \cdot (W_\varepsilon + \psi_\varepsilon) - \langle \nabla_vg \cdot (W_\varepsilon + \psi_\varepsilon) \rangle_\Omega + G_\varepsilon - \langle G_\varepsilon \rangle_\Omega \quad \text{in } \Omega \times \mathbb{R}_{>0}, \\ \frac{db_\varepsilon}{dt} &= \langle \nabla_{\mathbf{u}}g \cdot \alpha_\varepsilon \rangle_\Omega + \langle \nabla_vg \cdot b_\varepsilon \rangle_\Omega + \langle \nabla_vg \cdot (W_\varepsilon + \psi_\varepsilon) \rangle_\Omega + \langle G_\varepsilon \rangle_\Omega \quad \text{in } \Omega \times \mathbb{R}_{>0}, \\ W_\varepsilon(\cdot, 0) &= 0 \quad \text{in } \Omega, \quad b_\varepsilon(0) = 0, \quad \frac{\partial W_\varepsilon}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times \mathbb{R}_{>0}. \end{aligned} \quad (3.10)$$

We start estimating the term W_ε using semigroup estimates from [50, Lemma 1.3]. Denoting the right-hand side of the elliptic equation by R_ε , the solution W_ε may be written as

$$W_\varepsilon(\cdot, t) = \int_0^t S_\Delta((t - \tau)/\varepsilon) R_\varepsilon(\cdot, \tau) \, d\tau.$$

By Assumption 5, the Jacobian $\mathbf{J} \in L^\infty(\Omega \times \mathbb{R}_{\geq 0})^{(m+1) \times (m+1)}$ is uniformly bounded. We infer from decay estimate (2.5) for ψ_ε and $\langle R_\varepsilon \rangle_\Omega = 0$ that

$$\begin{aligned} \|W_\varepsilon(\cdot, t)\|_{L^p(\Omega)} &\leq C \int_0^t e^{-\lambda_1(t-\tau)/\varepsilon} \left(\|\alpha_\varepsilon(\cdot, \tau)\|_{L^p(\Omega)^m} + |\Omega|^{1/p} |b_\varepsilon(\tau)| \right. \\ &\quad \left. + \|(W_\varepsilon + \psi_\varepsilon)(\cdot, \tau)\|_{L^p(\Omega)} + \|G_\varepsilon(\cdot, \tau)\|_{L^p(\Omega)} \right) d\tau. \end{aligned} \quad (3.11)$$

This estimate implies that W_ε can always be estimated in terms of $\alpha_\varepsilon, b_\varepsilon$ and powers of ε . Hence, it remains to control the terms α_ε and b_ε . The components $\alpha_\varepsilon, b_\varepsilon$ satisfy a shadow problem whose solution is given by

$$\begin{pmatrix} \alpha_\varepsilon(\cdot, t) \\ b_\varepsilon(t) \end{pmatrix} = \int_0^t \mathbf{W}(t, \tau) \begin{pmatrix} \mathbf{B}(\cdot, \tau)(W_\varepsilon + \psi_\varepsilon)(\cdot, \tau) + \mathbf{F}_\varepsilon(\cdot, \tau) \\ \langle D(\cdot, \tau)(W_\varepsilon + \psi_\varepsilon)(\cdot, \tau) + G_\varepsilon(\cdot, \tau) \rangle_\Omega \end{pmatrix} d\tau \quad (3.12)$$

where we used the evolution system \mathcal{W} induced by the linearization of the shadow problem. This system is given by evolution operators $\mathbf{W}(t, s)$ for $t, s \in \mathbb{R}_{\geq 0}, s \leq t$, defined by

$$\xi(\cdot, t) = \mathbf{W}(t, s)\xi(\cdot, s), \quad \xi(\cdot, 0) = \begin{pmatrix} \xi_1^0 \\ \langle \xi_2^0 \rangle_\Omega \end{pmatrix}, \quad (3.13)$$

where $\xi \in C(\mathbb{R}_{\geq 0}; L^p(\Omega)^m \times \mathbb{R})$ is the unique solution of the homogeneous linear shadow problem

$$\begin{aligned} \frac{\partial \xi_1}{\partial t} &= \mathbf{A}(x, t)\xi_1 + \mathbf{B}(x, t)\xi_2 & \text{in } \Omega \times \mathbb{R}_{>0}, & \quad \xi_1(\cdot, 0) = \xi_1^0 \text{ in } \Omega, \\ \frac{d\xi_2}{dt} &= \langle \mathbf{C}(\cdot, t)\xi_1 \rangle_\Omega + \langle D(\cdot, t)\xi_2 \rangle_\Omega & \text{in } \mathbb{R}_{>0}, & \quad \xi_2(0) = \langle \xi_2^0 \rangle_\Omega. \end{aligned}$$

In order to control the growth of the errors $\alpha_\varepsilon, b_\varepsilon$, we assume the following stability condition.

Assumption 6 (Stability of linearized shadow system). *Let the evolution system \mathcal{W} be uniformly bounded for some $1 \leq p \leq \infty$, i.e., there is a constant $C > 0$ independent of time such that for all $s, t \in \mathbb{R}_{\geq 0}, s \leq t$,*

$$\|\mathbf{W}(t, s)\xi^0\|_{L^p(\Omega)^m \times \mathbb{R}} \leq C \|\xi^0\|_{L^p(\Omega)^m \times \mathbb{R}} \quad \forall \xi^0 \in L^p(\Omega)^m \times \mathbb{R}.$$

We refer to Section B.1 for a verification of Assumption 6 in the case of a stationary shadow limit. Furthermore, in case of dissipative systems, we refer to Section 4. Model 5.1 shows that space-dependent problems require more assumptions than just uniform stability for the ODE subsystem \mathcal{U} , which is assumed in [32]. This stability condition leads to the following error estimate.

Proposition 3.4. *Let Assumption 1–6 hold for some $1 \leq p \leq \infty$. Then for any $\alpha \in (0, 1], \delta_0 \in (0, 1/2]$ with $\gamma := 2\delta_0 + (\alpha - 1) \in (0, 1]$ there exist constants $C, \varepsilon_0 > 0$ independent of time T and diffusion $1/\varepsilon$ such that for all times $T \leq \varepsilon^{\alpha-1}$ and all $\varepsilon \leq \varepsilon_0$ the solution $(\alpha_\varepsilon, \beta_\varepsilon)$ of system (3.8) satisfies*

$$\sup_{t \in [0, T]} (\|\alpha_\varepsilon(\cdot, t)\|_{L^p(\Omega)^m} + \|\beta_\varepsilon(\cdot, t)\|_{L^p(\Omega)}) \leq C\varepsilon^\gamma. \quad (3.14)$$

Proof. We already estimated W_ε and received a relation to $\alpha_\varepsilon, b_\varepsilon$ in inequality (3.11). In view of estimate (3.7) for G_ε , where $\bar{\beta}_\varepsilon = b_\varepsilon + W_\varepsilon + \psi_\varepsilon$, and Assumption 4 on ψ_ε , we observe for each $\delta_0 \geq 0$

$$\begin{aligned} \|W_\varepsilon(\cdot, t)\|_{L^p(\Omega)} &\leq C \int_0^t e^{-\lambda_1(t-\tau)/\varepsilon} \left(\|\alpha_\varepsilon(\cdot, \tau)\|_{L^p(\Omega)^m} + |\Omega|^{1/p} |b_\varepsilon(\tau)| \right) d\tau \\ &\quad + C \int_0^t e^{-\lambda_1(t-\tau)/\varepsilon} \|W_\varepsilon(\cdot, \tau)\|_{L^p(\Omega)} d\tau + C|\Omega|^{1/p}\varepsilon. \end{aligned} \quad (3.15)$$

To obtain a corresponding inequality for $\alpha_\varepsilon, b_\varepsilon$, we consider the explicit formula (3.12). Applying Assumption 6 on \mathcal{W} , Assumption 4 and estimate (3.7) for the truncation $\mathbf{H}_\varepsilon = (\mathbf{F}_\varepsilon, G_\varepsilon)$ to this integral representation yields

$$\begin{aligned} \|\alpha_\varepsilon(\cdot, t)\|_{L^p(\Omega)^m} + |\Omega|^{1/p}|b_\varepsilon(t)| &\leq C \int_0^t \|(W_\varepsilon + \psi_\varepsilon)(\cdot, \tau)\|_{L^p(\Omega)} d\tau + C \int_0^t \|\mathbf{H}_\varepsilon(\cdot, \tau)\|_{L^p(\Omega)^{m+1}} d\tau \\ &\leq C \int_0^t \|W_\varepsilon(\cdot, \tau)\|_{L^p(\Omega)} d\tau + C \int_0^t \mathbf{1}_{\{\tau \leq -\varepsilon \log \varepsilon / \lambda_1\}}(\tau) |\Omega|^{1/p} |b_\varepsilon(\tau)| d\tau \\ &\quad + C |\Omega|^{1/p} (\varepsilon(1+t) + \varepsilon^{2\delta_0} t). \end{aligned}$$

We restrict ourselves to $\delta_0 \leq 1/2$. Since $-\varepsilon \log \varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$ by L'Hospital's rule, we absorb b_ε on the left-hand side and obtain

$$\sup_{t \in [0, T]} \left(\|\alpha_\varepsilon(\cdot, t)\|_{L^p(\Omega)^m} + |\Omega|^{1/p} |b_\varepsilon(t)| \right) \leq C \int_0^T \|W_\varepsilon(\cdot, \tau)\|_{L^p(\Omega)} d\tau + C |\Omega|^{1/p} (\varepsilon + \varepsilon^{2\delta_0} T). \quad (3.16)$$

Estimate (3.16) combined with inequality (3.15) for W_ε leads to an estimate for W_ε , more precisely,

$$\sup_{t \in [0, T]} \|W_\varepsilon(\cdot, t)\|_{L^p(\Omega)} \leq C \sup_{t \in [0, T]} \int_0^t \left(e^{-\lambda_1(t-\tau)/\varepsilon} + \varepsilon \right) \|W_\varepsilon(\cdot, \tau)\|_{L^p(\Omega)} d\tau + C |\Omega|^{1/p} \varepsilon (1 + \varepsilon^{2\delta_0} T).$$

Taking the supremum over t yields

$$\sup_{t \in [0, T]} \|W_\varepsilon(\cdot, t)\|_{L^p(\Omega)} \leq C \varepsilon (1 + T) \sup_{t \in [0, T]} \|W_\varepsilon(\cdot, t)\|_{L^p(\Omega)} + C \varepsilon (1 + \varepsilon^{2\delta_0} T).$$

Absorption of terms on the left-hand side for all $T \leq \varepsilon^{\alpha-1}$ implies for $\varepsilon \leq \varepsilon_0$ that

$$\sup_{t \in [0, T]} \|W_\varepsilon(\cdot, t)\|_{L^p(\Omega)} \leq C \varepsilon^{\min\{1, 2\delta_0 + \alpha\}}.$$

Using estimate (3.16) for α_ε , for each $\gamma = 2\delta_0 + (\alpha - 1) \in (0, 1]$ there holds

$$\sup_{t \in [0, T]} \left(\|\alpha_\varepsilon(\cdot, t)\|_{L^p(\Omega)^m} + |\Omega|^{1/p} |b_\varepsilon(t)| \right) \leq C (\varepsilon^{2\delta_0} T + \varepsilon T) \leq C \varepsilon^\gamma$$

for $\varepsilon \leq \varepsilon_0 = \varepsilon_0(\gamma)$. As a consequence, $\gamma > 0$ implies $2\delta_0 + \alpha > 1$ and $\|W_\varepsilon\|_{L^p(\Omega)} \leq C \varepsilon$. The relation $\beta_\varepsilon = W_\varepsilon + b_\varepsilon$ implies an estimate for β_ε . \square

If Assumption 6 holds with $p = \infty$, we already obtain a corresponding estimate in $L^\infty(\Omega_T)$ for solutions to the truncated problem (3.8). If this is not the case, Hölder's inequality yields bounds for the norms of $\alpha_\varepsilon, \beta_\varepsilon$ in the parabolic space $L_{p,r}(\Omega_T)$ for each $1 \leq r < \infty$, for instance,

$$\|\alpha_\varepsilon\|_{p,r} \leq T^{1/r} \|\alpha_\varepsilon\|_{p,\infty} \leq C \varepsilon^{\gamma - \frac{1}{r}(1-\alpha)}$$

and similarly for β_ε . Actually the latter inequalities imply, either for $p \geq 1 = n$ or for $p > n/2 \geq 1$, an $L^\infty(\Omega_T)$ estimate for the diffusing component β_ε using Proposition A.3. For non-diffusing components, we apply a further stability assumption according to [32].

Therefore, let us focus on the corresponding ODE subsystem induced by the submatrix $\mathbf{A}(x, t)$ of the above Jacobian \mathbf{J} defined in (3.9). To control the growth of the ODE components, we consider solutions to the initial value problem

$$\frac{\partial \xi}{\partial t} = \mathbf{A}(\cdot, t) \xi \quad \text{in } \Omega \times \mathbb{R}_{>0}, \quad \xi(\cdot, 0) = \xi^0 \quad \text{in } \Omega.$$

Using standard techniques of ordinary differential equations from [6, Ch. III, §1], existence of a unique mild solution $\xi \in C(\mathbb{R}_{\geq 0}; L^\infty(\Omega)^m)$ is assured. The solution is given implicitly by the Duhamel formula

$$\xi(\cdot, t) = \xi(\cdot, s) + \int_s^t \mathbf{A}(\cdot, \tau) \xi(\cdot, \tau) d\tau \quad \forall 0 \leq s \leq t.$$

The solution defines an evolution system \mathcal{U} consisting of a family of evolution operators $\mathbf{U}(t, s)$ on $L^\infty(\Omega)^m$:

$$\xi(\cdot, t) = \mathbf{U}(t, s) \xi(\cdot, s) \quad \forall s, t \in \mathbb{R}_{\geq 0}, s \leq t.$$

As shown in [32, Theorem 1], uniform stability of the evolution system \mathcal{U} is essential for showing convergence.

Assumption 7 (Stability of ODE subsystem). *Let the evolution system \mathcal{U} be uniformly bounded, i.e., there is a constant $C > 0$ independent of time such that*

$$\|\mathbf{U}(t, s) \xi^0\|_{L^\infty(\Omega)^m} \leq C \|\xi^0\|_{L^\infty(\Omega)^m} \quad \forall \xi^0 \in L^\infty(\Omega)^m, s, t \in \mathbb{R}_{\geq 0}, s \leq t.$$

This condition is optimal in the sense that there are examples where uniform convergence on long-time intervals may not be achievable in the absence of uniform boundedness in $L^\infty(\Omega)^m$ [32, Example 2]. Under this additional assumption, we obtain the following result.

Proposition 3.5. *Let Assumption 1–7 hold for some finite $p \geq 1 = n$ or $p > n/2$ for $n \geq 2$ and choose r as in relation (A.7). Then there exist triples $(\alpha, \delta_0, r) \in (0, 1] \times (0, 1/2] \times (1, \infty)$ and constants $C, \varepsilon_0 > 0$ independent of T, ε such that for all $T \leq \varepsilon^{\alpha-1}$ and $\varepsilon \leq \varepsilon_0$ there holds*

$$\|\alpha_\varepsilon\|_{L^\infty(\Omega_T)^m} \leq C \varepsilon^{-2(1-\alpha)} \left(\varepsilon^\gamma + \varepsilon^{(2-\alpha)/r} \right), \quad \|\beta_\varepsilon\|_{L^\infty(\Omega_T)} \leq C \varepsilon^{-(1-\alpha)} \left(\varepsilon^\gamma + \varepsilon^{(2-\alpha)/r} \right) \quad (3.17)$$

for $\gamma = 2\delta_0 + (\alpha - 1) \in (0, 1]$ from Proposition 3.4. If Assumption 6 holds with $p = \infty$, estimate (3.14) is true without requiring Assumption 7, and we may choose ε^γ as a convergence rate for each component.

Proof. If $p = \infty$, there is nothing to show, see Proposition 3.4 and estimate (3.14). Note that uniform boundedness of the evolution system \mathcal{W} of the entire shadow system in $L^\infty(\Omega)^m \times \mathbb{R}$ is sufficient, thus no boundedness of the ODE subsystem stated in Assumption 7 is needed.

Let us assume $p < \infty$ and consider the diffusive component β_ε of the truncated problem (3.8) solving

$$\frac{\partial \beta_\varepsilon}{\partial t} - \frac{1}{\varepsilon} \Delta \beta_\varepsilon = R_\varepsilon(x, t).$$

Proposition A.3 yields a constant $C > 0$ such that

$$\|\beta_\varepsilon\|_{L^\infty(\Omega_T)} \leq CT^{1-1/r} \|R_\varepsilon\|_{p,r}$$

for some $r > 1$ defined by (A.7). This constant C is independent of time T and only depends on a lower bound for the diffusion and parameters of the systems. Thus, it remains to find an estimate for $\|R_\varepsilon\|_{p,r}$. We infer from the right-hand side of the truncated system (3.8) that

$$\|R_\varepsilon\|_{p,r} \leq C (\|\alpha_\varepsilon\|_{p,r} + \|\beta_\varepsilon\|_{p,r} + \|\psi_\varepsilon\|_{p,r} + \max\{\|\mathbf{F}_\varepsilon\|_{p,r}, \|G_\varepsilon\|_{p,r}\}).$$

Assumption 4 on ψ_ε and Lemma 3.2 with $\bar{\beta}_\varepsilon = \beta_\varepsilon + \psi_\varepsilon$ yields

$$\|R_\varepsilon\|_{p,r} \leq C \left(\|\alpha_\varepsilon\|_{p,r} + \|\beta_\varepsilon\|_{p,r} + \varepsilon^{1/r} + \varepsilon^{2\delta_0} T^{1/r} \right).$$

Using Proposition 3.4, above a priori estimates for $\alpha_\varepsilon, \beta_\varepsilon$ in $L_{p,r}(\Omega_T)$ and $T \leq \varepsilon^{\alpha-1}$ for some $\alpha \in (0, 1]$ imply

$$\|\beta_\varepsilon\|_{L^\infty(\Omega_T)} \leq C \varepsilon^{-(1-\frac{1}{r})(1-\alpha)} \left(\varepsilon^{\gamma-\frac{1}{r}(1-\alpha)} + \varepsilon^{1/r} + \varepsilon^{2\delta_0-\frac{1}{r}(1-\alpha)} \right).$$

Since $\gamma = 2\delta_0 + (\alpha - 1) \leq 2\delta_0$, we further estimate

$$\|\beta_\varepsilon\|_{L^\infty(\Omega_T)} \leq C\varepsilon^{-(1-\alpha)} \left(\varepsilon^\gamma + \varepsilon^{(2-\alpha)/r} \right).$$

To obtain an analog estimate for α_ε , we restrict our consideration to Assumption 7 for the subsystem of non-diffusing components. Using the notion of Assumption 7 and the first equation of system (3.8), this error component is given by

$$\alpha_\varepsilon(\cdot, t) = \int_0^t \mathbf{U}(t, \tau) \mathbf{S}_\varepsilon(\cdot, \tau) d\tau,$$

for some right-hand side $\mathbf{S}_\varepsilon \in L^\infty(\Omega_T)^m$ which we already estimated. Using uniform boundedness of the evolutionary system \mathbf{U} , we obtain

$$\|\alpha_\varepsilon\|_{L^\infty(\Omega_T)^m} \leq C \int_0^T \|\mathbf{S}_\varepsilon(\cdot, \tau)\|_{L^\infty(\Omega)^m} d\tau \leq C\varepsilon^{-2(1-\alpha)} \left(\varepsilon^\gamma + \varepsilon^{(2-\alpha)/r} \right).$$

Specifically, we employed the estimate for β_ε and that \mathbf{F}_ε can be estimated by $\bar{\beta}_\varepsilon = \beta_\varepsilon + \psi_\varepsilon$, too. \square

3.2. Convergence results

We are now in a position to draw a conclusion for the original errors $(\mathbf{U}_\varepsilon, V_\varepsilon)$ using estimates for the truncated problem in the last section. In order to dispose of truncation, we infer from results of Proposition 3.5 that the truncated solution $(\alpha_\varepsilon, \beta_\varepsilon)$, for sufficiently large diffusion, is located in a neighborhood of Θ where the cut-off is not required.

Theorem 3.6. *Let the assumptions 1–7 hold for some p with $p \geq 1 = n$ or $p > n/2$ if $n \geq 2$ and let $r \in (1, \infty)$ given by (A.7). Then there exist bounds $\alpha_0 = \alpha_0(r) \in (0, 1)$, $\varepsilon_0 > 0$ and a constant $C > 0$ such that for any $\alpha \in [\alpha_0, 1)$, $\varepsilon \leq \varepsilon_0$ and times $T \leq \varepsilon^{\alpha-1}$ we have the uniform estimates*

$$\begin{aligned} \|\mathbf{u}_\varepsilon - \mathbf{u}\|_{L^\infty(\Omega_T)^m} &\leq C\varepsilon^{3(1-\alpha)}, \\ \|v_\varepsilon - v - \psi_\varepsilon\|_{L^\infty(\Omega_T)} &\leq C\varepsilon^{4(1-\alpha)}, \\ \|\langle v_\varepsilon \rangle_\Omega - v\|_{L^\infty((0, T))} &\leq C\varepsilon^{4(1-\alpha)}. \end{aligned} \tag{3.18}$$

For $p = \infty$, without requiring Assumption 7, similar estimates hold true with a convergence rate $\varepsilon^{(1-\alpha)}$ for any component.

Proof. To get rid of truncation, it suffices to show

$$\|\alpha_\varepsilon\|_{L^\infty(\Omega_T)^m} + \|\beta_\varepsilon\|_{L^\infty(\Omega_T)} \leq \varepsilon^{\delta_0} \quad \forall \varepsilon \leq \varepsilon_0 \tag{3.19}$$

since in that case Θ is not needed anymore in the definition $\bar{\mathbf{H}}_\varepsilon$ and we obtain $\bar{\mathbf{H}}_\varepsilon = \mathbf{H}$. By considering the two cases, above and below the critical time $t^* := -\delta_0\varepsilon \log \varepsilon/\lambda_1$ for construction (3.6), we deduce $\mathbf{H}_\varepsilon = \mathbf{H}$ in the following.

- If $t \leq t^*$, we obtain $\rho\left(\frac{2\lambda_1 t}{-\varepsilon \log \varepsilon}\right) = 1$ by $2\delta_0 \leq L$ and thus, $\mathbf{H}_\varepsilon(\alpha_\varepsilon, \beta_\varepsilon, x, t) = \rho\left(\frac{|\bar{\beta}_\varepsilon|}{L}\right) \mathbf{H}(\alpha_\varepsilon, \beta_\varepsilon, x, t)$. Additionally, there holds $\rho\left(\frac{|\bar{\beta}_\varepsilon|}{L}\right) = 1$ since by definition of $L = C_{v_0} + 2$

$$|\bar{\beta}_\varepsilon| \leq |\beta_\varepsilon| + |\psi_\varepsilon| \leq \varepsilon^{\delta_0} + C_g\varepsilon + (L-2)e^{-\lambda_1 t/\varepsilon} \leq L$$

for all $\varepsilon \leq \varepsilon_0$ small enough.

- If $t > t^*$, then $e^{-\lambda_1 t/\varepsilon} \leq \varepsilon^{\delta_0}$ and thus, for small ε

$$|\bar{\beta}_\varepsilon| \leq |\beta_\varepsilon| + |\psi_\varepsilon| \leq \varepsilon^{\delta_0} + C_g\varepsilon + (L-2)\varepsilon^{\delta_0} \leq L\varepsilon^{\delta_0}.$$

Clearly, $|\bar{\beta}_\varepsilon| \leq |\varepsilon^{-\delta_0}\bar{\beta}_\varepsilon| \leq L$ and we find once again by definition (3.6) that

$$\mathbf{H}_\varepsilon(\alpha_\varepsilon, \beta_\varepsilon, x, t) = \bar{\mathbf{H}}_\varepsilon(\alpha_\varepsilon, \beta_\varepsilon, x, t) = \mathbf{H}(\alpha_\varepsilon, \beta_\varepsilon, x, t).$$

This means that the cut-off does not affect right-hand sides of the truncated problem (3.8) if one restricts to the trajectory of the solution $(\alpha_\varepsilon, \beta_\varepsilon)$. By uniqueness of solutions to problem (3.8), we conclude $(\alpha_\varepsilon, \beta_\varepsilon) = (\mathbf{U}_\varepsilon, V_\varepsilon)$ and estimates (3.17) from Proposition 3.5 are also valid for the original error functions $(\mathbf{U}_\varepsilon, V_\varepsilon)$ on the domain Ω_T .

In the case of $p = \infty$, estimate (3.14) holds true and it remains to satisfy $\gamma > \delta_0$ such that inequality (3.19) is valid. The latter is equivalent to $\alpha > 1 - \delta_0$ and choosing $\alpha_0 = 1 - \delta_0$ yields the estimate

$$\|\mathbf{U}_\varepsilon\|_{L^\infty(\Omega_T)^m}, \|V_\varepsilon\|_{L^\infty(\Omega_T)} \leq C\varepsilon^{(1-\alpha)}.$$

For the case of $p < \infty$, in view of the results of Proposition 3.5, it remains to find triples (α, δ_0, r) such that

$$\delta_0 < 2(\alpha - 1) + \gamma \quad \text{and} \quad \delta_0 < 2(\alpha - 1) + (2 - \alpha)\frac{1}{r}$$

is satisfied where $\gamma = 2\delta_0 + (\alpha - 1)$. For existence of such triples, we define the following two restrictive curves (one depends on the parameter r as well)

$$\alpha > \ell(\delta_0) := 1 - \frac{1}{3}\delta_0 \quad \text{and} \quad \alpha > \ell_r(\delta_0) := \frac{r}{2r-1}\delta_0 + 1 - \frac{1}{2r-1}.$$

While ℓ is strictly monotone decreasing with $\ell(1/2) = 5/6 < 1$, the function ℓ_r is strictly increasing with $\ell_r(0) \in (0, 1)$ since $1 < r < \infty$. Thus, we always find such triples for small enough $\delta_0 > 0$.

Since these conditions are quite opaque, our goal is to further simplify inequalities (3.17) under consideration of the particular case $\delta_0 \leq 1/(2r)$. Recall that the intersection of ℓ_r and ℓ is given by one point for $\bar{\delta}_0 = \frac{3}{5r-1}$. In this case we have $\delta_0 < \bar{\delta}_0$ and the only restriction is given by $\ell(\delta_0) < \alpha < 1$. Notice that $\alpha > \ell(\delta_0)$ is equivalent to $\delta_0 > 3(1 - \alpha)$ and the assertion follows from estimates in Proposition 3.5. \square

Remark 3.7. Since the evolution systems \mathcal{U} and \mathcal{W} are uniformly bounded, the growth in time in estimates of Theorem 3.6 is polynomial. Similar estimates can be derived if the evolution systems are only uniformly bounded by some polynomial, as defined in [9, Definition 1.15], [14, Definition 2.7], we refer to [31, p. 63].

Remark 3.8. Similar to calculations in [39, Corollary 2, Proposition 5], we can obtain estimates for first-order derivatives of the errors. As this is a natural consequence of the weak formulation, we omit details.

Theorem 3.6 is an extension of [39, Theorem 3] to long time ranges that scale with the diffusion parameter $1/\varepsilon$. The above result enables us to check various models for uniform approximation on long-time scales, see for instance the model examples in Section 5 or [29]. In view of estimate (A.3), the mean value correction ψ_ε is of order ε for large times $t \geq T(\varepsilon)$ where

$$T(\varepsilon) := \max \left\{ 0, \frac{-\varepsilon \log (\|v^0 - \langle v^0 \rangle_\Omega\|_{L^\infty(\Omega)} \varepsilon)}{\lambda_1} \right\}.$$

This property allows to compare v_ε and v for large times. Moreover, the component v_ε becomes almost spatially homogeneous as time grows, compare [29, Theorem 3].

Corollary 3.9. *Let $(\mathbf{u}_\varepsilon, \mathbf{v}_\varepsilon)$ and (\mathbf{u}, v) be uniformly bounded, globally defined diffusive solutions of system (1.1) and a shadow solution of system (1.2), respectively. If $g - \langle g \rangle_\Omega$ is uniformly bounded in $\bar{\Omega} \times \mathbb{R}_{\geq 0}$, the error $V_\varepsilon = v_\varepsilon - v - \psi_\varepsilon$ satisfies the uniform estimate*

$$\|V_\varepsilon - \langle V_\varepsilon \rangle_\Omega\|_{L^\infty(\Omega \times \mathbb{R}_{\geq 0})} \leq C\varepsilon \tag{3.20}$$

for some $C > 0$ independent of diffusion.

Proof. Recall that $W_\varepsilon := V_\varepsilon - \langle V_\varepsilon \rangle_\Omega$ solves

$$\frac{\partial W_\varepsilon}{\partial t} - \frac{1}{\varepsilon} \Delta W_\varepsilon = g(\mathbf{u}_\varepsilon, v_\varepsilon, x, t) - g(\mathbf{u}, v, x, t) - \langle g(\mathbf{u}_\varepsilon, v_\varepsilon, \cdot, t) - g(\mathbf{u}, v, \cdot, t) \rangle_\Omega \quad \text{in } \Omega \times \mathbb{R}_{> 0}$$

endowed with homogeneous zero flux boundary and zero initial conditions. Uniform boundedness of the right-hand side in the latter equation yields estimate (3.20) by Proposition A.2 and decay estimate (A.3). \square

We emphasize that, in general, the stability Assumption 6–7 is required to obtain long-time estimates. To see this, we refer to Model 5.1 and [32, Example 2]. As shown in [32, Example 3], a convergence result on long-time scales as in Theorem 3.6 does not have to extend to a global convergence result. However, a stronger version of Assumption 6 leads to global-in-time error estimates. Consider again the evolution system \mathcal{W} defined by equation (3.13) on $L^p(\Omega)^m \times \mathbb{R}$ due to the linearization at the shadow solution.

Assumption 8 (Exponential stability). *Let the shadow evolution system \mathcal{W} defined by (3.13) be uniformly exponentially stable for some $1 \leq p \leq \infty$. This means that there exists a constant $C > 0$ and an exponent $\sigma > 0$, both independent of time, such that for all $s, t \in \mathbb{R}_{\geq 0}$, $s \leq t$ there holds*

$$\|\mathbf{W}(t, s)\xi^0\|_{L^p(\Omega)^m \times \mathbb{R}} \leq Ce^{-\sigma(t-s)}\|\xi^0\|_{L^p(\Omega)^m \times \mathbb{R}} \quad \forall \xi^0 \in L^p(\Omega)^m \times \mathbb{R}.$$

For the case of $p = \infty$, this assumption leads to global error estimates as the following analog of Proposition 3.4 already indicates.

Corollary 3.10. *Let assumptions 1–8 hold with some $1 \leq p \leq \infty$ and uniform exponential stability exponent $\sigma > 0$. Then, for any $\delta_0 \in (0, 1/2]$, there exist constants $C, \varepsilon_0 > 0$ independent of diffusion such that for all $\varepsilon \leq \varepsilon_0$ the solution $(\alpha_\varepsilon, \beta_\varepsilon)$ of the truncated problem (3.8) satisfies*

$$\|\alpha_\varepsilon\|_{L^p(\Omega \times \mathbb{R}_{\geq 0})^m} + \|\beta_\varepsilon\|_{L^p(\Omega \times \mathbb{R}_{\geq 0})} \leq C\varepsilon^{2\delta_0}. \quad (3.21)$$

Proof. We already estimated the function $W_\varepsilon = V_\varepsilon - \langle V_\varepsilon \rangle_\Omega$ in inequality (3.15). The corresponding inequality for $\alpha_\varepsilon, b_\varepsilon$ reads

$$\begin{aligned} \|\alpha_\varepsilon(\cdot, t)\|_{L^p(\Omega)^m} + |\Omega|^{1/p}|b_\varepsilon(t)| &\leq C \int_0^t e^{-\sigma(t-\tau)} (\|(W_\varepsilon + \psi_\varepsilon)(\cdot, \tau)\|_{L^p(\Omega)} + \|\mathbf{H}_\varepsilon(\cdot, \tau)\|_{L^p(\Omega)^{m+1}}) d\tau \\ &\leq C \int_0^t e^{-\sigma(t-\tau)} \|W_\varepsilon(\cdot, \tau)\|_{L^p(\Omega)} d\tau + C|\Omega|^{1/p}\varepsilon^{2\delta_0} \\ &\quad + C \int_0^t e^{-\sigma(t-\tau)} \mathbb{1}_{\{\tau \leq -\varepsilon \log \varepsilon / \lambda_1\}} |\Omega|^{1/p}|b_\varepsilon(\tau)| d\tau. \end{aligned}$$

Absorption of the function b_ε follows as in Proposition 3.4. This results in

$$\sup_{t \in [0, T]} \left(\|\alpha_\varepsilon(\cdot, t)\|_{L^p(\Omega)^m} + |\Omega|^{1/p}|b_\varepsilon(t)| \right) \leq C \int_0^T e^{-\sigma(t-\tau)} \|W_\varepsilon(\cdot, \tau)\|_{L^p(\Omega)} d\tau + C|\Omega|^{1/p}\varepsilon^{2\delta_0},$$

where we restrict ourselves to exponents $\delta_0 \leq 1/2$ and $\varepsilon \leq \varepsilon_0$. Combining above inequalities leads to an estimate for W_ε , more precisely, $\sup_{t \in [0, T]} \|W_\varepsilon(\cdot, t)\|_{L^p(\Omega)} \leq C\varepsilon$. Since $T > 0$ is arbitrary, we obtain a global estimate for the truncated solution. \square

The global analog of Theorem 3.6 is given next.

Theorem 3.11. *Let Assumption 1–8 hold for some p with $p \geq 1 = n$ or $p > n/2$ if $n \geq 2$ and some uniform exponential stability exponent $\sigma > 0$ of the evolution system \mathcal{W} . Let r be given by (A.7). Then there exist bounds $\alpha_0 = \alpha_0(r) \in (0, 1)$, $\varepsilon_0 > 0$ and a constant $C > 0$ such that for any $\alpha \in [\alpha_0, 1)$, $\varepsilon \leq \varepsilon_0$ and $T \leq \varepsilon^{\alpha-1}$ we have*

$$\begin{aligned} \|\mathbf{u}_\varepsilon - \mathbf{u}\|_{L^\infty(\Omega_T)^m} &\leq C\varepsilon^{3(1-\alpha)}, \\ \|v_\varepsilon - v - \psi_\varepsilon\|_{L^\infty(\Omega_T)} &\leq C\varepsilon^{4(1-\alpha)}, \\ \|\langle v_\varepsilon \rangle_\Omega - v\|_{L^\infty((0, T))} &\leq C\varepsilon^{4(1-\alpha)}. \end{aligned} \quad (3.22)$$

Concerning $p = \infty$, we have global error estimates

$$\|\mathbf{u}_\varepsilon - \mathbf{u}\|_{L^\infty(\Omega \times \mathbb{R}_{\geq 0})^m} + \|v_\varepsilon - v - \psi_\varepsilon\|_{L^\infty(\Omega \times \mathbb{R}_{\geq 0})} \leq C\varepsilon \quad (3.23)$$

without requiring Assumption 7.

Proof. To get rid of truncation, it is sufficient to verify (3.19) as in the proof of Theorem 3.6. If $p = \infty$, we can choose $2\delta_0 = 1$ in estimate (3.21) and remove the cut-off due to $2\delta_0 > \delta_0$ to obtain the global estimate. For the case of $p < \infty$, we restrict our time interval in view of the parabolic bootstrap method from Proposition A.3. By Corollary 3.10, we obtain

$$\|\beta_\varepsilon\|_{L^\infty(\Omega_T)} \leq \varepsilon^{-(1-\alpha)} \left(\varepsilon^{2\delta_0} + \varepsilon^{(2-\alpha)/r} \right), \quad \|\alpha_\varepsilon\|_{L^\infty(\Omega_T)} \leq \varepsilon^{-(1-\alpha)} \|\beta_\varepsilon\|_{L^\infty(\Omega_T)}$$

for all times $T \leq \varepsilon^{\alpha-1}$. To dispose of truncation, it remains to find triples (α, δ_0, r) such that the inequality

$$\delta_0 < 2(\alpha - 1) + 2\delta_0 \quad \text{and} \quad \delta_0 < 2(\alpha - 1) + (2 - \alpha)\frac{1}{r}$$

is satisfied. Existence of such triples can be shown as in the proof of Theorem 3.6. \square

Remark 3.12. If the evolution system \mathcal{U} satisfies an additional exponential stability condition in Assumption 7, then all components of \mathbf{U}_ε have the same convergence rate as V_ε . Recall that the restriction in time in above estimates is due to exponential stability for finite $p < \infty$. However, the case $p = \infty$ can be regarded in many applications, see Section 5.

Remark 3.13. We refer to [31, Ch. 7] for possible directions of the presented theory to shadow systems with other boundary conditions or other differential operators. Due to flexibility of the semigroup framework used in this work, we may also consider classical shadow systems and reaction-diffusion systems in which only some diffusion parameters tend to infinity, see Section 4.

4. Reaction-diffusion systems

We want to emphasize that the same stability principle applies to (classical) shadow systems where some components of the shadow limit \mathbf{u} diffuse. Classical shadow systems are considered in [15, 27, 29, 41] as well as recently in [23] in the context of control theory.

Let $\mathbf{D} \in \mathbb{R}_{>0}^{m \times m}$ be a diagonal matrix with non-negative entries $D_i \geq 0$ for $i = 1, \dots, m$ and consider instead of (1.2) the shadow system

$$\begin{aligned} \frac{\partial \mathbf{u}}{\partial t} - \mathbf{D} \Delta \mathbf{u} &= \mathbf{f}(\mathbf{u}, v, x, t) & \text{in } \Omega_T, & \quad \mathbf{u}(\cdot, 0) = \mathbf{u}^0 \text{ in } \Omega, & \quad \frac{\partial \mathbf{u}}{\partial \nu} = 0 \text{ on } \partial\Omega \times (0, T), \\ \frac{dv}{dt} &= \langle g(\mathbf{u}(\cdot, t), v(t), \cdot, t) \rangle_\Omega & \text{in } (0, T), & \quad v(0) = \langle v^0 \rangle_\Omega. \end{aligned} \quad (4.1)$$

As in the proof of Theorem 2.2, under Assumption 1-2, a Picard iteration yields a unique mild solution of the integral equations for $i = 1, \dots, m$

$$\begin{aligned} \mathbf{u}_i(\cdot, t) &= S_\Delta(D_i t) \mathbf{u}_i^0 + \int_0^t S_\Delta(D_i(t-s)) \mathbf{f}_i(\mathbf{u}(\cdot, s), v(s), \cdot, s) ds, \\ v(t) &= \langle v^0 \rangle_\Omega + \int_0^t \langle g(\mathbf{u}(\cdot, s), v(s), \cdot, s) \rangle_\Omega ds. \end{aligned}$$

This solution satisfies $\mathbf{u} \in L^\infty(\Omega_T)^m$, $v \in C([0, T])$ for all $T < T_{\max}$, where T_{\max} is the maximal time of existence of (\mathbf{u}, v) . Let us note that, in general, diffusive components are not continuous up to $t = 0$ with respect to $L^\infty(\Omega)$ [46, Part I, Lemma 2]. We additionally make Assumption 3-4.

The reaction-diffusion problem has a similar structure. We modify the components of \mathbf{u}_ε for $i = 1, \dots, m$,

$$\mathbf{u}_{\varepsilon, i}(\cdot, t) = S_\Delta(D_i t) \mathbf{u}_i^0 + \int_0^t S_\Delta(D_i(t-s)) f_i(\Psi(\cdot, s), \cdot, s) ds,$$

as well as the diffusion matrix \mathbf{D}_ε in problem (2.1). Thus, a unique local-in-time mild solution exists as in Theorem 2.2 with $(\mathbf{u}_\varepsilon, v_\varepsilon) \in L^\infty(\Omega_T)^{m+1}$ for all $T < T_{\max}(\varepsilon)$.

The cut-off can be performed in the same way under Assumption 5. The evolution system \mathcal{W} induced by the linearization is determined by the unique solution of the homogeneous linear shadow problem

$$\begin{aligned} \frac{\partial \xi_1}{\partial t} - \mathbf{D}\Delta \xi_1 &= \mathbf{A}(x, t)\xi_1 + \mathbf{B}(x, t)\xi_2 & \text{in } \Omega \times \mathbb{R}_{>0}, & \quad \xi_1(\cdot, 0) = \xi_1^0 \text{ in } \Omega, \\ \frac{d\xi_2}{dt} &= \langle \mathbf{C}(\cdot, t)\xi_1 \rangle_\Omega + \langle D(\cdot, t)\xi_2 \rangle_\Omega & \text{in } \mathbb{R}_{>0}, & \quad \xi_2(0) = \langle \xi_2^0 \rangle_\Omega \end{aligned} \quad (4.2)$$

endowed with zero Neumann boundary conditions for diffusive components of ξ_1 . Using Assumption 6, we obtain the result of Proposition 3.4. As all diffusive components can be treated with the bootstrap argument which makes use of Proposition A.3, Assumption 7 may be reduced to the subsystem of non-diffusive components. Therefore, we delete all rows and columns of $\mathbf{A}(\cdot, t)$ for which the diffusion is positive to obtain $\tilde{\mathbf{A}}(\cdot, t)$. In this way, we obtain the following analogon of the reaction-diffusion-ODE case.

Theorem 4.1. *Let the assumptions 1–6 hold for some p with $p \geq 1 = n$ or $p > n/2$ if $n \geq 2$ and let $r \in (1, \infty)$ given by (A.7). Furthermore, let Assumption 7 hold for the submatrix $\tilde{\mathbf{A}}(\cdot, t)$ of non-diffusive components. Then the results of Theorem 3.6 remain valid for system (4.1). Under the stronger exponential stability assumption 6, estimates established in Theorem 3.11 remain valid.*

Remark 4.2. Note that in the classical case, where all components of \mathbf{u} diffuse, we assume only Assumption 6. Furthermore, Corollary 3.9 can be adapted accordingly.

Proposition 3.4 and Theorem 3.6 can be proven in a similar way for the Hilbertian case $p = 2$ using energy estimates. The method uses error estimates for L^2 dissipative shadow systems – a smaller class of systems for which uniform boundedness of the corresponding evolution system \mathcal{W} is often easier to verify. In particular, one reduces considerations to the linearized shadow system without diffusion. This consideration is especially helpful when we look at classical shadow systems. Let us first consider the evolutionary subsystem \mathcal{U} from Assumption 7 to describe the principle of dissipativity. Concerning contractivity in the time-independent case, [11, Ch. III, Theorem 2.7] applies the condition

$$\|\mathbf{y}\|_{L^p(\Omega)^m} \leq \|(I - \lambda \mathbf{A})\mathbf{y}\|_{L^p(\Omega)^m} \quad \forall \lambda \in \mathbb{R}_{>0}, \mathbf{y} \in \mathbf{L}^p(\Omega)^m. \quad (4.3)$$

The time-dependent case can be treated in the same way, following [35]. Therefore, consider the bounded multiplication operators $\mathbf{A}(\cdot, t)$ in the setting (3.9), and let us assume

Assumption 9 (Dissipativity of ODE subsystem). *Let $1 \leq p \leq \infty$ and $\kappa : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ be a continuous function with $\kappa \in L^1(\mathbb{R}_{\geq 0})$ such that*

$$(1 - \lambda \kappa(t))\|\mathbf{y}\|_{L^p(\Omega)^m} \leq \|(I - \lambda \mathbf{A}(\cdot, t))\mathbf{y}\|_{L^p(\Omega)^m} \quad (4.4)$$

is satisfied for all $\mathbf{y} \in L^p(\Omega)^m$, $\lambda \in \mathbb{R}_{>0}$, and $t \in \mathbb{R}_{\geq 0}$.

Note that $\kappa \equiv 0$ corresponds to [26, Theorem 1]. Before providing equivalent formulations of dissipativity condition (4.4), let us state the well-known implication of this assumption.

Proposition 4.3. *Let $\mathbf{A} : \Omega \times \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}^{m \times m}$ be a measurable, locally bounded matrix-valued function satisfying Assumption 9. Then the corresponding evolution system \mathcal{U} induced by $\mathbf{A}(\cdot, t)$ is uniformly bounded on $L^p(\Omega)^m$ for all exponents $1 \leq p \leq \infty$.*

Proof. See [44, Remark 2.2, p. 119]. □

By [44, Remark 1.2], condition (4.4) is equivalent to dissipativity of $\mathbf{A}(\cdot, t) - \kappa(t)I$ on $L^p(\Omega)^m$ in the sense of inequality (4.3) for each time $t \in \mathbb{R}_{\geq 0}$. It turns out that there is even a simpler criterion for dissipativity

of multiplication operators on $L^p(\Omega)^m$ which is independent of the exponent $1 \leq p \leq \infty$. More precisely, inequality (4.4) can be verified via pointwise estimates of the corresponding quadratic form

$$Q(x, t) : \mathbb{R}^m \rightarrow \mathbb{R}, \quad \bar{\mathbf{y}} \mapsto \bar{\mathbf{y}}^T (\mathbf{A}(x, t) - \kappa(t)I) \bar{\mathbf{y}}. \quad (4.5)$$

Such a condition is already used in the time-independent case in [42, Propositions 6, 7]. Moreover, pointwise estimates of the latter quadratic form are a well-known technique in the context of classical solutions to preserve contractivity of the corresponding evolution system [33, Theorem 2.3].

Lemma 4.4. *Let κ be given by Assumption 9 for some measurable, bounded function $\mathbf{A} : \Omega \times \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}^{m \times m}$ and let Q be defined as above in (4.5). Then, dissipativity condition (4.4) on $L^p(\Omega)^m$ for some $1 \leq p \leq \infty$ is equivalent to $Q(x, t) \leq 0$ for a.e. $(x, t) \in \Omega \times \mathbb{R}_{\geq 0}$. Moreover, inequality (4.4) holds for all $1 \leq p \leq \infty$ if and only if it holds for one exponent p .*

Proof. By [44, Remark 1.2], estimate (4.4) is equivalent to dissipativity of $\mathbf{A}(\cdot, t) - \kappa(t)I$. The latter means that for all $\lambda \in \mathbb{R}_{>0}$, $\mathbf{y} \in L^p(\Omega)^m$, and $t \in \mathbb{R}_{\geq 0}$ there holds

$$\|\mathbf{y}\|_{L^p(\Omega)^m} \leq \|(I - \lambda(\mathbf{A}(\cdot, t) - \kappa(t)I))\mathbf{y}\|_{L^p(\Omega)^m}. \quad (4.6)$$

A further characterization of dissipativity via the duality map J as in [11, Ch. II, Proposition 3.23] can be used to rewrite this condition. For $1 < p < \infty$, the duality set is just a singleton by [11, Ch. II, Example 3.26 (ii)]. More precisely, $J(\mathbf{y}) = \{\mathbf{y}^*\}$ for $\mathbf{y}^* \in L^q(\Omega)^m$ where $\mathbf{y}^* = \mathbf{0}$ for $\mathbf{y} = \mathbf{0}$ and

$$\mathbf{y}^* = \frac{\mathbf{y}|\mathbf{y}|^{p-2}}{\|\mathbf{y}\|_{L^p(\Omega)^m}^{p-2}} \quad \text{for } \mathbf{y} \neq \mathbf{0}.$$

Remember that the dual pairing satisfies $\langle \mathbf{y}^*, \mathbf{y} \rangle = \|\mathbf{y}^*\|_{L^q(\Omega)^m}^2 = \|\mathbf{y}\|_{L^p(\Omega)^m}^2$ where p and q are conjugate exponents. Hence, inequalities (4.4)–(4.6) are equivalent to integral condition

$$\int_{\Omega} (\mathbf{y}^*)^T (\mathbf{A}(\cdot, t) - \kappa(t)I) \mathbf{y} \, dx \leq 0 \quad \forall \mathbf{y} \in L^p(\Omega)^m, t \in \mathbb{R}_{\geq 0}. \quad (4.7)$$

Let us first assume $Q \leq 0$ for a.e. $(x, t) \in \Omega \times \mathbb{R}_{\geq 0}$. We multiply the inequality $Q \leq 0$ with a symmetric choice of vectors $\bar{\mathbf{y}}(x)$ instead of \mathbf{y}^* and \mathbf{y} and obtain inequality (4.7) for $1 < p < \infty$ after integration. For $p \in \{1, \infty\}$, we use continuity of the L^p norm with respect to p since we already established estimate (4.6) for all $1 < p < \infty$. Since $\mathbf{A} - \kappa I$ is bounded, $I - \lambda(\mathbf{A}(t) - \kappa(t)I)$ is invertible for small $\lambda > 0$, and by [11, Ch. II, Proposition 3.14], the latter operator is invertible for all $\lambda > 0$ and estimate (4.6) yields

$$\|(I - \lambda(\mathbf{A}(\cdot, t) - \kappa(t)I))^{-1}\mathbf{y}\|_{L^p(\Omega)^m} \leq \|\mathbf{y}\|_{L^p(\Omega)^m}$$

for all $\mathbf{y} \in L^p(\Omega)^m$ and $1 < p < \infty$. The result follows by letting $p \rightarrow 1$ or $p \rightarrow \infty$ where [1, Theorem 2.14] applies. Thus, $Q \leq 0$ implies dissipativity.

Now, let dissipativity inequality (4.4) be fulfilled and assume Q is not non-positive, i.e., there is a set $\Omega_1 \subset \Omega$ with $|\Omega_1| > 0$ and some time point $t \geq 0$ as well as $\bar{\mathbf{y}} \in \mathbb{R}^m \setminus \{\mathbf{0}\}$ such that

$$Q(x, t)\bar{\mathbf{y}} = \bar{\mathbf{y}}^T (\mathbf{A}(x, t) - \kappa(t)I)\bar{\mathbf{y}} > 0$$

holds for a.e. $x \in \Omega_1$. Using measure theory, we find uniform bounds $0 < Q_0 \leq Q(\cdot, t)\bar{\mathbf{y}} \leq Q_1 < \infty$ almost everywhere on a possibly smaller set $\Omega_2 \subset \Omega_1$ with positive measure. Let us consider $\mathbf{y}_p \in L^p(\Omega)^m$ for $p < \infty$ given by $\mathbf{y}_p(x) := \mathbb{1}_{\Omega_2}(x)(Q(x, t)\bar{\mathbf{y}})^{-1/p}\bar{\mathbf{y}}$. This bounded vector-valued function satisfies

$$\int_{\Omega} (\mathbf{y}_p^*)^T (\mathbf{A}(\cdot, t) - \kappa(t)I) \mathbf{y}_p \, dx = \int_{\Omega_2} |\bar{\mathbf{y}}|^{p-2} \|\mathbf{y}_p\|_{L^p(\Omega_2)^m}^{2-p} \, dx = \|(Q(\cdot, t)\bar{\mathbf{y}})^{-1}\|_{L^p(\Omega_2)^m}^{2-p} > 0$$

which is a contradiction to condition (4.7), and thus to (4.4) for $1 < p < \infty$. For $p \in \{1, \infty\}$, let us consider $\mathbf{y}_2 \in L^\infty(\Omega)^m$ in preceding definition. Then

$$|(I - \lambda(\mathbf{A}(\cdot, t) - \kappa(t)I))\mathbf{y}_2|^2 = |\mathbf{y}_2|^2 + \lambda^2 |(\mathbf{A}(\cdot, t) - \kappa(t)I)\mathbf{y}_2|^2 - 2\lambda$$

holds on the set Ω_2 . For small enough $\lambda > 0$, the right-hand side of the latter equation is smaller than $|\mathbf{y}_2|^2$ since $\mathbf{A}, \kappa, \mathbf{y}_2$ are bounded functions on Ω_2 . This leads to a contradiction to condition (4.6). \square

Hence, Assumption 9 is independent of $1 \leq p \leq \infty$. Note that Assumption 7 only considers the subsystem of \mathbf{A} of non-diffusing components, for which we can verify dissipativity in $L^p(\Omega)^m$ for some (and hence all) $1 \leq p \leq \infty$. Since \mathbf{A} is non-symmetric in general, one can verify definiteness of the corresponding quadratic form Q defined in (4.5) by looking equivalently on the real eigenvalues of the symmetric part

$$\frac{1}{2} (\mathbf{A}(x, t) + \mathbf{A}(x, t)^T) - \kappa(t)I \in \mathbb{R}^{m \times m}.$$

Non-positivity of its eigenvalues $\lambda(x, t)$ pointwise for a.e. $(x, t) \in \Omega \times \mathbb{R}_{\geq 0}$ implies the condition $Q \leq 0$.

A similar dissipativity assumption can be assumed for the linearized shadow evolution system \mathcal{W} . Using the notation $\mathbf{D}^S := \text{diag}(\mathbf{D}, 0)$, let us rewrite the shadow problem (4.2) as an ordinary differential equation in the Banach space $L^p(\Omega)^m \times \mathbb{R}$,

$$\frac{d}{dt} \xi = \mathbf{D}^S \Delta \xi + \mathbf{L}_0(t) \xi \quad \text{in } \mathbb{R}_{>0}, \quad \xi(0) = \begin{pmatrix} \xi_1^0 \\ \langle \xi_2^0 \rangle_\Omega \end{pmatrix}.$$

By Assumption 5, $(\mathbf{L}_0(t))_{t \in \mathbb{R}_{\geq 0}}$ is a family of bounded shadow operators on $L^p(\Omega)^m \times \mathbb{R}$. The full operator $\mathbf{D}^S \Delta + \mathbf{L}_0(t)$ can be seen as a perturbation of the matrix operator $\mathbf{D}^S \Delta$ which generates a contraction semigroup on $L^p(\Omega)^m \times \mathbb{R}$ for all $1 \leq p \leq \infty$. Let us assume in analogy to Assumption 9

Assumption 10 (Dissipativity of shadow system). *Let $\varrho : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ be a continuous function such that $\varrho \in L^1(\mathbb{R}_{\geq 0})$ and*

$$(1 - \lambda \varrho(t)) \|\mathbf{y}\|_{L^p(\Omega)^m \times \mathbb{R}} \leq \|(I - \lambda \mathbf{L}_0(t))\mathbf{y}\|_{L^p(\Omega)^m \times \mathbb{R}} \quad (4.8)$$

is satisfied for all $\mathbf{y} \in L^p(\Omega)^m \times \mathbb{R}$, $\lambda \in \mathbb{R}_{>0}$, and $t \in \mathbb{R}_{\geq 0}$.

Using the duality map, one obtains an equivalent integral inequality of the form (4.7) for $1 < p < \infty$. The latter has a quite convenient form for L^2 energy estimates:

$$\int_{\Omega} \mathbf{y}^T (\mathbf{L}_0(\cdot, t) - \varrho(t)I) \mathbf{y} \, dx \leq 0 \quad \forall \mathbf{y} \in L^2(\Omega)^m \times \mathbb{R}, t \in \mathbb{R}_{\geq 0}$$

Proposition 4.5. *Let the linear operators $\mathbf{L}_0(t) : L^p(\Omega)^m \times \mathbb{R} \rightarrow L^p(\Omega)^m \times \mathbb{R}$ defined above for each $t \in \mathbb{R}_{\geq 0}$ satisfy Assumption 10 for some $1 \leq p \leq \infty$ and uniformly bounded coefficients $\mathbf{A}, \mathbf{B}, \mathbf{C}$ and D . Then the corresponding shadow evolution system \mathcal{W} induced by $\mathbf{D}^S \Delta + \mathbf{L}_0(t)$ is uniformly bounded on $L^p(\Omega)^m \times \mathbb{R}$ for the same exponent p , thus \mathcal{W} satisfies Assumption 6. Moreover, Assumption 7 is satisfied for all $1 \leq p \leq \infty$.*

Proof. By the same reasoning as in the proof of Proposition 4.3, we obtain uniform boundedness of the evolution system \mathcal{W} for the chosen exponent p . Let us now focus on the evolutionary subsystem \mathcal{U} . In view of Lemma 4.4 and Proposition 4.3, it remains to check dissipativity condition for the chosen value p . For $1 < p < \infty$, we infer from condition (4.7) that dissipativity of $\mathbf{L}_0(t) - \varrho(t)I$ in $L^p(\Omega)^m \times \mathbb{R}$ implies dissipativity of the corresponding subsystem $\mathbf{A}(\cdot, t) - \varrho(t)I$ in $L^p(\Omega)^m$ since the duality set is given by $\mathbf{y}^* = (\mathbf{y}_1^*, \mathbf{y}_2^*)^T$ for $\mathbf{y} = (\mathbf{y}_1, \mathbf{y}_2)^T \in L^p(\Omega)^m \times \mathbb{R}$. For $p \in \{1, \infty\}$, we follow the proof of Lemma 4.4. In this way, condition (4.6) can be shown by contradiction, assuming $Q \leq 0$ does not hold almost everywhere. \square

Let us conclude this section with a comment on exponential stability used in Theorem 3.11.

Remark 4.6. Assume that for some $\sigma \in \mathbb{R}_{>0}$ and non-negative $\kappa \in C(\mathbb{R}_{\geq 0}) \cap L^1(\mathbb{R}_{\geq 0})$ the stronger condition

$$(1 - \lambda(\varrho(t) - \sigma)) \|\mathbf{y}\|_{L^p(\Omega)^m \times \mathbb{R}} \leq \|(I - \lambda \mathbf{L}_0(t))\mathbf{y}\|_{L^p(\Omega)^m \times \mathbb{R}} \quad (4.9)$$

is satisfied for all $\mathbf{y} \in L^p(\Omega)^m \times \mathbb{R}$, $\lambda \in \mathbb{R}_{>0}$, and $t \in \mathbb{R}_{\geq 0}$. This condition implies uniform exponential stability of the evolutionary system \mathcal{W} in $L^p(\Omega)^m \times \mathbb{R}$ with exponent $\sigma > 0$ in Assumption 8.

5. Model examples

In this section, we provide two examples. The first example is a linear model which shows necessity of the stability assumption 6 in the case of a space-dependent shadow solution. The second model which is treated analytically is of predator-prey-type and exemplifies the global convergence result in Theorem 3.11.

5.1. Linear model

Let us focus on an equation for v_ε only,

$$\frac{\partial v_\varepsilon}{\partial t} - \frac{1}{\varepsilon} \Delta v_\varepsilon = D(x)v_\varepsilon \quad \text{in } \Omega \times \mathbb{R}_{>0}, \quad v_\varepsilon(\cdot, 0) = v^0 \quad \text{in } \Omega, \quad \frac{\partial v_\varepsilon}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times \mathbb{R}_{>0}, \quad (5.1)$$

for a space-dependent coefficient $D := w_1 + w_1^2 \in L^\infty(\Omega)$. Here, we take the eigenfunction $v^0 := w_1$ which corresponds to the first positive eigenvalue λ_1 of $-\Delta$ on $\Omega = (0, 1)$, i.e., $w_1(x) = \sqrt{2} \cos(\pi x)$ by equation (A.2). Of course, equation (5.1) can be extended to a full linear reaction-diffusion-ODE system by setting $A, B, C = 0$ in (3.9). The corresponding shadow limit is given by $v = 0$ since $\langle v^0 \rangle_\Omega = 0$ and $\langle D \rangle_\Omega = 1$. For this reason, Assumption 6 is not satisfied. We will verify that the error $V_\varepsilon = v_\varepsilon - \psi_\varepsilon$ grows exponentially in time. By Hölder's inequality, it remains to show exponential growth of $\langle V_\varepsilon \rangle_\Omega = \langle v_\varepsilon \rangle_\Omega$. The function v_ε is given by

$$v_\varepsilon(x, t) = S_\Delta(t/\varepsilon)v^0(x) + \int_0^t S_\Delta((t - \tau)/\varepsilon)D(x)v_\varepsilon(x, \tau) \, d\tau.$$

This implicit integral equation can be solved by a Picard iteration. According to [46, Part II, Theorem 1], we define approximations $v_\varepsilon^{(j)}(\cdot, t) \in L^\infty(\Omega)$ recursively given by

$$\begin{aligned} v_\varepsilon^{(1)}(\cdot, t) &= S_\Delta(t/\varepsilon)v^0 = e^{-\lambda_1 t/\varepsilon}w_1, \\ v_\varepsilon^{(j+1)}(\cdot, t) &= S_\Delta(t/\varepsilon)v^0 + \int_0^t S_\Delta((t - \tau)/\varepsilon) \left[D(\cdot)v_\varepsilon^{(j)}(\cdot, \tau) \right] \, d\tau. \end{aligned}$$

Using trigonometry, we write $D = w_1 + w_1^2 = w_0 + w_1 + \sqrt{2}^{-1}w_2$. We iteratively multiply the coefficient D with $v_\varepsilon^{(j)}$ and use that products $w_j w_i$ can be written as a linear combination of w_{j+i} and $w_{|j-i|}$. This procedure yields

$$\begin{aligned} v_\varepsilon^{(2)}(\cdot, t) &= e^{-\lambda_1 t/\varepsilon}w_1 + \int_0^t S_\Delta((t - \tau)/\varepsilon)f_1^{(1)}(\tau) \left[w_1 + w_1^2 + \sqrt{2}^{-1}w_1w_2 \right] \, d\tau \\ &= \left(\int_0^t f_1^{(1)}(\tau) \, d\tau \right) w_0 + \left(e^{-\lambda_1 t/\varepsilon} + \int_0^t e^{-\lambda_1(t-\tau)/\varepsilon} f_1^{(1)}(\tau) \, d\tau \right) w_1 \\ &\quad + \int_0^t S_\Delta((t - \tau)/\varepsilon)f_1^{(1)}(\tau)h^{(2)} \, d\tau, \end{aligned}$$

where $v_\varepsilon^{(1)}(\cdot, t) =: f_1^{(1)}(t)w_1$, $h^{(2)} = (w_1 + \sqrt{2}w_2 + w_3)/2$ and $w_0 \equiv 1$. To understand the next step, let us rewrite the second approximation as

$$v_\varepsilon^{(2)}(\cdot, t) = f_0^{(2)}(t)w_0 + f_1^{(2)}(t)w_1 + f_2^{(2)}(t)h^{(2)}$$

and note that the coefficients of the eigenfunctions in $h^{(2)}$ are all positive and $h^{(2)}$ includes w_1 as well. Considering spatial means, $\langle w_j \rangle_\Omega = 0$ for all $j \in \mathbb{N}$ implies $\langle v_\varepsilon^{(1)} \rangle_\Omega = 0$ and

$$\langle v_\varepsilon^{(2)}(\cdot, t) \rangle_\Omega = \int_0^t e^{-\lambda_1 \tau / \varepsilon} d\tau.$$

Using again $d = w_0 + w_1 + \sqrt{2}^{-1} w_2$, this leads to the third approximation of the form

$$\begin{aligned} v_\varepsilon^{(3)}(\cdot, t) &= \left(\int_0^t f_0^{(2)}(\tau) + f_1^{(2)}(\tau) d\tau \right) w_0 \\ &\quad + \left(e^{-\lambda_1 t / \varepsilon} + \int_0^t e^{-\lambda_1(t-\tau) / \varepsilon} (f_0^{(2)}(\tau) + f_1^{(2)}(\tau)) d\tau \right) w_1 + f_3^{(3)}(t) h^{(3)}, \end{aligned}$$

where $h^{(3)}$ is a sum of positive multiples of w_j for $j = 0, \dots, 4$ and $f_3^{(3)} \geq 0$ is a continuous function in time. Estimating from below, we successively gain for all $j \in \mathbb{N}$ (by setting $f_0^{(1)} \equiv 0$)

$$f_0^{(j+2)}(t) \geq \int_0^t f_0^{(j+1)}(\tau) + f_1^{(j+1)}(\tau) d\tau \geq \int_0^t \left(e^{-\lambda_1 \tau / \varepsilon} + \int_0^\tau f_0^{(j)}(r) + f_1^{(j)}(r) dr \right) d\tau.$$

Starting from the innermost double integral and applying Fubini's rule inductively, this yields

$$f_0^{(j+2)}(t) \geq \int_0^t f_0^{(j+1)}(\tau) + f_1^{(j+1)}(\tau) d\tau \geq \int_0^t \sum_{i=0}^j \frac{(t-\tau)^i}{i!} e^{-\lambda_1 \tau / \varepsilon} d\tau.$$

Since $v_\varepsilon^{(j)}$ converges to v_ε in $L^\infty(\Omega_T)$ as shown in [46, Part II, Theorem 1], we obtain a lower bound due to

$$\int_0^t \sum_{i=0}^j \frac{(t-\tau)^i}{i!} e^{-\lambda_1 \tau / \varepsilon} d\tau \leq f_0^{(j+2)}(t) \leq \langle v_\varepsilon^{(j+2)}(\cdot, t) \rangle_\Omega \rightarrow \langle v_\varepsilon(\cdot, t) \rangle_\Omega.$$

The theorem of monotone convergence leads to exponential growth of $\langle V_\varepsilon \rangle_\Omega$ since

$$\langle v_\varepsilon(\cdot, t) \rangle_\Omega \geq \int_0^t e^{t-\tau} e^{-\lambda_1 \tau / \varepsilon} d\tau \geq C\varepsilon (e^t - 1).$$

This induces exponential growth of $t \mapsto \|v_\varepsilon(\cdot, t)\|_{L^\infty(\Omega)}$ by Hölder's inequality.

5.2. Predator-prey model

Consider a closed system describing predator-prey dynamics, with a predator denoted by u_ε and a mobile prey v_ε . In fresh-water ecology, a biological example can be given by Hydra and Daphnia where the predator Hydra is sedentary, i.e., $D = 0$ [40, Example (b)]. The following model adapted from [40] includes both cases $D = 0$ and $D > 0$. The differential equations read

$$\begin{aligned} \frac{\partial u_\varepsilon}{\partial t} - D\Delta u_\varepsilon &= -pu_\varepsilon + bv_\varepsilon & \text{in } \Omega_T, & \quad u_\varepsilon(\cdot, 0) = u^0 \text{ in } \Omega, \\ \frac{\partial v_\varepsilon}{\partial t} - \frac{1}{\varepsilon}\Delta v_\varepsilon &= (d - au_\varepsilon - cv_\varepsilon)v_\varepsilon & \text{in } \Omega_T, & \quad v_\varepsilon(\cdot, 0) = v^0 \text{ in } \Omega, \quad \frac{\partial v_\varepsilon}{\partial \nu} = 0 \text{ on } \partial\Omega \times (0, T), \end{aligned} \tag{5.2}$$

where u_ε is endowed with a zero flux boundary condition if $D > 0$. Here, $a, b, c, d, p > 0$ are constants and the initial values $u^0, v^0 \geq 0$ satisfy Assumption 2 as well as non-negativity almost everywhere in Ω with $\langle v^0 \rangle_\Omega > 0$. The corresponding shadow limit is given by

$$\begin{aligned} \frac{\partial u}{\partial t} - D\Delta u &= -pu + bv & \text{in } \Omega_T, & \quad u(\cdot, 0) = u^0 \text{ in } \Omega, \\ \frac{dv}{dt} &= (d - a\langle u \rangle_\Omega - cv)v & \text{in } (0, T), & \quad v(0) = \langle v^0 \rangle_\Omega \end{aligned} \tag{5.3}$$

where u is endowed with a zero flux boundary condition if $D > 0$.

Let us first study dynamics of the shadow problem. If we integrate the mild solution u of system (5.3) over Ω , we obtain an ODE system for the masses $(\langle u \rangle_\Omega, v)$. This system admits the global attractor (\bar{u}, \bar{v}) where

$$\bar{u} = \frac{dp}{cp + ab} \quad \text{and} \quad b\bar{v} = p\bar{u}.$$

Convergence to the equilibrium is a consequence of the radially unbounded Lyapunov functional

$$L(\langle u \rangle_\Omega, v) = \frac{a}{2}(\langle u \rangle_\Omega - \bar{u})^2 + b(v - \bar{v} - \bar{v} \log(v/\bar{v}))$$

adapted from [40], where L is dissipative on trajectories, i.e.,

$$\frac{dL}{dt} = -ap(\langle u \rangle_\Omega - \bar{u})^2 - bc(v - \bar{v})^2 \leq 0.$$

Thus, we obtain the asymptotic behavior $(\langle u \rangle_\Omega, v) \rightarrow (\bar{u}, \bar{v})$ as $t \rightarrow \infty$ as well as $u \rightarrow \bar{u}$, since

$$u(\cdot, t) - \langle u \rangle_\Omega(t) = (S_\Delta(Dt)u^0 - \langle u^0 \rangle_\Omega)e^{-pt} \rightarrow 0 \quad \text{for } t \rightarrow \infty.$$

Hence, Assumption 1–4 is satisfied. For application of Theorem 3.11, it remains to compute the linearization

$$\mathbf{J}(x, t) = \begin{pmatrix} -p & b \\ -av(t) & d - 2cv(t) - au(x, t) \end{pmatrix}$$

at the shadow limit (u, v) . The shadow evolution system \mathcal{W} defined in (3.13) is induced by the operators

$$\begin{aligned} \mathbf{D}^S \Delta + \mathbf{L}_0(t) &: L^p(\Omega) \times \mathbb{R} \rightarrow L^p(\Omega) \times \mathbb{R}, \\ \mathbf{L}_0(t) \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} (x) &= \begin{pmatrix} -p\xi_1(x) + b\xi_2 \\ -av(t)\langle \xi_1 \rangle_\Omega + (d - 2cv(t) - a\langle u(\cdot, t) \rangle_\Omega)\xi_2 \end{pmatrix}, \quad x \in \Omega, t \in \mathbb{R}_{\geq 0}, \end{aligned}$$

where $\mathbf{D}^S = \text{diag}(D, 0) \in \mathbb{R}_{\geq 0}^{2 \times 2}$ is a diagonal matrix.

Lemma 5.1. *Let (u, v) be a shadow solution of system (5.3) for bounded initial conditions $u^0, v^0 \geq 0$ satisfying $\langle v^0 \rangle_\Omega > 0$. Then, Assumption 6–7 is satisfied for $p = \infty$. Moreover, the corresponding evolution system \mathcal{U} and \mathcal{W} is uniformly exponentially stable for the exponent $\eta = p > 0$ and some $\sigma > 0$, respectively.*

Proof. Assumption 7 is satisfied since $(U(t))_{t \in \mathbb{R}_{\geq 0}}$ with $U(t) = S_\Delta(Dt)e^{-pt}$ is uniformly exponentially stable with exponent $\eta = p > 0$. This is a consequence of contractivity of the heat semigroup $(S_\Delta(t))_{t \in \mathbb{R}_{\geq 0}}$, see Proposition A.1. Concerning the evolution system \mathcal{W} in Assumption 6, let us split the shadow operator as $\mathbf{L}_0(t) = \mathbf{L}_\infty + \mathbf{B}(t)$ for operator matrices

$$\begin{aligned} \mathbf{L}_\infty, \mathbf{B}(t) &: L^p(\Omega) \times \mathbb{R} \rightarrow L^p(\Omega) \times \mathbb{R}, \\ \mathbf{L}_\infty \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} (x) &= \begin{pmatrix} -p\xi_1(x) + b\xi_2 \\ -a\bar{v}\langle \xi_1 \rangle_\Omega - c\bar{v}\xi_2 \end{pmatrix} =: \begin{pmatrix} A_*\xi_1(x) + B_*\xi_2 \\ C_*\langle \xi_1 \rangle_\Omega + D_*\xi_2 \end{pmatrix}, \\ \mathbf{B}(t) \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} (x) &= \begin{pmatrix} 0 \\ -a(v(t) - \bar{v})\langle \xi_1 \rangle_\Omega + [-2c(v(t) - \bar{v}) - a(\langle u(\cdot, t) \rangle_\Omega - \bar{u})]\xi_2 \end{pmatrix}. \end{aligned}$$

Since $\lim_{t \rightarrow \infty} \mathbf{B}(t) = \mathbf{0}$ with respect to the operator norm on $L^\infty(\Omega) \times \mathbb{R}$, evolution systems induced by $\mathbf{D}^S \Delta + \mathbf{L}_0(t)$ and $\mathbf{D}^S \Delta + \mathbf{L}_\infty$ are asymptotically comparable. We will show that it remains to consider the latter semigroup for exponential stability of the former evolution system. To recognize this, we start from the definition of \mathcal{W} in condition 6. This evolution system is given by evolution operators $\mathbf{W}(t, s)$ for $t, s \in \mathbb{R}_{\geq 0}, s \leq t$, defined by equation (3.13), where $\xi \in C(\mathbb{R}_{\geq 0}; L^p(\Omega)^m \times \mathbb{R}^k)$ is the unique solution of the shadow problem

$$\frac{\partial \xi}{\partial t} - (\mathbf{D}^S \Delta + \mathbf{L}_\infty)\xi = \mathbf{B}(t)\xi \quad \text{in } \Omega \times \mathbb{R}_{> 0}$$

endowed with zero flux boundary conditions if necessary. We split the full operator into a time-independent, possibly unbounded part $\mathbf{L} = \mathbf{D}^S \Delta + \mathbf{L}_\infty$ and the bounded time-varying operator family $(\mathbf{B}(t))_{t \in \mathbb{R}_{\geq 0}}$. We are able to compare both evolution systems, the system \mathcal{W}_∞ induced by a semigroup $(\mathbf{W}_\infty(t))_{t \in \mathbb{R}_{\geq 0}}$ which is generated by the operator \mathbf{L} and the full evolution system \mathcal{W} , using the integral representation

$$\mathbf{W}(t, s)\xi^0 = \mathbf{W}_\infty(t-s)\xi^0 + \int_s^t \mathbf{W}_\infty(t-\tau)\mathbf{B}(\tau)\mathbf{W}(\tau, s)\xi^0 \, d\tau \quad \forall 0 \leq s \leq t$$

from [11, Ch. VI, Theorem 9.19]. Once we have shown uniform exponential stability for $(\mathbf{W}_\infty(t))_{t \in \mathbb{R}_{\geq 0}}$, estimations in $L^p(\Omega) \times \mathbb{R}$ for $1 \leq p \leq \infty$ yield

$$\|\mathbf{W}(t, s)\xi^0\|_{L^p(\Omega) \times \mathbb{R}} \leq C e^{-\sigma_\infty(t-s)} \|\xi^0\|_{L^p(\Omega) \times \mathbb{R}} + \int_s^t C e^{-\sigma_\infty(t-\tau)} \|\mathbf{B}(\tau)\|_{L^\infty(\Omega) \times \mathbb{R}} \|\mathbf{W}(\tau, s)\xi^0\|_{L^p(\Omega) \times \mathbb{R}} \, d\tau.$$

Gronwall's inequality results in the estimate

$$\|\mathbf{W}(t, s)\xi^0\|_{L^p(\Omega) \times \mathbb{R}} \leq C e^{-\sigma_\infty(t-s)} \exp\left(\int_s^t C \|\mathbf{B}(\tau)\|_{L^\infty(\Omega) \times \mathbb{R}} \, d\tau\right) \|\xi^0\|_{L^p(\Omega) \times \mathbb{R}}.$$

Although the theory of Bohl exponents was established for bounded operators in [6, Ch. III, pp. 118], the same estimates used to prove [6, Corollary 4.2] apply to the above estimate in the context of semigroup theory, see further [47, Corollary 4.2], [7, Theorem 5]. More precisely, since $\lim_{t \rightarrow \infty} \mathbf{B}(t) = \mathbf{0}$, for each $\gamma \in (0, 1)$ there is a $t_0 > 0$ such that $C \|\mathbf{B}(t)\|_{L^\infty(\Omega) \times \mathbb{R}} \leq \gamma \sigma_\infty$ for all $t \geq t_0$. This implies the estimate

$$\|\mathbf{W}(t, s)\xi^0\|_{L^p(\Omega) \times \mathbb{R}} \leq \tilde{C} e^{-\gamma \sigma_\infty(t-s)} \|\xi^0\|_{L^p(\Omega) \times \mathbb{R}}$$

for $\tilde{C} = C \exp(\int_0^{t_0} C \|\mathbf{B}(\tau)\|_{L^\infty(\Omega) \times \mathbb{R}} \, d\tau)$. Hence, uniform exponential stability carries over from the evolution system \mathcal{W}_∞ to the full evolution system \mathcal{W} on $L^p(\Omega) \times \mathbb{R}$, provided $\lim_{t \rightarrow \infty} \mathbf{B}(t) = \mathbf{0}$.

Using the spectral mapping theorem [11, Ch. IV, Corollary 3.12] for analytical semigroups, it is well-known that exponential stability of the semigroup $(\mathbf{W}_\infty(t))_{t \in \mathbb{R}_{\geq 0}}$ can be verified via the spectrum of its generator $\mathbf{L} = \mathbf{D}^S \Delta + \mathbf{L}_\infty$. It remains to show $s(\mathbf{D}^S \Delta + \mathbf{L}_\infty) < 0$ for uniform exponential stability of the evolution system \mathcal{W}_∞ resp. \mathcal{W} [11, Ch. V, Theorem 1.10]. We infer from Proposition B.3 that in case of $D = 0$

$$\sigma(\mathbf{L}) = \sigma(\mathbf{L}_\infty) = \{A_*\} \cup \Sigma,$$

where Σ consists of all eigenvalues of the constant coefficient matrix

$$\mathbf{J}_\infty = \begin{pmatrix} A_* & B_* \\ C_* & D_* \end{pmatrix}.$$

Note that $A_* = -p < 0$ and both eigenvalues of \mathbf{J}_∞ have negative real parts since $\text{tr}(\mathbf{J}_\infty) = -p - c\bar{v} < 0$ and $\det(\mathbf{J}_\infty) = (pc + ab)\bar{v} > 0$.

From the reasoning in Lemma B.1 we know that $\sigma(\mathbf{D}^S \Delta + \mathbf{L}_\infty)$ is a discrete set for $D > 0$. The semigroup $(\mathbf{W}_\infty(t))_{t \in \mathbb{R}_{\geq 0}}$ is defined by the solution $\xi = (w, z)$ of $\partial_t \xi - \mathbf{D}^S \Delta \xi = \mathbf{L}_\infty \xi$. While $w(\cdot, t) - \langle w \rangle_\Omega(t) = (S_\Delta(Dt)w^0 - \langle w^0 \rangle_\Omega) e^{-pt}$, integration yields

$$\langle \xi \rangle_\Omega = \begin{pmatrix} \langle w \rangle_\Omega \\ z \end{pmatrix} (t) = \exp(\mathbf{J}_\infty t) \begin{pmatrix} \langle w^0 \rangle_\Omega \\ \langle z^0 \rangle_\Omega \end{pmatrix}.$$

It is well-known from the theory of ODEs that $\langle \xi \rangle_\Omega$ decays exponentially to zero since \mathbf{J}_∞ is a stable matrix [6, Ch. I, Theorem 4.1]. Choosing $\sigma \in \mathbb{R}_{>0}$ such that $\sigma < \min\{p, \min_{\lambda \in \sigma(\mathbf{J}_\infty)} |\text{Re } \lambda|\}$ yields an estimation of both expressions. This results in

$$\|\mathbf{W}_\infty(t)\xi^0\|_{L^\infty(\Omega) \times \mathbb{R}} = \|\xi(\cdot, t)\|_{L^\infty(\Omega) \times \mathbb{R}} \leq C_\sigma e^{-\sigma t} \|\xi^0\|_{L^\infty(\Omega) \times \mathbb{R}}$$

for some $C_\sigma > 0$. Thus, Assumption 8 is satisfied for $p = \infty$ and each $D \geq 0$. \square

In summary, Theorem 3.11 yields global estimates

$$\|u_\varepsilon - u\|_{L^\infty(\Omega \times \mathbb{R}_{\geq 0})} + \|v_\varepsilon - v - \psi_\varepsilon\|_{L^\infty(\Omega \times \mathbb{R}_{\geq 0})} \leq C\varepsilon.$$

Note that the results on Lyapunov functions in [20, Proposition 2.1] is also applicable to partly diffusing systems. The same Lyapunov function which is known from the theory of ODEs can be extended to the reaction-diffusion case. Consequently, (\bar{u}, \bar{v}) is the only positive attractor for the diffusive system (5.2) and (\bar{u}, \bar{v}) is globally (for positive initial data) asymptotically stable by Lyapunov's direct method.

Acknowledgments

This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy EXC 2181/1 - 390900948 (the Heidelberg STRUCTURES Excellence Cluster and SFB1324 (B05) and the Klaus Tschira Foundation, Germany (00.277.2015).

A. Parabolic theory

The preceding results are based on properties of the heat semigroup $(S_\Delta(t))_{t \in \mathbb{R}_{\geq 0}}$ defined in [8, 43]. We study basic properties of this semigroup on the spaces $L^p(\Omega)$ for $1 \leq p < \infty$. Additionally, $L^\infty(\Omega_T)$ estimates are derived for solutions of the inhomogeneous heat equation with explicit dependence on time T .

Proposition A.1. *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with $\partial\Omega \in C^{0,1}$, $z^0 \in L^2(\Omega)$. Then, the homogeneous heat equation*

$$\frac{\partial z}{\partial t} - \Delta z = 0 \quad \text{in } \Omega_T, \quad z(\cdot, 0) = z^0 \quad \text{in } \Omega, \quad \frac{\partial z}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times (0, T)$$

has a unique variational solution $z \in C(\mathbb{R}_{\geq 0}; L^2(\Omega))$ satisfying $z \in L^2(0, T; H^1(\Omega))$ for each $T > 0$. The solution is given by the Fourier expansion

$$z(x, t) = (S_\Delta(t)z^0)(x) = \sum_{j \in \mathbb{N}_0} e^{-\lambda_j t} (z^0, w_j)_{L^2(\Omega)} w_j(x), \quad x \in \Omega, t \in \mathbb{R}_{\geq 0}, \quad (\text{A.1})$$

where $(\lambda_j, w_j)_{j \in \mathbb{N}_0}$ is a spectral basis of $-\Delta$ on $L^2(\Omega)$ solving problem (A.2).

Moreover, the heat semigroup $(S_\Delta(t))_{t \in \mathbb{R}_{\geq 0}}$ defined by (A.1) can be extended to a contraction semigroup on $L^p(\Omega)$ for each $1 \leq p \leq \infty$, which is strongly continuous for $1 \leq p < \infty$ and analytic for $1 < p < \infty$.

Proof. Existence and contractivity of the heat semigroup $(S_\Delta(t))_{t \in \mathbb{R}_{\geq 0}}$ is shown in [8, Theorem 1.3.9]. By [8, Theorems 1.4.1, 1.4.2], $(S_\Delta(t))_{t \in \mathbb{R}_{\geq 0}}$ is a strongly continuous semigroup for each $1 \leq p < \infty$, which even can be extended analytically to some sector in the complex plane for $p > 1$. The solution $z(t) = S_\Delta(t)z^0$ for $z^0 \in L^2(\Omega)$ satisfies $z \in C(\mathbb{R}_{\geq 0}; L^2(\Omega))$ due to strong continuity of the semigroup. Although the solution might lose its differentiability at $t = 0$, we obtain higher regularity for $t > 0$ by analyticity of the semigroup. This yields $z \in C(\mathbb{R}_{> 0}; \mathcal{D}(\Delta^\ell))$ for all integers $k, \ell \in \mathbb{N}_0$, where $\mathcal{D}(\Delta^0) := L^2(\Omega)$ and $\mathcal{D}(\Delta)$ is the domain of the generator of the heat semigroup defined on $L^2(\Omega)$ [4, Theorem 7.7]. Hence, the solution $z(\cdot, t)$ lies in $H^1(\Omega)$ for each $t > 0$ and the boundary condition is satisfied in the sense of distributions by the trace operator $H^1(\Omega) \hookrightarrow W^{-1/2, 2}(\partial\Omega)$ from [13, Theorem 1.5.1.2].

To determine the Fourier coefficients, we recall that the unique solution z solves the variational equation

$$\frac{d}{dt} \int_\Omega z(x, t) \varphi(x) dx + \int_\Omega \nabla z(x, t) \nabla \varphi(x) dx = 0 \quad \forall \varphi \in H^1(\Omega), t \in \mathbb{R}_{> 0}.$$

Since $z(\cdot, t) \in L^2(\Omega)$, we can expand this function in a Fourier series using a spectral basis of $-\Delta$ from equation (A.2). This leads to the series representation (A.1).

Concerning the regularity $z \in L^2(0, T; H^1(\Omega))$, we note that the partial sums

$$z_m(\cdot, t) = \sum_{j=0}^m e^{-\lambda_j t} (z^0, w_j)_{L^2(\Omega)} w_j$$

of the Fourier series form a Cauchy sequence in this space. Indeed, for $\ell > m \geq 0$

$$\|z_m - z_\ell\|_{L^2(0,T;H^1(\Omega))}^2 = \sum_{j=m+1}^{\ell} \int_0^T (1 + \lambda_j) e^{-2\lambda_j t} dt |(z^0, w_j)_{L^2(\Omega)}|^2 \leq C \sum_{j=m+1}^{\ell} |(z^0, w_j)_{L^2(\Omega)}|^2$$

tends to 0 as $m, \ell \rightarrow \infty$, due to Parseval's equality for $z^0 \in L^2(\Omega)$ and $C = 1/2 + 1/(2\lambda_1)$. \square

As used above, Galerkin's approximation is based on a spectral basis $(\lambda_j, w_j)_{j \in \mathbb{N}_0}$ of $-\Delta$ satisfying

$$-\Delta w_j = \lambda_j w_j \quad \text{in } \Omega, \quad \frac{\partial w_j}{\partial \nu} = 0 \quad \text{on } \partial\Omega \quad (\text{A.2})$$

in the weak sense [21, Theorem 1.2.8]. The principal eigenvalue $\lambda_0 = 0$ has eigenfunction $w_0 = |\Omega|^{-1/2}$. The other eigenvalues are strictly positive and tend to infinity as $j \rightarrow \infty$. The eigenfunctions $(w_j)_{j \in \mathbb{N}_0} \subset H^1(\Omega)$ form an orthonormal basis for $L^2(\Omega)$ and an orthogonal basis for $H^1(\Omega)$.

The first positive eigenvalue $\lambda_1 > 0$ of $-\Delta$ is also crucial for the decay estimate of the heat semigroup which is used in this work. For $\langle z^0 \rangle_\Omega = 0$, [50, Lemma 1.3] implies existence of a constant $\bar{C} > 0$ independent of τ and the initial conditions for which we have the following decay in time:

$$\|z(\cdot, t)\|_{L^p(\Omega)} \leq \bar{C} e^{-\lambda_1 t} \|z^0\|_{L^p(\Omega)} \quad \forall t \in \mathbb{R}_{\geq 0} \quad (\text{A.3})$$

Unfortunately, [50] uses estimates of the heat kernel for a bounded domain with boundary $\partial\Omega \in C^{2,\alpha}$ for some $\alpha \in (0, 1)$. In order to relax this condition, we use heat kernel estimates for $\partial\Omega \in C^{0,1}$ from [8, Theorem 3.2.9], which imply decay estimate (A.3) by the same steps as in the proof of [50].

By Duhamel's formula, we obtain the following result for the inhomogeneous heat equation.

Proposition A.2. *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with $\partial\Omega \in C^{0,1}$, $z^0 \in L^2(\Omega)$ and $R \in L^2(\Omega_T)$ be given. Then the inhomogeneous heat equation*

$$\frac{\partial z}{\partial t} - \Delta z = R(x, t) \quad \text{in } \Omega_T, \quad z(\cdot, 0) = z^0 \quad \text{in } \Omega, \quad \frac{\partial z}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times (0, T) \quad (\text{A.4})$$

has a unique mild solution $z \in L^2(0, T; H^1(\Omega)) \cap C([0, T]; L^2(\Omega))$, given by the separation of variables formula

$$\begin{aligned} z(\cdot, t) &= S_\Delta(t) z^0 + \int_0^t S_\Delta(t-s) R(\cdot, s) ds \\ &= \sum_{j \in \mathbb{N}_0} e^{-\lambda_j t} (z^0, w_j)_{L^2(\Omega)} w_j(x) + \int_0^t \sum_{j \in \mathbb{N}_0} e^{-\lambda_j(t-s)} (R(\cdot, s), w_j)_{L^2(\Omega)} w_j(x) ds. \end{aligned} \quad (\text{A.5})$$

In addition, $z^0 \in H^1(\Omega)$ implies a weak solution $z \in L^\infty(0, T; H^1(\Omega))$ with weak derivative $\partial_t z \in L^2(\Omega_T)$, and the weak formulation

$$(\partial_t z(\cdot, t), \varphi)_{L^2(\Omega)} + (\nabla z(\cdot, t), \nabla \varphi)_{L^2(\Omega)} = (R(\cdot, t), \varphi)_{L^2(\Omega)} \quad \forall \varphi \in H^1(\Omega)$$

holds for a.e. $t \in (0, T)$.

Proof. By [45, Section 4.2], there exists a unique mild solution $z \in C([0, T]; L^2(\Omega))$ of problem (A.4), which is given by the integral formula (A.5). To show that z is an element of $L^2(0, T; H^1(\Omega))$ resp. $L^\infty(0, T; H^1(\Omega))$, it is again sufficient to prove the Cauchy property of the partial sums given by expression (A.5). For $\ell > m \geq 0$, we obtain

$$\begin{aligned} \|(z_m - z_\ell)(\cdot, t)\|_{H^1(\Omega)}^2 &\leq \sum_{j=m+1}^{\ell} (1 + \lambda_j) \left[e^{-2\lambda_j t} |(z^0, w_j)_{L^2(\Omega)}|^2 + \left(\int_0^t e^{-\lambda_j(t-s)} (R(\cdot, s), w_j)_{L^2(\Omega)} ds \right)^2 \right] \\ &\leq \sum_{j=m+1}^{\ell} (1 + \lambda_j) |(z^0, w_j)_{L^2(\Omega)}|^2 + (T+1) \int_0^t \sum_{j=m+1}^{\ell} |(R(\cdot, s), w_j)_{L^2(\Omega)}|^2 ds \end{aligned}$$

by Hölder's inequality. The same reasoning as in the proof of Proposition A.1 applies to obtain a Cauchy sequence, hence, we omit details. Concerning the weak formulation, we use Galerkin's approximation [12, Section 7.1]. Following this classical approach from [12, §7.1.3, Theorem 5], one establishes the result for more regular initial data $z^0 \in H^1(\Omega)$. \square

Once we obtained a solution of problem (A.4) for $z^0 = 0$ in Proposition 3.5, we used an $L^\infty(\Omega_T)$ estimate for the solution of the inhomogeneous heat equation with explicit dependence on time T . This can be developed as in [34, Ch. III, §7], which essentially uses parabolic $L_{p,r}(\Omega_T)$ estimates in combination with the well-known truncation method of Stampacchia. However, we will use semigroup theory to prove such a result. Let us consider a bounded weak solution $z \in L^2(0, T; H^1(\Omega)) \cap C([0, T]; L^2(\Omega))$ which solves

$$\frac{\partial z}{\partial t} - d\Delta z = R(x, t) \quad \text{in } \Omega_T, \quad z(\cdot, 0) = 0 \quad \text{in } \Omega, \quad \frac{\partial z}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times (0, T) \quad (\text{A.6})$$

with right-hand side $R \in L_{p,r}(\Omega_T)$ and diffusion parameter d . The Lebesgue space $L_{p,r}(\Omega_T)$ is defined by all measurable functions ψ on Ω_T with finite mixed norm

$$\|\psi\|_{p,r} := \left(\int_0^T \left(\int_\Omega |\psi(x, t)|^p dx \right)^{r/p} dt \right)^{1/r} \quad \text{for } 1 \leq p, r < \infty$$

and an obvious modification for $r = \infty$ [34, Chapters I, II, §1 in both cases]. If we do not specify the region of integration within the notation $\|\cdot\|_{p,r}$, we assume to integrate over Ω_T .

The aim is to show $L^\infty(\Omega_T)$ estimates for the solution z which depend explicitly on time T and the mixed norm $\|R\|_{p,r}$. Within this procedure, the exponent p is restricted due to Sobolev embeddings by $p > n/2$ for $n \geq 2$. Let us choose a parameter $1 \leq r \leq \infty$ according to [34, Ch. III, §7], i.e., we assume the relation

$$0 \leq \frac{1}{r} + \frac{n}{2p} < 1 \quad (\text{A.7})$$

for given $1 \leq p, r \leq \infty$. Then we obtain the following result.

Proposition A.3. *Let $z \in L^2(0, T; H^1(\Omega)) \cap C([0, T]; L^2(\Omega))$ be a bounded weak solution of the initial boundary value problem (A.6) with right-hand side $R \in L_{p,r}(\Omega_T)$ and parameter values p, r satisfying relation (A.7). Then there exists a constant $C > 0$ which only depends on Ω, n, p, r and a lower bound of the diffusion d such that the diffusive component z satisfies the estimate*

$$\|z\|_{L^\infty(\Omega_T)} \leq CT^{1-1/r} \|R\|_{p,r}. \quad (\text{A.8})$$

Proof. The solution z is given by the integral representation (A.5), i.e.,

$$z(\cdot, t) = \int_0^t S_\Delta(d(t-s))R(\cdot, s) ds = \int_0^t S_\Delta(d(t-s)) (R(\cdot, s) - \langle R(\cdot, s) \rangle_\Omega) ds + \int_0^t \langle R(\cdot, s) \rangle_\Omega ds.$$

The first integral can be estimated with a certain estimate of the heat semigroup from [50, Lemma 1.3] similar to inequality (A.3). In fact, we obtain

$$\begin{aligned} \|S_\Delta(d(t-s)) (R(\cdot, s) - \langle R(\cdot, s) \rangle_\Omega)\|_{L^\infty(\Omega)} &\leq C \left(1 + (d(t-s))^{-\frac{n}{2p}} \right) \|R(\cdot, s) - \langle R(\cdot, s) \rangle_\Omega\|_{L^p(\Omega)} \\ &\leq 2C \left(1 + (d(t-s))^{-\frac{n}{2p}} \right) \|R(\cdot, s)\|_{L^p(\Omega)} \end{aligned}$$

for a constant $C > 0$ which depends on Ω only. Condition (A.7) yields that the function $\tau \mapsto 1 + (d\tau)^{-\frac{n}{2p}}$ is in $L^{\hat{r}}((0, T))$ where $\frac{1}{\hat{r}} + \frac{1}{r} = 1$. Since the second integral satisfies

$$\left| \int_0^T \langle R(\cdot, s) \rangle_\Omega ds \right| \leq |\Omega|^{-1/p} T^{1/\hat{r}} \|R\|_{p,r},$$

Hölder's inequality implies the desired inequality (A.8). \square

B. Spectral theory

To prove convergence results, we applied a stabilizing effect to ensure that diffusive solutions of system (1.1) stay nearby its shadow limit. Such a result is achieved via linearization at the shadow solution. In case of a bounded stationary shadow solution, we will give a complete characterization of the spectrum of the corresponding linearized shadow operator. This allows not only to verify stability Assumption 6 but also allows general stability results for bounded stationary solutions of the nonlinear shadow system (1.2). Note that the characterization in Proposition B.3 below is not limited to eigenvalues of the linearized operator \mathbf{L} as considered in [37, Appendix B]. Furthermore, the following results are valid for systems with a vector-valued shadow component $v \in \mathbb{R}^k$, see [31, Section 5.2.1].

B.1. Shadow operator

Let us consider a bounded steady state $(\bar{\mathbf{u}}, \bar{v}) \in L^\infty(\Omega)^m \times \mathbb{R}$ of shadow system (1.2), i.e., a bounded solution of the problem

$$\mathbf{0} = \mathbf{f}(\mathbf{u}, v, x) \quad \text{in } \Omega, \quad 0 = \langle g(\mathbf{u}, v, \cdot) \rangle_\Omega.$$

Stability or instability of stationary solutions is often verified using linearization at a steady state of the nonlinear problem [6, Ch. VII, Theorems 2.1, 2.3]. We consider a linearization in $L^p(\Omega)^m \times \mathbb{R}$ which we need to verify Assumption 6 in the time-independent case $(\mathbf{u}, v) = (\bar{\mathbf{u}}, \bar{v})$.

Let \mathbf{f}, g be once continuously differentiable with respect to the unknown variables \mathbf{u}, v according to Assumption 4. Using the Fréchet-derivative, the linearization is given by the operator $\mathbf{L} \in \mathcal{L}(L^p(\Omega)^m \times \mathbb{R})$,

$$(\mathbf{L}\xi)(x) = \begin{pmatrix} \mathbf{A}(x)\xi_1(x) + \mathbf{B}(x)\xi_2 \\ \langle \mathbf{C}(\cdot)\xi_1 \rangle_\Omega + \langle D(\cdot)\xi_2 \rangle_\Omega \end{pmatrix} \quad \forall \xi = (\xi_1, \xi_2) \in L^p(\Omega)^m \times \mathbb{R}, x \in \Omega. \quad (\text{B.1})$$

Here, we used the same notation as in the Jacobian (3.9) for the uniformly bounded entries $\mathbf{A}, \mathbf{B}, \mathbf{C}, D$ being the parts of the Jacobian of (\mathbf{f}, g) evaluated at the shadow steady state $(\bar{\mathbf{u}}, \bar{v})$. Note that \mathbf{L} is the generator of a uniformly continuous semigroup $(e^{\mathbf{L}t})_{t \in \mathbb{R}_{\geq 0}}$ by [11, Ch. II, Theorem 1.4]. Norm-continuity of this semigroup implies validity of the spectral mapping theorem for $1 \leq p \leq \infty$ [11, Ch. IV, Corollary 3.12]. As a consequence, the growth bound of the semigroup equals the spectral bound $s(\mathbf{L}) := \sup\{\text{Re } \lambda \mid \lambda \in \sigma(\mathbf{L})\}$ of the generator \mathbf{L} . By [11, Ch. V, Theorem 1.10], uniform exponential stability can be deduced from a negative spectral bound. As in the finite-dimensional case, one has to be careful if the spectral bound of \mathbf{L} is zero [9, Ch. III, Theorem 1.11]. Concerning the stability condition we are faced with in Assumption 6, it is clear that in the stationary case (B.1) the evolution system \mathcal{W} consists of the evolution operators

$$\mathbf{W}(t, s) = e^{\mathbf{L}(t-s)} \quad \forall 0 \leq s \leq t \in \mathbb{R}_{\geq 0}.$$

Hence, to determine their growth, it is important to locate the spectrum $\sigma(\mathbf{L})$ in the complex plane. To study invertibility of the operator $\lambda I - \mathbf{L}$ for some $\lambda \in \mathbb{C}$, we focus on the following system of equations

$$(\lambda I - \mathbf{L})\xi = \psi \quad \Leftrightarrow \quad \begin{cases} (\lambda I - \mathbf{A})\xi_1 - \mathbf{B}\xi_2 = \psi_1, \\ -\langle \mathbf{C}\xi_1 \rangle_\Omega + (\lambda I - \langle D \rangle_\Omega)\xi_2 = \psi_2. \end{cases}$$

Let \mathbf{A} also denote the bounded multiplication operator induced by the matrix $\mathbf{A}(x)$ on $L^p(\Omega)^m$, see Appendix B.2. If $\lambda \notin \sigma(\mathbf{A})$, the first equation can be solved and we obtain the following result.

Lemma B.1. *Let \mathbf{L} be the shadow operator defined in (B.1) on $L^p(\Omega)^m \times \mathbb{R}$ for bounded coefficient matrices $\mathbf{A}, \mathbf{B}, \mathbf{C}$ and D and an exponent $1 \leq p \leq \infty$. Then*

$$\Sigma := \sigma(\mathbf{L}) \cap \rho(\mathbf{A}) \subset \sigma_p(\mathbf{L})$$

is a discrete (probably empty) set of eigenvalues of \mathbf{L} . Moreover, $\sigma(\mathbf{L}) \subset \sigma(\mathbf{A}) \dot{\cup} \Sigma$.

Proof. Provided $\lambda \in \rho(\mathbf{A})$, we already have

$$(\lambda I - \mathbf{L})\xi = \psi \quad \Leftrightarrow \quad \xi_1 = (\lambda I - \mathbf{A})^{-1} [\psi_1 + \mathbf{B}\xi_2], \quad H(\lambda)\xi_2 = \tilde{\psi}(\lambda)$$

where

$$H(\lambda) = \lambda - \langle D \rangle_\Omega - \langle \mathbf{C}(\lambda I - \mathbf{A})^{-1} \mathbf{B} \rangle_\Omega, \quad \tilde{\psi}(\lambda) = \psi_2 + \langle \mathbf{C}(\lambda I - \mathbf{A})^{-1} \psi_1 \rangle_\Omega.$$

We apply an analytic Fredholm theorem [16, Theorem 4.34] to show discreteness of the remaining spectrum Σ in $\rho(\mathbf{A})$. Let us first note that the subsystem for ξ_2 can be solved for sufficiently large $\lambda > 0$. To see this, recall that \mathbf{A} is a bounded multiplication operator and hence, for $\lambda \in \rho(\mathbf{A})$ with $|\lambda| > 2\|\mathbf{A}\|$, we obtain an estimate for $(\lambda I - \mathbf{A})^{-1}$ by Neumann's series which is of the form $2/|\lambda|$. This implies that $H(\lambda) > 0$ for all sufficiently large $\lambda > 0$. Moreover, analyticity of the resolvent map $\lambda \mapsto (\lambda I - \mathbf{A})^{-1}$ implies analyticity of the complex-valued function H on $\rho(\mathbf{A})$. It is well-known that the set of zeros of the analytic function H is a discrete set $\Sigma \subset \rho(\mathbf{A})$. Consequently, $H(\lambda)$ is invertible for all $\lambda \in \rho(\mathbf{A}) \setminus \Sigma$. For values $\lambda \in \Sigma$ we infer an eigenfunction $(\xi_1, \xi_2) \in L^\infty(\Omega)^m \times \mathbb{R}$ of the eigenvalue equation $(\lambda I - \mathbf{L})\xi = \mathbf{0}$ due to $(\lambda I - \mathbf{A})^{-1} \mathbf{B} \in L^\infty(\Omega)^m$ by [18, Proposition 2.2]. This shows $\Sigma \subset \sigma_p(\mathbf{L})$. \square

Remark B.2. Note that the discrete set Σ is not necessarily closed. Indeed, a sequence of eigenvalues μ_j in $\Sigma \subset \sigma_p(\mathbf{L})$ has corresponding eigenfunctions such that the singular sequence of normalized eigenfunctions implies $\lim_{j \rightarrow \infty} \mu_j \in \sigma(\mathbf{L})$ which is a subset of $\sigma(\mathbf{A}) \cup \Sigma$. Hence, $\lim_{j \rightarrow \infty} \mu_j \in \sigma(\mathbf{A})$ and all accumulation points of Σ are included in $\sigma(\mathbf{A})$.

The multiplication operator induced by the ODE subsystem \mathbf{A} on $L^p(\Omega)^m$ causes problems while inverting the operator $\lambda I - \mathbf{L}$. As shown in [37, Theorem B.1], there holds $\sigma_p(\mathbf{A}) \subset \sigma_p(\mathbf{L})$. A characterization of the spectrum of the multiplication operator \mathbf{A} is given in Proposition B.4. The fact that the spectrum $\sigma(\mathbf{A})$ is essential enables us to verify that $\sigma(\mathbf{A})$ is a part of the spectrum of the shadow operator \mathbf{L} .

Proposition B.3. *Let $\mathbf{A}, \mathbf{B}, \mathbf{C}, D$ be matrix-valued functions with entries in $L^\infty(\Omega)$ according to the shadow operator \mathbf{L} defined by (B.1) on $L^p(\Omega)^m \times \mathbb{R}$ for some $1 \leq p \leq \infty$. Then there holds*

$$\sigma(\mathbf{L}) = \sigma(\mathbf{A}) \dot{\cup} \Sigma,$$

where $\Sigma \subset \sigma_p(\mathbf{L})$ is the discrete (possibly empty) set defined in Lemma B.1.

Proof. It remains to show $\sigma(\mathbf{A}) \subset \sigma(\mathbf{L})$, since from considerations in Lemma B.1 we already obtained

$$\rho(\mathbf{A}) \cap \Sigma = \Sigma \subset \sigma(\mathbf{L}) \subset \sigma(\mathbf{A}) \dot{\cup} \Sigma.$$

We apply [2, Theorem 2.2] to show equality of the essential Wolf spectrum of \mathbf{L} and \mathbf{A} , i.e., $\sigma_{\text{ess}}(\mathbf{A}) = \sigma_{\text{ess}}(\mathbf{L})$. The result of Proposition B.4 shows

$$\sigma(\mathbf{A}) = \sigma_{\text{ess}}(\mathbf{A}) := \{\lambda \in \mathbb{C} \mid \lambda I - \mathbf{A} \text{ is not a Fredholm operator}\}.$$

Hence, it remains to show $\sigma_{\text{ess}}(\mathbf{A}) = \sigma_{\text{ess}}(\mathbf{L})$. In order to apply [2, Theorem 2.2], we permute the operator matrix \mathbf{L} in (B.1). Let us consider the invertible permutation matrix

$$\mathbf{P} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{with} \quad \mathbf{P}^2 = I.$$

Then $\lambda I - \mathbf{L}$ is Fredholm if and only if $\lambda I - \tilde{\mathbf{L}}$ is Fredholm where $\tilde{\mathbf{L}} = \mathbf{P}^{-1} \mathbf{L} \mathbf{P} \in \mathcal{L}(\mathbb{R} \times L^p(\Omega)^m)$, hence $\sigma_{\text{ess}}(\mathbf{L}) = \sigma_{\text{ess}}(\tilde{\mathbf{L}})$. This is a consequence of the fact that $\mathbf{P} \in \mathcal{L}(\mathbb{R} \times L^p(\Omega)^m; L^p(\Omega)^m \times \mathbb{R})$ is Fredholm by invertibility and $\lambda I - \tilde{\mathbf{L}} = \mathbf{P}^{-1}(\lambda I - \mathbf{L})\mathbf{P}$ is a composition of Fredholm operators [4, Ch. 6]. We apply the results of [2] to the operator $\tilde{\mathbf{L}} \in \mathcal{L}(\mathbb{R} \times L^p(\Omega)^m)$ given by

$$\tilde{\mathbf{L}} := \begin{pmatrix} \tilde{A} & \tilde{B} \\ \tilde{C} & \tilde{D} \end{pmatrix}$$

where we take the bounded multiplication operators $\tilde{D} := \mathbf{A}$ on $X_2 := L^p(\Omega)^m$ and $\tilde{A} := \langle D \rangle_\Omega$ on $X_1 := \mathbb{R}$, where the latter operator has a compact resolvent on \mathbb{R} . The operators $\tilde{B} := \langle \mathbf{C} \cdot \rangle_\Omega : X_2 \rightarrow X_1$ and $\tilde{C} := \mathbf{B} : X_1 \rightarrow X_2$ consist of bounded integral and multiplication operators. Note that the operators $S(\mu) = S_0 + M(\mu)$ for $\mu \in \rho(\tilde{A})$ in assumption (e) of their paper is given by $S_0 = \tilde{D} = \mathbf{A}$, and the operator $M(\mu) = -\tilde{C}(\mu I - \tilde{A})^{-1}\tilde{B}$ which is compact for each $\mu \in \rho(\tilde{A})$ by [4, Proposition 6.3]. Since \mathbf{L} is bounded, the equality $\sigma_{\text{ess}}(\tilde{\mathbf{L}}) = \sigma_{\text{ess}}(S_0)$ is a consequence of [2, Theorem 2.2]. This shows $\sigma_{\text{ess}}(\mathbf{L}) = \sigma_{\text{ess}}(\mathbf{A})$. \square

B.2. Multiplication operator

Each matrix $\mathbf{A} \in L^\infty(\Omega)^{m \times m}$ induces a corresponding multiplication operator

$$\mathbf{M}_{\mathbf{A}} : L^p(\Omega)^m \rightarrow L^p(\Omega)^m, \quad \mathbf{z} \mapsto \mathbf{A}\mathbf{z}$$

where $(\mathbf{A}\mathbf{z})(x) := \mathbf{A}(x)\mathbf{z}(x)$ for each $\mathbf{z} \in L^p(\Omega)^m, x \in \Omega$. Since $\|\mathbf{M}_{\mathbf{A}}\| \leq \|\mathbf{A}\|_\infty$, this is a bounded, linear operator for each $1 \leq p \leq \infty$. We simply write \mathbf{A} instead of $\mathbf{M}_{\mathbf{A}}$. As Proposition B.3 shows, knowledge of the spectrum of the multiplication operator \mathbf{A} allows us to characterize the spectrum of the shadow operator \mathbf{L} defined in (B.1). We refer to [10, Ch. IX] and [18, Sections 1-3] for several characterizations of the spectrum $\sigma(\mathbf{A})$ of the multiplication operator \mathbf{A} on $L^p(\Omega)^m$ for $1 \leq p < \infty$. The following result concerning the essential spectrum is known for the case $p = 2$ by [18, Proposition 3.2, Corollary 3.4] and for the scalar case by [48, Proposition 3]. A generalization to arbitrary exponents $1 \leq p \leq \infty$ is given next.

Proposition B.4. *Let $\mathbf{A} \in L^\infty(\Omega)^{m \times m}, m \in \mathbb{N}$, and let \mathbf{A} denote its corresponding multiplication operator on $L^p(\Omega)^m$ for some $1 \leq p \leq \infty$. Then there exists a null set $N \subset \Omega$ such that*

$$\sigma(\mathbf{A}) = \overline{\bigcup_{x \in \Omega \setminus N} \sigma(\mathbf{A}(x))}. \quad (\text{B.2})$$

Moreover, the whole spectrum is essential in the sense of Wolf, i.e.,

$$\sigma(\mathbf{A}) = \sigma_{\text{ess}}(\mathbf{A}) = \{\lambda \in \mathbb{C} \mid \lambda I - \mathbf{A} \text{ is not a Fredholm operator}\}.$$

Proof. Boundedness of the multiplication operator leads to a non-empty resolvent set $\rho(\mathbf{A}) \neq \emptyset$. For $1 \leq p < \infty$, [10, Ch. IX, Theorem 2.4] states

$$\sigma(\mathbf{A}) = \{\lambda \in \mathbb{C} \mid |N_{\lambda, \varepsilon}| > 0 \quad \forall \varepsilon > 0\} =: \text{ess-}\sigma(\mathbf{A}(\Omega)),$$

for measurable sets

$$N_{\lambda, \varepsilon} := \{x \in \Omega \mid \text{dist}(\lambda, \sigma(\mathbf{A}(x))) < \varepsilon\}.$$

On the one hand, along the same lines of that proof, $\sigma(\mathbf{A}) \subset \text{ess-}\sigma(\mathbf{A}(\Omega))$ also holds for $p = \infty$. On the other hand, the proof of $\sigma(\mathbf{A}) \supset \text{ess-}\sigma(\mathbf{A}(\Omega))$ given in [10, Ch. IX, Theorem 2.4] does not apply for $p = \infty$. In order to prove the above representation (B.2) of the spectrum, it remains to show the inclusion $\text{ess-}\sigma(\mathbf{A}(\Omega)) \subset \sigma(\mathbf{A})$ and [10, Ch. IX, Remark 2.3] yields the result.

Using the idea of [18, Theorem 3.3], we show that each $\lambda \in \text{ess-}\sigma(\mathbf{A}(\Omega))$ is in the spectrum of \mathbf{A} . As the characteristic polynomial of the matrix $\mathbf{A}(x)$ factorizes with eigenvalues $\lambda_i(x) \in \mathbb{C}$, we obtain

$$|\det(\lambda I - \mathbf{A}(x))| = \prod_{i=1}^m |\lambda - \lambda_i(x)| \geq \text{dist}(\lambda, \sigma(\mathbf{A}(x)))^m \quad (\text{B.3})$$

for a.e. $x \in \Omega$. This estimate yields the inclusion

$$\Gamma_{\lambda, \varepsilon} := \{x \in \Omega \mid |\det(\lambda I - \mathbf{A}(x))| < \varepsilon^m\} \subset N_{\lambda, \varepsilon}$$

and we conclude that

$$\Gamma_\lambda := \{x \in \Omega \mid \det(\lambda I - \mathbf{A}(x)) = 0\} \quad (\text{B.4})$$

satisfies $0 \leq |\Gamma_\lambda| \leq \lim_{\varepsilon \rightarrow 0} |N_{\lambda, \varepsilon}|$ as the limit of the above subsets of $N_{\lambda, \varepsilon}$ as $\varepsilon \rightarrow 0$. The sequence $(|\Gamma_{\lambda, \varepsilon}|)_{\varepsilon > 0}$ of non-negative numbers is non-increasing as $\varepsilon \rightarrow 0$ with a limit which is either positive or zero. In the former case, we conclude that Γ_λ defined in (B.4) has positive measure which is equivalent to $\lambda \in \sigma_p(\mathbf{A})$ using [24, Theorem 2.1] or [18, Theorem 2.5]. In the latter case, $|\Gamma_\lambda| = \lim_{\varepsilon \rightarrow 0} |\Gamma_{\lambda, \varepsilon}| = 0$, we show that the injective operator $\lambda I - \mathbf{A}$ is not bounded from below, hence $\lambda \in \sigma(\mathbf{A})$. Although we know from $\lambda \in \text{ess-}\sigma(\mathbf{A}(\Omega))$ that $|N_{\lambda, \varepsilon}| > 0$ for all $\varepsilon > 0$, there are still two possibilities for the zero sequence $(|\Gamma_{\lambda, \varepsilon}|)_{\varepsilon > 0}$: either $|\Gamma_{\lambda, \varepsilon}| > 0$ for all $\varepsilon > 0$ or the sequence becomes stationary in the sense that $|\Gamma_{\lambda, \varepsilon}| = 0$ for all $0 < \varepsilon \leq \varepsilon_0$ and some $\varepsilon_0 > 0$. In both cases we construct a sequence $(\mathbf{f}_j)_{j \in \mathbb{N}} \subset L^p(\Omega)^m$ with $\|\mathbf{f}_j\|_{L^p(\Omega)^m} = 1$ for which $\|(\lambda I - \mathbf{A})\mathbf{f}_j\|_{L^p(\Omega)^m} \rightarrow 0$ as $j \rightarrow \infty$, hence $\lambda I - \mathbf{A}$ can not be bounded from below.

- Let $|\Gamma_{\lambda, \varepsilon}| > 0$ for all $\varepsilon > 0$. Thus, we are able to extract a decreasing subsequence $(\Gamma_{\lambda, \varepsilon_j})_{j \in \mathbb{N}}$ with $\varepsilon_j \rightarrow 0$ as $j \rightarrow \infty$ such that

$$|\Gamma_{\lambda, \varepsilon_j}| > 0, \quad \Gamma_{\lambda, \varepsilon_{j+1}} \subset \Gamma_{\lambda, \varepsilon_j} \quad \text{and} \quad |\Gamma_{\lambda, \varepsilon_j} \setminus \Gamma_{\lambda, \varepsilon_{j+1}}| > 0.$$

By choosing measurable sets $M_j \subset \Gamma_{\lambda, \varepsilon_j} \setminus \Gamma_{\lambda, \varepsilon_{j+1}}$ with $|M_j| > 0$ for all $j \in \mathbb{N}$, we obtain the estimate

$$\varepsilon_{j+1}^m \leq |\det(\lambda I - \mathbf{A}(x))| < \varepsilon_j^m \quad \forall x \in M_j. \quad (\text{B.5})$$

This enables us to apply [18, Lemma 3.1] to the matrix $(\lambda I - \mathbf{A}(x))^{-1}$. Consequently, we find measurable vector-valued functions $\mathbf{v}_j : M_j \rightarrow \mathbb{C}^m$ satisfying

$$|\mathbf{v}_j(x)|_2 = 1 \quad \text{and} \quad |(\lambda I - \mathbf{A}(x))^{-1} \mathbf{v}_j(x)|_2 = |(\lambda I - \mathbf{A}(x))^{-1}|_2$$

for all $x \in M_j$, where we used $|\cdot|_2$ for the Euclidean norm on \mathbb{C}^m and for the induced matrix norm. Define $\mathbf{u}_j(x) = (\lambda I - \mathbf{A}(x))^{-1} \mathbf{v}_j(x)$ as well as functions $\mathbf{f}_j \in L^p(\Omega)^m$ by

$$\mathbf{f}_j(x) = c_p(j) \frac{\mathbf{u}_j(x)}{|\mathbf{u}_j(x)|_2} \mathbf{1}_{M_j}(x)$$

where $c_p(j) = |M_j|^{-1/p}$ for $p < \infty$ and $c_p(j) = 1$ for $p = \infty$. This implies the normalization $\|\mathbf{f}_j\|_{L^p(\Omega)^m}^p = \int_\Omega |\mathbf{f}_j(x)|_2^p dx = 1$, with an obvious modification for $p = \infty$. Applying $\lambda I - \mathbf{A}$ to \mathbf{f}_j yields

$$(\lambda I - \mathbf{A}(x))\mathbf{f}_j(x) = c_p(j) \mathbf{1}_{M_j}(x) \mathbf{v}_j(x) |(\lambda I - \mathbf{A}(x))^{-1}|_2^{-1}.$$

From the invertibility condition (B.5) we infer

$$|(\lambda I - \mathbf{A}(x))^{-1}|_2^{-1} \leq \text{dist}(\lambda, \sigma(\mathbf{A}(x))) \quad \forall x \in M_j$$

where we used [11, Ch. IV, Corollary 1.14]. A combination of estimates (B.3) and (B.5) yields $\text{dist}(\lambda, \sigma(\mathbf{A}(x))) < \varepsilon_j$ for $x \in M_j$, which implies $\|(\lambda I - \mathbf{A})\mathbf{f}_j\|_{L^p(\Omega)^m} \leq \varepsilon_j$. Since $\varepsilon_j \rightarrow 0$, λ is an approximate eigenvalue of \mathbf{A} .

- Let $|\Gamma_{\lambda, \varepsilon}| = 0$ for all $0 < \varepsilon \leq \varepsilon_0$. The definition of $\Gamma_{\lambda, \varepsilon}$ yields the pointwise invertibility condition

$$|\det(\lambda I - \mathbf{A}(x))| \geq \varepsilon_0^m > 0 \quad \text{for a.e. } x \in \Omega.$$

Taking $M_j := N_{\lambda, \varepsilon_j} \subset \Omega$ with $|M_j| > 0$ for any zero sequence $(\varepsilon_j)_{j \in \mathbb{N}}$, we find, similar to the above reasoning, a sequence $(\mathbf{f}_j)_{j \in \mathbb{N}} \subset L^p(\Omega)^m$ satisfying $\|(\lambda I - \mathbf{A})\mathbf{f}_j\|_{L^p(\Omega)^m} \leq \varepsilon_j$. Since $\varepsilon_j \rightarrow 0$, λ is an approximate eigenvalue of \mathbf{A} . Note that in this case, $N_{\lambda, \varepsilon_j}$ cannot become stationary since then M_j and \mathbf{f}_j would become stationary which implies $(\lambda I - \mathbf{A})\mathbf{f}_j = \mathbf{0}$ – a contradiction to $\mathbf{f}_j \neq \mathbf{0}$.

It remains to show that $\lambda I - \mathbf{A}$ is not Fredholm for all $\lambda \in \sigma(\mathbf{A})$. To do so, we prove that for each $\lambda \in \sigma(\mathbf{A})$ either $\lambda I - \mathbf{A}$ has no closed range or an infinite-dimensional kernel.

If $\lambda \in \sigma_p(\mathbf{A})$, note that the results [18, Lemma 2.4, Theorem 2.5] hold independently of $1 \leq p \leq \infty$. Hence,

the first part of the proof of [18, Proposition 3.2] is still applicable: we infer $\sigma_p(\mathbf{A}) \subset \sigma_{\text{ess}}(\mathbf{A})$ from an infinite-dimensional kernel of $\lambda I - \mathbf{A}$ containing a subspace isomorphic to $L^p(\Gamma_\lambda)$ [18, Corollary 2.6].

If $\lambda \in \sigma(\mathbf{A}) \setminus \sigma_p(\mathbf{A})$, we necessarily have $|\Gamma_\lambda| = \lim_{\varepsilon \rightarrow 0} |\Gamma_{\lambda, \varepsilon}| = 0$. From above reasoning we know that $\lambda I - \mathbf{A}$ is not bounded from below. Thus, the injective operator $\lambda I - \mathbf{A}$ cannot have closed range by [4, Theorem 2.19, Remark 18] and $\lambda I - \mathbf{A}$ is not a Fredholm operator. \square

Note that the above proof may be shortened extremely for the cases $1 \leq p < \infty$. One can essentially use the same method of proof from [18, Proposition 3.2] for the case $p = 2$ having the characterization for the dual multiplication operator from [10, Ch. IX, Proposition 1.4] in mind.

References

- [1] R.A. Adams, J.F. Fournier, Sobolev spaces. Academic Press Amsterdam, 2003.
- [2] F.V. Atkinson, H. Langer, R. Mennicken, A.A. Shkalikov, The essential spectrum of some matrix operators. *Math. Nachr.*, Vol. 167, No. 1 (1994), pp. 5-20.
- [3] A. Bobrowski, M. Kunze, Irregular convergence of mild solutions of semilinear equations. *J. Math. Anal. Appl.*, Vol. 472, No. 2 (2019), pp. 1401-1419.
- [4] H. Brezis, Functional analysis, Sobolev spaces and partial differential equations. Springer-Verlag New York, 2010.
- [5] J. Carr, Applications of centre manifold theory. Springer-Verlag New York, 1981.
- [6] Ju.L. Daleckiĭ, M.G. Kreĭn, Stability of solutions of differential equations in Banach spaces. *Translations in Math. Monographs*, Vol. 43, Amer. Math. Soc. Providence, 1974.
- [7] R. Datko, Uniform asymptotic stability of evolutionary processes in a Banach space. *SIAM J. Math. Anal.*, Vol. 3, No. 3 (1972), pp. 428-445.
- [8] E.B. Davies, Heat kernels and spectral theory. *Cambridge Tracts in Math.*, Vol. 92, Univ. Press Cambridge, 1989.
- [9] T. Eisner, Stability of operators and operator semigroups. *Operator Theory: Advances and Applications*, Vol. 209, Birkhäuser-Verlag Basel, 2010.
- [10] K.-J. Engel, Operator matrices and systems of evolution equations. *Habilitationschrift*, Universität Tübingen, 1995.
- [11] K.-J. Engel, R. Nagel, One-parameter semigroups for linear evolution equations. Springer-Verlag New York, 2000.
- [12] L.C. Evans, Partial differential equations. 2nd Edition. Amer. Math. Soc. Providence, 2010.
- [13] P. Grisvard, Elliptic problems in nonsmooth domains. *Classics in Applied Mathematics*, Siam New York, 2011.
- [14] P.V. Hai, Polynomial stability of evolution cocycles and Banach function spaces. *Bull. Belg. Math. Soc. Simon Stevin*, Vol. 26, No. 2 (2019), pp. 299-314.
- [15] J.K. Hale, K. Sakamoto, Shadow systems and attractors in reaction-diffusion equations. *Appl. Anal.*, Vol. 32, No. 3-4 (1989), pp. 287-303.
- [16] G.W. Hanson, A.B. Yakovlev, Operator theory for electromagnetics: an introduction. Springer-Verlag New York, 2002.
- [17] M. Haragus, G. Iooss: Local bifurcations, center manifolds, and normal forms in infinite-dimensional dynamical systems. Springer-Verlag London, 2011.
- [18] V. Hardt, E. Wagenführer, Spectral properties of a multiplication operator. *Math. Nachr.*, Vol. 178, No. 1 (1996), pp. 135-156.
- [19] S. Härtling, A. Marciniak-Czochra, I. Takagi, Stable patterns with jump discontinuity in systems with Turing instability and hysteresis. *Disc. Cont. Dyn. Syst. - A*, Vol. 37, No. 2 (2017), pp. 757-800.
- [20] K. Hattaf, N. Yousfi, Global stability for reaction-diffusion equations in biology. *Comp. Math. Appl.*, Vol. 66, No. 8 (2013), pp. 1488-1497.
- [21] A. Henrot, Extremum problems for eigenvalues of elliptic operators. *Frontiers in Math.*, Birkhäuser-Verlag Basel, 2006.
- [22] V. Hernández-Santamaría, K. Le Balc'h, Local-controllability of the one-dimensional nonlocal Gray-Scott model with moving controls, *J. Evol. Equ.* (2021).
- [23] V. Hernández-Santamaría, E. Zuazua, Controllability of shadow reaction-diffusion systems. *J. Diff. Eq.*, Vol. 268, No. 7 (2019), pp. 3781-3818.
- [24] R. Heymann, Eigenvalues and stability properties of multiplication operators and multiplication semigroups. *Math. Nachr.*, Vol. 287, No. 5-6 (2014), pp. 574-584.
- [25] S. Hock, Y. Ng, J. Hasenauer, D. Wittmann, D. Lutter, D. Trümbach, W. Wurst, N. Prakash, F.J. Theis, Sharpening of expression domains induced by transcription and microRNA regulation within a spatio-temporal model of mid-hindbrain boundary formation. *BMC Syst. Biol.*, Vol. 7, No. 48 (2013).
- [26] T. Kato, Integration of the equation of evolution in a Banach space. *J. Math. Soc. Japan*, Vol. 5, No. 2 (1953), pp. 208-234.
- [27] J.P. Keener, Activators and inhibitors in pattern formation. *Studies in Appl. Math.*, Vol. 59, No. 1 (1978).
- [28] V. Klika, R.E. Baker, D. Headon, E.A. Gaffney, The influence of receptor-mediated interactions on reaction-diffusion mechanisms of cellular self-organization. *Bull. Math. Biol.*, Vol. 74 (2012), pp. 935-957.
- [29] S. Kondo, M. Mimura, A reaction-diffusion system and its shadow system describing harmful algal blooms. *Tamkang J. Math.*, Vol. 47, No. 1 (2016), pp. 71-92.
- [30] A. Köthe, A. Marciniak-Czochra, I. Takagi, Hysteresis-driven pattern formation in reaction-diffusion-ODE systems. *Disc. Cont. Dyn. Syst. - A*, Vol. 40, No. 6 (2020), pp. 3595-3627.
- [31] C. Kowall, Uniform shadow limit reduction for reaction-diffusion-ODE systems. *Dissertation*, Heidelberg University, 2021.

- [32] C. Kowall, A. Marciniak-Czochra, A. Mikelić, Long-time shadow limit for a reaction-diffusion-ODE system. *Appl. Math. Letters*, Vol. 112 (2021) 106790.
- [33] G.I. Kresin, V.G. Maz'ya, Criteria for validity of the maximum modulus principle for solutions of linear parabolic systems. *Ark. Mat.*, Vol. 32, No. 1 (1994), pp. 121-155.
- [34] O.A. Ladyzenskaja, V.A. Solonnikov, N.N. Ural'ceva, Linear and quasi-linear equations of parabolic type. *Translations of Mathematical Monographs Reprint*, Vol. 23, Amer. Math. Soc. Providence, 1968.
- [35] D.L. Lovelady, R.H. Martin, Jr., A global existence theorem for a nonautonomous differential equation in a Banach space. *Proc. Amer. Math. Soc.*, Vol. 35, No. 2 (1972), pp. 445-449.
- [36] A. Marasco, A. Iuorio, F. Cartení, G. Bonanomi, D.M. Tartakovsky, S. Mazzoleni, F. Giannino, Vegetation pattern formation due to interactions between water availability and toxicity in plant-soil feedback. *Bull. Math. Biol.*, Vol. 76 (2014), pp. 2866-2883.
- [37] A. Marciniak-Czochra, S. Härtling, G. Karch, K. Suzuki, Dynamical spike solutions in a nonlocal model of pattern formation. *Nonlinearity*, Vol. 31, No. 5 (2018).
- [38] A. Marciniak-Czochra, M. Kimmel, Reaction-diffusion model of early carcinogenesis: the effects of influx of mutated cells. *Math. Model. Nat. Phenom.*, Vol. 3, No. 7 (2008), pp. 90-114.
- [39] A. Marciniak-Czochra, A. Mikelić, Shadow limit using renormalization group method and center manifold method. *Vietnam J. Math.*, Vol. 45 (2017), pp. 103-125.
- [40] P. De Mottoni, F. Rothe, Convergence to homogeneous equilibrium state for generalized Volterra-Lotka systems with diffusion. *SIAM J. Appl. Math.*, Vol. 37, No. 3 (1979), pp. 648-663.
- [41] Y. Nishiura, Global structure of bifurcating solutions of some reaction-diffusion systems. *SIAM J. Math. Anal.*, Vol. 13, No. 4 (1982), pp. 555-593.
- [42] E.M. Ouhabaz, L^p contraction semigroups for vector valued functions. *Positivity*, Vol. 3 (1999), pp. 83-93.
- [43] E.M. Ouhabaz, *Analysis of heat equations on domains*. Princeton Univ. Press Princeton, 2005.
- [44] N.H. Pavel, Semilinear equations with dissipative time-dependent domain perturbations. *Israel J. Math.*, Vol. 46, No. 1-2 (1983), pp. 103-122.
- [45] A. Pazy, *Semigroups of linear operators and applications to partial differential equations*. Springer-Verlag New York, 1983.
- [46] F. Rothe, *Global solutions of reaction-diffusion systems*. *Lecture Notes in Math.*, Vol. 1072, Springer-Verlag Berlin, 1984.
- [47] R. Schnaubelt, Sufficient conditions for exponential stability and dichotomy of evolution equations. *Forum Math.*, Vol. 11, No. 5 (1999), pp. 543-566.
- [48] H. Takagi, Fredholm weighted composition operators. *Int. Equ. Oper. Theory*, Vol. 16 (1993), pp. 267-276.
- [49] I. Takagi, Point condensation for a reaction-diffusion system. *J. Diff. Equ.*, Vol. 61, No. 2 (1986), pp. 208-249.
- [50] M. Winkler, Aggregation vs. global diffusive behavior in the higher-dimensional Keller-Segel model. *J. Diff. Equ.*, Vol. 248, No. 12 (2010), pp. 2889-2905.
- [51] A. Yagi, *Abstract parabolic evolution equations and their applications*. Springer Monographs in Mathematics, Springer-Verlag Berlin, 2010.