

PARACANONICAL BASE LOCUS, ALBANESE MORPHISM, AND SEMI-ORTHOGONAL INDECOMPOSABILITY OF DERIVED CATEGORIES

FEDERICO CAUCCI

ABSTRACT. Motivated by an indecomposability criterion of Xun Lin for the bounded derived category of coherent sheaves on a smooth projective variety X , we study the paracanonical base locus of X , that is the intersection of the base loci of $\omega_X \otimes P_\alpha$, for all $\alpha \in \text{Pic}^0 X$. We prove that this is equal to the relative base locus of ω_X with respect to the Albanese morphism of X . As an application, we get that bounded derived categories of Hilbert schemes of points on certain surfaces do not admit non-trivial semi-orthogonal decompositions. We also have a consequence on the indecomposability of bounded derived categories in families. Finally, our viewpoint allows to unify and extend some results recently appearing in the literature.

1. INTRODUCTION

In this note we are interested in finding hypothesis ensuring the *indecomposability* of the bounded derived category of a variety X , that is the nonexistence of non-trivial semi-orthogonal decompositions of $\mathbf{D}^b(X)$ (see §2.1 for definitions). In recent years, bounded derived categories of projective varieties – and their semi-orthogonal decompositions – have been much studied. The interest in their indecomposability mainly rests on its conjectural relation with the minimal model program (see [KO] and §2.1):

Conjecture 1.1. Let X be a smooth projective variety. If $\mathbf{D}^b(X)$ has no non-trivial semi-orthogonal decompositions, then X is minimal, i.e., the canonical line bundle ω_X is nef.

Examples of varieties whose derived categories admit no non-trivial semiorthogonal decompositions are, for instance, varieties with trivial, or, more generally, algebraically trivial canonical bundle ([Br] and [KO, Corollary 1.7], respectively), curves of genus ≥ 1 [Ok], varieties whose Albanese morphism is finite [Pi, Theorem 1.4].¹ It is well known that the converse direction in Conjecture 1.1 is false (a counterexample is furnished by Enriques surfaces [Zu]), but we have the following folklore (see [BGL, Conjecture 1.6], or [BBOR, Question E]):

Conjecture 1.2. Let X be a smooth projective variety. If ω_X is nef and effective, then $\mathbf{D}^b(X)$ has no non-trivial semi-orthogonal decompositions.

At the moment of writing, this conjecture (as even the above Conjecture 1.1) is widely open in general and there are just some (classes of) varieties for which it has been verified: see [KO, BBOR], besides the references already quoted above. A particularly interesting case is given by symmetric

Key words and phrases. Derived categories, semi-orthogonal decompositions, Albanese morphism, canonical bundle, paracanonical base locus.

The author is supported by the ERC Consolidator Grant ERC-2017-CoG-771507-StabCondEn.

¹To be precise, in [Pi, Theorem 1.4], Pirozhkov proved that such varieties are noncommutatively stably semiorthogonally indecomposable, which is a stronger notion than indecomposability.

products of curves. It has been conjectured by Belmans-Galkin-Mukhopadhyay [BGM, Conjecture 1.1], and independently by Biswas-Gómez-Lee [BGL, Conjecture 1.4], that the bounded derived category of the n -th symmetric product of a smooth projective curve of genus $g \geq 2$ has no non-trivial semi-orthogonal decompositions, if $n \leq g - 1$. Very recently, Lin [Li, Theorem 1.4] nicely proved it by using a criterion that we will recall below (see Theorem 1.4). This complements some previous partial results due to [BGL] and [BBOR].

A result of Kawatani-Okawa [KO] is particularly inspiring for us: these authors established a relation between the base locus $\text{Bs}|\omega_X|$ of the canonical line bundle of X and the non-existence of semi-orthogonal decompositions (*SODs* for short) of $\mathbf{D}^b(X)$. Among other things, they proved [KO, Corollary 1.5] the following

Theorem 1.3 (Kawatani-Okawa). *If $\text{Bs}|\omega_X|$ is a finite set (possibly empty), then $\mathbf{D}^b(X)$ has no non-trivial SODs.*

So, in particular, a weak version of Conjecture 1.2 is known to be true: *the global generation of ω_X implies the nonexistence of SODs of $\mathbf{D}^b(X)$* . In [Li], Lin focused instead on the *paracanonical base locus* of a variety X , that is the closed subset

$$\text{PBs}|\omega_X| := \bigcap_{\alpha \in \text{Pic}^0 X} \text{Bs}|\omega_X \otimes P_\alpha|,$$

where ω_X is the canonical line bundle of X , and P_α denotes the topologically trivial line bundle on X corresponding to the (closed) point α of the Picard variety $\text{Pic}^0 X$. Lin refined Kawatani-Okawa Theorem as follows:

Theorem 1.4 (Lin). *If $\text{PBs}|\omega_X|$ is a finite set (possibly empty), then $\mathbf{D}^b(X)$ has no non-trivial SODs.*

Now the study of paracanonical base loci has its own interest. For instance, their emptiness (and related properties) has been implicitly studied in, e.g., [PPIII, BLNP, Ji], where several results of quite different flavor are obtained (see also [PP] and the references therein). Note that, since ω_X is an invertible sheaf, by definition $\text{PBs}|\omega_X|$ equals the support of the cokernel of the sum of evaluation maps

$$\bigoplus_{\alpha \in \text{Pic}^0 X} H^0(X, \omega_X \otimes P_\alpha) \otimes P_\alpha^\vee \rightarrow \omega_X.$$

Here we observe that, using some standard results of generic vanishing theory due to Chen, Jiang, Pareschi, Popa and Schnell (that will be recalled in §2.2 for reader's convenience), it is possible, quite easily, to give a different account of this locus, introducing a “relative” point of view (see Theorem 1.5 below). This allows to generalize Theorem 1.3 to a relative setting where the Albanese morphism

$$a_X: X \rightarrow \text{Alb } X$$

of X appears. More precisely, Lin's criterion can be read as a relative version of Kawatani and Okawa's one (see Corollary 1.7 below). Before giving the statement, let us fix some other notations. We denote as usual by

$$a_X^* a_{X*} \omega_X \rightarrow \omega_X$$

the adjunction morphism. The corresponding relative base ideal is

$$\mathfrak{b}_X := \mathfrak{b}(\omega_X, a_X) = \text{Im}[a_X^* a_{X*} \omega_X \otimes \omega_X^\vee \rightarrow \mathcal{O}_X],$$

and the *relative base locus of ω_X with respect to the Albanese morphism a_X* is, by definition, the closed subset cut out by the relative base ideal \mathfrak{b}_X .

Then our main result is

Theorem 1.5. *Let X be a smooth projective variety. Then the paracanonical base locus $\text{PBs}|\omega_X|$ equals the relative base locus of ω_X with respect to the Albanese morphism of X .*

In particular, we get that ω_X is a_X -relatively globally generated (this means that the natural morphism $a_X^*a_{X*}\omega_X \rightarrow \omega_X$ is surjective, i.e., $\mathfrak{b}_X = \mathcal{O}_X$) if and only if $\text{PBs}|\omega_X|$ is empty. Since a_X finite implies $a_X^*a_{X*}\omega_X \twoheadrightarrow \omega_X$, we have that

Corollary 1.6. *The paracanonical base locus of a variety with finite Albanese morphism is empty.*

Moreover, from Theorems 1.5 and 1.4, it follows

Corollary 1.7. *If \mathfrak{b}_X defines a finite set (possibly empty), then $\mathbf{D}^b(X)$ has no non-trivial SODs.*

This partially generalizes the above-mentioned result of Pirozhkov on varieties having finite Albanese morphism. It also allows to give an alternative proof of [Li, Theorem 1.4] on indecomposability of derived categories of symmetric products of curves (see §4.2), and to put both these results under the same perspective.

Our second result concerns indecomposability of derived categories of punctual Hilbert scheme on surfaces. Let S be a smooth projective surface and $S^{[n]}$ the Hilbert scheme of points of length n on S . In [BBOR, Proposition 5.1], the authors proved that, if $\text{Bs}|\omega_S|$ is empty, then, for all $n \geq 1$, $\text{Bs}|\omega_{S^{[n]}}|$ is empty too. Hence, by Theorem 1.3, $\mathbf{D}^b(\omega_{S^{[n]}})$ has no non-trivial SODs. It is natural to ask if the same holds true taking into account the paracanonical base locus instead of the usual canonical base locus. Theorem 1.5 turns out to be useful in this case. Indeed, by using it, we prove

Theorem 1.8. *Let $n > 0$ be a natural number. If $\text{PBs}|\omega_S| = \emptyset$, then*

$$\text{PBs}|\omega_{S^{[n]}}| = \emptyset.$$

In particular, $\mathbf{D}^b(\omega_{S^{[n]}})$ has no non-trivial SODs, if $\text{PBs}|\omega_S|$ is empty.

This should be seen as a counterpart to the symmetric product of curves situation mentioned above (see also §4.2 and Remark 4.4). It adds a new class of examples to the list of varieties appearing in the literature, whose derived categories are indecomposable. For instance, it applies to surfaces having finite Albanese morphism, but also to second symmetric products of curves (see Proposition 4.1). Indeed, the following result generalizes, respectively, [BGL, Theorem 1.3], and partially [Pi, Theorem 1.4] in dimension 2:

Corollary 1.9. a) *Let C be a smooth projective curve of genus ≥ 3 , and $C^{(2)}$ be its second symmetric product. Then the bounded derived category*

$$\mathbf{D}^b((C^{(2)})^{[n]})$$

of the Hilbert scheme of points of length n on $C^{(2)}$ has no non-trivial SODs, for any $n \geq 1$.

b) *Let S be a surface with finite Albanese morphism. Then $\mathbf{D}^b(S^{[n]})$ has no non-trivial SODs, for any $n \geq 1$.*

The proof of Theorem 1.8 will be given in §4.3. The last statement follows from Lin's criterion (Theorem 1.4).

Lastly, we consider indecomposability in families and we give a variant of [BBOR, Corollary B], where the authors proved that, if $f: \mathcal{X} \rightarrow T$ is a smooth projective family of varieties, with T is

an irreducible algebraic variety over \mathbb{C} , and there exists a point $t_0 \in T$ such that $\text{Bs}|\omega_{\mathcal{X}_{t_0}}|$ is finite (possibly empty), then $\mathbf{D}^b(\mathcal{X}_t)$ has no non-trivial SODs for all $t \in T$. Here, as usual, \mathcal{X}_t denotes the variety $f^{-1}(t)$. By assuming that the Albanese morphisms of the varieties appearing in the family f vary “smoothly”, we can prove that the same result is true, if the paracanonical base locus $\text{PBs}|\omega_{\mathcal{X}_{t_0}}|$ is finite (possibly empty) for a point $t_0 \in T$. Namely, in order to precisely state our result, let us recall that, given a smooth projective family $f: \mathcal{X} \rightarrow T$, there exists a commutative diagram

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{\varphi} & \text{Alb } \mathcal{X}/T \\ f \downarrow & \searrow \eta & \\ T & & \end{array} \quad (1.1)$$

where η is a smooth morphism such that, for all $t \in T$, the fiber $(\text{Alb } \mathcal{X}/T)_t$ is $\text{Alb } \mathcal{X}_t$, and the restriction $\varphi_t: \mathcal{X}_t \rightarrow (\text{Alb } \mathcal{X}/T)_t$ is the Albanese morphism $a_{\mathcal{X}_t}: \mathcal{X}_t \rightarrow \text{Alb } \mathcal{X}_t$ of \mathcal{X}_t . The morphism φ is a universal morphism from the family f to families of abelian varieties over T' , for any morphism $T' \rightarrow T$ (see [Gr2, n. 236], or [Fu, Ca] and the reference therein): φ is called the *relative Albanese morphism* for f and $\text{Alb } \mathcal{X}/T$ is the *relative Albanese variety* for f . Thanks to Theorem 1.5, we have the following result whose proof is the content of §5.

Theorem 1.10. *Let $f: \mathcal{X} \rightarrow T$ be a smooth projective family of varieties as above, and assume that its relative Albanese morphism φ is smooth. If $\text{PBs}|\omega_{\mathcal{X}_{t_0}}|$ is finite (possibly empty) for a certain point $t_0 \in T$, then $\mathbf{D}^b(\mathcal{X}_t)$ has no non-trivial SODs for all $t \in T$.*

We believe (and hope) that Theorem 1.5 could also be useful in some other context, possibly far from those considered in this note.

Notations. All varieties are assumed to be smooth, projective, irreducible and defined over \mathbb{C} . A coherent sheaf on a variety is simply called a *sheaf*. Let \mathcal{P} be the normalized Poincaré line bundle on $\text{Alb } X \times \text{Pic}^0 X$. Given a closed point $\alpha \in \text{Pic}^0 X$, we denote by $P_\alpha := \mathcal{P}|_{\text{Alb } X \times \{\alpha\}}$ the corresponding line bundle on $\text{Alb } X$. Recall that $\text{Pic}^0(\text{Alb } X) \cong \text{Pic}^0 X$ via the pullback along the Albanese morphism $a_X: X \rightarrow \text{Alb } X$ of X . So, by a slight abuse of notation, the line bundle $a_X^* P_\alpha$ on X corresponding to $\alpha \in \text{Pic}^0 X$ is simply denoted by P_α .

Acknowledgements. I thank Beppe Pareschi for answering a question of mine. I’m especially grateful to Xun Lin for very valuable comments on a first draft of this paper, and to Ángel D. R. Ortiz for our always fruitful conversations.

2. PREREQUISITES

Here we recall the definition of semi-orthogonal decomposition of a derived category, and some notions and results concerning the generic vanishing theory.

2.1. Semi-orthogonal decompositions. Let $\mathbf{D}^b(X)$ be the bounded derived category of a smooth projective variety X . If there exist full triangulated subcategories \mathcal{A} and \mathcal{B} of $\mathbf{D}^b(X)$ such that

$$\mathbf{D}^b(X) = \langle \mathcal{A}, \mathcal{B} \rangle, \quad (2.1)$$

i.e., \mathcal{A} and \mathcal{B} generate the triangulated category $\mathbf{D}^b(X)$, and

$$\text{Hom}(b, a) = 0$$

for any object $a \in \mathcal{A}$ and $b \in \mathcal{B}$, then we say that (2.1) is a *semi-orthogonal decomposition* (SOD for short) of $\mathbf{D}^b(X)$.

Definition 2.1. A semi-orthogonal decomposition of $\mathbf{D}^b(X)$ is said to be *non-trivial* if \mathcal{A} and \mathcal{B} are both non-zero. If $\mathbf{D}^b(X)$ has no non-trivial semi-orthogonal decompositions, it is said to be *indecomposable*.

We refer the reader to [Ku] for an overview on SODs in algebraic geometry.

The indecomposability of the derived category of a variety is (conjecturally) strongly related to its birational geometry. Indeed, the *DK-hypothesis* of [Ka2] (see also [Ka3]) predicts that, if there exists a K -inequality between two smooth projective varieties X and Y , say $X \leq_K Y$,² then there is a fully faithful functor of triangulated categories $\mathbf{D}^b(X) \hookrightarrow \mathbf{D}^b(Y)$. This would imply the existence of a semi-orthogonal decomposition of $\mathbf{D}^b(Y)$

$$\mathbf{D}^b(Y) = \left\langle \mathbf{D}^b(X)^\perp, \mathbf{D}^b(X) \right\rangle,$$

where $\mathbf{D}^b(X)^\perp$ is the right orthogonal complement of $\mathbf{D}^b(X)$ inside $\mathbf{D}^b(Y)$. See [Ka3], and the references therein, for more information about this (and related) ideas.

2.2. Generic vanishing. Let A be an abelian variety, and \mathcal{F} be a sheaf on A . We denote by

$$V^i(A, \mathcal{F}) := \{\alpha \in \text{Pic}^0 A \mid h^i(A, \mathcal{F} \otimes P_\alpha) > 0\}$$

the i -th *cohomological support loci* of \mathcal{F} . These are closed algebraic subset of $\text{Pic}^0 A$ by semicontinuity.

Definition 2.2. The sheaf \mathcal{F} is said to be:

- i) GV*, if $\text{codim}_{\text{Pic}^0 A} V^i(A, \mathcal{F}) \geq i$, for all $i \geq 0$;
- ii) M-regular*, if $\text{codim}_{\text{Pic}^0 A} V^i(A, \mathcal{F}) > i$, for all $i > 0$.

We refer the reader to the surveys [Pa] and [PP] (and the references therein) for the main results surrounding such notions.

For us, the central property is the following:

Proposition 2.3 ([PPI], Proposition 2.13). *Let \mathcal{F} be an M-regular sheaf on A . Then, for any non-empty open subset $U \subseteq \text{Pic}^0 A$, the sum of evaluation maps*

$$\bigoplus_{\alpha \in U} H^0(A, \mathcal{F} \otimes P_\alpha) \otimes P_\alpha^\vee \rightarrow \mathcal{F} \tag{2.2}$$

is surjective.

A sheaf satisfying the condition (2.2) is said to be *continuously globally generated*. This notion was introduced by Pareschi and Popa in [PPI], and it has to be considered as a “continuous” variant of the usual notion of generation by global sections.

Now, let $a: X \rightarrow A$ be a morphism from a smooth projective variety X to an abelian variety A . Then the higher direct images $R^j a_* \omega_X$ give non-trivial examples of *GV* sheaves (see [Ha]). Recently, it has been observed that indeed much more is true:

²This means that there exists a third smooth projective variety Z with birational morphisms $f: Z \rightarrow X$ and $g: Z \rightarrow Y$ such that $g^* \omega_Y \otimes f^* \omega_X^\vee$ is effective.

Theorem 2.4 (Chen-Jiang decomposition). *Given a morphism $a: X \rightarrow A$ and an integer $j \geq 0$, there exists a canonical decomposition*

$$R^j a_* \omega_X = \bigoplus_i \pi_i^* \mathcal{F}_i \otimes P_{\alpha_i}, \quad (2.3)$$

where $\pi_i: A \rightarrow A_i$ are quotient of abelian varieties with connected fibers, the \mathcal{F}_i 's are M -regular sheaves on A_i , and $\alpha_i \in \text{Pic}^0 A$ are torsion points.

This powerful result was proved by Chen-Jiang [CJ], assuming a generically finite, and soon after by Pareschi-Popa-Schnell [PPS] in complete generality. See also [Vi] for a simplified proof of Theorem 2.4 following the lines of [CJ]. The Chen-Jiang decomposition also holds true for the pushforwards of pluricanonical bundles $a_* \omega_X^{\otimes m}$, $m \geq 1$, by Lombardi-Popa-Schnell [LPS].

In particular, by [PPI, Proposition 2.13], the \mathcal{F}_i 's appearing in Theorem 2.4 are continuously globally generated sheaves on A_i . This fact – along with the decomposition (2.3) itself – is more than sufficient to prove Theorem 1.5.

3. PROOF OF THEOREM 1.5

We are now ready to give the proof of our first result. Let $a_X: X \rightarrow \text{Alb } X$ be the Albanese morphism of X , and

$$a_{X*} \omega_X = \bigoplus_i \pi_i^* \mathcal{F}_i \otimes P_{\alpha_i} \quad (3.1)$$

be the Chen-Jiang decomposition of $a_{X*} \omega_X$ (see §2.2). Hence, for each index i , $\pi_i: \text{Alb } X \rightarrow A_i$ is a quotient of abelian varieties with connected fibers, \mathcal{F}_i is an M -regular sheaf on A_i , and $\alpha_i \in \text{Pic}^0 X$ is a torsion point. The sheaves \mathcal{F}_i are continuously globally generated by [PPI, Proposition 2.13]. This means, in particular, that the sum of evaluation maps

$$\bigoplus_{\alpha \in \text{Pic}^0 A_i} H^0(A_i, \mathcal{F}_i \otimes P_\alpha) \otimes P_\alpha^\vee \rightarrow \mathcal{F}_i \quad (3.2)$$

is surjective. Note that the dual morphism $\widehat{\pi}_i: \text{Pic}^0 A_i \hookrightarrow \text{Pic}^0 X$ is injective, hence we simply denote by $\alpha = \widehat{\pi}_i(\alpha) \in \text{Pic}^0 X$ the image of an element $\alpha \in \text{Pic}^0 A_i$. Moreover, by the projection formula,

$$H^0(\text{Alb } X, \pi_i^* \mathcal{F}_i \otimes P_\alpha) = \begin{cases} H^0(A_i, \mathcal{F}_i \otimes P_\alpha) & \text{if } \alpha \in \text{Pic}^0 A_i, \\ 0 & \text{otherwise.} \end{cases} \quad (3.3)$$

Applying π_i^* to (3.2) and tensoring by P_{α_i} , we get that

$$\bigoplus_{\alpha \in \text{Pic}^0 X} H^0(\text{Alb } X, \pi_i^* \mathcal{F}_i \otimes P_\alpha) \otimes P_\alpha^\vee \otimes P_{\alpha_i} = \bigoplus_{\alpha \in \text{Pic}^0 A_i} H^0(\text{Alb } X, \pi_i^* \mathcal{F}_i \otimes P_\alpha) \otimes P_\alpha^\vee \otimes P_{\alpha_i} \rightarrow \pi_i^* \mathcal{F}_i \otimes P_{\alpha_i}$$

is surjective too, where the equality comes from (3.3). Now we change notation as follows: we call $\alpha_i - \alpha =: -\beta \in \text{Pic}^0 X$. So we have that the map

$$\bigoplus_{\beta \in \text{Pic}^0 X} H^0(\text{Alb } X, \pi_i^* \mathcal{F}_i \otimes P_{\alpha_i} \otimes P_\beta) \otimes P_\beta^\vee \rightarrow \pi_i^* \mathcal{F}_i \otimes P_{\alpha_i} \quad (3.4)$$

is surjective, and this holds true for any index i appearing in the Chen-Jiang decomposition of $a_{X*} \omega_X$. Therefore, taking the sum over i in (3.4) – and using (3.1) –, we get the following

Lemma 3.1. *Let X be a smooth projective variety. Then the sum of evaluation maps*

$$\bigoplus_{\beta \in \text{Pic}^0 X} H^0(\text{Alb } X, a_{X*}\omega_X \otimes P_\beta) \otimes P_\beta^\vee \rightarrow a_{X*}\omega_X \quad (3.5)$$

is surjective.

Now, if we apply a_X^* to (3.5) and post-compose with the natural morphism $a_X^*a_{X*}\omega_X \rightarrow \omega_X$, we recover the evaluation morphism

$$\bigoplus_{\beta \in \text{Pic}^0 X} H^0(X, \omega_X \otimes P_\beta) \otimes P_\beta^\vee \longrightarrow a_X^*a_{X*}\omega_X \longrightarrow \omega_X, \quad (3.6)$$

where we used that $H^0(\text{Alb } X, a_{X*}\omega_X \otimes P_\beta) = H^0(X, \omega_X \otimes P_\beta)$ for all $\beta \in \text{Pic}^0 X$. The proof of Theorem 1.5 follows from (3.6). Indeed, if $x \in X$ is not contained in the relative base locus of ω_X with respect to a_X , then the sum of evaluation maps

$$\bigoplus_{\beta \in \text{Pic}^0 X} H^0(X, \omega_X \otimes P_\beta) \otimes P_\beta^\vee \rightarrow \omega_X,$$

is surjective at x by (3.6). Therefore, by definition, $x \notin \text{PBs } |\omega_X|$. The other inclusion is completely similar (and indeed easier, since it does not need Lemma 3.1). \square

Remark 3.2. Let $a: X \rightarrow A$ be a morphism from a smooth projective variety X to an abelian variety A . The properties of $a_{X*}\omega_X$ we used in the proof of Theorem 1.5, hold also true for $a_*\omega_X$ (see §2.2). Therefore, if we define the *a-paracanonical base locus* as

$$\text{PBs }_a|\omega_X| := \bigcap_{\alpha \in \text{Pic}^0 A} \text{Bs } |\omega_X \otimes a^*P_\alpha|,$$

we have that this is equal to the relative base locus of ω_X with respect to the morphism $a: X \rightarrow A$. Note that $\text{PBs } |\omega_X| \subseteq \text{PBs }_a|\omega_X|$, for any a .

More generally, for any $m \geq 1$, the same happens for $a_*\omega_X^m$ by [LPS] (see the comment below Theorem 2.4), once one defines the *a-pluriparacanonical base locus* likewise.

4. EXAMPLES

In this section we present some non-trivial examples where the hypothesis of Corollary 1.7 can be (quickly) checked.

4.1. Varieties with finite Albanese morphism. As already observed, it is well known that, if a variety X has finite Albanese morphism a_X , then its canonical bundle ω_X is a_X -relatively globally generated. Thus, $\mathbf{D}^b(X)$ is indecomposable if a_X is finite. We refer the reader to the paper [Ka1] for some fundamental structural results on varieties with finite Albanese morphism, and to [Pi] for a stronger theorem on indecomposability, in this case.

4.2. Symmetric products of curves. Let C be a smooth projective curve of genus $g \geq 2$. Let $J(C)$ be its Jacobian variety. We denote by $C^{(n)}$ the n -th symmetric product of C . Let

$$a_n := a_{C^{(n)}} : C^{(n)} \rightarrow J(C), \quad D \mapsto \mathcal{O}_C(D - np_0) \quad (4.1)$$

be the Albanese morphism of $C^{(n)}$, where $p_0 \in C$ is a fixed point.

In [Li], by algebraically moving techniques, Lin proved that PBs $|\omega_{C^{(n)}}|$ is empty if $n \leq g-1$. In particular, $\mathbf{D}^b(C^{(n)})$ has no non-trivial SODs in that range. By making use of Theorem 1.5, we see how this fact also follows from a classical description of the symmetric product of curves appearing in [Sc].³ Namely, we have

Proposition 4.1. *Let $n \leq g-1$. Then the canonical bundle $\omega_{C^{(n)}}$ is a_n -relatively globally generated.*

Proof. Let

$$i_{n-1} : C^{(n-1)} \hookrightarrow C^{(n)}, \quad D \mapsto D + p_0,$$

and denote its image by $x = i_{n-1}(C^{(n-1)}) \subseteq C^{(n)}$. Let Θ be the theta divisor on $J(C)$, and $\theta \in \text{NS}(C^{(n)})$ the class of the pullback of Θ via a_n . We recall that algebraic and numerical equivalence of divisors coincide on $C^{(n)}$, and that

$$\omega_{C^{(n)}} = \theta + (g - n - 1)x$$

holds in $\text{NS}(C^{(n)})$ (see, e.g., [BGL, Lemma 2.1]). Now, it follows from [Sc, Proposition 10] that the natural morphism $a_n^* a_{n*} \mathcal{O}_{C^{(n)}}(x) \rightarrow \mathcal{O}_{C^{(n)}}(x)$ is surjective. Assume $m := g - n - 1 \geq 1$, then, by induction on m , we have that $a_n^* a_{n*} \mathcal{O}_{C^{(n)}}(mx) \rightarrow \mathcal{O}_{C^{(n)}}(mx)$ is surjective too. Indeed, the following diagram is commutative

$$\begin{array}{ccc} a_n^*(a_{n*} \mathcal{O}_{C^{(n)}}((m-1)x) \otimes a_{n*} \mathcal{O}_{C^{(n)}}(x)) & \xrightarrow{\text{ev} \otimes \text{ev}} & \mathcal{O}_{C^{(n)}}((m-1)x) \otimes \mathcal{O}_{C^{(n)}}(x) = \mathcal{O}_{C^{(n)}}(mx) \\ \downarrow & & \parallel \\ a_n^*(a_{n*} \mathcal{O}_{C^{(n)}}(mx)) & \xrightarrow{\text{ev}} & \mathcal{O}_{C^{(n)}}(mx) \end{array}$$

Therefore,

$$a_n^* a_{n*} \mathcal{O}_{C^{(n)}}(\theta + (g - n - 1)x) \rightarrow \mathcal{O}_{C^{(n)}}(\theta + (g - n - 1)x)$$

is surjective if $g - n - 1 \geq 0$, because $\mathcal{O}_{C^{(n)}}(\theta)$ is a pullback line bundle via a_n and the morphism a_n has connected fibers, hence $a_n^* a_{n*} \mathcal{O}_{C^{(n)}}(\theta) \cong \mathcal{O}_{C^{(n)}}(\theta)$ by the projection formula. It follows that $\omega_{C^{(n)}}$ is a_n -relatively globally generated. \square

4.3. Hilbert schemes of points on surfaces. Here we prove Theorem 1.8. Let S be a smooth projective surface. Given an integer $n > 0$, we denote by $S^{[n]}$ the Hilbert scheme of points of length n on S , and by $S^{(n)}$ the n -fold symmetric product. It is well-known (see [Fo1]) that $S^{[n]}$ is smooth and that the Hilbert-Chow morphism $\pi : S^{[n]} \rightarrow S^{(n)}$ is a canonical resolution of singularities of $S^{(n)}$. Moreover,

$$a_{S^{[n]}} : S^{[n]} \xrightarrow{\pi} S^{(n)} \xrightarrow{f_n} \text{Alb } S \tag{4.2}$$

is the Albanese morphisms of $S^{[n]}$, where f_n is the morphism induced by a_S by addition on $\text{Alb } S$. In particular, $\text{Alb}(S^{[n]}) \cong \text{Alb } S$ (see [Fo2], we also refer to [Hu1, Hu2] for a detailed account on these morphisms). Let us point out that (4.2) is a natural generalization of the morphism (4.1).

The Hilbert-Chow morphism π is a crepant resolution, i.e., $\omega_{S^{[n]}} = \pi^* \omega_{S^{(n)}}$, and $S^{(n)}$ is a Gorenstein variety, so that $\omega_{S^{(n)}}$ is a line bundle. It is easy to see that $\omega_{S^{[n]}}$ is $a_{S^{[n]}}$ -relatively globally generated, if $\omega_{S^{(n)}}$ is f_n -relatively globally generated. Indeed, since $\pi_* \mathcal{O}_{S^{[n]}} = \mathcal{O}_{S^{(n)}}$, by the projection formula we have that

$$a_{S^{[n]}}^* a_{S^{[n]}} \omega_{S^{[n]}} = \pi^* f_n^* f_{n*} \pi_* \pi^* \omega_{S^{(n)}} = \pi^* f_n^* f_{n*} \omega_{S^{(n)}}.$$

³Interestingly enough, the main result of [Sc] was also used by Belmans and Krug (see Remark 4.4 below) to give a simpler proof of a result of Toda constructing a non-trivial semi-orthogonal decomposition of $\mathbf{D}^b(C^{(n)})$, when $n \geq g \geq 2$.

Hence, if

$$f_n^* f_{n*} \omega_{S^{(n)}} \rightarrow \omega_{S^{(n)}}$$

is surjective, by applying π^* to it we get that

$$a_{S^{[n]}}^* a_{S^{[n]*}} \omega_{S^{[n]}} \rightarrow \omega_{S^{[n]}}$$

is surjective too.

So we reduced to prove

Proposition 4.2. *If ω_S is a_S -relatively globally generated, then*

$$f_n^* f_{n*} \omega_{S^{(n)}} \rightarrow \omega_{S^{(n)}}$$

is surjective.

Proof. Let S^n be the usual product variety. We consider the following commutative diagram

$$\begin{array}{ccc} S^n & \xrightarrow{q} & S^{(n)} \\ \downarrow a_{S^n} & & \downarrow f_n \\ (\text{Alb } S)^n & \xrightarrow{\sigma} & \text{Alb } S \end{array} \quad (4.3)$$

where q is the quotient morphism by the action of the symmetric group \mathfrak{S}_n on S^n , $a_{S^n} = (a_S)^n$ is the Albanese morphism of S^n , and σ is the addition morphism.

We make use of the following technical fact:

Proposition 4.3. *Let X be a smooth projective variety such that $\text{PBs } |\omega_X| = \emptyset$. Let $n \geq 1$ be a natural number. If $\sigma_n: \text{Alb } X^n \rightarrow \text{Alb } X$ denotes the addition morphism, then the sum of evaluation maps*

$$ev_n: \bigoplus_{\alpha \in \text{Pic}^0 X} H^0(X^n, \omega_{X^n} \otimes \sigma_n^* P_\alpha) \otimes \sigma_n^* P_\alpha^\vee \rightarrow \omega_{X^n}$$

*is surjective.*⁴

Granting it for the moment, we apply this to the surface $X = S$. So the locus

$$\bigcap_{\alpha \in \text{Pic}^0 S} \text{Bs } |\omega_{S^n} \otimes \sigma^* P_\alpha| = \emptyset.$$

Since S^n is quasi-compact, there exists a finite number N such that, for some $\alpha_1, \dots, \alpha_N \in \text{Pic}^0 S$, the intersection $\bigcap_{k=1}^N \text{Bs } |\omega_{S^n} \otimes \sigma^* P_{\alpha_k}|$ is empty, i.e., the sum of evaluation maps

$$\bigoplus_{k=1}^N H^0(S^n, \omega_{S^n} \otimes \sigma^* P_{\alpha_k}) \otimes \sigma^* P_{\alpha_k}^\vee \rightarrow \omega_{S^n} \quad (4.4)$$

is surjective. Let $q_*^{\mathfrak{S}_n} := (-)^{\mathfrak{S}_n} \circ q_*$ be the equivariant pushforward (see [Gr1, Chapitre V]). Since $q^* \omega_{S^{(n)}} = \omega_{S^n}$, and by definition $q_*^{\mathfrak{S}_n} \mathcal{O}_{S^n} = \mathcal{O}_{S^{(n)}}$ (see [Mu, pp. 66-67]), one has $q_*^{\mathfrak{S}_n} \omega_{S^n} = \omega_{S^{(n)}}$ thanks to the equivariant projection formula. Moreover, looking at the diagram (4.3),

$$q_*^{\mathfrak{S}_n} a_{S^n}^* \sigma^* P_\alpha = q_*^{\mathfrak{S}_n} q^* f_n^* P_\alpha = f_n^* P_\alpha,$$

⁴Otherwise said, if $\text{PBs } |\omega_X|$ is empty, then ω_{X^n} is relatively globally generated with respect to the composition $a_n := \sigma_n \circ a_{X^n}: X^n \rightarrow \text{Alb } X$, i.e., $\text{PBs } a_n^* |\omega_{X^n}| = \emptyset$ (see Remark 3.2).

for all $\alpha \in \text{Pic}^0 S$. Here the former equality comes from the commutativity of (4.3) and the latter again from the equivariant projection formula. Since q is finite, by taking the equivariant pushforward $q_*^{\mathfrak{S}_n}$ in (4.4), we see that the morphism

$$\bigoplus_{k=1}^N H^0(S^n, \omega_{S^n} \otimes \sigma^* P_{\alpha_k}) \otimes f_n^* P_{\alpha_k}^\vee \rightarrow \omega_{S^{(n)}} \quad (4.5)$$

is surjective too. Indeed, if \mathcal{R} is the kernel of (4.4), one has $R^1 q_*^{\mathfrak{S}_n} \mathcal{R} = (R^1 q_* \mathcal{R})^{\mathfrak{S}_n} = 0$, where the first equality follows from the Grothendieck spectral sequence and the fact that the invariant functor $(-)^{\mathfrak{S}_n}$ is exact. Now, by adjunction, the surjective morphism (4.5) factorizes as

$$\bigoplus_{k=1}^N H^0(S^n, \omega_{S^n} \otimes \sigma^* P_{\alpha_k}) \otimes f_n^* P_{\alpha_k}^\vee \rightarrow f_n^* f_{n*} \omega_{S^{(n)}} \rightarrow \omega_{S^{(n)}}.$$

So to conclude the proof of Proposition 4.2 – and hence of Theorem 1.8 – it only remains to give the

Proof of Proposition 4.3. The proof is essentially the same of [PPI, Proposition 2.12] and it uses the method of “reducible sections”. We proceed by induction on n . For $n = 1$ there is nothing to prove. Let us denote by $\pi_i: X^n \rightarrow X$ the i -th projection. Since $\alpha \in \text{Pic}^0 X$, we have

$$\sigma_n^* P_\alpha = \pi_1^* P_\alpha \otimes \dots \otimes \pi_n^* P_\alpha =: P_\alpha^{\boxtimes n},$$

where $\sigma_n: \text{Alb } X^n \rightarrow \text{Alb } X$ is the addition morphism. This can be easily proved from the see-saw principle (see, e.g., [Mu, pp. 74-75]) and induction on n .

Since we are assuming that $\text{PBs } |\omega_X| = \emptyset$, it certainly exists a set J such that, for any surjective function $f: J \rightarrow \text{Pic}^0 X$, one has that the sum of evaluation maps

$$\bigoplus_{j \in J} H^0(X, \omega_X \otimes P_{f(j)}) \otimes P_{f(j)}^\vee \rightarrow \omega_X$$

is surjective. Let $p: X^n = X^{n-1} \times X \rightarrow X^{n-1}$ be first projection. We consider the following commutative diagram, where the maps are given by evaluation:

$$\begin{array}{ccc} \bigoplus_{j \in J} H^0(p^* \omega_X^{\boxtimes n-1} \otimes p^* P_{f(j)}^{\boxtimes n-1}) \otimes H^0(\pi_n^* \omega_X \otimes \pi_n^* P_{f(j)}) \otimes \sigma_n^* P_{f(j)}^\vee & \xlongequal{\quad} & \bigoplus_{j \in J} H^0(\omega_{X^n} \otimes \sigma_n^* P_{f(j)}) \otimes \sigma_n^* P_{f(j)}^\vee \\ \downarrow & & \downarrow \\ \bigoplus_{j \in J} H^0(p^* \omega_X^{\boxtimes n-1} \otimes p^* P_{f(j)}^{\boxtimes n-1}) \otimes p^*(P_{f(j)}^\vee)^{\boxtimes n-1} \otimes \pi_n^* \omega_X & \longrightarrow & \omega_{X^n} \end{array}$$

By construction and the inductive hypothesis, the bottom horizontal morphism is surjective.⁵ Therefore, the support of the cokernel of the right vertical morphism, that equals $\text{Supp}(\text{Coker}(ev_n))$, is contained in $\bigcup_{j \in J} \text{Bs } |\pi_n^*(\omega_X \otimes P_{f(j)})|$. Since this holds true for any surjective function $f: J \rightarrow \text{Pic}^0 X$, it follows that

$$\text{Supp}(\text{Coker}(ev_n)) \subseteq \bigcap_{f: J \rightarrow \text{Pic}^0 X} \bigcup_{j \in J} \text{Bs } |\pi_n^*(\omega_X \otimes P_{f(j)})|. \quad (4.6)$$

⁵Indeed, the projection formula gives

$$H^0(p^* \omega_X^{\boxtimes n-1} \otimes p^* P_{f(j)}^{\boxtimes n-1}) = H^0(p_* \mathcal{O}_{X^n} \otimes \omega_X^{\boxtimes n-1} \otimes P_{f(j)}^{\boxtimes n-1}) = H^0(\omega_{X^{n-1}} \otimes \sigma_{n-1}^* P_{f(j)}).$$

But $\bigcap_{\alpha \in \text{Pic}^0 X} \text{Bs} |\pi_n^*(\omega_X \otimes P_\alpha)| = \pi_n^{-1}(\text{PBs} |\omega_X|) = \emptyset$, and this implies that the set on the right side of (4.6) is empty. Therefore, ev_n is surjective. \square

This concludes the proof of Theorem 1.8. \square

Remark 4.4. As observed, the case of surfaces treated in §4.3 is the natural generalization of the one of curves of §4.2. However, comparing Proposition 4.1 and Theorem 1.8, it appears a visible difference on the index n in the two cases. The reason behind that has already been properly explained in [BBOR, Remark 5.2].

In fact, by [To, Corollary 5.11], there exists a non-trivial semi-orthogonal decomposition of $\mathbf{D}^b(C^{(n)})$, where C is a curve of genus $g \geq 2$, if $n \geq g$ (see also [BK, Theorem D] and [JL, Corollary 3.8]). Note that this is what is expected by Conjecture 1.1, because $C^{(n)}$ is not minimal in the range $n \geq g \geq 2$.

5. INDECOMPOSABILITY IN FAMILIES: PROOF OF THEOREM 1.10

In this section we consider indecomposability of derived categories in certain families of varieties. Namely, taking notations as in the Introduction, we assume that $f: \mathcal{X} \rightarrow T$ is a smooth projective family of varieties, parametrized by a complex algebraic variety T , such that the relative Albanese morphism for f , denoted by

$$\varphi: \mathcal{X} \rightarrow \text{Alb } \mathcal{X}/T,$$

is a smooth morphism over T .

What we need to prove is the following

Proposition 5.1. *Under the above assumption, the function*

$$u: T \rightarrow \mathbb{N} \cup \{-\infty\}, \quad t \mapsto u(t) := \dim \text{PBs} |\omega_{\mathcal{X}_t}|$$

is upper semi-continuous.

Proof. The Proposition can be proved quite similarly to [BBOR, Proposition 2.5]. We give a sketch of it. Let us consider the exact sequence

$$\varphi^* \varphi_* \omega_f \otimes \omega_f^\vee \rightarrow \mathcal{O}_{\mathcal{X}} \rightarrow \mathcal{O}_{\mathcal{PB}} \rightarrow 0, \quad (5.1)$$

where $\omega_f := \det \Omega_{\mathcal{X}/T}$ is the relative canonical line bundle of the smooth morphism $f: \mathcal{X} \rightarrow T$, and $\mathcal{PB} \subseteq \mathcal{X}$ is the closed subscheme defined by the ideal sheaf $\text{Im}[\varphi^* \varphi_* \omega_f \otimes \omega_f^\vee \rightarrow \mathcal{O}_{\mathcal{X}}]$. Given any $t \in T$, we can restrict (5.1) to the fiber \mathcal{X}_t of f over t , obtaining the exact sequence

$$a_{\mathcal{X}_t}^* a_{\mathcal{X}_t*} \omega_{\mathcal{X}_t} \otimes \omega_{\mathcal{X}_t}^\vee \rightarrow \mathcal{O}_{\mathcal{X}_t} \rightarrow \mathcal{O}_{\mathcal{PB}_t} \rightarrow 0.$$

Indeed, from the cartesian diagram

$$\begin{array}{ccc} \mathcal{X}_t & \hookrightarrow & \mathcal{X} \\ \downarrow \varphi_t & & \downarrow \varphi \\ (\text{Alb } \mathcal{X}/T)_t & \xhookrightarrow{\iota} & \text{Alb } \mathcal{X}/T \end{array}$$

we get that

$$(\varphi^* \varphi_* \omega_f)|_{\mathcal{X}_t} = \varphi_t^* \iota^* \varphi_* \omega_f = \varphi_t^* \varphi_{t*} \omega_{\mathcal{X}_t} = a_{\mathcal{X}_t}^* a_{\mathcal{X}_t*} \omega_{\mathcal{X}_t},$$

where the first isomorphism follows by the commutativity of the diagram, and the last one by the definition of the relative Albanese morphism for f . The central isomorphism comes from the base change of the smooth morphism φ (see [BO, Lemma 1.3], or [Po, p. 276]). Indeed, from the formula in *op.cit.* we have

$$L\iota^*R\varphi_*\omega_f = R\varphi_{t*}\omega_{\mathcal{X}_t},$$

where L and R denote the left and right derived functors, respectively. By taking cohomology in degree 0, we get $L^0\iota^*(R\varphi_*\omega_f) = \varphi_{t*}\omega_{\mathcal{X}_t}$. So we need to prove that

$$L^0\iota^*(R\varphi_*\omega_f) = \iota^*\varphi_*\omega_f. \quad (5.2)$$

Note that, by definition and the projection formula, $R\varphi_*\omega_f = R\varphi_*(\omega_\varphi \otimes \varphi^*\omega_\eta) = R\varphi_*\omega_\varphi \otimes \omega_\eta$ (notations as in the diagram (1.1)). Since φ is a smooth morphism, the higher direct images $R^q\varphi_*\omega_\varphi$ are locally free sheaves for all q by Hodge theory (see, e.g., [EV, Theorem 10.21]). Hence, $R^q\varphi_*\omega_f$ are locally free too, and $L^p\iota^*(R^q\varphi_*\omega_f) = 0$ for $p > 0$. So the spectral sequence (see [Huy, (3.10), p. 81])

$$E_2^{p,q} = L^p\iota^*(R^q\varphi_*\omega_f) \Rightarrow L^{p+q}\iota^*(R\varphi_*\omega_f)$$

degenerates, and we get (5.2).

Therefore, thanks to the Theorem 1.5, we have that $\mathcal{PB}_t = \text{PBs}|\omega_{\mathcal{X}_t}|$ for any $t \in T$, and, since the fiber dimension of the proper morphism $\mathcal{PB} \rightarrow T$ is upper semi-continuous, this concludes the proof. \square

Proof of Theorem 1.10. At this point, the proof of Theorem 1.10 is exactly the same of [BBOR, Theorem B], where one uses Proposition 5.1 instead of [BBOR, Proposition 2.5] and Theorem 1.4 instead of [BBOR, Theorem 1.1]. \square

REFERENCES

- [BLNP] M. A. Barja, M. Lahoz, J. C. Naranjo and G. Pareschi, On the bicanonical map of irregular varieties, *J. Algebraic Geom.* **21** (2012), no. 3, 445–471.
- [BBOR] F. Bastianelli, P. Belmans, S. Okawa and A.T. Ricolfi, Indecomposability of derived categories in families, preprint arXiv:2007.00994v1 (2020).
- [BGM] P. Belmans, S. Galkin and S. Mukhopadhyay, Semiorthogonal decompositions for moduli of sheaves on curves, *Oberwolfach Reports 24/2018, Oberwolfach Workshop 1822, Interactions between Algebraic Geometry and Noncommutative Algebra*, 1473–1476.
- [BK] P. Belmans and A. Krug, Derived categories of (nested) Hilbert schemes, preprint arXiv:1909.04321v2 (2019).
- [BGL] I. Biswas, T. Gómez and K.-S. Lee, Semi-orthogonal decomposition of symmetric products of curves and canonical system, *Rev. Mat. Iberoam.* **37** (2021), no. 5, 1885–1896.
- [Br] T. Bridgeland, Equivalences of triangulated categories and Fourier-Mukai transforms, *Bull. London Math. Soc.* **31** (1999), no. 1, 25–34.
- [BO] A. Bondal and D. Orlov, Semiorthogonal decomposition for algebraic varieties arXiv:alg-geom/9506012 (1995).
- [Ca] F. Campana, Réduction d’Albanèse d’un morphisme propre et faiblement kählérien, I, *Compositio Math.* **54** (1985), no. 3, 373–398.
- [CJ] J.A. Chen and Z. Jiang, Positivity in varieties of maximal Albanese dimension, *J. Reine Angew. Math.* **736** (2018), 225–253.
- [EV] H. Esnault and E. Viehweg, *Lectures on vanishing theorems*, DMV Seminar, **20**, Birkhäuser Verlag, Basel, 1992.
- [Fo1] J. Fogarty, Algebraic families on an algebraic surface, *Amer. J. Math.* **90** (1968), 511–521.
- [Fo2] J. Fogarty, Algebraic families on an algebraic surface, II, The Picard scheme of the punctual Hilbert scheme, *Amer. J. Math.* **95** (1973), 660–687.
- [Fu] A. Fujiki, Relative algebraic reduction and relative Albanese map for a fiber space in \mathcal{C} , *Publ. Res. Inst. Math. Sci.* **19** (1983), no. 1, 207–236.
- [Gr1] A. Grothendieck, Sur quelques points d’algèbre homologique, *Tohoku Math. J. (2)* **9** (1957), 119–221.

- [Gr2] A. Grothendieck, *Fondements de la géométrie algébrique (Extraits du Séminaire Bourbaki 1957-1962)*, Paris 1962.
- [Ha] Ch. Hacon, A derived category approach to generic vanishing, *J. Reine Angew. Math.* **575** (2004), 173–187.
- [Hu1] M. E. Huibregtse, The Albanese mapping for a punctual Hilbert scheme: I. Irreducibility of the fibers, *Trans. Amer. Math. Soc.* **251** (1979), 267–285.
- [Hu2] M. E. Huibregtse, The Albanese mapping for a punctual Hilbert scheme: II. Symmetrized differentials and singularities, *Trans. Amer. Math. Soc.* **274** (1982), no. 1, 109–140.
- [Huy] D. Huybrechts, *Fourier-Mukai transforms in algebraic geometry*, Clarendon Press - Oxford, 2006.
- [Ji] Z. Jiang, On Severi type inequalities, *Math. Ann.* **379** (2021), no. 1-2, 133–158.
- [JL] Q. Jiang and N. C. Leung, Derived category of projectivization and flops, preprint arXiv:1811.12525v4 (2019).
- [Ka1] Y. Kawamata, Characterization of abelian varieties, *Compositio Math.* **43** (1981), no. 2, 253–276.
- [Ka2] Y. Kawamata, D-equivalence and K-equivalence, *J. Diff. Geom.* **61** (2002), 147–171.
- [Ka3] Y. Kawamata, Birational geometry and derived categories, in *Surveys in differential geometry 2017. Celebrating the 50th anniversary of the Journal of Differential Geometry*, *Surv. Differ. Geom.*, **22** (Int. Press, Somerville, MA, 2018), 291–317.
- [KO] K. Kawatani and S. Okawa, Nonexistence of semiorthogonal decompositions and sections of the canonical bundle, preprint arXiv:1508.00682v2 (2018).
- [Ku] A. Kuznetsov, Semiorthogonal decompositions in algebraic geometry, in *Proceedings of the International Congress of Mathematicians-Seoul 2014*, Vol. II, 635–660, Kyung Moon Sa, Seoul, 2014.
- [Li] X. Lin, On nonexistence of semi-orthogonal decompositions in algebraic geometry, preprint arXiv:2107.09564v2 (2021).
- [LPS] L. Lombardi, M. Popa and Ch. Schnell, Pushforwards of pluricanonical bundles under morphisms to abelian varieties, *J. Eur. Math. Soc.* **22** (2020), no. 8, 2511–2536.
- [Mu] D. Mumford, *Abelian varieties*, Tata Institute of Fundamental Research Studies in Mathematics **5**, Hindustan Book Agency, New Delhi, 2008 (With appendices by C. P. Ramanujam and Yuri Manin. Corrected reprint of the second (1974) edition).
- [Ok] S. Okawa, Semi-orthogonal decomposability of the derived category of a curve, *Adv. Math.* **228** (2011), no. 5, 2869–2873.
- [Pa] G. Pareschi, Basic results on irregular varieties via Fourier-Mukai methods, in *Current Developments in Algebraic Geometry*, *Math. Sci. Res. Inst. Publ.*, vol. 59 (Cambridge Univ. Press, Cambridge, 2012), 379–403.
- [PPI] G. Pareschi and M. Popa, Regularity on abelian varieties I, *J. Amer. Math. Soc.* **16** (2003), no. 2, 285–302.
- [PP] G. Pareschi and M. Popa, M-regularity and the Fourier-Mukai transform, *Pure Appl. Math. Q.* **4** (2008), no. 3, Special Issue: In honor of Fedor Bogomolov. Part 2, 587–611.
- [PPIII] G. Pareschi and M. Popa., Regularity on abelian varieties III: relationship with generic vanishing and applications, in *Grassmannians, moduli spaces and vector bundles*, 141–167, *Clay Math. Proc.* **14**, Amer. Math. Soc., Providence, RI, 2011.
- [PPS] G. Pareschi, M. Popa and C. Schnell, Hodge modules on complex tori and generic vanishing for compact Kähler manifolds, *Geom. Topol.* **21** (2017), no. 4, 2419–2460.
- [Pi] D. Pirozhkov, Stably semiorthogonally indecomposable varieties, preprint arXiv:2011.12743v2 (2021).
- [Po] A. Polishchuk, *Abelian varieties, theta functions and the Fourier transform*, Cambridge Univ. Press, 2002.
- [Sc] R. L. E. Schwarzenberger, Jacobians and symmetric products, *Illinois J. Math.* **7** (1963), 257–268.
- [To] Y. Toda, Semiorthogonal decompositions of stable pair moduli spaces via d-critical flips, *J. Eur. Math. Soc.* **23** (2021), no. 5, 1675–1725.
- [Vi] M. B. Villadsen, Chen-Jiang decompositions for projective varieties, without Hodge modules, published online at *Math. Z.* doi.org/10.1007/s00209-021-02851-2 (2021).
- [Zu] S. Zube, Exceptional vector bundles on Enriques surfaces, *Math. Notes* **61** (1997), no. 5-6, 693–699.

DIPARTIMENTO DI MATEMATICA “FEDERIGO ENRIQUES”, UNIVERSITÀ DEGLI STUDI DI MILANO, VIA CESARE SARDINI 50, 20133 MILANO – ITALY

Email address: federico.caucci@unimi.it