

COMPARISON RESULTS FOR POISSON EQUATION WITH MIXED BOUNDARY CONDITION ON MANIFOLDS

HAIQING CHENG, TENGFEI MA, AND KUI WANG

ABSTRACT. In this article, we study Talenti's comparison results for Poisson equation with mixed boundary condition on manifolds. Precisely, using symmetrization techniques and isoperimetric inequalities, we establish a L^1 estimate for solutions to Poisson equation with mixed boundary condition, on complete noncompact manifolds with nonnegative Ricci curvature and compact manifolds with positive Ricci curvature respectively. Particularly, on Riemann surfaces we obtain a Talenti's comparison result. Our results generalize the main theorems in [2] to Riemannian setting.

1. INTRODUCTION

Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with smooth boundary, $\Omega^\sharp \subset \mathbb{R}^n$ be a round ball with the same volume as Ω , $f(x)$ be a nonnegative function on Ω and f^\sharp be the Schwarz rearrangement of f (see Definition 2.1). If $u(x)$ and $v(x)$ are solutions to

$$-\Delta u(x) = f(x), \quad x \in \Omega,$$

and

$$-\Delta v(x) = f^\sharp(x), \quad x \in \Omega^\sharp,$$

with Dirichlet boundary conditions. Talenti [19] proved that

$$(1.1) \quad u^\sharp(x) \leq v(x), \quad x \in \Omega^\sharp,$$

where $u^\sharp(x)$ is the Schwarz rearrangement of u , and Kesavan [16] proved that equality holds in (1.1) if only if $f(x)$ is a radial function and Ω is a round ball. The proof of (1.1) is based on symmetrization techniques and the isoperimetric inequality. Talenti's comparison results play an important role in both partial differential equations and geometry problems, for instance there provide L^∞ estimate for solutions to PDEs and Faber-Krahn type inequality for the first Dirichlet eigenvalue, see [8, 14, 16]. Talenti's comparison results were generalized to nonlinear elliptic and parabolic equations with Dirichlet boundary condition (see for instance [4, 5, 6, 20] and references therein), to compact Riemannian manifolds with positive Ricci curvature [11], and to noncompact manifold with nonnegative Ricci curvature and positive asymptotic volume ratio [8] as well. We also refer the reader to the excellent books [15, 17] for related topics.

Recently, Alvino, Nitsch and Trombetti [3] proved (1.1) for solutions to Poisson equation on planer domains with Robin boundary condition with positive Robin parameter, and a comparison principle for dimension $n \geq 3$. These results were generalized to Riemannian

2020 *Mathematics Subject Classification.* 53C42, 53C44.

Key words and phrases. Talenti's comparison, Robin boundary, Riemannian manifolds.

The research of the third author is supported by NSFC No.11601359.

manifolds [9], to Robin boundary with nonconstant Robin parameter [2], and to the torsion problem for the Hermite operator as well, see [10] and [12].

The purpose of the present paper is to study Talent's comparison and related estimates for solutions to Poisson equation with mixed boundary on manifolds. In particular, we generalize Alvino-Chiacchio-Nitsch-Trombetti's result [2] to compact manifolds with positive Ricci curvature and noncompact manifolds with nonnegative Ricci curvature.

Let (M, g) be a complete Riemannian manifold of dimension n , which is either compact with $\text{Ric} \geq (n-1)\kappa$ for $\kappa > 0$ or noncompact with nonnegative Ricci curvature and positive asymptotic volume ratio. Denote by

$$(1.2) \quad \theta := \begin{cases} \lim_{r \rightarrow \infty} \frac{|B_p(r)|}{\omega_n r^n}, & \kappa = 0, \\ \frac{|M|}{|M_\kappa|}, & \kappa > 0, \end{cases}$$

where $B_p(r)$ is the geodesic ball centered at p with radius r in M , M_κ is the n dimensional space form of constant sectional curvature κ , ω_n is the volume of the unit ball in \mathbb{R}^n , and $|M|$ denotes the volume of M . It follows from the volume comparison that $\theta \leq 1$. Let $\Omega \subset M$ be a bounded domain with smooth boundary, $f(x)$ be a smooth and nonnegative function not identically zero on Ω , and $\beta(x)$ be a smooth and positive function on $\partial\Omega$. We consider the following Poisson equation with mixed boundary

$$(1.3) \quad \begin{cases} -\Delta u(x) = f(x), & x \in \Omega, \\ \frac{\partial u}{\partial \nu} + \beta(x)u = 0, & x \in \partial\Omega, \end{cases}$$

where Δ denotes the Laplace-Beltrami operator and ν denotes the outward unit normal to $\partial\Omega$. Since $\beta(x) > 0$ and $f \geq 0$, it then follows easily from maximum principle that the solution to (1.3) is positive on Ω , see Proposition 2.1. Let Ω^\sharp be a geodesic ball in M_κ satisfying $\theta|\Omega^\sharp| = |\Omega|$, f^\sharp defined on Ω^\sharp be the Schwarz rearrangement of f , and $v(x)$ be the solution to the following Schwarz rearrangement equation of (1.3)

$$(1.4) \quad \begin{cases} -\Delta v(x) = f^\sharp(x), & x \in \Omega^\sharp, \\ \frac{\partial v}{\partial \nu} + \bar{\beta}v = 0, & x \in \partial\Omega^\sharp, \end{cases}$$

where $\bar{\beta}$ is a positive constant defined by

$$(1.5) \quad \bar{\beta} = \frac{\theta|\partial\Omega^\sharp|}{\int_{\partial\Omega} \frac{1}{\beta(x)} dA}.$$

Throughout the paper dA denotes the induced measure on boundary $\partial\Omega$ and $\partial\Omega^\sharp$. Since the boundary value problem (1.4) is radially symmetric, then the solution $v(x)$ is radially symmetric as well. Moreover $v(x)$ is monotone decreasing along the radial direction, see Proposition 2.2.

Our first result is concerning a L^1 comparison between u and v . Precisely, we prove

Theorem 1.1. *Let (M, g) be a complete Riemannian manifold of dimension n , which is either compact with $\text{Ric} \geq (n-1)\kappa$ for $\kappa > 0$ or noncompact with nonnegative Ricci curvature and positive asymptotic volume ratio. Let $u(x)$ and $v(x)$ be the solutions to (1.3) and (1.4) respectively. If $n \geq 3$, we assume further that for all measurable $E \subset \Omega$,*

$$(1.6) \quad \int_E f(x) dx \leq \frac{|E|^{\frac{n-2}{n}}}{|\Omega|^{\frac{n-2}{n}}} \int_\Omega f(x) dx.$$

Then

$$(1.7) \quad \|u\|_{L^1(\Omega)} \leq \theta \|v\|_{L^1(\Omega^\sharp)}.$$

Remark 1.2. *In this paper, we mainly focus on L^1 comparison (1.6) on manifolds and we then set up Theorem 1.1 in smooth case. In fact, Theorem 1.1 still remains valid under the assumptions that Ω is a bounded domain with Lipschitz boundary, $f(x) \in L^2(\Omega)$ and $\beta(x)$ is a positive measurable function defined on $\partial\Omega$, see [2].*

Similarly as in [2], when $n = 2$ and f is a constant, we show a Talenti's comparison, which reads

Theorem 1.3. *Let (M, g) be a complete Riemannian surface, which is either compact with $\text{Ric} \geq \kappa > 0$ or noncompact with nonnegative Ricci curvature and positive asymptotic volume ratio. Suppose $f(x) \equiv 1$. Then*

$$(1.8) \quad u^\sharp(x) \leq v(x), \quad x \in \Omega^\sharp.$$

Equality holds if and only if Ω is isometric to Ω^\sharp and $\beta(x)$ is a constant on $\partial\Omega$.

This paper is organized as follows. In Section 2 we recall the Schwarz rearrangements, isoperimetric inequalities for manifolds, and some properties on solutions to Poisson equation with Robin boundary condition. In Section 3, we prove Theorem 1.1 and Theorem 1.2.

2. PRELIMINARIES

2.1. Notations and Schwarz rearrangement. Let (M, g) be a complete Riemannian manifold of dimension n with $\text{Ric} \geq (n-1)\kappa$, which is either noncompact with $\kappa = 0$ and positive asymptotic volume ratio or compact with $\kappa > 0$. Let M_κ be the n dimensional space form of constant sectional curvature κ . Without loss of generality, we assume $\kappa = 1$ if $\kappa > 0$ throughout this paper. Let Ω be a bounded smooth domain in M and Ω^\sharp be a geodesic ball in M_κ (i.e. \mathbb{R}^n or \mathbb{S}^n) with volume $|\Omega|/\theta$, where θ is a constant defined by (1.2).

We recall the definitions and properties of the Schwarz rearrangement of nonnegative functions on manifolds, see also [8, Section 2] and [11, Section 2].

Definition 2.1. *Let $h(x)$ be a nonnegative measurable function on Ω . Denote by $\Omega_{h,t} = \{x \in \Omega : h(x) > t\}$ and $\mu_h(t) = |\Omega_{h,t}|$, the decreasing rearrangement h^* of h is defined by*

$$(2.1) \quad h^*(s) = \begin{cases} \text{ess sup}_{x \in \Omega} h(x), & s = 0, \\ \inf\{t \geq 0 : \mu_h(t) < s\}, & s > 0, \end{cases}$$

for $s \in [0, |\Omega|]$. The Schwarz rearrangement of h is defined by

$$(2.2) \quad h^\sharp(x) = h^*(\theta\omega_n r^n(x)), \quad x \in \Omega^\sharp,$$

where $r(x)$ is the distance function from the center of Ω^\sharp in M_κ , and ω_n is the volume of unit ball in \mathbb{R}^n .

It follows directly from (2.1) and (2.2) that

$$(2.3) \quad \mu_h(t) = \theta\mu_{h^\sharp}(t)$$

for $t \geq 0$. Meanwhile, the Fubini's theorem gives

$$(2.4) \quad \int_{\Omega} h^p(x) dx = \int_0^{|\Omega|} (h^*)^p(s) ds = \theta \int_{\Omega^\sharp} (h^\sharp)^p(x) dx,$$

for $h \in L^p(\Omega)$, $p \geq 1$. Moreover for any nonnegative functions $f(x)$ and $g(x)$, the following inequality, known as Hardy-Littlewood inequality, holds true

$$(2.5) \quad \int_{\Omega} f(x)g(x) dx \leq \int_0^{|\Omega|} f^*(s)g^*(s) ds.$$

Taking $g(x)$ as the characteristic function of $\Omega_{h,t}$ in above inequality yields

$$(2.6) \quad \int_{\Omega_{h,t}} f(x) dx \leq \int_0^{\mu_h(t)} f^*(s) ds.$$

2.2. Isoperimetric inequality on manifolds. To prove Theorem 1.1 and Theorem 1.3, we require the following isoperimetric inequality on manifolds with Ricci curvature bounded from below.

Theorem 2.2. *With M , θ , Ω and Ω^\sharp as above, there holds*

$$(2.7) \quad |\partial\Omega| \geq \theta|\partial\Omega^\sharp|,$$

where $|\partial\Omega|$ denotes the $(n-1)$ -dimensional area of $\partial\Omega$. The equality holds if and only if Ω is isometric to Ω^\sharp .

When $\text{Ric} \geq n-1$, inequality (2.7) was shown by Lévy and Gromov [13], see also Theorem 2.1 of [18]. When M is noncompact and $\text{Ric} \geq 0$, inequality (2.7) was shown in dimension three by Agostiniani, Fogagnolo and Mazzieri [1] and in all dimensions by Brendle [7]. Meanwhile, Brendle proved in [7] that (2.7) also holds true when Ω is a compact minimal submanifold of M of dimension $n+2$ with nonnegative sectional curvature.

2.3. Poisson equation with Robin boundary. In this subsection, we collect some known facts about solutions to Poisson equation with Robin boundary condition.

Proposition 2.1. *Let $u(x)$ be the solution to (1.3). Assume $f(x)$ is nonnegative and not identically zero on Ω , and $\beta(x) > 0$ on $\partial\Omega$. Then*

$$u(x) > 0$$

for $x \in \Omega$.

Proof. Set $\bar{u}(x) = \frac{1}{2}(u(x) - |u(x)|)$, then using integration by parts we estimate that

$$\begin{aligned} \int_{\Omega} \langle \nabla \bar{u}, \nabla u \rangle dx &= \int_{\partial\Omega} \bar{u} \frac{\partial u}{\partial \nu} dA - \int_{\Omega} \bar{u} \Delta u dx \\ &= - \int_{\partial\Omega} \beta u \bar{u} dA + \int_{\Omega} \bar{u} f dx \\ &\leq 0, \end{aligned}$$

where we used equation (1.3) in the second equality and nonnegativity of f in the last step. Observing

$$\int_{\Omega} \langle \nabla \bar{u}, \nabla u \rangle dx = \int_{\Omega} |\nabla \bar{u}|^2 dx,$$

we then obtain

$$\int_{\Omega} |\nabla \bar{u}|^2 dx = 0,$$

gives $\bar{u} = 0$, hence $u(x) \geq 0$. Since $u(x)$ is sup-harmonic by equation (1.3) and f is not identically zero, then $u(x) > 0$ on Ω . \square

Proposition 2.2. *Let $v(x)$ be the solution to (1.4), written as $\varphi(r(x))$. If $f(x)$ is non-negative and not identically zero on Ω , and $\bar{\beta} > 0$, then*

$$(2.8) \quad \varphi'(r) < 0$$

for $r \in (0, R_0)$, where $\Omega^\sharp = B_o(R_0)$.

Proof. We rewrite $f^\sharp(x)$ as $h(r(x))$ and denote by

$$\text{sn}_\kappa(r) = \begin{cases} \sin r, & \kappa = 1, \\ r, & \kappa = 0, \end{cases}$$

then equation (1.4) gives

$$\varphi''(r) + (n-1) \frac{\text{sn}'_\kappa(r)}{\text{sn}_\kappa(r)} \varphi'(r) = -h(r).$$

Recalling $h \geq 0$, we then have

$$(\text{sn}_\kappa^{n-1} \varphi')' = -\text{sn}_\kappa^{n-1}(r)h(r) \leq 0,$$

for $r \in (0, R_0)$. Moreover there exists a $\varepsilon_0 < R_0$ such that $h(r) > 0$ for $r < \varepsilon_0$, since h is monotone nonincreasing and not identically zero. Thus $\varphi'(r) < 0$ for $0 < r < \varepsilon_0$, hence $\varphi'(r) < 0$ for all $r \in (0, R_0)$. \square

3. PROOF OF THEOREM 1.1 AND THEOREM 1.3

In this section, we will prove the main theorems. For simplicity, we write $\Omega_{u,t}$ as Ω_t , and $\Omega_{v,t}^\sharp$ as Ω_t^\sharp for short, for a.e. $t > 0$. Let

$$a(s) = \frac{s^{\frac{n-1}{n}}}{|\partial B_s|}$$

for $s \in (0, |\Omega^\sharp|)$, where B_s is a round geodesic ball in M_κ with volume s . It can be easily checked that $a(s) = n^{-1}\omega_n^{-1/n}$ if $\kappa = 0$, and $a(s)$ is monotone increasing in s if $\kappa = 1$. We prove the following lemma, essentially due to isoperimetric inequality (2.7), which will be used later.

Lemma 3.1. *Under the hypotheses of Theorem 1.1, we have*

$$(3.1) \quad \theta^2 \left(\frac{\mu_u(t)}{\theta} \right)^{\frac{2n-2}{n}} \leq a^2 \left(\frac{\mu_u(t)}{\theta} \right) \left(-\mu'_u(t) + \int_{\partial\Omega_t \cap \partial\Omega} \frac{1}{\beta(x)u(x)} dA \right) \int_0^{\mu_u(t)} f^*(s) ds$$

and

$$(3.2) \quad \mu_v^{\frac{2n-2}{n}}(t) = a^2(\mu_v(t)) \left(-\mu'_v(t) + \int_{\partial\Omega_t^\sharp \cap \partial\Omega^\sharp} \frac{1}{\bar{\beta}v(x)} dA \right) \int_0^{\mu_v(t)} (f^\sharp)^*(s) ds$$

for a.e. $t > 0$.

Proof. By using the Morse-Sard theorem, we have

$$\partial\Omega_t = \{x \in \Omega : u(x) = t\} \cup \{x \in \partial\Omega : u(x) \geq t\}$$

for a.e. $t > 0$. Let

$$g(x) = \begin{cases} |\nabla u|, & x \in \partial\Omega_t \cap \Omega, \\ -\frac{\partial u}{\partial \nu}, & x \in \partial\Omega_t \cap \partial\Omega, \end{cases}$$

and using the divergence theorem and equation (1.3) we have

$$\int_{\partial\Omega_t} g(x) dA = \int_{\partial\Omega_t} -\frac{\partial u}{\partial \nu} dA = - \int_{\Omega_t} \Delta u dx = \int_{\Omega_t} f(x) dx,$$

so using (2.6), we obtain

$$(3.3) \quad \int_{\Omega_t} g(x) dA \leq \int_0^{\mu_u(t)} f^*(s) ds.$$

By using Hölder inequality, inequality (3.3) and the coarea formula, we estimate that

$$(3.4) \quad \begin{aligned} |\partial\Omega_t|^2 &\leq \int_{\partial\Omega_t} g(x) dA \int_{\partial\Omega_t} \frac{1}{g(x)} dA \\ &\leq \int_0^{\mu_u(t)} f^*(s) ds \left(\int_{\partial\Omega_t \cap \Omega} \frac{1}{|\nabla u|} dA + \int_{\partial\Omega_t \cap \partial\Omega} \frac{1}{\beta(x)u(x)} dA \right) \\ &= \int_0^{\mu_u(t)} f^*(s) ds \left(-\mu'_u(t) + \int_{\partial\Omega_t \cap \partial\Omega} \frac{1}{\beta(x)u(x)} dA \right). \end{aligned}$$

On the other hand, the isoperimetric inequality (2.7) yields

$$(3.5) \quad |\partial\Omega_t| \geq \theta |\partial(\Omega_t^\sharp)| = \frac{\theta}{a(\mu_u(t)/\theta)} \left(\frac{\mu_u(t)}{\theta} \right)^{\frac{n-1}{n}}.$$

Combining (3.4) and (3.5), we have

$$\frac{\theta^2}{a^2(\mu_u(t)/\theta)} \mu_u^{\frac{2n-2}{n}}(t) \leq \int_0^{\mu_u(t)} f^*(s) ds \left(-\mu'_u(t) + \int_{\partial\Omega_t \cap \partial\Omega} \frac{1}{\beta(x)u(x)} dA \right).$$

This proves (3.1).

If $v(x)$ is the solution to (1.4), $v(x)$ is a radial function on Ω^\sharp and decreasing along the radial direction (see (2.8)), and then Ω_t^\sharp is a round ball. Therefore all previous inequalities hold as equalities if we replace u by v , hence

$$\mu_v^{\frac{2n-2}{n}}(t) = a^2(\mu_v(t)) \left(-\mu'_v(t) + \int_{\partial\Omega_t^\sharp \cap \partial\Omega^\sharp} \frac{1}{\beta v(x)} dA \right) \int_0^{\mu_v(t)} (f^\sharp)^*(s) ds$$

for all $t > 0$, proving (3.2). \square

Lemma 3.2. *Suppose u and v are solutions to (1.3) and (1.4). Then both u and v attain their minima on $\partial\Omega$ and $\partial\Omega^\sharp$. Denote by u_0 and v_0 the minima of u and v , then*

$$(3.6) \quad u_0 \leq v_0.$$

Proof. Recall that f is nonnegative, then $-\Delta u \geq 0$ and $-\Delta v \geq 0$. Therefore u and v attain their minima on $\partial\Omega$ and $\partial\Omega^\sharp$. Moreover $u(x) > u_0$ for $x \in \Omega$, $v(x) > v_0$ for $x \in \Omega^\sharp$ unless f is identically zero.

Noticing $v(x) = v_0$ on $\partial\Omega^\sharp$, and we compute by using integration by parts that

$$\begin{aligned}
v_0|\partial\Omega^\sharp|^2 &= \int_{\partial\Omega^\sharp} \frac{1}{\bar{\beta}} dA \int_{\partial\Omega^\sharp} \bar{\beta}v(x) dA \\
&= \int_{\partial\Omega^\sharp} \frac{1}{\bar{\beta}} dA \int_{\Omega^\sharp} -\Delta v dx \\
&= \frac{1}{\theta} \int_{\partial\Omega} \frac{1}{\beta(x)} dA \int_{\Omega^\sharp} f^\sharp(x) dx \\
&= \frac{1}{\theta^2} \int_{\partial\Omega} \frac{1}{\beta(x)} dA \int_{\Omega} -\Delta u(x) dx \\
&= \frac{1}{\theta^2} \int_{\partial\Omega} \frac{1}{\beta(x)} dA \int_{\partial\Omega} \beta(x)u(x) dA,
\end{aligned}$$

where we used (1.3) (1.4), and (1.5) in the above equalities. Recall from Hölder inequality that

$$\left(\int_{\partial\Omega} \sqrt{u(x)} dA \right)^2 \leq \int_{\partial\Omega} \frac{1}{\beta(x)} dA \int_{\partial\Omega} \beta(x)u(x) dA,$$

then we have

$$v_0|\partial\Omega^\sharp|^2 \geq \frac{1}{\theta^2} \left(\int_{\partial\Omega} \sqrt{u(x)} dA \right)^2 \geq \frac{u_0}{\theta^2} |\partial\Omega|^2 \geq u_0 |\partial\Omega^\sharp|^2,$$

which implies (3.6). Where we used isoperimetric inequality (2.7) in the last inequality. \square

Now we turn to prove Theorem 1.1.

Proof of Theorem 1.1. Set $\tilde{\mu}_u(t) = \frac{\mu_u(t)}{\theta}$. Dividing inequality (3.1) by $\tilde{\mu}_u(t)^{\frac{2}{n}-1}$ and then integrating over $[0, \tau]$, yields

$$\begin{aligned}
\theta^2 \int_0^\tau \tilde{\mu}_u(t) dt &\leq \int_0^\tau \left(a^2(\tilde{\mu}_u(t)) \tilde{\mu}_u(t)^{\frac{2}{n}-1} \left(\int_0^{\mu_u(t)} f^*(s) ds \right) (-\mu'_u(t)) \right) dt \\
&\quad + \int_0^\tau a^2(\tilde{\mu}_u(t)) \tilde{\mu}_u(t)^{\frac{2}{n}-1} \int_0^{\mu_u(t)} f^*(s) ds \left(\int_{\partial\Omega_t \cap \partial\Omega} \frac{1}{\beta(x)} \frac{1}{u(x)} dA \right) dt \\
&\leq \theta^2 \int_0^\tau \left(a^2(\tilde{\mu}_u(t)) \tilde{\mu}_u(t)^{\frac{2}{n}-1} \int_0^{\tilde{\mu}_u(t)} (f^\sharp)^*(s) ds \right) (-d\tilde{\mu}_u(t)) \\
&\quad + \theta a^2(|\Omega^\sharp|) |\Omega^\sharp|^{\frac{2}{n}-1} \int_0^{|\Omega^\sharp|} (f^\sharp)^*(s) ds \int_0^\tau \left(\int_{\partial\Omega_t \cap \partial\Omega} \frac{1}{\beta(x)} \frac{1}{u(x)} dA \right) dt,
\end{aligned}$$

where we used

$$\mu_u(t)^{\frac{2}{n}-1} \int_0^{\mu_u(t)} f^*(s) ds \leq |\Omega|^{\frac{2}{n}-1} \int_{\Omega} f(x) dx = |\Omega|^{\frac{2}{n}-1} \int_0^{|\Omega|} f^*(s) ds$$

by assumption (1.6) and the fact $a(s)$ is monotone increasing in s . Using Fubini's theorem, we estimate

$$(3.7) \quad \int_0^\tau \int_{\partial\Omega_t \cap \partial\Omega} \frac{1}{\beta(x)u(x)} dA dt \leq \int_0^\infty \int_{\partial\Omega_t \cap \partial\Omega} \frac{1}{\beta(x)u(x)} dA dt = \int_{\partial\Omega} \frac{1}{\beta(x)} dA,$$

so then we conclude

$$(3.8) \quad \theta^2 \int_0^\tau \tilde{\mu}_u(t) dt \leq -\theta^2 F(\tilde{\mu}_u(\tau)) + \theta a^2(|\Omega^\sharp|)|\Omega^\sharp|^{\frac{2}{n}-1} \int_0^{|\Omega^\sharp|} (f^\sharp)^*(s) ds \int_{\partial\Omega} \frac{1}{\beta(x)} dA,$$

where

$$F(s) := \int_0^s a^2(\sigma)\sigma^{\frac{2}{n}-1} \int_0^\sigma (f^\sharp)^*(r) dr d\sigma.$$

Substituting identity (1.5) to inequality (3.8) yields

$$(3.9) \quad \int_0^\tau \tilde{\mu}_u(t) dt \leq -F(\tilde{\mu}_u(\tau)) + a^2(|\Omega^\sharp|)|\Omega^\sharp|^{\frac{2}{n}-1} \int_0^{|\Omega^\sharp|} (f^\sharp)^*(s) ds \int_{\partial\Omega^\sharp} \frac{1}{\beta} dA.$$

For $\tau > v_0$, we have $\Omega_\tau^\sharp \subset\subset \Omega^\sharp$, and then equality (3.2) becomes to

$$\mu_v^{\frac{2n-2}{n}}(t) = a^2(\mu_v(t)) \left(-\mu'_v(t) \right) \int_0^{\mu_v(t)} (f^\sharp)^*(s) ds,$$

implying

$$(3.10) \quad \int_0^\tau \mu_v(t) dt + F(\mu_v(\tau)) = \int_0^{v_0} \mu_v(t) dt + F(\mu_v(v_0)).$$

Dividing equation (3.2) by $\mu_v(t)^{\frac{2-n}{n}}$ and integrating over $[0, v_0]$ yields

$$\int_0^{v_0} \mu_v(t) dt + F(\mu_v(v_0)) = a^2(|\Omega^\sharp|)|\Omega^\sharp|^{\frac{2}{n}-1} \int_0^{|\Omega^\sharp|} (f^\sharp)^*(s) ds \int_{\partial\Omega^\sharp} \frac{1}{\beta} dA,$$

and substituting above equality to (3.10), we obtain

$$(3.11) \quad \int_0^\tau \mu_v(t) dt = -F(\mu_v(\tau)) + a^2(|\Omega^\sharp|)|\Omega^\sharp|^{\frac{2}{n}-1} \int_0^{|\Omega^\sharp|} (f^\sharp)^*(s) ds \int_{\partial\Omega^\sharp} \frac{1}{\beta} dA$$

for all $\tau > v_0$. Combining inequality (3.9) and equality (3.11), we get

$$(3.12) \quad \int_0^\tau \tilde{\mu}_u(t) dt - \int_0^\tau \mu_v(t) dt \leq -F(\tilde{\mu}_u(t)) + F(\mu_v(t)),$$

thus we get

$$\int_0^\infty \mu_u(t) dt \leq \int_0^\infty \theta \mu_v(t) dt,$$

by letting $\tau \rightarrow \infty$, where we used $\lim_{\tau \rightarrow \infty} \mu_u(\tau) = \lim_{\tau \rightarrow 0} \mu_v(\tau) = 0$. Hence (1.7) holds true. \square

Now we deal with the case $n = 2$ and $f \equiv 1$.

Proof of Theorem 1.3. When $n = 2$ and $f(x) \equiv 1$, inequality (3.1) gives

$$(3.13) \quad \theta \leq \left(-\mu'_u(t) + \int_{\partial\Omega_t \cap \partial\Omega} \frac{1}{\beta(x)u(x)} dA \right) a^2(|\Omega^\sharp|),$$

where we used the fact that $a(s)$ is monotone increasing in s . Integrating inequality (3.13) over $[0, \tau]$ yields

$$(3.14) \quad \begin{aligned} \theta\tau &\leq \left(-\mu_u(\tau) + |\Omega| + \int_0^\tau \int_{\partial\Omega_t \cap \partial\Omega} \frac{1}{\beta(x)u(x)} dA dt \right) a^2(|\Omega^\sharp|) \\ &\leq \left(-\mu_u(\tau) + |\Omega| + \int_{\partial\Omega} \frac{1}{\beta(x)} dA \right) a^2(|\Omega^\sharp|), \end{aligned}$$

where we used (3.7) in the last inequality.

Analogously, by using equality (3.2) we have

$$(3.15) \quad \tau = \left(-\mu_v(\tau) + |\Omega^\sharp| + \int_{\partial\Omega^\sharp} \frac{1}{\beta} dA \right) a^2(|\Omega^\sharp|)$$

for $\tau \geq v_0$, then putting (3.14) and (3.15) together yields

$$\mu_u(\tau) - \theta\mu_v(\tau) \leq 0.$$

For $\tau \leq v_0$, it follows

$$\mu_u(\tau) \leq |\Omega| = \theta|\Omega^\sharp| = \theta\mu_v(\tau),$$

thus

$$\mu_u(\tau) - \theta\mu_v(\tau) \leq 0$$

holds for all $\tau > 0$, hence (1.8) holds.

If (1.8) holds as equality, then inequality (3.14) implies $\Omega_\tau \subset\subset \Omega$, and Ω_τ is a geodesic ball in M_κ for $\tau > v_0$ since isoperimetric inequality (2.7) holds as equality for Ω_τ . Therefore Ω is a geodesic ball in M_κ , $u(x)$ is a radial function, and then $\beta(x)$ is a constant on $\partial\Omega$. \square

REFERENCES

- [1] Virginia Agostiniani, Mattia Fogagnolo, and Lorenzo Mazzieri. Sharp geometric inequalities for closed hypersurfaces in manifolds with nonnegative Ricci curvature. *Invent. Math.*, 222(3):1033–1101, 2020.
- [2] A. Alvino, F. Chiacchio, C. Nitsch, and C. Trombetti. Sharp estimates for solutions to elliptic problems with mixed boundary conditions. *J. Math. Pures Appl. (9)*, 152:251–261, 2021.
- [3] A. Alvino, C. Nitsch, and C. Trombetti. A talenti comparison result for solutions to elliptic problems with robin boundary conditions. *to appear on Comm. Pure Appl. Math.*, *arXiv:1909.11950*, 2019.
- [4] A. Alvino, G. Trombetti, and P.-L. Lions. Comparison results for elliptic and parabolic equations via Schwarz symmetrization. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 7(2):37–65, 1990.
- [5] Angelo Alvino, Vincenzo Ferone, Guido Trombetti, and Pierre-Louis Lions. Convex symmetrization and applications. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 14(2):275–293, 1997.
- [6] Catherine Bandle. On symmetrizations in parabolic equations. *J. Analyse Math.*, 30:98–112, 1976.
- [7] Simon Brendle. Sobolev inequalities in manifolds with nonnegative curvature. *to appear on Comm. Pure Appl. Math.*, *arXiv:2009.13717*, 2020.
- [8] Daguang Chen and Haizhong Li. Talenti’s comparison theorem for poisson equation and applications on riemannian manifold with nonnegative ricci curvature. *arXiv:2104.05568*, 2021.
- [9] Daguang Chen, Haizhong Li, and Yilun Wei. Comparison results for solutions of poisson equations with robin boundary on complete riemannian manifolds. *arXiv:2110.0605*, 2021.
- [10] Francesco Chiacchio, Nunzia Gavitone, Carlo Nitsch, and Cristina Trombetti. Sharp estimates for the gaussian torsional rigidity with robin boundary conditions. *arXiv:2109.10117*, 2021.
- [11] Don Colladay, Jeffrey J. Langford, and Patrick McDonald. Comparison results, exit time moments, and eigenvalues on Riemannian manifolds with a lower Ricci curvature bound. *J. Geom. Anal.*, 28(4):3906–3927, 2018.
- [12] Francesco Della Pietra, Carlo Nitsch, Riccardo Scala, and Cristina Trombetti. An optimization problem in thermal insulation with Robin boundary conditions. *Comm. Partial Differential Equations*, 46(12):2288–2304, 2021.
- [13] Misha Gromov. *Metric structures for Riemannian and non-Riemannian spaces*, volume 152 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1999. Based on the 1981 French original [MR0682063 (85e:53051)], With appendices by M. Katz, P. Pansu and S. Semmes, Translated from the French by Sean Michael Bates.
- [14] A. Hurtado, S. Markvorsen, and V. Palmer. Estimates of the first Dirichlet eigenvalue from exit time moment spectra. *Math. Ann.*, 365(3-4):1603–1632, 2016.
- [15] Bernhard Kawohl. *Rearrangements and convexity of level sets in PDE*, volume 1150 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1985.

- [16] S. Kesavan. Some remarks on a result of Talenti. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 15(3):453–465 (1989), 1988.
- [17] S. Kesavan. *Symmetrization & applications*, volume 3 of *Series in Analysis*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2006.
- [18] Lei Ni and Kui Wang. Isoperimetric comparisons via viscosity. *J. Geom. Anal.*, 26(4):2831–2841, 2016.
- [19] Giorgio Talenti. Elliptic equations and rearrangements. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 3(4):697–718, 1976.
- [20] Giorgio Talenti. Nonlinear elliptic equations, rearrangements of functions and Orlicz spaces. *Ann. Mat. Pura Appl. (4)*, 120:160–184, 1979.

SCHOOL OF MATHEMATICAL SCIENCES, SOOCHOW UNIVERSITY, SUZHOU, 215006, CHINA

Email address: `chq4523@163.com`

SCHOOL OF MATHEMATICAL SCIENCES, SOOCHOW UNIVERSITY, SUZHOU, 215006, CHINA

Email address: `1829401184@stu.suda.edu.cn`

SCHOOL OF MATHEMATICAL SCIENCES, SOOCHOW UNIVERSITY, SUZHOU, 215006, CHINA

Email address: `kuiwang@suda.edu.cn`