

# Commutativity of quantization with conic reduction for torus actions on compact CR manifolds

Andrea Galasso\*

## Abstract

We define conic reduction  $X_\nu^{\text{red}}$  for torus actions on the boundary  $X$  of a strictly pseudo-convex domain and for a given weight  $\nu$  labelling a unitary irreducible representation. There is a natural residual circle action on  $X_\nu^{\text{red}}$ . We have two natural decompositions of the corresponding Hardy spaces  $H(X)$  and  $H(X_\nu^{\text{red}})$ . The first one is given by the ladder of isotypes  $H(X)_{k\nu}$ ,  $k \in \mathbb{Z}$ , the second one is given by the  $k$ -th Fourier components  $H(X_\nu^{\text{red}})_k$  induced by the residual circle action. The aim of this paper is to prove that they are isomorphic. Furthermore, we sketch the proof of a similar result for spaces of  $(0, q)$ -forms with  $L^2$ -coefficient when  $X$  is a CR manifold with non-degenerate Levi-curvature for  $k$  sufficiently large.

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Preliminaries</b>	<b>3</b>
2.1	Geometric setting . . . . .	3
2.2	Hardy spaces . . . . .	6
<b>3</b>	<b>Proof of Theorem 1.1</b>	<b>10</b>
3.1	Ohsawa–Takegoshi extension theorem . . . . .	10
3.2	The proof . . . . .	10
<b>4</b>	<b>Proof of Theorem 1.2</b>	<b>11</b>

**Keywords:** Hardy space, torus action, CR conic reduction

**Mathematics Subject Classification:** 32A2

## 1 Introduction

Let  $X$  be the boundary of a strictly pseudo-convex domain  $D$  in  $\mathbb{C}^{n+1}$ . Then  $(X, T^{1,0}X)$  is a contact manifold of dimension  $2n + 1$ ,  $n \geq 1$ , where  $T^{1,0}X$  is the sub-bundle of  $TX \otimes \mathbb{C}$  defining the CR structure. We denote by  $\omega_0 \in \mathcal{C}^\infty(X, T^*X)$  the contact 1-form whose kernel is the horizontal bundle  $HX \subset TX$ , we refer to Section 2.1 for definitions. Associated with this data we can define the Hardy space  $H(X)$ , it is the space of boundary values of holomorphic functions in  $D$  which lie in  $L^2(X)$ , the Hilbert space of square

---

\***Address:** Room 407, Chee-Chun Leung Cosmology Hall, National Taiwan University; **ORCID iD:** 0000-0002-5792-1674; **e-mail:** andrea.galasso@ncts.ntu.edu.tw

integrable functions on  $X$ . Suppose a contact and CR action of a  $t$ -dimensional torus  $\mathbb{T}^t$  is given; we denote by  $\mu : X \rightarrow \mathfrak{t}^*$  the associated CR moment map, we identify  $\mathfrak{t} \cong \iota\mathbb{R}^t$  with its dual  $\mathfrak{t}$ . Fix a weight  $\iota\nu$  in the lattice  $\iota\mathbb{Z}^t \subset \mathfrak{t}^*$ , if  $0 \in \mathfrak{t}^*$  does not lie in the image of the moment map the isotypes

$$H(X)_{k\nu} = \{f \in H(X) : (e^{\iota\theta} \cdot f)(x) = e^{\iota k\langle\nu,\theta\rangle} f(x), \theta \in \mathbb{R}^t\}, \quad k \in \mathbb{Z},$$

are finite dimensional.

Suppose that the ray  $\iota\mathbb{R}_+ \cdot \nu \in \mathfrak{t}^*$  is transversal to  $\mu$ , then  $X_\nu := \mu^{-1}(\iota\mathbb{R}_+ \cdot \nu)$  is a sub-manifold of  $X$  of codimension  $t - 1$ . There is a well-defined locally free action of  $\mathbb{T}_\nu^{t-1} := \exp_{\mathbb{T}^t}(\iota \ker \nu)$  on  $X_\nu$ , the resulting orbifold  $X_\nu^{\text{red}}$  is called *conic reduction* of  $X$  with respect to the weight  $\nu$ . The locus  $X_\nu$  is  $\mathbb{T}$ -invariant, we will always assume that the action of  $\mathbb{T}$  on  $X_\nu$  is locally free. After replacing  $\mathbb{T}$  with its quotient by a finite subgroup, we may and will assume without loss of generality that the action is generically free. In Section 2.1, we show that  $X_\nu^{\text{red}}$  is a strictly pseudo-convex domain of dimension  $2n - 2t + 3$ . There is a natural locally free circle action  $\mathbb{T}_\nu^1 := \exp_{\mathbb{T}^t}(\iota\nu)$  on  $X_\nu^{\text{red}}$  which induces an action on the Hardy space  $H(X_\nu^{\text{red}})$ . We denote by  $H(X_\nu^{\text{red}})_k$  the corresponding  $k$ -th Fourier component and we call the action of  $\mathbb{T}_\nu^1$  on  $X_\nu^{\text{red}}$  *residual circle action*. The aim of this paper is to prove the following theorem.

**Theorem 1.1.** *Suppose  $X_\nu \neq \emptyset$  and assume  $\mathbb{T}_\nu^1$  defines a transversal locally free CR circle action on  $X$ . There is a natural isomorphism of vector spaces  $\delta_k : H(X)_{k\nu} \rightarrow H(X_\nu^{\text{red}})_k$  for each  $k$ .*

From the point of view of symplectic geometry given a Hamiltonian and holomorphic action with moment map  $\Phi$  of a compact Lie Group  $G$  on a Hodge manifold  $(M, \omega)$  with quantizing circle bundle  $\pi : X \rightarrow M$ , one can always define an infinitesimal action of the Lie algebra  $\mathfrak{g}$  on  $X$ . If it can be integrated to an action of the whole group  $G$ , then one has a representation of  $G$  on  $H(X)$ . In [GS2] it was observed that associated to group actions one can define reductions  $M_{\text{red}}^\Theta$  by pulling back a  $G$ -invariant and proper sub-manifold  $\Theta$  of  $\mathfrak{g}^*$  via the moment map  $\Phi$ . When  $\Theta$  is chosen to be a cone through a co-adjoint orbit  $C(\mathcal{O}_\nu)$  one has associated reduction whose Hardy space of its quantization is

$$H(X_{\text{red}}^{C(\mathcal{O}_\nu)}) = \bigoplus_{k \in \mathbb{Z}} H(X_{\text{red}}^{C(\mathcal{O}_\nu)})_k,$$

where  $k$  labels an irreducible representation of the residual circle action described above. Now, it is natural to ask if this spaces are related with the decomposition induced by  $G$  on  $H(X)$ . When  $G = \mathbb{T}^r$  Theorem 1.1 states that they are isomorphic to  $H_{k\nu}(X)$ ; the canonical decomposition of the Hardy space  $H(X)$  of the quantitation by the built-in circle action does not play any role. The semi-classical parameter  $k$  is the one induced by the ladder  $k\nu$ ,  $k = 0, 1, 2, \dots$ , labelling unitary irreducible representations.

Further geometrical motivations for this theorem are explained in paper [P2], where it is proved that  $\delta_k$  is an isomorphism for  $k$  large enough in the setting when  $X$  is the circle bundle of a polarized Hodge manifold whose Grauert tube is  $D$ ; we also refer to [P1] for examples and the explicit expression of the leading term of the asymptotic expansion of  $\dim H(X)_{k\nu}$  as  $k$  goes to infinity. The proof of Theorem 1.1 is an application of the Ohsawa–Takegoshi extension theorem.

We prove Theorem 1.1 in the more general setting for spaces of  $(0, q)$ -forms when  $k$  is large. In fact, suppose more generally that the Levi form is non-degenerate and the number of either positive or negative eigenvalues of the Levi form is not equal to  $n$ , then the Szegő projector for functions  $S : L^2(X) \rightarrow H(X)$  is a smoothing operator, see

[Hs]. Hence, in the presence of a transversal locally free circle action, the  $k$ -th Fourier component  $S_k$  has a rapidly decreasing asymptotic expansion as  $k$  goes to infinity. From the standpoint of semi-classical analysis it is natural to look for a replacement of these spaces. We then consider  $(0, q)$  forms with  $L^2$  coefficients and the corresponding projector  $S^{(q)}$  onto the kernel of the Köhn Laplacian  $H^q(X)$ .

Let  $X_\nu^{\text{red}} := X_\nu / \mathbb{T}_\nu^{t-1}$ , defined in the same way as before. We will show that  $X_\nu^{\text{red}}$  is a CR manifold with natural CR structure induced by  $T^{1,0}X$  of dimension  $2n - 2(t - 1) + 1$ . Let  $L_{X_\nu^{\text{red}}}$  be the Levi form on  $X_\nu^{\text{red}}$  induced naturally from the Levi form  $L$  on  $X$ . Note that the bilinear form  $b$  defined in Section 2.1 is non-degenerate on  $X_\nu$ . For a given subspace  $\mathfrak{s}$  of  $\mathfrak{t}$ , we denote  $\mathfrak{s}_X$  the subspace of infinitesimal vector fields on  $X$ . Hence, on

$$\ker \nu_X \times \ker \nu_X \subseteq TX_\nu \times TX_\nu,$$

$b$  has constant signature, suppose  $b$  has  $r$  negative and  $t - 1 - r$  positive eigenvalues on  $X_\nu$ . Hence  $L_{X_\nu^{\text{red}}}$  has  $q - r$  negative and  $n - t + 1 - q + r$  positive eigenvalues at each point of  $X_\nu^{\text{red}}$ . We refer to Section 2.1 for definitions, we have:

**Theorem 1.2.** *Assume  $\mathbb{T}_\nu^1$  defines a transversal locally free CR circle action on a compact orientable CR manifold  $X$  with non-degenerate Levi form. Then  $X_\nu^{\text{red}}$  is a compact CR manifold with non-degenerate Levi form having  $q - r$ -negative eigenvalues. There is a natural isomorphism of vector spaces  $\sigma_k : H^q(X)_{k\nu} \rightarrow H^{q-r}(X_\nu^{\text{red}})_k$  for  $k$  sufficiently large.*

We only give a sketch of the proof in Section 4, which can be deduced as in [HH] and it is a consequence of the microlocal properties of the projector  $S_{k\nu}^{(q)}$  described in Section 2.2 and calculus of Fourier integral operators of complex type, see [MS].

## 2 Preliminaries

### 2.1 Geometric setting

We recall some notations concerning CR and contact geometry. Let  $(X, T^{1,0}X)$  be a compact, connected and orientable CR manifold of dimension  $2n + 1$ ,  $n \geq 1$ , where  $T^{1,0}X$  is a CR structure of  $X$ . There is a unique sub-bundle  $HX$  of  $TX$  such that  $HX \otimes \mathbb{C} = T^{1,0}X \oplus T^{0,1}X$ . Let  $J : HX \rightarrow HX$  be the complex structure map given by  $J(u + \bar{u}) = u - i\bar{u}$ , for every  $u \in T^{1,0}X$ . By complex linear extension of  $J$  to  $TX \otimes \mathbb{C}$ , the  $i$ -eigenspace of  $J$  is  $T^{1,0}X$ . We shall also write  $(X, HX, J)$  to denote a CR manifold.

Since  $X$  is orientable, there always exists a real non-vanishing 1-form  $\omega_0 \in C^\infty(X, T^*X)$  so that  $\langle \omega_0(x), u \rangle = 0$ , for every  $u \in H_xX$ , for every  $x \in X$ ;  $\omega_0$  is called contact form and it naturally defines a volume form on  $X$ . For each  $x \in X$ , we define a quadratic form on  $HX$  by

$$L_x(U, V) = \frac{1}{2} d\omega_0(JU, V), \quad \forall U, V \in H_xX.$$

Then, we extend  $L$  to  $HX \otimes \mathbb{C}$  by complex linear extension; for  $U, V \in T_x^{1,0}X$ ,

$$L_x(U, \bar{V}) = \frac{1}{2} d\omega_0(JU, \bar{V}) = -\frac{1}{2i} d\omega_0(U, \bar{V}).$$

The Hermitian quadratic form  $L_x$  on  $T_x^{1,0}X$  is called Levi form at  $x$ . In the case when  $X$  is the circle bundle of an Hodge manifold  $(M, \omega)$ , the positivity of  $\omega$  implies that the number of negative eigenvalues of the Levi form is equal to  $n$ . The Reeb vector field  $R \in C^\infty(X, TX)$  is defined to be the non-vanishing vector field determined by

$$\omega_0(R) = -1, \quad d\omega_0(R, \cdot) \equiv 0 \quad \text{on } TX.$$

Fix a smooth Hermitian metric  $\langle \cdot | \cdot \rangle$  on  $\mathbb{C}TX$  so that  $T^{1,0}X$  is orthogonal to  $T^{0,1}X$ ,  $\langle u | v \rangle$  is real if  $u, v$  are real tangent vectors,  $\langle R | R \rangle = 1$  and  $R$  is orthogonal to  $T^{1,0}X \oplus T^{0,1}X$ . For  $u \in \mathbb{C}TX$ , we write  $|u|^2 := \langle u | u \rangle$ . Denote by  $T^{*1,0}X$  and  $T^{*0,1}X$  the dual bundles of  $T^{1,0}X$  and  $T^{0,1}X$ , respectively. They can be identified with sub-bundles of the complexified cotangent bundle  $\mathbb{C}T^*X$ .

Assume that  $X$  admits an action of  $t$ -dimensional torus  $\mathbb{T}^t$ . In this work, we assume that the  $\mathbb{T}^t$ -action preserves  $\omega_0$  and  $J$ ; that is,  $t^*\omega_0 = \omega_0$  on  $X$  and  $t_*J = Jt_*$  on  $HX$ . Let  $\mathfrak{t}$  denote the Lie algebra of  $T$ , we identify  $\mathfrak{t}$  with its dual  $\mathfrak{t}^*$  by means of the scalar product  $\langle \cdot, \cdot \rangle$ . For any  $\xi \in \mathfrak{t}$ , we write  $\xi_X$  to denote the vector field on  $X$  induced by  $\xi$ . The moment map associated to the form  $\omega_0$  is the map  $\mu : X \rightarrow \mathfrak{t}^*$  such that, for all  $x \in X$  and  $\xi \in \mathfrak{t}$ , we have

$$\langle \mu(x), \xi \rangle = \omega_0(\xi_X(x)). \quad (1)$$

We can also define a non-degenerate bilinear form  $b$  on  $HX$  such that

$$b(\cdot, \cdot) = d\omega_0(\cdot, J\cdot). \quad (2)$$

Fix a maximal weight  $\nu \neq 0$  in the lattice inside  $\mathfrak{t}^*$ . Suppose that  $i\mathbb{R}_+ \cdot \nu$  is transversal to  $\mu$ , so we have

$$\mathfrak{t} = i\mathbb{R}_+ \cdot \nu \oplus d_p\mu(T_pX) \quad (3)$$

and  $X_\nu$  is a sub-manifold of  $X$  of codimension  $2n + 2 - t$ . We claim that the action of  $\mathbb{T}_\nu^{t-1}$  on  $X_\nu$  is locally free. In fact, by the contrary suppose that there exists  $\xi \in \mathfrak{t}$  such that  $\xi_X(x) = 0$  on  $T_xX_\nu$ ,  $x \in X_\nu$ . For each  $v \in T_xX$ , we have

$$(d_x\mu(v))(\xi) = d_x\omega_0(\xi_X, v) = 0$$

which contradicts (3). For simplicity, in the next Lemma we assume the action of  $\mathbb{T}_\nu^{t-1}$  on  $X_\nu$  to be free.

**Lemma 2.1.** *The manifold  $X_\nu/\mathbb{T}_\nu^{t-1}$  is a CR manifold. Let  $\pi : X_\nu \rightarrow X_\nu^{\text{red}}$  and  $\iota : X_\nu \hookrightarrow X$  be the natural quotient and inclusion, respectively, then there is a unique induced contact form  $\omega_0^{\text{red}}$  on  $X_\nu/\mathbb{T}_\nu^{t-1}$  such that*

$$\pi^*\omega_0^{\text{red}} = \iota^*\omega_0.$$

In particular, set  $HX_\nu = TX_\nu \cap HX$ , we have

$$HX = HX_\nu \oplus J\nu_X^\perp \quad \text{and} \quad HX_\nu = \nu_X^\perp \oplus d\pi^*HX_\nu^{\text{red}}.$$

*Proof.* It is an easy consequence of

$$v \in T_xX_\nu \Leftrightarrow d_x\mu(v)(\xi) = 0 \quad \text{for all } \xi \in \nu^\perp, v \in T_xX$$

which proves that

$$\text{val}_x(\nu_X^\perp(x)) = (T_xX_\nu)^{d_x\omega_0}.$$

□

We will also assume that  $\mathbb{T}_\nu^1$ -action is transversal CR, that is, the infinitesimal vector field

$$(\nu_X u)(x) = \frac{\partial}{\partial t} (u(\exp(it\nu) \circ x))|_{t=0}, \quad \text{for any } u \in C^\infty(X),$$

preserves the CR structure  $T^{1,0}X$ , so that  $\nu_X$  and  $T^{1,0}X \oplus T^{0,1}X$  generate the complex tangent bundle to  $X$ ,

$$\mathbb{C}T_xX = \mathbb{C}\nu_X(x) \oplus \mathbb{C}T_x^{1,0}X \oplus \mathbb{C}T_x^{0,1}X \quad (x \in X).$$

We define local coordinates that will be useful later. Recall that  $X$  admits a CR and transversal  $\mathbb{T}_\nu^1$ -action which is locally free on  $X_\nu$ ,  $T \in \mathcal{C}^\infty(X, TX)$  denotes the global real vector field given by this infinitesimal circle action. We will take  $T$  to be our Reeb vector field  $R$ . In a similar way as in Theorem 3.6 in [HH], there exist local coordinates  $v = (v_1, \dots, v_{t-1})$  of  $\mathbb{T}_\nu^{t-1}$  in a small neighborhood  $V_0$  of the identity  $e$  with  $v(e) = (0, \dots, 0)$ , local coordinates  $x = (x_1, \dots, x_{2n+1})$  defined in a neighborhood  $U_1 \times U_2$  of  $p \in X_\nu$ , where  $U_1 \subseteq \mathbb{R}^{t-1}$  (resp.  $U_2 \subseteq \mathbb{R}^{2n+2-t}$ ) is an open set of  $0 \in \mathbb{R}^{t-1}$  (resp.  $0 \in \mathbb{R}^{2n+2-t}$ ) and  $p \equiv 0 \in \mathbb{R}^{2n+1}$ , and a smooth function  $\gamma = (\gamma_1, \dots, \gamma_{t-1}) \in \mathcal{C}^\infty(U_2, U_1)$  with  $\gamma(0) = 0$  such that

$$\begin{aligned} & (v_1, \dots, v_{t-1}) \circ (\gamma(x_t, \dots, x_{2n+1}), x_t, \dots, x_{2n+1}) \\ &= (v_1 + \gamma_1(x_t, \dots, x_{2n+1}), \dots, v_{t-1} + \gamma_{t-1}(x_t, \dots, x_{2n+1}), x_t, \dots, x_{2n+1}) \end{aligned}$$

for each  $(v_1, \dots, v_{t-1}) \in V_0$  and  $(x_t, \dots, x_{2n+1}) \in U_2$ . Furthermore, we have

$$\mathfrak{t} = \text{span} \left\{ \partial_{x_j} \right\}_{j=1, \dots, t-1}, \quad \mu^{-1}(\iota \mathbb{R}_\nu \cdot \nu) \cap U = \{x_{2d-t+1} = \dots = x_{2d} = 0\},$$

on  $\mu^{-1}(\iota \mathbb{R}_\nu \cdot \nu) \cap U$  there exist smooth functions  $a_j$ 's with  $a_j(0) = 0$  for every  $0 \leq j \leq t-1$  and independent on  $x_1, \dots, x_{2(t-1)}, x_{2n+1}$  such that

$$J(\partial_{x_j}) = \partial_{x_{t-1+j}} + a_j(x) \partial_{x_{2n+1}} \quad j = 1, \dots, t-1,$$

the Levi form  $L_p$ , the Hermitian metric  $\langle \cdot | \cdot \rangle$  and the 1-form  $\omega_0$  can be written

$$L_p(Z_j, \bar{Z}_k) = \mu_j \delta_{j,k}, \quad \langle Z_j | \bar{Z}_k \rangle = \delta_{j,k} \quad (1 \leq j, k \leq n),$$

and

$$\begin{aligned} \omega_0(x) = & (1 + O(|x|)) dx_{2n+1} + \sum_{j=1}^{t-1} 4\mu_j x_{t-1+j} dx_j + \sum_{j=t}^n 2\mu_j x_{2j} dx_{2j-1} \\ & - \sum_{j=t}^n 2\mu_j x_{2j-1} dx_{2j} + \sum_{j=r}^{2n} b_j x_{2n+1} dx_j + O(|x|^2) \end{aligned}$$

where  $b_r, \dots, b_{2n} \in \mathbb{R}$ ,

$$T_p^{1,0} X = \text{span}\{Z_1, \dots, Z_n\}$$

and

$$\begin{aligned} Z_j &= \frac{1}{2} (\partial_{x_j} - \iota \partial_{x_{t-1+j}})(p) \quad (j = 1, \dots, t-1), \\ Z_j &= \frac{1}{2} (\partial_{x_{2j-1}} - \iota \partial_{x_{2j}})(p) \quad (j = t, \dots, n). \end{aligned}$$

We need to define in local coordinates we just introduced the phase function of the  $\mathbb{T}_\nu^{t-1}$ -invariant Szegő kernel  $\Phi_-(x, y) \in \mathcal{C}^\infty(U \times U)$  which is independent of  $(x_1, \dots, x_{t-1})$  and  $(y_1, \dots, y_{t-1})$ . Hence, we write  $\Phi_-(x, y) = \Phi_-((0, x''), (0, y'')) := \Phi_-(x'', y'')$ . Moreover, there is a constant  $c > 0$  such that

$$\text{Im} \Phi_-(x'', y'') \geq c (|\hat{x}''|^2 + |\hat{y}''|^2 + |\hat{x}'' - \hat{y}''|^2), \quad \text{for all } ((0, x''), (0, y'')) \in U \times U. \quad (4)$$

Furthermore,

$$\begin{aligned}
\Phi_-(x'', y'') &= -x_{2n+1} + y_{2n+1} + 2i \sum_{j=1}^{t-1} |\mu_j| y_{t-1+j}^2 + 2i \sum_{j=1}^{t-1} |\mu_j| x_{t-1+j}^2 + i \sum_{j=t}^n |\mu_j| |z_j - w_j|^2 \\
&+ \sum_{j=t}^n i \mu_j (\bar{z}_j w_j - z_j \bar{w}_j) + \sum_{j=1}^d (-b_{t-1+j} x_{d+j} x_{2n+1} + b_{t-1+j} y_{t-1+j} y_{2n+1}) \\
&+ \sum_{j=t}^n \frac{1}{2} (b_{2j-1} - i b_{2j}) (-z_j x_{2n+1} + w_j y_{2n+1}) + \sum_{j=t}^n \frac{1}{2} (b_{2j-1} + i b_{2j}) (-\bar{z}_j x_{2n+1} + \bar{w}_j y_{2n+1}) \\
&+ (x_{2n+1} - y_{2n+1}) f(x, y) + O(|(x, y)|^3),
\end{aligned} \tag{5}$$

where  $z_j = x_{2j-1} + i x_{2j}$ ,  $w_j = y_{2j-1} + i y_{2j}$ ,  $j = t, \dots, n$ ,  $\mu_j$ ,  $j = 1, \dots, n$ ,  $f$  is smooth and satisfies  $f(0, 0) = 0$ ,  $f(x, y) = \bar{f}(y, x)$ .

We now consider  $\mathbb{T}_\nu^1$  circle action on  $X$ . Let  $p \in \mu^{-1}(i\mathbb{R}_+ \cdot \nu)$ , there exist local coordinates  $v = (v_1, \dots, v_{t-1})$  of  $\mathbb{T}^{t-1}$  in a small neighborhood  $V_0$  of  $e$  with  $v(e) = (0, \dots, 0)$ , local coordinates  $x = (x_1, \dots, x_{2n+1})$  defined in a neighborhood  $U_1 \times U_2$  of  $p$ , where  $U_1 \subseteq \mathbb{R}^{t-1}$  (resp.  $U_{t-1} \subseteq \mathbb{R}^{2n+t}$ ) is an open set of  $0 \in \mathbb{R}^{t-1}$  (resp.  $0 \in \mathbb{R}^{2n+t}$ ) and  $p \equiv 0 \in \mathbb{R}^{2n+1}$ , and a smooth function  $\gamma = (\gamma_1, \dots, \gamma_t) \in \mathcal{C}^\infty(U_2, U_1)$  with  $\gamma(0) = 0$  such that  $T = -\frac{\partial}{\partial x_{2n+1}}$  and all the properties for the local coordinates defined before hold. The phase function  $\Psi$  satisfies  $\Psi(x, y) = -x_{2n+1} + y_{2n+1} + \hat{\Psi}(\mathring{x}'', \mathring{y}'')$ , where  $\hat{\Psi}(\mathring{x}'', \mathring{y}'') \in \mathcal{C}^\infty(U \times U)$ ,  $\mathring{x}'' = (x_t, \dots, x_{2n})$ ,  $\mathring{y}'' = (y_t, \dots, y_{2n})$ , and  $\Psi$  satisfies (5).

## 2.2 Hardy spaces

We denote by  $L_{(0,q)}^2(X)$  the completion of  $\Omega^{0,q}(X)$  with respect to  $(\cdot | \cdot)$ . We extend  $(\cdot | \cdot)$  to  $L_{(0,q)}^2(X)$  in the standard way. We extend  $\bar{\partial}_b$  to  $L_{(0,r)}^2(X)$ ,  $r = 0, 1, \dots, n$ , by

$$\bar{\partial}_b : \text{Dom } \bar{\partial}_b \subset L_{(0,r)}^2(X) \rightarrow L_{(0,r+1)}^2(X),$$

where  $\text{Dom } \bar{\partial}_b := \{u \in L_{(0,r)}^2(X); \bar{\partial}_b u \in L_{(0,r+1)}^2(X)\}$  and, for any  $u \in L_{(0,r)}^2(X)$ ,  $\bar{\partial}_b u$  is defined in the sense of distributions. We also write

$$\bar{\partial}_b^* : \text{Dom } \bar{\partial}_b^* \subset L_{(0,r+1)}^2(X) \rightarrow L_{(0,r)}^2(X)$$

to denote the Hilbert space adjoint of  $\bar{\partial}_b$  in the  $L^2$  space with respect to  $(\cdot | \cdot)$ . There is a well-defined orthogonal projection

$$S^{(q)} : L_{(0,q)}^2(X) \rightarrow \text{Ker } \square_b^q \tag{6}$$

with respect to the  $L^2$  inner product  $(\cdot | \cdot)$  and let

$$S^{(q)}(x, y) \in \mathcal{D}'(X \times X, T^{*0,q} X \boxtimes (T^{*0,q} X)^*)$$

denote the distribution kernel of  $S^{(q)}$ . We will write  $H^q(X)$  for  $\text{Ker } \square_b^q$  and for functions we simply write  $H(X)$  to mean  $H^0(X)$ . The distributional kernel  $S^{(q)}(x, y)$  was studied in [Hs], before recalling an explicit description describing the oscillatory integral defining the distribution  $S^{(q)}(x, y)$  we need to fix some notation. Firstly, we recall Hörmander symbol space. Let  $D \subset X$  be a local coordinate patch with local coordinates  $x = (x_1, \dots, x_{2n+1})$ .

**Definition 2.1.** For every  $m \in \mathbb{R}$ , we denote with

$$S_{1,0}^m(D \times D \times \mathbb{R}_+, T^{*0,q}X \boxtimes (T^{*0,q}X)^*) \subseteq \mathcal{C}^\infty(D \times D \times \mathbb{R}_+, T^{*0,q}X \boxtimes (T^{*0,q}X)^*) \quad (7)$$

the space of all  $a$  such that, for all compact  $K \Subset D \times D$  and all  $\alpha, \beta \in \mathbb{N}_0^{2n+1}$ ,  $\gamma \in \mathbb{N}_0$ , there is a constant  $C_{\alpha,\beta,\gamma} > 0$  such that

$$|\partial_x^\alpha \partial_y^\beta \partial_t^\gamma a(x, y, t)| \leq C_{\alpha,\beta,\gamma} (1 + |t|)^{m-\gamma}, \quad \text{for every } (x, y, t) \in K \times \mathbb{R}_+, t \geq 1.$$

For simplicity we denote with  $S_{1,0}^m$  the spaces defined in (7); furthermore, we write

$$S^{-\infty}(D \times D \times \mathbb{R}_+, T^{*0,q}X \boxtimes (T^{*0,q}X)^*) := \bigcap_{m \in \mathbb{R}} S_{1,0}^m(D \times D \times \mathbb{R}_+, T^{*0,q}X \boxtimes (T^{*0,q}X)^*).$$

Let  $a_j \in S_{1,0}^{m_j}$ ,  $j = 0, 1, 2, \dots$  with  $m_j \rightarrow -\infty$ , as  $j \rightarrow \infty$ . Then there exists  $a \in S_{1,0}^{m_0}$  unique modulo  $S^{-\infty}$ , such that

$$a - \sum_{j=0}^{k-1} a_j \in S_{1,0}^{m_k}(D \times D \times \mathbb{R}_+, T^{*0,q}X \boxtimes (T^{*0,q}X)^*)$$

for  $k = 0, 1, 2, \dots$ . If  $a$  and  $a_j$  have the properties above, we write  $a \sim \sum_{j=0}^{\infty} a_j$  in  $S_{1,0}^{m_0}$ .

It is known that the characteristic set of  $\square_b^q$  is given by

$$\Sigma = \Sigma^- \cup \Sigma^+, \quad \Sigma^- = \{(x, \lambda \omega_0(x)) \in T^*X; \lambda < 0\},$$

and  $\Sigma^+$  is defined similarly for  $\lambda > 0$ . We recall the following theorem (see [Hs, Theorem 1.2]).

**Theorem 2.1.** *Since the Levi form is non-degenerate,  $\square_b^q$  has  $L^2$  closed range. Then, there exist continuous operators  $S_-, S_+ : L_{(0,q)}^2(X) \rightarrow \text{Ker } \square_b^q$  such that*

$$S^{(q)} = S_- + S_+, \quad S_+ \equiv 0 \quad \text{if } q \neq n_+$$

and

$$\text{WF}'(S_-) = \text{diag}(\Sigma^- \times \Sigma^-), \quad \text{WF}'(S_+) = \text{diag}(\Sigma^+ \times \Sigma^+) \quad \text{if } q = n_- = n_+,$$

where  $\text{WF}'(S_-) = \{(x, \xi, y, \eta) \in T^*X \times T^*X; (x, \xi, y, -\eta) \in \text{WF}(S_-)\}$ ,  $\text{WF}(S_-)$  is the wave front set of  $S_-$  in the sense of Hörmander.

Moreover, consider any small local coordinate patch  $D \subset X$  with local coordinates  $x = (x_1, \dots, x_{2n+1})$ , then  $S_-(x, y)$ ,  $S_+(x, y)$  satisfy

$$S_{\mp}(x, y) \equiv \int_0^\infty e^{i\varphi_{\mp}(x,y)t} s_{\mp}(x, y, t) dt \quad \text{on } D,$$

with

$$s_{\mp}(x, y, t) \sim \sum_{j=0}^{\infty} s_{\mp}^j(x, y) t^{n-j} \quad \text{in } S_{1,0}^n(D \times D \times \mathbb{R}_+, T^{*0,q}X \boxtimes (T^{*0,q}X)^*),$$

$s_+(x, y, t) = 0$  if  $q \neq n_+$  and  $s_-^0(x, x) \neq 0$  for all  $x \in D$ . The phase functions  $\varphi_-, \varphi_+$  satisfy

$$\varphi_+, \varphi_- \in \mathcal{C}^\infty(D \times D), \quad \text{Im } \varphi_{\mp}(x, y) \geq 0, \quad \varphi_-(x, x) = 0, \quad \varphi_-(x, y) \neq 0 \quad \text{if } x \neq y,$$

and

$$d_x \varphi_-(x, y)|_{x=y} = -\omega_0(x), \quad d_y \varphi_-(x, y)|_{x=y} = \omega_0(x), \quad -\bar{\varphi}_+(x, y) = \varphi_-(x, y).$$

**Remark 2.1.** Kohn [Kohn] proved that if  $q = n_- = n_+$  or  $|n_- - n_+| > 1$  then  $\square_b^q$  has  $L^2$  closed range. For a description of the phase function in local coordinates see chapter 8 of part I in [Hs].

Now we focus on the decomposition induced by the group action on  $H^q(X)$ . Fix  $t \in \mathbb{T}^r$  and let  $t^* : \Lambda_x^r(\mathbb{C}T^*X) \rightarrow \Lambda_{t^{-1} \circ x}^r(\mathbb{C}T^*X)$  be the pull-back map. Since  $\mathbb{T}^r$  preserves  $J$ , we have  $t^* : T_x^{*0,q}X \rightarrow T_{g^{-1} \circ x}^{*0,q}X$ , for all  $x \in X$ . Thus, for  $u \in \Omega^{0,q}(X)$ , we have  $t^*u \in \Omega^{0,q}(X)$ . Put

$$\Omega^{0,q}(X)_{k\nu} := \{u \in \Omega^{0,q}(X); (e^{i\theta})^*u = e^{ik\langle \nu, \theta \rangle}u, \quad \forall \theta \in \mathbb{R}^r\}.$$

Since the Hermitian metric  $\langle \cdot | \cdot \rangle$  on  $\mathbb{C}TX$  is  $\mathbb{T}^r$ -invariant, the  $L^2$  inner product  $(\cdot | \cdot)$  on  $\Omega^{0,q}(X)$  induced by  $\langle \cdot | \cdot \rangle$  is  $\mathbb{T}^r$ -invariant. Let  $u \in L_{(0,q)}^2(X)$  and  $t \in \mathbb{T}^r$ , we can also define  $t^*u$  in the standard way. We introduce the following notation

$$L_{(0,q)}^2(X)_{k\nu} := \{u \in L_{(0,q)}^2(X); (e^{i\theta})^*u = e^{ik\langle \nu, \theta \rangle}u, \quad \forall \theta \in \mathbb{R}^r\},$$

and put

$$(\text{Ker } \square_b^q)_{k\nu} := \text{Ker } \square_b^q \cap L_{(0,q)}^2(X)_{k\nu}.$$

The equivariant Szegő projection is the orthogonal projection

$$S_{k\nu}^{(q)} : L_{(0,q)}^2(X) \rightarrow (\text{Ker } \square_b^q)_{k\nu}$$

with respect to  $(\cdot | \cdot)$ . Let  $S_{k\nu}^{(q)}(x, y) \in \mathcal{D}'(X \times X, T^{*0,q}X \boxtimes (T^{*0,q}X)^*)$  be the distribution kernel of  $S_{k\nu}^{(q)}$ . The asymptotic expansion for the distributional kernel of the projector  $S_{k\nu}^{(0)}$  was studied in [P1] when  $X$  is the quantizing circle bundle of a given Hodge manifold in Heisenberg local coordinates. Using similar ideas as in [HH] one can generalize the results for  $(0, q)$ -forms in the following theorem, we give a sketch of the proof.

**Theorem 2.2.** *Suppose that  $\square_b^q$  has  $L^2$  closed range. Then, there exist continuous operators  $S_{k\nu}^-, S_{k\nu}^+ : L_{(0,q)}^2(X) \rightarrow (\text{Ker } \square_b^q)_{k\nu}$  such that*

$$S_{k\nu}^{(q)} = S_{k\nu}^- + S_{k\nu}^+, \quad S_+ \equiv 0 \quad \text{if } q \neq n_+$$

*Suppose for simplicity that  $q \neq n_+$ . If  $q \neq n_-$ , then  $S_{k\nu}^{(q)} \equiv O(k^{-\infty})$  on  $X$ .*

*Suppose  $q = n_-$  and let  $D$  be an open set in  $X$  such that the intersection  $\mu^{-1}(i\mathbb{R}_+ \cdot \nu) \cap D = \emptyset$ . Then  $S_{k\nu}^{(q)} \equiv O(k^{-\infty})$  on  $D$ .*

*Let  $p \in \mu^{-1}(i\mathbb{R}_+ \cdot \nu)$  and let  $U$  a local neighborhood of  $p$  with local coordinates  $(x_1, \dots, x_{2n+1})$ . Then, if  $q = n_-$ , for every fix  $y \in U$ , we consider  $S_{k\nu}^{(q)}(x, y)$  as a  $k$ -dependent smooth function in  $x$ , then*

$$S_{k\nu}^{(q)}(x, y) = e^{ik\|\nu\|\Psi(x,y)} b(x, y, k\|\nu\|) + O(k^{-\infty}).$$

*for every  $x \in U$ . The phase function  $\Psi(x, y)$  is defined in local coordinates in the end of Section 2.1, the symbol satisfies*

$$b(x, y, k\|\nu\|) \in S_{\text{loc}}^{n+(1-t)/2}(1, U \times U, T^{*(0,q)}X \boxtimes (T^{*(0,q)}X)^*)$$

*and the leading term of  $b(x, x, \|\nu\|)$  is non-zero.*

*Proof.* Suppose  $q = n_-$ , on small local neighbourhood  $D$  of a point  $p \in X_\nu$  we have

$$S_{k\nu}^{(q)}(x, y) = \frac{1}{(2\pi)^t} \int_{-\pi}^{\pi} \dots \int_{-\pi}^{\pi} e^{ik\langle \nu, \theta \rangle} S^{(q)}(x, e^{i\theta} \cdot y) d\theta$$



where  $\theta = (\theta_1, \dots, \theta_t)$ . We set  $\theta_t$  the variable for circle action of  $\mathbb{T}_\nu^1$  and  $(\theta_1, \dots, \theta_{t-1})$  the variables for  $\mathbb{T}_\nu^{t-1}$ . We can then use local coordinates defined in 2.1, we have

$$S_{k\nu}^{(q)}(x, y) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ik\|\nu\|\theta_t - i kx_{2n+1} + i ky_{2n+1}} S_{\mathbb{T}^{t-1}}^{(q)}(\hat{x}, e^{i\theta_t} \cdot \hat{y}) d\theta_t$$

where

$$S_{\mathbb{T}^{t-1}}^{(q)}(x, y) = \frac{1}{(2\pi)^{t-1}} \int_{-\pi}^{\pi} \cdots \int_{-\pi}^{\pi} S^{(q)}(x, e^{i \text{diag}(\theta_1, \dots, \theta_{t-1})} \cdot y) d\theta_1 \cdots d\theta_{t-1}, \quad (8)$$

and  $\hat{x} = (x_1, \dots, x_n, 0)$  and  $\hat{y} = (y_1, \dots, y_n, 0)$ . The action of  $\mathbb{T}^t$  restricts to an action of  $\mathbb{T}^{t-1}$  whose moment map is given by

$$\mu|_{\mathbb{T}^{r-1}} = p_\nu \circ \mu, \quad \text{where } p_\nu : \mathfrak{t}^* \rightarrow \mathfrak{t}_\nu^*$$

is the canonical projection onto the  $t-1$ -dimensional subspace  $\nu^\perp$  in  $\mathfrak{t}^*$ . Since we are assuming  $X_\nu \neq \emptyset$  we have that  $0 \in \mathfrak{t}_{t-1}^*$  does lie in the image of the moment map  $\mu|_{\mathbb{T}^{r-1}}$ . We can apply Theorem 1.5 in [HH] for the action of  $\mathbb{T}^{r-1}$  with moment map  $\mu|_{\mathbb{T}^{r-1}}$  to the  $\mathbb{T}^{r-1}$ -invariant Szegő kernel in (8). The  $k\nu$  Fourier components are the one induced by the transversal  $\mathbb{T}_\nu^1$ -action and can be computed in a similar way as in Theorem 1.8 of [HH]. Thus the proof can be deduced by applying Theorem 1.5 and 1.8 in [HH].  $\square$

For ease of notation we introduce a definition to say that an operator behaves microlocally as equivariant Szegő projector we just studied. For simplicity we assume  $q = n_-$ .

**Definition 2.2** (Equivariant Szegő type operator). Suppose that  $q = n_-$  and consider  $H : \Omega^{0,q}(X) \rightarrow \Omega^{0,q}(X)$  be a continuous operator with distribution kernel

$$H(x, y) \in \mathcal{D}'(X \times X, T^{*0,q}X \boxtimes (T^{*0,q}X)^*).$$

We say that  $H$  is a complex Fourier integral operator of equivariant Szegő type of order  $n + (1-t)/2 \in \mathbb{Z}$  if  $H$  is smoothing away  $\mu^{-1}(i\mathbb{R}_+ \cdot \nu)$  and for given  $p \in \mu^{-1}(i\mathbb{R}_+ \cdot \nu)$  let  $D$  a local neighborhood of  $p$  with local coordinates  $(x_1, \dots, x_{2n+1})$ . Then, the distributional kernel of  $H$  satisfies

$$H_k(x, y) = e^{ik\Psi(x, y)} a(x, y, k) + O(k^{-\infty}) \quad \text{on } D$$

where  $a \in S_{1,0}^{n+(1-t)/2}(D \times D \times \mathbb{R}_+, T^{*0,q}X \boxtimes (T^{*0,q}X)^*)$ , and  $\Psi$  is as in the end of Section 2.1.

For  $k = 0$ ,  $H_0^{(q)}(X)$  is the space of  $\mathbb{T}$ -fixed vectors, by Theorem 1.5 in [HH] we have  $\dim H_0^{(q)}(X) < +\infty$  since when  $0 \notin \mu(X)$  the projector  $S_{\mathbb{T}^t}^{(q)}$  is smoothing. Now, given  $k_1, k_2 \in \mathbb{Z}$  with  $k_1, k_2 \neq 0$ , consider  $f_1 \in H_{k_1\nu}^{(q)}(X)$  and  $f_2 \in H_{k_2\nu}^{(q)}(X)$ , we have  $f_1 \cdot f_2 \in H_{(k_1+k_2)\nu}^{(q)}(X)$ ; we can study  $\dim H_{k\nu}^{(q)}(X)$  for  $k$  large, we have

$$\dim H_{k\nu}^{(q)}(X) = \int_X S_{k\nu}^{(q)}(x, x) dV_X(x).$$

These dimensions can be studied as  $k \rightarrow +\infty$  by using the microlocal properties of the Szegő kernel and Stationary Phase Lemma in a similar way as in Corollary 1.3 in [P1] we have  $\dim H_{k\nu}^{(q)}(X) = O(k^{d+1-t})$ . So, we have the following generalization of Theorem A.3 in [GS1], where it is proved for spaces of functions  $H(X)_{k\nu}$ .

**Lemma 2.2.** *If  $0 \notin \mu(X)$ , then  $H_{k\nu}^q(X)$  are finite dimensional.*

### 3 Proof of Theorem 1.1

#### 3.1 Ohsawa–Takegoshi extension theorem

We recall the Ohsawa–Takegoshi extension theorem from [GZ] in the form we need in this paper. Before stating the theorem we need some notation. Since the domain  $D$  is strictly pseudo-convex and the action of  $\mathbb{T}_\nu^1$  is transversal for  $(X, T^{1,0}X)$  we get a complex manifold  $N := X/\mathbb{T}_\nu^1$  and an ample line bundle  $(B, h)$  over it such that the curvature of the unique compatible and holomorphic connection is  $-2\nu\eta_\nu$ , where  $d\omega_0 = 2\pi^*\eta_\nu$  and  $\pi : X \rightarrow N$ ,  $X \subset B^*$ . In the situation we are considering here the transversal circle action is the one induced by restriction of the action of  $\mathbb{T}^r$  to  $\mathbb{T}_\nu^1$ . Notice that when  $X$  is the boundary of the Grauert tube of a given Hodge manifold  $M$ , one also has a built-in circle action; we will not discuss this situation and we instead refer to [P1] and [P2].

The sub-manifold  $X_\nu \subset X$  defines a circle bundle over a real sub-manifold  $\iota_N : N_\nu \hookrightarrow N$ , which is co-isotropic with respect to  $\eta_\nu$ . Then the space of leaves  $N_\nu^{\text{red}}$  of the characteristic foliation of  $\iota_N^*\eta_\nu$  inherits a unique symplectic form  $\eta_\nu^{\text{red}}$  such that

$$\pi_N^*\omega_0^{\text{red}} = \iota_N^*\omega_0,$$

where  $\pi_N : N_\nu \rightarrow N_\nu^{\text{red}}$  is the natural projection.

**Theorem 3.1** ([GZ]). *Let  $N$  be a complex manifold of dimension  $n$  and let  $N_\nu^{\text{red}}$  be a complex sub-manifold of codimension  $r - 1$ . Let  $h$  be a smooth metric on the holomorphic ample line bundle  $B$ . Then, there exists a uniform constant  $C$ , such that, for any holomorphic section  $s$  of  $B|_{N_\nu^{\text{red}}}$  on  $N_\nu^{\text{red}}$ , there exists a holomorphic section  $S$  of  $B$  on  $N$  satisfying  $S = s$  on  $N_\nu^{\text{red}}$  and*

$$\int_N |S|_h^2 dV_N \leq C \int_{N_\nu^{\text{red}}} |f|_h^2 dV_{N_\nu^{\text{red}}}.$$

#### 3.2 The proof

Consider the restriction to  $X_\nu$  of a function  $f \in H(X)_{k\nu}$ . Then, it is  $\mathbb{T}_\nu^{r-1}$ -invariant on  $X_\nu$ ; in fact for each  $\theta \in \nu^\perp$  we have

$$e^{i\theta} \cdot f = t^{ik\langle \nu, \theta \rangle} f = f;$$

thus, there is a natural injection

$$\delta_k : H(X)_{k\nu} \hookrightarrow H(X_\nu^{\text{red}}).$$

Furthermore, notice that the map  $\delta_k$  induces a map onto  $H_k(X_\nu^{\text{red}})$  by composing with

$$F_k : L^2(X_\nu^{\text{red}}) \rightarrow L_k^2(X_\nu^{\text{red}})$$

which takes the  $k$ -th component of  $L^2(X_\nu^{\text{red}})$  for the  $\mathbb{T}_\nu^1$ -action, since it is CR transversal.

Now, given  $f \in H_k(X_\nu^{\text{red}})$ , we recall that  $X_\nu^{\text{red}}$  is the total space of the circle bundle over  $N_\nu^{\text{red}}$ , thus  $f$  is a holomorphic section of the  $k$ -th tensor power of a line bundle over  $N_\nu^{\text{red}}$  by the usual identification. The complex manifold  $N_\nu^{\text{red}}$  lies naturally in  $N := X/\mathbb{T}_\nu^1$ , it is a complex sub-orbifold of complex codimension  $r - 1$ . By the Ohsawa–Takegoshi extension theorem there is a global holomorphic section  $F$  in  $H_{k\nu}(X)$  whose restriction to  $X_\nu^{\text{red}}$  equals  $f$ . The Theorem is thus concluded.

## 4 Proof of Theorem 1.2

In this section we shall explain how to prove asymptotic commutativity for quantization and reduction for spaces of  $(0, q)$ -forms. We recall that  $X_\nu^{\text{red}}$  is a CR manifold whose Levi form is non-degenerate, with  $r$  negative eigenvalues; let us denote with  $S^{(q)}$  the corresponding Szegő kernel  $S_{\text{red}}^{(q)}$  for  $(0, q)$ -forms whose  $k$ -th Fourier components are the one induced by the  $\mathbb{T}_\nu^1$ -action on  $X_\nu^{\text{red}}$ . Since  $\nu_X^\perp := \text{val}(\nu^\perp)$  is orthogonal to  $H_x X_\nu \cap J_x H_x X_\nu$  and  $H_x X_\nu \cap J_x H_x X_\nu \subset (\nu_X^\perp)^{\perp b}$  for every  $x \in X_\nu$ , we can find a  $\mathbb{T}$ -invariant orthonormal basis  $\{Z_1, \dots, Z_n\}$  of  $T^{*0,1}X$  on  $X_\nu$  such that for each  $j, k = 1, \dots, n$ ,

$$L_x(Z_j(x), \overline{Z}_j(x)) = \delta_{jk} \lambda_j(x),$$

where

$$\begin{aligned} Z_j(x) &\in (\nu_{X,x}^\perp + \iota J \nu_{X,x}^\perp) \text{ for each } j = 1, \dots, r-1, \\ Z_j(x) &\in H_x X_\nu \cap J_x(H_x X_\nu) \text{ for each } j = r, \dots, n. \end{aligned}$$

Let  $\{Z_1^*, \dots, Z_n^*\}$  denote the orthonormal basis of  $T^{*0,1}X$  on  $X_\nu$ , dual to  $Z_1, \dots, Z_n$ . Fix  $s = 0, 1, 2, \dots, n-r+1$ . For  $x \in X_\nu$ , put

$$B_x^{*0,s} X = \left\{ \sum_{r \leq j_1 < \dots < j_s \leq n} a_{j_1, \dots, j_s} Z_{j_1}^*, \dots, Z_{j_s}^*; a_{j_1, \dots, j_s} \in \mathbb{C}, \right\}$$

and let  $B^{*0,s} X$  be the vector bundle of  $X_\nu$  with fibre  $B_x^{*0,s}$ ,  $x \in X_\nu$ . Let  $C^\infty(X_\nu, B^{*0,s} X)^\mathbb{T}$  denote the set of all  $\mathbb{T}$ -invariant sections of  $X_\nu$  with values in  $B_x^{*0,s} X$ ; let

$$\iota_{\text{red}} : C^\infty(X_\nu, B^{*0,s} X)^\mathbb{T} \rightarrow \Omega^{0,s}(X_\nu^{\text{red}})$$

be the natural identification. Before defining the map between  $H^q(X)_{k\nu}$  and  $H^{q-r}(X_\nu^{\text{red}})_k$  we need one more piece of notation. We can assume that  $\lambda_1 < 0, \dots, \lambda_r < 0$  and also  $\lambda_t < 0, \dots, \lambda_{n-r+t-1} < 0$ . For  $x \in X_\nu$ , set

$$\hat{\mathcal{N}}(x, n_-) = \{c Z_t^* \wedge \dots \wedge Z_{n_- - r + t - 1}^*, c \in \mathbb{C}\}$$

and let  $\hat{p} : \mathcal{N}(x, n_-) \rightarrow \hat{\mathcal{N}}(x, n_-)$  to be

$$p(c Z_1^* \wedge \dots \wedge Z_r^* \wedge Z_t^* \wedge \dots \wedge Z_{n_- - r + t - 1}^*) := c Z_t^* \wedge \dots \wedge Z_{n_- - r + t - 1}^*$$

where  $c$  is a complex number. Put  $\iota_\nu : X_\nu \rightarrow X$  be the natural inclusion, let  $\iota_\nu^* : \Omega^{0,q}(X) \rightarrow \Omega^{0,q}(X_\nu)$  be the pull-back of  $\iota_\nu$ .

Let  $q = n_-$ ; now, inspired by [HH], we define the map

$$\sigma_{k\nu} : H_{k\nu}^{(q)}(X) \rightarrow H(X_\nu^{\text{red}})_k$$

given by

$$\sigma_{k\nu}(u) := k^{(t-1)/4} S_{\text{red},k}^{(q-r)} \circ \iota_{\text{red}} \circ \hat{p} \circ \tau_{x,n_-} \circ e \circ \iota_\nu^* \circ S_{k\nu}^{(q)}(u), \quad (9)$$

here,  $e(x)$  is a  $\mathbb{T}$ -invariant smooth function on  $X_\nu$  that can be found explicitly. We recall the idea of the proof which then follows from calculation of complex Fourier integral operators. Notice that operators  $S_{\text{red},k}^{(q-r)}$  and  $S_{k\nu}^{(q)}$  appearing in (9) are known explicitly. Micro-locally  $S_{\text{red},k}^{(q-r)}$  is the  $k$ -th Fourier component of the standard Szegő kernel for the CR manifold  $X_\nu^{\text{red}}$ , on the other hand  $S_{k\nu}^{(q)}$  is described in Section 2.2. In fact,  $\sigma_{k\nu}$  is an isomorphism for  $k$  large if we can prove that

$$\sigma_{k\nu} : H^q(X)_{k\nu} \rightarrow H^{q-r}(X_\nu)_k \quad \text{and} \quad \sigma_{k\nu}^* : H^{q-r}(X_\nu^{\text{red}})_k \rightarrow H^q(X)_{k\nu}$$

are injective for  $k$  large.

The injectivity of the map  $\sigma_{k\nu}$  is a consequence of the following equality of operators

$$\sigma_{k\nu}^* \sigma_{k\nu} \equiv c_0 (1 + R_k) S_{k\nu}^{(q)}$$

where  $c_0$  is a positive constant and  $1 + R_k : \Omega^{0,q}(X) \rightarrow \Omega^{0,q}(X)$  is an injective integral operator of *equivariant Szegő type*, see Definition 2.2. The injectivity of operator  $1 + R_k$  follows by direct computations of the leading symbol of  $R_k$  which vanishes at  $\text{diag}(X_\nu \times X_\nu)$ . In fact, as a consequence of this, by Lemma 6.8 in [HH] we have that  $\|R_k u\| \leq \epsilon_k \|u\|$  for all  $u \in \Omega^{0,q}(X)$ , for all  $k \in \mathbb{N}$  where  $\epsilon_k$  is a sequence with  $\lim_{k \rightarrow +\infty} \epsilon_k = 0$ ; this in turn implies, if  $k$  is large enough, that the map  $1 + R_k$  is injective.

The injectivity of the map  $\sigma_{k\nu}^*$  follows by studying  $\sigma_{k\nu} \sigma_{k\nu}^*$ . We don't repeat the proof here since it is an application of the stationary phase formula for Fourier integral operators of complex type, see [MS], and it follows in a similar way as in [HH].

**Acknowledgements:** This project was started during the author's postdoctoral fellowship at the National Center for Theoretical Sciences in Taiwan; the author thanks the Center for the support.

## References

- [GZ] Q. Guan, X. Zhou, *A solution of an  $L^2$  extension problem with an optimal estimate and applications*, Ann. of Math. (2), 181, 1139–1208 (2015)
- [GS1] V. Guillemin, S. Sterneberg *Geometric Quantization and Multiplicities of Group Representations*, Invent. math. 67, 515-538 (1982)
- [GS2] V. Guillemin, S. Sterneberg *Homogeneous Quantization and Multiplicities of Group Representations*, Journal of functional analysis 41, 344-380 (1982)
- [Hs] C.-Y. Hsiao, *Projections in several complex variables*, Mémoires de la Société Mathématique de France **123**, 1-136 (2010)
- [HH] C.-Y. Hsiao, R.-T. Huang, *G-invariant Szegő kernel asymptotics and CR reduction*, Calc. Var. Partial Differential Equations 60, no. 1 (2021)
- [Kohn] J. J. Kohn, *The range of the tangential Cauchy-Riemann operator*, Duke Math. J. **53**, No. 2, 307–562 (1986)
- [MS] A. Melin and J. Sjöstrand, *Fourier integral operators with complex valued phase functions*, Springer Lecture Notes in Math **459** (1074), pp. 120-223, (1975)
- [P1] R. Paoletti, *Asymptotics of Szegő kernels under Hamiltonian torus actions*, Isr. J. Math. 191 (1) 363–403, (2012)
- [P2] R. Paoletti, *Polarized orbifolds associated to quantized Hamiltonian torus actions*, Journal of Geometry and Physics, (170), (2021)