

DECAY ESTIMATES FOR FOURTH-ORDER SCHRÖDINGER OPERATORS IN DIMENSION TWO

PING LI, AVY SOFFER AND XIAOHUA YAO

ABSTRACT. In this paper we study the decay estimates of the fourth order Schrödinger operator $H = \Delta^2 + V(x)$ on \mathbb{R}^2 with a bounded decaying potential $V(x)$. We first deduce the asymptotic expansions of resolvent of H near the zero threshold in the presence of resonances or eigenvalue, and then use them to establish the $L^1 - L^\infty$ decay estimates of e^{-itH} generated by the fourth order Schrödinger operator H . Finally, we classify these zero resonance functions as the distributional solutions of $H\phi = 0$ in suitable weighted spaces. Our methods depend on Littlewood-Paley decomposition and oscillatory integral theory. Moreover, due to the degeneracy of Δ^2 at zero threshold and the lower even dimension (i.e. $n = 2$), we remark that the asymptotic expansions of resolvent $R_V(\lambda^4)$ and the classifications of resonances are more involved than Schrödinger operator $-\Delta + V$ in dimension two.

1. INTRODUCTION

1.1. Backgrounds. In this paper we consider the time decay estimates of e^{-itH} generated by the fourth order Schrödinger operator on $L^2(\mathbb{R}^2)$

$$H = H_0 + V(x), \quad H_0 = \Delta^2,$$

where $V(x)$ is a real measurable function satisfying $|V(x)| \lesssim (1 + |x|)^{-\beta}$ for some $\beta > 0$.

Recall that free propagator e^{-itH_0} on $L^2(\mathbb{R}^n)$ can be expressed by

$$e^{-itH_0} f = \mathfrak{F}^{-1}(e^{-it|\xi|^4} \hat{f}) = \int_{\mathbb{R}^n} I_0(t, x - y) f(y) dy, \quad (1.1)$$

where \hat{f} (or $\mathfrak{F}(f)$) denotes Fourier transform of f , $\mathfrak{F}^{-1}(f)$ denotes its inverse Fourier transform, and $I_0(t, x) := \mathfrak{F}^{-1}(e^{-it|\xi|^4})(x)$ is the kernel of e^{-itH_0} . It was well-known that the following sharp point-wise estimates hold (see e.g. [BKS00]),

$$|D^\alpha I_0(t, x)| \lesssim |t|^{-\frac{n+\alpha}{4}} (1 + |t|^{-\frac{1}{4}} |x|)^{-\frac{n-|\alpha|}{3}}, \quad |t| \neq 0, \quad x \in \mathbb{R}^n, \quad (1.2)$$

for any $\alpha \in \mathbb{N}^n$. Then by the equality (1.1), the estimate (1.2) immediately implies the following $L^1 - L^\infty$ estimates by Young's inequality:

$$\|D^\alpha e^{-itH_0}\|_{L^1 \rightarrow L^\infty} \lesssim |t|^{-\frac{n+\alpha}{4}}, \quad |\alpha| \leq n. \quad (1.3)$$

Recently, several works were devoted to the study of the fourth order Schrödinger operator $H = \Delta^2 + V$ with a decaying potential V , focusing on the time decay estimates of e^{-itH} generated by H . Feng, Soffer and Yao [FSY18] first gave the asymptotic expansion of the

Date: October 15, 2021.

Key words and phrases. Decay estimates, Fourth-order Schrödinger operators, Asymptotic expansion of resolvent, Zero resonance, Dimension two .

resolvent $R_V = (H - z)^{-1}$ around zero threshold assume that zero is a regular point of H for $n \geq 5$ and $n = 3$, and proved that the Kato-Jensen decay estimate of e^{-itH} is $(1 + |t|)^{-n/4}$ for $n \geq 5$ and $L^1 - L^\infty$ decay estimate is $O(|t|^{-1/2})$ for $n = 3$ in the regular case.

Somewhat later, based on the higher energy estimates of resolvent in [FSY18], Erdoğan, Green and Toprak [EGT19] for $n = 3$ and Green-Toprak [GT19] for $n = 4$, derived the asymptotic expansion of $R_V(z)$ near zero with the presence of resonance or eigenvalue, and proved that the time decay rate of $L^1 - L^\infty$ estimates is $|t|^{-n/4}$ for $n = 3, 4$ if zero is a regular point, and the time decay rate will be changed as zero energy resonance occurs. Soffer, Wu and Yao [SWY21] proved the time decay rate of $L^1 - L^\infty$ estimates is $|t|^{-1/4}$ for $n = 1$ whatever zero is a regular point or a zero resonance. It should be emphasized that presences of zero resonance do not change the optimal time decay rate of e^{-itH} in dimension one, at the cost of faster decay rate of potential. Moreover, we also note that there exist some interesting works [GG21] and [EG21] on the L^p bounds of wave operators for $n \geq 3$ in the regular case.

Besides the bi-Laplace operator Δ^2 , we remark that Kato-Jensen estimates and asymptotic expansion of resolvent have been first established by Feng, et al [FSWY20] for poly-Schrödinger operators $(-\Delta)^m + V$ with $n \geq 2m \geq 4$. In the previous work, Murata [Mur82] have generalized Kato and Jensen's works [JK79, Jen80] to a certain class of $P(D) + V$ assume that $P(D)$ satisfies

$$(\nabla P)(\xi_0) = 0, \det[\partial_i \partial_j P(\xi)] \Big|_{\xi_0} \neq 0. \quad (1.4)$$

However, the poly-harmonic operators $H_0 = (-\Delta)^m$ do not satisfy the nondegenerate condition (1.4) at $\xi = 0$ but the case $m = 1$. Hence the results in [FWY18] and [FSWY20] were not covered by [Mur82]. In particular, the degeneracy of $(-\Delta)^m$ at zero energy leads to much more complicated zero resonances classifications, and asymptotic expansion of the resolvent of $(-\Delta)^m + V$ as $m \geq 2$. These are actually some of the main difficult points for the higher order estimates.

In this paper, we are devoted to establish the $L^1 - L^\infty$ estimates of the fourth order Schrödinger operator $\Delta^2 + V$ in dimension two. We first deduce the asymptotic expansion of $R_V(z)$ near zero with the presence of resonance or eigenvalue, and then study the $L^1 - L^\infty$ estimates for each kind of zero resonance. In particular, we show that the $L^1 - L^\infty$ estimates of e^{-itH} has the optimal decay rate $O(|t|^{-\frac{1}{2}})$ if zero is a regular point and the first type of resonance. Moreover, we classify these zero resonance functions as the distributional solutions of $H\phi = 0$ in suitable weighted spaces. Due to the degeneracy of Δ^2 at zero threshold and the lower even dimension, we remark that the asymptotic expansions of resolvent $R_V(\lambda^4)$ at $\lambda = 0$ are much complicated compared with Schrödinger operator $-\Delta + V$ in dimension two (see e.g. [JN01, JN04]). One can see Section 4 for more details.

Finally, we notice that the studies of higher order elliptic operator are largely motivated by Schrödinger operator. It is well-known that the decay estimates of Schrödinger operator have been one of the active topics in the last thirty years, and applied broadly to nonlinear Schrödinger equations. For instance, Journé, Soffer and Sogge [JSS91] first established the $L^1 - L^\infty$ dispersive estimates in regular case when $n \geq 3$. Later, Weder [Wed00], Rodnianski and Schlag [RS04], Schlag and Goldberg [GS04], Schlag [Sch05] have made further contributions for $n \leq 3$. In particular, Yajima in [Yaj95] has established the $L^1 - L^\infty$ estimates for $-\Delta + V$

by wave operator method. For many further studies, one can refer to [Wed99, ES04, ES06, GG15, GG17, Sch07, Sch21] and therein references.

1.2. Main results. In this subsection, we will state our main results. We first introduce some notations and zero resonances.

For $a \in \mathbb{R}$, we write $a \pm$ to mean $a \pm \epsilon$ for a small, but fixed $\epsilon > 0$. $[a]$ denotes the largest integer less than or equal to a . For $a, b \in \mathbb{R}^+$, $a \lesssim b$ (or $a \gtrsim b$) means that there exists some constant $c > 0$ such that $a \leq cb$ (or $a \geq cb$). Moreover, if $a \lesssim b$ and $b \lesssim a$, then we write $a \sim b$. Let $\langle \cdot \rangle = (1 + |\cdot|^2)^{\frac{1}{2}}$, we define the weighted L^p spaces

$$L_s^p(\mathbb{R}^2) = \{f : (1 + |\cdot|)^s f \in L^p(\mathbb{R}^2)\}.$$

We use the smooth, even low energy cut-off χ defined by $\chi(\lambda) = 1$ if $|\lambda| < \lambda_0 \ll 1$ and $\chi(\lambda) = 0$ when $|\lambda| > 2\lambda_0$ for some sufficiently small constant $0 < \lambda_0 \ll 1$. In analyzing the high energy we utilize the complementary cut-off $\tilde{\chi}(\lambda) = 1 - \chi(\lambda)$.

In this paper, we can give the precise definitions of different types of zero resonances in the terms of the projection operators S_j corresponding to the kernel spaces of operators T_j (see Definition 2.4 of Section 2 below). Equivalently, we have characterizations of the zero resonance types in terms of the solutions to $H\phi = 0$ in distributional sense, see Theorem 5.7 of Section 5 below.

More specifically, we say that zero is a *resonance of the first kind* if $\exists 0 \neq \phi \in L_{-1}^\infty(\mathbb{R}^2)$ but have no nonzero $\phi \in L^\infty(\mathbb{R}^2)$ such that $H\phi = 0$, the *second kind* if $\phi \in L^\infty(\mathbb{R}^2)$ but have no nonzero $\phi \in L_1^\infty(\mathbb{R}^2)$, the *third kind* if $\phi \in L_1^\infty(\mathbb{R}^2)$ but have no nonzero $\phi \in L^2(\mathbb{R}^2)$, and a *resonance of the fourth kind* if $\exists 0 \neq \phi \in L^2(\mathbb{R}^2)$ (i.e. the fourth kind resonance is exactly the zero eigenvalue). If zero is neither a resonance or an eigenvalue, then we say zero is a *regular point* of H .

Now we are ready to present the main results as follows.

Theorem 1.1. *Let $|V(x)| \lesssim (1 + |x|)^{-\beta}$ for $x \in \mathbb{R}^2$ and some $\beta > 0$. Assume that $H = \Delta^2 + V(x)$ has no positive eigenvalues, and $P_{ac}(H)$ denotes the projection onto the absolutely continuous spectrum space of H . Then the following conclusions hold:*

(i) *If zero is the regular point and $\beta > 10$, then*

$$\|e^{-itH} P_{ac}(H)\|_{L^1 \rightarrow L^\infty} \lesssim |t|^{-\frac{1}{2}}. \quad (1.5)$$

(ii) *If zero is the first kind resonance and $\beta > 14$, then*

$$\|e^{-itH} P_{ac}(H)\|_{L^1 \rightarrow L^\infty} \lesssim |t|^{-\frac{1}{2}}. \quad (1.6)$$

(iii) *If zero is the second kind resonance and $\beta > 18$, then*

$$\|H^{\frac{1}{2}} e^{-itH} P_{ac}(H) \chi(H)\|_{L^1 \rightarrow L^\infty} + \|e^{-itH} \tilde{\chi}(H)\|_{L^1 \rightarrow L^\infty} \lesssim |t|^{-\frac{1}{2}}. \quad (1.7)$$

(iv) *If zero is the third or fourth kind resonance and $\beta > 18$, then*

$$\|H^{\frac{3}{2}} e^{-itH} P_{ac}(H) \chi(H)\|_{L^1 \rightarrow L^\infty} + \|e^{-itH} \tilde{\chi}(H)\|_{L^1 \rightarrow L^\infty} \lesssim |t|^{-\frac{1}{2}}. \quad (1.8)$$

Here, $\chi(\lambda)$ and $\tilde{\chi}(\lambda)$ in the (1.7) and (1.8) denote the low and high energy smooth cut-off functions, respectively.

Remark 1.2. *Some further comments on Theorem 1.1 :*

(i) *If zero is the regular point or the first kind resonance, then the decay estimates (1.5) and (1.6) of e^{-itH} are optimal in view of the fact*

$$\|e^{-it\Delta^2}\|_{L^1(\mathbb{R}^2)\rightarrow L^\infty(\mathbb{R}^2)} \lesssim |t|^{-1/2}.$$

Moreover, in view of the decay estimates (1.3) with a derivatives term for $n = 2$, we remark that it is possible to use the present methods to improve the estimates (1.5) and (1.6) to $O(|t|^{-(2+\alpha)/4})$ by adding a regular term $H^{\alpha/4}$ for $0 \leq |\alpha| \leq 2$. For such related results, one can refer to [SWY21] for dimension $n = 1$ (also see [Hill20] for the regular case) and [LWY21] for dimension $n = 3$.

(ii) *If zero is the second, third or fourth kind resonance, then the high energy part $e^{-itH}\tilde{\chi}(H)$ has always the sharp time decay estimates $O(|t|^{-1/2})$.*

For low energy part, however, since the zero singularity of the asymptotic expansions of $(M^\pm(\lambda))^{-1}$ are of too high order to estimate $e^{-itH}P_{ac}(H)\chi(H)$ (See Section 2 below), we put suitable regularity terms H^α into the low energy part $e^{-itH}P_{ac}(H)\chi(H)$ to temper zero singularities, as shown in the above decay estimates (1.7) and (1.8).

Besides the conditions of zero resonance, Theorem 1.1 also assumes that $H = \Delta^2 + V$ has no positive eigenvalues embedded into the absolutely continuous spectrum, which has been the indispensable assumption in dispersive estimates. For Schrödinger operator $-\Delta + V$, Kato in [Kat59] showed the absence of positive eigenvalues for $H = -\Delta + V$ with the decay potentials $V = O(|x|^{-1})$ as $|x| \rightarrow \infty$.

However, the situation for $H = \Delta^2 + V$ seems to more complicated than the second order cases. There exist compactly supported smooth potentials such that the $H = \Delta^2 + V$ has some positive eigenvalues (see [FSWY20]). On the other hand, more interestingly, [FSWY20] has showed that $H = \Delta^2 + V$ has no positive eigenvalues, assuming that the potential V is bounded and satisfies the repulsive condition (i.e. $(x \cdot \nabla)V \leq 0$). Moreover, we also notice that for a general self-adjoint operator \mathcal{H} on $L^2(\mathbb{R}^n)$, even if \mathcal{H} has a simple embedded eigenvalue, Costin and Soffer in [CS01] have proved that $\mathcal{H} + \epsilon W$ can kick off the eigenvalue located in a small interval under generic small perturbation of potential.

1.3. The outline of proof. In order to show Theorem 1.1, we first recall that from functional calculus and Stone's formula

$$H^\alpha e^{-itH} P_{ac}(H) f(x) = \frac{2}{\pi i} \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} [R_V^+(\lambda^4) - R_V^-(\lambda^4)] f(x) d\lambda. \quad (1.9)$$

Hence we need to study carefully the boundary behaviour of the resolvent operators $R_V(z) = (H - z)^{-1}$ by perturbations of the free resolvent $(\Delta^2 - z)^{-1}$, which in turn depends on the representation:

$$R_0(z) := (\Delta^2 - z)^{-1} = \frac{1}{2z^{\frac{1}{2}}} (R(-\Delta; z^{\frac{1}{2}}) - R(-\Delta; -z^{\frac{1}{2}})), \quad z \in \mathbb{C} \setminus [0, \infty). \quad (1.10)$$

Here the resolvent $R(-\Delta; z^{\frac{1}{2}}) := (-\Delta - z^{\frac{1}{2}})^{-1}$.

For $\lambda > 0$, if we define the incoming and outgoing resolvent operators by

$$R_0^\pm(\lambda^4) := R_0^\pm(\lambda^4 \pm i0) = \lim_{\epsilon \rightarrow 0} (\Delta^2 - (\lambda^4 \pm i\epsilon))^{-1}, \quad (1.11)$$

then we have

$$R_0^\pm(\lambda^4) = \frac{1}{2\lambda^2} (R^\pm(-\Delta; \lambda^2) - R(-\Delta; -\lambda^2)), \quad \lambda > 0. \quad (1.12)$$

It was well-known that by the limiting absorption principle (see e.g. [Agm75]), $R^\pm(-\Delta; \lambda^2)$ are well-defined as the bounded operators of $B(L_s^2, L_{-s}^2)$ for any $s > 1/2$, therefore $R_0^\pm(\lambda^4)$ are also well-defined between these weighted L^2 -space. These bounded properties can be extended to $R_V^\pm(\lambda^4)$ for $\lambda > 0$ for certain decay bounded potentials V , see [FSY18].

In order to estimate the integral (1.9), we need to study the properties of $R_V^\pm(\lambda^4)$ near $\lambda = 0$ and $+\infty$, which corresponds to the low energy estimate and high energy estimate respectively. For lower energy parts, since $R_V^\pm(\lambda^4)$ are related to $R_0^\pm(\lambda^4)$ by the following symmetric resolvent formula:

$$R_V^\pm(\lambda^4) = R_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)v(M^\pm(\lambda))^{-1}vR_0^\pm(\lambda^4),$$

where $M^\pm(\lambda) = U + vR_0^\pm(\lambda^4)v$, $U(x) = \text{sign}(V(x))$ and $v(x) = |V(x)|^{1/2}$. Then we turn to analyze the expansion of $(M^\pm(\lambda))^{-1}$ at zero (see Theorem 2.6 below).

By using the expansions of free resolvent of $R_0^\pm(\lambda^4)$ in (1.12), we can obtain that

$$M^\pm(\lambda) = \frac{a_\pm}{\lambda^2}P + g_0^\pm(\lambda)vG_{-1}v + T_0 + c_\pm\lambda^2vG_1v + \tilde{g}_2^\pm(\lambda)\lambda^4vG_2v + \lambda^4vG_3v + E_2^\pm(\lambda),$$

where $P, vG_i v (i = -1, 0, 1, 2, 3)$ are the specific bounded operators on $L^2(\mathbb{R}^2)$ and $E_2^\pm(\lambda)$ are the error terms, see Section 2 below for specific descriptions. Note that the expansion of the $M^\pm(\lambda)$ includes two terms with logarithm factors $g_0^\pm(\lambda)$ and $\tilde{g}_2^\pm(\lambda)$, which make the calculation of the inversion processes $(M^\pm(\lambda))^{-1}$ to be quite complicated. We need use the operator matrix Lemma 4.2 twice to complete the whole inversion processes in Section 4.

Firstly, in the regular case, we need to show that the inverse operator

$$d = \left(g_0^\pm(\lambda)(Q - S_0)vG_{-1}v(Q - S_0) + (Q - S_0)(T_0 - T_0D_1T_0)(Q - S_0) \right)^{-1}$$

exist on $(Q - S_0)L^2$ for enough small λ . Since $\dim(Q - S_0)L^2 = 2$, so the operators

$$(Q - S_0)vG_{-1}v(Q - S_0)$$

and

$$(Q - S_0)(T_0 - T_0D_1T_0)(Q - S_0)$$

are essentially two 2×2 -matrix linear transforms acting on $(Q - S_0)L^2$. Hence it suffices to prove the matrix $(Q - S_0)vG_{-1}v(Q - S_0)$ is invertible on the subspace $(Q - S_0)L^2$ in order to obtain the existence of the inverse operator d on $(Q - S_0)L^2$ as λ is small. This will be proved on Lemma 4.3.

Secondly, in the second kind of resonance, we will need to prove that another inverse operator

$$d_1 := \left(g_0^\pm(\lambda)^2L_1 + g_0^\pm(\lambda)L_2 + L_3 \right)^{-1}$$

exists on $(S_2 - S_3)L^2$ for enough small λ , where

$$\begin{aligned} L_1 &= (S_2 - S_3) \left(vG_{-1}vPvG_{-1}v - vG_{-1}vT_0D_4T_0vG_{-1}v \right) (S_2 - S_3), \\ L_2 &= (S_2 - S_3) \left(vG_{-1}vT_0 + T_0vG_{-1}v - vG_{-1}vT_0D_4(T_0^2 - c_+a_+vG_1v) \right. \\ &\quad \left. - (T_0^2 - c_+a_+vG_1v)D_4T_0vG_{-1}v \right) (S_2 - S_3), \\ L_3 &= (S_2 - S_3) \left(T_0^2 - c_+a_+vG_1v - (T_0^2 - c_+a_+vG_1v)D_4(T_0^2 - c_+a_+vG_1v) \right) (S_2 - S_3). \end{aligned}$$

Since $(S_2 - S_3)L^2$ is actually one dimensional space, hence it suffices to prove that $\text{tr}(L_1) \neq 0$ in order that L_1 is invertible acting on $(S_2 - S_3)L^2$. These calculations are very complicate, one can see Lemma 4.6 for the more details.

In order to obtain the decay estimates in Theorem 1.1, we will use the following Littlewood-Paley decompositions:

$$\begin{aligned} e^{-itH} P_{ac}(H)(x, y) &= \sum_{N=-\infty}^{\infty} \frac{2}{\pi i} \int_0^{\infty} e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) [R_V^+(\lambda^4) - R_V^-(\lambda^4)](x, y) d\lambda \\ &= \sum_{N=-\infty}^{\infty} K_N(t, x, y), \end{aligned} \tag{1.13}$$

where $\sum_{N=-\infty}^{\infty} \varphi_0(2^{-N}\lambda) = 1$ for $\lambda > 0$, and use oscillatory integrals methods to establish the bound of each dyadic kernel $K_N(t, x, y)$.

For each each dyadic kernel $K_N(t, x, y)$, we will make use of the asymptotic expansion of $R_V(\lambda^4)$ near zero for $N \ll 0$ (i.e. Theorem 2.6) and the higher energy estimates of resolvent $R_V(\lambda^4)$ for $N \gtrsim 0$ (see [FSY18] or Lemma 3.16 below). By the scaling process and suitable reductions for each N , we will reduce the estimate of $K_N(t, x, y)$ to study the following oscillatory integral:

$$\int_0^{\infty} e^{-it2^{4N}s^4} e^{\pm i2^N s \Phi(z)} \Psi(2^N s, z) \varphi_0(s) ds \tag{1.14}$$

where $\Phi(z)$ is some phase function on \mathbb{R}^m and $\Psi(s, z)$ is some amplitude functions $\mathbb{R}^+ \times \mathbb{R}^m$. Under the suitable conditions of $\Phi(z)$ and $\Psi(s, z)$, the stationary phase method can be applied to estimate the oscillatory integral (1.14), see Lemma 3.3 below for the details.

Noting that the oscillatory integrals (1.14) have similarly appeared in one dimensional case [SWY21], however, the estimates established in [SWY21] can't work in dimension two. Here we need to make more accurate estimates for the oscillatory integrals (1.14), i.e. Lemma 3.3, which allow us to work well on the two dimensional problems and even other dimensional cases (see e.g. [LWY21]).

The paper is organized as follows. In section 2, we first compute the expansions for the free resolvent, and then we state the asymptotic expansion of $(M^\pm(\lambda))^{-1}$ when λ is near zero. In section 3, we are devoted to proving Theorem 1.1. In section 4, we prove Theorem 2.6. In section 5, we characterize the resonance spaces for each kind of zero resonance.

2. RESOLVENT ASYMPTOTIC EXPANSIONS NEAR ZERO

In this section, we first study the free resolvent asymptotic expansions of $R_0^\pm(\lambda^4)$, then we give the asymptotic expansions of $M^\pm(\lambda)^{-1}$.

We first gathered some frequently used notations throughout the paper. For an operator $\mathcal{E}(\lambda)$, we write $\mathcal{E}(\lambda) = O_1(\lambda^{-s})$ if it's kernel $\mathcal{E}(\lambda)(x, y)$ has the property

$$\sup_{x, y \in \mathbb{R}^2} [\lambda^s |\mathcal{E}(\lambda)(x, y)| + \lambda^{s+1} |\partial_\lambda \mathcal{E}(\lambda)(x, y)|] < \infty.$$

Similarly, we use the notation $\mathcal{E}(\lambda) = O_1(\lambda^{-s}g(x, y))$ if $\mathcal{E}(\lambda)(x, y)$ satisfies

$$|\mathcal{E}(\lambda)(x, y)| + \lambda |\partial_\lambda \mathcal{E}(\lambda)(x, y)| < \lambda^{-s}g(x, y).$$

2.1. The free resolvent. In this subsection, by using the identity (1.12) and the Bessel function representation of the Schrödinger free resolvents $R^\pm(-\Delta; \lambda^2)$, we now derive the asymptotic expansions of the fourth order Schrödinger operators free resolvent $R_0^\pm(\lambda^4)$.

Recall first the free resolvent of Laplacian (see e.g. [EG13]):

$$R^\pm(-\Delta; \lambda^2) = \pm \frac{i}{4} H_0^\pm(\lambda|x-y|), \quad (2.1)$$

where H_0^\pm are Hankel functions of order zero:

$$H_0^\pm(z) = J_0(z) \pm iY_0(z), \quad (2.2)$$

and $H_0^-(z) = \overline{H_0^+(z)}$. From the series expansions for Bessel functions (see e.g. [AS64]), as $z \rightarrow 0$ we have

$$J_0(z) = 1 - \frac{1}{4}z^2 + \frac{1}{64}z^4 - \frac{1}{2304}z^6 + O(z^8), \quad (2.3)$$

$$Y_0(z) = \frac{2}{\pi} \left(\ln\left(\frac{z}{2}\right) + \gamma \right) J_0(z) + \frac{2}{\pi} \left(\frac{1}{4}z^2 - \frac{3}{128}z^4 + \frac{11}{13824}z^6 + O(z^8) \right), \quad (2.4)$$

where γ is Euler's constant.

Further, for $|z| > 1$, we have the representation (see e.g. [AS64])

$$H_0^\pm(z) = e^{\pm iz} w_\pm(z), \quad |w_\pm^{(\ell)}(z)| \lesssim (1 + |z|)^{-\frac{1}{2}-\ell}, \ell = 0, 1, 2, \dots. \quad (2.5)$$

Hence, if $\lambda|x-y| \gtrsim 1$, noting that

$$R_0^\pm(\lambda^4) = \frac{1}{2\lambda^2} (R^\pm(-\Delta; \lambda^2) - R(-\Delta; -\lambda^2)), \quad \lambda > 0,$$

by using (2.1) and (2.5), we obtain that

$$R_0^\pm(\lambda^4)(x, y) = \frac{i}{8\lambda^2} \left[\pm e^{\pm i\lambda|x-y|} w_\pm(\lambda|x-y|) - e^{-\lambda|x-y|} w_+(i\lambda|x-y|) \right]. \quad (2.6)$$

Now we compute the representation of $R_0^\pm(\lambda^4)(x, y)$ when $\lambda|x-y| \lesssim 1$. Let G_k denote integral operators with the kernel $G_k(x, y)$ which defined as follows:

$$\begin{aligned} G_{-1}(x, y) &= |x-y|^2, \quad G_0(x, y) = a_0|x-y|^2 \ln|x-y|, \quad G_1(x, y) = |x-y|^4, \\ G_2(x, y) &= |x-y|^6, \quad G_3(x, y) = a_2|x-y|^6 \ln|x-y|, \end{aligned} \quad (2.7)$$

where the coefficients $a_0 = \frac{1}{8\pi}$ and $a_2 = \frac{1}{4608\pi}$.

Lemma 2.1. *If $\lambda|x - y| \ll 1$, then*

$$R_0^\pm(\lambda^4)(x, y) = \frac{b_\pm}{\lambda^2} + g_0^\pm(\lambda)G_{-1}(x, y) + G_0(x, y) + c_\pm\lambda^2G_1(x, y) + \tilde{g}_2^\pm(\lambda)\lambda^4G_2(x, y) \quad (2.8)$$

$$+ \lambda^4G_3(x, y) + O(\lambda^{6-\epsilon}|x - y|^{8-\epsilon}),$$

where $b_\pm = \pm \frac{i}{8}$, $c_\pm = \pm \frac{i}{512}$, $g_0^\pm(\lambda) = a_0 \ln \lambda + \alpha_\pm$, $\tilde{g}_2^\pm(\lambda) = \lambda^4(a_2 \ln \lambda + \beta_\pm)$, and $\alpha_\pm, \beta_\pm \in \mathbb{C}$.

Proof. Let $r = |x - y|$. By using (2.2)-(2.4), we have

$$R^\pm(-\Delta; \lambda^2) = \frac{i}{4}H_0^\pm(\lambda r)$$

$$= \left(-\frac{1}{2\pi} \ln \lambda \pm \frac{i}{4} + \frac{\ln 2}{2\pi} - \frac{\gamma}{2\pi} \right) - \frac{1}{2\pi} \ln r + \left(\frac{1}{8\pi} \ln \lambda \mp \frac{i}{16} - \frac{\ln 2 - \gamma + 1}{8\pi} \right) \lambda^2 r^2$$

$$+ \frac{1}{8\pi} \lambda^2 r^2 \ln r + \left(-\frac{1}{128\pi} \ln \lambda \pm \frac{i}{256} + \frac{2 \ln 2 - 2\gamma + 3}{256\pi} \right) \lambda^4 r^4 - \frac{1}{128\pi} \lambda^4 r^4 \ln r$$

$$+ \left(\frac{1}{4608\pi} \ln \lambda \mp \frac{i}{9216} - \frac{\ln 2}{4608\pi} + \frac{\gamma}{4608\pi} - \frac{11}{27648\pi} \right) \lambda^6 r^6 + \frac{1}{4608\pi} \lambda^6 r^6 \ln r + O((\lambda r)^{8-\epsilon}),$$

$$R^+(-\Delta; -\lambda^2) = \frac{i}{4}H_0^+(i\lambda r)$$

$$= \left(-\frac{1}{2\pi} \ln \lambda + \frac{\ln 2 - \gamma}{2\pi} \right) - \frac{1}{2\pi} \ln r + \left(-\frac{1}{8\pi} \ln \lambda + \frac{\ln 2 - \gamma + 1}{8\pi} \right) \lambda^2 r^2 - \frac{1}{8\pi} \lambda^2 r^2 \ln r$$

$$+ \left(-\frac{1}{128\pi} \ln \lambda + \frac{2 \ln 2 - 2\gamma + 3}{256\pi} \right) \lambda^4 r^4 - \frac{1}{128\pi} \lambda^4 r^4 \ln r + \left(-\frac{1}{4608\pi} \ln \lambda + \frac{\ln 2 - \gamma}{4608\pi} \right.$$

$$\left. + \frac{11}{27648\pi} \right) \lambda^6 r^6 - \frac{1}{4608\pi} \lambda^6 r^6 \ln r + O((\lambda r)^{8-\epsilon}).$$

Noting that the identity,

$$R_0^\pm(\lambda^4) = \frac{1}{2\lambda^2} (R^\pm(-\Delta; \lambda^2) - R(-\Delta; -\lambda^2)), \quad \lambda > 0,$$

then the statement is clear by (2.1) and (2.7).

The proof of this lemma is completed. □

Lemma 2.2. *We obtain the following expansions of $R_0^\pm(\lambda^4)(x, y)$:*

$$R_0^\pm(\lambda^4)(x, y) = b_\pm \frac{1}{\lambda^2} + g_0^\pm(\lambda)G_{-1}(x, y) + G_0(x, y) + E_0^\pm(\lambda)(x, y), \quad (2.9)$$

where the error term satisfies $E_0^\pm(\lambda)(x, y) = O_1(\lambda^\ell |x - y|^{2+\ell})$ for $0 < \ell \leq 2$.

Proof. Let $r = |x - y|$. If $\lambda r \ll 1$, then the statement is clear by (2.8). If $\lambda r \gtrsim 1$, note that

$$g_0^\pm(\lambda)G_{-1}(x, y) + G_0(x, y) = |x - y|^2 \left(a_0 \ln(\lambda|x - y|) + \alpha_\pm \right),$$

by using (2.6) and (2.8), one has for $\ell > 0$,

$$\begin{aligned}
 & \left| R_0^+(\lambda^4)(x, y) - \left(b_+ \frac{1}{\lambda^2} + g_0^+(\lambda)G_{-1}(x, y) + G_0(x, y) \right) \right| \\
 &= \left| \frac{1}{\lambda^2} \left(\frac{i}{8} e^{i\lambda r} w_+(\lambda r) - \frac{i}{8} e^{-\lambda r} w_+(i\lambda r) - b_+ \right) - \left(g_0^+(\lambda)G_{-1}(x, y) + G_0(x, y) \right) \right| \\
 &\lesssim \frac{1}{\lambda^2} \left(|w_+(\lambda r)| + |w_+(i\lambda r)| + c \right) + \lambda^\ell r^{2+\ell} \\
 &\lesssim \frac{1}{\lambda^2 (1 + \lambda r)^{\frac{1}{2}}} + \frac{1}{\lambda^2} + \lambda^\ell r^{2+\ell} \lesssim \lambda^\ell r^{2+\ell}.
 \end{aligned} \tag{2.10}$$

Similarly, we obtain that for $0 < \ell \leq 2$,

$$\begin{aligned}
 |\partial_\lambda (R_0^+(\lambda^4)(x, y))| &\lesssim |\partial_\lambda (R_0^+(\lambda^4)(x, y))\chi(\lambda|x-y|)| + |\partial_\lambda (R_0^+(\lambda^4)(x, y))\tilde{\chi}(\lambda|x-y|)| \\
 &\lesssim \lambda^{-1} r^2 + \lambda r^4 \lesssim \lambda^{\ell-1} r^{2+\ell}.
 \end{aligned}$$

The proof of this lemma is completed. \square

2.2. The resolvent expansions near zero. In this subsection, we study carefully the asymptotic expansions of the perturbed resolvent in a neighborhood of the threshold.

Let $U(x) = \text{sign}(V(x))$ and $v(x) = |V(x)|^{1/2}$, then $V = Uv^2$. We write symmetric resolvent identity as follows:

$$R_V^\pm(\lambda^4) = R_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)v(M^\pm(\lambda))^{-1}vR_0^\pm(\lambda^4), \tag{2.11}$$

where $M^\pm(\lambda) = U + vR_0^\pm(\lambda^4)v$. So, we will turn to establish the expansions for $(M^\pm(\lambda))^{-1}$.

Let $T_0 = U + vG_0v$ and $P = \|V\|_{L^1}^{-1}v\langle v, \cdot \rangle$ denote the orthogonal projection onto the span of v . Then we have the following expansions of $M^\pm(\lambda)$.

Lemma 2.3. *Let $a_\pm = \pm \frac{i}{8} \|V\|_1$ and $M^\pm(\lambda) = U + vR_0^\pm(\lambda^4)v$. If $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with some $\beta > 0$, then the following expansions for $M^\pm(\lambda)$ in L^2 hold:*

(i) *If $\beta > 10$, then we have*

$$M^\pm(\lambda) = \frac{a_\pm}{\lambda^2} P + g_0^\pm(\lambda)vG_{-1}v + T_0 + E_0^\pm(\lambda), \tag{2.12}$$

where the error term satisfies $E_0^\pm(\lambda) = O_1(v(x)\lambda^2|x-y|^4v(y))$.

(ii) *If $\beta > 14$, then we have*

$$M^\pm(\lambda) = \frac{a_\pm}{\lambda^2} P + g_0^\pm(\lambda)vG_{-1}v + T_0 + c_\pm \lambda^2 vG_1v + E_1^\pm(\lambda), \tag{2.13}$$

where the error term satisfies $E_1^\pm(\lambda) = O_1(v(x)\lambda^{4-\epsilon}|x-y|^{6-\epsilon}v(y))$.

(iii) *If $\beta > 18$, then we have*

$$M^\pm(\lambda) = \frac{a_\pm}{\lambda^2} P + g_0^\pm(\lambda)vG_{-1}v + T_0 + c_\pm \lambda^2 vG_1v + \tilde{g}_2^\pm(\lambda)\lambda^4 vG_2v + \lambda^4 vG_3v + E_2^\pm(\lambda), \tag{2.14}$$

where the error term satisfies $E_2^\pm(\lambda) = O_1(v(x)\lambda^{6-\epsilon}|x-y|^{8-\epsilon}v(y))$.

Proof. We only prove (2.12), the other cases are similar. Let $r = |x - y|$. If $\lambda r \ll 1$, then the statement is clear by (2.8). If $\lambda r \gtrsim 1$, using (2.6) and (2.8), by the same argument with the proof of Lemma 2.2, we have

$$\left| M^\pm(\lambda) - \frac{a_\pm}{\lambda^2} P - g_0^\pm(\lambda) v G_{-1} v - T_0 \right| \lesssim \lambda^2 v(x) v(y) |x - y|^4, \quad (2.15)$$

and

$$\left| \partial_\lambda \left(M^\pm(\lambda) - \frac{a_\pm}{\lambda^2} P - g_0^\pm(\lambda) v G_{-1} v - T_0 \right) \right| \lesssim \lambda v(x) v(y) |x - y|^4. \quad (2.16)$$

Since the error term is Hilbert-Schmidt operator if $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 10$, which implies (2.12). \square

Now we define the type of resonances that may occur at the zero threshold energy as following.

Let $Q = I - P$ and S_0 be the orthogonal projection onto the kernel of $QvG_{-1}vQ$ on QL^2 . Then $S_0 \neq 0$ by Lemma 2.5(i). Hence, $QvG_{-1}vQ$ is not invertible on QL^2 . But $QvG_{-1}vQ + S_0$ is invertible on QL^2 , in this case, we define $D_0 = (QvG_{-1}vQ + S_0)^{-1}$ as an operator on QL^2 .

Definition 2.4. (i) If $S_0T_0S_0$ is invertible on S_0L^2 , then we say that zero is regular point of the spectrum of H . In this case, we define $D_1 = (S_0T_0S_0)^{-1}$ as an operator on S_0L^2 .

(ii) Assume that $S_0T_0S_0$ is not invertible on S_0L^2 . Let S_1 be the Riesz projection onto the kernel of $S_0T_0S_0$. Then $S_0T_0S_0 + S_1$ is invertible on S_0L^2 . In this case, we define $D_1 = (S_0T_0S_0 + S_1)^{-1}$ as an operator on S_0L^2 , this doesn't conflict with the previous definition since $S_1 = 0$ when zero is regular. We say zero is a resonance of the first kind if the operator

$$T_1 := -S_1T_0(Q - S_0) \left((Q - S_0)vG_{-1}v(Q - S_0) \right)^{-1} (Q - S_0)T_0S_1 \quad (2.17)$$

is invertible on S_1L^2 , where the inverse $\left((Q - S_0)vG_{-1}v(Q - S_0) \right)^{-1}$ is well-defined on the subspace $(Q - S_0)L^2$ by the (4.10) of Lemma 4.3.

(iii) Assume that T_1 is not invertible on S_1L^2 . Let S_2 be the Riesz projection onto the kernel of T_1 . Then $T_1 + S_2$ is invertible on S_1L^2 . In this case, we define $D_2 = (T_1 + S_2)^{-1}$ as an operator on S_1L^2 . Let $T_2 := S_2vG_{-1}vS_2$. then T_2 is not invertible on S_2L^2 by Lemma 2.5(iv). Let S_3 be the Riesz projection onto the kernel of T_2 , then $T_2 + S_3$ is invertible on S_2L^2 , in this case, we define $D_3 = (T_2 + S_3)^{-1}$ as an operator on S_2L^2 . We say zero is a resonance of the second kind if

$$T_3 := S_3T_0^2S_3 - c_\pm a_\pm S_3vG_1vS_3 \quad (2.18)$$

is invertible on S_3L^2 .

(iv) Assume that T_3 is not invertible on S_3L^2 . Let S_4 be the Riesz projection onto the kernel of T_3 . Then $T_3 + S_4$ is invertible on S_3L^2 . In this case, we define $D_4 = (T_3 + S_4)^{-1}$ as an operator on S_3L^2 . We say zero is a resonance of the third kind if

$$T_4 := S_4vG_2vS_4 \quad (2.19)$$

is invertible on S_4L^2 .

(v) Finally if T_4 is not invertible on S_4L^2 , we say there is a resonance of the fourth kind at zero. In this case, the operator $T_5 := S_4vG_3vS_4$ is always invertible on S_5L^2 where S_5 be the Riesz projection onto the kernel of T_4 , see Lemma 4.8. We define $D_5 = (T_4 + S_5)^{-1}$ as an operator on S_4L^2 .

From definitions above, it is clear that $S_0L^2 \supseteq S_1L^2 \supseteq S_2L^2 \supseteq S_3L^2 \supseteq S_4L^2 \supseteq S_5L^2$, and these projection spaces in Definition 2.4 can directly describe the zero energy resonance types of H as follows:

- (i) zero is a regular point of H if and only if $S_1L^2(\mathbb{R}^2) = \{0\}$;
- (ii) zero is a first kind resonance of H if and only if $S_1L^2(\mathbb{R}^2) \neq \{0\}$ and $S_2L^2(\mathbb{R}^2) = \{0\}$;
- (iii) zero is a second kind resonance of H if and only if $S_2L^2(\mathbb{R}^2) \neq \{0\}$ and $S_4L^2(\mathbb{R}^2) = \{0\}$;
- (iii) zero is a third kind resonance of H if and only if $S_4L^2(\mathbb{R}^2) \neq \{0\}$ and $S_5L^2(\mathbb{R}^2) = \{0\}$;
- (iv) zero is an eigenvalue of H (i.e. the fourth kind resonance) if and only if $S_5L^2(\mathbb{R}^2) \neq \{0\}$.

In view of the statements above, we have noticed that the space S_0L^2 and S_3L^2 do not directly correspond to the resonance types, but as transient paces, they actually will play important roles in establishing the expansions of $M(\lambda)^{-1}$ when zero is the regular point and second resonance type (see e.g. Section 4 for the more details).

In addition, we have the following characterizations of projection spaces S_jL^2 .

Lemma 2.5. (i) $S_0L^2 = \{f \in QL^2 \mid \langle x_j v, f \rangle = 0, j = 1, 2\}$.

(ii) $S_1L^2 = \{f \in S_0L^2 \mid S_0T_0f = 0\}$.

(iii) $S_2L^2 = \{f \in S_1L^2 \mid QT_0f = 0\}$.

(iv) $S_3L^2 = \{f \in S_2L^2 \mid \langle |x|^2 v, f \rangle = 0\}$.

(v) $S_4L^2 = \{f \in S_3L^2 \mid \langle x_i x_j v, f \rangle = 0, i, j = 1, 2, \text{ and } PT_0f = 0\}$.

(vi) $S_5L^2 = \{f \in S_4L^2 \mid \langle x_i x_j x_k v, f \rangle = 0, i, j, k = 1, 2\}$.

Proof. (i) Let $f \in S_0L^2$. Then we have

$$0 = \langle QvG_{-1}vQf, f \rangle = \langle vG_{-1}vf, f \rangle.$$

By using (2.7) and the orthogonality $\langle v, f \rangle = 0$, one has

$$\begin{aligned} 0 &= \langle vG_{-1}vf, f \rangle = \int_{\mathbb{R}^2} v(x)(G_{-1}vf)(x)\overline{f(x)}dx \\ &= \int_{\mathbb{R}^2} v(x)\overline{f(x)} \int_{\mathbb{R}^2} |x-y|^2 v(y)f(y)dydx \\ &= \int_{\mathbb{R}^2} v(x)\overline{f(x)} \int_{\mathbb{R}^2} (|x|^2 - 2(x_1y_1 + x_2y_2) + |y|^2)v(y)f(y)dydx \\ &= -2 \left(\int_{\mathbb{R}^2} x_1 v(x)f(x)dx \right)^2 - 2 \left(\int_{\mathbb{R}^2} x_2 v(x)f(x)dx \right)^2, \end{aligned}$$

which leads to $\langle x_j v, f \rangle = 0, j = 1, 2$.

On the other hand, if $\langle x_j v, f \rangle = 0 (j = 1, 2)$ and $f \in QL^2$, we can obtain $QvG_{-1}vQf = 0$.

(ii) It is obvious by the definition of S_1 .

(iii) Let $f \in S_2L^2 \subset S_1L^2$, then $S_0T_0f = 0$. Since $QT_0f = QT_0f - S_0T_0f = (Q - S_0)T_0f$, we only need to prove $(Q - S_0)T_0f = 0$. Indeed, if $f \in S_2L^2$, we have

$$\begin{aligned} 0 &= \langle -T_1f, f \rangle = \left\langle S_1T_0(Q - S_0)((Q - S_0)vG_{-1}v(Q - S_0))^{-1}(Q - S_0)T_0S_1f, f \right\rangle \\ &= \left\langle ((Q - S_0)vG_{-1}v(Q - S_0))^{-1}(Q - S_0)T_0f, (Q - S_0)T_0f \right\rangle. \end{aligned} \quad (2.20)$$

Since $((Q - S_0)vG_{-1}v(Q - S_0))^{-1} > 0$ on the space $(Q - S_0)L^2$ by the (4.10) of Lemma 4.3, so we must have $(Q - S_0)T_0f = 0$.

On the other hand, if $f \in QL^2$ and $\langle x_j v, f \rangle = 0 (j = 1, 2)$ and $QT_0f = 0$, then it is easy to check that $\langle T_1f, f \rangle = 0$.

(iv) Let $f \in S_3L^2 := \ker(T_2) = \ker(S_2vG_{-1}vvG_{-1}vS_2)$. Then we have

$$0 = \langle S_2vG_{-1}vvG_{-1}vS_2f, f \rangle = \langle vG_{-1}vf, vG_{-1}vf \rangle.$$

Note that the orthogonality $\langle v, f \rangle = 0$ and $\langle x_j v, f \rangle = 0 (j = 1, 2)$, by (2.7) we have

$$\begin{aligned} 0 &= \langle vG_{-1}vf, vG_{-1}vf \rangle = \int_{\mathbb{R}^2} \left| v(x)(G_{-1}vf)(x) \right|^2 dx \\ &= \int_{\mathbb{R}^2} \left| v(x) \left(\int_{\mathbb{R}^2} |x - y|^2 v(y)f(y) dy \right) \right|^2 dx \\ &= \int_{\mathbb{R}^2} |v(x)|^2 \left(\int_{\mathbb{R}^2} |y|^2 v(y)f(y) dy \right)^2 dx, \\ &= \langle |x|^2 v, f \rangle^2 \|V\|_{L^1}, \end{aligned}$$

which leads to $\langle |x|^2 v, f \rangle = 0$.

On the other hand, if $f \in S_2L^2$ and $\langle |x|^2 v, f \rangle = 0$, we can obtain $T_2f = 0$.

(v) Let $f \in S_4L^2$. Note that $QT_0f = 0$ and $P = \frac{\langle v, \cdot \rangle}{\|V\|_{L^2}}v$, we have

$$S_3T_0^2S_3f(x) = S_3T_0(PT_0f)(x) = \frac{\langle v, T_0f \rangle}{\|V\|_{L^1}}S_3(T_0v)(x),$$

which leads to

$$\langle S_3T_0^2S_3f, f \rangle = \frac{\langle v, T_0f \rangle}{\|V\|_{L^1}} \langle S_3(T_0v), f \rangle = \frac{\langle v, T_0f \rangle}{\|V\|_{L^1}} \langle v, T_0S_3f \rangle = \frac{\langle v, T_0f \rangle^2}{\|V\|_{L^1}}. \quad (2.21)$$

Since

$$|x - y|^4 = |x|^4 - 4|x|^2(x_1y_1 + x_2y_2) - 4|y|^2(x_1y_1 + x_2y_2) + 4(x_1^2y_1^2 + x_2^2y_2^2 + 2x_1x_2y_1y_2) + 2|x|^2|y|^2 + |y|^4,$$

and (2.7), one has

$$\begin{aligned} \langle S_3vG_1vS_3f, f \rangle &= \langle vG_1vf, f \rangle \\ &= \int_{\mathbb{R}^2} \left(\int_{\mathbb{R}^2} |x - y|^4 v(y)f(y) dy \right) v(x) \overline{f(x)} dx \\ &= 4\langle x_1^2 v, f \rangle^2 + 4\langle x_2^2 v, f \rangle^2 + 8\langle x_1x_2 v, f \rangle^2. \end{aligned} \quad (2.22)$$

Combing (2.21) and (2.22), we have

$$0 = \langle T_3 f, f \rangle = \frac{\langle v, T_0 f \rangle^2}{\|V\|_{L^1}} + \frac{\|V\|_{L^1}}{1024} \langle x_1^2 v, f \rangle^2 + \frac{\|V\|_{L^1}}{1024} \langle x_2^2 v, f \rangle^2 + \frac{\|V\|_{L^1}}{512} \langle x_1 x_2 v, f \rangle^2 \geq 0,$$

which implies that $\langle v, T_0 f \rangle = 0$ and $\langle x_i x_j v, f \rangle = 0 (i, j = 1, 2)$. Note that, $\langle v, T_0 f \rangle = 0$ means $PT_0 f = 0$ by $P = \frac{\langle v, \cdot \rangle}{\|V\|_{L^1}} v$, we obtain that $T_0 f = PT_0 f + QT_0 f = 0$.

On the other hand, if $f \in S_3 L^2$ and $PT_0 f = 0$, it is easy to check that $f \in S_4 L^2$.

(vi) Let $f \in S_5 L^2$. We only need to prove $\langle x_i x_j x_k v, f \rangle = 0$, $i, j, k = 1, 2$. Indeed, notice that

$$\begin{aligned} |x - y|^6 = & |x|^6 + 3|x|^4|y|^2 - 6|x|^4x \cdot y + 12|x|^2(x \cdot y)^2 - 12|x|^2|y|^2x \cdot y - 8(x \cdot y)^3 \\ & + 12|y|^2(x \cdot y)^2 - 6|y|^4x \cdot y + 3|x|^2|y|^4 + |y|^6, \end{aligned}$$

by using $\langle v, f \rangle = \langle x_i v, f \rangle = \langle x_i x_j v, f \rangle = 0, 1 \leq i, j \leq 2$, we have

$$\begin{aligned} 0 = & \langle S_4 v G_2 v S_4 f, f \rangle = \langle v G_2 v f, f \rangle \\ = & -12 \int_{\mathbb{R}^2} v(x) \int_{\mathbb{R}^2} |x|^2 |y|^2 (x \cdot y) v(y) f(y) dy f(x) dx - 8 \int_{\mathbb{R}^2} v(x) \int_{\mathbb{R}^2} (x \cdot y)^3 v(y) f(y) dy f(x) dx. \end{aligned}$$

Note that $(x \cdot y)^2 = x_1^2 y_1^2 + 2x_1 x_2 y_1 y_2 + x_2^2 y_2^2$, $(x \cdot y)^3 = x_1^3 y_1^3 + 3x_1^2 y_1^2 x_2 y_2 + 3x_1 y_1 x_2^2 y_2^2 + x_2^3 y_2^3$, then

$$\begin{aligned} 0 = & -12 \sum_{j=1}^2 \left(\int_{\mathbb{R}^2} |x|^2 x_j v(x) f(x) dx \right)^2 - 8 \sum_{j=1}^2 \left(\int_{\mathbb{R}^2} x_j^3 v(x) f(x) dx \right)^2 \\ & - 24 \left(\int_{\mathbb{R}^2} x_1^2 x_2 v(x) f(x) dx \right)^2 - 24 \left(\int_{\mathbb{R}^2} x_1 x_2^2 v(x) f(x) dx \right)^2, \end{aligned}$$

which leads to $\langle x_i x_j x_k v, f \rangle = 0, 1 \leq i, j, k \leq 2$.

On the other hand, if $f \in S_4 L^2$ and $\langle x_i x_j x_k v, f \rangle = 0 (1 \leq i, j, k \leq 2)$, it is easy to check that $S_4 v G_2 v S_4 f = 0$.

The proof of this lemma is completed. \square

Throughout the paper, $\Gamma_{i,j}^k = \Gamma_{i,j}^k(\lambda)$ denote bounded operators that may change in each occurrence, and satisfy

$$\|\Gamma_{i,j}^k(\lambda)\|_{L^2 \rightarrow L^2} = O_1(1), \lambda \ll 1.$$

For the asymptotic expansions of $(M^\pm(\lambda))^{-1}$ in L^2 for λ near zero, we have the theorem as following.

Theorem 2.6. *Let $g_0^\pm(\lambda)$ and $\tilde{g}_2^\pm(\lambda)$ be functions defined in Lemma 2.1. Suppose that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with some $\beta > 0$. Then for $0 < \lambda \ll 1$, we have the conclusions as follows.*

(i) *If zero is a regular point and $\beta > 10$, then*

$$\begin{aligned} (M^\pm(\lambda))^{-1} = & S_0 D_1 S_0 + g_0^\pm(\lambda)^{-1} Q \Gamma_{0,1}^0 Q + g_0^\pm(\lambda)^2 \lambda^2 S_0 \Gamma_{2,1}^0 S_0 \\ & + g_0^\pm(\lambda) \lambda^2 \left(S_0 \Gamma_{2,2}^0 Q + Q \Gamma_{2,3}^0 S_0 \right) + O_1(\lambda^2). \end{aligned} \tag{2.23}$$

(ii) If zero is a resonance of the first kind and $\beta > 14$, then

$$\begin{aligned} (M^\pm(\lambda))^{-1} = & g_0^\pm(\lambda)S_1D_2S_1 + \left(S_0\Gamma_{0,1}^1S_0 + S_1\Gamma_{0,2}^1Q + Q\Gamma_{0,3}^1S_1\right) + g_0^\pm(\lambda)^{-1}Q\Gamma_{0,4}^1Q \\ & + g_0^\pm(\lambda)^4\lambda^2S_1\Gamma_{2,1}^1S_1 + g_0^\pm(\lambda)^3\lambda^2\left(S_1\Gamma_{2,2}^1Q + Q\Gamma_{2,3}^1S_1\right) + g_0^\pm(\lambda)^2\lambda^2 \\ & \left(Q\Gamma_{2,4}^1Q + S_1\Gamma_{2,5}^1 + \Gamma_{2,6}^1S_1\right) + g_0^\pm(\lambda)\lambda^2\left(Q\Gamma_{2,7}^1 + \Gamma_{2,8}^1Q\right) + O_1(\lambda^2). \end{aligned} \quad (2.24)$$

(iii) If zero is a resonance of the second kind and $\beta > 18$, then

$$\begin{aligned} (M^\pm(\lambda))^{-1} = & \frac{h^\pm(\lambda)}{\lambda^2}S_2\Gamma_{-2,1}^2S_2 + g_0^\pm(\lambda)^5\left(S_2\Gamma_{0,1}^2 + \Gamma_{0,2}^2S_2 + S_1\Gamma_{0,3}^2Q + Q\Gamma_{0,4}^2S_1\right) \\ & + g_0^\pm(\lambda)^{10}\lambda^2\left(S_2\Gamma_{2,1}^2 + \Gamma_{2,2}^2S_2 + S_1\Gamma_{2,3}^2Q + Q\Gamma_{2,4}^2S_1\right) \\ & + g_0^\pm(\lambda)\lambda^2\left(S_2\Gamma_{2,5}^2 + \Gamma_{2,6}^2S_2 + S_1\Gamma_{2,7}^2Q + Q\Gamma_{2,8}^2S_1\right) + O_1(\lambda^2), \end{aligned} \quad (2.25)$$

where $h^\pm(\lambda) = \left(c_1 + g_0^\pm(\lambda)^{-1}c_2 + g_0^\pm(\lambda)^{-2}c_3\right)^{-1}$, which c_1 and c_2 are constants defined in Lemma 4.6. Moreover, $h^\pm(\lambda) = O_1(1)$ when λ enough small.

(iv) If zero is a resonance of the third kind and $\beta > 18$, then

$$(M^\pm(\lambda))^{-1} = \frac{\tilde{g}_2^\pm(\lambda)^{-1}}{\lambda^4}S_4\Gamma_{-4,1}^3S_4 - \frac{\tilde{g}_2^\pm(\lambda)^{-2}}{\lambda^4}S_4\Gamma_{-4,2}^3S_4 + O_1(\lambda^{-4}(\ln \lambda)^{-3}). \quad (2.26)$$

(v) If zero is a resonance of the fourth kind and $\beta > 18$, then

$$\begin{aligned} (M^\pm(\lambda))^{-1} = & \frac{1}{\lambda^4}S_5D_6S_5 + \frac{\tilde{g}_2^\pm(\lambda)^{-1}}{\lambda^4}\left(S_4\Gamma_{-4,1}^4S_4\right) + \frac{\tilde{g}_2^\pm(\lambda)^{-2}}{\lambda^4}S_4\Gamma_{-4,2}^4S_4 \\ & + O_1(\lambda^{-4}(\ln \lambda)^{-3}). \end{aligned} \quad (2.27)$$

As we will see, since the degenerate term $g_0^\pm(\lambda)G_{-1}(x, y)$ with the logarithm factor occurs in the expansions of the free resolvent $R_0^\pm(\lambda^4)(x, y)$ in \mathbb{R}^2 , it leads to quite complicate calculations for the inversion expansion of $(M^\pm(\lambda))^{-1}$. We will give the proof of this theorem in Section 4.

3. THE PROOF OF THEOREM 1.1

In this section, we are devoted to proving Theorem 1.1. Choosing a fixed even function $\varphi \in C_c^\infty(\mathbb{R})$ such that $\varphi(s) = 1$ for $|s| \leq \frac{1}{2}$ and $\varphi(s) = 0$ for $|s| \geq 1$. Let $\varphi_N(s) = \varphi(2^{-N}s) - \varphi(2^{-N+1}s)$, $N \in \mathbb{Z}$. Then $\varphi_N(s) = \varphi_0(2^{-N}s)$, $\text{supp}\varphi_0 \subset [\frac{1}{4}, 1]$, and

$$\sum_{N=-\infty}^{\infty} \varphi_0(2^{-N}s) = 1, \quad s \in \mathbb{R} \setminus \{0\}.$$

By using the functional calculus and Stone's formula, one has

$$\begin{aligned}
 H^\alpha e^{-itH} P_{ac}(H) f &= \frac{2}{\pi i} \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} [R_V^+(\lambda^4) - R_V^-(\lambda^4)] f d\lambda \\
 &= \frac{2}{\pi i} \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} \sum_{N=-\infty}^\infty \varphi_0(2^{-N}\lambda) [R_V^+(\lambda^4) - R_V^-(\lambda^4)] f d\lambda \\
 &= \frac{2}{\pi i} \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} \chi(\lambda) [R_V^+(\lambda^4) - R_V^-(\lambda^4)] f d\lambda \\
 &\quad + \frac{2}{\pi i} \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} \tilde{\chi}(\lambda) [R_V^+(\lambda^4) - R_V^-(\lambda^4)] f d\lambda,
 \end{aligned} \tag{3.1}$$

where $\chi(\lambda) = \sum_{N=-\infty}^{N'} \varphi_0(2^{-N}\lambda)$ and $\tilde{\chi}(\lambda) = \sum_{N=N'+1}^{+\infty} \varphi_0(2^{-N}\lambda)$ for some $N' < 0$ (fixed later but depending on specific zero energy type).

In order to prove Theorem 1.1, it suffices to show the following two theorems.

Theorem 3.1. (low energy decay estimate)

Let $|V(x)| \lesssim (1+|x|)^{-\beta}$ with some $\beta > 0$. Assume that $H = \Delta^2 + V$ has no positive embedded eigenvalues and $P_{ac}(H)$ denote the projection onto absolutely continuous spectrum space of H . Then the following statements hold:

(i) If zero is a regular point and $\beta > 10$, then

$$\|e^{-itH} P_{ac}(H) \chi(H)\|_{L^1 \rightarrow L^\infty} \lesssim |t|^{-\frac{1}{2}}. \tag{3.2}$$

(ii) If zero is a resonance of the first kind and $\beta > 14$, then

$$\|e^{-itH} P_{ac}(H) \chi(H)\|_{L^1 \rightarrow L^\infty} \lesssim |t|^{-\frac{1}{2}}. \tag{3.3}$$

(iii) If zero is a resonance of the second kind and $\beta > 18$, then

$$\|H^{\frac{1}{2}} e^{-itH} P_{ac}(H) \chi(H)\|_{L^1 \rightarrow L^\infty} \lesssim |t|^{-\frac{1}{2}}. \tag{3.4}$$

(iv) If zero is a resonance of the third and fourth kind and $\beta > 18$, then

$$\|H^{\frac{3}{2}} e^{-itH} P_{ac}(H) \chi(H)\|_{L^1 \rightarrow L^\infty} \lesssim |t|^{-\frac{1}{2}}. \tag{3.5}$$

Theorem 3.2. (high energy decay estimate)

Let $|V(x)| \lesssim (1+|x|)^{-\beta}$ with $\beta > 5$. Assume that H has no positive eigenvalues and $P_{ac}(H)$ denote the projection onto absolutely continuous spectrum space of H . Then

$$\|e^{-itH} P_{ac}(H) \tilde{\chi}(\lambda)\|_{L^1 \rightarrow L^\infty} \lesssim |t|^{-\frac{1}{2}}. \tag{3.6}$$

The proofs of the low and high energy parts will be given in the sequent subsections. To proceed, let's first estimate the free group $e^{-it\Delta^2}$ in dimension two by resolvent methods, which displays the basic techniques how to prove the potential cases.

3.1. The decay estimates for the free case. We first give a lemma which will be used frequently later.

Lemma 3.3. *Let A be a subset of \mathbb{Z} . Suppose that $\Psi(s, z)$ is a function on $\mathbb{R} \times \mathbb{R}^m$ which is smooth for the first variable s , and satisfies for any s, z ,*

$$|\partial_s^k \Psi(2^N s, z)| \lesssim 1, \quad k = 0, 1, \quad N \in A \subset \mathbb{Z}.$$

Suppose that $\Phi(z)$ is a nonnegative real valued function of \mathbb{R}^m and $N_0 = \lceil \frac{1}{3} \log_2 \frac{\Phi(z)}{t} \rceil$. Then

$$\left| \int_0^\infty e^{-it2^{4N}s^4} e^{\pm i2^N s \Phi(z)} \Psi(2^N s, z) \varphi_0(s) ds \right| \lesssim \begin{cases} (1 + |t|2^{4N})^{-\frac{1}{2}}, & \text{if } |N - N_0| \leq 2, \\ (1 + |t|2^{4N})^{-1}, & \text{if } |N - N_0| > 2. \end{cases} \quad (3.7)$$

As a consequence, we also have

$$\sup_{z \in \mathbb{R}^m} \left| \int_0^\infty e^{-it2^{4N}s^4} e^{\pm i2^N s \Phi(z)} \Psi(2^N s, z) \varphi_0(s) ds \right| \lesssim (1 + |t|2^{4N})^{-\frac{1}{2}}. \quad (3.8)$$

Proof. Without loss of generality, we assume that $t \geq 0$. We write

$$K_N^\pm(t, z) = \int_0^\infty e^{-it2^{4N}s^4} e^{\pm i2^N s \Phi(z)} \Psi(2^N s, z) \varphi_0(s) ds.$$

Now we shall divide into $2^N \Phi(z) < 1$ and $2^N \Phi(z) \geq 1$ two cases to compute the integral above.

Case 1. If $2^N \Phi(z) < 1$, then we divide into $|t|2^{4N} < 1$ and $|t|2^{4N} \geq 1$ two cases again. For $|t|2^{4N} < 1$. Note that $s \in \text{supp} \varphi_0 \subset [\frac{1}{4}, 1]$, one has

$$|K_N^\pm(t, z)| \lesssim \int_{\frac{1}{4}}^1 \left| \Psi(2^N s, z) \varphi_0(s) \right| ds \lesssim (1 + |t| \cdot 2^{4N})^{-1}. \quad (3.9)$$

For $|t|2^{4N} \geq 1$. Since $\text{supp} \varphi_0 \in [\frac{1}{4}, 1]$, by integration by parts, one has

$$K_N^\pm(t, z) = \frac{1}{4it2^{4N}} \int_{\frac{1}{4}}^1 e^{-it2^{4N}s^4} \partial_s (s^{-3} \varphi_0(s) e^{\pm i2^N s \Phi(z)} \Psi(2^N s, z)) ds.$$

Notice that, we have for $k = 0, 1$,

$$|\partial_s^k (e^{\pm i2^N s \Phi(z)})| \lesssim 1, \quad |\partial_s^k \Psi(2^N s, z)| \lesssim 1.$$

Thus, we obtain that $|K_N^\pm(t, z)|$ is controlled by $(1 + |t|2^{4N})^{-1}$ uniformly in z .

Case 2. If $2^N \Phi(z) \geq 1$, then we consider $K_N^-(t, z)$ and $K_N^+(t, z)$ case by case. We begin to estimate $K_N^-(t, z)$. If $|t|2^{4N} < 1$, then we can obtain that $|K_N^-(t, z)|$ is bounded by $(1 + |t|2^{4N})^{-1}$ uniformly in z . If $|t|2^{4N} \geq 1$, then we write

$$K_N^-(t, z) = \int_{\frac{1}{4}}^1 e^{-iU_-(t, s, z, N)} V(s, z, N) ds.$$

where $U_-(t, s, z, N) = t2^{4N}s^4 + 2^N s \Phi(z)$ and $V(s, z, N) = \Psi(2^N s, z) \varphi_0(s)$. Since

$$|\partial_s U_-(t, s, z, N)| = |4t2^{4N}s^3 + 2^N \Phi(z)| \gtrsim 1 + |t|2^{4N}.$$

Thus, $U_-(t, s, z, N)$ doesn't have critical point. Using integration by parts we obtain that $|K_N^-(t, z)|$ is controlled by $(1 + |t| \cdot 2^{4N})^{-1}$ uniformly in z .

Next we turn to the estimate of $K_N^+(t, z)$. If $|t|2^{4N} < 1$, then it is clear that $|K_N^+(t, z)|$ is controlled by $(1 + |t| \cdot 2^{4N})^{-1}$ uniformly in z . If $|t|2^{4N} \geq 1$, set $U_+(t, s, z, N) = t2^{4N}s^4 - 2^N s\Phi(z)$ and $V(s, z, N) = \Psi(2^N s, z)\varphi_0(s)$, then

$$K_N^+(t, x, y) = \int_{\frac{1}{4}}^1 e^{-iU_+(t, s, z, N)} V(s, z, N) ds.$$

Since $\partial_s U_+(t, s, z, N) = 4t2^{4N}s^3 - 2^N\Phi(z)$, thus $U_+(t, s, z, N)$ may have a critical point s_0 which satisfies $2^N\Phi(z) = 4t2^{4N}s_0^3$.

If $|N - N_0| < 2$, then $-2 + N < \frac{1}{3}\log_2 \frac{\Phi(z)}{t} < 3 + N$, that is, $\frac{1}{2^6}t2^{4N} < 2^N\Phi(z) < 2^9t2^{4N}$, which implies the critical point s_0 exists. Since

$$|\partial_s^2 u_+(t, s, z, N)| \gtrsim |t|2^{4N}, \quad s \in [1/4, 1],$$

hence by Van der Corput lemma (see [Ste93], p.334), we obtain that

$$|K_N^+(t, z)| \lesssim (|t|2^{4N})^{-\frac{1}{2}} \left(V(1, z, N) + \int_{\frac{1}{4}}^1 |\partial_s V(s, z, N)| ds \right).$$

Note that $V(1, z, N) = 0$ and $|\partial_s V(s, z, N)| \lesssim 1$, we have

$$|K_N^+(t, x, y)| \lesssim (|t|2^{4N})^{-\frac{1}{2}} \lesssim (1 + |t|2^{4N})^{-\frac{1}{2}}.$$

If $|N - N_0| \geq 2$, then $2^N\Phi(z) \leq \frac{1}{2^6}t2^{4N}$ or $2^N\Phi(z) \geq 2^9t2^{4N}$, thus the critical point s_0 doesn't exist. Using integration by parts again, we obtain that $|K_N^+(t, z)|$ is controlled by $(1 + |t|2^{4N})^{-1}$.

Combining **Case 1** and **Case 2**, we immediately obtain that (3.7) holds.

Finally, by the proof of (3.7), we obtain that (3.8) is bounded by $(1 + |t|2^{4N})^{-\frac{1}{2}}$ uniformly in z . \square

Next we turn to the decay estimates of the free case. by using the functional calculus and Stone's formula, one has

$$e^{-it\Delta^2} f = \frac{2}{\pi i} \int_0^\infty e^{-it\lambda^4} \lambda^3 [R_0^+(\lambda^4) - R_0^-(\lambda^4)] f d\lambda.$$

Hence we need to analyze the free resolvent $R_0^\pm(\lambda^4)$. We split the free resolvent into low and high parts based on the size of $\lambda|x - y|$. In particular, we write

$$R_0^\pm(\lambda^4)(x, y) = \chi(\lambda|x - y|)R_0^\pm(\lambda^4)(x, y) + \tilde{\chi}(\lambda|x - y|)R_0^\pm(\lambda^4)(x, y).$$

Let

$$R_0^\pm(\lambda^4)(x, y) = \frac{e^{\pm i\lambda|x - y|}}{\lambda^2} \tilde{R}^\pm(\lambda|x - y|),$$

then we have by (2.6) and Lemma 2.1,

$$\begin{aligned} \tilde{R}^\pm(p) &= e^{\mp ip} \left[b_\pm + a_0 p^2 \ln p + \alpha_\pm p^2 + c_\pm p^4 + a_2 p^6 \ln p + \beta_\pm p^6 + O_1(p^{8-\epsilon}) \right] \chi(p) \\ &+ \frac{1}{8} \left(\pm i w_\pm(p) - i e^{\mp ip} e^{-p} w_+(ip) \right) \tilde{\chi}(p). \end{aligned} \quad (3.10)$$

Proposition 3.4. *Let $N_0 = \lceil \frac{1}{3} \log_2 \frac{|x-y|}{t} \rceil$ and $N \in \mathbb{Z}$. Then*

$$\left| \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) R_0^\pm(\lambda^4)(x, y) d\lambda \right| \lesssim 2^{2N} \begin{cases} (1 + |t|2^{4N})^{-\frac{1}{2}}, & \text{if } |N - N_0| \leq 2 \\ (1 + |t|2^{4N})^{-1}, & \text{if } |N - N_0| > 2. \end{cases}$$

As a consequence,

$$\sup_{x, y \in \mathbb{R}^2} \left| \int_0^\infty e^{-it\lambda^4} \lambda^3 R_0^\pm(\lambda^4)(x, y) d\lambda \right| \lesssim |t|^{-\frac{1}{2}}, \quad (3.11)$$

which immediately implies that $\|e^{-it\Delta^2}\|_{L^1 \rightarrow L^\infty} \lesssim |t|^{-\frac{1}{2}}$.

Proof. We write

$$K_{0,N}^\pm(t, x, y) = \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) R_0^\pm(\lambda^4)(x, y) d\lambda.$$

Set

$$R_0^\pm(\lambda^4)(x, y) = \frac{e^{\pm i\lambda|x-y|}}{\lambda^2} \tilde{R}(\lambda|x-y|).$$

Let $2^N s = \lambda$, then we have

$$\begin{aligned} K_{0,N}^\pm(t, x, y) &= \int_0^\infty e^{-it\lambda^3} \lambda \varphi_0(2^{-N}\lambda) e^{\pm i\lambda|x-y|} \tilde{R}^\pm(\lambda|x-y|) d\lambda \\ &= 2^{2N} \int_0^\infty e^{-it2^{4N}s^4} e^{\pm i2^N s|x-y|} s \varphi_0(s) \tilde{R}^\pm(2^N s|x-y|) ds. \end{aligned}$$

Note that $s \in \text{supp} \varphi_0 \subset [\frac{1}{4}, 1]$, by (3.10) it is easy check that for any x, y

$$|\partial_s^k \tilde{R}^\pm(2^N s|x-y|)| \lesssim 1, \quad k = 0, 1.$$

By Lemma 3.3 with $z = (x, y)$, $\Phi(z) = |x-y|$ and

$$\Psi(2^N s; z) = \tilde{R}^\pm(2^N s|x-y|),$$

we have $|K_{0,N}^\pm(t, x, y)|$ is controlled by $2^{2N}(1 + |t|2^{4N})^{-1/2}$ when $|N - N_0| \leq 2$, controlled by $2^{2N}(1 + |t|2^{4N})^{-1}$ when $|N - N_0| > 2$.

Note that $\sum_{N=-\infty}^\infty \varphi_0(2^{-N}s) = 1$, then we have

$$K_0^\pm(t, x, y) := \int_0^\infty e^{-it\lambda^4} \lambda^3 R_0^\pm(\lambda^4)(x, y) d\lambda = \sum_{N=-\infty}^{+\infty} K_{0,N}^\pm(t, x, y).$$

Since for $t \neq 0$, there exists a constant $N'_0 \in \mathbb{Z}$ such that $|t|2^{4N'_0} \sim 1$. Hence we have for x, y ,

$$\begin{aligned}
 |K_0^\pm(t, x, y)| &\leq \sum_{|N-N_0| \leq 2} 2^{2N} (1 + |t|2^{4N})^{-\frac{1}{2}} + \sum_{N=-\infty}^{+\infty} 2^{2N} (1 + |t|2^{4N})^{-1} \\
 &\lesssim \sum_{|N-N_0| \leq 2} |t|^{-\frac{1}{2}} + \sum_{N=-\infty}^{N'_0} 2^{2N} (1 + |t|2^{4N})^{-1} + \sum_{N=N'_0+1}^{+\infty} 2^{2N} (1 + |t|2^{4N})^{-1} \\
 &\lesssim |t|^{-\frac{1}{2}} + \sum_{N=-\infty}^{N'_0} 2^{2N} + |t|^{-1} \sum_{N=N'_0+1}^{+\infty} 2^{-2N} \\
 &\lesssim |t|^{-\frac{1}{2}}.
 \end{aligned}$$

Thus the proof of this lemma is completed. \square

3.2. Low energy decay estimates. In this section, we are devoted to proving low energy dispersive estimates (3.2)-(3.5) case by case. The following lemma plays an important role to make use of cancellations of projection operators S_j appeared in the asymptotic expansions of resolvent $R_V^\pm(z)$ near zero in order to obtain the low energy decay estimates for all cases.

Lemma 3.5. *Assume that $\lambda > 0$, $F(p) \in C^\infty(\mathbb{R})$, $\bar{y} = (-y_2, y_1) \in \mathbb{R}^2$. Let $w \equiv w(x) = \frac{x}{|x|}$ for $x \neq 0$ and $w(x) = 0$ for $x = 0$. For $\theta \in [0, 1]$, let $|y| \cos \alpha = \langle y, w(x - \theta y) \rangle$ where $\alpha \equiv \alpha(x, y, \theta)$ is interpreted as the angle between y and $x - \theta y$. Then*

(i) *We have*

$$F(\lambda|x - y|) = F(\lambda|x|) - \lambda|y| \int_0^1 F'(\lambda|x - \theta y|) \cos \alpha d\theta. \quad (3.12)$$

(ii) *If $F'(0) = 0$, then*

$$\begin{aligned}
 F(\lambda|x - y|) &= F(\lambda|x|) - \lambda F'(\lambda|x|) \langle w(x), y \rangle \\
 &\quad + \lambda^2 |y|^2 \int_0^1 (1 - \theta) \left(F''(\lambda|x - \theta y|) \cos^2 \alpha + \frac{F'(\lambda|x - \theta y|)}{\lambda|x - \theta y|} \sin^2 \alpha \right) d\theta.
 \end{aligned}$$

(iii) *If $F'(0) = F''(0) = 0$, then*

$$\begin{aligned}
 F(\lambda|x - y|) &= F(\lambda|x|) - \lambda F'(\lambda|x|) \langle w(x), y \rangle \\
 &\quad + \frac{\lambda^2}{2} \left(\frac{F'(\lambda|x|)}{\lambda|x|} \langle w(x), \bar{y} \rangle^2 + F''(\lambda|x|) \langle w(x), y \rangle^2 \right) + \frac{\lambda^3 |y|^3}{2} \int_0^1 (1 - \theta)^2 \\
 &\quad \left[\left(\frac{F'(\lambda|x - \theta y|)}{\lambda^2 |x - \theta y|^2} - \frac{F''(\lambda|x - \theta y|)}{\lambda|x - \theta y|} \right) 3 \cos \alpha \sin^2 \alpha - F'''(\lambda|x - \theta y|) \cos^3 \alpha \right] d\theta.
 \end{aligned}$$

Here $\langle w(x), \bar{y} \rangle^2 = |y|^2 - \langle w(x), y \rangle^2$.

(iv) If $F^{(k)}(0) = 0, k = 1, 2, 3$, then

$$\begin{aligned} F(\lambda|x-y|) &= F(\lambda|x|) - \lambda F'(\lambda|x|)\langle w(x), y \rangle + \frac{\lambda^2}{2} \left(\frac{F'(\lambda|x|)}{\lambda|x|} \langle w(x), \bar{y} \rangle^2 + F''(\lambda|x|)\langle w(x), y \rangle^2 \right) \\ &\quad + \frac{\lambda^3}{3!} \left[\left(\frac{F'(\lambda|x|)}{\lambda^2|x|^2} - \frac{F''(\lambda|x|)}{\lambda|x|} \right) 3\langle w(x), y \rangle \langle w(x), \bar{y} \rangle^2 - F'''(\lambda|x|)\langle w(x), y \rangle^3 \right] \\ &\quad + \frac{\lambda^4|y|^4}{3!} \int_0^1 (1-\theta)^3 \left[\left(\frac{F'(\lambda|x-\theta y|)}{\lambda^3|x-\theta y|^3} - \frac{F''(\lambda|x-\theta y|)}{\lambda^2|x-\theta y|^2} \right) (15 \cos^2 \alpha \sin^2 \alpha \right. \\ &\quad \left. - 3 \sin^2 \alpha) + \frac{F'''(\lambda|x-\theta y|)}{\lambda|x-\theta y|} 6 \cos^2 \alpha \sin^2 \alpha + F^{(4)}(\lambda|x-\theta y|) \cos^4 \alpha \right] d\theta. \end{aligned}$$

Proof. Let $G_\varepsilon(y) = F(\lambda\sqrt{\varepsilon^2 + |x-y|^2})$, $\varepsilon \neq 0$. Then $G_\varepsilon(y) \in C^{k+1}(\mathbb{R})$ for $\varepsilon \neq 0$ and $F(\lambda|x-y|) = \lim_{\varepsilon \rightarrow 0} G_\varepsilon(y)$. By Taylor expansions, we have

$$G_\varepsilon(y) = \sum_{|\alpha| < k} \frac{\partial^\alpha G_\varepsilon(0)}{\alpha!} y^\alpha + k \int_0^1 (1-\theta)^{k-1} \sum_{|\alpha|=k} \frac{\partial^\alpha G_\varepsilon(\theta y) y^\alpha}{\alpha!} d\theta. \quad (3.13)$$

(i) Observe that

$$\partial_{y_j} G_\varepsilon(y) = \frac{-\lambda(x_j - y_j)}{(\varepsilon^2 + |x-y|^2)^{\frac{1}{2}}} G'(\lambda\sqrt{\varepsilon^2 + |x-y|^2}), \quad j = 1, 2.$$

Since there exists a constants $C = C(\lambda, x, y)$ such that $|\partial_{y_i} G_\varepsilon(\theta y)| \leq C (i = 1, 2)$ for $0 \leq \theta \leq 1$ and $0 < \varepsilon \leq 1$, then by Lebesgue's dominated convergence theorem, we have for $x - \theta y \neq 0$,

$$\lim_{\varepsilon \rightarrow 0} \int_0^1 \partial_{y_i} G_\varepsilon(\theta y) d\theta = \int_0^1 \frac{-\lambda(x_j - \theta y_j)}{|x - \theta y|} F'(\lambda|x - \theta y|) d\theta, \quad j = 1, 2,$$

and $\lim_{\varepsilon \rightarrow 0} \int_0^1 \partial_{y_i} G_\varepsilon(\theta y) d\theta = 0, j = 1, 2$ for $x - \theta y = 0$. From Taylor expansions (3.13) with $k = 1$, we obtain that

$$F(\lambda|x-y|) = F(\lambda|x|) - \lambda|y| \int_0^1 F'(\lambda|x-\theta y|) \cos \alpha d\theta.$$

(ii) Note that

$$\begin{aligned} \partial_{y_j}^2 G_\varepsilon(y) &= \frac{\lambda^2 \varepsilon^2 + \lambda^2(|x-y|^2 - (x_j - y_j)^2)}{(\varepsilon^2 + |x-y|^2)} \frac{F'(\lambda\sqrt{\varepsilon^2 + |x-y|^2})}{\lambda(\varepsilon^2 + |x-y|^2)^{\frac{1}{2}}} \\ &\quad + \frac{\lambda^2(x_j - y_j)^2}{\varepsilon^2 + |x-y|^2} F''(\lambda\sqrt{\varepsilon^2 + |x-y|^2}), \quad j = 1, 2, \end{aligned}$$

$$\partial_{y_1 y_2}^2 G_\varepsilon(y) = \frac{\lambda^2(x_1 - y_1)(x_2 - y_2)}{\varepsilon^2 + |x-y|^2} \left(F''(\lambda\sqrt{\varepsilon^2 + |x-y|^2}) - \frac{F'(\lambda\sqrt{\varepsilon^2 + |x-y|^2})}{\lambda(\varepsilon^2 + |x-y|^2)^{\frac{1}{2}}} \right).$$

Hence, we have for $x \neq \theta y$,

$$\lim_{\varepsilon \rightarrow 0} \partial_{y_j}^2 G_\varepsilon(\theta y) = \lambda^2 \left(1 - \frac{(x_j - \theta y_j)^2}{|x - \theta y|^2} \right) \frac{F'(\lambda|x - \theta y|)}{\lambda|x - \theta y|} + \frac{\lambda^2(x_j - \theta y_j)^2}{|x - \theta y|^2} F''(\lambda|x - \theta y|), \quad j = 1, 2.$$

Note that $F'(0) = 0$, by Hospital's formula we have for $x = \theta y$,

$$\lim_{\varepsilon \rightarrow 0} \partial_{y_j}^2 G_\varepsilon(\theta y) = \lambda^2 \lim_{\varepsilon \rightarrow 0} \frac{F'(\lambda\varepsilon)}{\lambda\varepsilon} = \lambda^2 \lim_{\varepsilon \rightarrow 0} F''(\lambda\varepsilon) = \lambda^2 F''(0), \quad j = 1, 2.$$

Similarly, we obtain that

$$\lim_{\varepsilon \rightarrow 0} \partial_{y_1 y_2}^2 G_\varepsilon(\theta y) = \lambda^2 w(x_1 - \theta y_1) w(x_2 - \theta y_2) \left(F''(\lambda|x - \theta y|) - \frac{F'(\lambda|x - \theta y|)}{\lambda|x - \theta y|} \right).$$

It is easy to check that

$$\lim_{\varepsilon \rightarrow 0} \partial_{y_j} G_\varepsilon(0) = -\lambda w(x_j) F'(\lambda|x|), \quad j = 1, 2.$$

Note that $\sin^2 \alpha = 1 - |\langle w(y), w(x - \theta y) \rangle|^2 = |\langle w(\bar{y}), w(x - \theta y) \rangle|^2$, by Taylor's expansions and Lebesgue's dominated convergence theorem, we obtain that

$$\begin{aligned} F(\lambda|x - y|) &= F(\lambda|x|) - \lambda F'(\lambda|x|) \langle w(x), y \rangle \\ &\quad + \lambda^2 |y|^2 \int_0^1 (1 - \theta) \left(\frac{F'(\lambda|x - \theta y|)}{\lambda|x - \theta y|} \sin^2 \alpha + F''(\lambda|x - \theta y|) \cos^2 \alpha \right) d\theta. \end{aligned}$$

Similarly, we can prove that (iii) and (iv) hold. \square

Remark 3.6. To use Taylor's expansion above, we need to know some facts related to $R_0^\pm(\lambda)$.

(i) Let $R_0^\pm(\lambda^4)(x, y) = \frac{1}{\lambda^2} F^\pm(\lambda|x - y|)$. Then, by (2.6) and Lemma 2.1 we have

$$\begin{aligned} F^\pm(p) &= (b_\pm + a_0 p^2 \ln p + \alpha_\pm p^2 + c_\pm p^4 + a_2 p^6 \ln p + \beta_\pm p^6 + O_1(p^{8-\epsilon})) \chi(p) \\ &\quad + \left(\pm \frac{i}{8} e^{\pm ip} w_\pm(p) - \frac{i}{8} e^{-p} w_+(ip) \right) \tilde{\chi}(p). \end{aligned} \quad (3.14)$$

(ii) By (3.14) we have

$$[R_0^+(\lambda^4) - R_0^-(\lambda^4)](x, y) = \frac{1}{\lambda^2} (F^+(\lambda|x - y|) - F^-(\lambda|x - y|)) := \frac{1}{\lambda^2} \bar{F}(\lambda|x - y|),$$

where

$$\bar{F}(p) = \frac{i}{4} \left(1 - \frac{1}{4} p^2 + \frac{1}{64} p^4 - \frac{1}{234} p^6 + O(p^8) \right) \chi(p) + \frac{i}{8} (e^{ip} w_+(p) + e^{-ip} w_-(p)) \tilde{\chi}(p). \quad (3.15)$$

Moreover, $\bar{F}(p) \in C^\infty(\mathbb{R})$, $\bar{F}'(0) = 0$ and $\bar{F}''(0) \neq 0$.

(iii) Let $\tilde{F}(p) = \bar{F}(p) - \frac{i}{16} p^2$. Then we have

$$\tilde{F}(p) = \frac{i}{4} \left(1 + \frac{1}{64} p^4 - \frac{1}{234} p^6 + O(p^8) \right) \chi(p) + \frac{i}{8} (e^{ip} w_+(p) + e^{-ip} w_-(p) - \frac{p^2}{2}) \tilde{\chi}(p). \quad (3.16)$$

Moreover, $\tilde{F}(p) \in C^\infty(\mathbb{R})$, $\tilde{F}^{(k)}(0) = 0$ ($k = 1, 2, 3$), and $\tilde{F}^{(4)}(0) \neq 0$.

3.2.1. Regular case. In order to show Theorem 3.1 in regular case, recall that Stone's formula

$$\begin{aligned} H^\alpha e^{-itH} P_{ac}(H) \chi(H) f &= \frac{2}{\pi i} \int_0^\infty e^{-it\lambda^4} \chi(\lambda) \lambda^{3+4\alpha} [R_V^+(\lambda^4) - R_V^-(\lambda^4)] f d\lambda \\ &= \sum_{N=-\infty}^{N'} \sum_{\pm} \frac{\pm 2}{\pi i} \int_0^\infty e^{-it\lambda^4} \varphi_0(2^{-N}\lambda) \lambda^{3+4\alpha} R_V^\pm(\lambda^4) f d\lambda. \end{aligned} \quad (3.17)$$

If zero is a regular point, using (2.11) and (2.23), we have

$$\begin{aligned} R_V^\pm(\lambda^4) &= R_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)v(S_0D_1S_0)vR_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)v(g_0^\pm(\lambda)^{-1}Q\Gamma_{0,1}^0Q) \\ &\quad \times vR_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)v(g_0^\pm(\lambda)^2\lambda^2S_0\Gamma_{2,1}^0S_0)vR_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)v \\ &\quad \times (g_0^\pm(\lambda)\lambda^2(S_0\Gamma_{2,2}^0Q + Q\Gamma_{2,3}^0S_0))vR_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)(vO_1(\lambda^2)v)R_0^\pm(\lambda^4). \end{aligned} \quad (3.18)$$

Hence, to prove Theorem 3.1 in regular case, combining with the Proposition 3.4, by (3.17) and (3.18) it suffices to prove the following Propositions 3.7–3.10.

Proposition 3.7. *Assume that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 10$. If $N \in \mathbb{Z}_-$ and $N \leq N'$, then*

$$\sup_{x,y \in \mathbb{R}^2} \left| \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) [R_0^\pm(\lambda^4)v(S_0D_1S_0)vR_0^\pm(\lambda^4)](x,y)d\lambda \right| \lesssim 2^{2N}(1 + |t|2^{4N})^{-1/2}.$$

Proof. We write

$$K_{1,N}^{0,\pm}(t; x, y) = \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) [R_0^\pm(\lambda^4)vS_0D_1S_0vR_0^\pm(\lambda^4)](x, y)d\lambda.$$

Let $R_0^\pm(\lambda^4)(x, y) = \frac{1}{\lambda^2}F^\pm(\lambda|x - y|)$. Using the orthogonality $S_0v = 0$ and Lemma 3.5(i) we have

$$\begin{aligned} & [R_0^\pm(\lambda^4)vD_1S_0vR_0^\pm(\lambda^4)](x, y) \\ &= \frac{1}{\lambda^4} \int_{\mathbb{R}^4} F^\pm(\lambda|x - u_2|)[vS_0D_1S_0v](u_2, u_1)F^\pm(\lambda|y - u_1|)du_1du_2 \\ &= \frac{1}{\lambda^2} \int_{\mathbb{R}^4} \int_0^1 \int_0^1 (F^\pm)'(\lambda|x - \theta_2u_2|)(F^\pm)'(\lambda|y - \theta_1u_1|) \cos \alpha_2 \cos \alpha_1 d\theta_1 d\theta_2 \\ &\quad |u_1||u_2|[vS_0D_1S_0v](u_2, u_1)du_1du_2, \end{aligned}$$

where $\cos \alpha_1 = \cos \alpha(y, u_1, \theta_1)$ and $\cos \alpha_2 = \cos \alpha(x, u_2, \theta_2)$.

Thus, we have

$$\begin{aligned} K_{1,N}^{0,\pm}(t; x, y) &= \int_{\mathbb{R}^4} \int_0^1 \int_0^1 \left(\int_0^\infty e^{-it\lambda^4} \lambda \varphi_0(2^{-N}\lambda) (F^\pm)'(\lambda|x - \theta_2u_2|)(F^\pm)'(\lambda|y - \theta_1u_1|)d\lambda \right) \\ &\quad \cos \alpha_2 \cos \alpha_1 d\theta_1 d\theta_2 |u_1||u_2|[vS_0D_1S_0v](u_2, u_1)du_1du_2. \end{aligned}$$

Let

$$E_{1,N}^{0,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2) = \int_0^\infty e^{-it\lambda^4} \lambda \varphi_0(2^{-N}\lambda) (F^\pm)'(\lambda|x - \theta_2u_2|)(F^\pm)'(\lambda|y - \theta_1u_1|)d\lambda.$$

Then

$$\begin{aligned} |K_{1,N}^{0,\pm}(t; x, y)| &\lesssim \int_{\mathbb{R}^4} \int_0^1 \int_0^1 |E_{1,N}^{0,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2)| d\theta_1 d\theta_2 \\ &\quad |u_1||u_2|[vS_0D_1S_0v](u_2, u_1)|du_1du_2. \end{aligned} \quad (3.19)$$

Now we begin to estimate $E_{1,N}^{0,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2)$. In fact, let

$$(F^\pm)'(p) = e^{\pm ip}F_1^\pm(p), \quad p \geq 0. \quad (3.20)$$

By (3.14) we have

$$F_1^\pm(p) = \frac{1}{8} \left(-w_\pm(p) \pm iw'_\pm(p) + ie^{\mp ip} e^{-p} (w_+(ip) - w'_+(ip)) \right), \quad p \gg 1, \quad (3.21)$$

$$F_1^\pm(p) = e^{\mp ip} \left(2a_0 p \ln p + (a_0 + 2\alpha_\pm)p + 4c_\pm p^3 + 6a_2 p^5 \ln p + (a_2 + 6\beta_\pm)p^5 + O(p^{7-\epsilon}) \right), \quad p \ll 1. \quad (3.22)$$

Let $2^N s = \lambda$, we have

$$\begin{aligned} & E_{1,N}^{0,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2) \\ &= \int_0^\infty e^{-it\lambda^4} \lambda \varphi_0(2^{-N}\lambda) e^{\pm i\lambda|x-\theta_2 u_2|} e^{\pm i\lambda|y-\theta_1 u_1|} F_1^\pm(\lambda|x-\theta_2 u_2|) F_1^\pm(\lambda|y-\theta_1 u_1|) d\lambda \\ &= 2^{2N} \int_0^\infty e^{-it2^{4N}s^4} s \varphi_0(s) e^{\pm i2^N s(|x-\theta_2 u_2|+|y-\theta_1 u_1|)} F_1^\pm(2^N s|x-\theta_2 u_2|) F_1^\pm(2^N s|y-\theta_1 u_1|) ds. \end{aligned}$$

Note that $s \in [1/4, 1]$ and $N \in \mathbb{Z}^-$, by (3.21) and (3.22) it is easy to check that for $k = 0, 1$,

$$|\partial_s^k F_1^\pm(2^N s|x-\theta_2 u_2|)| \lesssim 1, \quad \text{and} \quad |\partial_s^k F_1^\pm(2^N s|y-\theta_1 u_1|)| \lesssim 1.$$

Let $z = (x, y, \theta_1, \theta_2, u_1, u_2)$, $\Phi(z) = |x-\theta_2 u_2| + |y-\theta_1 u_1|$ and

$$\Psi(2^N s, z) = F_1^\pm(2^N s|x-\theta_2 u_2|) F_1^\pm(2^N s|y-\theta_1 u_1|).$$

By Lemma 3.3, we immediately obtain that for any z ,

$$|E_{1,N}^{0,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2)| \lesssim 2^{2N} (1 + |t|2^{4N})^{-1/2}.$$

Finally, from (3.19), by Hölder inequality we obtain that for any x, y ,

$$\begin{aligned} |K_{1,N}^{0,\pm}(t; x, y)| &\lesssim 2^{2N} (1 + |t|2^{4N})^{-1/2} \|u_1 v(u_1)\|_{L_{u_1}^2} \|u_2 v(u_2)\|_{L_{u_2}^2} \|S_0 D_1 S_0\|_{L^2 \rightarrow L^2} \\ &\lesssim 2^{2N} (1 + |t|2^{4N})^{-1/2}. \end{aligned}$$

The proof of this proposition is completed. \square

Proposition 3.8. *Assume that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 10$. If $N \in \mathbb{Z}_-$ and $N \leq N'$, then*

$$\sup_{x,y} \left| \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) [R_0^\pm(\lambda^4) v(g_0^\pm(\lambda)^{-1} Q \Gamma_{0,1}^0(\lambda) Q) v R_0^\pm(\lambda^4)](x, y) d\lambda \right| \lesssim \frac{2^{2N}}{(1 + |t|2^{4N})^{\frac{1}{2}}}.$$

Proof. Denote

$$K_{2,N}^{0,\pm}(t; x, y) = \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) [R_0^\pm(\lambda^4) v(g_0^\pm(\lambda)^{-1} Q \Gamma_{0,1}^0(\lambda) Q) v R_0^\pm(\lambda^4)](x, y) d\lambda.$$

Let

$$R_0^\pm(\lambda^2)(x, y) = \frac{1}{\lambda^2} F^\pm(\lambda|x-y|).$$

Using the orthogonality $Qv = 0$, by Lemma 3.5(i) we have

$$\begin{aligned}
& [R_0^\pm(\lambda^4)v(g_0^\pm(\lambda)^{-1}Q\Gamma_{0,1}^0(\lambda)Q)vR_0^\pm(\lambda^4)](x, y) \\
&= \frac{g_0^\pm(\lambda)^{-1}}{\lambda^4} \int_{\mathbb{R}^4} F^\pm(\lambda|x - u_2|)[vQ\Gamma_{0,1}^0(\lambda)Qv](u_2, u_1)F^\pm(\lambda|y - u_1|)du_1du_2 \\
&= \frac{g_0^\pm(\lambda)^{-1}}{\lambda^2} \int_{\mathbb{R}^4} \int_0^1 \int_0^1 (F^\pm)'(\lambda|x - \theta_2u_2|)(F^\pm)'(\lambda|y - \theta_1u_1|) \cos \alpha_2 \cos \alpha_1 d\theta_1 d\theta_2 \\
& \quad |u_1||u_2|[vQ\Gamma_{0,1}^0(\lambda)Qv](u_2, u_1)du_1du_2.
\end{aligned}$$

Thus, by Fubini's theorem we have

$$\begin{aligned}
& K_{2,N}^{0,\pm}(t; x, y) \\
&= \int_0^1 \int_0^1 \left(\int_0^\infty e^{-it\lambda^4} \lambda \varphi_0(2^{-N}\lambda) g_0^\pm(\lambda)^{-1} \int_{\mathbb{R}^4} (F^\pm)'(\lambda|x - \theta_2u_2|)(F^\pm)'(\lambda|y - \theta_1u_1|) \right. \\
& \quad \left. |u_1||u_2|v(u_1)v(u_2)[Q\Gamma_{0,1}^0(\lambda)Q](u_2, u_1)du_1du_2d\lambda \right) \cos \alpha_2 \cos \alpha_1 d\theta_1 d\theta_2.
\end{aligned}$$

Let

$$\begin{aligned}
E_{2,N}^{0,\pm}(t; x, y, \theta_1, \theta_2) &= \int_0^\infty e^{-it\lambda^4} \lambda^{1+\alpha} \varphi_0(2^{-N}\lambda) g_0^\pm(\lambda)^{-1} \int_{\mathbb{R}^4} (F^\pm)'(\lambda|x - \theta_2u_2|) \\
& \quad (F^\pm)'(\lambda|y - \theta_1u_1|)|u_1||u_2|v(u_1)v(u_2)[Q\Gamma_{0,1}^0(\lambda)Q](u_2, u_1)du_1du_2d\lambda.
\end{aligned}$$

Thus, we have

$$|K_{2,N}^{0,\pm}(t; x, y)| \lesssim \int_0^1 \int_0^1 |E_{2,N}^{0,\pm}(t; x, y, \theta_1, \theta_2)| d\theta_1 d\theta_2 \quad (3.23)$$

If for any $x, y \in \mathbb{R}^2$, and $0 \leq \theta_i \leq 1, i = 1, 2$, then

$$|E_{2,N}^{0,\pm}(t; x, y, \theta_1, \theta_2)| \lesssim 2^{(2+\alpha)N} (1 + |t|2^{4N})^{-1/2}. \quad (3.24)$$

By (3.23) we immediately obtain that $|K_{2,N}^{0,\pm}(t; x, y)|$ is bounded by $2^{2N} (1 + |t|2^{4N})^{-1/2}$ uniformly in x, y .

Finally, it remains the proof of (3.24). Indeed, let

$$(F^\pm)'(p) = e^{\pm ip} F_1^\pm(p), \quad p \geq 0.$$

Set $2^N s = \lambda$, we have

$$\begin{aligned}
& E_{2,N}^{0,\pm}(t; x, y, \theta_1, \theta_2) \\
&= 2^{2N} \int_0^\infty e^{-it2^{2N}s^4} s \varphi_0(s) g_0^\pm(2^N s)^{-1} \int_{\mathbb{R}^4} e^{\pm i2^N s|x - \theta_2u_2|} e^{\pm i2^N s|y - \theta_1u_1|} F_1^\pm(2^N s|x - \theta_2u_2|) \\
& \quad F_1^\pm(2^N s|y - \theta_1u_1|)|u_1||u_2|v(u_1)v(u_2)[Q\Gamma_{0,1}^0(2^N s)Q](u_2, u_1)du_1du_2ds \\
&= 2^{2N} \int_0^\infty e^{-it2^{2N}s^4} e^{\pm i2^N s(|x|+|y|)} s \varphi_0(s) \left(g_0^\pm(2^N s)^{-1} \int_{\mathbb{R}^4} e^{\pm i2^N s(|x - \theta_2u_2| - |x|)} e^{\pm i2^N s(|y - \theta_1u_1| - |y|)} \right. \\
& \quad \left. F_1^\pm(2^N s|x - \theta_2u_2|) F_1^\pm(2^N s|y - \theta_1u_1|)|u_1||u_2|v(u_1)v(u_2)[Q\Gamma_{0,1}^0(2^N s)Q](u_2, u_1)du_1du_2 \right) ds.
\end{aligned}$$

Let $z = (x, y, \theta_1, \theta_1)$, $\Phi(z) = |x| + |y|$ and

$$\begin{aligned} \Psi(2^N s, z) &= g_0^\pm(2^N s)^{-1} \int_{\mathbb{R}^4} e^{\pm i 2^N s(|x - \theta_2 u_2| - |x|)} e^{\pm i 2^N s(|y - \theta_1 u_1| - |y|)} F_1^\pm(2^N s|x - \theta_2 u_2|) \\ &\quad F_1^\pm(2^N s|y - \theta_1 u_1|) |u_1| |u_2| v(u_1) v(u_2) [Q\Gamma_{0,1}^0(2^N s)Q](u_2, u_1) du_1 du_2. \end{aligned}$$

By (3.21) and (3.22) it is easy to check that for $k = 0, 1$,

$$\begin{aligned} |\partial_s^k F_1^\pm(2^N s|x - \theta_2 u_2|)| &\lesssim 1, \quad |\partial_s^k F_1^\pm(2^N s|y - \theta_1 u_1|)| \lesssim 1, \quad |\partial_s^k (g_0^\pm(2^N s)^{-1})| \lesssim 1. \\ |\partial_s^k e^{\pm i 2^N s(|x - \theta_2 u_2| - |x|)}| &\lesssim |u_2|^k, \quad |\partial_s^k e^{\pm i 2^N s(|y - \theta_1 u_1| - |y|)}| \lesssim |u_1|^k. \end{aligned}$$

Note that $\|\Gamma_{0,2}^0(\lambda)\|_{L^2 \rightarrow L^2} = O_1(1)$, by Hölder's inequality we have

$$|\Psi(2^N s, z)| \lesssim \|u_1 v(u_1)\|_{L_{u_1}^2} \|u_2 v(u_2)\|_{L_{u_2}^2}.$$

Since

$$2^N \|Q\partial_s \Gamma_{0,1}^0(2^N s)Q\|_{L^2 \rightarrow L^2} \lesssim 2^N \|\partial_s \Gamma_{0,1}^0(2^N s)\|_{L^2 \rightarrow L^2} \lesssim 2^N \cdot (2^N s)^{-1} \lesssim 1,$$

by Hölder's inequality, we have

$$\begin{aligned} |\partial_s \Psi(2^N s, z)| &\lesssim (\|u_1 v(u_1)\|_{L_{u_1}^2} \|u_2 v(u_2)\|_{L_{u_2}^2} + \| |u_1|^2 v(u_1) \|_{L_{u_1}^2} \|u_2 v(u_2)\|_{L_{u_2}^2} \\ &\quad + \|u_1 v(u_1)\|_{L_{u_1}^2} \| |u_2|^2 v(u_2) \|_{L_{u_2}^2}) \lesssim 1. \end{aligned}$$

By Lemma 3.3, we obtain that the inequality (3.24) holds. \square

By using the same method with the proof of Proposition 3.8, we obtain the following propositions.

Proposition 3.9. *Assume that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 10$. Let $\Omega^\pm(\lambda)$ be one of three operators $g_0^\pm(\lambda)^2 \lambda^2 S_0 \Gamma_{2,1}^0(\lambda) S_0$, $g_0^\pm(\lambda) \lambda^2 S_0 \Gamma_{2,2}^0(\lambda) Q$, $g_0^\pm(\lambda) \lambda^2 Q \Gamma_{2,3}^0(\lambda) S_0$. If $N \in \mathbb{Z}_-$ and $N \leq N'$, then*

$$\sup_{x,y} \left| \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) [R_0^\pm(\lambda^4) v \Omega^\pm(\lambda) v R_0^\pm(\lambda^4)](x, y) d\lambda \right| \lesssim 2^{2N} (1 + |t| 2^{4N})^{-1/2}.$$

Proposition 3.10. *Let $\Gamma^0(\lambda) = O_1(\lambda^2)$ be the error term of regular case in the expansions of $(M^\pm(\lambda))^{-1}$. Suppose that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 10$. If $N \in \mathbb{Z}_-$ and $N \leq N'$, then*

$$\sup_{x,y} \left| \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) [R_0^\pm(\lambda^4) v \Gamma^0(\lambda) v R_0^\pm(\lambda^4)](x, y) d\lambda \right| \lesssim 2^{2N} (1 + |t| 2^{4N})^{-1/2}.$$

Proof. According to Riesz's theorem, it suffices to prove that for any $f, g \in L^1$,

$$\left| \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) \langle [v \Gamma^0(\lambda) v](R_0^\pm(\lambda^4) f), (R_0^\pm(\lambda^4))^* g \rangle d\lambda \right|$$

is controlled by $2^{2N} (1 + |t| 2^{4N})^{-1/2} \|f\|_{L^1} \|g\|_{L^1}$. Thus, it is enough to show that the following kernel is bounded uniformly in x, y by $2^{2N} (1 + |t| 2^{4N})^{-1/2}$:

$$K_{3,N}^{0,\pm}(t; x, y) := \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) \langle [v \Gamma^0(\lambda) v](R_0^\pm(\lambda^4)(*, y))(\cdot), (R_0^\pm)^*(\lambda^4)(x, \cdot) \rangle d\lambda.$$

Let $R_0^\pm(\lambda^4)(x, y) = \frac{e^{\pm i\lambda|x-y|}}{\lambda^2} \tilde{R}^\pm(\lambda|x-y|)$. We have

$$\begin{aligned} & \langle [v\Gamma^0(\lambda)v](R_0^\pm(\lambda^4)(*, y))(\cdot), R_0^\mp(\lambda)(x, \cdot) \rangle \\ &= \frac{1}{\lambda^4} \langle [v\Gamma^0(\lambda)v](e^{\pm i\lambda|*-y|} \tilde{R}^\pm(\lambda|*-y|))(\cdot), (e^{\mp i\lambda|x-\cdot|} \tilde{R}^\mp(\lambda|x-\cdot|)) \rangle \\ &= \frac{1}{\lambda^4} e^{\pm i\lambda(|x|+|y|)} \langle [v\Gamma^0(\lambda)v](e^{\pm i\lambda(|*-y|-|y|)} \tilde{R}^\pm(\lambda|*-y|))(\cdot), (e^{\mp i\lambda(|x-\cdot|-|x|)} \tilde{R}^\mp(\lambda|x-\cdot|)) \rangle \\ &:= \frac{1}{\lambda^4} e^{\pm i\lambda(|x|+|y|)} E_{3,N}^{0,\pm}(\lambda; x, y). \end{aligned}$$

Let $2^N s = \lambda$, then

$$K_{3,N}^{0,\pm}(t; x, y) = \int_0^\infty e^{-it2^{4N}s^4} s^{-1} \varphi_0(s) e^{\pm i2^N s(|x|+|y|)} E_{3,N}^{0,\pm}(2^N s; x, y) ds.$$

By (3.10) it is easy to check that for $k = 0, 1$,

$$\begin{aligned} |\partial_s^k (e^{\pm i\lambda(|\cdot-y|-|y|)} \tilde{R}^\pm(\lambda|\cdot-y|))| &\lesssim \langle \cdot \rangle^k, \\ |\partial_s^k (e^{\pm i\lambda(|x-\cdot|-|x|)} \tilde{R}^\pm(\lambda|x-\cdot|))| &\lesssim \langle \cdot \rangle^k. \end{aligned}$$

Since $\|\Gamma^0(\lambda)\|_{L^2 \rightarrow L^2} = O_1(\lambda^2)$ (e.g. choosing $\lambda < 2^{N'}$ for some large $N' \in \mathbb{N}_-$), then we have

$$\|\partial_s^k (\Gamma^0(2^N s))\|_{L^2 \rightarrow L^2} \lesssim 2^{2N} s^{2-k}, \quad k = 0, 1.$$

Hence, we have

$$|\partial_s^k (E_{3,N}^{0,\pm}(2^N s; x, y))| \lesssim \|\partial_s^k (\Gamma^0(2^N s))\|_{L^2 \rightarrow L^2} \lesssim 2^{2N} s^{2-k}, \quad k = 0, 1.$$

Let $z = (x, y)$, $\Phi(z) = |x| + |y|$, and $\Psi(2^N s; z) = E_{3,N}^{0,\pm}(2^N s; x, y)$. Using Lemma 3.3 again, we obtain that $|K_{3,N}^{0,\pm}(t; x, y)|$ is bounded by $2^{2N}(1 + |t|2^{4N})^{-1/2}$ uniformly in x, y . \square

3.2.2. The first kind of resonance. If zero is a resonance of the first kind, by using (2.11) and (2.24) one has

$$\begin{aligned} R_V^\pm(\lambda^4) &= R_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)v(g_0^\pm(\lambda)S_1D_2S_1)vR_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)v(S_0\Gamma_{0,1}^1S_0 \\ &\quad + S_1\Gamma_{0,2}^1Q + Q\Gamma_{0,3}^1S_1)vR_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)v(g_0^\pm(\lambda)^{-1}Q\Gamma_{0,4}^1Q)vR_0^\pm(\lambda^4) \\ &\quad - R_0^\pm(\lambda^4)v(g_0^\pm(\lambda)^4\lambda^2S_1\Gamma_{2,1}^1S_1)vR_0^\pm(\lambda^4) - g_0^\pm(\lambda)^3\lambda^2R_0^\pm(\lambda^4)v(S_1\Gamma_{2,2}^1Q \\ &\quad + Q\Gamma_{2,3}^1S_1)vR_0^\pm(\lambda^4) - g_0^\pm(\lambda)^2\lambda^2R_0^\pm(\lambda^4)v(Q\Gamma_{2,4}^1Q + S_1\Gamma_{2,5}^1 + \Gamma_{2,6}^1(\lambda)S_1) \\ &\quad \times vR_0^\pm(\lambda^4) - g_0^\pm(\lambda)\lambda^2R_0^\pm(\lambda^4)v(Q\Gamma_{2,7}^1(\lambda) + \Gamma_{2,8}^1Q)vR_0^\pm(\lambda^4) \\ &\quad - R_0^\pm(\lambda^4)(vO_1(\lambda^2)v)R_0^\pm(\lambda^4). \end{aligned} \tag{3.25}$$

In order to prove Theorem 3.1(ii), comparing with the analysis of regular case, it is enough to consider the term

$$R_0^\pm(\lambda^4)v(g_0^\pm(\lambda)S_1D_2S_1)vR_0^\pm(\lambda^4) := \Omega_1^\pm(\lambda).$$

For this term, we will make use of the resolvent difference $R_V^+(\lambda^4) - R_V^-(\lambda^4)$ in the Stone's formula

$$e^{-itH} P_{ac}(H) \chi(H) f = \frac{2}{\pi i} \int_0^\infty e^{-it\lambda^4} \chi(\lambda) \lambda^3 [R_V^+(\lambda^4) - R_V^-(\lambda^4)] f d\lambda. \quad (3.26)$$

Since

$$\begin{aligned} \Omega_1^+(\lambda) - \Omega_1^-(\lambda) &= (R_0^+(\lambda^4) - R_0^-(\lambda^4)) v \left(g_0^+(\lambda) S_1 D_2 S_1 \right) v R_0^+(\lambda^4) \\ &\quad + R_0^-(\lambda^4) v \times \left(g_0^-(\lambda) S_1 D_2 S_1 \right) v (R_0^+(\lambda^4) - R_0^-(\lambda^4)) \\ &\quad + R_0^-(\lambda^4) v \left((g_0^+(\lambda) - g_0^-(\lambda)) S_1 D_2 S_1 \right) v R_0^+(\lambda^4) \\ &:= \Omega_{1,1}(\lambda) + \Omega_{1,2}(\lambda) + \Omega_{1,3}(\lambda). \end{aligned} \quad (3.27)$$

Hence, to prove Theorem 3.1(ii), we only need to prove the following proposition.

Proposition 3.11. *Let $\Omega_{1,i}(\lambda)$ ($i = 1, 2, 3$) be operators defined in (3.27). Assume that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 14$. If $N \in \mathbb{Z}_-$ and $N \leq N'$, then*

$$\sup_{x,y} \left| \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) \Omega_{1,i}(\lambda)(x, y) d\lambda \right| \lesssim 2^{2N} (1 + |t| 2^{4N})^{-1/2}, \quad i = 1, 2, 3.$$

Proof. We first prove the case of $\Omega_{1,1}(\lambda)$. Denote

$$K_{1,N}^{1,\pm}(t; x, y) = \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) [(R_0^+(\lambda^4) - R_0^-(\lambda^4)) v (g_0^\pm(\lambda) S_1 D_2 S_1) v R_0^+(\lambda^4)](x, y) d\lambda.$$

Let

$$R_0^+(\lambda^2)(x, y) = \frac{1}{\lambda^2} F^+(\lambda|x - y|), \quad \text{and} \quad [R_0^+(\lambda^4) - R_0^-(\lambda^4)](x, y) = \frac{1}{\lambda^2} \bar{F}(\lambda|x - y|).$$

By Remark 3.6, then $\bar{F}(p) \in C^3(\mathbb{R})$, $\bar{F}'(0) = 0$. Using the orthogonality $S_0 v = 0$ and $S_0(x_j v) = 0$ ($j = 1, 2$), by Lemma 3.5(i) and (ii) we have

$$\begin{aligned} &[(R_0^+(\lambda^4) - R_0^-(\lambda^4)) v (g_0^\pm(\lambda) S_1 D_2 S_1) v R_0^+(\lambda^4)](x, y) \\ &= \frac{g_0^\pm(\lambda)}{\lambda^4} \int_{\mathbb{R}^4} \bar{F}(\lambda|x - u_2|) [v S_1 D_2 S_1 v](u_2, u_1) F^+(\lambda|y - u_1|) du_1 du_2 \\ &= \frac{g_0^\pm(\lambda)}{\lambda} \int_{\mathbb{R}^4} \int_0^1 \int_0^1 \left(\bar{F}''(\lambda|x - \theta_2 u_2|) \cos^2 \alpha_2 + \frac{\bar{F}'(\lambda|x - \theta_2 u_2|)}{\lambda|x - \theta_2 u_2|} \sin^2 \alpha_2 \right) (F^+)'(\lambda|y - \theta_1 u_1|) \\ &\quad (1 - \theta_2) \cos \alpha_1 d\theta_1 d\theta_2 |u_1| |u_2|^2 [v S_1 D_2 S_1 v](u_2, u_1) du_1 du_2. \end{aligned}$$

Thus, we have

$$\begin{aligned} K_{1,N}^{1,\pm}(t; x, y) &= \int_{\mathbb{R}^4} \int_0^1 \int_0^1 \left(\int_0^\infty e^{-it\lambda^4} \varphi_0(2^{-N}\lambda) \lambda^2 g_0^\pm(\lambda) \left(\bar{F}''(\lambda|x - \theta_2 u_2|) \cos^2 \alpha_2 \right. \right. \\ &\quad \left. \left. + \frac{\bar{F}'(\lambda|x - \theta_2 u_2|)}{\lambda|x - \theta_2 u_2|} \sin^2 \alpha_2 \right) (F^+)'(\lambda|y - \theta_1 u_1|) d\lambda \right) (1 - \theta_2) \cos \alpha_1 d\theta_1 d\theta_2 \\ &\quad |u_1| |u_2|^2 [v S_1 D_2 S_1 v](u_2, u_1) du_1 du_2, \end{aligned}$$

Set

$$E_{1,N}^{1,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2) = \int_0^\infty e^{-it\lambda^4} \varphi_0(2^{-N}\lambda) \lambda^2 g_0^\pm(\lambda) \left(\bar{F}''(\lambda|x - \theta_2 u_2|) \cos^2 \alpha_2 + \frac{\bar{F}'(\lambda|x - \theta_2 u_2|)}{\lambda|x - \theta_2 u_2|} \sin^2 \alpha_2 \right) (F^+)''(\lambda|y - \theta_1 u_1|) d\lambda.$$

Then we have

$$|K_{1,N}^{1,\pm}(t; x, y)| \lesssim \int_{\mathbb{R}^4} \int_0^1 \int_0^1 |E_{1,N}^{1,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2)| (1 - \theta_2) d\theta_1 d\theta_2 |u_1| |u_2|^2 |v S_1 \Gamma_{0,1}^1 S_1 v|(u_2, u_1) du_1 du_2. \quad (3.28)$$

Now we begin to estimate $E_{1,N}^{1,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2)$. In fact, let

$$(F^+)''(p) = e^{ip} F_1^+(p), \quad \text{and} \quad \frac{\bar{F}'(p) \sin^2 \alpha_2}{p} + \bar{F}''(p) \cos^2 \alpha_2 = e^{ip} \bar{F}_1(p).$$

By (3.15) one has

$$\begin{aligned} \bar{F}_1(p) &= \frac{1}{8p} \left[-w_+(p) + iw'_+(p) + e^{-2ip}(w_-(p) + iw'_-(p)) \right] \sin^2 \alpha_2 + \left[\frac{-i}{8}(w_+(p) \right. \\ &\quad \left. + w_-(p)) + \frac{1}{4} e^{-2ip}(w'_+(p) + w'_-(p)) \right] \cos^2 \alpha_2, \quad p \gg 1, \end{aligned} \quad (3.29)$$

$$\begin{aligned} \bar{F}_1(p) &= e^{-ip} \left[\frac{i \sin^2 \alpha_2}{4} \left(-\frac{1}{2}p + \frac{1}{16}p^2 - \frac{1}{39}p^4 \right) + \frac{i \cos^2 \alpha_2}{4} \left(-p + \frac{3}{16}p^2 - \frac{5}{39}p^4 \right) \right. \\ &\quad \left. + O(p^6) \right], \quad p \ll 1. \end{aligned} \quad (3.30)$$

Let $2^N s = \lambda$, we have

$$\begin{aligned} &E_{1,N}^{1,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2) \\ &= \int_0^\infty e^{-it\lambda^4} \varphi_0(2^{-N}\lambda) \lambda^2 g_0^\pm(\lambda) e^{i\lambda|x - \theta_2 u_2|} e^{i\lambda|y - \theta_1 u_1|} \bar{F}_1(\lambda|x - \theta_2 u_2|) F_1^+(\lambda|y - \theta_1 u_1|) d\lambda \\ &= 2^{2N} \int_0^\infty e^{-it2^{4N}s^4} s \varphi_0(s) e^{i2^N s(|x - \theta_2 u_2| + |y - \theta_1 u_1|)} (2^N s) g_0^\pm(2^N s) \bar{F}_1(2^N s|x - \theta_2 u_2|) \\ &\quad F_1^+(2^N s|y - \theta_1 u_1|) ds. \end{aligned}$$

Let $z = (x, y, \theta_1, \theta_2, u_1, u_2)$, $\Phi(z) = |x - \theta_2 u_2| + |y - \theta_1 u_1|$, and

$$\Psi(2^N s, z) = (2^N s) g_0^\pm(2^N s) \bar{F}_1(2^N s|x - \theta_2 u_2|) F_1^+(2^N s|y - \theta_1 u_1|).$$

By (3.21), (3.22), (3.29) and (3.30), for any z it is easy check that

$$|\partial_s^k \Psi(2^N s, z)| \lesssim 1, \quad k = 0, 1.$$

By Lemma 3.3 again, we immediately obtain that

$$|E_{1,N}^{1,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2)| \lesssim 2^{2N} (1 + |t|2^{4N})^{-1/2}.$$

Finally, by (3.28) and Hölder's inequality we obtain that $|K_{1,N}^{1,\pm}(t; x, y)|$ is bounded by $2^{2N} (1 + |t|2^{4N})^{-1/2}$ uniformly in x, y .

For the case $\Omega_{1,2}(\lambda)$, similar to the case of $\Omega_{1,1}(\lambda)$, we obtain that the conclusion holds.

For the case $\Omega_{1,3}(\lambda)$, since $\Omega_{1,3}(\lambda) = -\frac{i}{16}R_0^-(\lambda^4)v\left(S_1D_2S_1\right)vR_0^+(\lambda^4)$, we obtain that the conclusion holds by the same argument with the proof of Proposition 3.7. \square

3.2.3. The second kind of resonance. If zero is the second kind of resonance of H , using (2.11) and (2.25) one has

$$\begin{aligned} R_V^\pm(\lambda^4) &= R_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)v\left(\frac{h^\pm(\lambda)}{\lambda^2}S_2\Gamma_{-2,1}^2S_2\right)vR_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)v\left(g_0^\pm(\lambda)^5(S_2\Gamma_{0,1}^2\right. \\ &\quad \left.+ \Gamma_{0,2}^2S_2 + S_1\Gamma_{0,3}^2Q + Q\Gamma_{0,4}^2S_1)\right)vR_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)v\left(g_0^\pm(\lambda)^{10}\lambda^2(S_2\Gamma_{2,1}^2\right. \\ &\quad \left.+ \Gamma_{2,2}^2S_2 + S_1\Gamma_{2,3}^2Q + Q\Gamma_{2,4}^2S_1)\right)vR_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)v\left(g_0^\pm(\lambda)\lambda^2(S_2\Gamma_{2,5}^2\right. \\ &\quad \left.+ \Gamma_{2,6}^2S_2 + S_1\Gamma_{2,7}^2Q + Q\Gamma_{2,8}^2S_1)\right)vR_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)\left(vO_1(\lambda^2)v\right)R_0^\pm(\lambda^4). \end{aligned}$$

In order to prove Theorem 3.1(iii), comparing with the proofs of regular case and the first resonance, it suffices to estimate the bound of the term

$$R_0^\pm(\lambda^4)v\left(\frac{h^\pm(\lambda)}{\lambda^2}S_2\Gamma_{-2,1}^2S_2\right)vR_0^\pm(\lambda^4) := \Omega_2^\pm(\lambda).$$

Since

$$\begin{aligned} \Omega_2^+(\lambda) - \Omega_2^-(\lambda) &= (R_0^+(\lambda^4) - R_0^-(\lambda^4))v\left(\frac{h^+(\lambda)}{\lambda^2}S_2\Gamma_{-2,1}^2(\lambda)S_2\right)vR_0^+(\lambda^4) \\ &\quad + R_0^-(\lambda^4)v\left(\frac{h^-(\lambda)}{\lambda^2}S_2\Gamma_{-2,1}^2(\lambda)S_2\right)v(R_0^+(\lambda^4) - R_0^-(\lambda^4)) \\ &\quad + R_0^-(\lambda^4)v\left(\frac{h^+(\lambda) - h^-(\lambda)}{\lambda^2}S_2\Gamma_{-2,1}^2(\lambda)S_2\right)vR_0^+(\lambda^4) \\ &:= \Omega_{2,1}(\lambda) + \Omega_{2,3}(\lambda) + \Omega_{2,3}(\lambda). \end{aligned} \tag{3.31}$$

Hence, it suffices to prove the following proposition.

Proposition 3.12. *Let $\Omega_{2,i}(\lambda)$ ($i = 1, 2, 3$) be operators defined in (3.31). Assume that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 18$. If $N \in \mathbb{Z}_-$ and $N \leq N'$, then*

$$\sup_{x,y} \left| \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} \varphi_0(2^{-N}\lambda) \Omega_{2,i}(\lambda)(x, y) d\lambda \right| \lesssim 2^{(1+4\alpha)N} (1 + |t|2^{4N})^{-1/2}, \quad i = 1, 2;$$

$$\sup_{x,y} \left| \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} \varphi_0(2^{-N}\lambda) \Omega_{2,3}(\lambda)(x, y) d\lambda \right| \lesssim 2^{4\alpha N} (1 + |t|2^{4N})^{-1/2}.$$

In particular, for $\alpha = \frac{1}{2}$,

$$\sup_{x,y} \left| \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} \Omega_{2,i}(\lambda)(x, y) d\lambda \right| \lesssim |t|^{-\frac{1}{2}}, \quad i = 1, 2, 3.$$

Proof. For the case of $\Omega_{2,1}(\lambda)$, we use the orthogonality $S_2v = S_2(x_1v) = S_2(x_2v) = 0$ for S_2 on the left hand of $\Omega_{2,1}(\lambda)$, use the orthogonality $S_2v = 0$ for S_2 on the right hand of $\Omega_{2,1}(\lambda)$, by the same argument with the proof of Proposition 3.11, it is easy to check that the statement holds. For the case of $\Omega_{2,2}(\lambda)$, the proof proceeds identically.

Next we turn to the case of $\Omega_{2,3}(\lambda)$, its proof is similar to the proof of Proposition 3.8. Denote

$$\begin{aligned} & K_{1,N}^{2,\pm}(t; x, y) \\ &= \frac{h^+(\lambda - h^-(\lambda))}{\lambda^2} \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} \varphi_0(2^{-N}\lambda) [R_0^-(\lambda^4)vS_2\Gamma_{-2,1}^2(\lambda)S_2vR_0^+(\lambda^4)](x, y)d\lambda. \end{aligned}$$

Let $R_0^\pm(\lambda^2)(x, y) = \frac{1}{\lambda^2}F^\pm(\lambda|x - y|)$. Using the orthogonality $S_2v = 0$, by Lemma 3.5(i) we have

$$\begin{aligned} & \frac{h^+(\lambda - h^-(\lambda))}{\lambda^2} [R_0^-(\lambda^4)vS_2\Gamma_{-2,1}^2(\lambda)S_2vR_0^+(\lambda^4)](x, y) \\ &= \frac{h^+(\lambda - h^-(\lambda))}{\lambda^6} \int_{\mathbb{R}^4} F^-(\lambda|x - u_2|)[vS_2\Gamma_{-2,1}^2(\lambda)S_2v](u_2, u_1)F^+(\lambda|y - u_1|)du_1du_2 \\ &= \frac{h^+(\lambda - h^-(\lambda))}{\lambda^4} \int_{\mathbb{R}^4} \int_0^1 \int_0^1 (F^-)'(\lambda|x - \theta_2u_2|)(F^+)'(\lambda|y - \theta_1u_1|) \cos \alpha_2 \cos \alpha_1 d\theta_1 d\theta_2 \\ & \quad |u_1||u_2|[vS_2\Gamma_{-2,1}^2(\lambda)S_2v](u_2, u_1)du_1du_2. \end{aligned}$$

Thus, we have

$$\begin{aligned} & K_{1,N}^{2,\pm}(t; x, y) \\ &= \int_0^1 \int_0^1 \left(\int_0^\infty e^{-it\lambda^4} \lambda^{4\alpha} \varphi_0(2^{-N}\lambda) \frac{h^+(\lambda - h^-(\lambda))}{\lambda} \int_{\mathbb{R}^4} (F^-)'(\lambda|x - \theta_2u_2|)(F^+)'(\lambda|y - \theta_1u_1|) \right. \\ & \quad \left. |u_1||u_2|v(u_1)v(u_2)[S_2\Gamma_{-2,1}^2(\lambda)S_2](u_2, u_1)du_1du_2d\lambda \right) \cos \alpha_2 \cos \alpha_1 d\theta_1 d\theta_2. \end{aligned}$$

Notice that $h^+(\lambda - h^-(\lambda)) = O_1((\ln \lambda)^{-1})$, we have

$$|K_{1,N}^{2,\pm}(t; x, y)| \lesssim \int_0^1 \int_0^1 |E_{1,N}^{2,\pm}(t; x, y, \theta_1, \theta_2)| d\theta_1 d\theta_2, \quad (3.32)$$

where

$$\begin{aligned} |E_{1,N}^{2,\pm}(t; x, y, \theta_1, \theta_2)| &= \int_0^\infty e^{-it\lambda^4} \lambda^{4\alpha} \varphi_0(2^{-N}\lambda) \frac{1}{\lambda \ln \lambda} \int_{\mathbb{R}^4} (F^-)'(\lambda|x - \theta_2u_2|) \\ & \quad (F^+)'(\lambda|y - \theta_1u_1|)|u_1||u_2|v(u_1)v(u_2)[S_2\Gamma_{-2,1}^2(\lambda)S_2](u_2, u_1)du_1du_2d\lambda. \end{aligned}$$

Suppose that for any $x, y \in \mathbb{R}^2$, and $0 \leq \theta_i \leq 1, i = 1, 2$, then

$$|E_{2,N}^{0,\pm}(t; x, y, \theta_1, \theta_2)| \lesssim 2^{4\alpha N} (1 + |t|2^{4N})^{-1/2}. \quad (3.33)$$

From (3.32), we immediately obtain that $|K_{2,N}^{0,\pm}(t; x, y)|$ is controlled by $2^{4\alpha N} (1 + |t|2^{4N})^{-1/2}$ uniformly in x, y .

Finally, it remains the proof of (3.33). Indeed, let

$$(F^\pm)'(p) = e^{\pm ip} F_1^\pm(p), \quad p \geq 0.$$

Set $2^N s = \lambda$, we have

$$\begin{aligned}
 & E_{1,N}^{2,\pm}(t; x, y, \theta_1, \theta_2) \\
 &= 2^{4\alpha N} \int_0^\infty e^{-it2^{2N}s^4} \varphi_0(s) s^{-1+4\alpha} (\ln 2^N s)^{-1} \int_{\mathbb{R}^4} e^{-i2^N s|x-\theta_2 u_2|} e^{i2^N s|y-\theta_1 u_1|} F_1^-(2^N s|x-\theta_2 u_2|) \\
 & \quad F_1^+(2^N s|y-\theta_1 u_1|) |u_1| |u_2| v(u_1) v(u_2) [S_2 \Gamma_{-2,1}^2(2^N s) S_2](u_2, u_1) du_1 du_2 ds \\
 &= 2^{4\alpha N} \int_0^\infty e^{-it2^{2N}s^4} e^{-i2^N s(|x|-|y|)} s^{-1+4\alpha} \varphi_0(s) \left((\ln 2^N s)^{-1} \int_{\mathbb{R}^4} e^{-i2^N s(|x-\theta_2 u_2|-|x|)} e^{i2^N s(|y-\theta_1 u_1|-|y|)} \right. \\
 & \quad \left. F_1^-(2^N s|x-\theta_2 u_2|) F_1^+(2^N s|y-\theta_1 u_1|) |u_1| |u_2| v(u_1) v(u_2) [S_2 \Gamma_{-2,1}^2(2^N s) S_2](u_2, u_1) du_1 du_2 \right) ds.
 \end{aligned}$$

Let $z = (x, y, \theta_1, \theta_2)$, $\Phi(z) = |x| - |y|$ and

$$\begin{aligned}
 \Psi(2^N s, z) &= (\ln 2^N s)^{-1} \int_{\mathbb{R}^4} e^{-i2^N s(|x-\theta_2 u_2|-|x|)} e^{i2^N s(|y-\theta_1 u_1|-|y|)} F_1^-(2^N s|x-\theta_2 u_2|) \\
 & \quad F_1^+(2^N s|y-\theta_1 u_1|) |u_1| |u_2| v(u_1) v(u_2) [S_2 \Gamma_{-2,1}^2(2^N s) S_2](u_2, u_1) du_1 du_2.
 \end{aligned}$$

By the same argument with the proof of Proposition 3.11, by Lemma 3.3 we obtain that the inequality (3.33) holds. \square

3.2.4. The third or fourth kind resonance. If zero is the third kind resonance of H , using (2.11) and (2.26) one has

$$\begin{aligned}
 R_V^\pm(\lambda^4) &= R_0^\pm(\lambda^4) - R_0^\pm(\lambda^4) v \left(\frac{\tilde{g}_2^\pm(\lambda)^{-1}}{\lambda^4} S_4 \Gamma_{-4,1}^3 S_4 \right) v R_0^\pm(\lambda^4) - R_0^\pm(\lambda^4) v \\
 & \quad \times \left(\frac{\tilde{g}_2^\pm(\lambda)^{-2}}{\lambda^4} S_4 \Gamma_{-4,1}^3 S_4 \right) v R_0^\pm(\lambda^4) - R_0^\pm(\lambda^4) \left(v O_1(\lambda^{-4}(\ln \lambda)^{-3}) v \right) R_0^\pm(\lambda^4).
 \end{aligned} \tag{3.34}$$

If zero is the fourth kind resonance of H , using (2.11) and (2.27) one has

$$\begin{aligned}
 & R_V^\pm(\lambda^4) \\
 &= R_0^\pm(\lambda^4) - R_0^\pm(\lambda^4) v \left(\frac{1}{\lambda^4} S_5 D_6 S_5 \right) v R_0^\pm(\lambda^4) - R_0^\pm(\lambda^4) v \left(\frac{\tilde{g}_2^+(\lambda)^{-1}}{\lambda^4} S_4 \Gamma_{-4,1}^4 S_4 \right) v R_0^\pm(\lambda^4) \\
 & \quad - R_0^\pm(\lambda^4) v \left(\frac{\tilde{g}_2^+(\lambda)^{-2}}{\lambda^4} S_4 \Gamma_{-4,2}^4 S_4 \right) v R_0^\pm(\lambda^4) - R_0^\pm(\lambda^4) \left(v O_1(\lambda^{-4}(\ln \lambda)^{-3}) v \right) R_0^\pm(\lambda^4).
 \end{aligned} \tag{3.35}$$

In order to prove Theorem 3.1(iv), we plug (3.34) and (3.35) into the Stone's formula (3.1). Combining with Proposition 3.4, it suffices to show the following Propositions 3.13, 3.14.

Proposition 3.13. *Let $\Omega_{3,1}^\pm(\lambda) := \frac{\tilde{g}_2^\pm(\lambda)^{-1}}{\lambda^4} S_4 \Gamma_{-4,1}^3 S_4$ and $\Omega_{4,1}^\pm(\lambda) := \frac{1}{\lambda^4} S_5 D_6 S_5$. Assume that $V(x) \lesssim (1 + |x|)^{-\beta}$ with $\beta > 18$. If $N \in \mathbb{Z}_-$ and $N \leq N'_0$, then for $i = 3, 4$,*

$$\sup_{x,y} \left| \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} \varphi_0(2^{-N}\lambda) [R_0^\pm(\lambda^4) v \Omega_{i,1}^\pm(\lambda) v R_0^\pm(\lambda^4)](x, y) d\lambda \right| \lesssim 2^{(-2+4\alpha)N} (1 + |t| 2^{4N})^{-1/2}.$$

In particular, for $\alpha = 1$,

$$\sup_{x,y} \left| \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} [R_0^\pm(\lambda^4) v \Omega_{i,1}^\pm(\lambda) v R_0^\pm(\lambda^4)](x, y) d\lambda \right| \lesssim |t|^{-1/2}, \quad i = 3, 4.$$

Proof. We first prove the case $\Omega_{3,1}^\pm(\lambda)$. Since

$$\begin{aligned} & R_0^+(\lambda^4)v\Omega_{3,1}^+(\lambda)vR_0^+(\lambda^4) - R_0^-(\lambda^4)v\Omega_{3,1}^+(\lambda)vR_0^-(\lambda^4) \\ &= (R_0^+(\lambda^4) - R_0^-(\lambda^4))v\left(\frac{\tilde{g}_2^+(\lambda)^{-1}}{\lambda^4}S_4\Gamma_{-4,1}^3S_4\right)vR_0^+(\lambda^4) + R_0^-(\lambda^4)v\left(\frac{\tilde{g}_2^-(\lambda)^{-1}}{\lambda^4}S_4\Gamma_{-4,1}^3S_4\right) \\ & \quad \times v(R_0^+(\lambda^4) - R_0^-(\lambda^4)) + R_0^-(\lambda^4)v\left(\frac{\tilde{g}_2^+(\lambda)^{-1} - \tilde{g}_2^-(\lambda)^{-1}}{\lambda^4}S_4\Gamma_{-4,1}^3S_4\right)vR_0^+(\lambda^4) \\ & := \Omega_{31}(\lambda) + \Omega_{32}(\lambda) + \Omega_{33}(\lambda). \end{aligned}$$

For the first term $\Omega_{31}(\lambda)$. We will use the orthogonality $S_4v = 0, S_4(x_iv) = S_4(x_ix_jv) = 0(i, j = 1, 2)$ for S_4 on the left right of $\Omega_{31}(\lambda)$, and the orthogonality $S_4v = 0$ for S_4 on the right hand of $\Omega_{31}(\lambda)$. We write

$$\begin{aligned} & K_{1,N}^{3,\pm}(t; x, y) \\ &= \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} \varphi_0(2^{-N}\lambda) \frac{\tilde{g}_2^+(\lambda)^{-1}}{\lambda^4} [(R_0^+(\lambda^4) - R_0^-(\lambda^4))vS_4\Gamma_{-4,1}^3S_4vR_0^+(\lambda^4)](x, y)d\lambda. \end{aligned}$$

Let

$$R_0^+(\lambda^2)(x, y) = \frac{1}{\lambda^2}F^+(\lambda|x - y|), \quad \text{and} \quad [R_0^+(\lambda^4) - R_0^-(\lambda^4)](x, y) = \frac{1}{\lambda^2}\bar{F}(\lambda|x - y|).$$

Using the orthogonality $S_4v = 0, S_4(x_iv) = S_4(x_ix_jv) = 0, i, j = 1, 2$, we have

$$\begin{aligned} & [(R_0^+(\lambda^4) - R_0^-(\lambda^4))vS_4\Gamma_{-4,1}^3S_4vR_0^+(\lambda^4)](x, y) \\ &= \frac{1}{\lambda^4} \int_{\mathbb{R}^4} \bar{F}(\lambda|x - u_2|)[vS_4\Gamma_{-4,1}^3S_4v](u_2, u_1)F^+(\lambda|y - u_1|)du_1du_2 \\ &= \frac{1}{\lambda^4} \int_{\mathbb{R}^4} (\bar{F}(\lambda|x - u_2|) - \frac{i}{16}\lambda^2|x - u_2|^2)[vS_4\Gamma_{-4,1}^3S_4v](u_2, u_1)F^+(\lambda|y - u_1|)du_1du_2 \\ &= \frac{1}{\lambda^4} \int_{\mathbb{R}^4} \tilde{F}(\lambda|x - u_2|)[vS_4\Gamma_{-4,1}^3S_4v](u_2, u_1)F^+(\lambda|y - u_1|)du_1du_2, \end{aligned}$$

where $\tilde{F}(p) = \bar{F}(p) - \frac{i}{16}p^2, p \in \mathbb{R}$.

By Remark 3.6, $\tilde{F}(p) \in C^4(\mathbb{R}), \tilde{F}^{(k)}(0) = 0, k = 1, 2, 3$. By the orthogonality $S_4v(x) = 0, S_4(x_jv) = S_4(x_jx_kv) = 0, j, k = 1, 2$, and Lemma 3.5(i) and (iii) we have

$$\begin{aligned} & [(R_0^+(\lambda^4) - R_0^-(\lambda^4))vS_4\Gamma_{-4,1}^3S_4vR_0^+(\lambda^4)](x, y) \\ &= \frac{1}{2} \int_{\mathbb{R}^4} \int_0^1 \int_0^1 \left[\left(\frac{\tilde{F}'(\lambda|x - \theta_2u_2|)}{\lambda^2|x - \theta_2u_2|^2} - \frac{\tilde{F}''(\lambda|x - \theta_2u_2|)}{\lambda|x - \theta_2u_2|} \right) 3 \cos \alpha_2 \sin^2 \alpha_2 \right. \\ & \quad \left. - \tilde{F}'''(\lambda|x - \theta_2u_2|) \cos^3 \alpha_2 \right] (F^+)'(\lambda|y - \theta_1u_1|)(1 - \theta_2)^2 \cos \alpha_1 d\theta_1 d\theta_2 \\ & \quad |u_1||u_2|^3 [vS_4\Gamma_{-4,1}^3S_4v](u_2, u_1) du_1 du_2. \end{aligned}$$

Thus, by Fubini's theorem we have

$$\begin{aligned}
 & K_{1,N}^{3,\pm}(t; x, y) \\
 &= \frac{1}{2} \int_{\mathbb{R}^4} \int_0^1 \int_0^1 \left(\int_0^\infty e^{-it\lambda^4} \varphi_0(2^{-N}\lambda) \lambda^{-1+4\alpha} \tilde{g}_2^+(\lambda)^{-1} \left[\left(\frac{\tilde{F}'(\lambda|x - \theta_2 u_2|)}{\lambda^2|x - \theta_2 u_2|^2} - \frac{\tilde{F}''(\lambda|x - \theta_2 u_2|)}{\lambda|x - \theta_2 u_2|} \right) \right. \right. \\
 &\quad \left. \left. \times 3 \cos \alpha_2 \sin^2 \alpha_2 - \tilde{F}'''(\lambda|x - \theta_2 u_2|) \cos^3 \alpha_2 \right] (F^+)'(\lambda|y - \theta_1 u_1|) d\lambda \right) (1 - \theta_2)^2 \cos \alpha_1 d\theta_1 d\theta_2 \\
 &\quad |u_1| |u_2|^3 [v S_4 \Gamma_{-4,1}^3 S_4 v](u_2, u_1) du_1 du_2.
 \end{aligned}$$

Let

$$\begin{aligned}
 & E_{1,N}^{3,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2) \\
 &= \int_0^\infty e^{-it\lambda^4} \varphi_0(2^{-N}\lambda) \lambda^{-1+4\alpha} \tilde{g}_2^+(\lambda)^{-1} \left[\left(\frac{\tilde{F}'(\lambda|x - \theta_2 u_2|)}{\lambda^2|x - \theta_2 u_2|^2} - \frac{\tilde{F}''(\lambda|x - \theta_2 u_2|)}{\lambda|x - \theta_2 u_2|} \right) 3 \cos \alpha_2 \sin^2 \alpha_2 \right. \\
 &\quad \left. - \tilde{F}'''(\lambda|x - \theta_2 u_2|) \cos^3 \alpha_2 \right] (F^+)'(\lambda|y - \theta_1 u_1|) d\lambda.
 \end{aligned}$$

Then we have

$$\begin{aligned}
 |K_{1,N}^{3,\pm}(t; x, y)| &\lesssim \int_{\mathbb{R}^4} \int_0^1 \int_0^1 |E_{1,N}^{3,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2)| (1 - \theta_2)^2 d\theta_1 d\theta_2 \\
 &\quad |u_1| |u_2|^3 [v S_4 \Gamma_{-4,1}^3 S_4 v](u_2, u_1) du_1 du_2.
 \end{aligned} \tag{3.36}$$

Now we begin to estimate $E_{1,N}^{3,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2)$. Indeed, we write

$$(F^+)'(p) = e^{ip} F_1^+(p), \quad \left(\frac{\tilde{F}'(p)}{p^2} - \frac{\tilde{F}''(p)}{p} \right) 3 \cos \alpha_2 \sin^2 \alpha_2 - \tilde{F}'''(p) \cos^3 \alpha_2 = e^{ip} \tilde{F}_1(p). \tag{3.37}$$

Let $2^N s = \lambda$, we have

$$\begin{aligned}
 E_{1,N}^{3,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2) &= 2^{4\alpha N} \int_0^\infty e^{-it2^{4N}s^4} \varphi_0(s) s^{-1+4\alpha} (\tilde{g}_0^\pm(2^N s))^{-1} e^{i2^N s(|x - \theta_2 u_2| + |y - \theta_1 u_1|)} \\
 &\quad \tilde{F}_1(2^N s|x - \theta_2 u_2|) F_1^+(2^N s|y - \theta_1 u_1|) ds.
 \end{aligned}$$

Let $z = (x, y, \theta_1, \theta_2, u_1, u_2)$, $\Phi(z) = |x - \theta_2 u_2| + |y - \theta_1 u_1|$, and

$$\Psi(2^N s, z) = (\tilde{g}_0^+(2^N s))^{-1} \tilde{F}_1(2^N s|x - \theta_2 u_2|) F_1^+(2^N s|y - \theta_1 u_1|).$$

Using (3.16), (3.21), (3.22) and (3.37), it is easy check that for any z ,

$$|\partial_s^k \Psi(2^N s, z)| \lesssim 1, \quad k = 0, 1.$$

Using Lemma 3.3, we immediately obtain that

$$|E_{1,N}^{3,\pm}(t; x, y, \theta_1, \theta_2, u_1, u_2)| \lesssim 2^{4\alpha N} (1 + |t|2^{4N})^{-1/2}.$$

By Hölder's inequality we obtain that $|K_{1,N}^{3,\pm}(t; x, y)|$ is bounded by $2^{4\alpha N} (1 + |t|2^{4N})^{-1/2}$ uniformly in x, y .

For the second term $\Omega_{32}(\lambda)$, the proof proceeds identically with $\Omega_{32}(\lambda)$.

For the last term $\Omega_{33}(\lambda)$. Denote

$$\begin{aligned} & K_{2,N}^{3,\pm}(t; x, y) \\ &= \frac{\tilde{g}_2^+(\lambda)^{-1} - \tilde{g}_2^-(\lambda)^{-1}}{\lambda^4} \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} \varphi_0(2^{-N}\lambda) [R_0^-(\lambda^4)vS_4\Gamma_{-4,1}^3 S_4vR_0^+(\lambda^4)](x, y)d\lambda \\ &= -\frac{i}{9216\lambda^4} \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} \varphi_0(2^{-N}\lambda) [R_0^-(\lambda^4)vS_4\Gamma_{-4,1}^3 S_4vR_0^+(\lambda^4)](x, y) \end{aligned}$$

Here we use the orthogonality $S_4v = 0$, by the same argument with the proof of Proposition 3.8, we get $|K_{2,N}^{3,\pm}(t; x, y)|$ is controlled by $2^{(-2+\alpha)N}(1 + |t|2^{4N})^{-1/2}$ uniformly in x, y .

Next we prove the case $\Omega_{4,1}^\pm(\lambda)$. Since

$$\begin{aligned} & R_0^+(\lambda^4)v\left(\frac{1}{\lambda^4}S_5D_6S_5\right)vR_0^+(\lambda^4) - R_0^-(\lambda^4)v\left(\frac{1}{\lambda^4}S_5D_6S_5\right)vR_0^-(\lambda^4) \\ &= (R_0^+(\lambda^4) - R_0^-(\lambda^4))v\left(\frac{1}{\lambda^4}S_5D_6S_5\right)vR_0^+(\lambda^4) \\ &+ R_0^-(\lambda^4)v\left(\frac{1}{\lambda^4}S_5D_6S_5\right)v(R_0^+(\lambda^4) - R_0^-(\lambda^4)), \end{aligned} \tag{3.38}$$

using the orthogonality

$$S_5v(x) = 0, S_5(x_iv) = S_5(x_ix_jv) = S_5(x_ix_jx_kv) = 0, i, j, k = 1, 2,$$

by the same argument with the proof of the case $\Omega_{3,1}^\pm(\lambda)$ we get the statement holds. \square

Proposition 3.14. *Let $\Gamma^3(\lambda) = O_1(\lambda^{-4}(\ln \lambda)^{-3})$ be the error term of the third or fourth kind resonance in the expansions of $(M^\pm(\lambda))^{-1}$. Assume that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 18$. If $N \in \mathbb{Z}_-$ and $N \leq N'$, then*

$$\sup_{x,y} \left| \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} \varphi_0(2^{-N}\lambda) [R_0^\pm(\lambda^4)v\Gamma^3(\lambda)vR_0^\pm(\lambda^4)](x, y)d\lambda \right| \lesssim 2^{(-4+4\alpha)N}(1 + |t|2^{4N})^{-1/2}.$$

In particular, for $\alpha = \frac{3}{2}$,

$$\sup_{x,y} \left| \int_0^\infty e^{-it\lambda^4} \lambda^{3+4\alpha} [R_0^\pm(\lambda^4)v\Gamma^3(\lambda)vR_0^\pm(\lambda^4)](x, y)d\lambda \right| \lesssim |t|^{-1/2}.$$

Proof. By the same argument with the proof of Proposition 3.10, we can prove that this proposition holds. \square

3.3. Large energy decay estimates. In this subsection, we will show Theorem 3.6. To complete the proof, we use the formula,

$$\begin{aligned} & e^{-itH}\tilde{\chi}(H)f(x) \\ &= \frac{2}{\pi i} \int_{\mathbb{R}^2} \left(\int_0^\infty e^{-it\lambda^4} \lambda^3 \tilde{\chi}(\lambda) [R_V^+(\lambda^4) - R_V^-(\lambda)](x, y)d\lambda \right) f(y)dy, \end{aligned} \tag{3.39}$$

and the resolvent identity,

$$R_V^\pm(\lambda^4) = R_0^\pm(\lambda^4) - R_0^\pm(\lambda^4)VR_0^\pm(\lambda^4) + R_0^\pm(\lambda^4)VR_V^\pm(\lambda^4)VR_0^\pm(\lambda^4). \tag{3.40}$$

Combing with Proposition 3.4, it is enough to prove the following Propositions 3.15, 3.17.

Proposition 3.15. *Assume that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 3$. Then*

$$\left\| \int_0^\infty e^{-it\lambda^4} \lambda^3 \tilde{\chi}(\lambda) R_0^\pm(\lambda^4) V R_0^\pm(\lambda^4) d\lambda \right\|_{L^1 \rightarrow L^\infty} \lesssim |t|^{-\frac{1}{2}}. \quad (3.41)$$

Proof. Let

$$L_{1,N}^\pm(t; x, y) = \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) [R_0^\pm(\lambda^4) V R_0^\pm(\lambda^4)](x, y) d\lambda,$$

and $R_0^\pm(\lambda^4)(x, y) := \frac{e^{\pm i\lambda|x-y|}}{\lambda^2} \tilde{R}^\pm(\lambda|x-y|)$. Set $2^N s = \lambda$, then we have

$$\begin{aligned} L_{1,N}^\pm(t; x, y) &= \int_{\mathbb{R}^2} \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) R_0^\pm(\lambda^4)(x, u_1) V(u_1) R_0^\pm(\lambda^4)(u_1, y) d\lambda du_1 \\ &= \int_{\mathbb{R}^2} \int_0^\infty e^{-it2^{4N}s^4} e^{\pm i2^N s(|x|+|y|)} s^{-1} \varphi_0(s) \left(e^{\pm i2^N s(|x-u_1|-|x|)} e^{\pm i2^N s(|y-u_1|-|y|)} \right. \\ &\quad \left. \tilde{R}^\pm(2^N|x-u_1|) V(u_1) \tilde{R}^\pm(2^N|x-u_1|) \right) d\lambda du_1. \end{aligned}$$

By (3.10), it is easy to check that for $k = 0, 1$,

$$|\partial_s^k e^{\pm i2^N s(|x-u_1|-|x|)}| \lesssim |u_1|^k, \quad |\partial_s^k e^{\pm i2^N s(|x-u_1|-|x|)}| \lesssim |u_1|^k,$$

$$|\partial_s^k (\tilde{R}^\pm(2^N|x-u_1|) V(u_1) \tilde{R}^\pm(2^N|x-u_1|))| \lesssim |V(u_1)|.$$

Let $z = (x, y, u_1)$, $\Phi(z) = |x| + |y|$ and

$$\Psi(2^N s; z) = e^{\pm i2^N s(|x-u_1|-|x|)} e^{\pm i2^N s(|y-u_1|-|y|)} \tilde{R}^\pm(2^N|x-u_1|) V(u_1) \tilde{R}^\pm(2^N|x-u_1|).$$

Let $N_0 = \lceil \frac{1}{3} \log_2 \frac{|x|+|y|}{t} \rceil$ and $N > N'$, by using Lemma 3.3, we immediately obtain that $|L_{1,N}^\pm(t; x, y)|$ is bounded by $2^{2N}(1+|t|2^{4N})^{-\frac{1}{2}}$ if $|N-N_0| \leq 2$, and bounded by $2^{2N}(1+|t|2^{4N})^{-1}$ if $|N-N_0| > 2$.

Note that $\sum_{N=-\infty}^\infty \varphi_0(2^{-N}s) = 1$, then we have

$$L_1^\pm(t; x, y) := \int_0^\infty e^{-it\lambda^4} \lambda^3 \tilde{\chi}(\lambda) [R_0^\pm(\lambda^4) V R_0^\pm(\lambda^4)](x, y) d\lambda = \sum_{N=N'+1}^{+\infty} L_{1,N}^\pm(t; x, y).$$

Since for $t \neq 0$, there exists a constant $N'_0 \in \mathbb{Z}$ such that $|t|2^{4N'_0} \sim 1$. Hence we have for x, y ,

$$\begin{aligned} |L_1^\pm(t; x, y)| &\leq \sum_{|N-N_0| \leq 2, N > N'} 2^{2N}(1+|t|2^{4N})^{-\frac{1}{2}} + \sum_{|N-N_0| > 2, N > N'} 2^{2N}(1+|t|2^{4N})^{-1} \\ &\lesssim \sum_{|N-N_0| \leq 2} |t|^{-\frac{1}{2}} + \sum_{N=-\infty}^{+\infty} 2^{2N}(1+|t|2^{4N})^{-1} \\ &\lesssim |t|^{-\frac{1}{2}}, \end{aligned}$$

which implies the conclusion (3.41). \square

Before we deal with the term $R_0^\pm(\lambda^4) V R_V^\pm(\lambda^4) V R_0^\pm(\lambda^4)$, we state a lemma as follows, see [FSY18].

Lemma 3.16. *Let $k \geq 0$ and $|V(x)| \lesssim (1 + |x|)^{-k-1-}$ such that $H = \Delta^2 + V$ has no embedded positive eigenvalues. Then for any $\sigma > k + \frac{1}{2}$, $R_V^\pm(\lambda) \in \mathcal{B}(L_\sigma^2(\mathbb{R}^d), L_{-\sigma}^2(\mathbb{R}^d))$ are C^k -continuous for all $\lambda > 0$. Furthermore,*

$$\|\partial_\lambda R_V^\pm(\lambda)\|_{L_\sigma^2(\mathbb{R}^2) \rightarrow L_{-\sigma}^2(\mathbb{R}^2)} = O(|\lambda|^{\frac{-3(k+1)}{4}}), \quad k = 0, 1, \quad \text{as } \lambda \rightarrow +\infty.$$

Proposition 3.17. *Assume that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 5$. Then*

$$\left\| \int_0^\infty e^{-it\lambda^4} \lambda^3 \tilde{\chi}(\lambda) R_0^\pm(\lambda^4) V R_V^\pm(\lambda^4) V R_0^\pm(\lambda^4) d\lambda \right\|_{L^1 \rightarrow L^\infty} \lesssim |t|^{-\frac{1}{2}}. \quad (3.42)$$

Proof. The proof is similar to the proof of Proposition 3.10. To prove (3.42), it is enough to prove that for any $f, g \in L^1$,

$$\left| \int_0^\infty e^{-it\lambda^4} \lambda^3 \tilde{\chi}(\lambda) \langle V R_V^\pm(\lambda^4) V (R_0^\pm(\lambda^4) f, (R_0^\pm)^*(\lambda^4) g) \rangle d\lambda \right|$$

is bounded $|t|^{-\frac{1}{2}} \|f\|_{L^1} \|g\|_{L^1}$. Thus, it suffices to show that

$$L_{2,N}^\pm(t; x, y) := \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) \langle V R_V^\pm(\lambda^4) V (R_0^\pm(\lambda^4)(*, y))(\cdot), (R_0^\pm(\lambda^4))^*(x, \cdot) \rangle d\lambda$$

is bounded by $2^{2N}(1 + |t|2^{4N})^{-1}$ if $|N - N_0| > 2$, bounded by $2^{2N}(1 + |t|2^{4N})^{-1/2}$ if $|N - N_0| \leq 2$, where $N_0 = \lceil \frac{1}{3} \log_2 \frac{|x| + |y|}{t} \rceil$ and $N > N'$.

Let $R_0^\pm(\lambda^4)(x, y) = \frac{e^{\pm i\lambda|x-y|}}{\lambda^2} \tilde{R}^\pm(\lambda|x-y|)$. Thus, we have

$$\begin{aligned} & \langle V R_V^\pm(\lambda^4) V (R_0^\pm(\lambda^4)(*, y))(\cdot), R_0^\mp(\lambda)(x, \cdot) \rangle \\ &= \frac{1}{\lambda^4} \langle V R_V^\pm(\lambda^4) V (e^{\pm i\lambda|*-y|} \tilde{R}^\pm(\lambda|*-y|))(\cdot), (e^{\mp i\lambda|x-\cdot|} \tilde{R}^\mp(\lambda|x-\cdot|)) \rangle \\ &= \frac{1}{\lambda^4} e^{\pm i\lambda(|x|+|y|)} \langle V R_V^\pm(\lambda^4) V (e^{\pm i\lambda(|*-y|-|y|)} \tilde{R}^\pm(\lambda|*-y|))(\cdot), (e^{\mp i\lambda(|x-\cdot|-|x|)} \tilde{R}^\mp(\lambda|x-\cdot|)) \rangle \\ &:= \frac{1}{\lambda^4} e^{\pm i\lambda(|x|+|y|)} E_{2,N}^\pm(\lambda; x, y). \end{aligned}$$

Let $2^N s = \lambda$, then

$$L_{2,N}^\pm(t; x, y) = \int_0^\infty e^{-it2^{4N}s^4} s^{-1} \varphi_0(s) e^{\pm i2^N s(|x|+|y|)} E_{2,N}^\pm(2^N s; x, y) ds.$$

By Lemma 3.16, for $\sigma > k + \frac{1}{2}$ we have

$$\|\partial_\lambda^k R_V^\pm(2^{4N}s^4)\|_{L_\sigma^2 \rightarrow L_{-\sigma}^2} \lesssim 2^{kN} (2^N s)^{-3}, \quad k = 0, 1.$$

Furthermore, Since $|V(x)| \lesssim (1 + |x|)^{-5-}$, we obtain that for $\sigma > k + \frac{1}{2}$,

$$\|\partial_\lambda E_{2,N}^\pm(2^N s; x, y)\| \lesssim \|V(\cdot) \langle \cdot \rangle^{\sigma+k}\|_{L^2}^2 \|\partial_\lambda^k R_V^\pm(2^{4N}s^4)\|_{L_\sigma^2 \rightarrow L_{-\sigma}^2} \lesssim 2^{kN} (2^N s)^{-3}, \quad k = 0, 1.$$

By (3.10), then for $k = 0, 1$,

$$|\partial_s^k (e^{\pm i\lambda(|\cdot|-y|-|y|)} \tilde{R}^\pm(\lambda|\cdot-y|))| \lesssim \langle \cdot \rangle^k, \quad |\partial_s^k (e^{\pm i\lambda(|x-\cdot|-|x|)} \tilde{R}^\pm(\lambda|x-\cdot|))| \lesssim \langle \cdot \rangle^k.$$

Let $N_0 = \lceil \frac{1}{3} \log_2 \frac{|x| + |y|}{t} \rceil$ and $N > N'$, using Lemma 3.3 with $z = (x, y)$,

$$\Phi(z) = |x| + |y|, \quad \Psi(2^N s; z) = E_{3,N}^{0,\pm}(2^N s; x, y),$$

we immediately obtain that for $|N - N_0| > 2$,

$$|L_{2,N}^\pm(t; x, y)| \lesssim 2^{(k-3)N}(1 + |t|2^{4N})^{-1} \lesssim 2^{2N}(1 + |t|2^{4N})^{-1},$$

and $|L_{2,N}^\pm(t; x, y)|$ is controlled by $2^{2N}(1 + |t|2^{4N})^{-1/2}$ for $|N - N_0| \leq 2$.

Note that $\sum_{N=-\infty}^{\infty} \varphi_0(2^{-N}s) = 1$, then we have

$$\begin{aligned} L_2^\pm(t; x, y) &:= \int_0^\infty e^{-it\lambda^4} \lambda^3 \varphi_0(2^{-N}\lambda) \langle V R_V^\pm(\lambda^4) V (R_0^\pm(\lambda^4) f, (R_0^\pm)^*(\lambda^4) g) \rangle d\lambda \\ &= \sum_{N=N'+1}^{+\infty} L_{2,N}^\pm(t; x, y). \end{aligned}$$

Hence we have for x, y ,

$$\begin{aligned} |L_2^\pm(t; x, y)| &\leq \sum_{|N-N_0| \leq 2, N > N'} 2^{2N}(1 + |t|2^{4N})^{-\frac{1}{2}} + \sum_{|N-N_0| > 2, N > N'} 2^{2N}(1 + |t|2^{4N})^{-1} \\ &\lesssim |t|^{-\frac{1}{2}}, \end{aligned}$$

which implies the conclusion (3.42). \square

4. THE PROOF OF THEOREM 2.6

In this section, we are devoted to showing Theorem 2.6, i.e. computing the expansions of $(M^\pm(\lambda))^{-1}$ for λ near zero case by case, its proof is quite complicated and long due to terms with logarithm in the expansions of the free resolvent $R_0^\pm(\lambda^4)$.

Before computing the expansions of $(M^\pm(\lambda))^{-1}$ as $\lambda \rightarrow 0$, we first state the following lemmas which are used frequently.

Lemma 4.1. ([JN01, Lemma 2.1]) *Let A be a closed operator and S be a projection. Suppose $A + S$ has a bounded inverse. Then A has a bounded inverse if and only if*

$$a := S - S(A + S)^{-1}S \tag{4.1}$$

has a bounded inverse in SH , and in this case

$$A^{-1} = (A + S)^{-1} + (A + S)^{-1}Sa^{-1}S(A + S)^{-1}. \tag{4.2}$$

Lemma 4.2. ([JN01, Lemma 2.3]) *Let A be an operator matrix on $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$:*

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, \quad a_{ij} : \mathcal{H}_j \rightarrow \mathcal{H}_i, 1 \leq i, j \leq 2,$$

where a_{11}, a_{22} are closed and a_{12}, a_{21} are bounded. Suppose a_{22} has a bounded inverse. Then A has a bounded inverse if and only if $d \equiv (a_{11} - a_{12}a_{22}^{-1}a_{21})^{-1}$ exist and is bounded. Furthermore, we have

$$A^{-1} = \begin{pmatrix} d & -da_{12}a_{22}^{-1} \\ -a_{22}^{-1}a_{21}d & a_{22}^{-1}a_{21}da_{12}a_{22}^{-1} + a_{22}^{-1} \end{pmatrix}.$$

Now we begin to compute the asymptotic expansions of $(M^\pm(\lambda))^{-1}$ for $\lambda \rightarrow 0$.

Let $M^\pm(\lambda) = \frac{a_\pm}{\lambda^2} \widetilde{M}^\pm(\lambda)$. Hence we need to compute the asymptotic expansion of $(\widetilde{M}^\pm(\lambda))^{-1}$ when λ near zero. By using Lemma 2.3, we have

$$\begin{aligned} \widetilde{M}^\pm(\lambda) = & P + \frac{g_0^\pm(\lambda)}{a_\pm} \lambda^2 v G_{-1} v + \frac{1}{a_\pm} \lambda^2 T_0 + \frac{c_\pm}{a_\pm} \lambda^4 v G_1 v \\ & + \frac{\tilde{g}_2^\pm(\lambda)}{a_\pm} \lambda^6 v G_2 v + \frac{\lambda^6}{a_\pm} v G_3 v + O(\lambda^8). \end{aligned} \quad (4.3)$$

Note that P is not invertible on L^2 , by Lemma 4.1 we know that $\widetilde{M}^\pm(\lambda)$ is invertible on L^2 if and only if

$$M_1^\pm(\lambda) := Q - Q(\widetilde{M}^\pm(\lambda) + Q)^{-1}Q$$

is invertible on QL^2 . Thus, we need to compute $(\widetilde{M}^\pm(\lambda) + Q)^{-1}$ on QL^2 . Let $Q = 1 - P$, then $\widetilde{M}^\pm(\lambda) + Q$ is invertible on QL^2 , and we have by Neumann series

$$\begin{aligned} & (\widetilde{M}^\pm(\lambda) + Q)^{-1} \\ = & I - g_0^\pm(\lambda) \lambda^2 B_1^\pm - \lambda^2 B_2^\pm - g_0^\pm(\lambda)^2 \lambda^4 B_3^\pm - g_0^\pm(\lambda) \lambda^4 B_4^\pm - \lambda^4 B_5^\pm - g_0^\pm(\lambda)^3 \lambda^6 B_6^\pm \\ & - g_0^\pm(\lambda)^2 \lambda^6 B_7^\pm - g_0^\pm(\lambda) \lambda^6 B_8^\pm - \tilde{g}_2^\pm(\lambda) \lambda^6 B_9^\pm - \lambda^6 B_{10}^\pm + O(\lambda^{8-\epsilon}), \end{aligned} \quad (4.4)$$

where $B_j^\pm (1 \leq j \leq 10)$ are bounded operators in L^2 as follows:

$$\begin{aligned} B_1^\pm &= \frac{1}{a_\pm} v G_{-1} v, \quad B_2^\pm = \frac{1}{a_\pm} T_0, \quad B_3^\pm = -\frac{1}{a_\pm^2} (v G_{-1} v)^2, \quad B_4^\pm = -\frac{1}{a_\pm^2} (v G_{-1} v T_0 + T_0 v G_{-1} v), \\ B_5^\pm &= \frac{c_\pm}{a_\pm} v G_1 v - \frac{1}{a_\pm^2} T_0^2, \quad B_6^\pm = \frac{1}{a_\pm^3} (v G_{-1} v)^3, \quad B_7^\pm = \frac{1}{a_\pm^3} [(v G_{-1} v)^2 T_0 + T_0 (v G_{-1} v)^2], \\ B_8^\pm &= \frac{1}{a_\pm^3} (v G_{-1} v T_0^2 + T_0^2 v G_{-1} v) - \frac{c_\pm}{a_\pm^2} (v G_{-1} v \cdot v G_1 v + v G_1 v \cdot v G_{-1} v), \\ B_9^\pm &= \frac{1}{a_\pm} v G_2 v, \quad B_{10}^\pm = \frac{1}{a_\pm} v G_3 v + \frac{1}{a_\pm^3} T_0^3 - \frac{c_\pm}{a_\pm^2} (T_0 v G_1 v + v G_1 v T_0). \end{aligned}$$

We list the orthogonality relations of various operators and projections which will be used frequently.

$$\begin{aligned} QD_0 &= D_0Q = D_0; \\ S_i D_j &= D_j S_i = S_i, i \geq j, \quad S_i D_j = D_j S_i = D_i, i < j, \quad D_i D_j = D_j D_i = D_i, i > j; \end{aligned} \quad (4.5)$$

$$\begin{aligned} QvG_{-1}vS_0 &= S_0QvG_{-1}vS_0 = 0, \quad S_1T_0S_0 = S_0T_0S_1 = 0, \quad S_2T_0Q = QT_0S_2 = 0; \\ S_3vG_{-1}v &= vG_{-1}vS_3 = 0, \quad S_4T_0P = PT_0S_4 = 0, \quad S_4T_0 = T_0S_4 = 0; \\ S_4vG_1vS_0 &= S_0vG_1vS_4 = 0, \quad S_5vG_1vQ = QvG_1vS_5 = 0. \end{aligned} \quad (4.6)$$

Moreover, we have

$$\begin{aligned} QB_1^\pm S_0 &= S_0 B_1^\pm Q = S_3 B_1^\pm = B_1^\pm S_3 = S_3 B_3^\pm = B_3^\pm S_3 = S_3 B_4^\pm S_3 = 0, \quad S_4 B_8^\pm S_4 = 0; \\ S_4 B_i^\pm &= B_i^\pm S_4 = 0 (i = 1, 2, 3, 4, 6, 7), \quad S_4 B_5^\pm S_0 = S_0 B_5^\pm S_4 = S_5 B_5^\pm Q = QB_5^\pm S_5 = 0. \end{aligned} \quad (4.7)$$

Now we turn to the proof of Theorem 2.6 case by case. We only prove the assertion for the case + sign, the case - sign proceeds identically. By Lemma 4.1, we know that $\widetilde{M}^\pm(\lambda)$ is

invertible on L^2 if and only if

$$M_1^+(\lambda) := Q - Q(\widetilde{M}^+(\lambda) + Q)^{-1}Q$$

is invertible on QL^2 . Moreover, by (4.4) we get

$$M_1^+(\lambda) := \frac{\lambda^2}{a_+} \widetilde{M}_1^+(\lambda) = \frac{\lambda^2}{a_+} \left(A^+(\lambda) + a_+ W^+(\lambda) \right), \quad (4.8)$$

where $A^+(\lambda) = g_0^+(\lambda)QvG_{-1}vQ + QT_0Q$, and

$$\begin{aligned} W^+(\lambda) = & g_0^+(\lambda)^2 \lambda^2 Q B_3^+ Q + g_0^+(\lambda) \lambda^2 Q B_4^+ Q + \lambda^2 Q B_5^+ Q + g_0^+(\lambda)^3 \lambda^4 Q B_6^+ Q \\ & + g_0^+(\lambda)^2 \lambda^4 Q B_7^+ Q + g_0^+(\lambda) \lambda^4 Q B_8^+ Q + \widetilde{g}_2^+(\lambda) \lambda^4 Q B_9^+ Q + \lambda^4 Q B_{10}^+ Q + O(\lambda^{6-\epsilon}). \end{aligned}$$

In order to compute $(M_1^+(\lambda))^{-1}$, by (4.8) we need to obtain the expansions of $(\widetilde{M}_1^+(\lambda))^{-1}$ on QL^2 when λ near zero. Note that $A^+(\lambda) = g_0^+(\lambda)QvG_{-1}vQ + QT_0Q$ with logarithm factor $g_0^+(\lambda)$, and $QvG_{-1}vQ$ is not invertible on QL^2 by Lemma 2.5(i), hence we will use Lemma 4.2 to consider the inverse of $A^+(\lambda)$.

To compute the inverse of $A^+(\lambda)$ by Lemma 4.2, we first give a lemma as follows.

Lemma 4.3. *Suppose that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with some $\beta > 10$. Then the inverse operator*

$$d = \left(g_0^+(\lambda)(Q - S_0)vG_{-1}v(Q - S_0) + (Q - S_0)(T_0 - T_0D_1T_0)(Q - S_0) \right)^{-1} \quad (4.9)$$

exists on $(Q - S_0)L^2(\mathbb{R}^2)$ for $\lambda \ll 1$, and

$$d = g_0^+(\lambda)^{-1}(Q - S_0)F(\lambda)(Q - S_0),$$

where $F(\lambda) = (A + g_0^+(\lambda)^{-1}B)^{-1}$ with $A = (Q - S_0)vG_{-1}v(Q - S_0)$ and $B = (Q - S_0)(T_0 - T_0D_1T_0)(Q - S_0)$.

Moreover, A and B are bounded operators on $(Q - S_0)L^2$ with $\text{rank}A = \text{rank}B = 2$, and $\|F(\lambda)\|_{(Q-S_0)L^2} = O_1(1)$ for small enough $\lambda > 0$.

Proof. Since $QL^2(\mathbb{R}^2) = \text{span}\{v\}^\perp$ and $S_0L^2(\mathbb{R}^2) = \text{span}\{v, x_1v(x), x_2v(x)\}^\perp$, then

$$(Q - S_0)L^2(\mathbb{R}^2) = \overline{\text{span}\{Q(x_1v), Q(x_2v)\}}.$$

Moreover, $\text{rank}(Q - S_0) = 2$. Since

$$\begin{aligned} d = & g_0^+(\lambda)^{-1}(Q - S_0) \left[(Q - S_0)vG_{-1}v(Q - S_0) \right. \\ & \left. + g_0^+(\lambda)^{-1}(Q - S_0)(T_0 - T_0D_1T_0)(Q - S_0) \right]^{-1} (Q - S_0), \end{aligned}$$

hence, in order to prove the operator d exists, we only need to prove that $(Q - S_0)vG_{-1}v(Q - S_0)$ is invertible on $(Q - S_0)L^2$ by Neumann series when λ is small enough.

Without loss of generality, we assume that Qx_1v and Qx_2v are orthogonal on $(Q - S_0)L^2$. If $f \in (Q - S_0)L^2$, then $f = a_1Qx_1v + a_2Qx_2v$. Since $(Q - S_0)(Q(x_i v)) = Q(x_i v)$ ($i = 1, 2$),

then we have

$$\begin{aligned} & (Q - S_0)vG_{-1}v(Q - S_0)(Q(x_1v))(x) = (Q - S_0)vG_{-1}v(Q(x_1v))(x) \\ & = (Q - S_0)v(x) \int_{\mathbb{R}^2} |x - y|^2 v(y) Q(x_1v)(y) dy \\ & = -2(Q - S_0)x_1v(x) \langle x_1v, Qx_1v \rangle - 2(Q - S_0)x_2v(x) \langle x_2v, Qx_1v \rangle. \end{aligned}$$

Since $\langle x_2v, Qx_1v \rangle = \langle Qx_2v, Qx_1v \rangle = 0$ and $(Q - S_0)x_iv(x) = Q(x_iv)(i = 1, 2)$, then we get

$$(Q - S_0)vG_{-1}v(Q - S_0)(Qx_1v)(x) = -2Q(x_1v)(x) \|Qx_1v\|_{L^2}^2.$$

Similarly, we have

$$(Q - S_0)vG_{-1}v(Q - S_0)(Qx_2v)(x) = -2Q(x_2v)(x) \|Qx_2v\|_{L^2}^2,$$

which leads to

$$\begin{aligned} & (Q - S_0)vG_{-1}v(Q - S_0)f = -2a_1Qx_1v \|Qx_1v\|_{L^2}^2 - 2a_2Qx_2v \|Qx_1v\|_{L^2}^2 \\ & = (f_1 \ f_2) \begin{pmatrix} -2\|Qx_1v\|_{L^2}^2 & 0 \\ 0 & -2\|Qx_2v\|_{L^2}^2 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} := (f_1 \ f_2) A \begin{pmatrix} a_1 \\ a_2 \end{pmatrix}. \end{aligned} \quad (4.10)$$

Since $\det A = 4\|Qx_1v\|_{L^2}^2 \|Qx_2v\|_{L^2}^2 \geq 0$, so we need only to prove $\det A \neq 0$. Indeed, if $\det A = 0$, then $Qx_1v = 0$ or $Qx_2v = 0$, it is contradictory with $\text{rank}(Q - S_0) = 2$. Thus, $(Q - S_0)vG_{-1}v(Q - S_0)$ is invertible and strictly positive on the space $(Q - S_0)L^2$, and then d exists on $(Q - S_0)L^2$ for small λ .

Let $A = (Q - S_0)vG_{-1}v(Q - S_0)$, $B = (Q - S_0)(T_0 - T_0D_1T_0)(Q - S_0)$ and $F(\lambda) = [A + g_0^+(\lambda)^{-1}B]^{-1}$, we obtain that

$$d = g_0^+(\lambda)^{-1}(Q - S_0)F(\lambda)(Q - S_0).$$

Finally, we need to prove that $\|F(\lambda)\|_{(Q-S_0)L^2} = O_1(1)$ for enough small λ . Indeed, let $G(\lambda) = A + g_0^+(\lambda)^{-1}B$, then $F(\lambda) = G(\lambda)^{-1}$. Note that $G(\lambda)' = -\lambda^{-1}g_0^+(\lambda)^{-2}$, we have

$$\frac{d}{d\lambda}F(\lambda) = \frac{d}{d\lambda}(G(\lambda)^{-1}) = -\left(A + g_0^+(\lambda)^{-1}B\right)^{-1} \cdot \lambda^{-1}g_0^+(\lambda)^{-2} \cdot \left(A + g_0^+(\lambda)^{-1}B\right)^{-1}.$$

Observe that

$$\|F(\lambda)\|_{(Q-S_0)L^2 \rightarrow (Q-S_0)L^2} \leq C, \quad \lambda \ll 1,$$

then

$$\left\| \frac{d}{d\lambda}F(\lambda) \right\|_{(Q-S_0)L^2 \rightarrow (Q-S_0)L^2} \lesssim \lambda^{-1}(\ln \lambda)^{-2} \lesssim \lambda^{-1}, \quad \lambda \ll 1,$$

Hence, we have $\|F(\lambda)\|_{(Q-S_0)L^2 \rightarrow (Q-S_0)L^2} = O_1(1)$ for enough small λ . \square

Now we begin to compute the inverse of $A^+(\lambda)$, we have the following lemma.

Lemma 4.4. *Suppose that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with some $\beta > 10$. If zero is a regular point of the spectrum of H , then for $0 < |\lambda| \ll 1$,*

$$(A^+(\lambda))^{-1} = D_1 + g_0^+(\lambda)^{-1}S(\lambda), \quad (4.11)$$

where

$$S(\lambda) = S = \begin{pmatrix} (Q - S_0)F(\lambda)(Q - S_0) & -(Q - S_0)F(\lambda)(Q - S_0)T_0D_1 \\ -D_1T_0(Q - S_0)F(\lambda)(Q - S_0) & D_1T_0(Q - S_0)F(\lambda)(Q - S_0)T_0D_1 \end{pmatrix}$$

with $F(\lambda) = (A + g_0^+(\lambda)^{-1}B)^{-1}$ and $D_1 = (S_0T_0S_0)^{-1}$. Moreover, $\|S(\lambda)\|_{L^2 \rightarrow L^2} = O_1(1)$ for enough small λ .

Moreover, if zero is not regular point, the same formula (4.11) holds for $(A^+(\lambda) + S_1)^{-1}$ with $D_1 = (S_0T_0S_0 + S_1)^{-1}$.

Proof. We prove only the statement when $S_1 \neq 0$, the proof of the regular case is identical. Observe that

$$\begin{aligned} A^+(\lambda) + S_1 &= g_0^+(\lambda)(Q - S_0)vG_{-1}v(Q - S_0) + (Q - S_0)T_0(Q - S_0) + (Q - S_0)T_0S_0 \\ &\quad + S_0T_0(Q - S_0) + S_0T_0S_0 + S_1, \end{aligned}$$

by block format, we write $A^+(\lambda) + S_1$ as follows:

$$A^+(\lambda) + S_1 = \begin{pmatrix} g_0^+(\lambda)(Q - S_0)vG_{-1}v(Q - S_0) + (Q - S_0)T_0(Q - S_0) & (Q - S_0)T_0S_0 \\ S_0T_0(Q - S_0) & S_0T_0S_0 + S_1 \end{pmatrix}.$$

Note that $QL^2 = (Q - S_0)L^2 \oplus S_0L^2$, and $S_0T_0S_0 + S_1$ is invertible on S_0L^2 , by Lemma 4.2, then $A^+(\lambda) + S_1$ has a bounded inverse on S_0L^2 if and only if d defined in (4.9) exists on $(Q - S_0)L^2$ and is bounded for enough small λ .

Let $D_1 = (S_0T_0S_0 + S_1)^{-1} : S_0L^2 \rightarrow S_0L^2$, by Lemma 4.3, we know that the operator d exists on $(Q - S_0)L^2$ and

$$d = g_0^+(\lambda)^{-1}(Q - S_0)F(\lambda)(Q - S_0).$$

Using Lemma 4.2 again, we have $A^+(\lambda) + S_1$ is invertible on S_0L^2 , and

$$\begin{aligned} &(A^+(\lambda) + S_1)^{-1} \\ &= g_0^+(\lambda)^{-1} \begin{pmatrix} (Q - S_0)F(\lambda)(Q - S_0) & -(Q - S_0)F(\lambda)(Q - S_0)T_0D_1 \\ -D_1T_0(Q - S_0)F(\lambda)(Q - S_0) & D_1T_0(Q - S_0)F(\lambda)(Q - S_0)T_0D_1 + g_0^+(\lambda)D_1 \end{pmatrix} \\ &:= D_1 + g_0^+(\lambda)^{-1}S(\lambda), \end{aligned}$$

where

$$S(\lambda) = S = \begin{pmatrix} (Q - S_0)F(\lambda)(Q - S_0) & -(Q - S_0)F(\lambda)(Q - S_0)T_0D_1 \\ -D_1T_0(Q - S_0)F(\lambda)(Q - S_0) & D_1T_0(Q - S_0)F(\lambda)(Q - S_0)T_0D_1 \end{pmatrix}.$$

It is obvious $\|S(\lambda)\|_{L^2 \rightarrow L^2} = O_1(1)$ for $\lambda \ll 1$. \square

Remark 4.5. We list some orthogonal relations as follows:

$$S_1S = -S_1T_0(Q - S_0)F(\lambda)(Q - S_0) + S_1T_0(Q - S_0)F(\lambda)(Q - S_0)T_0D_1, \quad (4.12)$$

$$SS_1 = -(Q - S_0)F(\lambda)(Q - S_0)T_0S_1 + D_1T_0(Q - S_0)F(\lambda)(Q - S_0)T_0S_1, \quad (4.13)$$

$$S_1SS_1 = S_1T_0(Q - S_0)F(\lambda)(Q - S_0)T_0S_1, \quad (4.14)$$

$$S_2T_0(Q - S_0) = (Q - S_0)T_0S_2 = 0, \quad (4.15)$$

$$S_2S = SS_2 = 0, \quad S_2(A^+(\lambda) + S_1)^{-1} = (A^+(\lambda) + S_1)^{-1}S_2 = S_2. \quad (4.16)$$

(I) Regular point case. If zero is a regular point of the spectrum of H , then $S_0T_0S_0$ is invertible on S_0L^2 . Note that the identity

$$\left(\widetilde{M}_1^+(\lambda)\right)^{-1} = \left(A^+(\lambda)\right)^{-1} \left(I + a_+W^+(\lambda)\left(A^+(\lambda)\right)^{-1}\right)^{-1},$$

by Lemma 4.4 and Neumann series, we have

$$\begin{aligned} \left(\widetilde{M}_1^+(\lambda)\right)^{-1} &= D_1 + g_0^+(\lambda)^{-1}S - a_+g_0^+(\lambda)^2\lambda^2D_1B_3^+D_1 \\ &\quad - a_+g_0^+(\lambda)\lambda^2(D_1B_4^+D_1 + D_1B_3^+QS + SQB_3^+D_1) + O_1(\lambda^2). \end{aligned}$$

Using (4.8), then $(M_1^+(\lambda))^{-1} = \frac{a_+}{\lambda^2}(\widetilde{M}_1^+(\lambda))^{-1}$. Hence, we have

$$\begin{aligned} (M_1^+(\lambda))^{-1} &= \frac{a_+}{\lambda^2}D_1 + \frac{a_+g_0^+(\lambda)^{-1}}{\lambda^2}S - a_+^2g_0^+(\lambda)^2D_1B_3^+D_1 - a_+^2g_0^+(\lambda)(D_1B_4^+D_1 \\ &\quad + D_1B_3^+QS + SQB_3^+D_1) + O_1(1). \end{aligned}$$

Using the orthogonality $QD_1 = D_1Q = D_1$, by (4.4) and Lemma 4.1, we get

$$\begin{aligned} \left(\widetilde{M}^+(\lambda)\right)^{-1} &= \frac{a_+}{\lambda^2}D_1 + \frac{a_+g_0^+(\lambda)^{-1}}{\lambda^2}QSQ - a_+^2g_0^+(\lambda)^2D_1B_3^+D_1 \\ &\quad - a_+^2g_0^+(\lambda)(D_1B_4^+D_1 + D_1B_3^+QSQ + QSQB_3^+D_1) + O_1(1). \end{aligned}$$

Since $(M^+(\lambda))^{-1} = \frac{\lambda^2}{a_+}(\widetilde{M}^+(\lambda))^{-1}$, using the orthogonality $S_0D_1 = D_1S_0 = D_1$, we have

$$\begin{aligned} (M^+(\lambda))^{-1} &= S_0D_1S_0 + g_0^+(\lambda)^{-1}Q\Gamma_{0,1}^0Q + g_0^+(\lambda)^2\lambda^2S_0\Gamma_{2,1}^0S_0 \\ &\quad + g_0^+(\lambda)\lambda^2\left(S_0\Gamma_{2,2}^0Q + Q\Gamma_{2,3}^0S_0\right) + O_1(\lambda^2). \end{aligned}$$

The expansions of $(M^+(\lambda))^{-1}$ in regular case is completed. \square

If zero is not regular point of the spectrum of H . Let S_1 be the Riesz projection onto the kernel of $S_0T_0S_0$. Then it is easy to check that $S_0T_0S_0 + S_1$ is invertible on S_0L^2 . In this case, we define $D_1 = (S_0T_0S_0 + S_1)^{-1}$ as a bounded operator on S_0L^2 . From (4.8), by Lemma 4.4 and Neumann series one has

$$\begin{aligned} &\left(\widetilde{M}_1^+(\lambda) + S_1\right)^{-1} \\ &= D_1 + g_0^+(\lambda)^{-1}S - g_0^+(\lambda)^2\lambda^2\widetilde{B}_1 - g_0^+(\lambda)\lambda^2\widetilde{B}_2 - \lambda^2\widetilde{B}_3 - g_0^+(\lambda)^{-1}\lambda^2\widetilde{B}_4 \\ &\quad - g_0^+(\lambda)^{-2}\lambda^2\widetilde{B}_5 - g_0^+(\lambda)^4\lambda^4\widetilde{B}_6 - g_0^+(\lambda)^3\lambda^4\widetilde{B}_7 - g_0^+(\lambda)^2\lambda^4\widetilde{B}_8 - g_0^+(\lambda)\lambda^4\widetilde{B}_9 \\ &\quad - \widetilde{g}_2^+(\lambda)\lambda^4\widetilde{B}_{10} - \lambda^4\widetilde{B}_{11} - \widetilde{g}_2^+(\lambda)g_0^+(\lambda)^{-1}\lambda^4\widetilde{B}_{12} - g_0^+(\lambda)^{-1}\lambda^4\widetilde{B}_{13} \\ &\quad - \widetilde{g}_2^+(\lambda)g_0^+(\lambda)^{-2}\lambda^4\widetilde{B}_{14} - g_0^+(\lambda)^{-2}\lambda^4\widetilde{B}_{15} - g_0^+(\lambda)^{-3}\lambda^4\widetilde{B}_{16} + O(\lambda^{6-\epsilon}), \end{aligned} \tag{4.17}$$

where $\widetilde{B}_j(1 \leq j \leq 15)$ are bounded operators on L^2 as follows:

$$\begin{aligned} \widetilde{B}_1 &= a_+D_1B_3D_1, \widetilde{B}_2 = a_+(D_1B_4D_1 + D_1B_3QS + SQB_3D_1), \\ \widetilde{B}_3 &= a_+[D_1B_5D_1 + D_1B_4QS + SQB_4D_1 + SQB_3QS], \\ \widetilde{B}_4 &= a_+[D_1B_5QS + SQB_5D_1 + SQB_4QS], \widetilde{B}_5 = a_+SQB_5QS, \widetilde{B}_6 = -a_+^2D_1B_3D_1B_3D_1, \end{aligned}$$

$$\begin{aligned}
 \tilde{B}_7 &= a_+ D_1 B_6 D_1 - a_+^2 [D_1 B_3 D_1 B_4 D_1 + D_1 B_4 D_1 B_3 D_1 + D_1 B_3 D_1 B_3 Q S + S Q B_3 D_1 B_3 D_1 \\
 &\quad + D_1 B_3 Q S Q B_3 D_1], \\
 \tilde{B}_8 &= a_+ [D_1 B_7 D_1 + D_1 B_6 Q S + S Q B_6 D_1] - a_+^2 [D_1 B_4 D_1 B_4 D_1 \\
 &\quad + D_1 B_3 D_1 B_5 D_1 + D_1 B_5 D_1 B_3 D_1 + D_1 B_3 D_1 B_4 Q S + S Q B_4 D_1 B_3 D_1 \\
 &\quad + D_1 B_4 D_1 B_3 Q S + S Q B_3 D_1 B_4 D_1 + D_1 B_3 Q S Q B_4 D_1 + D_1 B_4 Q S Q B_3 D_1 \\
 &\quad + D_1 B_3 Q S Q B_3 Q S + S Q B_3 Q S Q B_3 D_1], \\
 \tilde{B}_9 &= a_+ [D_1 B_8 D_1 + D_1 B_7 Q S + S Q B_7 D_1 + S Q B_6 Q S] - a_+^2 [D_1 B_4 D_1 B_5 D_1 + D_1 B_5 D_1 B_4 D_1 \\
 &\quad + D_1 B_3 D_1 B_5 Q S + S Q B_5 D_1 B_3 D_1 + S Q B_3 D_1 B_4 Q S + S Q B_4 D_1 B_3 Q S \\
 &\quad + S Q B_3 D_1 B_5 D_1 + D_1 B_5 D_1 B_3 Q S + D_1 B_3 Q S Q B_5 D_1 + D_1 B_5 Q S Q B_3 D_1 \\
 &\quad + D_1 B_3 Q S Q B_4 Q S + S Q B_4 Q S Q B_3 D_1 + D_1 B_4 Q S Q B_3 Q S + S Q B_3 Q S Q B_4 D_1 \\
 &\quad + D_1 B_4 Q S Q B_4 D_1 + S Q B_3 Q S Q B_3 Q S], \tilde{B}_{10} = a_+ D_1 B_9 D_1, \\
 \tilde{B}_{11} &= a_+ [D_1 B_{10} D_1 + D_1 B_8 Q S + Q S B_8 D_1 + S Q B_7 Q S] - a_+^2 [D_1 B_5 D_1 B_5 D_1 \\
 &\quad + D_1 B_5 D_1 B_4 Q S + S Q B_4 D_1 B_5 D_1 + D_1 B_4 D_1 B_5 Q S + S Q B_5 D_1 B_4 D_1 \\
 &\quad + D_1 B_4 Q S Q B_5 D_1 + D_1 B_5 Q S Q B_4 D_1 + D_1 B_3 Q S Q B_5 Q S + S Q B_5 Q S Q B_3 D_1 \\
 &\quad + D_1 B_4 Q S Q B_4 Q S + Q S B_4 Q S Q B_4 D_1 + D_1 B_5 Q S Q B_3 Q S + S Q B_3 Q S Q B_5 D_1 \\
 &\quad + S Q B_3 D_1 B_5 Q S + S Q B_5 D_1 B_3 Q S + S Q B_3 Q S Q B_4 Q S + S Q B_4 Q S Q B_3 Q S \\
 &\quad + S Q B_4 D_1 B_4 Q S], \tilde{B}_{12} = a_+ [D_1 B_9 Q S + S Q B_9 D_1], \\
 \tilde{B}_{13} &= a_+ [D_1 B_{10} Q S + S Q B_{10} D_1 + S Q B_8 Q S] - a_+^2 [D_1 B_5 Q S Q B_5 D_1 + S Q B_5 D_1 B_5 D_1 \\
 &\quad + D_1 B_5 D_1 B_5 Q S + D_1 B_4 Q S Q B_5 Q S + S Q B_5 Q S Q B_4 D_1 + S Q B_4 Q S Q B_5 D_1 \\
 &\quad + D_1 B_5 Q S Q B_4 Q S + S Q B_5 D_1 B_4 Q S + S Q B_4 D_1 B_5 Q S + S Q B_3 Q S Q B_5 Q S \\
 &\quad + S Q B_5 Q S Q B_3 Q S + S Q B_4 Q S Q B_4 Q S], \tilde{B}_{14} = a_+ S Q B_9 Q S, \\
 \tilde{B}_{15} &= a_+ S Q B_{10} Q S - a_+^2 [S Q B_5 D_1 B_5 Q S + D_1 B_5 Q S Q B_5 Q S + S Q B_5 Q S Q B_5 D_1 \\
 &\quad + S Q B_4 Q S Q B_5 Q S + S Q B_5 Q S Q B_4 Q S], \tilde{B}_{16} = -a_+^2 S Q B_5 Q S Q B_5 Q S.
 \end{aligned}$$

According to Lemma 4.1, $\tilde{M}_1^+(\lambda)$ has bounded inverse if and only if

$$M_2^+(\lambda) := S_1 - S_1 (\tilde{M}_1^+(\lambda) + S_1)^{-1} S_1 \quad (4.18)$$

has bounded inverse on $S_1 L^2$. Using the orthogonality in Remark 4.5, one has

$$\begin{aligned}
 M_2^+(\lambda) &= g_0^+(\lambda)^{-1} \left(\tilde{T}_1 + g_0^+(\lambda)^3 \lambda^2 S_1 \tilde{B}_1 S_1 + g_0^+(\lambda)^2 \lambda^2 S_1 \tilde{B}_2 S_1 + g_0^+(\lambda) \lambda^2 S_1 \tilde{B}_3 S_1 + \lambda^2 S_1 \tilde{B}_4 S_1 \right. \\
 &\quad + g_0^+(\lambda)^{-1} \lambda^2 S_1 \tilde{B}_5 S_1 + g_0^+(\lambda)^5 \lambda^4 S_1 \tilde{B}_6 S_1 + g_0^+(\lambda)^4 \lambda^4 S_1 \tilde{B}_7 S_1 + g_0^+(\lambda)^3 \lambda^4 S_1 \tilde{B}_8 S_1 \\
 &\quad + g_0^+(\lambda)^2 \lambda^4 S_1 \tilde{B}_9 S_1 + \tilde{g}_2^+(\lambda) g_0^+(\lambda) \lambda^4 S_1 \tilde{B}_{10} S_1 + g_0^+(\lambda) \lambda^4 S_1 \tilde{B}_{11} S_1 + \tilde{g}_2^+(\lambda) \lambda^4 S_1 \tilde{B}_{12} S_1 \\
 &\quad + \lambda^4 S_1 \tilde{B}_{13} S_1 + \tilde{g}_2^+(\lambda) g_0^+(\lambda)^{-1} \lambda^4 S_1 \tilde{B}_{14} S_1 + g_0^+(\lambda)^{-1} \lambda^4 S_1 \tilde{B}_{15} S_1 \\
 &\quad \left. + g_0^+(\lambda)^{-2} \lambda^4 S_1 \tilde{B}_{16} S_1 + O(\lambda^{6-\epsilon}) \right) \\
 &:= g_0^+(\lambda)^{-1} (\tilde{T}_1 + W_1^+(\lambda)) := g_0^+(\lambda)^{-1} \tilde{M}_2^+(\lambda),
 \end{aligned}$$

where

$$\widetilde{T}_1 = -S_1 T_0 (Q - S_0) (A + g_0^+(\lambda)^{-1} B)^{-1} (Q - S_0) T_0 S_1. \quad (4.19)$$

Let

$$T_1 = -S_1 T_0 (Q - S_0) A^{-1} (Q - S_0) T_0 S_1,$$

then \widetilde{T}_1 is invertible on $S_1 L^2$ for enough small λ if and only if T_1 is invertible on $S_1 L^2$.

(II) The first kind of resonance. If there is a resonance of first kind at zero, then the operator T_1 is invertible on $S_1 L^2$, in this case we define $D_2 = T_1^{-1}$ as an operator on $S_1 L^2$. Moreover, \widetilde{T}_1 is invertible on $S_1 L^2$ and $D_2 = \widetilde{T}_1^{-1}$ for enough small λ . Note that

$$(\widetilde{M}_2^+(\lambda))^{-1} = D_2 (I + W_1^+(\lambda) D_2)^{-1}$$

and $(M_2^+(\lambda))^{-1} = g_0^+(\lambda) (\widetilde{M}_2^+(\lambda))^{-1}$, by Neumann series one has

$$\begin{aligned} (M_2^+(\lambda))^{-1} &= g_0^+(\lambda) D_2 + g_0^+(\lambda)^4 \lambda^2 D_2 \widetilde{B}_1 D_2 + g_0^+(\lambda)^3 \lambda^2 D_2 \widetilde{B}_2 D_2 + g_0^+(\lambda)^2 \lambda^2 D_2 \widetilde{B}_3 D_2 \\ &\quad + g_0^+(\lambda) \lambda^2 D_2 \widetilde{B}_4 D_2 + \lambda^2 D_2 \widetilde{B}_5 D_2 + O_1(\lambda^{4-\epsilon}). \end{aligned}$$

Using Lemma 4.1 and (4.17) one has

$$\begin{aligned} (\widetilde{M}_1^+(\lambda))^{-1} &= g_0^+(\lambda) S_1 \Gamma_{0,1}^1 S_1 + \left(Q \Gamma_{0,2}^1 S_1 + S_1 \Gamma_{0,3}^1 Q \right) + g_0^+(\lambda)^{-1} Q \Gamma_{0,5}^1 Q \\ &\quad + g_0^+(\lambda)^{-2} Q \Gamma_{0,6}^1 Q + g_0^+(\lambda)^4 \lambda^2 S_1 \Gamma_{2,1}^1 S_1 + g_0^+(\lambda)^3 \lambda^2 Q \Gamma_{2,1}^1 Q \\ &\quad + g_0^+(\lambda)^2 \lambda^2 Q \Gamma_{2,1}^1 Q + g_0^+(\lambda) \lambda^2 Q \Gamma_{2,1}^1 Q + O_1(\lambda^2). \end{aligned}$$

By the same argument with the proof of Theorem 2.6(i) one has

$$\begin{aligned} (M^+(\lambda))^{-1} &= g_0^+(\lambda) S_1 D_2 S_1 + \left(S_0 \Gamma_{0,1}^1 S_0 + S_1 \Gamma_{0,2}^1 Q + Q \Gamma_{0,3}^1 S_1 \right) + g_0^+(\lambda)^{-1} Q \Gamma_{0,4}^1 Q \\ &\quad + g_0^+(\lambda)^4 \lambda^2 S_1 \Gamma_{2,1}^1 S_1 + g_0^+(\lambda)^3 \lambda^2 \left(S_1 \Gamma_{2,2}^1 Q + Q \Gamma_{2,3}^1 S_1 \right) + g_0^+(\lambda)^2 \lambda^2 \\ &\quad \times \left(Q \Gamma_{2,4}^1 Q + S_1 \Gamma_{2,5}^1 + \Gamma_{2,6}^1 S_1 \right) + g_0^+(\lambda) \lambda^2 \left(Q \Gamma_{2,7}^1 + \Gamma_{2,8}^1 Q \right) + O_1(\lambda^2). \end{aligned}$$

The expansions of $(M^+(\lambda))^{-1}$ in the first kind resonance is completed. \square

If zero is not the first kind of resonance, then T_1 is not invertible on $S_1 L^2$. Let S_2 is the Riesz transform onto the kernel of T_1 , then $T_1 + S_2$ is invertible on $S_1 L^2$, in this case, we define $D_2 = (T_1 + S_2)^{-1}$ as an operator on $S_1 L^2$. Moreover, $D_2 = (\widetilde{T}_1 + S_2)^{-1}$. Then $S_1 D_2 = D_2 S_1 = D_2$, $S_2 D_2 = D_2 S_2 = S_2$ and $D_2 D_1 = D_1 D_2 = D_2$. Note that

$$(\widetilde{M}_2^+(\lambda) + S_2)^{-1} = D_2 \left(I + W_1^+(\lambda) D_2 \right)^{-1}.$$

By using Neumann series and (4.19) one has

$$\begin{aligned} &(\widetilde{M}_2^+(\lambda) + S_2)^{-1} \\ &= D_2 - g_0^+(\lambda)^3 \lambda^2 D_2 \widetilde{B}_1 D_2 - g_0^+(\lambda)^2 \lambda^2 D_2 \widetilde{B}_2 D_2 - g_0^+(\lambda) \lambda^2 D_2 \widetilde{B}_3 D_2 - \lambda^2 D_2 \widetilde{B}_4 D_2 \\ &\quad - g_0^+(\lambda)^{-1} \lambda^2 D_2 \widetilde{B}_5 D_2 - g_0^+(\lambda)^6 \lambda^4 B_1^1 - g_0^+(\lambda)^5 \lambda^4 B_2^1 - g_0^+(\lambda)^4 \lambda^4 B_3^1 - g_0^+(\lambda)^3 \lambda^4 B_4^1 \quad (4.20) \\ &\quad - g_0^+(\lambda)^2 \lambda^4 B_5^1 - \widetilde{g}_2^+(\lambda) g_0^+(\lambda) \lambda^4 B_6^1 - g_0^+(\lambda) \lambda^4 B_7^1 - \widetilde{g}_2^+(\lambda) \lambda^4 B_8^1 - \lambda^4 B_9^1 \\ &\quad - \widetilde{g}_2^+(\lambda) g_0^+(\lambda)^{-1} \lambda^4 B_{10}^1 - g_0^+(\lambda)^{-1} \lambda^4 B_{11}^1 - g_0^+(\lambda)^{-2} \lambda^4 B_{12}^1 + O(\lambda^{6-\epsilon}), \end{aligned}$$

where $B_j^1 (1 \leq j \leq 12)$ are bounded operators on L^2 as follows:

$$\begin{aligned}
 B_1^1 &= -D_2 \tilde{B}_1 D_2 \tilde{B}_1 D_2, \quad B_2^1 = D_2 \tilde{B}_6 D_2 - [D_2 \tilde{B}_1 D_2 \tilde{B}_2 D_2 + D_2 \tilde{B}_2 D_2 \tilde{B}_1 D_2], \\
 B_3^1 &= D_2 \tilde{B}_7 D_2 - [D_2 \tilde{B}_2 D_2 \tilde{B}_2 D_2 + D_2 \tilde{B}_1 D_2 \tilde{B}_3 D_2 + D_2 \tilde{B}_3 D_2 \tilde{B}_1 D_2], \\
 B_4^1 &= D_2 \tilde{B}_8 D_2 - [D_2 \tilde{B}_1 D_2 \tilde{B}_4 D_2 + D_2 \tilde{B}_4 D_2 \tilde{B}_1 D_2 + D_2 \tilde{B}_2 D_2 \tilde{B}_3 D_2 + D_2 \tilde{B}_3 D_2 \tilde{B}_2 D_2], \\
 B_5^1 &= D_2 \tilde{B}_9 D_2 - [D_2 \tilde{B}_3 D_2 \tilde{B}_3 D_2 + D_2 \tilde{B}_2 D_2 \tilde{B}_4 D_2 + D_2 \tilde{B}_4 D_2 \tilde{B}_2 D_2 + D_2 \tilde{B}_1 D_2 \tilde{B}_5 D_2 \\
 &\quad + D_2 \tilde{B}_5 D_2 \tilde{B}_1 D_2], \quad B_6^1 = D_2 \tilde{B}_{10} D_2, \\
 B_7^1 &= D_2 \tilde{B}_{11} D_2 - [D_2 \tilde{B}_2 D_2 \tilde{B}_5 D_2 + D_2 \tilde{B}_5 D_2 \tilde{B}_2 D_2 + D_2 \tilde{B}_3 D_2 \tilde{B}_4 D_2 + D_2 \tilde{B}_4 D_2 \tilde{B}_3 D_2], \\
 B_8^1 &= D_2 \tilde{B}_{12} D_2, \quad B_9^1 = D_2 \tilde{B}_{13} D_2 - [D_2 \tilde{B}_4 D_2 \tilde{B}_4 D_2 + D_2 \tilde{B}_3 D_2 \tilde{B}_5 D_2 + D_2 \tilde{B}_5 D_2 \tilde{B}_3 D_2], \\
 B_{10}^1 &= D_2 \tilde{B}_{14} D_2, \quad B_{11}^1 = D_2 \tilde{B}_{15} D_2 - [D_2 \tilde{B}_4 D_2 \tilde{B}_5 D_2 + D_2 \tilde{B}_5 D_2 \tilde{B}_4 D_2], \\
 B_{12}^1 &= D_2 \tilde{B}_{16} D_2 - D_2 \tilde{B}_5 D_2 \tilde{B}_5 D_2.
 \end{aligned}$$

According to Lemma 4.1, $\tilde{M}_2^+(\lambda)$ has bounded inverse on $S_1 L^2$ if and only if

$$M_3^+(\lambda) := S_2 - S_2 (\tilde{M}_2^+(\lambda) + S_2)^{-1} S_2 \quad (4.21)$$

has bounded inverse on $S_2 L^2(\mathbb{R}^2)$. By Neumann series we have

$$\begin{aligned}
 M_3^+(\lambda) &= g_0^+(\lambda)^3 \lambda^2 S_2 \tilde{B}_1 S_2 + g_0^+(\lambda)^2 \lambda^2 S_2 \tilde{B}_2 S_2 + g_0^+(\lambda) \lambda^2 S_2 \tilde{B}_3 S_2 + \lambda^2 S_2 \tilde{B}_4 S_2 \\
 &\quad + g_0^+(\lambda)^{-1} \lambda^2 S_2 \tilde{B}_5 S_2 + g_0^+(\lambda)^6 \lambda^4 S_2 B_1^1 S_2 + g_0^+(\lambda)^5 \lambda^4 S_2 B_2^1 S_2 + g_0^+(\lambda)^4 \lambda^4 S_2 B_3^1 S_2 \\
 &\quad + g_0^+(\lambda)^3 \lambda^4 S_2 B_4^1 S_2 + g_0^+(\lambda)^2 \lambda^4 S_2 B_5^1 S_2 + \tilde{g}_2^+(\lambda) g_0^+(\lambda) \lambda^4 S_2 B_6^1 S_2 + g_0^+(\lambda) \lambda^4 S_2 B_7^1 S_2 \\
 &\quad + \tilde{g}_2^+(\lambda) \lambda^4 S_2 B_8^1 S_2 + \lambda^4 S_2 B_9^1 S_2 + \tilde{g}_2^+(\lambda) g_0^+(\lambda)^{-1} \lambda^4 S_2 B_{10}^1 S_2 + g_0^+(\lambda)^{-1} \lambda^4 S_2 B_{11}^1 S_2 \\
 &\quad + g_0^+(\lambda)^{-2} \lambda^4 S_2 B_{12}^1 S_2 + O(\lambda^{6-\epsilon}).
 \end{aligned}$$

Note that, $S_2 S = S S_2 = 0$, $QvG_{-1}vS_0 = 0$, $QT_0S_2 = 0$, by (4.17) we have

$$S_2 \tilde{B}_1 S_2 = -\frac{1}{a_+} S_2 v G_{-1} v v G_{-1} v S_2, \quad S_2 \tilde{B}_2 S_2 = -\frac{1}{a_+} (S_2 v G_{-1} v T_0 S_2 + S_2 T_0 v G_{-1} v S_2),$$

$$S_2 \tilde{B}_3 S_2 = c_+ S_2 v G_1 v S_2 - \frac{1}{a_+} S_2 T_0^2 S_2,$$

$$S_2 \tilde{B}_4 S_2 = S_3 \tilde{B}_5 S_2 = S_2 B_8^1 S_2 = S_2 B_{10}^1 S_2 = S_2 B_{12}^1 S_2 = 0.$$

Moreover, we get

$$\begin{aligned}
 &M_3^+(\lambda) \\
 &= -\frac{1}{a_+} g_0^+(\lambda) \lambda^2 \left(A_1^+(\lambda) - g_0^+(\lambda)^5 \lambda^2 a_+ S_2 B_1^1 S_2 - g_0^+(\lambda)^4 \lambda^2 a_+ S_2 B_2^1 S_2 \right. \\
 &\quad - g_0^+(\lambda)^3 \lambda^2 a_+ S_2 B_3^1 S_2 - g_0^+(\lambda)^2 \lambda^2 a_+ S_2 B_4^1 S_2 - g_0^+(\lambda) \lambda^2 a_+ S_2 B_5^1 S_2 \\
 &\quad - \tilde{g}_2^+(\lambda) \lambda^2 a_+ S_2 B_6^1 S_2 - \lambda^2 a_+ S_2 B_7^1 S_2 - g_0^+(\lambda)^{-1} \lambda^2 a_+ S_2 B_9^1 S_2 \\
 &\quad \left. - g_0^+(\lambda)^{-2} \lambda^2 a_+ S_2 B_{11}^1 S_2 + O(\lambda^{4-\epsilon}) \right) \\
 &:= -\frac{1}{a_+} g_0^+(\lambda) \lambda^2 (A_1^+(\lambda) + W_2^+(\lambda)) := -\frac{1}{a_+} g_0^+(\lambda) \lambda^2 \tilde{M}_3^+(\lambda),
 \end{aligned} \quad (4.22)$$

where

$$A_1^+(\lambda) = g_0^+(\lambda)^2 S_2 v G_{-1} v v G_{-1} v S_2 + g_0^+(\lambda) (S_2 v G_{-1} v T_0 S_2 + S_2 T_0 v G_{-1} v S_2) + (S_2 T_0^2 S_2 - c_+ a_+ S_2 v G_1 v S_2).$$

To compute $(M_3^+(\lambda))^{-1}$, by (4.22) we need to compute the expansions of $(\widetilde{M}_3^+(\lambda))^{-1}$ on $S_2 L^2$. Noting that $A_1^+(\lambda)$ with logarithm factor, and by Lemma 2.5(iv) we know that T_2 is not invertible on $S_2 L^2$, hence we will use Lemma 4.2 to study the inverse of $A_1^+(\lambda)$.

To compute the inverse of $A_1^+(\lambda)$ by Lemma 4.2, we need the following lemma.

Lemma 4.6. *Suppose that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with some $\beta > 18$. Let*

$$\begin{aligned} L_1 &= (S_2 - S_3) \left(v G_{-1} v P v G_{-1} v - v G_{-1} v T_0 D_4 T_0 v G_{-1} v \right) (S_2 - S_3), \\ L_2 &= (S_2 - S_3) \left(v G_{-1} v T_0 + T_0 v G_{-1} v - v G_{-1} v T_0 D_4 (T_0^2 - c_+ a_+ v G_1 v) \right. \\ &\quad \left. - (T_0^2 - c_+ a_+ v G_1 v) D_4 T_0 v G_{-1} v \right) (S_2 - S_3), \\ L_3 &= (S_2 - S_3) \left(T_0^2 - c_+ a_+ v G_1 v - (T_0^2 - c_+ a_+ v G_1 v) D_4 (T_0^2 - c_+ a_+ v G_1 v) \right) (S_2 - S_3). \end{aligned}$$

Then the inverse operator

$$d_1 := g_0^+(\lambda)^{-2} \left(L_1 + g_0^+(\lambda)^{-1} L_2 + g_0^+(\lambda)^{-2} L_3 \right)^{-1}$$

exists on $(S_2 - S_3)L^2$ for enough small $\lambda > 0$, and

$$d_1 = g_0^+(\lambda)^{-2} h^+(\lambda) (S_2 - S_3),$$

where $h^+(\lambda) = \left(c_1 + g_0^+(\lambda)^{-1} c_2 + g_0^+(\lambda)^{-2} c_3 \right)^{-1}$ with $c_1 = \text{tr}(L_1)$, $c_2 = \text{tr}(L_2)$ and $c_3 = \text{tr}(L_3)$. Moreover, $h^+(\lambda) = O_1(1)$.

Proof. We first show that L_1 is invertible on $(S_2 - S_3)L^2$. Indeed, by Lemma 2.5, we know that $(S_2 - S_3)L^2 = \text{span}\{S_2(|x|^2 v)\}$, thus it suffices to prove $\langle |x|^2 v, L^1(|x|^2 v) \rangle > 0$. Using $S_2 v = S_2 x_i v = 0$ ($i = 1, 2$) and $S_3 v = S_3 x_i v = S_3 |x|^2 v = 0$ ($i = 1, 2$), one has

$$\begin{aligned} P v G_{-1} v S_2(|x|^2 v)(x) &= P v(x) \int_{\mathbb{R}^2} (|x|^2 - 2x \cdot y + |y|^2) v(y) S_2(|x|^2 v)(y) dy \\ &= P \langle |x|^2 v, S_2(|x|^2 v) \rangle v(x) = \|S_2(|x|^2 v)\|_{L^2}^2 v(x). \end{aligned} \tag{4.23}$$

Let $\phi(x) = P v G_{-1} v S_2(|x|^2 v)(x)$, by $S_3 |x|^2 v = 0$ and (4.23), one has

$$\begin{aligned} (S_2 - S_3) v G_{-1} v P v G_{-1} v (S_2 - S_3)(|x|^2 v)(x) &= (S_2 - S_3) v G_{-1} v \phi(x) \\ &= (S_2 - S_3) v(x) \int_{\mathbb{R}^2} (|x|^2 - 2x \cdot y + |y|^2) v(y) \phi(y) dy \\ &= \|S_2(|x|^2 v)\|_{L^2}^2 (S_2 - S_3) |x|^2 v(x) \int_{\mathbb{R}^2} v(y) v(y) dy \\ &= \|V\|_{L^1} \|S_2(|x|^2 v)\|_{L^2}^2 v(x). \end{aligned}$$

Furthermore,

$$\langle |x|^2 v, (S_2 - S_3) v G_{-1} v P v G_{-1} v (S_2 - S_3)(|x|^2 v) \rangle = \|V\|_{L^1} \|S_2(|x|^2 v)\|_{L^2}^4. \tag{4.24}$$

Using (4.23) and $S_3(|x|^2v) = 0$, we have

$$\begin{aligned} & \langle |x|^2v, (S_2 - S_3)vG_{-1}vT_0D_4T_0vG_{-1}vS_2(|x|^2v) \rangle \\ &= \langle D_4T_0PvG_{-1}vS_2(|x|^2v), T_0PvG_{-1}vS_2(|x|^2v) \rangle = \|S_2(|x|^2v)\|_{L^2}^4 \langle D_4T_0v, T_0v \rangle. \end{aligned} \quad (4.25)$$

Combing (4.24) and (4.25), one has

$$\langle |x|^2v, L^1(|x|^2v) \rangle = \|S_2(|x|^2v)\|_{L^2}^4 \|V\|_{L^1} \left(1 - \frac{\langle D_4T_0v, T_0v \rangle}{\|V\|_{L^1}}\right). \quad (4.26)$$

Let $\eta = D_4S_3T_0v$ and $\xi = S_3T_0v$, then $\xi = D_4^{-1}\eta = (T_3 + S_4)\eta$. By $S_3D_4 = D_4S_3 = D_4$, thus we have

$$1 - \frac{\langle D_4T_0v, T_0v \rangle}{\|V\|_{L^1}} = 1 - \frac{\langle S_3D_4T_0v, S_3T_0v \rangle}{\|V\|_{L^1}} = 1 - \frac{\langle \eta, \xi \rangle}{\|V\|_{L^1}}.$$

Nothing that

$$T_3 + S_4 = (-c_+a_+S_3vG_1vS_3 + S_4) + S_3T_0^2S_3 := A_1 + B_1,$$

by (2.22) we know that for $f \in S_3L^2$,

$$\langle f, A_1f \rangle = \frac{\|V\|_{L^1}}{1024} \left(\langle x_1^2v, f \rangle^2 + \langle x_2^2v, f \rangle^2 + 2\langle x_1x_2v, f \rangle^2 \right) + \|S_4f\|_{L^2}^2 > 0,$$

since $\eta = D_4\xi = S_3D_4\xi \in S_3L^2$, thus $\langle \eta, A_1\eta \rangle > 0$. Furthermore,

$$0 < \langle \eta, A_1\eta \rangle = \langle \eta, (T_3 + S_4)\eta - B_1\eta \rangle = \langle \eta, \xi \rangle - \langle \eta, B_1\eta \rangle. \quad (4.27)$$

By $S_3T_0Q = QT_0S_3 = 0$, then

$$B_1\eta = S_3T_0^2S_3\eta = S_3T_0PT_0S_3\eta = \frac{\langle T_0S_3\eta, v \rangle S_3T_0v}{\|V\|_{L^1}} = \frac{\langle \eta, S_3T_0v \rangle S_3T_0v}{\|V\|_{L^1}} = \frac{\langle \eta, \xi \rangle \xi}{\|V\|_{L^1}},$$

using (4.27), one has

$$0 < \langle \eta, A_1\eta \rangle = \langle \eta, \xi \rangle - \frac{|\langle \eta, \xi \rangle|^2}{\|V\|_{L^1}},$$

which implies that $\langle \eta, \xi \rangle > 0$ and $1 - \frac{\langle \eta, \xi \rangle}{\|V\|_{L^1}} > 0$. Thus, $1 - \frac{\langle D_4T_0v, T_0v \rangle}{\|V\|_{L^1}} > 0$. Furthermore, we obtain that $\langle |x|^2v, L^1(|x|^2v) \rangle > 0$. Thus, L_1 is invertible on $(S_2 - S_3)L^2$. Furthermore, d_1 exists on $(S_2 - S_3)L^2$ for enough small λ .

Note that $(S_2 - S_3)L^2 = \text{span}\{S_2(|x|^2v)\}$. Thus, $\text{rank}(S_2 - S_3) = 1$. Suppose that $f \in L^2$, then $L_1f = (S_2 - S_3)L_1f \in (S_2 - S_3)L^2$. Thus, $L_1f = \langle |x|^2v(x), L_1f \rangle |x|^2v(x)$. Moreover, $L_1 = \text{tr}(L_1) := c_1 \neq 0$ by $\langle |x|^2v, L^1(|x|^2v) \rangle > 0$. Similarly, $L_2 = \text{tr}(L_2) := c_2$ and $L_3 = \text{tr}(L_3) := c_3$. Thus, we have

$$\begin{aligned} d_1 &= \left(g_0^+(\lambda)^2 L_1 + g_0^+(\lambda) L_2 + L_3 \right)^{-1} \\ &= g_0^+(\lambda)^{-2} (S_2 - S_3) \left(L_1 + g_0^+(\lambda)^{-1} L_2 + g_0^+(\lambda)^{-2} L_3 \right)^{-1} (S_2 - S_3) \\ &= g_0^+(\lambda)^{-2} (S_2 - S_3) \left(c_1 + g_0^+(\lambda)^{-1} c_2 + g_0^+(\lambda)^{-2} c_3 \right)^{-1} (S_2 - S_3) \\ &:= g_0^+(\lambda)^{-2} h^+(\lambda) (S_2 - S_3), \end{aligned}$$

where $h^+(\lambda) = \left(c_1 + g_0^+(\lambda)^{-1} c_2 + g_0^+(\lambda)^{-2} c_3 \right)^{-1}$.

It is easy to check that $h^+(\lambda) = O_1(1)$. □

Now we start to compute the inverse of $A_1^+(\lambda)$ as follows:

Lemma 4.7. *Suppose that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with some $\beta > 18$. If zero is the second kind of resonance of the spectrum of H , for $0 < |\lambda| \ll 1$, then*

$$(A_1^+(\lambda))^{-1} = D_4 + h^+(\lambda)\tilde{S}(\lambda),$$

where $D_4 = T_3^{-1}$ and

$$\tilde{S}(\lambda) = \tilde{S} = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix},$$

which $h(\lambda)$ is defined in Lemma 4.6, and $b_{ij} = b_{ij}(\lambda)$ are bounded operators for enough small λ as follows:

$$\begin{aligned} b_{11} &= g_0^+(\lambda)^{-2}(S_2 - S_3), \\ b_{12} &= - (S_2 - S_3) \left(g_0^+(\lambda)^{-1}vG_{-1}vT_0 + g_0^+(\lambda)^{-2}(T_0^2 - c_+a_+vG_1v) \right) D_4, \\ b_{21} &= - D_4 \left(g_0^+(\lambda)^{-1}T_0vG_{-1}v + g_0^+(\lambda)^{-2}(T_0^2 - c_+a_+vG_1v) \right) (S_2 - S_3), \\ b_{22} &= D_4 \left(T_0vG_{-1}v + g_0^+(\lambda)^{-1}(T_0^2 - c_+a_+vG_1v) \right) (S_2 - S_3) \\ &\quad \times \left(vG_{-1}vT_0 + g_0^+(\lambda)^{-1}(T_0^2 - c_+a_+vG_1v) \right) D_4. \end{aligned} \tag{4.28}$$

Moreover, $\|\tilde{S}(\lambda)\|_{L^2 \rightarrow L^2} = O_1(1)$ for enough small λ .

In addition, if zero is not the first kind of resonance, the same formula holds for $(A_1^+(\lambda) + S_4)^{-1}$ with $D_4 = (T_3 + S_4)^{-1}$.

Proof. We only prove the case of $S_4 \neq 0$, the case of $S_4 = 0$ is identical. In this case, $T_3 + S_4$ is invertible on S_3L^2 . Observe that $vG_{-1}vS_3 = 0$, we have

$$\begin{aligned} A_1^+(\lambda) + S_4 &= \left((S_2 - S_3) + S_3 \right) \left(g_0^+(\lambda)^2vG_{-1}vPvG_{-1}v + g_0^+(\lambda)(vG_{-1}vT_0 + T_0vG_{-1}v) \right. \\ &\quad \left. + T_0^2 - c_+a_+vG_1v \right) \left((S_2 - S_3) + S_3 \right). \end{aligned}$$

By block format we write $A_1^+(\lambda) + S_4$ as follows:

$$A_1^+(\lambda) + S_4 = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}, \tag{4.29}$$

where $a_{ij} = a_{ij}(\lambda)$ are bounded operators for enough small λ as follows:

$$\begin{aligned} a_{11} &= (S_2 - S_3) \left(g_0^+(\lambda)^2vG_{-1}vPvG_{-1}v \right. \\ &\quad \left. + g_0^+(\lambda)(vG_{-1}vT_0 + T_0vG_{-1}v) + T_0^2 - c_+a_+vG_1v \right) (S_2 - S_3), \\ a_{12} &= (S_2 - S_3) \left(g_0^+(\lambda)vG_{-1}vT_0 + T_0^2 - c_+a_+vG_1v \right) S_3, \\ a_{21} &= S_3 \left(g_0^+(\lambda)T_0vG_{-1}v + T_0^2 - c_+a_+vG_1v \right) (S_2 - S_3), \\ a_{22} &= S_3T_0^2S_3 - c_+a_+S_3vG_1vS_3 + S_4 = T_3 + S_4. \end{aligned}$$

By Lemma 4.2, then $A^+(\lambda) + S_4$ has a bounded inverse if and only if

$$d_1 := (a_{11} - a_{12}a_{22}^{-1}a_{21})^{-1}$$

exist and is bounded. Let $D_4 = (T_3 + S_4)^{-1} : S_3L^2 \rightarrow S_3L^2$. Note that $\text{rank}(S_2 - S_3) = 1$, and $S_3D_4 = D_4S_3 = D_4$. By Lemma 4.6, the inverse operator

$$d_1 := g_0^+(\lambda)^{-2} \left(L_1 + g_0^+(\lambda)^{-1}L_2 + g_0^+(\lambda)^{-2}L_3 \right)^{-1}$$

exists on $(S_2 - S_3)L^2$ for enough small $\lambda > 0$, and

$$d_1 = g_0^+(\lambda)^{-2}h^+(\lambda)(S_2 - S_3).$$

Using Lemma 4.2 again, $A_1^+(\lambda) + S_4$ is invertible, and

$$(A_1^+(\lambda) + S_4)^{-1} = D_4 + h^+(\lambda)\tilde{S}(\lambda),$$

where

$$\tilde{S}(\lambda) = \tilde{S} = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix},$$

where $b_{i,j}(\lambda)$ are bounded operators defined in (4.28).

It is easy to check that $\|\tilde{S}(\lambda)\|_{L^2 \rightarrow L^2} = O_1(1)$ for enough small λ . \square

(III) The second kind of resonance. If zero is the second kind of resonance, then T_3 is invertible on S_3L^2 , let $D_4 = T_3^{-1}$. Since

$$(\widetilde{M}_3^+(\lambda))^{-1} = (A_1^+(\lambda))^{-1} \left(I + W_2^+(\lambda)(A_1^+(\lambda))^{-1} \right)^{-1}.$$

Note that $D_4B_1^1 = B_1^1D_4 = 0$, $D_4B_2^1D_4 = 0$, by Lemma 4.7 and (4.22), we have

$$\begin{aligned} (\widetilde{M}_3^+(\lambda))^{-1} &= h^+(\lambda)S_2\Gamma_{0,1}^2S_2 + g_0^+(\lambda)^5\lambda^2S_2\Gamma_{2,1}^2S_2 + \lambda^2S_2\Gamma_{2,2}^2S_2 \\ &\quad + g_0^+(\lambda)^{10}\lambda^4S_2\Gamma_{4,1}^2S_2 + \lambda^4S_2\Gamma_{4,2}^2S_2 + O_1(\lambda^4 \ln \lambda). \end{aligned}$$

By (4.22) one has

$$\begin{aligned} (M_3^+(\lambda))^{-1} &= \frac{g_0^+(\lambda)^{-1}h^+(\lambda)}{\lambda^2}S_2\Gamma_{-2,1}^2S_2 + g_0^+(\lambda)^4S_2\Gamma_{0,1}^2S_2 + S_2\Gamma_{0,2}^2S_2 \\ &\quad + g_0^+(\lambda)^9\lambda^2S_2\Gamma_{2,1}^2S_2 + \lambda^2S_2\Gamma_{2,1}^2S_2 + O_1(\lambda^2(\ln \lambda)^{-1}). \end{aligned}$$

According to Lemma 4.1 and (4.20),

$$\begin{aligned} (\widetilde{M}_2^+(\lambda))^{-1} &= \frac{h^+(\lambda)}{\lambda^2}S_2\Gamma_{-2,1}^2S_2 + g_0^+(\lambda)^5S_2\Gamma_{0,1}^2S_2 + g_0^+(\lambda)S_2\Gamma_{0,2}^2S_2 + g_0^+(\lambda)^{10}\lambda^2S_2\Gamma_{2,1}^2S_2 \\ &\quad + g_0^+(\lambda)\lambda^2S_2\Gamma_{2,1}^2S_2 + O_1(\lambda^2). \end{aligned}$$

By the same argument with the proof of the first kind of resonance, we immediately get

$$\begin{aligned} (M^\pm(\lambda))^{-1} &= \frac{h^\pm(\lambda)}{\lambda^2}S_2\Gamma_{-2,0}^2S_2 + g_0^\pm(\lambda)^5 \left(S_2\Gamma_{0,1}^2 + \Gamma_{0,2}^2S_2 + S_1\Gamma_{0,3}^2Q + Q\Gamma_{0,4}^2S_1 \right) \\ &\quad + g_0^\pm(\lambda)^{10}\lambda^2 \left(S_2\Gamma_{2,1}^2 + \Gamma_{2,2}^2S_2 + S_1\Gamma_{2,3}^2Q + Q\Gamma_{2,4}^2S_1 \right) \\ &\quad + g_0^\pm(\lambda)\lambda^2 \left(S_2\Gamma_{2,5}^2 + \Gamma_{2,6}^2S_2 + S_1\Gamma_{2,7}^2Q + Q\Gamma_{2,8}^2S_1 \right) + O_1(\lambda^2). \end{aligned}$$

The expansions of $(M^+(\lambda))^{-1}$ of the second kind resonance is completed. \square

If zero is not the second kind of resonance. Then T_3 is not invertible on S_3L^2 . Let S_4 is the Riesz projection onto the kernel of T_3 , in this case, $T_3 + S_4$ is invertible on S_3L^2 . Let $D_4 = (T_3 + S_4)^{-1}$. Noting that the identity,

$$\left(\widetilde{M}_3^+(\lambda) + S_4\right)^{-1} = \left(A_1^+(\lambda) + S_4\right)^{-1} \left(I + W_2^+(\lambda) \left(A_1^+(\lambda) + S_4\right)^{-1}\right)^{-1},$$

using Lemma 4.7 and (4.22), we have by Neumann series

$$\begin{aligned} & \left(\widetilde{M}_3^+(\lambda) + S_4\right)^{-1} \\ &= D_4 + \widetilde{S} - g_0^+(\lambda)^5 \lambda^2 B_1^2 - g_0^+(\lambda)^4 \lambda^2 B_2^2 - g_0^+(\lambda)^3 \lambda^2 B_3^2 - g_0^+(\lambda)^2 \lambda^2 B_4^2 \\ & \quad - g_0^+(\lambda) \lambda^2 B_5^2(\lambda) - \widetilde{g}_2^+(\lambda) \lambda^2 B_6^2 - \lambda^2 B_7^2 - g_0^+(\lambda)^{-1} \lambda^2 B_9^2 - g_0^+(\lambda)^{-2} \lambda^2 B_{11}^2 \\ & \quad + O_1(\lambda^{4-\epsilon}), \end{aligned} \tag{4.30}$$

where $B_j^2 = -a_+(D_4 + h^+(\lambda)\widetilde{S})S_2B_j^1S_2(D_4 + h^+(\lambda)\widetilde{S})$, $1 \leq j \leq 11$.

According to Lemma 4.1, $\widetilde{M}_3^+(\lambda)$ has bounded inverse if and only if

$$M_4^+(\lambda) := S_4 - S_4 \left(\widetilde{M}_3^+(\lambda) + S_4\right)^{-1} S_4 \tag{4.31}$$

has bounded inverse on S_4L^2 . Since

$$S_4\widetilde{S} = \widetilde{S}S_4 = 0, \quad S_4 \left(A_1^+(\lambda) + S_4\right)^{-1} = \left(A_1^+(\lambda) + S_4\right)^{-1} S_4 = S_4,$$

thus $S_4B_j = B_jS_4 = 0$ ($j = 1, 2, 3, 4, 6, 7$), $S_4\widetilde{B}_j = \widetilde{B}_jS_4 = 0$ ($j = 1, 2, 3, 7$). Moreover,

$$S_4B_j^2S_4 = 0 \quad (j = 1, \dots, 5, 8, 10, 11),$$

$$S_4B_6^2S_4 = -a_+S_4vG_2vS_4, \quad S_4B_7^2S_4 = -a_+S_4vG_3vS_4,$$

$$S_4B_9^2S_4 = c_+^2a_+ \left(S_4vG_1vQSQvG_1vS_4 + S_4vG_1vQSD_2SQvG_1vS_4 \right).$$

Hence, by (4.30) and (4.31) one has

$$\begin{aligned} M_4^+(\lambda) &= -a_+\widetilde{g}_2^+(\lambda)\lambda^2 \left(S_4vG_2vS_4 + \widetilde{g}_2^+(\lambda)^{-1}S_4vG_3vS_4 - c_+^2\widetilde{g}_2^+(\lambda)^{-1}g_0^+(\lambda)^{-1} \right. \\ & \quad \left. \times \left(S_4vG_1vQSQvG_1vS_4 + S_4vG_1vQSD_2SQvG_1vS_4 \right) + O(\lambda^{2-\epsilon}) \right) \\ &:= -a_+\widetilde{g}_2^+(\lambda)\lambda^2\widetilde{M}_4^+(\lambda). \end{aligned}$$

Let $T_4 = S_4vG_2vS_4$. Then

$$\widetilde{M}_4^+(\lambda) = T_4 + W_3^+(\lambda), \tag{4.32}$$

where

$$\begin{aligned} W_3^+(\lambda) &= \widetilde{g}_2^+(\lambda)^{-1}S_4vG_3vS_4 - \widetilde{g}_2^+(\lambda)^{-1}g_0^+(\lambda)^{-1}c_+^2 \left(S_4vG_1vQSQvG_1vS_4 \right. \\ & \quad \left. + S_4vG_1vQSD_2SQvG_1vS_4 \right) + O(\lambda^{2-\epsilon}). \end{aligned}$$

(IV) The third kind of resonance. If zero is the third kind of resonance of H . Then T_4 is invertible on S_4L^2 , let $D_5 = T_4^{-1}$. Since

$$\left(\widetilde{M}_4^+(\lambda)\right)^{-1} = D_5 \left(I + W_3^+(\lambda)D_5 \right)^{-1}, \quad \left(M_4^+(\lambda)\right)^{-1} = -\frac{\widetilde{g}_2^+(\lambda)^{-1}}{a_+\lambda^2} \left(\widetilde{M}_4^+(\lambda)\right)^{-1},$$

by Neumann series one has

$$(M_4^+(\lambda))^{-1} = \frac{-\tilde{g}_2^+(\lambda)^{-1}D_5}{a_+\lambda^2} + \frac{\tilde{g}_2^+(\lambda)^{-2}D_5vG_3vD_5}{a_+\lambda^2} + O_1(\lambda^{-2}(\ln \lambda)^{-3}).$$

using Lemma 4.1 and (4.30), we have

$$(\widetilde{M}_3^+(\lambda))^{-1} = \frac{\tilde{g}_2^+(\lambda)^{-1}g_0^+(\lambda)^{-1}D_5}{\lambda^4} - \frac{\tilde{g}_2^+(\lambda)^{-2}g_0^+(\lambda)^{-1}D_5vG_3vD_5}{\lambda^4} + O_1(\lambda^{-4}(\ln \lambda)^{-4}).$$

Using (4.22) one has

$$(M_3^+(\lambda))^{-1} = \frac{\tilde{g}_2^+(\lambda)^{-1}g_0^+(\lambda)^{-1}D_5}{\lambda^4} - \frac{\tilde{g}_2^+(\lambda)^{-2}g_0^+(\lambda)^{-1}D_5vG_3vD_5}{\lambda^4} + O_1(\lambda^{-4}(\ln \lambda)^{-4}).$$

By the same argument with the proof of the second kind of resonance, we immediately obtain that

$$\begin{aligned} (M^+(\lambda))^{-1} &= \frac{\tilde{g}_2^+(\lambda)^{-1}}{\lambda^4}D_5 - \frac{\tilde{g}_2^+(\lambda)^{-2}}{\lambda^4}D_5vG_3vD_5 + O_1(\lambda^{-4}(\ln \lambda)^{-3}) \\ &:= \frac{\tilde{g}_2^+(\lambda)^{-1}}{\lambda^4}S_4\Gamma_{-4,1}^3S_4 + \frac{\tilde{g}_2^+(\lambda)^{-2}}{\lambda^4}S_4\Gamma_{-4,2}^3S_4 + O_1(\lambda^{-4}(\ln \lambda)^{-3}). \end{aligned}$$

The expansions of $(M^+(\lambda))^{-1}$ of the third kind resonance is completed. \square

Before computing the expansions of $(M^+(\lambda))^{-1}$ of the fourth kind resonance, we first give a lemma as follows.

Lemma 4.8. *$\ker(S_5vG_3vS_5) = \{0\}$, thus the operator $T_5 = S_5vG_3vS_5$ is invertible on S_5L^2 .*

Proof. If $f \in \ker(S_5vG_3vS_5)$, then $S_5vG_3vS_5f = 0$ for $f \in S_5L^2$, thus

$$0 = \langle S_5vG_3vS_5f, f \rangle = \langle G_3vf, vf \rangle.$$

Since

$$R_0^+(\mu^4) = \frac{b_+}{\lambda^2}I + g_0^+(\lambda)G_{-1} + G_0 + c_+\lambda^2G_1 + \tilde{g}_2^+(\lambda)G_2 + \lambda^4G_3 + O(\lambda^{6-\epsilon}|x-y|^{8-\epsilon}),$$

then,

$$G_3 = \frac{R_0^+(\lambda^4) - \frac{b_+}{\lambda^2}I - g_0^+(\lambda)G_{-1} - G_0 - c_+\lambda^2G_1 - \tilde{g}_2^+(\lambda)G_2 + O(\lambda^{6-\epsilon}|x-y|^{8-\epsilon})}{\lambda^4}.$$

Noting that for $f \in S_5L^2$,

$$\langle G_2vf, vf \rangle = \langle G_1vf, vf \rangle = \langle G_{-1}vf, vf \rangle = 0,$$

$$\langle Ivf, vf \rangle = \langle v(x)Iv(y)f(y), f(y) \rangle = \langle Pf, f \rangle = 0,$$

we get

$$\begin{aligned} 0 &= \langle S_5vG_3vS_5f, f \rangle = \langle G_3vf, vf \rangle \\ &= \lim_{\lambda \rightarrow 0} \left\langle \frac{R_0^+(\lambda^4) - \frac{b_+}{\lambda^2}I - g_0^+(\lambda)G_{-1} - G_0 - c_+\lambda^2G_1 - \tilde{g}_2^+(\lambda)G_2 + O(\lambda^{6-\epsilon}|x-y|^{8-\epsilon})}{\lambda^4}vf, vf \right\rangle \\ &= \lim_{\lambda \rightarrow 0} \frac{1}{\lambda^4} \langle (R_0^+(\lambda^4) - G_0)vf, vf \rangle. \end{aligned}$$

Notice that $G_0 = (\Delta^2)^{-1}$, by using the Fourier transform one has

$$\begin{aligned} 0 &= \lim_{\lambda \rightarrow 0} \frac{1}{\lambda^4} \left\langle \left(\frac{1}{\xi^4 - \lambda^4} - \frac{1}{\xi^4} \right) \widehat{vf}(\xi), \widehat{vf}(\xi) \right\rangle \\ &= \lim_{\lambda \rightarrow 0} \left\langle \frac{\widehat{vf}(\xi)}{(\xi^4 - \lambda^4)\xi^4}, \widehat{vf}(\xi) \right\rangle = \left\langle \frac{\widehat{vf}(\xi)}{\xi^8}, \widehat{vf}(\xi) \right\rangle \\ &= \int_{\mathbb{R}^2} \frac{|\widehat{vf}(\xi)|^2}{\xi^8} d\xi = \langle G_0 vf, G_0 vf \rangle, \end{aligned}$$

where we use the monotone convergence theorem as $\lambda \rightarrow 0$ with choosing $\lambda \in \mathbb{C}$ such that $\lambda^4 < 0$. Thus, $vf = 0$ since $vf \in L^1$. Noting that $f \in S_5 L^2$, hence $f = -UvG_0vf = 0$. \square

(V) The fourth kind of resonance. If zero is not the fourth kind of resonance, then T_4 is not invertible on $S_4 L^2$. Let S_5 is the Riesz projection onto the kernel of T_4 , then $T_4 + S_5$ is invertible on $S_4 L^2$, let $D_5 = (T_4 + S_5)^{-1}$. Noting that the identity

$$\left(\widetilde{M}_4^+(\lambda) + S_5 \right)^{-1} = D_5 \left(I + W_3^+(\lambda) D_5 \right)^{-1},$$

by (4.32) and Neumann series, we have

$$\begin{aligned} \left(\widetilde{M}_4^+(\lambda) + S_5 \right)^{-1} &= D_5 - \widetilde{g}_2^+(\lambda)^{-1} D_5 v G_3 v D_5 + \widetilde{g}_2^+(\lambda)^{-1} g_0^+(\lambda)^{-1} c_+^2 \left(D_5 v G_1 v Q S Q v G_1 v D_5 \right. \\ &\quad \left. + D_5 v G_1 v Q S D_2 S Q v G_1 v D_5 \right) + O((\ln \lambda)^{-3}). \end{aligned}$$

According to Lemma 4.1, $\widetilde{M}_4^+(\lambda)$ has bounded inverse if and only if

$$M_5^+(\lambda) := S_5 - S_5 \left(\widetilde{M}_4^+(\lambda) + S_5 \right)^{-1} S_5 \tag{4.33}$$

has bounded inverse on $S_5 L^2(\mathbb{R}^2)$. Note that $S_5 v G_1 v Q = Q v G_1 v S_5 = 0$, by Neumann series one has

$$M_5(\lambda) = \widetilde{g}_2^+(\lambda)^{-1} S_5 v G_3 v S_5 + O((\ln \lambda)^{-2}).$$

By Lemma 4.8, $T_5 = S_5 v G_3 v S_5$ is invertible on $S_5 L^2$, let $D_6 = T_5^{-1}$. Then we have

$$\left(M_5(\lambda) \right)^{-1} = \widetilde{g}_2^+(\lambda) D_6 + O((\ln \lambda)^{-2}).$$

By the same argument with the proof of the third kind of resonance, we immediately obtain that

$$\left(M^+(\lambda) \right)^{-1} = \frac{1}{\lambda^4} S_5 D_6 S_5 + \frac{\widetilde{g}_2^+(\lambda)^{-1}}{\lambda^4} \left(S_4 \Gamma_{-4,1}^4 S_4 \right) + \frac{\widetilde{g}_2^+(\lambda)^{-2}}{\lambda^4} S_4 \Gamma_{-4,2}^4 S_4 + O_1(\lambda^{-4} (\ln \lambda)^{-3}).$$

Thus, the expansions of $\left(M^+(\lambda) \right)^{-1}$ of the fourth kind resonance is completed. \square

5. CLASSIFICATION OF THRESHOLD SPECTRAL SUBSPACES

From Definition 2.4 we know that the four projection subspaces $S_i L^2 (i = 1, 2, 4, 5)$ define the four zero resonance types. In this section, we will identify these four subspaces by the distributional solutions of $H\phi = 0$ in the weighted spaces L_s^∞ with some $s \in \mathbb{R}$.

We define $\ln^+ t = \ln t$ if $t > 1$ and $\ln^+ t = 0$ if $0 < t \leq 1$; $\ln^- t = 0$ if $t > 1$ and $\ln^- t = -\ln t$ if $0 < t \leq 1$. Then $\ln t = \ln^+ t - \ln^- t$.

Proposition 5.1. *Assume that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 14$. Then $f \in S_1 L^2(\mathbb{R}^2) \setminus \{0\}$ if and only if $f = Uv\phi$ with $\phi \in L_{-1}^\infty(\mathbb{R}^2)$ such that $H\phi = 0$ in the distributional sense, and*

$$\phi = -G_0 v f + c_3 x_2 + c_2 x_1 + c_1,$$

$$\text{where } c_3 = \frac{\langle f_2, T_0 f \rangle}{\|f_2\|_{L^2(\mathbb{R}^2)}^2}, \quad c_2 = -\frac{\langle f_2, T_0 f \rangle \langle x_2 v, f_1 \rangle}{\|f_2\|_{L^2(\mathbb{R}^2)}^2 \|f_1\|_{L^2(\mathbb{R}^2)}^2} + \frac{\langle f_1, T_0 f \rangle}{\|f_1\|_{L^2(\mathbb{R}^2)}^2},$$

$$c_1 = \frac{\langle f_2, T_0 f \rangle \langle x_2 v, f_1 \rangle \langle x_1 v, v \rangle}{\|f_2\|_{L^2(\mathbb{R}^2)}^2 \|f_1\|_{L^2(\mathbb{R}^2)}^2 \|V\|_{L^1(\mathbb{R}^2)}} - \frac{\langle f_1, T_0 f \rangle \langle x_1 v, v \rangle}{\|f_1\|_{L^2(\mathbb{R}^2)}^2 \|V\|_{L^1(\mathbb{R}^2)}} + \frac{\langle v, T_0 f \rangle}{\|V\|_{L^1(\mathbb{R}^2)}}$$

$$\text{with } f_1 = xv - \frac{\langle xv, v \rangle}{\|V\|_{L^1(\mathbb{R}^2)}} v, \quad f_2 = x_2 v - \frac{\langle x_2 v, f_1 \rangle}{\|f_1\|_{L^2(\mathbb{R}^2)}^2} f_1 - \frac{\langle x_2 v, v \rangle}{\|V\|_{L^1(\mathbb{R}^2)}} v.$$

Proof. Assume that $f \in S_1 L^2$, one has $S_0(U + vG_0v)f = 0$. Note that

$$0 = S_0(U + vG_0v)f = (I - \tilde{P})(U + vG_0v)f = Uf + vG_0vf - \tilde{P}T_0f,$$

where \tilde{P} is the projection onto the span $\{v, x_1v, x_2v\} = (S_0 L^2)^\perp$. In fact, note that v, f_1 and f_2 are the orthogonal basis of $(S_0 L^2)^\perp$. Hence, we have

$$\begin{aligned} \tilde{P}T_0f &= \frac{\langle f_2, T_0f \rangle}{\|f_2\|_{L^2}^2} f_2 + \frac{\langle f_1, T_0f \rangle}{\|f_1\|_{L^2}^2} f_1 + \frac{\langle v, T_0f \rangle}{\|V\|_{L^1}} v \\ &= x_2 v \frac{\langle f_2, T_0f \rangle}{\|f_2\|_{L^2}^2} + x_1 v \left(-\frac{\langle f_2, T_0f \rangle \langle x_2 v, f_1 \rangle}{\|f_2\|_{L^2}^2 \|f_1\|_{L^2}^2} + \frac{\langle f_1, T_0f \rangle}{\|f_1\|_{L^2}^2} \right) \\ &\quad + v \left(\frac{\langle f_2, T_0f \rangle \langle x_2 v, f_1 \rangle \langle x_1 v, v \rangle}{\|f_2\|_{L^2}^2 \|f_1\|_{L^2}^2 \|V\|_{L^1}} - \frac{\langle f_1, T_0f \rangle \langle x_1 v, v \rangle}{\|f_1\|_{L^2}^2 \|V\|_{L^1}} + \frac{\langle v, T_0f \rangle}{\|V\|_{L^1}} \right) \\ &:= c_3 x_2 v + c_2 x_1 v + c_1 v. \end{aligned}$$

Furthermore, we have

$$Uf = -vG_0vf + \tilde{P}T_0f = -vG_0vf + c_3 x_2 v + c_2 x_1 v + c_1 v.$$

Since $U^2 = 1$, then

$$f = U^2 f = Uv(-G_0vf + c_3 x_2 + c_2 x_1 + c_1) := Uv\phi.$$

Thus, $v f = v U v \phi = V \phi$.

To show that $H\phi = (\Delta^2 + V)(-G_0vf + c_3 x_2 + c_2 x_1 + c_1) = 0$, noting that $\Delta^2(c_3 x_2 + c_2 x_1 + c_1) = 0$ and $G_0 = (\Delta^2)^{-1}$, one has $(\Delta^2 + V)\phi = -vf + V\phi = 0$.

Next we show that $\phi \in L_{-1}^\infty := \{\langle \cdot \rangle^{-1} \phi \in L^\infty\}$. Since $c_3 x_2 + c_2 x_1 + c_1 \in L_{-1}^\infty$, it suffices to show that

$$G_0 v f(x) = \frac{1}{8\pi} \int_{\mathbb{R}^2} |x - y|^2 \ln |x - y| v(y) f(y) dy \in L_{-1}^\infty.$$

In fact, if $f \in S_0L^2$, by the orthogonality $\langle x_1v, f \rangle = \langle x_2v, f \rangle = \langle v, f \rangle = 0$, then

$$\int_{\mathbb{R}^2} |x|^2 \ln |x| v(y) f(y) dy = \int_{\mathbb{R}^2} x \cdot y \ln |x| v(y) f(y) dy = 0,$$

which leads to

$$\begin{aligned} G_0 v f(x) &= \frac{1}{8\pi} \int_{\mathbb{R}^2} |x|^2 (\ln |x-y| - \ln |x|) v(y) f(y) dy \\ &\quad - \frac{1}{4\pi} \int_{\mathbb{R}^2} x \cdot y (\ln |x-y| - \ln |x|) v(y) f(y) dy \\ &\quad + \frac{1}{8\pi} \int_{\mathbb{R}^2} |y|^2 \ln |x-y| v(y) f(y) dy := I_1 + I_2 + I_3. \end{aligned} \quad (5.1)$$

For I_1 , we divide into $|x| > 2|y|$ and $|x| \leq 2|y|$ two cases to prove I_1 . If $|x| > 2|y|$, noting that $\frac{|x-y|-|x|}{|x|} < 1$, using Taylor formula we have

$$\ln \frac{|x-y|}{|x|} = \ln \left(1 + \frac{|x-y|-|x|}{|x|} \right) = O\left(\frac{|y|}{|x|}\right).$$

By Hölder's inequality, one has

$$|I_1| \lesssim |x| \int_{\mathbb{R}^2} |y v(y)| \cdot |f(y)| dy \lesssim |x| \in L_{-1}^\infty.$$

If $|x| \leq 2|y|$, we have

$$\begin{aligned} |I_1| &\lesssim |x|^2 \int_{|x| \leq 2|y|} \left| \ln^+ |x-y| - \ln^+ |x| \right| \cdot |v(y) f(y)| dy \\ &\quad + |x|^2 \int_{|x| \leq 2|y|} \left| \ln^- |x-y| - \ln^- |x| \right| \cdot |v(y) f(y)| dy := I_{11} + I_{12}. \end{aligned}$$

Since $\ln^+ |x-y| \lesssim \ln^+ |x| + \ln^+ \langle y \rangle$, for $|x| > 1$, then

$$\left| \ln^+ |x-y| - \ln^+ |x| \right| \lesssim \left| \ln^+ |x| + \ln^+ \langle y \rangle - \ln^+ |x| \right| = \ln^+ \langle y \rangle \lesssim \langle y \rangle^{0+}, \quad |x| > 1.$$

Note that $1 \lesssim \frac{|y|}{|x|}$, by Hölder's inequality, one has

$$|I_{11}| \lesssim |x|^2 \int_{|x| \leq 2|y|} \frac{|y|}{|x|} \langle y \rangle^{0+} \cdot |v(y) f(y)| dy \lesssim |x| \in L_{-1}^\infty.$$

By Hölder's inequality again, we have

$$\begin{aligned} |I_{12}| &\lesssim |x|^2 \int_{|x| \leq 2|y|} \left| \ln^- |x-y| \right| \cdot |v(y) f(y)| dy + |x|^2 \int_{|x| \leq 2|y|} \left| \ln^- |x| \right| \cdot |v(y) f(y)| dy \\ &\lesssim |x|^2 \int_{|x| \leq 2|y|, |x-y| < 1} -\ln |x-y| \cdot |v(y) f(y)| dy + |x|^2 \int_{|x| \leq 2|y|, |x| < 1} -\ln |x| \cdot |v(y) f(y)| dy \\ &\lesssim |x|^2 \int_{|x-y| < 1} \frac{|y|}{|x|} \frac{1}{|x-y|^{0+}} \cdot |v(y) f(y)| dy + |x|^2 \int_{|x| \leq 2|y|} \frac{|y|}{|x|} \frac{1}{|x|^{0+}} \cdot |v(y) f(y)| dy \\ &\lesssim |x| \int_{|x-y| < 1} \frac{1}{|x-y|^{0+}} \cdot |y v(y) f(y)| dy + |x|^{1-} \int_{|x| \leq 2|y|} |y v(y) f(y)| dy \\ &\lesssim |x| + |x|^{1-} \in L_{-1}^\infty. \end{aligned}$$

Combining this with the computation above we obtain that $I_1 \in L_{-1}^\infty$.

Similarly, we can get $I_2 \in L_{-1}^\infty$.

We turn to estimate I_3 . Note that $\int_{\mathbb{R}^2} \ln^+ |x| v(y) f(y) dy = 0$, one has

$$\begin{aligned} |I_3| &\lesssim \left| \int_{\mathbb{R}^2} \ln |x-y| \cdot |y|^2 v(y) f(y) dy \right| \\ &= \left| \int_{|x-y|<1} \ln |x-y| \cdot |y|^2 v(y) f(y) dy + \int_{\mathbb{R}^2} (\ln^+ |x-y| - \ln^+ |x|) \cdot |y|^2 v(y) f(y) dy \right| \\ &\lesssim \left| \int_{|x-y|<1} \frac{|y|^2 |v(y) f(y)|}{|x-y|^{0+}} dy + \left| \int_{\mathbb{R}^2} (\ln^+ |x-y| - \ln^+ |x|) \cdot |y|^2 v(y) f(y) dy \right|. \end{aligned}$$

Note that, if $|x| \geq 1$, then \ln^+ is a increasing function, and $|x-y| \leq |x|(1+|y|)$, thus

$$\ln^+ |x-y| \leq \ln^+ |x| + \ln^+ |1+y| \lesssim \ln^+ |x| + \ln^+ \langle y \rangle.$$

By Young's inequalities and Hölder's inequalities, we have

$$|I_3| \lesssim \int_{|x-y|<1} \frac{|y|^2 |v(y) f(y)|}{|x-y|^{0+}} dy + \int_{\mathbb{R}^2} \langle y \rangle^{2+} |v(y) f(y)| dy \lesssim c \in L_{-1}^\infty.$$

Combining with all estimates above, we obtain that $G_0 v f \in L_{-1}^\infty$. Moreover, $\phi \in L_{-1}^\infty$.

On the other hand, if there exists nonzero $\phi \in L_{-1}^\infty$ such that $H\phi = 0$ and $f = Uv\phi$, then

$$\Delta^2(\phi + G_0 v f) = \Delta^2(\phi + (\Delta^2)^{-1} v f) = \Delta^2(\phi + Uv f) = (\Delta^2 + V)(\phi) = 0,$$

since $\phi \in L^\infty$ and $G_0 v f \in L_{-1}^\infty$, then $\phi + G_0 v f$ is a polynomial and its degree is less than 1. Hence we assume that $\phi = -G_0 v f + c_3 x_2 + c_2 x_1 + c_1 \in L_{-1}^\infty$. Furthermore, we have

$$\begin{aligned} \phi(x) &= -\frac{1}{8\pi} \int_{\mathbb{R}^2} (|x|^2 - 2x \cdot y + |y|^2) \ln |x-y| v(y) f(y) dy + c_3 x_2 + c_2 x_1 + c_1 \\ &:= -\frac{1}{8\pi} \phi_2(x) + c_3 x_2 + c_2 x_1 + c_1. \end{aligned} \tag{5.2}$$

Since $\phi \in L_{-1}^\infty$ and $c_3 x_2 + c_2 x_1 + c_1 \in L_{-1}^\infty$, which implies that $\phi_2 \in L_{-1}^\infty$. Since

$$\begin{aligned} \phi_2(x) &= \int_{\mathbb{R}^2} (|x|^2 - 2x \cdot y) (\ln |x-y| - \ln |x|) v(y) f(y) dy \\ &\quad + 2 \sum_{i=1}^2 x_i \ln |x| \int_{\mathbb{R}^2} y_i v(y) f(y) dy + |x|^2 \ln |x| \int_{\mathbb{R}^2} v(y) f(y) dy \\ &\quad + \int_{\mathbb{R}^2} |y|^2 \ln |x-y| v(y) f(y) dy, \end{aligned} \tag{5.3}$$

note that the first and fourth integral terms in (5.3) belong to L_{-1}^∞ by the estimates of $I_i (i = 1, 2, 3)$ in (5.1), thus the second and third integral terms in (5.3) are in L_{-1}^∞ , since $x_i \ln |x| \notin L_{-1}^\infty (i = 1, 2)$ and $|x|^2 \ln |x| \notin L_{-1}^\infty$, which implies that

$$\langle v, f \rangle = \langle x_1 v, f \rangle = \langle x_2 v, f \rangle = 0.$$

Hence, $f \in S_0 L^2$. From $f = Uv\phi$ and $H\phi = 0$, we obtain that

$$\begin{aligned} S_0 T_0 S_0 f &= S_0 T_0 (Uv\phi) = S_0 (U + vG_0 f) (Uv\phi) = S_0 (vG_0 v\phi + v\phi) \\ &= S_0 vG_0 (\Delta^2 + V)\phi = S_0 vG_0 H\phi = 0. \end{aligned}$$

Hence, we obtain that $f \in S_1L^2$. \square

Remark 5.2. Assume that $|V(x)| \lesssim (1 + |x|)^{-\beta-}$. For any $p \geq 1$, if $\beta > 2(p + 1)$, then one has the improved estimate for large $|x|$

$$\begin{aligned} \left| \int_{|x| < 2|y|} (\ln|x-y| - \ln|x|)v(y)f(y)dy \right| &\lesssim \int_{\mathbb{R}^2} (\langle y \rangle^{0+} + \ln^-|x-y|) \cdot |v(y)f(y)|dy \\ &\lesssim \frac{1}{|x|^p} \int_{\mathbb{R}^2} (\langle y \rangle^{0+} + \ln^-|x-y|) \langle y \rangle^{p-\beta-} |f(y)|dy \\ &\lesssim \frac{1}{|x|^p}. \end{aligned}$$

Proposition 5.3. Assume that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 18$. Then $f \in S_2L^2(\mathbb{R}^2) \setminus \{0\}$ if and only if $f = Uv\phi$ with $\phi \in L^\infty(\mathbb{R}^2)$ such that $H\phi = 0$ in the distributional sense, and

$$\phi = -G_0vf + \frac{\langle v, T_0f \rangle}{\|V\|_{L^1(\mathbb{R}^2)}}.$$

Proof. Since $S_2 \leq S_1$, by Proposition 5.1, we know that $f \in S_1L^2$ if and only if $f = Uv\phi$ with $\phi \in L^\infty_{-1}$ such that $H\phi = 0$ in the distributional sense and

$$\phi = -G_0vf + c_3x_2 + c_2x_1 + c_1.$$

Thus, we just need to prove that $c_1 = \frac{\langle v, T_0f \rangle}{\|V\|_{L^1}}$, $c_2 = c_3 = 0$, and $\phi \in L^\infty$. Indeed, by the proof of Proposition 5.1, we know that

$$f_1 = xv - \frac{\langle xv, v \rangle}{\|V\|_{L^1}}v \in QL^2, \quad f_2 = x_2v - \frac{\langle x_2v, f_1 \rangle}{\|f_1\|_{L^2}^2}f_1 - \frac{\langle x_2v, v \rangle}{\|V\|_{L^1}}v \in QL^2.$$

If $f \in S_2L^2$, then $QT_0f = 0$. Hence

$$\langle f_i, T_0f \rangle = \langle Qf_i, T_0f \rangle = \langle f_i, QT_0f \rangle = 0, \quad i = 1, 2,$$

which leads to $c_1 = \frac{\langle v, T_0f \rangle}{\|V\|_{L^1}}$, and $c_2 = c_3 = 0$.

Next we show that $\phi \in L^\infty$. Since $\frac{\langle v, T_0f \rangle}{\|V\|_{L^1}} \in L^\infty$, it suffices to show that

$$G_0vf(x) = \frac{1}{8\pi} \int_{\mathbb{R}^2} |x-y|^2 \ln|x-y|v(y)f(y)dy \in L^\infty.$$

If $f \in S_2L^2 \subset S_1L^2$, using the orthogonality $S_2v = 0, S_2x_i v = 0 (i = 1, 2)$, then

$$\int_{\mathbb{R}^2} |x|^2 \ln|x|v(y)f(y)dy = \int_{\mathbb{R}^2} x \cdot y \ln|x|v(y)f(y)dy = \int_{\mathbb{R}^2} \frac{x \cdot y}{|x|^2}v(y)f(y)dy = 0.$$

Furthermore, we have

$$\begin{aligned} G_0vf(x) &= \frac{1}{16\pi} \int_{\mathbb{R}^2} |x|^2 \left[\ln \left(\frac{|x-y|^2}{|x|^2} \right) + 2 \frac{x \cdot y}{|x|^2} \right] v(y)f(y)dy \\ &\quad - \frac{1}{4\pi} \int_{\mathbb{R}^2} x \cdot y (\ln|x-y| - \ln|x|)v(y)f(y)dy \\ &\quad + \frac{1}{8\pi} \int_{\mathbb{R}^2} |y|^2 \ln|x-y|v(y)f(y)dy := I_1 + I_2 + I_3. \end{aligned} \tag{5.4}$$

For I_1 , we divide into $|x| > 4|y|$ and $|x| \leq 4|y|$ two cases to prove I_1 . If $|x| > 4|y|$, by Taylor formula, one has

$$\ln\left(\frac{|x-y|^2}{|x|^2}\right) + 2\frac{x \cdot y}{|x|^2} = \ln\left(1 + \frac{|x-y|^2 - |x|^2}{|x|^2}\right) + 2\frac{x \cdot y}{|x|^2} = \frac{|y|^2}{|x|^2} + O\left(\frac{|y|^{2+}}{|x|^{2+}}\right).$$

By Hölder's inequality, we obtain

$$\begin{aligned} |I_1| &\lesssim \left| \int_{|x|>4|y|} |x|^2 \left[\ln\left(\frac{|x-y|^2}{|x|^2}\right) + 2\frac{x \cdot y}{|x|^2} \right] v(y)f(y)dy \right| \\ &\lesssim \int_{\mathbb{R}^2} |y|^2 |v(y)| \cdot |f(y)| dy \lesssim c. \end{aligned}$$

If $|x| \leq 4|y|$, we have

$$\begin{aligned} |I_1| &\lesssim \left| \int_{|x|\leq 4|y|} |x|^2 \left[\ln\left(\frac{|x-y|^2}{|x|^2}\right) + 2\frac{x \cdot y}{|x|^2} \right] v(y)f(y)dy \right| \\ &\lesssim |x|^2 \left| \int_{|x|\leq 4|y|} (\ln|x-y| - \ln|x|)v(y)f(y)dy \right| + |x|^2 \int_{|x|\leq 4|y|} \frac{|y|^2}{|x|^2} |v(y)f(y)| dy \\ &\lesssim |x|^2 \int_{|x|\leq 4|y|} \left| \ln^+ |x-y| - \ln^+ |x| \right| \cdot |v(y)f(y)| dy \\ &\quad + |x|^2 \int_{|x|\leq 4|y|} \left| \ln^- |x-y| - \ln^- |x| \right| \cdot |v(y)f(y)| dy + \int_{|x|\leq 4|y|} |y|^2 |v(y)f(y)| dy. \end{aligned}$$

Noting that $1 \lesssim \frac{|y|^2}{|x|^2}$, by the same argument with the proof of the terms I_{11} and I_{12} in Proposition 5.1, we get that $I_1 \in L^\infty$ in the case of $|x| \leq 4|y|$. Moreover, we get $I_1 \in L^\infty$.

For I_2 and I_3 , by using the same argument with Proposition 5.1, we obtain that both I_2 and I_3 belong to L^∞ .

Combining all computations above, we obtain that $G_0vf \in L^\infty$. Moreover, $\phi \in L^\infty$.

On the other hand, if there exists nonzero $\phi \in L^\infty$ such that $H\phi = 0$ and $f = Uv\phi$, then

$$\Delta^2(\phi + G_0vf) = \Delta^2(\phi + (\Delta^2)^{-1}vf) = \Delta^2(\phi + Uvf) = 0.$$

Since $\phi \in L^\infty$ and $G_0vf \in L_{-1}^\infty$, then $\phi + G_0vf$ is a polynomial and its degree is zero, hence we may assume that $\phi = -G_0vf + c \in L^\infty$. Furthermore, we have

$$\begin{aligned} \phi(x) &= -\frac{1}{8\pi} \int_{\mathbb{R}^2} (|x|^2 - 2x \cdot y + |y|^2) \ln|x-y| v(y)f(y) dy + \frac{\langle v, T_0 f \rangle}{\|V\|_{L^1}} \\ &:= -\frac{1}{8\pi} \phi_2(x) + c. \end{aligned}$$

Since $\phi \in L^\infty$ and $c \in L^\infty$, which implies that $\phi_2 \in L^\infty$. Since

$$\begin{aligned} \phi_2(x) &= \int_{\mathbb{R}^2} |x|^2 \left[\ln \left(\frac{|x-y|^2}{|x|^2} \right) + 2 \frac{x \cdot y}{|x|^2} \right] v(y) f(y) dy \\ &\quad - \int_{\mathbb{R}^2} 2x \cdot y (\ln |x-y| - \ln |x|) v(y) f(y) dy + 2 \sum_{i=1}^2 x_i \ln |x| \int_{\mathbb{R}^2} y_i v(y) f(y) dy \\ &\quad + |x|^2 \ln |x| \int_{\mathbb{R}^2} v(y) f(y) dy + \int_{\mathbb{R}^2} |y|^2 \ln |x-y| v(y) f(y) dy \\ &\quad + |x|^2 \ln |x|^2 \int_{\mathbb{R}^2} v(y) f(y) dy - 2 \int_{\mathbb{R}^2} (x \cdot y) v(y) f(y) dy, \end{aligned} \quad (5.5)$$

note that the first, second and fifth integral terms in (5.5) are in L^∞ by the estimates of $I_i (i = 1, 2, 3)$ in (5.4); thus the other integral terms in (5.5) are in L^∞ , while these functions $x_i \ln |x| \notin L^\infty (i = 1, 2)$ and $|x|^2 \ln |x|$ are not in L^∞ . So we have

$$\langle v, f \rangle = \langle x_1 v, f \rangle = \langle x_2 v, f \rangle = 0.$$

The proof of this lemma is completed. \square

Proposition 5.4. *Assume that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 18$. Then $f \in S_4 L^2(\mathbb{R}^2) \setminus \{0\}$ if and only if $f = Uv\phi$ with $\phi \in L_1^\infty(\mathbb{R}^2)$ such that $H\phi = 0$ in the distributional sense, and*

$$\phi = -G_0 v f.$$

Proof. Since $S_4 \leq S_3 \leq S_2$, by Proposition 5.3, we know that $f \in S_2 L^2$ if and only if $f = Uv\phi$ with $\phi \in L^\infty$ such that $H\phi = 0$ in the distributional sense and

$$\phi = -G_0 v f + \frac{\langle v, T_0 f \rangle}{\|V\|_{L^1}}.$$

Thus, we only need to prove that $\langle v, T_0 f \rangle = 0$ and $\phi \in L_1^\infty$. Indeed, if $f \in S_4 L^2$, then $\frac{\langle v, T_0 f \rangle}{\|V\|_{L^1}} = 0$. Moreover, $\phi = -G_0 v f$.

Next we begin to prove $\phi = -G_0 v f \in L_1^\infty$. Indeed, if $f \in S_4 L^2$, then

$$\int_{\mathbb{R}^2} x \cdot y \ln |x| v(y) f(y) dy = \int_{\mathbb{R}^2} |x|^2 \ln |x| v(y) f(y) dy = \int_{\mathbb{R}^2} \ln |x| |y|^2 v(y) f(y) dy = 0,$$

which leads to

$$\begin{aligned} G_0 v f(x) &= \frac{1}{16\pi} \int_{\mathbb{R}^2} |x|^2 \left[\ln \left(\frac{|x-y|^2}{|x|^2} \right) + 2 \frac{x \cdot y}{|x|^2} - \frac{|y|^2}{|x|^2} \right] v(y) f(y) dy \\ &\quad - \frac{1}{4\pi} \int_{\mathbb{R}^2} x \cdot y \left[\ln \left(\frac{|x-y|^2}{|x|^2} \right) + 2 \frac{x \cdot y}{|x|^2} \right] v(y) f(y) dy \\ &\quad + \frac{1}{8\pi} \int_{\mathbb{R}^2} |y|^2 (\ln |x-y| - \ln |x|) v(y) f(y) dy \\ &:= I_1 + I_2 + I_3. \end{aligned} \quad (5.6)$$

For I_1 . We divide into $|x| > 4|y|$ and $|x| \leq 4|y|$ two cases to estimate I_1 . If $|x| > 4|y|$. By Taylor formula, one has

$$\ln \left(\frac{|x-y|^2}{|x|^2} \right) + 2 \frac{x \cdot y}{|x|^2} - \frac{|y|^2}{|x|^2} = O\left(\frac{|y|^3}{|x|^3}\right).$$

Using Hölder inequality, one has

$$I_1 \lesssim |x|^2 \int_{|x|>4|y|} \frac{|y|^3}{|x|^3} |v(y)f(y)| dy \lesssim \frac{1}{|x|} \int_{\mathbb{R}^2} |y|^3 |v(y)f(y)| dy \lesssim \frac{1}{|x|}.$$

If $|x| \leq 4|y|$. Using Remark 5.2, for large $|x|$ we have

$$\left| \int_{|x| \leq 4|y|} \ln \left(\frac{|x-y|^2}{|x|^2} \right) v(y)f(y) dy \right| \lesssim \frac{1}{|x|^3}. \quad (5.7)$$

Note that $|x \cdot y| \leq |y|^2$ and $1 \leq \frac{|y|^2}{|x|^2}$, using (5.7) and Hölder's inequality, one has

$$\begin{aligned} |I_1| &\lesssim |x|^2 \left| \int_{|x| \leq 4|y|} \left[\ln \left(\frac{|x-y|^2}{|x|^2} \right) + 2 \frac{x \cdot y}{|x|^2} - \frac{|y|^2}{|x|^2} \right] v(y)f(y) dy \right| \\ &\lesssim |x|^2 \frac{1}{|x|^3} + \int_{|x| \leq 4|y|} |y|^2 |v(y)f(y)| dy \\ &\lesssim \frac{1}{|x|} + \int_{|x| \leq 4|y|} \frac{|y|}{|x|} |y|^2 |v(y)f(y)| dy \lesssim \frac{1}{|x|}. \end{aligned}$$

Hence, combining the two cases above, we get that I_1 is controlled by $|x|^{-1} \in L_1^\infty$.

Similarly, we obtain that both I_2 and I_3 are controlled by $|x|^{-1}$ which belongs to L_1^∞ .

On the other hand, if there exists nonzero $\phi \in L_1^\infty$ such that $H\phi = 0$ and $f = Uv\phi$, then

$$\Delta^2(\phi + G_0vf) = \Delta^2(\phi + (\Delta^2)^{-1}vf) = \Delta^2(\phi + Uvf) = 0,$$

hence $\phi + G_0vf$ is a polynomial and its degree is less than 3. Since $\phi \in L_1^\infty$ and $G_0vf \in L^\infty$, thus we have $\phi = -G_0vf \in L_1^\infty$. Furthermore,

$$\phi(x) = \frac{1}{8\pi} \int_{\mathbb{R}^2} |x-y|^2 \ln |x-y| v(y)f(y) dy := \frac{1}{8\pi} \phi_3(x).$$

Since

$$\begin{aligned} \phi_3(x) &= \frac{1}{2} \int_{\mathbb{R}^2} |x|^2 \left[\ln \left(\frac{|x-y|^2}{|x|^2} \right) + 2 \frac{x \cdot y}{|x|^2} - \frac{|y|^2}{|x|^2} \right] v(y)f(y) dy - \int_{\mathbb{R}^2} x \cdot y \\ &\quad \times \left[\ln \left(\frac{|x-y|^2}{|x|^2} \right) + 2 \frac{x \cdot y}{|x|^2} \right] v(y)f(y) dy + \frac{1}{2} \int_{\mathbb{R}^2} |y|^2 (\ln |x-y| - \ln |x|) \\ &\quad \times v(y)f(y) dy + |x|^2 \ln |x| \int_{\mathbb{R}^2} v(y)f(y) dy - \sum_{i=1}^2 x_i \int_{\mathbb{R}^2} y_i v(y)f(y) dy \\ &\quad + \frac{1}{2} \int_{\mathbb{R}^2} |y|^2 v(y)f(y) dy - 2 \sum_{i=1}^2 x_i \ln |x| \int_{\mathbb{R}^2} y_i v(y)f(y) dy \\ &\quad + 2 \sum_{i=1}^2 \frac{x_i^2}{|x|^2} \int_{\mathbb{R}^2} y_i^2 v(y)f(y) dy + \frac{4x_1x_2}{|x|^2} \int_{\mathbb{R}^2} y_1y_2 v(y)f(y) dy \\ &\quad - \frac{1}{2} \ln |x| \int_{\mathbb{R}^2} |y|^2 v(y)f(y) dy, \end{aligned} \quad (5.8)$$

note that the first three integral terms in (5.8) are in L_1^∞ by the estimates of $I_i (i = 1, 2, 3)$ in (5.6), thus the other integral terms in (5.8) are also in L_1^∞ , but these functions $|x|^2 \ln |x|$,

$\ln|x|$, c , x_i , $x_i \ln|x|$, $\frac{x_i^2}{|x|^2}$, $\frac{x_i x_j}{|x|^2}$ ($i, j = 1, 2$) are not in L_1^∞ , hence we have

$$\langle v, f \rangle = \langle x_i v, f \rangle = \langle x_i x_j v, f \rangle = 0, \quad i, j = 1, 2.$$

Since $f = Uv\phi = -UvG_0vf$, which implies

$$0 = f + UvG_0vf = U(U + vG_0v)f = UT_0f.$$

Hence $T_0f = 0$, that is, $f \in S_4L^2$. \square

Proposition 5.5. *Assume that $|V(x)| \lesssim (1 + |x|)^{-\beta}$ with $\beta > 18$. Then $f \in S_5L^2(\mathbb{R}^2) \setminus \{0\}$ if and only if $f = Uv\phi$ with $\phi \in L_2^\infty(\mathbb{R}^2)$ such that $H\phi = 0$ in the distributional sense, and*

$$\phi = -G_0vf.$$

Remark 5.6. *Noting that $L_2^\infty(\mathbb{R}^2) \subset L^2(\mathbb{R}^2)$, hence the fourth kind zero resonance is actually an eigenvalue of H . Conversely, let $\phi \in L^2(\mathbb{R}^2)$ such that $H\phi = 0$, then we can adapt the proof of Proposition 5.5 to show the eigenfunction $\phi \in L_2^\infty(\mathbb{R}^2)$, which implies that zero eigenvalue is exactly the fourth kind resonance from Definition 2.4.*

Proof. Since $S_5 \leq S_4$, by Proposition 5.4, we know that $f \in S_4L^2$ if and only if $f = Uv\phi$ with $\phi \in L_1^\infty$ such that $H\phi = 0$ in the distributional sense and

$$\phi = -G_0vf.$$

Thus, we just need to prove that $\phi = -G_0vf \in L_2^\infty$.

Now we begin to prove $\phi = -G_0vf \in L_2^\infty$. Indeed, if $f \in S_5L^2$, then

$$\langle v, f \rangle = \langle x_j v, f \rangle = \langle x_i x_j v, f \rangle = \langle x_i x_j x_k v, f \rangle = 0, \quad i, j, k = 1, 2.$$

Moreover, we obtain that

$$\begin{aligned} G_0vf(x) &= \frac{1}{32\pi}|x|^2 \int_{\mathbb{R}^2} \left[\ln\left(\frac{|x-y|^4}{|x|^4}\right) + 4\frac{x \cdot y}{|x|^2} - 2\frac{|y|^2}{|x|^2} + 4\frac{(x \cdot y)^2}{|x|^4} \right] v(y)f(y)dy \\ &\quad - \frac{1}{4\pi} \int_{\mathbb{R}^2} x \cdot y \left[\ln\left(\frac{|x-y|^2}{|x|^2}\right) + 2\frac{x \cdot y}{|x|^2} - \frac{|y|^2}{|x|^2} \right] v(y)f(y)dy \\ &\quad + \frac{1}{16\pi} \int_{\mathbb{R}^2} |y|^2 \left[\ln\left(\frac{|x-y|^2}{|x|^2}\right) + 2\frac{x \cdot y}{|x|^2} \right] v(y)f(y)dy \\ &:= I_1 + I_2 + I_3. \end{aligned} \tag{5.9}$$

For I_1 , we divide into $|x| > 16|y|$ and $|x| \leq 16|y|$ two cases to compute I_1 . If $|x| > 16|y|$, since

$$|x-y|^4 = |x|^4 + |y|^4 - 4x \cdot y|y|^2 - 4x \cdot y|x|^2 + 2|x|^2|y|^2 + 4(x \cdot y)^2,$$

then

$$\left| |x-y|^4 - |x|^4 \right| = \left| |y|^4 - 4x \cdot y|y|^2 - 4x \cdot y|x|^2 + 2|x|^2|y|^2 + 4(x \cdot y)^2 \right| < |x|^4.$$

By Taylor formula we have

$$\begin{aligned} \ln\left(\frac{|x-y|^4}{|x|^4}\right) &= \ln\left(1 + \frac{|x-y|^4 - |x|^4}{|x|^4}\right) \\ &= -4\frac{x \cdot y}{|x|^2} + 2\frac{|y|^2}{|x|^2} + 4\frac{(x \cdot y)^2}{|x|^4} + O\left(\frac{|y|^4}{|x|^4}\right). \end{aligned} \tag{5.10}$$

Thus, we have

$$\ln\left(\frac{|x-y|^4}{|x|^4}\right) + 4\frac{x \cdot y}{|x|^2} - 2\frac{|y|^2}{|x|^2} + 4\frac{(x \cdot y)^2}{|x|^4} = O\left(\frac{|y|^4}{|x|^4}\right).$$

By Hölder inequality, we obtain

$$\begin{aligned} |I_1| &\lesssim |x|^2 \left| \int_{|x|>16|y|} \left[\ln\left(\frac{|x-y|^4}{|x|^4}\right) + 4\frac{x \cdot y}{|x|^2} - 2\frac{|y|^2}{|x|^2} + 4\frac{(x \cdot y)^2}{|x|^4} \right] v(y)f(y)dy \right| \\ &\lesssim |x|^2 \int_{|x|>16|y|} \frac{|y|^4}{|x|^4} |v(y)f(y)| dy \lesssim \frac{1}{|x|^2}. \end{aligned}$$

If $|x| \leq 16|y|$, using remark 5.2 we have

$$\left| \int_{|x|\leq 4|y|} \ln\left(\frac{|x-y|^2}{|x|^2}\right) v(y)f(y)dy \right| \lesssim \frac{1}{|x|^4}. \quad (5.11)$$

Note that $|x \cdot y| \leq |y|^2$ and $1 \leq \frac{|y|^4}{|x|^4}$, then using (5.11) and Hölder inequality, we obtain that I_1 is also controlled by $|x|^{-2}$ in this case.

Combining with the computations above, we obtain that I_1 is bounded by $|x|^{-2}$.

We obtain that I_2 and I_3 are controlled by $|x|^{-2}$ by using similar method with the proof of Proposition 5.4. Hence G_0vf are bounded by $|x|^{-2}$ which belongs to L_2^∞ .

On the other hand, if there exists nonzero $\phi \in L_2^\infty$ such that $H\phi = 0$ and $f = Uv\phi$, then we have $\phi = -G_0vf \in L_2^\infty$ by the same argument with the proof of Proposition 5.4. Furthermore,

$$\phi(x) = \frac{1}{8\pi} \int_{\mathbb{R}^2} |x-y|^2 \ln|x-y| v(y)f(y)dy := \frac{1}{8\pi} \phi_4(x).$$

Noting that

$$\begin{aligned} \phi_4(x) &= \frac{1}{4}|x|^2 \int_{\mathbb{R}^2} \left[\ln\left(\frac{|x-y|^4}{|x|^4}\right) + 4\frac{x \cdot y}{|x|^2} - 2\frac{|y|^2}{|x|^2} + 4\frac{(x \cdot y)^2}{|x|^4} \right] v(y)f(y)dy \\ &\quad - \int_{\mathbb{R}^2} x \cdot y \left[\ln\left(\frac{|x-y|^2}{|x|^2}\right) + 2\frac{x \cdot y}{|x|^2} - \frac{|y|^2}{|x|^2} \right] v(y)f(y)dy \\ &\quad + \frac{1}{2} \int_{\mathbb{R}^2} |y|^2 \left[\ln\left(\frac{|x-y|^2}{|x|^2}\right) + 2\frac{x \cdot y}{|x|^2} \right] v(y)f(y)dy + |x|^2 \ln|x| \int_{\mathbb{R}^2} v(y)f(y)dy \\ &\quad - \int_{\mathbb{R}^2} x \cdot y v(y)f(y)dy + \frac{1}{2} \int_{\mathbb{R}^2} |y|^2 v(y)f(y)dy - \frac{1}{|x|^2} \int_{\mathbb{R}^2} (x \cdot y)^2 v(y)f(y)dy \quad (5.12) \\ &\quad - 2 \ln|x| \int_{\mathbb{R}^2} x \cdot y v(y)f(y)dy + \frac{2}{|x|^2} \int_{\mathbb{R}^2} (x \cdot y)^2 v(y)f(y)dy \\ &\quad - \frac{1}{|x|^2} \int_{\mathbb{R}^2} x \cdot y |y|^2 v(y)f(y)dy - \ln|x| \int_{\mathbb{R}^2} |y|^2 v(y)f(y)dy \\ &\quad - \frac{1}{|x|^2} \int_{\mathbb{R}^2} x \cdot y v(y)f(y)dy, \end{aligned}$$

since the first three integral terms in (5.12) are in L_2^∞ by the estimates of I_i ($i = 1, 2, 3$) in (5.9), thus the other integral terms in (5.12) are in L_2^∞ , which leads to

$$\langle v, f \rangle = \langle x_j v, f \rangle = \langle x_i x_j v, f \rangle = \langle x_i x_j x_k v, f \rangle = 0, \quad i, j, k = 1, 2.$$

Hence the proof of this lemma is completed. \square

As the consequence of Definition 2.4 and Propositions 5.1–5.5, we immediately have the following theorem.

Theorem 5.7. *Let $H = \Delta^2 + V$ on $L^2(\mathbb{R}^2)$ and $|V(x)| \lesssim (1 + |x|)^{-\beta}$ for some $\beta > 0$. Then we have following statements:*

(i) *If $\beta > 10$, then zero is the regular point of H if and only if there exists only zero solution $\phi(x) \in L_{-1}^\infty(\mathbb{R}^2)$ such that $H\phi = 0$ in the distributional sense,*

(ii) *If $\beta > 14$, then zero is the first kind resonance of H if and if there exists some nonzero $\phi(x) \in L_{-1}^\infty(\mathbb{R}^2)$ but no nonzero $\phi \in L^\infty(\mathbb{R}^2)$ such that $H\phi = 0$.*

(iii) *If $\beta > 18$, then zero is the second kind resonance of H if and only if there exists some nonzero $\phi(x) \in L^\infty(\mathbb{R}^2)$ but no nonzero $\phi \in L_1^\infty(\mathbb{R}^2)$ such that $H\phi = 0$.*

(iv) *If $\beta > 18$, then zero is the third kind resonance of H if and only if there exists some nonzero $\phi(x) \in L_1^\infty(\mathbb{R}^2)$ but no nonzero $\phi \in L^2(\mathbb{R}^2)$ such that $H\phi = 0$.*

(v) *If $\beta > 18$, then zero is the fourth kind resonance (i.e. eigenvalue) of H if and only if there exists some nonzero $\phi(x) \in L^2(\mathbb{R}^2)$ such that $H\phi = 0$.*

Let $\phi \in C^\infty(\mathbb{R}^2)$ be a positive function which is equal to $c \ln |x| + d$ for $|x| > 1$, where c, d are some positive constants. Then we can easily check that $H\phi(x) = 0$ when taking $V(x) = -(\Delta^2\phi)/\phi$. It is obvious that $V(x) \in C_0^\infty(\mathbb{R}^2)$ and $\phi(x) \in L_{-1}^\infty(\mathbb{R}^2)$, hence by Theorem 5.7, we know that the zero is a resonance point of H . On the other hand, as shown in Remark 1.2 of [SWY21], zero cannot be an eigenvalue of H in one dimensional case, assuming that V has fast enough decay. However, it would be an interesting problem to find whether there exist some sufficient decay potentials V such that zero is an eigenvalue of $H = \Delta^2 + V$ in \mathbb{R}^2 .

Acknowledgements: A. Soffer is partially supported by the Simon’s Foundation (No.395767). X. H. Yao and Ping Li are partially supported by NSFC (No.11771165 and 12171182). We would like to thank Dr. Z. Wu for his useful discussion.

REFERENCES

- [AS64] M. Abramowitz, I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards Applied Mathematics Series, vol. 55, U.S. Government Printing Office, Washington, D.C., 1964. For sale by Superintendent of Documents.
- [Agm75] Shmuel Agmon, *Spectral properties of Schrödinger operators and scattering theory*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **2** (1975), no. 2, 151–218. MR 0397194
- [BKS00] Matania Ben-Artzi, Herbert Koch, and Jean-Claude Saut, *Dispersion estimates for fourth order Schrödinger equations*, C. R. Acad. Sci. Paris Sér. I Math. **330** (2000), no. 2, 87–92. MR 1745182
- [CS01] O. Costin and A. Soffer, *Resonance theory for Schrödinger operators*, Comm. Math. Phys. **224** (2001), no. 1, 133–152, Dedicated to Joel L. Lebowitz. MR 1868995
- [EG13] M. Burak Erdoğan and William R. Green, *Dispersive estimates for Schrödinger operators in dimension two with obstructions at zero energy*, Trans. Amer. Math. Soc. **365** (2013), no. 12, 6403–6440. MR 3105757
- [EG21] M. Burak Erdoğan and William R. Green, *The L^p -continuity of wave operator for higher order Schrödinger operators*, <https://arxiv.org/pdf/2107.09620.pdf>.

- [EGT19] M. Burak Erdoğan, William R. Green, and Ebru Toprak, *On the fourth order schrödinger equation in three dimensions: dispersive estimates and zero energy resonance*, J. Differential Equations, **217**(2021), 152-185.
- [ES04] M. Burak Erdoğan and Wilhelm Schlag, *Dispersive estimates for Schrödinger operators in the presence of a resonance and/or an eigenvalue at zero energy in dimension three. I*, Dyn. Partial Differ. Equ. **1** (2004), no. 4, 359–379. MR 2127577
- [ES06] M. Burak Erdoğan and Wilhelm Schlag, *Dispersive estimates for Schrödinger operators in the presence of a resonance and/or an eigenvalue at zero energy in dimension three. II*, J. Anal. Math. **99** (2006), 199–248. MR 2279551
- [FSWY20] Hongliang Feng, Avy Soffer, Zhao Wu and Xiaohua Yao, *Decay estimates for higher-order elliptic operators*, Trans. Amer. Math. Soc. **373** (2020), no. 4, 2805-2859. MR 3724151
- [FSY18] Hongliang Feng, Avy Soffer, and Xiaohua Yao, *Decay estimates and Strichartz estimates of fourth-order Schrödinger operator*, J. Funct. Anal. **274** (2018), no. 2, 605–658. MR 3724151
- [FWY18] Hongliang Feng, Zhao Wu, and Xiaohua Yao, *Time asymptotic expansions of solution for fourth-order schrödinger equation with zero resonance or eigenvalue*, <https://arxiv.org/abs/1812.00223> (2018).
- [GG15] Michael Goldberg and William R. Green, *Dispersive estimates for higher dimensional Schrödinger operators with threshold eigenvalues I: The odd dimensional case*, J. Funct. Anal. **269** (2015), no. 3, 633–682. MR 3350725
- [GG17] Michael Goldberg and William R. Green, *Dispersive estimates for higher dimensional Schrödinger operators with threshold eigenvalues II. The even dimensional case*, J. Spectr. Theory **7** (2017), no. 1, 33–86. MR 3629407
- [GG21] Michael Goldberg and William R. Green, *On the L^p boundedness of the wave operators for fourth order Schrödinger operator*, Trans. Amer. Math. Soc. **374** (2021), no. 6, 4075–4092.
- [GS04] M. Goldberg and W. Schlag, *Dispersive estimates for Schrödinger operators in dimensions one and three*, Comm. Math. Phys. **251** (2004), no. 1, 157–178. MR 2096737
- [GT19] William R. Green and Ebru Toprak, *On the fourth order schrödinger equation in four dimensions: dispersive estimates and zero energy resonance*, J. Differential Equations, **267**(2019) 1899-1954.
- [Hill20] Thomas Hill, *Dispersive estimates of Schrödinger and Schrödinger-like equations in one dimension*, Thesis (Ph.D.)–University of Cincinnati. 2020. MR 4187680
- [Jen80] Arne Jensen, *Spectral properties of Schrödinger operators and time-decay of the wave functions results in $L^2(\mathbf{R}^m)$, $m \geq 5$* , Duke Math. J. **47** (1980), no. 1, 57–80. MR 563367
- [JK79] Arne Jensen and Tosio Kato, *Spectral properties of Schrödinger operators and time-decay of the wave functions*, Duke Math. J. **46** (1979), no. 3, 583–611. MR 544248
- [JN01] Arne Jensen and Gheorghe Nenciu, *A unified approach to resolvent expansions at thresholds*, Rev. Math. Phys. **13** (2001), no. 6, 717–754. MR 1841744
- [JN04] ———, *Erratum: “A unified approach to resolvent expansions at thresholds” [Rev. Math. Phys. **13** (2001), no. 6, 717–754; mr1841744]*, Rev. Math. Phys. **16** (2004), no. 5, 675–677. MR 2079122
- [JSS91] J.-L. Journé, A. Soffer, and C. D. Sogge, *Decay estimates for Schrödinger operators*, Comm. Pure Appl. Math. **44** (1991), no. 5, 573–604. MR 1105875
- [Kat59] Tosio Kato, *Growth properties of solutions of the reduced wave equation with a variable coefficient*, Comm. Pure Appl. Math. **12** (1959), 403–425. MR 0108633
- [LWY21] Ping Li, Zijun Wan and Xiaohua Yao, *The improved decay estimates for the fourth-order Schrödinger operators in dimension three*, Preprint.
- [Mur82] Minoru Murata, *Asymptotic expansions in time for solutions of Schrödinger-type equations*, J. Funct. Anal. **49** (1982), no. 1, 10–56. MR 680855
- [RS04] Igor Rodnianski and Wilhelm Schlag, *Time decay for solutions of Schrödinger equations with rough and time-dependent potentials*, Invent. Math. **155** (2004), no. 3, 451–513. MR 2038194
- [Sch05] W. Schlag, *Dispersive estimates for Schrödinger operators in dimension two*, Comm. Math. Phys. **257** (2005), no. 1, 87–117. MR 2163570
- [Sch07] W. Schlag, *Dispersive estimates for Schrödinger operators: a survey*, Mathematical aspects of nonlinear dispersive equations, Ann. of Math. Stud., vol. 163, Princeton Univ. Press, Princeton, NJ, 2007, pp. 255–285. MR 2333215

- [Sch21] W. Schlag, *On pointwise decay of waves*, J. Math. Phys. **62** (2021), no. 6, Paper No. 061509, 27 pp. MR 4276283
- [SWY21] Avy Soffer, Zhao Wu and Xiaohua Yao, *Decay estimates for bi-Schrödinger operators in dimension one*, <https://arxiv.org/pdf/2106.15966>.
- [Ste93] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series, vol. 43, Princeton University Press, Princeton, NJ, 1993, With the assistance of Timothy S. Murphy, Monographs in Harmonic Analysis, III. MR 1232192
- [Wed99] Ricardo A. Weder, *The $W^{k,p}$ -continuity of the Schrödinger wave operators on the line*, Comm. Math. Phys. **208** (1999), no. 2, 507–520. MR 1729096
- [Wed00] Ricardo A. Weder, *$L^p - L^{p'}$ estimates for the Schrödinger equation on the line and inverse scattering for the non linear Schrödinger equation with a potential*, J. Funct. Analysis **170**(2000), 37–68.
- [Yaj95] Kenji Yajima, *The $W^{k,p}$ -continuity of wave operators for Schrödinger operators*, J. Math. Soc. Japan **47** (1995), no. 3, 551–581. MR 1331331

PING LI, SCHOOL OF INFORMATIONS AND MATHEMATICS , YANGTZE UNIVERSITY, JINGZHOU 434000, CHINA

Email address: liping@whu.edu.cn

AVY SOFFER, MATHEMATICS DEPARTMENT, RUTGERS UNIVERSITY, NEW BRUNSWICK, NJ, 08903, USA

Email address: soffer@math.rutgers.edu

XIAOHUA YAO, DEPARTMENT OF MATHEMATICS AND HUBEI PROVINCE KEY LABORATORY OF MATHEMATICAL PHYSICS, CENTRAL CHINA NORMAL UNIVERSITY, WUHAN, 430079, P.R. CHINA

Email address: yaoxiaohua@mail.ccnu.edu.cn