

SHUFFLE ALGEBRAS, LATTICE PATHS AND THE COMMUTING SCHEME

ALEXANDR GARBALI AND PAUL ZINN-JUSTIN

ABSTRACT. The commutative trigonometric shuffle algebra A is a space of symmetric rational functions satisfying certain wheel conditions [FO97, FHH⁺09]. We describe a ring isomorphism between A and the center of the Hecke algebra using a realization of the elements of A as partition functions of coloured lattice paths associated to the R -matrix of $\mathcal{U}_{t^{1/2}}(\widehat{\mathfrak{gl}}_\infty)$. As an application, we compute under certain conditions the Hilbert series of the *commuting scheme* and identify it with a particular element of the shuffle algebra A , thus providing a combinatorial formula for it as a “domain wall” type partition function of coloured lattice paths.

1. INTRODUCTION

1.1. **Center of Hecke and shuffle products.** Consider the *Hecke algebra* \mathcal{H}_n ($n \in \mathbb{Z}_{\geq 0}$), which we define for our purposes to be the algebra (over a field extension \mathbb{F} of $\mathbb{Q}(t)$) to be specified below) with generators T_1, \dots, T_{n-1} and relations

$$T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1} \quad T_i^2 = (t-1)T_i + t \quad T_i T_j = T_j T_i \quad (|i-j| \geq 2)$$

The standard basis $(T_w)_{w \in \mathcal{S}_n}$ of \mathcal{H}_n is obtained by writing $T_w := T_{i_1} \dots T_{i_k}$ if $w = s_{i_1} \dots s_{i_k}$ is a reduced word in the elementary transpositions s_i . Denote $|w| = k$ the length of w .

Combine the centers $Z(\mathcal{H}_n)$ into a graded algebra

$$Z := \bigoplus_{n \geq 0} Z(\mathcal{H}_n)$$

by defining a shuffle product

$$(1) \quad * : Z(\mathcal{H}_k) \otimes Z(\mathcal{H}_\ell) \rightarrow Z(\mathcal{H}_{k+\ell})$$

$$x \mapsto \sum_{w \in \mathcal{S}^{k,\ell}} t^{-|w|} T_w x T_{w^{-1}}$$

where $\mathcal{H}_k \otimes \mathcal{H}_\ell$ is embedded inside $\mathcal{H}_{k+\ell}$ in the obvious way, and $\mathcal{S}^{k,\ell}$ is the set of shortest representatives of cosets in $\mathcal{S}_{k+\ell}/(\mathcal{S}_k \times \mathcal{S}_\ell)$.

Let us also consider Λ , the algebra of *symmetric functions* with coefficients in \mathbb{F} . One can for example define it as a polynomial ring in countably many variables: $\Lambda := \mathbb{F}[p_1, p_2, \dots]$ where $\deg p_r = r$.

There is a graded algebra isomorphism Φ between Z and Λ [WW15], which will be reviewed in §2; it is a deformation of the classical Frobenius map in the context of the symmetric group.

In what follows we fix the base field to be

$$\mathbb{F} = \mathbb{Q}(q, t)$$

Date: October 15, 2021.

We thank Allen Knutson, Peter MacNamara, Andrei Negut, Gufang Zhao for discussions.

Define $\mathbb{F}[x_1^\pm, \dots, x_n^\pm]_0$ to be the space of Laurent polynomials with specific degree bounds:
(2)

$$\mathbb{F}[x_1^\pm, \dots, x_n^\pm]_0 = \mathbb{F}\text{-span of } \left\{ x_1^{i_1} \dots x_n^{i_n}, i_1, \dots, i_n \in \mathbb{Z} : \left| \sum_{j=1}^r i_j \right| \leq r(n-r), 0 \leq r \leq n \right\}$$

and $\mathbb{F}[x_1^\pm, \dots, x_n^\pm]_0^{\mathcal{S}^n}$ to be its subspace of symmetric Laurent polynomials.

Consider¹

$$A_n = \{P \in \mathbb{F}[x_1^\pm, \dots, x_n^\pm]_0^{\mathcal{S}^n} : P(x, qx, tx, \dots) = P(tqx, qx, tx, \dots) = 0\}$$

One can again introduce a shuffle product

$$* : A_k \otimes A_\ell \rightarrow A_{k+\ell}$$

$$P \otimes Q \mapsto \sum_{w \in \mathcal{S}^{k,\ell}} \left[P(x_{w(1)}, \dots, x_{w(k)}) Q(x_{w(k+1)}, \dots, x_{w(k+\ell)}) \prod_{\substack{1 \leq i \leq k \\ k+1 \leq j \leq k+\ell}} \tilde{\omega}(x_{w(i)}/x_{w(j)}) \right]$$

where $\tilde{\omega}(x) = \frac{(1-q^{-1}x)(1-qt^{-1}x)(1-tx)}{-x(1-x)}$ making


$$A = \bigoplus_{n \geq 0} A_n$$

a graded algebra. There is another graded algebra isomorphism Υ from A to Λ [FHH⁺09], which will be reviewed in §3.3.

It is natural to ask if there is an interpretation of the isomorphism $\Upsilon^{-1} \circ \Phi$ from Z to A . Note however that Λ possesses graded algebra automorphisms, and in particular we shall allow ourselves to work modulo the automorphisms that rescale the variables p_r .

1.2. Lattice paths. Consider a $n \times n$ square grid, and a set of n paths entering from the left external edges of the grid and exiting at the top. We label incoming paths on the left $1, \dots, n$ from top to bottom. We say that a set of paths P has connectivity $\text{conn}(P) = w \in \mathcal{S}_n$ if the path labelled i (i.e., the i^{th} incoming path on the left counted from the top) exits at the $w^{-1}(i)^{\text{th}}$ location at the top (counted from the left).

Example 1. Here are all the paths with connectivity 2431:



To each such set of lattice paths P we associate a (Boltzmann) weight defined as the product over each vertex of the square grid of a factor that depends on the configuration

¹ A_n is not quite the shuffle algebra as normally defined, but is more convenient for us; the standard shuffle algebra, denoted \mathcal{A}_n , will be given in §3.

around that vertex:

$$(3) \quad \text{wt}(P) = \prod_{i,j=1}^n \left\{ \begin{array}{ll} 1-t & \begin{array}{c} \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \end{array} \\ q^{-1}x_i x_j^{-1}(1-t) & \begin{array}{c} \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \end{array} \\ t(1-q^{-1}x_i x_j^{-1}) & \begin{array}{c} \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \end{array} \\ 1-q^{-1}x_i x_j^{-1} & \begin{array}{c} \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \end{array} \\ 1-tq^{-1}x_i x_j^{-1} & \begin{array}{c} \text{---} \text{---} \\ | \quad | \\ \text{---} \text{---} \end{array} \end{array} \right.$$

where the convention is that colours can be substituted as long as one preserves the ordering; that is, if two paths are present at a vertex, red (resp. green) stands for the one with the lower (resp. greater) label.

To any element $c = \sum_{v \in \mathcal{S}_n} c_v T_v \in Z(\mathcal{H}_n)$ ($c_v \in \mathbb{F}$) of the center of the Hecke algebra we associate the *partition function*

$$(4) \quad f(c) := \alpha_n \sum_{\substack{\text{lattice paths } P \\ \text{on the } n \times n \text{ grid}}} \text{wt}(P) c_{\text{conn}(P)}$$

where $\alpha_n = (q/t)^{\frac{n(n-1)}{2}}(1-t)^{-n}$ is a constant which we introduce for convenience.

We can now formulate our first theorem:

Theorem 1. *There is a commuting square of graded algebra morphisms*


$$\begin{array}{ccc} Z & \xrightarrow{\Phi} & \Lambda \\ \downarrow f & & \downarrow \sigma_{q^{-1}\sigma_t^{-1}} \\ \mathbb{A} & \xrightarrow{\Upsilon} & \Lambda \end{array}$$

where $\sigma_{q^{-1}\sigma_t^{-1}}$ is the automorphism that sends p_r to $(-1)^{r-1} \frac{(1-q)^r(1-t^r)}{(1-q^r)(1-t)^r} p_r$.

1.3. Partition function with connectivity $12 \dots n$. We consider here the special case of the unit $1_n \in Z(\mathcal{H}_n)$ of \mathcal{H}_n which corresponds to the partition function with connectivity $12 \dots n$

$$f(1_n) = \alpha_n \sum_{\substack{\text{lattice paths } P \\ \text{on the } n \times n \text{ grid} \\ \text{conn}(P)=1_n}} \text{wt}(P)$$

Example 2. *Here are the identity paths at $n = 3$:*



Theorem 2. *The partition function $f(1_n)$, as an element of $\mathbb{F}[x_1^\pm, \dots, x_n^\pm]_0^{\mathcal{S}^n}$, is determined by $f(1_0) = 1$ and the recurrence relation*

$$(5) \quad (x_1 + \dots + x_n)f(1_n) = \sum_{i=1}^n x_i f(1_{n-1})[\hat{x}_i] \prod_{\substack{j=1 \\ j \neq i}}^n \frac{(1 - q_1 x_j/x_i)(1 - q_2 x_j/x_i)(1 - q_3 x_i/x_j)}{-x_j/x_i(1 - x_j/x_i)}$$

where $f(1_{n-1})[\hat{x}_i] \in \mathbb{F}[x_1^\pm, \dots, x_{i-1}, x_{i+1}, \dots, x_n^\pm]_0^{\mathcal{S}^n}$.

In §2.4 we will show that the partition functions $f(c)$ can be interpreted using the R -matrices of $\mathcal{U}_{t^{1/2}}(\widehat{\mathfrak{gl}}_\infty)$. This means that one can study them using the quantum inverse scattering method (QISM) [KBI97] (see also [Lam18]). We note, however, that the recurrence relation (5) has an unusual form from the point of view of QISM due to the division by $x_1 + \dots + x_j$ at every j^{th} step of the recursive computation of $f(1_n)$. It is not clear to us if one can prove (5) or find an alternative recursion relation for $f(1_n)$ using QISM. Our proof of (5) is indirect; it follows as a corollary of our proof of Theorem 1 given in §4.

1.4. The commuting scheme. We now give a geometric interpretation of the partition function $f(1_n)$ of the previous section in terms of the commuting scheme. (The geometric interpretation of other $f(c)$ is left for future work.)

Given $n \in \mathbb{Z}_{\geq 0}$, define the *commuting scheme* \mathfrak{C}_n to be the affine scheme in $\mathfrak{gl}_n(\mathbb{C})^2$ consisting of pairs of commuting $n \times n$ matrices:

$$(6) \quad \mathfrak{C}_n := \{(A, B) \in \mathfrak{gl}_n(\mathbb{C}) \times \mathfrak{gl}_n(\mathbb{C}) : [A, B] = 0\}$$

\mathfrak{C}_n is known to be irreducible, of dimension $n(n+1)$ [MT55], smooth in codimension 1 [Pop08], and we have the conjecture

Conjecture 1. *\mathfrak{C}_n is Cohen–Macaulay.*

which would imply that \mathfrak{C}_n is reduced and normal. This conjecture will not be addressed in the present paper.

The group $GL_2(\mathbb{C}) \times GL_n(\mathbb{C})$ acts on $\mathfrak{gl}_n(\mathbb{C})^2$ by

$$(f, g) \in GL_2(\mathbb{C}) \times GL_n(\mathbb{C}) : \begin{pmatrix} A \\ B \end{pmatrix} \mapsto f \begin{pmatrix} gAg^{-1} \\ gBg^{-1} \end{pmatrix}$$

and leaves \mathfrak{C}_n invariant. In particular, the maximal torus $T = (\mathbb{C}^\times)^{2+n}$ induces a \mathbb{Z}^{2+n} -grading of the corresponding coordinate rings; explicitly,

$$\deg A_{ij} = \varepsilon_1 + \varepsilon_{i+2} - \varepsilon_{j+2} \quad \deg B_{ij} = \varepsilon_2 + \varepsilon_{i+2} - \varepsilon_{j+2} \quad i, j = 1, \dots, n$$

where ε_i is the unit vector with nonzero coordinate i .

We consider the Hilbert series $\chi(\mathfrak{C}_n)$ of the commuting scheme with the multigrading above (i.e., the T -character of its coordinate ring); in fact, it is convenient to consider instead the K -polynomial (or Poincaré polynomial) of \mathfrak{C}_n , defined by

$$K_n = \frac{\chi(\mathfrak{C}_n)}{\chi(\mathfrak{gl}_n(\mathbb{C})^2)}$$

Explicitly, introducing formal variables $q_1, q_2, x_1, \dots, x_n$ corresponding to $(\mathbb{C}^\times)^{2+n}$, one has

$$\chi(\mathfrak{gl}_n(\mathbb{C})^2) = \prod_{i,j=1}^n \frac{1}{(1 - q_1 x_i x_j^{-1})(1 - q_2 x_i x_j^{-1})}$$

K_n is known to be a Laurent polynomial in $\mathbb{Z}[q_1^\pm, q_2^\pm, x_1^\pm, \dots, x_n^\pm]$. Furthermore, because of the $GL_2(\mathbb{C}) \times GL_n(\mathbb{C})$ -invariance of \mathfrak{C}_n , K_n is invariant under the action of the corresponding Weyl group, i.e., it is a *symmetric* Laurent polynomial in the variables q_1, q_2 and x_1, \dots, x_n separately.

Finally, one can perform the substitution $q_i \mapsto 1 - q_i, x_i \mapsto 1 - x_i$ in K_n , expand in power series and keep only the lowest degree terms in those variables. We obtain this way a homogeneous polynomial of degree $\text{codim } \mathfrak{C}_n = n(n-1)$ in $\mathbb{Z}[q_1, q_2, x_1, \dots, x_n]$ (with its ordinary grading), called the *multidegree* of \mathfrak{C}_n and denoted

$$D_n := \text{mdeg } \mathfrak{C}_n$$

Our first result on the commuting scheme is the following:

Theorem 3. *Assuming Conjecture 1, the following formula holds for the K -polynomial of the commuting scheme:*

$$(7) \quad K_n = (q_1 q_2)^{\frac{n(n-1)}{2}} f(1_n)$$

with the identification $q = q_1^{-1}, t = (q_1 q_2)^{-1}$.

Performing the substitution of variables above, it is convenient to slightly rearrange the Boltzmann weights of §1.2 to:

$$(8) \quad \text{wt}_K(P) = \prod_{i,j=1}^n \left\{ \begin{array}{ll} 1 - q_1 q_2 & \begin{array}{c} \text{---} \text{---} \text{---} \\ | \quad | \quad | \\ \text{---} \text{---} \end{array} \\ q_1(1 - q_1 q_2) & \begin{array}{c} \text{---} \\ | \\ \text{---} \end{array} \\ 1 - q_1 x_i x_j^{-1} & \begin{array}{c} \text{---} \\ | \\ \text{---} \end{array} \\ q_1 q_2 (1 - q_1 x_i x_j^{-1}) & \begin{array}{c} \text{---} \\ | \\ \text{---} \end{array} \\ q_2(1 - q_1 x_i x_j^{-1}) & \begin{array}{c} \text{---} \\ | \\ \text{---} \end{array} \\ x_i x_j^{-1} (1 - q_2 x_i^{-1} x_j) & \begin{array}{c} \text{---} \\ | \\ \text{---} \end{array} \end{array} \right.$$

Then (7) can be formulated equivalently as

$$K_n = (1 - q_1 q_2)^{-n} \sum_{\substack{\text{lattice paths } P \\ \text{on the } n \times n \text{ grid} \\ \text{with identity connectivity}}} \text{wt}_K(P)$$

Example 3. At $n = 2$, the weights are

$$\begin{aligned} \text{wt}_K \left(\begin{array}{|c|c|c|} \hline \color{red}{\downarrow} & \color{green}{\downarrow} & \color{green}{\downarrow} \\ \hline \color{green}{\leftarrow} & \color{green}{\leftarrow} & \color{green}{\leftarrow} \\ \hline \color{green}{\uparrow} & \color{green}{\uparrow} & \color{green}{\uparrow} \\ \hline \end{array} \right) &= (1 - q_1 q_2)^3 (1 - q_2) \\ \text{wt}_K \left(\begin{array}{|c|c|c|} \hline \color{red}{\downarrow} & \color{green}{\downarrow} & \color{green}{\downarrow} \\ \hline \color{red}{\leftarrow} & \color{green}{\leftarrow} & \color{green}{\leftarrow} \\ \hline \color{green}{\uparrow} & \color{green}{\uparrow} & \color{green}{\uparrow} \\ \hline \end{array} \right) &= q_2 (1 - q_1 q_2)^2 (1 - q_1 x_1 x_2^{-1}) (1 - q_1 x_2 x_1^{-1}) \end{aligned}$$

and one can check that their sum divided by the common factor $(1 - q_1 q_2)^2$ indeed reproduces the K -polynomial of \mathfrak{C}_2

$$K_2 = 1 + q_1^2 q_2 - q_1 q_2 x_1 x_2^{-1} - q_1 q_2 - q_1 q_2 x_1^{-1} x_2 + q_1 q_2^2$$

Define another (Boltzmann) weight in a similar fashion:

$$(9) \quad \text{wt}_H(P) = \prod_{i,j=1}^n \begin{cases} q_1 + q_2 & \color{red}{\downarrow}, \color{red}{\leftarrow}, \color{green}{\leftarrow}, \color{green}{\downarrow} \\ q_1 + x_i - x_j & \color{green}{\downarrow}, \color{green}{\leftarrow}, \color{red}{\leftarrow}, \color{red}{\downarrow} \\ q_2 - x_i + x_j & \color{green}{\downarrow}, \color{green}{\leftarrow}, \color{red}{\leftarrow}, \color{red}{\downarrow} \end{cases}$$

where colours of paths can be substituted freely.

We can state our second result:

Theorem 4. *The following formula holds for the multidegree of the commuting scheme:*

$$(10) \quad D_n := (q_1 + q_2)^{-n} \sum_{\substack{\text{lattice paths } P \\ \text{on the } n \times n \text{ grid} \\ \text{with identity connectivity}}} \text{wt}_H(P)$$

Note that Conjecture 1 and Theorem 3 imply Theorem 4, as can be easily checked. However Theorem 4 can be proven independently of Conjecture 1.

1.5. Plan of the paper. The rest of the paper is organized as follows. In §2, we provide details on the Hecke algebra construction of lattice paths partition functions, and on the map Φ from the center of Hecke to symmetric functions. In §3, we give details on the shuffle algebra, the map Υ and derive the recurrence relations which are required for proving Theorems 1-3. In §4, we prove our Theorem 1 by an explicit comparison of Φ and $\Upsilon \circ f$, and obtain Theorem 2 as a corollary of all that precedes. In §5, we turn to the geometry of the commuting scheme and prove Theorems 3 and 4, ending with some comments.

2. HECKE ALGEBRA AND LATTICE PATHS

2.1. The Hecke algebra and its center. As in §1.1, we consider the Hecke algebra \mathcal{H}_n , a \mathbb{F} -algebra with generators T_1, \dots, T_{n-1} and relations

$$T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1} \quad T_i^2 = (t-1)T_i + t \quad T_i T_j = T_j T_i \quad (|i-j| \geq 2)$$

In this paper we choose \mathbb{F} to be $\mathbb{Q}(q, t)$, though we shall not need the variable q until §2.3.

We have the simple lemma:

Lemma 1.

$$T_i T_w = \begin{cases} T_{s_i w} & s_i w > w \\ (t-1)T_w + tT_{s_i w} & s_i w < w \end{cases}$$

$$T_w T_i = \begin{cases} T_{ws_i} & ws_i > w \\ (t-1)T_w + tT_{ws_i} & ws_i < w \end{cases}$$

expressing the action of T_i on the standard basis $(T_w)_{w \in \mathcal{S}_n}$, where $T_w := T_{i_1} \dots T_{i_k}$ if $w = s_{i_1} \dots s_{i_k}$ is a reduced word in the elementary transpositions s_i .

We shall often need two particular elements of the center $Z(\mathcal{H}_n)$, namely the complete symmetrizer

$$(11) \quad S_n = \sum_{w \in \mathcal{S}_n} T_w$$

and the complete antisymmetrizer

$$(12) \quad A_n = \sum_{w \in \mathcal{S}_n} (-t)^{-|w|} T_w$$

Both are primitive idempotents up to normalization.

Next, we consider the shuffle product structure:

Proposition 1. *The product $*$ in (1) is a well-defined map, making $Z = \bigoplus_{n \geq 0} Z(\mathcal{H}_n)$ a graded (associative) algebra.*

Proof. Associativity of the product is obvious: an m -fold product can be expressed as

$$(13) \quad x_1 * \dots * x_m = \sum_{w \in \mathcal{S}^{\lambda_1, \dots, \lambda_m}} t^{-|w|} T_w (x_1 \otimes \dots \otimes x_m) T_w^{-1}$$

where $x_i \in \mathcal{H}_{\lambda_i}$ and $\mathcal{S}^{\lambda_1, \dots, \lambda_m}$ is the set of shortest representatives in $\mathcal{S}_n / \mathcal{S}_{\lambda_1, \dots, \lambda_m}$, $\mathcal{S}_{\lambda_1, \dots, \lambda_m} = \mathcal{S}_{\lambda_1} \times \dots \times \mathcal{S}_{\lambda_m}$.

The only nontrivial part is to show that $*$ is well-defined as a map from $Z(\mathcal{H}_k) \otimes Z(\mathcal{H}_\ell)$ to $Z(\mathcal{H}_{k+\ell})$, i.e., we need to show that the r.h.s. of (1) is central, or equivalently that it commutes with all T_i s. The proof can be formulated by repeatedly applying Lemma 1; instead, we shall illustrate it by using the standard diagrammatic way to depict the Hecke algebra, as acting on n strands:

$$T_i = \underbrace{\downarrow \quad \dots \quad \downarrow}_{i-1} \quad \begin{array}{c} \nearrow \quad \searrow \\ \searrow \quad \nearrow \end{array} \quad \underbrace{\downarrow \quad \dots \quad \downarrow}_{n-i-1}$$

The arrows keep track of the ordering of operators: moving forward w.r.t. the orientation of a line is reading an expression right to left. Sometimes we will omit arrows; then all lines are implicitly oriented downwards. (The diagrammatic calculus that follows can be viewed as warming up for the considerably more difficult proof of our Theorem 1, which will also be diagrammatic.)

We recall that $\mathcal{S}_{k,\ell}$ is the set of shortest representatives in $\mathcal{S}_{k+\ell} / (\mathcal{S}_k \times \mathcal{S}_\ell)$; equivalently, it is the set of (Grassmannian) permutations of size $k + \ell$ with a unique possible descent at $(k, k + 1)$.

Fixing $i = 1, \dots, n-1$, we distinguish four subsets of $\mathcal{S}_{k,\ell}$ according to the preimages of i and $i+1$:

$$\mathcal{S}_{\leq, \leq} = \{w \in \mathcal{S}^{k,\ell} : w^{-1}(i) \leq k + 1/2, w^{-1}(i+1) \leq k + 1/2\}$$

Note that $w \mapsto s_i w$ induces a bijection between $\mathcal{S}_{<, >}$ and $\mathcal{S}_{>, <}$. We now discuss the corresponding terms in (1) separately:

- If $w \in \mathcal{S}_{<, <}$, then $T_w(a \otimes b)T_{w^{-1}}$ commutes with T_i provided $a \in Z(\mathcal{H}_k)$, as shown diagrammatically:

$$\begin{aligned} T_w(a \otimes b)T_{w^{-1}}T_i &= \begin{array}{c} T_{w^{-1}} \\ \left\{ \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \end{array} \right\} \\ T_w \end{array} = \begin{array}{c} \text{Diagram 3} \\ \text{Diagram 4} \end{array} \\ &= \begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \end{array} = \begin{array}{c} \text{Diagram 7} \\ \text{Diagram 8} \end{array} = T_i T_w(a \otimes b)T_{w^{-1}} \end{aligned}$$

- Similarly, if $w \in \mathcal{S}_{>, >}$, then $T_w(a \otimes b)T_{w^{-1}}$ commutes with T_i provided $b \in Z(\mathcal{H}_\ell)$.
- Finally, we group together the two terms corresponding to $w \in \mathcal{S}_{<, >}$ and $s_i w \in \mathcal{S}_{>, <}$ to obtain:

$$\begin{aligned} (T_w(a \otimes b)T_{w^{-1}} + t^{-1}T_{s_i w}(a \otimes b)T_{w^{-1}s_i})T_i &= \begin{array}{c} T_{w^{-1}} \\ \left\{ \begin{array}{c} \text{Diagram 9} \\ \text{Diagram 10} \end{array} \right\} \\ T_w \end{array} + t^{-1} \begin{array}{c} T_{w^{-1}s_i} \\ \left\{ \begin{array}{c} \text{Diagram 11} \\ \text{Diagram 12} \end{array} \right\} \\ T_{s_i w} \end{array} \\ &= \begin{array}{c} \text{Diagram 13} \\ \text{Diagram 14} \end{array} + \begin{array}{c} \text{Diagram 15} \\ \text{Diagram 16} \end{array} + (1-t^{-1}) \begin{array}{c} \text{Diagram 17} \\ \text{Diagram 18} \end{array} \\ &= T_w(a \otimes b)T_{w^{-1}s_i} + T_{s_i w}(a \otimes b)T_{w^{-1}} + (1-t^{-1})T_{s_i w}(a \otimes b)T_{w^{-1}s_i} \end{aligned}$$

Clearly, the same result is obtained by multiplying by T_i on the left.

By multiplying by $t^{-|w|}$ and summing over w , we obtain the desired commutation property for the r.h.s. of (1). \square

We define an algebra morphism Ψ from $\Lambda = \mathbb{F}[p_1, p_2, \dots]$ to Z as follows. Define Jucys–Murphy elements in \mathcal{H}_n (see e.g. [Ram97]) to be

$$J_j = \sum_{i=1}^{j-1} t^{i-j+1} T_i T_{i+1} \dots T_{j-1} T_{j-2} \dots T_i, \quad j = 2, \dots, n$$

They form a commutative subalgebra, and it is well-known that symmetric polynomials in the J_j are central. Then Ψ sends p_n to the element

$$\Psi(p_n) = [n]_{t^{-1}} \prod_{j=2}^n J_j$$

where $[n]_{t^{-1}} = 1 + \dots + t^{-(n-1)}$.

Proposition 2 ([WW15]). *Ψ is a graded algebra isomorphism from Λ to Z .*

Its inverse is the Φ of §1.1. As a corollary, Z is commutative.

Denote

$$(14) \quad \tilde{h}_n = \Phi(1_n)$$

$$(15) \quad \tilde{e}_n = \Phi(t^{-\frac{n(n-1)}{2}} T_{w_0}^2)$$

$$(16) \quad h_n = \Phi(S_n)$$

$$(17) \quad e_n = \Phi(A_n)$$

where 1_n is the identity of \mathcal{H}_n , and w_0 the longest element of \mathcal{S}_n . We can identify these symmetric functions. Given $u \in \mathbb{F}$, define σ_u to be the automorphism of Λ

$$(18) \quad \sigma_u : p_r \mapsto \frac{(1-u)^r}{1-u^r} p_r$$

Proposition 3 ([Las06]). *h_n (resp. e_n) is the complete (resp. elementary) symmetric function of degree n .*

Furthermore, $\tilde{h}_n = \sigma_t(h_n)$ and $\tilde{e}_n = \sigma_t(e_n)$.

We also denote $\sigma_\infty : p_r \mapsto (-1)^{r-1} p_r$. Note $\sigma_{u^{-1}} = \sigma_\infty \sigma_u$. It is well-known that σ_∞ exchanges h_n and e_n (and similarly for \tilde{h}_n and \tilde{e}_n).

2.2. Partition functions from the Hecke algebra. We now work inside the algebra \mathcal{H}_{2n} . There are two obvious embeddings of \mathcal{H}_n into \mathcal{H}_{2n} , given by $T_i \mapsto T_i$ and $T_i \mapsto T_{i+n}$ respectively, and we denote the corresponding subalgebras $\mathcal{H}_n^{(1)}$ and $\mathcal{H}_n^{(2)}$. Similarly, denote $\mathcal{S}_n^{(1)}$ and $\mathcal{S}_n^{(2)}$ the two subgroups of \mathcal{S}_{2n} (both isomorphic to \mathcal{S}_n) acting nontrivially on $\{1, \dots, n\}$ (resp. $\{n+1, \dots, 2n\}$) only.

Also define the *R-matrix*

$$(19) \quad \check{R}_i(u) = 1 - t + (1-u)T_i \quad i = 1, \dots, 2n-1$$

as an element of $\mathcal{H}_{2n}[u^\pm]$.

Extending the diagrammatic calculus introduced in the proof of Proposition 1, we depict *R*-matrices as (flat) crossings of two lines carrying variables (so-called spectral parameters), the parameter of the *R*-matrix being the ratio of spectral parameters:


$$\check{R}_i(u) = \underbrace{\left| \begin{array}{c} \vdots \\ \vdots \end{array} \right|}_{i-1} \quad \begin{array}{c} u' \nearrow \\ \times \\ \nwarrow u'' \end{array} \quad \underbrace{\left| \begin{array}{c} \vdots \\ \vdots \end{array} \right|}_{2n-i-1} \quad u = u''/u'$$

so that (19) can be expressed as:

$$\begin{array}{c} u' \swarrow \searrow \\ \times \\ \swarrow \searrow \\ u'' \end{array} = - \begin{array}{c} \swarrow \searrow \\ \times \\ \swarrow \searrow \end{array} - u \begin{array}{c} \swarrow \searrow \\ \times \\ \swarrow \searrow \end{array} \quad u = u''/u'$$


The *Yang–Baxter equation* is known to hold:

$$(20) \quad \check{R}_i(u)\check{R}_{i+1}(uv)\check{R}_i(v) = \check{R}_{i+1}(v)\check{R}_i(uv)\check{R}_{i+1}(u) \quad i = 1, \dots, 2n - 2$$




as an identity in $\mathcal{H}_{2n}[u^\pm, v^\pm]$, as well as the *unitarity equation*

$$(21) \quad \check{R}_i(u^{-1})\check{R}_i(u) = (1 - tu)(1 - tu^{-1})$$



The central object of study of this section is the “formal partition function”, which is the following element of $\mathcal{H}_{2n}[x_1^\pm, \dots, x_n^\pm, y_1^\pm, \dots, y_n^\pm]$:

$$(22) \quad Z(x_1, \dots, x_n, y_1, \dots, y_n) = \check{R}_n(x_n/y_1)\check{R}_{n-1}(x_{n-1}/y_1)\check{R}_{n+1}(x_n/y_2) \dots \check{R}_n(x_1/y_n)$$



We also write $Z = Z(x_1, \dots, x_n, y_1, \dots, y_n)$ when there is no need to emphasize the dependence on the variables.

Z has an expansion of the form

$$(23) \quad Z = \sum_{w \in \mathcal{S}_{2n}} Z_w T_w$$


where each $Z_w \in \mathbb{Z}[t^\pm, x_1^\pm, \dots, x_n^\pm, y_1^\pm, \dots, y_n^\pm]$.

2.3. Exchange relation and symmetry. Consider the operator τ_i (resp. τ'_j) that permutes variables x_i and x_{i+1} (resp. y_j and y_{j+1}), $i, j = 1, \dots, n - 1$.


We have the following fundamental lemma:

Lemma 2. Z satisfies the exchange relations:

$$(24) \quad (\tau_i Z) \check{R}_{n+i}(x_{i+1}/x_i) = \check{R}_i(x_{i+1}/x_i) Z \quad i = 1, \dots, n-1$$



$$(25) \quad \check{R}_{n+j}(y_j/y_{j+1}) \tau'_j Z = Z \check{R}_j(y_j/y_{j+1}) \quad j = 1, \dots, n-1$$



Proof. Both equalities result from repeated application of the Yang–Baxter equation (20). \square

We now consider the specialization $y_i = qx_i$, $i = 1, \dots, n$. We first note

Lemma 3. $Z_w|_{y_i=qx_i} \in \mathbb{F}[x_1^\pm, \dots, x_n^\pm]_0$.

Proof. Recall that each of the n^2 factors of Z corresponds to the vertex (i, j) on the diagram of (22), and is a polynomial of degree 1 in the parameter $x_i/(qx_j)$. Now fix $r \in \{0, \dots, n\}$ and consider the degree $i_1 + \dots + i_r$ in the first r variables x_1, \dots, x_r . Among the factors of Z , the ones corresponding on the diagram to the top-left $r \times r$ square have degree zero, and similarly for the bottom $(n-r) \times (n-r)$ square. The ones for the top-right $r \times (n-r)$ square can have monomials of degree 0 or 1, whereas for the bottom-left $(n-r) \times r$ square they can be of degree 0 or -1 . Summing these degrees, we find the desired inequality $|i_1 + \dots + i_r| \leq r(n-r)$. \square

Of course the same proof would work with any r -subset of variables, suggesting that we should study the behaviour of $Z(x_1, \dots, x_n, qx_1, \dots, qx_n)$ under the interchange of the variables x_i . We have the following condition:

Proposition 4. Let χ be a linear form on \mathcal{H}_{2n} . $\chi(Z(x_1, \dots, x_n, qx_1, \dots, qx_n))$ is a symmetric Laurent polynomial in the x_i s provided χ satisfies

$$(26) \quad \chi(ab) = \chi(ba) \quad \forall a \in \mathcal{H}_{2n}, b \in \mathcal{H}_n^{(1)} \otimes \mathcal{H}_n^{(2)}$$

Proof. We assume $n \geq 2$, otherwise the Proposition is trivial. By definition exchanging x_i and x_{i+1} in $\chi(Z(x_1, \dots, x_n, qx_1, \dots, qx_n))$ amounts to considering $\chi(\tau_i \tau'_i Z)$ and then substituting $y_i = qx_i$. One has

$$\chi(\tau_i \tau'_i Z) = \chi(\check{R}_i(x_{i+1}/x_i) \check{R}_{n+i}(y_i/y_{i+1})^{-1} Z \check{R}_i(y_i/y_{i+1}) \check{R}_{n+i}(x_{i+1}/x_i)^{-1})$$

Assuming the property (26) of the Proposition, one can rewrite this

$$\chi(\tau_i \tau'_i Z) = \chi(\check{R}_i(y_i/y_{i+1}) \check{R}_i(x_{i+1}/x_i) Z \check{R}_{n+i}(y_i/y_{i+1})^{-1} \check{R}_{n+i}(x_{i+1}/x_i)^{-1})$$

Substituting $y_i = qx_i$ and simplifying using unitarity equation (21), we obtain the desired symmetry of $\chi(Z(x_1, \dots, x_n, qx_1, \dots, qx_n))$. \square

In what follows, we restrict ourselves to the following class of linear forms. Denote $\langle \bullet \rangle$ the linear form that extracts the coefficient of 1 in a linear combination of T_w , i.e., $\langle T_w \rangle := \delta_{w,1}$. One checks that

$$\langle ab \rangle = \langle ba \rangle = \sum_{w \in \mathcal{S}_{2n}} t^{|w|} a_w b_{w^{-1}}, \quad a = \sum_{w \in \mathcal{S}_{2n}} a_w T_w, \quad b = \sum_{w \in \mathcal{S}_{2n}} b_w T_w$$

Given $c \in Z(\mathcal{H}_n)$, we then define

$$(27) \quad f(c) := \alpha_n \langle (c \otimes S_n) Z(x_1, \dots, x_n, qx_1, \dots, qx_n) \rangle$$

where S_n was defined in (11), and $\alpha_n = (q/t)^{\frac{n(n-1)}{2}} (1-t)^{-n}$ is the same constant as in the introduction. Clearly, $\chi = \langle (c \otimes S_n) \bullet \rangle$ fulfils the condition of Proposition 4.

Example 4. *Three important examples are:*

- *the identity: $f(1_n)$ is the object of §1.3 and §5.*
- *the complete antisymmetrizer: $f(A_n)$ will be computed in §4.*
- *the complete symmetrizer: $f(S_n)$ is also interesting and related to the famous Izergin determinant [Ize87].*

We finally introduce a lattice path representation of $f(c)$, reconnecting it to the content of §1.2.

2.4. Lattice path representation. Consider the following representation of \mathcal{H}_{2n} . As a vector space, $\mathcal{M} = (\mathbb{F}^{n+1})^{\otimes 2n}$. Letting $(e_i)_{i=1, \dots, n+1}$ be the standard basis of \mathbb{F}^{n+1} , define the action of \mathcal{H}_{2n} by

$$T_i(e_{k_1} \otimes \dots \otimes e_{k_{2n}}) = \sum_{k'_i, k'_{i+1}=1}^{n+1} \begin{bmatrix} & k_i & \\ k'_i & & k_{i+1} \\ & k'_{i+1} & \end{bmatrix} e_{k_1} \otimes \dots \otimes e_{k_{i-1}} \otimes e_{k'_i} \otimes e_{k'_{i+1}} \otimes e_{k_{i+2}} \otimes \dots \otimes e_{k_{2n}}$$

where

$$(28) \quad \begin{bmatrix} & i & \\ j & & k \\ & \ell & \end{bmatrix} = \begin{cases} t-1 & i = j > k = \ell \\ 1 & j = k > i = \ell \\ t & j = k \leq i = \ell \\ 0 & \text{else} \end{cases}$$

Given $v \in \mathcal{S}_{2n}$, define

$$|v\rangle = e_{\omega(v(1))} \otimes \dots \otimes e_{\omega(v(2n))}$$

where $\omega(i) = i$ for $1 \leq i \leq n$ and $\omega(i) = n + 1$ for $n + 1 \leq i \leq 2n$. Note that $|v\rangle$ only depends on the class of v inside $\mathcal{S}_n^{(2)} \setminus \mathcal{S}_{2n}$; we define this way a basis indexed by $\mathcal{S}_n^{(2)} \setminus \mathcal{S}_{2n}$ of the submodule of \mathcal{M} of interest to us.

Similarly, define dual basis elements

$$\langle v | = e_{\omega(v(1))}^* \otimes \cdots \otimes e_{\omega(v(2n))}^*$$

in \mathcal{M}^* .

We have the easy lemma

Lemma 4. *For all $w \in \mathcal{S}_{2n}$,*

- (a) $\langle 1 | T_w = t^{|w|} \langle w |$.
 (b) $T_w |1\rangle = t^{|w_2|} |w^{-1}\rangle$ where $w^{-1} = w_1 w_2$, $w_1 \in \mathcal{S}_n^{(2)}$ and w_2 minimal representative in $\mathcal{S}_n^{(2)} \setminus \mathcal{S}_{2n}$.

Proof. (a) Write $w = w_1 w_2$ where $w_1 \in \mathcal{S}_n^{(2)}$ and w_2 minimal representative in $\mathcal{S}_n^{(2)} \setminus \mathcal{S}_{2n}$. One has directly $\langle 1 | T_{w_1} = t^{|w_1|} \langle 1 |$. Then use a reduced decomposition of w_2 and note that in (28), one only ever uses the case $j < \ell$ which leads to $\langle 1 | T_{w_2} = t^{|w_2|} \langle w_2 |$ inductively. At both stages, the only nonzero contributions come from the third line of (28). Case (b) is similar except the first (resp. second) stage involves the second (resp. third) line of (28). \square

Define, noting that $\mathcal{S}_n^{(1)}$ naturally sits inside $\mathcal{S}_n^{(2)} \setminus \mathcal{S}_{2n}$,

$$(29) \quad F_v(x_1, \dots, x_n, y_1, \dots, y_n) = \langle 1 | Z(x_1, \dots, x_n, y_1, \dots, y_n) |v\rangle \quad v \in \mathcal{S}_n^{(1)}$$

This is a slightly more general definition than we need for the interpretation of the r.h.s. of (4); indeed, we have:

Proposition 5.

$$\sum_{\substack{\text{lattice paths } P \\ \text{on the } n \times n \text{ grid} \\ \text{conn}(P)=v}} \text{wt}(P) = F_v(x_1, \dots, x_n, qx_1, \dots, qx_n)$$

Proof. Consider the diagrammatic representation (22) of Z and apply the representation described above; this amounts to assigning to each edge a label in $\{1, \dots, n + 1\}$. F_v corresponds, in accordance to (29), to further imposing that labels of external edges be fixed: $1, \dots, n$ on the left side, $v(1), \dots, v(n)$ on the top side, and $n + 1$ on the right and bottom sides. Now identify labels $1, \dots, n$ with the corresponding lattice paths, and $n + 1$ with “empty”. Then this produces precisely a lattice path with connectivity P . Finally, what needs to be checked is that the weight assigned to a lattice path coincide with the entry of Z . This can be done locally at the level of each individual crossing (R -matrix); and indeed, combining (28) with (19) and setting $y_i = qx_i$ leads to the weights in (3). \square

Finally, we can show that the definition (27) given above is equivalent to the one (4) in the introduction:

Lemma 5. *If $c = \sum_{v \in \mathcal{S}_n} c_v T_v \in Z(\mathcal{H}_n)$, then $c_v = c_{v^{-1}}$ for all $v \in \mathcal{S}_n$ and*

$$f(c) = \alpha_n \sum_{v \in \mathcal{S}_n} c_v F_v(x_1, \dots, x_n, qx_1, \dots, qx_n)$$

Proof. First we observe that

$$(30) \quad \langle x(1 \otimes S_n) \rangle = \langle 1|x|1 \rangle \quad \forall x \in \mathcal{H}_{2n}$$

by testing this on $x = T_w$, $w \in \mathcal{S}_{2n}$, and noting that both sides of the equality are 0 unless $w \in \mathcal{S}_{2n}^{(2)}$, in which case they are equal to $t^{|w|}$ (using the definition (11) for the l.h.s. and Lemma 4 for the r.h.s.). We now start from the definition (27) of $f(c)$ and apply the equality above, as well as Lemma 4 (b):

$$f(c) = \alpha_n \langle 1|Zc|1 \rangle = \alpha_n \sum_{v \in \mathcal{S}_n} c_{v-1} F_v(x_1, \dots, x_n, qx_1, \dots, qx_n)$$

This is almost (4), except a small simplification was made: any central element satisfies $c_v = c_{v-1}$ (short proof: it is obviously true for the generators S_n of Z , and $*$ preserves this property). \square

We conclude this section by reformulating more explicitly the exchange relations of Lemma 2 in terms of the F_v . The next proposition will not be used in what follows, but is included to reconnect to the existing literature [BW18].

Proposition 6. For $v \in \mathcal{S}_n^{(1)}$ and $i, j = 1, \dots, n-1$,

$$(1 - tx_{i+1}/x_i)\tau_i F_v = \begin{cases} (1-t)F_v + (1-x_{i+1}/x_i)F_{s_i v} & s_i v > v \\ (1-t)(x_{i+1}/x_i)F_v + t(1-x_{i+1}/x_i)F_{s_i v} & s_i v < v \end{cases}$$

$$(1 - ty_j/y_{j+1})\tau'_j F_v = \begin{cases} (1-t)F_v + (1-y_j/y_{j+1})F_{vs_j} & vs_j > v \\ (1-t)(y_j/y_{j+1})F_v + t(1-y_j/y_{j+1})F_{vs_j} & vs_j < v \end{cases}$$

Proof. First, note that (23), (29), and Lemma 4 (a) imply

$$(31) \quad F_v = \sum_{w: \dot{w}=v} t^{|w|} Z_w$$

where we have used the notation \dot{w} for the class of $w \in \mathcal{S}_{2n}$ in $\mathcal{S}_n^{(2)} \setminus \mathcal{S}_{2n}$.

We then start from the identity (24) of Lemma 2.

Using (23) and Lemma 1, we obtain for the r.h.s.:

$$\check{R}_i(x_{i+1}/x_i)Z = \sum_{w \in \mathcal{S}_{2n}} Z_w \begin{cases} (1-t)T_w + (1-x_{i+1}/x_i)T_{s_i w} & s_i w > w \\ (1-t)(x_{i+1}/x_i)T_w + t(1-x_{i+1}/x_i)T_{s_i w} & s_i w < w \end{cases}$$

We now take the bra-ket $\langle 1|\cdot|v \rangle$ and note that only w such that either $\dot{w} = v$ or $\dot{w} = s_i v$ contribute. Summing over the coset, we get for $s_i v > v$

$$\begin{aligned} \langle 1|\check{R}_i(x_{i+1}/x_i)Z|v \rangle &= (1-t) \sum_{w: \dot{w}=v} t^{|w|} Z_w + t(1-x_{i+1}/x_i) \sum_{w: \dot{w}=s_i v} t^{|s_i w|} Z_w \\ &= (1-t)F_v + (1-x_{i+1}/x_i)F_{s_i v} \end{aligned}$$

and for $s_i v < v$

$$\begin{aligned} \langle 1|\check{R}_i(x_{i+1}/x_i)Z|v \rangle &= (1-t)(x_{i+1}/x_i) \sum_{w: \dot{w}=v} t^{|w|} Z_w + (1-x_{i+1}/x_i) \sum_{w: \dot{w}=s_i v} t^{|s_i w|} Z_w \\ &= (1-t)(x_{i+1}/x_i)F_v + t(1-x_{i+1}/x_i)F_{s_i v} \end{aligned}$$

3.1. The shuffle algebra \mathcal{A}^+ . The shuffle algebra \mathcal{A}^+ is a vector space whose elements are symmetric rational functions of the form

$$(34) \quad P(x_1, \dots, x_n) = \frac{p(x_1, \dots, x_n)}{\prod_{1 \leq i < j \leq n} (x_i - x_j)^2} \quad p(x_1, \dots, x_n) \in \mathbb{F}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]^{\mathcal{S}_n}$$

where $p(x_1, \dots, x_n)$ satisfies the *wheel condition*

$$(35) \quad p(x_1, \dots, x_n) = 0 \quad \text{if} \quad (x_i, x_j, x_k) = (x, q_1 x, q_1 q_2 x) \quad \text{or} \quad (x_i, x_j, x_k) = (x, q_2 x, q_1 q_2 x)$$

The number of arguments (x_1, \dots, x_n) gives a grading

$$\mathcal{A}^+ = \bigoplus_{n \geq 0} \mathcal{A}_n^+$$

We set $\mathcal{A}_0^+ = \mathbb{F}$ and $\mathcal{A}_1^+ = \mathbb{F}[x^{\pm 1}]$. The shuffle algebra \mathcal{A}^+ is closed under the *shuffle product* $*$, defined for two elements $F \in \mathcal{A}_k^+$ and $G \in \mathcal{A}_l^+$ by

$$(36) \quad (F * G)(x_1, \dots, x_{k+l}) = \frac{1}{k!l!} \text{Sym} \left[F(x_1, \dots, x_k) G(x_{k+1}, \dots, x_{k+l}) \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq l}} \omega(x_{k+j}, x_i) \right]$$

where Sym is the symmetrization over all arguments x_i

$$(37) \quad \text{Sym} [f(x_1, \dots, x_n)] = \sum_{w \in \mathcal{S}_n} f(x_{w(1)}, \dots, x_{w(n)})$$

and $\omega(x, y)$ is defined as follows

$$(38) \quad \omega(x, y) = \frac{(x - q_1 y)(x - q_2 y)(x - q_3 y)}{(x - y)^3}$$

It was shown in [Neg14] that the algebra \mathcal{A}^+ is generated by the elements $x^i \in \mathcal{A}_1^+$ for $i \in \mathbb{Z}$.

3.2. The commutative subalgebra \mathcal{A} . Consider a subalgebra $\mathcal{A} \subset \mathcal{A}^+$ of the elements $P \in \mathcal{A}_n$ for which the two limits

$$(39) \quad \lim_{\xi \rightarrow 0} P(\xi x_1, \dots, \xi x_r, x_{r+1}, \dots, x_n),$$

$$(40) \quad \lim_{\xi \rightarrow \infty} P(\xi x_1, \dots, \xi x_r, x_{r+1}, \dots, x_n)$$

exist and coincide for all fixed $r = 1, \dots, n$. This subalgebra splits into components of fixed degree

$$\mathcal{A} = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} \mathcal{A}_n$$

Remark 1. The shuffle algebras \mathcal{A}_n and \mathbb{A}_n are isomorphic. For $P(x_1, \dots, x_n) \in \mathcal{A}_n$ and $Q(x_1, \dots, x_n) \in \mathbb{A}_n$, s.t. $Q \mapsto P$, we have

$$(41) \quad P(x_1, \dots, x_n) = \frac{1}{V_n(x_1, \dots, x_n)} Q(x_1, \dots, x_n)$$

where $V_n = V_n(x_1, \dots, x_n)$ is defined by

$$(42) \quad V_n := \prod_{1 \leq i \neq j \leq n} (1 - x_i/x_j)$$

Proof. The product $V_n^{-1}Q(x_1, \dots, x_n)$ is of the form (34) and the wheel conditions of \mathcal{A}_n^+ and A_n coincide. The fact that the limiting conditions (39) and (40) of \mathcal{A}_n are equivalent to the degree constraints (2) of A_n was shown in [FHH⁺09]. \square

Another important result of [FHH⁺09] is the following proposition.

Proposition 7. *The algebra $(\mathcal{A}, *)$ is commutative and the dimension of the graded subspace \mathcal{A}_n is equal to the number of partitions of n .*

There are several distinguished bases in the space \mathcal{A} . These bases are built using the shuffle elements

$$(43) \quad \epsilon_n(x_1, \dots, x_n; q_k) = \prod_{1 \leq i < j \leq n} \frac{(x_i - q_k x_j)(x_i - q_k^{-1} x_j)}{(x_i - x_j)^2} \quad k = 1, 2, 3$$

In the following we will abbreviate them as $\epsilon_n(q_k) = \epsilon_n(x_1, \dots, x_n; q_k)$. Due to commutativity the products $\epsilon_i(q_k) * \epsilon_j(q_k)$ can be ordered such that $i > j$. Thus for a partition λ the elements

$$(44) \quad \epsilon_\lambda(q_k) = \epsilon_{\lambda_1}(q_k) * \epsilon_{\lambda_2}(q_k) * \dots * \epsilon_{\lambda_{\ell(\lambda)}}(q_k)$$

give three bases for each $k = 1, 2, 3$.

3.3. Macdonald operators and the isomorphism Υ . Define the Heisenberg algebra \mathfrak{h} with generators $a_{\pm n}$, $n > 0$, satisfying the relations

$$(45) \quad [a_m, a_n] = \delta_{m, -n} m \frac{1 - q^{|m|}}{1 - t^{|m|}}$$

We have two commutative subalgebras generated by $\{a_{\pm n}\}_{n>0}$ and containing the elements $a_{\pm \lambda} = a_{\pm \lambda_1} \dots a_{\pm \lambda_{\ell(\lambda)}}$. Define the Fock space \mathcal{F} by introducing the vacuum vector $|\emptyset\rangle$ which is annihilated by the positive modes $a_n |\emptyset\rangle = 0$, for $n > 0$. The action of the negative modes creates a basis

$$(46) \quad |a_\lambda\rangle = a_{-\lambda} |\emptyset\rangle$$

for all partitions λ .

Let $x = (x_1, x_2, \dots)$ be the alphabet for the ring of symmetric functions Λ with the base field \mathbb{F} . Recall the power sum basis of Λ

$$(47) \quad p_\lambda = p_{\lambda_1} \dots p_{\lambda_{\ell(\lambda)}} \quad p_n = x_1^n + x_2^n + \dots$$

The Fock space \mathcal{F} is isomorphic to Λ which is explicitly given by

$$|a_\lambda\rangle \mapsto p_\lambda$$

In Λ a symmetric function is considered in the power sum basis and the above isomorphism replaces it with a vector in the Fock space. A distinguished basis in Λ is given by the Macdonald symmetric functions P_λ (see [Mac98]). We view it as a basis in the Fock space denoted by $|P_\lambda\rangle$.

The Macdonald operators are defined as the operators which act diagonally on P_λ . The *first* Macdonald operator E [Mac98, Ch. VI] acts as follows

$$(48) \quad E |P_\lambda\rangle = \sum_{i \geq 1} (q^{\lambda_i} - 1) t^{-i} |P_\lambda\rangle$$

This operator is realized on the Fock space with the help of the vertex operator [Shi06]

$$(49) \quad \eta(z) := \exp\left(\sum_{n>0} \frac{1-t^{-n}}{n} a_{-n} z^n\right) \exp\left(-\sum_{n>0} \frac{1-t^n}{n} a_n z^{-n}\right)$$

The Fourier modes of this operator η_n ($n \in \mathbb{Z}$) are given by

$$\eta(z) = \sum_{n \in \mathbb{Z}} \eta_n z^{-n}$$

and have a well defined action on \mathcal{F} . The operator E is given by the zero mode [Shi06]

$$(50) \quad \eta_0 = (t-1)E + 1$$

We can write the eigenvalue equation as follows

$$\frac{t^{-1}}{1-t^{-1}} \eta_0 |P_\lambda\rangle = u_\lambda(e_1) |P_\lambda\rangle$$

where

$$u_\lambda(x_i) = q^{\lambda_i} t^{-i}$$

and e_1 is the first elementary symmetric function in the infinite alphabet $e_1 = x_1 + x_2 + \dots$. One can build higher order operators in a systematic way using the algebra \mathcal{A} and the operators η .

The product of Heisenberg operators is *normally ordered* if all positive modes are on the right and negative modes on the left. The product $\eta(z)\eta(w)$ can be ordered at the expense of a rational function

$$(51) \quad \eta(z)\eta(w) = \frac{(1-w/z)(1-qt^{-1}w/z)}{(1-qw/z)(1-t^{-1}w/z)} : \eta(z)\eta(w) :$$

where the operators inside $::$ are normally ordered. For $f \in \mathcal{A}_n$ define the mapping $\mathcal{O} : \mathcal{A} \rightarrow \text{End}_{\mathbb{F}}(\mathcal{F})$ by

$$(52) \quad \mathcal{O}(f) = \frac{1}{(t-1)^n n!} \oint \frac{dz_1 \cdots dz_n}{z_1 \cdots z_n} \frac{f(z_1, \dots, z_n)}{\epsilon_n(z_1, \dots, z_n; q_1) \epsilon_n(z_1, \dots, z_n; q_3)} : \eta(z_1) \cdots \eta(z_n) :$$

where the integration contours are given by $|z_1| = \dots = |z_n| = 1$ and $|t|, |q| < 1$. It was shown in [FHH⁺09] that \mathcal{O} is compatible with the shuffle product, i.e.

$$(53) \quad \mathcal{O}(f * g) = \mathcal{O}(f)\mathcal{O}(g)$$

Since \mathcal{A} is commutative this implies the commutativity of the operators $[\mathcal{O}(f), \mathcal{O}(g)] = 0$ for any two elements $f, g \in \mathcal{A}$. The map \mathcal{O} was used in [FHH⁺09] to define an isomorphism $\Upsilon' : \mathcal{A} \rightarrow \Lambda$:

Proposition 8. *Given $f \in \mathcal{A}_n$, there exists a $S \in \Lambda_n$ such that for any partition λ ,*

$$(54) \quad \mathcal{O}(f) |P_\lambda\rangle = u_\lambda(S) |P_\lambda\rangle$$

$\Upsilon' : f \mapsto S$ is a ring isomorphism from \mathcal{A} to Λ .

By inserting a V_n^{-1} in (54), cf Remark 1, we obtain the isomorphism Υ from \mathcal{A} to Λ that is stated in the introduction and used in Theorem 1. The symmetric functions of Λ which correspond to the shuffle elements $\epsilon_n(q_k)$, $k = 1, 2, 3$ are all given by a plethystic substitution applied to the elementary symmetric functions e_n . Recalling (18) we set

$$(55) \quad e_n^{(1)} = e_n \quad e_n^{(2)} = \sigma_{q^{-1}} \sigma_{qt^{-1}}^{-1}(e_n) \quad e_n^{(3)} = \sigma_{q^{-1}} \sigma_t^{-1}(e_n)$$

Note that the plethystic substitution in $e_n^{(3)}$ appears in the statement of Theorem 1. This element will play a special role. We have [FHH⁺09]

$$(56) \quad \Upsilon'(\epsilon_n(q_k)) = e_n^{(k)} \quad k = 1, 2, 3$$

or equivalently

$$(57) \quad \mathcal{O}(\epsilon_n(q_k)) |P_\lambda\rangle = u_\lambda(e_n^{(k)}) |P_\lambda\rangle \quad k = 1, 2, 3$$

The eigenvalue equation (57) for $k = 1$ shows that the operator $\mathcal{O}(\epsilon_n(q_1))$ coincides with the n^{th} Macdonald operator [Mac98]. Thus one shows (57) by matching the Macdonald difference operators E_n with $\mathcal{O}(\epsilon_n(q_1))$. Based on this result one can show the other two eigenvalue equations for $k = 2, 3$ using the Wronski relations

$$(58) \quad \sum_{j=0}^n (q_k^j - q_l^{n-j}) \frac{(1 - q_l)^j}{(1 - q_k)^j} \epsilon_{n-j}(q_k) * \epsilon_j(q_l) = 0 \quad k, l = 1, 2, 3$$

In [FHH⁺09] the authors proved these Wronski relations. Applying \mathcal{O} to (58) with $l = 1, k = 2, 3$ gives a relation between the eigenvalues of $\mathcal{O}(\epsilon_n(q_1))$ and $\mathcal{O}(\epsilon_n(q_k))$ which determines the eigenvalue in (57) for $k = 2, 3$.

We finish this section by giving a lemma which will be used at a later stage.

Lemma 6. *The following identities hold*

$$(59) \quad (x_1 + \cdots + x_n) \epsilon_n(q_k) = \frac{1}{1 - q_k} (x_1 * \epsilon_{n-1}(q_k) - q_k \epsilon_{n-1}(q_k) * x_1) \quad k = 1, 2, 3$$

Proof. For simplicity we fix $k = 1$. Let us denote the right hand side of (59) by A_n and rewrite it using the definition of the shuffle product (36)

$$A_n = \frac{1}{1 - q_1} \sum_{i=1}^n x_i \epsilon_{n-1}(q_1)[\hat{x}_i] \left(\prod_{\substack{j=1 \\ j \neq i}}^n \frac{(x_j - q_1 x_i)(x_j - q_2 x_i)(x_j - q_3 x_i)}{(x_j - x_i)^3} - q_1 \prod_{\substack{j=1 \\ j \neq i}}^n \frac{(x_i - q_1 x_j)(x_i - q_2 x_j)(x_i - q_3 x_j)}{(x_i - x_j)^3} \right)$$

where $\epsilon_{n-1}(q_1)[\hat{x}_i]$ depends on the $n - 1$ arguments $(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$. Noting that

$$\begin{aligned} \epsilon_{n-1}(q_1)[\hat{x}_i] \prod_{\substack{j=1 \\ j \neq i}}^n \frac{(x_j - q_1 x_i)}{(x_j - x_i)} &= \epsilon_n(q_1) \prod_{\substack{j=1 \\ j \neq i}}^n \frac{(x_j - x_i)}{(x_j - q_1^{-1} x_i)} \\ \epsilon_{n-1}(q_1)[\hat{x}_i] \prod_{\substack{j=1 \\ j \neq i}}^n \frac{(x_i - q_1 x_j)}{(x_i - x_j)} &= \epsilon_n(q_1) \prod_{\substack{j=1 \\ j \neq i}}^n \frac{(x_i - x_j)}{(x_i - q_1^{-1} x_j)} \end{aligned}$$

leads to

$$A_n = \epsilon_n(q_1) \frac{1}{1 - q_1} \sum_{i=1}^n x_i \left(\prod_{\substack{j=1 \\ j \neq i}}^n \frac{(x_j - q_2 x_i)(x_j - q_3 x_i)}{(x_j - q_1^{-1} x_i)(x_j - x_i)} - q_1 \prod_{\substack{j=1 \\ j \neq i}}^n \frac{(x_i - q_2 x_j)(x_i - q_3 x_j)}{(x_i - q_1^{-1} x_j)(x_i - x_j)} \right)$$

The factor next to $\epsilon_n(q_1)$ does not have poles at $x_i = x_j$ or $x_i = q_1^{-1} x_j$ because of the symmetry in x_i and vanishing residues at $x_i = q_1^{-1} x_j$ which can be easily verified. It means

that this factor must be a symmetric polynomial of degree 1. Thus it is proportional to the elementary symmetric polynomial e_1 in n arguments

$$\frac{1}{1 - q_1} \sum_{i=1}^n x_i \left(\prod_{\substack{j=1 \\ j \neq i}}^n \frac{(x_i - q_2 x_j)(x_i - q_3 x_j)}{(x_i - q_1^{-1} x_j)(x_i - x_j)} - q_1 \prod_{\substack{j=1 \\ j \neq i}}^n \frac{(x_j - q_2 x_i)(x_j - q_3 x_i)}{(x_j - q_1^{-1} x_i)(x_j - x_i)} \right) = C \sum_{i=1}^n x_i$$

We need to show that the proportionality factor $C = 1$. This can be verified by setting $x_i = 0$ for $i > 1$ in the above equation. This completes the proof of (59). \square

3.4. Shuffle elements of \mathcal{A} and specializations. In this section we look at the problem of determining the shuffle algebra elements of \mathcal{A} recursively via specializations of the arguments x_i . This problem was investigated in [Neg14, Neg16] using a coproduct map of \mathcal{A}^+ . Our treatment of this problem is appropriate for the goal of matching shuffle elements with partition functions in §4.

Recall that the shuffle elements of \mathcal{A} have the form

$$(60) \quad P(x_1, \dots, x_n) = \frac{p(x_1, \dots, x_n)}{\prod_{1 \leq i < j \leq n} (x_i - x_j)^2}$$

where $p(x_1, \dots, x_n)$ is a symmetric polynomial. We start with a lemma which gives us information about the degree of $p(x_1, \dots, x_n)$.

Lemma 7. *Let $P(x_1, \dots, x_n)$ be an element of \mathcal{A}_n as in (60). Let x be an indeterminate and $c_1, \dots, c_r \in \mathbb{F}$ then $p(c_1 x, \dots, c_r x, x_{r+1}, \dots, x_n) \in \mathbb{F}[x, x_{r+1}, \dots, x_n]$ is of the form*

$$(61) \quad x^{r(r-1)} G(x, x_{r+1}, \dots, x_n; c_1, \dots, c_r)$$

and $G(x, x_{r+1}, \dots, x_n; c_1, \dots, c_r)$ is of degree at most $2r(n-r)$ in x .

Proof. Consider the specialization $x_k = c_k x$ ($k = 1, \dots, r$) of (60)

$$\begin{aligned} P(c_1 x, \dots, c_r x, x_{r+1}, \dots, x_n) \\ = \frac{p(c_1 x, \dots, c_r x, x_{r+1}, \dots, x_n)}{x^{r(r-1)} \prod_{i=1}^r \prod_{j=r+1}^n (c_i x - x_j)^2 \prod_{1 \leq i < j \leq r} (c_i - c_j)^2 \prod_{r+1 \leq i < j \leq n} (x_i - x_j)^2} \end{aligned}$$

The existence of the limit (39) implies that, for $x \rightarrow 0$, the polynomial in the numerator $p(c_1 x, \dots, c_r x, x_{r+1}, \dots, x_n)$ contains an overall factor $x^{r(r-1)}$ to compensate the same factor in the denominator. Thus we established the form (61) and

$$\begin{aligned} P(c_1 x, \dots, c_r x, x_{r+1}, \dots, x_n) \\ = \frac{G(x, x_{r+1}, \dots, x_n; c_1, \dots, c_r)}{\prod_{i=1}^r \prod_{j=r+1}^n (c_i x - x_j)^2 \prod_{1 \leq i < j \leq r} (c_i - c_j)^2 \prod_{r+1 \leq i < j \leq n} (x_i - x_j)^2} \end{aligned}$$

The existence of the limit (40) implies that, for $x \rightarrow \infty$, the degree in x of the polynomial $G(x, x_{r+1}, \dots, x_n)$ is controlled by $\prod_{i=1}^r \prod_{j=r+1}^n (c_i x - x_j)^2$, i.e. bounded by $2r(n-r)$. \square

In the following we will only be interested in the situations when $c_i = q_k^{i-1}$. The next statement gives a recipe for a recursive computation of shuffle elements of \mathcal{A} .

Proposition 9. *Let $r \in \{1, \dots, n\}$, and $P(x_1, \dots, x_n)$ be an element of \mathcal{A}_n as in (60). Fix $k \in \{1, 2, 3\}$, then $P(x_1, \dots, x_n)$ is determined by the following specializations:*

- (1) $\lim_{x_i \rightarrow 0} \cdots \lim_{x_1 \rightarrow 0} P(x_1, \dots, x_i, x_{i+1}, \dots, x_n)$, $i = 1, \dots, r-1$.
- (2) $P(x, q_k x, \dots, q_k^{r-1} x, x_{r+1}, \dots, x_n)$.

Proof. We fix $k', k'' \in \{1, 2, 3\}$ such that k, k', k'' are all distinct. The proof is by induction on r . The case $r = 1$ is trivial. Now assume the property true at $r \geq 1$ and consider the case $r + 1 \geq 2$. We need to show that we are able to compute

$$P(x, q_k x, \dots, q_k^{r-1} x, x_{r+1}, \dots, x_n)$$

from specializations (1) and (2) where r is replaced by $r + 1$. According to Lemma 7 with $c_i = q_k^{i-1}$ for $1 \leq i \leq r$ we have

$$\begin{aligned} P(x, q_k x, \dots, q_k^{r-1} x, x_{r+1}, \dots, x_n) \\ = \frac{G(x, x_{r+1}, \dots, x_n)}{\prod_{i=1}^r \prod_{j=r+1}^n (q_k^{i-1} x - x_j)^2 \prod_{1 \leq i < j \leq r} (q_k^{i-1} - q_k^{j-1})^2 \prod_{r+1 \leq i < j \leq n} (x_i - x_j)^2} \end{aligned}$$

where G is a polynomial of degree $2r(n - r)$ in x . According to the wheel conditions (35), it has the following factorization:

$$G(x, x_{r+1}, \dots, x_n) = \prod_{i=0}^{r-2} \prod_{j=r+1}^n (q_{k'} x_j - q_k^i x) (q_{k''} x_j - q_k^i x) H(x, x_r, \dots, x_n)$$

with the degree of H in x being $2r(n - r) - 2(r - 1)(n - r) = 2(n - r)$. To determine H , we therefore need to know its values at $2(n - r) + 1$ distinct specializations of x . The value of H at $x = 0$ gives one specialization and can be computed from the knowledge of

$$\lim_{x_r \rightarrow 0} \cdots \lim_{x_1 \rightarrow 0} P(x_1, \dots, x_r, x_{r+1}, \dots, x_n)$$

The specialization (2), using the fact that P is a symmetric polynomial, is equivalent to computing H at $x = q_k^{-r} x_j$ and $x = q_k x_j$, $j = r + 1, \dots, n$. This is the required number of specializations, which means G is determined and with it $P(x, q_k x, \dots, q_k^{r-1} x, x_{r+1}, \dots, x_n)$. \square

Let us apply this result to the computation of the basis elements $\epsilon_\lambda(q_k)$ with $\ell(\lambda) = m$ and $\lambda_1 + \cdots + \lambda_m = n$. For convenience we only consider the case $k = 3$ and denote

$$\epsilon_\lambda = \epsilon_\lambda(q_3) = \epsilon_\lambda(t)$$

We apply Proposition 9 with $r = n$ and $k = 3$ to the problem of computing ϵ_λ . This means that in order to determine ϵ_λ we need to know the values of

$$(62) \quad \lim_{x_i \rightarrow 0} \cdots \lim_{x_1 \rightarrow 0} \epsilon_\lambda(x_1, \dots, x_i, x_{i+1}, \dots, x_n), \quad i = 1, \dots, n - 1$$

$$(63) \quad \epsilon_\lambda(x, tx, \dots, t^{n-1}x)$$

One can compute the limits in (62) using the following lemma.

Lemma 8. *The elements $\epsilon_{\lambda_1, \dots, \lambda_m}$, with $\lambda_1 + \cdots + \lambda_m = n$, satisfy the recurrence relation*

$$(64) \quad \lim_{x_n \rightarrow 0} \epsilon_{\lambda_1, \dots, \lambda_m}(x_1, \dots, x_{n-1}, x_n) = \sum_{i=1}^m \epsilon_{\lambda_1, \dots, \lambda_i-1, \dots, \lambda_m}(x_1, \dots, x_{n-1})$$

Proof. We start by writing $\epsilon_{\lambda_1, \dots, \lambda_m}$ explicitly as a shuffle product of ϵ_{λ_i}

$$\begin{aligned} \epsilon_{\lambda_1, \dots, \lambda_m} &= \sum_{w \in \mathcal{S}_n / \mathcal{S}_{\lambda_1, \dots, \lambda_m}} \epsilon_{\lambda_1}(x_{w(1)}, \dots, x_{w(\lambda_1)}) \cdots \epsilon_{\lambda_m}(x_{w(\lambda_1 + \cdots + \lambda_{m-1} + 1)}, \dots, x_{w(n)}) \\ (65) \quad &\prod_{1 \leq a < b \leq m} \prod_{i \in I_a, j \in I_b} \omega(x_{w(j)}, x_{w(i)}) \end{aligned}$$

where we defined the subsets I_a by

$$(66) \quad I_a := \left\{ \sum_{s=1}^{a-1} \lambda_s + 1, \dots, \sum_{s=1}^a \lambda_s \right\}$$

From definitions (43) and (38) we have two obvious identities

$$\epsilon_k(x_1, \dots, x_{k-1}, 0) = \epsilon_{k-1}(x_1, \dots, x_{k-1}) \quad \omega(x, 0) = \omega(0, x) = 1$$

Fix $r \in \{1, \dots, m\}$ and consider the summands of (65) with w such that $n \in w(I_r)$. The factor ϵ_{λ_r} will be replaced with $\epsilon_{\lambda_{r-1}}$ upon setting $x_n = 0$ while the ω factors containing x_n will disappear. The summation over $w \in \mathcal{S}_n / \mathcal{S}_{\lambda_1, \dots, \lambda_m}$ (with $n \in w(I_r)$) reduces to the summation over $w' \in \mathcal{S}_{n-1} / \mathcal{S}_{\lambda_1, \dots, \lambda_{r-1}, \dots, \lambda_m}$ which is equal to the shuffle product of $\epsilon_{\lambda_1}, \dots, \epsilon_{\lambda_{r-1}}, \dots, \epsilon_{\lambda_m}$ similarly to (65). Treating in this way every summand with $n \in w(I_r)$ for $r \in \{1, \dots, m\}$ gives rise to the formula (64). \square

It is clear that computing (62) can be done inductively using (64). The specializations in (63) are not difficult to compute and in fact follow from more general specializations considered in [FHH⁺09]. We have

$$(67) \quad \epsilon_\lambda(x, tx, \dots, t^{n-1}x) = 0, \quad \lambda \neq (1, 1, \dots, 1)$$

$$(68) \quad \epsilon_{(1,1,\dots,1)}(x, tx, \dots, t^{n-1}x) = \prod_{1 \leq i < j \leq n} \frac{(qt^i - t^j)(t^{i+1} - qt^j)(t^i - t^{j+1})}{qt(t^i - t^j)^3}$$

Let us explain (68). We use the explicit expression (65) with $\lambda_i = 1$ in which case all $\epsilon_{\lambda_i} = 1$ and all sets I_a have a single element. Due to the vanishing of $\omega(tx, x) = 0$ only one term in the summation is non-vanishing. This term must have x 's with fully ordered indices as follows

$$\prod_{1 \leq i < j \leq n} \omega(x_i, x_j)|_{x_k = t^{k-1}x}$$

which is equal to the product on the right hand side of (68).

3.5. The shuffle element κ_n . The results of this section are gathered in the following proposition:

Proposition 10. *Let $\kappa_0 = 1 \in \mathcal{A}_0$. Each of the following formulas uniquely determines the same sequence of shuffle algebra elements $\kappa_n \in \mathcal{A}_n$:*

$$(69) \quad \Upsilon'(\kappa_n) = \sigma_q(e_n)$$

$$(70) \quad \kappa_n = \sum_{j=0}^n q_k^{-n} (q_k - 1)^{n-j} \kappa_j * \epsilon_{n-j}(q_k) \quad k = 1, 2, 3$$

$$(71) \quad (x_1 + \dots + x_n) \kappa_n = \kappa_{n-1} * x_1$$

Proof. It is clear that each of these equations uniquely determines the element κ_n , we only need to show their consistency. We can take (71) together with $\kappa_0 = 1$ as the definition of κ_n and prove that it implies (70) and (69).

The derivation of (70) is recursive. The case $n = 0$ of (70) obviously holds. Using (71) we will derive (70) for κ_n assuming that (70) holds for $n - 1$. We act with $*x_1$ on both sides of

(70) where n is replaced by $n - 1$

$$(72) \quad \kappa_{n-1} * x_1 = \sum_{j=0}^{n-1} q_k^{-n+1} (q_k - 1)^{n-1-j} \kappa_j * \epsilon_{n-j-1}(q_k) * x_1$$

On the left hand side we apply (71) and on the right hand side (59)

$$\begin{aligned} \sum_{i=1}^n x_i \kappa_n &= \sum_{j=0}^{n-1} q_k^{-n} (q_k - 1)^{n-j} \kappa_j * \left(\sum_{i=1}^{n-j} x_i \epsilon_{n-j}(q_k) \right) \\ &\quad + \sum_{j=0}^{n-1} q_k^{-n} (q_k - 1)^{n-j-1} \kappa_j * x_1 * \epsilon_{n-j-1}(q_k) \end{aligned}$$

On the right hand side we apply (71) to $\kappa_j * x_1$

$$\begin{aligned} \sum_{i=1}^n x_i \kappa_n &= \sum_{j=0}^{n-1} q_k^{-n} (q_k - 1)^{n-j} \kappa_j * \left(\sum_{i=1}^{n-j} x_i \epsilon_{n-j}(q_k) \right) \\ &\quad + \sum_{j=0}^{n-1} q_k^{-n} (q_k - 1)^{n-j-1} \left(\sum_{i=1}^{j+1} x_i \kappa_{j+1} \right) * \epsilon_{n-j-1}(q_k) \end{aligned}$$

In the last summation we can shift the index j

$$(73) \quad \begin{aligned} \sum_{i=1}^n x_i \kappa_n &= \sum_{j=0}^{n-1} q_k^{-n} (q_k - 1)^{n-j} \kappa_j * \left(\sum_{i=1}^{n-j} x_i \epsilon_{n-j}(q_k) \right) \\ &\quad + \sum_{j=1}^n q_k^{-n} (q_k - 1)^{n-j} \left(\sum_{i=1}^j x_i \kappa_j \right) * \epsilon_{n-j}(q_k) \end{aligned}$$

From the definition of the shuffle product (36) it is easy to check that

$$\kappa_j * \left(\sum_{i=1}^{n-j} x_i \epsilon_{n-j}(q_k) \right) + \left(\sum_{i=1}^j x_i \kappa_j \right) * \epsilon_{n-j}(q_k) = \left(\sum_{i=1}^n x_i \right) \kappa_j * \epsilon_{n-j}(q_k)$$

Using this identity in (73) for every pair of summands with $j = 1, \dots, n - 1$ leads to

$$\begin{aligned} \sum_{i=1}^n x_i \kappa_n &= \sum_{i=1}^n x_i q_k^{-n} (q_k - 1)^n \kappa_0 * \epsilon_n(q_k) + \sum_{i=1}^n x_i q_k^{-n} \kappa_n * \epsilon_0(q_k) \\ &\quad + \sum_{i=1}^n x_i \sum_{j=1}^{n-1} q_k^{-n} (q_k - 1)^{n-j} \kappa_j * \epsilon_{n-j}(q_k) \end{aligned}$$

After removing the common factor $\sum_{i=1}^n x_i$ and putting all terms together on the right hand side we obtain (70) for κ_n .

Next we show that (69) holds. For that we will use (70). It is sufficient to consider $k = 1$ in (70). We apply the map Υ' to this equation

$$(74) \quad \Upsilon'(\kappa_n) = \sum_{j=0}^n q^n (q^{-1} - 1)^{n-j} \Upsilon'(\kappa_j) \Upsilon'(\epsilon_{n-j}(q^{-1}))$$

From (56) we have $\Upsilon'(\epsilon_{n-j}(q^{-1})) = e_{n-j}$. Let us denote $e'_n = \Upsilon'(\kappa_n)$, then (74) becomes

$$(75) \quad e'_n = \sum_{j=0}^n q^j (1-q)^{n-j} e'_j e_{n-j}$$

We need to show that $e'_n = \sigma_q(e_n)$. Note that, together with the initial condition, (75) uniquely determines e'_n . Let us solve (75) using generating functions, set

$$(76) \quad E'(z) := \sum_{n=0}^{\infty} e'_n z^n \quad E(z) := \sum_{n=0}^{\infty} e_n z^n = \exp\left(-\sum_{j>0} \frac{(-1)^j}{j} p_j z^j\right)$$

Multiply both sides of (75) by z^n and sum over $n \geq 0$

$$E'(z) = E'(qz)E((1-q)z)$$

By iterating this equation we find

$$\begin{aligned} E'(z) &= \prod_{k=0}^{\infty} E((1-q)q^k z) = \exp\left(-\sum_{k \geq 0} \sum_{j > 0} \frac{(-1)^j (1-q)^j}{j} p_j q^{jk} z^j\right) \\ &= \exp\left(-\sum_{j > 0} \frac{(-1)^j (1-q)^j}{j(1-q^j)} p_j z^j\right) = \exp\left(-\sum_{j > 0} \frac{(-1)^j}{j} \sigma_q(p_j) z^j\right) = \sum_{n=0}^{\infty} \sigma_q(e_n) z^n \end{aligned}$$

Comparing the last result with (76) shows that $e'_n = \sigma_q(e_n)$. \square

4. MATCHING THE PARTITION FUNCTIONS WITH SHUFFLE ELEMENTS

We now investigate the partition functions associated to the elements $A_{\lambda_1} * \cdots * A_{\lambda_m}$ (where, by commutativity, λ can be chosen to be a partition) that were studied in the previous section on the shuffle algebra side.

Proposition 11. *Given $\lambda_1, \dots, \lambda_m \in \mathbb{Z}_{>0}$ with $n = \sum_{i=1}^m \lambda_i$, one has the limit*

$$(77) \quad \lim_{x_n \rightarrow 0} x_n^{n-1} f(A_{\lambda_1} * \cdots * A_{\lambda_m}) = (-1)^{n-1} \prod_{i=1}^{n-1} x_i \sum_{i=1}^m f(A_{\lambda_1} * \cdots * A_{\lambda_{i-1}} * A_{\lambda_m})$$

Proof. We start from the definition of $A_{\lambda_1} * \cdots * A_{\lambda_m}$ (cf (13)):

$$A_{\lambda_1} * \cdots * A_{\lambda_m} = \sum_{v \in \mathcal{S}_{\lambda_1, \dots, \lambda_m}} \sum_{w \in \mathcal{S}^{\lambda_1, \dots, \lambda_m}} (-1)^{|v|} t^{-(|v|+|w|)} T_w T_v T_{w^{-1}}$$

By definition of the groups involved, one has $T_w T_v = T_{wv}$ (and similarly $T_v T_{w^{-1}} = T_{vw^{-1}}$, but we choose to use the first identity).

Plugging this into (27), we have

$$\begin{aligned} \alpha_n^{-1} f(A_{\lambda_1} * \cdots * A_{\lambda_m}) &= \sum_{v \in \mathcal{S}_{\lambda_1, \dots, \lambda_m}} \sum_{w \in \mathcal{S}^{\lambda_1, \dots, \lambda_m}} (-1)^{|v|} t^{-(|v|+|w|)} \langle Z|_{y_i=qx_i} (T_w T_v T_{w^{-1}} \otimes S_n) \rangle \\ &= \sum_{v \in \mathcal{S}_{\lambda_1, \dots, \lambda_m}} \sum_{w \in \mathcal{S}^{\lambda_1, \dots, \lambda_m}} (-1)^{|v|} t^{-(|v|+|w|)} \langle T_{w^{-1}} Z|_{y_i=qx_i} T_{wv} (1 \otimes S_n) \rangle \end{aligned}$$

Using (30), we can rewrite this as

$$\begin{aligned}
\alpha_n^{-1} f(A_{\lambda_1} * \cdots * A_{\lambda_m}) &= \sum_{v \in \mathcal{S}_{\lambda_1, \dots, \lambda_m}} \sum_{w \in \mathcal{S}^{\lambda_1, \dots, \lambda_m}} (-1)^{|v|} t^{-(|v|+|w|)} \langle 1 | T_{w^{-1}} Z|_{y_i=qx_i} T_{wv} | 1 \rangle \\
(78) \qquad \qquad \qquad &= \sum_{v \in \mathcal{S}_{\lambda_1, \dots, \lambda_m}} \sum_{w \in \mathcal{S}^{\lambda_1, \dots, \lambda_m}} (-t)^{-|v|} \langle w^{-1} | Z|_{y_i=qx_i} | v^{-1} w^{-1} \rangle
\end{aligned}$$

This has the following interpretation: $\alpha_n^{-1} f(A_{\lambda_1} * \cdots * A_{\lambda_m})$ is the sum over v and w of partition functions of lattice paths, where the paths are labelled $w^{-1}(1), \dots, w^{-1}(n)$ on the left side and $v^{-1}(w^{-1}(1)), \dots, v^{-1}(w^{-1}(n))$ on the top side (with an extra weight $(-t)^{-|v|}$).

We can now take the limit $x_n \rightarrow 0$. In this paragraph we denote $a = w^{-1}$, $b = v^{-1}w^{-1}$, which are two arbitrary permutations, and consider $\langle a | Z|_{y_i=qx_i} | b \rangle$ as $x_n \rightarrow 0$. We are only interested for the purposes of deriving (77) in the terms of order x_n^{-n+1} ; from Lemma 3 we know that this is the minimum order in x_n and in fact, as in the proof of the lemma, we know that the only dependence on x_n comes from the last row and column of the diagram of Z . Let us examine these in more detail with our boundary conditions $\langle a |$ and $| b \rangle$. We look at the destiny of the path $b(n)$ which ends at the top right: to maximize the degree in x_n^{-1} , we must pick up a x_n^{-1} at each vertex (i, n) , $i = 1, \dots, n-1$; this means, according to (3) that the first type of vertex (a path making a bend when the right edge is either empty or has a higher labelled path) must be excluded. Since all right edges are empty in the last column, one can prove inductively that the paths must go straight through every vertex (i, n) , $i = 1, \dots, n-1$, incurring a weight of $-q^{-1}x_i x_n^{-1}$. At vertex (n, n) , because both bottom and right edges are empty, we must have the first type of vertex in (3), incurring a $1-t$. From there, the path can only go straight left to its destination with label $a(n)$, incurring a weight t^{n-1} and imposing $a(n) = b(n)$. Combining all this, we find

$$\begin{aligned}
(79) \quad \langle a | Z(x_1, \dots, x_n, qx_1, \dots, qx_n) | b \rangle &= \delta_{a(n), b(n)} (1-t) t^{n-1} x_n^{-n+1} \prod_{i=1}^{n-1} (-q^{-1}x_i) \\
&\langle a|_{1, \dots, n-1} | Z(x_1, \dots, x_{n-1}, qx_1, \dots, qx_{n-1}) | b|_{1, \dots, n-1} \rangle + O(x_n^{-n+2}) \quad a, b \in \mathcal{S}_n^{(1)}
\end{aligned}$$

We apply this formula to $f(A_{\lambda_1} * \cdots * A_{\lambda_m})$. We fix $w \in \mathcal{S}_{\lambda_1, \dots, \lambda_m}$ and denote $k = w^{-1}(n)$. Because of the form of w , k must be the last element of its block I_a , $1 \leq a \leq m$ (I_a was defined in (66)). According to (79), the summation can be restricted to v such that $v(k) = k$. Consider now the permutation w' with $w'(i) = w(i)$, $i = 1, \dots, k-1$, $w'(i) = w'(i+1)$, $i = k, \dots, n-1$, and the permutation v' with $v'(i) = i$ for $i = 1, \dots, k-1$ and $v'(i) = v(i+1) - 1$ for $i = k, \dots, n-1$. Because k is last of its block, $|v'| = |v|$ and we have

$$\begin{aligned}
\alpha_n^{-1} f(A_{\lambda_1} * \cdots * A_{\lambda_m}) &\stackrel{x_n \rightarrow 0}{\sim} (1-t) t^{n-1} x_n^{-n+1} \prod_{i=1}^{n-1} (-q^{-1}x_i) \sum_{v' \in \mathcal{S}_{\lambda_1, \dots, \lambda_a-1, \dots, \lambda_m}} (-t)^{-|v'|} \sum_{w' \in \mathcal{S}^{\lambda_1, \dots, \lambda_a-1, \dots, \lambda_m}} \\
&\langle w'^{-1}(1), \dots, w'^{-1}(n-1) | Z(x_1, \dots, x_{n-1}, qx_1, \dots, qx_{n-1}) | v'^{-1}(w'^{-1}(1)), \dots, v'^{-1}(w'^{-1}(n-1)) \rangle
\end{aligned}$$


Comparing with (78) and using $\alpha_n^{-1} \alpha_{n-1} = (1-t)(t/q)^{n-1}$, we finally find the desired limit (77). \square

In what follows, we shall use the diagrammatic language. We shall repeatedly use the following simple identities:


We now go back to $A_{\lambda_1} * \dots * A_{\lambda_m}$. Starting from the expression

$$f(A_{\lambda_1} * \dots * A_{\lambda_m}) = \alpha_n \sum_{w \in \mathcal{S}^{\lambda_1, \dots, \lambda_m}} t^{-|w|} \langle (T_w(A_{\lambda_1} \otimes \dots \otimes A_{\lambda_m})T_{w^{-1}} \otimes S_n)Z \rangle |_{y_i=qx_i, i=1, \dots, n}$$

and using the cyclicity of $\langle \bullet \rangle$, we can describe the expression inside it as the diagram: (compare with (83))



As a side remark, using the idempotency of A_n and S_n , we can introduce the more symmetric diagram



with $\tilde{X}_w = \langle X_w \rangle A_{\lambda_1} \otimes \dots \otimes A_{\lambda_m} \otimes S_n$. We shall not use \tilde{X}_w in what follows.

Proposition 12. • For $m < n$, one has the vanishing condition:

(85)
$$f(A_{\lambda_1} * \dots * A_{\lambda_m})(x, \dots, t^m x, x_{m+2}, \dots, x_n) = 0$$

• The partition function $f(A_n)$ has the fully factorized form

(86)
$$f(A_n) = t^{-\frac{n(n-1)}{2}} \prod_{\substack{i,j=1 \\ i \neq j}}^n (1 - tx_i/x_j)$$

- One has the specialization

$$(87) \quad f(A_1 * \cdots * A_1)(1, t, \dots, t^{n-1}) = [n]_{t^{-1}}! \prod_{1 \leq i < j \leq n} (q - t^{j-i})(q - t^{i-j+1})$$

Proof. We first consider $m = 1$, and investigate the effect of setting $x_1 = tx$, $x_2 = x$ in X_w , where here $w = 1$. Noting that $\check{R}_1(t) = (1 - t)(1 + T_1) = (1 - t)S_2$, and using (81), we can insert a crossing at the right of the top two horizontal lines and then apply the Yang–Baxter equation repeatedly to move the crossing to the left, resulting in zero according to (82):

$$\begin{aligned}
 X_1|_{x_1=tx, x_2=x} &= \frac{1}{1+t} \\
 &= \frac{1}{1+t} \\
 &= 0
 \end{aligned}$$

We conclude that $x_1 - tx_2$ divides $f(A_n)$, and by symmetry, $f(A_n)$ is a multiple of $\prod_{i \neq j} (1 - tx_i/x_j)$. The latter has degree range $[-(n-1), (n-1)]$ in each x_i , which is the maximum allowed according to (2), so only remains a constant in \mathbb{F} to be determined. The latter is determined inductively by using Proposition 11.

We proceed identically for $1 < m < n$ (on the example picture, $m = 3$). We set $x_1 = t^m x$, \dots , $x_m = tx$, $x_{m+1} = x$. We note that the R -matrix associated to the longest element of S_{m+1} and parameters $t^m, \dots, 1$ is nothing but $(1 - t)^{\frac{m(m+1)}{2}} \prod_{i=1}^m [i]_t! S_{m+1}$ (see e.g. [Las06,


Lem. 3] or [IO09, 3.1]), and move it across to the left using the Yang–Baxter equation:

$$X_w|_{x_1=t^m x, \dots, x_{m+1}=x} = \frac{1}{[m+1]_t!}$$


The pigeonhole principle tells us there will be two numbers between 1 and $m + 1$ whose preimages under w lie in the same block. Since $w \in \mathcal{S}^{\lambda_1, \dots, \lambda_m}$, all numbers whose preimages are in the same block are consecutive, and so we can apply (82) to conclude that the result is zero. By summing over w and using symmetry in the exchange of variables, we obtain the result (85) as stated in the Proposition.

The only case left is $m = n$, i.e., $A_1 * \dots * A_1$. We set $x_1 = t^{n-1}x, \dots, x_{n-1} = tx, x_n = x$ but this time the result does not vanish. However, noting once again that S_n is up to normalization the product of R -matrices associated to the longest permutation w_0 of \mathcal{S}_n , we can move it to the left and absorb any crossing from T_w :


$$X_w|_{x_1=t^{n-1}x, \dots, x_n=x} =$$



where in the last line we find it convenient to reintroduce the longest element w_0 , before moving back S_n to the right. Performing the summation over w (one could instead absorb the top crossings as well, but the result would be the same), we find



We then impose $y_i = qx_i$ and note that up to normalization, the S_n at the top is nothing but the R -matrix associated to the longest element of $\mathcal{S}_n^{(2)}$; we can pull it to the bottom:



Finally, the bracket is

$$f(A_1 * \cdots * A_1) = \alpha_n t^{-\frac{n(n-1)}{2}} [n]_t! \left\langle \begin{array}{c} \text{Diagram} \end{array} \right\rangle$$

This is essentially the same situation that was considered at the end of §2. First we expand each R -matrix in the upper-left triangle as $\tilde{R}_i(u) = u(1-t) + t(1-u)T_i^{-1}$ and note that the only term that has a nonzero identity coefficient is

$$f(A_1 * \cdots * A_1) = \alpha_n [n]_t! (1-t)^n \prod_{\substack{1 \leq i, j \leq n \\ i+j < n+1}} (1 - q^{-1} t^{n-i-j-1}) \left\langle \begin{array}{c} \text{Diagram} \end{array} \right\rangle$$

Secondly, we absorb all the crossings of the lower-right triangle into S_n , resulting in a weight of $\prod_{\substack{1 \leq i, j \leq n \\ i+j > n+1}} (1 - q^{-1} t^{n-i-j+2})$. Recombining and simplifying, we obtain (87). \square

We are now in a position to prove Theorem 1. First, combining Lemma 3, Proposition 4 and Lemma 10, we find that $f(c)$, from its definition (27), is α_n times an element of A_n . α_n contains a factor of $(1-t)^{-n}$, but it is compensated by the fact that any lattice path entering from the left and exiting from the top must have a bend, resulting in a factor of $(1-t)^n$ according to (3) in the expression (4) of $f(c)$. Therefore, f is a map from $Z(\mathcal{H}_n)$ to A_n .

Next, we identify $f(A_{\lambda_1} * \cdots * A_{\lambda_m})$. We use the characterization of Proposition 9, so that $\epsilon_{\lambda_1, \dots, \lambda_m}$ is entirely determined by Lemma 8 and (68). On the other hand, it is easy to check that Proposition 12 implies the exact same relations for $V_n^{-1} f(A_{\lambda_1} * \cdots * A_{\lambda_m})$ (where V_n is defined in (42), and implements the isomorphism $V_n \times$ from \mathcal{A}_n to A_n). Therefore,

$$f(A_{\lambda_1} * \cdots * A_{\lambda_m}) = V_n \epsilon_{\lambda_1, \dots, \lambda_m}$$

We finally compare $\Upsilon(f(A_{\lambda_1} * \cdots * A_{\lambda_m}))$ and $\Phi(A_{\lambda_1} * \cdots * A_{\lambda_m})$.

According to the ring map property of Υ and (56) applied to $\epsilon_r = \epsilon_r^{(3)}$,

$$\begin{aligned} \Upsilon(f(A_{\lambda_1} * \cdots * A_{\lambda_m})) &= \Upsilon(V_n \epsilon_{\lambda_1}^{(3)} * \cdots * \epsilon_{\lambda_m}^{(3)}) \\ &= \prod_{i=1}^m \Upsilon(V_{\lambda_i} \epsilon_{\lambda_i}^{(3)}) \\ &= \prod_{i=1}^m \sigma_{q^{-1}} \sigma_t^{-1}(e_{\lambda_i}) \end{aligned}$$

On the other hand, from (17) and the ring map property of Φ ,

$$\Phi(A_{\lambda_1} * \cdots * A_{\lambda_m}) = \prod_{i=1}^m e_{\lambda_i}$$

We have therefore checked the statement of Theorem 1 on a basis of \mathbb{Z} .

Corollary 1. *We have the following identity*

$$(88) \quad f(1_n) = V_n \kappa_n$$

Proof. According to (69) and Proposition 8, $\Upsilon(V_n \kappa_n) = \Upsilon'(\kappa_n) = \sigma_q(e_n)$. On the other hand, according to Proposition 3, $\Phi(1_n) = \tilde{h}_n = \sigma_t(h_n) = \sigma_t \sigma_\infty(e_n)$. Applying Theorem 1, we conclude that (88) holds. \square

Using the definition of the shuffle product (36) we can write

$$(89) \quad (x_1 + \cdots + x_n) \kappa_n(x_1, \dots, x_n) = \sum_{i=1}^n x_i \kappa_{n-1}(x_1, \dots, \hat{x}_i, \dots, x_n) \prod_{\substack{j=1 \\ j \neq i}}^n \frac{(x_i - q_1 x_j)(x_i - q_2 x_j)(x_i - q_3 x_j)}{(x_i - x_j)^3}$$

The recurrence relation for $f(1_n)$ stated in (5) follows from (88), (89) and (42) thus proving Theorem 2.

5. APPLICATION TO THE COMMUTING SCHEME

5.1. Proof of Theorem 3. In this section, all schemes are over \mathbb{C} , as in the introduction.

Proposition 13. *Assuming Conjecture 1, K_n satisfies the following recurrence relation:*

$$(90) \quad (x_1 + \cdots + x_n) K_n = \sum_{i=1}^n x_i K_{n-1}[\hat{x}_i] \prod_{\substack{j=1 \\ j \neq i}}^n \frac{(1 - q_1 x_j/x_i)(1 - q_2 x_j/x_i)(1 - q_1 q_2 x_i/x_j)}{1 - x_j/x_i}$$

where $K_{n-1}[\hat{x}_i] = K_{n-1}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$.

Proof. For the purposes of this proof, we need more geometry than in the rest of the paper. We think of the multi-graded Hilbert series as the pushforward to a point in localized equivariant K -theory. Equivalently, the K -polynomial of \mathfrak{C}_n is viewed as (the pushforward of) the K_T -class $[\mathcal{O}_{\mathfrak{C}_n}]$ in the ambient space $\mathfrak{gl}_n \times \mathfrak{gl}_n$. Note that $K_T(\mathfrak{gl}_n \times \mathfrak{gl}_n) \cong K_T(pt) \cong \mathbb{Z}[q_1^\pm, q_2^\pm, x_1^\pm, \dots, x_n^\pm]$ where the variables $q_1, q_2, x_1, \dots, x_n$ are the *inverses* of the equivariant parameters attached to the torus T .

Consider

$$\mathfrak{F}_n = \{(v, A, B) \in \mathbb{P}^{n-1} \times \mathfrak{gl}_n \times \mathfrak{gl}_n : [A, B] = 0 \text{ and } v \text{ eigenvector of } A \text{ and } B\}$$

There is a natural projective morphism

$$\begin{aligned} \mathfrak{F}_n &\xrightarrow{p} \mathfrak{C}_n \\ (v, A, B) &\mapsto (A, B) \end{aligned}$$

which is generically n -to-1 (choice of one eigenvector among n). There is also the affine morphism $f : \mathfrak{F}_n \rightarrow \mathbb{P}^{n-1}$. Since all fibers are isomorphic by GL_n -action, the dimension of a fiber is $\dim \mathfrak{F}_n - \dim \mathbb{P}^{n-1} = n^2 + 1$.

The computation is in two parts:

- Computing $\pi_*[\mathcal{O}_{\mathfrak{F}_n}(1)]$ by equivariant localization (which relates it to $\pi_*[\mathcal{O}_{\mathfrak{C}_{n-1}}]$ using $(v, A, B) \mapsto (A|_{\mathbb{C}^n/v}, B|_{\mathbb{C}^n/v})$).
- Comparing $p_*[\mathcal{O}_{\mathfrak{F}_n}(1)]$ and $[\mathcal{O}_{\mathfrak{C}_n}]$.

5.1.1. *Equivariant localization.* Torus fixed points in \mathbb{P}^{n-1} are basis vectors e_i , $i = 1, \dots, n$. Consider such a fixed point, which up to \mathcal{S}_n action, we can choose to be $v = e_n$. The fiber is

$$f^{-1}(v) = \{v\} \times X, \quad X = \left\{ \left(A = \begin{pmatrix} A' & 0 \\ \alpha & a \end{pmatrix}, B = \begin{pmatrix} B' & 0 \\ \beta & b \end{pmatrix} \right) : AB = BA \right\}$$

More explicitly, the equations are

$$(91) \quad A'B' = B'A'$$

$$(92) \quad a\beta + \alpha B' = b\alpha + \beta A'$$

In particular note that $(A', B') \in \mathfrak{C}_{n-1}$. There are $2n$ extra variables α, β, a, b and $n-1$ extra equations, and $\dim \mathfrak{C}_{n-1} + 2n - (n-1) = n^2 + 1 = \dim f^{-1}(v)$, so assuming Conjecture 1 (Cohen–Macaulay property for \mathfrak{C}_{n-1} , which implies it for $\mathfrak{C}_{n-1} \times \mathbb{A}^{2n+2}$), the unmixedness theorem asserts that the extra equations (92) form a regular sequence.

We then conclude by standard exact sequence arguments that

$$(93) \quad [\mathcal{O}_X] = [\mathcal{O}_{\mathfrak{C}_{n-1}}] \prod_{j=1}^{n-1} (1 - q_1 x_j / x_n) (1 - q_2 x_j / x_n) (1 - q_1 q_2 x_n / x_j)$$

where $[\mathcal{O}_X]$ is the K_T class of \mathcal{O}_X in $\mathfrak{gl}_n \times \mathfrak{gl}_n$, whereas $[\mathcal{O}_{\mathfrak{C}_{n-1}}]$ is the K_T class of \mathfrak{C}_{n-1} in $\mathfrak{gl}_{n-1} \times \mathfrak{gl}_{n-1}$ (where the T -action on $\mathfrak{gl}_{n-1} \times \mathfrak{gl}_{n-1}$ has a one-dimensional kernel leading to the obvious inclusion $K_{T/\mathbb{C}^\times} \subset K_T$, i.e., as a polynomial $[\mathcal{O}_{\mathfrak{C}_{n-1}}] = K_{n-1}$ is independent of x_n). The factors $1 - q_1 x_j / x_n$ (resp. $1 - q_2 x_j / x_n$) come from the zero column of A (resp. B) whereas the factors $1 - q_1 q_2 x_n / x_j$ come from the equations (92).

We have thus managed to compute $[f^{-1}(v)] = [v][\mathcal{O}_X]$.

The weights at the tangent space of e_i are x_i/x_j , $j \neq i$ (recalling that the x_i are the inverses of the equivariant parameters). Pulling back the relation

$$1 = \sum_{i=1}^n \frac{[e_i]}{\prod_{\substack{j=1 \\ j \neq i}}^n (1 - x_j/x_i)}$$

in localized equivariant K -theory of \mathbb{P}^{n-1} to \mathfrak{F}_n and multiplying by $[\mathcal{O}_{\mathfrak{F}_n}(1)]$, we obtain

$$[\mathcal{O}_{\mathfrak{F}_n}(1)] = \sum_{i=1}^n x_i [e_i] [\mathcal{O}_{\mathfrak{C}_{n-1}}](\hat{x}_i) \prod_{\substack{j=1 \\ j \neq i}}^n \frac{(1 - q_2 x_j/x_i)(1 - q_1 x_j/x_i)(1 - q_1 q_2 x_i/x_j)}{1 - x_j/x_i}$$

Finally, pushing forward to $\mathfrak{gl}_n \times \mathfrak{gl}_n$ leads to:

$$(94) \quad p_*[\mathcal{O}_{\mathfrak{F}_n}(1)] = \sum_{i=1}^n x_i [\mathcal{O}_{\mathfrak{C}_{n-1}}](\hat{x}_i) \prod_{\substack{j=1 \\ j \neq i}}^n \frac{(1 - q_2 x_j/x_i)(1 - q_1 x_j/x_i)(1 - q_1 q_2 x_i/x_j)}{1 - x_j/x_i}$$

5.1.2. *Pushforward.* Now let us try to compute directly the pushforward of $[\mathcal{O}_{\mathfrak{F}_n}(1)]$ under $p : \mathfrak{F}_n \rightarrow \mathfrak{C}_n$. Write $\mathfrak{C}_n = \text{Spec } R$ and $\mathfrak{F}_n = \text{Proj } S$ where $S_0 = R$. Since \mathfrak{C}_n is affine, the pushforward is simply given in terms of the sheaf cohomology groups of $\mathcal{O}_{\mathfrak{F}_n}(1)$ (viewed as modules over R). We claim the following

$$H^i(\mathfrak{F}_n, \mathcal{O}_{\mathfrak{F}_n}(1)) \cong \begin{cases} S_1 & i = 0 \\ 0 & i > 0 \end{cases}$$

Proof. Let us first write the equations of \mathfrak{F}_n explicitly:

$$(95) \quad AB = BA \quad \text{minors of size 2 of } (v, Av) \text{ and } (v, Bv) = 0$$

Now define the patches $U_i = \{v_i \neq 0\}$. Let us consider $U_1 \cap U_2$ inside U_1, U_2 . Denote $z = v_1/v_2$, $r_i = v_i/v_2$, $i > 2$ and collectively $r = (r_i)$. A section on U_1 is given by a polynomial $P(z, r, A, B)$ modulo the relations (95) rewritten in terms of z, r, A, B . Let us call these equations $q_i(z)$ where dependence on r, A, B is suppressed. Note that the $q_i(z)$ are of degree less or equal to 2 in z . Similarly a section on U_2 is given by a polynomial $Q(z^{-1}, rz^{-1}, A, B)$ modulo the same equations expressed in terms of z^{-1}, rz^{-1}, A, B . The $\mathcal{O}(1)$ matching condition on $U_1 \cap U_2$ is

$$(96) \quad P(z) - zQ(z^{-1}) = \sum_i c_i(z)q_i(z)$$

where the $c_i(z)$ are Laurent polynomials in z . Now assume that P is of degree greater than 1 in z . Then its top coefficient is equal to the top coefficient of $Q := \sum_i c_i^{pol}(z)q_i(z)$ where $c_i^{pol}(z)$ is the part of $c_i(z)$ with nonnegative powers of z . By subtracting Q from P , we can reduce the degree of P until it is at most 1.

This way, one can show that the restriction of a global section to U_i is of degree in v_i/v_j at most 1 for all $i \neq j$; this immediately implies that the v_i generate the global sections of $\mathcal{O}(1)$. Linear independence is obvious (it can be tested at the generic point).

(96) also implies that the map $(P, Q) \mapsto P(z) - zQ(z^{-1})$ is surjective, so that $H^1(\mathcal{O}_{\mathfrak{F}_n}(1)) = 0$. Vanishing of higher cohomology follows similarly. \square

Since $H^0(\mathcal{O}_{\mathfrak{F}_n}(1)) \cong S_1$ is a free R -module with basis (v_1, \dots, v_n) and higher sheaf cohomology vanishes, we conclude that

$$(97) \quad p_*[\mathcal{O}_{\mathfrak{F}_n}(1)] = [\mathcal{O}_{\mathfrak{C}_n}] \sum_{j=1}^n x_j$$

where x_j is the weight of the basis element v_j .

Combining (94) and (97) leads to the recurrence relation (90). \square

We are now in a position to prove Theorem 3 by relating the shuffle element κ_n and the K -polynomial K_n . K_n is uniquely determined by (90) and the initial condition $K_0 = 1$.

Similarly κ_n can be computed recursively with (89). By comparing (89) with (90) (and checking the trivial initial condition $\kappa_0 = K_0 = 1$) we conclude that

$$(98) \quad \kappa_n(x_1, \dots, x_n) = \frac{t^{n(n-1)/2}}{V_n(x)} K_n(x_1, \dots, x_n)$$

Combining (88) with (98) gives us the statement of Theorem 3.

5.2. Proof of Theorem 4. Theorem 4 involves computing the multidegree of \mathfrak{C}_n , which means one needs to work in equivariant cohomology. The situation is much simpler than in equivariant K -theory (which was needed for Theorem 3), and we only sketch the corresponding geometric argument. By abuse of notation, we use the same symbols $q_1, q_2, x_1, \dots, x_n$ for the cohomology equivariant parameters. This is consistent with our passage from K -polynomial to multidegree, which was defined in the introduction as the substitution $q_i \mapsto 1 - q_i$ and $x_i \mapsto 1 - x_i$ and then expanding at leading order. We refer to this limit in what follows as the GRR limit.

View the flag variety \mathcal{B}_n as the variety of Borel subalgebras of \mathfrak{gl}_n and consider $\tilde{\mathfrak{F}}_n = \{(\mathfrak{b}, A, B) \in \mathcal{B}_n \times \mathfrak{gl}_n \times \mathfrak{gl}_n : [A, B] = 0, A, B \in \mathfrak{b}\}$. There is a projection map p from $\tilde{\mathfrak{F}}_n$ to \mathfrak{C}_n which is generically $n!$ -to-1 (generic matrices are diagonalizable); one then obtains by equivariant localization techniques (similar to 5.1.1) the formula

$$(99) \quad D_n = \frac{1}{n!} \prod_{\substack{i,j=1 \\ i \neq j}}^n (q_1 + x_i - x_j)(q_2 + x_i - x_j) \sum_{w \in \mathcal{S}_n} w \left(\prod_{1 \leq i < j \leq n} \frac{q_1 + q_2 + x_i - x_j}{(x_j - x_i)(q_1 + x_i - x_j)(q_2 + x_i - x_j)} \right)$$

$$= \text{Sym} \left(\prod_{1 \leq i < j \leq n} \frac{(q_1 + q_2 + x_i - x_j)(q_1 - x_i + x_j)(q_2 - x_i + x_j)}{x_j - x_i} \right)$$

where the symmetric group acts by permuting variables, and Sym denotes symmetrization w.r.t. to it.

One can think of this expression as the GRR limit of the shuffle element $\frac{1}{n!} f(A_1 * \dots * A_1)$ (studied in §4).²

At $t = 1$, from (1), one has the identity in the center of the symmetric group algebras $A_1 * \dots * A_1 = n! 1_n$, so that in the GRR limit, $\frac{1}{n!} f(A_1 * \dots * A_1)$ and $f(1_n)$ coincide. As was already remarked in §1.4, expanding the weights (8), which correspond to $f(1_n)$, in the GRR limit, result in the weights (9). This leads to the formula of Theorem 4.

A few comments are in order. In [DFZJ06, KZJ07] another expression for the multidegree of the commuting variety was obtained, using apparently unrelated methods. There is actually a connection: consider Proposition 6 and take the GRR limit. One recovers one of the exchange relations of the work above (cf [DFZJ06, Eq. (3.10)] or [KZJ07, Prop. 6]). The expression for the multidegree of \mathfrak{C}_n there is essentially a repeated application of these identities; it is not obvious if it is more or less useful than our (new) formula (99). (The extension of these ideas to K -theory will be investigated elsewhere [KZJ].)

The expression of Theorem 4, in contrast, has many desirable properties; for instance, it is division- and subtraction-free. Regarding positivity, let us only consider the bidegree

²It should be noted that the element $f(A_1 * \dots * A_1)$ itself has a geometric interpretation [Gin11] as the Hilbert series of the coordinate ring of the normalization of the isospectral commuting variety $\mathfrak{X}_{\text{norm}}$. What we are finding here is that to go from K -theory to cohomology, instead of using the structure sheaf of \mathfrak{C}_n one can use the combinatorially simpler structure sheaf of $\mathfrak{X}_{\text{norm}}$.

REFERENCES

- [BW18] A. Borodin and M. Wheeler, *Coloured stochastic vertex models and their spectral theory*, 2018, [arXiv:1808.01866](#).
- [DFZJ06] P. Di Francesco and P. Zinn-Justin, *Inhomogeneous model of crossing loops and multidegrees of some algebraic varieties*, *Comm. Math. Phys.* **262** (2006), no. 2, 459–487, [arXiv:math-ph/0412031](#). MR2200268.
- [FHH⁺09] B. Feigin, K. Hashizume, A. Hoshino, J. Shiraishi, and S. Yanagida, *A commutative algebra on degenerate $\mathbb{C}\mathbb{P}^1$ and Macdonald polynomials*, *J. Math. Phys.* **50** (2009), no. 9, 095215, [arXiv:0904.2291](#).
- [FO97] B. Feigin and A. Odesskii, *A family of elliptic algebras*, *Int. Math. Res. Not.* **1997** (1997), no. 11, 531–539.
- [FT11] B. Feigin and A. Tsymbaliuk, *Equivariant K-theory of Hilbert schemes via shuffle algebra*, *Kyoto J. Mat.* **51** (2011), no. 4, 831–854, [arXiv:0904.1679](#).
- [Gin11] V. Ginzburg, *Isospectral commuting variety, the Harish-Chandra D-module, and principal nilpotent pairs*, 2011, [arXiv:1108.5367](#).
- [IO09] A. P. Isaev and O. V. Ogievetsky, *Braids, shuffles and symmetrizers*, *Journal of Physics A: Mathematical and Theoretical* **42** (2009), no. 30, 304017, [arXiv:0812.3974](#), [doi:10.1088/1751-8113/42/30/304017](#).
- [Ize87] A. Izergin, *Partition function of a six-vertex model in a finite volume*, *Dokl. Akad. Nauk SSSR* **297** (1987), no. 2, 331–333. MR919260.
- [KBI97] V. Korepin, N. Bogoliubov, and A. Izergin, *Quantum inverse scattering method and correlation functions*, vol. 3, Cambridge University Press, 1997.
- [KZJ] A. Knutson and P. Zinn-Justin, work in progress.
- [KZJ07] ———, *A scheme related to the Brauer loop model*, *Adv. Math.* **214** (2007), no. 1, 40–77, [arXiv:math.AG/0503224](#). MR2348022.
- [Lam18] J. Lamers, *The functional method for the domain-wall partition function*, *SIGMA Symmetry Integrability Geom. Methods Appl.* **14** (2018), paper 64, 23, [arXiv:1801.09635](#), [doi:10.3842/SIGMA.2018.064](#).
- [Las06] A. Lascoux, *The Hecke algebra and structure constants of the ring of symmetric polynomials*, 2006, [arXiv:math/0602379](#).
- [Mac98] I. Macdonald, *Symmetric functions and Hall polynomials*, Oxford University Press, 1998.
- [MT55] T. S. Motzkin and O. Taussky, *Pairs of matrices with property L. II*, *Transactions of the American Mathematical Society* **80** (1955), no. 2, 387–401, [doi:10.2307/1992996](#).
- [Neg14] A. Neguț, *The shuffle algebra revisited*, *Int. Math. Res. Not.* **2014** (2014), no. 22, 6242–6275, [arXiv:1209.3349](#).
- [Neg16] ———, *The m/n Pieri rule*, *Int. Math. Res. Not.* **2016** (2016), no. 1, 219–257, [arXiv:1407.5303](#).
- [Pop08] V. L. Popov, *Irregular and singular loci of commuting varieties*, *Transformation Groups* **13** (2008), 819–837, [arXiv:0801.3074](#), [doi:10.1007/s00031-008-9018-9](#).
- [Ram97] A. Ram, *Seminormal representations of Weyl groups and Iwahori–Hecke algebras*, *Proceedings of the London Mathematical Society* **75** (1997), no. 1, 99–133, [arXiv:math/9511223](#).
- [Shi06] J. Shiraishi, *A family of integral transformations and basic hypergeometric series*, *Comm. Math. Phys.* **263** (2006), no. 2, 439–460, [arXiv:math/0501251](#).
- [SV13] O. Schiffmann and E. Vasserot, *The elliptic Hall algebra and the K-theory of the Hilbert scheme of \mathbb{A}^2* , *Duke Math. J.* **162** (2013), no. 2, 279–366, [arXiv:0905.2555](#).
- [WW15] J. Wan and W. Wang, *Frobenius map for the centers of Hecke algebras*, *Trans. Amer. Math. Soc.* **367** (2015), no. 8, 5507–5520, [arXiv:1208.4446](#).

A. GARBALI, P. ZINN-JUSTIN, SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, VICTORIA 3010, AUSTRALIA.

Email address: alexandr.garbali@unimelb.edu.au

Email address: pzinn@unimelb.edu.au