

ON HOMOGENEOUS AND OSCILLATING RANDOM WALKS ON THE INTEGERS

Julien Brémont

Université Paris-Est Créteil, octobre 2021

Abstract

Considering homogeneous and oscillating random walks on the integers, we simplify classical works on recurrence of Spitzer and Kemperman, respectively. Some links with renewal theory are discussed.

1 Introduction

A primary question in the study of the asymptotic behaviour of a discrete time Markov chain on \mathbb{Z}^d is that of recurrence. In the homogeneous case, i.e. when the jumping law is the same everywhere, this problem concerns Birkhoff sums $S_n = X_1 + \dots + X_n$, where the (X_i) are \mathbb{Z}^d -valued random variables, independent and identically distributed (*i.i.d.*), with common law μ . Improving a former result of Chung and Fuchs, Spitzer's analytical recurrence criterion (1957, cf [11], T2) then states that transience is equivalent to the integrability of $Re(1/(1 - \hat{\mu}))$ on the unit cube of \mathbb{R}^d . Importantly, this holds in full generality, without any moment condition for example. For a general model of inhomogeneous random walk on \mathbb{Z}^d , the transition law at a point $x \in \mathbb{Z}^d$ is given by some probability measure μ_x . The question of the recurrence is more complex and cannot be reduced to that of *i.i.d.* sums. One has to understand how the Markov chain perceives the space given by the couple $(\mathbb{Z}^d, (\mu_x)_{x \in \mathbb{Z}^d})$.

A situation, of intermediate difficulty, where the *i.i.d.* case helps is the following one. Fix a Markov chain on \mathbb{Z}^d , starting at 0. Given a subset F with $\{0\} \subset F \subset \mathbb{Z}^d$, then the random walk is recurrent at 0 if and only if, first, it visits F infinitely often and, next, the random walk restricted to times of visits (itself a Markov chain) is recurrent at 0. This cuts the problem in two parts. If ever able to solve the first one, then remains to study a Markov chain living in the smaller set F . Inevitably however, if F is too small, the latter has heavy-tailed jumps. This naive strategy nevertheless works for instance for a nearest-neighbour model in \mathbb{Z}^2 , when the transition laws only depend on the second coordinate. This is an example of random walk in a stratified environment, as considered in [3, 1, 2]. Taking $F = \mathbb{Z} \times \{0\} \subset \mathbb{Z}^2$, in this particular situation the vertical component of the random walk (in restriction to vertical movements) is a nearest-neighbour one-dimensional Markov chain. The condition for its recurrence is well-known, for example in the theory of birth and death processes. When assuming recurrence to $\mathbb{Z} \times \{0\}$ of the initial random walk, the induced random walk in $\mathbb{Z} \times \{0\}$ is heavy-tailed. Nonetheless, the invariance of the environment by horizontal translations makes it *i.i.d.*. The analysis is then

AMS 2020 subject classifications : 60G17, 60J10.

Key words and phrases : Homogeneous random walk, oscillating random walk, recurrence criterion.

naturally orientated into precisising the jumping law of this random walk and next using Spitzer's analytical recurrence criterion for \mathbb{Z} -valued *i.i.d.* sums (cf [1, 2]).

In order to expand this line of research, one has to develop results around the cornerstone that constitutes Spitzer's theorem. A preliminary step of clarification is necessary, as the known proof of Spitzer's theorem, available in [11], is not linear and substantially requires the apparatus of Potential Theory for discrete Markov chains. It is also disseminated in [11] and a little painful to reconstitute (Kesten-Spitzer [8] is helpful). The present work provides a short proof of Spitzer's result and next discusses some links with renewal theory. In the same way we also consider oscillating random walks on the integers and reprove known results of Kemperman [7] on the recurrence of this model, using a combinatorial remark simplifying the analysis.

The material and the results in this text are essentially not new. Our effort has been concentrated on the exposition, which tries to be in straight line and self-contained. Many unanswered questions are addressed along the way, suggesting further work. The essential ones concern inner products of probabilistic Green functions and their translations in Harmonic analysis.

2 Homogeneous random walk on the integers

We fix \mathbb{Z} as state space. Let us first recall classical facts and notations.

2.1 Preliminaries

- *Laws and characteristic functions.* Let X be a \mathbb{Z} -valued random variable, with law denoted by $\mathcal{L}(X) = \mu$, and $(X_n)_{n \geq 1}$ be *i.i.d.* copies. We suppose X non constant, with $\gcd(\text{Supp}(\mu)) = 1$. Introduce the characteristic function $\hat{\mu}(t) = \mathbb{E}(e^{itX})$, $t \in \mathbb{R}$, which is 2π -periodic. Defining $d = \gcd(\text{Supp}(\mathcal{L}(X_1 - X_2))) \geq 1$, then :

$$\hat{\mu}(t) = 1 \text{ iff } t \in 2\pi\mathbb{Z} \text{ and } |\hat{\mu}(t)| = 1 \text{ iff } t \in (2\pi/d)\mathbb{Z}.$$

We write $S_n = \sum_{i=1}^n X_i$, with $S_0 = 0$. Since $|\hat{\mu}(t)| < 1$ except for finitely t , $\mathcal{L}(S_n)$ does not concentrate at any point, as $n \rightarrow +\infty$:

$$\forall y \in \mathbb{Z}, P(S_n = y) = \frac{1}{2\pi} \int_0^{2\pi} e^{-ity} (\hat{\mu}(t))^n dt \rightarrow_{n \rightarrow +\infty} 0.$$

Also, for some $\alpha > 0$ and small t , $\text{Re}(1 - \hat{\mu}(t)) \geq \alpha t^2$. Indeed, take $M > 0$ so that $P(0 < |X| < M) > 0$. Then for $|t| \leq \pi/M$, $\text{Re}(1 - \hat{\mu}(t)) = 2\mathbb{E}(\sin^2(tX/2)) \geq 2\pi^{-2}t^2\mathbb{E}(X^2 1_{|X| < M})$.

- *Markov chains and Green functions.* For any Markov chain (S_n) on \mathbb{Z} , P_x and \mathbb{E}_x stand for x as starting point. The Green function is $G(x, y) = \mathbb{E}_x(\sum_{n \geq 0} 1_{S_n=y})$. Let also $G_N(x, y) = \mathbb{E}_x(\sum_{0 \leq n < N} 1_{S_n=y})$, $N \geq 1$. For $y \in \mathbb{Z}$, introduce $T_y = \min\{n \geq 1, S_n = y\}$. For any $x \neq y$:

$$G_N(x, y) = \sum_{1 \leq k < N} P_x(T_y = k) G_{N-k}(y, y). \quad (1)$$

Hence, $G_N(x, y) \leq G_N(y, y)$ and this gives $G(x, y) = P_x(T_y < \infty)G(y, y)$. Also, for $x \in \mathbb{Z}$:

$$G(x, x) = \sum_{n \geq 0} P_x(T_x < \infty)^n = 1/(1 - P_x(T_x < \infty)),$$

so the recurrence of x , i.e. the property $P_x(T_x < \infty) = 1$, is equivalent to $G(x, x) = +\infty$. For any $x \neq y$ with $P_x(T_y < \infty) > 0$, note that $P_x(T_x < T_y) < 1$. Then, in the same way :

$$\mathbb{E}_x \left(\sum_{n=0}^{T_y-1} 1_{S_n=x} \right) = 1 + \sum_{n \geq 1} P_x(T_x < T_y)^n = \frac{1}{1 - P_x(T_x < T_y)} < \infty.$$

Still for any $x \neq y$, we have :

$$\begin{aligned} G_N(x, x) &= \mathbb{E}_x \left(\sum_{n=0}^{T_y \wedge N-1} 1_{S_n=x} \right) + \mathbb{E}_x \left(1_{T_y < N} \sum_{n=T_y}^{N-1} 1_{S_n=x} \right) \\ &= \mathbb{E}_x \left(\sum_{n=0}^{T_y \wedge N-1} 1_{S_n=x} \right) + \sum_{k=1}^{N-1} P_x(T_y = k) G_{N-k}(y, x). \end{aligned} \quad (2)$$

Thus, $0 \leq G_N(x, x) - G_N(y, x) \leq \mathbb{E}_x(\sum_{n=x}^{T_y \wedge N-1} 1_{S_n=x})$. Hence we deduce the important remark : *Claim* : for $x \neq y$ with $P_x(T_y < \infty) > 0$, then $(G_N(x, x) - G_N(y, x))_{N \geq 0}$ is bounded.

In the particular situation when the Markov chain is homogeneous, $G(x, y) = G(x - y, 0)$ and $G_N(x, y) = G_N(x - y, 0)$. For $x \neq 0$, notice also that $P_0(T_0 < T_x) = P_0(T_0 < T_{-x})$, as :

$$\begin{aligned} P_0(T_0 < T_x) &= \sum_{k \geq 1} P_0(S_k = 0, S_l \notin \{0, x\}, 0 < l < k) \\ &= \sum_{k \geq 1} P_0(S_k = 0, S_k - S_l \notin \{0, -x\}, 0 < l < k) = P_0(T_0 < T_{-x}). \end{aligned}$$

In the homogeneous case, with a step of law μ , we often put μ as a superscript and write S_n^μ , $G^\mu(x, y)$, $G_N^\mu(x, y)$. When $\mathcal{L}(X) = \mu$, we also write $\mathbb{E}^\mu(f(X))$ for $\int_{\mathbb{Z}} f d\mu$.

2.2 Spitzer's analytical recurrence criterion

We consider an homogeneous random walk (S_n) on \mathbb{Z} with step μ , not Dirac, supposing $\gcd(\text{Supp}(\mu)) = 1$. On $(0, 2\pi)$, $t \mapsto \text{Re}(1/(1 - \hat{\mu}(t)))$ is > 0 , continuous and symmetric under $t \mapsto 2\pi - t$. Hence, integrability on $(0, 2\pi)$ holds iff it holds on $(0, \varepsilon)$, for some $\varepsilon > 0$.

Theorem 2.1. (*Spitzer, 1957*)

The point 0 is transient for (S_n) iff $\int_0^{2\pi} \text{Re}(1/(1 - \hat{\mu}(t))) dt < +\infty$.

This follows from the following proposition, where constants are optimal.

Proposition 2.2. *We have $G(0, 0) \leq \frac{1}{\pi} \int_0^{2\pi} \text{Re}(1/(1 - \hat{\mu}(t))) dt \leq 2G(0, 0)$.*

Proof of the proposition :

Set $a_N(y) = G_N(0, 0) - G_N(0, y)$. We show that $(a_N(y))_{N \geq 0}$ is bounded. Note that $a_N(y) = G_N(0, 0) - G_N(-y, 0)$. Take $y \neq 0$. This is evident if 0 is transient, whereas if it is recurrent, then $P_0(T_{-y} < \infty) = 1$, as $\gcd(\text{Supp}(\mu)) = 1$. Thus $(a_N(y))_{N \geq 0}$ is bounded, by the *Claim* above.

Step 1. Let $x > 0$. We show that $\Delta(x) := \lim_{N \rightarrow +\infty} (a_N(x) + a_N(-x))$ exists. We have :

$$a_N(x) + a_N(-x) = \frac{1}{2\pi} \int_0^{2\pi} (2 - e^{-itx} - e^{itx}) \sum_{n=0}^{N-1} (\hat{\mu}(t))^n dt = \frac{1}{\pi} \int_0^{2\pi} \frac{1 - \cos(tx)}{1 - \hat{\mu}(t)} (1 - (\hat{\mu}(t))^N) dt.$$

Since $|1 - \hat{\mu}(t)| \geq \operatorname{Re}(1 - \hat{\mu}(t)) \geq \alpha t^2$ and x is fixed, $(1 - \cos(tx))/(1 - \hat{\mu}(t))$ is integrable. As $|\hat{\mu}(t)| < 1$ except for finitely many t , the required limit exists and satisfies :

$$\Delta(x) = \frac{1}{\pi} \int_0^{2\pi} \frac{1 - \cos(tx)}{1 - \hat{\mu}(t)} dt = \frac{1}{\pi} \int_0^{2\pi} (1 - \cos(tx)) \operatorname{Re}((1 - \hat{\mu}(t))^{-1}) dt. \quad (3)$$

Step 2. For $x > 0$, we give a probabilistic expression for $\Delta(x)$. First, if $y \neq 0$, using (1) and (2) :

$$a_N(y) = \mathbb{E}_0 \left(\sum_{n=0}^{T_y \wedge N-1} 1_{S_n=0} \right) + \sum_{k=1}^{N-1} P_0(T_y = k) (G_{N-k}(y, 0) - G_{N-k}(y, y)).$$

By homogeneity, $a_N(y) = \mathbb{E}_0(\sum_{n=0}^{T_y \wedge N-1} 1_{S_n=0}) - \sum_{1 \leq k < N} P_0(T_y = k) a_{N-k}(-y)$. Taking $y = x$ and adding $a_N(-x)$ we obtain :

$$a_N(x) + a_N(-x) = \mathbb{E}_0 \left(\sum_{n=0}^{T_x \wedge N-1} 1_{S_n=0} \right) + \sum_{k=1}^{N-1} P_0(T_x = k) (a_N(-x) - a_{N-k}(-x)) + P_0(T_x \geq N) a_N(-x).$$

Consider the terms on the right, when $N \rightarrow +\infty$. The first one tends to $\mathbb{E}_0(\sum_{n=0}^{T_x-1} 1_{S_n=0})$. As $P_0(S_n = y) \rightarrow_{n \rightarrow +\infty} 0$, we get $\lim_{N \rightarrow +\infty} a_{N+1}(y) - a_N(y) = 0$ and thus $\lim_{N \rightarrow +\infty} a_N(-x) - a_{N-k}(-x) = 0$ for fixed k . By dominated convergence, the second term goes to zero, as $(a_N(-x))_{N \geq 0}$ is bounded. The latter also implies that the third term goes to zero in case of recurrence and to $P_0(T_x = \infty)G(0, 0)(1 - P_0(T_{-x} < \infty))$ in case of transience. Thus, if $x > 0$:

$$\Delta(x) = \mathbb{E}_0 \left(\sum_{n=0}^{T_x-1} 1_{S_n=0} \right) + 1_{TR} G(0, 0) P_0(T_x = \infty) P_0(T_{-x} = \infty). \quad (4)$$

Step 3. By (4)=(3), $\mathbb{E}_0(\sum_{0 \leq n < T_x} 1_{S_n=0}) \leq (1/\pi) \int_0^{2\pi} \operatorname{Re}((1 - \hat{\mu}(t))^{-1}) dt$. Letting $x \rightarrow +\infty$, we obtain the first inequality. Next, from (3)=(4), for any $\delta > 0$:

$$\frac{1}{\pi} \int_{\delta}^{2\pi-\delta} (1 - \cos(tx)) \operatorname{Re}((1 - \hat{\mu}(t))^{-1}) dt \leq 2G(0, 0).$$

If $x \rightarrow +\infty$, $\pi^{-1} \int_{\delta}^{2\pi-\delta} \operatorname{Re}((1 - \hat{\mu}(t))^{-1}) dt \leq 2G(0, 0)$, by the Riemann-Lebesgue lemma. Finally we let $\delta \rightarrow 0$ to conclude the proof. \square

Remark. — When transience holds, constants in Prop. 2.2 are optimal. When $\operatorname{Supp}(\mu) \subset \mathbb{N}^*$, then $G(0, 0) = 1$ and $\Delta(x) = 2 - P_0(T_x < \infty) \rightarrow 2 - 1/\mathbb{E}(X)$, as $x \rightarrow +\infty$, by renewal theory (this is reproved in the next section). As $\lim_{x \rightarrow +\infty} \Delta(x) = \pi^{-1} \int_0^{2\pi} \operatorname{Re}((1 - \hat{\mu}(t))^{-1}) dt$, the conclusion comes from the fact that $\mathbb{E}(X)$ can take any value in $[1, +\infty]$.

Remark. — Approaching $G(0, 0)$ in two steps, first by the always finite $\lim_{N \rightarrow +\infty} (2G_N(0, 0) - G_N(0, x) - G_N(0, -x))$ and next the limit as $x \rightarrow +\infty$, is a simple but profound idea. One may observe that a similar one is used by Riemann in the first chapters of the theory of trigonometric

series. This kind of “second derivative at infinity” suggests some general link between transience and a condition of positive curvature at infinity (where the terms are to be redefined).

Remark. — The weak form of the theorem, due to Chung and Fuchs (1951), can be reduced to the following observation, where interversion is direct for $0 < s < 1$:

$$\begin{aligned} G^\mu(0, 0) &= \lim_{s \uparrow 1} \sum_{n \geq 0} s^n P^\mu(S_n = 0) = \lim_{s \uparrow 1} \sum_{n \geq 0} \frac{1}{2\pi} \int_0^{2\pi} s^n (\hat{\mu}(t))^n dt \\ &= \lim_{s \uparrow 1} \frac{1}{2\pi} \int_0^{2\pi} \operatorname{Re} \left(\frac{1}{1 - s\hat{\mu}(t)} \right) dt. \end{aligned} \quad (5)$$

The finiteness of the right-hand side is thus a transience criterion for the random walk. Notice that the operation $s \uparrow 1$ is not natural in this problem, as the level sets of $z \mapsto \operatorname{Re}(1/(1-z))$ in the unit disk are horocycles (Euclidean circles). There is no monotony in the limit and indeed, as seen above, the right-hand side may differ from $(2\pi)^{-1} \int_0^{2\pi} \operatorname{Re}((1 - \hat{\mu}(t))^{-1}) dt$.

Remark. — The theorem has been extended to general countable discrete Abelian groups by Kesten and Spitzer [8], to \mathbb{R}^d by Ornstein [9] and Port and Stone [10]. Recall also that in any case $\lim_{N \rightarrow +\infty} a_N(x)$ exists and is called the potential kernel; see Spitzer [11], chap. 7.

Remark. — In the second step of the proof of the proposition and in the transient case, one can directly write $\Delta(x) = G(0, 0)(2 - P_0(T_x < \infty) - P_0(T_{-x} < \infty))$, when $x > 0$. It is interesting to check equality with (4) in this case, i.e. that for $x \neq 0$:

$$G(0, 0)(2 - P_0(T_x < \infty) - P_0(T_{-x} < \infty)) = \frac{1}{1 - P_0(T_0 < T_x)} + G(0, 0)P_0(T_x = \infty)P_0(T_{-x} = \infty).$$

Equivalently, for fixed $x \neq 0$:

$$G(0, 0) = \frac{1}{1 - P_0(T_0 < T_x)} \times \frac{1}{1 - P_0(T_x < \infty)P_0(T_{-x} < \infty)}. \quad (6)$$

This relation, valid for a general Markov chain (transient or not), corresponds to the following decomposition of $G(0, 0)$. Let $T^{(0)} = 0$ and next $T^{(k+1)}$ be the first time $> T^{(k)}$ of passage at zero after having visited x at least once, for $k \geq 0$. Then (the k^{th} term being 0 if $T^{(k)} = +\infty$) :

$$G(0, 0) = \sum_{k \geq 0} \mathbb{E}_0 \left(\sum_{n=T^{(k)}}^{T^{(k+1)}-1} 1_{S_n=0} \right).$$

Call A_k the generic term in the above sum. Then :

$$A_0 = \mathbb{E}_0 \left(\sum_{n=0}^{T^{(1)}-1} 1_{S_n=0} \right) = \mathbb{E}_0 \left(\sum_{n=0}^{T_x-1} 1_{S_n=0} \right) = \frac{1}{1 - P_0(T_0 < T_x)}.$$

Next, for $k \geq 1$, $A_k = P_0(T^{(k)} < \infty)A_0 = (P_0(T^{(1)} < \infty))^k A_0$. Now, $P_0(T^{(1)} < \infty) = P_0(T_x < \infty)P_0(T_{-x} < \infty)$ and this gives the announced formula when summing on $k \geq 0$. Remark also that the last product is related with loops, more precisely :

$$P_0(T_x < \infty)P_0(T_{-x} < \infty) = \frac{P_0(T_x < T_0 < \infty)}{1 - P_0(T_0 < T_x)}.$$

The left-hand side is $P_0(\text{reach } x \text{ and come back at } 0)$. Decomposing it as the probability of making first $n \geq 0$ loops at 0 without touching x , then going directly to x and finally coming back to 0, this equals $\sum_{n \geq 0} P_0(T_0 < T_x)^n P_0(T_x < T_0 < \infty)$, so the right-hand side expression.

Notice furthermore that one may readily derive from (6), convening that $1/0^+ = +\infty$, the following always valid form for $\Delta(x)$, $x > 0$:

$$\Delta(x) = \mathbb{E}_0 \left(\sum_{n=0}^{T_x-1} 1_{S_n=0} \right) \left(1 + \left(1 + \frac{P_0(T_x < \infty)}{P_0(T_x = \infty)} + \frac{P_0(T_{-x} < \infty)}{P_0(T_{-x} = \infty)} \right)^{-1} \right).$$

Finally, observe that the second term in the right-hand side of (6) is symmetric with respect to 0 and x . This raises the question, for a general irreducible aperiodic Markov chain on \mathbb{Z} , whether $\lim_{N \rightarrow +\infty} G_N(x, x)/G_N(y, y) = (1 - P_y(T_y < T_x))/(1 - P_x(T_x < T_y))$.

3 Fourier transform of probability measures on \mathbb{N}^*

We study $1/(1 - \hat{\mu}_+)$ when μ_+ is a probability measure on \mathbb{N}^* . As the random walk with step μ_+ is transient, $Re(1/(1 - \hat{\mu}_+)) \in L^1(0, 2\pi)$. In fact $(1/(2\pi)) \int_0^{2\pi} Re(1/(1 - \hat{\mu}_+(t))) dt \leq 1$, directly by relation (5), as $G^{\mu_+}(0, 0) = 1$. The exact value would follow easily from the considerations of Proposition 2.2 combined with renewal theory.

We instead make the computation using complex analysis with an Herglotz type argumentation. This allows to derive the renewal theorem directly from the Riemann-Lebesgue lemma. We next show that the Fourier coefficients of $Re(1/(1 - \hat{\mu}_+))$ have an interesting probabilistic interpretation.

Lemma 3.1.

Let μ_+ , with $Supp(\mu_+) \subset \mathbb{N}^*$ and $gcd(Supp(\mu_+)) = 1$. Then $t \mapsto Re(1/(1 - \hat{\mu}_+(t)))$ is real, positive, even and in $L^1(0, 2\pi)$, with :

$$\frac{1}{2\pi} \int_0^{2\pi} Re \left(\frac{1}{1 - \hat{\mu}_+(t)} \right) dt = 1 - \frac{1}{2E^{\mu_+}(X)}. \quad (7)$$

Proof of the lemma :

Introduce $f(z) = 1/(1 - \mathbb{E}^{\mu_+}(z^X))$, holomorphic in $\Delta = \{|z| < 1\}$. For $0 \leq r \leq 1$, the map $z \mapsto Re(f(rz))$ is > 0 and harmonic on Δ . Thus, fixing $0 < r < 1$ and using the Poisson kernel :

$$Re(f(rz)) = \frac{1}{2\pi} \int_0^{2\pi} Re \left(\frac{e^{i\theta} + z}{e^{i\theta} - z} \right) Re(f(re^{i\theta})) d\theta, \quad z \in \Delta. \quad (8)$$

By holomorphic extension (using also that $f(0) = 1 \in \mathbb{R}$) :

$$f(rz) = \frac{1}{2\pi} \int_0^{2\pi} \frac{e^{i\theta} + z}{e^{i\theta} - z} Re(f(re^{i\theta})) d\theta, \quad z \in \Delta. \quad (9)$$

Taking $z = 0$ in (8), we get $1 = \frac{1}{2\pi} \int_0^{2\pi} Re(f(re^{i\theta})) d\theta$, so the positive measures $(\nu_r)_{0 < r < 1}$ on the torus $\mathbb{R} \setminus 2\pi\mathbb{Z}$ with density $Re(f(re^{i\theta}))$ have constant mass 2π . Let us take a cluster value ν of ν_r , as $r \uparrow 1$, for the weak-* topology. We get from (9), fixing first $z \in \Delta$:

$$f(z) = \frac{1}{2\pi} \int_0^{2\pi} \frac{e^{i\theta} + z}{e^{i\theta} - z} d\nu(\theta) = \frac{1}{2\pi} \int_0^{2\pi} \left(1 + 2 \sum_{n \geq 1} z^n e^{-in\theta} \right) d\nu(\theta).$$

Permuting the sum and the integral in the last expression, the Fourier coefficients of ν are uniquely determined by the development in series of f around 0. Hence ν is unique and we conclude that (ν_r) converges to ν , as $r \uparrow 1$. We shall now determine this measure.

First, when $\theta \in \mathbb{R} \setminus 2\pi\mathbb{Z}$ is fixed, then $\hat{\mu}_+(\theta) \neq 1$, so $\lim_{r \uparrow 1} Re(f(re^{i\theta})) = Re(1/(1 - \hat{\mu}_+(\theta)))$. Thus ν is locally $Re(1/(1 - \hat{\mu}_+(\theta)))d\theta$, giving $\nu = Re(1/(1 - \hat{\mu}_+(\theta)))d\theta + \alpha_0\delta_0$. Hence :

$$\frac{1}{1 - \mathbb{E}^{\mu_+}(z^X)} = \frac{1}{2\pi} \int_0^{2\pi} \left(\frac{e^{i\theta} + z}{e^{i\theta} - z} \right) Re \left(\frac{1}{1 - \hat{\mu}_+(\theta)} \right) d\theta + \frac{\alpha_0}{2\pi} \left(\frac{1+z}{1-z} \right), \quad z \in \Delta.$$

To determine α_0 , take $z = e^{-u}$, with a real $u \downarrow 0$, and multiply both sides by $1 - e^{-u}$:

$$\frac{1 - e^{-u}}{1 - \mathbb{E}^{\mu_+}(e^{-uX})} = \frac{1}{2\pi} \int_0^{2\pi} (1 - e^{-u}) \left(\frac{e^{i\theta} + e^{-u}}{e^{i\theta} - e^{-u}} \right) Re \left(\frac{1}{1 - \hat{\mu}_+(\theta)} \right) d\theta + \frac{\alpha_0}{2\pi} (1 + e^{-u}).$$

The left-hand side goes to $1/E^{\mu_+}(X)$. As $(1 - e^{-u}) \times (e^{i\theta} + e^{-u})/(e^{i\theta} - e^{-u})$ stays bounded by 2, the first term on the right-hand side tends to 0 by dominated convergence. Finally we get $\alpha_0/\pi = 1/E^{\mu_+}(X)$ and therefore the relation :

$$\frac{1}{1 - \mathbb{E}^{\mu_+}(z^X)} = \frac{1}{2\pi} \int_0^{2\pi} \left(\frac{e^{i\theta} + z}{e^{i\theta} - z} \right) Re \left(\frac{1}{1 - \hat{\mu}_+(\theta)} \right) d\theta + \frac{1}{2\mathbb{E}^{\mu_+}(X)} \left(\frac{1+z}{1-z} \right), \quad z \in \Delta.$$

Expression (7) is now given by $z = 0$. □

Proposition 3.2.

Let μ_+ , with $Supp(\mu_+) \subset \mathbb{N}^*$ and $gcd(Supp(\mu_+)) = 1$.

i) For $x \geq 1$:

$$\frac{1}{\pi} \int_0^{2\pi} \cos(tx) Re \left(\frac{1}{1 - \hat{\mu}_+(t)} \right) dt = P_0^{\mu_+}(T_x < \infty) - \frac{1}{E^{\mu_+}(X)}.$$

As a result (renewal theorem, Erdős-Feller-Pollard [6]), $\lim_{x \rightarrow +\infty} P_0^{\mu_+}(T_x < \infty) = 1/E^{\mu_+}(X)$.

ii) The function $t \mapsto t/|1 - \hat{\mu}_+(t)|$ belongs to $L^2(0, \pi)$. The function $t \mapsto Im(1/(1 - \hat{\mu}_+(t)))$ is real and odd; it does not belong to $L^1(0, \pi)$, whenever $E^{\mu_+}(X) < \infty$. Also, for $x \geq 1$:

$$\frac{1}{\pi} \int_0^{2\pi} \sin(tx) Im \left(\frac{1}{1 - \hat{\mu}_+(t)} \right) dt = P_0^{\mu_+}(T_x < \infty).$$

iii) We have $|1 - \hat{\mu}_+|^{-1} \in L^\gamma(0, 2\pi)$, $0 < \gamma < 1$. Also $t^\varepsilon |1 - \hat{\mu}_+(t)|^{-1} \in L^1(0, \pi)$, $\varepsilon > 0$.

Proof of the proposition :

i) Start as in Prop. 2.2. Fixing $x \geq 1$, we first have, for $N \geq 1$:

$$2G_N^{\mu_+}(0, 0) - G_N^{\mu_+}(0, x) - G_N^{\mu_+}(0, -x) = \frac{1}{\pi} \int_0^{2\pi} \frac{1 - \cos(tx)}{1 - \hat{\mu}_+(t)} (1 - (\hat{\mu}_+(t))^N) dt.$$

We can again take the limit as $N \rightarrow +\infty$ in the right-hand side and next the real part. The limit of the left-hand side is trivial, so for any $x \geq 1$:

$$2 - P_0^{\mu_+}(T_x < \infty) = \frac{1}{\pi} \int_0^{2\pi} (1 - \cos(tx)) \operatorname{Re} \left(\frac{1}{1 - \hat{\mu}_+(t)} \right) dt. \quad (10)$$

The fact that $\operatorname{Re}(1/(1 - \hat{\mu}_+)) \in L^1(0, 2\pi)$ can be recovered when minoring the right-hand side by $\pi^{-1} \int_\delta^{2\pi-\delta}$, $\delta > 0$, letting $x \rightarrow +\infty$ with the Riemann-Lebesgue lemma and finally $\delta \rightarrow 0$.

Subtracting (10) to twice (7), we obtain the desired relation for $x \geq 1$. Then the renewal theorem is a consequence of the Riemann-Lebesgue lemma. Mention in passing another proof, even simpler. By (10) and the Riemann-Lebesgue lemma, $\lim_{x \rightarrow +\infty} P_0(T_x < \infty)$ exists; then in Spitzer [11], P3, the limit is identified as $1/\mathbb{E}^{\mu_+}(X)$.

ii) Let us place on $(0, \pi)$. As $1 - \operatorname{Re}(\hat{\mu}_+(t)) \geq \alpha t^2$, we have :

$$\frac{t}{|1 - \hat{\mu}_+(t)|} = \frac{t}{\sqrt{1 - \operatorname{Re}(\hat{\mu}_+(t))}} \times \frac{\sqrt{1 - \operatorname{Re}(\hat{\mu}_+(t))}}{|1 - \hat{\mu}_+(t)|} \in L^2(0, \pi),$$

as the first term is bounded and the square of the second one is $\operatorname{Re}(1/(1 - \hat{\mu}_+(t))) \in L^1(0, \pi)$. For the imaginary part, let us write, fixing $x \geq 1$ and taking $N \geq 1$:

$$\begin{aligned} G_N^{\mu_+}(0, x) - G_N^{\mu_+}(0, -x) &= \frac{1}{2\pi} \int_0^{2\pi} (e^{-itx} - e^{itx}) \sum_{n=0}^{N-1} (\hat{\mu}_+(t))^n dt \\ &= -\frac{i}{\pi} \int_0^{2\pi} \sin(tx) \left(\frac{1 - (\hat{\mu}_+(t))^N}{1 - \hat{\mu}_+(t)} \right) dt. \end{aligned} \quad (11)$$

As $\sin(tx) \operatorname{Im}(1/(1 - \hat{\mu}_+(t)))$ is integrable, we can let $N \rightarrow +\infty$ and then take the imaginary part inside the integral. The left-hand side limit being obvious, we obtain for $x \geq 1$:

$$P_0^{\mu_+}(T_x < \infty) = \frac{1}{\pi} \int_0^{2\pi} \sin(tx) \operatorname{Im} \left(\frac{1}{1 - \hat{\mu}_+(t)} \right) dt.$$

Whenever $E^{\mu_+}(X) < \infty$, the left-hand side goes to $1/E^{\mu_+}(X) > 0$, hence the Riemann-Lebesgue lemma is not verified, giving $\operatorname{Im}(1/(1 - \hat{\mu}_+(t))) \notin L^1(0, 2\pi)$.

iii) The holomorphic function $f(z) = 1/(1 - \mathbb{E}^{\mu_+}(z^X))$, $z \in \Delta$, has a positive and harmonic real part. The latter thus is in $h^1(\Delta)$. By Duren [5], Theorem 4.2, $f \in H^\gamma(\Delta)$, $0 < \gamma < 1$, i.e. :

$$\sup_{0 < r < 1} \int_0^{2\pi} |f(re^{i\theta})|^\gamma d\theta < \infty.$$

By Fatou's lemma, as $r \uparrow 1$, we get $\int_0^{2\pi} |1 - \hat{\mu}_+(\theta)|^{-\gamma} d\theta < \infty$. For the last point, write :

$$\frac{t^\varepsilon}{|1 - \hat{\mu}_+(t)|} = \frac{t^\varepsilon}{|1 - \hat{\mu}_+(t)|^{\varepsilon/2}} \times \frac{1}{|1 - \hat{\mu}_+(t)|^{1-\varepsilon/2}} \in L^1(0, \pi),$$

as the first term on the right-hand side is bounded. This ends the proof of the proposition. \square

Corollary 3.3.

i) $\operatorname{Re}((1 - \hat{\mu}_+)^{-1}) \in L^2(0, 2\pi)$ iff $(G^{\mu_+}(0, x) - 1/\mathbb{E}^{\mu_+}(X))_{x \geq 0} \in l^2$.

ii) $\operatorname{Im}((1 - \hat{\mu}_+)^{-1}) \in L^2(0, 2\pi)$ iff $(G^{\mu_+}(0, x))_{x \geq 0} \in l^2$. In this case $|1 - \hat{\mu}_+|^{-1} \in L^2(0, 2\pi)$.

Proof of the corollary :

Point *i*) is clear as $Re((1 - \hat{\mu}_+)^{-1}) \in L^1(0, 2\pi)$, hence the $(G^{\mu_+}(0, x) - 1/\mathbb{E}^{\mu_+}(X))$ are its Fourier coefficients. Idem, when $Im((1 - \hat{\mu}_+)^{-1}) \in L^2(0, 2\pi)$ then the $(G^{\mu_+}(0, x))_{x \geq 0}$ are its Fourier coefficients and thus belong to l^2 . Reciprocally, if $(G^{\mu_+}(0, x))_{x \geq 0}$ is l^2 , define the L^2 odd function $f(t) = \sum_{x \geq 1} G^{\mu_+}(0, x) \sin(xt)$. For all $x \in \mathbb{Z}$, we thus have :

$$\int_0^{2\pi} \frac{\sin(tx)}{\sin t} [\sin t (Im((1 - \hat{\mu}_+(t))^{-1}) - f(t))] dt = 0.$$

The function inside the brackets belongs to L^2 and is even. Writing $\sin(t(1+x)) + \sin(t(1-x)) = 2\sin(t)\cos(tx)$, for $x \geq 0$, the latter is thus orthogonal to all $\cos(tx)$, $x \geq 0$, hence equals zero a.e.. Hence $Im((1 - \hat{\mu}_+)^{-1}) = f$, a.e., and thus belongs to L^2 .

Finally, when $Im((1 - \hat{\mu}_+)^{-1}) \in L^2(0, 2\pi) \subset L^1(0, 2\pi)$, then $\mathbb{E}^{\mu_+}(X) = \infty$ by Proposition 3.2. The conditions on Fourier coefficients in order to belong to $L^2(0, 2\pi)$, for $Re((1 - \hat{\mu}_+)^{-1})$ and $Im((1 - \hat{\mu}_+)^{-1})$, are now identical. □

Remark. — In Proposition 3.2 the Fourier coefficients of $Re((1 - \hat{\mu}_+)^{-1})$ and $Im((1 - \hat{\mu}_+)^{-1})$ are probabilistic quantities. Those of $Re((1 - \hat{\mu}_+)^{-1})$ exactly measure the error in the renewal theorem. One further question is whether $t^{1/2+\varepsilon}/|1 - \hat{\mu}_+(t)|$ belongs to $L^2(0, \pi)$, for $\varepsilon > 0$. It is in $L^1(0, \pi)$, by the Cauchy-Schwarz inequality, since $t^{-1/2+\varepsilon}$ and $t/|1 - \hat{\mu}_+(t)|$ are in $L^2(0, \pi)$. Notice also that $\sum_{x \geq 0} G^{\mu_+}(0, x) = +\infty$, hence $(G^{\mu_+}(0, x))_{x \geq 0}$ is never l^1 . The question whether $(G^{\mu_+}(0, x))_{x \geq 0} \in l^2$ is related to the transience of some symmetric oscillating random walk, as detailed in the next section.

Remark. — For complex numbers a and b , write $\langle a, b \rangle = Re(a\bar{b})$ for the real inner product of the vectors in \mathbb{R}^2 with affixes a and b . As a corollary of Proposition 3.2, although $1/|1 - \hat{\mu}_+|$ may not belong to $L^1(0, 2\pi)$, we have $\langle (1 - \hat{\mu}_+(t))^{-1}, e^{itx} \rangle \in L^1(0, 2\pi)$, for all $x \in \mathbb{Z}$, with :

$$\begin{cases} \frac{1}{\pi} \int_0^{2\pi} \langle (1 - \hat{\mu}_+(t))^{-1}, e^{itx} \rangle dt = 2G^{\mu_+}(0, x) - 1/E^{\mu_+}(X), & x \geq 0, \\ \frac{1}{\pi} \int_0^{2\pi} \langle (1 - \hat{\mu}_+(t))^{-1}, e^{-itx} \rangle dt = -1/E^{\mu_+}(X), & x \geq 1. \end{cases}$$

Let $x \geq 1$. Even if this is not always true, in general $Im(\hat{\mu}_+(t)) \geq 0$ for small $t > 0$, so in this case $1/(1 - \hat{\mu}_+(t))$ is in the first quadrant, as well as e^{itx} and contrary to e^{-itx} . Hence it is natural that the first integral above is larger than the second one.

To conclude this section, let us present a variation on Lemma 3.1.

Lemma 3.4.

Let μ_+ , with $Supp(\mu_+) \subset \mathbb{N}^*$ and $gcd(Supp(\mu_+)) = 1$. Then :

$$\frac{1}{2\pi} \int_0^{2\pi} Re \left(\frac{1}{1 - \hat{\mu}_+(t)} \right) \frac{1}{1 + \left[\frac{\sin(t/2)}{\sinh(1/2)} \right]^2} dt = \frac{\tanh(1/2)}{1 - \mathbb{E}^{\mu_+}(e^{-X})} - \frac{1}{2\mathbb{E}^{\mu_+}(X)}.$$

Proof of the lemma :

Introduce the homography $\rho(z) = (1-z)/(1+z)$, exchanging the open unit disk Δ and the half plane $Re > 0$. Let $f(z) = 1/(1 - \mathbb{E}^{\mu_+}(e^{-\rho(z)X}))$, $z \in \Delta$. It is holomorphic in Δ and $Re(f)$ is > 0 and harmonic on Δ . By harmonicity, for $0 < r < 1$:

$$1/(1 - \mathbb{E}^{\mu_+}(e^{-X})) = \frac{1}{2\pi} \int_0^{2\pi} \operatorname{Re}(f(re^{i\theta}))d\theta.$$

Proceeding as in Lemma 3.1, we get :

$$f(z) = \frac{1}{2\pi} \int_0^{2\pi} \frac{e^{i\theta} + z}{e^{i\theta} - z} d\nu(\theta), \quad z \in \Delta,$$

where ν is the limit as $r \uparrow 1$ of the positive measures ν_r on $\mathbb{R} \setminus 2\pi\mathbb{Z}$ with density $\operatorname{Re}(f(re^{i\theta}))$. In order to detail ν , note first that when $\theta \in \mathbb{R} \setminus 2\pi\mathbb{Z}$ is fixed, then :

$$\lim_{r \uparrow 1} \rho(re^{i\theta}) = \frac{1 - e^{i\theta}}{1 + e^{i\theta}} = -i \tan(\theta/2).$$

When $\theta \notin \{\pi\} \cup \{2 \arctan(2k\pi), k \in \mathbb{Z}\}$, then $\mathbb{E}^{\mu_+}(e^{i \tan(\theta/2)X}) \neq 1$ and ν is locally $g(\theta)d\theta$, with $g(\theta) = \operatorname{Re}(1/(1 - \mathbb{E}^{\mu_+}(e^{i \tan(\theta/2)X})))$. As result ν decomposes as follows :

$$\nu = \operatorname{Re}(1/(1 - \mathbb{E}^{\mu_+}(e^{i \tan(t/2)X}))dt + \alpha_\pi \delta_\pi + \sum_{k \in \mathbb{Z}} \alpha_k \delta_{\theta_k},$$

where $\theta_k = 2 \arctan(2\pi k)$, for non-negative α_π and (α_k) . In order to determine these coefficients, start from the relation, for $z \in \Delta$:

$$\frac{1}{1 - \mathbb{E}^{\mu_+}(e^{-\rho(z)X})} = \frac{1}{2\pi} \int_0^{2\pi} \left(\frac{e^{i\theta} + z}{e^{i\theta} - z} \right) g(\theta)d\theta + \frac{\alpha_\pi}{2\pi} \left(\frac{1 - z}{1 + z} \right) + \sum_{k \in \mathbb{Z}} \frac{\alpha_k}{2\pi} \left(\frac{e^{i\theta_k} + z}{e^{i\theta_k} - z} \right).$$

Recall that g is integrable and $\sum_k \alpha_k < \infty$ (the total mass of ν is $2\pi/(1 - \mathbb{E}^{\mu_+}(e^{-X}))$). Take $z = -r$ above, as $r \uparrow 1$, and multiply first both sides by $1 - r$. Notice that for $\theta \neq \pi \pmod{2\pi}$, $(1 - r) \times (e^{i\theta} - r)/(e^{i\theta} + r)$ stays bounded by 2 and converges to 0. Hence as $r \uparrow 1$, by dominated convergence, the right-hand side converges to $\frac{\alpha_\pi}{2\pi}(1 + 1)$, whereas the left-hand side is equivalent to $(1 - r)/P^{\mu_+}(X > 0)$ and therefore goes to 0. We obtain $\alpha_\pi = 0$.

Fixing $k \in \mathbb{Z}$, take now $z = re^{i\theta_k}$ and let $r \uparrow 1$, after multiplying both sides by $(1 - r)$. Idem, for $\theta \neq \theta_k \pmod{2\pi}$, $(1 - r) \times (e^{i\theta} + re^{i\theta_k})/(e^{i\theta} - re^{i\theta_k})$ stays bounded by 2 and converges to 0. By dominated convergence the right-hand side converges to α_k/π , and this equals :

$$\lim_{r \uparrow 1} \frac{1 - r}{1 - \mathbb{E}^{\mu_+}(e^{-\rho(re^{i\theta_k})X})}.$$

To determine the limit, note first that $\lim_{u \downarrow 0^+} u/(1 - \mathbb{E}^{\mu_+}(e^{-uX})) = 1/\mathbb{E}^{\mu_+}(X)$, by monotone convergence. Next, $\rho(e^{i\theta_k}) = -i \tan(\theta_k/2) = -2ik\pi$, so the denominator is :

$$1 - \mathbb{E}^{\mu_+}(e^{-(\rho(re^{i\theta_k}) - \rho(e^{i\theta_k}))X}).$$

We next have, decomposing in real and imaginary parts :

$$\begin{aligned} \rho(re^{i\theta_k}) - \rho(e^{i\theta_k}) &= \frac{2(1 - r)e^{i\theta_k}}{(1 + re^{i\theta_k})(1 + e^{i\theta_k})} = \frac{1 - r}{\cos(\theta_k/2)} \frac{(1 + r) \cos(\theta_k/2) + i(1 - r) \sin(\theta_k/2)}{(1 + r)^2 \cos^2(\theta_k/2) + (1 - r)^2 \sin^2(\theta_k/2)} \\ &= A(r) + iB(r). \end{aligned}$$

- Case 1 : $\mathbb{E}^{\mu_+}(X) = +\infty$. Then, as $A(r)/(1 - r) \rightarrow_{r \uparrow 1} 1/(2 \cos^2(\theta_k/2)) > 0$:

$$\left| \frac{1-r}{1 - \mathbb{E}^{\mu_+}(e^{-\rho(re^{i\theta_k})X})} \right| \leq \frac{1-r}{1 - \mathbb{E}^{\mu_+}(e^{-A(r)X})} \xrightarrow{r \uparrow 1} 0.$$

- Case 2 : $\mathbb{E}^{\mu_+}(X) < +\infty$. Then :

$$\frac{1 - \mathbb{E}^{\mu_+}(e^{-(\rho(re^{i\theta_k}) - \rho(e^{i\theta_k}))X})}{1-r} = \frac{1 - \mathbb{E}^{\mu_+}(e^{-A(r)X})}{1-r} + \mathbb{E}^{\mu_+}(e^{-A(r)X}(1 - e^{-iB(r)X})/(1-r)).$$

As $r \uparrow 1$, the first term on the right-hand side tends to $\mathbb{E}^{\mu_+}(X)/(2 \cos^2(\theta_k/2))$. Since $t \mapsto e^{it}$ is 1-Lipschitz on \mathbb{R} , $|1 - e^{-iB(r)X}|/(1-r) \leq |B(r)|X/(1-r)$. As $|B(r)| \leq C(1-r)^2$, the previous quantity is both bounded by $C'X$ and tends to 0 as $r \uparrow 1$. Since $\mathbb{E}^{\mu_+}(X) < \infty$, by dominated convergence the second term goes to 0 as $r \uparrow 1$. Finally, $\alpha_k/\pi = 2 \cos^2(\theta_k/2)/\mathbb{E}^{\mu_+}(X)$. As $\cos^2(\theta_k/2) = 1/(1 + 4\pi^2 k^2)$, this leads to :

$$\begin{aligned} \frac{1}{1 - \mathbb{E}^{\mu_+}(e^{-\rho(z)X})} &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \left(\frac{e^{i\theta} + z}{e^{i\theta} - z} \right) \operatorname{Re} \left(\frac{1}{1 - \mathbb{E}^{\mu_+}(e^{i \tan(\theta/2)X})} \right) d\theta \\ &+ \frac{1}{\mathbb{E}^{\mu_+}(X)} \sum_{k \in \mathbb{Z}} \left(\frac{e^{i\theta_k} + z}{e^{i\theta_k} - z} \right) \frac{1}{1 + 4\pi^2 k^2}, \quad z \in \Delta. \end{aligned}$$

Taking $z = 0$ and making the change of variable $\theta = 2 \arctan t$ in the first integral :

$$\begin{aligned} \frac{1}{1 - \mathbb{E}^{\mu_+}(e^{-X})} &= \frac{1}{\pi} \int_{-\infty}^{\infty} \operatorname{Re} \left(\frac{1}{1 - \hat{\mu}_+(t)} \right) \frac{1}{1+t^2} dt + \frac{1}{\mathbb{E}^{\mu_+}(X)} \sum_{k \in \mathbb{Z}} \frac{1}{1 + 4\pi^2 k^2} \\ &= \frac{1}{\pi} \int_0^{2\pi} \operatorname{Re} \left(\frac{1}{1 - \hat{\mu}_+(t)} \right) \sum_{k \in \mathbb{Z}} \frac{1}{1 + (t + 2k\pi)^2} dt + \frac{1}{\mathbb{E}^{\mu_+}(X)} \sum_{k \in \mathbb{Z}} \frac{1}{1 + 4\pi^2 k^2}. \end{aligned}$$

Finally, for a real a and a complex number z , we have (cf Cartan [4], ex. 4, p172) :

$$\frac{\pi}{a} \frac{\sinh(2\pi a)}{\cosh(2\pi a) - \cos(2\pi z)} = \sum_{k \in \mathbb{Z}} \frac{1}{a^2 + (z + k)^2}.$$

Taking $z = t/(2\pi)$ and $a = 1/(2\pi)$, we get $\sum_{k \in \mathbb{Z}} 1/(1 + (t + 2k\pi)^2) = (1/2) \frac{\sinh(1)}{\cosh(1) - \cos t}$ and we arrive at the announced formula. \square

Remark. — In the function $z \mapsto 1/(1 - \mathbb{E}^{\mu_+}(e^{-\rho(z)X}))$, $z \in \Delta$, the term $\mathbb{E}^{\mu_+}(e^{-\rho(z)X})$ could be replaced by $\mathbb{E}^{\mu_+}(h(z)^X)$ or $\mathbb{E}^{\mu_+}(h(z^X))$, for any h holomorphic in Δ with $|h| < 1$ in Δ , for example an automorphism of Δ . This would give new relations. However, $\lim_{r \uparrow 1} \mathbb{E}^{\mu_+}(h(re^{i\theta})^X)$ is often delicate to determine concretely.

4 Kemperman's oscillating random walk

We move to a non-homogeneous context on \mathbb{Z} and suppose that the random walk, written as (S_n) , jumps on \mathbb{Z} according to probability measures μ on $(-\infty, 0]$ and ν on $[1, +\infty)$, respectively. For later application, no moment assumption is made on either μ or ν . This model of "oscillating random walk" has been introduced by Kemperman in [7].

We associate to μ and ν one-sided sub-probability measures μ_+ and ν_- , respectively, with $\text{Supp}(\mu_+) \subset \mathbb{N}^*$ and $\text{Supp}(\nu_-) \subset -\mathbb{N}^*$, in the following way. For any $k \geq 1$:

$$\mu_+(k) = \mathbb{P}_0^\mu(S_n \text{ enters } \mathbb{N}^* \text{ at } k) \text{ and } \nu_-(-k) = \mathbb{P}_0^\nu(S_n \text{ enters } -\mathbb{N}^* \text{ at } -k). \quad (12)$$

The question whether 0 is recurrent for (S_n) will naturally appear as a property of the sole couple (μ_+, ν_-) . We first discuss the relation between μ and μ_+ (or ν and ν_-).

4.1 Link between μ and μ_+

For a measure w on \mathbb{Z} , let the restrictions $w^- = w|_{\leq 0}$ and $w^+ = w|_{\geq 1}$. We place in the commutative Banach algebra of signed measures on \mathbb{Z} , with convolution as product, written as $w_1 w_2$. Recall the fundamental property of the exponential, $\exp(w_1 + w_2) = \exp(w_1) \exp(w_2)$. Also, if w is a non-negative measure with mass < 1 :

$$\delta_0 - w = \exp(-L_w), \text{ where } L_w = \sum_{n \geq 1} \frac{w^n}{n}.$$

Given a probability measure w on \mathbb{Z} , we write (S_n^w) for the *i.i.d.* random walk with step w , setting $S_0^w = 0$. When several (S_n^w) appear, corresponding to different probability measures, they are supposed to be independent.

Proposition 4.1.

Let μ be a probability measure on \mathbb{Z} and μ_+ defined as in (12). Then μ_+ is a probability measure iff $\sum_{n \geq 1} \mu^n(\mathbb{N}^*)/n = +\infty$. When $\hat{\mu}_+(t) \neq 1$:

$$\frac{1}{1 - \hat{\mu}_+(t)} = \lim_{s \uparrow 1} e^{\sum_{n \geq 1} s^n \widehat{(\mu^n)^+}(t)/n}.$$

Proof of the proposition :

Let $0 < s < 1$ and define $L_\mu^\pm = \sum_{n \geq 1} s^n (\mu^n)^\pm / n$. Then $\delta_0 - s\mu = \exp(-L_\mu^+) \exp(-L_\mu^-)$. Set $N = \min\{n \geq 1, S_n^\mu \geq 1\}$, $\eta_0 = \delta_0$ and $\eta_n(A) = P^\mu(N \geq n, S_n \in A)$, $n \geq 1$. Let $\eta = \sum_{n \geq 0} s^n \eta_n$.

By definition, $\eta_{n+1} = (\eta_n)^- \mu$. Summing on $n \geq 0$ with coefficients s^{n+1} , we get $\eta - \delta_0 = \eta^- s\mu$. This gives $\eta^- (\delta_0 - s\mu) = \delta_0 - \eta^+$ and therefore $\eta^- \exp(-L_\mu^-) = (\delta_0 - \eta^+) \exp(L_\mu^+)$. The left-hand side is a measure on $(-\infty, 0]$ and the right-hand side on $[0, +\infty)$, with mass at 0 equal to one.

Hence $\eta^- \exp(-L_\mu^-) = (\delta_0 - \eta^+) \exp(L_\mu^+) = \delta_0$. This gives $\delta_0 - \eta^+ = \exp(-L_\mu^+)$ or equivalently $\sum_{n \geq 0} (\eta^+)^n = \exp(L_\mu^+)$, from which the assertions follow (observing that $\mu_+ = \lim_{s \uparrow 1} \eta^+$). \square

Remark. — Can we write the limit as $e^{\sum_{n \geq 1} \widehat{(\mu^n)^+}(t)/n}$ for $0 < t < 2\pi$, when μ_+ is a probability and $|\hat{\mu}| < 1$ on $(0, 2\pi)$, hence suppressing the disgraceful $\lim_{s \uparrow 1}$?

4.2 The concentrated Markov chain

Lemma 4.2.

i) If either $\mu_+(\mathbb{N}^) < 1$ or $\nu_-(-\mathbb{N}^*) < 1$, then (S_n) is transient.*

ii) Let $\mu_+(\mathbb{N}^) = \nu_-(-\mathbb{N}^*) = 1$ and call (Z_n) the Markov chain jumping with μ_+ on $(-\infty, 0]$ and ν_- on $[1, +\infty)$. Then 0 is recurrent for (S_n) is recurrent if and only if 0 is recurrent for (Z_n) .*

Proof of the lemma :

i) If $\mu_+(\mathbb{N}) < 1$, then (S_n^μ) a.-s. makes only finitely many records in the right direction, giving $S_n^\mu \rightarrow -\infty$, a.-s.. Thus $P_0(S_n^\mu \rightarrow -\infty, \text{ with } S_k^\mu \leq 0, \forall k \geq 0) > 0$ and so $P_0(S_n \rightarrow -\infty \text{ and } S_k \leq 0, \forall k \geq 0) > 0$. Hence (S_n) is transient. The situation $\nu_-(\mathbb{N}^*) < 1$ is treated similarly.

ii) Let $\mu_+(\mathbb{N}^*) = \nu_-(\mathbb{N}^*) = 1$. Then (S_n) visits both $(-\infty, 0]$ and $[1, +\infty)$ infinitely often, a.-s.. Start (Z_n) at 0. Idem, start (S_n) at 0 and let τ be its a.-s. finite entrance time in $[1, +\infty)$. Then S_τ has the law of Z_1 . Looking now at $(S_{\tau+n})_{n \geq 0}$ at left record times on $[1, +\infty)$ and right record times on $(-\infty, 0]$, then $(S_{\tau+n})_{n \geq 0}$ a.-s. comes back to 0 iff $(Z_n)_{n \geq 1}$ does. \square

We now suppose that $\mu_+(\mathbb{N}^*) = \nu_-(\mathbb{N}^*) = 1$. Notice that this property may not be sufficient for recurrence (rarely, very large jumps across 0 may occur, ensuring $|S_n| \rightarrow +\infty$). Using the previous lemma, we focus on (Z_n) . This random walk is rather particular as it can essentially be reduced to the two sequences of positive and negative jumps (written in the order they appear).

Lemma 4.3.

i) Let $0 \leq k \leq n$ and $x, y \in \mathbb{Z}$. The sequences $(l_i^+)_{1 \leq i \leq k}$ and $(-l_j^-)_{1 \leq j \leq n-k}$ are respectively the ordered sequences of positive and negative jumps of a trajectory (which is then unique) of $(Z_m)_{0 \leq m \leq n}$, with $Z_0 = x$ and $Z_n = y$, if and only if $\sum_{1 \leq i \leq k} l_i^+ - \sum_{1 \leq j \leq n-k} l_j^- = y - x$ and $(l_k^+ \geq y, \text{ if } k \geq 1 \ \& \ y > 1)$ and $(-l_{n-k}^- \leq y - 1, \text{ if } n - k \geq 1 \ \& \ y \leq -1)$.

ii) We have $P_x(Z_n = y) = \sum_{k=0}^n P(S_k^{\mu_+} + S_{n-k}^{\nu_-} = y - x, X_k^{\mu_+} \geq y, X_{n-k}^{\nu_-} \leq y - 1)$, for any $n \geq 0$ and $x, y \in \mathbb{Z}$, where the second and third conditions are omitted if $k = 0$ or $n - k = 0$.

Proof of the lemma :

i) Starting from a trajectory of $(Z_m)_{0 \leq m \leq n}$, denote by $(l_i^+)_{1 \leq i \leq k}$ and $(-l_j^-)_{1 \leq j \leq n-k}$, for some $0 \leq k \leq n$, the ordered sequences of positive and negative jumps. From these two sequences we recover the trajectory, simply noting that the current position of the walker gives the direction of the next jump. Hence, starting for example from $x \leq 0$, use first the (l_i^+) until reaching $[1, +\infty)$, next the (l_j^-) until coming back to $(-\infty, 0]$, etc, until exhausting the two lists.

Starting from x and arriving at y , we have $\sum_i l_i^+ - \sum_j l_j^- = y - x$. Suppose that $k \geq 1$ and $y > 1$ (the cases $n - k \geq 1$ and $y \leq -1$ would be treated in the same way). When running the exhaustion process of the lists, two cases may occur :

- the (l_i^+) are finished first. When this happens, the position is $> y$ and the last positive jump must have been $> y$. The path to y is ended with the remaining negative jumps.
- the list (l_i^-) is ended first. The trajectory terminates with positive jumps (each with a starting point in $(-\infty, 0]$) and the last one has to be $\geq y$.

Reciprocally, suppose the conditions satisfied and for instance $k \geq 1 \ \& \ y \geq 1$. Starting from x , run the exhaustion process of the lists. If the (l_j^-) finish first, only positive jumps remain. As the last one is $\geq y$, this last sequence of jumps will have non-positive starting points, so the trajectory will be “admissible”. If the (l_i^+) are ended first, we are $> y$ when this happens. Only remain negative jumps for going to y , hence the trajectory is also “admissible”.

ii) Let independent $(X_k^{\mu_+}, X_l^{\nu_-})_{k, l \geq 0}$, with $\mathcal{L}(X_k^{\mu_+}) = \mu_+$, $\mathcal{L}(X_l^{\nu_-}) = \nu_-$. When running from some fixed x the exhaustion process with the two lists $(X_k^{\mu_+})_{k \geq 0}$ and $(X_l^{\nu_-})_{l \geq 0}$, we obtain a realization of $(Z_n)_{n \geq 0}$, with $Z_0 = x$. By *i)*, $\{Z_n = y, \text{ with } k \text{ positive jumps}\} = \{S_k^{\mu_+} + S_{n-k}^{\nu_-} = y - x, X_k^{\mu_+} \geq y, X_{n-k}^{\nu_-} \leq y - 1\}$. Take the probability and sum on $0 \leq k \leq n$ to get the result. \square

The Green function of (Z_n) appears to be related to the l^2 -inner product of the Green functions G^{μ_+} and G^{ν_-} . The next result recovers Theorems 4.6 and 4.8 of Kemperman [7].

Proposition 4.4.

i) The Green function G^Z of (Z_n) verifies $G^Z(0, 0) = \sum_{m \geq 0} G^{\mu_+}(0, m)G^{\nu_-}(0, -m)$ and :

$$\overline{G^Z}(0, 0) = \lim_{s \uparrow 1} \frac{1}{2\pi} \int_0^{2\pi} \operatorname{Re} \left((1 - s\hat{\mu}_+(t))^{-1} (1 - s\hat{\nu}_-(t))^{-1} \right) dt.$$

ii) When $\nu_-(A) = \mu_+(-A)$, for $A \subset \mathbb{Z}$, then $G^Z(0, 0) = (2\pi)^{-1} \int_0^{2\pi} |1 - \hat{\mu}_+(t)|^{-2} dt$.

Proof of the proposition :

i) By Lemma 4.3, $P_0(Z_n = 0) = \sum_{k=0}^n P(S_k^{\mu_+} + S_{n-k}^{\nu_-} = 0) = \sum_{k=0}^n (\mu_+^k \nu_-^{n-k})(0)$. Thus :

$$G^Z(0, 0) = \sum_{k, l \geq 0} (\mu_+^k \nu_-^l)(0) = \sum_{m \geq 0} \sum_{k, l \geq 0} \mu_+^k(m) \nu_-^l(-m) = \sum_{m \geq 0} G^{\mu_+}(0, m) G^{\nu_-}(0, -m).$$

Taking $0 < s < 1$, we next have :

$$\sum_{k, l \geq 0} s^{k+l} (\mu_+^k \nu_-^l)(0) = \frac{1}{2\pi} \int_0^{2\pi} \sum_{k, l \geq 0} (s^k \hat{\mu}_+^k s^l \hat{\nu}_-^l)(t) dt = \frac{1}{2\pi} \int_0^{2\pi} \operatorname{Re} \left(\frac{1}{1 - s\hat{\mu}_+(t)} \frac{1}{1 - s\hat{\nu}_-(t)} \right) dt.$$

As $G^Z(0, 0) = \lim_{s \uparrow 1} \sum_{k, l \geq 0} s^{k+l} (\mu_+^k \nu_-^l)(0)$, we get the desired expression.

ii) In this case, $\hat{\mu}_+(t)$ is the conjugate of $\hat{\nu}_-(t)$, $t \in \mathbb{R}$. By i), $2\pi G^Z(0, 0)$ equals :

$$\lim_{s \uparrow 1} \int_0^{2\pi} |1 - s\hat{\mu}_+(t)|^{-2} dt = \lim_{s \uparrow 1} s^{-2} \int_0^{2\pi} |s^{-1} - \hat{\mu}_+(t)|^{-2} dt = \int_0^{2\pi} |1 - \hat{\mu}_+(t)|^{-2} dt,$$

where monotone convergence is used at the end (this does not work in general). □

Remark. — Point i) is the analog of the Chung-Fuchs result. One may now ask whether $(\operatorname{Re}((1 - \hat{\mu}_+)^{-1}(1 - \hat{\nu}_-)^{-1}))^- \in L^1(0, 2\pi)$, so that $\int_0^{2\pi} \operatorname{Re}((1 - \hat{\mu}_+(t))^{-1}(1 - \hat{\nu}_-(t))^{-1}) dt$ always has a meaning, and then if $G^Z(0, 0)$ has the same order as the integral. Note also in passing the relation $G^Z(0, 0) = \mathbb{E}(\sum_{k, l \geq 0} 1_{S_k^{\mu_+} = -S_l^{\nu_-}}) = \mathbb{E}(\operatorname{card}(\{S_k^{\mu_+}, k \geq 0\} \cap \{-S_l^{\nu_-}, l \geq 0\}))$.

4.3 Characterization of the positive recurrence of (Z_n)

Proposition 4.5.

Let $\mu_+(\mathbb{N}^*) = \nu_-(-\mathbb{N}^*) = 1$. Suppose that $\gcd(\operatorname{Supp}(\mu_+)) = \gcd(\operatorname{Supp}(\nu_-)) = 1$ and that (Z_n) is aperiodic and irreducible on \mathbb{Z} , with $Z_0 = 0$.

i) The random walk (Z_n) is positive recurrent iff $\mathbb{E}^{\mu_+}(X) < \infty$ and $\mathbb{E}^{\nu_-}(X) > -\infty$.

ii) If among $\mathbb{E}^{\mu_+}(X)$ and $\mathbb{E}^{\nu_-}(X)$ exactly one is finite, then (Z_n) is null recurrent.

iii) If $\int_0^{2\pi} |1 - \hat{\mu}_+(t)|^{-2} dt < \infty$ and $\int_0^{2\pi} |1 - \hat{\nu}_-(t)|^{-2} dt < \infty$, then (Z_n) , hence (S_n) , is transient.

Proof of the proposition :

Notice that (S_n) also has to be irreducible. For iii), Proposition 4.4 gives that $(G^{\mu_+}(0, m))_{m \geq 0}$ and $(G^{\nu_-}(0, -m))_{m \geq 0}$ are in l^2 , so $(G^{\mu_+}(0, m)G^{\nu_-}(0, -m))_{m \geq 0} \in l^1$, by the Cauchy-Schwarz inequality. Using Proposition 4.4 i), this proves transience.

If $\mathbb{E}^{\mu_+}(X) < \infty$, then $\lim_{m \rightarrow +\infty} G^{\mu_+}(0, m) = 1/\mathbb{E}^{\mu_+}(X) > 0$, by the renewal theorem. By Proposition 4.4, (Z_n) is recurrent iff $\sum_{m \geq 0} G^{\nu_-}(0, -m) = +\infty$, which is true. The case $\mathbb{E}^{\nu_-}(X) > -\infty$ is treated similarly.

To discuss the positivity/nullity of the recurrence of (Z_n) , define $V_n = \sum_{0 \leq k \leq n} \mu_+^k \nu_-^{n-k}$, $n \geq 0$. By Lemma 4.3, $\forall l \in \mathbb{Z}$, $V_n(l) = P_{-l}(Z_n = 0) \leq 1$. This also follows from $\bar{V}_n(l) = P(l \in \{S_k^{\mu_+} + S_{n-k}^{\nu_-}, 0 \leq k \leq n\})$, as $k \mapsto S_k^{\mu_+} + S_{n-k}^{\nu_-}$ is a.s. increasing. As (Z_n) is irreducible and aperiodic, $P_x(Z_n = y)$ converges for all x and y , as $n \rightarrow +\infty$; see for instance Woess [12], Theorem 3.48. We set $w(l) = \lim_{n \rightarrow +\infty} V_n(l)$, $l \in \mathbb{Z}$. From the relations $V_n = \nu_-^n + \mu_+ V_{n-1} = \mu_+^n + \nu_- V_{n-1}$ and dominated convergence, we get :

$$\forall y \in \mathbb{Z}, w(y) = \sum_{k \geq 1} \mu_+(k)w(y-k) = \sum_{k \leq -1} \nu_-(k)w(y-k).$$

Hence w is a bounded harmonic function, for both μ_+ and ν_- . By Woess [12], relation (8.3), in the proof of Theorem 8.2, if $k \in \text{Supp}(\mu_+) \cup \text{Supp}(\nu_-)$, then $\forall y \in \mathbb{Z}$, $w(y) = w(y-k)$. From the hypothesis on the supports, w equals some constant $c \in [0, 1]$.

Next, from Lemma 4.3 that for $n \geq 1$ and $y \geq 1$, with obvious notations :

$$P_x(Z_n = y) = \nu_-^n(y-x) + (V_{n-1}\mu_{+, \geq y})(y-x), \quad (13)$$

as well as $P_x(Z_n = y) = \mu_+^n(y-x) + (V_{n-1}\nu_{-, \leq y-1})(y-x)$, when $y \leq 0$. Taking now $n \geq 1$ and writing $1 = \sum_{y \in \mathbb{Z}} P_0(Z_n = y)$, we obtain :

$$\begin{aligned} 1 &= \sum_{y \geq 0} \sum_{l \in \mathbb{Z}} V_{n-1}(l) \mu_{+, \geq y}(y-l) + \sum_{y \leq -1} \sum_{l \in \mathbb{Z}} V_{n-1}(l) \nu_{-, \leq y-1}(y-l) \\ &= \sum_{l \in \mathbb{Z}} V_{n-1}(l) \sum_{y \geq 0} \mu_{+, \geq y}(y-l) + \sum_{l \in \mathbb{Z}} V_{n-1}(l) \sum_{y \leq -1} \nu_{-, \leq y-1}(y-l) \\ &= \sum_{l \leq 0} V_{n-1}(l) \mu_+[-l, +\infty) + \sum_{l \geq 1} V_{n-1}(l) \nu_-(-\infty, -l-1]. \end{aligned} \quad (14)$$

Hence, for any $L > 0$, we get $1 \geq c \sum_{-L \leq l \leq 0} \mu_+[-l, +\infty) + c \sum_{1 \leq l \leq L} \nu_-(-\infty, -l-1]$. If either $\mathbb{E}^{\mu_+}(X) = \infty$ or $\mathbb{E}^{\nu_-}(X) = -\infty$, then $c = 0$. By (13), this gives $P_x(Z_n = y) \rightarrow 0$, for any x and y . Therefore (Z_n) is either transient or null recurrent. This proves *ii*), as recurrence already holds if one of the expectations is finite.

If (Z_n) is positive recurrent, then $c > 0$ and $\sum_{-L \leq l \leq 0} \mu_+[-l, +\infty) + c \sum_{1 \leq l \leq L} \nu_-(-\infty, -l-1] \leq 1/c$, for any $L > 0$, giving $\mathbb{E}^{\mu_+}(X) < \infty$ and $\mathbb{E}^{\nu_-}(X) > -\infty$. Reciprocally, if $\mathbb{E}^{\mu_+}(X) < \infty$ and $\mathbb{E}^{\nu_-}(X) > -\infty$, then, by dominated convergence in (14) :

$$1 = c \sum_{l \leq 0} \mu_+[-l, +\infty) + c \sum_{l \geq 1} \nu_-(-\infty, -l-1],$$

giving $c > 0$. As $c = \lim_{n \rightarrow +\infty} P_y(Z_n = 0)$, for any y , we obtain that (Z_n) in this case is positive recurrent. This shows point *i*) of the proposition. \square

Remark. — When $\mu_+ = \delta_1$ and $\nu_- = \delta_{-1}$, then (Z_n) is not aperiodic (nor irreducible on \mathbb{Z}) and the period is two. One has $V_n = \sum_{k=0}^n \delta_{2k-n}$, which does not converge pointwise. Conditions of aperiodicity and irreducibility are discussed in [7], Lemmas 4.1 and 4.3.

Remark. — To complete the study, the remaining case is $\mathbb{E}^{\mu_+}(X) = \mathbb{E}^{\nu_-}(-X) = +\infty$, with $|1 - \hat{\mu}_+(t)|^{-1}$ or $|1 - \hat{\nu}_-(t)|^{-1}$ not in $L^2(0, 2\pi)$. In this case, by Proposition 3.2, $Re((1 - \hat{\mu}_+)^{-1})$

and $Re((1 - \hat{\nu}_-)^{-1})$ have non-negative real cosine coefficients. A related question of harmonic analysis is then the following one. Let real functions $f > 0$ and $g > 0$ on $(0, 2\pi)$ be L^1 , even and with real Fourier coefficients $\hat{f}(n) \geq 0$, $\hat{g}(n) \geq 0$, $n \geq 0$. Are $\int_0^{2\pi} f(t)g(t)dt$ and $\sum_{n \geq 0} \hat{f}(n)\hat{g}(n)$ infinite simultaneously ? One direction is clear. Introducing the Fejer kernel K_n , we have :

$$(K_n * f)(t) = \sum_{j=-n}^n \left(1 - \frac{|j|}{n+1}\right) \hat{f}(j)e^{ijt}.$$

Then, for $M > 0$, $\frac{1}{2\pi} \int_0^{2\pi} (K_n * f)(g \wedge M)dt \leq \frac{1}{2\pi} \int_0^{2\pi} (K_n * f)gdt = \sum_{j=-n}^n (1 - \frac{|j|}{n+1}) \hat{f}(j)\hat{g}(j)$. Using that $K_n * f \rightarrow f$ in L^1 for the term on the left and monotone convergence for the last sum, we obtain $\frac{1}{2\pi} \int_0^{2\pi} f(g \wedge M)dt \leq \sum_{n \geq 0} \hat{f}(n)\hat{g}(n)$. Letting $M \rightarrow +\infty$, we get one inequality.

References

- [1] J. BRÉMONT, *On planar random walks in environments invariant by horizontal translations*, Markov Processes and Related Fields 2016, issue 2, vol. 22, 267-310.
- [2] J. BRÉMONT, *Markov chains in a stratified environment*, ALEA, Lat. Am. J. Probab. Math. Stat. 14, 751-798 (2017).
- [3] M. CAMPANINO, D. PETRITIS, *Random walks on randomly oriented lattices*, Markov Processes and Related Fields 9 (2003), no. 3, 391-412.
- [4] H. CARTAN, *Théorie élémentaire des fonctions analytiques d'une ou plusieurs variables complexes*, Hermann, Paris, 1961, 232 pages.
- [5] P. L. DUREN, *Theory of H^p spaces*. New York ; London : Academic Press, 1970.
- [6] P. ERDÖS, W. FELLER AND H. POLLARD, *A property of power series with positive coefficients*. Bull. Amer. Math. Soc. 55 (2) : 201-204 (February 1949).
- [7] J. H. B. KEMPERMAN, *The oscillating random walk*. Stochastic Processes and their Applications. Volume 2, Issue 1, January 1974, 1-29.
- [8] H. KESTEN, F. SPITZER, *Random walks on countably infinite abelian groups*. Acta Math. 114: 237-265 (1965).
- [9] D. ORNSTEIN, *Random walks I and II*. Trans. Amer. Math. Soc. 138, 1969, 1-43; 45-60.
- [10] S. C. PORT AND C. J. STONE, *Potential theory of random walks on abelian groups*. Acta Math. 122, 1969, 19-114.
- [11] F. SPITZER, *Principles of random walks*. Second edition. Graduate Texts in Mathematics, vol. 34. Springer-Verlag, New-York Heidelberg, 1976.
- [12] W. WOESS, *Denumerable Markov chains*. EMS Textbooks in Mathematics, 2009. 351 pages.

UNIV PARIS EST CRETEIL, CNRS, LAMA, F-94010 CRETEIL, FRANCE
 UNIV GUSTAVE EIFFEL, LAMA, F-77447 MARNE-LA-VALLÉE, FRANCE
E-mail address : julien.bremont@u-pec.fr