

Covariant derivatives for Ehresmann connections

G.E. Prince and D.J. Saunders

October 15, 2021

Abstract. We deal with the construction of covariant derivatives for some quite general Ehresmann connections on fibre bundles. We show how the introduction of a vertical endomorphism allows construction of covariant derivatives separately on both the vertical and horizontal distributions of the connection which can then be glued together on the total space.

Subject Classification (2020): Primary 53C05, 53B05 Secondary 34A26

Keywords: Ehresmann connections, covariant derivatives, torsion, linear connections, Hopf bundle, sprays

1 Introduction

In this short paper we address the question of the existence of covariant derivatives for Ehresmann, or ‘nonlinear’, connections [2]. We will show how quite simple additional requirements can yield the familiar covariant derivative of the linear case.

We shall take as our definition of an Ehresmann connection the minimalist version, using only a horizontal distribution on the tangent bundle of a fibre bundle complementary to the vertical sub-bundle: although other authors sometimes impose additional conditions involving completeness ([3], see also [4]), we shall not need them.

First of all we give a construction of a covariant derivative on a manifold from covariant derivatives on vector submodules or distributions. This has very recently been presented in [5]. Then we show how to construct covariant derivatives on pairs of distributions using certain endomorphisms. In the context of Ehresmann connections on a fibre bundle $\pi : E \rightarrow B$, we then show how to construct covariant derivatives on the corresponding vertical and horizontal distributions in quite general cases, not necessarily principal or vector bundles, glueing them together to give a covariant derivative on E . Finally, we present a range of examples and applications from the Hopf bundle to second order ordinary differential equations.

We remark that a method of constructing a covariant derivative from an Ehresmann connection on a general fibred manifold was given in [1], but in that approach the operator was defined only on projectable vector fields, mapping them to vertical vector fields.

2 The construction of covariant derivatives I

In proposition 2.1 and theorem 2.2 we present an apparently new construction of a covariant derivative on a manifold from covariant derivatives on vector submodules or distributions [5].

We begin by demonstrating, as a first step, an extension of these submodule derivatives. In what follows we use submodule and distribution interchangeably even though they are distinct constructs.

Let \mathcal{M} be a manifold and P a projector onto a given submodule of vector fields (alternatively, distribution). Suppose that $\text{im}(P)$ has a covariant derivative ∇^P , that is $\nabla_X^P Y$ has the usual properties for $X, Y \in \text{im}(P)$ over the smooth functions, $\mathfrak{F}(\mathcal{M})$, on \mathcal{M} , namely \mathbb{R} -linearity in both arguments and

$$\nabla_{fX}^P Y = f\nabla_X^P Y \text{ and } \nabla_X^P(fY) = X(f)Y + f\nabla_X^P Y \quad (1)$$

(it's not required that $\nabla_X^P Y \in \text{im}(P)$). ∇^P can be partially extended to the module of vector fields $\mathfrak{X}(\mathcal{M})$ as follows:

Proposition 2.1. Define $\bar{\nabla}_X^P Y$ for $X \in \mathfrak{X}(\mathcal{M}), Y \in \text{im}(P)$ over $\mathfrak{F}(\mathcal{M})$ by

$$\bar{\nabla}_X^P Y := \nabla_{P(X)}^P Y + P([X - P(X), Y]). \quad (2)$$

Then $\bar{\nabla}_X^P Y$ has the covariant derivative properties.

Proof. The \mathbb{R} -linearity properties are obvious.

$$\begin{aligned} \bar{\nabla}_{fX}^P Y &= \nabla_{P(fX)}^P Y + P([fX - P(fX), Y]) \\ &= f(\nabla_{P(X)}^P Y + P([X - P(X), Y])) - Y(f)P(X - P(X)) \\ &= f\bar{\nabla}_X^P Y \quad \text{since } P^2 = P. \end{aligned}$$

$$\begin{aligned} \bar{\nabla}_X^P(fY) &= \nabla_{P(X)}^P(fY) + P([X - P(X), fY]) \\ &= P(X)(f)Y + f\nabla_{P(X)}^P Y + fP([X - P(X), Y]) + (X - P(X))(f)P(Y) \\ &= X(f)Y + f\bar{\nabla}_X^P Y \quad \text{since } Y \in \text{im}(P). \end{aligned}$$

□

Now suppose that $T\mathcal{M}$ has a direct sum decomposition into N submodules (of possibly different dimensions) with corresponding unique projectors $P_A, A = 1, \dots, N$. If each submodule has a covariant derivative ∇^A with extension $\bar{\nabla}^A$ as defined in proposition 2.1 then

Theorem 2.2. For $X, Y \in \mathfrak{X}(\mathcal{M})$

$$\nabla_X Y := \sum_{A=1}^N \bar{\nabla}_X^A(P_A(Y))$$

has the covariant derivative properties over $\mathfrak{F}(\mathcal{M})$

Proof. Again, \mathbb{R} -linearity is obvious.

$$\begin{aligned} \nabla_{fX} Y &:= \sum_{A=1}^N \bar{\nabla}_{fX}^A(P_A(Y)) \\ &= \sum_{A=1}^N f\bar{\nabla}_X^A(P_A(Y)) = f\nabla_X Y. \end{aligned}$$

$$\begin{aligned}
\nabla_X(fY) &:= \sum_{A=1}^N \bar{\nabla}_X^A(P_A(fY)) \\
&= \sum_{A=1}^N (f\bar{\nabla}_X^A(P_A(Y)) + X(f)P_A(Y)) \\
&= f\nabla_X Y + X(f)Y \quad \text{since } \sum_{A=1}^N P_A = I_{T\mathcal{M}}.
\end{aligned}$$

□

The following corollary will be useful in the direct sum situation of theorem 2.2.

Corollary 2.3. $\nabla P_B = 0$ for all $B \in \{1, \dots, N\}$ if and only if $\nabla_X^B Y \in \text{im}(P_B)$ for all $X, Y \in \text{im}(P_B)$ and for each $B \in \{1, \dots, N\}$.

Proof.

$$\begin{aligned}
(\nabla_X P_B)(Y) &= \nabla_X(P_B(Y)) - P_B(\nabla_X Y) \\
&= \sum_{A=1}^N \bar{\nabla}_X^A(P_A(P_B(Y))) - P_B\left(\sum_{A=1}^N \bar{\nabla}_X^A(P_A(Y))\right) \\
&= \bar{\nabla}_X^B(P_B(Y)) - P_B\left(\sum_{A=1}^N \left(\nabla_{P_A(X)}^A(P_A(Y)) + P_A([X - P_A(X), P_A(Y)])\right)\right) \\
&= \bar{\nabla}_X^B(P_B(Y)) - P_B\left(\sum_{A=1}^N \nabla_{P_A(X)}^A(P_A(Y))\right) - P_B([X - P_B(X), P_B(Y)]) \\
&= \nabla_{P_B(X)}^B(P_B(Y)) - P_B\left(\sum_{A=1}^N \nabla_{P_A(X)}^A(P_A(Y))\right)
\end{aligned}$$

So, if $\nabla_X^A Y \in \text{im}(P_A)$ for all $X, Y \in \text{im}(P_A)$ then the right-hand side is zero. Conversely, if $(\nabla_X P_B)(Y) = 0$ for all B, X, Y then for fixed C and for $X, Y \in \text{im}(P_C)$ we have for all $B \neq C$

$$0 = -P_B(\nabla_{P_C(X)}^C(P_C(Y))) = -P_B(\nabla_X^C Y).$$

Hence $\nabla_X^C Y \in \text{im}(P_C)$ for all $X, Y \in \text{im}(P_C)$, recalling that $I_{T\mathcal{M}} = \sum_{A=1}^N P_A$.

□

Proposition 2.1 and theorem 2.2 provide the building blocks for the covariant derivatives we will construct in the remaining sections.

3 The construction of covariant derivatives II

Now suppose that we have a pair of disjoint distributions V, H of equal constant rank on \mathcal{M} . (We don't assume that $T\mathcal{M} = V \oplus H$.) Let \bar{V}, \bar{H} denote fixed complements of V, H in $T\mathcal{M}$, defining projectors P_V, P_H . Now assume the existence of a *vertical endomorphism*, a vertical-valued 1-form S on $T\mathcal{M}$ with $\ker(S) = \bar{H}, \text{im}(S) = V$. This determines a unique horizontal-valued 1-form (*horizontal endomorphism*) Q with $\ker(Q) = \bar{V}, \text{im}(Q) = H$ and $Q \circ S = P_H$ and $S \circ Q = P_V$.

Theorem 3.1. *The derivations ∇^V, ∇^H on V and H respectively defined by*

$$\nabla_X^V Y := S([X, Q(Y)]), \quad X, Y \in V; \quad \nabla_X^H Y := Q([X, S(Y)]), \quad X, Y \in H$$

have the covariant derivative properties over $\mathfrak{F}(M)$.

Proof. It suffices to show that ∇^V has the properties. The \mathbb{R} -linearity property of $\nabla_X^V Y$ is obvious. For $X, Y \in V$ and $f \in \mathfrak{F}(M)$

$$\begin{aligned} \nabla_{fX}^V Y &= S([fX, Q(Y)]) = S(f[X, Q(Y)] - Q(Y)(f)X) \\ &= fS([X, Q(Y)]) \text{ since } \ker(S) = \bar{H} \\ &= f\nabla_X^V Y \end{aligned}$$

and

$$\begin{aligned} \nabla_X^V (fY) &= S([X, fQ(Y)]) = S(X(f)Q(Y) + f[X, Q(Y)]) \\ &= X(f)Y + fS([X, Q(Y)]) \text{ since } P_V = S \circ Q \\ &= X(f)Y + \nabla_X^V Y \end{aligned}$$

□

One generalisation of this result that will be useful in the case of Ehresmann connections is as follows.

Suppose M has disjoint distributions H_1, \dots, H_N and V of equal constant rank. Let $\bar{V}, \bar{H}_A, A = 1, \dots, N$, denote fixed complements of V, H_A in $T\mathcal{M}$ and denote the corresponding projectors by P_V, P_A .

Now assume the existence of N vertical endomorphisms, S_1, \dots, S_N , on $T\mathcal{M}$ with $\ker(S_A) = \bar{H}_A, \text{im}(S_A) = V$. These determine M unique horizontal endomorphisms Q_A with $\ker(Q_A) = \bar{V}, \text{im}(Q_A) = H_A$ and $Q_A \circ S_A = P_A$ and $S_A \circ Q_A = P_V$; notice that $S_A \circ Q_B = 0, A \neq B$. Further, set $S := \frac{1}{\sqrt{N}} \sum_A S_A$ and $Q := \frac{1}{\sqrt{N}} \sum_A Q_A$ and note that $S \circ Q = P_V$.

Proposition 3.2. *The derivations ∇^V, ∇^A on V and H_A respectively defined by*

$$\begin{aligned} \nabla_X^V Y &:= S([X, Q(Y)]), \quad X, Y \in V \\ \nabla_X^A Y &:= Q_A([X, S_A(Y)]), \quad X, Y \in H_A, \end{aligned}$$

have the covariant derivative properties over $\mathfrak{F}(M)$.

Note that $\nabla_X^V Y := S_A([X, Q_A(Y)])$ for any A also has the desired property but is less canonical.

4 Special Ehresmann Connections

Suppose that $\pi : E \rightarrow B$ is a fibred manifold with an Ehresmann connection: V is the vertical sub-bundle of TE comprised of tangent spaces to the fibres, F , and H is the horizontal sub-bundle of TE complementary to V . Remembering that pointwise TE is a direct sum of V and H , if both V and H have covariant derivatives ∇^V, ∇^H , a covariant derivative on E can be

constructed using theorem 2.2. Of course this begs the question of the construction of these separate derivatives; in geometric cases V will have a covariant derivative by virtue of the geometry of F but for H this requires additional structure.

The significant question is whether such covariant derivatives on the total space E are restricted to the vector bundle case. We will give examples in the next section.

If $\dim(V) = \dim(H)$, that is, the dimensions of the tangent spaces to F and B coincide, for example when $E = TB$, the existence of a vertical endomorphism is sufficient to construct a covariant derivative (see theorem 4.1 below). In the case where $\dim(V) < \dim(H)$ a vertical endomorphism can still be used if $H = H_1 \oplus H_2$ with $\dim(H_1) = \dim(V)$ and H_2 is equipped with a covariant derivative. While this sounds a little artificial, it is exactly the case in the geometry of time-dependent second order ordinary differential equations. Indeed, in that case the vertical endomorphism is central and V, H_1 and H_2 naturally appear as eigenspaces of a related endomorphism, see [5, 6]. We treat the simpler, time-independent case in example 4.

The following result follows directly from proposition 2.1 and theorems 2.2, 3.1.

Theorem 4.1. *Let $\pi : E \rightarrow B$ be a fibred manifold with fibres F and base B having the same constant dimension. Assume an Ehresmann connection on E with vertical and horizontal distributions V, H and corresponding projectors P_V, P_H . Further suppose that S and Q are vertical and horizontal endomorphisms (so that $\text{im}(S) = V = \text{Ker}(S)$ and $\text{im}(Q) = H = \text{ker}(Q)$.) Then E has covariant derivative*

$$\begin{aligned} \nabla_X Y &:= \bar{\nabla}_X^V(P_V(Y)) + \bar{\nabla}_X^H(P_H(Y)) \\ &:= S([P_V(X), Q(Y)]) + P_V([P_H(X), P_V(Y)]) \\ &\quad + Q([P_H(X), S(Y)]) + P_H([P_V(X), P_H(Y)]). \end{aligned} \tag{3}$$

Recalling that the curvature of an Ehresmann connection is $R(X, Y) = P_V([P_H(X), P_H(Y)])$ it is straightforward to see that the torsion of this covariant derivative also carries this information: $P_V \circ T(P_H(X), P_H(Y)) = R(X, Y)$; this will appear again in the examples.

Another straightforward case is when $\dim(H)$ is an integer multiple of $\dim(V)$.

Theorem 4.2. *Let $\pi : E \rightarrow B$ be a fibred manifold with fibres F and base B , both of constant dimension and with $\dim(B)$ being an integer multiple, N , of $\dim(F)$. Assume an Ehresmann connection on E with vertical subbundle V and horizontal subbundle $H = H_1 \oplus \dots \oplus H_N$ with $\dim(H_A) = \dim(V)$ for all A . The corresponding projectors, vertical and horizontal endomorphisms are P_V, H_A, S_A, Q_A , $S := \frac{1}{\sqrt{N}} \sum_A S_A$, $Q := \frac{1}{\sqrt{N}} \sum_A Q_A$. Then E has covariant derivative*

$$\begin{aligned} \nabla_X Y &:= \bar{\nabla}_X^V(P_V(Y)) + \sum_A \bar{\nabla}_X^A(P_A(Y)) \\ &:= S([P_V(X), Q(Y)]) + \sum_A P_V([P_A(X), P_V(Y)]) \\ &\quad + \sum_A (Q_A([P_A(X), S_A(Y)]) + P_A([X - P_A(X), P_A(Y)])) . \end{aligned} \tag{4}$$

The proof follows from proposition 2.1, theorem 2.2 and proposition 3.2.

As a result of corollary 2.3 the covariant derivatives of theorems 4.1 and 4.2 have the properties $\nabla P = 0$ for all the projectors. In addition, in the case of the covariant derivative of theorem 4.1 $\nabla S = 0 = \nabla Q$. This is not so in the case of theorem 4.2.

4.1 Applications and Examples

Example 1. Take $B = \mathbb{R}^2$ with coordinates x, y and $F = S^1$ with co-ordinate θ on \mathbb{R} and vertical distribution $V(E) = \text{Sp}\{V := \frac{\partial}{\partial\theta}\}$. Define the horizontal lift through $(x_0, y_0, \theta_0) \in E$ of the coordinate curve $\gamma(s) := (x_0 + s, y_0)$ through $(x_0, y_0) \in B$ to be $\gamma^h(s) := (x_0 + s, y_0, \theta_0 \exp(s))$. Do the same for the y co-ordinate curves. This generates the horizontal lifts $(\frac{\partial}{\partial x})^H := \frac{\partial}{\partial x} + \theta \frac{\partial}{\partial\theta}$ and $(\frac{\partial}{\partial y})^H := \frac{\partial}{\partial y} + \theta \frac{\partial}{\partial\theta}$, so we have a horizontal distribution

$$H(E) = \text{Sp} \left\{ H_1 := \frac{\partial}{\partial x} + \theta \frac{\partial}{\partial\theta}, H_2 := \frac{\partial}{\partial y} + \theta \frac{\partial}{\partial\theta} \right\}.$$

It is straightforward to show that H satisfies the conditions for an Ehresmann connection.

The vertical and horizontal endomorphisms of theorem 4.2 are $S_1 := dx \otimes V, S_2 := dy \otimes V$ and $Q_1 := \psi \otimes H_1, Q_2 := \psi \otimes H_2$ with $\psi := d\theta - \theta(dx + dy)$. The components of ∇ are

$$\nabla_V V = 0, \nabla_V H_A = 0, \nabla_{H_A} H_A = -H_A, \nabla_{H_A} H_B = 0, \nabla_{H_A} V = -V.$$

Example 2. Affine connection with torsion

Consider an affine connection with torsion on a manifold M^n , with connection coefficients Γ_{bc}^a in local coordinates x^a . This induces an n -dimensional horizontal distribution on the tangent bundle TM (with local adapted coordinates (x^a, u^a)) spanned by

$$H_a := \frac{\partial}{\partial x^a} - \Gamma_{ab}^c u^b \frac{\partial}{\partial u^c}.$$

With vertical distribution $\text{Sp}\{V_c := \frac{\partial}{\partial u^c}\}$, the basis dual to $\{H_a, V_b\}$ is

$$\{dx^a, \phi^b := du^b + \Gamma_{cd}^b u^d dx^c\}.$$

The non-zero bracket relations are

$$[H_a, V_b] = \Gamma_{ab}^c V_c, [H_a, H_b] = \left[\left(\frac{\partial \Gamma_{ad}^c}{\partial x^b} - \frac{\partial \Gamma_{bd}^c}{\partial x^a} \right) + (\Gamma_{ad}^e \Gamma_{be}^c - \Gamma_{bd}^e \Gamma_{ae}^c) \right] u^d V_c$$

Notice that $[H_a, H_b] = -R_{dab}^c u^d V_c$ where R_{dab}^c are the components of the Riemann curvature of the connection on M^n .

It is straightforward to verify that the (1,1) tensors

$$S = dx^a \otimes V_a \text{ and } Q = \phi^a \otimes H_a$$

are intrinsic (tensorial). Using these in the definitions of ∇^V, ∇^H from theorem 3.1, we find

$$\nabla_{V_a}^V V_b = 0, \nabla_{H_a}^H H_b = \Gamma_{ab}^c H_c.$$

Constructing the covariant derivative ∇ of theorems 2.2 and 4.1 gives

$$\nabla_{V_a} V_b = 0, \nabla_{V_a} H_b = 0, \nabla_{H_a} V_b = \Gamma_{ab}^c V_c, \nabla_{H_a} H_b = \Gamma_{ab}^c H_c.$$

A straightforward calculation gives the only non-zero torsion components as

$$T(H_a, H_b) = (\Gamma_{ab}^c - \Gamma_{ba}^c) H_c - \left[\left(\frac{\partial \Gamma_{ad}^c}{\partial x^b} - \frac{\partial \Gamma_{bd}^c}{\partial x^a} \right) + (\Gamma_{ad}^e \Gamma_{be}^c - \Gamma_{bd}^e \Gamma_{ae}^c) \right] u^d V_c$$

which capture, as horizontal and vertical lifts respectively, the torsion and curvature of the original affine connection on M . Note also that the Ehresmann curvature is the vertical part of the torsion (up to a sign).

Example 3. Nonlinear connection on TM

The case where $E = TM$ is a standard one and we apply 4.1 directly. As in the example above, TM has coordinates (x^a, u^b) . Put $V = \text{Sp}\{V_a := \frac{\partial}{\partial u^a}\}$ and $H = \text{Sp}\{H_a := \frac{\partial}{\partial x^a} - \Gamma_a^b \frac{\partial}{\partial u^b}\}$ so that $S = dx^a \otimes V_a$ and $Q = \phi^a \otimes H_a$, where $\{dx^a, \phi^b := du^b + \Gamma_c^b dx^c\}$ is dual to $\{H_a, V_b\}$. The results appear similar to those in the affine connection case with Γ_b^a replacing $\Gamma_{bc}^a u^c$. We get

$$\nabla_{V_a}^V V_b = S([V_a, H_b]) = 0, \quad \nabla_{H_a}^H H_b = Q([H_a, V_b]) = V_b(\Gamma_a^c)H_c.$$

and

$$\nabla_{V_a} V_b = 0, \quad \nabla_{V_a} H_b = 0, \quad \nabla_{H_a} V_b = V_b(\Gamma_a^c)V_c, \quad \nabla_{H_a} H_b = V_b(\Gamma_a^c)H_c.$$

More interesting is that the torsion acts non-trivially only on H with

$$T(H_a, H_b) = Q([H_a, V_b] - [H_b, V_a]) - [H_a, H_b];$$

this means that the horizontal component is

$$P_H(T(H_a, H_b)) = (V_b(\Gamma_a^c) - V_a(\Gamma_b^c))H_c$$

and the vertical component is the Ehresmann curvature $R(H_a, H_b)$. This horizontal component is zero if and only if there are a local functions f^c such that $\Gamma_a^c = V_a(-\frac{1}{2}f^c)$. The reason for the numerical coefficient, and a global explanation of the phenomenon, may be seen in the next Example.

Example 4. SODE connection on TM

Again we take $E := TM$ with standard vertical endomorphism $S = dx^a \otimes V_a$. A second order differential equation field (SODE) is a vector field Γ on TM satisfying $S(\Gamma) = \Delta$, where Δ is the canonical dilation field on the vector bundle $\tau_M : TM \rightarrow M$. The reason for the name is that such a vector field defines a family of curves on M satisfying second order equations of the form $\ddot{x}^a = f^a(x, \dot{x})$.

Every SODE gives rise to a nonlinear connection on TM with horizontal projector $P_H = \frac{1}{2}(I - \mathcal{L}_\Gamma S)$; this is called the SODE connection of Γ . In coordinates, if

$$\Gamma = u^a \frac{\partial}{\partial x^a} + f^a \frac{\partial}{\partial u^a}$$

then

$$P_H = dx^a \otimes \left(\frac{\partial}{\partial x^a} - \Upsilon_a^b \frac{\partial}{\partial u^b} \right), \quad \Upsilon_a^b = -\frac{1}{2} \frac{\partial f^b}{\partial u^a}.$$

We might therefore ask when a general nonlinear connection on TM was derived in this way from a SODE, and we can use theorem 4.1 as in the previous example. So suppose such a nonlinear connection has horizontal vector fields

$$H_a = \frac{\partial}{\partial x^a} - \Gamma_a^b \frac{\partial}{\partial u^b}.$$

Regard the connection as a splitting of the short exact sequence of vector bundles over TM

$$0 \longrightarrow V\tau_M \longrightarrow TTM \xrightarrow{\quad h \quad} \tau_M^* TM \longrightarrow 0$$

where, if $v, w \in T_p M$, the splitting map h gives the horizontal lift of v to $h(v, w) \in T_w T M$, so that

$$v = v^a \frac{\partial}{\partial x^a} \Big|_p, \quad h(v, w) = v^a H_a|_w = v^a \left(\frac{\partial}{\partial x^a} \Big|_w - \Gamma_a^b(w) \frac{\partial}{\partial u^b} \Big|_w \right).$$

Define a vector field Γ on $T M$ by $\Gamma_v = h(v, v) \in T_v T M$, so that

$$\Gamma_v = v^a \left(\frac{\partial}{\partial x^a} \Big|_v - \Gamma_a^b(v) \frac{\partial}{\partial u^b} \Big|_v \right) = u^a(v) \left(\frac{\partial}{\partial x^a} \Big|_v - \Gamma_a^b(v) \frac{\partial}{\partial u^b} \Big|_v \right)$$

and therefore that

$$\Gamma = u^a H_a = u^a \frac{\partial}{\partial x^a} - u^a \Gamma_a^b \frac{\partial}{\partial u^b}.$$

We see that Γ is a SODE with $f^b := -u^a \Gamma_a^b$, giving rise to a SODE connection. Put

$$\Upsilon_c^b = -\frac{1}{2} \frac{\partial f^b}{\partial u^c} = \frac{1}{2} \frac{\partial}{\partial u^c} (u^a \Gamma_a^b) = \frac{1}{2} (\Gamma_c^b + u^a V_c(\Gamma_a^b)).$$

Now suppose that the original nonlinear connection is ‘symmetric’ (that is, that the torsion of ∇ is vertical, as in the previous example) and that the connection coefficients Γ_a^b are homogeneous (in other words that $\Delta(\Gamma_a^b) = \Gamma_a^b$, where $\Delta = u^c V_c$). We see that in these circumstances

$$u^a V_c(\Gamma_a^b) = u^a V_a(\Gamma_c^b) = \Delta(\Gamma_c^b) = \Gamma_c^b$$

so that $\Gamma_c^b = \Upsilon_c^b$ and the two connections are identical. Note, incidentally, that

$$\Delta(f^b) = -u^a \Gamma_a^b - u^a \Delta(\Gamma_a^b)$$

so that Γ is a spray, with $\Delta(f^b) = 2f^b$, precisely when the connection coefficients are homogeneous.

Now return to the affine connection on $T M$. We find that

$$\nabla_{H_a} \Gamma = (\Delta(\Gamma_a^b) - \Gamma_a^b) H_b, \quad \nabla_{V_a} \Gamma = H_a$$

so that the connection coefficients are homogeneous precisely when $\nabla_{X^H} \Gamma = 0$ for any vector field X on M , and similarly

$$\nabla_{H_a} \Delta = (\Delta(\Gamma_a^b) - \Gamma_a^b) V_b, \quad \nabla_{V_a} \Delta = \nabla_{V_a} (u^b V_b) = V_a$$

so that the connection coefficients are also homogeneous precisely when $\nabla_{X^H} \Delta = 0$ for any vector field X on M . Thus

$$\nabla_{X^H} \Delta = 0, \quad P_H \circ T = 0$$

is a sufficient condition for the nonlinear connection to be a SODE connection, and it will then be the SODE connection of a spray.

Example 5. The Hopf bundle

One application of theorem 4.2 is to the Hopf bundle $\pi : S^3 \rightarrow S^2$. If we regard S^3 as the submanifold of \mathbb{R}^4 given by $x^2 + y^2 + z^2 + w^2 = 1$ then one choice for the definition of π is given by

$$\pi(x, y, z, w) = (x^2 + y^2 - z^2 - w^2, 2(xw + yz), 2(yw - xz)) \in S^2 \subset \mathbb{R}^3.$$

Now consider the infinitesimal rotations in the six coordinate 2-planes of \mathbb{R}^4 ,

$$\begin{aligned}\theta &= y\frac{\partial}{\partial x} - x\frac{\partial}{\partial y}, & \phi &= z\frac{\partial}{\partial x} - x\frac{\partial}{\partial z}, & \psi &= w\frac{\partial}{\partial x} - x\frac{\partial}{\partial w}, \\ \xi &= w\frac{\partial}{\partial z} - z\frac{\partial}{\partial w}, & \eta &= y\frac{\partial}{\partial w} - w\frac{\partial}{\partial y}, & \zeta &= z\frac{\partial}{\partial y} - y\frac{\partial}{\partial z}\end{aligned}$$

so that these restrict to vector fields tangent to S^3 , and put

$$V = \theta - \xi, \quad \Lambda = \phi - \eta, \quad \Sigma = \psi - \zeta$$

as vector fields on S^3 . We may check that (Λ, Σ, V) form a global frame for S^3 defining an absolute parallelism, and that V is vertical with respect to π , so that (Λ, Σ) span the horizontal sub-bundle of an Ehresmann connection on the Hopf bundle with

$$[\Sigma, \Lambda] = 2V, \quad [\Lambda, V] = 2\Sigma, \quad [V, \Sigma] = 2\Lambda.$$

Writing $H_1 = \langle \Lambda \rangle$ and $H_2 = \langle \Sigma \rangle$, we can apply theorem 4.2 to obtain a covariant derivative ∇ on S^3 . It is straightforward to check that if $X, Y \in \{\Lambda, \Sigma, V\}$ then $\nabla_X Y = 0$. This linear connection is not symmetric, but if we subtract half the torsion to obtain a symmetric connection with the same geodesics then the result is just the Levi-Civita connection of the standard metric on S^3 induced by its embedding in \mathbb{R}^4 .

References

- [1] M. Crampin and L.A. Ibort. Graded Lie algebras of derivations and Ehresmann connections, *J. Math. Pures et Appl.* **66** (1987) 113–125.
- [2] C. Ehresmann. Les connexions infinitésimales dans un espace fibré différentiable, *Colloque de Topologie, Bruxelles* (1950) 29–55.
- [3] W. Greub, S. Halperin, R. Vanstone. *Connections, curvature and cohomology*, Academic Press 1972.
- [4] I. Kolář, P. Michor and J. Slovák. *Natural operators in differential geometry*, Springer-Verlag 1993.
- [5] G.E. Prince, M. Farré Puiggalí, D.J. Saunders, D. Martín de Diego. Linear connections and shape maps for second order ODEs with and without constraints, *J. Geom. Phys.* **170** (2021) 104390.
- [6] O. Krupková and G.E. Prince. Second order ordinary differential equation in jet bundles and the inverse problem of the calculus of variation in: *Handbook of Global Analysis*, edited by D. Krupka and D. Saunders, Elsevier 2008.

G.E. Prince

Department of Mathematics and Statistics, La Trobe University,
Victoria 3086, Australia

Email: g.prince@latrobe.edu.au

D.J. Saunders

Lepage Research Institute,
17. novembra 1, 081 16 Prešov, Slovakia

Email: david@symplectic.email