

On the classification of Landsberg spherically symmetric Finsler metrics

S. G. ELGENDI

ABSTRACT. In this paper, as an application of the inverse problem of calculus of variations, we investigate two compatibility conditions on the spherically symmetric Finsler metrics. By making use of these conditions, we focus our attention on the Landsberg spherically symmetric Finsler metrics. We classify all spherically symmetric manifolds of Landsberg or Berwald types. For the higher dimensions $n \geq 3$, we prove that: all Landsberg spherically symmetric manifolds are either Riemannian or their geodesic sprays have a specific formula; all regular Landsberg spherically symmetric metrics are Riemannian; all (regular or non-regular) Berwald spherically symmetric metrics are Riemannian. Moreover, we establish new unicorns, i.e., new explicit examples of non-regular non-Berwaldian Landsberg metrics are obtained. For the two-dimensional case, we characterize all Berwald or Landsberg spherically symmetric surfaces.

1. INTRODUCTION

A spray S on an n -dimensional manifold M is a vector field on the tangent bundle TM and arises from a system of second-order ordinary differential equations and conversely, a spray can be associated with a system of second-order ordinary differential equations. The inverse problem for sprays (or, the so-called Finsler metrization problem) is the problem of deciding, for a given spray S , whether or not there exists a Lagrangian or a Finsler function such that S is its geodesic spray. The inverse problem for sprays is a special case of the inverse problem of the calculus of variations for arbitrary systems of second-order ordinary differential equations. Therefore, a Finsler function F has a unique geodesic spray but for a given spray, to decide if it is coming from a Finsler function, one has to solve the inverse problem.

In a Berwald manifold (M, F) and for any piecewise smooth curve $c(t)$ connecting two points $p, q \in M$, the Berwald parallel translation P_c is linear isometry between (T_pM, F_p) and (T_qM, F_q) and this is equivalent to that the geodesic spray of F is quadratic. But, in a Landsberg manifold the parallel translation P_c along c preserves the induced Riemannian metrics on the slit tangent spaces, that is, $P_c : (T_pM \setminus \{0\}, g_p) \rightarrow (T_qM \setminus \{0\}, g_q)$ is an isometry. This is equivalent to that the horizontal covariant derivative of the metric tensor of F with respect to the Berwald connection vanishes. It is clear that every Berwald manifold is Landsbergian. Whether there exists a regular Landsberg manifold, which is not Berwaldian, is a long-standing question in Finsler geometry, which is still open. To the best of our knowledge, it is not known a concrete example of a regular non-Berwaldian Landsberg manifold. In [1], G. S. Asanov introduced a class of examples, arising from general relativity, of non-Berwaldian Landsberg manifolds, of dimension at least 3. In [10], Z. Shen generalized Asanov's class to the (α, β) -metrics of Landsberg type. The difficulty of finding a regular non-Berwaldian Landsberg space leads Bao [2] to describe them as the unicorns of Finsler geometry. For more details and more examples of non-regular non-Berwaldian Landsberg metrics, we refer to, [3, 4, 5]. In dimension two, Zhou [12] provides non-regular examples of non-Berwaldian Landsberg spherically symmetric surfaces. But it was shown in [6] that these surfaces are Berwaldian. At the end of this paper we discuss this class in some details.

Let Ω be a convex domain in \mathbb{R}^n . A Finsler metric F on Ω is said to be spherically symmetric if the orthogonal group acts as isometries of F , that is, (Ω, F) is invariant under all rotations in \mathbb{R}^n . In other words, F is spherically symmetric if it is orthogonally invariant in the sense that $F(Ax, Ay) = F(x, y)$, where $A \in O(n)$, for more details see [7]. Spherically symmetric metrics were introduced by Rutz [9] where she generalized the classic Birkhoff theorem in general relativity to the

2020 *Mathematics Subject Classification.* 53C60, 53B40, 58B20.

Key words and phrases. spherically symmetric Finsler metrics; Landsberg metrics; Berwald metrics; inverse problem.

Finslerian case. Recently, spherically symmetric Finsler metrics have been paid a lot of attention, for example, we refer to [8, 12].

Let $|\cdot|$ and $\langle \cdot, \cdot \rangle$ be the standard Euclidean norm and the inner product on \mathbb{R}^n . A Finsler metric F on $\mathbb{B}^n(r_0) \subset \mathbb{R}^n$ is spherically symmetric if and only if there exists a positive C^∞ function ϕ such that $F(x, y) = u\phi(r, s)$ where $r = |x|$, $u = |y|$ and $s = \frac{\langle x, y \rangle}{|y|}$ and $(x, y) \in T\mathbb{B}^n(r_0) \setminus \{0\}$. The geodesic spray S of F is given by

$$G^i = uPy^i + u^2Qx^i.$$

Now, since a spherically symmetric metric F is written in the form $F = u\phi$ then we can talk about ϕ rather than F . For a given function ϕ we have the geodesic spray and hence the functions P and Q . But if P and Q are given, then we have to solve the inverse problem to decide if there exists a function ϕ such that the geodesic spray is given by P and Q .

In this paper, we solve the inverse problem for the spherically symmetric metrics and find compatibility conditions on P , Q and ϕ , that is for given P and Q then the compatibility conditions provide the function ϕ where the geodesic spray is given by P and Q (see Proposition 4.1).

Depending on the compatibility conditions, we classify all Landsberg spherically symmetric Finsler metrics; that is, we prove that a Landsberg spherically symmetric Finsler metric of dimension $n \geq 3$ is either Riemannian or the geodesic spray is given by the functions

$$P = c_1s + \frac{c_2}{r^2}\sqrt{r^2 - s^2}, \quad Q = \frac{1}{2}c_0s^2 - \frac{c_2s}{r^4}\sqrt{r^2 - s^2} + c_3,$$

where c_0, c_1, c_2, c_3 are arbitrary functions of r (see Theorem 5.1). We find out new non-regular Landsberg spherically symmetric metrics which are not Berwaldian. These metrics are given by $F = u\phi(r, s)$, where ϕ is given by

$$\begin{aligned} \phi(r, s) &= (5s^2 - r^2)^{\frac{1}{3}}(2s^2 - r^2)^{\frac{1}{6}} \exp\left(\frac{1}{3} \operatorname{arctanh}\left(\frac{\sqrt{5}(\sqrt{5}s + 5r)}{10\sqrt{r^2 - s^2}}\right) - \frac{1}{6} \operatorname{arctanh}\left(\frac{\sqrt{2}(\sqrt{2}s + 2r)}{2\sqrt{r^2 - s^2}}\right)\right) \\ &\quad + \frac{1}{3} \operatorname{arctanh}\left(\frac{\sqrt{5}(\sqrt{5}s - 5r)}{2\sqrt{r^2 - s^2}}\right) - \frac{1}{6} \operatorname{arctanh}\left(\frac{\sqrt{2}(\sqrt{2}s - 2r)}{2\sqrt{r^2 - s^2}}\right), \\ \phi(r, s) &= \frac{1}{r}(r^4 - 5r^2s^2 + 5s^4)^{\frac{1}{4}} \left(\frac{A}{B}\right)^{\frac{5-\sqrt{5}}{20}} \left(\frac{C}{D}\right)^{\frac{5+\sqrt{5}}{20}} \exp\left(-\frac{\sqrt{5}}{10} \operatorname{arctanh}\left(\frac{\sqrt{5}(r^2 - 2s^2)}{r^2}\right)\right), \end{aligned}$$

where

$$\begin{aligned} A &:= (2\sqrt{r^2 - s^2} - 2r + (\sqrt{5} - 1)s)\sqrt{r^2 - s^2} - (\sqrt{5} - 1)rs, \\ B &:= (2\sqrt{r^2 - s^2} - 2r - (\sqrt{5} - 1)s)\sqrt{r^2 - s^2} + (\sqrt{5} - 1)rs, \\ C &:= (2\sqrt{r^2 - s^2} - 2r - (\sqrt{5} + 1)s)\sqrt{r^2 - s^2} + (\sqrt{5} + 1)rs, \\ D &:= (2\sqrt{r^2 - s^2} - 2r + (\sqrt{5} + 1)s)\sqrt{r^2 - s^2} - (\sqrt{5} + 1)rs. \end{aligned}$$

For more details about the above two metrics, see Examples 1 and 2. We show that all (regular or non-regular) Berwald spherically symmetric metrics of dimension $n \geq 3$ are Riemannian (see Theorem 5.4). Also, we prove that all regular Landsberg spherically symmetric metrics of dimension $n \geq 3$ are Riemannian (see Theorem 5.5).

Finally, in Section 6, we focus our attention on the two-dimensional case. We characterize all Berwaldian and Landsbergian spherically symmetric surfaces. We prove that a spherically symmetric Finsler surface is Berwald if and only if

$$\begin{aligned} P &= b_1s + \frac{b_2}{\sqrt{r^2 - s^2}} + \frac{b_3(r^2 - 2s^2)}{\sqrt{r^2 - s^2}}, \\ Q &= b_0s^2 + \frac{1}{2}b_1 + \frac{b_2s(r^2 - 2s^2)}{r^4\sqrt{r^2 - s^2}} - \frac{b_3s(3r^2 - 2s^2)}{r^2\sqrt{r^2 - s^2}} - \frac{a}{r^2}s\sqrt{r^2 - s^2}, \end{aligned}$$

where a, b_0, b_1, b_2, b_3 are arbitrary functions of r and to be chosen such that the compatibility conditions are satisfied. We discuss the Landsberg surfaces. Moreover, we give a note on a class of Landsberg surfaces mentioned in [12] which, in fact, is Berwaldian.

2. PRELIMINARIES

Let M be an n -dimensional manifold and (TM, π, M) be its tangent bundle. We denote by $\mathcal{TM} := TM \setminus \{0\}$ the set of nonzero tangent vectors. Also, (x^i) denote the local coordinates on the base manifold M and (x^i, y^i) are the induced coordinates on TM . The vector 1-form J on TM defined by $J = \frac{\partial}{\partial y^i} \otimes dx^i$ is called the natural almost-tangent structure of TM . The vertical vector field $\mathcal{C} = y^i \frac{\partial}{\partial y^i}$ on TM is called the canonical or the Liouville vector field.

A vector field $S \in \mathfrak{X}(TM)$ is called a spray if $JS = \mathcal{C}$ and $[\mathcal{C}, S] = S$. Locally, a spray can be expressed as follows

$$S = y^i \frac{\partial}{\partial x^i} - 2G^i \frac{\partial}{\partial y^i},$$

where the spray coefficients $G^i = G^i(x, y)$ are 2-homogeneous functions in y .

A nonlinear connection on TM is defined by an n -dimensional distribution $H : z \in \mathcal{TM} \rightarrow H_z(TM)$ which is supplementary to the vertical distribution. This means that for all $z \in \mathcal{TM}$, we have

$$T_z(TM) = H_z(TM) \oplus V_z(TM).$$

Every spray S induces a canonical nonlinear connection through the corresponding horizontal and vertical projectors,

$$h = \frac{1}{2}(Id + [J, S]), \quad v = \frac{1}{2}(Id - [J, S])$$

With respect to the induced nonlinear connection, we have two associated projectors h and v which are expressed as follows

$$h = \frac{\delta}{\delta x^i} \otimes dx^i, \quad v = \frac{\partial}{\partial y^i} \otimes \delta y^i,$$

$$\frac{\delta}{\delta x^i} = \frac{\partial}{\partial x^i} - G^j_i(x, y) \frac{\partial}{\partial y^j}, \quad \delta y^i = dy^i + G^i_j(x, y) dx^j, \quad G^j_i(x, y) = \frac{\partial G^j}{\partial y^i}.$$

If $X \in \mathfrak{X}(M)$, i_X and \mathcal{L}_X denote the interior product by X and the Lie derivative in the direction of X , respectively. Every vector ℓ -form L defines two graded derivations i_L and d_L of the exterior algebra of M such that

$$i_L f = 0, \quad i_L df = df \circ L, \quad d_L := [i_L, d] = i_L \circ d - (-1)^{\ell-1} di_L,$$

where $f \in C^\infty(M)$ and df is the exterior derivative of f . If $X \in \mathfrak{X}(M)$ is a vector field, then i_X is simply the interior product by X and $d_X = \mathcal{L}_X$ the Lie derivative with respect to X .

A Finsler function on M is defined as follows:

Definition 2.1. A Finsler manifold of dimension n is a pair (M, F) , where M is a differentiable manifold of dimension n and F is a map, called Finsler function or Finsler metric,

$$F : TM \longrightarrow \mathbb{R},$$

such that:

- (a): F is smooth and strictly positive on TM and $F(x, y) = 0$ if and only if $y = 0$,
- (b): F is positively homogeneous of degree 1 in the direction argument y : $d_C F = \mathcal{L}_C F = F$,
- (c): The metric tensor $g_{ij} = \dot{\partial}_i \dot{\partial}_j E$ has maximal rank on TM , where $E := \frac{1}{2} F^2$ is the energy function.

In this case (M, F) is called a regular Finsler space. If F is not smooth or even not defined in some directions, then we call the Finsler function non-regular. In other words, by a non-regular Finsler metric F we mean that F does not satisfy the condition (a) or (c) on certain directions of TM .

Since the 2-form $dd_J E$ is non-degenerate, the Euler-Lagrange equation

$$i_S dd_J E = -dE$$

uniquely determines a spray S on TM . This spray is called the *geodesic spray* of the Finsler function F .

Definition 2.2. A spray S on a manifold M is called *Finsler metrizable* if there exists a Finsler function F such that the geodesic spray of the Finsler manifold (M, F) is S .

It is known that a spray S is Finsler metrizable if and only if there exists a non-degenerate solution F for the system

$$(2.1) \quad d_h F = 0, \quad d_C F = F,$$

where h is the horizontal projector associated to S .

So, to solve the inverse problem for a given spray S , we have to find a non-degenerate solution F for the system (2.1).

3. SPHERICALLY SYMMETRIC METRICS

Let $|\cdot|$ and $\langle \cdot, \cdot \rangle$ be the standard Euclidean norm and inner product on \mathbb{R}^n . A Finsler metric F on $\mathbb{B}^n(r_0) \subset \mathbb{R}^n$ is spherically symmetric if and only if there exists a function $\phi : [0, r_0) \times \mathbb{R}^n \rightarrow \mathbb{R}$ such that

$$F(x, y) = |y|\phi\left(|x|, \frac{\langle x, y \rangle}{|y|}\right)$$

where $(x, y) \in T\mathbb{B}^n(r_0) \setminus \{0\}$. Or simply we write $F = u\phi(r, s)$ where $r = |x|$, $u = |y|$ and $s = \frac{\langle x, y \rangle}{|y|}$.

The spherically symmetric metrics are a special case of the general (α, β) -metrics. The general (α, β) -metrics are introduced and studied by C. Yu and H. Zhu [11]. Therefore, the spherically symmetric metrics $F = u\phi(r, s)$ on $\mathbb{B}^n(r_0)$ are regular if and only if ϕ is positive and C^∞ function and

$$(3.1) \quad \phi - s\phi_s > 0, \quad \phi - s\phi_s + (r^2 - s^2)\phi_{ss} > 0$$

when $n \geq 3$ or $\phi - s\phi_s + (r^2 - s^2)\phi_{ss} > 0$ when $n = 2$ for all $|s| \leq r < r_0$. Throughout, the subscript s (resp. r) refers to the derivative with respect to s (rep. r). The spherically symmetric Finsler metrics are studied in many papers for example, we refer to, [7, 8, 12].

Remark 3.1. It should be noted that throughout, the results and formulae work and are valid for both regular and non-regular spherically symmetric metrics unless it is specified. So we consider the most general case, that is, the non-regular case and when the situation needs, we mention clearly that the metric is regular.

The components g_{ij} of the metric tensor of the spherically symmetric metric $F = u\phi(r, s)$ are given by

$$(3.2) \quad g_{ij} = \sigma_0 \delta_{ij} + \sigma_1 \delta_{ik} \delta_{hj} x^k x^h + \frac{\sigma_2}{u} (\delta_{ik} \delta_{hj} + \delta_{jk} \delta_{hi}) x^h y^k + \frac{\sigma_3}{u^2} \delta_{ik} \delta_{hj} y^k y^h,$$

where

$$\sigma_0 = \phi(\phi - s\phi_s), \quad \sigma_1 = \phi_s^2 + \phi\phi_{ss}, \quad \sigma_2 = (\phi - s\phi_s)\phi_s - s\phi\phi_{ss}, \quad \sigma_3 = s^2\phi\phi_{ss} - s(\phi - s\phi_s)\phi_s.$$

The above formula can be found in [12] and to keep the indices consistent in both sides of a tensorial equation, we write for example, $\delta_{ik}y^k$ when there is a y^i and i is a lower index.

The components g^{ij} of the inverse metric tensor are given by [12] as follows

$$(3.3) \quad g^{ij} = \rho_0 \delta^{ij} + \frac{\rho_1}{u^2} y^i y^j + \frac{\rho_2}{u} (x^i y^j + x^j y^i) + \rho_3 x^i x^j,$$

where

$$\rho_0 = \frac{1}{\phi(\phi - s\phi_s)}, \quad \rho_1 = \frac{(s\phi + (r^2 - s^2)\phi_s)(\phi\phi_s - s\phi_s^2 - s\phi_{ss})}{\phi^3(\phi - s\phi_s)(\phi - s\phi_s + (r^2 - s^2)\phi_{ss})},$$

$$\rho_2 = -\frac{\phi\phi_s - s\phi_s^2 - s\phi_{ss}}{\phi^2(\phi - s\phi_s)(\phi - s\phi_s + (r^2 - s^2)\phi_{ss})}, \quad \rho_3 = -\frac{\phi_{ss}}{\phi(\phi - s\phi_s)(\phi - s\phi_s + (r^2 - s^2)\phi_{ss})}.$$

Moreover, the geodesic spray coefficients G^i of F are given by

$$(3.4) \quad G^i = uPy^i + u^2Qx^i,$$

where the functions P and Q have the following formulae

$$(3.5) \quad Q := \frac{1}{2r} \frac{-\phi_r + s\phi_{rs} + r\phi_{ss}}{\phi - s\phi_s + (r^2 - s^2)\phi_{ss}}, \quad P := -\frac{Q}{\phi} (s\phi + (r^2 - s^2)\phi_s) + \frac{1}{2r\phi} (s\phi_r + r\phi_s).$$

For the geodesic spray of the spherically symmetric metric F , we have the associated nonlinear connection whose components G_j^i are defined by $G_j^i := \frac{\partial G^i}{\partial y^j}$. Using the following identities:

$$(3.6) \quad \frac{\partial r}{\partial x^j} = \frac{1}{r} \delta_{jk} x^k, \quad \frac{\partial u}{\partial y^j} = \frac{1}{u} \delta_{jk} y^k, \quad \frac{\partial u}{\partial x^j} = 0, \quad \frac{\partial r}{\partial y^j} = 0,$$

$$\frac{\partial s}{\partial x^j} = \frac{1}{u} \delta_{jk} y^k, \quad \frac{\partial s}{\partial y^j} = \frac{1}{u} \delta_{jk} x^k - \frac{s}{u^2} \delta_{jk} y^k,$$

the components G_j^i can be calculated as follows

$$(3.7) \quad G_j^i = uP\delta_j^i + P_s \delta_{kj} x^k y^i + \frac{1}{u} (P - sP_s) \delta_{kj} y^k y^i + uQ_s \delta_{jk} x^i x^k + (2Q - sQ_s) \delta_{jk} x^i y^k.$$

The components G_{jkl}^i of the Berwald curvature are defined by

$$G_{jkl}^i = \frac{\partial^2}{\partial y^\ell \partial y^k} G_j^i.$$

For a spherically symmetric Finsler metric $F = u \phi(r, s)$, we use the identities (3.6) and (3.7) to find the components G_{jkl}^i as follows

$$\begin{aligned}
G_{jkl}^i &= \frac{P_{ss}}{u} (\delta_j^i \delta_{hk} \delta_{t\ell} + \delta_\ell^i \delta_{hk} \delta_{tj} + \delta_k^i \delta_{hj} \delta_{t\ell}) x^h x^t + \frac{1}{u} (P - sP_s) (\delta_j^i \delta_{k\ell} + \delta_k^i \delta_{j\ell} + \delta_\ell^i \delta_{jk}) \\
&\quad - \frac{sP_{ss}}{u^2} (\delta_j^i (\delta_{hk} \delta_{t\ell} + \delta_{h\ell} \delta_{tk}) + \delta_k^i (\delta_{hj} \delta_{t\ell} + \delta_{h\ell} \delta_{tj}) + \delta_\ell^i (\delta_{hk} \delta_{tj} + \delta_{hj} \delta_{tk})) x^h y^t \\
&\quad - \frac{sP_{ss}}{u^2} (\delta_{jk} \delta_{h\ell} + \delta_{j\ell} \delta_{kh} + \delta_{k\ell} \delta_{jh}) x^h y^i + \frac{1}{u} (Q_s - sQ_{ss}) (\delta_{jk} \delta_{h\ell} + \delta_{j\ell} \delta_{kh} + \delta_{k\ell} \delta_{jh}) x^i x^h \\
&\quad + \frac{1}{u^3} (s^2 P_{ss} + sP_s - P) ((\delta_j^i \delta_{kh} \delta_{t\ell} + \delta_k^i \delta_{jh} \delta_{t\ell} + \delta_\ell^i \delta_{kh} \delta_{tj}) y^h y^t + (\delta_{jk} \delta_{\ell h} + \delta_{j\ell} \delta_{kh} + \delta_{k\ell} \delta_{jh}) y^h y^i) \\
&\quad + \frac{1}{u^5} (3P - s^3 P_{sss} - 6s^2 P_{ss} - 3sP_s) \delta_{jh} \delta_{km} \delta_{\ell t} y^h y^m y^t y^i + \frac{P_{sss}}{u^2} \delta_{jh} \delta_{km} \delta_{\ell t} x^h x^m x^t y^i \\
&\quad + \frac{1}{u^4} (s^2 P_{sss} + 3sP_{ss}) (\delta_{jm} \delta_{kh} \delta_{\ell t} + \delta_{jm} \delta_{kt} \delta_{\ell h} + \delta_{\ell m} \delta_{kh} \delta_{jt}) x^t y^m y^h y^i \\
&\quad - \frac{1}{u^3} (P_{ss} + sP_{sss}) (\delta_{km} \delta_{jh} \delta_{\ell t} + \delta_{jm} \delta_{kh} \delta_{\ell t} + \delta_{jm} \delta_{\ell h} \delta_{kt}) x^t x^m y^h y^i + \frac{Q_{sss}}{u} \delta_{km} \delta_{jh} \delta_{\ell t} x^h x^m x^t x^i \\
&\quad + \frac{1}{u^3} (s^2 Q_{sss} + sQ_{ss} - Q_s) (\delta_{km} \delta_{jh} \delta_{\ell t} + \delta_{\ell m} \delta_{kh} \delta_{jt} + \delta_{\ell h} \delta_{km} \delta_{jt}) x^h x^i y^m y^t - \frac{sQ_{sss}}{u^2} (\delta_{\ell m} \delta_{jh} \delta_{kt} \\
&\quad + \delta_{\ell t} \delta_{kh} \delta_{jm} + \delta_{\ell m} \delta_{kh} \delta_{jt}) x^h x^i x^m y^t + \frac{1}{u^4} (3sQ_s - 3s^2 Q_{ss} - s^3 Q_{sss}) \delta_{\ell m} \delta_{kh} \delta_{jt} y^t y^m y^h x^i \\
&\quad + \frac{1}{u^2} (s^2 Q_{ss} - sQ_s) (\delta_{k\ell} \delta_{hj} + \delta_{j\ell} \delta_{hk} + \delta_{kj} \delta_{h\ell}) y^h x^i.
\end{aligned}$$

The above formula of the Berwald curvature can be found, for example, in [8]. By straightforward calculations, the components $E_{ij} := G_{ijh}^h$ of the mean Berwald curvature can be calculated as follows:

$$\begin{aligned}
(3.8) \quad E_{ij} &= \frac{1}{u} ((n+1)(P - sP_s) + (r^2 - s^2)(Q_s - sQ_{ss})) \delta_{ij} + \frac{1}{u^3} ((n+1)(s^2 P_{ss} + sP_s - P) \\
&\quad + r^2 (s^2 Q_{sss} + sQ_{ss} - Q_s) + 3s^2 Q_s - 3s^3 Q_{ss} - s^4 Q_{sss}) \delta_{ih} \delta_{jk} y^h y^k \\
&\quad + \frac{1}{u} ((n+1)P_{ss} + 2(Q_s - sQ_{ss}) + (r^2 - s^2)Q_{sss}) \delta_{ih} \delta_{jk} x^h x^k \\
&\quad - \frac{s}{u^2} ((n+1)P_{ss} + 2(Q_s - sQ_{ss}) + (r^2 - s^2)Q_{sss}) (\delta_{ih} \delta_{jk} + \delta_{ik} \delta_{jh}) x^h y^k.
\end{aligned}$$

One can rewrite E_{ij} as follows

$$(3.9) \quad E_{ij} = \frac{H}{u} \delta_{ij} - \frac{1}{u^3} (sH_s + H) \delta_{hi} \delta_{kj} y^h y^k + \frac{H_s}{su^2} (s(\delta_{hi} \delta_{kj} + \delta_{ki} \delta_{hj}) x^h y^k - u \delta_{hi} \delta_{kj} x^h x^k),$$

where

$$H := (n+1)(P - sP_s) + (r^2 - s^2)(Q_s - sQ_{ss}).$$

Now, we have the following proposition.

Proposition 3.2. *If the mean Berwald curvature vanishes, then either $n = 2$ or $H = 0$.*

Proof. Assume that $E_{ij} = 0$, then contracting (3.9) by δ^{ij} , and using the facts that $\delta_{ij} y^i y^j = u^2$, $\delta_{ij} x^i x^j = r^2$ and $\delta_{ij} x^i y^j = \langle x, y \rangle = su$, we have

$$s(n-1)H - (r^2 - s^2)H_s = 0.$$

Contracting (3.9) by $x^i x^j$ implies

$$sH - (r^2 - s^2)H_s = 0.$$

Subtracting the above two equations, yields

$$s(n-2)H = 0.$$

Which holds for all s , then we obtain that $n = 2$ or $H = 0$ and hence the proof is completed. \square

To avoid confusions, we have to clarify a point mentioned in [12, Proposition 2.1] by providing the following remark.

Remark 3.3. It is clear that if $H = 0$, then the mean Berwald curvature vanishes, that is $E_{ij} = 0$. By making use of the above proposition, when $n = 2$, the condition $E_{ij} = 0$ does not imply $H = 0$ contrary to what mentioned in [12] and discussed in [6]. For the higher dimensions $n \geq 3$, $E_{ij} = 0$ if and only if $H = 0$.

The scalar trace E of the mean Berwald curvature is $E = g^{ij} E_{ij}$. For a spherically symmetric Finsler metric, the scalar trace of the mean Berwald curvature is given as follows.

Proposition 3.4. *The scalar trace E of the mean Berwald curvature of a spherically symmetric Finsler metric is given by*

$$E = \frac{1}{su} (s((n-1)\rho_0 + \rho_3(r^2 - s^2))H - (r^2 - s^2)(\rho_0 + \rho_3(r^2 - s^2))H_s).$$

Proof. By using (3.3) and the fact that $y^i E_{ij} = 0$, the scalar trace of the mean Berwald curvature is calculated as follows

$$\begin{aligned} E &= g^{ij} E_{ij} \\ &= \frac{\rho_0}{su} \left((n-1)sH - (r^2 - s^2)H_s \right) + \frac{\rho_3}{su} (s(r^2 - s^2)H - (r^2 - s^2)^2 H_s) \\ &= \frac{1}{su} (s((n-1)\rho_0 + \rho_3(r^2 - s^2))H - (r^2 - s^2)(\rho_0 + \rho_3(r^2 - s^2))H_s). \end{aligned}$$

□

The components L_{jkl} of the Landsberg curvature are defined by

$$L_{jkl} = -\frac{1}{2} F G_{jkl}^h \frac{\partial F}{\partial y^h}.$$

For a spherically symmetric Finsler metric $F = u \phi(r, s)$, the components L_{jkl} are given by

$$\begin{aligned} (3.10) \quad L_{jkl} &= -\frac{\phi}{2} \left(L_1 \delta_{jh} \delta_{km} \delta_{\ell t} x^h x^m x^t + \frac{3sL_2 - s^3 L_1}{u^3} \delta_{jh} \delta_{km} \delta_{\ell t} y^h y^m y^t + L_2 (\delta_{kl} \delta_{jh} + \delta_{jl} \delta_{kh} \right. \\ &\quad \left. + \delta_{kj} \delta_{\ell h}) x^h - \frac{sL_2}{u} (\delta_{k\ell} \delta_{hj} + \delta_{kh} \delta_{\ell j} + \delta_{kj} \delta_{h\ell}) y^h - \frac{sL_1}{u} (\delta_{mk} \delta_{\ell t} \delta_{hj} + \delta_{kh} \delta_{mj} \delta_{\ell t} \right. \\ &\quad \left. + \delta_{mj} \delta_{h\ell} \delta_{kt}) x^m x^t y^h + \frac{s^2 L_1 - L_2}{u^2} (\delta_{mk} \delta_{\ell t} \delta_{hj} + \delta_{kh} \delta_{mj} \delta_{\ell t} + \delta_{mj} \delta_{h\ell} \delta_{kt}) y^m y^t x^h \right) \end{aligned}$$

where

$$\begin{aligned} L_1 &= 3\phi_s P_{ss} + \phi P_{sss} + (s\phi + (r^2 - s^2)\phi_s) Q_{sss}, \\ L_2 &= -s\phi P_{ss} + \phi_s (P - sP_s) + (s\phi + (r^2 - s^2)\phi_s) (Q_s - sQ_{ss}). \end{aligned}$$

It is well known that a Finsler metric F is said to be Berwaldain if and only if its Berwald curvature vanishes, that is, $G_{ijk}^h = 0$ and this is equivalent to the fact that the geodesic spray of F is quadratic in y . Also, F is called Landsbergain if and only if its Landsberg curvature vanishes, that is, $L_{ijk} = 0$. Paying more attention to the spherically symmetric Finsler metrics of Landsberg type, we have the following proposition.

Proposition 3.5. *If a spherically symmetric Finsler metric $F = u\phi$ is Landsbergian then either $n = 2$ or $L_1 = L_2 = 0$.*

Proof. Let F be a Landsberg spherically symmetric Finsler metric, then $L_{jkl} = 0$. Contracting (3.10) by $x^j \delta^{k\ell}$ and using the facts that $\delta_{ij} y^i y^j = u^2$, $\delta_{ij} x^i x^j = r^2$ and $\delta_{ij} x^i y^j = \langle x, y \rangle = su$, we have

$$(r^2 - s^2)L_1 + (n+1)L_2 = 0.$$

Similarly, contracting (3.10) by $x^j x^k x^\ell$, we get

$$(r^2 - s^2)L_1 + 3L_2 = 0.$$

Subtracting the above two equations implies

$$(n-2)L_2 = 0.$$

Therefore, $n = 2$ or $L_2 = 0$ and this completes the proof. □

By the above proposition together with the fact that if $L_1 = L_2 = 0$ then $L_{jkl} = 0$, one can see that a Landsbergian spherically symmetric Finsler metric of dimension $n \geq 3$ is characterized by the following two conditions

$$\begin{aligned} (3.11) \quad L_1 &= 3\phi_s P_{ss} + \phi P_{sss} + (s\phi + (r^2 - s^2)\phi_s) Q_{sss} = 0, \\ L_2 &= -s\phi P_{ss} + \phi_s (P - sP_s) + (s\phi + (r^2 - s^2)\phi_s) (Q_s - sQ_{ss}) = 0. \end{aligned}$$

Again, to avoid confusions in the class of the spherically symmetric Finsler metrics of Landsberg type, we have the following remark.

Remark 3.6. It is clear that if $L_1 = L_2 = 0$ then $L_{jkl} = 0$ and hence the metric F is Landsbergian. By the above proposition, when $n = 2$, $L_{jkl} = 0$ does not yield $L_1 = L_2 = 0$ as be discussed in Section 6. For the higher dimensions $n \geq 3$, $L_{jkl} = 0$ if and only if $L_1 = L_2 = 0$.

4. THE COMPATIBILITY CONDITIONS

For a non-regular spherically symmetric metric $F = u\phi(r, s)$, the case $\phi - s\phi_s = 0$ is excluded. If $\phi - s\phi_s = 0$, then $\phi = f(r)s$ and hence $F = f(r)us = f(r)\langle x, y \rangle$ which is linear in y for an arbitrary function $f(r)$. If the Finsler function is linear in y , then the metric tensor is degenerate everywhere and this is impossible. Moreover, if $\phi_s = 0$, then $F = f(r)u$ which is the Euclidean metric or a conformal transformation of the Euclidean metric and this case also is excluded.

Proposition 4.1. *Let $F = u\phi(r, s)$ be a spherically symmetric metric, then the geodesic spray of F is not defined if and only if $\phi - s\phi_s + (r^2 - s^2)\phi_{ss} = 0$ or equivalently ϕ is given by*

$$\phi(r, s) = f_1(r)s + f_2(r)\sqrt{r^2 - s^2},$$

where $f_1(r)$ and $f_2(r)$ are arbitrary functions of r .

Proof. By (3.5), it is clear that the function Q is not defined and so the geodesic spray if and only if $\phi - s\phi_s + (r^2 - s^2)\phi_{ss} = 0$. Since the above formula holds for all r and s , then we can write

$$\frac{\phi_{ss}}{\phi - s\phi_s} = -\frac{1}{r^2 - s^2}.$$

Since $(\phi - s\phi_s)_s = -s\phi_{ss}$, integrating the above equation with respect to s gives

$$\phi - s\phi_s = \frac{f_1(r)}{\sqrt{r^2 - s^2}}.$$

Which has the solution

$$\phi(r, s) = f_1(r)s + f_2(r)\sqrt{r^2 - s^2}.$$

□

For a given spherically symmetric Finsler metric $F = u\phi$, then the geodesic spray (3.4) is given by the functions P and Q . But if the functions P and Q are given, then to find the Finsler metric whose geodesic spray is determined by P and Q , we have to solve the inverse problem or the metrizable problem. So, we investigate the following two compatibility conditions.

Proposition 4.2. *Let $P(r, s)$ and $Q(r, s)$ be given, then the Finsler function $F = u\phi(r, s)$ whose geodesic spray given by P and Q is determined by the function ϕ provided that ϕ satisfies the following compatibility conditions:*

$$(4.1) \quad \begin{aligned} C_1 &:= (1 + sP - (r^2 - s^2)(2Q - sQ_s))\phi_s + (sP_s - 2P - s(2Q - sQ_s))\phi = 0, \\ C_2 &:= \frac{1}{r}\phi_r - (P + Q_s(r^2 - s^2))\phi_s - (P_s + sQ_s)\phi = 0. \end{aligned}$$

Proof. Let $P(r, s)$ and $Q(r, s)$ be two given functions. To find the Finsler function $F = u\phi(r, s)$ whose geodesic spray determined by P and Q , we have to solve the system $d_h F = 0$ (2.1). Locally, the system $d_h F = 0$ gives rise to

$$(4.2) \quad \frac{\partial(u\phi)}{\partial x^j} - G_j^i \frac{\partial(u\phi)}{\partial y^i} = 0.$$

We have the following properties

$$\frac{\partial\phi}{\partial x^i} = \frac{\phi_r}{r}\delta_{ik}x^k + \frac{\phi_s}{u}\delta_{ik}y^k, \quad \frac{\partial\phi}{\partial y^i} = \frac{\phi_s}{u}\delta_{ik}x^k - \frac{s\phi_s}{u^2}\delta_{ik}y^k.$$

Moreover, since $u\phi$ is homogeneous of degree one in y , then $y^j \frac{\partial(u\phi)}{\partial y^j} = u\phi$. Also, we have $x^j \frac{\partial(u\phi)}{\partial y^j} = s\phi + (r^2 - s^2)\phi_s$. Now, substituting the above formulae into the system (4.2), we have

$$\begin{aligned} \frac{\partial(u\phi)}{\partial x^j} - G_j^i \frac{\partial(u\phi)}{\partial y^i} &= \frac{u\phi_r}{r}\delta_{jk}x^k + \phi_s\delta_{jk}y^k - \left(uP\delta_j^i + P_s\delta_{kj}x^k y^i + \frac{1}{u}(P - sP_s)\delta_{kj}y^k y^i \right. \\ &\quad \left. + uQ_s\delta_{jk}x^i x^k + (2Q - sQ_s)\delta_{jk}x^i y^k \right) \frac{\partial(u\phi)}{\partial y^i} \\ &= u \left(\frac{1}{r}\phi_r - P\phi_s - \phi P_s - Q_s(s\phi + (r^2 - s^2)\phi_s) \right) \delta_{jk}x^k \\ &\quad + (\phi_s - 2P\phi + s\phi P_s + sP\phi_s - (2Q - sQ_s)(s\phi + (r^2 - s^2)\phi_s)) \delta_{jk}y^k \\ &= 0. \end{aligned}$$

Now, if there exist two functions μ and η such that

$$\mu\delta_{jk}x^k + \eta\delta_{jk}y^k = 0.$$

Then by contracting the above equation by x^j , resp. by y^j we get the following two equations

$$r^2\mu + s\eta = 0, \quad s\mu + u\eta = 0.$$

Then, we have $(r^2 - s^2)\mu = 0$ which holds for all r and s , that is, $\mu = 0$ and hence $\eta = 0$. Using this property, we get the conditions C_1 and C_2 .

Conversely, assume that the conditions C_1 and C_2 are satisfied. Rewriting C_1 and C_2 as follows

$$(4.3) \quad \phi_s + s(P_s\phi + P\phi_s) - 2P\phi - (s\phi + (r^2 - s^2)\phi_s)(2Q - sQ_s) = 0,$$

$$(4.4) \quad \frac{1}{r}\phi_r - (P\phi_s + P_s\phi) - (s\phi + (r^2 - s^2)\phi_s)Q_s = 0.$$

Adding (4.3) to the multiplication of (4.4) by s , we have

$$\phi_s + \frac{s}{r}\phi_r - 2P\phi - 2Q(s\phi + (r^2 - s^2)\phi_s) = 0.$$

Which implies

$$P = \frac{Q}{\phi}(s\phi + (r^2 - s^2)\phi_s) + \frac{1}{2r\phi}(r\phi_s + s\phi_r).$$

By the above formula of P , we get

$$P\phi_s + \phi P_s = \frac{1}{2r}(\phi_r + s\phi_{rs} + r\phi_{ss} + 2rQ(s\phi_s - \phi - (r^2 - s^2)\phi_{ss})) - Q_s(s\phi + (r^2 - s^2)\phi_s).$$

Now, substituting $P\phi_s + \phi P_s$ into (4.4), we obtain that

$$Q = \frac{1}{2r} \frac{-\phi_r + s\phi_{rs} + r\phi_{ss}}{\phi - s\phi_s + (r^2 - s^2)\phi_{ss}}.$$

This completes the proof. \square

5. LANDSBERG SPHERICALLY SYMMETRIC METRICS

In this section we focus our attention to the spherically symmetric metrics of Landsberg type. We start by the following result.

Theorem 5.1. *A Landsberg spherically symmetric Finsler metric of dimension $n \geq 3$ is either Riemannian or the geodesic spray is determined by the functions*

$$(5.1) \quad P = c_1s + \frac{c_2}{r^2}\sqrt{r^2 - s^2}, \quad Q = \frac{1}{2}c_0s^2 - \frac{c_2s}{r^4}\sqrt{r^2 - s^2} + c_3,$$

where c_0, c_1, c_2, c_3 are arbitrary functions of r .

Proof. Suppose that $F = u\phi(r, s)$ be a spherically symmetric metric of dimension $n \geq 3$, then by Proposition 3.5, the function ϕ and the functions P and Q of the geodesic spray satisfy the Landsberg conditions (3.11), that is, we have $L_1 = L_2 = 0$. The condition $L_2 = 0$ implies that

$$Q_s - sQ_{ss} = -\frac{-s\phi P_{ss} + \phi_s(P - sP_s)}{s\phi + (r^2 - s^2)\phi_s}.$$

It should be noted that if $s\phi + (r^2 - s^2)\phi_s = 0$, then by taking the derivative with respect to s implies $\phi - s\phi_s + (r^2 - s^2)\phi_{ss} = 0$ which is impossible by Proposition 4.1. Now, making use of the property that $(Q_s - sQ_{ss})_s = -sQ_{sss}$, then differentiating the above equation with respect to s , we get Q_{sss} . Substituting Q_{sss} into the Landsberg condition $L_1 = 0$ and straightforward simplifications, we have

$$((r^2 - s^2)P_{ss} + (P - sP_s))(s\phi_s^2 + s\phi\phi_{ss} - \phi\phi_s) = 0.$$

If $s\phi_s^2 + s\phi\phi_{ss} - \phi\phi_s = 0$, then we can write

$$\frac{-s\phi_{ss}}{\phi - s\phi_s} = -\frac{\phi_s}{\phi}.$$

By integration with respect to s , we have

$$\phi - s\phi_s = \frac{f_1(r)}{\phi}.$$

Using the substitution $\phi = s\psi$, then we can rewrite the above equation as follows

$$\psi\psi_s = -\frac{f_1(r)}{s^3}.$$

Which implies

$$\phi = \sqrt{f_1(r) + f_2(r)s^2}.$$

That is, the Finsler function given by $F = u\phi = \sqrt{f_1(r)u^2 + f_2(r)\langle x, y \rangle^2}$ is Riemannian.

If $(r^2 - s^2)P_{ss} + (P - sP_s) = 0$, then by assuming that $P - sP_s \neq 0$ we can write

$$\frac{-sP_{ss}}{P - sP_s} = \frac{s}{r^2 - s^2}.$$

Since $(P - sP_s)_s = -sP_{ss}$, then integrating the above equation with respect to s gives rise to

$$P - sP_s = \frac{c_1(r)}{\sqrt{r^2 - s^2}}.$$

Which has the solution

$$P = c_2(r)s + \frac{c_1(r)}{r^2}\sqrt{r^2 - s^2}.$$

Taking the fact that $s\phi + (r^2 - s^2)\phi_s \neq 0$, then substituting by the function P into the Landsberg condition $L_2 = 0$, implies that

$$Q_s - sQ_{ss} = -\frac{c_1(r)}{(r^2 - s^2)^{3/2}}.$$

Using the substitution $s\psi = Q_s$, then we can write

$$s^2\psi_s = \frac{c_2(r)}{(r^2 - s^2)^{3/2}}.$$

Which yields the solution

$$Q_s = c_0(r)s - \frac{c_2(r)(r^2 - 2s^2)}{r^4\sqrt{r^2 - s^2}}.$$

Which has the solution

$$Q = \frac{1}{2}c_0(r)s^2 - \frac{c_2(r)s}{r^4}\sqrt{r^2 - s^2} + c_3(r).$$

In the case where $P - sP_s = 0$ then we have $(r^2 - s^2)P_{ss} = 0$ for all r and s and hence we obtain that $P_{ss} = 0$ and also the Landsberg condition $L_2 = 0$, implies that $Q_s - sQ_{ss} = 0$ and hence $Q_{sss} = 0$. Therefore, $P = f_1(r)s$ and $Q = f_2(r)s^2 + f_3(r)$ which are a special case of the given formulae for P and Q . Hence the proof is completed. \square

Based on the above proof, we provide the following remark.

Remark 5.2. It should be noted that if $P - sP_s = 0$ and $Q_s - sQ_{ss} = 0$, then $P_{ss} = 0$ and $Q_{sss} = 0$. That is the Berwald curvature vanishes and hence the metric is Berwaldian. Moreover, if $n \geq 3$ and the Berwald curvature vanishes, then we get $P - sP_s = 0$ and $Q_s - sQ_{ss} = 0$. That is, as it was mentioned in [7, 8], the Berwaldain spherically symmetric Finsler metrics are characterized by $P - sP_s = 0$ and $Q_s - sQ_{ss} = 0$ provided that $n \geq 3$.

Theorem 5.3. *There exist non-Berwaldian Landsberg spherically symmetric Finsler metrics of dimension $n \geq 3$.*

Proof. Let $F = u\phi$ be a non-Riemannian Landsberg spherically symmetric Finsler metric of dimension $n \geq 3$, then by Theorem 5.1 the functions P and Q are given by

$$P = c_1s + \frac{c_2}{r^2}\sqrt{r^2 - s^2}, \quad Q = \frac{1}{2}c_0s^2 - \frac{c_2s}{r^4}\sqrt{r^2 - s^2} + c_3.$$

Now, the functions P , Q and ϕ must satisfy the compatipility conditons (4.1). Since ϕ is non-zero then dividing the conditions $C_1 = 0$ and $C_2 = 0$ in (4.1) by ϕ and substituting by P and Q , straightforward calculations imply the following two equations:

$$\begin{aligned} \frac{\phi_s}{\phi}(r^2 + (c_1 + 2c_3)r^2s^2 - 2c_3r^4 + 2c_2s\sqrt{r^2 - s^2}) - (c_1 + 2c_3)sr^2 - 2c_2\sqrt{r^2 - s^2} &= 0, \\ \frac{\phi_s}{\phi}(-c_0r^4s(r^2 - s^2) - c_1sr^4 - 2c_2s^2\sqrt{r^2 - s^2}) + r^3\frac{\phi_r}{\phi} - c_0s^2r^4 - c_1r^4 + 2c_2s\sqrt{r^2 - s^2} &= 0. \end{aligned}$$

Solving the above equations algebraically for $\frac{\phi_s}{\phi}$ and $\frac{\phi_r}{\phi}$, we obtain that

$$(5.2) \quad \begin{aligned} \frac{\phi_s}{\phi} &= \frac{(c_1 + 2c_3)r^2s + 2c_2\sqrt{r^2 - s^2}}{r^2 + (c_1 + 2c_3)r^2s^2 - 2c_3r^4 + 2c_2s\sqrt{r^2 - s^2}}, \\ \frac{\phi_r}{\phi} &= \frac{(2c_0c_2r^4 + 4(c_1 + c_3)c_2r^2 - 2c_2)s\sqrt{r^2 - s^2}}{r(r^2 + (c_1 + 2c_3)r^2s^2 - 2c_3r^4 + 2c_2s\sqrt{r^2 - s^2})} \\ &\quad + \frac{c_0c_1r^6s^2 + (c_0 + 4c_1c_3 + 2c_1^2)r^4s^2 - 2c_1c_3r^6 + c_1r^4}{r(r^2 + (c_1 + 2c_3)r^2s^2 - 2c_3r^4 + 2c_2s\sqrt{r^2 - s^2})}. \end{aligned}$$

It should be noted that the above two formulas are obtained in [8, Eq. (4.9)] by a long and completely different way.

Now, ϕ must satisfy the condition $\phi_{rs} = \phi_{sr}$ so we have to find what conditions on the functions c_0 , c_1 , c_2 and c_3 so that $\phi_{rs} = \phi_{sr}$. For this purpose we have

$$\left(\frac{\phi_s}{\phi}\right)_r - \left(\frac{\phi_r}{\phi}\right)_s = \frac{\phi(\phi_{sr} - \phi_{rs})}{\phi^2}.$$

That is, $\phi_{rs} = \phi_{sr}$ if and only if $\left(\frac{\phi_s}{\phi}\right)_r = \left(\frac{\phi_r}{\phi}\right)_s$. Applying the condition $\left(\frac{\phi_s}{\phi}\right)_r = \left(\frac{\phi_r}{\phi}\right)_s$ on (5.2), we get the following

$$\begin{aligned} & \frac{r^2}{\sqrt{r^2 - s^2}(r^2 + (c_1 + 2c_3)r^2s^2 - 2c_3r^4 + 2c_2s\sqrt{r^2 - s^2})^2} \left(s\sqrt{r^2 - s^2}(c_1' r^2 + 2c_3' r^2 + 8c_3^2 r^3 \right. \\ & - 2c_0 r^3 - 2c_1^2 r^3 + 4c_0 c_1 c_3 r^7 + 4c_0 c_3 r^5 + 4c_1^2 c_3 r^5 + 8c_1 c_3^2 r^5 - 2c_0 c_1 r^5 + 2c_1 c_3' r^4 \\ & - 2c_1' c_3 r^4) + (4c_0 c_2 c_3 r^7 + 4c_1 c_2 c_3 r^5 - 8c_2 c_3^2 s^2 r^3 + 2c_0 c_2 s^2 r^3 + 2c_1 c_2 s^2 r - 4c_2 c_3' s^2 r^2 \\ & - 4c_2 c_3 s^2 r + 4c_2' c_3 s^2 r^2 + 2c_2' r^2 - 2c_2^2 s^2 - 4c_0 c_2 c_3 s^2 r^5 - 4c_1 c_2 c_3 s^2 r^3 + 8c_2 c_3^2 r^5 - 2c_0 c_2 r^5 \\ & \left. - 2c_1 c_2 r^3 + 4c_2 c_3' r^3 - 4c_2' c_3 r^4 + 4c_2 c_3 r^3) \right) = 0, \end{aligned}$$

where c_1' (resp. c_2' , c_3') is the derivative of c_1 (resp. c_2 , c_3) with respect to r . Straightforward simplifications, we have

$$(5.3) \quad s\sqrt{r^2 - s^2}A(r) + (r^2 - s^2)B(r) = 0,$$

where

$$\begin{aligned} A(r) &:= r^2(4c_0 c_1 c_3 r^5 + 2(2c_0 c_3 + 2c_1^2 c_3 + 4c_1 c_3^2 - c_0 c_1)r^3 + 2(c_1 c_3' - c_1' c_3)r^2 \\ & \quad + 2(4c_3^2 - c_0 - c_1^2)r + 2c_3' + c_1') \\ B(r) &:= 4c_0 c_2 c_3 r^5 + 2c_2(2c_1 c_3 + 4c_3^2 - c_0)r^3 + 4(c_2 c_3' - c_2' c_3)r^2 + 2c_2(2c_3 - c_1)r + 2c_2'. \end{aligned}$$

Since the equation (5.3) is satisfied for all s , then we must have

$$A(r) = 0, \quad B(r) = 0.$$

Without loss of generality, we can assume that c_1 and c_3 are arbitrary functions in r and solve $A(r) = 0$ algebraically for c_0 we have

$$(5.4) \quad c_0 = -\frac{4c_1 c_3(2c_3 + c_1)r^3 - 2(c_1' c_3 - c_1 c_3')r^2 + 2(4c_3^2 - c_1^2)r + c_1' + 2c_3'}{2r(c_1 r^2 + 1)(2c_3 r^2 - 1)}.$$

By substituting c_0 into $B(r) = 0$ and solving it for c_2 we obtain that

$$(5.5) \quad c_2 = c\sqrt{(c_1 r^2 + 1)(2c_3 r^2 - 1)},$$

where c is an arbitrary real constant. So if $c_1 = -\frac{1}{r^2}$ or $c_3 = \frac{1}{2r^2}$, then $c_2 = 0$ and this implies that the spray coefficients are quadratic and hence the space is Berwaldian. Moreover, according to Proposition 3.1, we have to exclude the case where ϕ satisfies the equation

$$\phi - s\phi_s + (r^2 - s^2)\phi_{ss} = 0.$$

Moreover, dividing the above equation by ϕ , we have

$$(5.6) \quad 1 - \frac{s\phi_s}{\phi} + (r^2 - s^2)\frac{\phi_{ss}}{\phi} = 0.$$

By using the property that $\left(\frac{\phi_s}{\phi}\right)_s = \frac{\phi_{ss}}{\phi} - \left(\frac{\phi_s}{\phi}\right)^2$ and substituting from (5.2) into (5.6), we get that

$$(5.7) \quad -\frac{r^4(c_1 r^2 + 1)(2c_3 r^2 - 1)}{(r^2 + (c_1 + 2c_3)r^2s^2 - 2c_3r^4 + 2c_2s\sqrt{r^2 - s^2})^2} = 0.$$

That is, the choices $c_1 = -\frac{1}{r^2}$ or $c_3 = \frac{1}{2r^2}$ must be excluded since it implies the non-existence of the function Q . Also, the case where $c_1 + 2c_3 = 0$ implies that $c_2 = c\sqrt{-(2c_3 r^2 - 1)^2}$ which is a contradiction because c_2 is a real-valued function of r . That is, we have the conditions

$$(5.8) \quad c_1 \neq -\frac{1}{r^2}, \quad c_3 \neq \frac{1}{2r^2}, \quad c_1 + 2c_3 \neq 0.$$

Taking the equations (5.4), (5.5) and (5.8) into account, one can find explicit examples of non-Berwaldian Landsberg metrics of dimension $n \geq 3$. This can be seen by the Examples 1 and 2. \square

Example 1. To find an explicit example of a non-Berwaldian Landsberg spherically symmetric metric, consider the functions c_1 and c_3 are given by

$$c_1 = \frac{1}{r^2}, \quad c_3 = \frac{1}{r^2}.$$

Then the functions c_0 and c_2 are calculated by the help of (5.4) and (5.5) as follows

$$c_0 = -\frac{3}{r^4}, \quad c_2 = \sqrt{2}c = \frac{1}{2},$$

where we choose the constant $c = \frac{1}{2\sqrt{2}}$. By substituting the functions c_0, c_1, c_2 and c_3 into (5.2), we have

$$\frac{\phi_s}{\phi} = \frac{3s + \sqrt{r^2 - s^2}}{-r^2 + 3s^2 + s\sqrt{r^2 - s^2}}, \quad \frac{\phi_r}{\phi} = -\frac{r}{-r^2 + 3s^2 + s\sqrt{r^2 - s^2}}.$$

Solving the above equations for ϕ , we obtain that

$$(5.9) \quad \phi(r, s) = (5s^2 - r^2)^{\frac{1}{3}}(2s^2 - r^2)^{\frac{1}{6}} \exp \left(\frac{1}{3} \operatorname{arctanh} \left(\frac{\sqrt{5}(\sqrt{5}s + 5r)}{10\sqrt{r^2 - s^2}} \right) - \frac{1}{6} \operatorname{arctanh} \left(\frac{\sqrt{2}(\sqrt{2}s + 2r)}{2\sqrt{r^2 - s^2}} \right) + \frac{1}{3} \operatorname{arctanh} \left(\frac{\sqrt{5}(\sqrt{5}s - 5r)}{2\sqrt{r^2 - s^2}} \right) - \frac{1}{6} \operatorname{arctanh} \left(\frac{\sqrt{2}(\sqrt{2}s - 2r)}{2\sqrt{r^2 - s^2}} \right) \right).$$

Therefore, the metric $F = u\phi(r, s)$ is a non-Berwaldian Landsberg metric with the geodesic spray obtained by the the functions P and Q (3.5) and ϕ is given by (5.9).

Another example can be obtained as the following.

Example 2. Let the functions c_1 and c_3 be given by

$$c_1 = 0, \quad c_3 = \frac{1}{r^2}.$$

Then the functions c_0 and c_2 are calculated the help of (5.4) and (5.5) as follows

$$c_0 = -\frac{2}{r^4}, \quad c_2 = c = \frac{1}{2},$$

where we choose the constant $c = \frac{1}{2}$. By substituting the functions c_0, c_1, c_2 and c_3 into (5.2), we have

$$(5.10) \quad \frac{\phi_s}{\phi} = \frac{2s + \sqrt{r^2 - s^2}}{-r^2 + 2s^2 + s\sqrt{r^2 - s^2}}, \quad \frac{\phi_r}{\phi} = \frac{-2s^2 - s\sqrt{r^2 - s^2}}{r(-r^2 + 2s^2 + s\sqrt{r^2 - s^2})}.$$

Solving the first equation for ϕ by integrating with respect to s , we have

$$\phi(r, s) = a(r)(r^4 - 5r^2s^2 + 5s^4)^{\frac{1}{4}} \left(\frac{A}{B} \right)^{\frac{5-\sqrt{5}}{20}} \left(\frac{C}{D} \right)^{\frac{5+\sqrt{5}}{20}} \exp \left(-\frac{\sqrt{5}}{10} \operatorname{arctanh} \left(\frac{\sqrt{5}(r^2 - 2s^2)}{r^2} \right) \right),$$

where

$$\begin{aligned} A &:= \left(2\sqrt{r^2 - s^2} - 2r + (\sqrt{5} - 1)s \right) \sqrt{r^2 - s^2} - (\sqrt{5} - 1)rs, \\ B &:= \left(2\sqrt{r^2 - s^2} - 2r - (\sqrt{5} - 1)s \right) \sqrt{r^2 - s^2} + (\sqrt{5} - 1)rs, \\ C &:= \left(2\sqrt{r^2 - s^2} - 2r - (\sqrt{5} + 1)s \right) \sqrt{r^2 - s^2} + (\sqrt{5} + 1)rs, \\ D &:= \left(2\sqrt{r^2 - s^2} - 2r + (\sqrt{5} + 1)s \right) \sqrt{r^2 - s^2} - (\sqrt{5} + 1)rs. \end{aligned}$$

The function $a(r)$ that satisfies the second equation of (5.10) is $a(r) = \frac{1}{r}$. That is, the function ϕ is given by

$$(5.11) \quad \phi(r, s) = \frac{1}{r}(r^4 - 5r^2s^2 + 5s^4)^{\frac{1}{4}} \left(\frac{A}{B} \right)^{\frac{5-\sqrt{5}}{20}} \left(\frac{C}{D} \right)^{\frac{5+\sqrt{5}}{20}} \exp \left(-\frac{\sqrt{5}}{10} \operatorname{arctanh} \left(\frac{\sqrt{5}(r^2 - 2s^2)}{r^2} \right) \right).$$

Therefore, the metric $F = u\phi(r, s)$ is a non-Berwaldian Landsberg metric with the geodesic spray obtained by the functions P and Q and ϕ is given by (5.11).

We end this section by the following two theorems.

Theorem 5.4. *All Berwaldian spherically symmetric metrics of dimension $n \geq 3$ are Riemannian.*

Proof. Let F be a Berwald spherically symmetric of dimension $n \geq 3$. Since every Berwald metric is Landsbergian, then the geodesic spray of F is given by the functions P and Q (5.1);

$$P = c_1 s + \frac{c_2}{r^2} \sqrt{r^2 - s^2}, \quad Q = \frac{1}{2} c_0 s^2 - \frac{c_2 s}{r^4} \sqrt{r^2 - s^2} + c_3.$$

Moreover, since F is Berwaldian then the mean curvature E_{ij} (3.8) vanishes. Substituting by

$$P - sP_s = \frac{c_2}{\sqrt{r^2 - s^2}}, \quad Q_s - sQ_{ss} = -\frac{c_2}{(r^2 - s^2)^{3/2}},$$

$$P_{ss} = -\frac{c_2}{(r^2 - s^2)^{3/2}}, \quad Q_{sss} = \frac{3c_2}{(r^2 - s^2)^{5/2}}$$

into the equation $E_{ij} = 0$ implies

$$\frac{nc_2}{u\sqrt{r^2 - s^2}} \left(\delta_{ij} - \frac{r^2}{u^2(r^2 - s^2)} \delta_{ih} \delta_{jk} y^i y^j - \frac{1}{r^2 - s^2} \delta_{ih} \delta_{jk} x^i x^j + \frac{s}{u(r^2 - s^2)} (\delta_{ih} \delta_{jk} + \delta_{ik} \delta_{jh}) x^h y^k \right) = 0.$$

Contracting the above equation by δ^{ij} and using the facts that $\delta_{ij} y^i y^j = u^2$, $\delta_{ij} x^i x^j = r^2$ and $\delta_{ij} x^i y^j = \langle x, y \rangle$, we have

$$n(n-2)c_2 \sqrt{r^2 - s^2} = 0.$$

From which together with the facts that $n \geq 3$ and the equation holds for all r and s , we must have $c_2 = 0$. That is, we have

$$P = c_1 s, \quad Q = \frac{1}{2} c_0 s^2 + c_3.$$

Now, the equation (5.2) becomes

$$(5.12) \quad \begin{aligned} \frac{\phi_s}{\phi} &= \frac{(c_1 + 2c_3)s}{1 + (c_1 + 2c_3)s^2 - 2c_3r^2}, \\ \frac{\phi_r}{\phi} &= \frac{c_0 c_1 r^6 s^2 + (c_0 + 4c_1 c_3 + 2c_1^2)r^4 s^2 - 2c_1 c_3 r^6 + c_1 r^4}{r(r^2 + (c_1 + 2c_3)r^2 s^2 - 2c_3 r^4)}, \end{aligned}$$

where c_0 is given by (5.4). Integrating $\frac{\phi_s}{\phi}$ with respect to s implies

$$\phi = a(r) \sqrt{(c_1 + 2c_3)s^2 - 2c_3r^2 + 1},$$

where $a(r)$ is to be chosen such that both formulae of (5.12) are satisfied, that is, calculating $\frac{\phi_r}{\phi}$ and equating it with the second formula of (5.12) we obtain the formula of $a(r)$ as follows

$$a(r) = \exp \int \frac{r(2c_1 c_3 r^2 - c_3' r - c_1 - 2c_3)}{2c_3 r^2 - 1} dr.$$

Consequently, the metric $F = u\phi = a(r)u\sqrt{(c_1 + 2c_3)s^2 - 2c_3r^2 + 1}$ is Riemannian. \square

Theorem 5.5. *All regular Landsberg spherically symmetric metrics of dimension $n \geq 3$ are Riemannian.*

Proof. Let $F = u\phi$ be a regular Landsberg spherically symmetric metric of dimension $n \geq 3$. Now, by the regularity condition (3.1) together with the fact that ϕ is positive, then we can write

$$1 - s \frac{\phi_s}{\phi} + (r^2 - s^2) \frac{\phi_{ss}}{\phi} > 0.$$

Moreover, by (5.7) we get

$$1 - s \frac{\phi_s}{\phi} + (r^2 - s^2) \frac{\phi_{ss}}{\phi} = -\frac{r^4(c_1 r^2 + 1)(2c_3 r^2 - 1)}{(r^2 + (c_1 + 2c_3)r^2 s^2 - 2c_3 r^4 + 2c_2 s \sqrt{r^2 - s^2})^2}.$$

By (5.5), we must have $(c_1 r^2 + 1)(2c_3 r^2 - 1) > 0$ and hence we have

$$\phi - s\phi_s + (r^2 - s^2)\phi_{ss} < 0$$

which is a contradiction and hence the metric is Berwaldian. Thus the result follows by Theorem 5.4. \square

6. THE TWO-DIMENSIONAL CASE

This section is devoted to the two-dimensional spherically symmetric Finsler metrics. We characterize all spherically symmetric surfaces of Berwald or Landsberg types and then we determine all Berwald surfaces. We end this section by a note on a class of surfaces mentioned in [12].

We start by the following proposition.

Proposition 6.1. *For a spherically symmetric surface the mean Berwald curvature E_{ij} and the scalar trace E of the mean Berwald curvature vanish if and only if $sH - (r^2 - s^2)H_s = 0$.*

Proof. Assume that E_{ij} vanishes then contracting (3.9) by δ^{ij} together with the fact that $n = 2$ imply

$$sH - (r^2 - s^2)H_s = 0.$$

Also, let the scalar E vanish then by Proposition 3.4, we have

$$(\rho_0 + \rho_3(r^2 - s^2))(sH - (r^2 - s^2)H_s) = 0.$$

By substituting by the formulae of ρ_0 and ρ_3 given in (3.3), we obtain that

$$\frac{sH - (r^2 - s^2)H_s}{\phi(\phi - s\phi_s + (r^2 - s^2)\phi_{ss})} = 0.$$

Therefore, $sH - (r^2 - s^2)H_s = 0$.

Conversely, assume that $sH - (r^2 - s^2)H_s = 0$, then by Proposition 3.4 it is clear that $E = 0$. Now, taking the fact that $r^2 - s^2 = \frac{(x_1y_2 - x_2y_1)^2}{u^2}$ into account, we can conclude that the components E_{11} , E_{12} and E_{22} vanish. For example, we have

$$E_{11} = \frac{y_2^2}{su^3}(su^2H - (x_1y_2 - x_2y_1)^2H_s) = \frac{y_2^2}{su}(sH - (r^2 - s^2)H_s) = 0.$$

Similarly, we can calculate E_{12} and E_{22} . Consequently, the proof is completed. \square

Proposition 6.2. *A spherically symmetric surface is Landsbergian if and only if $(r^2 - s^2)L_1 + 3L_2 = 0$.*

Proof. Assume that the surface is Landsbergian, then $L_{ijk} = 0$. Contracting (3.10) by $x^j\delta^{k\ell}$, we have

$$(r^2 - s^2)L_1 + 3L_2 = 0.$$

Conversely, assume that $(r^2 - s^2)L_1 + 3L_2 = 0$, one can see that the components $L_{ijk} = 0$. For example, we have

$$L_{111} = \frac{\phi}{2u^3}(-x_1^3u^3 + 3x_1^2y_1su^2 - 3x_1y_1^2s^2u + s^3y_1^3)L_1 + 3(y_1su^2 - x_1u^3 - sy_1^3 + x_1y_1^2u)L_2.$$

Using the formulae of u , r and s when $n = 2$, one can see that

$$L_{111} = \frac{y_2^3\phi}{2u^4}(x_1y_2 - x_2y_1)((r^2 - s^2)L_1 + 3L_2) = 0.$$

Similarly, we can calculate L_{112} , L_{122} and L_{222} . As required. \square

Rewriting the Landsberg condition for spherically symmetric surfaces and straightforward calculations imply that

$$(6.1) \quad (r^2 - s^2)L_1 + 3L_2 = \lambda_1\phi_s + \lambda_2\phi,$$

where

$$\lambda_1 := \frac{1}{s}(sH - (r^2 - s^2)H_s), \quad \lambda_2 := (r^2 - s^2)K_s - 3sK, \quad K := P_{ss} - Q_s + sQ_{ss}.$$

Now, we announce our first result in this section.

Theorem 6.3. *A spherically symmetric Finsler surface is Berwaldian if and only if*

$$P = b_1s + \frac{b_2}{\sqrt{r^2 - s^2}} + \frac{b_3(r^2 - 2s^2)}{\sqrt{r^2 - s^2}},$$

$$Q = b_0s^2 + \frac{1}{2}b_1 + \frac{b_2s(r^2 - 2s^2)}{r^4\sqrt{r^2 - s^2}} - \frac{b_3s(3r^2 - 2s^2)}{r^2\sqrt{r^2 - s^2}} - \frac{a}{r^2}s\sqrt{r^2 - s^2},$$

where a , b_0 , b_1 , b_2 , b_3 are arbitrary functions of r and to be chosen such that the compatibility conditions are satisfied.

Proof. Let F be a Berwald spherically symmetric Finsler surface, then the mean curvature E_{ij} and the scalar trace E vanish. Then, by Proposition 6.1, we have $sH - (r^2 - s^2)H_s = 0$ which has the solution

$$(6.2) \quad H = \frac{a_0(r)}{\sqrt{r^2 - s^2}}.$$

Moreover, since every Berwald metric is Landsbergian, then F is Landsbergian. Now, by Proposition 6.2, we have $(r^2 - s^2)L_1 + 3L_2 = 0$ and since $sH - (r^2 - s^2)H_s = 0$, then (6.1) implies

$$\lambda_2 = (r^2 - s^2)K_s - 3sK = 0.$$

That is, we have

$$\frac{K_s}{K} = \frac{P_{sss} + sQ_{sss}}{P_{ss} - Q_s + sQ_{ss}} = \frac{3s}{r^2 - s^2}.$$

Then, we have the solution

$$(6.3) \quad K = P_{ss} - Q_s + sQ_{ss} = \frac{b_0(r)}{(r^2 - s^2)^{3/2}}.$$

One can see that $P_{ss} - Q_s + sQ_{ss} = P_{ss} + (sQ_s)_s - 2Q_s$, then integrating the above equation with respect to s , we have

$$(6.4) \quad P_s + sQ_s - 2Q = \frac{b_0(r)s}{r^2\sqrt{r^2 - s^2}} + b(r).$$

Now, by using the definition of H together with the equations (6.2) and (6.3), we can write

$$(r^2 - s^2)P_{ss} - 3sP_s + 3P = \frac{a_1(r)}{\sqrt{r^2 - s^2}}.$$

The solution of the above equation can be written in the form

$$(6.5) \quad P = b_1s + \frac{b_2}{\sqrt{r^2 - s^2}} + \frac{b_3(r^2 - 2s^2)}{\sqrt{r^2 - s^2}},$$

where b_1, b_2, b_3 are arbitrary functions of r . To find the function Q , we rewrite (6.4) as follows

$$s^3 \left(\frac{Q}{s^2} \right)_s = -P_s + \frac{b_0(r)s}{r^2\sqrt{r^2 - s^2}}.$$

Hence, by integration with respect to s we can write the function Q as follows

$$(6.6) \quad Q = b_0s^2 + \frac{1}{2}b_1 + \frac{b_2s(r^2 - 2s^2)}{r^4\sqrt{r^2 - s^2}} - \frac{b_3s(3r^2 - 2s^2)}{r^2\sqrt{r^2 - s^2}} - \frac{a}{r^2}s\sqrt{r^2 - s^2},$$

where b_0, a are arbitrary functions of r .

Conversely, if P and Q are given by the formulae (6.5) and (6.6), then straightforward calculations imply that the Berwald curvature vanishes. Or instead, one can see that $(r^2 - s^2)L_1 + 3L_2 = 0$ and $sH - (r^2 - s^2)H_s = 0$ which means that the surface is Landsbergian and weakly Berwald ($E_{ij} = 0$). But it is known that a Landsbergian weakly Berwald surface is Berwaldian and hence the proof is completed. \square

It should be noted that, in [12], L. Zhou provided a class of Landsberg spherically symmetric surfaces and he claimed that this class is not Berwaldian. In fact, this class is Berwaldian as it is shown in [6]. Also, this can be seen easily since the corresponding functions P and Q are a special case of Theorem 6.3. Moreover, we are going to discuss another point of Zhou's class related to the compatibility conditions.

In what follows, to make the comparison easier and to avoid confusions, we use the terminology and notations of [12]. In the class of metrics mentioned in [12, Theorem 3.3] the functions ϕ, P and Q are not compatible, i.e., the compatibility conditions are not satisfied. We will provide an example showing this non-compatibility soon. The functions P and Q of the class provided in [12, Theorem 3.3] are given by

$$(6.7) \quad P = -\frac{s}{r^2} + \frac{c}{r^2}\sqrt{r^2 - s^2}, \quad Q = c_0 - \frac{4r^4c_0^2 + 2r^3c_0' + c^2}{2r^4(2r^2c_0 - 1)}s^2 - \frac{s}{r^4}\sqrt{r^2 - s^2},$$

where c_0 is function of r and $c \neq 3$ is a constant, c_0' denotes the derivative of c_0 with respect to r . By the formula of Q , it is clear that the case where $c_0 = \frac{1}{2r^2}$ is excluded.

Plugging the functions P and Q given by (6.7) into the compatibility conditions (4.1), then straightforward calculations by hand or by using Maple program, we have the following two equations:

$$\frac{\phi_s}{\phi}(r^2 - s^2)((2c_0r^2 - 1)\sqrt{r^2 - s^2} - (c + 1)s) + (2c_0r^2 - 1)s\sqrt{r^2 - s^2} + 2cr^2 - (c + 1)s^2 = 0,$$

$$\begin{aligned} & \frac{\phi_r}{\phi}(2c_0r^2 - 1)r^2\sqrt{r^2 - s^2} + \frac{\phi_s}{\phi}\left(\sqrt{r^2 - s^2}((r^2 - s^2)(4c_0^2sr^4 + sc^2) + (2c_0r^2 - 1)sr^2)\right. \\ & \left. + 3s^2r^2 + 2c_0cs^2r^4 + 2c_0r^6 + cr^4 - 2c_0cr^6 - cr^2s^2 - 6c_0s^2r^4 + 4c_0s^4r^2 - r^4 - 2s^4\right) \\ & + \sqrt{r^2 - s^2}(4c_0^2s^2r^4 + 2c_0r^4 + c^2s^2 - r^2) + 2c_0csr^4 - sr^2 + 2c_0sr^4 - csr^2 - 4c_0s^3r^2 + 2s^3 = 0. \end{aligned}$$

By solving the above equations for $\frac{\phi_s}{\phi}$ and $\frac{\phi_r}{\phi}$, we have

$$(6.8) \quad \begin{aligned} \frac{\phi_s}{\phi} &= -\frac{(c+1)s^2 - (2r^2c_0 - 1)s\sqrt{r^2 - s^2} - 2r^2c}{(r^2 - s^2)((2r^2c_0 - 1)\sqrt{r^2 - s^2} - (c+1)s)}, \\ \frac{\phi_r}{\phi} &= -\frac{\sqrt{r^2 - s^2}(2c(2c_0r^2 - 1)(1-c)(r^2 - s^2) - r^2 + 2s^2 + 4c_0r^4 + 8c_0^2r^4s^2 - 4c_0^2r^6 - 8c_0r^2s^2)}{r(r^2 - s^2)(2r^2c_0 - 1)((c+1)s - (2r^2c_0 - 1)\sqrt{r^2 - s^2})} \\ &+ \frac{5cr^2s - 2c^3r^2s - 4cc_0r^5s + 12cc_0r^2s^3 - 14cc_0r^4s + 4cc_0r^3s^3 + r^2s - 4cs^3 - 2c_0r^4s}{r(r^2 - s^2)(2r^2c_0 - 1)((c+1)s - (2r^2c_0 - 1)\sqrt{r^2 - s^2})} \\ &- \frac{4c_0r^2s^3 + 2c^3s^3 - 2s^3}{r(r^2 - s^2)(2r^2c_0 - 1)((c+1)s - (2r^2c_0 - 1)\sqrt{r^2 - s^2})}. \end{aligned}$$

Now, consider the following example.

Example 3. This example is constructed from [12, Theorem 3.3] and shows a non-compatibility between the functions P , Q and ϕ . Let $c = -1$ and $c_0 = \frac{1}{r^2}$, then, by substitution in the formula of $a(r)$, we have

$$a(r) = \exp\left(\int -\frac{2c_0r^2 - 1 + 2c^2 - 2c}{r(2c_0r^2 - 1)}dr\right) = \exp\left(\int -\frac{5}{r}dr\right) = \frac{1}{r^5}.$$

Now, we calculate F as follows

$$\begin{aligned} F &= u \exp\left(\int \frac{(c+1)s^2 - (2r^2c_0 - 1)s\sqrt{r^2 - s^2} - 2r^2c}{(r^2 - s^2)((2c_0r^2 - 1)\sqrt{r^2 - s^2} - (c+1)s)}ds\right) a(r) \\ &= \frac{u}{r^5} \exp\left(\int \frac{-s\sqrt{r^2 - s^2} + 2r^2}{(r^2 - s^2)(\sqrt{r^2 - s^2})}ds\right) \\ &= \frac{u}{r^5} \sqrt{r^2 - s^2} \exp\left(\frac{2s}{\sqrt{r^2 - s^2}}\right). \end{aligned}$$

Using (3.5) and the above formula of $\phi(r, s)$, we obtain that

$$(6.9) \quad P = -\frac{s}{r^2} - \frac{3}{4r^2} \sqrt{r^2 - s^2}, \quad Q = \frac{7}{8r^2} - \frac{3s^2}{r^4} - \frac{3s}{4r^4} \sqrt{r^2 - s^2}$$

On the other hand, by (6.7), the choice $c = -1$ and $c_0 = \frac{1}{r^2}$ leads to

$$(6.10) \quad P = -\frac{s}{r^2} - \frac{1}{r^2} \sqrt{r^2 - s^2}, \quad Q = \frac{1}{r^2} - \frac{s^2}{2r^4} - \frac{s}{r^4} \sqrt{r^2 - s^2}.$$

It is clear that the functions P and Q in the equations (6.9) and (6.10) are different. This is because the functions P , Q and ϕ do not satisfy the compatibility conditions, one can see that the compatibility conditions are $C_1 = 0$ and $C_2 = \frac{1}{r^2}\phi$.

The problem of the non-compatibility in the previous example comes from the factor $a(r)$. In [12], it is mentioned that the function $a(r)$ is given by $\phi(r, 0) = a(r)$ and it is not clear how this formula is obtained, in the above example, one can see that $\phi(r, 0) = \frac{1}{r^4} \neq a(r)$.

In the previous example, by substitution by $c = -1$ and $c_0 = \frac{1}{r^2}$ into (6.8), we have

$$\frac{\phi_s}{\phi} = -\frac{r^2s - s^3 - 2r^2\sqrt{r^2 - s^2}}{(r^2 - s^2)^2}, \quad \frac{\phi_r}{\phi} = -\frac{5r^4 - 11r^2s^2 + 6s^4 - 2r^2s\sqrt{r^2 - s^2}}{r(r^2 - s^2)^2}.$$

Solving the above equations, we get that

$$\phi = \frac{1}{r^6} \sqrt{r^2 - s^2} \exp\left(\frac{2s}{\sqrt{r^2 - s^2}}\right).$$

That is, $a(r) = \frac{1}{r^6}$. Moreover, one can see that the above formula of ϕ together with (3.5) yield the same formulae (6.10) of P and Q .

REFERENCES

- [1] G. S. Asanov, *Finsleroid-Finsler spaces of positive-definite and relativistic types*, Rep. Math. Phys., **58** (2006), 275–300.
- [2] D. Bao, *On two curvature-driven problems in Riemann-Finsler geometry*, Adv. Stud. Pure Math., **48** (2007), 19–71.
- [3] S. G. Elgendi, *On the problem of non-Berwaldian Landsberg spaces*, Bull. Aust. Math. Soc., **102**, (2020), 331–341.
- [4] S. G. Elgendi, *Solutions for the Landsberg unicorn problem in Finsler geometry*, J. Geom. Phys., **159**, (2021). arXiv:1908.10910 [math.DG].
- [5] S. G. Elgendi and L. Kozma, *(α, β) -metrics satisfying T -condition or σT -condition*, J. Geom. Anal. (2020).
- [6] S. G. Elgendi and Nabil L. Youssef, *A note on a result of L. Zhou's on Landsberg surfaces with $K = 0$ and $J = 0$* , Differ. Geom. Appl. **77** (2021). arXiv:2103.08550 [math.DG]
- [7] E. Guo and X. Mo, *The geometry of spherically symmetric Finsler manifolds*, Springer, 2018.
- [8] X. Mo and L. Zhou, *The curvatures of spherically symmetric Finsler metrics in \mathbb{R}^n* , arXiv: 1202.4543v4 [math.DG].
- [9] S. F. Rutz, *Symmetry in Finsler spaces*, Contemp. Math. **196** (1996), 289–300.
- [10] Z. Shen, *On a class of Landsberg metrics in Finsler geometry*, Canad. J. Math., **61** (2009), 1357-1374.
- [11] C. Yu, H. Zhu, *On a new class of Finsler metrics*, Differ. Geom. Appl. **29** (2011), 244–554.
- [12] L. Zhou, *The Finsler surface with $K = 0$ and $J = 0$* , Differ. Geom. Appl. **35** (2014),370-380. arXiv: 1202.4543v4 [math.DG].

S. G. ELGENDI, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, BENHA UNIVERSITY, EGYPT,
INSTITUTE OF MATHEMATICS, UNIVERSITY OF DEBRECEN, DEBRECEN, HUNGARY
Email address: salah.ali@fsci.bu.edu.eg, salahelgendi@yahoo.com.