

Quantum Rényi divergences and the strong converse exponent of state discrimination in operator algebras

Fumio Hiai¹ and Milán Mosonyi^{2,3}

¹ Graduate School of Information Sciences, Tohoku University,
Aoba-ku, Sendai 980-8579, Japan

² MTA-BME Lendület Quantum Information Theory Research Group

³ Mathematical Institute, Budapest University of Technology and Economics,
Egry József u 1., Budapest, 1111 Hungary

Abstract

The sandwiched Rényi α -divergences of two finite-dimensional quantum states play a distinguished role among the many quantum versions of Rényi divergences as the tight quantifiers of the trade-off between the two error probabilities in the strong converse domain of state discrimination. In this paper we show the same for the sandwiched Rényi divergences of two normal states on an injective von Neumann algebra, thereby establishing the operational significance of these quantities. Moreover, we show that in this setting, again similarly to the finite-dimensional case, the sandwiched Rényi divergences coincide with the regularized measured Rényi divergences, another distinctive feature of the former quantities. Our main tool is an approximation theorem (martingale convergence) for the sandwiched Rényi divergences, which may be used for the extension of various further results from the finite-dimensional to the von Neumann algebra setting.

We also initiate the study of the sandwiched Rényi divergences of pairs of states on a C^* -algebra, and show that the above operational interpretation, as well as the equality to the regularized measured Rényi divergence, holds more generally for pairs of states on a nuclear C^* -algebra.

2020 Mathematics Subject Classification: 81P45, 81P18, 94A17, 46L52, 46L53, 81R15, 62H15

Keywords and phrases: Quantum Rényi divergence, injective von Neumann algebra, nuclear C^* -algebra, martingale convergence, state discrimination, quantum hypothesis testing, strong converse exponent.

Contents

1	Introduction	2
2	Preliminaries	4
2.1	Haagerup's L^p -spaces and Kosaki's L^p -spaces	4
2.2	Generalized conditional expectations	5
2.3	Sandwiched Rényi divergences	6

¹*E-mail:* hiai.fumio@gmail.com

²*E-mail:* milan.mosonyi@gmail.com

3	The von Neumann algebra case	7
3.1	Martingale convergence for sandwiched Rényi divergences	7
3.2	The strong converse exponent in injective von Neumann algebras	10
4	The C^*-algebra case	17
4.1	Sandwiched and standard Rényi divergences in C^* -algebras	17
4.2	The strong converse exponent in nuclear C^* -algebras	22
A	Strong converse exponent in the finite-dimensional case	25
B	Boundary values of convex functions on $(0, 1)$	26
C	Proof of Theorem 4.3	27
	References	31

1 Introduction

Rényi's α -divergences [58] give a one-parameter family of pseudo-distances on probability measures, which play a central role in information theory as quantifiers of the trade-off between the relevant operational quantities in many information theoretic tasks; see, e.g., [13]. In quantum information theory, the non-commutativity of density operators allows infinitely many different extensions of the classical Rényi divergences to pairs of finite-dimensional quantum states; among others, the standard (or Petz-type) Rényi divergences [53], the sandwiched Rényi divergences [48, 68], their common generalization, the Rényi (α, z) -divergences [5, 32], or the maximal (or geometric) Rényi divergences [40, 55]. Many of these notions have also been extended to pairs of density operators on infinite-dimensional Hilbert spaces, or even to pairs of normal states on a von Neumann algebra [8, 24, 25, 26, 34, 35, 41, 52]. The study of these quantities has been motivating extensive research in the fields of matrix analysis and operator algebras.

On the other hand, in quantum information theory the relevant problem is to identify the quantum Rényi divergences with operational significance, i.e., those which appear as natural quantifiers of the trade-off relations between the quantities describing a problem, like error probabilities or coding rates. This has been established for the standard Rényi divergences with parameter values $\alpha \in (0, 1)$ in the context of binary state discrimination (hypothesis testing) of finite-dimensional quantum states in a series of works [4, 20, 21, 28, 42, 50, 51], and was also extended to the von Neumann algebra setting in [33]. Complementary to this, the sandwiched Rényi divergences were shown to have operational significance for the parameter values $\alpha > 1$ in the strong converse problem of binary state discrimination of finite-dimensional quantum states [21, 44, 45], and in the strong converse problem of classical-quantum channel coding [46, 47]. Apart from the standard and the sandwiched Rényi α -divergences mentioned above, no other quantum Rényi divergence has been shown to have a direct operational interpretation so far.

It is also a problem of central importance how much the distinguishability of two states, as measured by a quantum divergence, changes under quantum operations, in particular, under quantum measurements. While there is no known explicit formula for the optimal post-measurement Rényi α -divergence (called the measured Rényi divergence), it is known to be

strictly smaller than the standard Rényi α -divergence [7, 27]. Interestingly, if the measured Rényi α -divergence is evaluated on several copies of the states, and normalized by the number of copies, then the asymptotic limit of these quantities, called the regularized measured Rényi α -divergence, turns out to coincide with the sandwiched Rényi α -divergence for all $\alpha \in [1/2, +\infty)$. This is another feature distinguishing the sandwiched Rényi divergences among the multitude of different quantum Rényi divergences.

The proof of the hypothesis testing interpretation of the sandwiched Rényi divergences in [44] goes via replacing the quantum i.i.d. problem with a non-i.i.d. classical hypothesis testing problem by block-diagonalizing (pinching) large tensor powers of the first state by the spectral projections of the same tensor powers of the second state. It can be shown that the resulting classical problem has the same optimal error asymptotics as the original quantum problem, by using the pinching inequality [19] and the fact that the number of distinct eigen-values of the n th tensor power of a density operator grows only polynomially in n , even though the dimension of the underlying Hilbert space grows exponentially. The same technique can be used to show the equality of the regularized measured Rényi divergences and the sandwiched Rényi divergences [44]. While the pinching technique is very simple, it is also very powerful (see, e.g., [60] for further applications), but its applicability is obviously limited to the finite-dimensional case. Hence, even though the sandwiched Rényi divergences have been defined for pairs of normal states on a von Neumann algebra already some time ago [8, 34, 35], it has been an open problem (as proposed in [8]) whether they have an operational significance similar to the finite-dimensional case. This has been confirmed very recently in [41] in the simplest case where the von Neumann algebra is the space of bounded operators on an infinite-dimensional Hilbert space, using a finite-dimensional approximation technique, in particular, by showing that the Rényi divergences of the restrictions of the states to finite-dimensional subspaces converge to the Rényi divergences of the original states as the subspaces increase to the whole space.

In this paper we extend the above results about the sandwiched Rényi divergences to considerably more general settings, including injective, i.e., approximately finite-dimensional (AFD) von Neumann algebras. Our main tool is again finite-dimensional approximation. More generally, we show in Theorem 3.1 that the sandwiched Rényi divergences have the martingale convergence property for every $\alpha \in [1/2, +\infty) \setminus \{1\}$, i.e., if an increasing net of von Neumann subalgebras generates the whole algebra then the sandwiched Rényi α -divergences of the restrictions of two states converge to the sandwiched Rényi α -divergence of the original states. The proof is based on variational representations of the sandwiched Rényi divergences [26, 35], and the martingale convergence of generalized conditional expectations from [30]. Using this result, we show in Theorem 3.7 that the strong converse exponents of discriminating two normal states of an injective algebra are equal to their Hoeffding anti-divergences, analogously to the finite-dimensional case [44] and the case where the algebra is $\mathcal{B}(\mathcal{H})$ [41]. Based on this result, in Theorem 3.11 we give a direct operational representation of the sandwiched Rényi α -divergences as generalized cutoff rates, following Csiszár's approach [13]. Finally, using again the martingale convergence property, in Proposition 3.13 we show that for any $\alpha \in [1/2, +\infty) \setminus \{1\}$, the sandwiched Rényi α -divergence of two states on an injective algebra coincides with their regularized measured Rényi α -divergence.

Moreover, we also initiate the study of the sandwiched Rényi divergences for states of a C^* -algebra. In Theorem 4.3 we show that for any two states on a C^* -algebra, and any representation of the algebra that admits normal extensions of the states to the generated von Neumann algebra, the sandwiched Rényi α -divergence of the extensions is independent

of the specific representation for any $\alpha \in [1/2, +\infty)$, and hence it gives a well-defined notion of sandwiched Rényi α -divergence of the original states. We also show the same statement for the standard α -divergence and every $\alpha \in [0, +\infty) \setminus \{1\}$. In Proposition 4.5 we establish the basic properties of these extensions: joint lower semi-continuity, monotonicity under composition with unital positive maps (Schwarz maps in the case of the standard Rényi divergences), the inequality between the sandwiched and the standard Rényi divergences, and the martingale convergence property for both. In Theorem 4.12 we show that the sandwiched Rényi divergences on nuclear C^* -algebras have the same operational interpretation as in the case of injective von Neumann algebras, i.e., we show the equality of the strong converse exponents and the Hoeffding anti-divergences, from which the generalized cutoff rate representation also follows immediately. Finally, in Proposition 4.14 we show that the sandwiched Rényi divergences coincide with the regularized measured Rényi divergences on nuclear C^* -algebras.

The main text is accompanied by three appendices. In Appendix A, we fill a gap in the proof of the equality of the strong converse exponents and the Hoeffding anti-divergences in the finite-dimensional case [44], and show the same equality for a slightly modified definition of the strong converse exponent. Appendix B contains a simple observation about the boundary values of convex functions, needed in the proof of Theorem 3.7. Finally, Appendix C contains the rather technical proof of Theorem 4.3.

2 Preliminaries

2.1 Haagerup's L^p -spaces and Kosaki's L^p -spaces

Let \mathcal{M} be a general von Neumann algebra with the predual \mathcal{M}_* , and \mathcal{M}_*^+ be the positive part of \mathcal{M}_* consisting of normal positive linear functionals on \mathcal{M} .

For each $p \in (0, +\infty]$, Haagerup's L^p -space [65] is denoted by $L^p(\mathcal{M})$, and its positive part by $L^p(\mathcal{M})_+$. In particular, $L^\infty(\mathcal{M}) = \mathcal{M}$. There exists an order isomorphism $\mathcal{M}_* \cong L^1(\mathcal{M})$, given as $\psi \in \mathcal{M}_* \mapsto h_\psi \in L^1(\mathcal{M})$, so that $\text{tr}(h_\psi) := \psi(1)$, $\psi \in \mathcal{M}_*$, defines a positive linear functional tr on $L^1(\mathcal{M})$. For $1 \leq p < +\infty$ the L^p -norm $\|a\|_p$ of $a \in L^p(\mathcal{M})$ is given by $\|a\|_p := \text{tr}(|a|^p)^{1/p}$, and the L^∞ -norm $\|\cdot\|_\infty$ is the operator norm $\|\cdot\|$ on \mathcal{M} . For $1 \leq p < +\infty$, $L^p(\mathcal{M})$ is a Banach space with the norm $\|\cdot\|_p$, whose dual Banach space is $L^q(\mathcal{M})$, where $1/p + 1/q = 1$, by the duality

$$(a, b) \in L^p(\mathcal{M}) \times L^q(\mathcal{M}) \longmapsto \text{tr}(ab) (= \text{tr}(ba)).$$

In particular, $L^2(\mathcal{M})$ is a Hilbert space with the inner product $\langle a, b \rangle = \text{tr}(a^*b) (= \text{tr}(ba^*))$. Then

$$(\mathcal{M}, L^2(\mathcal{M}), J = *, L^2(\mathcal{M})_+)$$

becomes a *standard form* of \mathcal{M} [18], where \mathcal{M} is represented on $L^2(\mathcal{M})$ by the left multiplication. Each $\rho \in \mathcal{M}_*^+$ is represented as

$$\rho(x) = \text{tr}(xh_\rho) = \langle h_\rho^{1/2}, xh_\rho^{1/2} \rangle, \quad x \in \mathcal{M},$$

with the vector representative $h_\rho^{1/2} \in L^2(\mathcal{M})_+$. Note that the support projection $s(\rho) (\in \mathcal{M})$ of the functional ρ coincides with that of the operator h_ρ .

Assume that \mathcal{M} is σ -finite, and let a faithful $\omega \in \mathcal{M}_*^+$ be given with $h_\omega \in L^1(\mathcal{M})_+$. Consider an embedding \mathcal{M} into $L^1(\mathcal{M})$ by $x \mapsto h_\omega^{1/2} x h_\omega^{1/2}$. Defining $\|h_\omega^{1/2} x h_\omega^{1/2}\|_\infty := \|x\|$ on $h_\omega^{1/2} \mathcal{M} h_\omega^{1/2}$ we have a pair $(h_\omega^{1/2} \mathcal{M} h_\omega^{1/2}, L^1(\mathcal{M}))$ of compatible Banach spaces (see, e.g., [6]). For $1 < p < +\infty$ Kosaki's (symmetric) L^p -space $L^p(\mathcal{M}, \omega)$ with respect to ω is the complex interpolation space

$$C_{1/p}(h_\omega^{1/2} \mathcal{M} h_\omega^{1/2}, L^1(\mathcal{M}))$$

equipped with the interpolation norm $\|\cdot\|_{p,\omega}$ ($= \|\cdot\|_{C_{1/p}}$) [6]. Moreover, $L^1(\mathcal{M}, \omega) := L^1(\mathcal{M})$ with $\|\cdot\|_{1,\omega} = \|\cdot\|_1$ and $L^\infty(\mathcal{M}, \omega) := h_\omega^{1/2} \mathcal{M} h_\omega^{1/2}$ ($\cong \mathcal{M}$) with $\|\cdot\|_{\infty,\omega} = \|\cdot\|$. Kosaki's theorem [38, Theorem 9.1] says that for every $p \in [1, +\infty]$ and $1/p + 1/q = 1$,

$$L^p(\mathcal{M}, \omega) = h_\omega^{\frac{1}{2q}} L^p(\mathcal{M}) h_\omega^{\frac{1}{2q}} (\subseteq L^1(\mathcal{M})), \quad (2.1)$$

$$\|h_\omega^{\frac{1}{2q}} a h_\omega^{\frac{1}{2q}}\|_{p,\omega} = \|a\|_p, \quad a \in L^p(\mathcal{M}), \quad (2.2)$$

that is, $L^p(\mathcal{M}) \cong L^p(\mathcal{M}, \omega)$ by the isometry $a \mapsto h_\omega^{\frac{1}{2q}} a h_\omega^{\frac{1}{2q}}$. Interpolation L^p -spaces were introduced in [38] in terms of more general embedding $x \in \mathcal{M} \mapsto h_\omega^\eta x h_\omega^{1-\eta} \in L^1(\mathcal{M})$ with $0 \leq \eta \leq 1$. (The $\eta = 1/2$ case is the above symmetric $L^1(\mathcal{M}, \omega)$.) When \mathcal{M} is general and $\omega \in \mathcal{M}_*^+$ is not faithful with the support projection $e := s(\omega) \in \mathcal{M}$, Kosaki's L^p -space $L^p(\mathcal{M}, \omega)$ with respect to ω is still defined over $e\mathcal{M}e$ so that (2.1) and (2.2) hold with $eL^p(\mathcal{M})e$ in place of $L^p(\mathcal{M})$. Brief descriptions of interpolation L^p -spaces and Haagerup's $L^p(\mathcal{M})$ can be found in [26, Appendix A].

2.2 Generalized conditional expectations

Let \mathcal{M} and \mathcal{N} be von Neumann algebras, whose standard forms are $(\mathcal{M}, \mathcal{H}, J, \mathcal{P})$ and $(\mathcal{N}, \mathcal{H}_0, J_0, \mathcal{P}_0)$, respectively. Let $\Phi : \mathcal{N} \rightarrow \mathcal{M}$ be a unital normal positive map and $\omega \in \mathcal{M}_*^+$ be given. Assume that ω is faithful and $\omega \circ \Phi \in \mathcal{N}_*^+$ is also faithful. In this setting, it was shown in [1] that there exists a unique unital normal positive map $\Phi_\omega^* : \mathcal{M} \rightarrow \mathcal{N}$ such that

$$\langle Jx\Omega, \Phi(y)\Omega \rangle = \langle J_0\Phi_\omega^*(x)\Omega_0, y\Omega_0 \rangle, \quad x \in \mathcal{M}, y \in \mathcal{N}, \quad (2.3)$$

where $\Omega \in \mathcal{P}$ and $\Omega_0 \in \mathcal{P}_0$ are the vector representatives of ω and $\omega \circ \Phi$, respectively. We moreover have

$$\omega \circ \Phi \circ \Phi_\omega^* = \omega$$

and Φ_ω^* is completely positive if and only if so is Φ . This map Φ_ω^* is called the ω -dual map of Φ , or the *Petz recovery map* (see [54]), whose definition by (2.3) is independent of the choice of the standard forms of \mathcal{M}, \mathcal{N} . In terms of Haagerup's L^1 -elements h_ω and $h_{\omega \circ \Phi}$, we note [26, Lemma 8.3] that the map Φ_ω^* is determined by

$$\Phi_*(h_\omega^{1/2} x h_\omega^{1/2}) = h_{\omega \circ \Phi}^{1/2} \Phi_\omega^*(x) h_{\omega \circ \Phi}^{1/2}, \quad x \in \mathcal{M}, \quad (2.4)$$

where $\Phi_* : L^1(\mathcal{M}) \rightarrow L^1(\mathcal{N})$ is the predual map of Φ via $\mathcal{M}_* \cong L^1(\mathcal{M})$ and $\mathcal{N}_* \cong L^1(\mathcal{N})$, i.e., $\Phi_*(h_\psi) = h_{\psi \circ \Phi}$, $\psi \in \mathcal{M}_*$. Note that the construction of Φ_ω^* is possible even when ω and/or $\omega \circ \Phi$ are not faithful (see [26, Theorem 6.1 and Lemma 8.3]), though the above setting is sufficient for our present purpose.

In particular, let \mathcal{N} be a von Neumann subalgebra of \mathcal{M} containing the unit of \mathcal{M} , and $\omega \in \mathcal{M}_*^+$ be faithful. The ω -dual map Φ_ω^* of the injection $\Phi : \mathcal{N} \hookrightarrow \mathcal{M}$ is called the *generalized*

conditional expectation with respect to ω [1], which we denote by $\mathcal{E}_{\mathcal{N},\omega} : \mathcal{M} \rightarrow \mathcal{N}$. Note that $\mathcal{E}_{\mathcal{N},\omega}$ coincides with the genuine conditional expectation from \mathcal{M} onto \mathcal{N} with respect to ω if the latter exists (i.e., if \mathcal{N} is globally invariant under the modular automorphism group of \mathcal{M} with respect to ω [61]).

2.3 Sandwiched Rényi divergences

The notion of sandwiched Rényi divergences with parameter $\alpha \in [1/2, +\infty) \setminus \{1\}$, introduced first in [48, 68] for pairs of finite-dimensional density operators, was generalized by Berta–Scholz–Tomamichel [8] and Jenčová [34, 35] to the general von Neumann algebra setting. The definitions are different from each other between the three papers [8, 34, 35] but their equivalence was proved in [34, 35] (also [26, Sec. 3.3]). Here we work with the definition in [34] based on Kosaki’s L^p -spaces. Let $\rho, \sigma \in \mathcal{M}_*^+$ with representing $h_\rho, h_\sigma \in L^1(\mathcal{M})_+$. For every $\alpha \in (1, +\infty)$ define

$$Q_\alpha^*(\rho\|\sigma) := \begin{cases} \|h_\rho\|_{\alpha,\sigma}^\alpha & \text{if } h_\rho \in L^\alpha(\mathcal{M}, \sigma), \\ +\infty & \text{otherwise,} \end{cases}$$

where $L^\alpha(\mathcal{M}, \sigma)$ has been described in the last of Section 2.1. Note here that the condition $h_\rho \in L^\alpha(\mathcal{M}, \sigma)$ contains, in particular, $s(\rho) \leq s(\sigma)$. Then the *sandwiched Rényi α -divergence* $D_\alpha^*(\rho\|\sigma)$ is

$$D_\alpha^*(\rho\|\sigma) := \frac{1}{\alpha - 1} \log Q_\alpha^*(\rho\|\sigma).$$

On the other hand, for every $\alpha \in [1/2, 1)$, in view of [35, Theorem 3.1] and [26, Theorem 3.11], we define $D_\alpha^*(\rho\|\sigma)$ as

$$Q_\alpha^*(\rho\|\sigma) := \text{tr} \left(h_\sigma^{\frac{1-\alpha}{2\alpha}} h_\rho h_\sigma^{\frac{1-\alpha}{2\alpha}} \right)^\alpha = \|h_\sigma^{\frac{1-\alpha}{2\alpha}} h_\rho^{1/2}\|_{2\alpha}^{2\alpha}, \quad D_\alpha^*(\rho\|\sigma) := \frac{1}{\alpha - 1} \log Q_\alpha^*(\rho\|\sigma). \quad (2.5)$$

It is customary to define D_α^* with a normalization like

$$D_\alpha^*(\rho\|\sigma) = \frac{1}{\alpha - 1} \log \frac{Q_\alpha^*(\rho\|\sigma)}{\rho(1)}$$

(or with restricting ρ to states) but we use D_α^* without this normalization because that is the natural choice for the study of the strong converse exponent in Section 3.2; see, e.g., (3.7).

When ρ is a state, note that $\alpha \mapsto D_\alpha^*(\rho\|\sigma)$ is monotone increasing on $[1/2, 1) \cup (1, +\infty)$,

$$\lim_{\alpha \nearrow 1} D_\alpha^*(\rho\|\sigma) = D(\rho\|\sigma), \quad (2.6)$$

and if, in addition, $D_\alpha^*(\rho\|\sigma) < +\infty$ for some $\alpha > 1$, then

$$\lim_{\alpha \searrow 1} D_\alpha^*(\rho\|\sigma) = D(\rho\|\sigma), \quad (2.7)$$

where $D(\rho\|\sigma)$ is the *relative entropy* [2, 3, 39, 67]. Moreover, for any $\rho, \sigma \in \mathcal{M}_*^+$,

$$\lim_{\alpha \rightarrow +\infty} D_\alpha^*(\rho\|\sigma) = D_{\max}(\rho\|\sigma), \quad (2.8)$$

where

$$D_{\max}(\rho\|\sigma) := \log \min\{\Phi \geq 0 : \rho \leq \Phi\sigma\}$$

(= $+\infty$ if no such Φ exists), called the *max-relative entropy* [14, 57]. For these properties of D_α^* , see [48, 68] for the finite-dimensional case and [8, 34] (also a concise survey in [26, Sec. 3.3]) for the von Neumann algebra case.

The next variational formulas shown in [35, Proposition 3.4] and [26, Lemma 3.19] are the von Neumann algebra versions of [17, Lemma 4], which will play a crucial role in the next section. Here, \mathcal{M}_+ is the set of positive operators in \mathcal{M} and \mathcal{M}_{++} is the set of invertible $x \in \mathcal{M}_+$.

Proposition 2.1 ([26, 35]). *For any $\rho, \sigma \in \mathcal{M}_*^+$ the following hold:*

(i) *For every $\alpha \in (1, +\infty)$,*

$$Q_\alpha^*(\rho\|\sigma) = \sup_{x \in \mathcal{M}_+} \left[\alpha \rho(x) - (\alpha - 1) \text{tr} \left(h_\sigma^{\frac{\alpha-1}{2\alpha}} x h_\sigma^{\frac{\alpha-1}{2\alpha}} \right)^{\frac{\alpha}{\alpha-1}} \right]. \quad (2.9)$$

(ii) *For every $\alpha \in [1/2, 1)$,*

$$Q_\alpha^*(\rho\|\sigma) = \inf_{x \in \mathcal{M}_{++}} \left[\alpha \rho(x) + (1 - \alpha) \text{tr} \left(h_\sigma^{\frac{1-\alpha}{2\alpha}} x^{-1} h_\sigma^{\frac{1-\alpha}{2\alpha}} \right)^{\frac{\alpha}{1-\alpha}} \right]. \quad (2.10)$$

The older version of quantum Rényi divergences is the *standard* (or *Petz type*) *Rényi divergence* $D_\alpha(\rho\|\sigma)$ [37, 52], defined for every $\rho, \sigma \in \mathcal{M}_*^+$ and any $\alpha \in [0, +\infty) \setminus \{1\}$ in terms of the *relative modular operator* $\Delta_{\rho, \sigma}$ [3]. The following brief overview is based on [26]. When $0 \leq \alpha < 1$, note that $h_\sigma^{1/2} \in \mathcal{D}(\Delta_{\rho, \sigma}^{\alpha/2})$ and define

$$Q_\alpha(\rho\|\sigma) := \|\Delta_{\rho, \sigma}^{\alpha/2} h_\sigma^{1/2}\|^2.$$

When $\alpha > 1$,

$$Q_\alpha(\rho\|\sigma) := \begin{cases} \|\Delta_{\rho, \sigma}^{\alpha/2} h_\sigma^{1/2}\|^2 & \text{if } s(\rho) \leq s(\sigma) \text{ and } h_\sigma^{1/2} \in \mathcal{D}(\Delta_{\rho, \sigma}^{\alpha/2}), \\ +\infty & \text{otherwise.} \end{cases}$$

Then $D_\alpha(\rho\|\sigma)$ is given as

$$D_\alpha(\rho\|\sigma) := \frac{1}{\alpha - 1} \log Q_\alpha(\rho\|\sigma).$$

Properties of $D_\alpha(\rho\|\sigma)$ in the von Neumann algebra case were summarized in [24, Proposition 5.3], and a handy description of $D_\alpha(\rho\|\sigma)$ in terms of h_ρ, h_σ was given in [26, Theorem 3.6]; in particular, $D_\alpha(\rho\|\sigma) = \frac{1}{\alpha-1} \log \text{tr}(h_\rho^\alpha h_\sigma^{1-\alpha})$ when $\alpha \in [0, 1)$. (Compare this with (2.5).) Note ([8, Theorem 1.2], [34, Corollary 3.6]) that the inequality

$$D_\alpha^*(\rho\|\sigma) \leq D_\alpha(\rho\|\sigma) \quad (2.11)$$

holds for every $\rho, \sigma \in \mathcal{M}_*^+$ and any $\alpha \in [1/2, +\infty) \setminus \{1\}$, while equality holds here when ρ, σ “commute” (see [26, Remark 3.18 (2)] for the precise statement).

3 The von Neumann algebra case

3.1 Martingale convergence for sandwiched Rényi divergences

Let \mathcal{M} be a von Neumann algebra and $\{\mathcal{M}_i\}_{i \in \mathcal{I}}$ be an increasing net (on a directed set \mathcal{I}) of von Neumann subalgebras of \mathcal{M} containing the unit of \mathcal{M} , such that \mathcal{M} is generated by

$\bigcup_{i \in \mathcal{I}} \mathcal{M}_i$, i.e.,

$$\mathcal{M} = \left(\bigcup_{i \in \mathcal{I}} \mathcal{M}_i \right)''.$$

Let $\rho, \sigma \in \mathcal{M}_*^+$ and $\rho_i := \rho|_{\mathcal{M}_i}$, $\sigma_i := \sigma|_{\mathcal{M}_i}$ for each $i \in \mathcal{I}$.

The next theorem provides the *martingale convergence* for the sandwiched Rényi divergence D_α^* . It will play an essential role repeatedly in our later discussions.

Theorem 3.1. *Let $\{\mathcal{M}_i\}_{i \in \mathcal{I}}$ be as stated above, and assume that \mathcal{M} is σ -finite. Then for every $\rho, \sigma \in \mathcal{M}_*^+$ and for every $\alpha \in [1/2, +\infty) \setminus \{1\}$ we have*

$$D_\alpha^*(\rho\|\sigma) = \lim_i D_\alpha^*(\rho_i\|\sigma_i) \quad \text{increasingly.} \quad (3.1)$$

To prove the theorem, we first give two lemmas. First, we state the martingale convergence for the generalized conditional expectations in [30, Theorem 3] as a lemma, which was given in [30] in a slightly more general setting.

Lemma 3.2 ([30]). *In the situation stated above, assume that σ is faithful and for each $i \in \mathcal{I}$ let $\mathcal{E}_{\mathcal{M}_i, \sigma} : \mathcal{M} \rightarrow \mathcal{M}_i$ be the generalized conditional expectation with respect to σ (see Section 2.2). Then for every $x \in \mathcal{M}$ we have $\mathcal{E}_{\mathcal{M}_i, \sigma}(x) \rightarrow x$ strongly.*

The next lemma is indeed a special case of [34, Proposition 3.12]. The argument is also found in [36, Sec. 2.7]. We supply a sketchy proof for the convenience of the reader.

Lemma 3.3 ([34]). *Assume that $\sigma \in \mathcal{M}_*^+$ is faithful. Let \mathcal{N} be a von Neumann subalgebra of \mathcal{M} containing the unit of \mathcal{M} , and $\sigma_0 := \sigma|_{\mathcal{N}}$. Let $\mathcal{E}_{\mathcal{N}, \sigma} : \mathcal{M} \rightarrow \mathcal{N}$ be the generalized conditional expectation with respect to σ . Then for every $p \in [1, +\infty)$ and $x \in \mathcal{M}_+$ we have*

$$\mathrm{tr}(h_{\sigma_0}^{\frac{1}{2p}} \mathcal{E}_{\mathcal{N}, \sigma}(x) h_{\sigma_0}^{\frac{1}{2p}})^p \leq \mathrm{tr}(h_\sigma^{\frac{1}{2p}} x h_\sigma^{\frac{1}{2p}})^p,$$

where h_{σ_0} is the element of $L^1(\mathcal{N})_+$ corresponding to σ_0 , and $h_\sigma \in L^1(\mathcal{M})_+$ corresponds to σ .

Proof (sketch). We utilize Kosaki's (symmetric) interpolation L^p -space $L^p(\mathcal{M}, \sigma)$ with the norm $\|\cdot\|_{p, \sigma}$ for $p \in [1, +\infty]$; see Section 2.1. Let $\Psi : L^1(\mathcal{M}) \rightarrow L^1(\mathcal{N})$ be the predual map of the injection $\mathcal{N} \hookrightarrow \mathcal{M}$. In the present situation, the description of Φ_ω^* in (2.4) shows that Ψ restricted to $L^\infty(\mathcal{M}, \sigma)$ is given as

$$\Psi(h_\sigma^{1/2} x h_\sigma^{1/2}) = h_{\sigma_0}^{1/2} \mathcal{E}_{\mathcal{N}, \sigma}(x) h_{\sigma_0}^{1/2}, \quad x \in \mathcal{M},$$

that is, the map $\mathcal{E}_{\mathcal{N}, \sigma}$ coincides with $\Phi_\sigma : \mathcal{M} \rightarrow \mathcal{N}$ given in [34]. Hence Ψ is contractive from $L^\infty(\mathcal{M}, \sigma)$ to $L^\infty(\mathcal{N}, \sigma_0)$ with respect to $\|\cdot\|_{\infty, \sigma}$. It follows from the complex interpolation method (the Riesz–Thorin theorem) that Ψ is a contraction from $L^p(\mathcal{M}, \sigma)$ to $L^p(\mathcal{N}, \sigma_0)$ with respect to $\|\cdot\|_{p, \sigma}$ for any $p \in (1, +\infty)$. For $h_\sigma^{1/2} x h_\sigma^{1/2} \in L^\infty(\mathcal{M}, \sigma) \subseteq L^p(\mathcal{M}, \sigma)$ we have

$$\|h_{\sigma_0}^{1/2} \mathcal{E}_{\mathcal{N}, \sigma}(x) h_{\sigma_0}^{1/2}\|_{p, \sigma} \leq \|h_\sigma^{1/2} x h_\sigma^{1/2}\|_{p, \sigma}.$$

Noting by (2.1) and (2.2) that

$$\|h_\sigma^{1/2} x h_\sigma^{1/2}\|_{p, \sigma} = \|h_\sigma^{\frac{1}{2p}} x h_\sigma^{\frac{1}{2p}}\|_p = \left[\mathrm{tr}(h_\sigma^{\frac{1}{2p}} x h_\sigma^{\frac{1}{2p}})^p \right]^{1/p}$$

and similarly for $\|h_{\sigma_0}^{1/2} \mathcal{E}_{\mathcal{N},\sigma}(x) h_{\sigma_0}^{1/2}\|_{p,\sigma}$, we have

$$\mathrm{tr}\left(h_{\sigma_0}^{\frac{1}{2p}} \mathcal{E}_{\mathcal{N},\sigma}(x) h_{\sigma_0}^{\frac{1}{2p}}\right)^p \leq \mathrm{tr}\left(h_{\sigma}^{\frac{1}{2p}} x h_{\sigma}^{\frac{1}{2p}}\right)^p,$$

as desired. \square

Proof of Theorem 3.1. From the monotonicity property of D_{α}^* proved in [8, 34] it follows that $D_{\alpha}^*(\rho_i|\sigma_i) \leq D_{\alpha}^*(\rho|\sigma)$ and $i \in \mathcal{I} \mapsto D_{\alpha}^*(\rho_i|\sigma_i)$ is increasing. Hence, to show (3.1), it suffices to prove that

$$D_{\alpha}^*(\rho|\sigma) \leq \sup_{i \in \mathcal{I}} D_{\alpha}^*(\rho_i|\sigma_i). \quad (3.2)$$

To do this, we may assume that σ is faithful. Indeed, assume that (3.2) has been shown when σ is faithful. For general $\sigma \in \mathcal{M}_*^+$, since \mathcal{M} is σ -finite, there exists a $\sigma_0 \in \mathcal{M}_*^+$ with $s(\sigma_0) = 1 - s(\sigma)$ and let $\sigma^{(n)} := \sigma + n^{-1}\sigma_0$, $\sigma_i^{(n)} := \sigma^{(n)}|_{\mathcal{M}_i}$. From the lower semi-continuity and the order relation of D_{α}^* (see [34], [26, Theorem 3.16]) it follows that

$$D_{\alpha}^*(\rho|\sigma) \leq \liminf_{n \rightarrow \infty} D_{\alpha}^*(\rho|\sigma^{(n)}) \leq \liminf_{n \rightarrow \infty} \sup_i D_{\alpha}^*(\rho_i|\sigma_i^{(n)}) \leq \sup_i D_{\alpha}^*(\rho_i|\sigma_i),$$

proving (3.2) for general σ . Below we assume the faithfulness of σ and divide the proof into two cases $1 < \alpha < +\infty$ and $1/2 \leq \alpha < 1$.

Case $1 < \alpha < +\infty$. We need to prove that

$$Q_{\alpha}^*(\rho|\sigma) \leq \sup_{i \in \mathcal{I}} Q_{\alpha}^*(\rho_i|\sigma_i). \quad (3.3)$$

For every $x \in \mathcal{M}_+$ and $i \in \mathcal{I}$, by Lemma 3.3 we have

$$\begin{aligned} & \alpha\rho(x) - (\alpha - 1)\mathrm{tr}\left(h_{\sigma}^{\frac{\alpha-1}{2\alpha}} x h_{\sigma}^{\frac{\alpha-1}{2\alpha}}\right)^{\frac{\alpha}{\alpha-1}} \\ & \leq \alpha\rho(x) - (\alpha - 1)\mathrm{tr}\left(h_{\sigma_i}^{\frac{\alpha-1}{2\alpha}} \mathcal{E}_{\mathcal{M}_i,\sigma}(x) h_{\sigma_i}^{\frac{\alpha-1}{2\alpha}}\right)^{\frac{\alpha}{\alpha-1}} \\ & = \alpha\rho(x - \mathcal{E}_{\mathcal{M}_i,\sigma}(x)) + \alpha\rho_i(\mathcal{E}_{\mathcal{M}_i,\sigma}(x)) - (\alpha - 1)\mathrm{tr}\left(h_{\sigma_i}^{\frac{\alpha-1}{2\alpha}} \mathcal{E}_{\mathcal{M}_i,\sigma}(x) h_{\sigma_i}^{\frac{\alpha-1}{2\alpha}}\right)^{\frac{\alpha}{\alpha-1}} \\ & \leq \alpha\rho(x - \mathcal{E}_{\mathcal{M}_i,\sigma}(x)) + \sup_{j \in \mathcal{I}} Q_{\alpha}^*(\rho_j|\sigma_j), \end{aligned}$$

where the last inequality is due to the variational formula (2.9). Lemma 3.2 gives

$$\alpha\rho(x) - (\alpha - 1)\mathrm{tr}\left(h_{\sigma}^{\frac{\alpha-1}{2\alpha}} x h_{\sigma}^{\frac{\alpha-1}{2\alpha}}\right)^{\frac{\alpha}{\alpha-1}} \leq \sup_{j \in \mathcal{I}} Q_{\alpha}^*(\rho_j|\sigma_j), \quad x \in M_+.$$

By (2.9) again we have (3.3).

Case $1/2 \leq \alpha < 1$. We need to prove that

$$Q_{\alpha}^*(\rho|\sigma) \geq \inf_{i \in \mathcal{I}} Q_{\alpha}^*(\rho_i|\sigma_i). \quad (3.4)$$

For every $x \in \mathcal{M}_{++}$ and $i \in \mathcal{I}$, by Lemma 3.3 we have

$$\begin{aligned} & \alpha\rho(x) + (1 - \alpha)\mathrm{tr}\left(h_{\sigma}^{\frac{1-\alpha}{2\alpha}} x^{-1} h_{\sigma}^{\frac{1-\alpha}{2\alpha}}\right)^{\frac{\alpha}{1-\alpha}} \\ & \geq \alpha\rho(x) + (1 - \alpha)\mathrm{tr}\left(h_{\sigma_i}^{\frac{1-\alpha}{2\alpha}} \mathcal{E}_{\mathcal{M}_i,\sigma}(x^{-1}) h_{\sigma_i}^{\frac{1-\alpha}{2\alpha}}\right)^{\frac{\alpha}{1-\alpha}} \\ & \geq \alpha\rho(x) + (1 - \alpha)\mathrm{tr}\left(h_{\sigma_i}^{\frac{1-\alpha}{2\alpha}} \mathcal{E}_{\mathcal{M}_i,\sigma}(x)^{-1} h_{\sigma_i}^{\frac{1-\alpha}{2\alpha}}\right)^{\frac{\alpha}{1-\alpha}} \\ & = \alpha\rho(x - \mathcal{E}_{\mathcal{M}_i,\sigma}(x)) + \alpha\rho_i(\mathcal{E}_{\mathcal{M}_i,\sigma}(x)) + (1 - \alpha)\mathrm{tr}\left(h_{\sigma_i}^{\frac{1-\alpha}{2\alpha}} \mathcal{E}_{\mathcal{M}_i,\sigma}(x)^{-1} h_{\sigma_i}^{\frac{1-\alpha}{2\alpha}}\right)^{\frac{\alpha}{1-\alpha}} \\ & \geq \alpha\rho(x - \mathcal{E}_{\mathcal{M}_i,\sigma}(x)) + \inf_{j \in \mathcal{I}} Q_{\alpha}^*(\rho_j|\sigma_j), \end{aligned}$$

where the second inequality above follows from the Jensen inequality $\mathcal{E}_{\mathcal{M}_i, \sigma}(x^{-1}) \geq \mathcal{E}_{\mathcal{M}_i, \sigma}(x)^{-1}$ (see [11, Corollary 2.3]), and the last inequality is due to (2.10). By Lemma 3.2 and (2.10) we have (3.4). \square

From Theorem 3.1 we can easily obtain the following martingale type convergence for D_α^* under the restriction to reduced subalgebras $e_i \mathcal{M} e_i$ with $e_i \nearrow 1$. See [41] for related results in the case $\mathcal{M} = \mathcal{B}(\mathcal{H})$.

Proposition 3.4. *Assume that \mathcal{M} is σ -finite. Let $\{e_i\}_{i \in \mathcal{I}}$ be an increasing net of projections in \mathcal{M} such that $e_i \nearrow 1$. Then for every $\rho, \sigma \in \mathcal{M}_*^+$ and every $\alpha \in [1/2, +\infty) \setminus \{1\}$ we have*

$$D_\alpha^*(\rho \|\sigma) = \lim_i D_\alpha^*(e_i \rho e_i \|\| e_i \sigma e_i) \quad \text{increasingly,}$$

where $e_i \rho e_i$ is the restriction of ρ to the reduced von Neumann algebra $e_i \mathcal{M} e_i$ and similarly for $e_i \sigma e_i$.

Proof. Let $\mathcal{M}_i := e_i \mathcal{M} e_i \oplus \mathbb{C}(1 - e_i)$; then $\{\mathcal{M}_i\}_{i \in \mathcal{I}}$ is an increasing net of von Neumann subalgebras of \mathcal{M} containing the unit of \mathcal{M} with $\mathcal{M} = (\bigcup_i \mathcal{M}_i)''$. Note that $\rho_i := \rho|_{\mathcal{M}_i} = e_i \rho e_i \oplus \rho(1 - e_i)$ and similarly for $\sigma_i := \sigma|_{\mathcal{M}_i}$. Hence by the definition of Q_α^* or by using the variational formulas in Proposition 2.1, for any $\alpha \in [1/2, +\infty) \setminus \{1\}$ we have

$$Q_\alpha^*(\rho_i \|\sigma_i) = Q_\alpha^*(e_i \rho e_i \|\| e_i \sigma e_i) + \rho(1 - e_i)^\alpha \sigma(1 - e_i)^{1-\alpha}.$$

Below we will give the proof for the case $\alpha > 1$ (that for $1/2 \leq \alpha < 1$ is similar by using the reverse monotonicity of Q_α^*). Let $i, j \in \mathcal{I}$ be such that $i \leq j$. Noting that $e_i \mathcal{M} e_i \oplus \mathbb{C}(e_j - e_i) \subseteq e_j \mathcal{M} e_j$, by the monotonicity of Q_α^* we have

$$\begin{aligned} Q_\alpha^*(e_j \rho e_j \|\| e_j \sigma e_j) &\geq Q_\alpha^*(e_i \rho e_i \|\| e_i \sigma e_i) + \rho(e_j - e_i)^\alpha \sigma(e_j - e_i)^{1-\alpha} \\ &\geq Q_\alpha^*(e_i \rho e_i \|\| e_i \sigma e_i). \end{aligned}$$

Hence, $i \in \mathcal{I} \mapsto Q_\alpha^*(e_i \rho e_i \|\| e_i \sigma e_i)$ is increasing. Since Theorem 3.1 implies that

$$Q_\alpha^*(\rho \|\sigma) = \lim_i [Q_\alpha^*(e_i \rho e_i \|\| e_i \sigma e_i) + \rho(1 - e_i)^\alpha \sigma(1 - e_i)^{1-\alpha}],$$

the assertion follows if we show the following:

- when $Q_\alpha^*(\rho \|\sigma) = +\infty$, $\lim_i Q_\alpha^*(e_i \rho e_i \|\| e_i \sigma e_i) = +\infty$,
- when $Q_\alpha^*(\rho \|\sigma) < +\infty$, $\lim_i \rho(1 - e_i)^\alpha \sigma(1 - e_i)^{1-\alpha} = 0$.

These can indeed be proved in a similar way to the proof of [24, Theorem 4.5] by taking $\rho(1 - e_i)^\alpha \sigma(1 - e_i)^{1-\alpha}$ and $\rho(e_j - e_i)^\alpha \sigma(e_j - e_i)^{1-\alpha}$ in place of $\rho(1 - e_i)f(\rho(1 - e_i)/\sigma(1 - e_i))$ and $\rho(e_j - e_i)f(\rho(e_j - e_i)/\sigma(e_j - e_i))$, respectively, there. The details are omitted here. \square

3.2 The strong converse exponent in injective von Neumann algebras

Let \mathcal{M} be a von Neumann algebra and $\rho, \sigma \in \mathcal{M}_*^+$ be non-zero. For each $n \in \mathbb{N}$ let $\mathcal{M}^{\overline{\otimes} n}$ be the n -fold von Neumann algebra tensor product of \mathcal{M} , and $\rho_n := \rho^{\otimes n}$ (resp., $\sigma_n := \sigma^{\otimes n}$) be the n -fold tensor product of ρ (resp., σ), which are elements of $(\mathcal{M}^{\overline{\otimes} n})_*^+$; see [62, Chap. IV].

In this section we consider the simple hypothesis testing problem for the *null hypothesis* $H_0 : \rho$ versus the *alternative hypothesis* $H_1 : \sigma$, and extend the result [44] on the strong converse exponent to the injective von Neumann algebra case. For a test $T_n \in \mathcal{M}^{\otimes n}$ with $0 \leq T_n \leq 1$, $\rho_n(1 - T_n)$ and $\sigma_n(T_n)$ represent the *type I* and the *type II error probabilities*, respectively (hence $\rho_n(T_n)$ is the *type I success probability*) when ρ, σ are states. We begin by defining several forms of the strong converse exponents in the following:

Definition 3.5. For each type II error exponent $r \in \mathbb{R}$ we define the *strong converse exponents* of simple hypothesis testing for $H_0 : \rho$ vs. $H_1 : \sigma$ as follows:

$$\begin{aligned} \underline{sc}_r(\rho\|\sigma) &:= \inf_{\{T_n\}} \left\{ \liminf_{n \rightarrow \infty} -\frac{1}{n} \log \rho_n(T_n) : \liminf_{n \rightarrow \infty} -\frac{1}{n} \log \sigma_n(T_n) \geq r \right\}, \\ \overline{sc}_r(\rho\|\sigma) &:= \inf_{\{T_n\}} \left\{ \limsup_{n \rightarrow \infty} -\frac{1}{n} \log \rho_n(T_n) : \liminf_{n \rightarrow \infty} -\frac{1}{n} \log \sigma_n(T_n) \geq r \right\}, \\ sc_r(\rho\|\sigma) &:= \inf_{\{T_n\}} \left\{ \lim_{n \rightarrow \infty} -\frac{1}{n} \log \rho_n(T_n) : \liminf_{n \rightarrow \infty} -\frac{1}{n} \log \sigma_n(T_n) \geq r \right\}, \end{aligned}$$

where the infima are taken over all test sequences $\{T_n\}$ with $T_n \in \mathcal{M}^{\otimes n}$, $0 \leq T_n \leq 1$ ($n \in \mathbb{N}$) for which the indicated condition holds (and furthermore the limit exists for $sc_r(\rho\|\sigma)$). Also, we write $\underline{sc}_r^0(\rho\|\sigma)$, $\overline{sc}_r^0(\rho\|\sigma)$ and $sc_r^0(\rho\|\sigma)$ for the above infima when the condition $\liminf_{n \rightarrow \infty} -\frac{1}{n} \log \sigma_n(T_n) \geq r$ is replaced with $\liminf_{n \rightarrow \infty} -\frac{1}{n} \log \sigma_n(T_n) > r$.

For any $r \in \mathbb{R}$ it is obvious that

$$\begin{aligned} \underline{sc}_r(\rho\|\sigma) \leq \overline{sc}_r(\rho\|\sigma) \leq sc_r(\rho\|\sigma), \\ \underline{sc}_r^0(\rho\|\sigma) \leq \overline{sc}_r^0(\rho\|\sigma) \leq sc_r^0(\rho\|\sigma). \end{aligned} \tag{3.5}$$

The following is the definition of the Hoeffding anti-divergence [44, 41] in the von Neumann algebra setting.

Definition 3.6. For any $\rho, \sigma \in \mathcal{M}_*^+$ define

$$\begin{aligned} \psi^*(\rho\|\sigma|\alpha) &:= \log Q_\alpha^*(\rho\|\sigma) = (\alpha - 1)D_\alpha^*(\rho\|\sigma), \quad \alpha \in (1, +\infty), \\ \tilde{\psi}^*(\rho\|\sigma|u) &:= (1 - u)\psi^*(\rho\|\sigma|(1 - u)^{-1}), \quad u \in (0, 1). \end{aligned}$$

The *Hoeffding anti-divergence* of ρ and σ is then defined for each $r \in \mathbb{R}$ by

$$H_r^*(\rho\|\sigma) := \sup_{\alpha > 1} \frac{\alpha - 1}{\alpha} \{r - D_\alpha^*(\rho\|\sigma)\} = \sup_{u \in (0, 1)} \{ur - \tilde{\psi}^*(\rho\|\sigma|u)\}. \tag{3.6}$$

The aim of this section is to find whether the strong converse exponents in (3.5) are all equal to $H_r^*(\rho\|\sigma)$ for given r , as in the finite-dimensional case. To do so, we may assume without loss of generality that $\rho, \sigma \in \mathcal{M}_*^+$ are states. In fact, for any $\lambda, \mu > 0$ it is immediate to check that if $\varepsilon_r(\rho\|\sigma)$ is any strong converse exponent in (3.5), then

$$\varepsilon_r(\lambda\rho\|\mu\sigma) = \varepsilon_{r+\log \mu}(\rho\|\sigma) - \log \lambda.$$

On the other hand, since $Q_\alpha^*(\lambda\rho\|\mu\sigma) = \lambda^\alpha \mu^{1-\alpha} Q_\alpha^*(\rho\|\sigma)$, one easily sees that

$$H_r^*(\lambda\rho\|\mu\sigma) = H_{r+\log \mu}^*(\rho\|\sigma) - \log \lambda. \tag{3.7}$$

Thus, in the rest of the section, we will always assume that $\rho, \sigma \in \mathcal{M}_*^+$ are states. Under this assumption, when $r < 0$, any exponent in (3.5) is clearly equal to 0 by taking $T_n = 1$ for all n , while $H_r^*(\rho|\sigma)$ is also 0 whenever $D_\alpha^*(\rho|\sigma) < +\infty$ for some $\alpha > 1$, otherwise $H_r^*(\rho|\sigma) = -\infty$ for all $r \in \mathbb{R}$. Therefore, when $r < 0$, the conjecture holds in a trivial way, or otherwise it is not true. Furthermore, when $s(\rho) \not\leq s(\sigma)$, any exponent in (3.5) is 0 for all $r \in \mathbb{R}$ by taking $T_n = 1 - s(\sigma_n) = 1 - s(\sigma)^{\otimes n}$, while $H_r^*(\rho|\sigma) = -\infty$ for all r by definition. So the conjecture is not true for any r in this case. Therefore, we may restrict our consideration to the case where $r \geq 0$ and $s(\rho) \leq s(\sigma)$.

When \mathcal{M} is finite-dimensional and $s(\rho) \leq s(\sigma)$, it was proved in [44, Theorem 4.10] that $\underline{sc}_r(\rho|\sigma) = H_r^*(\rho|\sigma)$ for any $r \geq 0$. (Note that $\underline{sc}_r(\rho|\sigma)$ here was denoted by $B_e^*(r)$ in [44].) The result has recently been extended in [41, Theorem IV.5] to the infinite-dimensional $\mathcal{B}(\mathcal{H})$ setting in such a way that $\underline{sc}_r(\rho|\sigma) = \overline{sc}_r(\rho|\sigma) = H_r^*(\rho|\sigma)$ for any $r \in \mathbb{R}$ under the assumption that $D_\alpha^*(\rho|\sigma) < +\infty$ for some $\alpha > 1$. To further extend the result to the von Neumann algebra setting, let us next recall the fundamental notions of injectivity and AFD for von Neumann algebras.

A von Neumann algebra \mathcal{M} on a Hilbert space \mathcal{H} is *injective* if and only if there exists a (not necessarily normal) conditional expectation (i.e., a projection of norm one [66]) from $\mathcal{B}(\mathcal{H})$ onto \mathcal{M} ; see, e.g., [63, Corollary XV.1.3]. A fundamental result of Connes [12] (see also [63, Theorem XVI.1.9]) says that a von Neumann algebra \mathcal{M} of separable predual is injective if and only if \mathcal{M} is *AFD* (*approximately finite dimensional*), i.e., there exists an increasing sequence $\{\mathcal{M}_j\}_{j=1}^\infty$ of finite-dimensional *-subalgebras of \mathcal{M} such that $\mathcal{M} = (\bigcup_{j=1}^\infty \mathcal{M}_j)''$. In [16] the result was furthermore extended in such a way that a (general) von Neumann algebra \mathcal{M} is injective if and only if there is an increasing net $\{\mathcal{M}_i\}_{i \in \mathcal{I}}$ of finite-dimensional *-subalgebras of \mathcal{M} with $\mathcal{M} = (\bigcup_{i \in \mathcal{I}} \mathcal{M}_i)''$.

The main result of this section is the following:

Theorem 3.7. *Assume that \mathcal{M} is injective. Let $\rho, \sigma \in \mathcal{M}_*^+$ be states such that $D_\alpha^*(\rho|\sigma) < +\infty$ for some $\alpha > 1$ (in particular, $s(\rho) \leq s(\sigma)$). Then for every $r \in \mathbb{R}$ we have*

$$\underline{sc}_r(\rho|\sigma) = \overline{sc}_r(\rho|\sigma) = sc_r(\rho|\sigma) = \underline{sc}_r^0(\rho|\sigma) = \overline{sc}_r^0(\rho|\sigma) = sc_r^0(\rho|\sigma) = H_r^*(\rho|\sigma). \quad (3.8)$$

Proof. To prove this, we may and do assume that \mathcal{M} is σ -finite. Indeed, let $e := s(\rho) \vee s(\sigma)$. Then $e\mathcal{M}e$ is injective and all the quantities in (3.8) for any $r \in \mathbb{R}$ as well as $D_\alpha^*(\rho|\sigma)$ for any $\alpha \in [1/2, +\infty) \setminus \{1\}$ are left unchanged when ρ, σ are replaced with $\rho|_{e\mathcal{M}e}, \sigma|_{e\mathcal{M}e}$.

When $r < 0$, the quantities in (3.8) are all equal to 0 as explained in the paragraph after Definition 3.6. Hence we may assume that $r \geq 0$. By (3.5) it suffices to show the following two inequalities

$$\underline{sc}_r(\rho|\sigma) \geq H_r^*(\rho|\sigma), \quad (3.9)$$

$$sc_r^0(\rho|\sigma) \leq H_r^*(\rho|\sigma). \quad (3.10)$$

The inequality (3.9) follows in the same way (using Nagaoka's method [49]) as the proof of [44, Lemma 4.7] in the finite-dimensional case, as mentioned in [8, Sec. 5]. The injectivity assumption is unnecessary for this part. As for the inequality (3.10), we need to first prove it in the finite-dimensional case, because the exponent $sc_r^0(\rho|\sigma)$ was not treated in [44]. To make the proof of this part more understandable, we present it in Appendix A separately.

Now, by the injectivity assumption of \mathcal{M} , we can choose an increasing net $\{\mathcal{M}_i\}_{i \in \mathcal{I}}$ of finite-dimensional *-subalgebras containing the unit of \mathcal{M} such that $\mathcal{M} = (\bigcup_{i \in \mathcal{I}} \mathcal{M}_i)''$. For

each $i \in \mathcal{I}$ let $\rho_i := \rho|_{\mathcal{M}_i}$ and $\sigma_i := \sigma|_{\mathcal{M}_i}$. According to [44, Corollary 3.11], $\alpha \mapsto \psi^*(\rho_i|\sigma_i|\alpha)$ is a finite-valued convex function on $(1, +\infty)$, whence it is also continuous there. Thus, the function

$$u \mapsto \tilde{\psi}^*(\rho_i|\sigma_i|u) = (1-u)\psi^*(\rho_i|\sigma_i|(1-u)^{-1})$$

is also finite-valued, convex and continuous on $(0, 1)$. In particular, it can be extended to a convex and continuous function on $[0, 1]$ by

$$\tilde{\psi}^*(\rho_i|\sigma_i|0) := \lim_{u \searrow 0} \tilde{\psi}^*(\rho_i|\sigma_i|u) = 0, \quad \tilde{\psi}^*(\rho_i|\sigma_i|1) := \lim_{u \nearrow 1} \tilde{\psi}^*(\rho_i|\sigma_i|u) = D_{\max}(\rho_i|\sigma_i),$$

where the first equality is simple, and see (2.8) for the second one. We hence have

$$H_r^*(\rho_i|\sigma_i) = \sup_{u \in (0,1)} \{ur - \tilde{\psi}^*(\rho_i|\sigma_i|u)\} = \max_{u \in [0,1]} \{ur - \tilde{\psi}^*(\rho_i|\sigma_i|u)\}.$$

Also, from the monotonicity of D_α^* , note that $i \in \mathcal{I} \mapsto \psi^*(\rho_i|\sigma_i|\alpha)$ is increasing for any $\alpha > 1$, so that $i \in \mathcal{I} \mapsto ur - \tilde{\psi}^*(\rho_i|\sigma_i|u)$ is decreasing for any $u \in [0, 1]$. For every $r \geq 0$, since $sc_r^0(\rho|\sigma) \leq sc_r^0(\rho_i|\sigma_i)$ for all $i \in \mathcal{I}$ as immediately verified, we have

$$\begin{aligned} sc_r^0(\rho|\sigma) &\leq \inf_{i \in \mathcal{I}} sc_r^0(\rho_i|\sigma_i) = \inf_{i \in \mathcal{I}} H_r^*(\rho_i|\sigma_i) \\ &= \inf_{i \in \mathcal{I}} \max_{u \in [0,1]} \{ur - \tilde{\psi}^*(\rho_i|\sigma_i|u)\} \\ &= \max_{u \in [0,1]} \inf_{i \in \mathcal{I}} \{ur - \tilde{\psi}^*(\rho_i|\sigma_i|u)\} \\ &= \max_{u \in [0,1]} \left\{ ur - \sup_i \tilde{\psi}^*(\rho_i|\sigma_i|u) \right\}, \end{aligned} \quad (3.11)$$

where the above first equality is due to Proposition A.2 in Appendix A and the third equality is due to a minimax theorem in [41, Lemma II.3] (also see [42, Corollary A.2]).

Furthermore, Theorem 3.1 implies that

$$\sup_{i \in \mathcal{I}} \tilde{\psi}^*(\rho_i|\sigma_i|u) = \sup_{i \in \mathcal{I}} \frac{\alpha-1}{\alpha} D_\alpha^*(\rho_i|\sigma_i) = \frac{\alpha-1}{\alpha} D_\alpha^*(\rho|\sigma) = \tilde{\psi}^*(\rho|\sigma|u) \quad (3.12)$$

for every $u \in (0, 1)$ and $\alpha = (1-u)^{-1} \in (1, +\infty)$. By assumption, $\tilde{\psi}^*(\rho|\sigma|u) < +\infty$ for some $u \in (0, 1)$. Hence by Lemma B.1 we confirm that

$$\sup_{i \in \mathcal{I}} \tilde{\psi}^*(\rho_i|\sigma_i|0) = \lim_{u \searrow 0} \tilde{\psi}^*(\rho|\sigma|u), \quad \sup_{i \in \mathcal{I}} \tilde{\psi}^*(\rho_i|\sigma_i|1) = \lim_{u \nearrow 1} \tilde{\psi}^*(\rho|\sigma|1). \quad (3.13)$$

From (3.11)–(3.13) we conclude that

$$\begin{aligned} sc_r^0(\rho|\sigma) &\leq \sup_{u \in (0,1)} \left\{ ur - \tilde{\psi}^*(\rho|\sigma|u) \right\} \\ &= \sup_{\alpha > 1} \frac{\alpha-1}{\alpha} \{r - D_\alpha^*(\rho|\sigma)\} = H_r^*(\rho|\sigma), \end{aligned}$$

proving (3.10). \square

Remark 3.8. According to [41, Example III.39] (also [24, Remark 5.4(1)]), there exist commuting density operators ρ, σ on \mathcal{H} with $\dim \mathcal{H} = +\infty$ such that

$$D(\rho|\sigma) < +\infty, \quad D_\alpha^*(\rho|\sigma) = +\infty \quad \text{for all } \alpha \in (1, +\infty).$$

In this case, $H_r^*(\rho\|\sigma) = -\infty$ for all $r \in \mathbb{R}$, whereas any exponent in (3.5) is non-negative. See [41, Theorem IV.5] for a more precise result in this case. Therefore, the assumption of $D_\alpha^*(\rho\|\sigma) < +\infty$ for some $\alpha > 1$ is essential in Theorem 3.7, as well as in the convergence (2.7).

For given states $\rho, \sigma \in \mathcal{M}_*^+$ and any $r \geq 0$, the n th minimal type I error probability of Hoeffding type is defined as

$$\alpha_{e^{-nr}}^*(\rho_n\|\sigma_n) := \min_{0 \leq T_n \leq 1} \{\rho_n(1 - T_n) : \sigma_n(T_n) \leq e^{-nr}\}, \quad (3.14)$$

where the minimum is taken over all tests in $\mathcal{M}^{\otimes n}$ with $\sigma_n(T_n) \leq e^{-nr}$. Note that the minimum exists since the set of such tests is weakly compact. The n th maximal type I success probability is then given as

$$1 - \alpha_{e^{-nr}}^*(\rho_n\|\sigma_n) = \max_{0 \leq T_n \leq 1} \{\rho_n(T_n) : \sigma_n(T_n) \leq e^{-nr}\}.$$

In terms of this we can reformulate the assertion of Theorem 3.7 on the strong converse exponents as follows:

Theorem 3.9. *Under the same assumption as in Theorem 3.7, for every $r \geq 0$ we have*

$$\lim_{n \rightarrow \infty} -\frac{1}{n} \log \{1 - \alpha_{e^{-nr}}^*(\rho_n\|\sigma_n)\} = H_r^*(\rho\|\sigma).$$

Proof. For each n one can choose a test T_n in $\mathcal{M}^{\otimes n}$ such that $\sigma_n(T_n) \leq e^{-nr}$ and $\rho_n(T_n) = 1 - \alpha_{e^{-nr}}^*(\rho_n\|\sigma_n)$. Then it follows that

$$\liminf_{n \rightarrow \infty} -\frac{1}{n} \log \{1 - \alpha_{e^{-nr}}^*(\rho_n\|\sigma_n)\} = \liminf_{n \rightarrow \infty} -\frac{1}{n} \log \rho_n(T_n) \geq \underline{s}\mathcal{C}_r(\rho\|\sigma).$$

On the other hand, if a sequence of tests $\{T_n\}$ satisfies $\liminf_n -\frac{1}{n} \log \sigma_n(T_n) > r$, then we have $1 - \alpha_{e^{-nr}}^*(\rho_n\|\sigma_n) \geq \rho_n(T_n)$ for all sufficiently large n and hence

$$\limsup_{n \rightarrow \infty} -\frac{1}{n} \log \{1 - \alpha_{e^{-nr}}^*(\rho_n\|\sigma_n)\} \leq \limsup_{n \rightarrow \infty} -\frac{1}{n} \log \rho_n(T_n).$$

Therefore,

$$\limsup_{n \rightarrow \infty} -\frac{1}{n} \log \{1 - \alpha_{e^{-nr}}^*(\rho_n\|\sigma_n)\} \leq \overline{s}\mathcal{C}_r^0(\rho\|\sigma).$$

Since $\underline{s}\mathcal{C}_r(\rho\|\sigma) = \overline{s}\mathcal{C}_r^0(\rho\|\sigma) = H_r^*(\rho\|\sigma)$ by Theorem 3.7, we find that

$$\lim_{n \rightarrow \infty} -\frac{1}{n} \log \{1 - \alpha_{e^{-nr}}^*(\rho_n\|\sigma_n)\} = H_r^*(\rho\|\sigma),$$

as asserted. □

Using Theorem 3.7 we can give a direct operational interpretation of the sandwiched Rényi divergences as generalized cutoff rates, following the idea of Csiszár [13].

Definition 3.10. Let $\rho, \sigma \in \mathcal{M}_*^+$ be states, and $\kappa \in (0, 1)$. The *generalized κ -cutoff rate* $C_\kappa(\rho\|\sigma)$ is defined to be the infimum of all $r_0 \in \mathbb{R}$ such that $\underline{s}\mathcal{C}_r(\rho\|\sigma) \geq \kappa(r - r_0)$ holds for every $r \in \mathbb{R}$.

Theorem 3.11. *Assume that \mathcal{M} is injective. Let $\rho, \sigma \in \mathcal{M}_*^+$ be states such that $D_{\alpha_0}^*(\rho||\sigma) < +\infty$ for some $\alpha_0 > 1$. Then*

$$C_\kappa(\rho||\sigma) = D_{\frac{1}{1-\kappa}}^*(\rho||\sigma), \quad \text{or equivalently,} \quad D_\alpha^*(\rho||\sigma) = C_{\frac{\alpha-1}{\alpha}}(\rho||\sigma)$$

for every $\alpha \in (1, \alpha_0)$ and corresponding $\kappa = (\alpha - 1)/\alpha \in (0, \kappa_0)$, where $\kappa_0 := (\alpha_0 - 1)/\alpha_0$.

Proof. By Theorem 3.7 and (3.6),

$$\begin{aligned} \underline{s}C_r(\rho||\sigma) &= H_r^*(\rho||\sigma) = \sup_{u \in (0,1)} \{ur - \tilde{\psi}^*(\rho||\sigma|u)\} \\ &\geq \kappa r - \tilde{\psi}^*(\rho||\sigma|\kappa) = \kappa \left(r - D_{\frac{1}{1-\kappa}}^*(\rho||\sigma) \right), \end{aligned}$$

showing $C_\kappa(\rho||\sigma) \geq D_{\frac{1}{1-\kappa}}^*(\rho||\sigma)$. As we have seen in the proof of Theorem 3.7, $\tilde{\psi}^*(\rho_i||\sigma_i|\cdot)$ is convex and continuous on $(0, 1)$ for every i , and hence $\tilde{\psi}^*(\rho||\sigma|\cdot) = \sup_i \tilde{\psi}^*(\rho_i||\sigma_i|\cdot)$ is convex and lower semi-continuous on $(0, 1)$, where the equality is by Theorem 3.1. By assumption, and the monotonicity of D_α^* in α ([34, Proposition 3.7]), $\tilde{\psi}^*(\rho||\sigma|\cdot)$ is finite-valued on $(0, \kappa_0)$, and hence it has finite left- and right-derivatives at every $\kappa \in (0, \kappa_0)$. Thus, for any such κ and $r \in [\partial^- \tilde{\psi}^*(\rho||\sigma|\kappa), \partial^+ \tilde{\psi}^*(\rho||\sigma|\kappa)]$,

$$\begin{aligned} \overline{s}C_r(\rho||\sigma) &= H_r^*(\rho||\sigma) = \sup_{u \in (0,1)} \{ur - \tilde{\psi}^*(\rho||\sigma|u)\} \\ &= \kappa r - \tilde{\psi}^*(\rho||\sigma|\kappa) = \kappa \left(r - D_{\frac{1}{1-\kappa}}^*(\rho||\sigma) \right), \end{aligned}$$

where the first equality is again due to Theorem 3.7. This shows $C_\kappa(\rho||\sigma) \leq D_{\frac{1}{1-\kappa}}^*(\rho||\sigma)$, completing the proof. \square

In the rest of this section we consider the (regularized) measured Rényi divergences. Let us first recall these notions in the present setting. A measurement (POVM) in \mathcal{M} is given by a finite family $\mathfrak{M} = (M_j)_{1 \leq j \leq k}$ of $M_j \in \mathcal{M}_+$ ($j = 1, \dots, k$) such that $\sum_{j=1}^k M_j = 1$. For each $\rho \in \mathcal{M}_*^+$ let $\mathfrak{M}(\rho) := (\rho(M_j))_{1 \leq j \leq k}$ denote the post-measurement probability distribution on $\{1, \dots, k\}$.

Definition 3.12. The *measured Rényi divergence* $D_\alpha^{\text{meas}}(\rho||\sigma)$ and its *regularized version* $\overline{D}_\alpha^{\text{meas}}(\rho||\sigma)$ for $\alpha \in [1/2, +\infty) \setminus \{1\}$ are defined as

$$\begin{aligned} D_\alpha^{\text{meas}}(\rho||\sigma) &:= \sup \{D_\alpha(\mathfrak{M}(\rho)||\mathfrak{M}(\sigma)) : \mathfrak{M} \text{ a measurement in } M\}, \\ \overline{D}_\alpha^{\text{meas}}(\rho||\sigma) &:= \sup_{n \in \mathbb{N}} \frac{1}{n} D_\alpha^{\text{meas}}(\rho^{\otimes n}||\sigma^{\otimes n}). \end{aligned}$$

One might also consider more general notions of measurements in the definition, but that does not change the value of the (regularized) measured Rényi divergence, as was shown in [26, Proposition 5.2]. In the opposite direction, one might restrict measurements \mathfrak{M} to two-valued ones (tests), which leads to the notion of the *test-measured Rényi divergence* $D_\alpha^{\text{test}}(\rho||\sigma)$ and its regularized version $\overline{D}_\alpha^{\text{test}}(\rho||\sigma)$, defined as

$$\begin{aligned} D_\alpha^{\text{test}}(\rho||\sigma) &:= \sup_{T \in \mathcal{M}, 0 \leq T \leq 1} D_\alpha((\rho(T), \rho(1-T))||(\sigma(T), \sigma(1-T))), \\ \overline{D}_\alpha^{\text{test}}(\rho||\sigma) &:= \sup_{n \in \mathbb{N}} \frac{1}{n} D_\alpha^{\text{test}}(\rho^{\otimes n}||\sigma^{\otimes n}). \end{aligned}$$

It is obvious that

$$\begin{aligned} D_{\alpha}^{\text{test}}(\rho\|\sigma) &\leq \overline{D}_{\alpha}^{\text{test}}(\rho\|\sigma) \\ D_{\alpha}^{\text{meas}}(\rho\|\sigma) &\leq \overline{D}_{\alpha}^{\text{meas}}(\rho\|\sigma) \end{aligned} \quad (3.15)$$

for all $\alpha \in [1/2, +\infty) \setminus \{1\}$.

Let \mathcal{M} be injective, and $\{\mathcal{M}_i\}_{i \in \mathcal{I}}$ be an increasing net of finite-dimensional *-subalgebras of $\mathcal{M}_0 := e\mathcal{M}e$ with $\mathcal{M}_0 = (\bigcup_i \mathcal{M}_i)''$, where $e := s(\rho) \vee s(\sigma)$ (see the proof of Theorem 3.7). Note that for each $n \in \mathbb{N}$, $\mathcal{M}_0^{\otimes n} = (\bigcup_i \mathcal{M}_i^{\otimes n})''$, $(\rho|_{\mathcal{M}_0})^{\otimes n} = \rho^{\otimes n}|_{\mathcal{M}_0^{\otimes n}}$, and similarly for σ . Hence, for every $\alpha \in [1/2, +\infty) \setminus \{1\}$ and every $n \in \mathbb{N}$ one has by Theorem 3.1,

$$\begin{aligned} D_{\alpha}^*(\rho^{\otimes n}\|\sigma^{\otimes n}) &= \lim_i D_{\alpha}^*((\rho|_{\mathcal{M}_i})^{\otimes n}\|(\sigma|_{\mathcal{M}_i})^{\otimes n}) \\ &= \lim_i n D_{\alpha}^*(\rho|_{\mathcal{M}_i}\|\sigma|_{\mathcal{M}_i}) = n D_{\alpha}^*(\rho\|\sigma), \end{aligned} \quad (3.16)$$

where the second equality above follows from the additivity of D_{α}^* under tensor product in the finite-dimensional case. Indeed, the additivity of D_{α}^* under tensor product holds true in the general von Neumann algebra case, as observed in [8] (see the footnote of p. 1860) in the approach of Araki and Masuda's L^p -norms.

Proposition 3.13. *Assume that \mathcal{M} is injective. Then for every $\rho, \sigma \in \mathcal{M}_*^+$ and every $\alpha \in [1/2, +\infty) \setminus \{1\}$ we have*

$$D_{\alpha}^*(\rho\|\sigma) = \overline{D}_{\alpha}^{\text{meas}}(\rho\|\sigma) = \lim_{n \rightarrow \infty} \frac{1}{n} D_{\alpha}^{\text{meas}}(\rho^{\otimes n}\|\sigma^{\otimes n}), \quad (3.17)$$

and for every $\alpha > 1$,

$$D_{\alpha}^*(\rho\|\sigma) = \overline{D}_{\alpha}^{\text{test}}(\rho\|\sigma) = \lim_{n \rightarrow \infty} \frac{1}{n} D_{\alpha}^{\text{test}}(\rho^{\otimes n}\|\sigma^{\otimes n}). \quad (3.18)$$

Proof. For any $\alpha \in [1/2, +\infty) \setminus \{1\}$, by (3.16) and (3.15) note that

$$D_{\alpha}^*(\rho\|\sigma) \geq \overline{D}_{\alpha}^{\text{meas}}(\rho\|\sigma) \geq \overline{D}_{\alpha}^{\text{test}}(\rho\|\sigma).$$

To prove (3.17), it suffices to show that

$$\liminf_{n \rightarrow \infty} \frac{1}{n} D_{\alpha}^{\text{meas}}(\rho^{\otimes n}\|\sigma^{\otimes n}) \geq D_{\alpha}^*(\rho\|\sigma). \quad (3.19)$$

Let $e := s(\rho) \vee s(\sigma)$. Let $v < D_{\alpha}^*(\rho\|\sigma)$ be arbitrary. By Theorem 3.1 there exists a finite-dimensional *-subalgebra \mathcal{N}_0 of $e\mathcal{M}e$ such that $D_{\alpha}^*(\rho|_{\mathcal{N}_0}\|\sigma|_{\mathcal{N}_0}) > v$. Write $\rho_0 := \rho|_{\mathcal{N}_0}$ and $\sigma_0 := \sigma|_{\mathcal{N}_0}$. From [44, Theorem 3.7] and [21, Corollary 4] it follows that

$$D_{\alpha}^*(\rho_0\|\sigma_0) = \lim_{n \rightarrow \infty} \frac{1}{n} D_{\alpha}^{\text{meas}}(\rho_0^{\otimes n}\|\sigma_0^{\otimes n}).$$

Hence there exists an $n_0 \in \mathbb{N}$ such that $\frac{1}{n} D_{\alpha}^{\text{meas}}(\rho_0^{\otimes n}\|\sigma_0^{\otimes n}) > v$ for all $n \geq n_0$. Therefore,

$$\liminf_{n \rightarrow \infty} \frac{1}{n} D_{\alpha}^{\text{meas}}(\rho^{\otimes n}\|\sigma^{\otimes n}) \geq \liminf_{n \rightarrow \infty} \frac{1}{n} D_{\alpha}^{\text{meas}}(\rho_0^{\otimes n}\|\sigma_0^{\otimes n}) \geq v,$$

which implies (3.19) by letting $v \nearrow D_{\alpha}^*(\rho\|\sigma)$.

When $\alpha > 1$, the proof of (3.18) can proceed in the same way as above by appealing to [44, Corollary 4.6]. \square

Remark 3.14. The relation between $\overline{D}_{\alpha}^{\text{test}}(\rho\|\sigma)$ and $\overline{D}_{\alpha}^{\text{meas}}(\rho\|\sigma)$ for $\alpha < 1$ does not seem to have been considered even in the finite-dimensional case. We discuss it in detail in [43].

4 The C^* -algebra case

4.1 Sandwiched and standard Rényi divergences in C^* -algebras

In this section we first extend the notion of the sandwiched and the standard Rényi divergences to positive linear functionals on a general unital C^* -algebra. In the rest of the section let \mathcal{A} be a unital C^* -algebra. (The unitality assumption is not essential, since we can work with the unitization $\mathcal{A} \oplus \mathbb{C}$ if \mathcal{A} is non-unital.) Let \mathcal{A}_+^* be the set of positive linear functionals (automatically bounded) on \mathcal{A} . To define D_α^* and D_α for $\rho, \sigma \in \mathcal{A}_+^*$, let us consider the universal representation $\{\pi_u, \mathcal{H}_u\}$ of \mathcal{A} ; then ρ and σ have the respective normal extensions $\bar{\rho}$ and $\bar{\sigma}$ to $\pi_u(\mathcal{A})'' \cong \mathcal{A}^{**}$ such that $\rho = \bar{\rho} \circ \pi$ and $\sigma = \bar{\sigma} \circ \pi$; see, e.g., [62, Definition III.2.3, Theorem III.2.4].

Definition 4.1. For every $\rho, \sigma \in \mathcal{A}_+^*$ let $\bar{\rho}, \bar{\sigma}$ be as stated above. For every $\alpha \in [1/2, +\infty) \setminus \{1\}$ we define the *sandwiched Rényi α -divergence* of ρ and σ by

$$\widehat{D}_\alpha^*(\rho\|\sigma) := D_\alpha^*(\bar{\rho}\|\bar{\sigma}). \quad (4.1)$$

For $\alpha \in [0, +\infty) \setminus \{1\}$ define also the *standard Rényi α -divergence* of ρ and σ by

$$\widehat{D}_\alpha(\rho\|\sigma) := D_\alpha(\bar{\rho}\|\bar{\sigma}). \quad (4.2)$$

One question arises immediately: when \mathcal{A} itself is a von Neumann algebra and ρ, σ are normal functionals, do definitions (4.1) and (4.2) give the same notions of sandwiched and standard Rényi divergences (see Section 2.3 for the definitions), i.e., do the equalities

$$\widehat{D}_\alpha^*(\rho\|\sigma) = D_\alpha^*(\rho\|\sigma), \quad \widehat{D}_\alpha(\rho\|\sigma) = D_\alpha(\rho\|\sigma) \quad (4.3)$$

hold? Theorem 4.3 below shows that these are indeed the case. For stating it, let us introduce the following:

Definition 4.2. Let $\rho, \sigma \in \mathcal{A}_+^*$. We say that a representation $\pi : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ is (ρ, σ) -*normal*, if there exist $\rho_\pi, \sigma_\pi \in (\pi(\mathcal{A})'')_*^+$ such that $\rho = \rho_\pi \circ \pi$, $\sigma = \sigma_\pi \circ \pi$.

Theorem 4.3. *Let $\rho, \sigma \in \mathcal{A}_+^*$ and π be any (ρ, σ) -normal representation of \mathcal{A} . Then we have the following:*

(i) For any $\alpha \in [1/2, +\infty) \setminus \{1\}$,

$$\widehat{D}_\alpha^*(\rho\|\sigma) = D_\alpha^*(\rho_\pi\|\sigma_\pi), \quad i.e., \quad D_\alpha^*(\bar{\rho}\|\bar{\sigma}) = D_\alpha^*(\rho_\pi\|\sigma_\pi). \quad (4.4)$$

(ii) For any $\alpha \in [0, +\infty) \setminus \{1\}$,

$$\widehat{D}_\alpha(\rho\|\sigma) = D_\alpha(\rho_\pi\|\sigma_\pi), \quad i.e., \quad D_\alpha(\bar{\rho}\|\bar{\sigma}) = D_\alpha(\rho_\pi\|\sigma_\pi). \quad (4.5)$$

In particular, the equalities in (4.3) hold when \mathcal{A} is a von Neumann algebra and ρ, σ are normal functionals.

The proof of the theorem is somewhat technical based on Kosaki's interpolation L^p -spaces [38], so we defer it to Appendix C. Once (4.3) is confirmed by Theorem 4.3, we may and do rewrite $\widehat{D}_\alpha^*(\rho\|\sigma)$ and $\widehat{D}_\alpha(\rho\|\sigma)$ for $\rho, \sigma \in \mathcal{A}_+^*$ as $D_\alpha^*(\rho\|\sigma)$ and $D_\alpha(\rho\|\sigma)$, respectively.

Remark 4.4. Recall that $D_{1/2}^*(\rho\|\sigma) = -2\log F(\rho, \sigma)$ for $\rho, \sigma \in \mathcal{M}_*^+$, where \mathcal{M} is a von Neumann algebra and $F(\rho, \sigma)$ is the *fidelity* of ρ, σ . Theorem 4.3 (for $\alpha = 1/2$) shows that the fidelity of states $\rho, \sigma \in \mathcal{A}_+^*$ may be defined by $F(\rho_\pi, \sigma_\pi)$ via any (ρ, σ) -normal representation π of \mathcal{A} . In view of (2.6), Theorem 4.3 also shows that the relative entropy $D(\rho\|\sigma)$ of $\rho, \sigma \in \mathcal{A}_+^*$ can be defined by $D(\rho_\pi\|\sigma_\pi)$ as above; see [3, Sec. 5] and [29, Lemma 3.1].

Based on Definition 4.1 and Theorem 4.3 we can easily extend properties of the sandwiched and the standard Rényi divergences from the von Neumann setting to the C^* -algebra setting. For instance, some important ones are given in the next proposition.

Proposition 4.5. *Let \mathcal{A} be a unital C^* -algebra and $\rho, \sigma \in \mathcal{A}_+^*$.*

- (1) Joint lower semi-continuity: *The map $(\rho, \sigma) \in \mathcal{A}_+^* \times \mathcal{A}_+^* \mapsto D_\alpha^*(\rho\|\sigma)$ is jointly lower semi-continuous in the norm topology for every $\alpha \in (1, +\infty)$ and jointly continuous in the norm topology for every $\alpha \in [1/2, 1)$. The map $(\rho, \sigma) \in \mathcal{A}_+^* \times \mathcal{A}_+^* \mapsto D_\alpha(\rho\|\sigma)$ is jointly lower semi-continuous in the weak (i.e., $\sigma(\mathcal{A}^*, \mathcal{A}^{**})$ -) topology for every $\alpha \in [0, 2]$ and jointly continuous in the norm topology for every $\alpha \in [0, 1)$.*
- (2) Monotonicity (Data-processing inequality): *Let $\Phi : \mathcal{B} \rightarrow \mathcal{A}$ be a unital positive linear map between unital C^* -algebras. Then for every $\alpha \in [1/2, +\infty) \setminus \{1\}$,*

$$D_\alpha^*(\rho \circ \Phi \|\sigma \circ \Phi) \leq D_\alpha^*(\rho \|\sigma).$$

If Φ is, in addition, a Schwarz map, then for every $\alpha \in [0, 2] \setminus \{1\}$,

$$D_\alpha(\rho \circ \Phi \|\sigma \circ \Phi) \leq D_\alpha(\rho \|\sigma).$$

- (3) Inequality between D_α^* and D_α : *For every $\alpha \in [1/2, +\infty) \setminus \{1\}$,*

$$D_\alpha^*(\rho \|\sigma) \leq D_\alpha(\rho \|\sigma).$$

- (4) Martingale convergence: *Assume that $\{\mathcal{A}_i\}_{i \in \mathcal{I}}$ is an increasing net of C^* -subalgebras of \mathcal{A} containing the unit of \mathcal{A} such that $\bigcup_{i \in \mathcal{I}} \mathcal{A}_i$ is norm-dense in \mathcal{A} . Then for every $\alpha \in [0, 2] \setminus \{1\}$,*

$$D_\alpha(\rho|_{\mathcal{A}_i} \|\sigma|_{\mathcal{A}_i}) \nearrow D_\alpha(\rho \|\sigma).$$

Moreover, if there exists a (ρ, σ) -normal representation π of \mathcal{A} such that $\pi(\mathcal{A})''$ is σ -finite (this is automatic if \mathcal{A} is separable), then for every $\alpha \in [1/2, +\infty) \setminus \{1\}$,

$$D_\alpha^*(\rho|_{\mathcal{A}_i} \|\sigma|_{\mathcal{A}_i}) \nearrow D_\alpha^*(\rho \|\sigma).$$

Proof. (1) Let $\rho_n, \sigma_n \in \mathcal{A}_+^*$, $n \in \mathbb{N}$, be such that $\|\rho_n - \rho\| \rightarrow 0$ and $\|\sigma_n - \sigma\| \rightarrow 0$, with normal extensions $\bar{\rho}_n, \bar{\sigma}_n$ to $\pi_u(\mathcal{A})''$ as well as $\bar{\rho}, \bar{\sigma}$. By Kaplansky's density theorem, note that $\|\bar{\rho}_n - \bar{\rho}\| = \|\rho_n - \rho\|$ and $\|\bar{\sigma}_n - \bar{\sigma}\| = \|\sigma_n - \sigma\|$. Hence the assertions follow from the corresponding result in the von Neumann algebra case; see [34, Proposition 3.10] and [26, Theorem 3.16 (3)] for D_α^* , and [24, Proposition 5.3 (6)] and [26, Corollary 3.8] for D_α .

(2) As shown in the proof of [25, Proposition 7.4], Φ extends to a unital positive normal map $\bar{\Phi} : \pi_u(\mathcal{B})'' \rightarrow \pi_u(\mathcal{A})''$ in such a way that $\bar{\Phi} \circ \pi_u = \pi_u \circ \Phi$, where π_u on the left-hand side is the universal representation of \mathcal{B} and that of \mathcal{A} is on the right-hand side. Then $\bar{\rho} \circ \bar{\Phi}$ and $\bar{\sigma} \circ \bar{\Phi}$

are the normal extensions of $\rho \circ \Phi$ and $\sigma \circ \Phi$ to $\pi_u(\mathcal{B})''$, as seen in the proof of [25, Proposition 7.4]. By [34, Theorem 3.14] and [35, Theorem 4.1]³ we have $D_\alpha^*(\rho \circ \Phi \| \sigma \circ \Phi) \leq D_\alpha^*(\rho \| \sigma)$ for all $\alpha \in [1/2, +\infty) \setminus \{1\}$. In addition, assume that Φ is a Schwarz map. Then it is easy to see by Kaplansky's density theorem that $\bar{\Phi}$ is a Schwarz map again. Hence [24, Proposition 5.3 (9)] gives $D_\alpha(\rho \circ \Phi \| \sigma \circ \Phi) \leq D_\alpha(\rho \| \sigma)$ for all $\alpha \in [0, 2] \setminus \{1\}$.

(3) By [8, Theorem 12] (also [34, Corollary 3.6]) we have

$$D_\alpha^*(\rho \| \sigma) = D_\alpha^*(\bar{\rho} \| \bar{\sigma}) \leq D_\alpha(\bar{\rho} \| \bar{\sigma}) = D_\alpha(\rho \| \sigma).$$

(4) Let π be as stated in the latter assertion (for D_α^*). Then it is clear that $\{\pi(\mathcal{A}_i)''\}_{i \in \mathcal{I}}$ is an increasing net of von Neumann subalgebras of σ -finite $\pi(\mathcal{A})''$ containing the unit of $\pi(\mathcal{A})''$ such that $(\bigcup_i \pi(\mathcal{A}_i)'')'' = \pi(\mathcal{A})''$. By Theorems 3.1 and 4.3, for every $\alpha \in [1/2, +\infty) \setminus \{1\}$ we have

$$D_\alpha^*(\rho|_{\mathcal{A}_i} \| \sigma|_{\mathcal{A}_i}) = D_\alpha^*(\rho_\pi|_{\pi(\mathcal{A}_i)''} \| \sigma_\pi|_{\pi(\mathcal{A}_i)''}) \nearrow D_\alpha^*(\rho_\pi \| \sigma_\pi) = D_\alpha^*(\rho \| \sigma).$$

As for the first assertion, note that $Q_\alpha(\bar{\rho} \| \bar{\sigma})$ for $\alpha \in (1, 2]$ is the standard f -divergence of $\bar{\rho}, \bar{\sigma}$ for $f(t) = t^\alpha$ and $-Q_\alpha(\bar{\rho} \| \bar{\sigma})$ for $\alpha \in [0, 1)$ is that for $f(t) = -t^\alpha$. Since these functions f_α are operator convex on $(0, +\infty)$, the martingale convergence for D_α when $\alpha \in [0, 2] \setminus \{1\}$ follows from [24, Theorem 4.1 (v)]. \square

Example 4.6. Let $\mathcal{C}(\mathcal{H})$ be the compact operator ideal in $\mathcal{B}(\mathcal{H})$, and $\mathcal{T}(\mathcal{H})$ be the trace class on \mathcal{H} . Consider the unital C^* -algebra $\mathcal{A} := \mathcal{C}(\mathcal{H}) + \mathbb{C}1$. It is well known that $\mathcal{A}^* \cong \mathcal{T}(\mathcal{H})$ and $\mathcal{T}(\mathcal{H})^* \cong \mathcal{B}(\mathcal{H})$. Thus we can identify $\mathcal{A}_+^* = \mathcal{T}(\mathcal{H})_+ = \mathcal{B}(\mathcal{H})_+^*$. With this identification, for every $\rho, \sigma \in \mathcal{T}(\mathcal{H})_+$, the C^* -versions $D_\alpha^*(\rho \| \sigma)$ and $D_\alpha(\rho \| \sigma)$ of ρ, σ considered as elements of \mathcal{A}_+^* coincide with those of ρ, σ as elements of $\mathcal{B}(\mathcal{H})_+^*$. Recently in [41], the (α, z) -Rényi divergences, a more general notion than the sandwiched and the standard Rényi divergences, have been discussed for more general operators ρ, σ in $\mathcal{B}(\mathcal{H})_+$, as well as their strong converse exponents.

Example 4.7. Let (\mathcal{A}, G, τ) be a C^* -dynamical system, where G is a group and τ is an action of G as $*$ -automorphisms of \mathcal{A} . Let $\mathcal{S}_G(\mathcal{A})$ denote the set of τ -invariant states of \mathcal{A} . Let $\varphi \in \mathcal{S}_G(\mathcal{A})$ and $\{\pi_\varphi, \mathcal{H}_\varphi, \xi_\varphi\}$ be the cyclic representation of \mathcal{A} induced by φ . Then there is a unique unitary representation U_φ of G on \mathcal{H}_φ such that $U_\varphi(g)\xi_\varphi = \xi_\varphi$ and $\pi_\varphi(\tau_g(a)) = U_\varphi(g)\pi_\varphi(a)U_\varphi(g)^*$ for all $a \in \mathcal{A}, g \in G$. It is well known (see, e.g., [9, Corollary 4.3.11]) that \mathcal{A} is G -abelian, i.e., $\pi_\varphi(\mathcal{A})' \cap U_\varphi(G)'$ is abelian for all $\varphi \in \mathcal{S}_G(\mathcal{A})$ if and only if $\mathcal{S}_G(\mathcal{A})$ is a (Choquet) simplex. In this case, each $\varphi \in \mathcal{S}_G(\mathcal{A})$ has a unique maximal representing measure μ_φ on $\mathcal{S}_G(\mathcal{A})$, which is the $\pi_\varphi(\mathcal{A})' \cap U_\varphi(G)'$ -orthogonal measure of φ ; see [9, Proposition 4.3.3]. Define a unital positive linear map $\Phi : \mathcal{A} \rightarrow C(\mathcal{S}_G(\mathcal{A}))$ (consisting of all continuous functions

³In [34, 35] (also [26]) it was implicitly assumed that \mathcal{M} is σ -finite. But this assumption can be removed. Indeed, let $\Phi : \mathcal{N} \rightarrow \mathcal{M}$ be a unital positive normal map between von Neumann algebras and $\rho, \sigma \in \mathcal{M}_+^*$. Let $e := s(\rho) \vee s(\sigma)$ and $e_0 := s(\rho \circ \Phi) \vee s(\sigma \circ \Phi)$. Since $s(\rho)(1 - \Phi(e_0))s(\rho) = 0$, $s(\rho) = s(\rho)\Phi(e_0) = \Phi(e_0)s(\rho)$ and $s(\rho) \leq \Phi(e_0)$. Similarly, $s(\sigma) = s(\sigma)\Phi(e_0) = \Phi(e_0)s(\sigma)$ and $s(\sigma) \leq \Phi(e_0)$. Let $P := 1 - s(\rho), Q := 1 - s(\sigma)$ and $A := 1 - \Phi(e_0) \geq 0$; hence $P \geq A$ and $Q \geq A$. Note that $PQP \geq PAP = A, QPQPQ \geq QAQ = A$, and so on. Hence $(PQ)^n P \geq A$ for all $n \geq 1$. Since $(PQ)^n P \rightarrow P \wedge Q$ strongly, we have $P \wedge Q \geq A$, which means that $e \leq \Phi(e_0)$. We thus find that $\tilde{\Phi} := e\Phi(\cdot)e|_{e_0\mathcal{N}e_0}$ is a unital positive normal map from $e_0\mathcal{N}e_0$ to $e\mathcal{M}e$. Note that $e_0\mathcal{N}e_0$ and $e\mathcal{M}e$ are σ -finite. Moreover, $D_\alpha^*(\rho \| \sigma) = D_\alpha^*(\rho|_{e\mathcal{M}e} \| \sigma|_{e\mathcal{M}e})$ and

$$D_\alpha^*(\rho \circ \Phi \| \sigma \circ \Phi) = D_\alpha^*(\rho \circ \Phi|_{e_0\mathcal{N}e_0} \| \sigma \circ \Phi|_{e_0\mathcal{N}e_0}) = D_\alpha^*((\rho|_{e\mathcal{M}e}) \circ \tilde{\Phi} \| (\sigma|_{e\mathcal{M}e}) \circ \tilde{\Phi}).$$

on $\mathcal{S}_G(\mathcal{A})$ by $(\Phi a)(\omega) := \omega(a)$ for $a \in \mathcal{A}$, $\omega \in \mathcal{S}_G(\mathcal{A})$. Let $\rho, \sigma \in \mathcal{S}_G(\mathcal{A})$. Since $\rho = \mu_\rho \circ \Phi$ and $\sigma = \mu_\sigma \circ \Phi$, Proposition 4.5 (2) gives

$$\begin{aligned} D_\alpha^*(\rho\|\sigma) &\leq D_\alpha(\mu_\rho\|\mu_\sigma), & \alpha \in [1/2, +\infty) \setminus \{1\}, \\ D_\alpha(\rho\|\sigma) &\leq D_\alpha(\mu_\rho\|\mu_\sigma), & \alpha \in [0, 2] \setminus \{1\}, \end{aligned} \quad (4.6)$$

where $D_\alpha(\mu_\rho\|\mu_\sigma)$ is the classical Rényi relative entropy. Now assume that \mathcal{A} is G -central, i.e., $\pi_\varphi(\mathcal{A})' \cap U_\varphi(G)' \subseteq \pi_\varphi(\mathcal{A})''$ (hence $\pi_\varphi(\mathcal{A})' \cap U_\varphi(G)' \subseteq$ the center of $\pi_\varphi(\mathcal{A})''$) for all $\varphi \in \mathcal{S}_G(\mathcal{A})$, which is stronger than G -abeliannes and weaker than some other conditions of asymptotic abeliannes; see [9, Sec. 4.3], [15]. In this case, with $\varphi := (\rho + \sigma)/2$ there is an isomorphism $\theta_\varphi : L^\infty(\mathcal{S}_G(\mathcal{A}), \mu_\varphi) \rightarrow \pi_\varphi(\mathcal{A})' \cap U_\varphi(G)'$ such that

$$\langle \xi_\varphi, \theta_\varphi(f) \pi_\varphi(a) \xi_\varphi \rangle = \int f(\omega) \omega(a) d\mu_\varphi(\omega), \quad f \in L^\infty(\mathcal{S}_G(\mathcal{A}), \mu_\varphi). \quad (4.7)$$

For this, see [9, Proposition 41.22]. Since $\rho, \sigma \leq 2\varphi$, note [9, Corollary 4.1.17] that $\mu_\rho, \mu_\sigma \leq 2\mu_\varphi$ so that one can take $g_\rho := d\mu_\rho/d\mu_\varphi$ and $g_\sigma := d\mu_\sigma/d\mu_\varphi$ in $L^\infty(\mathcal{S}_G(\mathcal{A}), \mu_\varphi)_+$. Then for every $a \in \mathcal{A}$ one has

$$\rho(a) = \int g_\rho(\omega) \omega(a) d\mu_\varphi(\omega) = \langle \xi_\varphi, \theta_\varphi(g_\rho) \pi_\varphi(a) \xi_\varphi \rangle$$

thanks to (4.7). Hence the normal extension of ρ to $\pi_\varphi(\mathcal{A})''$ is given as

$$\tilde{\rho}(x) = \langle \xi_\varphi, \theta_\varphi(g_\rho) x \xi_\varphi \rangle, \quad x \in \pi_\varphi(\mathcal{A})'',$$

which implies that

$$\begin{aligned} \tilde{\rho}(\theta_\varphi(f)) &= \langle \xi_\varphi, \theta_\varphi(g_\rho f) \xi_\varphi \rangle = \int g_\rho(\omega) f(\omega) d\mu_\varphi(\omega) \quad (\text{by (4.7)}) \\ &= \int f(\omega) d\mu_\rho(\omega), \quad f \in L^\infty(\mathcal{S}_G(\mathcal{A}), \mu_\varphi). \end{aligned}$$

Hence $\tilde{\rho} \circ \theta_\varphi = \mu_\rho$ follows. Similarly, replacing ρ, g_ρ with σ, g_σ in the above argument, one has $\tilde{\sigma} \circ \theta_\varphi = \mu_\sigma$ for the normal extension $\tilde{\sigma}$ of σ to $\pi_\varphi(\mathcal{A})''$. From Proposition 4.5 (2) and Theorem 4.3 it follows that

$$\begin{aligned} D_\alpha(\mu_\rho\|\mu_\sigma) &\leq D_\alpha^*(\tilde{\rho}\|\tilde{\sigma}) = D_\alpha^*(\rho\|\sigma), & \alpha \in [1/2, +\infty) \setminus \{1\}, \\ D_\alpha(\mu_\rho\|\mu_\sigma) &\leq D_\alpha(\tilde{\rho}\|\tilde{\sigma}) = D_\alpha(\rho\|\sigma), & \alpha \in [0, 2] \setminus \{1\}. \end{aligned} \quad (4.8)$$

Therefore, when \mathcal{A} is G -central and $\rho, \sigma \in \mathcal{S}_G(\mathcal{A})$, it follows from (4.6) and (4.8) that $D_\alpha^*(\rho\|\sigma) = D_\alpha(\mu_\rho\|\mu_\sigma)$ for all $\alpha \in [1/2, +\infty) \setminus \{1\}$ and $D_\alpha(\rho\|\sigma) = D_\alpha(\mu_\rho\|\mu_\sigma)$ for all $\alpha \in [0, 2] \setminus \{1\}$, showing also $D(\rho\|\sigma) = D(\mu_\rho\|\mu_\sigma)$ as shown in [29, Theorem 3.2]. Furthermore, μ_ρ and μ_σ are singular if and only if ρ and σ are orthogonal in the sense of [9, Definition 4.1.20], i.e., $s(\vec{\rho}) \perp s(\vec{\sigma})$ [22]. In this case, $D_\alpha^*(\rho\|\sigma) = D_\alpha(\mu_\rho\|\mu_\sigma) = +\infty$ for all $\alpha \in [1/2, +\infty) \setminus \{1\}$.

Example 4.8. Let $\mathcal{B} = \bigotimes_1^\infty \mathbb{M}_d$ be a UHF (or one-dimensional spin) C^* -algebra, and τ be the action of S_∞ (the group of finite permutations on \mathbb{N}) given by $\tau(\bigotimes_1^\infty a_n) = \bigotimes_1^\infty a_{g(n)}$, $g \in S_\infty$. Let K be a compact group and u_k ($k \in K$) a continuous unitary representation of K on \mathbb{C}^d , so a product action β of K on \mathcal{B} is defined by $\beta_k := \bigotimes_1^\infty \text{Ad } u_k$. The β -fixed point C^* -subalgebra $\mathcal{A} := \mathcal{B}^\beta$ of \mathcal{B} is called a *gauge-invariant C^* -algebra*. Let $\mathcal{B}_n := \bigotimes_1^n \mathbb{M}_d$

and $\mathcal{A}_n := \mathcal{B}_n^\beta = \mathcal{A} \cap \mathcal{B}_n$ for $n \geq 1$; then $\bigcup_{n=1}^\infty \mathcal{A}_n$ is norm-dense in \mathcal{A} (so \mathcal{A} is an AF C^* -algebra.) Then $(\mathcal{B}, S_\infty, \tau)$ and $(\mathcal{A}, S_\infty, \tau|_{\mathcal{A}})$ are typical cases of Example 4.7, since \mathcal{B} and \mathcal{A} are asymptotically abelian with respect to S_∞ . Størmer's theorem [59] says that the extreme points of $\mathcal{S}_{S_\infty}(\mathcal{B})$ are the symmetric product states $\bigotimes_1^\infty \psi$, $\psi \in \mathcal{S}(\mathbb{M}_d)$. For every $\rho, \sigma \in \mathcal{S}_{S_\infty}(\mathcal{A})$, by Proposition 4.5 (4) and Example 4.7 one has

$$\begin{aligned} D_\alpha^*(\rho\|\sigma) &= \lim_{n \rightarrow \infty} D_\alpha^*(\rho|_{\mathcal{A}_n}\|\sigma|_{\mathcal{A}_n}) = D_\alpha(\mu_\rho\|\mu_\sigma), & \alpha \in [1/2, +\infty) \setminus \{1\}, \\ D_\alpha(\rho\|\sigma) &= \lim_{n \rightarrow \infty} D_\alpha(\rho|_{\mathcal{A}_n}\|\sigma|_{\mathcal{A}_n}) = D_\alpha(\mu_\rho\|\mu_\sigma), & \alpha \in [0, 2] \setminus \{1\}, \end{aligned}$$

where μ_ρ, μ_σ are the representing measures of ρ, σ on the extreme boundary of $\mathcal{S}_{S_\infty}(\mathcal{A})$. In particular, it follows that $D_\alpha^*(\rho\|\sigma) = D_\alpha(\rho\|\sigma)$ for any $\rho, \sigma \in \mathcal{S}_{S_\infty}(\mathcal{A})$ and all $\alpha \in [1/2, 2] \setminus \{1\}$ even though $D_\alpha^*(\rho|_{\mathcal{A}_n}\|\sigma|_{\mathcal{A}_n}) < D_\alpha(\rho|_{\mathcal{A}_n}\|\sigma|_{\mathcal{A}_n})$ unless $\rho|_{\mathcal{A}_n}$ and $\sigma|_{\mathcal{A}_n}$ commute. This phenomenon can naturally be understood by considering the asymptotic abelianness of \mathcal{A} with respect to S_∞ . Now let $\mathcal{T}(\mathcal{A})$ denote the set of trace states on \mathcal{A} . Recall [56, Theorem 3.2] that any extreme point of $\mathcal{T}(\mathcal{A})$ is the restriction to \mathcal{A} of a symmetric product state of \mathcal{B} , so that $\mathcal{T}(\mathcal{A}) \subseteq \mathcal{S}_{S_\infty}(\mathcal{A})$. Furthermore, note (see [64, Sec. 4]) that $\mathcal{T}(\mathcal{A})$ is a face of $\mathcal{S}_{S_\infty}(\mathcal{A})$. Hence, for every $\rho, \sigma \in \mathcal{T}(\mathcal{A})$ one has

$$D_\alpha(\rho\|\sigma) = \lim_{n \rightarrow \infty} D_\alpha(\rho|_{\mathcal{A}_n}\|\sigma|_{\mathcal{A}_n}) = D_\alpha(\mu_\rho\|\mu_\sigma), \quad \alpha \in [0, +\infty) \setminus \{1\}, \quad (4.9)$$

where the representing measures μ_ρ, μ_σ are supported on the extreme boundary of $\mathcal{T}(\mathcal{A})$ and $D_\alpha^*(\rho\|\sigma) = D_\alpha(\rho\|\sigma)$ holds since ρ, σ are trace states (hence they commute). For example, let $d = 2$, $K = \mathbb{T} = \{\zeta \in \mathbb{C} : |\zeta| = 1\}$ and $u_\zeta := \begin{bmatrix} 1 & 0 \\ 0 & \zeta \end{bmatrix}$, $\zeta \in \mathbb{T}$. Then $\mathcal{A} = \mathcal{B}^\beta$ is the so-called *GICAR algebra*, and note that

$$\mathcal{A}_n = \bigoplus_{k=0}^n \mathbb{M}_{\binom{n}{k}}, \quad n \geq 1.$$

It is well known that the extreme boundary of $\mathcal{T}(\mathcal{A})$ is parametrized by $\lambda \in [0, 1]$; to be precise, it is $\{\omega_\lambda : \lambda \in [0, 1]\}$, where $\omega_\lambda = (\bigotimes_1^\infty \psi_\lambda)|_{\mathcal{A}}$ with $\psi_\lambda := \text{Tr} \left(\begin{bmatrix} \lambda & 0 \\ 0 & 1 - \lambda \end{bmatrix} \cdot \right)$ on \mathbb{M}_2 . Let $\rho, \sigma \in \mathcal{T}(\mathcal{A})$ and decompose them as $\rho = \int_0^1 \omega_\lambda d\mu_\rho(\lambda)$, $\sigma = \int_0^1 \omega_\lambda d\mu_\sigma(\lambda)$ with unique probability measures μ_ρ, μ_σ on $[0, 1]$. Since

$$\omega_\lambda|_{\mathcal{A}_n} = \bigoplus_{k=0}^n \lambda^k (1 - \lambda)^{n-k} \text{Tr}_{\binom{n}{k}}$$

(with convention $\lambda^0 = 1$ for $\lambda = 0$), the convergence $\lim_{n \rightarrow \infty} Q_\alpha(\rho|_{\mathcal{A}_n}\|\sigma|_{\mathcal{A}_n}) = Q_\alpha(\mu_\rho\|\mu_\sigma)$ due to (4.9) can be rewritten in an explicit form as

$$\begin{aligned} & \lim_{n \rightarrow \infty} \sum_{k=0}^n \binom{n}{k} \left(\int_0^1 \lambda^k (1 - \lambda)^{n-k} d\mu_1(\lambda) \right)^\alpha \left(\int_0^1 \lambda^k (1 - \lambda)^{n-k} d\mu_2(\lambda) \right)^{1-\alpha} \\ &= \int_0^1 \left(\frac{d\mu_1}{d(\mu_1 + \mu_2)} \right)^\alpha \left(\frac{d\mu_2}{d(\mu_1 + \mu_2)} \right)^{1-\alpha} d(\mu_1 + \mu_2), \quad \alpha \in [0, +\infty), \end{aligned}$$

for any probability measures μ_1, μ_2 on $[0, 1]$. (It does not seem easy to verify this convergence formula in a direct manner.)

4.2 The strong converse exponent in nuclear C^* -algebras

Let \mathcal{A} and $\rho, \sigma \in \mathcal{A}_+^*$ be as in Section 4.1. For each $r \in \mathbb{R}$, we define the *Hoeffding anti-divergence* $H_r^*(\rho\|\sigma)$ in the same way as in Definition 3.6 with use of $D_\alpha^*(\rho\|\sigma)$, i.e., $\widehat{D}_\alpha^*(\rho\|\sigma)$ in (4.1) or (4.4). For any (ρ, σ) -normal representation π of \mathcal{A} , Theorem 4.3 gives

$$H_r^*(\rho\|\sigma) = H_r^*(\rho_\pi\|\sigma_\pi), \quad r \in \mathbb{R}. \quad (4.10)$$

We furthermore define strong converse exponents as in Definition 3.5 in the C^* -algebra setting. Here, recall (see, e.g., [62, Sec. IV.4]) that for C^* -algebras \mathcal{A} and \mathcal{B} , a C^* -cross-norm on the algebraic tensor product $\mathcal{A} \odot \mathcal{B}$ is not unique in general (though it exists always), and we have the smallest one $\|\cdot\|_{\min}$ and the largest one $\|\cdot\|_{\max}$. The *minimal (or spatial) C^* -tensor product* $\mathcal{A} \otimes_{\min} \mathcal{B}$ is the completion of $\mathcal{A} \odot \mathcal{B}$ with respect to $\|\cdot\|_{\min}$, and the *maximal C^* -tensor product* $\mathcal{A} \otimes_{\max} \mathcal{B}$ is that with respect to $\|\cdot\|_{\max}$. To discuss the simple hypothesis testing on a C^* -algebra, the minimal C^* -tensor product is suitable for the following reasons: When \mathcal{A} and \mathcal{B} are realized in $\mathcal{B}(\mathcal{H}_1)$ and $\mathcal{B}(\mathcal{H}_2)$ respectively, $\mathcal{A} \otimes_{\min} \mathcal{B}$ is isomorphic to the C^* -algebra generated by $\mathcal{A} \odot \mathcal{B}$ in $\mathcal{B}(\mathcal{H}_1 \otimes \mathcal{H}_2)$; see [62, Theorem IV.4.9 (iii)]. For any representations π_1 of \mathcal{A} and π_2 of \mathcal{B} , the tensor product representation $\pi_1 \otimes \pi_2$ of $\mathcal{A} \otimes_{\min} \mathcal{B}$ satisfies $(\pi_1 \otimes \pi_2)(\mathcal{A} \otimes_{\min} \mathcal{B})'' = \pi_1(\mathcal{A})'' \overline{\otimes} \pi_2(\mathcal{B})''$; see [62, Proposition IV.4.13].

Let $\rho, \sigma \in \mathcal{A}_+^*$. For each $n \in \mathbb{N}$ let $\mathcal{A}^{\otimes n}$ ($= \mathcal{A}^{\otimes_{\min} n}$) be the n -fold minimal C^* -tensor product of \mathcal{A} , and $\rho_n := \rho^{\otimes n}$ (resp., $\sigma_n := \sigma^{\otimes n}$) be the n -fold tensor product of ρ (resp., σ) on $\mathcal{A}^{\otimes n}$. The following are the strong converse exponents in the setting of the simple hypothesis testing for the null hypothesis $H_0 : \rho$ versus the alternative hypothesis $H_1 : \sigma$.

Definition 4.9. For each $r \in \mathbb{R}$ we define the *strong converse exponents* $\underline{sc}_r(\rho\|\sigma)$ etc. in the same expressions as in Definition 3.5, where $\{T_n\}$ in the present setting are taken as $T_n \in \mathcal{A}^{\otimes n}$, $0 \leq T_n \leq 1$, $n \in \mathbb{N}$. Then the relation (3.5) is obvious as before.

Remark 4.10. The most general notion of a test in the C^* -algebra setting is not covered by tests taken in Definition 4.9. For instance, one could say that a measurement is represented by a self-adjoint element X in \mathcal{A} , and the possible outcomes are the points of the spectrum of X . After measuring X , one could always make a classical post-processing, e.g., by dividing the spectrum of X into two disjoint sets B_0 and B_1 . If the measurement outcome falls into B_0 , we accept H_0 , otherwise we accept H_1 . The error probabilities corresponding to such a test may not be described by an element $T \in \mathcal{A}$ with $0 \leq T \leq I$. However, the next lemma shows that considering such more general tests does not change the strong converse exponents.

Lemma 4.11. *Let $\rho, \sigma \in \mathcal{A}_+^*$ and π be a (ρ, σ) -normal representation of \mathcal{A} . For every $r \in \mathbb{R}$, let $\varepsilon_r(\rho\|\sigma)$ be any strong converse exponent given in Definition 4.9, and $\varepsilon_r(\rho_\pi\|\sigma_\pi)$ be the same strong converse exponent in Definition 3.5 for ρ_π, σ_π . Then we have*

$$\varepsilon_r(\rho\|\sigma) = \varepsilon_r(\rho_\pi\|\sigma_\pi). \quad (4.11)$$

Proof. Let $\mathcal{M} := \pi(\mathcal{A})''$. For every $T_n \in \mathcal{A}^{\otimes n}$, $n \in \mathbb{N}$, with $0 \leq T_n \leq 1$, one has $\pi^{\otimes n}(T_n) \in \mathcal{M}^{\otimes n}$, $0 \leq \pi^{\otimes n}(T_n) \leq 1$, $\rho_{\pi, n}(\pi^{\otimes n}(T_n)) = \rho_n(T_n)$ and $\sigma_{\pi, n}(\pi^{\otimes n}(T_n)) = \sigma_n(T_n)$, where $\rho_{\pi, n} := (\rho_\pi)^{\otimes n}$ and $\sigma_{\pi, n} := (\sigma_\pi)^{\otimes n}$. Hence $\varepsilon_r(\rho_\pi\|\sigma_\pi) \leq \varepsilon_r(\rho\|\sigma)$ holds immediately. Conversely, for every $\tilde{T}_n \in \mathcal{M}^{\otimes n} = \pi^{\otimes n}(\mathcal{A}^{\otimes n})''$, $n \in \mathbb{N}$, with $0 \leq \tilde{T}_n \leq 1$, by Kaplansky's density theorem

and [31, Lemma IV.3.8] one can choose $T_n \in \mathcal{A}^{\otimes n}$, $n \in \mathbb{N}$, such that $0 \leq T_n \leq 1$ and

$$\begin{aligned} \rho_n(T_n) &= \rho_{\pi,n}(\pi^{\otimes n}(T_n)) \in \begin{cases} (e^{-1}\rho_{\pi,n}(\tilde{T}_n), e\rho_{\pi,n}(\tilde{T}_n)) & \text{if } \rho_{\pi,n}(\tilde{T}_n) > 0, \\ [0, 1/n^n] & \text{if } \rho_{\pi,n}(\tilde{T}_n) = 0, \end{cases} \\ \sigma_n(T_n) &= \sigma_{\pi,n}(\pi^{\otimes n}(T_n)) \in \begin{cases} (e^{-1}\sigma_{\pi,n}(\tilde{T}_n), e\sigma_{\pi,n}(\tilde{T}_n)) & \text{if } \sigma_{\pi,n}(\tilde{T}_n) > 0, \\ [0, 1/n^n] & \text{if } \sigma_{\pi,n}(\tilde{T}_n) = 0, \end{cases} \end{aligned}$$

which imply that

$$\begin{aligned} -\frac{1}{n} \log \rho_n(T_n) &\in \begin{cases} (-\frac{1}{n} - \frac{1}{n} \log \rho_{\pi,n}(\tilde{T}_n), \frac{1}{n} - \frac{1}{n} \log \rho_{\pi,n}(\tilde{T}_n)) & \text{if } \rho_{\pi,n}(\tilde{T}_n) > 0, \\ (\log n, +\infty) & \text{if } \rho_{\pi,n}(\tilde{T}_n) = 0, \end{cases} \\ -\frac{1}{n} \log \sigma_n(T_n) &\in \begin{cases} (-\frac{1}{n} - \frac{1}{n} \log \sigma_{\pi,n}(\tilde{T}_n), \frac{1}{n} - \frac{1}{n} \log \sigma_{\pi,n}(\tilde{T}_n)) & \text{if } \sigma_{\pi,n}(\tilde{T}_n) > 0, \\ (\log n, +\infty) & \text{if } \sigma_{\pi,n}(\tilde{T}_n) = 0. \end{cases} \end{aligned}$$

From these one can see that $\varepsilon_r(\rho\|\sigma) \leq \varepsilon_r(\tilde{\rho}\|\tilde{\sigma})$. Hence (4.11) follows. \square

Now we recall (see, e.g., [63, Chap. XV]) that \mathcal{A} is said to be *nuclear* if, for every C^* -algebra \mathcal{B} , there is a unique C^* -cross-norm on $\mathcal{A} \odot \mathcal{B}$, i.e., $\mathcal{A} \otimes_{\min} \mathcal{B} = \mathcal{A} \otimes_{\max} \mathcal{B}$. Concerning nuclear C^* -algebras, among many others, the most fundamental result is that \mathcal{A} is nuclear if and only if \mathcal{A}^{**} is injective. So, if \mathcal{A} is nuclear, then $\pi(\mathcal{A})''$ is injective for every representation π of \mathcal{A} . Typical examples of nuclear C^* -algebras are AF C^* -algebras, in particular, the compact operator ideal $\mathcal{C}(\mathcal{H})$ (or rather $\mathcal{C}(\mathcal{H}) + \mathbb{C}1$ in the present setting; see Example 4.6).

The next theorem is the C^* -algebra version of Theorem 3.7, where we restrict ρ, σ to states on \mathcal{A} for the same reason as before.

Theorem 4.12. *Let $\rho, \sigma \in \mathcal{A}_+^*$ be states such that $D_\alpha^*(\rho\|\sigma) < +\infty$ for some $\alpha > 1$. If there exists a (ρ, σ) -normal representation π of \mathcal{A} such that $\pi(\mathcal{A})''$ is injective, then all the equalities in (3.8) hold for every $r \in \mathbb{R}$ in the present C^* -algebra situation too. In particular, this is the case if \mathcal{A} is nuclear.*

Proof. Let π be as stated in the theorem. Then Theorem 3.7 says that the equalities in (3.8) for ρ_π, ρ_σ in place of ρ, σ hold for every $r \in \mathbb{R}$. Hence the assertion follows from Lemma 4.11 and (4.10). \square

We remark that the same formula as in Theorem 3.9 holds true under the assumption of Theorem 4.12, where $\alpha_{e^{-nr}}^*(\rho_n\|\sigma_n)$ is defined as in (3.14) with $T_n \in \mathcal{A}^{\otimes n}$, $0 \leq T_n \leq 1$. The proof is the same as that of Theorem 3.9. Also, the generalized κ -cutoff rate $C_\kappa(\rho\|\sigma)$ in Definition 3.10 makes sense for states $\rho, \sigma \in \mathcal{A}_+^*$, and Theorem 3.11 is extended to the C^* -algebra setting with the same proof.

In the rest of the section let us discuss the (regularized) measured and the (regularized) test-measured Rényi divergences in the C^* -algebra setting. A measurement in \mathcal{A} is given by $\mathfrak{M} = (M_j)_{1 \leq j \leq k}$ of $M_j \in \mathcal{A}_+$ such that $\sum_{j=1}^k M_j = 1$. Then $D_\alpha^{\text{meas}}(\rho\|\sigma)$ and $\overline{D}_\alpha^{\text{meas}}(\rho\|\sigma)$ for $\rho, \sigma \in \mathcal{A}_+^*$ are defined in the same way as in Definition 3.12 by taking measurements in \mathcal{A} instead of \mathcal{M} . The test-measured versions $D_\alpha^{\text{test}}(\rho\|\sigma)$ and $\overline{D}_\alpha^{\text{test}}(\rho\|\sigma)$ for $\rho, \sigma \in \mathcal{A}_+^*$ are also defined in the same way as just after Definition 3.12 with tests $T \in \mathcal{A}$, $0 \leq T \leq 1$.

Lemma 4.13. *Let $\rho, \sigma \in \mathcal{A}_+^*$ and π be any (ρ, σ) -normal representation of \mathcal{A} . Then for every $\alpha \in [1/2, +\infty) \setminus \{1\}$ we have*

$$D_\alpha^{\text{meas}}(\rho \parallel \sigma) = D_\alpha^{\text{meas}}(\rho_\pi \parallel \sigma_\pi), \quad \overline{D}_\alpha^{\text{meas}}(\rho \parallel \sigma) = \overline{D}_\alpha^{\text{meas}}(\rho_\pi \parallel \sigma_\pi), \quad (4.12)$$

$$D_\alpha^{\text{test}}(\rho \parallel \sigma) = D_\alpha^{\text{test}}(\rho_\pi \parallel \sigma_\pi), \quad \overline{D}_\alpha^{\text{test}}(\rho \parallel \sigma) = \overline{D}_\alpha^{\text{test}}(\rho_\pi \parallel \sigma_\pi), \quad (4.13)$$

Proof. Let $\mathcal{M} := \pi(\mathcal{A})''$. If $(M_i)_{1 \leq i \leq k}$ is a measurement in \mathcal{A} , then $(\pi(M_i))_{1 \leq i \leq k}$ is a measurement in \mathcal{M} . Hence $D_\alpha^{\text{meas}}(\rho \parallel \sigma) \leq D_\alpha^{\text{meas}}(\rho_\pi \parallel \sigma_\pi)$ holds immediately. Conversely, let $\widetilde{\mathfrak{M}} = (\widetilde{M}_i)_{1 \leq i \leq k}$ be a measurement in \mathcal{M} . Consider the representation $\pi_k = \pi \otimes \text{id}_k$ of $\mathbb{M}_k(\mathcal{A}) = \mathcal{A} \otimes \mathbb{M}_k$ on $\mathcal{H} \otimes \mathbb{C}^k$ (where \mathbb{M}_k is the $k \times k$ matrix algebra and \mathcal{H} is the representing Hilbert space of π). Noting that $\pi_k(\mathbb{M}_k(\mathcal{A}))'' = \mathbb{M}_k(\pi(\mathcal{A}))'' = \mathbb{M}_k(\mathcal{M})$, we define $\tilde{A} \in \mathbb{M}_k(\mathcal{M})$ by

$$\tilde{A} := \begin{bmatrix} \widetilde{M}_1^{1/2} & & \\ & \ddots & \\ & & \widetilde{M}_k^{1/2} \\ & & & 0 \end{bmatrix}.$$

Note that $\tilde{A}^* \tilde{A}$ is a projection with the $(1, 1)$ -block 1 and all other blocks 0. Hence \tilde{A} is a contraction. By Kaplansky's density theorem and [31, Lemma IV.3.8] there exists a net $\{A^{(\lambda)}\}$ of contractions in $\mathbb{M}_k(\mathcal{A})$ such that $\pi_k(A^{(\lambda)}) \rightarrow \tilde{A}$ in the strong* topology. Write $A^{(\lambda)} = [a_{ij}^{(\lambda)}]_{i,j=1}^k$ with $a_{ij}^{(\lambda)} \in \mathcal{A}$; then $\pi_k(A^{(\lambda)}) = [\pi(a_{ij}^{(\lambda)})]_{i,j=1}^k$ so that $\pi(a_{i1}^{(\lambda)}) \rightarrow \widetilde{M}_i^{1/2}$ in the strong* topology for $1 \leq i \leq k$. Set $M_i^{(\lambda)} := (a_{i1}^{(\lambda)})^* a_{i1}^{(\lambda)} \in \mathcal{A}_+$, $1 \leq i \leq k$. Then $\pi(M_i^{(\lambda)}) \rightarrow \widetilde{M}_i$ strongly for $1 \leq i \leq k$, and $\sum_{i=1}^k M_i^{(\lambda)} = \sum_{i=1}^k (a_{i1}^{(\lambda)})^* a_{i1}^{(\lambda)}$ is the $(1, 1)$ -block of $(A^{(\lambda)})^* A^{(\lambda)}$ so that $\sum_{i=1}^k M_i^{(\lambda)} \leq 1$. Therefore, one can define a net $\{\mathfrak{M}_\lambda\}$ of measurements in \mathcal{A} by $\mathfrak{M}_\lambda = (M_i^{(\lambda)})_{1 \leq i \leq k+1}$, where $M_{k+1}^{(\lambda)} := 1 - \sum_{i=1}^k M_i^{(\lambda)}$. Since $\pi(M_{k+1}^{(\lambda)}) \rightarrow 1 - \sum_{i=1}^k \widetilde{M}_i = 0$, we have

$$\mathfrak{M}_\lambda(\rho) = (\rho_\pi \circ \pi(M_i^{(\lambda)}))_{1 \leq i \leq k+1} \longrightarrow \widetilde{\mathfrak{M}}(\rho_\pi) \oplus 0,$$

and similarly $\mathfrak{M}_\lambda(\sigma) \rightarrow \widetilde{\mathfrak{M}}(\sigma_\pi) \oplus 0$. From the lower semi-continuity of (classical) Rényi divergence, it follows that

$$D_\alpha(\widetilde{\mathfrak{M}}(\rho_\pi) \parallel \widetilde{\mathfrak{M}}(\sigma_\pi)) \leq \liminf_\lambda D_\alpha(\mathfrak{M}_\lambda(\rho) \parallel \mathfrak{M}_\lambda(\sigma)) \leq D_\alpha^{\text{meas}}(\rho \parallel \sigma).$$

Hence $D_\alpha^{\text{meas}}(\rho_\pi \parallel \sigma_\pi) \leq D_\alpha^{\text{meas}}(\rho \parallel \sigma)$, implying the first equality of (4.12). Moreover, applying this to $\rho^{\otimes n}, \sigma^{\otimes n}$ and the representation $\pi^{\otimes n}$ of $\mathcal{A}^{\otimes n}$, we have for every $n \in \mathbb{N}$,

$$D_\alpha^{\text{meas}}(\rho^{\otimes n} \parallel \sigma^{\otimes n}) = D_\alpha^{\text{meas}}(\rho_\pi^{\otimes n} \parallel \sigma_\pi^{\otimes n}),$$

since $\rho_\pi^{\otimes n}, \sigma_\pi^{\otimes n}$ are the respective normal extensions of $\rho^{\otimes n}, \sigma^{\otimes n}$ to $\pi^{\otimes n}(\mathcal{A}^{\otimes n})'' = \mathcal{M}^{\otimes n}$. Thus, the second equality of (4.12) holds as well. The proof of (4.13) is similar (and easier) by approximating tests in \mathcal{M} with those in \mathcal{A} (similarly to the proof of Lemma 4.11). \square

Proposition 4.14. *Assume that there exists a (ρ, σ) -normal representation π of \mathcal{A} such that $\pi(\mathcal{A})''$ is injective (in particular, this is the case if \mathcal{A} is nuclear). Then (3.17) holds for $\alpha \in [1/2, +\infty) \setminus \{1\}$ and (3.18) holds for $\alpha > 1$.*

Proof. By Theorem 4.3 and Lemma 4.13, one can apply Proposition 3.13 to ρ_π, σ_π to obtain the assertion. \square

Acknowledgments

The work of M.M. was partially funded by the National Research, Development and Innovation Office of Hungary via the research grants K124152 and KH129601, and by the Ministry of Innovation and Technology and the National Research, Development and Innovation Office within the Quantum Information National Laboratory of Hungary.

A Strong converse exponent in the finite-dimensional case

In this appendix we assume that a von Neumann algebra \mathcal{M} is finite-dimensional, so $\mathcal{M} \subseteq \mathcal{B}(\mathcal{H})$ with a finite-dimensional Hilbert space \mathcal{H} . Note that \mathcal{M} is isomorphic to $\bigoplus_{i=1}^m \mathcal{B}(\mathcal{H}_i)$, a finite direct sum of finite-dimensional $\mathcal{B}(\mathcal{H}_i)$, so it is clear that all the arguments in [44] are valid with \mathcal{M} in place of $\mathcal{B}(\mathcal{H})$. Let Tr be the usual trace on \mathcal{M} (such that $\text{Tr}(e) = 1$ of all minimal projections $e \in \mathcal{M}$). Below, to designate states of \mathcal{M} , we use density operators ρ, σ with respect to Tr rather than positive functionals. Recall that both of the relative entropy $D(\rho\|\sigma)$ and the max-relative entropy $D_{\max}(\rho\|\sigma)$ showing up in (2.6)–(2.8) play an important role to describe $\psi(s) := \psi^*(\rho\|\sigma|s+1)$ and $H_r^*(\rho\|\sigma)$ in [44, Sec. 4].

The aim of this appendix is to give Proposition A.2, which is used in Section 3.2. The main assertion is that for finite-dimensional density operators $sc_r^0(\rho\|\sigma) \leq H_r^*(\rho\|\sigma)$, $r \geq 0$, which in turn can be obtained easily from the weaker inequalities $sc_r(\rho\|\sigma) \leq H_r^*(\rho\|\sigma)$, $r \geq 0$. The latter was proved in [44]; however, the proof contains a gap, as it is implicitly assumed there that $D(\rho\|\sigma) < D_{\max}(\rho\|\sigma)$. Our main contribution in Proposition A.2 is filling this gap; for this we give a characterization of the case $D(\rho\|\sigma) = D_{\max}(\rho\|\sigma)$, which may be of independent interest.

Lemma A.1. *For density operators ρ, σ in \mathcal{M} with $s(\rho) \leq s(\sigma)$, the following conditions are equivalent:*

- (a) $s \mapsto \psi(s) := \psi^*(\rho\|\sigma|s+1)$ is affine on $(0, +\infty)$;
- (b) $D(\rho\|\sigma) = D_{\max}(\rho\|\sigma)$;
- (c) ρ and σ commute, and $\rho\sigma^{-1} = \gamma s(\rho)$ for some constant $\gamma > 0$;
- (d) $s(\rho)\sigma = \sigma s(\rho)$ and $\rho = \gamma\sigma s(\rho)$ for some constant $\gamma > 0$.

Moreover, if the above hold, then we have $D(\rho\|\sigma) = \log \gamma$ and

$$H_r^*(\rho\|\sigma) = (r - D(\rho\|\sigma))_+, \quad r \geq 0. \quad (\text{A.1})$$

Proof. (a) \iff (b). Since $\psi(s)$ is a differentiable convex function on $[0, +\infty)$, this is clear from [44, Lemma 4.2].

(b) \implies (c). Consider $D_2(\rho\|\sigma) := \log \text{Tr} \rho^2 \sigma^{-1}$, the standard (or Petz-type) Rényi 2-divergence of ρ, σ . Note that

$$D(\rho\|\sigma) \leq D_2^*(\rho\|\sigma) \leq D_2(\rho\|\sigma) \leq D_{\max}(\rho\|\sigma),$$

where the first inequality is seen from the properties noted in Section 2.3, the second is due to the Araki–Lieb–Thirring inequality, and the last was shown in [10, Lemma 7]. Hence

(b) implies that $D_2^*(\rho\|\sigma) = D_2(\rho\|\sigma)$, i.e., $\text{Tr}(\sigma^{-1/4}\rho\sigma^{-1/4})^2 = \text{Tr}\sigma^{-1/2}\rho^2\sigma^{-1/2}$. Using [23, Theorem 2.1] we find that ρ, σ commute. Hence, (a) says that $\psi(s) = \log \text{Tr} \rho^{s+1}\sigma^{-s}$ is affine on $(0, +\infty)$. From [28, Lemma 3.2] for the commutative case, it follows that $\rho\sigma^{-1} = \gamma s(\rho)$, implying (c).

(c) \implies (d) is obvious.

(d) \implies (b). From condition (c) it easily follows that $D_{\max}(\rho\|\sigma) = \log \gamma$. Moreover, since

$$\begin{aligned} D(\rho\|\sigma) &= \text{Tr}(\rho \log \rho - \rho \log(s(\rho)\sigma)) \\ &= \text{Tr}(\rho \log \rho - \rho \log(\gamma^{-1}\rho)) = \log \gamma, \end{aligned}$$

(b) follows.

Finally, if (b) and hence (a) hold, then $\psi(s) = D(\rho\|\sigma)s$ for all $s > 0$, from which (A.1) follows immediately. \square

The next proposition is used in the proof of Theorem 3.7, while the former is a specialized case of the latter.

Proposition A.2. *For every density operators ρ, σ in \mathcal{M} with $s(\rho) \leq s(\sigma)$ and any $r \geq 0$ we have*

$$\underline{sc}_r(\rho\|\sigma) = sc_r^0(\rho\|\sigma) = H_r^*(\rho\|\sigma). \quad (\text{A.2})$$

Proof. Since $sc_r^0(\rho\|\sigma) \geq \underline{sc}_r(\rho\|\sigma) \geq H_r^*(\rho\|\sigma)$, where the second inequality is by [44, Lemma 4.7], it suffices to prove that $sc_r^0(\rho\|\sigma) \leq H_r^*(\rho\|\sigma)$, $r \geq 0$. Moreover, the last inequality follows if we can prove that $sc_r(\rho\|\sigma) \leq H_r^*(\rho\|\sigma)$, $r \geq 0$, since then

$$sc_r^0(\rho\|\sigma) \leq \inf_{r' > r} sc_{r'}(\rho\|\sigma) \leq \inf_{r' > r} H_{r'}^*(\rho\|\sigma) = H_r^*(\rho\|\sigma),$$

where the first inequality is obvious by definition, the second inequality is to be proved below, and the equality follows from the fact that $r \mapsto H_r^*(\rho\|\sigma)$ is a monotone increasing finite-valued convex function on \mathbb{R} , whence it is also continuous.

Let us therefore prove $sc_r(\rho\|\sigma) \leq H_r^*(\rho\|\sigma)$, $r \geq 0$. The proof of [44, Theorem 4.10] gives this when $D(\rho\|\sigma) < D_{\max}(\rho\|\sigma)$. Assume thus that $D(\rho\|\sigma) = D_{\max}(\rho\|\sigma) =: D$. For any $r \geq 0$, the test sequence $T_{n,r} := e^{-n(r-D)_+} s(\rho)^{\otimes n}$, $n \in \mathbb{N}$, yields

$$-\frac{1}{n} \log \text{Tr} \rho_n T_{n,r} = (r - D)_+ = H_r^*(\rho\|\sigma),$$

where the last equality is by (A.1), and

$$-\frac{1}{n} \log \text{Tr} \sigma_n T_{n,r} = -\frac{1}{n} \log(e^{-n(r-D)_+} \text{Tr}(\sigma s(\rho))^{\otimes n}) = D + (r - D)_+ \geq r,$$

where we have used that $\sigma s(\rho) = e^{-D}\rho$ by Lemma A.1. This proves $sc_r(\rho\|\sigma) \leq H_r^*(\rho\|\sigma)$. \square

B Boundary values of convex functions on $(0, 1)$

Let $\{\phi_i\}_{i \in \mathcal{I}}$ be a set of convex functions on $(0, 1)$ with values in $(-\infty, +\infty]$. Define

$$\phi(u) := \sup_{i \in \mathcal{I}} \phi_i(u), \quad u \in (0, 1),$$

which is obviously convex on $(0, 1)$ with values in $(-\infty, +\infty]$. We extend ϕ_i and ϕ to $[0, 1]$ by continuity as

$$\begin{aligned}\phi_i(u) &:= \lim_{u \searrow 0} \phi_i(u), & \phi_i(1) &:= \lim_{u \nearrow 1} \phi_i(u), \\ \phi(0) &:= \lim_{u \searrow 0} \phi(u), & \phi(1) &:= \lim_{u \nearrow 1} \phi(u).\end{aligned}$$

We then give the next lemma to use it in the proof of Theorem 3.7.

Lemma B.1. *In the situation stated above, if $\phi(u) < +\infty$ for some $u \in (0, 1)$, then*

$$\phi(0) = \sup_{i \in \mathcal{I}} \phi_i(0), \quad \phi(1) = \sup_{i \in \mathcal{I}} \phi_i(1).$$

Proof. By assumption we have a $u_0 \in (0, 1)$ with $\phi(u_0) < +\infty$. Obviously, $\phi_i(0) \leq \phi(0)$ and $\phi_i(1) \leq \phi(1)$ for all $i \in \mathcal{I}$. Hence it suffices to show that $\phi(0) \leq \sup_i \phi_i(0)$ and $\phi(1) \leq \sup_i \phi_i(1)$. Let us prove the first inequality (the proof of the latter is similar). Set $\xi := \sup_i \phi_i(0)$. If $\xi = +\infty$, the assertion holds trivially. So assume $\xi < +\infty$. By convexity, for every $i \in \mathcal{I}$ we have

$$\phi_i(u) \leq \frac{u_0 - u}{u_0} \xi + \frac{u}{u_0} \phi(u_0), \quad u \in (0, u_0),$$

so that

$$\phi(u) \leq \frac{u_0 - u}{u_0} \xi + \frac{u}{u_0} \phi(u_0), \quad u \in (0, u_0).$$

This implies that $\phi(0) \leq \xi = \sup_i \phi_i(0)$. □

C Proof of Theorem 4.3

Let $\rho, \sigma \in \mathcal{A}_+^*$ and π be any (ρ, σ) -normal representation of \mathcal{A} with $\tilde{\rho} = \rho_\pi$ and $\tilde{\sigma} = \sigma_\pi$, the normal extensions to $\mathcal{M} := \pi(\mathcal{A})''$. Also, let $\bar{\rho}, \bar{\sigma}$ be the normal extensions of ρ, σ to the enveloping von Neumann algebra \mathcal{A}^{**} and $\bar{\pi} : \mathcal{A}^{**} \rightarrow \mathcal{M}$ be the normal extension of π to \mathcal{A}^{**} (see [62, p. 121]). Let $s(\bar{\pi})$ be the support projection of $\bar{\pi}$. Concerning the support projections $s(\tilde{\rho})$ and $s(\bar{\rho})$ we have $s(\tilde{\rho}) = \bar{\pi}(s(\bar{\rho}))$ with $s(\bar{\rho}) \leq s(\bar{\pi})$. Therefore, $s(\tilde{\rho}) \leq s(\tilde{\sigma})$ is equivalent to $s(\bar{\rho}) \leq s(\bar{\sigma})$. This means that the condition $s(\tilde{\rho}) \leq s(\tilde{\sigma})$ is independent of the choice of a representation π as above. (The condition is called the *absolute continuity* of ρ with respect to σ [22].)

Now let $\hat{\pi}$ be another (ρ, σ) -normal representation of \mathcal{A} with $\hat{\rho} := \rho_{\hat{\pi}}$ and $\hat{\sigma} := \sigma_{\hat{\pi}}$, the normal extensions to $\hat{\mathcal{M}} := \hat{\pi}(\mathcal{A})''$. The next lemma is a main ingredient of the proof of Theorem 4.3.

Lemma C.1. *In the situation stated above, assume that $s(\tilde{\rho}) \leq s(\tilde{\sigma})$ (hence $s(\hat{\rho}) \leq s(\hat{\sigma})$ as well). Let z_0, \hat{z}_0 denote the central supports of $s(\tilde{\sigma}), s(\hat{\sigma})$, respectively. Then there exists an isomorphism $\Lambda : \mathcal{M}z_0 \rightarrow \hat{\mathcal{M}}\hat{z}_0$ for which we have*

$$\tilde{\rho}(x) = \hat{\rho} \circ \Lambda(x), \quad \tilde{\sigma}(x) = \hat{\sigma} \circ \Lambda(x), \quad x \in \mathcal{M}z_0, \quad (\text{C.1})$$

and for every $p \in [1, +\infty)$,

$$\text{tr}(h_{\tilde{\sigma}}^{1/2p} x h_{\tilde{\sigma}}^{1/2p})^p = \text{tr}(h_{\hat{\sigma}}^{1/2p} \Lambda(x) h_{\hat{\sigma}}^{1/2p})^p, \quad x \in \mathcal{M}z_0, \quad (\text{C.2})$$

where $h_{\tilde{\sigma}} \in L^1(\mathcal{M})_+$ and $h_{\hat{\sigma}} \in L^1(\hat{\mathcal{M}})_+$ are Haagerup's L^1 -elements corresponding to $\tilde{\sigma} \in \mathcal{M}_*^+$ and $\hat{\sigma} \in \hat{\mathcal{M}}_*^+$, respectively.

Proof. We will work in the standard forms

$$(\mathcal{M}, L^2(\mathcal{M}), J = *, L^2(\mathcal{M})_+), \quad (\hat{\mathcal{M}}, L^2(\hat{\mathcal{M}}), \hat{J} = *, L^2(\hat{\mathcal{M}})_+).$$

For brevity we write

$$\begin{cases} h_0 := h_{\tilde{\rho}} \in L^1(\mathcal{M})_+, \\ k_0 := h_{\tilde{\sigma}} \in L^1(\mathcal{M})_+, \\ e_0 := s(\tilde{\sigma}) = s(k_0) \in \mathcal{M}, \\ e'_0 := J e_0 J \in \mathcal{M}', \end{cases} \quad \begin{cases} \hat{h}_0 := h_{\hat{\rho}} \in L^1(\hat{\mathcal{M}})_+, \\ \hat{k}_0 := h_{\hat{\sigma}} \in L^1(\hat{\mathcal{M}})_+, \\ \hat{e}_0 := s(\hat{\sigma}) = s(\hat{k}_0) \in \hat{\mathcal{M}}, \\ \hat{e}'_0 := \hat{J} \hat{e}_0 \hat{J} \in \hat{\mathcal{M}}'. \end{cases}$$

Below the proof is divided into several steps.

Step 1. Note that

$$\overline{\pi(\mathcal{A})k_0^{1/2}} = \overline{\mathcal{M}k_0^{1/2}} = L^2(\mathcal{M})e_0 = e'_0 L^2(\mathcal{M})$$

and for every $a \in \mathcal{A}$,

$$\langle k_0^{1/2}, \pi(a)e'_0 k_0^{1/2} \rangle = \langle k_0^{1/2}, \pi(a)k_0^{1/2} \rangle = \tilde{\sigma} \circ \pi(a) = \sigma(a).$$

Hence $\{\pi(\cdot)e'_0, e'_0 L^2(\mathcal{M}), k_0^{1/2}\}$ is the cyclic representation of \mathcal{A} with respect to σ , and similarly $\{\hat{\pi}(\cdot)\hat{e}'_0, \hat{e}'_0 L^2(\hat{\mathcal{M}}), \hat{k}_0^{1/2}\}$ is the same. By the uniqueness (up to unitary conjugation) of the cyclic representation, there exists a unitary $V : L^2(\mathcal{M})e_0 \rightarrow L^2(\hat{\mathcal{M}})\hat{e}_0$ such that

$$V k_0^{1/2} = \hat{k}_0^{1/2}, \quad V(\pi(a)e'_0)V^* = \hat{\pi}(a)\hat{e}'_0, \quad a \in \mathcal{A}. \quad (\text{C.3})$$

We hence have an isomorphism $V \cdot V^* : \mathcal{M}e'_0 \rightarrow \hat{\mathcal{M}}\hat{e}'_0$.

Step 2. Since z_0 is the central support of e'_0 , note that $xz_0 \in \mathcal{M}z_0 \mapsto xe'_0 \in \mathcal{M}e'_0$ ($x \in \mathcal{M}$) is an isomorphism, and similarly so is $\hat{x}\hat{z}_0 \in \hat{\mathcal{M}}\hat{z}_0 \mapsto \hat{x}\hat{e}'_0 \in \hat{\mathcal{M}}\hat{e}'_0$ ($\hat{x} \in \hat{\mathcal{M}}$). Hence one can define an isomorphism $\Lambda : \mathcal{M}z_0 \rightarrow \hat{\mathcal{M}}\hat{z}_0$ as follows:

$$\Lambda : \mathcal{M}z_0 \cong \mathcal{M}e'_0 \cong \hat{\mathcal{M}}\hat{e}'_0 \cong \hat{\mathcal{M}}\hat{z}_0, \quad xz_0 \mapsto xe'_0 \mapsto V(xe'_0)V^* = \hat{x}\hat{e}'_0 \mapsto \hat{x}\hat{z}_0. \quad (\text{C.4})$$

Note [18, Lemma 2.6] that the standard forms of $\mathcal{M}z_0$ and $\hat{\mathcal{M}}\hat{z}_0$ are respectively given by

$$\begin{aligned} (\mathcal{M}z_0, z_0 L^2(\mathcal{M})z_0 = L^2(\mathcal{M})z_0, J = *, z_0 L^2(\mathcal{M})_+ z_0 = L^2(\mathcal{M})_+ z_0), \\ (\hat{\mathcal{M}}\hat{z}_0, \hat{z}_0 L^2(\hat{\mathcal{M}})\hat{z}_0 = L^2(\hat{\mathcal{M}})\hat{z}_0, \hat{J} = *, \hat{z}_0 L^2(\hat{\mathcal{M}})_+ \hat{z}_0 = L^2(\hat{\mathcal{M}})_+ \hat{z}_0). \end{aligned}$$

By the uniqueness (up to unitary conjugation) of the standard form (under isomorphism) [18, Theorem 2.3], there exists a unitary $U : L^2(\mathcal{M})z_0 \rightarrow L^2(\hat{\mathcal{M}})\hat{z}_0$ such that

$$\Lambda(x) = UxU^*, \quad x \in \mathcal{M}z_0, \quad (\text{C.5})$$

$$(U\xi)^* = U(\xi^*), \quad \xi \in z_0 L^2(\mathcal{M})z_0. \quad (\text{C.6})$$

$$U(L^2(\mathcal{M})_+ z_0) = L^2(\hat{\mathcal{M}})_+ \hat{z}_0. \quad (\text{C.7})$$

Step 3. Since $s(h_0) \leq e_0 \leq z_0$ by assumption, one has $h_0^{1/2}, k_0^{1/2} \in L^2(\mathcal{M})z_0$, and similarly $\hat{h}_0^{1/2}, \hat{k}_0^{1/2} \in L^2(\hat{\mathcal{M}})\hat{z}_0$. By (C.7) one has $Uh_0^{1/2}, Uk_0^{1/2} \in L^2(\hat{\mathcal{M}})\hat{z}_0$. We here confirm that

$$Uh_0^{1/2} = \hat{h}_0^{1/2}, \quad Uk_0^{1/2} = \hat{k}_0^{1/2}. \quad (\text{C.8})$$

To show this, for every $a \in \mathcal{A}$ we find that

$$\begin{aligned}
\langle Uh_0^{1/2}, (\hat{\pi}(a)\hat{z}_0)Uh_0^{1/2} \rangle &= \langle h_0^{1/2}, \Lambda^{-1}(\hat{\pi}(a)\hat{z}_0)h_0^{1/2} \rangle \quad (\text{by (C.5)}) \\
&= \langle h_0^{1/2}, (\pi(a)z_0)h_0^{1/2} \rangle \quad (\text{by (C.3) and (C.4)}) \\
&= \tilde{\rho} \circ \pi(a) = \rho(a) = \hat{\rho} \circ \hat{\pi}(a) \\
&= \langle \hat{h}_0^{1/2}, (\hat{\pi}(a)\hat{z}_0)\hat{h}_0^{1/2} \rangle,
\end{aligned}$$

which implies that $Uh_0^{1/2} = \hat{h}_0^{1/2}$. The proof of $Uk_0^{1/2} = \hat{k}_0^{1/2}$ is similar. By (C.5) and (C.8) we have also

$$\Lambda(x)\hat{h}_0^{1/2} = U(xh_0^{1/2}), \quad \Lambda(x)\hat{k}_0^{1/2} = U(xk_0^{1/2}), \quad x \in \mathcal{M}z_0. \quad (\text{C.9})$$

These imply (C.1). Furthermore, by (C.8) and (C.9) we have $\Lambda(e_0)\hat{k}_0^{1/2} = Uk_0^{1/2} = \hat{k}_0^{1/2}$, from which $\Lambda(e_0) \geq \hat{e}_0$ follows. Applying the same argument to $\Lambda^{-1}(\hat{x}) = U^*\hat{x}U$ ($\hat{x} \in \hat{\mathcal{M}}\hat{z}_0$) with $k_0^{1/2}, \hat{k}_0^{1/2}$ exchanged gives $\Lambda^{-1}(\hat{e}_0) \geq e_0$ as well. Therefore,

$$\Lambda(e_0) = \hat{e}_0. \quad (\text{C.10})$$

Step 4. We define

$$(\Lambda^{-1})_* : L^1(\mathcal{M}z_0) = L^1(\mathcal{M})z_0 \rightarrow L^1(\hat{\mathcal{M}}\hat{z}_0) = L^1(\hat{\mathcal{M}})\hat{z}_0$$

by transforming $\psi \in (\mathcal{M}z_0)_* \mapsto \psi \circ \Lambda^{-1} \in (\hat{\mathcal{M}}\hat{z}_0)_*$ via $L^1(\mathcal{M}z_0) \cong (\mathcal{M}z_0)_*$ and $L^1(\hat{\mathcal{M}}\hat{z}_0) \cong (\hat{\mathcal{M}}\hat{z}_0)_*$, that is, $(\Lambda^{-1})_* : h_\psi \in L^1(\mathcal{M}z_0) \mapsto h_{\psi \circ \Psi^{-1}} \in L^1(\hat{\mathcal{M}}\hat{z}_0)$ for $\psi \in (\mathcal{M}z_0)_*$. Of course, $(\Lambda^{-1})_*$ is an isometry with respect to $\|\cdot\|_1$. Now, Kosaki's (symmetric) interpolation L^p -spaces enter into our discussions. We here confirm that

$$(\Lambda^{-1})_*(k_0^{1/2}xk_0^{1/2}) = \hat{k}_0^{1/2}\Lambda(x)\hat{k}_0^{1/2}, \quad x \in \mathcal{M}z_0. \quad (\text{C.11})$$

Indeed, for every $x, y \in \mathcal{M}z_0$ we find that

$$\begin{aligned}
\text{tr}(\Lambda(y)(\Lambda^{-1})_*(k_0^{1/2}xk_0^{1/2})) &= \text{tr}(yk_0^{1/2}xk_0^{1/2}) = \langle (yk_0^{1/2})^*, xk_0^{1/2} \rangle \\
&= \langle U((yk_0^{1/2})^*), U(xk_0^{1/2}) \rangle \\
&= \langle (\Lambda(y)\hat{k}_0^{1/2})^*, \Lambda(x)\hat{k}_0^{1/2} \rangle \quad (\text{by (C.6) and (C.9)}) \\
&= \text{tr}(\Lambda(y)\hat{k}_0^{1/2}\Lambda(x)\hat{k}_0^{1/2}),
\end{aligned}$$

which yields (C.11).

Step 5. Thanks to (C.11) we see that the isometry $(\Lambda^{-1})_* : L^1(\mathcal{M}z_0) \rightarrow L^1(\hat{\mathcal{M}}\hat{z}_0)$ with respect to $\|\cdot\|_1$ is restricted to an isometry from $k_0^{1/2}(\mathcal{M}z_0)k_0^{1/2}$ (embedded into $L^1(\mathcal{M}z_0)$) onto $\hat{k}_0^{1/2}(\hat{\mathcal{M}}\hat{z}_0)\hat{k}_0^{1/2}$ (embedded into $L^1(\hat{\mathcal{M}}\hat{z}_0)$) with respect to $\|\cdot\|_\infty$, i.e.,

$$\|k_0^{1/2}xk_0^{1/2}\|_\infty = \|x\| = \|\Lambda(x)\| = \|\hat{k}_0^{1/2}\Lambda(x)\hat{k}_0^{1/2}\|_\infty, \quad x \in \mathcal{M}z_0.$$

By Kosaki's construction in [38] (or the Riesz–Thorin theorem) it follows that $(\Lambda^{-1})_*$ gives rise to an isometry

$$\begin{aligned}
(\Lambda^{-1})_* : L^p(\mathcal{M}z_0, \tilde{\sigma}) &= C_{1/p}((k_0^{1/2}(\mathcal{M}z_0)k_0^{1/2}, L^1(\mathcal{M}z_0)) \\
&\rightarrow L^p(\hat{\mathcal{M}}\hat{z}_0, \hat{\sigma}) = C_{1/p}(\hat{k}_0^{1/2}(\hat{\mathcal{M}}\hat{z}_0)\hat{k}_0^{1/2}, L^1(\hat{\mathcal{M}}\hat{z}_0))
\end{aligned}$$

with respect to the interpolation norms $\|\cdot\|_{p,\tilde{\sigma}}$ and $\|\cdot\|_{p,\hat{\sigma}}$ for any $p \in [1, +\infty)$. For every $x \in \mathcal{M}_{z_0}$, applying this to $k_0^{1/2} x k_0^{1/2}$ with (C.11) gives

$$\|k_0^{1/2} x k_0^{1/2}\|_{p,\tilde{\sigma}} = \|\hat{k}_0^{1/2} \Lambda(x) \hat{k}_0^{1/2}\|_{p,\hat{\sigma}}.$$

By [38, Theorem 9.1], for every $p \in [1, +\infty)$ the above equality is rephrased as Haagerup's L^p -norm equality

$$\|k_0^{1/2p} x k_0^{1/2p}\|_p = \|\hat{k}_0^{1/2p} \Lambda(x) \hat{k}_0^{1/2p}\|_p,$$

which is (C.2), as asserted. \square

We are now in a position to prove Theorem 4.3.

Proof of (i). We use the variational expressions in Proposition 2.1 based on Lemma C.1. Assume first that $\alpha > 1$. If $s(\tilde{\rho}) \not\leq s(\tilde{\sigma})$, then $s(\hat{\rho}) \not\leq s(\hat{\sigma})$ (as mentioned at the beginning of this appendix) so that both of $D_\alpha^*(\tilde{\rho}|\tilde{\sigma})$ and $D_\alpha^*(\hat{\rho}|\hat{\sigma})$ are $+\infty$. Hence we assume that $s(\tilde{\rho}) \leq s(\tilde{\sigma})$ (and $s(\hat{\rho}) \leq s(\hat{\sigma})$). Using (2.9) we have

$$\begin{aligned} Q_\alpha^*(\tilde{\rho}|\tilde{\sigma}) &= \sup_{x \in \mathcal{M}_+} \left[\alpha \tilde{\rho}(x) - (\alpha - 1) \operatorname{tr} \left(h_{\tilde{\sigma}}^{\frac{\alpha-1}{2\alpha}} x h_{\tilde{\sigma}}^{\frac{\alpha-1}{2\alpha}} \right)^{\frac{\alpha}{\alpha-1}} \right] \\ &= \sup_{x \in (\mathcal{M}_{z_0})_+} \left[\alpha \tilde{\rho}(x) - (\alpha - 1) \operatorname{tr} \left(h_{\tilde{\sigma}}^{\frac{\alpha-1}{2\alpha}} x h_{\tilde{\sigma}}^{\frac{\alpha-1}{2\alpha}} \right)^{\frac{\alpha}{\alpha-1}} \right] \quad (\text{since } s(\tilde{\rho}) \leq s(\tilde{\sigma}) \leq z_0) \\ &= \sup_{x \in (\mathcal{M}_{z_0})_+} \left[\alpha \hat{\rho}(\Lambda(x)) - (\alpha - 1) \operatorname{tr} \left(h_{\tilde{\sigma}}^{\frac{\alpha-1}{2\alpha}} \Lambda(x) h_{\tilde{\sigma}}^{\frac{\alpha-1}{2\alpha}} \right)^{\frac{\alpha}{\alpha-1}} \right] \quad (\text{by Lemma C.1}) \\ &= \sup_{\hat{x} \in (\hat{\mathcal{M}}_{\hat{z}_0})_+} \left[\alpha \hat{\rho}(\hat{x}) - (\alpha - 1) \operatorname{tr} \left(h_{\tilde{\sigma}}^{\frac{\alpha-1}{2\alpha}} \hat{x} h_{\tilde{\sigma}}^{\frac{\alpha-1}{2\alpha}} \right)^{\frac{\alpha}{\alpha-1}} \right] \\ &= Q_\alpha^*(\hat{\rho}|\hat{\sigma}). \end{aligned}$$

Next, assume that $1/2 \leq \alpha < 1$. When $s(\tilde{\rho}) \leq s(\tilde{\sigma})$, we use (2.10) as above to have

$$\begin{aligned} Q_\alpha^*(\tilde{\rho}|\tilde{\sigma}) &= \inf_{x \in (\mathcal{M}_{z_0})_{++}} \left[\alpha \tilde{\rho}(x) + (1 - \alpha) \operatorname{tr} \left(h_{\tilde{\sigma}}^{\frac{1-\alpha}{2\alpha}} x^{-1} h_{\tilde{\sigma}}^{\frac{1-\alpha}{2\alpha}} \right)^{\frac{\alpha}{1-\alpha}} \right] \\ &= \inf_{x \in (\mathcal{M}_{z_0})_{++}} \left[\alpha \hat{\rho}(\Lambda(x)) + (1 - \alpha) \operatorname{tr} \left(h_{\tilde{\sigma}}^{\frac{1-\alpha}{2\alpha}} \Lambda(x^{-1}) h_{\tilde{\sigma}}^{\frac{1-\alpha}{2\alpha}} \right)^{\frac{\alpha}{1-\alpha}} \right] \quad (\text{by Lemma C.1}) \\ &= \inf_{\hat{x} \in (\hat{\mathcal{M}}_{\hat{z}_0})_{++}} \left[\alpha \hat{\rho}(\hat{x}) + (1 - \alpha) \operatorname{tr} \left(h_{\tilde{\sigma}}^{\frac{1-\alpha}{2\alpha}} \hat{x}^{-1} h_{\tilde{\sigma}}^{\frac{1-\alpha}{2\alpha}} \right)^{\frac{\alpha}{1-\alpha}} \right] \quad (\text{since } \Lambda(x^{-1}) = \Lambda(x)^{-1}) \\ &= Q_\alpha^*(\hat{\rho}|\hat{\sigma}). \end{aligned}$$

For general $\rho, \sigma \in \mathcal{A}_+^*$, let $\sigma_\varepsilon := \sigma + \varepsilon\rho$ for any $\varepsilon > 0$. Then σ_ε has the normal extensions $\tilde{\sigma}_\varepsilon = \tilde{\sigma} + \varepsilon\tilde{\rho}$ to \mathcal{M} and $\hat{\sigma}_\varepsilon = \hat{\sigma} + \varepsilon\hat{\rho}$ to $\hat{\mathcal{M}}$. The above case yields $Q_\alpha^*(\tilde{\rho}|\tilde{\sigma}_\varepsilon) = Q_\alpha^*(\hat{\rho}|\hat{\sigma}_\varepsilon)$ for all $\varepsilon > 0$. From the continuity of Q_α^* on $\mathcal{M}_+^* \times \mathcal{M}_+^*$ in the norm topology when $1/2 \leq \alpha < 1$ (see [26, Theorem 3.16 (3)]), letting $\varepsilon \searrow 0$ gives $Q_\alpha^*(\tilde{\rho}|\tilde{\sigma}) = Q_\alpha^*(\hat{\rho}|\hat{\sigma})$, implying (i). \square

Proof of (ii). Assume first that $s(\tilde{\rho}) \leq s(\tilde{\sigma})$ (hence $s(\hat{\rho}) \leq s(\hat{\sigma})$). Below let us use the same symbols as in the proof of Lemma C.1. Recall [3] that the relative modular operator $\Delta_{\tilde{\rho},\tilde{\sigma}}$ is defined as $\Delta_{\tilde{\rho},\tilde{\sigma}} := S_{\tilde{\rho},\tilde{\sigma}}^* \overline{S_{\tilde{\rho},\tilde{\sigma}}}$, where $S_{\tilde{\rho},\tilde{\sigma}}$ is a closable conjugate linear operator defined by

$$S_{\tilde{\rho},\tilde{\sigma}}(x k_0^{1/2} + \zeta) := e_0 x^* h_0^{1/2}, \quad x \in \mathcal{M}, \quad \zeta \in (L^2(\mathcal{M})e_0)^\perp.$$

Similarly, $\Delta_{\hat{\rho}, \hat{\sigma}} := S_{\hat{\rho}, \hat{\sigma}}^* \overline{S_{\hat{\rho}, \hat{\sigma}}}$ is given, where

$$S_{\hat{\rho}, \hat{\sigma}}(\hat{x} \hat{k}_0^{1/2} + \hat{\zeta}) := \hat{e}_0 \hat{x}^* \hat{h}_0^{1/2}, \quad \hat{x} \in \hat{\mathcal{M}}, \quad \hat{\zeta} \in (L^2(\hat{\mathcal{M}}) \hat{e}_0)^\perp.$$

Since $s(h_0) \leq e_0 \leq z_0$, we can consider $S_{\bar{\rho}, \bar{\sigma}}$ and $\Delta_{\bar{\rho}, \bar{\sigma}}$ as operators on $L^2(\mathcal{M})z_0$ (they are zero operators on $(L^2(\mathcal{M})z_0)^\perp$). Similarly, $S_{\hat{\rho}, \hat{\sigma}}$ and $\Delta_{\hat{\rho}, \hat{\sigma}}$ are considered on $L^2(\hat{\mathcal{M}})\hat{z}_0$. Let us use an isomorphism $\Lambda : \mathcal{M}z_0 \rightarrow \hat{\mathcal{M}}\hat{z}_0$ and a unitary $U : L^2(\mathcal{M})z_0 \rightarrow L^2(\hat{\mathcal{M}})\hat{z}_0$. Since $\overline{\mathcal{M}k_0^{1/2}} = L^2(\mathcal{M})e_0$ and $\overline{\hat{\mathcal{M}}\hat{k}_0^{1/2}} = L^2(\hat{\mathcal{M}})\hat{e}_0$, it follows from (C.9) that $U(L^2(\mathcal{M})e_0) = L^2(\hat{\mathcal{M}})\hat{e}_0$ and hence $U((L^2(\mathcal{M})e_0)^\perp) = (L^2(\hat{\mathcal{M}})\hat{e}_0)^\perp$. For every $\hat{x} = \Lambda(x) \in \hat{\mathcal{M}}\hat{z}_0$ (with $x \in \mathcal{M}z_0$) and $\hat{\zeta} \in (L^2(\hat{\mathcal{M}})\hat{e}_0)^\perp$, we find that

$$\begin{aligned} S_{\hat{\rho}, \hat{\sigma}}(\hat{x} \hat{k}_0^{1/2} + \hat{\zeta}) &= U e_0 U^* U x^* U^* U h_0^{1/2} \quad (\text{by (C.10), (C.5) and (C.8)}) \\ &= U e_0 x^* h_0^{1/2} = U S_{\bar{\rho}, \bar{\sigma}}(x k_0^{1/2} + U^* \hat{\zeta}) \\ &= U S_{\bar{\rho}, \bar{\sigma}}(U^* \hat{x} k_0^{1/2} + U^* \hat{\zeta}) \quad (\text{by (C.9)}) \\ &= U S_{\bar{\rho}, \bar{\sigma}} U^*(\hat{x} \hat{k}_0^{1/2} + \hat{\zeta}). \end{aligned}$$

This implies that $S_{\hat{\rho}, \hat{\sigma}} = U S_{\bar{\rho}, \bar{\sigma}} U^*$ and hence $\Delta_{\hat{\rho}, \hat{\sigma}} = U \Delta_{\bar{\rho}, \bar{\sigma}} U^*$. Therefore, for every $\alpha \in [0, +\infty)$, $\hat{k}_0^{1/2}$ is in $\mathcal{D}(\Delta_{\hat{\rho}, \hat{\sigma}}^{\alpha/2})$ if and only if $k_0^{1/2} = U^* \hat{k}_0^{1/2}$ is in $\mathcal{D}(\Delta_{\bar{\rho}, \bar{\sigma}}^{\alpha/2})$, and in this case,

$$Q_\alpha(\tilde{\rho} \|\tilde{\sigma}) = \|\Delta_{\bar{\rho}, \bar{\sigma}}^{\alpha/2} k_0^{1/2}\|^2 = \|\Delta_{\hat{\rho}, \hat{\sigma}}^{\alpha/2} \hat{k}_0^{1/2}\|^2 = Q_\alpha(\hat{\rho} \|\hat{\sigma}).$$

Otherwise, $Q_\alpha(\tilde{\rho} \|\tilde{\sigma}) = Q_\alpha(\hat{\rho} \|\hat{\sigma}) = +\infty$.

Next assume that $s(\tilde{\rho}) \not\leq s(\tilde{\sigma})$, equivalently $s(\hat{\rho}) \not\leq s(\hat{\sigma})$. Then $Q_\alpha(\tilde{\rho} \|\tilde{\sigma}) = Q_\alpha(\hat{\rho} \|\hat{\sigma}) = +\infty$ for $\alpha > 1$. When $0 \leq \alpha < 1$, let $\sigma_\varepsilon := \sigma + \varepsilon \rho$ for every $\varepsilon > 0$. From the continuity of Q_α on $\mathcal{M}_*^+ \times \mathcal{M}_*^+$ [26, Corollary 3.8], the above case yields

$$Q_\alpha(\tilde{\rho} \|\tilde{\sigma}) = \lim_{\varepsilon \searrow 0} Q_\alpha(\tilde{\rho} \|\tilde{\sigma}_\varepsilon) = \lim_{\varepsilon \searrow 0} Q_\alpha(\hat{\rho} \|\hat{\sigma}_\varepsilon) = Q_\alpha(\hat{\rho} \|\hat{\sigma}),$$

implying (ii). □

Remark C.2. The notion of *standard f -divergences* $S_f(\rho \|\sigma)$ with a parametrization of operator convex functions f on $(0, +\infty)$ has been studied in [24] in the von Neumann algebra setting. From the above proof of (ii) we observe that $S_f(\rho \|\sigma)$ can be extended to $\rho, \sigma \in \mathcal{A}_+^*$ as

$$S_f(\rho \|\sigma) := S_f(\rho_\pi \|\sigma_\pi)$$

independently of the choice of a (ρ, σ) -normal representation π of \mathcal{A} . Then we can easily extend properties of $S_f(\rho \|\sigma)$ given in [24] to the C^* -algebra setting (like Proposition 4.5 for the sandwiched and the standard Rényi divergences).

References

- [1] L. Accardi and C. Cecchini, Conditional expectations in von Neumann algebras and a theorem of Takesaki, *J. Functional Analysis* **45** (1982), 245–273.
- [2] H. Araki, Relative entropy of states of von Neumann algebras, *Publ. Res. Inst. Math. Sci.* **11** (1976), 809–833.

- [3] H. Araki, Relative entropy for states of von Neumann algebras II, *Publ. Res. Inst. Math. Sci.* **13** (1977), 173–192.
- [4] K. M. R. Audenaert, J. Calsamiglia, Ll. Masanes, R. Muñoz-Tapia, A. Acín, E. Bagan and F. Verstraete, Discriminating states: the quantum Chernoff bound, *Phys. Rev. Lett.* **98** (2007), 160501.
- [5] K. M. R. Audenaert and N. Datta, α - z -Rényi relative entropies, *J. Math. Phys.* **56** (2015), 022202, 16 pp.
- [6] J. Bergh and J. Löfström, *Interpolation Spaces: An Introduction*, Springer-Verlag, Berlin-Heidelberg-New York, 1976.
- [7] M. Berta, O. Fawzi and M. Tomamichel, On variational expressions for quantum relative entropies, *Lett. Math. Phys.*, **107** (2017), 2239–2265.
- [8] M. Berta, V. B. Scholz and M. Tomamichel, Rényi divergences as weighted non-commutative vector valued L_p -spaces, *Ann. Henri Poincaré* **19** (2018), 1843–1867.
- [9] O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics 1*, 2nd ed., Springer-Verlag, New York, 1987.
- [10] F. Buscemi and N. Datta, The quantum capacity of channels with arbitrarily correlated noise, *IEEE Trans. Inform. Theory* **56** (2010), 1447–1460.
- [11] M.-A. Choi, A Schwarz inequality for positive linear maps on C^* -algebras, *Illinois J. Math.* **18** (1974), 565–574.
- [12] A. Connes, Classification of injective factors, Cases II_1 , II_∞ , III_λ , $\lambda \neq 1$, *Ann. of Math. (2)* **104** (1976), 73–115.
- [13] I. Csiszár, Generalized cutoff rates and Rényi’s information measures, *IEEE Trans. Inform. Theory*, **41** (1995), 26–34.
- [14] N. Datta, Min- and max-relative entropies and a new entanglement monotone, *IEEE Trans. Inform. Theory* **55** (2009), 2816–2826.
- [15] S. Doplicher, D. Kastler and E. Størmer, Invariant states and asymptotic abelianness, *J. Functional Analysis* **3** (1969), 419–434.
- [16] G. A. Elliott, On approximately finite-dimensional von Neumann algebras. II, *Canad. Math. Bull.* **21** (1978), 415–418.
- [17] R. L. Frank and E. H. Lieb, Monotonicity of a relative Rényi entropy, *J. Math. Phys.* **54** (2013), 122201, 5 pp.
- [18] U. Haagerup, The standard form of von Neumann algebras, *Math. Scand.* **37** (1975), 271–283.
- [19] M. Hayashi, Optimal sequence of quantum measurements in the sense of Stein’s lemma in quantum hypothesis testing, *J. Phys. A: Math. Gen.* **35** (2002), 10759–10773.
- [20] M. Hayashi, Error exponent in asymmetric quantum hypothesis testing and its application to classical-quantum channel coding, *Phys. Rev. A* **76** (2007), 062301.

- [21] M. Hayashi and M. Tomamichel, Correlation detection and an operational interpretation of the Rényi mutual information, *J. Math. Phys.* **57** (2016), 102201, 28 pp.
- [22] F. Hiai, Absolute continuity and disjointness of states in C^* -dynamical systems, *J. Operator Theory* **11** (1984), 319–331.
- [23] F. Hiai, Equality cases in matrix norm inequalities of Golden-Thompson type, *Linear and Multilinear Algebra* **36** (1994), 239–249.
- [24] F. Hiai, Quantum f -divergences in von Neumann algebras I. Standard f -divergences, *J. Math. Phys.* **59** (2018), 102202, 27 pp.
- [25] F. Hiai, Quantum f -divergences in von Neumann algebras II. Maximal f -divergences, *J. Math. Phys.* **60** (2019), 012203, 30 pp.
- [26] F. Hiai, *Quantum f -Divergences in von Neumann Algebras. Reversibility of Quantum Operations*, Mathematical Physics Studies, Springer, Singapore, 2021.
- [27] F. Hiai and M. Mosonyi, Different quantum f -divergences and the reversibility of quantum operations, *Rev. Math. Phys.*, **29** (2017), 1750023, 80 pp.
- [28] F. Hiai, M. Mosonyi and T. Ogawa, Error exponents in hypothesis testing for correlated states on a spin chain, *J. Math. Phys.* **49** (2008), 032112, 22 pp.
- [29] F. Hiai, M. Ohya and M. Tsukada, Sufficiency and relative entropy in $*$ -algebras with applications in quantum systems, *Pacific J. Math.* **107** (1983), 117–140.
- [30] F. Hiai and M. Tsukada, Strong martingale convergence of generalized conditional expectations on von Neumann algebras, *Trans. Amer. Math. Soc.* **282** (1984), 791–798.
- [31] R. B. Israel, *Convexity in the Theory of Lattice Gases*, Princeton University Press, Princeton, NJ, 1979.
- [32] V. Jaksic, Y. Ogata, Y. Pautrat and C.-A. Pillet, Entropic fluctuations in quantum statistical mechanics. An Introduction, *Quantum Theory from Small to Large Scales, August 2010*, Lecture Notes of the Les Houches Summer School **95**, 2012, pp. 213–410, Oxford University Press.
- [33] V. Jaksic, Y. Ogata, C.-A. Pillet and R. Seiringer, Quantum hypothesis testing and non-equilibrium statistical mechanics, *Rev. Math. Phys.* **24** (2012), 1230002, 67 pp.
- [34] A. Jenčová, Rényi relative entropies and noncommutative L_p -spaces, *Ann. Henri Poincaré* **19** (2018), 2513–2542.
- [35] A. Jenčová, Rényi relative entropies and noncommutative L_p -spaces II, *Ann. Henri Poincaré*, Online first, 2021.
- [36] M. Junge and N. Laracuate, Multivariate trace inequalities, p-fidelity, and universal recovery beyond tracial settings, Preprint, 2021, arXiv:2009.11866v2.
- [37] H. Kosaki, Interpolation theory and the Wigner–Yanase–Dyson–Lieb concavity, *Comm. Math. Phys.* **87** (1982), 315–329.
- [38] H. Kosaki, Applications of the complex interpolation method to a von Neumann algebra: non-commutative L^p -spaces, *J. Functional Analysis* **56** (1984), 29–78.

- [39] H. Kosaki, Relative entropy of states: a variational expression, *J. Operator Theory* **16** (1986), 335–348.
- [40] K. Matsumoto, A new quantum version of f -divergence, In: *Reality and Measurement in Algebraic Quantum Theory*, Springer Proc. Math. Stat. **261**, 2018, pp. 229–273, Springer, Singapore.
- [41] M. Mosonyi, The strong converse exponent of discriminating infinite-dimensional quantum states, Preprint, 2021, arXiv:2107.08036.
- [42] M. Mosonyi and F. Hiai, On the quantum Rényi relative entropies and related capacity formulas, *IEEE Trans. Inform. Theory* **57** (2011), 2474–2487.
- [43] M. Mosonyi and F. Hiai, Test-measured Rényi divergences, in preparation.
- [44] M. Mosonyi and T. Ogawa, Quantum hypothesis testing and the operational interpretation of the quantum Rényi relative entropies, *Comm. Math. Phys.* **334** (2015), 1617–1648.
- [45] M. Mosonyi and T. Ogawa, Two approaches to obtain the strong converse exponent of quantum hypothesis testing for general sequences of quantum states, *IEEE Trans. Inform. Theory*, **61** (2015), 6975–6994.
- [46] M. Mosonyi and T. Ogawa, Strong converse exponent for classical-quantum channel coding, *Comm. Math. Phys.*, **355** (2017), 373–426.
- [47] M. Mosonyi and T. Ogawa, Divergence radii and the strong converse exponent of classical-quantum channel coding with constant compositions, *IEEE Trans. Inform. Theory*, **67** (2021), 1668–1698.
- [48] M. Müller-Lennert, F. Dupuis, O. Szehr, S. Fehr and M. Tomamichel, On quantum Rényi entropies: a new generalization and some properties, *J. Math. Phys.* **54** (2013), 122203, 20 pp.
- [49] H. Nagaoka, Strong converse theorems in quantum information theory, *Proceedings of ERATO Workshop on Quantum Information Science*, 2001, page 33. (Also appeared in *Asymptotic Theory of Quantum Statistical Inference*, ed. M. Hayashi, World Scientific, 2005.)
- [50] H. Nagaoka, The converse part of the theorem for quantum Hoeffding bound, Preprint, 2016, arXiv:quant-ph/0611289.
- [51] M. Nussbaum and A. Szkoła, A lower bound of Chernoff type for symmetric quantum hypothesis testing; *Ann. Statist.* **37** (2009), 1040–1057.
- [52] D. Petz, Quasi-entropies for states of a von Neumann algebra, *Publ. Res. Inst. Math. Sci.* **21** (1985), 787–800.
- [53] D. Petz, Quasi-entropies for finite quantum systems; *Rep. Math. Phys.* **23** (1986), 57–65.
- [54] D. Petz, Sufficiency of channels over von Neumann algebras, *Quart. J. Math. Oxford Ser. (2)* **39** (1988), 97–108.
- [55] D. Petz and M. B. Ruskai, Contraction of generalized relative entropy under stochastic mappings on matrices, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **1** (1998), 83–89.

- [56] G. Price, Extremal traces on some group-invariant C^* -algebras *J. Functional Analysis* **49** (1982), 145–151.
- [57] R. Renner, *Security of Quantum Key Distribution*, PhD dissertation, Swiss Federal Institute of Technology Zurich, Diss. ETH No. 16242, (2005).
- [58] A. Rényi, On measures of entropy and information, *Proc. 4th Berkeley Sympos. Math. Statist. and Prob.* **1**, 1961, pp. 547–561, Univ. California Press, Berkeley, CA.
- [59] E. Størmer, Symmetric states of infinite tensor products of C^* -algebras, *J. Functional Analysis* **3** (1969), 48–68.
- [60] D. Sutter, M. Berta and M. Tomamichel, Multivariate trace inequalities, *Comm. Math. Phys.* **352** (2017), , 37–58.
- [61] M. Takesaki, Conditional expectations in von Neumann algebras, *J. Functional Analysis* **9** (1972), 306–321.
- [62] M. Takesaki, *Theory of Operator Algebras I*, Encyclopaedia of Mathematical Sciences, Vol. 124, Springer-Verlag, Berlin, 2002.
- [63] M. Takesaki, *Theory of operator algebras. III*, Encyclopaedia of Mathematical Sciences, Vol. 127, Springer-Verlag, Berlin, 2003.
- [64] M. Takesaki and M. Winnink, Local normality in quantum statistical mechanics, *Comm. Math. Phys.* **30** (1973), 129–152.
- [65] M. Terp, *L^p spaces associated with von Neumann algebras*, Notes, Copenhagen Univ., 1981.
- [66] J. Tomiyama, On the projection of norm one in W^* -algebras, *Proc. Japan Acad.* **33** (1957), 608–612.
- [67] H. Umegaki, Conditional expectation in an operator algebra, IV (entropy and information), *Kōdai Math. Sem. Rep.* **14** (1962), 59–85.
- [68] M. M. Wilde, A. Winter and D. Yang, Strong converse for the classical capacity of entanglement-breaking and Hadamard channels via a sandwiched Rényi relative entropy, *Comm. Math. Phys.* **331** (2014), 593–622.