

Low regularity extensions beyond Cauchy horizons

M. Lesourd* and E. Minguzzi†

October 15, 2021

Abstract

We prove that if in a C^0 spacetime a complete partial Cauchy hypersurface has a non-empty Cauchy horizon, then the horizon is caused by the presence of almost closed causal curves behind it or by the influence of points at infinity. This statement is related to strong cosmic censorship and a conjecture of Wald. In this light, Wald's conjecture can be formulated as a PDE problem about the location of Cauchy horizons inside black hole interiors.

1 Introduction

Penrose introduced strong cosmic censorship in the seminal paper [24]. In an elegant article appearing in the same volume, Geroch and Horowitz [11] elaborated on the differences between the strong and weak formulations of cosmic censorship, and sketched possible approaches one could take to prove such statements. In the context of Lorentzian geometry, they formulated a conjecture which they deemed to be in the spirit of strong cosmic censorship. This conjecture was picked up and sharpened by Wald in Chapter 12 of his landmark book [29].

WALD'S STRONG COSMIC CENSORSHIP. Let (S, h, K) be an initial data set for Einstein's equation, with (S, h) a complete Riemannian manifold and with the Einstein-matter equations in which matter is described by a quasi-linear, diagonal, second order hyperbolic system, and where the stress-energy tensor satisfies the dominant energy condition. Then, if the maximal Cauchy development of this initial data is extendible, for each $p \in \overline{H^+(S)}$ in any extension, either strong causality is violated at p or $\overline{J^-(p)} \cap S$ is non-compact.¹

*Black Hole Initiative, Harvard University, Cambridge, MA 02138. E-mail: mlesourd@fas.harvard.edu

†Università degli Studi di Firenze, Dipartimento di Matematica e Informatica "U. Dini", Via S. Marta 3, I-50139 Firenze, Italia. E-mail: ettore.minguzzi@unifi.it

¹In Wald's book $J^-(p)$ is replaced by $I^-(p)$, but the two formulations are easily shown to be equivalent.

Wald's conjecture seeks to pinpoint why Cauchy horizons form. That is, they are associated with one of two things: the formation of almost closed causal curves, or the possibility that some point on the horizon be influenced by spatial infinity. With regards to the basic examples, in Taub-NUT it is the former, and in Reissner-Nordstrom it is the latter. As for the Kerr-Newman family (with $a \neq 0$), both occur.

With time, the weak and strong conjectures have been translated into PDE problems about the Cauchy development of generic initial data.

MODERN STRONG COSMIC CENSORSHIP. Maximal Cauchy developments of *generic* initial data are inextendible within some *regularity* class.

Remarkable recent work on the subject [3, 6, 15, 16, 18, 19, 28] shows that the SCC is highly sensitive to the setting in which one poses the problem, and the specific definitions of *generic* and *regular* at play. The SCC is now more of a theme of plausible statements (as opposed to a fixed universal statement), each more or less natural relative to the particular setting, and each having varying chances of being true.

In spite of the intricacies that the modern picture has unveiled, the simplicity and clarity of Wald's conjecture, viewed as a structure theorem for Cauchy horizons, grants it an enduring significance. Moreover, with the current emphasis on low regularity extensions, a modern version of Wald's conjecture ought to permit for C^0 extensions.

Before going further, we note that Wald's conjecture cannot hold for positive cosmological constant. A counterexample is provided by the non-extremal Reissner-Nordström de Sitter spacetime. This is a vacuum solution of Einstein's equation with $\Lambda > 0$ which is strongly causal but for which we can find a space-like partial Cauchy hypersurface such that for some $p \in H^+(S)$, $\overline{J^-(p) \cap S}$ is compact (cf. Fig. 1).

Remark 1.1. In [7, Thm. 2.1] Etesi stated a weaker version in which $\overline{J^-(p) \cap S}$ is replaced by $J^-(p) \cap S$. The stronger version is clearly Wald's, since it is easier to prove that a set is non-compact if it can be non-closed. However, consider the next example.

Let (M, g) be the 1+1 Minkowski spacetime of coordinates (t, x) , $g = -dt^2 + dx^2$ with the timelike lines $x = -1$ and $x = +1$ identified and the point $r = (2, 0)$ removed. For $S = t^{-1}(0)$ and $p = (2.5, 0.5)$ we have $p \in H^+(S)$, $J^-(p) \cap S$ is compact but neither $J^-(p) \cap J^+(S)$ nor its closure are compact. The statement which Etesi claims to prove is thus false. The argument fails at [7, Lemma 2.3]: it is not possible to infer the non-compactness of $J^-(p) \cap S$ from that of $J^-(p) \cap J^+(S)$ or its closure by just using causality theory.

In recent years there have been numerous studies under a C^0 assumption. Fathi and Siconolfi studied the existence of time functions under such regularity [8], and Chruściel and Grant [4] presented the first investigation of causality theory under the C^0 assumption. Sbierski [26, 27] studied causality further and proved the C^0 inextendibility of Schwarzschild spacetime. Sämann [25] studied




Figure 1: The Carter-Penrose diagram of the maximal analytic extension of the non-extremal Reissner-Nordström de Sitter spacetime. The partial Cauchy hypersurface S admits a point $p \in H^+(S)$ such that $\overline{J^-(p) \cap J^+(S)}$ is non-compact while $\overline{J^-(p) \cap S}$ is compact. The double line is a singularity.

stable causality and global hyperbolicity and their characterizations. Galloway and Ling [9, 17] proved the C^0 inextendibility of AdS spacetime and showed extendibility through the Big Bang of Milne-like hyperbolic FLRW spacetimes. Chruściel and Klinger obtained a C^0 -inextendibility criterion for expanding singularities [5]. Galloway, Ling and Sbierski [10] proved that in globally hyperbolic spacetimes timelike completeness guarantees C^0 inextendibility, a result improved by Minguzzi and Suhr [23] who showed that global hyperbolicity could in fact be dropped. Bernard and Suhr [1, 2] and Minguzzi [20, 22] obtained several results on regularity of time functions and Cauchy hypersurfaces under even weaker conditions. Grant, Kunzinger, Sämann and Steinbauer clarified properties and pathologies of the chronological future under low regularity [12].

In our study, we shall make repeated use of the causality theory developed by the second author in [20]. The regularity permitted in [20] is based on the framework of closed and proper cone structures, and the results there are valid for Lorentz-Finsler theory under upper semi-continuity conditions on the cone distribution and Finsler fundamental function. Spacetimes endowed with C^0 metrics are special cases of proper cone structures, and so the results on Cauchy horizons developed in that work apply in our case. In fact, it is likely that the results of the present work could hold for the more general proper cone structures of [20], but owing to being motivated by PDEs, we do not see the need to consider lower than C^0 regularity.

Although Wald's formulation involves conditions on the initial data (which guarantees the existence of a unique, up to isometry, maximal Cauchy development), one can isolate the purely Lorentzian geometric component of the statement. In particular, what we prove is the following

Theorem 1.2. *Let (M, g) be a C^0 spacetime and let S be a complete Cauchy C^1*

spacelike hypersurface for M . Suppose that (M, g) has a C^0 extension (M', g') in which S is still acausal and that $H := H^+(S, M') \neq \emptyset$. Let p be a point in H . Then there is a past inextendible geodesic generator γ of the horizon with future endpoint p that either

- (a) it is past imprisoned in a compact subset of the horizon, and so accumulates in the past direction on a compact region where strong causality (actually distinction) is violated,
- (b) it escapes every compact set in the past direction and $\overline{J^-(p) \cap J^+(S)}$ is non-compact.

Note that although (a) implies that strong causality is violated somewhere on the horizon, this need not occur at p . Rather, p is connected to the imprisoning region by way of a null achronal generator.

As for (b), note that strong causality violation can occur without (a) happening at all. For example, every point beyond the Cauchy horizon of maximally extended Kerr is traversed by a closed timelike curve. This means that there cannot be arbitrarily small causally convex neighborhoods around $p \in H^+(S)$. Thus strong causality violation at p can occur in (b) without the imprisonment in (a) occurring.

Note also that we allow for the original spacetime (M, g) to be C^0 , whereas Wald's formulation presumes that (M, g) is at least C^2 . Our reason is simply because we can prove the result in such regularity, and that we find it convenient to use the same regularity for the original and extended spacetimes. In this way we need only to prove²

Theorem 1.3. *Let (M, g) be a C^0 spacetime and let S be a complete acausal C^1 spacelike hypersurface. Let $p \in H := H^+(S)$. Then there is a past inextendible geodesic generator γ of the horizon with future endpoint p that either*

- (a) it is past imprisoned in a compact subset of the horizon, and so accumulates in the past direction on a compact region where strong causality (actually distinction) is violated,
- (b) it escapes every compact set in the past direction and $\overline{J^-(p) \cap J^+(S)}$ is non-compact.

The proof of this statement, which largely follows from the theory developed in [20], is given in the next section. This statement is of course consistent with the examples discussed above, and it seems the best version one can obtain in the context of low regularity causality theory.

Of course, some of the terms need to be clarified. For instance, on S the metric g induces a C^0 Riemannian metric h . By *complete* we mean that the closed balls with respect to the associated distance are compact (we do not have at our disposal the Hopf-Rinow theorem in such low regularity). The notion of

²The spacetime previously denoted M' is now denoted M , that previously denoted M is now $\text{Int}D(S)$.

geodesic generator will also be clarified in the next section.

Finally, in the spirit of Wald's original conjecture, we will also give necessary and sufficient conditions for $\overline{J^-(p)} \cap S$ to be noncompact. We do this in the context of spacetimes to which we can associate a concept of *null infinity* \mathcal{J}^+ . This assumption is inspired by Dafermos-Luk's recent groundbreaking study of strong cosmic censorship in vacuum. Their paper deals with strong cosmic censorship as a problem about the stability of Cauchy horizons in a given class of black hole spacetimes, i.e., spacetimes already possessing a future complete \mathcal{J}^+ along with an event horizon $\partial I^-(\mathcal{J}^+) \neq \emptyset$. This allows them to sidestep the issue of global existence in the exterior region (i.e., weak cosmic censorship guaranteeing a future complete \mathcal{J}^+). In that setting we show the following, where the precise definitions involving \mathcal{J}^+ are given below.

Proposition 1.4. *Let (M, g) be a C^2 spacetime admitting a conformal embedding into the unphysical spacetime (\tilde{M}, \tilde{g}) with null boundary $\mathcal{J}^+ \subset \tilde{M}$ such that $\mathcal{J}^+ \subset \overline{D^+(S)}$ where the closure is taken in the unphysical spacetime (\tilde{M}, \tilde{g}) . Let (M', g') be a C^0 extension of (M, g) with $H^+(S, M') \neq \emptyset$ and $p \in H^+(S, M')$. Then $\overline{J^-(p)} \cap S$ is noncompact if and only if $J^-(p)$ intersects every neighborhood of \mathcal{J}^+ .*

Thus, the character of $\overline{J^-(p)} \cap S$ is really about the location of Cauchy horizons (forming dynamically within black holes) relative to \mathcal{J}^+ .

If, for example, we start with initial data S that differs from a Cauchy surface of the maximal Schwarzschild solution only within a compact subset of the black hole interior, then, even if a Cauchy horizon forms in the development of S , $\overline{J^-(p)} \cap S$ remains compact by domain of dependence.

In view of this, Wald's conjecture can be given a modern reformulation as follows.

CAUCHY HORIZON CONJECTURE. Let (S, g) be a complete Riemannian manifold, and let (S, g, k) be *generic* initial data for the Einstein system, possibly with non-zero cosmological constant Λ , and possibly with matter sources (in which case we may assume various admissibility conditions on T_{ab}). Suppose that the maximal Cauchy development $D(S) = M$ produces an event horizon $\partial I^-(\mathcal{J}^+) \neq \emptyset$. If M admits a C^0 extension $M' \supset M$ and a Cauchy horizon $H^+(S, M') \neq \emptyset$, then

$$\overline{J^-(H^+(S, M'))} \cap \partial I^-(\mathcal{J}^+) \text{ is noncompact}$$

This conjecture can be studied by setting up characteristic initial value problems in which one of the initial data hypersurfaces is an event horizon. This is exactly what is done in [6] (though in [6] they study two-ended data meaning that both hypersurfaces are event horizons), and note that the C^0 extensions constructed in [6] do indeed satisfy the above statement.

Finally, we note that since it is in principle possible that portions of $H^+(S)$ differ in the regularity of permitted extensions, the C^0 qualifier widens the scope

of the conjecture, and that the above statement does not distinguish between $\Lambda <, >, = 0$, though indeed its validity may depend on Λ .

2 Preliminaries and proof

Let us introduce some notations and conventions. The spacetime (M, g) is a smooth connected time-oriented Lorentzian manifold (without boundary) of dimension $n + 1$ (working with C^1 manifolds would have the same generality as every C^1 manifold has a unique smooth compatible structure (Whitney) [14, Thm. 2.9]). The signature of g is $(-, +, \dots, +)$. The spacetime is said to be C^k if g is C^k , $k \geq 0$. The symbol of inclusion is reflexive, $X \subset X$. The interior $\text{Int}X$ of a set $X \subset M$ might be denoted $\overset{\circ}{X}$ for shortness. The reader is referred to [20, 22] for the C^0 causality theory and to [21] for all the other conventions adopted without mention in this work.

Two events are chronologically related if there is a piecewise C^1 timelike curve connecting them. They are causally related if they coincide or there is a continuous causal curve connecting them. The chronological and causal relations are denoted I and J , respectively. We refer to [20, 26] for these concepts. It must be recalled that for C^0 spacetimes we have $I \subset \bar{J}$, and the equality holds for g locally Lipschitz [4, 20].

A set is *achronal* if no pair of points of it belongs to I , and *acausal* if no pair of points of it belongs to $J \setminus \Delta$ where $\Delta = \{(x, x) : x \in M\} \subset M \times M$ is the diagonal. Lightlike geodesics are defined as in [20, Def. 2.6] as continuous causal curves for which locally no pair of points of it belong to \bar{J} (this coincides with local achronality only under a locally Lipschitz assumption on g).

Sets of the form $J^\pm(p) \setminus I^\pm(p)$ have been termed *causal bubbles* [4]. It is now known that they do not exist iff the Kronheimer and Penrose's causal space condition (push up property) $I \circ J \cup J \circ I \subset I$ holds [20, Thm. 2.8] (see also [12, Thm. 2.12]), which is the case, for instance, under a locally Lipschitz condition on g [4, 20].

In the C^0 theory the notion of *edge* of an achronal hypersurface S does not differ from that of the C^2 theory [13, p. 202] [20, Sec. 2.18]. The proof that $\bar{S} \setminus S \subset \text{edge}(S) \subset \bar{S}$, cf. [20, Prop. 2.132] passes to the C^0 theory as it uses just the chronological relation and its openness.

An acausal edgeless (and hence closed) set is a *partial Cauchy hypersurface*. A *Cauchy hypersurface* is a partial Cauchy hypersurface intersected by every inextendible continuous causal curve.

Definition 2.1. Let (M, g) be a C^0 spacetime. The future *domain of dependence* or future *Cauchy development* $D^+(S)$ of a closed and achronal subset $S \subset M$, consists of those $p \in M$ such that every past inextendible continuous causal curve passing through p intersect S . The future *Cauchy horizon* is

$$H^+(S) = \overline{D^+(S)} \setminus I^-(D^+(S)).$$

We shall need the following result proved in [20, Thm. 2.32] for more general “proper cone structures”. The fact that a C^0 spacetime is a proper cone

structure is proved in [20, Prop. 2.4] (and similarly under a locally Lipschitz assumption).

Theorem 2.2. *(Cauchy horizons are generated by lightlike geodesics)*

Let (M, g) be a C^0 spacetime and let S be a closed and acausal hypersurface (possibly with edge). The set $H^+(S) \setminus S$ is an achronal locally Lipschitz topological hypersurface and every $p \in H^+(S)$ is the future endpoint of a lightlike geodesic γ contained in $H^+(S)$, either past inextendible or starting from some point in S . No two points $p, q \in H^+(S) \setminus S$ can be such that $(p, q) \in \overset{\circ}{J}$, thus every continuous causal curve contained in $H^+(S) \setminus S$ is a lightlike geodesic.

Notice that this result does not state that the generators cannot intersect other generators in their interior (as we have in the regular theory).

Now, the main strategy is to study the fate of this generator γ in the past direction. The first step is to show that it cannot reach the edge of S as S has no edge due to the completeness condition. This fact is non-trivial as S might be “lightlike” at the edge and more generally because, though S can be extended as an achronal hypersurface (hence, locally, a Lipschitz graph), it could be the case that no extension is C^1 , so that no sensible notion of space metric could make sense over the extension.³

Lemma 2.3. *Let S be a complete acausal C^1 spacelike hypersurface on a C^0 spacetime. Then $\text{edge}(S) = \emptyset$, thus S is closed.*

Proof. Assume, by contradiction, that there is $p \in \text{edge}(S)$. Let $\{x^a\}$ be local coordinates such that $g(p) = -(dx^0)^2 + \sum_{i=1}^n (dx^i)^2$, ∂_0 is future directed timelike, $x^a(p) = 0$ for every $a = 0, 1, \dots, n$. Observe that S cannot accumulate on any other point of the axis $x^i = 0$ because S is acausal hence achronal.

The bilinear form $\eta := b[-a(dx^0)^2 + \sum_{i=1}^n (dx^i)^2]$ evaluated at p is such that for $a < 1$ its light cone is narrower than that of $g(p)$, moreover, for $b > 1$ the hyperboloid $\{v : \eta(v, v) = 1\}$ does not intersect the corresponding unit hyperboloid for $g(p)$. Let, for instance, $b = a^{-1} = 2$. These properties extend by continuity to a sufficiently small coordinate cylindrical neighborhood C of p . Let $\pi : C \rightarrow D$ be the projection on the disk $D = C \cap \{x^0 = 0\}$, and let the disk have diameter $\delta > 0$ in the Euclidean coordinate metric. The properties imply that if $v \in TC$ is a g -spacelike vector then it is a η -spacelike vector on C and $\sqrt{g(v, v)} \leq \sqrt{\eta(v, v)} \leq |v^\perp|$ where $v^\perp = \pi_*(v)$ is the projection of v on the space orthogonal to ∂_0 and $|\cdot|$ is the coordinate Euclidean space norm. Let $q, r \in S \cap C$. If there is curve in $S \cap C$ connecting q to r then this curve has a length smaller than that of its projection to D .

Let A be a maximal achronal set containing S . Without loss of generality (take a smaller coordinate cylinder C if necessary) we can assume that the set $A \cap C$ is a graph over D (because A is an achronal boundary, so the result

³It should be mentioned that in low regularity there are results on the extension of partial Cauchy hypersurfaces to Cauchy hypersurfaces in globally hyperbolic spacetimes [22, Thm. 2.7]. These theorems cannot be used as they require that the hypersurface be extendible at least in a neighborhood of the edge while preserving causality and regularity properties.

follows from [20, Thm. 2.19]). Let $q, r \in S \cap C$, and let $\bar{q} = \pi(q)$, $\bar{r} = \pi(r)$. The segment $\bar{\sigma}$ connecting \bar{q} to \bar{r} can be lifted to a curve σ in A connecting q to r . If this curve is not entirely contained in S then there is a maximal segment τ with starting point q entirely contained in S and converging to some point in $\bar{S} \setminus S$ such that the projection $\bar{\tau}$ has length less than that of $\bar{\sigma}$ and hence of δ . However, this implies that τ has length less than δ which gives a contradiction with the completeness of S , as on S the closed ball of radius δ centered at q would be non-compact. We conclude that for every $q, r \in S \cap C$ the lift σ is actually contained in S which proves that $\rho(q, r) \leq \delta$ where ρ is the distance of (S, h) . However, completeness of S implies that $\rho(q, r) \rightarrow \infty$ for $r \rightarrow p$, which is not the case as the bound δ is independent of r , that is, it is uniform over $S \cap C$. The contradiction proves that $\text{edge}(S) = \emptyset$. \square

We are ready to prove Theorem 1.3 and hence Theorem 1.2.

Proof of Theorem 1.3. Since S is a complete acausal C^1 spacelike hypersurface, by Lemma 2.3, it is actually closed and $\text{edge}(S) = \emptyset$. This means that S is a topological hypersurface and so, by [20, Thm. 2.33] the generators of $H^+(S)$ do not reach S in the past direction. Also from [20, Thm. 2.34] the set $D^+(S) \setminus S$ is open and $H^+(S) \cap S = \emptyset$. By Theorem 2.2 for every $p \in H^+(S)$ there is a lightlike geodesic γ with future endpoint p which is past inextendible and contained in $H^+(S)$. If $J^-(p) \cap J^+(S)$ has compact closure, then the same is true for the set $J^-(p) \cap H \subset J^-(p) \cap J^+(S)$, then γ is past imprisoned in a compact set contained in the closed set H , and by [20, Thm. 2.22] it accumulates on a compact subset of H over which future and past distinction (and hence strong causality) are violated. \square

Proof of Proposition 1.4. For convenience, write

$$C \equiv \overline{J^-(p) \cap S} \tag{1}$$

By assumption, S has an end $S_{\text{end}} \subset S$, which we define to be that part of S contained in a neighborhood of \mathcal{J}^+ .

First consider \Leftarrow . If $J^-(p)$ intersects every neighborhood of \mathcal{J}^+ , then there is a sequence of points $\{p_i\} \in J^-(p) \cap S_{\text{end}}$ which diverges to infinity in S_{end} ; that is, p_i eventually escapes every compact set of S .

For the converse, we show that if there is a neighborhood of \mathcal{J}^+ that does not intersect $J^-(p)$, then C is compact. If such a neighborhood exists, then we can define a closed subset of S , C' , of the form $C' = S \setminus S_{\text{end}}$ such that $C \subset C'$. Both C and C' are closed, so we need only show that C' is compact. This follows from the fact that S is composed of a compact core and a union of ends. \square

Acknowledgments

M. Lesourd thanks the John Templeton and Gordon and Betty Moore Foundations for their support of Harvard's Black Hole Initiative. E. Minguzzi thanks GNFM of INDAM for partial support.

References

- [1] P. Bernard and S. Suhr. Lyapounov functions of closed cone fields: from Conley theory to time functions. *Commun. Math. Phys.*, 359:467–498, 2018.
- [2] P. Bernard and S. Suhr. Cauchy and uniform temporal functions of globally hyperbolic cone fields. *Proc. Amer. Math. Soc.*, 148:4951–4966, 2020.
- [3] V. Cardoso, J. L. Costa, K. Destounis, P. Hintz, and A. Jansen. Strong cosmic censorship in charged black-hole spacetimes: still subtle. *Phys. Rev. D*, 98:104007, 2018.
- [4] P. T. Chruściel and J. D. E. Grant. On Lorentzian causality with continuous metrics. *Class. Quantum Grav.*, 29:145001, 2012.
- [5] P. T. Chruściel and P. Klinger. The annoying null boundaries. *J. Phys.: Conf. Ser.*, 968:012003, 2018.
- [6] M. Dafermos and J. Luk. The interior of dynamical vacuum black holes I: The C^0 -stability of the Kerr Cauchy horizon. arXiv:1710.01722, 2017.
- [7] G. Etesi. A proof of the Geroch–Horowitz–Penrose formulation of the Strong Cosmic Censor conjecture motivated by computability theory. *Int. J. Theor. Phys.*, 52:946–960, 2013.
- [8] A. Fathi and A. Siconolfi. On smooth time functions. *Math. Proc. Camb. Phil. Soc.*, 152:303–339, 2012.
- [9] G. J. Galloway and E. Ling. Some remarks on the C^0 -(in)extendibility of spacetimes. *Ann. Henri Poincaré*, 18:3427–3447, 2017.
- [10] G. J. Galloway, E. Ling, and J. Sbierski. Timelike completeness as an obstruction to C^0 -extensions. *Commun. Math. Phys.*, 359:937–949, 2018.
- [11] R. Geroch and G. T. Horowitz. *Global structure of spacetimes*, volume General relativity: An Einstein centenary survey, pages 212–292. Cambridge University Press, Cambridge, 1979.
- [12] J. D. E. Grant, M. Kunzinger, C. Sämann, and R. Steinbauer. The future is not always open. *Lett. Math. Phys.*, 110:83–103, 2020. arXiv:1901.07996.
- [13] S. W. Hawking and G. F. R. Ellis. *The Large Scale Structure of Space-Time*. Cambridge University Press, Cambridge, 1973.
- [14] M. W. Hirsch. *Differential topology*. Springer-Verlag, New York, 1976.
- [15] C. Kehle. Diophantine approximation as cosmic censor for Kerr-Ads black holes. arXiv:2007.12614, 2020.
- [16] C. Kehle and M. Van de Moortel. Strong cosmic censorship in the presence of matter: the decisive effect of horizon oscillations on the black hole interior geometry. arXiv:2105.04604, 2021.

- [17] E. Ling. The Big Bang is a coordinate singularity for $k = -1$ inflationary FLRW spacetimes. *Found. Phys.*, 50:385–428, 2020.
- [18] J. Luk and S.-J. Oh. Proof of linear instability of the Reissner-Nordström Cauchy horizon under scalar perturbations. *Duke Math. J.*, 166:437–493, 2017.
- [19] J. Luk and S.-J. Oh. Strong cosmic censorship in spherical symmetry for two-ended asymptotically flat initial data I. The interior of the black hole region. *Ann. Math.*, 190:1–111, 2019.
- [20] E. Minguzzi. Causality theory for closed cone structures with applications. *Rev. Math. Phys.*, 31:1930001, 2019. arXiv:1709.06494.
- [21] E. Minguzzi. Lorentzian causality theory. *Living Rev. Relativ.*, 22:3, 2019.
- [22] E. Minguzzi. On the regularity of Cauchy hypersurfaces and temporal functions in closed cone structures. *Rev. Math. Phys.*, 32:2050033, 2020. arXiv:1909.09352.
- [23] E. Minguzzi and S. Suhr. Some regularity results for Lorentz-Finsler spaces. *Ann. Glob. Anal. Geom.*, 56:597–611, 2019. arXiv:1903.00842.
- [24] R. Penrose. *Singularities and time-asymmetry*, volume General relativity: An Einstein centenary survey, pages 581–638. Cambridge University Press, Cambridge, 1979.
- [25] C. Sämann. Global hyperbolicity for spacetimes with continuous metrics. *Ann. Henri Poincaré*, 17(6):1429–1455, 2016.
- [26] J. Sbierski. The C^0 -inextendibility of the Schwarzschild spacetime and the spacelike diameter in Lorentzian geometry. *J. Diff. Geom.*, 108:319–378, 2018.
- [27] J. Sbierski. On the proof of the C^0 -inextendibility of the Schwarzschild spacetime. *J. Phys. Conf. Ser.*, 968:012012, 2018.
- [28] M. Van de Moortel. Mass inflation and the c^2 -inextendibility of spherically symmetric charged scalar field dynamical black holes. *Commun. Math. Phys.*, 382:1263–1341, 2021.
- [29] R. M. Wald. *General Relativity*. The University of Chicago Press, Chicago, 1984.