

On some batch code properties of the simplex code*

Henk D.L. Hollmann¹, Karan Khathuria¹, Ago-Erik Riet², and Vitaly Skachek¹

¹Institute of Computer Science, University of Tartu, Tartu 50409, Estonia

{henk.d.l.hollmann,karan.khathuria,vitaly.skachek}@ut.ee

²Institute of Mathematics and Statistics, University of Tartu, Tartu 50409, Estonia
ago-erik.riet@ut.ee

October 22, 2021

Abstract

The binary k -dimensional simplex code is known to be a 2^{k-1} -batch code and is conjectured to be a 2^{k-1} -functional batch code. Here, we offer a simple, constructive proof of a result that is “in between” these two properties. Our approach is to relate these properties to certain (old and new) additive problems in finite abelian groups. We also formulate a conjecture for finite abelian groups that generalizes the above-mentioned conjecture.

Keywords: Batch codes - Simplex code - Switch codes - Functional batch code - Finite abelian groups

1 Introduction

A *t*-batch code is a method to store a data record in encoded form on multiple servers in such a way that the bit-values in any batch of t positions from the record can be retrieved by decoding the bit-values in t disjoint groups of positions.

Batch codes were initially introduced in [6] as a method to improve load-balancing in distributed data storage systems. Later, so-called *switch codes* (a special case of batch codes) were proposed in [11] as a method to increase the throughput rate in network switches.

In [9, 10], it was shown that the well-known binary simplex code, a code of length $2^k - 1$, dimension k , and minimum distance 2^{k-1} ($k \geq 1$ integer) is a 2^{k-1} -batch code. The proof of

*An extended abstract of part of this work has been submitted to WCC 2022 (the Twelfth International Workshop on Coding and Cryptography).

that result is somewhat cumbersome, and the algorithm resulting from the proof requires to store and use a database containing all the solutions for the cases where $k \leq 7$. More recently, in [15, 16] the authors conjecture that the k -dimensional simplex code is even a 2^{k-1} -functional batch code. (For precise definitions of this and other used notions, we refer to Section 2.)

In this paper, we give a simple, algorithmic proof of a result that falls halfway between the known result for the simplex code in [9, 10] and the conjecture in [15, 16]. Our approach is to relate the required properties of the simplex code to certain additive problems in finite abelian groups.

The contents of this paper are as follows. In Section 2, we provide precise definitions of all the notions mentioned above, together with precise statements of some known results and conjectures. In our approach, we deal with certain reformulations of these statements, as derived in Section 3. In Section 4 we describe a variation of an algorithm in abelian groups first discovered by Marshall Hall, Jr., with slightly simpler proofs than those given in [5], that we then use to demonstrate our main result.

The problems that we investigate here can be considered as special cases for groups of the form \mathbb{Z}_2^k of problems for general finite abelian groups. In Section 5, we discuss these relations. We formulate a new conjecture for finite abelian groups that would imply the functional batch conjectures if true for groups of the form \mathbb{Z}_2^k , and we prove that this new conjecture holds for the case of cyclic groups of prime order. We end with some conclusions in Section 6.

2 Preliminaries

All codes in this paper are binary and linear. We use \mathbb{F}_2 or \mathbb{Z}_2 to denote the finite field of two elements 0,1, with addition and multiplication modulo 2, and we write \mathbb{F}_2^k for the vector space of dimension k over \mathbb{F}_2 . Thus \mathbb{F}_2^k consists of all binary vectors of length k . The i th unit vector \mathbf{e}_i is the binary vector that has a one in position i and a zero in all other positions. We will write E_k to denote the set of the k unit vectors of length k . For convenience, we commonly number the positions with the integers $1, \dots, k$. The *support* of a vector $\mathbf{v} \in \mathbb{F}_2^k$, written as $\text{supp}(\mathbf{v})$, is the collection of positions where \mathbf{v} has a 1, and the (Hamming) *weight* $w(\mathbf{v})$ of \mathbf{v} is the size of $\text{supp}(\mathbf{v})$.

Definition 2.1 We say that a binary $k \times n$ matrix \mathbf{G} can *serve a request sequence* $\mathbf{r}_1, \dots, \mathbf{r}_t$ of (not necessarily distinct) nonzero vectors in \mathbb{F}_2^k if we can find pairwise disjoint subsets I_1, \dots, I_t of the set of column indices $\{1, 2, \dots, n\}$ of \mathbf{G}_k such that for $j = 1, \dots, t$, the columns of \mathbf{G}_k with index in I_j sum to \mathbf{r}_j .

Note that if the columns of \mathbf{G} with index in I sum up to \mathbf{r} , then for any code word $\mathbf{c} = \mathbf{a}\mathbf{G}$ in the binary linear code generated by \mathbf{G} , the linear combination (\mathbf{r}, \mathbf{a}) of the data bits encoded by \mathbf{c} can be obtained as the sum of the values c_i with $i \in I$. Therefore, Definition 2.1 expresses the property that a sequence of requests for the linear

combinations $(\mathbf{r}_1, \mathbf{a}), \dots, (\mathbf{r}_t, \mathbf{a})$ of the data \mathbf{a} encoded by a code word $\mathbf{c} = \mathbf{a}\mathbf{G}$ can be served *simultaneously* even if the value at each position of the code word can be read at most *once*. We will be interested in various properties of such matrices defined in terms of the particular request sequences that they can serve.

Definition 2.2 The binary $k \times n$ matrix \mathbf{G} (as well as the binary linear code generated by \mathbf{G}) is (i) a *t-PIR code*, (ii) a *t-batch code*, (iii) a *t-odd batch code*, or (iv) a *t-functional batch code* if \mathbf{G} can serve any request sequence of length t consisting of (i) the t -fold repetition of a unit vector in E_k , (ii) unit vectors in E_k only, (iii) vectors in \mathbb{F}_2^k of odd weight only, or (iv) nonzero vectors in \mathbb{F}_2^k , respectively.

The notions of t -PIR code and t -batch code are well known (but note that some authors employ a more general definition and refer to these codes as *multiset primitive*), and together with t -functional batch codes are defined, for example, in [15, 16]. For a recent overview of these and related types of codes, see [7]. The notion of t -odd batch code is new and is introduced here for convenience.

The (binary) *simplex code* of length $n = 2^k - 1$ has a $k \times (2^k - 1)$ generator matrix \mathbf{G}_k whose columns are the distinct nonzero vectors in \mathbb{F}_2^k . In [9] (see also [10]) it was shown that \mathbf{G}_k is a 2^{k-1} -batch code, but the proof is somewhat cumbersome. Recently, it was conjectured that \mathbf{G}_k is even a 2^{k-1} -functional batch code, and it was shown that \mathbf{G}_k is a t -functional batch code for $t = 2^{k-2} + 2^{k-4} + \lfloor 2^{k/2}/\sqrt{24} \rfloor$, again with a rather involved proof [15, 16]. After completion of this paper, we learned that this result was further improved in [13] and [12], where it was shown that G_k is a t -functional batch code for $t = \lfloor (2/3) \cdot 2^{k-1} \rfloor$ and $t = \lfloor (5/6) \cdot 2^{k-1} \rfloor - k$, respectively.

In this paper, we will provide a simple algorithmic proof that \mathbf{G}_k is a 2^{k-1} -odd batch code. In fact, we will prove slightly more.

Theorem 2.3 For every integer $k \geq 1$, the binary simplex code of length $n = 2^k - 1$ and dimension k , with generator matrix \mathbf{G}_k as above, is a 2^{k-1} -odd batch code. In addition, every sequence of 2^{k-1} odd-weight vectors from \mathbb{F}_2^k can be served with column subsets of size at most two.

Note that since every unit vector has odd weight (in fact, a weight equal to 1), Theorem 2.3 implies that the k -dimensional simplex code generated by \mathbf{G}_k is in fact a 2^{k-1} -batch code, a fact that was first proved in [9, 10].

Earlier, we mentioned the conjecture that the simplex code of length $2^k - 1$ is a 2^{k-1} -functional batch code. We believe that even a slightly stronger statement may be true.

Conjecture 2.4 For every integer $k \geq 1$, the binary simplex code of length $n = 2^k - 1$ and dimension k , with generator matrix \mathbf{G}_k as above, can serve every sequence of 2^{k-1} vectors from \mathbb{F}_2^k with column subsets of size at most two.

Now a request $\mathbf{r} \neq \mathbf{0}$ is served by one column \mathbf{x} or by two columns \mathbf{x}, \mathbf{y} from \mathbf{G}_k , precisely when there are two vectors \mathbf{x}, \mathbf{y} with $\mathbf{x} \neq \mathbf{0}$ such that $\mathbf{x} + \mathbf{r} = \mathbf{y}$. So it is easily seen that the above conjecture is in fact equivalent to the following.

Conjecture 2.5 Let $k \geq 1$ be an integer. For every sequence of nonzero vectors $\mathbf{r}_1, \dots, \mathbf{r}_{2^{k-1}}$ in \mathbb{F}_2^k , there are pairwise distinct nonzero vectors $\mathbf{x}_1, \dots, \mathbf{x}_{2^{k-1}}$ in \mathbb{F}_2^k such that the nonzero vectors among $\mathbf{y}_1 = \mathbf{x}_1 + \mathbf{r}_1, \dots, \mathbf{y}_{2^{k-1}} = \mathbf{x}_{2^{k-1}} + \mathbf{r}_{2^{k-1}}$ are pairwise distinct and distinct from $\mathbf{x}_1, \dots, \mathbf{x}_{2^{k-1}}$.

3 A reformulation

We will in fact prove the following slight generalization of Theorem 2.3.

Theorem 3.1 Let $k \geq 1$ be an integer, and let H be a $(k-1)$ -dimensional subspace of \mathbb{F}_2^k . The binary simplex code of length $n = 2^k - 1$ and dimension k , with generator matrix \mathbf{G}_k as above, can serve every sequence of 2^{k-1} vectors from the complement $\mathbb{F}_2^k \setminus H$ of H with column subsets of size at most two.

For later use, we now derive several equivalent formulations of Theorem 3.1. Note that if $\mathbf{x}, \mathbf{y} \in \mathbb{F}_2^k$ with $\mathbf{x} + \mathbf{y} = \mathbf{r}$ and $\mathbf{r} \in \mathbb{F}_2^k \setminus H$, then without loss of generality we may assume that $\mathbf{x} \in \mathbb{F}_2^k \setminus H$ (hence nonzero) and $\mathbf{y} \in H$. So it is easily seen that Theorem 3.1 is in fact equivalent to the following.

Theorem 3.2 Let $k \geq 1$ be an integer, and let H be a $(k-1)$ -dimensional subspace of \mathbb{F}_2^k . For every sequence of vectors $\mathbf{r}_1, \dots, \mathbf{r}_{2^{k-1}}$ in $\mathbb{F}_2^k \setminus H$, there are pairwise distinct vectors $\mathbf{x}_1, \dots, \mathbf{x}_{2^{k-1}}$ in $\mathbb{F}_2^k \setminus H$ such that the nonzero vectors among the vectors $\mathbf{x}_1 + \mathbf{r}_1, \dots, \mathbf{x}_{2^{k-1}} + \mathbf{r}_{2^{k-1}}$ in H are also pairwise distinct.

Of special interest is the case where H and $\mathbb{F}_2^k \setminus H$ are the collection of even and odd weight vectors in \mathbb{F}_2^k , respectively. Note that for this case, Theorem 3.1 reduces to Theorem 2.3.

We will now show the equivalence of Theorem 3.2 and the following.

Theorem 3.3 Let $k \geq 1$ be an integer. Given any sequence $\mathbf{r}_1, \dots, \mathbf{r}_{2^k}$ in \mathbb{F}_2^k , there exists a numbering $\mathbf{x}_1, \dots, \mathbf{x}_{2^k}$ of the vectors in \mathbb{F}_2^k such that the nonzero vectors in the sequence $\mathbf{x}_1 + \mathbf{r}_1, \dots, \mathbf{x}_{2^k} + \mathbf{r}_{2^k}$ are pairwise distinct.

To show this, we need some preparation. Let H be a $(k-1)$ -dimensional subspace of \mathbb{F}_2^k . Then H is of the form

$$H = \mathbf{u}^\perp := \{\mathbf{h} \in \mathbb{F}_2^k \mid h_1 u_1 + \dots + h_k u_k = 0\}$$

for some $\mathbf{u} = (u_1, \dots, u_k) \in \mathbb{F}_2^k \setminus \{0\}$. Now H is a subgroup of $(\mathbb{F}_2^k, +)$ isomorphic to $(\mathbb{F}_2^{k-1}, +)$. Indeed, if $u_j \neq 0$, then it is easily verified that $\varphi : H \rightarrow \mathbb{F}_2^{k-1}$ defined by $\varphi(\mathbf{h}) = (h_1, \dots, h_{j-1}, h_{j+1}, \dots, h_k)$ is an isomorphism. Fix some $\mathbf{a} \in \mathbb{F}_2^k \setminus H$, and extend φ to a linear map $\varphi : \mathbb{F}_2^k \rightarrow \mathbb{F}_2^{k-1}$ by defining $\varphi(\mathbf{a} + \mathbf{h}) = \varphi(\mathbf{h})$ for $\mathbf{h} \in H$. Note that φ also sets up a one-to-one correspondence between $\mathbb{F}_2^k \setminus H$ and \mathbb{F}_2^{k-1} .

Now let $k \geq 2$ be an integer. First, suppose Theorem 3.3 holds for $k-1$, let H be a $(k-1)$ -dimensional subspace of \mathbb{F}_2^k , and let $\mathbf{r}_1, \dots, \mathbf{r}_{2^{k-1}}$ be a sequence in $\mathbb{F}_2^k \setminus H$.

Let $\varphi : \mathbb{F}_2^k \rightarrow \mathbb{F}_2^{k-1}$ be a linear map, 1-1 on both H and $\mathbb{F}_2^k \setminus H$, constructed as discussed above. Put $\mathbf{r}'_i = \varphi(\mathbf{r}_i)$ ($i = 1, \dots, 2^{k-1}$). Applying Theorem 3.3 for $k-1$, we conclude that there is a numbering $\mathbf{x}'_1, \dots, \mathbf{x}'_{2^{k-1}}$ of the vectors in \mathbb{F}_2^{k-1} such that the nonzero vectors among $\mathbf{y}'_1 = \mathbf{x}'_1 + \mathbf{r}'_1, \dots, \mathbf{y}'_{2^{k-1}} = \mathbf{x}'_{2^{k-1}} + \mathbf{r}'_{2^{k-1}}$ are pairwise distinct. Let $\mathbf{x}_i \in \mathbb{F}_2^k \setminus H$, $\mathbf{r}_i \in \mathbb{F}_2^k \setminus H$, and $\mathbf{y}_i \in H$ ($i = 1, \dots, 2^{k-1}$) be the unique vectors such that $\varphi(\mathbf{x}_i) = \mathbf{x}'_i$, $\varphi(\mathbf{r}_i) = \mathbf{r}'_i$, and $\varphi(\mathbf{y}_i) = \mathbf{y}'_i$, respectively. By the linearity of φ , we have that $\mathbf{x}_i + \mathbf{r}_i = \mathbf{y}_i$ for all i ; moreover, since φ is one-to-one on H and on $\mathbb{F}_2^k \setminus H$, with $\mathbf{y}_i = \mathbf{0}$ if and only if $\varphi(\mathbf{y}_i) = \mathbf{y}'_i = 0$, the \mathbf{x}_i are pairwise distinct and the nonzero \mathbf{y}_i are also pairwise distinct. So we conclude that Theorem 3.2 holds for k .

Conversely, suppose that Theorem 3.2 holds for k , and let $\mathbf{r}'_1, \dots, \mathbf{r}'_{2^{k-1}}$ be in \mathbb{F}_2^{k-1} . Let $\mathbf{r}_1, \dots, \mathbf{r}_{2^{k-1}}$ be the unique vectors in $\mathbb{F}_2^k \setminus H$ for which $\varphi(\mathbf{r}_i) = \mathbf{r}'_i$ for all i . Applying Theorem 3.2, we conclude that there are pairwise distinct vectors $\mathbf{x}_1, \dots, \mathbf{x}_{2^{k-1}}$ in $\mathbb{F}_2^k \setminus H$ such that the nonzero vectors among $\mathbf{y}_1 = \mathbf{x}_1 + \mathbf{r}_1, \dots, \mathbf{y}_{2^{k-1}} = \mathbf{x}_{2^{k-1}} + \mathbf{r}_{2^{k-1}}$ are also pairwise distinct. Now let $\mathbf{x}'_j = \varphi(\mathbf{x}_j)$ and $\mathbf{y}'_j = \varphi(\mathbf{y}_j)$ for all j . By linearity of φ , we have $\mathbf{x}'_j + \mathbf{r}'_j = \mathbf{y}'_j$ for all j . Moreover, since φ is one-to-one both on H and on $\mathbb{F}_2^k \setminus H$, the \mathbf{x}'_j are pairwise distinct, hence they form a numbering of the vectors in \mathbb{F}_2^{k-1} , and the nonzero \mathbf{y}'_j are also pairwise distinct. So we conclude that Theorem 3.3 holds for $k-1$. We have proved the following.

Theorem 3.4 For every integer $k \geq 2$, Theorem 3.1, Theorem 3.2, and Theorem 2.3 are all equivalent to Theorem 3.3 (with k replaced by $k-1$).

4 A servicing algorithm

We will now describe an algorithm to solve the numbering problem inherent in Theorem 3.3. We first introduce some terminology.

Definition 4.1 Let $(G, +)$ be a finite abelian group. A *service* for a given sequence r_1, \dots, r_m in G is a collection of pairwise distinct $x_1, \dots, x_m \in G$ such that the m elements $y_1 = x_1 + r_1, \dots, y_m = x_m + r_m$ are also pairwise distinct in G .

We will often think of such a service as a collection of ordered triples

$$(x_1, y_1, r_1), \dots, (x_m, y_m, r_m)$$

with x_1, \dots, x_m pairwise distinct in G , y_1, \dots, y_m pairwise distinct in G , and $x_i + r_i = y_i$ for $i = 1, \dots, m$. The next result is crucial for our approach.

Theorem 4.2 Let $(G, +)$ be a finite abelian group. Given a service x_1, \dots, x_m for the sequence r_1, \dots, r_m in G with $0 \leq m \leq |G| - 2$, and some additional element $r_0 \in G$, we can find an element $x \in G$ distinct from x_1, \dots, x_m such that some permutation of x, x_1, \dots, x_m is a service for r_0, r_1, \dots, r_m .

Proof. We give a constructive proof of this theorem by describing an algorithm that extends a given service consisting of the triples $(x_1, y_1, r_1), \dots, (x_m, y_m, r_m)$ in G of length at most $|G| - 2$ as in the theorem, so with $x_j + r_j = y_j$ for $j = 1, \dots, m$. To this end, let y_{-1}, y_0 be two distinct elements outside $\{y_1, \dots, y_m\}$ (which is possible by our assumption that $m \leq |G| - 2$), define $x_0 = y_0 - r_0$, and set $c = x_0 + y_{-1}$. (Note that we may assume that $x_0 \in \{x_1, \dots, x_m\}$ since otherwise we could extend the service with the new triple (x_0, y_0, r_0) ; however, we will not use this information below.)

Assume that after t steps of our algorithm ($t = 0, 1, \dots$), we have found t triples $(x_1, y_1, r_1), \dots, (x_t, y_t, r_t)$ (after renumbering triples if necessary) from the given service such that the relations $x_j + r_{j-1} = y_{j-2}$ hold for $j = 1, \dots, t$, and in addition,

$$x_j + y_{j-1} = c \tag{1}$$

holds for $j = 0, \dots, t$, see Figure 1. Note that the case $t = 0$ describes the initial situation where we have no triples yet, no relations, and where $x_0 + y_{-1} = c$ by definition of c .




Figure 1: The extension algorithm

Then in step $t + 1$, to extend the list of triples from the given service, we proceed as follows. Define $x = y_{t-1} - r_t$. Note that from the “crossing edges”, we obtain the relation $r_t = y_t - x_t = y_{t-1} - x$, hence

$$x + y_t = x_t + y_{t-1} = c \tag{2}$$

by our assumptions. Depending on where x is situated, we now distinguish several cases. Case 1: $x \notin \{x_1, \dots, x_m\}$. Then the t new triples $(x_1, y_{-1}, r_0), \dots, (x_t, y_{t-2}, r_{t-1})$, the triple (x, y_{t-1}, r_t) , and the $m - t$ old triples $(x_{t+1}, y_{t+1}, r_{t+1}), \dots, (x_m, y_m, r_m)$ together constitute a service for r_0, r_1, \dots, r_m , and we are done.

Case 2: $x \in \{x_{t+1}, \dots, x_m\}$. Then after renumbering triples if necessary, we may assume that $x = x_{t+1}$, where $x_{t+1} + r_{t+1} = y_{t+1}$. By (2), we have $x_{t+1} + y_t = c$, and we have extended the configuration in Figure 1 from t to $t + 1$ triples.

Case 3: $x \in \{x_1, \dots, x_t\}$. We will show that this case cannot occur. Indeed, suppose that $x = x_j$ with $1 \leq j \leq t$. From (1) and (2), we have that $x_j + y_{j-1} = c = x_j + y_t$, hence $y_t = y_{j-1}$ for some $j = 1, \dots, t$, contradicting our assumptions.

Since in any configuration, $t \leq m$ must hold, the algorithm will eventually end in case 1, and thus will produce an extended service. \square

Remark 4.3 In the case where $G = \mathbb{F}_2^k$, we can say slightly more. Let us define a *special service* for a sequence of *nonzero* vectors $\mathbf{r}_1, \dots, \mathbf{r}_m$ in \mathbb{F}_2^k as a sequence of triples $(\mathbf{x}_i, \mathbf{y}_i, \mathbf{r}_i)$ with $\mathbf{x}_i + \mathbf{y}_i = \mathbf{r}_i$ for $i = 1, \dots, m$ for which $\mathbf{x}_1, \dots, \mathbf{x}_m, \mathbf{y}_1, \dots, \mathbf{y}_m$ are pairwise distinct. (Note that Conjecture 2.5 involves a service notion similar to that of a special service.) Consider again the algorithm in the proof of Theorem 4.2, now starting with a special service of length m with $2m \leq 2^k - 2$. For the newly constructed vector \mathbf{x} , again we have to distinguish various cases. In case 1, \mathbf{x} is not contained in $\{\mathbf{x}_1, \dots, \mathbf{x}_m, \mathbf{y}_{-1}, \mathbf{y}_0, \mathbf{y}_1, \dots, \mathbf{y}_m\}$ and we can extend our special service. In case 2, \mathbf{x} is contained in $\{\mathbf{x}_{t+1}, \dots, \mathbf{x}_m, \mathbf{y}_{t+1}, \dots, \mathbf{y}_m\}$; due to the symmetry between the \mathbf{x}_i and \mathbf{y}_i for $i > t$, we may assume (after renumbering triples if necessary) that $\mathbf{x} = \mathbf{x}_{t+1}$ and we can extend the configuration. In case 3, we assume that \mathbf{x} is contained in $\{\mathbf{x}_1, \dots, \mathbf{x}_t\}$, and just as before, we show that this case cannot occur. Finally, in case 4, we assume that $\mathbf{x} = \mathbf{y}_j$ for $-1 \leq j \leq t$. Now if $\mathbf{x} = \mathbf{y}_t$, then from the known relations we find that $\mathbf{x}_t = \mathbf{y}_{t-1}$, contradicting our assumptions. And if $\mathbf{x} = \mathbf{y}_j$ with $-1 \leq j \leq t - 1$, then combining known relations we find that $\mathbf{y}_j + \mathbf{y}_t = \mathbf{c} = \mathbf{x}_{j+1} + \mathbf{y}_j$, so $\mathbf{y}_t = \mathbf{x}_{j+1}$, contradicting our assumptions again, unless $j = -1$. We conclude that the only way in which our extension algorithm for a special service can go wrong is where $\mathbf{x} = \mathbf{y}_{-1}$, $\mathbf{y}_t = \mathbf{x}_0$, and $t \geq 1$ (since $\mathbf{r}_0 \neq 0$); it works “almost”, but not always. Unfortunately, we have not been able to use this property to find some other way to extend a special service.

Theorem 4.2 has an interesting consequence.

Theorem 4.4 Let $(G, +)$ be an finite abelian group of size n and let $r = r_1, \dots, r_n$ be a sequence in G . Then there is a numbering x_1, \dots, x_n of the elements of G such that $x_1 + r_1, \dots, x_n + r_n$ form a permutation of the elements of G if and only if $r_1 + \dots + r_n = 0$.

Proof. If (x_i, r_i, y_i) ($i = 1, \dots, n$) is a service for r , then the x_i and the y_i are both a permutation of G , so if $x_i + r_i = y_i$ for all i , then $\sum r_i = \sum y_i - \sum x_i = 0$. So the condition on the r_i is necessary. Conversely, suppose that $\sum r_i = 0$. Using Theorem 4.2, we can construct a service $(x_1, r_1, y_1), \dots, (x_{n-1}, r_{n-1}, y_{n-1})$ for r_1, \dots, r_{n-1} . Let g denote the sum of all the elements of G . If x_n and y_n are the elements in G that do not occur among x_i and y_i ($i = 1, \dots, n - 1$), respectively, then $x_n = g - \sum_{i \neq n} x_i$, $y_n = g - \sum_{i \neq n} y_i$ and $r_n = -\sum_{i \neq n} r_i$, and since $x_i + r_i = y_i$ for $i = 1, \dots, n - 1$, we conclude that in addition $x_n + r_n = y_n$. \square

This theorem (with a slightly different proof, but employing essentially the same algorithm) was first stated in [5]. We can also employ Theorem 4.2 to prove Theorem 3.3.

Proof of Theorem 3.3: Given a sequence $\mathbf{r}_1, \dots, \mathbf{r}_{2^k}$, we can use Theorem 4.2 for the group $(\mathbb{F}_2^k, +)$ repeatedly to construct pairwise distinct vectors $\mathbf{x}_1, \dots, \mathbf{x}_{2^k-1} \in \mathbb{F}_2^k$ such that (after renumbering the \mathbf{r}_i 's if necessary) the vectors $\mathbf{y}_1 = \mathbf{x}_1 + \mathbf{r}_1, \dots, \mathbf{y}_{2^k-1} = \mathbf{x}_{2^k-1} + \mathbf{r}_{2^k-1}$ are also pairwise distinct. Let \mathbf{x}_{2^k} denote the element such that $\{\mathbf{x}_1, \dots, \mathbf{x}_{2^k}\} = \mathbb{F}_2^k$. For every vector $\mathbf{a} \in \mathbb{F}_2^k$, the triples $(\mathbf{x}_i + \mathbf{a}, \mathbf{y}_i + \mathbf{a}, \mathbf{r}_i)$ for $i = 1, \dots, 2^k - 1$ again form a service for $\mathbf{r}_1, \dots, \mathbf{r}_{2^k-1}$, so by choosing $\mathbf{a} = \mathbf{x}_{2^k} + \mathbf{r}_{2^k}$ and replacing \mathbf{x}_i and \mathbf{y}_i by

$\mathbf{x}_i + \mathbf{a}$ and $\mathbf{y}_i + \mathbf{a}$, we may assume without loss of generality that $\mathbf{x}_{2^k} = \mathbf{r}_{2^k}$. Now define $\mathbf{y}_{2^k} = \mathbf{0}$ and add the triple $(\mathbf{r}_{2^k}, \mathbf{0}, \mathbf{r}_{2^k})$ as the last triple to complete the service to one for $\mathbf{r}_1, \dots, \mathbf{r}_{2^k}$. \square

We have now proved Theorem 3.3; in view of Theorem 3.4, this implies Theorem 3.2, Theorem 3.1, and our main result Theorem 2.3.

5 Relation with other additive problems in finite abelian groups

We have seen that Theorem 2.3 and its reformulation in Theorem 3.3 are closely related to Hall's result in Theorem 4.4 for finite abelian groups via the notion of a service. In Remark 4.3 we introduced the notion of a special service in the group \mathbb{F}_2^k , which is the kind of service required in Conjecture 2.5. We extend this definition to general abelian groups in the obvious way.

Definition 5.1 Let $(G, +)$ be a finite abelian group. A *special service* for a given sequence r_1, \dots, r_m in $G \setminus \{0\}$ is a collection of pairwise distinct $x_1, \dots, x_m \in G$ such that the m elements $y_1 = x_1 + r_1, \dots, y_m = x_m + r_m$ are pairwise distinct and also distinct from x_1, \dots, x_m .

We believe that the corresponding statement for special services in general abelian groups should be the following.

Conjecture 5.2 Let $(G, +)$ be a finite abelian group. There exists a special service for every sequence r_1, \dots, r_m in $G \setminus \{0\}$ provided that $2m \leq |G| - 1$.

As far as we know, this conjecture is new. The relevance of this conjecture is given by the following.

Theorem 5.3 Conjecture 5.2 for the group $(G, +) = (\mathbb{F}_2^k, +)$ implies Conjecture 2.5.

Proof. Suppose that Conjecture 5.2 holds for the case where $(G, +) = (\mathbb{F}_2^k, +)$. Let $\mathbf{r} = \mathbf{r}_1, \dots, \mathbf{r}_{2^{k-1}}$ be a sequence in $\mathbb{F}_2^k \setminus \{0\}$ as in the statement of Conjecture 2.5. According to our assumption, there are $\mathbf{x}_1, \dots, \mathbf{x}_{2^{k-1}-1}$ such that if $\mathbf{y}_i = \mathbf{x}_i + \mathbf{r}_i$ for $i = 1, \dots, 2^{k-1}-1$, then the vectors $\mathbf{x}_1, \dots, \mathbf{x}_{2^{k-1}-1}, \mathbf{y}_1, \dots, \mathbf{y}_{2^{k-1}-1}$ are pairwise distinct. Now by replacing, if necessary, \mathbf{x}_i by $\mathbf{x}_i + \mathbf{a}$ and \mathbf{y}_i by $\mathbf{y}_i + \mathbf{a}$ for all i , for a suitable $\mathbf{a} \in \mathbb{F}_2^k$, we may assume without loss of generality that $\mathbf{r}_{2^{k-1}}$ does not occur among the $2^k - 2$ vectors $\mathbf{x}_1, \dots, \mathbf{x}_{2^{k-1}-1}, \mathbf{y}_1, \dots, \mathbf{y}_{2^{k-1}-1}$. Then set $\mathbf{x}_{2^{k-1}} = \mathbf{r}_{2^{k-1}}$ and $\mathbf{y}_{2^{k-1}} = \mathbf{0}$. It is now easily verified that the sequence $\mathbf{x}_1, \dots, \mathbf{x}_{2^k}$ indeed satisfies the requirements in Conjecture 2.5. \square

Remark 5.4 In [12, Section VIII] the authors state they believe that every request sequence $r_1, \dots, r_{2^{k-1}}$ can be served by what they refer to as a *Hadamard solution*. This statement is equivalent to the claim that the simplex generator matrix G_k can serve every sequence of 2^{k-1} non-zero vectors from \mathbb{F}_2^k with column subsets of size one or two, of which as most two subsets have size one. Inspection of the proof of Theorem 5.3 shows that the truth of Conjecture 5.2 for the group $(G, +) = (\mathbb{F}_2^k, +)$ indeed also implies this stronger statement.

In the remainder of this section, we will discuss Conjecture 5.2 and its relation to other additive problems in abelian groups. Note that the statement in Conjecture 5.2 does not always hold when $2m = |G|$. Indeed, if $G = \mathbb{Z}_2 \oplus \mathbb{Z}_2$ and the sequence is given by $r_1 = (0, 1)$ and $r_2 = (1, 0)$, then without loss of generality we can choose $x_1 = (0, 0)$, but now there is no valid choice for x_2 . Note also that we have to require that every r_i is nonzero since otherwise x and $x + r_i$ will be equal for every $x \in G$. For later reference, we note the following simple result.

Theorem 5.5 The statement in Conjecture 5.2 holds under the weaker condition that $4m \leq |G| + 3$.

Proof. Let $r_1, r_2, \dots \in G \setminus \{0\}$, and suppose that for some integer $k \geq 1$, the sequence x_1, \dots, x_{k-1} is a special service for r_1, \dots, r_{k-1} . To extend this to a special service for r_1, \dots, r_k , we need to choose $x_k \in G$ such that $x_k \notin x_i + \{0, r_i, -r_k, r_i - r_k\}$ for $i = 1, \dots, k-1$, which is obviously possible if $4(k-1) < |G|$. We conclude that we can construct a special service for r_1, \dots, r_m by repeated extension provided that $4(m-1) < |G|$, or, equivalently, $4m \leq |G| + 3$. \square

Conjecture 5.2 seems somewhat similar in nature to Snevily's conjecture and its generalizations. Here Snevily's conjecture [8], first proved in [3], is the following.

Theorem 5.6 Let $(G, +)$ be a finite abelian group of odd order. For every $X \subseteq G$ with $|X| = m$ and for every set $\{r_1, \dots, r_m\} \subseteq G$, there is a numbering x_1, \dots, x_m of the elements of X such that the sums $x_1 + r_1, \dots, x_m + r_m$ are pairwise distinct.

First, Alon [2] (for the case $a = 1$) and then Dasgupta *et al.* [4] (for general a) proved the following generalization of Snevily's conjecture for certain groups.

Theorem 5.7 Let p be an odd prime, let a be a positive integer, and let G be \mathbb{Z}_{p^a} or $(\mathbb{Z}_p)^a$. Let r_1, \dots, r_m , $m < p$, be a sequence of (not necessarily distinct) elements in G . Then for every $X \subseteq G$ with $|X| = m$, there is a numbering x_1, \dots, x_m of the elements of X such that the sums $x_1 + r_1, \dots, x_m + r_m$ are pairwise distinct.

Alon also noted counterexamples to more general statements, notably the cases $G = \mathbb{Z}_{ms}$, $r_1 = \dots = r_{m-1} = 0$, $r_m = s$ and $X = \{0, s, \dots, (m-1)s\}$ and $G = \mathbb{Z}_m$, $X = G$, and $r_1 = \dots = r_{m-1} = 0$, $r_m = 1$. The proof by Alon and Dasgupta *et al.* of Theorem 5.7 uses the following powerful algebraic result called the ‘‘Combinatorial Nullstellensatz’’.

Theorem 5.8 (Alon [1]) Let \mathbb{F} be an arbitrary field and let $f = f(x_1, \dots, x_m)$ be a polynomial in $\mathbb{F}[x_1, \dots, x_m]$. Suppose that there is a monomial $\prod_{i=1}^m x_i^{t_i}$ of f such that $\sum t_i$ equals the total degree of f and whose coefficient in f is nonzero. Then, if S_1, \dots, S_m are subsets of \mathbb{F} with $|S_i| > t_i$, there are $s_1 \in S_1, \dots, s_m \in S_m$ such that $f(s_1, \dots, s_m) \neq 0$.

Here we mimic Alon's proof to show that Conjecture 5.2 holds in the case where G is cyclic of prime order.

Theorem 5.9 Conjecture 5.2 holds in the case where $G = \mathbb{Z}_p$ with p prime.

Proof. Let $\mathbb{F} = \mathbb{F}_p$, where p is a prime, and consider the polynomial f in $\mathbb{F}[x_1, \dots, x_m]$ defined by

$$f(x_1, \dots, x_m) = \prod_{1 \leq i < j \leq m} (x_i - x_j)(x_i - x_j + r_i - r_j) \prod_{\substack{i, j=1 \\ i \neq j}}^m (x_i - x_j - r_j), \quad (3)$$

where $r_1, \dots, r_m \in \mathbb{F} \setminus \{0\}$. Note that we have $f(x_1, \dots, x_m) \neq 0$ precisely when x_1, \dots, x_m are pairwise distinct, $y_1 = x_1 + r_1, \dots, y_m = x_m + r_m$ are pairwise distinct, and y_1, \dots, y_m are distinct from x_1, \dots, x_m , where $x_i \neq y_i$ since $r_i \neq 0$ ($i = 1, \dots, m$), that is, when x_1, \dots, x_m is a special service for r_1, \dots, r_m in the group $(\mathbb{F}, +)$. So it is sufficient to prove that f has a nonzero evaluation in \mathbb{F}^m . We have that

$$D = \deg(f) = \deg \prod_{1 \leq i < j \leq m} (x_i - x_j)^4 = 2m(m-1). \quad (4)$$

Thus the monomial $(x_1 \cdots x_m)^{2(m-1)}$ has degree D , and we can apply Theorem 5.8 with $t_i = 2(m-1)$ for all i . Now by a special case of Dyson's Conjecture, see [14] and the references therein, we know that the coefficient of $(x_1 \cdots x_m)^{2(m-1)}$ in f , hence in $\prod_{1 \leq i < j \leq m} (x_i - x_j)^4$, is equal to $(2m)!/2^m$. Since $(2m)!/2^m = 1 \cdot 3 \cdots (2m-1) \cdot m!$, we easily see that this coefficient is nonzero in \mathbb{F}_p precisely when $m = 1$ or $p \geq 2m + 1$. Taking $S_i = \mathbb{F}$, we have $p = |S_i| > t_i = 2m - 2$ for all i , and hence by Theorem 5.8, f indeed has a nonzero evaluation. \square

Unfortunately, in order to prove Conjecture 2.4, we need the case $G = \mathbb{F}_2^k$ of Conjecture 5.2, which requires the use of the field $\mathbb{F} = \mathbb{F}_{2^k}$; however the above proof clearly fails in characteristic 2.

Remark 5.10 In fact it is not difficult to determine exactly what can be said using the above proof technique in the field $\mathbb{F} = \mathbb{F}_{2^k}$. Indeed, by the famous Vandermonde identity

$$\prod_{1 \leq i < j \leq m} (x_i - x_j) = \det(x_i^{m-j}) = \sum_{\pi \in \mathfrak{S}_m} (-1)^{\sigma(\pi)} x_{\pi(1)}^{m-1} x_{\pi(2)}^{m-2} \cdots x_{\pi(m-1)}^1$$

(see, e.g., [14]), where \mathfrak{S}_m denotes the group of all permutations on $\{1, \dots, m\}$. Noting that the finite field \mathbb{F}_{2^k} has characteristic 2, we conclude from (4) that the monomials of the polynomial f in (3) all have the same total degree $2m(m-1)$ and are of the form

$$x_{\pi(1)}^{4(m-1)} x_{\pi(2)}^{4(m-2)} \cdots x_{\pi(m-1)}^4$$

for a permutation $\pi \in \mathfrak{S}_m$. Taking $t_i = 4(m-i)$ and $S_i = \mathbb{F}$ ($i = 1, \dots, m$), we see from Theorem 5.8 that f has a nonzero evaluation over \mathbb{F}_{2^k} provided that $2^k \geq 4(m-1)+1$. We conclude that this proof technique, when applied to the group $G = \mathbb{Z}_2^k$, cannot improve on the trivial result in Theorem 5.5.

Remark 5.11 Some progress towards Conjecture 5.2 can be made for groups of the form $(G, +) = (\mathbb{Z}_{p^k}, +)$ with p prime and $k \geq 1$ integer, by considering the coefficients of other monomials of the polynomial f in (3) over the field $\mathbb{F} = \mathbb{F}_q$ with $q = p^k$. For example, considering f as a polynomial over \mathbb{Q} , for $m = 3$, the monomial $x_1^6 x_2^5 x_3$ has coefficient 8 in f , and for $m = 4$, the monomial $x_1^8 x_2^8 x_3^7 x_4$ has coefficient -72 in f . As a consequence, for such groups Conjecture 5.2 holds for $m = 3$ if $p \geq 3$, and for $m = 4$ if $p \geq 5$.

Remark 5.12 Obviously Conjecture 5.2 holds for $m = 1$, and it also holds for $m = 2$. Indeed, given $r_1, r_2 \in G \setminus \{0\}$ and $x_1 \in G$, the collection x_1, x_2 serves r_1, r_2 provided that $x_1, x_2, x_1 + r_1, x_2 + r_2$ are pairwise distinct, which is the case if x_2 is not contained in $\{x_1, x_1 - r_2, x_1 + r_1, x_1 + r_1 - r_2\}$. So if $|G| \geq 5$, there remains at least one valid choice for x_2 . In a similar way, a tedious analysis shows that Conjecture 5.2 also holds for $m = 3$; we omit the details.

6 Conclusions

We related various known and conjectured batch-type properties of the simplex codes to certain additive problems in finite abelian groups. By applying known methods for these more general problems we obtained a simple, constructive proof of a generalization of the theorem that the k -dimensional binary simplex code is a 2^{k-1} -batch code. Recently, it was conjectured that the k -dimensional binary simplex code is even a 2^{k-1} -functional batch code. Here we proposed a new conjecture for finite abelian groups, resembling Snevily's conjecture (now a theorem), that includes a slightly stronger version of that simplex conjecture as a special case, and we proved this new conjecture for cyclic groups of prime order.

Acknowledgments

We thank Lev Yohananov and Eitan Yaakobi for sharing their unpublished work [12] with us.

This research was supported by the Estonian Research Council grants PRG49 and PSG114, and by the European Regional Development Fund via CoE project EXCITE.

References

- [1] N. Alon. Combinatorial nullstellensatz. *Combinatorics, Probability and Computing*, pages 7–29, 1999.
- [2] N. Alon. Additive Latin transversals. *Isr. J. Math.*, pages 125–130, 2000.
- [3] B. Arsovski. A proof of Snevily’s Conjecture. *Isr. J. Math.*, pages 505–508, 2011.
- [4] S. Dasgupta, Gy. Károlyi, O. Serra, and B. Szegedy. Transversals of additive Latin squares. *Israel J. Math.*, 126:17–28, 2001.
- [5] M. Hall, Jr. A combinatorial problem on abelian groups. *Proc. Amer. Math. Soc.*, 3(4):584–587, 1952.
- [6] Y. Ishai, E. Kushilevitz, R. Ostrovsky, and A. Sahai. Batch codes and their applications. In *Proc. 36th ACM Symposium on Theory of Computing (STOC)*, Chicago, pages 1057–1061, June 2004.
- [7] V. Skachek. Batch and PIR codes and their connections to locally repairable codes. In M. Greferath, M. O. Pavčević, N. Silberstein, and M. Ángeles Vázquez-Castro, editors, *Network Coding and Subspace Designs*, pages 427–442. Springer, 2018.
- [8] H. S. Snevily. The Cayley addition table of \mathbb{Z}_n . *Amer. Math Monthly*, pages 584–585, 1999.
- [9] Z. Wang, H. M. Kiah, and Y. Cassuto. Optimal binary switch codes with small query size. In *Proc. 2015 IEEE International Symposium on Information Theory (ISIT)*, pages 636–640, Hong Kong, 2015.
- [10] Z. Wang, H. M. Kiah, Y. Cassuto, and J. Bruck. Switch codes: Codes for fully parallel reconstruction. *IEEE Trans. on Inform. Theory*, 63(4):2061–2070, 2017.
- [11] Z. Wang, O. Shaked, Y. Cassuto, and J. Bruck. Codes for network switches. In *Proc. IEEE International Symposium on Information Theory (ISIT)*, pages 1057–1061, Istanbul, 2013.
- [12] L. Yohananov and E. Yaakobi. Almost optimal construction of functional batch codes using Hadamard codes. submitted to *IEEE Trans. on Inform. Theory*, 2021.
- [13] L. Yohananov and E. Yaakobi. Almost optimal construction of functional batch codes using Hadamard codes. In *Proc. IEEE Symp. Information Theory (ISIT)*, pages 3139–3144, Melbourne, 2021.
- [14] D. Zeilberger. A combinatorial proof of Dyson’s conjecture. *Discrete Math.*, pages 317–321, 1982.

- [15] Y. Zhang, T. Etzion, and E. Yaakobi. Bounds on the length of functional PIR and batch codes. *IEEE Trans. on Inform. Theory*, pages 4917–4934, 2020.
- [16] Y. Zhang, E. Yaakobi, and T. Etzion. Bounds on the length of functional PIR and batch codes. In *Proc. IEEE International Symposium on Information Theory*, pages 2129–2133, Paris, 2019.