

RESOLVENT EXPANSIONS FOR SELF-ADJOINT OPERATORS VIA BOUNDARY TRIPLETS

YURI LATUSHKIN AND SELIM SUKHTAIEV

ABSTRACT. In this paper we develop certain aspects of perturbation theory for self-adjoint operators subject to small variations of their domains. We use the abstract theory of boundary triplets to quantify such perturbations and give the second order asymptotic analysis for resolvents, spectral projections, discrete eigenvalues of the corresponding self-adjoint operators. In particular, we derive explicit formulas for the first variation and the Hessian of the eigenvalue curves bifurcating from a discrete eigenvalue of an unperturbed operator. An application is given to a matrix valued Robin Laplacian and more general Robin-type self-adjoint extensions.

1. INTRODUCTION

This paper concerns asymptotic perturbation theory for self-adjoint extensions of symmetric operators. Our main goal is to derive explicit second order Taylor expansions for resolvents, spectral projections, and eigenvalue curves for one-parameter families of self-adjoint extensions of a given symmetric operator. The type of admissible perturbations covers, in particular, variation of boundary conditions for quantum graph operators and, more generally, variation of domain of self-adjointness for extensions of abstract symmetric operators. Such non-additive perturbations arise, for example, in the geometric spectral theory, in which case ad-hoc methods are used to relate, say, the Hessian of parameter-dependent eigenvalues to the reduced Dirichlet-to-Neumann map, see, for example, recent manuscripts [3], [4], [5], [15]. We provide a complete operator theoretic treatment for the second order expansion problem which, in particular, yields an abstract version of the reduced Dirichlet-to-Neumann map and its relation to the Hessian of eigenvalue curves. The first and second order expansion formulas go back to Hadamard, Rayleigh, and Rellich and have been closely studied subsequently, cf. e.g. [1, 7, 14, 16, 17]

The first and second order asymptotic expansions discussed in this paper are also central in the Maslov index theory, cf. [6, 23], for partial differential operators on varying domains or subject to varying boundary conditions, a subject that recently attracted much attention, cf. [8, 9, 10, 11, 19, 20] and the bibliography therein. For example, Deng–Jones’s [11] foundational approach to the spectral count via Maslov index for Robin Laplacian essentially reduces to finding the sign of the derivative of the eigenvalue curves, see recent papers [8, 20]. Further developments of this approach, see [10], led to elaborate spectral flow formulas and provided new tools in geometric analysis of Laplace eigenfunctions.

Date: October 15, 2021.

This work naturally emerged from [19] and [5] as the result numerous stimulating discussions with Prof. G. Berkolaiko. We thank him for generously sharing his ideas as well as his notes on related topics. We wish to thank Prof. P. Kuchment for pointing out many references. Y.L. was supported by the NSF grants DMS-1710989 and DMS-2106157, and would like to thank the Courant Institute of Mathematical Sciences and especially Prof. Lai-Sang Young for the opportunity to visit CIMS.

We stress that the perturbation theory developed here covers not merely “additive” perturbations of a given self-adjoint operator but rather a “structural” perturbation of the dense subspace on which the perturbed operator is defined. More specifically, we focus on the second order Taylor expansion for resolvents, spectral projections, and eigenvalue curves of C^2 -regular one parameter families of self-adjoint extensions of a given symmetric operator in abstract Hilbert spaces. We use the abstract theory of boundary triples and Krein–Naimark-type resolvent formulas, a subject having rich and distinguished history [2, 13, 24]. Among applications is an explicit formula for the Hessian and the first derivative of an eigenvalue curve of a one dimensional Schrödinger operator with matrix-valued potential subject to varying Robin boundary conditions on a bounded interval.

To give this work yet another perspective, let us mention some closely related papers. In [9, 10], G. Cox, C. Jones, J. Marzuola utilized explicit formulas for the first derivative of eigenvalue curves of parameter dependent Robin-type elliptic operators to evaluate the spectral shift between the Dirichlet and Neumann realizations in terms of the Morse index of the Dirichlet-to-Neumann map, which yields a plethora of interesting insights in spectral geometry, e.g., a novel formula for the number of nodal domains of Dirichlet eigenfunctions, and a geometric proof Friedlander’s inequality, [12]. Two more interesting connections are to a recent paper by G. Berkolaiko, P. Kuchment [5], where the Hessian is computed for simple eigenvalues in the case of “additive” perturbations, and to the work of G. Berkolaiko, Y. Canzani, G. Cox, J. L. Marzuola [3], where the Hessian is computed for dispersion relations (i.e. eigenvalue surfaces parameterized by quasimomenta which arise in the Floquet-Bloch decomposition) of some periodic discrete graphs. It is worth noting that in some of these works, e.g. in [3] the parameter is higher-dimensional while we focus on one-parameter families. That said, we believe that the methods developed in the present paper could be applicable in multi-parameter setting.

This work should be considered as a sequel to [19] where a symplectic form of the Krein resolvent formula was derived in order to establish the first order expansion for resolvents and eigenvalues of self-adjoint extensions. We stress that the abstract setting considered in [19] is significantly more general than the one assumed in the current work but the expansion carried out in [19] is only to the first order. The major difference is in the type of trace maps used in [19] where they are assumed to be merely densely defined and with dense range while in the current work the traces are bounded (in particular, have full domain) and surjective. In short, in [19], $(\mathfrak{H}, \Gamma_0, \Gamma_1)$ is not required to be a classical boundary triplet, [2, 13, 24], but it is in the current manuscript.

Let $\mathcal{H}, \mathfrak{H}$ be complex Hilbert spaces. Let A be a densely defined, closed, symmetric operator acting in \mathcal{H} and having equal (possibly infinite) deficiency indices. Let $(\Gamma_0, \Gamma_1, \mathfrak{H})$ be a boundary triplet associated with A , see Hypothesis 2.1. It is well known that self-adjoint extensions of A are parametrized by Lagrangian planes (also known as self-adjoint linear relations) in $\mathfrak{H} \times \mathfrak{H}$, cf., e.g., [2, 13, 24], by means of the identity $[\Gamma_0, \Gamma_1]^\top(\text{dom}(\mathcal{A})) = \mathcal{F}$, where $A \subset \mathcal{A} = \mathcal{A}^* \subset A^*$ and $\mathcal{F} \subset \mathfrak{H} \times \mathfrak{H}$ is a Lagrangian plane. Firstly, we show that for a C^2 path of Lagrangian planes $t \mapsto \mathcal{F}_t$ and a family of self-adjoint extensions of A with $[\Gamma_0, \Gamma_1]^\top(\text{dom}(\mathcal{A}_t)) = \mathcal{F}_t$, the path $t \mapsto (\mathcal{A}_t - \mathbf{i})^{-1}$ is also C^2 -regular in uniform operator topology. Our argument is based on a Krein-type formula from [19], cf. (2.4), expressing resolvent difference in terms of the difference of orthogonal projections onto the respective planes. Secondly, in our main Theorems 2.5, 2.9, 2.10, we derive explicit expressions for the coefficients of the second order Taylor expansions for the resolvents of t -dependent

operators \mathcal{A}_t , the operators per se (more precisely, their conjugated restrictions onto spectral subspaces), and their eigenvalue curves. Next, in Theorem 2.11 we specialized to the case when the Lagrangian planes \mathcal{F}_t are given as kernels of the 1×2 block operator matrices $Z_t = [X_t, Y_t]$ in $\mathfrak{H} \times \mathfrak{H}$, and derive explicit formulas for the first and second derivatives of the eigenvalues of \mathcal{A}_t in terms of Z_t and its derivatives. In particular, Corollary 2.12 gives an expansion of the eigenvalue curves for the abstract Robin-type self-adjoint extensions of A up to the second order and demonstrates that the Hessian in this case features a remarkable similarity with that derived by G. Berkolaiko and P. Kuchement in [5, Eq. 40] for additive perturbations of a fixed self-adjoint operator with a simple eigenvalue.

2. RESOLVENT EXPANSIONS

Throughout, A denotes a densely defined, symmetric operator acting in a complex Hilbert space \mathcal{H} and having equal deficiency indices, i.e., $\dim \ker(A^* - \mathbf{i}) = \dim \ker(A^* + \mathbf{i}) \leq \infty$. We equip the vector space $\mathcal{H}_+ = \text{dom}(A^*)$ with the graph scalar product

$$\langle u, v \rangle_{\mathcal{H}_+} := \langle u, v \rangle_{\mathcal{H}} + \langle A^*u, A^*u \rangle_{\mathcal{H}}, \quad u, v \in \text{dom}(A^*).$$

Let $\mathcal{H}_- := (\mathcal{H}_+)^*$ be the dual to \mathcal{H}_+ space consisting of linear bounded functionals. We notice a chain of natural embeddings, $\mathcal{H}_+ \hookrightarrow \mathcal{H} \hookrightarrow \mathcal{H}_-$, where the first embedding is given by $\mathcal{H}_+ \ni u \mapsto u \in \mathcal{H}$, and the second embedding is given by $\mathcal{H} \ni v \mapsto \langle \cdot, v \rangle_{\mathcal{H}}$. Let $\Phi^{-1} : \mathcal{H}_+ \rightarrow \mathcal{H}_-$ be the conjugate linear Riesz isomorphism such that

$$\mathcal{H}_+ \langle u, \Phi^{-1}w \rangle_{\mathcal{H}_-} = \langle u, w \rangle_{\mathcal{H}_+} = \langle u, w \rangle_{\mathcal{H}} + \langle A^*u, A^*w \rangle_{\mathcal{H}}, \quad u, w \in \mathcal{H}_+.$$

Next, we fix the standard setting of boundary triplets, see [2, 24] for more details, the history of the subject and a vast bibliography. Below, we write $[a, b]^\top$ for a 2×1 column vector with entries a, b .

Hypothesis 2.1. *Assume that $(\Gamma_0, \Gamma_1, \mathfrak{H})$ is a boundary triplet. That is, we assume that we are given an auxiliary (boundary) Hilbert space \mathfrak{H} and a trace operator*

$$\mathsf{T} := [\Gamma_0, \Gamma_1]^\top \in \mathcal{B}(\mathcal{H}_+, \mathfrak{H} \times \mathfrak{H}),$$

which is bounded, surjective and satisfies the following abstract Green identity,

$$\langle A^*u, v \rangle_{\mathcal{H}} - \langle u, A^*v \rangle_{\mathcal{H}} = \langle \Gamma_1 u, \Gamma_0 v \rangle_{\mathfrak{H}} - \langle \Gamma_0 u, \Gamma_1 v \rangle_{\mathfrak{H}} \text{ for all } u, v \in \mathcal{H}_+. \quad (2.1)$$

Let us introduced a bounded symplectic form ω defined by the formulas

$$\begin{aligned} \omega((f_1, f_2)^\top, (g_1, g_2)^\top) &:= \langle f_2, g_1 \rangle_{\mathfrak{H}} - \langle f_1, g_2 \rangle_{\mathfrak{H}} \\ &= \langle J(f_1, f_2)^\top, (g_1, g_2)^\top \rangle_{\mathfrak{H} \times \mathfrak{H}}, \quad J := \begin{bmatrix} 0 & I_{\mathfrak{H}} \\ -I_{\mathfrak{H}} & 0 \end{bmatrix}, \quad f_k, g_k \in \mathfrak{H}, k = 1, 2, \end{aligned} \quad (2.2)$$

so that the right-hand side of (2.1) becomes $\omega(\mathsf{T}u, \mathsf{T}v)$. We denote the annihilator of a subspace $\mathcal{F} \subset \mathfrak{H} \times \mathfrak{H}$ by

$$\mathcal{F}^\circ := \{(f_1, f_2)^\top \in \mathfrak{H} \times \mathfrak{H} : \omega((f_1, f_2)^\top, (g_1, g_2)^\top) = 0 \text{ for all } (g_1, g_2)^\top \in \mathcal{F}\},$$

and recall that the subspace \mathcal{F} is called *Lagrangian* if $\mathcal{F} = \mathcal{F}^\circ$. Further, let $\Lambda(\mathfrak{H} \times \mathfrak{H})$ denote the metric space of Lagrangian subspaces of $\mathfrak{H} \times \mathfrak{H}$ equipped with the metric

$$d(\mathcal{F}_1, \mathcal{F}_2) := \|Q_1 - Q_2\|_{\mathcal{B}(\mathfrak{H} \times \mathfrak{H})}, \quad \mathcal{F}_1, \mathcal{F}_2 \in \Lambda(\mathfrak{H} \times \mathfrak{H}), \quad (2.3)$$

where Q_j is the orthogonal projection onto \mathcal{F}_j acting in $\mathfrak{H} \times \mathfrak{H}$, $j = 1, 2$. Let us also recall the classical, and central for this paper, fact that the Lagrangian planes in $\Lambda(\mathfrak{H} \times \mathfrak{H})$ are

in one-to-one correspondence with self-adjoint extensions of A , see [24]. That is, for every self-adjoint extension \mathcal{A} of A the subspace $\mathsf{T}(\text{dom}(\mathcal{A})) \subset \mathfrak{H} \times \mathfrak{H}$ is Lagrangian with respect to ω , and, conversely, every Lagrangian plane $\mathcal{F} \in \Lambda(\mathfrak{H} \times \mathfrak{H})$ gives rise to a self-adjoint extension of A with domain $\mathsf{T}^{-1}(\mathcal{F}) \subset \text{dom}(A^*)$, see [19] and the references therein for more details.

In this paper we are interested in explicit second order Taylor expansions for resolvents and eigenvalues of a family of self-adjoint extensions \mathcal{A}_t of A defined by a given family of Lagrangian planes $t \mapsto \mathcal{F}_t$ for t near some t_0 . We note that the regularity of a path of Lagrangian planes \mathcal{F}_t is determined by the regularity of the path $t \mapsto Q_t$ of orthogonal projections Q_t acting in $\mathfrak{H} \times \mathfrak{H}$ with $\text{ran}(Q_t) = \mathcal{F}_t$, in other words, the topology on $\Lambda(\mathfrak{H} \times \mathfrak{H})$ is determined by the metric in (2.3). Throughout, we use “dot” to denote t -derivative so that, in particular, $\dot{Q}_{t_0} = \frac{d}{dt}Q_t|_{t=t_0}$. Let us now record our assumptions.

Hypothesis 2.2. *Let $Q : [0, 1] \rightarrow \mathcal{B}(\mathfrak{H} \times \mathfrak{H})$, $t \mapsto Q_t$ be a one-parameter family of orthogonal projections and assume that $\text{ran}(Q_t) \in \Lambda(\mathfrak{H} \times \mathfrak{H})$ is a Lagrangian plane. Fix $t_0 \in [0, 1]$ and assume that*

$$Q_t \underset{t \rightarrow t_0}{=} Q_{t_0} + \dot{Q}_{t_0}(t - t_0) + \frac{1}{2}\ddot{Q}_{t_0}(t - t_0)^2 + o(t - t_0)^2 \text{ in } \mathcal{B}(\mathfrak{H} \times \mathfrak{H}).$$

Let \mathcal{A}_t be a family of self-adjoint extensions of A satisfying

$$\mathsf{T}(\text{dom}(\mathcal{A}_t)) = \text{ran}(Q_t).$$

Finally, denote $R_t(\zeta) := (\mathcal{A}_t - \zeta)^{-1} \in \mathcal{B}(\mathcal{H})$ for $\zeta \notin \text{Spec}(\mathcal{A}_t)$.

We recall from [19, Theorem 3.11] the following Krein-type resolvent formula.

Theorem 2.3. *Assume Hypothesis 2.2 and let $t, t_0 \in [0, 1]$, $\zeta \notin \text{Spec}(\mathcal{A}_t) \cup \text{Spec}(\mathcal{A}_{t_0})$. Then one has*

$$R_t(\zeta) - R_{t_0}(\zeta) = (\mathsf{TR}_t(\zeta))^*(Q_t - Q_{t_0})JQ_{t_0}\mathsf{TR}_{t_0}(\zeta). \quad (2.4)$$

Remark 2.4. 1) Due to Lagrangian properties of Q_t we have $Q_{t_0}JQ_{t_0} = 0$; in addition, since $\mathsf{T}(\text{ran}(R_{t_0}(\zeta))) = \mathsf{T}(\text{dom}(\mathcal{A}_{t_0})) = \text{ran}(Q_{t_0})$, we have $Q_{t_0}\mathsf{TR}_{t_0}(\zeta) = \mathsf{TR}_{t_0}(\zeta)$ and formula (2.4) can be written as follows,

$$R_t(\zeta) - R_{t_0}(\zeta) = (\mathsf{TR}_t(\zeta))^*Q_tJ\mathsf{TR}_{t_0}(\zeta). \quad (2.5)$$

2) The adjoint operator $(\mathsf{TR}_t(\zeta))^*$ in the right-hand side of (2.4) is a rather subtle object. It should be viewed as an operator in $\mathcal{B}(\mathfrak{H} \times \mathfrak{H}, \mathcal{H})$. Indeed, one has $\mathsf{T} \in \mathcal{B}(\mathcal{H}_+, \mathfrak{H} \times \mathfrak{H})$ and $\text{ran}(R_t(\zeta)) \subset \mathcal{H}_+$, so that the product $\mathsf{T}(R_t(\zeta))$ is well defined as a mapping from \mathcal{H} to $\mathfrak{H} \times \mathfrak{H}$. Moreover, we claim that $R_t(\zeta) \in \mathcal{B}(\mathcal{H}, \mathcal{H}_+)$, hence, in particular $\mathsf{T}(R_t(\zeta)) \in \mathcal{B}(\mathcal{H}, \mathfrak{H} \times \mathfrak{H})$ so its adjoint is indeed in $\mathcal{B}(\mathfrak{H} \times \mathfrak{H}, \mathcal{H})$ as claimed above. To check the inclusion $R_t(\zeta) \in \mathcal{B}(\mathcal{H}, \mathcal{H}_+)$ we notice that for some $c > 0$ and every $u \in \mathcal{H}$ one has

$$\begin{aligned} \|R_{t_0}(\zeta)u\|_{\mathcal{H}_+}^2 &= \|R_{t_0}(\zeta)u\|_{\mathcal{H}}^2 + \|A^*R_{t_0}(\zeta)u\|_{\mathcal{H}}^2 \\ &\leq \|R_{t_0}(\zeta)u\|_{\mathcal{H}}^2 + 2\|(\mathcal{A}_{t_0} - \zeta)R_{t_0}(\zeta)u\|_{\mathcal{H}}^2 + 2\|\zeta R_{t_0}(\zeta)u\|_{\mathcal{H}}^2 \\ &\leq \|R_{t_0}(\zeta)u\|_{\mathcal{H}}^2 + 2\|u\|_{\mathcal{H}}^2 + 2\|\zeta R_{t_0}(\zeta)u\|_{\mathcal{H}}^2 \leq c\|u\|_{\mathcal{H}}^2. \end{aligned}$$

3) It is possible to represent $(\mathsf{TR}_t(\zeta))^*$ as the product of two adjoints, $(R_t(\zeta))^*$ and T^* . In fact, the adjoint of the resolvent $(R_t(\zeta))^* \in \mathcal{B}(\mathcal{H}_-, \mathcal{H})$ is a natural extension of $R_t(\bar{\zeta})$, originally defined as a mapping from \mathcal{H} to itself, to a mapping from \mathcal{H}_- to \mathcal{H} , see [19, Proposition 2.4]. In this work, however, we chose not to unfold the product $(\mathsf{TR}_t(\zeta))^*$ and shall view $R_t(\zeta)$ as an element of $\mathcal{B}(\mathcal{H}, \mathcal{H}_+)$ whenever it precedes T .

4) From the outset we only have $\text{ran}((\text{TR}_t(\zeta))^*) \subset \mathcal{H}$. However, (2.5) suggests that $\text{ran}((\text{TR}_t(\zeta))^*) \subset \mathcal{H}_+$ as both resolvents in the left-hand side of (2.5) have ranges in \mathcal{H}_+ . It turns out, cf. Theorem 2.5 (3), that Krein's-type formula (2.5) produces a similar effect for the derivative $\dot{R}_{t_0}(\zeta)$ that satisfies $\text{ran}(\dot{R}_{t_0}(\zeta)) \subset \mathcal{H}_+$ although *a priori* one only has $\text{ran}(\dot{R}_{t_0}(\zeta)) \subset \mathcal{H}$.

Our first principal result gives Taylor expansions for the resolvent of $R_t(\zeta)$

Theorem 2.5. *Assume Hypothesis 2.2. Suppose that $\zeta_0 \notin \text{Spec}(\mathcal{A}_{t_0})$ for some $t_0 \in [0, 1]$ and $\zeta_0 \in \mathbb{C}$, and define*

$$\mathcal{U}_\varepsilon = \{(t, \zeta) \in [0, 1] \times \mathbb{C} : |t - t_0| \leq \varepsilon, |\zeta - \zeta_0| \leq \varepsilon\} \text{ for } \varepsilon > 0.$$

- (1) *There exists an $\varepsilon > 0$ such that if $(t, \zeta) \in \mathcal{U}_\varepsilon$ then $\zeta \notin \text{Spec}(\mathcal{A}_t)$.*
(2) *For $\varepsilon > 0$, $(t, \zeta) \in \mathcal{U}_\varepsilon$ as in part (1) above one has*

$$(\text{TR}_t(\zeta))^*(Q_t - Q_{t_0})JQ_{t_0}\text{TR}_{t_0}(\zeta) \in \mathcal{B}(\mathcal{H}, \mathcal{H}_+). \quad (2.6)$$

- (3) *For $\varepsilon > 0$, $\zeta \in \mathbb{C}$ as in part (1) above we define*

$$\dot{R}_{t_0}(\zeta) := (\text{TR}_{t_0}(\zeta))^*\dot{Q}_{t_0}J\text{TR}_{t_0}(\zeta). \quad (2.7)$$

Then one has

$$\dot{R}_{t_0}(\zeta) \in \mathcal{B}(\mathcal{H}, \mathcal{H}_+). \quad (2.8)$$

Moreover, the following asymptotic expansion holds uniformly for $|\zeta - \zeta_0| < \varepsilon$,

$$R_t(\zeta) = R_{t_0}(\zeta) + \dot{R}_{t_0}(\zeta)(t - t_0) + o(t - t_0) \text{ in } \mathcal{B}(\mathcal{H}, \mathcal{H}_+). \quad (2.9)$$

- (4) *For $\varepsilon > 0$ as in part (1) the following asymptotic expansion holds uniformly for $|\zeta - \zeta_0| < \varepsilon$,*

$$R_t(\zeta) \underset{t \rightarrow t_0}{=} R_{t_0}(\zeta) + \dot{R}_{t_0}(\zeta)(t - t_0) + \frac{1}{2}\ddot{R}_{t_0}(\zeta)(t - t_0)^2 + o(t - t_0)^2 \text{ in } \mathcal{B}(\mathcal{H}), \quad (2.10)$$

where $\dot{R}_{t_0}(\zeta)$ is given by (2.7) and

$$\ddot{R}_{t_0}(\zeta) := (\text{TR}_{t_0})^*\ddot{Q}_{t_0}J\text{TR}_{t_0} + 2(\text{TR}_{t_0})^*\dot{Q}_{t_0}J\text{TR}_{t_0}(\zeta). \quad (2.11)$$

Proof. Proof of (1). First, we show that for $\zeta = \mathbf{i}$ we have

$$\|R_t(\mathbf{i})\|_{\mathcal{B}(\mathcal{H}, \mathcal{H}_+)} \underset{t \rightarrow t_0}{=} \mathcal{O}(1). \quad (2.12)$$

Of course, such asymptotic formula holds in $\mathcal{B}(\mathcal{H})$ by self-adjointness of \mathcal{A}_t . A stronger version which we are after holds due to the fact that $R_t(\mathbf{i})$ is the resolvent of a self-adjoint restriction of A^* ; this observation is crucial throughout the proof.

Since $\|R_t(\mathbf{i})\|_{\mathcal{B}(\mathcal{H})} \leq 1$ there exists $c > 0$ such that for all $u \in \mathcal{H}_+$ and t near t_0 one has

$$\begin{aligned} \|R_t(\mathbf{i})u\|_{\mathcal{H}_+}^2 &= \|R_t(\mathbf{i})u\|_{\mathcal{H}}^2 + \|A^*R_t(\mathbf{i})u\|_{\mathcal{H}}^2 \\ &= \|R_t(\mathbf{i})u\|_{\mathcal{H}}^2 + 2\|(A_t - \mathbf{i})R_t(\mathbf{i})u\|_{\mathcal{H}}^2 + 2\|\mathbf{i}R_t(\mathbf{i})u\|_{\mathcal{H}}^2 \leq c\|u\|_{\mathcal{H}}^2 \end{aligned}$$

which confirms (2.12) (we used $(A_t - \mathbf{i})R_t(\mathbf{i}) = I$). Next, by Krein's formula (2.5) with $\zeta = \mathbf{i}$, we have

$$R_t(\mathbf{i}) = R_{t_0}(\mathbf{i}) + (\text{TR}_t(\mathbf{i}))^*(Q_t - Q_{t_0})J\text{TR}_{t_0}(\mathbf{i}).$$

Combining this with $\|TR_t(\mathbf{i})\|_{\mathcal{B}(\mathcal{H}, \mathfrak{H} \times \mathfrak{H})} \underset{t \rightarrow t_0}{=} \mathcal{O}(1)$ we get

$$\|R_t(\mathbf{i}) - R_{t_0}(\mathbf{i})\|_{\mathcal{B}(\mathcal{H})} \underset{t \rightarrow t_0}{=} \mathcal{O}(t - t_0), \quad (2.13)$$

so that $\mathcal{A}_t \rightarrow \mathcal{A}_{t_0}$, $t \rightarrow t_0$ in the norm resolvent sense. Then [21, Theorem VIII.23(a)] yields the assertion.

Proof of (2). By Krein's formula the operator in (2.6) is equal to the difference of the resolvents of two self-adjoint restrictions of A^* , each of which is an element of $\mathcal{B}(\mathcal{H}, \mathcal{H}_+)$ as discussed in Remark 2.4 part 2).

Proof of (3). We begin by obtaining an improved version of (2.13) given by

$$\|R_t(\mathbf{i}) - R_{t_0}(\mathbf{i})\|_{\mathcal{B}(\mathcal{H}, \mathcal{H}_+)} \underset{t \rightarrow t_0}{=} \mathcal{O}(t - t_0). \quad (2.14)$$

To that end we write

$$\begin{aligned} \|R_t(\mathbf{i})u - R_{t_0}(\mathbf{i})u\|_{\mathcal{H}_+}^2 &= \|R_t(\mathbf{i})u - R_{t_0}(\mathbf{i})u\|_{\mathcal{H}}^2 + \|A^*R_t(\mathbf{i})u - A^*R_{t_0}(\mathbf{i})u\|_{\mathcal{H}}^2 \\ &\leq \|R_t(\mathbf{i})u - R_{t_0}(\mathbf{i})u\|_{\mathcal{H}}^2 + 2\|(\mathcal{A}_t - \mathbf{i})R_t(\mathbf{i})u - (\mathcal{A}_{t_0} - \mathbf{i})R_{t_0}(\mathbf{i})u\|_{\mathcal{H}}^2 \\ &\quad + 2\|-\mathbf{i}R_t(\mathbf{i})u + \mathbf{i}R_{t_0}(\mathbf{i})u\|_{\mathcal{H}}^2 \\ &\leq 3\|R_t(\mathbf{i})u - R_{t_0}(\mathbf{i})u\|_{\mathcal{H}}^2 \underset{t \rightarrow t_0}{=} \mathcal{O}(t - t_0), \text{ uniformly for } \|u\|_{\mathcal{H}} \leq 1. \end{aligned}$$

We will need a slightly more general version of (2.14) in which \mathbf{i} is replaced by $\zeta \in \mathbb{C}$ near ζ_0 . Let us fix $\varepsilon_0 > 0$ such that $\mathbb{B}_{\varepsilon_0}(\zeta_0) \subset \mathbb{C} \setminus \text{Spec}(\mathcal{A}_{t_0})$ for the ε_0 -disc centered at ζ_0 . Then by (2.13) and [21, Theorem VIII.23] we have $\mathbb{B}_{\varepsilon_0}(\zeta_0) \cap \text{Spec}(\mathcal{A}_t) = \emptyset$ for t sufficiently close to t_0 . Hence,

$$\sup\{\|R_t(\zeta)\|_{\mathcal{B}(\mathcal{H})} : (t, \zeta) \in \mathcal{U}_\varepsilon\} < \infty \quad (2.15)$$

for a sufficiently small $\varepsilon \in (0, \varepsilon_0)$. Then combining the resolvent identity

$$R_t(\zeta) = R_t(\mathbf{i}) - (\mathbf{i} - \zeta)R_t(\mathbf{i})R_t(\zeta),$$

and (2.15) we infer that the asymptotic formula

$$\|R_t(\zeta) - R_{t_0}(\zeta)\|_{\mathcal{B}(\mathcal{H}, \mathcal{H}_+)} \underset{t \rightarrow t_0}{=} \mathcal{O}(t - t_0),$$

holds uniformly for $|\zeta - \zeta_0| < \varepsilon$.

We are now ready to turn to the proof of (2.8), (2.9). To that end we will first establish the same expansion but in the space $\mathcal{B}(\mathcal{H})$ possessing weaker topology. Employing Krein's formula (2.4) ones again we obtain

$$\begin{aligned} R_t(\zeta) &\underset{t \rightarrow t_0}{=} R_{t_0}(\zeta) + (TR_{t_0}(\zeta) + \mathcal{O}_{\mathcal{B}(\mathcal{H}, \mathfrak{H} \times \mathfrak{H})}(t - t_0))^* (\dot{Q}_{t_0}(t - t_0) + o(t - t_0)) JTR_{t_0}(\zeta) \\ &= R_{t_0}(\zeta) + \dot{R}_{t_0}(\zeta)(t - t_0) + o(t - t_0) \text{ in } \mathcal{B}(\mathcal{H}). \end{aligned} \quad (2.16)$$

Our next objective is to derive this asymptotic expansion in $\mathcal{B}(\mathcal{H}, \mathcal{H}_+)$ and to prove (2.8). To that end we pick any sequence $t_n \rightarrow t_0$, $n \rightarrow \infty$ and introduce a sequence of operators

$$\mathcal{D}_n(\zeta) := \frac{R_{t_n}(\zeta) - R_{t_0}(\zeta)}{t_n - t_0}, n \in \mathbb{N}.$$

First we notice that $\text{ran}(\mathcal{D}_n) \subset \text{dom}(A^*) = \mathcal{H}_+$ and, in fact, by Remark 2.4 (2) one has $\mathcal{D}_n \in \mathcal{B}(\mathcal{H}, \mathcal{H}_+)$. We claim that the sequence $\{\mathcal{D}_n(\zeta)\}$ is Cauchy in $\mathcal{B}(\mathcal{H}, \mathcal{H}_+)$. Let us observe that

$$A^*(R_{t_n}(\zeta) - R_{t_m}(\zeta)) = \zeta(R_{t_m}(\zeta) - R_{t_n}(\zeta)),$$

that is $A^*(\mathcal{D}_n(\zeta) - \mathcal{D}_m(\zeta)) = \zeta(\mathcal{D}_m(\zeta) - \mathcal{D}_n(\zeta))$, so for arbitrary $u \in \mathcal{H}$ we have

$$\begin{aligned} \|(\mathcal{D}_n(\zeta) - \mathcal{D}_m(\zeta))u\|_{\mathcal{H}_+}^2 &= \|(\mathcal{D}_n(\zeta) - \mathcal{D}_m(\zeta))u\|_{\mathcal{H}}^2 + \|A^*(\mathcal{D}_n(\zeta) - \mathcal{D}_m(\zeta))u\|_{\mathcal{H}}^2 \\ &= (1 + |\zeta|^2)\|(\mathcal{D}_n(\zeta) - \mathcal{D}_m(\zeta))u\|_{\mathcal{H}}^2, \end{aligned}$$

thus, one has

$$\|\mathcal{D}_n(\zeta) - \mathcal{D}_m(\zeta)\|_{\mathcal{B}(\mathcal{H}, \mathcal{H}_+)}^2 \leq (1 + |\zeta|^2)\|\mathcal{D}_n(\zeta) - \mathcal{D}_m(\zeta)\|_{\mathcal{B}(\mathcal{H})}^2 \xrightarrow{m, n \rightarrow \infty} 0,$$

where we used that $\{\mathcal{D}_n\}_{n \in \mathbb{N}}$ is Cauchy in $\mathcal{B}(\mathcal{H})$ by (2.16), (2.13). Thus, the sequence \mathcal{D}_n converges in $\mathcal{B}(\mathcal{H}, \mathcal{H}_+)$ to an operator $\mathcal{D}_\infty \in \mathcal{B}(\mathcal{H}, \mathcal{H}_+)$. By definition of $\|\cdot\|_{\mathcal{H}_+}$, we automatically also infer

$$\|\mathcal{D}_n(\zeta) - \mathcal{D}_\infty(\zeta)\|_{\mathcal{B}(\mathcal{H})} \rightarrow 0, n \rightarrow \infty.$$

At the same time, (2.16) show that

$$\|\mathcal{D}_n(\zeta) - \dot{R}_{t_0}(\zeta)\|_{\mathcal{B}(\mathcal{H})} \rightarrow 0, n \rightarrow \infty.$$

Hence $\dot{R}_{t_0}(\zeta) = \mathcal{D}_\infty(\zeta) \in \mathcal{B}(\mathcal{H}, \mathcal{H}_+)$ as asserted in (2.8). Also, $\mathcal{D}_\infty(\zeta)$ is independent of the choice of $t_n \rightarrow t_0$. Thus

$$R_t(\zeta) - R_{t_0}(\zeta) = \mathcal{D}_\infty(\zeta)(t - t_0) + o(t - t_0), \quad t \rightarrow t_0,$$

and (2.9) holds.

Proof of (4). The asymptotic formula (2.10) is obtained by substituting resolvent expansion (2.9) into Krein's formula (2.4) and grouping powers of $(t - t_0)$. This formal expansion is indeed justified since the operator $\text{TR}_{t_0}(\zeta)$ is well defined and, by (2.9), we have

$$\text{TR}_t(\zeta) = \text{TR}_{t_0}(\zeta) + \text{T}\dot{R}_{t_0}(\zeta)(t - t_0) + o(t - t_0) \text{ in } \mathcal{B}(\mathcal{H}, \mathfrak{H} \times \mathfrak{H}).$$

□

In what follows we will describe how a multiple isolated eigenvalue λ of \mathcal{A}_{t_0} bifurcates for t near t_0 .

Hypothesis 2.6. Fix $t_0 \in [0, 1]$, suppose that $\lambda = \lambda(t_0)$ is an isolated eigenvalue of \mathcal{A}_{t_0} with finite multiplicity $m \in \mathbb{N}$. Let

$$\gamma := \{z \in \mathbb{C} : 2|z - \lambda| = \text{dist}(\lambda, \text{Spec}(\mathcal{A}_{t_0}) \setminus \{\lambda\})\}.$$

We continue to assume Hypothesis 2.2. Then, by Theorem 2.5, there exists $\varepsilon > 0$ such that γ encloses the λ -group of m eigenvalues (not necessarily distinct) of the operator \mathcal{A}_t whenever $|t - t_0| < \varepsilon$ and $\varepsilon > 0$ is sufficiently small. For such t we let P_t denote the Riesz projection for \mathcal{A}_t corresponding to the λ -group,

$$P_t := \frac{-1}{2\pi\mathbf{i}} \int_{\gamma} R_t(\zeta) d\zeta, \quad \text{where } R_t(\zeta) = (\mathcal{A}_t - \zeta)^{-1}. \quad (2.17)$$

Let us recall from [18, Section III.6.5] the following Laurent expansion for the resolvent,

$$R_{t_0}(\zeta) = (\lambda - \zeta)^{-1} P_{t_0} + \sum_{n=0}^{\infty} (\zeta - \lambda)^n S_{t_0}^{n+1}, \quad (2.18)$$

where $S_{t_0} = \lim_{\zeta \rightarrow \lambda} R_{t_0}(\zeta)(I_{\mathcal{H}} - P_{t_0})$ is the reduced resolvent of the operator \mathcal{A}_{t_0} at the eigenvalue λ given by the formula

$$S_{t_0} := \frac{-1}{2\pi\mathbf{i}} \int_{\gamma} (\lambda - \zeta)^{-1} R_{t_0}(\zeta) d\zeta. \quad (2.19)$$

Remark 2.7. We note that by Theorem 2.5, part (3) one has

$$(\mathrm{TR}_{t_0}(\zeta))^* \dot{Q}_{t_0} J \mathrm{TR}_{t_0}(\zeta) \in \mathcal{B}(\mathcal{H}, \mathcal{H}_+).$$

Therefore, in particular

$$\mathrm{T}(\mathrm{TR}_{t_0}(\zeta))^* \dot{Q}_{t_0} J \mathrm{TR}_{t_0}(\zeta) \in \mathcal{B}(\mathcal{H}, \mathfrak{H} \times \mathfrak{H}).$$

Furthermore, since the integrals in (2.17) and (2.19) converge with respect to the norm of the space $\mathcal{B}(\mathcal{H}, \mathcal{H}_+)$, the two inclusions above also hold with the two copies of $R_{t_0}(\zeta)$ replaced by $S_{t_0}^2, S_{t_0}, P_{t_0}$ in any combination. In particular, the operators in (2.21), (2.22), (2.25), (2.26), (2.48), and in Remark 2.14 used below are all well defined and bounded in the respective spaces.

The following result gives the second order Taylor expansion for the Riesz projection just introduced.

Proposition 2.8. *Under Hypotheses 2.2 and 2.6, for the Riesz projection introduced in (2.17) one has*

$$P_t \underset{t \rightarrow t_0}{=} P_{t_0} + \dot{P}_{t_0}(t - t_0) + \frac{1}{2} \ddot{P}_{t_0}(t - t_0)^2 + o(t - t_0)^2, \quad \text{in } \mathcal{B}(\mathcal{H}), \quad (2.20)$$

with

$$\dot{P}_{t_0} := (\mathrm{TS}_{t_0})^* \dot{Q}_{t_0} J \mathrm{TP}_{t_0} + (\mathrm{TP}_{t_0})^* \dot{Q}_{t_0} J \mathrm{TS}_{t_0}, \quad (2.21)$$

and

$$\begin{aligned} \ddot{P}_{t_0} := & (\mathrm{TS}_{t_0})^* \ddot{Q}_{t_0} J \mathrm{TP}_{t_0} + (\mathrm{TP}_{t_0})^* \ddot{Q}_{t_0} J \mathrm{TS}_{t_0} \\ & + 2(\mathrm{T}(\mathrm{TS}_{t_0})^* \dot{Q}_{t_0} J \mathrm{TS}_{t_0})^* \dot{Q}_{t_0} J \mathrm{TP}_{t_0} + 2(\mathrm{T}(\mathrm{TS}_{t_0})^* \dot{Q}_{t_0} J \mathrm{TP}_{t_0})^* \dot{Q}_{t_0} J \mathrm{TS}_{t_0} \\ & + 2(\mathrm{T}(\mathrm{TP}_{t_0})^* \dot{Q}_{t_0} J \mathrm{TS}_{t_0})^* \dot{Q}_{t_0} J \mathrm{TS}_{t_0} - 2(\mathrm{T}(\mathrm{TS}_{t_0}^2)^* \dot{Q}_{t_0} J \mathrm{TP}_{t_0})^* \dot{Q}_{t_0} J \mathrm{TP}_{t_0} \\ & - 2(\mathrm{T}(\mathrm{TP}_{t_0})^* \dot{Q}_{t_0} J \mathrm{TS}_{t_0}^2)^* \dot{Q}_{t_0} J \mathrm{TP}_{t_0} - 2(\mathrm{T}(\mathrm{TP}_{t_0})^* \dot{Q}_{t_0} J \mathrm{TP}_{t_0})^* \dot{Q}_{t_0} J \mathrm{TS}_{t_0}^2. \end{aligned} \quad (2.22)$$

Proof. We plug (2.18) in (2.7), (2.11) and observe that the coefficients of the terms containing $(\lambda - \zeta)^{-1}$ in the Laurent expansions of $\dot{R}_{t_0}(\zeta), \ddot{R}_{t_0}(\zeta)$ are given by the right-hand sides of (2.21), (2.22) respectively. Upon using the right hand sides of (2.21), (2.22), in the Laurent expansions of $\dot{R}_{t_0}(\zeta), \ddot{R}_{t_0}(\zeta)$, we multiply both sides of (2.10) by $\frac{-1}{2\pi\mathbf{i}}$, integrate over γ , and employ (2.17) together with Cauchy's integral formula to arrive at (2.20). In other words, $\dot{P}_{t_0}, \ddot{P}_{t_0}$ are residues of the operator-valued functions $\zeta \mapsto -\dot{R}_{t_0}(\zeta), \zeta \mapsto -\ddot{R}_{t_0}(\zeta)$; the residues are given by (2.21), (2.22). \square

Our next objective is to derive an asymptotic expansion for the operator that is similar to $P_t \mathcal{A}_t P_t$ for t near t_0 but acts on a t -independent m -dimensional space $\mathrm{ran}(P_{t_0})$. To that end, we introduce the operator U_t mapping $\mathrm{ran}(P_{t_0})$ onto $\mathrm{ran}(P_t)$ unitarily, for t near t_0 , and defined by the formulas

$$\begin{aligned} U_t &:= (I - D_t^2)^{-1/2} ((I - P_t)(I - P_{t_0}) + P_t P_{t_0}), \\ U_t^{-1} &= ((I - P_{t_0})(I - P_t) + P_{t_0} P_t)(I - D_t^2)^{-1/2}, \end{aligned} \quad (2.23)$$

where $D_t := P_t - P_{t_0}$ is invertible (for t near t_0) since $\|D_t\|_{\mathcal{B}(\mathcal{H})} \underset{t \rightarrow t_0}{=} o(1)$ by (2.20). In particular, one has $U_t P_{t_0} = P_t U_t$. Given the auxiliary operators U_t we consider an m -dimensional operator $W_t := P_{t_0} U_t^{-1} \mathcal{A}_t U_t P_{t_0}$ and obtain its second order Taylor expansion next. Here and in what follows we identify W_t and its restriction to $\text{ran}(P_{t_0})$.

Theorem 2.9. *Assume Hypotheses 2.2 and 2.6 and let $W_t := P_{t_0} U_t^{-1} \mathcal{A}_t U_t P_{t_0}$ for t near t_0 . Then the following second order Taylor expansion holds,*

$$W_t \underset{t \rightarrow t_0}{=} W_{t_0} + \dot{W}_{t_0}(t - t_0) + \frac{1}{2} \ddot{W}_{t_0}(t - t_0)^2 + o(t - t_0)^2, \quad (2.24)$$

where $W_{t_0} := \lambda P_{t_0}$ and

$$\dot{W}_{t_0} := (T P_{t_0})^* J \dot{Q}_{t_0} T P_{t_0}, \quad (2.25)$$

$$\ddot{W}_{t_0} := (T P_{t_0})^* J \ddot{Q}_{t_0} T P_{t_0} - 2(\text{T}(T S_{t_0})^* \dot{Q}_{t_0} J T P_{t_0})^* \dot{Q}_{t_0} J T P_{t_0}. \quad (2.26)$$

Proof. Our strategy is to expand the function

$$t \mapsto P_{t_0} U_t^{-1} R_t(\zeta) U_t P_{t_0}, \quad (2.27)$$

uniformly for ζ near λ , to the second order in $(t - t_0)$, then multiply both sides of the obtained expansion by $\frac{-1}{2\pi i} \zeta$, integrate over γ , and use the formula

$$\mathcal{A}_t P_t = \frac{-1}{2\pi i} \int_{\gamma} \zeta R_t(\zeta) d\zeta. \quad (2.28)$$

In order to obtain the expansion for the operator in (2.27), we split it into four terms,

$$\begin{aligned} P_{t_0} U_t^{-1} R_t(\zeta) U_t P_{t_0} &= \\ &= P_{t_0} U_t^{-1} P_{t_0} R_t(\zeta) P_{t_0} U_t P_{t_0} + P_{t_0} U_t^{-1} (I - P_{t_0}) R_t(\zeta) P_{t_0} U_t P_{t_0} \\ &\quad + P_{t_0} U_t^{-1} P_{t_0} R_t(\zeta) (I - P_{t_0}) U_t P_{t_0} + P_{t_0} U_t^{-1} (I - P_{t_0}) R_t(\zeta) (I - P_{t_0}) U_t P_{t_0} \\ &= \tau_1(\zeta) + \tau_2(\zeta) + \tau_3(\zeta) + \tau_4(\zeta), \end{aligned} \quad (2.29)$$

where $\tau_k(\zeta)$ denotes the respective term in (2.29). The remaining part of the argument is divided into three steps. In Steps 1 and 2, we derive auxiliary expansions in $t - t_0$. Step 3 gives asymptotic expansion for integrated versions of $\tau_k(\zeta)$.

Step 1. Let us denote

$$\begin{aligned} \mathfrak{S}_0 &:= (T P_{t_0})^* \dot{Q}_{t_0} J T S_{t_0}, \\ \mathfrak{S}_1 &:= (\text{T}(T S_{t_0})^* \dot{Q}_{t_0} J T P_{t_0})^* \dot{Q}_{t_0} J T P_{t_0}, \quad \mathfrak{S}_2 := (\text{T}(T S_{t_0}^2)^* \dot{Q}_{t_0} J T P_{t_0})^* \dot{Q}_{t_0} J T P_{t_0}. \end{aligned} \quad (2.30)$$

Then one has

$$P_{t_0} P_t P_{t_0} = P_{t_0} - \mathfrak{S}_2 (t - t_0)^2 + o(t - t_0)^2, \quad (2.31)$$

$$P_{t_0} P_t = P_{t_0} + \mathfrak{S}_0 (t - t_0) + o(t - t_0). \quad (2.32)$$

Proof of (2.31), (2.32). To prove (2.31), we multiply both sides of (2.20) by P_{t_0} from the left and right sides and use (2.21), (2.22), and the identities $S_{t_0} P_{t_0} = 0$ and $P_{t_0} (\text{T} S_{t_0})^* = (\text{T} S_{t_0} P_{t_0})^*$. Similarly, one obtains (2.32). \square

Step 2. One has

$$P_{t_0} U_t P_{t_0} = P_{t_0} - \frac{1}{2} \mathfrak{S}_2 (t - t_0)^2 + o(t - t_0)^2, \quad (2.33)$$

$$P_{t_0}U_t^{-1}(I - P_{t_0}) = \mathfrak{S}_0(t - t_0) + o(t - t_0), \quad (2.34)$$

where we used notation (2.30).

Proof of (2.33), (2.34). Let us derive (2.33). First, recalling U_t from (2.23) we get

$$\begin{aligned} P_{t_0}U_tP_{t_0} &= P_{t_0}(I - D_t^2)^{-1/2}P_tP_{t_0} = P_{t_0}\left(I + \frac{1}{2}D_t^2 + o(t - t_0)^2\right)P_tP_{t_0} \\ &= P_{t_0}P_tP_{t_0} + \frac{1}{2}P_{t_0}D_t^2P_tP_{t_0} + o(t - t_0)^2. \end{aligned} \quad (2.35)$$

Next, since $D_t = P_t - P_{t_0} = \dot{P}_{t_0}(t - t_0) + o(t - t_0)$, we have

$$\begin{aligned} P_{t_0}D_t^2P_tP_{t_0} &= P_{t_0}(\dot{P}_{t_0}^2(t - t_0)^2 + o(t - t_0)^2)(P_{t_0} + \dot{P}_{t_0}(t - t_0) + o(t - t_0))P_{t_0} \\ &= P_{t_0}\dot{P}_{t_0}^2P_{t_0}(t - t_0)^2 + o(t - t_0)^2 = \mathfrak{S}_2(t - t_0)^2 + o(t - t_0)^2, \end{aligned} \quad (2.36)$$

where in the last equality we utilized (2.21) and $S_{t_0}P_{t_0} = 0$. Then (2.33) can be inferred from (2.31), (2.35), (2.36).

To prove (2.34), we use (2.23), (2.32) and get

$$\begin{aligned} P_{t_0}U_t^{-1}(I - P_{t_0}) &= P_{t_0}(I - P_{t_0} - P_t + 2P_{t_0}P_t + \mathcal{O}(t - t_0)^2)(I - P_{t_0}) \\ &= P_{t_0}P_t(I - P_{t_0}) + o(t - t_0) = \mathfrak{S}_0(t - t_0) + o(t - t_0), \end{aligned}$$

where we used $\mathfrak{S}_0(1 - P_{t_0}) = \mathfrak{S}_0$. □

Step 3. Let us denote

$$\mathfrak{Q}_1 := (\mathbb{T}P_{t_0})^*\dot{Q}_{t_0}J\mathbb{T}P_{t_0}, \quad \mathfrak{Q}_2 := (\mathbb{T}P_{t_0})^*\ddot{Q}_{t_0}J\mathbb{T}P_{t_0},$$

recall $\tau_k(\zeta)$ introduced in (2.29), and $\mathfrak{S}_0, \mathfrak{S}_1, \mathfrak{S}_2$ from (2.30). Then we have

$$\frac{-1}{2\pi\mathbf{i}} \int_{\gamma} \zeta \tau_1(\zeta) d\zeta = \lambda P_{t_0} - \mathfrak{Q}_1(t - t_0) - \left(\frac{1}{2}\mathfrak{Q}_2 + 2\lambda\mathfrak{S}_2 + \mathfrak{S}_1\right)(t - t_0)^2 + o(t - t_0)^2, \quad (2.37)$$

$$\frac{-1}{2\pi\mathbf{i}} \int_{\gamma} \zeta \tau_k(\zeta) d\zeta = \lambda \mathfrak{S}_2(t - t_0)^2 + o(t - t_0)^2, \quad k = 2, 3, \quad (2.38)$$

$$\frac{-1}{2\pi\mathbf{i}} \int_{\gamma} \zeta \tau_4(\zeta) d\zeta = o(t - t_0)^2. \quad (2.39)$$

Proof of (2.37), (2.38). To prove (2.37) we will combine the second order Taylor expansions (2.33) and (2.10) with respect to parameter t , collect powers of $(\lambda - \zeta)$, and contour integrate with respect to ζ as follows.

Introducing $\tilde{\mathfrak{R}}(\zeta) := (\mathbb{T}(\mathbb{T}R_{t_0}(\zeta)))^*\dot{Q}_{t_0}J\mathbb{T}P_{t_0})^*\dot{Q}_{t_0}J\mathbb{T}P_{t_0}$ and noting that by conjugation $P_{t_0}U_t^{-1}P_{t_0}$ admits the same asymptotic as $P_{t_0}U_tP_{t_0}$, we infer

$$\begin{aligned} \tau_1(\zeta) &= P_{t_0} \left(P_{t_0} - \frac{1}{2}\mathfrak{S}_2(t - t_0)^2 + o(t - t_0)^2 \right) P_{t_0} \times \\ &\times P_{t_0} \left((\lambda - \zeta)^{-1}P_{t_0} + (\lambda - \zeta)^{-2}\mathfrak{Q}_1(t - t_0) + \left(\frac{1}{2}(\lambda - \zeta)^{-2}\mathfrak{Q}_2 + (\lambda - \zeta)^{-2}\tilde{\mathfrak{R}}(\zeta)\right)(t - t_0)^2 \right) P_{t_0} \times \\ &\times \left(P_{t_0} - \frac{1}{2}\mathfrak{S}_2(t - t_0)^2 + o(t - t_0)^2 \right) P_{t_0}, \end{aligned}$$

hence,

$$\begin{aligned} \tau_1(\zeta) &= (\lambda - \zeta)^{-1}P_{t_0} + (\lambda - \zeta)^{-2}\mathfrak{Q}_1(t - t_0) \\ &+ \left(\frac{1}{2}(\lambda - \zeta)^{-2}\mathfrak{Q}_2 - (\lambda - \zeta)^{-1}\mathfrak{S}_2 + (\lambda - \zeta)^{-2}\tilde{\mathfrak{R}}(\zeta) \right) (t - t_0)^2 + o(t - t_0)^2. \end{aligned} \quad (2.40)$$

The Laurent expansion (2.18) yields

$$\frac{-1}{2\pi\mathbf{i}} \int_{\gamma} \zeta(\lambda - \zeta)^{-2}R_{t_0}(\zeta)d\zeta = -S_{t_0} - \lambda S_{t_0}^2,$$

and, consequently,

$$\frac{-1}{2\pi\mathbf{i}} \int_{\gamma} \zeta(\lambda - \zeta)^{-2}\tilde{\mathfrak{R}}(\zeta)d\zeta = -\mathfrak{S}_1 - \lambda\mathfrak{S}_2. \quad (2.41)$$

Multiplying both sides of (2.40) by $\frac{-1}{2\pi\mathbf{i}}\zeta$, integrating over γ , and using (2.41) we obtain (2.37).

Let us derive (2.38) for $k = 2$ (the case $k = 3$ follows by conjugation). Using Taylor expansions (2.10), (2.34) and denoting $\widehat{\mathfrak{R}}(\zeta) := (\mathrm{TR}_{t_0}(\zeta))^*\dot{Q}_{t_0}J\mathrm{TR}_{t_0}(\zeta)$ we arrive at

$$\begin{aligned} \tau_2(\zeta) &= (\mathfrak{S}_0(t - t_0) + o(t - t_0))(I - P_{t_0})((\lambda - \zeta)^{-1}\widehat{\mathfrak{R}}(\zeta)(t - t_0) + o(t - t_0))P_{t_0} \\ &\times \left(P_{t_0} - \frac{1}{2}\mathfrak{S}_2(t - t_0)^2 + o(t - t_0)^2 \right), \end{aligned}$$

and, consequently, infer

$$\tau_2(\zeta) = (\lambda - \zeta)^{-1}\mathfrak{S}_0\widehat{\mathfrak{R}}(\zeta)(t - t_0)^2 + o(t - t_0)^2. \quad (2.42)$$

Let us observe that

$$\frac{-1}{2\pi\mathbf{i}} \int_{\gamma} \zeta(\lambda - \zeta)^{-1}R_{t_0}(\zeta)d\zeta = -P_{t_0} + \lambda S_{t_0}, \quad S_{t_0}P_{t_0} = 0,$$

thus

$$\frac{-1}{2\pi\mathbf{i}} \int_{\gamma} \zeta(\lambda - \zeta)^{-1}\mathfrak{S}_0\widehat{\mathfrak{R}}(\zeta)d\zeta = \lambda\mathfrak{S}_2, \quad (2.43)$$

where we used $S_{t_0}P_{t_0} = 0$. Multiplying both sides of (2.42) by $\frac{-1}{2\pi\mathbf{i}}\zeta$, integrating over γ and using (2.43) we obtain

$$\frac{-1}{2\pi\mathbf{i}} \int_{\gamma} \zeta\tau_2(\zeta)d\zeta = \lambda\mathfrak{S}_2(t - t_0)^2 + o(t - t_0)^2.$$

Let us prove (2.39). Using the expansion of the resolvent (2.10) to the first order, (2.34) and its conjugate version, denoting $\mathfrak{R}(\zeta) = (\mathrm{TR}_{t_0}(\zeta))^*\dot{Q}_{t_0}J\mathrm{TR}_{t_0}(\zeta)$ we obtain

$$\begin{aligned} \tau_4(\zeta) &= (\mathfrak{S}_0(t - t_0) + o(t - t_0))((I - P_{t_0})R_{t_0}(\zeta)(I - P_{t_0}) + \mathfrak{R}(\zeta)(t - t_0) + o(t - t_0)) \\ &\times (\mathfrak{S}_0^*(t - t_0) + o(t - t_0)) = \mathfrak{S}_0(I - P_{t_0})R_{t_0}(\zeta)(I - P_{t_0})\mathfrak{S}_0^*(t - t_0)^2 + o(t - t_0)^2 \end{aligned}$$

Multiplying both sides of the above equation by $\frac{-1}{2\pi\mathbf{i}}\zeta$, integrating over γ and noticing that the function $\zeta \mapsto (I - P_{t_0})R_{t_0}(\zeta)(I - P_{t_0})$ is analytic with respect to ζ near λ we obtain (2.39). \square

Finally, to prove the main assertion, (2.24), one combines (2.37)–(2.39) (add up all equations) together with (2.29), (2.28). \square

We are now ready to present our next principal result, a version of what is sometimes called the Kato Selection Theorem, cf. [18, Theorem II.5.11], that allows one to compute the second order Taylor expansion of the appropriately chosen eigenvalue curves bifurcating from a multiple eigenvalue.

Theorem 2.10. *Assuming Hypotheses 2.2 and 2.6, we let $W_t = P_{t_0}U_t^{-1}\mathcal{A}_tU_tP_{t_0}$ and suppose that \dot{W}_{t_0} , considered as an operator in $\text{ran}(P_{t_0})$, has $m' \leq m$ distinct eigenvalues denoted by $\mu_1, \dots, \mu_{m'}$. We denote the corresponding to μ_i eigenprojection of \dot{W}_{t_0} in $\text{ran}(P_{t_0})$ by $P_{t_0}^{(i)}$, $1 \leq i \leq m'$. Furthermore, for each $i \leq m'$ we suppose that $P_{t_0}^{(i)}\ddot{W}_{t_0}P_{t_0}^{(i)}$ considered as an operator in $\text{ran}(P_{t_0}^{(i)})$ has $m'_i \leq m$ distinct eigenvalues denoted by $\nu_{i,1}, \dots, \nu_{i,m'_i}$. Then for t near t_0 there is a labeling $\lambda_{i,k}(t)$ of the eigenvalue curves of the operator \mathcal{A}_tP_t or W_t such that*

$$\lambda_{i,k}(t) = \lambda + \mu_i(t - t_0) + \frac{1}{2}\nu_{i,k}(t - t_0)^2 + o(t - t_0)^2, \quad k = 1, \dots, m'_i, i = 1, \dots, m'. \quad (2.44)$$

Moreover, there is an orthonormal set of eigenvectors $u_{i,k}$ of the operators $P_{t_0}^{(i)}\ddot{W}_{t_0}P_{t_0}^{(i)}$ (and therefore of the operators \dot{W}_{t_0} , W_{t_0} and \mathcal{A}_{t_0}),

$$\{u_{i,k} \in \text{ran}(P_{t_0}^{(i)}) : k = 1, \dots, m'_i, i = 1, \dots, m'\} \subset \text{ran}(P_{t_0}),$$

such that

$$\begin{aligned} \lambda_{i,k}(t) &= \lambda + \omega(\dot{Q}_{t_0}Tu_{i,k}, Tu_{i,k})(t - t_0) \\ &+ \frac{1}{2}\left(\omega(\ddot{Q}_{t_0}Tu_{i,k}, Tu_{i,k}) - 2\langle T(\mathbb{T}S_{t_0})^*\dot{Q}_{t_0}JTP_{t_0}u_{i,k}, \dot{Q}_{t_0}JTP_{t_0}u_{i,k}\rangle_{\mathfrak{H} \times \mathfrak{H}}\right)(t - t_0)^2 \\ &+ o(t - t_0)^2, \quad k = 1, \dots, m'_i, i = 1, \dots, m', \end{aligned} \quad (2.45)$$

where $\mathbb{T} \in \mathcal{B}(\mathcal{H}_+, \mathfrak{H} \times \mathfrak{H})$ is the trace operator, the symplectic form ω is defined in (2.2), S_{t_0} is the reduced resolvent of the operator \mathcal{A}_{t_0} at λ .

Proof. Taylor's expansion (2.24) yields

$$\frac{1}{(t - t_0)}\left(\frac{1}{(t - t_0)}(W_t - W_{t_0}) - \dot{W}_{t_0}\right) \underset{t \rightarrow t_0}{=} \frac{1}{2}\ddot{W}_{t_0} + o(1).$$

Multiplying this by $P_{t_0}^{(i)}$ from both sides and using the identities

$$P_{t_0}^{(i)}W_{t_0}P_{t_0}^{(i)} = \lambda I_{\text{ran}P_{t_0}^{(i)}}, \quad P_{t_0}^{(i)}\dot{W}_{t_0}P_{t_0}^{(i)} = \mu_i I_{\text{ran}P_{t_0}^{(i)}},$$

we obtain

$$\frac{1}{(t - t_0)}\left(\frac{1}{(t - t_0)}(P_{t_0}^{(i)}W_tP_{t_0}^{(i)} - \lambda) - \mu_i\right) \underset{t \rightarrow t_0}{=} \frac{1}{2}P_{t_0}^{(i)}\ddot{W}_{t_0}P_{t_0}^{(i)} + o(1). \quad (2.46)$$

By continuity of the spectrum, see, e.g., [22, Theorem VIII.24], every point in the set

$$\text{Spec}\left(\frac{1}{2}P_{t_0}^{(i)}\ddot{W}_{t_0}P_{t_0}^{(i)}\right) = \left\{\frac{1}{2}\nu_{i,1}, \dots, \frac{1}{2}\nu_{i,m'_i}\right\}$$

can be approximated by an eigenvalue of the operator in the left-hand side of (2.46) for t near t_0 . Such an eigenvalue is of the form

$$\frac{1}{(t - t_0)}\left(\frac{1}{(t - t_0)}(\lambda_{i,k}(t) - \lambda) - \mu_i\right), \quad \text{for some choice of } \lambda_{i,k}(t) \in \text{Spec}(\mathcal{A}_tP_t|_{\text{ran}(P_t)}) = \text{Spec}(W_t).$$

That is, we have

$$\frac{1}{(t-t_0)} \left(\frac{1}{(t-t_0)} (\lambda_{i,k}(t) - \lambda) - \mu_i \right) \underset{t \rightarrow t_0}{=} \frac{1}{2} \nu_{i,k} + o(1),$$

hence, (2.44) holds as asserted.

To prove (2.45), let $u_{i,k} \in \text{ran}(P_{t_0}^{(i)})$ be the normalized eigenvectors of \ddot{W}_{t_0} such that $\ddot{W}_{t_0} u_{i,k} = \nu_{i,k} u_{i,k}$. Since $u_{i,k} \in \text{ran}(P_{t_0}^{(i)})$ we have $\dot{W}_{t_0} u_{i,k} = \mu_i u_{i,k}$. Using these identities and (2.25), (2.26) we arrive at

$$\begin{aligned} \mu_i &= \langle \dot{W}_{t_0} u_{i,k}, u_{i,k} \rangle_{\mathcal{H}} = \omega(\dot{Q}_{t_0} \text{T}u_{i,k}, \text{T}u_{i,k}), \\ \nu_{i,k} &= \langle \ddot{W}_{t_0} u_{i,k}, u_{i,k} \rangle_{\mathcal{H}} = \omega(\ddot{Q}_{t_0} \text{T}u_{i,k}, \text{T}u_{i,k}) - 2 \langle \text{T}(\text{TS}_{t_0})^* \dot{Q}_{t_0} \text{J} \text{T}u_{i,k}, \dot{Q}_{t_0} \text{J} \text{T}P_{t_0} u_{i,k} \rangle_{\mathcal{H}}. \end{aligned} \quad (2.47)$$

Finally, combining (2.47) and (2.44) we obtain (2.45) \square

In applications, e.g. such as quantum graphs, spectral theory of differential operators, one often describes a Lagrangian plane $\mathcal{F} \in \Lambda(\mathfrak{H} \times \mathfrak{H})$ by means of two bounded operators $X, Y \in \mathcal{B}(\mathfrak{H})$, satisfying $XY^* = YX^*$, $0 \notin \text{Spec}(XX^* + YY^*)$, via

$$\mathcal{F} = \{(u, v) \in \mathfrak{H} \times \mathfrak{H} : Xu + Yv = 0\} = \ker(Z), \quad \text{where } Z = [X, Y] \in \mathcal{B}(\mathfrak{H} \times \mathfrak{H}, \mathfrak{H}).$$

It is therefore natural to derive the second order Taylor expansion in this setting, which is done in the following theorem.

Theorem 2.11. *Under hypotheses of Theorem 2.10, suppose that the Lagrangian planes $\text{ran}(Q_t) = \text{T}(\text{dom}(\mathcal{A}_t))$ are given by $\text{ran}(Q_t) = \ker Z_t$ where $Z_t := [X_t, Y_t]$ and $X_t, Y_t \in \mathcal{B}(\mathfrak{H})$, $X_t Y_t^* = Y_t X_t^*$, $0 \notin \text{Spec}(X_t X_t^* + Y_t Y_t^*)$, satisfy*

$$Z_t \underset{t \rightarrow t_0}{=} Z_{t_0} + \dot{Z}_{t_0}(t - t_0) + \frac{1}{2} \ddot{Z}_{t_0}(t - t_0)^2 + o(t - t_0)^2 \text{ in } \mathcal{B}(\mathfrak{H} \times \mathfrak{H}).$$

Then the eigenvalue curves $\lambda_{i,k}(t)$, defined as in Theorem 2.10, bifurcating from the isolated eigenvalue $\lambda \in \text{Spec}(\mathcal{A}_{t_0})$ admit the second order Taylor expansion

$$\begin{aligned} \lambda_{i,k}(t) \underset{t \rightarrow t_0}{=} & \lambda + \langle Z_{t_0} \text{J} \dot{Z}_{t_0}^* \phi_{i,k}, \phi_{i,k} \rangle_{\mathfrak{H}} (t - t_0) \\ & + \frac{1}{2} \left(\langle Z_{t_0} \text{J} \ddot{Z}_{t_0}^* \phi_{i,k}, \phi_{i,k} \rangle_{\mathfrak{H}} - 2 \langle \text{T}(\text{TS})^* \dot{Z}_{t_0}^* \phi_{i,k}, \dot{Z}_{t_0}^* \phi_{i,k} \rangle_{\mathfrak{H} \times \mathfrak{H}} \right) (t - t_0)^2 \\ & + o(t - t_0)^2, \end{aligned} \quad (2.48)$$

where S_{t_0} is the reduced resolvent of \mathcal{A}_{t_0} at λ , and we introduce notation

$$\phi_{i,k} = (Z_{t_0} Z_{t_0}^*)^{-1} Z_{t_0} \text{J} \text{T}u_{i,k}$$

for $u_{i,k}$ given in Theorem 2.10.

Proof. For brevity, we denote $u = u_{i,k}$, $\phi := \phi_{i,k}$ and suppress dependence of P_{t_0} , Z_{t_0} , Q_{t_0} and their derivatives on t_0 , e.g., \ddot{Z}_{t_0} will be denoted by \ddot{Z} . Let us record some relevant identities. First, the orthogonal projection Q onto the kernel of Z , i.e. onto the Lagrangian plane $\text{T}(\text{dom}(\mathcal{A}))$, can be written as

$$Q = -\text{J}Z^*(ZZ^*)^{-1}Z\text{J},$$

thus we have $\text{T}u = Q\text{T}u$, and, by $\phi = (ZZ^*)^{-1}Z\text{J}\text{T}u$, one also has

$$\text{T}u = -\text{J}Z^*\phi. \quad (2.49)$$

In addition, the identity $XY^* = YX^*$ yields $ZJZ^* = 0$, therefore, by differentiating we arrive at

$$\dot{Z}JZ^* = -ZJ\dot{Z}^*, \quad (2.50)$$

$$\ddot{Z}JZ^* = -2\dot{Z}J\dot{Z}^* - ZJ\ddot{Z}^*. \quad (2.51)$$

Furthermore, since $\text{ran}(Q) = \ker(Z)$ we have $ZQ = 0$, hence,

$$\begin{aligned} \dot{Z}Q + Z\dot{Q} &= 0, \\ Q\dot{Z}^* + \dot{Q}Z^* &= 0. \end{aligned} \quad (2.52)$$

Also, since $QJ + JQ = J$ we get

$$\dot{Q}J = -J\dot{Q}, \quad \ddot{Q}J = -J\ddot{Q}. \quad (2.53)$$

Let us rewrite the projection Q onto the Lagrangian plane as $Q = \alpha\beta\gamma$ with

$$\alpha := -JZ^* \in \mathcal{B}(\mathfrak{H}, \mathfrak{H} \times \mathfrak{H}), \quad \beta := (ZZ^*)^{-1} \in \mathcal{B}(\mathfrak{H}), \quad \gamma := ZJ \in \mathcal{B}(\mathfrak{H} \times \mathfrak{H}, \mathfrak{H}).$$

We proceed by introducing three more auxiliary identities (2.54), (2.55), (2.56) below. One has

$$\dot{Q}JTu = -Q\dot{Z}^*\phi. \quad (2.54)$$

To prove this, we note that $\gamma JTu = 0$, therefore

$$\dot{Q}JTu = \alpha\beta\dot{\gamma}JTu = -\alpha\beta\dot{Z}Tu \stackrel{(2.49)}{=} \alpha\beta\dot{Z}JZ^*\phi \stackrel{(2.50)}{=} -\alpha\beta ZJ\dot{Z}^*\phi = -Q\dot{Z}^*\phi.$$

We also have

$$\dot{Q}Tu = (1 - Q)T\dot{P}u. \quad (2.55)$$

One arrives at this identity by differentiating both sides of the equation $QTP = TP$ and noting that $Pu = u$. Next, we record the identity

$$\dot{Q}Tu = (1 - Q)T(TS)^*\dot{Q}JTu, \quad (2.56)$$

which follows from (2.55), explicit formula for \dot{P} in (2.21), and the fact that the second term in the RHS of (2.21) vanishes since $S_{t_0}u = 0$.

We are now ready to prove (2.48). To that end we employ (2.45) together with the auxiliary identities (2.49)-(2.56). It suffices to show that

$$\omega(\dot{Q}Tu, Tu) = \langle ZJ\dot{Z}^*\phi, \phi \rangle_{\mathfrak{H}}, \quad (2.57)$$

$$\begin{aligned} \omega(\ddot{Q}Tu, Tu) - 2\langle T(TS)^*\dot{Q}JTu, \dot{Q}JTu \rangle_{\mathfrak{H} \times \mathfrak{H}} &= \\ &= \langle ZJ\ddot{Z}^*\phi, \phi \rangle_{\mathfrak{H}} - 2\langle T(TS)^*\dot{Z}^*\phi, \dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}}. \end{aligned} \quad (2.58)$$

Let us first derive (2.57). One has

$$\begin{aligned} \omega(\dot{Q}Tu, Tu) &= \langle J\dot{Q}Tu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} \stackrel{(2.53)}{=} -\langle \dot{Q}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} \stackrel{(2.54)}{=} \langle Q\dot{Z}^*\phi, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ &= \langle \dot{Z}^*\phi, QTu \rangle_{\mathfrak{H} \times \mathfrak{H}} = \langle \dot{Z}^*\phi, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} \stackrel{(2.49)}{=} \langle \dot{Z}^*\phi, -JZ^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}} = \langle ZJ\dot{Z}^*\phi, \phi \rangle_{\mathfrak{H}}. \end{aligned}$$

The derivation of (2.58) is more subtle. Our immediate objective is to derive the following,

$$\omega(\ddot{Q}Tu, Tu) = \langle ZJ\ddot{Z}^*\phi, \phi \rangle_{\mathfrak{H}} - 2\omega(Q\dot{Z}^*\phi, \dot{Z}^*\phi), \quad (2.59)$$

$$\omega(Q\dot{Z}^*\phi, \dot{Z}^*\phi) = \langle T(TS)^*\dot{Z}^*\phi, (1 - Q)\dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}}, \quad (2.60)$$

$$\langle T(TS)^*\dot{Q}JTu, \dot{Q}JTu \rangle_{\mathfrak{H} \times \mathfrak{H}} = \langle T(TS)^*\dot{Z}^*\phi, Q\dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}}, \quad (2.61)$$

since (2.58) readily follows from (2.59)–(2.61). To begin the proof of (2.59) we differentiate $Q = \alpha\beta\gamma$ twice. Since $\gamma JTu = 0$ and infer

$$\ddot{Q}JTu = \alpha\beta\ddot{\gamma}JTu + 2(\dot{\alpha}\beta + \alpha\dot{\beta})\dot{\gamma}JTu.$$

Plugging this into the left-hand side of (2.59) we arrive at

$$\begin{aligned} \omega(\ddot{Q}Tu, Tu) &= -\langle \ddot{Q}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ &= -\langle \alpha\beta\ddot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} - 2\langle (\dot{\alpha}\beta + \alpha\dot{\beta})\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}}. \end{aligned} \quad (2.62)$$

Let us re-write both terms in the right-hand side of (2.62). The first term is

$$\begin{aligned} -\langle \alpha\beta\ddot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} &\stackrel{(2.49)}{=} -\langle \alpha\beta\ddot{Z}JZ^*\phi, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ &\stackrel{(2.51)}{=} 2\langle \alpha\beta\dot{Z}J\dot{Z}^*\phi, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} + \langle \alpha\beta ZJ\ddot{Z}^*\phi, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ &= 2\langle (ZZ^*)^{-1}\dot{Z}J\dot{Z}^*\phi, ZZ^*\phi \rangle_{\mathfrak{H}} + \langle \alpha\beta\gamma\ddot{Z}^*\phi, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ &= 2\langle J\dot{Z}^*\phi, \dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}} + \langle Q\ddot{Z}^*\phi, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} \stackrel{(2.49)}{=} \langle ZJ\ddot{Z}^*\phi, \phi \rangle_{\mathfrak{H}}, \end{aligned}$$

where we used $\langle J\dot{Z}^*\phi, \dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}} = 0$. In order to re-write the second term in the right-hand side of (2.62) we first notice that $\beta = (ZZ^*)^{-1}$ yields

$$\dot{\beta} = -\beta(\dot{Z}Z^* + Z\dot{Z}^*)\beta,$$

thus

$$\begin{aligned} &\langle (\dot{\alpha}\beta + \alpha\dot{\beta})\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ &= \langle \dot{\alpha}\beta\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} - \langle \alpha\beta\dot{Z}Z^*\beta\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} - \langle \alpha\beta Z\dot{Z}^*\beta\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ &= \langle \dot{\alpha}\beta\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} - \langle \alpha\beta\dot{Z}Z^*\beta\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} + \langle \alpha\beta ZJJ\dot{Z}^*\beta\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ &= \langle \dot{\alpha}\beta\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} - \langle \alpha\beta\dot{Z}Z^*\beta\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} - \langle \alpha\beta\gamma\dot{\alpha}\beta\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ &= \langle \dot{\alpha}\beta\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} - \langle \alpha\beta\dot{Z}Z^*\beta\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} - \langle \dot{\alpha}\beta\dot{\gamma}JTu, QTu \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ &= -\langle \alpha\beta\dot{Z}Z^*\beta\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}}, \end{aligned} \quad (2.63)$$

where we used $Q = \alpha\beta\gamma$, $QTu = Tu$. Now, let us further simplify the last expression. One has

$$\begin{aligned} -\langle \alpha\beta\dot{Z}Z^*\beta\dot{\gamma}JTu, Tu \rangle_{\mathfrak{H} \times \mathfrak{H}} &= -\langle \dot{Z}Z^*(ZZ^*)^{-1}\dot{Z}JJTu, (ZZ^*)^{-1}ZJTu \rangle_{\mathfrak{H}} \\ &\stackrel{(2.49)}{=} -\langle \dot{Z}Z^*(ZZ^*)^{-1}\dot{Z}JZ^*\phi, \phi \rangle_{\mathfrak{H}} \stackrel{(2.50)}{=} \langle \dot{Z}Z^*(ZZ^*)^{-1}ZJ\dot{Z}^*\phi, \phi \rangle_{\mathfrak{H}} \\ &= \langle J(-JZ)^*(ZZ^*)^{-1}ZJ\dot{Z}^*\phi, \dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}} = \omega(Q\dot{Z}^*\phi, \dot{Z}^*\phi). \end{aligned} \quad (2.64)$$

Combining (2.62), (2.63) and (2.64) one infers (2.59) as asserted.

Next, we prove (2.60). One has

$$\begin{aligned} \omega(Q\dot{Z}^*\phi, \dot{Z}^*\phi) &= \langle JQ\dot{Z}^*\phi, \dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}} \stackrel{(2.52)}{=} -\langle J\dot{Q}Z^*\phi, \dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}} \stackrel{(2.53)}{=} \langle \dot{Q}JZ^*\phi, \dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ &= -\langle \dot{Q}Tu, \dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}} \stackrel{(2.56)}{=} -\langle (1-Q)T(TS)^*\dot{Q}JTu, \dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ &\stackrel{(2.54)}{=} \langle (1-Q)T(TS)^*Q\dot{Z}^*\phi, \dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ &= \langle T(QTS)^*\dot{Z}^*\phi, (1-Q)\dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}} = \langle T(TS)^*\dot{Z}^*\phi, (1-Q)\dot{Z}^*\phi \rangle_{\mathfrak{H} \times \mathfrak{H}}, \end{aligned}$$

where we used $QTS = TS$ which holds since $\text{ran}(S) \subset \text{ran}(Q)$.

To complete the proof of the theorem we derive (2.61). To that end, we have

$$\begin{aligned} \langle T(TS)^* \dot{Q}JT u, \dot{Q}JT u \rangle_{\mathfrak{H} \times \mathfrak{H}} & \stackrel{(2.54)}{=} \langle T(TS)^* Q \dot{Z}^* \phi, Q \dot{Z}^* \phi \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ & = \langle T(QTS)^* \dot{Z}^* \phi, Q \dot{Z}^* \phi \rangle_{\mathfrak{H} \times \mathfrak{H}} = \langle T(TS)^* \dot{Z}^* \phi, Q \dot{Z}^* \phi \rangle_{\mathfrak{H} \times \mathfrak{H}}, \end{aligned}$$

where, as before, we used $QTS = TS$. \square

Corollary 2.12. *Under the hypotheses and general setting of Theorem 2.11 we assume that $X_t = \Theta_t$, $Y_t = -I$, that is, we assume that \mathcal{A}_t is a Robin-type self-adjoint extension of A with $\text{dom}(\mathcal{A}_t) = \{u \in \mathcal{H}_+ : \Gamma_1 u = \Theta_t \Gamma_0 u\}$, and that $\Theta_t = \Theta_t^*$. Then one has*

$$\begin{aligned} \lambda_{i,k}(t) \Big|_{t \rightarrow t_0} & = \lambda + \langle \dot{\Theta} \Gamma_0 u_0, \Gamma_0 u_0 \rangle_{\mathfrak{H}} (t - t_0) \\ & + \frac{1}{2} \left(\langle (\ddot{\Theta}_{t_0} - 2\dot{\Theta}_{t_0} \Gamma_0 (\Gamma_0 S_{t_0})^* \dot{\Theta}_{t_0}) \Gamma_0 u_{i,k}, \Gamma_0 u_{i,k} \rangle_{\mathfrak{H}} \right) (t - t_0)^2 \\ & + o(t - t_0)^2. \end{aligned} \quad (2.65)$$

Proof. Formula (2.65) follows directly from (2.48) with $Z_t = [\Theta_t, -I]$. \square

Example 2.13 (Matrix Robin Laplacian). Let $V \in L^\infty([a, b], \mathbb{C}^{n \times n})$ be a matrix valued bounded function with $V(x) = (V(x))^*$, $x \in [a, b]$, and let $A = -\frac{d^2}{dx^2} + V$ be the minimal operator with $\text{dom}(A) = H_0^2([a, b]; \mathbb{C}^n)$. Given $u \in H^2([a, b], \mathbb{C}^n)$ we denote $\gamma_D u = [u(a), u(b)]^\top$, $\gamma_N u = [u(a), -u'(b)]^\top$ and view γ_D, γ_N as bounded linear operators in $\mathcal{B}(H^2([a, b], \mathbb{C}^{2n}); \mathbb{C}^{2n})$. Let $\Theta \in C^2([0, 1], \mathbb{C}^{2n \times 2n})$ be a matrix-valued function with $\Theta_t = \Theta_t^*$, $t \in [0, 1]$, and define a one-parameter family of Robin extensions of A by the formula

$$\text{dom}(\mathcal{A}_t) = \{u \in H^2([a, b]; \mathbb{C}^n) : \gamma_N u = \Theta_t \gamma_D u\}.$$

Suppose that $\lambda \in \mathbb{R}$ is a simple eigenvalue of \mathcal{A}_{t_0} for some t_0 and suppose that $u_0 \in \ker(\mathcal{A}_{t_0} - \lambda)$ has unit norm. Then there is an eigenvalue curve bifurcating from λ and obeying the following asymptotic expansion,

$$\begin{aligned} \lambda(t) & = \lambda + \left\langle \dot{\Theta}_{t_0} \gamma_D u, \gamma_D u \right\rangle_{\mathbb{C}^{2n}} (t - t_0) \\ & + \frac{1}{2} \left\langle (\ddot{\Theta}_{t_0} - 2\dot{\Theta}_{t_0} \gamma_D (\gamma_D S)^* \dot{\Theta}_{t_0}) \gamma_D u, \gamma_D u \right\rangle_{\mathbb{C}^{2n}} (t - t_0)^2 + o(t - t_0)^2, t \rightarrow t_0. \end{aligned}$$

Remark 2.14. Our main results, Theorem 2.9, 2.10 and Theorem 2.11 can be extended to a more general setting in which the self-adjoint extensions are subject to further additive perturbations $\mathcal{A}_t + V_t$. To elaborate on this case, we augment Hypothesis 2.2 by fixing a one-parameter family of bounded self-adjoint operators $t \mapsto V_t \in \mathcal{B}(\mathcal{H})$ with

$$V_t \Big|_{t \rightarrow t_0} = V_{t_0} + \dot{V}_{t_0} (t - t_0) + \frac{1}{2} \ddot{V}_{t_0} (t - t_0)^2 + o(t - t_0)^2 \text{ in } \mathcal{B}(\mathcal{H}).$$

Let us introduce $H_t := \mathcal{A}_t + V_t$, denote its resolvent by $R_t(\zeta) := (H_t - \zeta)^{-1} \in \mathcal{B}(\mathcal{H})$ for $\zeta \notin \text{Spec}(H_t)$, and its reduced resolvent at the eigenvalue λ by S_{t_0} . Keeping general setting of Theorem 2.10 with \mathcal{A}_t replaced by H_t we record the first and second order terms in the Taylor expansion of the eigenvalue curves. The first order term is given by

$$\dot{\lambda}_{i,k} = \langle \dot{V}_{t_0} u_{i,k}, u_{i,k} \rangle_{\mathcal{H}} + \omega(\dot{Q}_{t_0} \Gamma u_{i,k}, \Gamma u_{i,k}).$$

The second order term is given by

$$\begin{aligned} \ddot{\lambda}_{i,k} = & \frac{1}{2} \left(\langle \ddot{V}_{t_0} u_{i,k}, u_{i,k} \rangle_{\mathcal{H}} + \omega(\ddot{Q}_{t_0} T u_{i,k}, T u_{i,k}) \right) - \langle T(TS_{t_0})^* \dot{Q}_{t_0} JTP_{t_0} u_{i,k}, \dot{Q}_{t_0} JTP_{t_0} u_{i,k} \rangle_{\mathfrak{H} \times \mathfrak{H}} \\ & + 2 \langle TS \dot{V}_{t_0} P u_{i,k}, \dot{Q}_{t_0} JTP_{t_0} u_{i,k} \rangle_{\mathfrak{H} \times \mathfrak{H}} - \langle \dot{V}_{t_0} S_{t_0} \dot{V}_{t_0} u_{i,k}, u_{i,k} \rangle_{\mathcal{H}}. \end{aligned}$$

The derivation relies on Krein's formula from [19],

$$R_t(\zeta) - R_{t_0}(\zeta) = R_t(\zeta)(V_{t_0} - V_t)R_{t_0}(\zeta) + (TR_t(\zeta))^*(Q_t - Q_{t_0})JQ_{t_0}TR_{t_0}(\zeta),$$

and resolvent expansions arguments presented above.

REFERENCES

- [1] C. BANDLE AND A. WAGNER, *Second domain variation for problems with Robin boundary conditions*, J. Optim. Theory Appl., 167 (2015), pp. 430–463.
- [2] J. BEHRNDT, S. HASSI, AND H. DE SNOO, *Boundary Value Problems, Weyl Functions, and Differential Operators*, Springer International Publishing, 2020.
- [3] G. BERKOLAIKO, Y. CANZANI, G. COX, AND J. L. MARZUOLA, *A local test for global extrema in the dispersion relation of a periodic graph*, arxiv: 2004.12931.
- [4] G. BERKOLAIKO, G. COX, AND J. L. MARZUOLA, *Nodal deficiency, spectral flow, and the Dirichlet-to-Neumann map*, Lett. Math. Phys., 109 (2019), pp. 1611–1623.
- [5] G. BERKOLAIKO AND P. KUCHMENT, *Spectral shift via “lateral” perturbation*, arxiv: 2011.11142.
- [6] B. BOOSS BAVNBEK AND C. ZHU, *The Maslov index in symplectic Banach spaces*, Mem. Amer. Math. Soc., 252 (2018), pp. x+118.
- [7] V. I. BURENKOV, P. D. LAMBERTI, AND M. LANTSA DE KRISTOFORIS, *Spectral stability of nonnegative selfadjoint operators*, Sovrem. Mat. Fundam. Napravl., 15 (2006), pp. 76–111.
- [8] G. COX, C. K. R. T. JONES, Y. LATUSHKIN, AND A. SUKHTAYEV, *The Morse and Maslov indices for multidimensional Schrödinger operators with matrix-valued potentials*, Trans. Amer. Math. Soc., 368 (2016), pp. 8145–8207.
- [9] G. COX, C. K. R. T. JONES, AND J. L. MARZUOLA, *A Morse index theorem for elliptic operators on bounded domains*, Comm. Partial Differential Equations, 40 (2015), pp. 1467–1497.
- [10] G. COX, C. K. R. T. JONES, AND J. L. MARZUOLA, *Manifold decompositions and indices of Schrödinger operators*, Indiana Univ. Math. J., 66 (2017), pp. 1573–1602.
- [11] J. DENG AND C. JONES, *Multi-dimensional Morse index theorems and a symplectic view of elliptic boundary value problems*, Trans. Amer. Math. Soc., 363 (2011), pp. 1487–1508.
- [12] L. FRIEDLANDER, *Some inequalities between Dirichlet and Neumann eigenvalues*, Arch. Rational Mech. Anal., 116 (1991), pp. 153–160.
- [13] V. I. GORBACHUK AND M. L. GORBACHUK, *Boundary value problems for operator differential equations*, vol. 48 of Mathematics and its Applications (Soviet Series), Kluwer Academic Publishers Group, Dordrecht, 1991. Translated and revised from the 1984 Russian original.
- [14] P. GRINFELD, *Hadamard's formula inside and out*, J. Optim. Theory Appl., 146 (2010), pp. 654–690.
- [15] B. HELFFER AND M. P. SUNDQVIST, *Spectral flow for pair compatible equipartitions*, Communications in Partial Differential Equations, 0 (2021), pp. 1–28.
- [16] A. HENROT, *Extremum problems for eigenvalues of elliptic operators*, Frontiers in Mathematics, Birkhäuser Verlag, Basel, 2006.
- [17] D. HENRY, *Perturbation of the boundary in boundary-value problems of partial differential equations*, vol. 318 of London Mathematical Society Lecture Note Series, Cambridge University Press, Cambridge, 2005. With editorial assistance from Jack Hale and Antônio Luiz Pereira.
- [18] T. KATO, *Perturbation theory for linear operators*, Classics in Mathematics, Springer-Verlag, Berlin, 1995. Reprint of the 1980 edition.
- [19] Y. LATUSHKIN AND S. SUKHTAYEV, *First-order asymptotic perturbation theory for extensions of symmetric operators*, arXiv:2012.00247.
- [20] Y. LATUSHKIN AND A. SUKHTAYEV, *Hadamard-type formulas via the Maslov form*, J. Evol. Equ., 17 (2017), pp. 443–472.

- [21] M. REED AND B. SIMON, *Methods of modern mathematical physics. I. Functional analysis*, Academic Press, New York-London, 1972.
- [22] ———, *Methods of modern mathematical physics. IV. Analysis of operators*, Academic Press, New York, 1978.
- [23] J. ROBBIN AND D. SALAMON, *The spectral flow and the Maslov index*, Bull. London Math. Soc., 27 (1995), pp. 1–33.
- [24] K. SCHMÜDGEN, *Unbounded self-adjoint operators on Hilbert space*, vol. 265 of Graduate Texts in Mathematics, Springer, Dordrecht, 2012.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA
Email address: `latushkiny@missouri.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, AUBURN UNIVERSITY, AUBURN, AL 36849, USA
Email address: `szs0266@auburn.edu`