

LINKING COEFFICIENTS AND THE KONTSEVICH INTEGRAL

JEAN-BAPTISTE MEILHAN

ABSTRACT. It is well known how the linking number and framing can be extracted from the degree 1 part of the (framed) Kontsevich integral. This note gives a general formula expressing any product of powers of these two invariants as combination of coefficients in the Kontsevich integral. This allows in particular to express the sum of all coefficients of a given degree in terms of the linking coefficients. The proofs are purely combinatorial.

1. INTRODUCTION

The *Kontsevich integral* is a strong invariant of framed oriented knots and links, which dominates all finite type and quantum invariants, in the sense that any other factors through it. It takes values in a certain space of *chord diagrams*, which are copies of the oriented unit circle, endowed with a number of chords, which are pairings of pairwise disjoint points on the circles; chord diagrams naturally come with a degree, which is given by the number of chords. Kontsevich defined this invariant in terms of iterated integrals, what can be seen as a far-reaching generalization of the Gauss integral for the linking number of two curves [5]. As a matter of fact, it is well-known that the linking number appears as the simplest coefficient in the Kontsevich integral. Specifically, given a framed link L , denoting by $C_L[D]$ the coefficient of a chord diagram D in the Kontsevich integral of L , and denoting by ℓ_{ij} the linking number of the i th and j th components, we have

$$(1_1) \quad \ell_{ij}(L) = C_L[\textcircled{i} \text{---} \textcircled{j}].$$

Denoting half the framing of the i th component of L by $\ell_{ii}(L)$, it is also well-known that

$$(2_1) \quad \ell_{ii}(K) = \frac{1}{2} fr_i(L) = C_L[\textcircled{\text{---}}_i].$$

Hence the degree 1 part of the Kontsevich integral of a link is fully characterized by the linking coefficients, *i.e.* the coefficients of the linking matrix. The main purpose of this note is a formula generalizing these two elementary results.

Let \mathcal{S}_m be the set of symmetric matrices of size m with coefficients in \mathbb{N} . Given $S = (s_{ij})_{i,j} \in \mathcal{S}_m$, we define $\mathcal{D}_S(m)$ as the set of all possible chord diagrams on m circles with exactly s_{ij} chords of type (i, j) for all i, j . Here, a *type (i, j) chord* is a chord whose endpoints sit on components i and j ; in particular, a type (i, i) chord has both endpoints on the i th component.

Theorem 1. Let L be an m -component framed oriented link in S^3 and let $S = (s_{ij})_{i,j} \in \mathcal{S}_m$. We have

$$\ell_S(L) := \prod_{1 \leq i \leq j \leq m} \frac{1}{s_{ij}!} \ell_{ij}(L)^{s_{ij}} = \sum_{D \in \mathcal{D}_S(m)} C_L[D].$$

Key words and phrases. Kontsevich integral, Jacobi diagrams, linking number, framing.

This general formula has several noteworthy consequences.

On one hand, if S has a single nonzero entry $n = s_{ij}$ with $i < j$, we obtain a generalization of (1₁) to all powers of the linking number:

$$(1_n) \quad \frac{1}{n!} \ell_{ij}(L)^n = \sum_{D \in \mathcal{M}_n^{ij}(m)} C_L[D],$$

where $\mathcal{M}_n^{ij}(m)$ denotes the set of all possible degree n chord diagrams on m circles whose n chords are of type (i, j) .

Similarly, if S has a single nonzero entry $n = s_{ii}$ on the diagonal, we obtain that

$$(2_n) \quad \frac{1}{n! 2^n} fr_i(L)^n = \sum_{D \in \mathcal{I}_n^i(m)} C_L[D],$$

where $\mathcal{I}_n^i(m)$ denotes the set of all degree n chord diagrams on m circles, such that all n chords are on the i th circle.

On the other hand, the set $\mathcal{D}_k(m)$ of all degree k chord diagrams on m circles is partitioned into the sets $\mathcal{D}_S(m)$ for all matrices S in \mathcal{S}_m with $|S| = k$, where $|S| = \sum_{1 \leq i < j \leq m} s_{ij}$ is the *degree* of S . Thus we have:

$$(3) \quad \sum_{D \in \mathcal{D}_k(m)} C_L[D] = \sum_{S \in \mathcal{S}_m : |S| = k} \ell_S(L).$$

This expresses the sum of *all* coefficients of degree k in the Kontsevich integral in terms of the linking coefficients.

More generally, Theorem 1 gives partial, but general (*i.e.* in all degrees) identifications of the coefficients of the Kontsevich integral in terms of the coefficients of the linking matrix. A number of works investigate, in a similar way, how combinations of coefficients in the Kontsevich integral identify with classical invariants of knot theory, see for example [10, 4, 8, 9, 2, 3], although such results are often only given for low degree terms.

Acknowledgments. This work is partially supported by the project AlMaRe (ANR-19-CE40-0001-01) of the ANR. The author wishes to thank Georges Abitbol and Benjamin Audoux for inspiring discussions.

2. THE FRAMED KONTSEVICH INTEGRAL IN A NUTSHELL


We briefly review the combinatorial definition of the framed Kontsevich integral, as given by Le and Murakami in [6]; see also [7, §6].

A *chord diagram* D on the disjoint union \bigcirc^m of m copies of the oriented circle, is a collection of copies of the unit interval, such that the set of all endpoints is embedded into \bigcirc^m . We call *chord* any of these copies of the interval. The *degree* of D is defined as its number of chords.

In figures, bold lines depict (portions of) \bigcirc^m , and dashed lines are used for chords.

We denote by $\mathcal{A}(m)$ the \mathbb{Q} -vector space generated by all chord diagrams on \bigcirc^m , modulo the $4T$ relation:

We now describe the source of the framed Kontsevich integral. A framed q -tangle is an oriented tangle, equipped with a framing and a parenthesization on both sets of boundary points. Any such tangle can be decomposed into copies of the q -tangles I , X_{\pm} , C_{\pm} and Λ_{\pm} shown in Figure 2.1, along with those obtained by reversing the orientation on any component. Such a decomposition is not unique, but a complete set of relations is known, relating any two possible decompositions,

FIGURE 2.1. The elementary q-tangles I , X_{\pm} , C_{\pm} and Λ_{\pm}

see [7, Thm. 6.5]. The *framed Kontsevich integral* Z can thus be determined by specifying its values on the above q-tangles so that all relations are satisfied. This is done as follows.

We set $Z(I)$ to be the portion of diagram \uparrow without chord.

For the positive and negative crossings X_{\pm} , we set

$$(2.1) \quad Z(X_{\pm}) = \sum_{k \geq 0} \frac{(\pm 1)^k}{2^k k!} X_k, \text{ where } X_k = \begin{array}{c} \uparrow \\ \text{---} \\ \times \\ \text{---} \\ \uparrow \end{array} \Big|_{k \text{ chords}}.$$

Next, set $Z(C_{\pm}) = \sqrt{\nu}$, where $\nu \in \mathcal{A}(1)$ is the Kontsevich integral of the 0-framed unknot U_0 , which was explicitly computed in [1] as follows:

$$(2.2) \quad \nu = \chi \left(\exp_{\sqcup} \left(\sum_{n \geq 1} b_{2n} W_{2n} \right) \right),$$

where W_{2n} is a *wheel*, that is a univalent diagram of the form  with $2n$

univalent vertices, b_{2n} is the coefficient of x^{2n} in the Taylor expansion of $\frac{1}{2} \ln \frac{\sinh(x/2)}{x/2}$, \exp_{\sqcup} stands for the exponential with respect to the disjoint union of diagrams, and the map χ takes the sum over all possible ways of attaching the univalent vertices of a (disjoint union of) wheel(s) to \circlearrowleft , then applies recursively the STU relation below to produce a combination of chord diagrams:

$$\begin{array}{c} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} - \begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \end{array}.$$

Lastly, we set $Z(\Lambda_{\pm}) = \Phi^{\pm 1}$, where Φ is a *Drinfeld associator*; we do not further discuss here this important ingredient of the construction, as it will play no role in our argument. The interested reader is referred to [7, App. D].

3. LINKING COEFFICIENTS AND THE KONTSEVICH INTEGRAL

The purpose of this section is to prove Theorem 1. This will be done in Subsection 3.3, after setting some notation and preliminary results in the next two subsections.

3.1. Some notation. Let A be an element of $\mathcal{A}(m)$. Let D be a chord diagram on m circles. We denote by $C[D](A)$ the coefficient of D in A . In particular, we set

$$C_L[D] := C[D](Z(L)),$$

for a framed oriented link L , that is, $C_L[D]$ denotes ‘the coefficient’ of diagram D in the Kontsevich integral of L . This quantity is of course in general not well-defined since an element of $\mathcal{A}(m)$ consists of diagrams subject to the $4T$ relation, but taking an appropriate combination of such coefficients shall yield a link invariant, see Claim 3.1 below.

We denote by $C[D]$ the assignment $A \mapsto C[D](A)$ and, abusing notation, we still denote by $C[D]$ the precomposition $L \mapsto C_L[D]$ with the Kontsevich integral.

Let $S = (s_{ij})_{i,j} \in \mathcal{S}_m$ be a symmetric matrix of size m with entries in \mathbb{N} . We set

$$\mathcal{L}_S := \sum_{D \in \mathcal{D}_S(m)} C[D],$$

where $\mathcal{D}_S(m)$ is the set of all chord diagrams on m circles with exactly s_{ij} chords of type (i, j) for all $i \leq j$, as defined in the introduction.

Claim 3.1. This formula yields a well-defined map on $\mathcal{A}(m)$, and in particular defines an m -component link invariant.

This is straightforwardly checked using the following general invariance criterion: given a collection \mathcal{D} of chord diagrams, the assignment $X := \sum_{D \in \mathcal{D}} C[D]$ defines a link invariant if and only if X vanishes on any linear combination of chord diagrams arising from a $4T$ relation. See for example [2].

We also recall from the introduction the link invariant associated with the symmetric matrix S ,

$$\ell_S = \prod_{1 \leq i < j \leq m} \frac{1}{s_{ij}!} \ell_{ij}^{s_{ij}},$$

where ℓ_{ij} is the linking number between components i and j if $i \neq j$, and $\ell_{ii} := \frac{1}{2} fr_i$ is half the framing of the i th component.

3.2. Crossing change formula for the invariant \mathcal{L}_S . Let us pick some indices a, b in $\{1, \dots, m\}$ (possibly with $a = b$). Consider two m -component links L_+ and L_- , that are identical away from a small 3-ball, where they look as follows:

$$L_+ = \begin{array}{c} \nearrow \\ \text{a} \times \text{b} \\ \searrow \end{array} \quad \text{and} \quad L_- = \begin{array}{c} \nwarrow \\ \text{a} \times \text{b} \\ \nearrow \end{array}.$$

We stress that this crossing change may involve two strands of either the same ($a = b$) or different ($a \neq b$) components.

Now let $S = (s_{ij})_{i,j} \in \mathcal{S}_m$. By the mere definition of the Kontsevich integral at a crossing, we have

$$\mathcal{L}_S(L_+) - \mathcal{L}_S(L_-) = \sum_{j \geq 0} \frac{1}{(2j+1)!2^{2j}} \mathcal{L}_S(D_{2j+1}),$$

where $D_k \in \mathcal{A}(m)$ is obtained from the Kontsevich integral of L_{\pm} by replacing the local contribution of the crossing involved in the crossing change, as given in (2.1), by the local diagram  with exactly k parallel chords.

Set $s := s_{ab} = s_{ba}$, the entry of the matrix S corresponding to our crossing change. In order to slightly simplify our notation, for any p such that $s \geq p \geq 0$ we denote by \mathcal{L}_p the invariant \mathcal{L}_{S_p} , where S_p is the matrix S with the coefficient s replaced by p ; in particular we have $\mathcal{L}_s = \mathcal{L}_S$.

Clearly, we have $\mathcal{L}_s(D_k) = 0$ for $k > s$, hence the variation formula

$$(3.1) \quad \mathcal{L}_S(L_+) - \mathcal{L}_S(L_-) = \sum_{j=0}^{\lfloor \frac{s+1}{2} \rfloor} \frac{1}{(2j+1)!2^{2j}} \mathcal{L}_s(D_{2j+1}).$$

When $k \leq s$, we have the following.

Claim 3.2. For all k such that $s \geq k \geq 0$, we have $\mathcal{L}_s(D_k) = \mathcal{L}_{s-k}(D_0)$.

Proof. A degree n chord diagram on m circles that contributes to $\mathcal{L}_s(D_k)$ necessarily contains k parallel chords of type (a, b) , as imposed by the definition of D_k . The set of all diagrams contributing to $\mathcal{L}_s(D_k)$ is thus obtained by adding $n - k$ chords, with exactly $s - k$ additional chords of type (a, b) , in all possible ways. But since these additional chords do not arise from the crossing change, they are attached

outside a disk containing the k parallel chords of type (a, b) . This is thus equivalent to taking the contribution of *all* chord diagrams in $\mathcal{D}_{S_{s-k}}$. \square

By Claim 3.2, computing the variation $\mathcal{L}_S(L_+) - \mathcal{L}_S(L_-)$ reduces to computing $\mathcal{L}_k(D_0)$ for all k . This is done in the next lemma.

Lemma 3.3. *We have $\mathcal{L}_k(D_0) = \sum_{p=0}^k \frac{(-1)^p}{p!2^p} \mathcal{L}_{k-p}(L_+)$ for all $k; s \geq k \geq 0$.*

Proof. The proof is by induction on k . The formula for $k = 0$ is clear: the invariant \mathcal{L}_0 vanishes on any diagram with a chord of type (a, b) , so that $\mathcal{L}_0(D_k) = 0$ for all $k \geq 1$, and by definition of the Kontsevich integral at a positive crossing (2.1), we thus have $\mathcal{L}_0(L_+) = \mathcal{L}_0(D_0)$. For the inductive step, again by definition of the Kontsevich integral at a positive crossing, we have

$$\begin{aligned} \mathcal{L}_k(D_0) &= \mathcal{L}_k(L_+) - \sum_{j=1}^k \frac{1}{j!2^j} \mathcal{L}_k(D_j) \\ &= \mathcal{L}_k(L_+) - \sum_{j=1}^k \frac{1}{j!2^j} \mathcal{L}_{k-j}(D_0) \\ &= \mathcal{L}_k(L_+) - \sum_{j=1}^k \frac{1}{j!2^j} \sum_{i=0}^{k-j} \frac{(-1)^i}{i!2^i} \mathcal{L}_{k-j-i}(L_+), \end{aligned}$$

where the second equality follows from Claim 3.2, while the third equality uses the induction hypothesis. For each p such that $0 \leq p \leq k$, the coefficient of $\mathcal{L}_{k-p}(L_+)$ in the above double sum is then given by

$$\begin{aligned} - \sum_{j=1}^p \frac{1}{j!2^j} \times \frac{(-1)^{p-j}}{(p-j)!2^{p-j}} &= \frac{-1}{p!2^p} \sum_{j=1}^p (-1)^{p-j} \binom{p}{j} \\ &= \frac{-1}{p!2^p} \underbrace{\left(\sum_{j=0}^p (-1)^{p-j} \binom{p}{j} \right)}_{=0} - (-1)^p \\ &= \frac{(-1)^p}{p!2^p}. \end{aligned}$$

This concludes the proof. \square

3.3. Proof of Theorem 1. We proceed by induction on the degree $|S| = \sum_{i \leq j} s_{ij}$ of S . The base case $n = 1$ corresponds to the case where S has a single nonzero entry $s_{ij} = 1$ ($i \leq j$), and is given by the well-known formulas (1₁) and (2₁) recalled in the introduction.

Now, assume that the formula holds for all matrices of \mathcal{S}_m of degree $< k$. Let $S \in \mathcal{S}_m$ be a degree k matrix. Choose some indices a, b such that, in the matrix S , the entry $s = s_{ab}$ is nonzero (possibly $a = b$). Let L_+ and L_- be two m -component links that differ by a crossing change between components a and b , as in Subsection 3.2. Combining (3.1) with Claim 3.2 and Lemma 3.3, we obtain

$$\mathcal{L}_S(L_+) - \mathcal{L}_S(L_-) = \sum_{j=0}^{\lfloor \frac{s+1}{2} \rfloor} \frac{1}{(2j+1)!2^{2j}} \sum_{k=0}^{s-2j-1} \frac{(-1)^k}{k!2^k} \mathcal{L}_{s-2j-k-1}(L_+).$$

By the induction hypothesis, we have

$$\mathcal{L}_{s-2j-k-1}(L_+) = \frac{1}{(s-2j-k-1)!} \ell_{ab}^{s-2j-k-1} \prod_{\{i,j\} \neq \{a,b\}} \frac{1}{s_{ij}!} \ell_{ij}^{s_{ij}}.$$

Setting $\ell_0 := \prod_{\{i,j\} \neq \{a,b\}} \frac{1}{s_{ij}!} \ell_{ij}^{s_{ij}}$, we thus have

$$\mathcal{L}_S(L_+) - \mathcal{L}_S(L_-) = \sum_{j=0}^{\lfloor \frac{s+1}{2} \rfloor} \frac{\ell_0}{(2j+1)!2^{2j}} \sum_{k=0}^{s-2j-1} \frac{(-1)^k}{k!(s-2j-k-1)!2^k} \ell_{ab}(L_+)^{s-2j-k-1}.$$

The coefficient of $\ell_{ab}(L_+)^{s-i}$ in the above formula is given by

$$\begin{aligned} \sum_{j=0}^{\lfloor \frac{i+1}{2} \rfloor} \frac{\ell_0}{(2j+1)!2^{2j}} \cdot \frac{(-1)^{i-2j-1}}{(i-2j-1)!(s-i)!2^{i-2j-1}} &= \frac{(-1)^{i+1}\ell_0}{2^{i-1}(s-i)!} \sum_{j=0}^{\lfloor \frac{i+1}{2} \rfloor} \frac{1}{i!} \binom{i}{2j+1} \\ &= \frac{(-1)^{i+1}\ell_0}{2^{i-1}i!(s-i)!} \underbrace{\sum_{j=0}^{\lfloor \frac{i+1}{2} \rfloor} \binom{i-1}{2j} + \binom{i-1}{2j+1}}_{=2^{i-1}}. \end{aligned}$$

This shows that

$$\mathcal{L}_S(L_+) - \mathcal{L}_S(L_-) = \ell_0 \sum_{i=1}^n \frac{(-1)^{i+1}}{i!(s-i)!} \ell_{ab}^{s-i}.$$

Now, this formula coincides with the variation of the linking invariant ℓ_S :

$$\ell_S(L_+) - \ell_S(L_-) = \ell_0 \sum_{i=1}^n \frac{(-1)^{i+1}}{i!(s-i)!} \ell_{ab}^{s-i}.$$

This is easily verified using the binomial formula, noting that $\ell_{ab}(L_-) = \ell_{ab}(L_+) - 1$ (in particular, if $a = b$, we indeed have $fr_a(L_-) = fr_a(L_+) - 2$).

Hence we showed that the invariants \mathcal{L}_S and ℓ_S have the same variation formula under a crossing change. By a sequence of such operations, any m -component link can be deformed into a split union of unknots, each with framing 0 or 1 depending on the parity of the framing of the component: it remains to check that both invariants take the same value on such links.

Denote by U_0 and U_1 the unknot with framing 0 or 1, respectively, and let L_0 be a split union of m unknots, such that the i th component is a copy of U_{ε_i} , $\varepsilon_i \in \{0, 1\}$. The framed Kontsevich integral of L_0 can be written in $\mathcal{A}(m)$ as a disjoint union $Z(L_0) = \sqcup_i Z(U_{\varepsilon_i})$; in particular, we may assume that it only contains type (i, i) chords for $1 \leq i \leq m$.

It follows that, if the matrix S contains a nonzero coefficient away from the diagonal, then both invariants \mathcal{L}_S and ℓ_S clearly vanish, and the proof is complete.

In the case where S is a diagonal matrix, the invariant \mathcal{L}_S of L_0 splits as

$$\mathcal{L}_S(L_0) = \prod_{i=1}^m \sum_{D \in \mathcal{S}_{s_{ii}}} C_{U_{\varepsilon_i}}[D],$$

where \mathcal{S}_k denotes the set of all possible chord diagrams on \bigcirc with k chords. The proof then follows readily from the following claim.

Claim 3.4. For all integer k , we have

$$\sum_{D \in \mathcal{S}_k} C_{U_0}[D] = 0 \quad \text{and} \quad \sum_{D \in \mathcal{S}_k} C_{U_1}[D] = \frac{1}{k!2^k}.$$

Indeed, these are the values taken by the invariant $\frac{1}{k!}(\frac{1}{2}fr)^k$ on both U_0 and U_1 , thus showing that the invariants \mathcal{L}_S and ℓ_S do also coincide on L_0 when S is diagonal. Hence it only remains to prove Claim 3.4 to complete the proof.

Proof of Claim 3.4. For simplicity, set $\mathcal{F}_k(K) := \sum_{D \in \mathcal{S}_k} C_K[D]$ for a knot K .

Let us first compute $\mathcal{F}_k(U_0)$. We recalled in (2.2) the computation of $Z(U_0) = \nu$ of [1]. Now, given a (disjoint union of) wheel(s) with k univalent vertices attached to \circlearrowleft in some way, applying recursively the STU relation to get a combination of chord diagrams, as prescribed by χ , produces an alternate sum with 2^k terms, where the coefficients add up to zero. This simple observation shows that $\mathcal{F}_k(U_0) = 0$.

We now consider $\mathcal{F}_k(U_1)$. It is well-known that $\sqrt{\nu}$ commutes with any chord endpoint in a chord diagram (this is a consequence of the 4T relation). This can be used to check that

$$Z(U_1) = Z(U_0) \sharp \left(\exp_{\sharp} \frac{1}{2} \circlearrowleft \right) = Z(U_0) \sharp \left(\sum_{k \geq 0} \frac{1}{k! 2^k} D_k \right),$$

where D_k denotes the chord diagram on \circlearrowleft with k parallel chords and where \sharp is the connected sum of chord diagrams. From the above computation of $\mathcal{F}_k(U_0)$, we obtain that the only term contributing to $\mathcal{F}_k(U_1)$ is the degree k diagram D_k arising from $\exp_{\sharp} \frac{1}{2} \circlearrowleft$. Summarizing, we obtain that $\mathcal{F}_k(U_1) = \frac{1}{k! 2^k}$. \square

REFERENCES

- [1] D. Bar-Natan, S. Garoufalidis, L. Rozansky, and D. P. Thurston. Wheels, wheeling, and the Kontsevich integral of the unknot. *Isr. J. Math.*, 119:217–237, 2000.
- [2] A. Casejuane. *Formules combinatoires pour les invariants de nœuds et de variétés de dimension 3*. PhD thesis, Université Grenoble Alpes, 2021.
- [3] A. Casejuane and J.-B. Meilhan. Universal invariants, the conway polynomial and the casson-walker-lescop invariant, 2020.
- [4] N. Habegger and G. Masbaum. The Kontsevich integral and Milnor’s invariants. *Topology*, 39(6):1253–1289, 2000.
- [5] M. Kontsevich. Vassiliev’s knot invariants. In *I. M. Gelfand seminar. Part 2: Papers of the Gelfand seminar in functional analysis held at Moscow University, Russia, September 1993*, pages 137–150. Providence, RI: American Mathematical Society, 1993.
- [6] T. Q. T. Le and J. Murakami. The universal Vassiliev-Kontsevich invariant for framed oriented links. *Compos. Math.*, 102(1):41–64, 1996.
- [7] T. Ohtsuki. *Quantum invariants. A study of knots, 3-manifolds, and their sets*. Singapore: World Scientific, 2002.
- [8] M. Okamoto. Vassiliev invariants of type 4 for algebraically split links. *Kobe J. Math.*, 14(2):145–196, 1997.
- [9] M. Okamoto. On Vassiliev invariants for algebraically split links. *J. Knot Theory Ramifications*, 7(6):807–835, 1998.
- [10] T. Stanford. Some computational results on mod 2 finite-type invariants of knots and string links. In *Invariants of knots and 3-manifolds. Proceedings of the workshop, Kyoto, Japan, September 17–21, 2001*, pages 363–376. Coventry: Geometry and Topology Publications, 2002.

UNIV. GRENoble ALPES, CNRS, IF, 38000 GRENoble, FRANCE
 Email address: `jean-baptiste.meilhan@univ-grenoble-alpes.fr`