

TAIL PROCESSES FOR STABLE-REGENERATIVE MODEL

SHUYANG BAI AND YIZAO WANG

ABSTRACT. We investigate a family of discrete-time stationary processes, known as stable-regenerative model, that may exhibit typical behaviors of short-range or long-range dependence, respectively, depending on the parameters. We elaborate the phase transition in terms of the tail processes that characterize local clustering of extremes. In particular, in the sub-critical regime, we compute the candidate extremal index and the extremal index, and they are not the same.

1. INTRODUCTION AND MAIN RESULTS

1.1. The model. The objective of this paper is to study the local behavior of extremes of a family of stationary stochastic processes known as stable-regenerative model that has attracted attentions in the studies of stochastic processes with long-range dependence. In an accompanying paper [3] the macroscopic/global limit of extremes are established in terms of convergence of the random sup-measures in the framework of O'Brien et al. [16], and a phase transition is revealed. Here, we characterize the microscopic/local limit of extremes, in terms of the tail processes as introduced by Basrak and Segers [4].

The family of processes of our interest has a tail parameter $\alpha \in (0, 2)$, a memory parameter $\beta \in (0, 1)$, and a multiplicity parameter $p \in \mathbb{N} = \{1, 2, \dots\}$. The representation is intrinsically related to renewal processes, for which we introduce some notations. Consider a discrete-time renewal process, with the consecutive renewal times denoted by $\tau := \{\tau_0, \tau_1, \dots\}$, and τ_0 is the initial renewal time. We let F denote the \mathbb{N} -valued renewal distribution, that is $F(x) = \mathbb{P}(\tau_i - \tau_{i-1} \leq x), i \in \mathbb{N}, x \geq 0$, with probability mass function $f(n) = \mathbb{P}(\tau_i - \tau_{i-1} = n), n \in \mathbb{N}$. Throughout, we assume

$$(1.1) \quad \overline{F}(x) = 1 - F(x) \sim C_F x^{-\beta} \text{ as } x \rightarrow \infty \quad \text{with} \quad \beta \in (0, 1),$$

which implies an infinite mean, and the following technical assumption

$$(1.2) \quad \sup_{n \in \mathbb{N}} \frac{nf(n)}{\overline{F}(n)} < \infty.$$

By default, a renewal process starts at renewal at time 0, and hence $\tau_0 = 0$. Note that our renewal processes may be *delayed*, that is, τ_0 is not necessarily zero, and may be a random variable in $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, and we shall be specific when this is the case. An important notion is the *stationary delayed distribution* of the renewal process, denoted by π . More precisely, π is supported on \mathbb{N}_0 with

$$\pi(k) \equiv \pi(\{k\}) = \overline{F}(k) = 1 - F(k), \quad k \in \mathbb{N}_0.$$

(For the sake of simplicity, we do not distinguish $\pi(k)$, the mass function at $k \in \mathbb{N}_0$, from $\pi(\{k\})$, the measure evaluated at the set $\{k\}$.) Note that the stationary delayed distribution π is a σ -finite and infinite measure on \mathbb{N} , since the renewal distribution has infinite mean. In words, a delayed renewal process τ with first renewal time following π remains shift-invariant in law. More details are provided in Section 2.

We then provide a series representation of the model. Consider

$$(1.3) \quad \sum_{i=1}^{\infty} \delta_{(y_i, d_i)} \stackrel{d}{=} \text{PPP}((0, \infty) \times \mathbb{N}_0, \alpha x^{-\alpha-1} dx d\pi),$$

2010 *Mathematics Subject Classification.* 60G70; Secondary, 60G52; 60K05.

Key words and phrases. multiple integral, stable regenerative set, regular variation, tail process, extremal index, short-range dependence, long-range dependence, renewal process.

where the right-hand side is understood as the Poisson point process on $(0, \infty) \times \mathbb{N}_0$ with intensity measure $\alpha x^{-\alpha-1} dx d\pi$. In addition, let $\{\tau^{(i, d_i)}\}_{i \in \mathbb{N}}$ denote, given the above Poisson point process, conditionally independent delayed renewal processes, each $\tau^{(i, d_i)}$ with initial renewal time delayed at $\tau_0 = d_i$ and common renewal distribution F .

In this paper, we consider the *stable-regenerative multiple-stable model* given by

$$(1.4) \quad \{X_k\}_{k \in \mathbb{N}_0} \stackrel{d}{=} \left\{ \sum_{0 < i_1 < \dots < i_p} [\varepsilon_{i_1}][y_{i_1}] \mathbf{1}_{\{k \in \bigcap_{r=1}^p \tau^{(i_r, d_{i_r})}\}} \right\}_{k \in \mathbb{N}_0},$$

where $\{\varepsilon_i\}$ are i.i.d. Rademacher random variables independent of the point process in (1.3), $[\varepsilon_i] = \varepsilon_{i_1} \times \dots \times \varepsilon_{i_p}$, and similar notation applies for $[y_i]$. The name ‘stable-regenerative’ comes from the fact each renewal process τ , say non-delayed, scales to a β -stable regenerative random closed sets [6, 12]. The name ‘multiple-stable’ comes from the fact that the series representation above corresponds to a *multiple-stable* process, as it has a representation in terms of *multiple-stable integrals* (see e.g. [13, 15, 21] for multiple-stable integrals, and see [2] for our model in multiple-stable-integral representation; we omit the explanation as such a representation is not used in our proofs).

We call $\{X_k\}_{k \in \mathbb{N}_0}$ in (1.4) a *stable-regenerative model* for the sake of simplicity. The case $p = 1$ corresponds to a family of stable processes with *long-range dependence* [5, 18, 22], which exhibits non-standard asymptotic behaviors in terms of limit theorems for sums and extremes as revealed in [17, 24]. For $p \geq 2$, a functional central limit theorem has been established recently in [2], under the assumption

$$(1.5) \quad \beta_p := p\beta - p + 1 \in (0, 1),$$

with a new family of self-similar multiple-stable processes with stationary increments arising in the limit (see also [1] for a variation of the model that scales to multiple-Gaussian processes). The functional central limit theorem indicates that under the assumption (1.5), the process exhibits behaviors of long-range dependence.

Our motivation is to understand the limit behavior of extremes for all possible values of β_p . Note that with $p = 1$ and necessarily $\beta_p = \beta \in (0, 1)$, there is only the regime corresponding to long-range dependence: an extremal limit theorem in terms of random sup-measures [16] has been established in [24], where the limit random sup-measures exhibit non-trivial dependence structure and in particular their marginal laws may no longer belong to the classical extreme-value distributions when $\beta > 1/2$.

It is expected that when $\beta_p < 0$ the process should no longer exhibit long-range dependence, and this is confirmed by our results. The macroscopic limit is investigated in terms of convergence of random sup-measure in the companion paper [3], and here we examine the microscopic limit of extremes by investigating *tail processes*. See Remark 1.2 for an overview of the phase transition for both types of limit theorems.

1.2. Tail processes. For a general stationary process with regularly varying tails $\{X_k\}_{k \in \mathbb{N}_0}$, tail processes characterize the possible local clustering of extremes. The tail process arises in a conditional limit theorem that may be stated as

$$(1.6) \quad \mathcal{L} \left(\frac{X_0}{x}, \dots, \frac{X_m}{x} \mid |X_0| > x \right) \rightarrow \mathcal{L}(Y_0, \dots, Y_m), \text{ for all } m \in \mathbb{N},$$

as $x \rightarrow \infty$. The left-hand side is understood as the conditional law of $(X_i/x)_{i=0, \dots, m}$ given $|X_0| > x$, and the right-hand side the law of (Y_0, \dots, Y_m) . If the above holds, then $\{Y_i\}_{i \in \mathbb{N}_0}$ is referred to as the *tail process* of $\{X_i\}_{i \in \mathbb{N}_0}$. The idea of tail processes dated back to at least Davis and Hsing [8], Davis and Mikosch [9] (and see references therein for some closely related earlier developments), and here we follow the modern formulation introduced by Basrak and Segers [4]. The tail processes reflect local asymptotic behaviors, and different stochastic models have different tail processes in general. The investigations of tail processes have received reinvigorated interests since [4]. See also Kulik and Soulier [14] for the state of the art on the topic.

There are several equivalent characterizations of the tail process. In particular, equivalently to (1.6), one has

$$(1.7) \quad \mathcal{L} \left(\frac{X_0}{X_0}, \dots, \frac{X_m}{X_0} \mid |X_0| > x \right) \rightarrow (\Theta_0, \dots, \Theta_m), \text{ for all } m \in \mathbb{N},$$

and the process $\Theta = \{\Theta_i\}_{i \in \mathbb{N}_0}$ is referred as the *spectral tail process*. In such a case, one can show that

$$\{Y_i\}_{i \in \mathbb{N}_0} \stackrel{d}{=} \{V_\alpha \Theta_i\}_{i \in \mathbb{N}_0},$$

where V_α is an α -Pareto random variable ($\mathbb{P}(V_\alpha > x) = x^{-\alpha}, x \geq 1$) independent from $\{\Theta_i\}_{i \in \mathbb{N}_0}$. Furthermore, Θ , when it exists, can be uniquely extended to a \mathbb{Z} -indexed stochastic process. The limit theorems (1.6) and (1.7) are equivalent to the fact that the process $\{X_k\}_{k \in \mathbb{N}_0}$ is a regularly varying stochastic process (i.e., all finite-dimensional distributions have multivariate regularly varying tails). It is well known that with $p = 1$ all stable processes are of such a type and hence their tail processes exist. We believe all multiple-stable processes with $p \geq 2$ are of such a type, but are unable to find a reference. Our results imply this fact for our model.

1.3. Main results: a phase transition. We first describe the spectral tail processes that will appear in the limit. Let $\Theta^* \equiv \{\Theta_k^*\}_{k \in \mathbb{N}_0}$ be a $\{0, 1\}$ -valued sequence defined as follows: let $\{\tau^{(i)}\}_{i=1, \dots, p}$ denote i.i.d. copies of a standard (non-delayed) renewal process, with the renewal distribution F as in (1.1), and consider

$$(1.8) \quad \Theta_k^* := \begin{cases} 1, & \text{if } k \in \boldsymbol{\eta}, \\ 0, & \text{otherwise,} \end{cases} \quad k = 0, 1, \dots \quad \text{with} \quad \boldsymbol{\eta} := \bigcap_{r=1}^p \tau^{(r)}.$$

Moreover, let ε a Rademacher random variable independent from Θ^* , we set

$$(1.9) \quad \Theta := \varepsilon \Theta^* = \{\varepsilon \Theta_i^*\}_{i \in \mathbb{N}_0}.$$

We emphasize that the law of the spectral tail process is completely determined by F and p . Note that $\boldsymbol{\eta}$ is a non-delayed renewal process ($0 \in \boldsymbol{\eta}$), and it is possibly terminating. This is the case when the renewal distribution has a mass at infinity, and we write $\boldsymbol{\eta} = \{0, \eta_1, \dots, \eta_k\}$ if the $(k+1)$ -th renewal time is with value infinity. The renewal process $\boldsymbol{\eta}$ is terminating (i.e., $\eta_1 = \infty$) if and only if $\beta_p < 0$, and in this case with probability one, $\Theta_k = 0$ for all k large enough. Some properties of $\boldsymbol{\eta}$ are summarized in Section 2.1.

The main result of the paper is the following.

Theorem 1.1. *For all $m \in \mathbb{N}$,*

$$\mathcal{L} \left(\frac{X_0}{X_0}, \dots, \frac{X_m}{X_0} \mid |X_0| > x \right) \rightarrow \mathcal{L}(\varepsilon \Theta_0^*, \dots, \varepsilon \Theta_m^*),$$

as $x \rightarrow \infty$, with the right-hand side as introduced in (1.9).

The following is a summary of the phase transition for the stable-regenerative model, combining Theorem 1.1 and our result in [3]. The summary depends some immediate consequences on extremal index of the stable-regenerative model, denoted by θ , which are explained in Section 1.4.

Remark 1.2. There are three different regimes regarding the limit of extremes for our model, and a brief summary of the extremal index θ and the random sup-measure in the macroscopic limit in each regime is in Table 1.

regime	extremal index θ (microscopic)	limit random sup-measure (macroscopic)
super-critical, $\beta_p > 0$	$\theta = 0$	with long-range dependence
critical, $\beta_p = 0$	$\theta = 0$	independently scattered
sub-critical, $\beta_p < 0$	$\theta \in (0, 1)$	independently scattered

TABLE 1. Summary of phase transition.

At microscopic level here, Theorem 1.1 reveals that the super-critical ($\beta_p > 0$) and critical ($\beta_p = 0$) regimes have the same type of asymptotic behavior, both with extremal index $\theta = 0$; while at macroscopic level as described in [3], the critical and sub-critical ($\beta_p \leq 0$) regimes have the same type of asymptotic

behavior, with independently scattered α -Fréchet random sup-measure arising in the limit. On the other hand, the limit random sup-measures in the regime $\beta_p > 0$ are of a new type, extending the one characterized in [24]. They again exhibit long-range dependence and, unless $p = 1, \beta < 1/2$, their marginal law goes beyond the family of classical extreme-value distributions due to an aggregation effect (see [25] for an explanation when $p = 1$).

So, the convergence of tail processes reveals a more delicate picture of local behaviors when $\beta_p \leq 0$. At the same time, the tail process at the super-critical regime $\beta_p > 0$ is still of a local nature, as it is unable to recover from the tail processes the random sup-measures that arise in the macroscopic limit obtained in [3]. It is also remarkable that at the critical regime, while the macroscopic limit is independently scattered, reflecting typical behavior of short-range dependence, the tail process has extremal index $\theta = 0$, reflecting the infinite size of the local cluster of extremes.

Remark 1.3. For both the original process $\{X_k\}_{k \in \mathbb{N}_0}$ and the spectral tail process Θ , they can be extended in a canonical way to two-sided ones. For the original process, this is explained in Section 2.3. For Θ , one may simply start independent renewal processes in the backward temporal direction, set their intersection as $\eta' \stackrel{d}{=} -\eta$, and define Θ_{-k} accordingly as (1.9). The details are omitted.

1.4. A notable example regarding extremal index. A widely investigated notion for regularly-varying stochastic process is the *extremal index*. A closely related notion is the *candidate extremal index*. For both, our reference is again Kulik and Soulier [14]. Our model in the sub-critical regime then provides an example so that the two indices are both within $(0, 1)$ and yet are *not the same*.

First, the candidate extremal index, denoted by ϑ , can be read from the local tail process Θ [14, Proposition 5.6.4] as

$$\vartheta = \mathbb{E} \left(\sup_{i \geq 0} |\Theta_i|^\alpha - \sup_{i \geq 1} |\Theta_i|^\alpha \right) = \mathfrak{q}_{F,p},$$

with

$$(1.10) \quad \mathfrak{q}_{F,p} := \mathbb{P}(\eta_1 = \infty) = \left(\sum_{n=0}^{\infty} u(n)^p \right)^{-1} = \lim_{n \rightarrow \infty} \overline{F}_p(n) \in (0, 1).$$

(Recall η in (1.8), and see more details in [3].) Next for the extremal index, denoted by θ , it has been proved in [3] that

$$\theta = D_{\beta,p} \mathfrak{q}_{F,p},$$

with

$$D_{\beta,p} := \sum_{s=q_{\beta,p}}^p \binom{p}{s} (-1)^{p-s} (-\beta_p)^{p-1} \quad \text{with} \quad q_{\beta,p} := \min\{q \in \mathbb{N} : \beta_q < 0\}.$$

Indeed, the extremal index θ , if it exists, can be defined as the unique value from $[0, 1]$ so that the following convergence holds

$$(1.11) \quad \mathbb{P} \left(\frac{1}{b_n} \max_{k=1, \dots, n} X_k \leq x \right) = \exp(-\theta x^{-\alpha}),$$

with b_n is such that $\lim_{n \rightarrow \infty} n\mathbb{P}(X_1 > b_n) = 1$. We have (see (3.6) below)

$$b_n \sim \left(\frac{1}{2} \frac{n \log^{p-1} n}{p!(p-1)!} \right)^{1/\alpha}.$$

Then, the definition (1.11) is implied in particular from the convergence of random sup-measure established in [3], formally

$$(1.12) \quad \frac{1}{b_n} \left\{ \max_{k/n \in G} X_k \right\}_{G \in \mathcal{G}} \Rightarrow (D_{\beta,p} \mathfrak{q}_{\beta,p})^{1/\alpha} \{ \mathcal{M}_\alpha^{\text{is}}(G) \}_{G \in \mathcal{G}},$$

where \mathcal{G} is the collection of all open intervals of $[0, 1]$ and $\mathcal{M}_\alpha^{\text{is}}$ is the *independently scattered* α -Fréchet random sup-measure. Now (1.11) follows by taking $G = [0, 1]$.

We are not aware of any other examples of a regularly varying stochastic process with short-range dependence so that the candidate extremal index and the extremal index are both between 0 and 1 and yet not the same. The proof of (1.11) established in [3] is, however, too sophisticated to see quickly why the two indices are not the same here. Therefore, we present a simpler and more accessible computation for the extremal index directly, for the case $p = 2$ only, in Section 5.

We conclude the introduction with a few remarks.

Remark 1.4. Note that $\theta = D_{\beta,p}\vartheta$, and here we collect a couple of facts on $D_{\beta,p}$:

- (i) for all $\beta \in (0, 1)$ so that $\beta_p < 0$, $D_{\beta,p} \in (0, 1)$,
- (ii) for all $\beta \in (0, 1/2)$, $D_{\beta,p} = 1 - p\beta^{p-1}$.

To see the first, introduce

$$f_p(x) = \frac{1}{(p-1)!} \sum_{s=0}^p \binom{p}{s} (-1)^{p-s} (s-x)_+^{p-1} = \frac{1}{(p-1)!} \sum_{s=0}^p (-1)^{p-s} \binom{p}{s} (x-s)_+^{p-1},$$

which is the probability density function of the so-called Irwin–Hall distribution, the one for the sum of n i.i.d. uniform random variables over $(0, 1)$. Then we can write

$$D_{\beta,p} = (p-1)!(1-\beta)^{p-1} f_p((1-\beta)^{-1}) > 0, \text{ for all } \beta \in (0, 1 - 1/p),$$

or exactly when $\beta_p < 0$. To show that $D_{\beta,p} < 1$, it is equivalent to show that $f_p(x) < ((p-1)!)^{-1} x^{p-1}$ for all $x > 1$. But, recall that $f_p(x)$ is the p -times convolution function of uniform density function over $(0, 1)$ evaluated at x . while $\frac{1}{(p-1)!} x^{p-1}$ is the p -times convolution of the indicator function over $[0, \infty)$ evaluated at x . The desired relation now follows.

To see the second, first recall that for any polynomial function $Q(s)$ with degree at most $p-1$, one has $\sum_{s=0}^p (-1)^{p-s} \binom{p}{s} Q(s) = 0$ since this corresponds to a p -times differencing operation. Then, take $Q(s) = (-\beta_s)^{p-1} = (s(1-\beta) - 1)^{p-1}$ here. For $\beta \in (0, 1/2)$, $q_{\beta,p} = 2$, and it then follows that

$$D_{\beta,p} = \sum_{s=2}^p (-1)^{p-s} \binom{p}{s} (-\beta_s)^{p-1} = -(-1)^p Q(0) - (-1)^{p-1} p Q(1) = 1 - p\beta^{p-1}.$$

Remark 1.5. A natural follow-up problem is to establish a point-process convergence of the extremes of our model. This is well-known to follow, under some technical assumptions, once the convergence of cluster point process is established. Most recent developments on the topic have been presented in Kulik and Soulier [14]. We shall establish the convergence of cluster point process in Section 4, by verifying the so-called *anti-clustering condition*. However, the classical approach to establish eventually the point-process convergence (e.g. [14, Theorem 7.3.1]) requires $\vartheta = \theta$ and hence does not apply here. One has to proceed differently in order to establish the point-process convergence. We leave this problem to a future investigation.

Remark 1.6. In the case $\beta_p = 0$, we have at the critical regime, in place of (1.12),

$$\mathbb{P} \left(\frac{1}{\tilde{b}_n} \max_{k=1, \dots, n} X_k \leq x \right) = \exp \left(-\tilde{\theta} x^{-\alpha} \right) \quad \text{with} \quad \tilde{b}_n = \left(\frac{n(\log \log n)^{p-1}}{\log n} \right)^{1/\alpha},$$

and the value $\tilde{\theta} > 0$ explicitly computed in [3]. This implies that the extremal index θ defined in the common notion via (1.11) (with normalization $b_n \gg \tilde{b}_n$ instead) is zero. One may argue that at the super-critical regime, the extremal index is again $\theta = 0$ in the same way.

The paper is organized as follows. Section 2 provides related background, notably on multiple-stable processes and renewal processes. Section 3 proves Theorem 1.1. Section 4 proves the anti-clustering condition and the convergence of cluster point process when $\beta_p < 0$. Section 5 provides a direct computation of the extremal index with $p = 2$, $\beta \in (0, 1/2)$.

Acknowledgements. The authors are grateful to Gennady Samorodnitsky for suggesting to investigate the tail processes for stable-regenerative multiple-stable processes and for several helpful discussions, and to Rafał Kulik for very detailed explanations regarding tail processes and extremal indices. YW's research was partially supported by Army Research Office, US (W911NF-20-1-0139).

2. PRELIMINARY RESULTS

2.1. Renewal processes with infinite mean. Throughout, our references on renewal processes are Giacomini [12, Appendix A.5] and Bingham et al. [7]. Besides the notations and properties of renewal processes in Section 1, we shall also use the renewal mass function of a standard renewal process (no delay) τ with renewal distribution F :

$$u(k) := \mathbb{P}(k \in \tau), \quad k \in \mathbb{N}_0.$$

It is well known that as $k \rightarrow \infty$,

$$(2.1) \quad u(k) \sim \frac{k^{\beta-1}}{C_F \Gamma(\beta) \Gamma(1-\beta)},$$

implies the assumption (1.1), and furthermore under the assumption (1.2) the two are equivalent (see [11]).

Note that τ is null-recurrent, and hence the stationary delayed distribution for F can be taken as $\pi(k) = \overline{F}(k), k \in \mathbb{N}_0$ (more generally one may set $\pi(k) = C\overline{F}(k)$, and we choose the constant $C = 1$ for simplicity). The meaning of stationary delayed distribution is standard and explained in the Section 2.3, where for the sake of completeness, we present a proof for a system of *two-sided renewal processes* to be shift-invariant.

Next, we recall some properties of the intersected renewal processes $\eta = \bigcap_{r=1}^p \tau^{(r)} = \{0, \eta_1, \eta_2, \dots\}$ as introduced in (1.8). We let F_p denote the cumulative distribution function of η_1 and we shall need the asymptotics of $\overline{F}_p(x) = 1 - F_p(x)$. In summary, we have the following:

$$\overline{F}_p(n) \sim \begin{cases} n^{-\beta_p} \frac{(C_F \Gamma(\beta) \Gamma(1-\beta))^p}{\Gamma(\beta_p) (\Gamma(1-\beta_p))}, & \text{if } \beta_p > 0, \\ \frac{(C_F \Gamma(\beta) \Gamma(1-\beta))^p}{\log n}, & \text{if } \beta_p = 0, \end{cases}$$

and when $\beta_p < 0$, the renewal process is terminating, with the probability $\mathbf{q}_{F,p} = \mathbb{P}(\eta_1 = \infty)$ as given in (1.10). All these follows from standard background on renewal processes [7, 12]. For more details, see [3].

2.2. An equivalent series representation. Here, we explain a different representation that shall be needed for our proofs, for finite-dimensional distributions of the process. We are interested in the joint law of (X_0, \dots, X_m) , for some m fixed. Then, for all those $i \in \mathbb{N}_0$ such that $\tau^{(i, d_i)} \cap \{0, \dots, m\} = \emptyset$, they do not contribute in the series representation. Therefore, by a standard thinning argument of Poisson point processes, it follows that

$$(2.2) \quad \sum_{i=1}^{\infty} \delta_{(\varepsilon_i \Gamma_i^{-1/\alpha}, \tau^{(i, d_i)} \cap \{0, \dots, m\})} \mathbf{1}_{\{\tau^{(i, d_i)} \cap \{0, \dots, m\} \neq \emptyset\}} \stackrel{d}{=} \sum_{i=1}^{\infty} \delta_{(w_m^{1/\alpha} \varepsilon_i \Gamma_i^{-1/\alpha}, R_{m,i})},$$

where on the right-hand side,

$$w_m = \sum_{k=0}^m \pi(k) = \sum_{k=0}^m \overline{F}(k) \sim \frac{C_F}{1-\beta} \cdot m^{1-\beta},$$

the random variables $\{\Gamma_i\}_{i \in \mathbb{N}}$ are consecutive arrival times of a standard Poisson process, $\{\varepsilon_i\}_{i \in \mathbb{N}}$ are i.i.d. Rademacher random variables, $\{R_{m,i}\}_{i \in \mathbb{N}}$ are i.i.d. random closed subsets of $\{0, \dots, m\}$ with the law $R_{m,i} \stackrel{d}{=} R_m$ described below, and all families are independent. Suppose τ^* is a shifted renewal process with the stationary delayed distribution π and renewal distribution F defined on a certain measurable space with respect to certain infinite measure μ^* (since π is an infinite measure). Then, one can introduce a probability measure μ_m on the same measurable space via

$$\frac{d\mu_m}{d\mu^*} = \frac{\mathbf{1}_{\{\tau^* \cap \{0, \dots, m\} \neq \emptyset\}}}{\mu^*(\{\tau^* : \tau^* \cap \{0, \dots, m\} \neq \emptyset\})} = \frac{\mathbf{1}_{\{\tau^* \cap \{0, \dots, m\} \neq \emptyset\}}}{w_m}.$$

Then, the law of R_m is the one induced by τ^* with respect to the probability measure μ_m . Moreover, it is immediately verified that

- (i) $\mathbb{P}(k \in R_m) = 1/w_m, k = 0, \dots, m$ (shift invariance).

(ii) $\mathbb{P}(\min(R_m \cap \{k+1, k+2, \dots\}) \leq k+j \mid k \in R_m) = F(j)$ (Markov/renewal property).

By (2.2), we shall work with the following equivalent representation of (1.4) for finite-dimensional distribution:

$$(2.3) \quad \{X_k\}_{k=0, \dots, m} \stackrel{d}{=} \left\{ w_m^{p/\alpha} \sum_{0 < i_1 < \dots < i_p} \frac{[\varepsilon_i]}{[\Gamma_i]^{1/\alpha}} \mathbf{1}_{\{k \in \cap_{r=1}^p R_{m, i_r}\}} \right\}_{k=0, \dots, m},$$

where on the right-hand side the notations are as in (2.2).

Remark 2.1. Note that the representation in (2.3) is slightly different from the ones used in [2, 3]: we include the time zero here, which is more convenient when studying the tail processes. Such a change does not effect the normalization or the limit object (recall that the original process is stationary in time). Moreover, in [2, 3] we eventually let $m \rightarrow \infty$, while here $m \in \mathbb{N}$ is fixed for all the discussions before Section 5.

2.3. A two-sided representation. We provide another series representation of $\{X_k\}_{k \in \mathbb{Z}}$, based on two-sided renewal processes. We do not need this in the rest of the paper. Such a representation is of its own interest, and helps illuminate the notion of delayed stationary distribution which is now defined in a two-sided manner. The results in this section may have been known in the literature but we were unable to find it.

Recall (1.3) and (1.4). In place of $\tau^{(i, d_i)}$ now we introduce $\tau^{(i, d_i, g_i)}$: this is a two-sided renewal processes to be introduced with the first renewal time to the right of the origin (included) is d_i , the first renewal time to the left of the origin (not included) is $-g_i$ (so $g_i \in \mathbb{N}$).

We first introduce

$$\sum_{i=1}^{\infty} \delta_{(y_i, d_i, g_i)} \stackrel{d}{=} \text{PPP}((0, \infty) \times \mathbb{N}_0 \times \mathbb{N}, \alpha x^{-\alpha-1} dx d\tilde{\pi}),$$

with the measure $\tilde{\pi}$ on $\mathbb{N}_0 \times \mathbb{N}$ determined by the following mass function

$$\tilde{\pi}(d, g) := \pi(d) \cdot \frac{f(d+g)}{\bar{F}(d)} = f(d+g), \quad d \in \mathbb{N}_0, g \in \mathbb{N}.$$

Note that the factor $f(d+g)/\bar{F}(d)$ is the probability mass function at $d+g$, $g \in \mathbb{N}$, of the conditional law of a renewal time with respect to F , given that the renewal time is strictly larger than d .

Now we attach independent renewal processes to each pair of (d_i, g_i) . Let each $\tau^{(i, \rightarrow)}$ be a copy of $\tau = \{\tau_0, \tau_1, \dots\}$, a renewal process starting from the origin (so $\tau_0 = 0$) with renewal distribution F : $\tau^{(i, \rightarrow)} \stackrel{d}{=} \{0, \tau_1, \tau_2, \dots\}$. Similarly, let each $\tau^{(i, \leftarrow)}$ be a time-reversed renewal process, starting from zero: so with τ as before, $\tau^{(i, \leftarrow)} \stackrel{d}{=} \{0, -\tau_1, -\tau_2, \dots\}$. All the renewal processes are assumed independent from everything else. Then, we set

$$(2.4) \quad \tau^{(i, d_i, g_i)} := (d_i + \tau^{(i, \rightarrow)}) \cup (-g_i + \tau^{(i, \leftarrow)}),$$

and $d_i + \tau^{(i, \rightarrow)}$, $-g_i + \tau^{(i, \leftarrow)}$ are understood as random subsets in \mathbb{Z} . More precisely, we present each two-sided renewal processes as the set of renewal times as a random closed subset of \mathbb{Z} , and work with the path space

$$S = \{\mathbf{t} = \{t_i\}_{i \in \mathbb{N}} : t_i \in \mathbb{Z}, \text{ all distinct}\}.$$

The ordering of $\{t_i\}_{i \in \mathbb{N}}$ is irrelevant. The two-sided series representation is then given by

$$(2.5) \quad \{X_k\}_{k \in \mathbb{Z}} \stackrel{d}{=} \left\{ \sum_{\mathbf{i} \in \mathcal{D}_p} [\varepsilon_{\mathbf{i}}] [y_{\mathbf{i}}] \mathbf{1}_{\{k \in \cap_{r=1}^p \tau^{(i_r, d_{i_r}, g_{i_r})}\}} \right\}_{k \in \mathbb{Z}}$$

with $\{\varepsilon_i\}$ as in (1.4) which is independent of everything else, and

$$\mathcal{D}_p = \{(i_1, \dots, i_p) \in \mathbb{N}^p : i_1 < \dots < i_p\}.$$

Lemma 2.2. *The two-sided representation (2.5) represents a stationary process $\{X_k\}_{k \in \mathbb{Z}}$ which restricted to $k \in \mathbb{N}_0$ has the representation (1.4).*

Proof. It is known that the multiple series in (2.5) converges almost surely and unconditionally (e.g., [23, Theorem 1.3 and Remark 1.5]). By construction, (2.5) restricted to $k \in \mathbb{N}_0$ is the same as (1.4). It remains to prove stationarity of the process $\{X_k\}_{k \in \mathbb{Z}}$ (it is well-known to be so when restricted $k \in \mathbb{N}_0$, but we shall prove it again here). This can be checked by the shift-invariance property of the intensity measure of the Poisson point process

$$\sum_{i=1}^{\infty} \delta_{(y_i, \tau^{(i, d_i, g_i)})}.$$

(We omit the discussions regarding $\{\varepsilon_i\}_{i \in \mathbb{N}}$ for the sake of simplicity.) Introduce the following shift operation: Consider the shift operation \mathbf{B} on $\mathbf{t} \in S$: $\mathbf{B}(\mathbf{t}) = \mathbf{t} + 1$. for $\mathbf{t} = \{t_i\}_{i \in \mathbb{N}} \in S$, $\mathbf{B}(\mathbf{t}) = \mathbf{t} + 1 = \{t_i + 1\}_{i \in \mathbb{Z}}$ (so that $d + \tau^{(i, \rightarrow)} = \mathbf{B}^d(\tau^{(i, \rightarrow)})$). Let P_{τ} denote the probability measure on S induced by the distribution of the renewal process τ as above. Then, the Poisson point process above is on the space $(0, \infty) \times S$, with its intensity measure denoted by $\alpha x^{-\alpha-1} dx dQ$, and Q is defined as the push-forward measure with respect to the construction (2.4) from the product measure $\tilde{\pi} \times P_{\tau} \times P_{\tau}$ on $(\mathbb{N}_0 \times \mathbb{N}) \times S \times S$.

Then, the stationarity of the process $\{X_k\}_{k \in \mathbb{Z}}$ follows from the following shift-invariance property for the measure Q on S :

$$(2.6) \quad Q \circ \mathbf{B} = Q \circ \mathbf{B}^{-1} = Q.$$

To prove the above, we start by writing for every $\mathbf{t} \in S$,

$$\mathbf{d}_k(\mathbf{t}) := \min\{t_i : t_i \geq k\} - k, \quad \text{and} \quad \mathbf{g}_k(\mathbf{t}) := k - \max\{t_i : t_i < k\}.$$

In words, $\mathbf{d}_k(\mathbf{t})$ represents the distance between k and the first renewal time to the right of k (included) of \mathbf{t} , and $\mathbf{g}_k(\mathbf{t})$ the distance between k and the first renewal time to the left of k (not included). We write $\{\mathbf{d}_k = d, \mathbf{g}_k = g\} = \{\mathbf{t} \in S : \mathbf{d}_k(\mathbf{t}) = d, \mathbf{g}_k(\mathbf{t}) = g\}$. Our construction tells that

$$(2.7) \quad Q(\mathbf{d}_0 = d, \mathbf{g}_0 = g) = \tilde{\pi}(d, g) = f(d + g) \text{ for all } k \in \mathbb{Z}, d \in \mathbb{N}_0, g \in \mathbb{N}.$$

In view of the renewal property, one can then complete the proof of (2.6) and hence the lemma, by showing that $Q(\mathbf{d}_1 = d, \mathbf{g}_1 = g) = f(d + g)$ for all $d \in \mathbb{N}_0, g \in \mathbb{N}$. First, it is easy to see

$$\{\mathbf{d}_1 = d, \mathbf{g}_1 = g\} = \{\mathbf{d}_0 = d + 1, \mathbf{g}_0 = g - 1\}, \text{ for all } d \geq 0, g \geq 2,$$

and that the desired identity follows from (2.7). It remains to consider $d \geq 0, g = 1$. We have this time, letting \mathbf{d}'_0 denote the second renewal time to the right of the origin (included),

$$\{\mathbf{d}_1 = d, \mathbf{g}_1 = 1\} = \{\mathbf{d}_0 = 0, \mathbf{d}'_0 = d + 1\}.$$

Then by construction, $Q(\mathbf{d}_0 = 0, \mathbf{d}'_0 = d + 1) = Q(\mathbf{d}_0 = 0) \mathbb{P}(\tau_1 = d + 1)$. Then $Q(\mathbf{d}_0 = 0) = \pi(0) = \overline{F}(0) = 1$ and $\mathbb{P}(\tau_1 = d + 1) = f(d + 1)$. This completes the proof. \square

3. CONVERGENCE FOR TAIL PROCESSES

We prove Theorem 1.1. Below C will denote a generic positive constant whose value may change from one expression to another.

In order to establish $(m + 1)$ -dimensional multivariate regular variation, we shall work with the representation (2.3) with m throughout fixed. We introduce some notations. Set

$$(3.1) \quad \begin{aligned} \tilde{\ell}_k(1) &:= \min\{i \in \mathbb{N} : k \in R_{m,i}\}, \\ \tilde{\ell}_k(s) &:= \min\{i > \tilde{\ell}_k(s-1) : k \in R_{m,i}\}, \quad s \geq 2, \quad k = 0, \dots, m, \end{aligned}$$

namely, $\tilde{\ell}_k(1), \tilde{\ell}_k(2), \dots$ are the successive i -indices such that $k \in R_{m,i}$. For $\mathbf{i} = (i_1, \dots, i_p) \in \mathcal{D}_p$, we write $\tilde{\ell}_k(\mathbf{i}) = (\tilde{\ell}_k(i_1), \dots, \tilde{\ell}_k(i_p))$. In this way, we write

$$(3.2) \quad X_k = w_m^{p/\alpha} \sum_{\mathbf{i} \in \mathcal{D}_p} \frac{[\varepsilon_{\tilde{\ell}_k(\mathbf{i})}]}{[\Gamma_{\tilde{\ell}_k(\mathbf{i})}]^{1/\alpha}} \quad \text{and} \quad T_k := w_m^{p/\alpha} \frac{[\varepsilon_{\tilde{\ell}_k((1, \dots, p))}]}{[\Gamma_{\tilde{\ell}_k((1, \dots, p))}]^{1/\alpha}}, \quad k = 0, \dots, m.$$

Here and below the notational convention is to write the product $\varepsilon_{\tilde{\ell}_k(i_1)} \cdots \varepsilon_{\tilde{\ell}_k(i_p)}$ as $[\varepsilon_{\tilde{\ell}_k(\mathbf{i})}]$, and similarly for $[\Gamma_{\tilde{\ell}_k(\mathbf{i})}]$. Note that the indicator functions are dropped in the representation (3.2). Moreover, in order to study the marginal distribution, taking into account of the thinning probability $\mathbb{P}(k \in R_{m,i}) = w_m^{-1}$ we shall work with the following marginal representation for single fixed $k \in \mathbb{N}_0$:

$$(3.3) \quad X_k \stackrel{d}{=} \sum_{\mathbf{i} \in \mathcal{D}_p} \frac{[\varepsilon_{\mathbf{i}}]}{[\Gamma_{\mathbf{i}}]^{1/\alpha}}, \quad T_k \stackrel{d}{=} \frac{[\varepsilon_{(1,\dots,p)}]}{[\Gamma_{(1,\dots,p)}]^{1/\alpha}}.$$

Recall that from [23], we have

$$(3.4) \quad \mathbf{q}_p(x) := \mathbb{P}\left([\Gamma_{1:p}]^{-1} > x\right) \sim \frac{x^{-1} \log^{p-1} x}{p!(p-1)!}.$$

as $x \rightarrow \infty$. Moreover we have the following.

Lemma 3.1. *We have $\mathbb{P}(\max_{(1,\dots,p) \neq \mathbf{i} \in \mathcal{D}_p} [\Gamma_{\mathbf{i}}]^{-1/\alpha} > x) = \mathbb{P}((\Gamma_1 \cdots \Gamma_{p-1} \Gamma_{p+1})^{-1/\alpha} > x)$ and*

$$(3.5) \quad \limsup_{x \rightarrow \infty} \frac{\mathbb{P}\left((\Gamma_1 \cdots \Gamma_{p-1} \Gamma_{p+1})^{-1/\alpha} > x\right)}{\mathbf{q}_{p-1}(x^\alpha)} \leq \mathbb{E}\Gamma_2^{-1} < \infty.$$

Moreover, as $x \rightarrow \infty$,

$$(3.6) \quad \mathbb{P}(|X_k| > x) \sim 2\mathbb{P}(X_k > x) \sim \mathbb{P}(|T_k| > x) = \mathbf{q}_p(x^\alpha) \sim \frac{\alpha^{p-1} x^{-\alpha} \log^{p-1}(x)}{p!(p-1)!},$$

and

$$(3.7) \quad \mathbb{P}(|X_k - T_k| > x) \leq C\mathbf{q}_{p-1}(x^\alpha).$$

In words, T_k is the product of Γ_i for the smallest p i -indices such that $k \in R_{m,i}$, and the tail of $|X_k|$ is determined by this single term alone. Then, the joint representation (3.2) allows us to deal with the joint law of T_0, \dots, T_m . The above can be read from [23], which deals with a more general setup. For the sake of convenience we include a proof here.

Proof of Lemma 3.1. The first claim (3.5) follows from monotonicity. Let $\tilde{\Gamma}_2 \stackrel{d}{=} \Gamma_2$ be a random variable independent from everything else. We then have

$$(\Gamma_1 \cdots \Gamma_{p-1} \Gamma_{p+1})^{-1/\alpha} \stackrel{d}{=} [\Gamma_{1:p-1}]^{-1/\alpha} \cdot \left(\Gamma_{p-1} + \tilde{\Gamma}_2\right)^{-1/\alpha} \leq [\Gamma_{1:p-1}]^{-1/\alpha} \tilde{\Gamma}_2^{-1/\alpha}.$$

The right-hand side is now a product of two independent random variables. The second part now follows from Breiman's lemma (e.g. [20]).

Notice that in view of (3.4), the relation (3.7) implies (3.6), and hence it is left to prove (3.7). Let M be a fixed integer such that $M \geq 2p/\alpha + q$. Note that the conclusion concerns only the marginal distribution, and hence we can work with the simplified representation of X_k in (3.3). By (3.5), any finite sum of terms of the form $[\varepsilon_{\mathbf{i}}][\Gamma_{\mathbf{i}}]^{-1/\alpha}$ with $\mathbf{i} \neq (1, \dots, p)$ will have lighter tails than $[\Gamma_{(1,\dots,p)}]^{-1/\alpha}$. Since the sets $\{\mathbf{i} \in \mathcal{D}_q : i_q \leq M\}$, $q = 0, \dots, p-1$ are finite (when $q = 0$ the set is understood as an empty set), it suffices to show the following: for any fixed $i_1 < \dots < i_q \leq M$ and $q = 0, 1, \dots, p-1$,

$$(3.8) \quad \mathbb{P}\left([\Gamma_{i_{1:q}}]^{-1/\alpha} \left| \sum_{M < i_{q+1} < \dots < i_p} \frac{[\varepsilon_{i_{q+1:p}}]}{[\Gamma_{i_{q+1:p}}]^{1/\alpha}} \right| > x\right) = o(x^{-\alpha} \log^{p-1} x), \quad x \rightarrow \infty,$$

here and below we write $i_{1:q} = (i_1, \dots, i_q)$, $[\Gamma_{i_{1:q}}] = \Gamma_{i_1} \times \dots \times \Gamma_{i_q}$, and similarly for other terms. If $q = 0$, in view of the inequality $\mathbb{E}\left([\Gamma_{i_{1:p}}]^{2/\alpha}\right) \leq c(i_1 \dots i_p)^{2/\alpha}$ which holds by the choice of M (cf. [23, Eq.(3.2)]), as well as the orthogonality induced by $[\varepsilon_{i_{1:p}}]$, one can verify that the second moment of the multiple series in

(3.8) is finite, and hence the tail decay is $O(x^{-2}) = o(x^{-\alpha} \log^{p-1} x)$ as $x \rightarrow \infty$. Suppose $q > 0$ below. Then the probability above is bounded by

$$(3.9) \quad \mathbb{P} \left([\Gamma_{(1, \dots, q)}]^{-1/\alpha} \left| \sum_{M < i_{q+1} < \dots < i_p} \frac{[\varepsilon_{i_{q+1:p}}]}{[\Gamma_{i_{q+1:p}}]^{1/\alpha}} \right| > x \right).$$

Using $[\Gamma_{i_{q+1:p}}]^{-1/\alpha} \leq \prod_{r=q+1}^p (\Gamma_{i_r} - \Gamma_q)^{-1/\alpha}$, conditioning on $\{\Gamma_i\}_{i \in \mathbb{N}}$ and applying the contraction principle for multilinear form of Rademacher random variables [10, Theorem 3.6], the probability in (3.9) is bounded, up to a multiplicative constant, by

$$\mathbb{P} \left([\Gamma_{(1, \dots, q)}]^{-1/\alpha} \left| \sum_{M < i_{q+1} < \dots < i_p} [\varepsilon_{i_{q+1:p}}] \prod_{r=q+1}^p (\Gamma_{i_r} - \Gamma_q)^{-1/\alpha} \right| > x \right).$$

Note that $[\Gamma_{(1, \dots, q)}]^{1/\alpha}$ is independent of the absolute value part above, which can be shown to have a finite second moment similarly as before using the fact $(\Gamma_i - \Gamma_q)_{i > M} \stackrel{d}{=} (\Gamma_i)_{i > M-q}$ with the choice of M . Therefore, by Breiman's lemma again, the probability above is of the same order as $\mathbb{P}([\Gamma_{(1, \dots, q)}]^{-1/\alpha} > x) \sim \mathbf{q}_q(x^\alpha) = o(x^{-\alpha} \log^{p-1} x)$ as $x \rightarrow \infty$ in view of (3.4). \square

Proof of Theorem 1.1. Write

$$T_m^* := w_m^{p/\alpha} \frac{[\varepsilon_{(1, \dots, p)}]}{[\Gamma_{(1, \dots, p)}]^{1/\alpha}},$$

and

$$(3.10) \quad H_k := \mathbf{1}_{\{\tilde{\ell}_k((1, \dots, p)) = (1, \dots, p)\}}, \quad k = 0, \dots, m.$$

Note that $T_k H_k = T_m^* H_k$. The proof consists of establishing the following asymptotic equivalence in law:

$$(3.11) \quad \mathcal{L} \left(\frac{X_0}{X_0}, \dots, \frac{X_m}{X_0} \left| |X_0| > x \right. \right) \sim \mathcal{L} \left(\frac{X_0}{X_0}, \dots, \frac{X_m}{X_0} \left| |T_m^*| > x, H_0 = 1 \right. \right)$$

$$(3.12) \quad \sim \mathcal{L} \left(\frac{X_0}{T_m^*}, \dots, \frac{X_m}{T_m^*} \left| |T_m^*| > x, H_0 = 1 \right. \right)$$

$$(3.13) \quad \sim \mathcal{L} \left(\frac{T_m^* H_0}{T_m^*}, \dots, \frac{T_m^* H_m}{T_m^*} \left| |T_m^*| > x, H_0 = 1 \right. \right)$$

$$(3.14) \quad \sim \mathcal{L}(H_0, \dots, H_m \mid H_0 = 1).$$

Here and below, $\mathcal{L}(\dots \mid \dots)$ denote the corresponding conditional law, and $\mathcal{L}(\dots) \sim \mathcal{L}(\dots)$ means that as $x \rightarrow \infty$, the two sides have same limit law, given that the limit law of one of them exists. We will leave the verification of first step (3.11) to the end. We first verify steps (3.12) and (3.13). Write, for each k ,

$$X_k = T_k + X_k - T_k = T_m^* H_k + T_k(1 - H_k) + X_k - T_k.$$

Observe that $|T_k|(1 - H_k) \leq w_m^{p/\alpha} \max_{(1, \dots, p) \neq i \in \mathcal{D}_p} [\Gamma_i]^{-1/\alpha}$. It then follows from Lemma 3.1 that

$$(3.15) \quad \mathbb{P}(|X_k| > x) \sim \mathbb{P}(|T_m^* H_k| > x) \sim \frac{w_m^p \alpha^{p-1} x^{-\alpha} \log^{p-1} x}{p!(p-1)!} \cdot \frac{1}{w_m^p} = \frac{\alpha^{p-1} x^{-\alpha} \log^{p-1}(x)}{p!(p-1)!},$$

and

$$(3.16) \quad \mathbb{P}(|X_k - T_m^* H_k| > x) = O(x^{-\alpha} \log^{p-2} x)$$

as $x \rightarrow \infty$. Next, we need the following facts: for random variables Y and Z , not necessarily independent, such that $\mathbb{P}(|Z| > \epsilon x) = o(\mathbb{P}(|Y| > x))$ as $x \rightarrow \infty$ for any $\epsilon > 0$, we have as $x \rightarrow \infty$ the convergences in distribution:

$$(3.17) \quad \mathcal{L} \left(\frac{Y}{Y+Z} \left| |Y| > x \right. \right) \rightarrow \mathcal{L}(1) \quad \text{and} \quad \mathcal{L} \left(\frac{Z}{Y} \left| |Y| > x \right. \right) \rightarrow \mathcal{L}(0),$$

where $\mathcal{L}(1)$ and $\mathcal{L}(0)$ denote the laws of constants 1 and 0, respectively. Indeed, the first relation follows from the second one, whereas the second holds since

$$\mathbb{P}(|Z/Y| > \epsilon \mid |Y| > x) \leq \mathbb{P}(|Z| > \epsilon x) / \mathbb{P}(|Y| > x) \rightarrow 0$$

for any $\epsilon > 0$. Now (3.12) follows by writing for each k ,

$$\frac{X_k}{X_0} = \frac{X_k}{T_m^*} \cdot \frac{T_m^*}{T_m^* + X_0 - T_m^*}$$

and then applying the first relation in (3.17) with $Y = T_m^*$, $Z = X_0 - T_m^*$, as well as the relations (3.15) and (3.16). The step (3.13) follows by writing for each k ,

$$\frac{X_k}{T_m^*} = \frac{T_m^* H_k}{T_m^*} + \frac{X_k - T_m^* H_k}{T_m^*},$$

and applying the second relation in (3.17).

The last step (3.14) follows from independence between $(H_k)_{k=0, \dots, m}$ and T_m^* . Comparing the definitions (3.10) (note that $\tilde{\ell}_k((1, \dots, p)) = (1, \dots, p)$ means $k \in R_{m,i}$, $i = 1, \dots, p$) and (1.9), one readily checks that

$$\mathcal{L}(H_0, \dots, H_m \mid H_0 = 1) = \mathcal{L}(\Theta_0^*, \dots, \Theta_m^*).$$

Now we return to verify the first step (3.11). Introduce $B_0(x) := \{|X_0| > x\}$, $B_m^*(x) := \{|T_m^*| > x\}$, and

$$E = \left\{ \left(\frac{X_0}{X_0}, \dots, \frac{X_m}{X_0} \right) \in A \right\},$$

where A is a Borel set in \mathbb{R}^{m+1} whose boundary is not charged by the limit law (3.14). It suffices to prove

$$(3.18) \quad \lim_{x \rightarrow \infty} \mathbb{P}(E \mid |X_0| > x) = \lim_{x \rightarrow \infty} \mathbb{P}(E \mid |T_m^*| > x, H_0 = 1).$$

It is clear that, from the established steps (3.12) to (3.14), the (limit of) right-hand side of (3.18) exists. For the left-hand side, write

$$\mathbb{P}(E \mid |X_0| > x) = \frac{\mathbb{P}(E \cap \{|X_0| > x, H_0 = 1\}) + \mathbb{P}(E \cap \{|X_0| > x, H_0 = 0\})}{\mathbb{P}(|X_0| > x)}.$$

In view of (3.6) and (3.7), we have

$$\mathbb{P}(E \cap \{|X_0| > x, H_0 = 0\}) \leq \mathbb{P}(|X_0| > x, H_0 = 0) = o(\mathbb{P}(|X_0| > x))$$

as $x \rightarrow \infty$. Therefore, with in addition (3.15), we see that $\mathbb{P}(E \mid |X_0| > x)$ has the same limit as

$$(3.19) \quad \frac{\mathbb{P}(E \cap \{|X_0| > x, H_0 = 1\})}{\mathbb{P}(|T_m^*| > x, H_0 = 1)}$$

as $x \rightarrow \infty$. For the numerator, we have the following upper and lower bounds:

$$\begin{aligned} \mathbb{P}(E \cap \{|X_0| > x, H_0 = 1\}) &\geq \mathbb{P}(E \cap \{|T_m^*| > (1 + \epsilon)x, H_0 = 1\}) - \mathbb{P}(|X_0 - T_0| > \epsilon x), \\ \mathbb{P}(E \cap \{|X_0| > x, H_0 = 1\}) &\leq \mathbb{P}(E \cap \{|T_m^*| > (1 - \epsilon)x, H_0 = 1\}) + \mathbb{P}(|X_0 - T_0| > \epsilon x). \end{aligned}$$

The limit of (3.19) with the numerator replaced by the upper and lower bound above, as $x \rightarrow \infty$, can be determined by applying (3.7), (3.15) and the relations (3.12) to (3.14) (with x replaced by $(1 \pm \epsilon)x$). Letting $\epsilon \downarrow 0$, we conclude that (3.19) has the same limit as

$$\frac{\mathbb{P}(E \cap \{|T_m^*| > x, H_0 = 1\})}{\mathbb{P}(|T_m^*| > x) \mathbb{P}(H_0 = 1)} = \mathbb{P}(E \mid |T_m^*| > x, H_0 = 1).$$

This completes the proof of (3.18). \square

4. CONVERGENCE OF CLUSTER POINT PROCESS WHEN $\beta_p < 0$

Assume $\beta_p < 0$ in this section. Based on the convergence of the tail processes we can actually prove the convergence of cluster point process, which is based on the verification of the so-called anti-clustering condition in Lemma 4.2 below. Notice that we can read the information of the candidate extremal index $\vartheta = \mathbf{q}_{F,p}$ from the following result, but not the extremal index $\theta = D_{\beta,p} \mathbf{q}_{F,p}$.

Proposition 4.1. *Assume $\beta_p < 0$. For $a_n \rightarrow \infty, r_n \rightarrow \infty$ and $r_n = o(\log a_n)$, with $M_{r_n} := \max_{k=1, \dots, r_n} |X_k|$, the limit of*

$$\mathcal{L} \left(\sum_{i=1}^{r_n} \delta_{X_i/a_n} \mid M_{r_n} > a_n \right)$$

in the space of $\mathfrak{M}_p(\overline{\mathbb{R}} \setminus \{0\})$ is $G\delta_\varepsilon$, a unit mass at a Rademacher random variable ε multiplied by a geometric random variable G with mean $1/\mathbf{q}_{F,p}$ and independent from ε .

Proof. By Theorem 1.1, [4, Theorem 4.3] and a continuous mapping argument (cf. [4, Remark 4.6]), the limit is a point measure

$$\sum_{i=0}^{\infty} \delta_{\varepsilon \Theta_i^*}$$

restricted to $\mathfrak{M}_p(\overline{\mathbb{R}} \setminus \{0\})$ given that the anti-clustering condition [4, Condition 4.1] holds, which is verified in Lemma 4.2 below. Note that because of the restriction, the point process above is actually

$$\left(\sum_{i=0}^{\infty} \mathbf{1}_{\{\Theta_i=1\}} \right) \delta_\varepsilon$$

(note that $\Theta_0^* = 1$ always), and the summation in the parenthesis is readily checked to be a geometric random variable with the desired law, by examining the definition of Θ^* in (1.8) and the renewal property. \square

Lemma 4.2 (Anti-clustering condition). *Assume $\beta_p < 0$. For $a_n \rightarrow \infty, r_n \rightarrow \infty$ and $r_n = o(\log a_n)$,*

$$\lim_{\ell \rightarrow \infty} \limsup_{n \rightarrow \infty} \mathbb{P} \left(\max_{\ell \leq |k| \leq r_n} |X_k| > a_n \eta \mid |X_0| > a_n \eta \right) = 0, \text{ for all } \eta > 0.$$

Proof. We shall compute, by stationarity and a union bound,

$$(4.1) \quad \left(\max_{\ell \leq |k| \leq r_n} |X_k| > a_n \eta \mid |X_0| > a_n \eta \right) \leq \frac{2}{\mathbb{P}(|X_0| > a_n \eta)} \sum_{k=\ell}^{r_n} \mathbb{P}(|X_0| > a_n \eta, |X_k| > a_n \eta).$$

Unlike in the proof of Theorem 1.1, we only need a series representation for two-dimensional (X_0, X_K) , for each $K \in \mathbb{N}$ fixed. This can be done in the same way as (2.3) is derived, with the only modification that the restriction $\tau^* \cap \{0, \dots, m\} \neq \emptyset$ is now replaced by $\tau^* \cap \{0, K\} \neq \emptyset$. More precisely, let $\{\tilde{R}_{K,i}\}_{i \in \mathbb{N}}$ be i.i.d. copies of \tilde{R}_K , of which the law, denoted by $\tilde{\mu}_K$, is determined by

$$\frac{d\tilde{\mu}_K}{d\mu^*} = \frac{\mathbf{1}_{\{\tau^* \cap \{0, K\} \neq \emptyset\}}}{\mu^*(\{\tau^* : \tau^* \cap \{0, K\} \neq \emptyset\})} = \frac{\mathbf{1}_{\{\tau^* \cap \{0, K\} \neq \emptyset\}}}{\tilde{w}_K},$$

and one can compute by inclusion–exclusion that $\tilde{w}_K = 2 - u(K)$. Then we arrive at

$$\{X_k\}_{k=0, K} \stackrel{d}{=} \left\{ \tilde{w}_K^{p/\alpha} \sum_{i \in \mathcal{D}_p} \frac{[\varepsilon_i]}{[\Gamma_i]^{1/\alpha}} \mathbf{1}_{\{k \in \tilde{R}_{K,i}\}} \right\}_{k=0, K},$$

where as usual $\{\tilde{R}_{K,i}\}_{i \in \mathbb{N}}$ are independent from $\{\varepsilon_i\}_{i \in \mathbb{N}}$ and $\{\Gamma_i\}_{i \in \mathbb{N}}$.

This time, introduce (similar to (3.1)) for $k = 0, K$,

$$\begin{aligned} \hat{\ell}_k(1) &:= \min \left\{ j \in \mathbb{N} : k \in \tilde{R}_{K,j} \right\}, \\ \hat{\ell}_k(s) &:= \min \left\{ j > \hat{\ell}_{K,k}(s-1) : k \in \tilde{R}_{K,j} \right\}, s \geq 2. \end{aligned}$$

We can write

$$(4.2) \quad \widehat{X}_k := \widetilde{w}_K^{p/\alpha} \sum_{\mathbf{i} \in \mathcal{D}_p} \frac{[\varepsilon_{\widehat{\ell}_k(\mathbf{i})}]}{[\Gamma_{\widehat{\ell}_k(\mathbf{i})}]^{1/\alpha}} \quad \text{with} \quad \widehat{T}_k := \widetilde{w}_K^{p/\alpha} \frac{[\varepsilon_{\widehat{\ell}_k((1, \dots, p))}]}{[\Gamma_{\widehat{\ell}_k((1, \dots, p))}]^{1/\alpha}}, \quad k = 0, K.$$

Then, $(\widehat{X}_0, \widehat{X}_K) \stackrel{d}{=} (X_0, X_K)$ and $(\widehat{X}_k, \widehat{T}_k) \stackrel{d}{=} (X_k, T_k)$, $k = 0, K$. Then

$$(4.3) \quad \begin{aligned} \mathbb{P}(|X_0| > a_n \eta, |X_K| > a_n \eta) &\leq \mathbb{P}\left(|\widehat{T}_0| > a_n \eta/2, |\widehat{T}_K| > a_n \eta/2\right) + 2\mathbb{P}\left(|\widehat{X}_0 - \widehat{T}_0| > a_n \eta/2\right) \\ &\leq \mathbb{P}\left(|\widehat{T}_0| > a_n \eta/2, |\widehat{T}_K| > a_n \eta/2\right) + C \mathfrak{q}_{p-1}((a_n \eta/2)^\alpha), \end{aligned}$$

where the last step follows from (3.7). Introduce

$$\widehat{H}_k := \mathbf{1}_{\{\widehat{\ell}_k((1, \dots, p)) = (1, \dots, p)\}}, \quad k = 0, K.$$

Therefore we have, using the representation (4.2) again,

$$(4.4) \quad \begin{aligned} &\mathbb{P}\left(|\widehat{T}_0| > \frac{a_n \eta}{2}, |\widehat{T}_K| > \frac{a_n \eta}{2}\right) \\ &\leq \mathbb{P}\left(\frac{(2 - u(K))^{p/\alpha}}{[\Gamma_{(1, \dots, p)}]^{1/\alpha}} > \frac{a_n \eta}{2}\right) \mathbb{P}\left(\widehat{H}_0 = \widehat{H}_K = 1\right) + \mathbb{P}\left(\frac{(2 - u(K))^{p/\alpha}}{(\Gamma_1 \cdots \Gamma_{p-1} \Gamma_{p+1})^{1/\alpha}} > \frac{a_n \eta}{2}\right) \\ &\leq C \mathfrak{q}_p((a_n \eta)^\alpha / 2^{p+\alpha}) \mathbb{P}\left(\widehat{H}_0 = \widehat{H}_K = 1\right) + C \mathfrak{q}_{p-1}((a_n \eta)^\alpha / 2^{p+\alpha}). \end{aligned}$$

It is straightforward to compute

$$(4.5) \quad \begin{aligned} \mathbb{P}\left(\widehat{H}_0 = \widehat{H}_K = 1\right) &= \mathbb{P}\left(\widehat{H}_0 = 1\right) \mathbb{P}\left(\widehat{H}_K = 1 \mid \widehat{H}_0 = 1\right) \\ &\leq \mathbb{P}\left(\widehat{H}_K = 1 \mid \widehat{H}_0 = 1\right) = u(K)^p. \end{aligned}$$

To sum up, by (4.3), (4.4) and (4.5), the right-hand side of (4.1) is further bounded from above by

$$\frac{C \mathfrak{q}_p((a_n \eta)^\alpha / 2^{p+\alpha})}{\mathfrak{q}_p((a_n \eta)^\alpha)} \sum_{k=\ell}^{\infty} u(k)^p + \frac{C r_n \mathfrak{q}_{p-1}((a_n \eta)^\alpha / 2^{p+\alpha})}{\mathfrak{q}_p((a_n \eta)^\alpha)} \leq C \sum_{k=\ell}^{\infty} u(k)^{-p} + C \frac{r_n}{\log a_n}.$$

Recall $u(k)$ in (2.1). Since $\beta_p < 0$, $\sum_{k=1}^{\infty} u(k)^p < \infty$. We have thus proved the desired result. \square

Remark 4.3. We expect the assumption $r_n = o(\log a_n)$ could be relaxed. According to our analysis in [3] (d_n is used for length of each block therein), there is no constraint on the upper bound of the rate $r_n \rightarrow \infty$ other than $r_n = o(n)$, which is typical. At two places (4.3) and (4.4) above, we expect the bound $\mathfrak{q}_{p-1}(C a_n^\alpha)$ can be improved. We do not pursue such an improvement as the current approach is limited if one is interested in proving a full point-process convergence as explained Remark 1.5.

5. EXTREMAL INDEX IN THE SUB-CRITICAL REGIME, $p = 2$

In this section, we restrict to the sub-critical regime

$$p = 2, \quad \beta \in (0, 1/2),$$

and compute the extremal index directly. Recall the series representation of $\{X_k\}_{k=0, \dots, n}$ in (2.3), and for convenience we repeat here

$$(5.1) \quad \{X_k\}_{k=0, \dots, n} \stackrel{d}{=} \{X_{n,k}\}_{0=1, \dots, n} \equiv \left\{ w_n^{2/\alpha} \sum_{1 \leq i_1 < i_2} \frac{\varepsilon_{i_1} \varepsilon_{i_2}}{\Gamma_{i_1}^{1/\alpha} \Gamma_{i_2}^{1/\alpha}} \mathbf{1}_{\{k \in R_{n,i_1} \cap R_{n,i_2}\}} \right\}_{k=0, \dots, n},$$

and we follow the same notations in earlier sections except that we now have n instead of m and will let $n \rightarrow \infty$. We also emphasize on the dependence on n of the series representation when writing $X_{n,k}$ instead of X_k . Introduce

$$b_n = \left(\frac{1}{4} n \log n \right)^{1/\alpha}.$$

The goal is to establish the following. Note that we work with partial maximum starting from $k = 1$ instead of $k = 0$, only for the sake of convention (consistent with [3]). The same limit theorem below holds with $\max_{k=1, \dots, n}$ replaced by $\max_{k=0, \dots, n}$ by the same proof.

Theorem 5.1. *Assume $\beta \in (0, 1/2)$. As $n \rightarrow \infty$, we have*

$$\frac{1}{b_n} \max_{k=1, \dots, n} X_k \Rightarrow \theta^{1/\alpha} Z_\alpha \quad \text{with} \quad \theta = (1 - 2\beta) \mathfrak{q}_{F,2},$$

where Z_α is a standard α -Fréchet distribution ($\mathbb{P}(Z_\alpha \leq x) = e^{-x^{-\alpha}}$, $x \geq 0$).

The proof below is different from the one presented in [3] and a non-trivial adaption is needed to extend the proof here to $p \geq 3$. We shall proceed with following approximation procedure. Fix throughout a sequence of increasing integers $\{m_n\}_{n \in \mathbb{N}}$ such that

$$(5.2) \quad \frac{w_n^2}{n \log^{2/(2-\alpha)} n} \ll m_n \ll \frac{w_n^2}{n}$$

as $n \rightarrow \infty$, where $a_n \ll b_n$ means $a_n/b_n \rightarrow 0$ as $n \rightarrow \infty$. Introduce the following truncated version of $X_{n,k}$ in (5.1):

$$\tilde{X}_{n,k} := w_n^{2/\alpha} \sum_{1 \leq i_1 < i_2 \leq m_n} \frac{\varepsilon_{i_1} \varepsilon_{i_2}}{\Gamma_{i_1}^{1/\alpha} \Gamma_{i_2}^{1/\alpha}} \mathbf{1}_{\{k \in R_{n,i_1} \cap R_{n,i_2}\}}.$$

Then Theorem 5.1 follows from the following two results.

Proposition 5.2. *As $n \rightarrow \infty$, we have*

$$\frac{1}{b_n} \max_{k=1, \dots, n} \tilde{X}_{n,k} \Rightarrow \theta^{1/\alpha} Z_\alpha,$$

Lemma 5.3. *As $n \rightarrow \infty$, we have*

$$\frac{1}{b_n} \max_{k=1, \dots, n} |X_{n,k} - \tilde{X}_{n,k}| \xrightarrow{\mathbb{P}} 0,$$

where $\xrightarrow{\mathbb{P}}$ stands for convergence in probability.

5.1. Proof of Proposition 5.2. Let $\{U_i\}_{i \in \mathbb{N}}$ be i.i.d. uniform random variables independent of everything else. Introduce

$$V_{i_1, i_2} := \varepsilon_{i_1} \varepsilon_{i_2} (U_{i_1} U_{i_2})^{-1/\alpha}, \quad R_{n, i_1, i_2} := R_{n, i_1} \cap R_{n, i_2}, \quad i_1, i_2 \in \mathbb{N}, \quad i_1 < i_2.$$

Then by a well-known relation between order statistics of uniform distribution and Poisson process, we have

$$\left\{ \tilde{X}_{n,k} \right\}_{k=1, \dots, n} \stackrel{d}{=} \left\{ \left(\frac{w_n}{\Gamma_{m_n+1}} \right)^{2/\alpha} \sum_{1 \leq i_1 < i_2 \leq m_n} V_{i_1, i_2} \mathbf{1}_{\{k \in R_{n, i_1, i_2}\}} \right\}_{k=1, \dots, n}$$

Noting that $\Gamma_{m_n+1} \sim m_n$ as $n \rightarrow \infty$ almost surely, we can instead work with

$$(5.3) \quad \left\{ X_{n,k}^* \right\}_{k=1, \dots, n} = \left\{ \left(\frac{w_n}{m_n} \right)^{2/\alpha} \sum_{1 \leq i_1 < i_2 \leq m_n} V_{i_1, i_2} \mathbf{1}_{\{k \in R_{n, i_1, i_2}\}} \right\}_{k=1, \dots, n}.$$

Lemma 5.4. *We have as $n \rightarrow \infty$,*

$$\rho_n := \mathbb{P}(R_{n,1,2} \neq \emptyset) \sim \mathfrak{q}_{F,2} \frac{n}{w_n^2}.$$

Proof. Introduce

$$\begin{aligned} q_{n,1} &:= \mathbb{P}(R_{n,1,2} \cap \{1, \dots, n\} \neq \emptyset, \max R_{n,1,2} \leq n), \\ q_{n,2} &:= \mathbb{P}(R_{n,1,2} \cap \{1, \dots, n\} \neq \emptyset, \max R_{n,1,2} > n). \end{aligned}$$

Then, $q_{n,1} \leq \rho_n \leq q_{n,1} + q_{n,2}$. For $q_{n,1}$ we can apply the last-renewal decomposition and Markov property:

$$q_{n,1} = \sum_{i=1}^n \mathbb{P}(\max R_{n,1,2} = i \mid i \in R_{n,1,2}) \mathbb{P}(i \in R_{n,1,2}) = \sum_{i=1}^n \mathfrak{q}_{F,2} \frac{1}{w_n^2} = \frac{\mathfrak{q}_{F,2} n}{w_n^2},$$

where we have used the fact that conditioning on $i \in R_{n,1,2}$, the count $\sum_{k=i}^{\infty} \mathbf{1}_{\{k \in R_{n,1,2}\}}$ follows a geometric distribution with mean $\mathfrak{q}_{F,2}^{-1}$. For $q_{n,2}$, write first by a similar decomposition based on the last renewal before time n ,

$$\begin{aligned} q_{n,2} &= \sum_{i=1}^n \mathbb{P}(\max(R_{n,1,2} \cap \{1, \dots, n\}) = i, \max R_{n,1,2} > n) \\ &\leq \sum_{i=1}^n \mathbb{P}(i \in R_{n,1,2}) \mathbb{P}(\max R_{n,1,2} > n \mid i \in R_{n,1,2}) \leq \sum_{i=1}^n \frac{1}{w_n^2} \sum_{j=n-i+1}^{\infty} u(j)^2. \end{aligned}$$

The last step above follows from the renewal property and then the union bound. With $v(i) := \sum_{j=i}^{\infty} u(j)^2 \downarrow 0$ as $i \rightarrow \infty$ due to the fact $u(j) \leq Cj^{\beta-1}$ with $\beta \in (0, 1/2)$, the last displayed expression becomes $w_n^{-2} \sum_{i=1}^n v(i) = nw_n^{-2} \sum_{i=1}^n (v(i)/n) = nw_n^{-2} o(1) = o(q_{n,1})$. So we conclude the whole proof. \square

Introduce the following two counting numbers:

$$(5.4) \quad N_n := \sum_{1 \leq i_1 < i_2 \leq m_n} \mathbf{1}_{\{R_{n,i_1,i_2} \cap \{1, \dots, n\} \neq \emptyset\}},$$

and

$$(5.5) \quad M_n := \sum_{\substack{1 \leq i_1 < i_2 \leq m_n, 1 \leq j_1 < j_2 \leq m_n, \\ \{i_1, i_2\} \neq \{j_1, j_2\}, \{i_1, i_2\} \cap \{j_1, j_2\} \neq \emptyset}} \mathbf{1}_{\{R_{n,i_1,i_2} \cap \{1, \dots, n\} \neq \emptyset, R_{n,j_1,j_2} \cap \{1, \dots, n\} \neq \emptyset\}}.$$

Lemma 5.5. *We have as $n \rightarrow \infty$ that*

$$(5.6) \quad \mathbb{E}N_n \sim \lambda_n := \frac{\mathfrak{q}_{F,2}}{2} \frac{nm_n^2}{w_n^2} \rightarrow \infty, \quad \frac{N_n}{\lambda_n} \xrightarrow{\mathbb{P}} 1$$

and

$$\mathbb{E}M_n \leq C \frac{n^2 m_n^3}{w_n^4} = o(\lambda_n).$$

Proof. For the first part, we have

$$\mathbb{E}N_n \sim \frac{m_n^2 \rho_n}{2} = \frac{\mathfrak{q}_{F,2}}{2} \frac{nm_n^2}{w_n^2},$$

which tends to ∞ due to $m_n \gg w_n^2 / (n \log^{2/(2-\alpha)} n)$, the first part of assumption (5.2). Next, note that

$$\begin{aligned} \rho'_n &:= \mathbb{P}(R_{n,1,2} \cap \{1, \dots, n\} \neq \emptyset, R_{n,1,3} \cap \{1, \dots, n\} \neq \emptyset) \\ &\leq \sum_{k=1}^n \sum_{k'=1}^n \mathbb{P}(k \in R_{n,1,2}, k' \in R_{n,1,3}) \leq \frac{2}{w_n^3} \sum_{k=1}^n \sum_{k'=k}^n u(k' - k) \leq C \frac{n^2}{w_n^4}. \end{aligned}$$

Hence

$$\mathbb{E}M_n \leq C \rho'_n m_n^3 \leq C \frac{n^2 m_n^3}{w_n^4} = o(\lambda_n).$$

where the last relation follows from $m_n \ll w_n^2/n$, the second part of (5.2).

Next, by a decomposition of the double sum over $1 \leq i_1 < i_2 \leq m_n$ and $1 \leq j_1 < j_2 \leq m_n$ according to $|\{i_1, i_2\} \cap \{j_1, j_2\}| = 0, 1, 2$, we have as $n \rightarrow \infty$,

$$\mathbb{E}N_n^2 = \binom{m_n}{2} \binom{m_n - 2}{2} \rho_n^2 + \mathbb{E}M_n + \mathbb{E}N_n = (\mathbb{E}N_n)^2(1 - o(1)) + O(\lambda_n),$$

and hence

$$\text{Var}(N_n) = o((\mathbb{E}N_n)^2),$$

which concludes the convergence in probability in (5.6). \square

Proof of Proposition 5.2. We shall work with $X_{n,k}^*$ in (5.3). The key underlying structure is that its partial maximum can be approximated by a collection of λ_n (see (5.6)) i.i.d. random variables, as summarized in (5.9) below, and this approximation alone explains why the extra factor $D_{2,\beta} = 1 - 2\beta$ shows up in the extremal index compared to the candidate extremal index.

We start by setting the random index set

$$J_n(k) := \{(i_1, i_2) \in \{1, \dots, m_n\}^2 : i_1 < i_2, k \in R_{n,i_1,i_2}\}, \quad k = 1, \dots, n.$$

It follows from (5.6) that with probability tending to 1, we have $J_n(k) \neq \emptyset$ for some $k = 1, \dots, n$. So

$$(5.7) \quad \lim_{n \rightarrow \infty} \mathbb{P} \left(\max_{k=1, \dots, n} X_{n,k}^* = \left(\frac{w_n^2}{m_n^2} \right)^{1/\alpha} \max_{k=1, \dots, n} \left(\sum_{(i_1, i_2) \in J_n(k)} V_{i_1, i_2} \right) \right) = 1,$$

where the sum over $(i_1, i_2) \in J_n(k)$ is understood as $-\infty$ if $J_n(k) = \emptyset$, and similarly for any sum or maximum that appears below whose the index set is empty. Define also

$$I_n(k) := \bigcup_{(i_1, i_2) \in J_n(k)} \{i_1, i_2\}.$$

In fact $|J_n(k)| = \binom{|I_n(k)|}{2}$, and hence $|J_n(k)|$ cannot take arbitrary integer values. But for simplicity of notation we shall still write a consecutive integer range for $|J_n(k)|$ below. Let

$$\mathcal{K}_n := \left\{ k \in \{1, \dots, n\} : |J_n(k)| = 1, \text{ and} \right. \\ \left. \text{either } J_n(k') = J_n(k) \text{ or } I_n(k') \cap I_n(k) = \emptyset \ \forall k' \in \{1, \dots, n\} \setminus \{k\} \right\},$$

and set $\mathcal{K}_n^c := \{1, \dots, n\} \setminus \mathcal{K}_n$. Then by independence,

$$(5.8) \quad \max_{k \in \mathcal{K}_n} \left(\sum_{(i_1, i_2) \in J_n(k)} V_{i_1, i_2} \right) \stackrel{d}{=} \max_{1 \leq \ell \leq \tilde{N}_n} V_\ell \quad \text{with} \quad \tilde{N}_n := |\cup_{k \in \mathcal{K}_n} J_n(k)|,$$

where $\{V_\ell\}_{\ell \in \mathbb{N}}$ are i.i.d. of the same distribution as $V_{1,2}$ and independent of everything else.

Next, observe that that $\tilde{N}_n \leq N_n$ with N_n in (5.4). In addition, any (i_1, i_2) satisfying $R_{n,i_1,i_2} \neq \emptyset$ and $1 \leq i_1 < i_2 \leq m_n$, but not included in $\cup_{k \in \mathcal{K}_n} J_n(k)$, must have been counted at least once by M_n in (5.5). So

$$0 \leq N_n - \tilde{N}_n \leq M_n.$$

This together with Lemma 5.5 implies that $\tilde{N}_n/\lambda_n \xrightarrow{\mathbb{P}} 1$ as $n \rightarrow \infty$ with λ_n in (5.6). Hence it follows from (5.7), (5.8) and the independence between \tilde{N}_n and $\{V_\ell\}_{\ell \in \mathbb{N}}$ that

$$(5.9) \quad \begin{aligned} \lim_{n \rightarrow \infty} \mathbb{P} \left(\frac{1}{b_n} \max_{k \in \mathcal{K}_n} X_{n,k}^* \leq x \right) &= \lim_{n \rightarrow \infty} \mathbb{P} \left(\left(\frac{w_n^2}{b_n^\alpha m_n^2} \right)^{1/\alpha} \max_{1 \leq \ell \leq \tilde{N}_n} V_\ell \leq x \right) \\ &= \lim_{n \rightarrow \infty} \mathbb{P} \left(\left(\frac{w_n^2}{b_n^\alpha m_n^2} \right)^{1/\alpha} \max_{1 \leq \ell \leq \lambda_n} V_\ell \leq x \right), \text{ for all } x > 0. \end{aligned}$$

Now we are working with classical extreme-value limit theorem for $[\lambda_n]$ i.i.d. random variables of regularly-varying tail (e.g., [19, Proposition 1.11]). The normalization, say d_n , for

$$\frac{1}{d_n} \max_{\ell=1, \dots, \lambda_n} V_\ell \Rightarrow Z_\alpha,$$

is well-known to be determined by $\lim_{n \rightarrow \infty} \lambda_n \mathbb{P}(V_1 > d_n) = 1$. It is elementary to verify that

$$(5.10) \quad \mathbb{P}(V_1 < -x) = \mathbb{P}(V_1 > x) = \frac{1}{2} \mathbb{P}(U_1 U_2 < x^{-\alpha}) \sim \frac{\alpha}{2} x^{-\alpha} \log x$$

as $x \rightarrow \infty$. Recall also $\lambda_n \sim (\mathfrak{q}_{F,2}/2) n m_n^2 / w_n^2$ and the requirement (5.2) which implies $\log(\lambda_n) \sim \log(w_n^2/n) \sim (1-2\beta) \log n$. So we have

$$\begin{aligned} d_n &\sim \left(\frac{1}{2} \lambda_n \log \lambda_n \right)^{1/\alpha} \sim \left(\frac{1-2\beta}{2} \lambda_n \log n \right)^{1/\alpha} \\ &\sim \left(\frac{b_n^\alpha m_n^2}{w_n^2} \right)^{1/\alpha} (\mathfrak{q}_{F,2} (1-2\beta))^{1/\alpha} = \left(\frac{b_n^\alpha m_n^2}{w_n^2} \right)^{1/\alpha} \theta^{1/\alpha}. \end{aligned}$$

The above and (5.9) now imply that

$$\frac{1}{b_n} \max_{k \in \mathcal{K}_n} X_{n,k}^* \Rightarrow \theta^{1/\alpha} Z_\alpha$$

as $n \rightarrow \infty$.

To complete the proof, it remains to show that as $n \rightarrow \infty$

$$\frac{1}{b_n} \max_{k \in \mathcal{K}_n^c} X_{n,k}^* \xrightarrow{\mathbb{P}} 0.$$

We fix p^* large enough so that $1 - (p^* + 1)\beta < 0$ (recall $\beta \in (0, 1/2)$). Then

$$\begin{aligned} \mathbb{P}(|J_n(k)| > p^* \text{ for some } k = 1, \dots, n) &\leq \sum_{k=1}^n \binom{m_n}{p^* + 1} w_n^{-p^* - 1} \\ &\leq C \frac{w_n^{p^* + 1}}{n^{p^*}} \leq C n^{1 - (p^* + 1)\beta} \rightarrow 0, \end{aligned}$$

where we have used (5.2) in the second inequality above. Hence with probability tending to 1 as $n \rightarrow \infty$, we have

$$\frac{1}{b_n} \max_{k \in \mathcal{K}_n^c} X_{n,k}^* = \left(\frac{w_n^2}{b_n^\alpha m_n^2} \right)^{1/\alpha} \max_{j=1, \dots, p^*} \max_{k=1, \dots, n} \sum_{(i_1, i_2) \in J_n(k), |J_n(k)|=j} V_{i_1, i_2}.$$

So it suffices to show for fixed $j = 1, \dots, p^*$ as $n \rightarrow \infty$ that

$$(5.11) \quad \left(\frac{w_n^2}{b_n^\alpha m_n^2} \right)^{1/\alpha} \max_{k=1, \dots, n} W_j(k) \xrightarrow{\mathbb{P}} 0 \quad \text{with} \quad W_j(k) = \sum_{(i_1, i_2) \in J_n(k), |J_n(k)|=j} |V_{i_1, i_2}|.$$

Note that $\{W_j(k)\}_{k=1, \dots, n}$ are identically distributed but possibly dependent random variables, and the total number of distinct $W_j(k)$'s, say $K_j(n)$, does not exceed $N_n - \tilde{N}_n \leq M_n$. On the other hand in view of (5.10),

$$(5.12) \quad \mathbb{P}(W_j(1) > x) \leq j \mathbb{P}(|V_{1,2}| > x/j) \leq C x^{-\alpha} \log x$$

for all $x > 0$ and some constant $C > 0$. Suppose $\{\widetilde{W}_j(\ell)\}_{\ell \in \mathbb{N}}$ are i.i.d. copies of $W_j(1)$ and independent of everything else. Then by the tail bound (5.12), the tail estimate (5.10), the extreme-value limit theorem for $|V_\ell|$ and the fact that $M_n/\lambda_n \xrightarrow{\mathbb{P}} 0$ (Lemma 5.5), we have

$$\mathbb{P}\left(\left(\frac{w_n^2}{b_n^\alpha m_n^2} \right)^{1/\alpha} \max_{\ell=1, \dots, K_j(n)} \widetilde{W}_j(\ell) > \epsilon \right) \leq C \mathbb{P}\left(\left(\frac{w_n^2}{b_n^\alpha m_n^2} \right)^{1/\alpha} \max_{\ell=1, \dots, M_n} |V_\ell| > \epsilon \right) \rightarrow 0$$

for any $\epsilon > 0$ as $n \rightarrow \infty$. Then (5.11) follows from [22, Proposition 9.7.3]. \square

5.2. Proof of Lemma 5.3. Lemma 5.3 follows if one establishes that

$$A_n := \frac{w_n^{2/\alpha}}{b_n} \max_{k=1, \dots, n} \sum_{1 \leq i_1 \leq m_n < i_2} \frac{\varepsilon_{i_1} \varepsilon_{i_2}}{\Gamma_{i_1}^{1/\alpha} \Gamma_{i_2}^{1/\alpha}} \mathbf{1}_{\{k \in R_{n, i_1, i_2}\}} \xrightarrow{\mathbb{P}} 0$$

and

$$B_n := \frac{w_n^{2/\alpha}}{b_n} \max_{k=1, \dots, n} \sum_{m_n < i_1 < i_2} \frac{\varepsilon_{i_1} \varepsilon_{i_2}}{\Gamma_{i_1}^{1/\alpha} \Gamma_{i_2}^{1/\alpha}} \mathbf{1}_{\{k \in R_{n, i_1, i_2}\}} \xrightarrow{\mathbb{P}} 0$$

as $n \rightarrow \infty$.

First, choose a fixed number m so that $m > 4/\alpha$. Assume without loss of generality that $m_n > 4/\alpha$. Then write

$$A_n = A_{n,1} + A_{n,2} := \frac{w_n^{2/\alpha}}{b_n} \max_{k=1, \dots, n} \sum_{m \leq i_1 \leq m_n < i_2} \dots + \frac{w_n^{2/\alpha}}{b_n} \max_{k=1, \dots, n} \sum_{1 \leq i_1 < m, i_2 > m_n} \dots.$$

Then by bounding the maximum of non-negative numbers with their sum, the orthogonality induced by $\varepsilon_{i_1} \varepsilon_{i_2}$ and [23, the inequality (3.2)], we have

$$\begin{aligned} \mathbb{E} A_{n,1}^2 &\leq b_n^{-2} w_n^{4/\alpha} \mathbb{E} \sum_{k=1}^n \left(\sum_{m \leq i_1 \leq m_n < i_2} \frac{\varepsilon_{i_1} \varepsilon_{i_2}}{\Gamma_{i_1}^{1/\alpha} \Gamma_{i_2}^{1/\alpha}} \mathbf{1}_{\{k \in R_{n, i_1, i_2}\}} \right)^2 \\ &= b_n^{-2} w_n^{4/\alpha} \sum_{m \leq i_1 \leq m_n < i_2} \mathbb{E} \left(\Gamma_{i_1}^{-2/\alpha} \Gamma_{i_2}^{-2/\alpha} \right) \left(\sum_{k=1}^n \mathbb{P}(k \in R_{n, i_1, i_2}) \right) \\ &\leq C n b_n^{-2} w_n^{4/\alpha-2} \sum_{m \leq i_1 \leq m_n < i_2} i_1^{-2/\alpha} i_2^{-2/\alpha} \leq C n b_n^{-2} w_n^{4/\alpha-2} m_n^{1-2/\alpha} \rightarrow 0, \end{aligned}$$

where we have used the relation $\frac{w_n^2}{n \log^{2/(2-\alpha)} n} \ll m_n$ in the first part of (5.2) in the last step.

For $A_{n,2}$, since i_1 takes finitely many values, it suffices to show for fixed i_1 that

$$b_n^{-2} w_n^{4/\alpha} \mathbb{E} \left| \max_{k=1, \dots, n} \sum_{i_2 > m_n} \Gamma_{i_2}^{-1/\alpha} \varepsilon_{i_2} \mathbf{1}_{\{k \in R_{n, i_1, i_2}\}} \right|^2 \leq C n b_n^{-2} w_n^{4/\alpha-2} m_n^{1-2/\alpha} \rightarrow 0,$$

which follows as above.

For B_n , similarly we have

$$\mathbb{E} B_n^2 \leq C n b_n^{-2} w_n^{4/\alpha-2} \sum_{m < i_1 < i_2} i_1^{-2/\alpha} i_2^{-2/\alpha} \leq C n b_n^{-2} w_n^{4/\alpha-2} m_n^{2-4/\alpha} \rightarrow 0.$$

REFERENCES

- [1] Bai, S. (2021). Limit theorems for conservative flows on multiple stochastic integrals. *Journal of Theoretical Probability*, to appear.
- [2] Bai, S., Owada, T., and Wang, Y. (2020). A functional non-central limit theorem for multiple-stable processes with long-range dependence. *Stochastic Process. Appl.*, 130(9):5768–5801.
- [3] Bai, S. and Wang, Y. (2021). Phase transition of extremes for a family of multiple-stable processes. preprint.
- [4] Basrak, B. and Segers, J. (2009). Regularly varying multivariate time series. *Stochastic Process. Appl.*, 119(4):1055–1080.
- [5] Beran, J., Feng, Y., Ghosh, S., and Kulik, R. (2013). *Long-memory processes*. Springer, Heidelberg. Probabilistic properties and statistical methods.
- [6] Bertoin, J. (1999). Subordinators: examples and applications. In *Lectures on probability theory and statistics (Saint-Flour, 1997)*, volume 1717 of *Lecture Notes in Math.*, pages 1–91. Springer, Berlin.
- [7] Bingham, N. H., Goldie, C. M., and Teugels, J. L. (1987). *Regular variation*, volume 27 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge.

- [8] Davis, R. A. and Hsing, T. (1995). Point process and partial sum convergence for weakly dependent random variables with infinite variance. *Ann. Probab.*, 23(2):879–917.
- [9] Davis, R. A. and Mikosch, T. (1998). The sample autocorrelations of heavy-tailed processes with applications to ARCH. *Ann. Statist.*, 26(5):2049–2080.
- [10] de la Peña, V. H., Montgomery-Smith, S. J., and Szulga, J. (1994). Contraction and decoupling inequalities for multilinear forms and U -statistics. *Ann. Probab.*, 22(4):1745–1765.
- [11] Doney, R. A. (1997). One-sided local large deviation and renewal theorems in the case of infinite mean. *Probab. Theory Related Fields*, 107(4):451–465.
- [12] Giacomin, G. (2007). *Random polymer models*. Imperial College Press, London.
- [13] Krakowiak, W. and Szulga, J. (1986). Random multilinear forms. *Ann. Probab.*, 14(3):955–973.
- [14] Kulik, R. and Soulier, P. (2020). *Heavy-tailed time series*. Springer.
- [15] Kwapien, S. a. and Woyczyński, W. A. (1992). *Random series and stochastic integrals: single and multiple*. Probability and its Applications. Birkhäuser Boston, Inc., Boston, MA.
- [16] O’Brien, G. L., Torfs, P. J. J. F., and Vervaat, W. (1990). Stationary self-similar extremal processes. *Probab. Theory Related Fields*, 87(1):97–119.
- [17] Owada, T. and Samorodnitsky, G. (2015). Functional central limit theorem for heavy tailed stationary infinitely divisible processes generated by conservative flows. *Ann. Probab.*, 43(1):240–285.
- [18] Pipiras, V. and Taqqu, M. S. (2017). *Long-range dependence and self-similarity*, volume 45 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press.
- [19] Resnick, S. I. (1987). *Extreme values, regular variation, and point processes*, volume 4 of *Applied Probability. A Series of the Applied Probability Trust*. Springer-Verlag, New York.
- [20] Resnick, S. I. (2007). *Heavy-tail phenomena*. Springer Series in Operations Research and Financial Engineering. Springer, New York. Probabilistic and statistical modeling.
- [21] Rosiński, J. and Samorodnitsky, G. (1999). Product formula, tails and independence of multiple stable integrals. In *Advances in stochastic inequalities (Atlanta, GA, 1997)*, volume 234 of *Contemp. Math.*, pages 169–194. Amer. Math. Soc., Providence, RI.
- [22] Samorodnitsky, G. (2016). *Stochastic processes and long range dependence*. Springer, Cham, Switzerland.
- [23] Samorodnitsky, G. and Szulga, J. (1989). An asymptotic evaluation of the tail of a multiple symmetric α -stable integral. *Ann. Probab.*, 17(4):1503–1520.
- [24] Samorodnitsky, G. and Wang, Y. (2019). Extremal theory for long range dependent infinitely divisible processes. *Ann. Probab.*, 47(4):2529–2562.
- [25] Wang, Y. (2020). Choquet random sup-measures with aggregations. *Extremes*, to appear. arXiv preprint arXiv:2005.13383.

SHUYANG BAI, DEPARTMENT OF STATISTICS, UNIVERSITY OF GEORGIA, 310 HERTY DRIVE, ATHENS, GA, 30602, USA.
 Email address: bsy9142@uga.edu

YIZAO WANG, DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF CINCINNATI, 2815 COMMONS WAY, CINCINNATI, OH, 45221-0025, USA.
 Email address: yizao.wang@uc.edu