

THE EXTENSION PROPERTY FOR DOMAINS WITH ONE SINGULAR POINT

PEKKA KOSKELA AND ZHENG ZHU

ABSTRACT. An arbitrary outward cuspidal domain is shown to be bi-Lipschitz equivalent to a Lipschitz outward cuspidal domain via a global transformation. This allows us to extend earlier Sobolev extension results on Lipschitz outward cuspidal domains from the work of Maz'ya and Poborchii to arbitrary outward cuspidal domains. We also establish a limit case of extension results on outward cuspidal domains.

1. INTRODUCTION

A domain $\Omega \subset \mathbb{R}^n$ is said to be a Sobolev (p, q) -extension domain for $1 \leq q \leq p \leq \infty$, if there exists a bounded extension operator

$$E : W^{1,p}(\Omega) \rightarrow W^{1,q}(\mathbb{R}^n),$$

such that for every $u \in W^{1,p}(\Omega)$, we have $E(u) \in W^{1,q}(\mathbb{R}^n)$ with

$$\|E(u)\|_{W^{1,q}(\mathbb{R}^n)} \leq C\|u\|_{W^{1,p}(\Omega)}$$

for a constant C independent of u . The smallest constant in the inequality above is denoted by $\|E\|$. In [3, 22], Calderón and Stein proved that if $\Omega \subset \mathbb{R}^n$ is a Lipschitz domain, then there exists a bounded linear extension operator $E : W^{k,p}(\Omega) \rightarrow W^{k,p}(\mathbb{R}^n)$, for all $k \geq 1$ and $1 \leq p \leq \infty$. Here $W^{k,p}(\Omega)$ is the class of those L^p -integrable functions whose weak derivatives up to order k belong to $L^p(\Omega)$. In [9], Jones introduced the notion of (ε, δ) -domains which are generalizations of Lipschitz domains. He proved that, for every (ε, δ) -domain, there exists a bounded linear extension operator $E : W^{k,p}(\Omega) \rightarrow W^{k,p}(\mathbb{R}^n)$, for all $k \geq 1$ and $1 \leq p \leq \infty$. This has motivated the search for geometric characterizations for Sobolev extension domains. A geometric characterization of simply connected planar Sobolev $(2, 2)$ -extension domains was obtained in [23]. By a more recent sequence of results in [10, 11, 12, 19], we understand the geometry of simply connected planar Sobolev (p, p) -extension domains, for all $1 \leq p \leq \infty$. Furthermore, by [20], geometric characterizations of planar simply connected extension domains are also known in the case of homogeneous Sobolev spaces $L^{k,p}(\Omega)$ for $2 < p < \infty$. Here $L^{k,p}(\Omega)$ is the seminormed space of those locally integrable functions whose k -th order distributional partial derivatives belong to $L^p(\Omega)$. However, no characterizations are available in the general setting.

2010 *Mathematics Subject Classification.* 46E35.

The authors have been supported by the Academy of Finland via Centre of Excellence in Analysis and Dynamics Research (Project #323960).

In this paper, we consider Sobolev extension properties for a class of quasiconvex Euclidean domains with only a single singular boundary point. A domain $\Omega \subset \mathbb{R}^n$ is called quasiconvex, if for every $x, y \in \Omega$ there exists a curve $\gamma_{x,y} \subset \Omega$ connecting x, y with

$$\text{length}(\gamma_{x,y}) \leq C|x - y|$$

for a constant C independent of x and y . We study the outward cuspidal domains defined by setting

$$(1.1) \quad \Omega_\psi^n := \{z = (t, x) \in (0, 1] \times \mathbb{R}^{n-1}; |x| < \psi(t)\} \cup \{z = (t, x) \in [1, 2) \times \mathbb{R}^{n-1}; |x| < \psi(1)\},$$

where $\psi: (0, 1] \rightarrow (0, \infty)$ is a left-continuous and increasing function. (Left-continuity is required just to ensure Ω_ψ^n to be open. The term “increasing” is used in the non-strict sense.) If the left-continuous and increasing function ψ is Lipschitz, Ω_ψ^n is called a Lipschitz outward cuspidal domain. This kind of model domains have been widely studied, see Maz’ya and Poborchi’s monograph [17] and references therein. From now on, every left-continuous and increasing function $\psi: (0, 1] \rightarrow (0, \infty)$ will be called a cuspidal function. Our class of domains Ω_ψ^n was introduced in [4]. It was shown in [4] that for an arbitrary cuspidal function ψ , the Sobolev space $W^{1,p}(\Omega_\psi^n)$ coincides with the Hajlasz-Sobolev space $M^{1,p}(\Omega_\psi^n)$ for all $1 < p \leq \infty$. See [6] for the definition of the Hajlasz-Sobolev space $M^{1,p}(\Omega)$. Returning to the problem of geometric characterization for Sobolev extension




FIGURE 1. An outward cuspidal domain Ω_ψ^n

domains, one may ask a natural question:

For which cuspidal functions ψ , the outward cuspidal domain Ω_ψ^n is a Sobolev (p, q) -extension domain for given $1 \leq q \leq p \leq \infty$?

The first observation is that Ω_ψ^n is a Sobolev (∞, ∞) -extension domain, for every cuspidal function ψ . This follows from a result in [7] since Ω_ψ^n is always quasiconvex for an arbitrary cuspidal function ψ .

In [15, 16, 17], for certain Lipschitz cuspidal functions ψ , Maz'ya and Poborchi used integrability conditions on ψ to characterize the Sobolev extension property for Ω_ψ^n . By transferring an outward cuspidal domain onto a Lipschitz outward cuspidal domain via a global bi-Lipschitz transformation, we obtain a more general version of their result.

Theorem 1.2. *Let $\psi : (0, 1] \rightarrow (0, \infty)$ be a cuspidal function such that $\psi|_{(0,1]} > 0$ and the function $\psi(t)/t$ is nondecreasing on $(0, 1]$ with $\lim_{t \rightarrow 0} \psi(t)/t = 0$. We have following statements.*

(1) : If

$$(1.3) \quad \int_0^1 \left(\frac{t^s}{\psi(t)} \right)^{\frac{n}{s-1}} \frac{dt}{t} < \infty,$$

then there exists a bounded linear extension operator E_1 from $W^{1,p}(\Omega_\psi^n)$ to $W^{1,q}(\mathbb{R}^n)$ whenever $\frac{1+(n-1)s}{n} \leq p < \infty$ and $1 \leq q \leq \frac{np}{1+(n-1)s}$, and a bounded linear extension operator E_2 from $W^{1,p}(\Omega_\psi^n)$ to $W^{1,q}(\mathbb{R}^n)$ whenever $\frac{1+(n-1)s}{2+(n-2)s} \leq p < \infty$ and $1 \leq q \leq \frac{(1+(n-1)s)p}{1+(n-1)s+(s-1)p}$.

(2) : If

$$(1.4) \quad \int_0^1 \left(\frac{t^s}{\psi(t)} \right)^{\frac{n}{s-1}} \left| \log \left(\frac{\psi(t)}{t} \right) \right|^{-\alpha} \frac{dt}{t} < \infty$$

with $\alpha = \frac{(n-2)p}{p+1-n}$, then there exists a bounded linear extension operator E_3 from $W^{1,p}(\Omega_\psi^n)$ to $W^{1,q}(\mathbb{R}^n)$ whenever $\frac{(n-1)^2s+(n-1)}{n} \leq p < \infty$ and $1 \leq q \leq n-1$.

Furthermore, under the doubling condition

$$(1.5) \quad \psi(2t) \leq C\psi(t), \text{ for } t \in \left(0, \frac{1}{2}\right),$$

on ψ , the statements converse to (1) and (2) also hold.

Given $1 < s < \infty$, we refer to the domain Ω_ψ^n with $\psi(t) = t^s$ by $\Omega_{t^s}^n$. Let $n \geq 3$. Theorem 1.2 yields that $\Omega_{t^s}^n$ is a Sobolev (p, q) -extension domain, whenever $1 \leq q < n-1$ and $(n-1)q/(n-1-q) \leq p \leq \infty$. Our bi-Lipschitz transformation method allows us to extend this result from the case of t^s to arbitrary cuspidal functions. This result can be regarded as a limit case of Theorem 1.2 and also of theorems by Maz'ya and Poborchi in [17].

Theorem 1.6. *Let $3 \leq n < \infty$ and $\psi : (0, 1] \rightarrow (0, \infty)$ be a cuspidal function. Then the corresponding outward cuspidal domain Ω_ψ^n is a Sobolev (p, q) -extension domain, whenever $1 \leq q < n-1$ and $(n-1)q/(n-1-q) \leq p \leq \infty$.*

The sharpness part of Theorem 1.2 also yields the sharpness of Theorem 1.6.

Proposition 1.7. *For arbitrary $n-1 \leq p < \infty$, there exists $1 < s_1 < \infty$ such that $\Omega_{t^s}^n$ is not a Sobolev $(p, n-1)$ -extension domain when $s > s_1$. For arbitrary $1 \leq q < n-1$ and $q \leq p < (n-1)q/(n-1-q)$, there exists $1 < s_2 < \infty$ such that $\Omega_{t^s}^n$ is not a Sobolev (p, q) -extension domain when $s > s_2$.*

The paper is organized as follows. Section 2 contains definitions and preliminary results. Section 3 contains proofs of all results presented above. Section 4 contains some further discussion and a conjecture.

2. DEFINITIONS AND PRELIMINARIES

In this note, $\Omega \subset \mathbb{R}^n$ is always a bounded domain. C will refer to constants that depend on various parameters and may differ even in a chain of inequalities. The Euclidean distance between points $x, y \in \mathbb{R}^n$ is denoted by $|x - y|$. The open n -dimensional ball of radius r centered at the point x is denoted by $B^n(x, r)$.

Let us give the definition of the Sobolev space $W^{1,p}(\Omega)$.

Definition 2.1. We define the first order Sobolev space $W^{1,p}(\Omega)$, $1 \leq p \leq \infty$, as the set

$$\{ u \in L^p(\Omega); \nabla u \in L^p(\Omega; \mathbb{R}^n) \}.$$

Here $\nabla u = \left(\frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n} \right)$ is the weak (or distributional) gradient of the integrable function u .

The Sobolev space $W^{1,p}(\Omega)$ is equipped with the norm:

$$\|u\|_{W^{1,p}(\Omega)} = \|u\|_{L^p(\Omega)} + \|\nabla u\|_{L^p(\Omega)}$$

for $1 \leq p \leq \infty$, where $\|f\|_{L^p(\Omega)}$ denotes the usual L^p -norm for $p \in [1, \infty]$. Let us give the definition of Sobolev extension domains.

Definition 2.2. Let $1 \leq q \leq p \leq \infty$. A bounded domain $\Omega \subset \mathbb{R}^n$ is said to be a Sobolev (p, q) -extension domain, if there is a bounded extension operator E from $W^{1,p}(\Omega)$ to $W^{1,q}(\mathbb{R}^n)$ such that, for every $u \in W^{1,p}(\Omega)$, there exists a function $E(u) \in W^{1,q}(\mathbb{R}^n)$ with $E(u)|_{\Omega} \equiv u$ and

$$\|E(u)\|_{W^{1,q}(\mathbb{R}^n)} \leq C\|u\|_{W^{1,p}(\Omega)}$$

for a positive constant C independent of u .

An outward cuspidal domain Ω_{ψ}^n has a singular point on the boundary. However, it still has some nice geometric properties. For example, it satisfies the following *segment condition*.

Definition 2.3. We say that a domain $\Omega \subset \mathbb{R}^n$ satisfies the segment condition if every $x \in \partial\Omega$ has a neighborhood U_x and a nonzero vector y_x such that if $z \in \bar{\Omega} \cap U_x$, then $z + ty_x \in \Omega$ for $0 < t < 1$.

The following lemma tells us that Sobolev functions on a domain with the segment condition can be approximated by globally smooth functions. See [2, Theorem 3.22].

Lemma 2.4. *If a domain $\Omega \subset \mathbb{R}^n$ satisfies the segment condition, then the set of restrictions to Ω of functions in $C_o^\infty(\mathbb{R}^n)$ is dense in $W^{1,p}(\Omega)$ for $1 \leq p < \infty$. In short, $C_o^\infty(\mathbb{R}^n) \cap W^{1,p}(\Omega)$ is dense in $W^{1,p}(\Omega)$ for $1 \leq p < \infty$.*

3. PROOFS OF THEOREM 1.2, THEOREM 1.6 AND PROPOSITION 1.7

As mentioned in the introduction, every outward cuspidal domain Ω_{ψ}^n is a Sobolev (∞, ∞) -extension domain. We show that every outward cuspidal domain is globally bi-Lipschitz equivalent to a Lipschitz outward cuspidal domain. For the proof of this result, we first introduce a Lipschitz cuspidal function $\hat{\psi}$ generated by a given cuspidal function ψ .

Lemma 3.1. *Let $\psi : (0, 1] \rightarrow (0, \infty)$ be an arbitrary cuspidal function. Then the following claims hold.*

(1) : *For every $0 < \hat{t} < 1$, there exists a unique pair $(t_{\hat{t}}, r_{\hat{t}})$ with $0 < t_{\hat{t}} < 1$, $\psi(t_{\hat{t}}) \leq r_{\hat{t}} \leq \lim_{s \rightarrow t_{\hat{t}}^+} \psi(s)$ and $t_{\hat{t}} + r_{\hat{t}} = (1 + \psi(1))\hat{t}$.*

(2) : *The function $\hat{\psi} : (0, 1] \rightarrow (0, \infty)$ defined by setting*

$$\hat{\psi}(\hat{t}) = r_{\hat{t}} \text{ for every } \hat{t} \in (0, 1]$$

is a Lipschitz cuspidal function.




FIGURE 2. Bi-Lipschitz transformations

Proof. Let $(t, r) \in (0, 1] \times (0, \psi(1))$ be a pair of positive numbers such that for every $x \in \mathbb{R}^{n-1}$ with $|x| = r$, we have $(t, x) \in \partial\Omega_\psi^n$. Define a function T on $\partial\Omega_\psi^n$ by setting

$$T(t, x) := t + |x|$$

for every $(t, x) \in \partial\Omega_\psi^n$. Since ψ is left-continuous and increasing on $(0, 1]$, we always have $T(t_1, x_1) < T(t_2, x_2)$ for every $(t_1, x_1), (t_2, x_2) \in \partial\Omega_\psi^n$ with $0 < t_1 < t_2 < 1$. By the same reason, we have $T(t, x_1) < T(t, x_2)$ for every $(t, x_1), (t, x_2) \in \partial\Omega_\psi^n$ with $0 < t < 1$ and

$$\psi(t) \leq |x_1| < |x_2| \leq \lim_{s \rightarrow t^+} \psi(s).$$

Hence, for every $0 < \hat{t} < 1$, there exists a unique pair $(t_{\hat{t}}, r_{\hat{t}})$ with $0 < t_{\hat{t}} < 1$, $\psi(t_{\hat{t}}) \leq r_{\hat{t}} \leq \lim_{s \rightarrow t_{\hat{t}}^+} \psi(s)$ and $t_{\hat{t}} + r_{\hat{t}} = (1 + \psi(1))\hat{t}$. Now, let us show $\hat{\psi}$ is a Lipschitz cuspidal function. By the definition, it is easy to see $\hat{\psi}$ is increasing. Hence, it suffices to show that it is Lipschitz. Define a measurable subset $A \subset (0, 1]$ by setting

$$A := \{\hat{t} \in (0, 1] : r_{\hat{t}} \geq t_{\hat{t}}\}.$$

Let $\hat{t}_1, \hat{t}_2 \in (0, 1]$ be arbitrary. According to their locations, we divide the following argument into three cases. First, let us assume $\hat{t}_1, \hat{t}_2 \in A$. Since $\hat{t} \sim r_{\hat{t}}$ for every $\hat{t} \in A$, by the definition, we have

$$(3.2) \quad |\hat{\psi}(\hat{t}_1) - \hat{\psi}(\hat{t}_2)| = |r_{\hat{t}_1} - r_{\hat{t}_2}| \leq C|\hat{t}_1 - \hat{t}_2|.$$

Next, we assume $\hat{t}_1, \hat{t}_2 \in (0, 1] \setminus A$. For every $\hat{t} \in (0, 1] \setminus A$, we have $\hat{t} \sim t_{\hat{t}}$. Hence, the triangle inequality implies

$$(3.3) \quad |\hat{\psi}(\hat{t}_1) - \hat{\psi}(\hat{t}_2)| = |r_{\hat{t}_1} - r_{\hat{t}_2}| \leq C|\hat{t}_1 - \hat{t}_2| + C|t_{\hat{t}_1} - t_{\hat{t}_2}| \leq C|\hat{t}_1 - \hat{t}_2|.$$

Finally, we assume $\hat{t}_1 \in A$ and $\hat{t}_2 \in (0, 1] \setminus A$. Since both $t_{\hat{t}}$ and $r_{\hat{t}}$ are continuous with respect to \hat{t} , there exists $\hat{s} \in (0, 1]$ between \hat{t}_1 and \hat{t}_2 with $t_{\hat{s}} = r_{\hat{s}}$. By (3.2) and (3.3), the triangle inequality implies

$$(3.4) \quad |\hat{\psi}(\hat{t}_1) - \hat{\psi}(\hat{t}_2)| \leq |\hat{\psi}(\hat{t}_1) - \hat{\psi}(\hat{s})| + |\hat{\psi}(\hat{s}) - \hat{\psi}(\hat{t}_2)| \leq C|\hat{t}_1 - \hat{s}| + C|\hat{s} - \hat{t}_2| \leq C|\hat{t}_1 - \hat{t}_2|.$$

By combining (3.2), (3.3) and (3.4), we conclude that $\hat{\psi}$ is Lipschitz. \square

We are ready to prove the following proposition which claims that an arbitrary outward cuspidal domain is globally bi-Lipschitz equivalent to a Lipschitz outward cuspidal domain.

Proposition 3.5. *For a given cuspidal function ψ , let $\hat{\psi}$ be the Lipschitz cuspidal function defined in Lemma 3.1. Then there exists a global bi-Lipschitz transformation $\mathcal{O} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ with $\mathcal{O}(\Omega_{\psi}^n) = \Omega_{\hat{\psi}}^n$.*

Proof. Let $\psi : (0, 1] \rightarrow (0, \infty)$ be a cuspidal function. Let $\hat{\psi} : (0, 1] \rightarrow (0, \infty)$ be the corresponding Lipschitz cuspidal function defined in Lemma 3.1. We extend the definition of ψ to the entire real line \mathbb{R} by setting $\psi(t) = 0$ for every $t \leq 0$. Without loss of generality, we may assume $2\psi(1) < 1$. Otherwise, we consider the cuspidal function $\tilde{\psi}$ defined by setting

$$\tilde{\psi}(t) := \frac{\psi(t)}{2\psi(1)} \text{ for } t \in (-\infty, 1).$$

Obviously, Ω_{ψ}^n and $\Omega_{\tilde{\psi}}^n$ are globally bi-Lipschitz equivalent.

We define

$$U_1 := \{(t, x) \in (-\infty, 1 + \psi(1)] \times \mathbb{R}^{n-1} : |x| \leq 1 + \psi(1) - t\}, U_2 := \Omega_{\tilde{\psi}}^n \setminus U_1,$$

$$U_3 := \{(t, x) \in (-\infty, \infty) \times \mathbb{R}^{n-1} : |x| \geq \max\{\psi(1), 1 + \psi(1) - t\}\},$$

$$U_4 := \{(t, x) \in [2, \infty) \times \mathbb{R}^{n-1} : |x| \leq \psi(1)\}.$$

Obviously, we have $\mathbb{R}^n = \bigcup_{i=1}^4 U_i$. Then we define a global transformation $\mathcal{O} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ by setting

$$(3.6) \quad \mathcal{O}(t, x) := \begin{cases} \left(\left(\frac{1}{1+\psi(1)} \right) (t + |x|), x \right), & (t, x) \in U_1, \\ \left(\left(\frac{t}{1+|x|-\psi(1)} + \frac{2(|x|-\psi(1))}{1+|x|-\psi(1)} \right), x \right), & (t, x) \in U_2, \\ (t + |x| - \psi(1), x), & (t, x) \in U_3, \\ (t, x), & (t, x) \in U_4. \end{cases}$$

The global homeomorphism \mathcal{O} is differentiable almost everywhere and there exists a positive constant $C > 1$ such that for almost every $z \in \mathbb{R}^n$, we have

$$\frac{1}{C} \leq |D\mathcal{O}(z)| \leq C.$$

Hence, $\mathcal{O} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a global bi-Lipschitz transformation. By a simple computation, we obtain

$$\mathcal{O}(U_1 \cap \Omega_\psi^n) = \{(s, y) \in (0, 1] \times \mathbb{R}^{n-1} : |y| < \hat{\psi}(s)\}$$

and

$$\mathcal{O}(U_2) = \{(s, y) \in (1, 2) \times \mathbb{R}^{n-1} : |y| < \hat{\psi}(1)\}.$$

Hence $\mathcal{O}(\Omega_\psi^n) = \Omega_\psi^n$. \square

Let us prove Theorem 1.2.

Proof of Theorem 1.2. Let $\psi : (0, 1] \rightarrow (0, \infty)$ be a cuspidal function such that $\psi(t)/t$ is nondecreasing in $(0, 1]$ with $\lim_{t \rightarrow 0} \psi(t)/t = 0$. Let $\hat{\psi}$ be the corresponding Lipschitz cuspidal function, defined as in Lemma 3.1. Following the notation from Lemma 3.1, for every $\hat{t} \in (0, 1]$, we have

$$t_{\hat{t}} + r_{\hat{t}} = (1 + \psi(1))\hat{t} \text{ and } \hat{\psi}(\hat{t}) = r_{\hat{t}}.$$

Since ψ is left-continuous and increasing, for every $\hat{t} \in \left(0, \frac{1}{1+\psi(1)}\right)$, we have

$$\psi(t_{\hat{t}}) \leq r_{\hat{t}} < \psi((1 + \psi(1))\hat{t}).$$

Hence, we have

$$\lim_{\hat{t} \rightarrow 0} \frac{\hat{\psi}(\hat{t})}{\hat{t}} \leq C \lim_{\hat{t} \rightarrow 0} \frac{\psi(\hat{t})}{\hat{t}} = 0$$

and

$$(3.7) \quad \frac{\hat{\psi}(\hat{t})}{\hat{t}} = \frac{r_{\hat{t}}}{\hat{t}} = \frac{(1 + \psi(1))\hat{t} - t_{\hat{t}}}{\hat{t}} = (1 + \psi(1)) \left(1 - \frac{t_{\hat{t}}}{t_{\hat{t}} + r_{\hat{t}}}\right).$$

By (3.7), to show that $\hat{\psi}(\hat{t})/\hat{t}$ is increasing, it suffices to show that

$$(3.8) \quad \frac{r_{\hat{t}_1}}{t_{\hat{t}_1}} \leq \frac{r_{\hat{t}_2}}{t_{\hat{t}_2}} \text{ for } 0 < \hat{t}_1 \leq \hat{t}_2 < 1.$$

Since ψ is non-decreasing and $\hat{t}_1 \leq \hat{t}_2$, we always have $r_{\hat{t}_1} \leq r_{\hat{t}_2}$. Hence, if $t_{\hat{t}_1} = t_{\hat{t}_2}$, we immediately obtain inequality (3.8). If $t_{\hat{t}_1} < t_{\hat{t}_2}$, the fact that ψ is non-decreasing implies $r_{\hat{t}_1} \leq \psi(t_{\hat{t}_2}) \leq r_{\hat{t}_2}$. Since ψ has at most countably many points of discontinuity, for every $\epsilon > 0$, we can find a point $t_\epsilon \in (t_{\hat{t}_1}, t_{\hat{t}_1} + \epsilon)$ of continuity of ψ with $r_{\hat{t}_1} \leq \psi(t_\epsilon)$. The fact that $\psi(t)/t$ is non-decreasing implies

$$\frac{r_{\hat{t}_1}}{t_{\hat{t}_1}} \leq \lim_{\epsilon \rightarrow 0} \frac{\psi(t_\epsilon)}{t_{\hat{t}_1} + \epsilon} \leq \lim_{\epsilon \rightarrow 0} \frac{\psi(t_\epsilon)}{t_\epsilon} \leq \frac{\psi(t_{\hat{t}_2})}{t_{\hat{t}_2}} \leq \frac{r_{\hat{t}_2}}{t_{\hat{t}_2}}.$$

(1): We show that the assumption that original cuspidal function ψ satisfies the integrability condition (1.3) implies that the corresponding Lipschitz cuspidal function $\hat{\psi}$ also satisfies the integrability condition (1.3). Define a measurable subset $B \subset (0, 1]$ by setting

$$B := \{\hat{t} \in (0, 1] : t_{\hat{t}} \geq r_{\hat{t}}\}.$$

Then, for every $\hat{t} \in B$, we have $\hat{t} \sim t_{\hat{t}}$ and $\hat{\psi}(\hat{t}) = r_{\hat{t}} \geq \psi(t_{\hat{t}})$. Hence, we have

$$(3.9) \quad \int_B \left(\frac{\hat{t}^s}{\hat{\psi}(\hat{t})} \right)^{\frac{n}{s-1}} \frac{d\hat{t}}{\hat{t}} \leq C \int_0^1 \left(\frac{t^s}{\psi(t)} \right)^{\frac{n}{s-1}} \frac{dt}{t} < \infty.$$

If $\hat{t} \in (0, 1] \setminus B$, we have $\hat{\psi}(\hat{t}) = r_{\hat{t}} \sim \hat{t}$. Then we have

$$(3.10) \quad \int_{(0,1] \setminus B} \left(\frac{\hat{t}^s}{\hat{\psi}(\hat{t})} \right)^{\frac{n}{s-1}} \frac{d\hat{t}}{\hat{t}} \leq C \int_0^1 \hat{t}^{n-1} d\hat{t} < \infty.$$

Hence, by combining the last two inequalities, we obtain that $\hat{\psi}$ satisfies the integrability condition that

$$\int_0^1 \left(\frac{\hat{t}^s}{\hat{\psi}(\hat{t})} \right)^{\frac{n}{s-1}} \frac{d\hat{t}}{\hat{t}} < \infty.$$

In conclusion, we have shown that $\hat{\psi}$ satisfies all assumptions of the theorems due to Maz'ya and Poborchi in [17, page 304 and 312]. Hence there exists a bounded linear extension operator $\tilde{E}_1 : W^{1,p}(\Omega_{\hat{\psi}}^n) \rightarrow W^{1,q}(\mathbb{R}^n)$ whenever $\frac{1+(n-1)s}{n} \leq p < \infty$ and $1 \leq q \leq \frac{np}{1+(n-1)s}$. Then, for $\frac{1+(n-1)s}{n} \leq p < \infty$, we define an extension operator E_1 on $W^{1,p}(\Omega_{\psi}^n)$ by setting

$$E_1(u)(x) := \tilde{E}_1(u \circ \mathcal{O}^{-1})(\mathcal{O}(x))$$

for every function $u \in W^{1,p}(\Omega_{\psi}^n)$ and every $x \in \mathbb{R}^n$. By the facts that a bi-Lipschitz transformation preserves first order Sobolev spaces and that $\tilde{E}_1 : W^{1,p}(\Omega_{\hat{\psi}}^n) \rightarrow W^{1,q}(\mathbb{R}^n)$ is a bounded linear extension operator whenever $\frac{1+(n-1)s}{n} \leq p < \infty$ and $1 \leq q \leq \frac{np}{1+(n-1)s}$, we obtain that $E_1 : W^{1,p}(\Omega_{\psi}^n) \rightarrow W^{1,q}(\mathbb{R}^n)$ is also a bounded linear extension operator whenever $\frac{1+(n-1)s}{n} \leq p < \infty$ and $1 \leq q \leq \frac{np}{1+(n-1)s}$. The theorem from [17, page 312] tells us that there exists a bounded linear extension operator $\tilde{E}_2 : W^{1,p}(\Omega_{\hat{\psi}}^n) \rightarrow W^{1,q}(\mathbb{R}^n)$ whenever $\frac{1+(n-1)s}{2+(n-2)s} \leq p < \infty$ and $1 \leq q \leq \frac{(1+(n-1)s)p}{1+(n-1)s+(s-1)p}$. Then, for every $\frac{1+(n-1)s}{2+(n-2)s} \leq p < \infty$, we define an extension operator E_2 on $W^{1,p}(\Omega_{\psi}^n)$ by setting

$$E_2(u)(x) := \tilde{E}_2(u \circ \mathcal{O}^{-1})(\mathcal{O}(x))$$

for every function $u \in W^{1,p}(\Omega_{\psi}^n)$ and $x \in \mathbb{R}^n$. By the same reason as above, $E_2 : W^{1,p}(\Omega_{\psi}^n) \rightarrow W^{1,q}(\mathbb{R}^n)$ is a bounded linear extension operator whenever $\frac{1+(n-1)s}{2+(n-2)s} \leq p < \infty$ and $1 \leq q \leq \frac{(1+(n-1)s)p}{1+(n-1)s+(s-1)p}$.

(2) : By an argument similar with the first case (1), we obtain that if the original cuspidal function ψ satisfies the integrability condition (1.4) then the corresponding Lipschitz cuspidal function $\hat{\psi}$ also satisfies the integrability condition (1.4). Hence, we have shown that $\hat{\psi}$ satisfies all the assumptions of the theorem from [17, page 308]. This theorem tells us that there exists a bounded linear extension

operator $\tilde{E}_3 : W^{1,p}(\Omega_\psi^n) \rightarrow W^{1,q}(\mathbb{R}^n)$ whenever $\frac{(n-1)^2s+(n-1)}{n} \leq p < \infty$ and $1 \leq q \leq n-1$. For every $\frac{(n-1)^2s+(n-1)}{n} \leq p < \infty$, we define an extension operator E_3 on $W^{1,p}(\Omega_\psi^n)$ by setting

$$E_3(u)(x) := \tilde{E}_3(u \circ \mathcal{O}^{-1})(\mathcal{O}(x))$$

for every function $u \in W^{1,p}(\Omega_\psi^n)$ and $x \in \mathbb{R}^n$. By the same reason as above, $E_3 : W^{1,p}(\Omega_\psi^n) \rightarrow W^{1,q}(\mathbb{R}^n)$ is a bounded linear extension operator for every $\frac{(n-1)^2s+(n-1)}{n} \leq p < \infty$ and $1 \leq q \leq n-1$.

Next, let us show the necessity of integrability conditions (1.3) and (1.4). First, let us show that if the doubling condition (1.5) holds for a cuspidal function ψ , it also holds for its corresponding Lipschitz cuspidal function $\hat{\psi}$. Since $\hat{\psi}$ is Lipschitz and increasing, it suffices to show that there exists a positive constant $C > 1$ such that for every $\hat{t} \in \left(0, \frac{1}{2(1+\psi(1))}\right]$, we have

$$(3.11) \quad \hat{\psi}(2\hat{t}) \leq C\hat{\psi}(\hat{t}).$$

Let $\hat{t} \in \left(0, \frac{1}{2(1+\psi(1))}\right]$ be arbitrary. There exists a unique pair $(t_{\hat{t}}, r_{\hat{t}})$ with

$$(3.12) \quad \psi(t_{\hat{t}}) \leq r_{\hat{t}} \leq \lim_{s \rightarrow t_{\hat{t}}^+} \psi(s)$$

and

$$(3.13) \quad t_{\hat{t}} + r_{\hat{t}} = (1 + \psi(1))\hat{t}.$$

Moreover, there exists a unique pair $(t_{2\hat{t}}, r_{2\hat{t}})$ with

$$(3.14) \quad \psi(t_{2\hat{t}}) \leq r_{2\hat{t}} \leq \lim_{s \rightarrow t_{2\hat{t}}^+} \psi(s)$$

and

$$(3.15) \quad t_{2\hat{t}} + r_{2\hat{t}} = (1 + \psi(1))2\hat{t}.$$

By the definition in Lemma 3.1, we have

$$\hat{\psi}(2\hat{t}) = r_{2\hat{t}} \text{ and } \hat{\psi}(\hat{t}) = r_{\hat{t}}.$$

If $r_{2\hat{t}} \leq 2r_{\hat{t}}$, then (3.11) holds with $C = 2$. Hence, we assume $r_{2\hat{t}} > 2r_{\hat{t}}$. By (3.13) and (3.15), we have $t_{2\hat{t}} < 2t_{\hat{t}}$. Since ψ is increasing and satisfies inequality (1.5), by (3.12) and (3.14), we have

$$r_{2\hat{t}} \leq \psi(2t_{\hat{t}}) \leq C\psi(t_{\hat{t}}) \leq Cr_{\hat{t}}.$$

We have showed inequality (3.11) holds for every $\hat{t} \in \left(0, \frac{1}{2(1+\psi(1))}\right]$. The fact that Ω_ψ^n is globally bi-Lipschitz equivalent to Ω_ψ^n implies that Ω_ψ^n and $\Omega_{\hat{\psi}}^n$ have the same Sobolev extension properties. By the results due to Maz'ya and Poborch in [17, pages 304 and 312], if Ω_ψ^n is a Sobolev (p, q) -extension domain with

$$\left(\frac{1 + (n-1)s}{n} \leq p < \infty, 1 \leq q \leq \frac{np}{1 + (n-1)s} \right)$$

or

$$\left(\frac{1 + (n-1)s}{2 + (n-2)s} \leq p < \infty, 1 \leq q \leq \frac{(1 + (n-1)s)p}{1 + (n-1)s + (s-1)p} \right),$$

then $\hat{\psi}$ satisfies the integrability condition

$$\int_0^1 \left(\frac{t^s}{\hat{\psi}(t)} \right)^{\frac{n}{s-1}} \frac{dt}{t} < \infty.$$

By the theorem from [17, page 308], if $\Omega_{\hat{\psi}}^n$ is a Sobolev (p, q) -extension domain with $\frac{(n-1)^2 s + (n-1)}{n} \leq p < \infty$ and $1 \leq q \leq n-1$, then $\hat{\psi}$ satisfies the integrability condition

$$\int_0^1 \left(\frac{t^s}{\hat{\psi}(t)} \right)^{\frac{n}{s-1}} \left| \log \left(\frac{\hat{\psi}(t)}{t} \right) \right|^{-\alpha} \frac{dt}{t} < \infty$$

for $\alpha = \frac{(n-2)p}{p+1-n}$. By the definition of $\hat{\psi}$, for every $t \in (0, 1]$, we always have

$$\psi(t) \geq \hat{\psi} \left(\frac{1}{1 + \psi(1)} t \right).$$

Hence, if the corresponding Lipschitz cuspidal function $\hat{\psi}$ satisfies the integrability conditions (1.3) and (1.4), the original cuspidal function ψ also satisfies them. \square

Let us prove Theorem 1.6.

Proof of Theorem 1.6. As we mentioned, every outward cuspidal domain is a Sobolev (∞, ∞) -extension domain. Hence, it suffices to deal with the case $1 \leq q \leq p < \infty$. We first prove the result for Lipschitz outward cuspidal domains and then extend the result to arbitrary outward cuspidal domains via the global bi-Lipschitz equivalence method established in Proposition 3.5.

Let $1 \leq q < n-1$ and $(n-1)q/(n-1-q) \leq p < \infty$ be fixed. Let ψ be a Lipschitz cuspidal function. We define a cylinder $\widehat{\mathcal{C}}_o$ by setting

$$(3.16) \quad \widehat{\mathcal{C}}_o := \{(t, x) \in [1, 3) \times \mathbb{R}^{n-1} : |x| < 2\psi(1)\}.$$

Then we define two sub-cylinders $\widehat{\mathcal{C}}_o^1$ and $\widehat{\mathcal{C}}_o^2$ of $\widehat{\mathcal{C}}_o$ by setting

$$\widehat{\mathcal{C}}_o^1 := \{(t, x) \in (1, 2) \times \mathbb{R}^{n-1} : |x| < 2\psi(1)\}$$

and

$$\widehat{\mathcal{C}}_o^2 := \{(t, x) \in (2, 3) \times \mathbb{R}^{n-1} : |x| < 2\psi(1)\}.$$

We also define a sub-cylinder \mathcal{C}_o^1 of $\widehat{\mathcal{C}}_o^1$ by setting

$$\mathcal{C}_o^1 := \{(t, x) \in (1, 2) \times \mathbb{R}^{n-1} : |x| < \psi(1)\}.$$

Then $A_{\mathcal{C}_o^1} := \widehat{\mathcal{C}}_o^1 \setminus \overline{\mathcal{C}}_o^1$ is an annular set. We define a reflection $\widetilde{\mathcal{R}}_1 : A_{\mathcal{C}_o^1} \rightarrow \mathcal{C}_o^1$ by setting

$$(3.17) \quad \widetilde{\mathcal{R}}_1(t, x) := \left(t, \left(\frac{3}{2}\psi(1) - \frac{|x|}{2} \right) \frac{x}{|x|} \right).$$

There exists a positive constant C such that for every $(t, x) \in A_{\mathcal{C}_o^1}$, we have

$$(3.18) \quad |D\widetilde{\mathcal{R}}_1(t, x)| \leq C \text{ and } \frac{1}{C} \leq |J_{\widetilde{\mathcal{R}}_1}(t, x)| \leq C.$$

We also define a cut-off function \tilde{L}_1 on the annular set $\overline{A_{C_1}}$ with $\tilde{L}_1 \equiv 0$ on $[1, 2] \times \partial B^{n-1}(0, 2\psi(1))$ and $\tilde{L}_1 \equiv 1$ on $[1, 2] \times \partial B^{n-1}(0, \psi(1))$ by setting

$$(3.19) \quad \tilde{L}_1(t, x) := 2 - \frac{|x|}{\psi(1)}.$$

There exists a positive constant C , such that for every $(t, x) \in A_{C_1}$, we have

$$(3.20) \quad |\nabla \tilde{L}_1(t, x)| \leq C$$

Next, we define a reflection $\widetilde{\mathcal{R}}_2 : \widehat{C}_o^2 \rightarrow \widehat{C}_o^1$ by setting

$$(3.21) \quad \widetilde{\mathcal{R}}_2(t, x) := (t - 2, x).$$

There exists a positive constant C , such that for every $(t, x) \in \widehat{C}_o^2$, we have

$$(3.22) \quad |D\widetilde{\mathcal{R}}_2(t, x)| \leq C \text{ and } \frac{1}{C} \leq |J_{\widetilde{\mathcal{R}}_2}(t, x)| \leq C.$$

Moreover we define a cut-off function \tilde{L}_2 on \widehat{C}_o^2 with $\tilde{L}_2 \equiv 1$ on $\{2\} \times \overline{B^{n-1}(0, 2\psi(1))}$ and $\tilde{L}_2 \equiv 0$ on $\{3\} \times \overline{B^{n-1}(0, 2\psi(1))}$ by setting

$$(3.23) \quad \tilde{L}_2(t, x) := 3 - t.$$

Then, for every $(t, x) \in \widehat{C}_o^2$, we have

$$(3.24) \quad |\nabla \tilde{L}_2(t, x)| \leq 2.$$

Next, we define a double outward cuspidal domain $\widehat{\Omega}_\psi^n$ by setting

$$\widehat{\Omega}_\psi^n := \{(t, x) \in (0, 1] \times \mathbb{R}^{n-1} : |x| < 2\psi(t)\} \cup \widehat{C}_o.$$

We will construct a bounded linear extension operator E from $W^{1,p}(\Omega_\psi^n)$ to $W^{1,q}(\mathbb{R}^n)$ such that for every function $u \in W^{1,p}(\Omega_\psi^n)$, we have $E(u) = 0$ on $\partial\widehat{\Omega}_\psi^n \setminus \{0\}$. We define an annular-type set by setting

$$A_\psi^n := \{(t, x) \in (0, 1] \times \mathbb{R}^{n-1} : \psi(t) < |x| < 2\psi(t)\}.$$

Moreover we define a reflection $\mathcal{R} : A_\psi^n \rightarrow \Omega_\psi^n$ by setting

$$(3.25) \quad \mathcal{R}(z) = \mathcal{R}(t, x) := \left(t, \left(\frac{-|x|}{2} + \frac{3}{2}\psi(t) \right) \frac{x}{|x|} \right).$$

Since ψ is Lipschitz, there exists a positive constant C such that for every $z = (t, x) \in A_\psi^n$, we have

$$(3.26) \quad |D\mathcal{R}(z)| \leq C \text{ and } \frac{1}{C} \leq |J_{\mathcal{R}}(z)| \leq C.$$

We define a cut-off function L on A_ψ^n by setting

$$(3.27) \quad L(t, x) := \frac{-|x|}{\psi(t)} + 2.$$

Since ψ is Lipschitz and

$$\psi(t) < |x| < 2\psi(t)$$

for every point $(t, x) \in A_\psi^n$, there exists a positive constant C such that

$$(3.28) \quad |\nabla L(t, x)| \leq \frac{C}{\psi(t)} \text{ for almost every } (t, x) \in A_\psi^n.$$

By Lemma 2.4, $C_o^\infty(\mathbb{R}^n) \cap W^{1,p}(\Omega_\psi^n)$ is dense in $W^{1,p}(\Omega_\psi^n)$. We first define a bounded linear extension operator from the dense subspace $C_o^\infty(\mathbb{R}^n) \cap W^{1,p}(\Omega_\psi^n)$ to $W^{1,q}(\mathbb{R}^n)$ and then extend it to the full space $W^{1,p}(\Omega_\psi^n)$. Let $u \in C_o^\infty(\mathbb{R}^n) \cap W^{1,p}(\Omega_\psi^n)$ be fixed. We define an extension $E(u)$ on $\widehat{\Omega}_\psi^n$ by setting

$$(3.29) \quad E(u)(z) := \begin{cases} u(z), & z \in \overline{\Omega}_\psi^n, \\ L(z)(u \circ \mathcal{R})(z), & z \in A_\psi^n, \\ \tilde{L}_1(z)(u \circ \tilde{\mathcal{R}}_1)(z), & z \in A_{C_1^o}, \\ \tilde{L}_2(z)(E(u) \circ \tilde{\mathcal{R}}_2)(z), & z \in \widehat{C}_o^2. \end{cases}$$

Then, $E(u)$ is continuous on $\widehat{\Omega}_\psi^n$ with $E(u) \equiv 0$ on $\partial\widehat{\Omega}_\psi^n \setminus \{0\}$. Hence, we can simply extend $E(u)$ to be zero outside the domain $\widehat{\Omega}_\psi^n$. First, let us estimate the L^q -norm of $E(u)$. By the definition of $E(u)$, the Hölder inequality implies

$$(3.30) \quad \left(\int_{\Omega_\psi^n} |E(u)(z)|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p dz \right)^{\frac{1}{p}}.$$

By (3.22) and the fact that $0 \leq \tilde{L}_2 \leq 1$ on \widehat{C}_o^2 , the change of variables formula implies

$$(3.31) \quad \int_{\widehat{C}_o^2} |E(u)(z)|^q dz \leq C \int_{\widehat{C}_o^2} |(E(u) \circ \tilde{\mathcal{R}}_2)(z)|^q dz \leq C \int_{\widehat{C}_o^1} |E(u)(z)|^q dz.$$

By (3.18) and the fact that $0 \leq \tilde{L}_1 \leq 1$ on $A_{C_1^o}$, the change of variables formula and the Hölder inequality imply

$$(3.32) \quad \left(\int_{\widehat{C}_o^1} |E(u)(z)|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{C_1^o} |E(u)(z)|^q dz \right)^{\frac{1}{q}} \\ + C \left(\int_{A_{C_1^o}} |E(u)(z)|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p dz \right)^{\frac{1}{p}}.$$

By combining (3.31) and (3.32), we obtain

$$(3.33) \quad \left(\int_{\widehat{C}_o} |E(u)(z)|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p dz \right)^{\frac{1}{p}}.$$

By (3.26) and the fact that $0 \leq L(z) \leq 1$ on A_ψ^n , the change of variables formula and the Hölder inequality imply

$$(3.34) \quad \left(\int_{A_\psi^n} |L(z)(u \circ \mathcal{R})(z)|^q dz \right)^{\frac{1}{q}} \leq \left(\int_{\Omega_\psi^n} |u(z)|^p dz \right)^{\frac{1}{p}}.$$

Combining inequalities (3.30), (3.33) and (3.34), we obtain

$$(3.35) \quad \left(\int_{\widehat{\Omega}_\psi^n} |E(u)(z)|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p dz \right)^{\frac{1}{p}}.$$

Now, let us estimate the L^q -norm of $|\nabla E(u)|$. First, the Hölder inequality implies

$$(3.36) \quad \left(\int_{\Omega_\psi^n} |\nabla E(u)(z)|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |\nabla u(z)|^p dz \right)^{\frac{1}{p}}.$$

The chain rule implies that for almost every $z \in A_{C_1^1}$, we have

$$(3.37) \quad |\nabla E(u)(z)| \leq \left| \nabla \tilde{L}_1(z)(u \circ \tilde{\mathcal{R}}_1)(z) \right| + \left| \tilde{L}_1(z) \nabla (u \circ \tilde{\mathcal{R}}_1)(z) \right|.$$

By (3.18) and (3.20), the change of variables formula and the Hölder inequality imply

$$(3.38) \quad \left(\int_{A_{C_1^1}} \left| \nabla \tilde{L}_1(z)(u \circ \tilde{\mathcal{R}}_1)(z) \right|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p dz \right)^{\frac{1}{p}}.$$

By (3.18) and the fact that $0 \leq \tilde{L}_1(z) \leq 1$ for every $z \in A_{C_1^1}$, the change of variables formula and the Hölder inequality imply

$$(3.39) \quad \left(\int_{A_{C_1^1}} \left| \tilde{L}_1(z) \nabla (u \circ \tilde{\mathcal{R}}_1)(z) \right|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |\nabla u(z)|^p dz \right)^{\frac{1}{p}}.$$

By combining (3.37), (3.38) and (3.39), we obtain

$$(3.40) \quad \left(\int_{A_{C_1^1}} |\nabla E(u)(z)|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p + |\nabla u(z)|^p dz \right)^{\frac{1}{p}}.$$

The chain rule implies that for almost every $z \in \widehat{C}_o^2$, we have

$$(3.41) \quad |\nabla E(u)(z)| \leq \left| \nabla \tilde{L}_2(z) \left(E(u) \circ \tilde{\mathcal{R}}_2 \right) (z) \right| + \left| \tilde{L}_2(z) \nabla \left(E(u) \circ \tilde{\mathcal{R}}_2 \right) (z) \right|.$$

By (3.22), (3.24) and (3.32), the change of variables formula and the Hölder inequality imply

$$(3.42) \quad \left(\int_{\widehat{\mathcal{C}}_o^2} \left| \nabla \tilde{L}_2(z) \left(E(u) \circ \tilde{\mathcal{R}}_2 \right) (z) \right|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\widehat{\mathcal{C}}_o^1} |E(u)(z)|^q dz \right)^{\frac{1}{q}} \\ \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p dz \right)^{\frac{1}{p}}.$$

By (3.22) and the fact that $0 \leq \tilde{L}_2(z) \leq 1$ for almost every $z \in \widehat{\mathcal{C}}_o^2$, the change of variables formula, (3.36) and (3.40) imply

$$(3.43) \quad \left(\int_{\widehat{\mathcal{C}}_o^2} \left| \tilde{L}_2(z) \nabla \left(E(u) \circ \tilde{\mathcal{R}}_2 \right) (z) \right|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\widehat{\mathcal{C}}_o^1} |\nabla E(u)(z)|^q dz \right)^{\frac{1}{q}} \\ \leq C \left(\int_{A_{\mathcal{C}_o^1}} |\nabla E(u)(z)|^q dz \right)^{\frac{1}{q}} + \left(\int_{\mathcal{C}_o^1} |\nabla E(u)(z)|^q dz \right)^{\frac{1}{q}} \\ \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p + |\nabla u(z)|^p dz \right)^{\frac{1}{p}}.$$

Hence, by combining (3.41), (3.42) and (3.43), we obtain

$$(3.44) \quad \left(\int_{\widehat{\mathcal{C}}_o^2} |\nabla E(u)(z)|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p + |\nabla u(z)|^p dz \right)^{\frac{1}{p}}.$$

Since $\mathcal{C}_o^1 \subset \Omega_\psi^n$, by combining (3.36), (3.40) and (3.44), we obtain

$$(3.45) \quad \left(\int_{\widehat{\mathcal{C}}_o} |\nabla E(u)(z)|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p + |\nabla u(z)|^p dz \right)^{\frac{1}{p}}.$$

The chain rule implies that for almost every $z \in A_\psi^n$, we have

$$(3.46) \quad |\nabla E(u)(z)| \leq |\nabla L(z)(u \circ \mathcal{R})(z)| + |L(z) \nabla(u \circ \mathcal{R})(z)|.$$

By (3.26) and the fact $0 \leq L(z) \leq 1$ on A_ψ^n , the change of variables formula and the Hölder inequality yield

$$(3.47) \quad \left(\int_{A_\psi^n} |L(z) \nabla(u \circ \mathcal{R})(z)|^q dz \right)^{\frac{1}{q}} \leq \left(\int_{\Omega_\psi^n} |\nabla u(z)|^p dz \right)^{\frac{1}{p}}.$$

The Hölder inequality implies

$$(3.48) \quad \left(\int_{A_\psi^n} |\nabla L(z)(u \circ \mathcal{R})(z)|^q dz \right)^{\frac{1}{q}} \leq \left(\int_{A_\psi^n} |\nabla L(z)|^{\frac{pq}{p-q}} dz \right)^{\frac{p-q}{pq}} \cdot \left(\int_{A_\psi^n} |(u \circ \mathcal{R})(z)|^p dz \right)^{\frac{1}{p}}.$$

By (3.26), the change of variables formula implies

$$(3.49) \quad \int_{A_\psi^n} |(u \circ \mathcal{R})(z)|^p dz \leq C \int_{\Omega_\psi^n} |u(z)|^p dz.$$

By (3.28) and the fact that ψ is Lipschitz, we have

$$(3.50) \quad \int_{A_\psi^n} |\nabla L(z)|^{\frac{pq}{p-q}} \leq C \int_0^1 \psi(t)^{n-1-\frac{pq}{p-q}} dt < C \int_0^1 t^{n-1-\frac{pq}{p-q}} dt < \infty$$

whenever $1 \leq q < n-1$ and $(n-1)q/(n-1-q) \leq p < \infty$. By combining inequalities (3.48), (3.49) and (3.50), we obtain

$$(3.51) \quad \left(\int_{A_\psi^n} |\nabla L(z)(u \circ \mathcal{R})(z)|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p dz \right)^{\frac{1}{p}}.$$

By combining (3.46), (3.47) and (3.51), we obtain

$$(3.52) \quad \left(\int_{A_\psi^n} |\nabla E(u)(z)|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p + |\nabla u(z)|^p dz \right)^{\frac{1}{p}}.$$

By combining inequalities (3.36), (3.44) and (3.52), we obtain

$$(3.53) \quad \left(\int_{\widehat{\Omega}_\psi^n} |\nabla E(u)(z)|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p + |\nabla u(z)|^p dz \right)^{\frac{1}{p}}.$$

Finally, by combining (3.35) and (3.53), we obtain the desired norm inequality

$$(3.54) \quad \left(\int_{\widehat{\Omega}_\psi^n} |E(u)(z)|^q + |\nabla E(u)(z)|^q dz \right)^{\frac{1}{q}} \leq C \left(\int_{\Omega_\psi^n} |u(z)|^p + |\nabla u(z)|^p dz \right)^{\frac{1}{p}}.$$

Hence, the extension operator E defined in (3.29) is a bounded linear extension operator from the dense subspace $C_o^\infty(\mathbb{R}^n) \cap W^{1,p}(\Omega_\psi^n)$ to $W^{1,q}(\mathbb{R}^n)$ whenever $1 \leq q < n-1$ and $(n-1)q/(n-1-q) \leq p < \infty$. For an arbitrary $u \in W^{1,p}(\Omega_\psi^n)$, there exists a Cauchy sequence $\{u_m\}_{m=1}^\infty \subset C_o^\infty(\mathbb{R}^n) \cap W^{1,p}(\Omega_\psi^n)$ which converges to u with respect to the $W^{1,p}$ -norm. Since E is a bounded linear extension operator from $C_o^\infty(\mathbb{R}^n) \cap W^{1,p}(\Omega_\psi^n)$ to $W^{1,q}(\mathbb{R}^n)$, there exists a subsequence of $\{u_m\}$ which converges to u almost everywhere on Ω_ψ^n and $\{E(u_m)\}$ is also a Cauchy sequence in $W^{1,q}(\mathbb{R}^n)$ which converges to some function $v \in W^{1,q}(\mathbb{R}^n)$. To simplify the notation, we still denote this subsequence by $\{u_m\}$. Then, we have $v|_{\Omega_\psi^n}(z) = u(z)$ for almost every $z \in \Omega_\psi^n$ and

$$\|v\|_{W^{1,q}(\mathbb{R}^n)} \leq \liminf_{m \rightarrow \infty} \|E(u_m)\|_{W^{1,q}(\mathbb{R}^n)} \leq C \liminf_{m \rightarrow \infty} \|u_m\|_{W^{1,p}(\Omega_\psi^n)} \leq C \|u\|_{W^{1,p}(\Omega_\psi^n)}.$$

Furthermore, by picking an extra subsequence if necessary, it follows that $v(z) = E(u)(z)$ for almost every $z \in \mathbb{R}^n$. Hence, E is a bounded linear extension operator from $W^{1,p}(\Omega_\psi^n)$ to $W^{1,q}(\mathbb{R}^n)$, whenever $1 \leq q < n-1$ and $(n-1)q/(n-1-q) \leq p < \infty$. In conclusion, we have proved that a Lipschitz outward cuspidal domain Ω_ψ^n is a Sobolev (p, q) -extension domain, whenever $1 \leq q < n-1$ and $(n-1)q/(n-1-q) \leq p < \infty$.

Next, we extend the result to an arbitrary outward cuspidal domain by using global bi-Lipschitz transformations. Let $\psi : (0, 1] \rightarrow (0, \infty)$ be an arbitrary cuspidal domain. By Proposition 3.5, there exists a Lipschitz cuspidal function $\hat{\psi}$ and a global bi-Lipschitz homeomorphism $\mathcal{O} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ with $\mathcal{O}(\Omega_\psi^n) = \Omega_{\hat{\psi}}^n$. Fix $1 \leq q < n-1$ and $(n-1)q/(n-1-q) \leq p < \infty$. Let $u \in W^{1,p}(\Omega_\psi^n)$ be arbitrary. Since \mathcal{O} is bi-Lipschitz with $\mathcal{O}(\Omega_\psi^n) = \Omega_{\hat{\psi}}^n$, we have $u \circ \mathcal{O}^{-1} \in W^{1,p}(\Omega_{\hat{\psi}}^n)$ with

$$(3.55) \quad \|u \circ \mathcal{O}^{-1}\|_{W^{1,p}(\Omega_{\hat{\psi}}^n)} \leq C \|u\|_{W^{1,p}(\Omega_\psi^n)}.$$

By the argument above, the Lipschitz outward cuspidal domain $\Omega_{\hat{\psi}}^n$ is a Sobolev (p, q) -extension domain, there exists a function $E(u \circ \mathcal{O}^{-1}) \in W^{1,q}(\mathbb{R}^n)$ with $E(u \circ \mathcal{O}^{-1})|_{\Omega_{\hat{\psi}}^n} \equiv u \circ \mathcal{O}^{-1}$ and

$$(3.56) \quad \|E(u \circ \mathcal{O}^{-1})\|_{W^{1,q}(\mathbb{R}^n)} \leq C \|u \circ \mathcal{O}^{-1}\|_{W^{1,p}(\Omega_{\hat{\psi}}^n)}.$$

By the fact that \mathcal{O} is bi-Lipschitz, we have $(E(u \circ \mathcal{O}^{-1})) \circ \mathcal{O} \in W^{1,q}(\mathbb{R}^n)$ with

$$(3.57) \quad \|(E(u \circ \mathcal{O}^{-1})) \circ \mathcal{O}\|_{W^{1,q}(\mathbb{R}^n)} \leq C \|E(u \circ \mathcal{O}^{-1})\|_{W^{1,q}(\mathbb{R}^n)}.$$

By the definitions, we have $(E(u \circ \mathcal{O}^{-1})) \circ \mathcal{O}|_{\Omega_\psi^n} \equiv u$. By combining (3.55), (3.56) and (3.57), we obtain the desired norm inequality

$$\|(E(u \circ \mathcal{O}^{-1})) \circ \mathcal{O}\|_{W^{1,q}(\mathbb{R}^n)} \leq C \|u\|_{W^{1,p}(\Omega_\psi^n)}.$$

Hence, Ω_ψ^n is a Sobolev (p, q) -extension domain whenever $1 \leq q < n-1$ and $(n-1)q/(n-1-q) \leq p < \infty$. \square

Let us prove Proposition 1.7 which will show the sharpness of Theorem 1.6.

Proof of Proposition 1.7. Let $n-1 \leq p < \infty$. Set

$$s_1 := \frac{np - (n-1)}{(n-1)^2}.$$

By Theorem 1.2, $\Omega_{t^s}^n$ is a Sobolev $(p, n-1)$ -extension domain if and only if $1 \leq s \leq s_1$. Let $1 \leq q < n-1$ and $q \leq p < \frac{(n-1)q}{n-1-q}$. Set

$$s_2 := \frac{pq + p - q}{pq + (n-1)(q-p)}.$$

By Theorem 1.2, $\Omega_{t^s}^n$ is a Sobolev (p, q) -extension domain if and only if $1 \leq s < s_2$. \square

4. FURTHER COMMENTS

In the monograph [17] and their papers referred to therein, Maz'ya and Poborchi also dealt with generalized outward cuspidal domains with Lipschitz base domains. To be more precise, for a bounded Lipschitz domain $U \in \mathbb{R}^{n-1}$ with $0 \in U$ and a cuspidal function $\psi : (0, 1] \rightarrow (0, \infty)$, the corresponding generalized outward cuspidal domain with the base domain U is defined by setting

$$(4.1) \quad U_\psi^n := \{(t, x) \in (0, 1] \times \mathbb{R}^{n-1} : x \in \psi(t)U\} \cup \{(t, x) \in (1, 2) \times \mathbb{R}^{n-1} : x \in \psi(1)U\}.$$

We have only discussed outward cuspidal domains whose base domains are the unit ball. Maz'ya and Poborchi established results for generalized Lipschitz outward cuspidal domains U_ψ^n with Lipschitz

base domains $U \subset \mathbb{R}^{n-1}$. Hence, our results in Theorem 1.2 only extend Maz'ya and Poborchi's results in a special case. For a full extension, one would need to establish a global bi-Lipschitz equivalence of Proposition 3.5 analog for general outward cuspidal domains U_ψ^n . This appears to be technically challenging but we expect it to be doable.

Conjecture 4.2. *For every generalized outward cuspidal domain U_ψ^n , there exists a Lipschitz cuspidal function $\hat{\psi}$, a Lipschitz base domain \hat{U} and a global bi-Lipschitz transformation $\mathcal{O} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ with $\mathcal{O}(U_\psi^n) = \hat{U}_{\hat{\psi}}^n$.*

REFERENCES

- [1] G. Acosta and I. Ojea, *Extension theorems for external cusps with minimal regularity*. Pacific J. Math. 259 (2012), no. 1, 1–39.
- [2] R. A. Adams and J. J. F. Fournier, *Sobolev space*. Second edition. Pure and Applied Mathematics (Amsterdam), 140. Elsevier/Academic Press, Amsterdam, 2003.
- [3] A. P. Calderón, *Lebesgue spaces of differentiable functions and distributions*, Proc. Symp. Pure Math. 4 (1961), 33–49.
- [4] S. Eriksson-Bique, P. Koskela, J. Malý and Z. Zhu, *Pointwise Inequalities for Sobolev Functions on Outward Cuspidal Domains*, International Mathematics Research Notices, , rnaa279, <https://doi.org/10.1093/imrn/rnaa279>
- [5] L. C. Evans and R. F. Gariepy, *Measure Theory and Fine Properties of Functions*, Revised ed., in: Textbooks in Mathematics, CRC Press, Boca Raton, FL, 2015.
- [6] P. Hajlasz, *Sobolev spaces on an arbitrary metric space*, Potential Anal. 5(1996), no. 4, 403–415.
- [7] P. Hajlasz, P. Koskela and H. Tuominen, *Sobolev embeddings, extensions and measure density condition*, J. Funct. Anal. 254 (2008), no. 5, 1217–1234.
- [8] P. Hajlasz, P. Koskela and H. Tuominen, *Measure density and extendability of Sobolev functions*. Rev. Mat. Iberoam. 24 (2008), no. 2, 645–669.
- [9] P. W. Jones, *Quasiconformal mappings and extendability of functions in Sobolev spaces*. Acta Math. 147 (1981) 71–78.
- [10] P. Koskela, *Extensions and imbeddings*. J. Funct. Anal. 159 (1998), no. 2, 369–383.
- [11] P. Koskela, T. Rajala and Y. Zhang, *A geometric characterization of planar Sobolev extension domains*. <https://arxiv.org/abs/1502.04139>.
- [12] P. Koskela, T. Rajala and Y. Zhang, *Planar $W^{1,1}$ -extension domains*. contained in a JYU-dissertation.
- [13] P. Koskela and Z. Zhu, *Product of extension domains is still an extension domain*. Indiana Univ. Math. J. 69 (2020), n0. 1, 137-150.
- [14] V. Maz'ya, *Sobolev spaces: with applications to elliptic partial differential equations*. Springer, Berlin/Heidelberg, 2010.
- [15] V. G. Maz'ya and S. V. Poborchi, *On extension of functions in Sobolev classes to the exterior of a domain with the vertex of a peak on the boundary*, Dokl. Akad. Nauk SSSR 275:5 (1984) 1066-1069 (Russian). English translation: Soviet Math. 29 (1984) 361-364.
- [16] V. G. Maz'ya and S. V. Poborchi, *Extension of functions in Sobolev classes to the exterior of a domain with the vertex of a peak on the boundary I*, Czech. Math. Journ. 36:111 (1986) 634-661 (Russian).
- [17] V. G. Maz'ya and S. V. Poborchi, *Differentiable functions on bad domains*, World Scientific Publishing Co., River Edge, NJ, 1997.
- [18] V. G. Maz'ya and S. V. Poborchi, *Extension of functions in Sobolev classes to the exterior of a domain with the vertex of a peak on the boundary II*, Czech. Math. Journ. 37:112 (1987) 128-150 (Russian).
- [19] P. Shvartsman, *On Sobolev extension domains in \mathbb{R}^n* . J. Funct. Anal. 258 (2010), no. 7, 2205–2245.
- [20] P. Shvartsman, N. Zobin, *On planar Sobolev L_p^m -extension domains*. Adv. Math., 287 (2016), 237–346.
- [21] E. M. Stein, *Singular integrals and differentiability properties of functions*. Princeton University Press, Princeton, 1971.

- [22] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, With the assistance of Timothy S. Murphy, Princeton Mathematical Series, 43. Monographs in Harmonic Analysis, III. Princeton University Press, Princeton, NJ, 1993.
- [23] S. K. Vodop'yanov, V. M. Gol'dstein, T. G. Latfullin, *A criterion for the extension of functions of the class L^1_2 from unbounded plane domains*. Siberian. Math. J., 20 (1979), 416–419.
- [24] W. P. Ziemer, *Weakly differentiable functions. Sobolev spaces and functions of bounded variation*. Graduate Texts in Mathematics, 120. Springer-Verlag, New York, 1989.

PEKKA KOSKELA, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ, P.O. Box 35 (MAD), FI-40014, JYVÄSKYLÄ, FINLAND
Email address: pekka.j.koskela@jyu.fi

ZHENG ZHU, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ, P.O. Box 35 (MAD), FI-40014, JYVÄSKYLÄ, FINLAND
Email address: zheng.z.zhu@jyu.fi