

THE CONDENSATION OF A TRAPPED DILUTE BOSE GAS WITH THREE-BODY INTERACTIONS

PHAN THÀNH NAM, JULIEN RICAUD, AND ARNAUD TRIAY

ABSTRACT. We consider a trapped dilute gas of N bosons in \mathbb{R}^3 interacting via a three-body interaction potential of the form $N V(N^{1/2}(x - y, x - z))$. In the limit $N \rightarrow \infty$, we prove that the ground state of the system exhibits the complete Bose–Einstein condensation, and that the condensate is the unique minimizer of the 3D energy-critical nonlinear Schrödinger functional where the nonlinear term is coupled with the scattering energy of the interaction potential.

CONTENTS

1. Introduction	1
1.1. Model	3
1.2. Main results	5
1.3. Ingredients of the proof	6
2. Scattering energy	12
2.1. General setting	12
2.2. Three-body symmetry	14
3. Dyson lemmas	15
3.1. Dyson lemma for non-radial potentials	15
3.2. Dyson lemma with the three-body symmetry	17
3.3. Many-body Dyson lemma	17
4. Reduction to softer interaction potentials	21
4.1. Binding inequality	21
4.2. Four-body estimate	21
4.3. Proof of Lemma 9	22
5. Conclusion of energy lower bound	28
6. Energy upper bound	36
6.1. The transformation U_N and the proof of Corollary 19	37
6.2. Computation of the upper bound: proof of Theorem 18	41
7. Convergence of ground states	48
References	49

1. INTRODUCTION

The Bose–Einstein condensation was predicted in 1924 [5, 13], but it was not until 1995 that it was first experimentally realized [1, 11], leading to the 2001 Nobel prize in Physics of Cornell, Wieman and Ketterle. While the theoretical works of Bose and Einstein [5, 13] focus only on the ideal gas, interactions among particles do not only make the analysis more involved but also lead to new phenomena, one of the

Date: October 29, 2021.

most famous being superfluidity [15, 4]. In dilute regimes, where collisions occur rarely, the interaction is most often described by an effective two-body potential $v_{2B}(x_1 - x_2)$ which is computed by integrating out the possible internal degrees of freedom of two particles placed at x_1 and x_2 , and in the ideal situation where all the others are infinitely far away. This approximation, enough for most applications, might break down if a third particle nearby affects strongly the internal structure of the two others. In this case, one can add a correction by using another effective interaction potential $v_{3B}(x_1 - x_2, x_1 - x_3)$ in the ideal situation where three-particles are close and the others infinitely far away, etc., namely

$$U(x_1, \dots, x_N) \simeq \sum_{1 \leq i < j \leq N} v_{2B}(x_i - x_j) + \sum_{1 \leq i < j < k \leq N} v_{3B}(x_i - x_j, x_i - x_k) + \dots$$

In this many-body expansion, the three- and higher-order corrections although small are not always negligible: three-body interactions account for 2% of the binding energy of liquid He⁴ [26], even 14% for water [25], making the two-body approximation fail to explain certain of their physical properties [32, 34]. In the realm of condensed matter, the Bose–Hubbard model with tunable two-body and three-body interactions was derived from cold polar molecules, where the two-body interaction can independently be tuned and even switched off [7]. Finally, in the study of ultra-cold gases, three-body interactions have received a strong interest with the hope of observing exotic states like self-trapped droplets or Pfaffian states [31]. In particular, a repulsive three-body interaction can stabilize a condensate against collapse due to an attractive two-body interaction, the competing contributions can lead to crystallization and even to some super-solid state [3, 6].

In this work, we consider a gas of N bosons in \mathbb{R}^3 trapped in a confining potential V_{ext} and interacting via three-body interactions. It would be also interesting to add two-body interactions to the model, but we omit that here in order to simplify the problem. We study the Gross–Pitaevskii like regime where the interaction potential scales like

$$V_N(x - y, x - z) = NV \left(N^{1/2}(x - y, x - z) \right) .$$

We will prove that the ground state energy and the ground state of the system are effectively described by the energy-critical nonlinear Schrödinger functional where the nonlinear term is coupled with the scattering energy of the interaction potential. To our knowledge, it is the first time that the ground state of a dilute Bose gas with three-body interactions was investigated rigorously. In fact, a simpler result with the weaker interaction potential of the form

$$N^{6\beta-2}V \left(N^\beta(x - y, x - z) \right)$$

with $\beta \geq 0$ small can be handled by following the method in [18]. In this case, the usual mean-field approximation is correct to the leading order. The critical case $\beta = 1/2$ that we consider here is more difficult since the three-body correlation between particles yields a leading order correction to the ground state energy, which is similar to the Gross–Pitaevskii regime of the two-body interaction studied in [24, 21, 22].

In the time-dependent problem, the potential $N^{6\beta-2}V(N^\beta(x - y, x - z))$ with $\beta \geq 0$ small already received some attention [10, 8, 35, 9, 16, 29, 19]. The critical regime $\beta = 1/2$ is again more difficult, and will be dealt with in a separate paper [30].

1.1. **Model.** Let us now explain the mathematical setting in detail. We consider a system of N identical bosons in \mathbb{R}^3 described by the Hamiltonian

$$H_N = \sum_{i=1}^N (-\Delta_{x_i} + V_{\text{ext}}(x_i)) + \sum_{1 \leq i < j < k \leq N} V_N(x_i - x_j, x_i - x_k) \quad (1)$$

which acts on the symmetric space

$$\mathfrak{H}^N = L_s^2((\mathbb{R}^3)^N) = \bigotimes_s^N L^2(\mathbb{R}^3).$$

Here we assume that the external trapping potential satisfies

$$V_{\text{ext}} \in L_{\text{loc}}^\infty(\mathbb{R}^3, \mathbb{R}), \quad V_{\text{ext}}(x) \geq C|x|^\alpha \quad \text{for some } C > 0, \alpha > 0. \quad (2)$$

Moreover, the interaction potential is chosen of the form

$$V_N(x, y) = NV(N^{1/2}(x, y)), \quad \forall x, y \in \mathbb{R}^3 \quad (3)$$

where $V : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is nonnegative, bounded and has the symmetry properties

$$V(x, y) = V(y, x), \quad V(x - y, x - z) = V(y - x, y - z) = V(z - y, z - x) \quad (4)$$

which guarantee that the interaction energy of three particles depends only on their relative distances. Thus H_N models a trapped dilute Bose gas where the range of the interaction potential is much smaller than the average distance between particles ($N^{-1/2} \ll N^{-1/3}$).

Under our conditions (2) and (4), the Hamiltonian H_N is well-defined and bounded from below with core domain $\bigotimes_s^N C_c^\infty(\mathbb{R}^3)$. Hence, it can be extended to be a self-adjoint operator on \mathfrak{H}^N by Friedrichs' method. The extension, still denoted by H_N , is bounded from below and has compact resolvent since $V_{\text{ext}}(x) \rightarrow \infty$ as $|x| \rightarrow \infty$.

In the present paper, we are interested in the ground state problem

$$E(N) = \inf_{\|\Psi\|_{\mathfrak{H}^N} = 1} \langle \Psi, H_N \Psi \rangle.$$

When N becomes large, it is in general impossible to compute, both analytically and numerically, the ground state energy $E(N)$ and the corresponding ground state from the full many-body description (1). Hence, for practical computations it is important to derive an effective theory which relies on less variables.

The usual mean-field approximation suggests to restrict to the complete condensation ansatz of N particles

$$\Psi_N \approx u^{\otimes N}(x_1, \dots, x_N) = u(x_1) \dots u(x_N), \quad u \in L^2(\mathbb{R}^3). \quad (5)$$

However, the correlation due to the strong interaction between particles at short distances will also play a leading order role. Since the gas is dilute, the correlation structure is encoded in the scattering problem associated to the three-body interaction potential $V_N(x - y, x - z)$. Let us consider the three-body operator

$$-\Delta_{x_1} - \Delta_{x_2} - \Delta_{x_3} + V_N(x_1 - x_2, x_1 - x_3) \quad \text{on } L^2((\mathbb{R}^3)^3).$$

We can remove the center of mass by change of variables:

$$r_1 = \frac{1}{3}(x_1 + x_2 + x_3), \quad r_2 = x_1 - x_2, \quad r_3 = x_1 - x_3. \quad (6)$$

Denoting the momentum variable $p_x = -i\nabla_x$ with $x \in \mathbb{R}^3$ (here $\mathbf{i}^2 = -1$) we have

$$p_{x_1} = \frac{1}{3}p_{r_1} + p_{r_2} + p_{r_3}, \quad p_{x_2} = \frac{1}{3}p_{r_1} - p_{r_2}, \quad p_{x_3} = \frac{1}{3}p_{r_1} - p_{r_3},$$

and hence

$$\begin{aligned}
& -\Delta_{x_1} - \Delta_{x_2} - \Delta_{x_3} + V_N(x_1 - x_2, x_1 - x_3) \\
&= \left(\frac{1}{3}p_{r_1} + p_{r_2} + p_{r_3}\right)^2 + \left(\frac{1}{3}p_{r_1} - p_{r_2}\right)^2 + \left(\frac{1}{3}p_{r_1} - p_{r_3}\right)^2 + V(r_1, r_2) \\
&= \frac{1}{3}p_{r_1}^2 + 2(p_{r_2}^2 + p_{r_3}^2 + p_{r_2}p_{r_3}) + V_N(r_2, r_3). \quad (7)
\end{aligned}$$

Thus after removing the center of mass, we are left with the two-body operator

$$2(p_x^2 + p_y^2 + p_x p_y) + V_N(x, y) = -2\Delta_{\mathcal{M}} + V_N(x, y) \quad \text{on } L^2((\mathbb{R}^3)^2)$$

where $-\Delta_{\mathcal{M}} = |\mathcal{M}\nabla_{\mathbb{R}^6}|^2$ and the matrix $\mathcal{M} : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}^3 \times \mathbb{R}^3$ is given by

$$\mathcal{M} := \left(\frac{1}{2} \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}\right)^{1/2} = \frac{1}{2\sqrt{2}} \begin{pmatrix} \sqrt{3} + 1 & \sqrt{3} - 1 \\ \sqrt{3} - 1 & \sqrt{3} + 1 \end{pmatrix}. \quad (8)$$

The operator $-2\Delta_{\mathcal{M}} + V_N(x, y)$ is associated to the (modified) *scattering energy*

$$b_{\mathcal{M}}(V_N) := \inf_{\omega \in \dot{H}^1(\mathbb{R}^6)} \int_{\mathbb{R}^6} (2|\mathcal{M}\nabla\omega(\mathbf{x})|^2 + V_N(\mathbf{x})|1 - \omega(\mathbf{x})|^2) \, d\mathbf{x} \quad (9)$$

Recall the standard scattering energy of a potential $W : \mathbb{R}^d \rightarrow \mathbb{R}$

$$b(W) := \inf_{\omega \in \dot{H}^1(\mathbb{R}^d)} \int_{\mathbb{R}^d} (2|\nabla\omega(\mathbf{x})|^2 + W(\mathbf{x})|1 - \omega(\mathbf{x})|^2) \, d\mathbf{x}. \quad (10)$$

By change of variables, the modified scattering energy can be written as

$$b_{\mathcal{M}}(V) = b(V(\mathcal{M}\cdot)) \det \mathcal{M}$$

(here $\det \mathcal{M} = 3\sqrt{3}/8$). Moreover, the specific choice in (3) ensures that

$$b_{\mathcal{M}}(V_N) = \frac{b_{\mathcal{M}}(V)}{N^2}.$$

In summary, to the leading order we expect that

$$V_N(x, y) \approx \frac{b_{\mathcal{M}}(V)}{N^2} \delta_{x=y=0}. \quad (11)$$

Putting the formal approximations (5) and (11) together, we arrive at

$$E(N) \approx Ne_{\text{GP}} = N \inf_{\|u\|_{L^2(\mathbb{R}^3)}=1} \mathcal{E}_{\text{GP}}(u) \quad (12)$$

where $\mathcal{E}_{\text{GP}}(u)$ is the 3D energy-critical nonlinear Schrödinger (NLS) functional

$$\mathcal{E}_{\text{GP}}(u) = \int_{\mathbb{R}^3} \left(|\nabla u(x)|^2 + V_{\text{ext}}(x)|u(x)|^2 + \frac{b_{\mathcal{M}}(V)}{6}|u(x)|^6 \right) \, dx. \quad (13)$$

Note that the effective expression (13) is analogous to the Gross–Pitaevskii functional with two-body interaction (for the latter the nonlinear term is $|u|^4$ instead of $|u|^6$). The validity of the Gross–Pitaevskii theory for the ground state energy of trapped dilute Bose gases with a two-body interaction potential of the form $N^2W(N(x - y))$ has been proved in seminal papers of Lieb, Seiringer and Yngvason [24, 21, 22]. Our aim here is to extend these results to the three-body interaction case.

1.2. **Main results.** Our first result is a rigorous justification of (12).

Theorem 1 (Ground state energy). *Let $0 \leq V \in L^\infty(\mathbb{R}^6)$ be compactly supported and satisfy the symmetry conditions (4). Let V_{ext} be as in (2). Then the ground state energy of H_N in (1) satisfies*

$$\lim_{N \rightarrow \infty} \frac{E(N)}{N} = e_{\text{GP}} = \inf_{\|u\|_{L^2(\mathbb{R}^3)}=1} \mathcal{E}_{\text{GP}}(u) \quad (14)$$

where the effective functional $\mathcal{E}_{\text{GP}}(u)$ is given in (13).

As a by product of our proof of Theorem 1, we also obtain the condensation of the ground state of H_N . Note that the approximation (5) is expected to hold not in the norm topology, but rather in a weaker topology defined by reduced density matrices. Recall that the one-body density matrix $\gamma_{\Psi_N}^{(1)}$ of a N -body normalized wave function Ψ_N is a nonnegative trace class operator on $L^2(\mathbb{R}^3)$ with kernel¹

$$\gamma_{\Psi_N}^{(1)}(x; y) = \int \Psi_N(x, x_2, \dots, x_N) \overline{\Psi_N(y, x_2, \dots, x_N)} dx_2 \cdots dx_N. \quad (15)$$

Equivalently, we can also write $\gamma_{\Psi_N}^{(1)} = \text{Tr}_{2 \rightarrow N} |\Psi\rangle\langle\Psi_N|$, where the partial trace is taken over all but the first variables. Thus the proper meaning of (5) is

$$\gamma_{\Psi_N}^{(1)} \approx |u\rangle\langle u| \quad (16)$$

which is often referred to as the complete Bose–Einstein condensation.

When $\gamma_{\Psi_N}^{(1)}$ is a ground state, or more generally an approximate ground state, of H_N , then the right side of (16) should be the minimizer of e_{GP} . The existence of a minimizer u_0 of e_{GP} follows straightforwardly from the direct variational method, where the compactness in $L^2(\mathbb{R}^3)$ of minimizing sequence is guaranteed by the trapping condition $V_{\text{ext}}(x) \rightarrow \infty$ as $|x| \rightarrow \infty$. Moreover, thanks to the diamagnetic inequality $\mathcal{E}_{\text{GP}}(u) \geq \mathcal{E}_{\text{GP}}(|u|)$ and the strict convexity of $0 \leq \rho \mapsto \mathcal{E}_{\text{GP}}(\sqrt{\rho})$, the minimizer u_0 is unique up to a complex phase and can be chosen such that $u_0(x) \geq 0$ pointwise. Moreover, it solves the nonlinear equation

$$(-\Delta + V_{\text{ext}}(x) + b_{\mathcal{M}}(V)|u_0(x)|^4/2 - \varepsilon_0) u_0(x) = 0, \quad x \in \mathbb{R}^3 \quad (17)$$

for some chemical potential $\varepsilon_0 \in \mathbb{R}^3$ (the Lagrange multiplier associated with the mass constraint $\|u\|_{L^2(\mathbb{R}^3)} = 1$). Moreover, the trapping condition implies exponential decay for u_0 (see for instance [24, Appendix A]), hence that $u_0 \in L^\infty(\mathbb{R}^3)$.

Our second result is a rigorous justification of (16) for the ground state, or more generally for approximate ground states, of H_N .

Theorem 2 (Condensation of approximate ground states). *Let V, V_{ext} be as in Theorem 1. Assume that Ψ_N is an approximate ground state for H_N , namely*

$$\|\Psi_N\|_{\mathfrak{H}^N} = 1, \quad \lim_{N \rightarrow \infty} \frac{\langle \Psi_N, H_N \Psi_N \rangle}{N} = e_{\text{GP}}.$$

Then we have

$$\lim_{N \rightarrow \infty} \gamma_{\Psi_N}^{(1)} = |u_0\rangle\langle u_0|$$

in trace class, where u_0 is the unique minimizer of e_{GP} .

¹In our convention, inner products in (complex) Hilbert spaces are linear in the second argument and anti-linear in the first.

From our approach, we also find that for every $0 < \beta < 1/2$, the ground state energy of the Hamiltonian

$$H_{N,\text{mf}} = \sum_{i=1}^N (-\Delta_{x_i} + V_{\text{ext}}(x_i)) + \sum_{1 \leq i < j < k \leq N} N^{6\beta-2} V(N^\beta(x_i - x_j, x_i - x_k)) \quad (18)$$

satisfies

$$\lim_{N \rightarrow \infty} \frac{E(N)}{N} = \inf_{\|u\|_{L^2(\mathbb{R}^3)}=1} \int_{\mathbb{R}^3} \left(|\nabla u(x)|^2 + V_{\text{ext}}(x)|u(x)|^2 + \frac{\widehat{V}(0)}{6}|u(x)|^6 \right) dx. \quad (19)$$

This is an analogue of (14) where the scattering energy $b_{\mathcal{M}}(V)$ is replaced by its first Born approximation $\widehat{V}(0) = \int_{\mathbb{R}^6} V$. Consequently, we also obtain the convergence of approximate ground states of $H_{N,\text{mf}}$ to the minimizer of the right side of (19). When $\beta > 0$ is small, (19) can be proved by using a standard mean-field technique, for example by using a quantitative quantum de Finetti as in [18]. The proof of (19) for the whole range $0 < \beta < 1/2$ is more difficult and can be obtained from a simplification of our method. The critical case $\beta = 1/2$ in Theorem 1 is the hardest case since the strong correlation yields a leading order correction to the mean-field approximation

In the context of quantum dynamics, it was proved in [9, 29] that for $\beta > 0$ small, the Schrödinger evolution $\Psi_N(t) = e^{itH_{N,\text{mf}}^0} \Psi_N(0)$ with initial condition $\gamma_{\Psi_N(0)}^{(1)} \approx |u(0)\rangle\langle u(0)|$ exhibits the complete condensation $\gamma_{\Psi_N(t)}^{(1)} \approx |u(t)\rangle\langle u(t)|$ for any time $t > 0$ and the condensate $u(t)$ is determined by the time-dependent equation

$$i\partial_t u(t, x) = \left(-\Delta + \frac{1}{2}\widehat{V}(0)|u(t, x)|^4 - \varepsilon(t) \right) u(t, x). \quad (20)$$

Here $H_{N,\text{mf}}^0$ is the Hamiltonian as in (18) but without the external potential. We expect that the same result holds for all $0 < \beta < 1/2$, and that in the critical case $\beta = 1/2$ the factor $\widehat{V}(0)$ in (20) will be replaced by the scattering energy $b_{\mathcal{M}}(V)$.

1.3. Ingredients of the proof. Now let us quickly explain some ideas of the proof. We will focus on Theorem 1. The implication of Theorem 2 from Theorem 1 follows from a standard Hellmann–Feynman argument as in [21, 22], which will be discussed in Section 7.

Upper bound. To get the leading order upper bound $E(N) \leq Ne_{\text{GP}} + o(N)$, the uncorrelated trial state as in (5) is insufficient. In fact, the energy per particles over the Hartree states $u^{\otimes N}$ is given by a functional similar to $\mathcal{E}_{\text{GP}}(u)$ but the scattering energy $b_{\mathcal{M}}(V)$ in front of the nonlinear term $|u|^6$ is replaced by $\widehat{V}(0) = \int_{\mathbb{R}^6} V$. Hence, it is important to take some correlation into account. Heuristically, we can think of the Jastrow–Dyson type state

$$\Psi_N(x_1, \dots, x_N) = \prod_{j=1}^N u_0(x_j) \prod_{p < k < \ell} f_N(x_p - x_k, x_p - x_\ell) \quad (21)$$

where u_0 is the minimizer of e_{GP} and $f_N : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is a function satisfying the symmetry (4) and solves the scattering equation

$$-2\Delta_{\mathbf{x}} f_N(\mathbf{x}) + V_N(\mathbf{x}) f_N(\mathbf{x}) = 0, \quad \forall \mathbf{x} \in \mathbb{R}^6$$

which is equivalent to

$$(-\Delta_x - \Delta_y - \Delta_z) f_N(x - y, x - z) + (V_N f_N)(x - y, x - z) = 0, \quad \forall x, y, z \in \mathbb{R}^3. \quad (22)$$

The existence of such a function is proved in Theorem 5. In particular, the scattering energy is encoded in f_N as

$$\int_{\mathbb{R}^6} V_N f_N = b_{\mathcal{M}}(V_N) = N^{-2} b_{\mathcal{M}}(V).$$

Unfortunately, we are not able to compute the ground state energy per particle of the trial state (21) to the leading order (the computation is significantly more complicated than the two-body interaction case).

Here we follow an alternative approach. First, for ease of computation, we extend H_N to the operator $\mathbb{H}_N = 0 \oplus \bigoplus_{M=1}^{\infty} H_{M,N}$ acting on the bosonic Fock space $\mathcal{F}(\mathfrak{H}) = \Omega \oplus \bigoplus_{M=1}^{\infty} \mathfrak{H}^M$ (here Ω is the vacuum) with

$$H_{M,N} = \sum_{j=1}^M (-\Delta_j + V_{\text{ext}}(x_j)) + \sum_{1 \leq i < j < k \leq M} NV(N^{1/2}(x_i - x_j, x_i - x_k)). \quad (23)$$

This extension can be written conveniently as

$$\begin{aligned} \mathbb{H}_N = \int_{\mathbb{R}^3} (\nabla_x a_x^* \nabla_x a_x + V_{\text{ext}}(x) a_x^* a_x) dx \\ + \frac{1}{6} \int_{(\mathbb{R}^3)^3} V_N(x-y, x-z) a_x^* a_y^* a_z^* a_x a_y a_z dx dy dz \end{aligned}$$

using the standard creation and annihilation operators a_x^*, a_x . To capture the condensation, we define the Weyl operator

$$W(f) = \exp(a^*(f) - a(f)), \quad \forall f \in L^2(\mathbb{R}^3)$$

which is a unitary operator on Fock space and satisfies

$$W(f)^* a(g) W(f) = a(g) + \langle g, f \rangle, \quad \forall f, g \in L^2(\mathbb{R}^3).$$

To create the desired correlation structure encoded in the scattering solution f_N in (22), we introduce another unitary transformation

$$U_N := \exp(\mathbb{1}(\mathcal{N} \leq L) B_1^* - B_1 \mathbb{1}(\mathcal{N} \leq L)),$$

where $\mathcal{N} = \int a_x^* a_x dx$ is the number operator on Fock space and

$$B_1^* = -\frac{1}{6} N^{\frac{3}{2}} \int_{(\mathbb{R}^3)^3} (1 - f_N)(x-y, x-z) u_0(x) u_0(y) u_0(z) a_x^* a_y^* a_z^* dx dy dz. \quad (24)$$

We choose the cut-off L to verify $1 \ll L \leq CN^{1/2}$ such that U_N does not create too many excited particles. We will prove (see Theorem 18 below) that

$$\left\langle \Omega, U_N^* W(\sqrt{N}\varphi)^* \mathbb{H}_N W(\sqrt{N}\varphi) U_N \Omega \right\rangle \leq Ne_{\text{GP}} + \mathcal{O}(N^{2/3}). \quad (25)$$

Note that the trial state $W(\sqrt{N}\varphi) U_N \Omega$ does not belong to the N -body space \mathfrak{H}^N , but it essentially lives on sections of $N + O(\sqrt{N})$ particle number. Following an idea of Solovej [33], by controlling the fluctuations of the particle number of $W(\sqrt{N}\varphi) U_N \Omega$, we are able to construct a trial state in the N -body Hilbert space and obtain the desired upper bound

$$E(N) \leq Ne_{\text{GP}} + o(N).$$

In the two-body interaction case, a similar trial state has been used [2, 27]. In this case, equation (24) is replaced by a kernel that is quadratic in terms of creation and annihilation operators, which simplifies the computation greatly since $U_N \Omega$ is a quasi-free state. In our case, the computation with the cubic kernel in equation (24)

is more complicated, but technically manageable. The details will be explained in Section (6).

Lower bound. The main difficulty of Theorem 1 lies on the lower bound. First, by following the arguments in [24, 21, 22], we will replace the singular potential V_N by a softer one, using a Dyson lemma. We will prove (see Theorem 7) that given the potential $V_N \geq 0$ supported in $|\mathbf{x}| \leq O(N^{-1/2})$ and $R \gg N^{-1/2}$, we can find a function $U \geq 0$ supported in $|x| \leq R$ such that U_R satisfies the symmetry (4),

$$U_R(x, y) \leq CR^{-6} \mathbf{1}(|x| \leq R) \mathbf{1}(|y| \leq R), \quad \int_{\mathbb{R}^6} U_R = b_{\mathcal{M}}(V)(1 + o(1)_{N \rightarrow \infty})$$

and

$$-2\mathcal{M}\nabla_{\mathbf{x}} \mathbf{1}_{\{|\mathbf{x}| \leq \sqrt{2}R\}} \mathcal{M}\nabla_{\mathbf{x}} + V_N(\mathbf{x}) \geq N^{-2}U_R(\mathbf{x}) \quad \text{on } L^2(\mathbb{R}^6). \quad (26)$$

More precisely, we will derive an improved version of (26), where only the high-momentum part $|p| \geq \varepsilon^{-1}$ of the kinetic energy is needed to replace V_N by $N^{-2}U_R$, and the low-momentum part $|p| \leq \varepsilon^{-1}$ is kept in order to recover the full nonlinear functional in (13).

Note that all existing proofs of the Dyson lemma and its generalizations rely on the radial symmetry of the potential (see [24, 23]), which is not available here. Our proof of (26) goes differently and works for a much larger class of potentials. We will derive (26) from a related result of the standard scattering energy (see Theorem 6) which is of general interest.

Now coming back to the Hamiltonian H_N , using (26) we obtain in Lemma 8 the following lower bound for all $1 > \varepsilon > 0$,

$$\begin{aligned} & (1 - \varepsilon)^{-1} H_N + C_\varepsilon R^3 N^2 \\ & \geq \sum_{i=1}^N h_i + \frac{1}{6N^2} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} U_R(x_i - x_j, x_i - x_k) \prod_{\ell \neq i, j, k} \theta_{2R} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right) \end{aligned} \quad (27)$$

where

$$h = \varepsilon p^2 + p^2 \mathbf{1}_{\{|p| \leq \varepsilon^{-1}\}} + V_{\text{ext}}(x) \quad \text{and} \quad \theta_{2R} = 1(|x| \geq 2R).$$

The cut-off $\theta_{2R}(x) = \mathbf{1}_{\{|x| \geq 2R\}}$ appears due to the fact that we do not include the event of having four particles within a distance $O(R)$. This is a disadvantage of using the Dyson lemma, and the four-body problem here is similar to the three-body problem for the two-body interaction case. The standard way to remove the cut-off θ_{2R} is to use Bernoulli's inequality

$$\begin{aligned} \prod_{\ell \neq i, j, k} \theta_{2R} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right) &= \prod_{\ell \neq i, j, k} \left(1 - \chi_{2R} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right) \right) \\ &\geq 1 - \sum_{\ell \neq i, j, k} \chi_{2R} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right) \end{aligned}$$

where $\chi_{2R}(x) = \mathbf{1}_{\{|x| < 2R\}}$. Consequently the interaction in (27) can be bounded from below as

$$\frac{1}{6N^2} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} U_R(x_i - x_j, x_i - x_k) \prod_{\ell \neq i, j, k} \theta_{2R} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right)$$

$$\begin{aligned} &\geq \frac{1}{6N^2} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} U_R(x_i - x_j, x_i - x_k) \\ &\quad - \frac{C}{N^2 R^6} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} \sum_{\ell \neq i, j, k} \mathbb{1}_{\{|x_i - x_j| \leq 4R\}} \mathbb{1}_{\{|x_i - x_k| \leq 4R\}} \mathbb{1}_{\{|x_i - x_\ell| \leq 4R\}}. \end{aligned} \quad (28)$$

Following the approach in [22], we can show (see Lemma 12) that up to a replacement of N by $M \approx N$ if necessary, the ground state Ψ_N of H_N satisfies the four-body collision estimate

$$\left\langle \Psi_N, \prod_{\ell=2}^4 \mathbb{1}(|x_1 - x_\ell| \leq R) \Psi_N \right\rangle \leq CR^9. \quad (29)$$

Therefore, the expectation against Ψ_N of the error term in (28) is bounded by CN^2R^3 , which coincides with the error in (27). In summary, from (27) and (28) we deduce that

$$\frac{E_N}{N} \geq \frac{1-\varepsilon}{N} \left\langle \Psi_N, \left(\sum_{i=1}^N h_i + \frac{1}{6N^2} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} U_R(x_i - x_j, x_i - x_k) \right) \Psi_N \right\rangle - C_\varepsilon NR^3 \quad (30)$$

where Ψ_N is the ground state of H_N . In order to keep the error of order $o(1)$, we need to take

$$N^{-1/3} \gg R \gg N^{-1/2}. \quad (31)$$

These constraints on R are optimal to make the Dyson lemma useful: the condition $R \gg N^{-1/2}$ ensures that we replace V_N by a less singular potential, while the condition $R \ll N^{-1/3}$ keeps us in the dilute regime where there are essentially no four-body interactions, and hence applying the Dyson lemma does not change the energy to the leading order.

So far we follow closely the existing analysis in the two-body interaction case. Now let us explain a crucial new difficulty in the three-body interaction case, which implicitly lies on the fact that we are dealing with an energy-critical problem here. In the two-body interaction case, as soon as we arrive at an analogue of (30), the right-hand side can be treated easily using standard mean-field techniques, e.g. using the coherent state method as in [22] or using the quantum de Finetti theorem as in [28]. A key ingredient needed in [22, 28] is the two-body inequality

$$|W(x-y)| \leq C \|W\|_{L^1(\mathbb{R}^3)} (1-\Delta_x)(1-\Delta_y) \quad (32)$$

on $L^2((\mathbb{R}^3)^2)$, which allows to control the interaction potential efficiently by the kinetic operator. The bound (32) also plays an essential role in the study of the Gross–Pitaevskii dynamics in [14]. In fact, as proved in [28], the following refinement of (32) holds true

$$|W(x-y)| \leq C_\eta \|W\|_{L^1(\mathbb{R}^3)} (1-\Delta_x)^{3/4+\eta} (1-\Delta_y)^{3/4+\eta} \quad (33)$$

for every $\eta > 0$. Roughly speaking, (33) can be interpreted as a variant of the Sobolev embedding Theorem $L^\infty(\mathbb{R}^3) \subset H^{3/2+2\eta}(\mathbb{R}^3)$, where the total $(3/2 + 2\eta)$ derivative on $x-y$ are divided equally for variables x and y . In the three-body interaction case, the analogue of (33) becomes

$$|W(x-y, x-z)| \leq C_\eta \|W\|_{L^1(\mathbb{R}^6)} (1-\Delta_x)^{1+\eta} (1-\Delta_y)^{1+\eta} (1-\Delta_z)^{1+\eta} \quad (34)$$

on $L^2((\mathbb{R}^3)^3)$ for every $\eta > 0$, which should be compared with the Sobolev embedding Theorem $L^2(\mathbb{R}^6) \subset H^{3+3\eta}(\mathbb{R}^6)$. In particular, there is no extension of (32) to the three-body case, namely we cannot take $\eta = 0$ in (34). For that reason, we are not able to apply directly the mean-field techniques from [22, 28] to handle the right-hand side of (30). More precisely, we can try to replace (34) by the following bound

$$|W(x-y, x-z)| \leq C_p \|W\|_{L^p(\mathbb{R}^6)} (1 - \Delta_x)(1 - \Delta_y)(1 - \Delta_z) \quad (35)$$

for $p > 1$, but $\|U_R\|_{L^p} \sim R^{-6(p-1)}$ is too large in application due to the constraint $R \ll N^{-1/3}$ in (31).

To overcome this difficulty, we will further replace U_R by softer potentials before applying the mean-field techniques. To be precise, by the bosonic symmetry of Ψ_N , we can rewrite the right-hand side of (30) as

$$\begin{aligned} & \frac{1}{N} \left\langle \Psi_N, \left(\sum_{i=1}^N h_i + \frac{1}{6N^2} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} U_R(x_i - x_j, x_i - x_k) \right) \Psi_N \right\rangle \\ & \approx \frac{1}{N_1} \left\langle \Psi_N \left(\sum_{i=1}^{N_1} h_i + \frac{1}{6N_1^2} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} U_R(x_i - x_j, x_i - x_k) \right) \Psi_N \right\rangle \quad (36) \end{aligned}$$

for every $N \geq N_1 \gg 1$. Then by using again the Dyson lemma, we can replace U_R by U_{R_1} , namely

$$\begin{aligned} & \frac{1}{N_1} \left\langle \Psi_N \left(\sum_{i=1}^{N_1} h_i + \frac{1}{6N_1^2} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} U_R(x_i - x_j, x_i - x_k) \right) \Psi_N \right\rangle \\ & \geq \frac{1-\varepsilon}{N_1} \left\langle \Psi_N \left(\sum_{i=1}^{N_1} h_i + \frac{1}{6N_1^2} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} U_{R_1}(x_i - x_j, x_i - x_k) \right) \Psi_N \right\rangle + o(1) \quad (37) \end{aligned}$$

as soon as

$$N_1^{-1/3} \gg R_1 \gg R \gg N_1^{-1/2}. \quad (38)$$

The latter constraints are comparable to (31).

Note that there is a subtle difference between (30) and (37). In our first use of the Dyson lemma to obtain (30), we sacrifice essentially all of the high momentum part of the kinetic energy. However, in our second use of the Dyson lemma to obtain (37), we only need a small fraction of the kinetic energy. More precisely, we have

$$-\varepsilon \mathcal{M} \nabla_{\mathbf{x}} \mathbb{1}_{\{|\mathbf{x}| \leq \sqrt{2}R\}} \mathcal{M} \nabla_{\mathbf{x}} + N_1^{-2} U_R(\mathbf{x}) \geq N_1^{-2} U_{R_1}(\mathbf{x}) (1 + o(1)_{N \rightarrow \infty}) \quad (39)$$

which follows from the same proof of (26) and the fact that the scattering energy of $W = \varepsilon^{-1} N_1^{-2} U_R$ is well approximated by its first Born approximation $\widetilde{W}(0)$. The latter fact is a consequence of the inequality

$$b_{\mathcal{M}}(W) \geq \|W\|_{L^1(\mathbb{R}^6)} - C \|W\|_{L^{3/2}(\mathbb{R}^6)}^2$$

where the error term $\|W\|_{L^{3/2}(\mathbb{R}^6)}^2$ is much smaller than $\|W\|_{L^1(\mathbb{R}^6)}$ when $R \gg N_1^{-1/2}$. Heuristically, (39) is more advantageous than (26) since the potential $N^{-2} U_R$ is less singular than the original potential V_N .

To summarize, using (36), (37) and the bosonic symmetry again we obtain

$$\begin{aligned}
& \frac{1}{N} \left\langle \Psi_N, \left(\sum_{i=1}^N h_i + \frac{1}{6N^2} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} U_R(x_i - x_j, x_i - x_k) \right) \Psi_N \right\rangle \\
& \geq \frac{1-\varepsilon}{N} \left\langle \Psi_N, \left(\sum_{i=1}^N h_i + \frac{1}{6N^2} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} U_{R_1}(x_i - x_j, x_i - x_k) \right) \Psi_N \right\rangle + o(1) \quad (40)
\end{aligned}$$

provided that

$$R^{2/3} \gg R_1 \gg R. \quad (41)$$

The condition (41) is obtained from (38) after an optimization over N_1 . Repeating this procedure, we find that for every $\ell \geq 1$,

$$\begin{aligned}
& \frac{1}{N} \left\langle \Psi_N, \left(\sum_{i=1}^N h_i + \frac{1}{6N^2} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} U_{R_\ell}(x_i - x_j, x_i - x_k) \right) \Psi_N \right\rangle \\
& \geq \frac{1-\varepsilon}{N} \left\langle \Psi_N, \left(\sum_{i=1}^N h_i + \frac{1}{6N^2} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} U_{R_{\ell+1}}(x_i - x_j, x_i - x_k) \right) \Psi_N \right\rangle + o(1) \quad (42)
\end{aligned}$$

provided that

$$R_\ell^{2/3} \gg R_{\ell+1} \gg R_\ell. \quad (43)$$

For every $\eta > 0$ small, we can take $R_\ell = N^\eta$ for some $\ell = \ell(\eta)$ sufficiently large. Putting all together, we deduce from (30) that

$$\begin{aligned}
& \frac{E_N}{N} \geq \\
& \frac{(1-\varepsilon)^{\ell+1}}{N} \left\langle \Psi_N, \left(\sum_{i=1}^N h_i + \frac{1}{6N^2} \sum_{\substack{1 \leq i, j, k \leq N \\ i \neq j \neq k \neq i}} U_{R_\ell}(x_i - x_j, x_i - x_k) \right) \Psi_N \right\rangle + o(1). \quad (44)
\end{aligned}$$

Finally, since the potential U_{R_ℓ} is not too singular, we can handle the right-hand side of (44) by the mean-field technique from [17]. The details will be discussed in Section 5. All this gives

$$\frac{E_N}{N} \geq (1-\varepsilon)^{\ell+1} \inf_{\|u\|_{L^2(\mathbb{R}^3)}=1} \left(\langle u, hu \rangle + \frac{b\mathcal{M}(V)}{6} \int_{\mathbb{R}^3} |u(x)|^6 dx \right) + o(1) \quad (45)$$

where $h = \varepsilon p^2 + p^2 \mathbf{1}_{\{|p| \leq \varepsilon^{-1}\}} + V_{\text{ext}}(x)$. Finally, taking $\varepsilon \rightarrow 0^+$ we obtain the desired lower bound

$$\frac{E_N}{N} \geq \inf_{\|u\|_{L^2(\mathbb{R}^3)}=1} \left(\langle u, (-\Delta + V_{\text{ext}})u \rangle + \frac{b\mathcal{M}(V)}{6} \int_{\mathbb{R}^3} |u(x)|^6 dx \right) + o(1). \quad (46)$$

This completes our sketch of the proof of Theorem 1.

Organization of the paper. In Section (2), we will discuss basic facts on the scattering energy in (10). Then we derive several versions of the Dyson lemma in Section 3, which will be used to replace the potential V_N by softer ones in Section (4). In Section 5, we conclude the energy lower bound in Theorem 1. The matching energy upper bound is proved in Section 6. Finally, the convergence of states in Theorem 2 is obtained in Section 7.

Acknowledgments. AT thanks Jonas Lampart for helpful discussions. We received funding from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy (EXC-2111-390814868).

Notation: from now on and for shortness, when there is no possible confusion we will denote by $\|\cdot\|_p$ the L^p -norm on whole space \mathbb{R}^d : $\|\cdot\|_p := \|\cdot\|_{L^p(\mathbb{R}^d)}$.

2. SCATTERING ENERGY

2.1. General setting. Let $d \geq 3$ and let $0 \leq v \in L^\infty(\mathbb{R}^d)$ be compactly supported. We define the *zero-scattering energy* of v by

$$b(v) := \inf_{\varphi \in \dot{H}^1(\mathbb{R}^d)} \int_{\mathbb{R}^d} (2|\nabla\varphi(\mathbf{x})|^2 + v(\mathbf{x})|1 - \varphi(\mathbf{x})|^2) \, d\mathbf{x}. \quad (47)$$

Here $\dot{H}^1(\mathbb{R}^d)$ is the space of functions $g : \mathbb{R}^d \rightarrow \mathbb{C}$ vanishing at infinity and $|\nabla g| \in L^2(\mathbb{R}^d)$ (denoted by $D^1(\mathbb{R}^d)$ in Lieb–Loss [20, Section 8.3]).

Theorem 3 (General scattering solution). *Let $d \geq 3$. Let $0 \leq v \in L^\infty(\mathbb{R}^d)$ be compactly supported. Then the variational problem (47) has a unique minimizer $\omega = (-2\Delta + v)^{-1}v$ and it solves the scattering equation*

$$-2\Delta\omega(\mathbf{x}) + v(\mathbf{x})(\omega(\mathbf{x}) - 1) = 0, \quad \forall \mathbf{x} \in \mathbb{R}^d.$$

We have the pointwise estimates

$$0 \leq \omega(\mathbf{x}) < 1, \quad \omega(\mathbf{x}) \leq \frac{C_{d,v}}{|\mathbf{x}|^{d-2} + 1}, \quad |\nabla\omega(\mathbf{x})| \leq \frac{C_{d,v}}{|\mathbf{x}|^{d-1} + 1}, \quad \forall \mathbf{x} \in \mathbb{R}^d. \quad (48)$$

Moreover, the scattering energy satisfies

$$b(v) = \int_{\mathbb{R}^d} v(\mathbf{x})(1 - \omega(\mathbf{x})) \, d\mathbf{x}, \quad 0 \leq \int_{\mathbb{R}^d} v - b(v) \leq C_d \|v\|_{L^{2d/(d+2)}}^2.$$

Proof. Let $\{\varphi_n\}_{n=1}^\infty \subset \dot{H}^1(\mathbb{R}^d)$ be a minimizing sequence for the functional

$$\mathcal{E}[\varphi] = \int_{\mathbb{R}^d} (2|\nabla\varphi(\mathbf{x})|^2 + v(\mathbf{x})|1 - \varphi(\mathbf{x})|^2) \, d\mathbf{x}.$$

We can assume that φ_n is real-valued since we can ignore the imaginary part of φ_n without increasing $\mathcal{E}[\varphi_n]$. Moreover, we can assume that $0 \leq \varphi_n \leq 1$ since we can replace φ_n by $\min(\max(\varphi_n, 0), 1)$ without increasing $\mathcal{E}[\varphi_n]$. Since $\mathcal{E}[\varphi_n]$ is bounded, φ_n is bounded in $\dot{H}^1(\mathbb{R}^d)$ and $\sqrt{v}(1 - \varphi_n)$ is bounded in $L^2(\mathbb{R}^d)$. By the Banach–Alaoglu theorem, up to a subsequence we can assume that $\varphi_n \rightharpoonup \omega$ weakly in $\dot{H}^1(\mathbb{R}^d)$ and $\sqrt{v}(1 - \varphi_n) \rightarrow \sqrt{v}(1 - \omega)$ weakly in $L^2(\mathbb{R}^d)$. By Fatou's lemma, we conclude that ω is a minimizer. The minimizer ω is unique since the functional $\varphi \mapsto \mathcal{E}[\varphi]$ is strictly convex.

From the above proof, we have $0 \leq \omega \leq 1$. Moreover, for any function $\varphi \in C_c^\infty(\mathbb{R}^d)$, $\mathcal{E}[\omega] \leq \mathcal{E}[\omega + t\varphi]$ for all $t \geq 0$. Hence

$$0 \leq \frac{d}{dt} \Big|_{t=0} \mathcal{E}[\omega + t\varphi] = 2 \int_{\mathbb{R}^d} (2\nabla\omega \cdot \nabla\varphi + v(\omega - 1)\varphi).$$

Thus

$$-2\Delta\omega + v(\omega - 1) = 0 \quad (49)$$

in the distributional sense. Since $0 \leq v(1 - \omega) \leq v \in L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$, we get $\omega \in H^2(\mathbb{R}^d) \cap C^1(\mathbb{R}^d)$ by the standard elliptic regularity [20, Theorem 10.2]. Thus equation (49) holds in the pointwise sense.

The scattering equation can be written as

$$\omega(\mathbf{x}) = (-2\Delta)^{-1}[v(1 - \omega)](\mathbf{x}) = \frac{1}{2|\mathbb{S}^{d-1}|(d-2)} \int_{\mathbb{R}^d} \frac{v(\mathbf{y})(1 - \omega(\mathbf{y})) \, d\mathbf{y}}{|\mathbf{x} - \mathbf{y}|^{d-2}} \quad (50)$$

where $|\mathbb{S}^{d-1}|$ is the surface area of the $(d-1)$ dimensional sphere \mathbb{S}^{d-1} . Since $v(1 - \omega) \in L^1(\mathbb{R}^d)$ and it has compact support, we deduce from (50) that $\omega(\mathbf{x})$ decays as $\mathcal{O}(|\mathbf{x}|^{2-d})$ as $|\mathbf{x}| \rightarrow \infty$. Since $0 \leq \omega \leq 1$ everywhere, we conclude that

$$\omega(\mathbf{x}) \leq \frac{C_{d,v}}{|\mathbf{x}|^{d-2} + 1}, \quad \forall \mathbf{x} \in \mathbb{R}^d.$$

Moreover, note that ω is a subharmonic function in \mathbb{R}^d since $2\Delta\omega = v(1 - \omega) \geq 0$. Therefore, from $\omega \leq 1$ everywhere and $\omega(\mathbf{x}) \rightarrow 0$ as $|\mathbf{x}| \rightarrow \infty$ we find that $\omega < 1$ everywhere by the mean-value theorem [?, Section 9.2].

From (50), we also obtain

$$\nabla\omega(\mathbf{x}) = -\frac{1}{2|\mathbb{S}^{d-1}|} \int_{\mathbb{R}^d} \frac{v(\mathbf{y})(1 - \omega(\mathbf{y}))(\mathbf{x} - \mathbf{y}) \, d\mathbf{y}}{|\mathbf{x} - \mathbf{y}|^d}. \quad (51)$$

This implies that $|\nabla\omega(\mathbf{x})|$ decays as $\mathcal{O}(|\mathbf{x}|^{1-d})$ as $|\mathbf{x}| \rightarrow \infty$. Moreover, since $\omega \in C^1(\mathbb{R}^d)$ we conclude that

$$|\nabla\omega(\mathbf{x})| \leq \frac{C_{d,v}}{|\mathbf{x}|^{d-1} + 1}, \quad \forall x \in \mathbb{R}^d.$$

Finally, since ω is a minimizer for (47) we have

$$b(v) = \int_{\mathbb{R}^d} (2|\nabla\omega(\mathbf{x})|^2 + v(\mathbf{x})|1 - \omega(\mathbf{x})|^2) \, d\mathbf{x}.$$

On the other hand, from the scattering equation we have

$$\int_{\mathbb{R}^d} (2|\nabla\omega(\mathbf{x})|^2 + v(\mathbf{x})(\omega(\mathbf{x}) - 1)\omega(\mathbf{x})) \, d\mathbf{x} = 0.$$

Thus we can rewrite

$$b(v) = \int_{\mathbb{R}^d} (v(\mathbf{x})|1 - \omega(\mathbf{x})|^2 - v(\mathbf{x})(\omega(\mathbf{x}) - 1)\omega(\mathbf{x})) \, d\mathbf{x} = \int_{\mathbb{R}^d} v(\mathbf{x})(1 - \omega(\mathbf{x})) \, d\mathbf{x}.$$

Using the scattering equation $\omega = (-2\Delta + v)^{-1}v$ and the Hardy–Littlewood–Sobolev inequality, we can estimate

$$\begin{aligned} 0 \leq \int_{\mathbb{R}^d} v - b(v) &= \int_{\mathbb{R}^d} v\omega = \int_{\mathbb{R}^d} v(-2\Delta + v)^{-1}v \\ &\leq \frac{1}{2} \int_{\mathbb{R}^d} v(-\Delta)^{-1}v \leq C_d \|v\|_{L^{2d/(d+2)}}^2. \quad \square \end{aligned}$$

Remark 4. By repeatedly using scattering equation $\omega = (-2\Delta + v)^{-1}v$ and the resolvent formula we can write the scattering energy as a Born series expansion

$$b(v) = \int_{\mathbb{R}^d} v - \int_{\mathbb{R}^d} v(-2\Delta + v)^{-1}v = \int_{\mathbb{R}^d} v - \int_{\mathbb{R}^d} v(-2\Delta)^{-1}v + \dots$$

If $v \geq 0$ and $v \not\equiv 0$, then $b(v) < \int_{\mathbb{R}^d} v$ since $(-2\Delta + v)^{-1} > 0$ on $L^2(\mathbb{R}^d)$. \triangle

2.2. Three-body symmetry. Let $V : \mathbb{R}^6 \rightarrow \mathbb{R}_+$ satisfy the three-body symmetry (4). One can check that (4) is equivalent to

$$V(x, y) = V(y, x) = V(x - y, -y), \quad \forall (x, y) \in \mathbb{R}^3 \times \mathbb{R}^3. \quad (52)$$

Put differently, $V = V(g \cdot)$ with $g : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}^3 \times \mathbb{R}^3$ given by

$$S := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{or} \quad A := \begin{pmatrix} 1 & -1 \\ 0 & -1 \end{pmatrix}.$$

Note that both S and A are symmetries ($S^2 = A^2 = 1$) and $SAS = ASA$. Thus the group generated by A and S is finite, which is

$$\mathcal{G} = \{I, S, A, AS, SA, ASA\}. \quad (53)$$

In summary, the symmetry (4) is equivalent to the fact that $V = V(g \cdot)$ for all $g \in \mathcal{G}$, namely V is invariant under the actions of \mathcal{G} .

Next, let us consider the scattering problem associated to the three-body interaction potential $V(x - y, x - z)$. Consider the operator

$$-\Delta_{x_1} - \Delta_{x_2} - \Delta_{x_3} + V(x_1 - x_2, x_1 - x_3) \quad \text{on } L^2((\mathbb{R}^3)^3).$$

After removing the center of mass, we are left with the two-body operator

$$-2\Delta_{\mathcal{M}} + V(x, y) \quad \text{on } L^2((\mathbb{R}^3)^2)$$

where $-\Delta_{\mathcal{M}} = |\mathcal{M}\nabla_{\mathbb{R}^6}|^2$ with the matrix $\mathcal{M} : \mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}^3 \times \mathbb{R}^3$ given by

$$\mathcal{M} := \left(\frac{1}{2} \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \right)^{1/2} = \frac{1}{2\sqrt{2}} \begin{pmatrix} \sqrt{3} + 1 & \sqrt{3} - 1 \\ \sqrt{3} - 1 & \sqrt{3} + 1 \end{pmatrix}. \quad (54)$$

We define the *modified scattering energy*

$$b_{\mathcal{M}}(V) := \inf_{\varphi \in \dot{H}^1(\mathbb{R}^d)} \int_{\mathbb{R}^d} (2|\mathcal{M}\nabla\varphi(\mathbf{x})|^2 + V(\mathbf{x})|1 - \varphi(\mathbf{x})|^2) \, d\mathbf{x} \quad (55)$$

As we will see, by change of variables we can transfer the results in the previous section on the standard scattering energy $b(V)$ defined in (47) to understand $b_{\mathcal{M}}(V)$ in (55). To be precise, from Theorem 3 we have

Theorem 5 (Modified scattering solution). *Let $0 \leq V \in L^\infty(\mathbb{R}^6)$ be compactly supported and satisfy the symmetry (4). Then the variational problem (55) has a unique minimizer $\omega = (-2\Delta_{\mathcal{M}} + V)^{-1}V$. The function ω satisfies the symmetry (4) and it solves the modified scattering equation*

$$-2\Delta_{\mathcal{M}}\omega(\mathbf{x}) + V(\mathbf{x})(\omega(\mathbf{x}) - 1) = 0, \quad \forall \mathbf{x} \in \mathbb{R}^6$$

which is equivalent to

$$(-\Delta_x - \Delta_y - \Delta_z)\omega(x - y, x - z) + (V(\omega - 1))(x - y, x - z) = 0, \quad \forall x, y, z \in \mathbb{R}^3.$$

We have the pointwise estimates

$$0 \leq \omega(\mathbf{x}) < 1, \quad \omega(\mathbf{x}) \leq \frac{C_V}{|\mathbf{x}|^4 + 1}, \quad |\nabla\omega(\mathbf{x})| \leq \frac{C_V}{|\mathbf{x}|^5 + 1}, \quad \forall \mathbf{x} \in \mathbb{R}^6. \quad (56)$$

The modified scattering energy satisfies

$$b_{\mathcal{M}}(V) = \int_{\mathbb{R}^6} V(\mathbf{x})(1 - \omega(\mathbf{x})) \, d\mathbf{x}, \quad 0 \leq \int_{\mathbb{R}^6} V - b_{\mathcal{M}}(V) \leq C\|V\|_{L^{3/2}(\mathbb{R}^6)}^2. \quad (57)$$

Moreover $b_{\mathcal{M}}(V) = b(V(\mathcal{M}\cdot)) \det \mathcal{M}$ with b defined in (47). Here $\det \mathcal{M} = 3\sqrt{3}/8$.

Proof. Note that for every $\varphi \in \dot{H}^1(\mathbb{R}^6)$ we have

$$\begin{aligned} \int_{\mathbb{R}^6} 2|\mathcal{M}\nabla\varphi|^2 + V|1 - \varphi|^2 &= \int_{\mathbb{R}^6} (2|\mathcal{M}(\nabla\varphi)(\mathcal{M}\cdot)|^2 + V(\mathcal{M}\cdot)|1 - \varphi(\mathcal{M}\cdot)|^2) \det \mathcal{M} \\ &= \det \mathcal{M} \int_{\mathbb{R}^6} 2|\nabla(\varphi(\mathcal{M}\cdot))|^2 + V(\mathcal{M}\cdot)|1 - \varphi(\mathcal{M}\cdot)|^2. \end{aligned} \quad (58)$$

Moreover, it is obvious that $\varphi \in \dot{H}^1(\mathbb{R}^6)$ if and only if $\varphi(\mathcal{M}\cdot) \in \dot{H}^1(\mathbb{R}^6)$. Therefore,

$$\begin{aligned} b_{\mathcal{M}}(V) &= \inf_{\varphi \in \dot{H}^1(\mathbb{R}^d)} \int_{\mathbb{R}^d} 2|\nabla_{\mathcal{M}}\varphi|^2 + V|1 - \varphi|^2 \\ &= \det \mathcal{M} \inf_{\varphi \in \dot{H}^1(\mathbb{R}^d)} \int_{\mathbb{R}^6} 2|\nabla(\varphi(\mathcal{M}\cdot))|^2 + V(\mathcal{M}\cdot)|1 - \varphi(\mathcal{M}\cdot)|^2 \\ &= \det \mathcal{M} \inf_{\varphi \in \dot{H}^1(\mathbb{R}^d)} \int_{\mathbb{R}^6} 2|\nabla\varphi|^2 + V(\mathcal{M}\cdot)|1 - \varphi|^2 = b(V(\mathcal{M}\cdot)) \det \mathcal{M}. \end{aligned}$$

Thanks to (58), it is obvious that the minimizer of $b_{\mathcal{M}}(V)$ in (55) is $\omega = \tilde{\omega}(\mathcal{M}^{-1}\cdot)$ with $\tilde{\omega}$ the unique minimizer of $b(V(\mathcal{M}\cdot))$ defined in (47). Thus most of statements in Theorem 5 follow from Theorem 3. From the equation

$$-2\Delta_{\mathcal{M}}\omega(\mathbf{x}) + V(\mathbf{x})(\omega(\mathbf{x}) - 1) = 0, \quad \forall \mathbf{x} \in \mathbb{R}^6$$

we can also deduce that

$$(-\Delta_x - \Delta_y - \Delta_z)\omega(x - y, x - z) + (V(\omega - 1))(x - y, x - z) = 0, \quad \forall x, y, z \in \mathbb{R}^3.$$

by removing the center of mass similarly to (7).

Finally let us prove that ω is invariant under the actions of \mathcal{G} . Since $\omega = (-2\Delta_{\mathcal{M}} + V)^{-1}V$ and V is invariant under the actions of \mathcal{G} , it remains to check that $-\Delta_{\mathcal{M}}$ is also invariant under the actions of \mathcal{G} . For every $\varphi \in C_c^\infty(\mathbb{R}^6)$ and $g \in \mathcal{G}$, using

$$g^\dagger \mathcal{M}^2 g = g \mathcal{M}^2 g^\dagger = \mathcal{M}^2 \quad (59)$$

and $\det g = 1$ we have

$$\begin{aligned} \langle \varphi(g\cdot), -\Delta_{\mathcal{M}}\varphi(g\cdot) \rangle_{L^2(\mathbb{R}^6)} &= \int_{\mathbb{R}^6} |\mathcal{M}\nabla(\varphi(g\mathbf{x}))|^2 d\mathbf{x} = \int_{\mathbb{R}^6} |\mathcal{M}g^\dagger(\nabla\varphi)(g\mathbf{x})|^2 d\mathbf{x} \\ &= \int_{\mathbb{R}^6} |\mathcal{M}(\nabla\varphi)(g\mathbf{x})|^2 (\det g) d\mathbf{x} = \int_{\mathbb{R}^6} |\mathcal{M}(\nabla\varphi)(\mathbf{x})|^2 d\mathbf{x} \\ &= \langle \varphi, -\Delta_{\mathcal{M}}\varphi \rangle_{L^2(\mathbb{R}^6)}. \end{aligned}$$

Thus $\omega = \omega(g\cdot)$ for all $g \in \mathcal{G}$, and hence it satisfies the symmetry (4). \square

3. DYSON LEMMAS

3.1. Dyson lemma for non-radial potentials.

Theorem 6 (Dyson lemma for non-radial potentials). *Let $d \geq 3$. Let $0 \leq v \in L^\infty(\mathbb{R}^d)$ with $\text{Supp } v \subset \{|\mathbf{x}| \leq R_0\}$. Let $0 \leq U \in C(\mathbb{R}^d)$ be radial, $\text{Supp } U \subset \{R_1 \leq |\mathbf{x}| \leq R_2\}$ and $\int_{\mathbb{R}^d} U = 1$. Then we have the operator inequality*

$$-2\nabla_{\mathbf{x}} \mathbf{1}_{\{|\mathbf{x}| \leq R_2\}} \nabla_{\mathbf{x}} + v(\mathbf{x}) \geq b(v) \left(1 - \frac{C_d R_0}{R_1}\right) U(\mathbf{x}) \quad \text{on } L^2(\mathbb{R}^d)$$

with a constant $C_d > 0$ depending only on the dimension d .

Proof. If $R_1 < 2R_0$, then we can take $C_d = 2$ and the desired inequality holds trivially since the right-hand side is negative. Thus it remains to consider the case $R_1 \geq 2R_0$.

Let ω be the scattering solution to $b(v)$ as in Theorem 3. Then the function $f := 1 - \omega$ satisfies

$$0 < f \leq 1, \quad -2\Delta f + vf = 0 \quad \text{and} \quad \lim_{|\mathbf{x}| \rightarrow \infty} f(\mathbf{x}) = 1.$$

Now we take an arbitrary function $\varphi \in C_c^\infty(\mathbb{R}^d)$ and denote $\eta = \varphi/f$. For every $R \in [R_1, R_2]$, integrating by part and using the scattering equation we can bound

$$\begin{aligned} & \int_{B(0, R_2)} \left(2|\nabla\varphi|^2 + v|\varphi|^2 \right) \geq \int_{B(0, R)} \left(2|\nabla\varphi|^2 + v|\varphi|^2 \right) \\ &= \int_{B(0, R)} \left(2f^2|\nabla\eta|^2 + 2|\eta|^2|\nabla f|^2 + 2f\nabla f\nabla(|\eta|^2) + vf^2|\eta|^2 \right) \\ &= \int_{B(0, R)} 2f^2|\nabla\eta|^2 + \int_{B(0, R)} 2|\eta|^2f(-\Delta f + vf) + \int_{\partial B(0, R)} 2|\eta|^2f\nabla f \cdot \vec{n} \\ &= \int_{B(0, R)} 2f^2|\nabla\eta|^2 + \int_{\partial B(0, R)} 2|\varphi|^2f^{-1}\nabla f \cdot \vec{n} \geq 2 \int_{\partial B(0, R)} |\varphi|^2f^{-1}\nabla f \cdot \vec{n} \quad (60) \end{aligned}$$

where $\vec{n}_{\mathbf{x}} = \mathbf{x}/|\mathbf{x}|$ is the outward unit normal vector on the sphere $\partial B(0, R)$.

Now let us compute $f^{-1}\nabla f \cdot \vec{n}$ on the sphere $\partial B(0, R)$. Recall from (51) that

$$\nabla f(\mathbf{x}) \cdot \vec{n}_{\mathbf{x}} = \frac{1}{2|\mathbb{S}^{d-1}|} \int_{\mathbb{R}^d} v(\mathbf{y})f(\mathbf{y}) \frac{(\mathbf{x} - \mathbf{y})}{|\mathbf{x} - \mathbf{y}|^d} \cdot \vec{n}_{\mathbf{x}} \, d\mathbf{y}.$$

For every $|\mathbf{x}| \geq 2R_0$ and $|\mathbf{y}| \leq R_0$, by the triangle inequality we have

$$\left| \frac{\mathbf{x} - \mathbf{y}}{|\mathbf{x} - \mathbf{y}|^d} - \frac{\mathbf{x}}{|\mathbf{x}|^d} \right| \leq C_d \frac{R_0}{|\mathbf{x}|^d}, \quad \text{hence} \quad \frac{\mathbf{x} - \mathbf{y}}{|\mathbf{x} - \mathbf{y}|^d} \cdot \vec{n}_{\mathbf{x}} \geq \frac{1}{|\mathbf{x}|^{d-1}} \left(1 - C_d \frac{R_0}{|\mathbf{x}|} \right).$$

Since $\text{Supp } v \subset B(0, R_0)$ and $b(v) = \int_{\mathbb{R}^d} v f$, we have for all $|\mathbf{x}| \geq 2R_0$ that

$$\nabla f(\mathbf{x}) \cdot \vec{n}_{\mathbf{x}} \geq \frac{1}{2|\mathbb{S}^{d-1}|} \int_{\mathbb{R}^d} \frac{v(\mathbf{y})f(\mathbf{y})}{|\mathbf{x}|^{d-1}} \left(1 - C_d \frac{R_0}{|\mathbf{x}|} \right) \, d\mathbf{y} = \frac{b(v)}{2|\mathbb{S}^{d-1}||\mathbf{x}|^{d-1}} \left(1 - \frac{C_d R_0}{|\mathbf{x}|} \right). \quad (61)$$

Consequently, for every $R \in [R_1, R_2]$ such that $1 - C_d R_0/R \geq 0$, inserting (61) in (60) and using $f^{-1} \geq 1$ we get

$$\int_{B(0, R_2)} \left(2|\nabla\varphi|^2 + v|\varphi|^2 \right) \geq \frac{b(v)}{|\mathbb{S}^{d-1}|R^{d-1}} \left(1 - \frac{C_d R_0}{R} \right) \int_{\partial B(0, R)} |\varphi|^2. \quad (62)$$

On the other hand, if $1 - C_d R_0/R < 0$, then (62) holds trivially since the left-hand side is always nonnegative. Thus (62) holds true for all $R \in [R_1, R_2]$. Integrating both sides of (62) against $|\mathbb{S}^{d-1}|R^{d-1}U(R)$ with $R \in [R_1, R_2]$ and using $\int_{\mathbb{R}^d} U = 1$ for the left-hand side, we conclude that

$$\int_{B(0, R_2)} \left(2|\nabla\varphi|^2 + v|\varphi|^2 \right) \geq b(v) \left(1 - \frac{C_d R_0}{R_1} \right) \int_{\mathbb{R}^d} U |\varphi|^2.$$

Since the latter bound holds true for all $\varphi \in C_c^\infty(\mathbb{R}^d)$, we obtain the desired operator inequality. \square

3.2. Dyson lemma with the three-body symmetry. Now we have the following variant of Theorem 6 for the interaction potential with the three-body symmetry.

Theorem 7 (Dyson lemma with modified scattering energy). *Let $0 \leq V \in L^\infty(\mathbb{R}^6)$ be supported in $B(0, R_0)$ and satisfy the symmetry (4). Let $0 \leq \tilde{U} \in C(\mathbb{R}^6)$ be radial, $\text{Supp } \tilde{U} \subset \{R_1 \leq |\mathbf{x}| \leq R_2\}$ and $\int_{\mathbb{R}^6} \tilde{U} = 1$. Define*

$$U := \frac{1}{6} \sum_{g \in \mathcal{G}} \tilde{U}(\mathcal{M}^{-1}g) \det(\mathcal{M}^{-1}). \quad (63)$$

with \mathcal{G} and \mathcal{M} are given in (53) and (54). Then $0 \leq U \in C(\mathbb{R}^6)$, $\int_{\mathbb{R}^6} U = 1$, $\text{Supp } U \subset \{\sqrt{2/3}R_1 \leq |\mathbf{x}| \leq \sqrt{2}R_2\}$ and U satisfies the symmetry (4). Moreover, we have the operator inequality

$$-2\mathcal{M}\nabla_{\mathbf{x}}\mathbb{1}_{\{|\mathbf{x}| \leq \sqrt{2}R_2\}}\mathcal{M}\nabla_{\mathbf{x}} + V(\mathbf{x}) \geq b_{\mathcal{M}}(V) \left(1 - \frac{CR_0}{R_1}\right) U(\mathbf{x}) \quad \text{on } L^2(\mathbb{R}^6).$$

Here $C > 0$ is a universal constant (independent of V, U, R_0, R_1, R_2).

Proof. From the definition (63), it is clear that $U(\mathbf{x}) = U(g\mathbf{x})$ for all $g \in \mathcal{G}$. Thus U satisfies the symmetry (4). Using (59) and $\sqrt{1/2} \leq \mathcal{M} \leq \sqrt{3/2}$ we have

$$|\mathcal{M}^{-1}g\mathbf{x}| = |\mathcal{M}^{-1}\mathbf{x}| \in [\sqrt{2/3}|\mathbf{x}|, \sqrt{2}|\mathbf{x}|], \quad \forall \mathbf{x} \in \mathbb{R}^6.$$

Therefore, from the assumption $\text{Supp } \tilde{U} \subset \{R_1 \leq |\mathbf{x}| \leq R_2\}$ we deduce that

$$\text{Supp } U \subset \{\sqrt{2/3}R_1 \leq |\mathbf{x}| \leq \sqrt{2}R_2\}.$$

Moreover, $\int_{\mathbb{R}^6} U = \int_{\mathbb{R}^6} \tilde{U} = 1$ by change of variables and using $\det g = 1$ for $g \in \mathcal{G}$.

Next, we prove the operator inequality. We start by applying Theorem 6 for $V(\mathcal{M}\cdot)$. Note that $\text{Supp } V(\mathcal{M}\cdot) \subset B(0, \sqrt{3/2}R_0)$ since $\text{Supp } V \subset B(0, R_0)$ and $\sqrt{1/2} \leq \mathcal{M} \leq \sqrt{3/2}$. Hence, Theorem 6 gives

$$-2\nabla_{\mathbf{x}}\mathbb{1}_{\{|\mathbf{x}| \leq R_2\}}\nabla_{\mathbf{x}} + V(\mathcal{M}\mathbf{x}) \geq \left(1 - \frac{CR_0}{R_1}\right) b(V(\mathcal{M}\cdot))\tilde{U}(\mathbf{x}) \quad \text{on } L^2(\mathbb{R}^6).$$

Since $V = V(g\cdot)$ for $g \in \mathcal{G}$, the change of variable $\mathbf{x} = \mathcal{M}^{-1}g\mathbf{y}$ and (59) give

$$-2\mathcal{M}\nabla_{\mathbf{y}}\mathbb{1}_{\{|\mathcal{M}^{-1}\mathbf{y}| \leq R_2\}}\mathcal{M}\nabla_{\mathbf{y}} + V(\mathbf{y}) \geq \left(1 - \frac{CR_0}{R_1}\right) b(V(\mathcal{M}\cdot))\tilde{U}(\mathcal{M}^{-1}g\mathbf{y}) \quad \text{on } L^2(\mathbb{R}^6).$$

On the left-hand side we can use $\mathbb{1}_{\{|\mathcal{M}^{-1}\mathbf{y}| \leq R_2\}} \leq \mathbb{1}_{\{|\mathbf{y}| \leq \sqrt{2}R_2\}}$ as $|\mathcal{M}^{-1}\mathbf{y}| \geq \sqrt{1/2}|\mathbf{y}|$. On the right-hand side, we average over $g \in \mathcal{G}$ and use $b(V(\mathcal{M}\cdot)) = b_{\mathcal{M}}(V) \det \mathcal{M}^{-1}$ (see Theorem 5). The proof is therefore complete as it yields

$$-2\mathcal{M}\nabla_{\mathbf{y}}\mathbb{1}_{\{|\mathbf{y}| \leq \sqrt{2}R_2\}}\mathcal{M}\nabla_{\mathbf{y}} + V(\mathbf{y}) \geq \left(1 - \frac{CR_0}{R_1}\right) b_{\mathcal{M}}(V)U(\mathbf{y}) \quad \text{on } L^2(\mathbb{R}^6). \quad \square$$

3.3. Many-body Dyson lemma. We have the following many-body version of the Dyson lemma.

Lemma 8 (Many-body Dyson lemma). *Let $0 \leq W \in L^\infty(\mathbb{R}^6)$ be supported in $B(0, R_0)$ and satisfy the symmetry (4). Let $0 \leq \tilde{U} \in L^\infty(\mathbb{R}^6)$ be radial, supported*

in $\{1/8 \leq |\mathbf{x}| \leq 1/4\}$ and satisfy $\int_{\mathbb{R}^6} \tilde{U} = 1$. Define U as in (63) and $U_R = R^{-6}U(R^{-1}\cdot)$. Then for all $s > 0$ and $0 < \varepsilon < 1$ we have

$$\begin{aligned} \sum_{i=1}^M p_i^2 \mathbb{1}_{\{|p_i| > s\}} + \frac{1}{6} \sum_{\substack{1 \leq i, j, k \leq M \\ i \neq j \neq k \neq i}} W(x_i - x_j, x_i - x_k) \\ \geq \frac{b_{\mathcal{M}}(W)(1 - \varepsilon)}{6} \left(1 - \frac{CR_0}{R}\right) \\ \times \sum_{\substack{1 \leq i, j, k \leq M \\ i \neq j \neq k \neq i}} U(x_i - x_j, x_i - x_k) \prod_{\ell \neq i, j, k} \theta_{2R} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right) \\ - C\varepsilon^{-1} s^5 R^3 M^2. \end{aligned} \quad (64)$$

Here $C > 0$ is a universal constant (independent of $W, U, M, R, \varepsilon, s$).

Note that in (64) we only use the high-momentum part of the kinetic energy on the left-hand side. This is important for our application, since we need the low-momentum part to recover the NLS functional. If we use fully the kinetic energy, then the bound becomes simpler:

$$\begin{aligned} \sum_{i=1}^M p_i^2 + \frac{1}{6} \sum_{\substack{1 \leq i, j, k \leq M \\ i \neq j \neq k \neq i}} W(x_i - x_j, x_i - x_k) \\ \geq \frac{b_{\mathcal{M}}(W)}{6} \left(1 - \frac{CR_0}{R}\right) \\ \times \sum_{\substack{1 \leq i, j, k \leq M \\ i \neq j \neq k \neq i}} \tilde{U}(x_i - x_j, x_i - x_k) \prod_{\ell \neq i, j, k} \theta_{2R} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right). \end{aligned} \quad (65)$$

The latter bound follows from (64) by taking $s \rightarrow 0$ and then $\varepsilon \rightarrow 0$.

Proof. Denote $\chi_R = \mathbb{1}_{\{|x| \leq R\}} = 1 - \theta_R$ with $x \in \mathbb{R}^3$. For $(x_1, \dots, x_M) \in (\mathbb{R}^3)^M$ and $i, j, k \in \{1, 2, \dots, M\}$ with $i \neq j \neq k \neq i$, we denote

$$F_{ijk} := \chi_R(x_i - x_j) \chi_R(x_i - x_k) \chi_R(x_j - x_k) \prod_{\ell \neq i, j, k} \theta_{2R} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right). \quad (66)$$

Clearly $F_{ijk} \in \{0, 1\}$. Moreover, by the triangle inequality

$$F_{ijk} \leq \chi_R(x_i - x_j) \chi_R(x_i - x_k) \prod_{\ell \neq i, j, k} \theta_R(x_i - x_\ell).$$

Hence, for every $1 \leq i \leq M$, there are at most one pair j, k such that $F_{ijk} = F_{ikj} \neq 0$. Thus we have the ‘‘no four-body collision’’ bound

$$\sum_{\substack{1 \leq j, k \leq M \\ i \neq j \neq k \neq i}} F_{ijk} \leq 2. \quad (67)$$

Multiplying the above inequalities to the right by $p_i \mathbb{1}_{\{|p_i|>s\}}$ and to the left by $p_i \mathbb{1}_{\{|p_i|>s\}}$, where $p_i = -i\nabla_{x_i}$, then summing over i , we obtain

$$\sum_{i=1}^M p_i^2 \mathbb{1}_{\{|p_i|>s\}} \geq \frac{1}{2} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} \mathbb{1}_{\{|p_i|>s\}} p_i F_{ijk} p_i \mathbb{1}_{\{|p_i|>s\}}. \quad (68)$$

Let us now remove the momentum cut-off on the right-hand side of (68). By decomposing $\mathbb{1}_{\{|p_i|>s\}} = 1 - \mathbb{1}_{\{|p_i| \leq s\}}$ and using the Cauchy–Schwarz inequality, we obtain for all $0 < \varepsilon < 1$

$$\mathbb{1}_{\{|p_i|>s\}} p_i F_{ijk} p_i \mathbb{1}_{\{|p_i|>s\}} \geq (1 - \varepsilon) p_i F_{ijk} p_i - \varepsilon^{-1} \mathbb{1}_{\{|p_i| \leq s\}} p_i F_{ijk} p_i \mathbb{1}_{\{|p_i| \leq s\}}.$$

Hence, we deduce from (68) that

$$\sum_{i=1}^M p_i^2 \mathbb{1}_{\{|p_i|>s\}} \geq \frac{1 - \varepsilon}{2} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} p_i F_{ijk} p_i - \frac{\varepsilon^{-1}}{2} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} \mathbb{1}_{\{|p_i| \leq s\}} p_i F_{ijk} p_i \mathbb{1}_{\{|p_i| \leq s\}}. \quad (69)$$

For every $1 \leq i, j \leq M$ with $i \neq j$, we have

$$\sum_{\substack{1 \leq k \leq M \\ k \neq i,j}} F_{ijk} \leq \chi_R(x_i - x_j).$$

On the other hand, using

$$\|f(x)g(p)\|_{\mathfrak{S}^2(L^2(\mathbb{R}^d))} = (2\pi)^{-d/2} \|f\|_{L^2(\mathbb{R}^d)} \|g\|_{L^2(\mathbb{R}^d)},$$

where $\mathfrak{S}^2(L^2(\mathbb{R}^d))$ is the space of Hilbert–Schmidt operators, we find that

$$\begin{aligned} 0 &\leq \mathbb{1}_{\{|p_i| \leq s\}} p_i \mathbb{1}_{\{|x_i - x_j| < R\}} p_i \mathbb{1}_{\{|p_i| \leq s\}} \\ &\leq \|\mathbb{1}_{\{|p_i| \leq s\}} p_i \mathbb{1}_{\{|x_i - x_j| < R\}} p_i \mathbb{1}_{\{|p_i| \leq s\}}\| \\ &\leq \|\mathbb{1}_{\{|p_i| \leq s\}} p_i \mathbb{1}_{\{|x_i - x_j| < R\}}\|_{\mathfrak{S}^2(L^2(\mathbb{R}^3, dx_i))}^2 \\ &= (2\pi)^{-3} \|\mathbb{1}_{\{|p_i| \leq s\}} |p_i|\|_{L^2(\mathbb{R}^3, dp_i)}^2 \|\mathbb{1}_{\{|x_i - x_j| < R\}}\|_{L^2(\mathbb{R}^3, dx_i)}^2 \leq C s^5 R^3. \end{aligned}$$

Thus we can bound the last term on the right-hand side of (69) as

$$\begin{aligned} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} \mathbb{1}_{\{|p_i| \leq s\}} p_i F_{ijk} p_i \mathbb{1}_{\{|p_i| \leq s\}} &\leq \sum_{\substack{1 \leq i,j \leq M \\ i \neq j}} \mathbb{1}_{\{|p_i| \leq s\}} p_i \chi_R(x_i - x_j) p_i \mathbb{1}_{\{|p_i| \leq s\}} \\ &\leq C s^5 R^3 M^2. \end{aligned}$$

Hence, (69) reduces to

$$\sum_{i=1}^M p_i^2 \mathbb{1}_{\{|p_i|>s\}} \geq \frac{1 - \varepsilon}{2} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} p_n F_{ijk} p_n - C \varepsilon^{-1} s^5 R^3 M^2$$

which is equivalent to

$$\sum_{i=1}^M p_i^2 \mathbb{1}_{\{|p_i|>s\}} + \frac{1 - \varepsilon}{6} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} W(x_i - x_j, x_i - x_k)$$

$$\geq \frac{1-\varepsilon}{6} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} \left[\sum_{n \in \{i,j,k\}} p_n F_{ijk} p_n + W(x_i - x_j, x_i - x_k) \right] - C\varepsilon^{-1} s^5 R^3 M^2. \quad (70)$$

Now let us show that for any given (i, j, k) with $i \neq j \neq k \neq i$, we have

$$\begin{aligned} & \sum_{n \in \{i,j,k\}} p_n F_{ijk} p_n + W(x_i - x_j, x_i - x_k) \\ & \geq \left(1 - \frac{C}{M^{1/2}R}\right) b_{\mathcal{M}}(W) U(x_i - x_j, x_i - x_k) \prod_{\ell \neq i,j,k} \left(\frac{x_i + x_j + x_k}{3} - x_\ell\right). \end{aligned} \quad (71)$$

We do the change of variables similarly to (6):

$$r_i = \frac{1}{3}(x_i + x_j + x_k), \quad r_j = x_i - x_j, \quad r_k = x_i - x_k.$$

Then

$$p_{x_i} = \frac{1}{3}p_{r_i} + p_{r_j} + p_{r_k}, \quad p_{x_j} = \frac{1}{3}p_{r_i} - p_{r_j}, \quad p_{x_k} = \frac{1}{3}p_{r_i} - p_{r_k},$$

hence

$$\begin{aligned} \sum_{n \in \{i,j,k\}} p_n F_{ijk} p_n &= p_{x_i} F_{ijk} p_{x_i} + p_{x_j} F_{ijk} p_{x_j} + p_{x_k} F_{ijk} p_{x_k} \\ &= \left(\frac{1}{3}p_{r_i} + p_{r_j} + p_{r_k}\right) F_{ijk} \left(\frac{1}{3}p_{r_i} + p_{r_j} + p_{r_k}\right) \\ &\quad + \left(\frac{1}{3}p_{r_i} - p_{r_j}\right) F_{ijk} \left(\frac{1}{3}p_{r_i} - p_{r_j}\right) + \left(\frac{1}{3}p_{r_i} - p_{r_k}\right) F_{ijk} \left(\frac{1}{3}p_{r_i} - p_{r_k}\right) \\ &= \frac{1}{3}p_{r_i} F_{ijk} p_{r_i} + 2p_{r_j} F_{ijk} p_{r_j} + 2p_{r_k} F_{ijk} p_{r_k} + p_{r_j} F_{ijk} p_{r_k} + p_{r_k} F_{ijk} p_{r_j}. \end{aligned}$$

We can remove $p_{r_i} F_{ijk} p_{r_i} \geq 0$ for a lower bound. Moreover, by introducing the notation $\mathbf{r}_{jk} = (r_j, r_k) \in \mathbb{R}^6$ we have

$$F_{ijk} = \chi_R(r_j) \chi_R(r_k) \chi_R(r_j - r_k) \prod_{\ell \neq i,j,k} \theta_{2R}(r_i - x_\ell) \geq \mathbf{1}_{\{|\mathbf{r}_{jk}| \leq R/2\}} \prod_{\ell \neq i,j,k} \theta_{2R}(r_i - x_\ell).$$

Thus

$$\begin{aligned} & \sum_{n \in \{i,j,k\}} p_n F_{ijk} p_n \\ & \geq 2p_{r_j} F_{ijk} p_{r_j} + 2p_{r_k} F_{ijk} p_{r_k} + p_{r_j} F_{ijk} p_{r_k} + p_{r_k} F_{ijk} p_{r_j} = 2\mathcal{M} p_{\mathbf{r}_{ij}} F_{ijk} \mathcal{M} p_{\mathbf{r}_{ij}} \\ & \geq 2\mathcal{M} p_{\mathbf{r}_{ij}} \mathbf{1}_{\{|\mathbf{r}_{jk}| \leq R/2\}} \mathcal{M} p_{\mathbf{r}_{ij}} \theta_{2R}(r_i - x_\ell). \end{aligned}$$

Since $W \geq 0$, we have the obvious bound

$$W(x_i - x_j, x_i - x_k) = W(\mathbf{r}_{jk}) \geq W(\mathbf{r}_{jk}) \prod_{\ell \neq i,j,k} \theta_{2R}(r_i - x_\ell).$$

Using Theorem 7 we obtain

$$\begin{aligned} & \sum_{n \in \{i,j,k\}} p_n F_{ijk} p_n + W(x_i - x_j, x_i - x_k) \\ & \geq \left[2\mathcal{M} p_{\mathbf{r}_{ij}} \mathbf{1}_{\{|\mathbf{r}_{jk}| \leq R/2\}} \mathcal{M} p_{\mathbf{r}_{ij}} + W(\mathbf{r}_{jk})\right] \prod_{\ell \neq i,j,k} \theta_{2R}(r_i - x_\ell) \\ & \geq (1 - CR_0/R) b_{\mathcal{M}}(W) U_R(\mathbf{r}_{jk}) \theta_{2R}(r_i - x_\ell). \end{aligned}$$

Thus (71) holds true. This completes the proof of (64). \square

4. REDUCTION TO SOFTER INTERACTION POTENTIALS

Lemma 9 (Reduction to softer interaction potentials). *Let $s > 1 > \varepsilon, \beta > 0$. Let $0 \leq \tilde{U} \in L^\infty(\mathbb{R}^6)$ be radial, supported in $\{1/8 \leq |\mathbf{x}| \leq 1/4\}$ and satisfy $\int_{\mathbb{R}^6} \tilde{U} = 1$. Define U as in (63) and $U_R = R^{-6}U(R^{-1}\cdot)$. Then there exist $N \geq M \geq (1 - \varepsilon)N$ and $N^{-\beta/2} \geq R \geq N^{-\beta}$ such that*

$$E_N \geq \inf \sigma_{L^2_s(\mathbb{R}^{3M})} \left(\sum_{i=1}^M (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M^2} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} U_R(x_i - x_j, x_i - x_k) \right) - C_{\varepsilon,s,\beta} R^{2/7} - \varepsilon C_\beta$$

where $h_{\varepsilon,s} = \varepsilon p^2 + \mathbf{1}_{\{|p| \leq s\}} p^2 + V_{\text{ext}}(x)$, and $b_\varepsilon = b_{\mathcal{M}}(V)(1 - \varepsilon)^4$.

4.1. Binding inequality. Consider the Hamiltonian

$$H_{M,N} = \sum_{i=1}^M (-\Delta_{x_i} + V_{\text{ext}}(x_i)) + \sum_{1 \leq i < j < k \leq M} NV(N^{1/2}(x_i - x_j), N^{1/2}(x_i - x_k)) \quad (72)$$

and denote by $E(M, N)$ its ground state energy and $\Psi_{M,N}$ its ground state.

Lemma 10. *Let $1 > \varepsilon > 0$. Then there exists $M = M(N, \varepsilon)$ such that $N \geq M \geq (1 - \varepsilon)N$ and*

$$E(M, N) - E(M - 4, N) \leq C\varepsilon^{-1}. \quad (73)$$

Here $C > 0$ is a constant independent of N, ε .

Proof. Denote

$$Z := \inf_{m \in [(1-\varepsilon)N, N]} (E(m, N) - E(m - 4, N)).$$

Since $V_{\text{ext}}, V \geq 0$, $E(m, N)$ is nonnegative and increasing in m . Hence,

$$4E(N, N) \geq \sum_{(1-\varepsilon)N \leq m \leq N} (E(m, N) - E(m - 4, N)) \geq Z(\varepsilon N - 1).$$

Combined with the obvious upper bound $E(N, N) \leq C_0 N$, it gives $Z \leq C\varepsilon^{-1}$. \square

Remark 11. Using the convexity of the NLS functional and following the argument in [22], we can find $N \geq M \geq N + o(N)$ such that

$$E(M, N) - E(M - 4, N) \leq C$$

for all N . However, this stronger conclusion is not really needed for our purpose. \triangle

4.2. Four-body estimate.

Lemma 12. *Let $M = M(N, \varepsilon)$ be as in Lemma 10. Then the ground state $\Psi_{M,N}$ of $H_{M,N}$ satisfies*

$$\left\langle \prod_{i=2}^4 \mathbf{1}_{\{|x_1 - x_i| \leq R\}} \right\rangle_{\Psi_{M,N}} \leq C_\varepsilon R^9. \quad (74)$$

Proof. The technique is the same as in [22]. Using $V_{\text{ext}}, V \geq 0$ and the monotonicity of the heat kernel, we find that

$$\langle \xi(x_1, x_2, x_3, x_4) \rangle_{\Psi_{M,N}} \leq e^{(E(M,N) - E(M-4,N))} \left\| \xi \prod_{i=1}^4 e^{\Delta_{x_i}} \xi \right\|$$

for any measurable function $\xi(x_1, x_2, x_3, x_4) \geq 0$. Here we used the operator norm in $L^2(\mathbb{R}^{12})$ on the right side. In particular, applying this bound to

$$\xi(x_1, x_2, x_3, x_4) = \prod_{i=2}^4 \mathbb{1}_{\{|x_1 - x_i| \leq R\}}, \quad x_i \in \mathbb{R}^3$$

and using $E(M, N) - E(M - 4, N) \leq C\varepsilon^{-1}$ we find that

$$\begin{aligned} \left\langle \prod_{i=1}^3 \mathbb{1}_{\{|x_i - x_{i+1}| \leq R\}} \right\rangle_{\Psi_{M,N}} &\leq C_\varepsilon \left\| \prod_{i=2}^4 \mathbb{1}_{\{|x_1 - x_i| \leq R\}} \prod_{i=1}^4 e^{\Delta_{x_i}} \prod_{i=2}^4 \mathbb{1}_{\{|x_1 - x_i| \leq R\}} \right\| \\ &\leq C_\varepsilon \prod_{i=2}^4 \left\| \mathbb{1}_{\{|x_1 - x_i| \leq R\}} e^{-\frac{1}{2}p_i^2} \right\|_{\mathfrak{S}^2(L^2(\mathbb{R}^3, dx_i))}^2 \\ &= C_\varepsilon \prod_{i=2}^4 (2\pi)^{-3} \left\| \mathbb{1}_{\{|x_1 - x_i| \leq R\}} \right\|_{L^2(\mathbb{R}^3, dx_i)}^2 \left\| e^{-\frac{1}{2}p_i^2} \right\|_{L^2(\mathbb{R}^3, dp_i)}^2 \\ &\leq C_\varepsilon R^9. \quad \square \end{aligned}$$

4.3. Proof of Lemma 9. Now we are ready to conclude

Proof of Lemma 9. Step 1. Let $s > 1 > \varepsilon > 0$ be independent of N . Let $N \geq M \geq (1 - \varepsilon)N$ be as in Lemma 10 and let $\Psi_{M,N}$ be the ground state for $H_{M,N}$ in (72). Let

$$M^{-1/2} \ll R_0 \ll M^{-1/3}. \quad (75)$$

Applying Lemma 8 with $W = NV(N^{1/2} \cdot)$, which is supported in $B(0, CN^{-1/2})$ and has $b_{\mathcal{M}}(W) = b_{\mathcal{M}}(V)/N^2$, we have the operator inequality on $L_s^2(\mathbb{R}^{3M})$

$$\begin{aligned} \sum_{i=1}^M p_i^2 \mathbb{1}_{\{|p_i| > s\}} + \frac{1}{6} \sum_{\substack{1 \leq i, j, k \leq M \\ i \neq j \neq k \neq i}} NV(N^{1/2}(x_i - x_j), N^{1/2}(x_i - x_k)) \\ \geq \frac{b_{\mathcal{M}}(V)(1 - \varepsilon)}{6N^2} \left(1 - \frac{C}{N^{1/2}R_0} \right) \\ \times \sum_{\substack{1 \leq i, j, k \leq M \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \prod_{\ell \neq i, j, k} \theta_{2R_0} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right) \\ - C\varepsilon^{-1} s^5 R_0^3 M^2. \end{aligned}$$

Multiplying both sides with $(1 - \varepsilon)/M$ and using $N \geq M \geq (1 - \varepsilon)N$, we find that

$$\begin{aligned} \frac{H_{M,N}}{M(1 - \varepsilon)} &\geq \frac{b_\varepsilon}{6M^3} \left(1 - \frac{C}{M^{1/2}R_0} \right) \\ &\times \sum_{\substack{1 \leq i, j, k \leq M \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \prod_{\ell \neq i, j, k} \theta_{2R_0} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right) \end{aligned}$$

$$+ \frac{1}{M} \sum_{i=1}^M (h_{\varepsilon,s})_i - C_{\varepsilon,s} M R_0^3. \quad (76)$$

where $h_{\varepsilon,s} = \varepsilon p^2 + \mathbb{1}_{\{|p| \leq s\}} p^2 + V_{\text{ext}}(x)$ and $b_\varepsilon = b_{\mathcal{M}}(V)(1 - \varepsilon)^4$.

Next, we remove the four-body cut-off in the interaction term on the right-hand side of (76). Recall that $\chi_R(x) = \mathbb{1}_{\{|x| \leq R\}} = 1 - \theta_R(x)$. Using $\text{Supp } U_R \subset B(0, R)$ and Bernoulli's inequality, we have for every $1 \leq i, j, k \leq M$ with $i \neq j \neq k \neq i$,

$$\begin{aligned} 0 &\leq U_{R_0}(x_i - x_j, x_i - x_k) \left(1 - \prod_{\ell \neq i, j, k} \theta_{2R_0} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right) \right) \\ &\leq \frac{C}{R_0^6} \chi_{R_0}(x_i - x_j) \chi_{R_0}(x_i - x_k) \left(1 - \prod_{\ell \neq i, j, k} \theta_{2R_0} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right) \right) \\ &\leq \frac{C}{R_0^6} \chi_{R_0}(x_i - x_j) \chi_{R_0}(x_i - x_k) \left(1 - \prod_{\ell \neq i, j, k} \theta_{4R_0}(x_i - x_\ell) \right) \\ &= \frac{C}{R_0^6} \chi_{R_0}(x_i - x_j) \chi_{R_0}(x_i - x_k) \left(1 - \prod_{\ell \neq i, j, k} (1 - \chi_{4R_0}(x_i - x_\ell)) \right) \\ &\leq \frac{C}{R_0^6} \sum_{\ell \neq i, j, k} \chi_{4R_0}(x_i - x_j) \chi_{4R_0}(x_i - x_k) \chi_{4R_0}(x_i - x_\ell). \end{aligned}$$

Combining with Lemma 12, we find that

$$\begin{aligned} &\frac{1}{M^3} \sum_{\substack{1 \leq i, j, k \leq M \\ i \neq j \neq k \neq i}} \left\langle U_{R_0}(x_i - x_j, x_i - x_k) \left(1 - \prod_{\ell \neq i, j, k} \theta_{2R_0} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right) \right) \right\rangle_{\Psi_{M,N}} \\ &\leq \frac{C}{M^3 R_0^6} \sum_{\substack{1 \leq i, j, k \leq M \\ i \neq j \neq k \neq i}} \sum_{\ell \neq i, j, k} \left\langle \chi_{4R_0}(x_i - x_j) \chi_{4R_0}(x_i - x_k) \chi_{4R_0}(x_i - x_\ell) \right\rangle_{\Psi_{M,N}} \\ &\leq C_\varepsilon M R_0^3. \end{aligned}$$

Thus under the condition (75), we deduce from (76) that

$$\begin{aligned} \frac{E_{M,N}}{M(1 - \varepsilon)} &= \frac{1}{M(1 - \varepsilon)} \langle H_{M,N} \rangle_{\Psi_{M,N}} \\ &\geq \frac{1}{M} \left\langle \sum_{i=1}^M (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M^2} \left(1 - \frac{C}{M^{1/2} R_0} \right) \sum_{\substack{1 \leq i, j, k \leq M \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\ &\quad - C_{\varepsilon,s} M R_0^3. \end{aligned} \quad (77)$$

Combining with the upper bound $E_{M,N} \leq CM$ we get

$$\frac{b_\varepsilon}{6M^3} \left\langle \sum_{\substack{1 \leq i, j, k \leq M \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \leq C_{\varepsilon,s}. \quad (78)$$

Hence, (77) can be simplified to

$$\begin{aligned} \frac{E_{M,N}}{M(1-\varepsilon)} &\geq \frac{1}{M} \left\langle \sum_{i=1}^M (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M^2} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\ &\quad - C_{\varepsilon,s} \left(\frac{1}{M^{1/2}R_0} + MR_0^3 \right). \end{aligned}$$

The optimal choice of R_0 is determined by $1/(M^{1/2}R_0) = MR_0^3$, namely

$$R_0 = M^{-3/8}.$$

The condition (75) is clearly satisfied. With this choice, we have

$$\begin{aligned} \frac{E_{M,N}}{M(1-\varepsilon)} &\geq \frac{1}{M} \left\langle \sum_{i=1}^M (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M^2} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\ &\quad - C_{\varepsilon,s} M^{-1/8}. \quad (79) \end{aligned}$$

Step 2. We continue to replace the potential U_{R_0} by a softer one U_{R_1} . Unlike Step 1, now we can use only a small fraction of the kinetic energy. For that reason we will replace the particle number M by a smaller parameter M_1 in order to improve the error of removing the four-body cut-off. The parameter M_1 and R_1 will be chosen such that (recall that $R_0 = M^{-3/8}$)

$$M_1^{-1/3} \gg R_1 \gg R_0 \gg M_1^{-1/2}. \quad (80)$$

To be precise, by the bosonic symmetry of $\Psi_{M,N}$, the main term in (79) can be written as

$$\begin{aligned} &\frac{1}{M} \left\langle \sum_{i=1}^M (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M^2} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\ &= \frac{1}{M_1} \left\langle \sum_{i=1}^{M_1} (h_{\varepsilon,s})_i + \frac{b_\varepsilon(M-1)(M-2)}{6M^2(M_1-1)(M_1-2)} \sum_{\substack{1 \leq i,j,k \leq M_1 \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\ &\geq \frac{1}{M_1} \left\langle \sum_{i=1}^{M_1} (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M_1^2} \sum_{\substack{1 \leq i,j,k \leq M_1 \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}}. \quad (81) \end{aligned}$$

Next, we take a small parameter $\delta_\varepsilon \in (0, \varepsilon)$ and apply the Dyson lemma to the potential $W = \delta_\varepsilon b_\varepsilon M_1^{-2} U_{R_0}$, which is supported in $B(0, R_0)$ and has the scattering energy

$$b_{\mathcal{M}}(W) \geq \|W\|_{L^1(\mathbb{R}^6)} - C \|W\|_{L^{3/2}(\mathbb{R}^6)}^2 \geq \frac{b_\varepsilon}{\delta_\varepsilon M_1^2} \left(1 - \frac{C_\varepsilon}{M_1^2 R_0^4} \right).$$

Here we have used (57) in the latter estimate (the condition $R_0 \gg M_1^{-1/2}$ ensures that the scattering energy of W is well approximated by its first Born approximation). Thus from (65), after multiplying both sides with δ_ε , we have the operator inequality on $L_s^2(\mathbb{R}^{3M_1})$

$$\begin{aligned}
& \delta_\varepsilon \sum_{i=1}^{M_1} p_i^2 + \frac{b_\varepsilon}{6M_1^2} \sum_{\substack{1 \leq i,j,k \leq M_1 \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \\
& \geq \frac{b_\varepsilon}{6M_1^2} \left(1 - \frac{C_\varepsilon}{M_1^2 R_0^4}\right) \left(1 - \frac{CR_0}{R_1}\right) \\
& \quad \times \sum_{\substack{1 \leq i,j,k \leq M_1 \\ i \neq j \neq k \neq i}} U_{R_1}(x_i - x_j, x_i - x_k) \prod_{\ell \neq i,j,k} \theta_{2R_1} \left(\frac{x_i + x_j + x_k}{3} - x_\ell\right).
\end{aligned}$$

Thanks to Lemma 12, we can remove the four-body cut-off in the interaction term similarly to Step 1, namely

$$\begin{aligned}
& \frac{1}{M_1^3} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} \left\langle U_{R_1}(x_i - x_j, x_i - x_k) \left(1 - \prod_{\ell \neq i,j,k} \theta_{2R_1} \left(\frac{x_i + x_j + x_k}{3} - x_\ell\right)\right) \right\rangle_{\Psi_{M,N}} \\
& \leq \frac{C}{M_1^3 R_1^6} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} \sum_{\ell \neq i,j,k} \left\langle \chi_{4R_1}(x_i - x_j) \chi_{4R_1}(x_i - x_k) \chi_{4R_1}(x_i - x_\ell) \right\rangle_{\Psi_{M,N}} \\
& \leq C_\varepsilon M_1 R_1^3.
\end{aligned}$$

Hence,

$$\begin{aligned}
& \frac{1}{M_1} \left\langle \sum_{i=1}^{M_1} \delta_\varepsilon p_i^2 + \frac{b_\varepsilon}{6M_1^2} \sum_{\substack{1 \leq i,j,k \leq M_1 \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\
& \geq \frac{b_\varepsilon}{6M_1^3} \left(1 - \frac{C_\varepsilon}{M_1^2 R_0^4}\right) \left(1 - \frac{CR_0}{R_1}\right) \left\langle \sum_{\substack{1 \leq i,j,k \leq M_1 \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\
& \quad - C_\varepsilon M_1 R_1^3.
\end{aligned}$$

We can take $\delta_\varepsilon = \varepsilon^2$ and using $\delta_\varepsilon p^2 \leq \varepsilon h_{\varepsilon,s}$ for simplicity. Inserting the latter bound in (81), we deduce from (79) that

$$\begin{aligned}
\frac{E_{M,N}}{M(1-\varepsilon)^2} & \geq \frac{1}{M_1} \left\langle \sum_{i=1}^{M_1} (h_{\varepsilon,s})_i \right\rangle_{\Psi_{M,N}} \\
& + \frac{b_\varepsilon}{6M_1^2} \left(1 - \frac{C_\varepsilon}{M_1^2 R_0^4}\right) \left(1 - \frac{CR_0}{R_1}\right) \left\langle \sum_{\substack{1 \leq i,j,k \leq M_1 \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\
& \quad - C_{\varepsilon,s} M^{-1/8} - C_\varepsilon M_1 R_1^3. \quad (82)
\end{aligned}$$

Combined with the bound $E_{M,N} \leq CM$, it gives the following analogue of (78):

$$\frac{b_\varepsilon}{6M_1^3} \left\langle \sum_{\substack{1 \leq i,j,k \leq M_1 \\ i \neq j \neq k \neq i}} U_{R_1}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \leq C_{\varepsilon,s}.$$

Thus (82) can be simplified into

$$\begin{aligned} \frac{E_{M,N}}{M(1-\varepsilon)^2} &\geq \frac{1}{M_1} \left\langle \sum_{i=1}^{M_1} (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M_1^2} \sum_{\substack{1 \leq i,j,k \leq M_1 \\ i \neq j \neq k \neq i}} U_{R_0}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\ &\quad - C_{\varepsilon,s} M^{-1/8} - C_\varepsilon \left(M_1 R_1^3 + \frac{1}{M_1^2 R_0^4} + \frac{R_0}{R_1} \right). \end{aligned} \quad (83)$$

We can choose M_1 and R_1 such that

$$M_1 R_1^3 = \frac{1}{M_1^2 R_0^4} = \frac{R_0}{R_1} = \left((M_1 R_1^3)^2 \frac{1}{M_1^2 R_0^4} \left(\frac{R_0}{R_1} \right)^6 \right)^{1/9} = R_0^{2/9}$$

namely

$$R_1 = R_0^{7/9}, \quad M_1 = \frac{R_0^{2/9}}{R_1^3} = R_1^{-19/7}.$$

The condition (80) is clearly satisfied. We can take $\delta_\varepsilon = \varepsilon^2/2$ for simplicity and arrive at

$$\begin{aligned} \frac{E_{M,N}}{M(1-\varepsilon)^2} &\geq \frac{1}{M_1} \left\langle \sum_{i=1}^{M_1} (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M_1^2} \sum_{\substack{1 \leq i,j,k \leq M_1 \\ i \neq j \neq k \neq i}} U_{R_1}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\ &\quad - C_{\varepsilon,s} R_1^{2/7}. \end{aligned} \quad (84)$$

Step 3. We can iterate the argument in Step 2 to reach softer potentials. Denote

$$R_j = R_{j-1}^{2/9}, \quad M_j = R_j^{-19/7}, \quad \forall j \geq 1. \quad (85)$$

Then by induction, we can prove that for every $J \geq 1$,

$$\begin{aligned} \frac{E_{M,N}}{M(1-\varepsilon)^{J+1}} &\geq \frac{1}{M_J} \left\langle \sum_{i=1}^{M_J} (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M_J^2} \sum_{\substack{1 \leq i,j,k \leq M_J \\ i \neq j \neq k \neq i}} U_{R_J}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\ &\quad - C_{\varepsilon,s,J} R_J^{2/7}. \end{aligned} \quad (86)$$

Indeed, the case $J = 1$ has been handled in Step 2. The general case is very similar and we only mention some main estimates for the reader's convenience. Assuming that (86) holds for $J - 1$, by the bosonic symmetry we can write

$$\begin{aligned} &\frac{E_{M,N}}{M(1-\varepsilon)^J} \\ &\geq \frac{1}{M_{J-1}} \left\langle \sum_{i=1}^{M_{J-1}} (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M_{J-1}^2} \sum_{\substack{1 \leq i,j,k \leq M_{J-1} \\ i \neq j \neq k \neq i}} U_{R_{J-1}}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} - C_{\varepsilon,s} R_{J-1}^{2/7} \\ &\geq \frac{1}{M_J} \left\langle \sum_{i=1}^{M_J} (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M_J^2} \sum_{\substack{1 \leq i,j,k \leq M_J \\ i \neq j \neq k \neq i}} U_{R_{J-1}}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} - C_{\varepsilon,s} R_{J-1}^{2/7}. \end{aligned} \quad (87)$$

Applying the Dyson lemma to $W = \delta_\varepsilon b_\varepsilon M_J^{-2} U_{R_{J-1}}$, we deduce from (65) that

$$\begin{aligned} & \delta_\varepsilon \sum_{i=1}^{M_J} p_i^2 + \frac{b_\varepsilon}{6M_J^2} \sum_{\substack{1 \leq i,j,k \leq M_J \\ i \neq j \neq k \neq i}} U_{R_{J-1}}(x_i - x_j, x_i - x_k) \\ & \geq \frac{b_\varepsilon}{6M_J^2} \left(1 - \frac{C_\varepsilon}{M_J^2 R_{J-1}^4}\right) \left(1 - \frac{C R_{J-1}}{R_J}\right) \\ & \quad \times \sum_{\substack{1 \leq i,j,k \leq M_J \\ i \neq j \neq k \neq i}} U_{R_J}(x_i - x_j, x_i - x_k) \prod_{\ell \neq i,j,k} \theta_{2R_J} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right). \end{aligned}$$

The four-body cut-off can be removed by Lemma 12, leading to

$$\begin{aligned} & \frac{1}{M_J^3} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} \left\langle U_{R_J}(x_i - x_j, x_i - x_k) \left(1 - \prod_{\ell \neq i,j,k} \theta_{2R_J} \left(\frac{x_i + x_j + x_k}{3} - x_\ell \right)\right) \right\rangle_{\Psi_{M,N}} \\ & \leq \frac{C}{M_1^3 R_1^6} \sum_{\substack{1 \leq i,j,k \leq M_J \\ i \neq j \neq k \neq i}} \sum_{\ell \neq i,j,k} \left\langle \chi_{4R_1}(x_i - x_j) \chi_{4R_1}(x_i - x_k) \chi_{4R_J}(x_i - x_\ell) \right\rangle_{\Psi_{M,N}} \\ & \leq C_\varepsilon M_J R_J^3. \end{aligned}$$

Moreover, combining with the upper bound $E_{M,N} \leq CM$, we get

$$\frac{b_\varepsilon}{6M_J^3} \left\langle \sum_{\substack{1 \leq i,j,k \leq M_J \\ i \neq j \neq k \neq i}} U_{R_J}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \leq C_{\varepsilon,s}. \quad (88)$$

Hence,

$$\begin{aligned} & \frac{1}{M_J} \left\langle \delta_\varepsilon \sum_{i=1}^{M_J} p_i^2 + \frac{b_\varepsilon}{6M_J^2} \sum_{\substack{1 \leq i,j,k \leq M_J \\ i \neq j \neq k \neq i}} U_{R_{J-1}}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\ & \geq \frac{b_\varepsilon}{6M_J^3} \left\langle \sum_{\substack{1 \leq i,j,k \leq M_J \\ i \neq j \neq k \neq i}} U_{R_J}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\ & \quad - C_{\varepsilon,s} \left(M_J R_J^3 + \frac{C R_{J-1}}{R_J} + \frac{1}{M_J^2 R_{J-1}^4} \right). \end{aligned}$$

Thus from (87), by choosing $\delta_\varepsilon = \varepsilon^2$ and using $\delta_\varepsilon p^2 \leq \varepsilon h_{\varepsilon,s}$, we obtain

$$\begin{aligned} \frac{E_{M,N}}{M(1-\varepsilon)^{J+1}} & \geq \frac{1}{M_J} \left\langle \sum_{i=1}^{M_J-1} (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M_J^2} \sum_{\substack{1 \leq i,j,k \leq M_J \\ i \neq j \neq k \neq i}} U_{R_J}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\ & \quad - C_{\varepsilon,s} R_{J-1}^{2/7} - C_{\varepsilon,s} \left(M_J R_J^3 + \frac{1}{M_J^2 R_{J-1}^4} + \frac{R_{J-1}}{R_J} \right). \quad (89) \end{aligned}$$

With the choice in (85), we have

$$M_J R_J^3 = \frac{1}{M_J^2 R_{J-1}^4} = \frac{R_{J-1}}{R_J} = R_{J-1}^{2/9} = R_J^{2/7}$$

and the desired estimate (86) follows.

Step 4. To conclude, we choose $J \geq 1$ depending only on β such that

$$\frac{\beta}{2} < \frac{3}{8} \left(\frac{7}{9} \right)^J \leq \beta.$$

Then

$$N^{-\beta/2} \gg R_J = (R_0)^{\left(\frac{7}{9}\right)^J} = M^{-\frac{3}{8}\left(\frac{7}{9}\right)^J} \geq N^{-\beta}.$$

From (86) and (88) we have

$$\begin{aligned} & \frac{E_{M,N}}{M(1-\varepsilon)^{J+1}} \\ & \geq \frac{1}{M_J} \left\langle \sum_{i=1}^{M_J} (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M_J^2} \sum_{\substack{1 \leq i,j,k \leq M_J \\ i \neq j \neq k \neq i}} U_{R_J}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} - C_{\varepsilon,s} R_J^{2/7} \\ & = \frac{1}{M} \left\langle \sum_{i=1}^M (h_{\varepsilon,s})_i + \frac{b_\varepsilon (M_J - 1)(M_J - 2)}{6M_J^2 (M - 1)(M - 2)} \sum_{\substack{1 \leq i,j,k \leq M_J \\ i \neq j \neq k \neq i}} U_{R_J}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\ & \quad - C_{\varepsilon,s} R_J^{2/7} \\ & = \frac{1}{M} \left\langle \sum_{i=1}^M (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M^2} \sum_{\substack{1 \leq i,j,k \leq M_J \\ i \neq j \neq k \neq i}} U_{R_J}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} - C_{\varepsilon,s} R_J^{2/7} \\ & \quad + \frac{b_\varepsilon}{6M^3} \left(\frac{(M_J - 1)(M_J - 2)M^2}{(M - 1)(M - 2)M_J^2} - 1 \right) \left\langle \sum_{\substack{1 \leq i,j,k \leq M_J \\ i \neq j \neq k \neq i}} U_{R_J}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} \\ & \geq \frac{1}{M} \left\langle \sum_{i=1}^M (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M^2} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} U_{R_J}(x_i - x_j, x_i - x_k) \right\rangle_{\Psi_{M,N}} - C_{\varepsilon,s} R_J^{2/7}. \end{aligned}$$

In the last estimate we have used the bosonic symmetry for the main term, and the choice $M_J = R_J^{-19/7}$ for the error term. Since $N \geq M \geq N(1-\varepsilon)$ and $CN \geq E_N \geq E_{M,N}$, we have the obvious bound

$$\frac{E_{M,N}}{M(1-\varepsilon)^{J+1}} \leq \frac{E_N}{N(1-\varepsilon)^{J+2}} \leq \frac{E_N}{N} + \varepsilon C_J.$$

The desired conclusion of Lemma 9 follows. \square

5. CONCLUSION OF ENERGY LOWER BOUND

This section is devoted to the proof of the following theorem.

Theorem 13 (Energy lower bound). *Suppose $L_{\text{loc}}^\infty(\mathbb{R}^3) \ni V_{\text{ext}}(x) \geq c_0|x|^\alpha$ for some $c_0, \alpha > 0$. Then*

$$\lim_{N \rightarrow \infty} \frac{E_N}{N} \geq e_{\text{GP}} = \inf_{\|u\|_2=1} \mathcal{E}_{\text{GP}}(u) = \inf_{\|u\|_2=1} \int_{\mathbb{R}^3} \left(|\nabla u|^2 + V_{\text{ext}}|u|^2 + \frac{b_{\mathcal{M}}(V)}{6}|u|^6 \right).$$

Let $s > 1 > \varepsilon > 0$. Note that

$$h_{\varepsilon,s} := \varepsilon p^2 + \mathbf{1}_{\{|p| \leq s\}} p^2 + V_{\text{ext}}(x) \geq \varepsilon p^2 + c_0 |x|^\alpha \quad \text{on } L^2(\mathbb{R}^3).$$

Thanks to the Lieb–Thirring inequality in [12, Theorem 1], we have

$$\text{Tr}((-\Delta + |x|^\alpha + 1)^{-q}) \leq \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{1}{(|2\pi k|^2 + |x|^\alpha + 1)^q} dk dx < \infty$$

for any $q > q_*(\alpha) := \frac{3}{2} + \frac{3}{\alpha}$. Thus $\text{Tr}(h_{\varepsilon,s}^{-q}) \leq C_\varepsilon$. Therefore, if we introduce the projections

$$P = \mathbf{1}(h_{\varepsilon,s} \leq L), \quad Q = 1 - P = \mathbf{1}(h_{\varepsilon,s} > L)$$

for some parameter $L > 0$, then we have

$$\text{Tr} P = \text{Tr} \mathbf{1}(h_{\varepsilon,s} \leq L) \leq \text{Tr} \left(\frac{L^q}{h_{\varepsilon,s}^q} \right) \leq C_\varepsilon L^q. \quad (90)$$

Let $\beta > 0$ small (depending on q). From Lemma 9, we consider the Hamiltonian

$$\tilde{H}_M := \sum_{i=1}^M (h_{\varepsilon,s})_i + \frac{b_\varepsilon}{6M^2} \sum_{\substack{1 \leq i,j,k \leq M \\ i \neq j \neq k \neq i}} U_R(x_i - x_j, x_i - x_k) \quad (91)$$

for some $N \geq M \geq (1-\varepsilon)N$ and $N^{-\beta/2} \geq R \geq N^{-\beta}$. Recall $b_\varepsilon = b_{\mathcal{M}}(V)(1-\varepsilon)^4$ and $U_R = R^{-6}U(R^{-1}\cdot)$ with a fixed function $0 \leq U \in L^\infty(\mathbb{R}^6)$ satisfying the three-body symmetry (4) and $\int U = 1$.

Let Ψ_M be a ground state for \tilde{H}_M . Its k -body density matrix is an operator

$$\gamma_M^{(k)} := \text{Tr}_{k+1 \rightarrow M}[\gamma_M]$$

on $L_s^2(\mathbb{R}^{3k})$ with kernel

$$\begin{aligned} \gamma_M^{(k)}(x_1, \dots, x_k; y_1, \dots, y_k) &= \int_{\mathbb{R}^{3(M-k)}} \Psi_M(x_1, \dots, x_k, x_{k+1}, \dots, x_M) \times \dots \\ &\quad \dots \times \overline{\Psi_M(y_1, \dots, y_k, x_{k+1}, \dots, x_M)} dx_{k+1} \dots dx_M. \end{aligned}$$

Thus $\gamma_M^{(k)} \geq 0$ and $\text{Tr} \gamma_M^{(k)} = 1$. Using this notation, we can write

$$\frac{\tilde{E}_M}{M} := \frac{1}{M} \inf \sigma_{L_s^2(\mathbb{R}^{3M})} \tilde{H}_M = \frac{\langle \Psi_M, \tilde{H}_M \Psi_M \rangle}{M} = \text{Tr} [h_{\varepsilon,s} \gamma_M^{(1)}] + \frac{b_\varepsilon}{6} \text{Tr} [U_R \gamma_M^{(3)}].$$

Notice that, since $U_R \geq 0$ and $\gamma_M^{(3)} \geq 0$, this implies in particular that

$$\text{Tr} [h_{\varepsilon,s} \gamma_M^{(1)}] \leq \frac{\tilde{E}_M}{M} < +\infty. \quad (92)$$

Now let us impose the finite dimensional cut-off P and use the quantitative quantum de Finetti theorem. We bound from below the one-body term as follows

$$\begin{aligned} 3 \text{Tr} [h_{\varepsilon,s} \gamma_M^{(1)}] &= \text{Tr} [(h_1 + h_2 + h_3) \gamma_M^{(3)}] \\ &\geq \text{Tr} \left[\left(\sum_{i=1}^3 h_i \right) P^{\otimes 3} \gamma_M^{(3)} P^{\otimes 3} \right] \\ &= \text{Tr} \left[\left(\sum_{i=1}^3 h_i \right) \tilde{\gamma}_M^{(3)} \right] + \text{Tr} \left[\left(\sum_{i=1}^3 h_i \right) (P^{\otimes 3} \gamma_M^{(3)} P^{\otimes 3} - \tilde{\gamma}_M^{(3)}) \right] \end{aligned}$$

$$\geq \frac{1}{1+\varepsilon} \operatorname{Tr} \left[\left(\sum_{i=1}^3 h_i \right) \tilde{\gamma}_M^{(3)} \right] + \operatorname{Tr} \left[\left(\sum_{i=1}^3 h_i \right) \left(P^{\otimes 3} \gamma_M^{(3)} P^{\otimes 3} - \tilde{\gamma}_M^{(3)} \right) \right] \quad (93)$$

for a $\tilde{\gamma}^{(3)} = P^{\otimes 3} \tilde{\gamma}^{(3)} P^{\otimes 3}$ that will be chosen later (for the Quantum de Finetti theorem) and where we used the shortened notation $h_i := (h_{\varepsilon, s})_i$.

For the three-body term, we apply the Cauchy–Schwarz inequality, with $0 < \varepsilon < 1$, as follows

$$\begin{aligned} U_R &= (\mathbb{1}^{\otimes 3} - P^{\otimes 3} + P^{\otimes 3}) U_R (\mathbb{1}^{\otimes 3} - P^{\otimes 3} + P^{\otimes 3}) \\ &= P^{\otimes 3} U_R P^{\otimes 3} + (\mathbb{1}^{\otimes 3} - P^{\otimes 3}) U_R P^{\otimes 3} + P^{\otimes 3} U_R (\mathbb{1}^{\otimes 3} - P^{\otimes 3}) \\ &\quad + (\mathbb{1}^{\otimes 3} - P^{\otimes 3}) U_R (\mathbb{1}^{\otimes 3} - P^{\otimes 3}) \\ &\geq \left(1 - \frac{\varepsilon}{1+\varepsilon}\right) P^{\otimes 3} U_R P^{\otimes 3} + \left(1 - \frac{1+\varepsilon}{\varepsilon}\right) (\mathbb{1}^{\otimes 3} - P^{\otimes 3}) U_R (\mathbb{1}^{\otimes 3} - P^{\otimes 3}) \\ &\geq \frac{1}{1+\varepsilon} P^{\otimes 3} U_R P^{\otimes 3} - \varepsilon^{-1} \frac{C}{R^6} (\mathbb{1}^{\otimes 3} - P^{\otimes 3}), \end{aligned}$$

where we used for the second inequality that

$$(\mathbb{1}^{\otimes 3} - P^{\otimes 3}) U_R (\mathbb{1}^{\otimes 3} - P^{\otimes 3}) \leq \|U_R\|_{\infty} (\mathbb{1}^{\otimes 3} - P^{\otimes 3})^2 \leq \frac{C}{R^6} (\mathbb{1}^{\otimes 3} - P^{\otimes 3}).$$

Moreover, since

$$\begin{aligned} \mathbb{1}^{\otimes 3} &= Q \otimes \mathbb{1}^{\otimes 2} + P \otimes \mathbb{1}^{\otimes 2} =: Q_1 + P \otimes \mathbb{1}^{\otimes 2} \\ &= Q_1 + P \otimes Q \otimes \mathbb{1} + P \otimes P \otimes Q + P^{\otimes 3} \leq Q_1 + Q_2 + Q_3 + P^{\otimes 3}, \end{aligned}$$

we have

$$U_R \geq \frac{1}{1+\varepsilon} P^{\otimes 3} U_R P^{\otimes 3} - \varepsilon^{-1} \frac{C}{R^6} (Q_1 + Q_2 + Q_3).$$

Using now

$$\begin{aligned} \operatorname{Tr} \left[P^{\otimes 3} U_R P^{\otimes 3} \gamma_M^{(3)} \right] &= \operatorname{Tr} \left[U_R P^{\otimes 3} \gamma_M^{(3)} P^{\otimes 3} \right] \\ &= \operatorname{Tr} \left[U_R \tilde{\gamma}_M^{(3)} \right] + \operatorname{Tr} \left[U_R \left(P^{\otimes 3} \gamma_M^{(3)} P^{\otimes 3} - \tilde{\gamma}_M^{(3)} \right) \right] \end{aligned}$$

and

$$\operatorname{Tr} \left[(Q_1 + Q_2 + Q_3) \gamma_M^{(3)} \right] = 3 \operatorname{Tr} \left[Q \gamma_M^{(1)} \right] \leq 3 \operatorname{Tr} \left[\frac{h_{\varepsilon, s}}{L} \gamma_M^{(1)} \right] \leq \frac{3}{L} \frac{\tilde{E}_M}{M}$$

by (92), we obtain

$$\operatorname{Tr} \left[U_R \gamma_M^{(3)} \right] \geq \frac{1}{1+\varepsilon} \operatorname{Tr} \left[U_R \tilde{\gamma}_M^{(3)} \right] + \frac{1}{1+\varepsilon} \operatorname{Tr} \left[U_R \left(P^{\otimes 3} \gamma_M^{(3)} P^{\otimes 3} - \tilde{\gamma}_M^{(3)} \right) \right] - \frac{3C}{\varepsilon L R^6} \frac{\tilde{E}_M}{M}. \quad (94)$$

Combining (93) and (94), we have

$$\begin{aligned} \frac{\tilde{E}_M}{M} &\geq \frac{1}{1+\varepsilon} \operatorname{Tr} \left[\left(\frac{h_1 + h_2 + h_3}{3} + \frac{b_{\varepsilon}}{6} U_R \right) \tilde{\gamma}_M^{(3)} \right] \\ &\quad + \frac{1}{3} \operatorname{Tr} \left[\left(\sum_{i=1}^3 h_i \right) \left(P^{\otimes 3} \gamma_M^{(3)} P^{\otimes 3} - \tilde{\gamma}_M^{(3)} \right) \right] \\ &\quad + \frac{1}{1+\varepsilon} \frac{b_{\varepsilon}}{6} \operatorname{Tr} \left[U_R \left(P^{\otimes 3} \gamma_M^{(3)} P^{\otimes 3} - \tilde{\gamma}_M^{(3)} \right) \right] - \frac{C}{2} \frac{b_{\varepsilon}}{\varepsilon L R^6} \frac{\tilde{E}_M}{M} \end{aligned}$$

hence

$$\begin{aligned} \left(1 + \frac{C}{2} \frac{b_\varepsilon}{\varepsilon L R^6}\right) \frac{\tilde{E}_M}{M} &\geq \frac{1}{1+\varepsilon} \operatorname{Tr} \left[\left(\frac{h_1 + h_2 + h_3}{3} + \frac{b_\varepsilon}{6} U_R \right) \tilde{\gamma}_M^{(3)} \right] \\ &- \left(\frac{\|P^{\otimes 3}(h_1 + h_2 + h_3)P^{\otimes 3}\|}{3} + \frac{b_\varepsilon}{1+\varepsilon} \frac{C}{6R^6} \right) \operatorname{Tr} \left| P^{\otimes 3} \gamma_M^{(3)} P^{\otimes 3} - \tilde{\gamma}_M^{(3)} \right|. \end{aligned}$$

In order to deal with the term $\operatorname{Tr} \left| P^{\otimes 3} \gamma_M^{(3)} P^{\otimes 3} - \tilde{\gamma}_M^{(3)} \right|$ we can now use the Quantum de Finetti Theorem. We recall its formulation [18, Theorem 3.1] for the convenience of the reader.

Theorem 14 (Quantitative quantum de Finetti in finite dimension). *Let \mathcal{K} be a finite dimensional Hilbert space and $k \in \mathbb{N}$. For every state G_k on $\mathcal{K}^k := \bigotimes_s^k \mathcal{K}$ and for every $p = 1, 2, \dots, k$ we have*

$$\operatorname{Tr}_{\mathcal{K}^p} \left| \operatorname{Tr}_{p+1 \rightarrow k} [G_k] - \int_{SK} |u^{\otimes p}\rangle \langle u^{\otimes p}| d\mu_{G_k}(u) \right| \leq \frac{4p \dim \mathcal{K}}{k} \operatorname{Tr} [G_k] \quad (95)$$

where

$$d\mu_{G_k}(u) := \dim \mathcal{K}^k \langle u^{\otimes k}, G_k u^{\otimes k} \rangle du \quad (96)$$

with du being the normalized uniform (Haar) measure on the unit sphere SK .

Following exactly [18, Lemma 3.4], which one deals with two-body density matrices, we derive now a localized version for the three-body density matrices.

Let $\mathfrak{H} := L^2(\mathbb{R}^3)$ and projections $P_- := P$ and $P_+ := Q$. Let γ_N be an arbitrary N -body (mixed) state. Then there exist localized states $G_N^- = G_N$ and $G_N^+ = G_N^\perp$ (we use both notations) in the Fock space $\mathcal{F}(\mathfrak{H}) = \mathbb{C} \oplus \mathfrak{H} \oplus \mathfrak{H}^2 \oplus \dots$ of the form

$$G_N^\pm = G_{N,0}^\pm \oplus G_{N,1}^\pm \oplus \dots \oplus G_{N,N}^\pm \oplus 0 \oplus \dots \quad (97)$$

whose reduced density matrices satisfy, for any $0 \leq n \leq N$,

$$P_\pm^{\otimes n} \gamma_N^{(n)} P_\pm^{\otimes n} = (G_N^\pm)^{(n)} = \binom{N}{n}^{-1} \sum_{k=n}^N \binom{k}{n} \operatorname{Tr}_{n+1 \rightarrow k} [G_{N,k}^\pm]. \quad (98)$$

As reminded in the aforementioned paper, the relations (98) determine uniquely the localized states G_N and G_N^\perp and ensure that they are (mixed) states on the Fock spaces $\mathcal{F}(P\mathfrak{H})$ and $\mathcal{F}(Q\mathfrak{H})$, respectively:

$$\sum_{k=0}^N \operatorname{Tr} [G_{N,k}] = \sum_{k=0}^N \operatorname{Tr} [G_{N,k}^\perp] = 1. \quad (99)$$

We now apply the quantitative de Finetti Theorem 14 to $\mathcal{K} = P\mathfrak{H}$, $G_k = G_{N,k}$ and $p = 3$. We obtain the following

Lemma 15 (Quantitative quantum de Finetti for the localized state). *Let γ_N be an arbitrary N -body (mixed) state. We have*

$$\operatorname{Tr}_{\mathfrak{H}^3} \left| P^{\otimes 3} \gamma_N^{(3)} P^{\otimes 3} - \int_{SP\mathfrak{H}} |u^{\otimes 3}\rangle \langle u^{\otimes 3}| d\mu_N(u) \right| \leq \frac{12 \operatorname{Tr} P}{N}, \quad (100)$$

where

$$d\mu_N(u) = \sum_{k=3}^N \frac{k!(N-3)!}{N!(k-3)!} d\mu_{N,k}(u), \quad d\mu_{N,k}(u) = \dim(P\mathfrak{H})_s^k \langle u^{\otimes k}, G_{N,k} u^{\otimes k} \rangle du. \quad (101)$$

Proof. The proof follows the lines of the one of [18, Lemma 3.4]. Applying Theorem 14 to $\mathcal{K} = P\mathfrak{H}$, $G_k = G_{N,k}$ and $p = 3$, we have

$$\begin{aligned} \mathrm{Tr}_{(P\mathfrak{H})^3} \left| \mathrm{Tr}_{4 \rightarrow k}[G_{N,k}] - \int_{SP\mathfrak{H}} |u^{\otimes 3}\rangle\langle u^{\otimes 3}| d\mu_{N,k}(u) \right| \\ \leq \frac{12 \dim(P\mathfrak{H})}{k} \mathrm{Tr}[G_{N,k}] = \frac{12 \mathrm{Tr} P}{k} \mathrm{Tr}[G_{N,k}] \end{aligned}$$

where we notice that on the left-hand side we can replace $\mathrm{Tr}_{(P\mathfrak{H})^3}$ by $\mathrm{Tr}_{\mathfrak{H}^3}$. Combining this and (98), the triangle inequality gives

$$\begin{aligned} \mathrm{Tr}_{\mathfrak{H}^3} \left| P^{\otimes 3} \gamma_N^{(3)} P^{\otimes 3} - \int_{SP\mathfrak{H}} |u^{\otimes 3}\rangle\langle u^{\otimes 3}| d\mu_N(u) \right| \\ \leq \sum_{k=3}^N \binom{N}{3}^{-1} \binom{k}{3} \frac{12 \mathrm{Tr} P}{k} \mathrm{Tr}[G_{N,k}] \leq 12 \frac{\mathrm{Tr} P}{N} \sum_{k=3}^N \mathrm{Tr}[G_{N,k}] \leq 12 \frac{\mathrm{Tr} P}{N}, \end{aligned}$$

where the last inequality is due to (99). \square

We return to our lower-bound and applies Lemma 15 to γ_M and therefore chose

$$\tilde{\gamma}_M^{(3)} := \int_{SP\mathfrak{H}} |u^{\otimes 3}\rangle\langle u^{\otimes 3}| d\mu_M(u),$$

in our ongoing lower bound. It verifies $P^{\otimes 3} \tilde{\gamma}_M^{(3)} P^{\otimes 3} = \tilde{\gamma}_M^{(3)}$ as required. This yields

$$\begin{aligned} \left(1 + \frac{C}{2} \frac{b_\varepsilon}{\varepsilon LR^6}\right) \frac{\tilde{E}_M}{M} &\geq \frac{1}{1+\varepsilon} \int_{SP\mathfrak{H}} \tilde{\mathcal{E}}_{\varepsilon,R}(u) d\mu_M(u) - 12C_\varepsilon \left(L + \frac{b_\varepsilon}{1+\varepsilon} \frac{C}{6R^6}\right) \frac{L^q}{M} \\ &\geq \frac{\tilde{e}_{\varepsilon,R}}{1+\varepsilon} \int_{SP\mathfrak{H}} d\mu_M(u) - 12C_\varepsilon \left(L + \frac{b_\varepsilon}{1+\varepsilon} \frac{C}{6R^6}\right) \frac{L^q}{M}, \end{aligned}$$

where we defined the functional $\tilde{\mathcal{E}}_{\varepsilon,R}$ and the associated groundstate energy $\tilde{e}_{\varepsilon,R}$ by

$$\tilde{e}_{\varepsilon,R} := \inf_{SP\mathfrak{H}} \tilde{\mathcal{E}}_{\varepsilon,R}(u) := \inf_{SP\mathfrak{H}} \left(\langle u, h_{\varepsilon,s} u \rangle + \frac{b_\varepsilon}{6} \langle u^{\otimes 3}, U_R u^{\otimes 3} \rangle \right)$$

and we used that

$$\mathrm{Tr} \left[\left(\frac{h_1 + h_2 + h_3}{3} + \frac{b_\varepsilon}{6} U_R \right) \tilde{\gamma}_M^{(3)} \right] = \int_{SP\mathfrak{H}} \tilde{\mathcal{E}}_{\varepsilon,R}(u) d\mu_M(u).$$

Here the tilde highlights the fact that $\tilde{e}_{\varepsilon,R}$ is minimized on the subset $SP\mathfrak{H} := \{u \in P\mathfrak{H} \mid \|u\|_2 = 1\}$ of $S\mathfrak{H} := \{u \in \mathfrak{H} \mid \|u\|_2 = 1\}$, onto which e_{GP} or $e_\varepsilon(s)$, defined below, will be minimized.

To go further, we will use the following four ingredients. First, the following lemma (proved at the end of this section)

Lemma 16. *The measure $d\mu_N$ in (101) satisfies*

$$1 \geq \int_{SP\mathfrak{H}} d\mu_N(u) \geq 1 - 3 \mathrm{Tr}(Q\gamma_N^{(1)}) - \frac{12 \mathrm{Tr}(P)}{N}.$$

In particular, if $\gamma_M = |\Psi_M\rangle\langle\Psi_M|$ is the ground state for the Hamiltonian \tilde{H}_M in (91), then thanks to the kinetic bound (92) we get

$$\mathrm{Tr}(Q\gamma_M^{(1)}) \leq L^{-1} \mathrm{Tr} \left(h_{\varepsilon,s} \gamma_M^{(1)} \right) \leq C_\varepsilon L^{-1}.$$

Moreover, recall from (90) that $\text{Tr } P \leq C_\varepsilon L^q$. Hence, if L is chosen appropriately then the bound in Lemma 16 gives us

$$1 \geq \int_{SP\mathfrak{H}} d\mu_N(u) \geq 1 - C_\varepsilon L^{-1} - C_\varepsilon \frac{L^q}{N} \geq 1 + o(1).$$

Second, defining the functional $\mathcal{E}_{\varepsilon,s}$ and its groundstate energy $e_\varepsilon(s)$ by

$$e_\varepsilon(s) := \inf_{\|u\|_2=1} \mathcal{E}_{\varepsilon,s}(u) := \inf_{\|u\|_2=1} \langle u, h_{\varepsilon,s} u \rangle + \frac{b_\varepsilon}{6} \|u\|_{L^6(\mathbb{R}^3)}^6, \quad \forall s > 1 > \varepsilon \geq 0, \quad (102)$$

we use this other lemma (also proved at the end of this section)

Lemma 17. *For any fix $\varepsilon, s > 0$, there exists $C_{\varepsilon,s} \geq 0$ such that*

$$\inf_{SP\mathfrak{H}} \mathcal{E}_{\varepsilon,s} \geq \tilde{e}_{\varepsilon,R} \geq -C_{\varepsilon,s} R + \inf_{SP\mathfrak{H}} \mathcal{E}_{\varepsilon,s}.$$

By the variational principle, it gives

$$\tilde{e}_{\varepsilon,R} \geq -C_{\varepsilon,s} R + \inf_{SP\mathfrak{H}} \geq -C_{\varepsilon,s} R + e_\varepsilon(s).$$

Third, since $\varepsilon > 0$ and $\frac{b_{\mathcal{M}}(V)}{6} \int_{\mathbb{R}^3} |u|^6 \leq \mathcal{E}_{0,s}(u)$ —because $V_{\text{ext}} \geq 0$ —, we have

$$\mathcal{E}_{\varepsilon,s}(u) \geq \langle u, h_{0,s} u \rangle + (1 - \varepsilon)^4 \frac{b_{\mathcal{M}}(V)}{6} \int_{\mathbb{R}^3} |u|^6 \geq (1 - \mathcal{O}(\varepsilon)) \mathcal{E}_{0,s}(u) \geq (1 - \mathcal{O}(\varepsilon)) e_0(s)$$

hence

$$\tilde{e}_{\varepsilon,R} \geq -C_{\varepsilon,s} R + (1 - \mathcal{O}(\varepsilon)) e_0(s).$$

Fourth and finally, Lemma 9 and $M \geq (1 - \varepsilon)N$ give

$$\frac{E_N}{N} \geq (1 - \varepsilon) \frac{\tilde{E}_M}{M} - C_{\varepsilon,s,\beta} R^{2/7} M^{-1}.$$

Combining these four ingredients, we obtain, with $\tilde{C}_\varepsilon := (1 - \varepsilon)C_\varepsilon$, that

$$\begin{aligned} \frac{E_N}{N} &\geq \frac{(1 - \mathcal{O}(\varepsilon)) (1 - o(1)_{M \rightarrow \infty})}{1 + \frac{C}{2} \frac{b_\varepsilon}{\varepsilon L R^6}} e_0(s) \\ &- C_{\varepsilon,s} \frac{(1 - \mathcal{O}(\varepsilon)) (1 - o(1)_{M \rightarrow \infty})}{1 + \frac{C}{2} \frac{b_\varepsilon}{\varepsilon L R^6}} R - 12 \tilde{C}_\varepsilon \frac{L + \frac{b_\varepsilon}{1 + \varepsilon} \frac{C}{6 R^6} L^q}{1 + \frac{C}{2} \frac{b_\varepsilon}{\varepsilon L R^6}} \frac{L^q}{M} - C_{\varepsilon,s,\beta} \frac{R^{2/7}}{M}. \end{aligned} \quad (103)$$

The conditions $N \geq M \geq (1 - \varepsilon)N$ and $N^{-\beta/2} \geq R \geq N^{-\beta}$ imply immediately that the fourth term of the right-hand side vanishes (for $s > 1 > \varepsilon > 0$ fixed and when $R^{-1}, M, L, N \rightarrow +\infty$). Moreover, these conditions together with the definition $b_\varepsilon = b_{\mathcal{M}}(V)(1 - \varepsilon)^4$, give that

$$\frac{b_\varepsilon}{\varepsilon L R^6} \leq b_{\mathcal{M}}(V) \frac{(1 - \varepsilon)^4 N^{6\beta}}{\varepsilon L}.$$

Hence, in order for the denominator $1 + \frac{C}{2} \frac{b_\varepsilon}{\varepsilon L R^6}$ to converge to 1 (for $s > 1 > \varepsilon > 0$ fixed and when $R^{-1}, M, L, N \rightarrow +\infty$), we need to chose L such that $N^{6\beta} = o(L)$. With this assumption, we have that the second term of the right-hand side vanishes (for $s > 1 > \varepsilon > 0$ fixed and when $R^{-1}, M, L, N \rightarrow +\infty$). This assumption on L , combined with the aforementioned conditions, also implies that the numerator of the third term of the right-hand side of (103) verifies

$$\tilde{C}_\varepsilon \left(L + \frac{b_\varepsilon}{1 + \varepsilon} \frac{C}{6 R^6} \right) \frac{L^q}{M} \leq \frac{\tilde{C}_\varepsilon}{1 - \varepsilon} \left(L + \frac{C}{6} b_{\mathcal{M}}(V) \frac{(1 - \varepsilon)^4 N^{6\beta}}{1 + \varepsilon} \right) \frac{L^q}{N} \leq \frac{2\tilde{C}_\varepsilon}{1 - \varepsilon} \frac{L^{q+1}}{N}.$$

Thus, this third term vanishes as soon as $L = o\left(N^{\frac{1}{1+q}}\right)$. The existence of an L is therefore equivalent to $6\beta < 1/(q+1)$ and, since the computations hold for any $q > q_*(\alpha) := \frac{3}{2} + \frac{3}{\alpha}$, it is equivalent to $\beta < \frac{1}{3} \frac{\alpha}{5\alpha+6} \in (0, 1/15)$.

We have therefore obtained for any $s > 1 > \varepsilon > 0$ that

$$\lim_{N \rightarrow \infty} \frac{E_N}{N} \geq (1 - \mathcal{O}(\varepsilon)) e_0(s) \quad (104)$$

and, recalling the definition (102) of e_0 , we can pass to the limits $s \rightarrow +\infty$ and $\varepsilon \rightarrow 0$ in (104) to obtain the lower-bound claimed in Theorem 13:

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{E_N}{N} &\geq \lim_{s \rightarrow \infty} e_0(s) = e_{\text{GP}} := \inf_{\|u\|_2=1} \mathcal{E}_{\text{GP}}(u) \\ &:= \inf_{\|u\|_2=1} \int_{\mathbb{R}^3} |\nabla u|^2 + V_{\text{ext}} |u|^2 + \frac{b_{\mathcal{M}}(V)}{6} |u|^6. \end{aligned}$$

We conclude this section with the proofs of Lemmas 16 and 17.

Proof of Lemma 16. Note that from (101) we obtain that $\int d\mu_N \leq 1$ since every measure $\mu_{N,k}$ is normalized. It remains to prove the lower bound. From (100) and the triangle inequality, we have

$$\int d\mu_N \geq \text{Tr}(P^{\otimes 3} \gamma_N^{(3)} P^{\otimes 3}) - \frac{12 \text{Tr}(P)}{N}.$$

On the other hand, by the cyclic property of the trace, we can decompose

$$\begin{aligned} 1 &= \text{Tr}(\gamma_N^{(3)}) = \text{Tr}((P_1 + Q_1) \gamma_N^{(3)} (P_1 + Q_1)) = \text{Tr}(P_1 \gamma_N^{(3)} P_1) + \text{Tr}(Q_1 \gamma_N^{(3)} Q_1), \\ &= \text{Tr}(P_1 (P_2 + Q_2) \gamma_N^{(3)} (P_2 + Q_2) P_1) + \text{Tr}(Q_1 \gamma_N^{(3)} Q_1) \\ &= \text{Tr}(P_1 P_2 \gamma_N^{(3)} P_2 P_1) + \text{Tr}(P_1 Q_2 \gamma_N^{(3)} Q_2 P_1) + \text{Tr}(Q_1 \gamma_N^{(3)} Q_1), \\ &= \text{Tr}(P_1 P_2 P_3 \gamma_N^{(3)} P_3 P_2 P_1) + \text{Tr}(P_1 P_2 Q_3 \gamma_N^{(3)} Q_3 P_2 P_1) + \text{Tr}(P_1 Q_2 \gamma_N^{(3)} Q_2 P_1) \\ &\quad + \text{Tr}(Q_1 \gamma_N^{(3)} Q_1) \\ &\leq \text{Tr}(P^{\otimes 3} \gamma_N^{(3)} P^{\otimes 3}) + 3 \text{Tr}(Q \gamma_N^{(1)}). \quad \square \end{aligned}$$

For Lemma 17, note that we will follow the proof of [18, Lemma 4.1] but we do not need to assume something similar to their assumption $|x|w(x) \in L^1(\mathbb{R}^3)$ since by Lemma 9 we have in our case $U \in L^1(\mathbb{R}^6) \cap L^\infty(\mathbb{R}^6)$ and is compactly supported and, therefore, $z \mapsto z \|U(\cdot, z)\|_{L^\infty(\mathbb{R}^3)} \in L^1(\mathbb{R}^3)$: $\left\| z \|U(\cdot, z)\|_{L^\infty(\mathbb{R}^3)} \right\|_{L^1(\mathbb{R}^3)} \leq r \|U\|_{L^\infty(\mathbb{R}^6)}$ where r is such that $\text{Supp } U \subset B(0, r)$.

Proof of Lemma 17. We have

$$\mathcal{E}_{\varepsilon, s}(u) - \tilde{\mathcal{E}}_{\varepsilon, R}(u) = \frac{b_\varepsilon}{6} \left(\|u\|_{L^6(\mathbb{R}^3)}^6 - \langle u^{\otimes 3}, U_R u^{\otimes 3} \rangle \right).$$

First we notice by Hölder inequality and since $U \geq 0$ and $\int_{\mathbb{R}^6} U_R = \int_{\mathbb{R}^6} U = 1$ that

$$\begin{aligned} \langle u^{\otimes 3}, U_R u^{\otimes 3} \rangle &\leq \left(\int_{\mathbb{R}^9} |u(x)|^6 U_R(x-y, x-z) \right)^{\frac{1}{3}} \left(\int_{\mathbb{R}^9} |u(y)|^6 U_R(y-x, y-z) \right)^{\frac{1}{3}} \\ &\quad \times \left(\int_{\mathbb{R}^9} |u(z)|^6 U_R(z-y, z-x) \right)^{\frac{1}{3}} = \|u\|_{L^6(\mathbb{R}^3)}^6. \end{aligned}$$

We also used the symmetries of U . Since $b_\varepsilon \geq 0$, it implies $\mathcal{E}_{\varepsilon,s}(u) \geq \tilde{\mathcal{E}}_{\varepsilon,R}(u)$ hence the claim

$$\inf_{SP\mathfrak{H}} \mathcal{E}_{\varepsilon,s} \geq \tilde{\varepsilon}_{\varepsilon,R}.$$

We continue our proof by rewriting $\langle u^{\otimes 3}, U_R u^{\otimes 3} \rangle$ by change of variables:

$$\begin{aligned} \langle u^{\otimes 3}, U_R u^{\otimes 3} \rangle &= \int_{\mathbb{R}^9} |u(x)|^2 |u(y)|^2 |u(z)|^2 R^{-6} U(R^{-1}(x-y), R^{-1}(x-z)) \, dx \, dy \, dz \\ &= \int_{\mathbb{R}^9} |u(x)|^2 |u(x-Ry)|^2 |u(x-Rz)|^2 U(y,z) \, dx \, dy \, dz, \end{aligned}$$

while, since $\int_{\mathbb{R}^6} U = 1$, we have

$$\|u\|_{L^6(\mathbb{R}^3)}^6 = \left(\int_{\mathbb{R}^6} U \right) \int_{\mathbb{R}^3} |u(x)|^6 \, dx = \int_{\mathbb{R}^9} U(y,z) |u(x)|^6 \, dx \, dy \, dz,$$

and consequently

$$\begin{aligned} \langle u^{\otimes 3}, U_R u^{\otimes 3} \rangle - \|u\|_{L^6(\mathbb{R}^3)}^6 &= \int_{\mathbb{R}^9} |u(x)|^2 U(y,z) (|u(x-Ry)|^2 |u(x-Rz)|^2 - |u(x)|^4) \, dx \, dy \, dz. \end{aligned}$$

We now write the term in parenthesis as an integral:

$$\begin{aligned} |u(x-Ry)|^2 |u(x-Rz)|^2 - |u(x)|^4 &= \int_0^1 \frac{d}{dt} (|u(x-tRy)|^2 |u(x-tRz)|^2) \, dt \\ &= \int_0^1 \nabla |u|^2(x-tRy) \cdot Ry |u(x-tRz)|^2 \, dt \\ &\quad + \int_0^1 |u(x-tRy)|^2 \nabla (|u|^2)(x-tRz) \cdot Rz \, dt. \end{aligned}$$

Using the property $U(y,z) = U(z,y)$, then $|U(y,z)| \leq \|U(\cdot, z)\|_{L^\infty(\mathbb{R}^3)}$ for any $z \in \mathbb{R}^3$, then

$$\int_{\mathbb{R}^3} |u(x)|^2 |\nabla |u|^2(x-tRz)| \, dx = [|u|^2 \star |\nabla |u|^2(\cdot)|] (tRz),$$

we have

$$\begin{aligned} &\left| \langle u^{\otimes 3}, U_R u^{\otimes 3} \rangle - \|u\|_{L^6(\mathbb{R}^3)}^6 \right| \\ &\leq 2R \|u\|_2^2 \| |u|^2 \star |\nabla |u|^2(\cdot)| \|_{L^\infty(\mathbb{R}^3)} \int_{\mathbb{R}^3} \int_0^1 |z| \|U(\cdot, z)\|_{L^\infty(\mathbb{R}^3)} \, dt \, dz \\ &= 2R \|u\|_2^2 \| |u|^2 \star |\nabla |u|^2(\cdot)| \|_{L^\infty(\mathbb{R}^3)} \left\| z \|U(\cdot, z)\|_{L^\infty(\mathbb{R}^3)} \right\|_{L^1(\mathbb{R}^3)}. \end{aligned}$$

We now use the same bound as in [18, Lemma 4.1], and the convexity inequality for gradients (see [20, Theorem 7.8]), and obtain

$$0 \leq \|u\|_6^6 - \langle u^{\otimes 3}, U_R u^{\otimes 3} \rangle \leq 2R \|u\|_2^2 \|u\|_6^3 \|\nabla u\|_2 \|z\| \|U(\cdot, z)\|_\infty \|1\|_1.$$

On $SP\mathfrak{H}$, using Sobolev inequality, we have

$$0 \leq \mathcal{E}_{\varepsilon,s}(u) - \tilde{\mathcal{E}}_{\varepsilon,R}(u) \leq R \frac{b_\varepsilon}{3} \|z\| \|U(\cdot, z)\|_\infty \|1\|_1 \left(1 + \|\nabla u\|_2^2\right)^2, \quad \forall u \in SP\mathfrak{H},$$

and, since $\|\nabla u\|_2^2 \leq 2\varepsilon^{-1}\tilde{\mathcal{E}}_{\varepsilon,R}(u)$, we obtain

$$\inf_{SP\mathfrak{H}} \mathcal{E}_{\varepsilon,s} \leq \mathcal{E}_{\varepsilon,s}(u) \leq \tilde{\mathcal{E}}_{\varepsilon,R}(u) + R\frac{b_\varepsilon}{3} \|z\|U(\cdot, z)\|_\infty\|_1 \left(1 + 2\varepsilon^{-1}\tilde{\mathcal{E}}_{\varepsilon,R}(u)\right)^2, \quad \forall u \in SP\mathfrak{H}.$$

Applying this to a minimizing sequence $\{u_n\}_n \subset SP\mathfrak{H}$ to $\tilde{e}_{\varepsilon,R}$ and passing to the limit gives

$$\inf_{SP\mathfrak{H}} \mathcal{E}_{\varepsilon,s} \leq \tilde{e}_{\varepsilon,R} + R\frac{b_\varepsilon}{3} \|z\|U(\cdot, z)\|_\infty\|_1 \left(1 + 2\varepsilon^{-1}\tilde{e}_{\varepsilon,R}\right)^2 \leq \tilde{e}_{\varepsilon,R} + C_{\varepsilon,s}R, \quad \forall u \in SP\mathfrak{H},$$

where $C_{\varepsilon,s} := \frac{b_\varepsilon}{3} \|z\|U(\cdot, z)\|_\infty\|_1 \left(1 + 2\varepsilon^{-1} \inf_{SP\mathfrak{H}} \mathcal{E}_{\varepsilon,s}\right)^2$ is independent of R . \square

6. ENERGY UPPER BOUND

Let us recall that $\omega := 1 - f$, where f is the solution to the scattering equation of V , satisfies

$$0 \leq \omega \leq 1, \quad \omega(X) \leq \frac{C}{|X|^4 + 1} \quad \text{and} \quad |\nabla \omega(X)| \leq \frac{C}{|X|^5 + 1}, \quad \forall X \in \mathbb{R}^6. \quad (105)$$

We define $\omega_N := \omega(N^{1/2}\cdot)$ and $f_N := f(N^{1/2}\cdot)$, and recall that $V_N := NV(N^{1/2}\cdot)$.

For $M \geq 1$, let us define the following Hamiltonian

$$H_{M,N} = \sum_{j=1}^M (-\Delta_j + V_{\text{ext}}(x_j)) + \sum_{1 \leq i < j < k \leq M} NV(N^{1/2}(x_i - x_j, x_i - x_k)) \quad (106)$$

and

$$\begin{aligned} \mathbb{H}_N &= \bigoplus_{M \geq 1} H_{M,N} = \int_{\mathbb{R}^3} (\nabla_x a_x^* \nabla_x a_x + V(x) a_x^* a_x) \, dx \\ &\quad + \frac{1}{6} \int_{(\mathbb{R}^3)^3} V_N(x - y, x - z) a_x^* a_y^* a_z^* a_x a_y a_z \, dx \, dy \, dz. \end{aligned}$$

For $f \in L^2(\mathbb{R}^3)$, we define the Weyl operator $W(f) = \exp(a^*(f) - a(f))$ which is a unitary operator and, for $g \in L^2(\mathbb{R}^3)$, satisfies

$$W(f)^* a(g) W(f) = a(g) + \langle g, f \rangle \quad \text{and} \quad W(f)^* a^*(g) W(f) = a^*(g) + \langle f, g \rangle.$$

We also define, for $\varphi \in H^2(\mathbb{R}^3)$, $B_1 = B_1[\varphi]$ and $b = b[\varphi]$ through

$$\begin{aligned} B_1^* &= \alpha N^{\frac{3}{2}} \int_{(\mathbb{R}^3)^3} b(x, y, z) a_x^* a_y^* a_z^* \, dx \, dy \, dz \\ &= \alpha N^{\frac{3}{2}} \int_{(\mathbb{R}^3)^3} \omega_N(x, y, z) \varphi(x) \varphi(y) \varphi(z) a_x^* a_y^* a_z^* \, dx \, dy \, dz \end{aligned} \quad (107)$$

where $\alpha = -\frac{1}{6}$ —that is the unique solution to $6\alpha^2 + 2\alpha + 1/6 = 0$ —, as well as $L = N^{1/2}$, $\chi^{\leq L} = \mathbf{1}_{[0,L]}(\mathcal{N})$ and

$$U_N := \exp(\chi^{\leq L} B_1^* - B_1 \chi^{\leq L}).$$

This section is devoted to the following theorem and its Corollary.

Theorem 18 (Energy upper bound). *Let $\varphi \in H^2(\mathbb{R}^3)$ with $\|\varphi\|_{L^2} = 1$, we have*

$$\left\langle \Omega, U_N^* W(\sqrt{N}\varphi)^* \mathbb{H}_N W(\sqrt{N}\varphi) U_N \Omega \right\rangle \leq N\mathcal{E}_{GP}(\varphi) + \mathcal{O}(N^{1/2}).$$

Corollary 19. $E(N) \leq Ne_{GP} + \mathcal{O}(N^{2/3})$.

In the sequel, for shortness, we will use the notation

$$B := \chi^{\leq L} B_1^* - B_1 \chi^{\leq L}, \quad \text{hence} \quad U_N = e^{-B}. \quad (108)$$

6.1. The transformation U_N and the proof of Corollary 19. Let $p \geq 1$ and denote $\sharp = (\sharp_1, \dots, \sharp_p) \in \{*, *\}^p$. Then one can easily check that

$$[a_{x_1}^{\sharp_1} \dots a_{x_p}^{\sharp_p}, \mathcal{N}] = \kappa(\sharp) \mathcal{N},$$

where $\kappa(\sharp) = 2\#\{i | \sharp_i = *\} - p$. Also, if $\kappa(\sharp) \wedge 3 = 1$ then

$$\langle \Omega, U_N^* a_{x_1}^{\sharp_1} \dots a_{x_p}^{\sharp_p} U_N \Omega \rangle = 0. \quad (109)$$

We have the following lemmas.

Lemma 20. *Let $\varphi \in C^\infty(\mathbb{R}^3)$ and B_1 be as in (107). There exists $C_\varphi > 0$ such that*

$$\forall N \in \mathbb{N} \setminus \{0\}, \quad \pm([B_1, B_1^*] - \mathcal{Q}(\varphi)) \leq C_\varphi \left(\frac{\mathcal{N}}{N^{1/2}} + \frac{\mathcal{N}^2}{N} \right),$$

where

$$\mathcal{Q}(\varphi) = \frac{3\alpha^2}{2} \int_{\mathbb{R}^9} N^3 \omega_N(x, y, z)^2 |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz \geq 0.$$

Moreover, there exists $C > 0$ such that

$$\forall N \in \mathbb{N} \setminus \{0\}, \forall \varphi \in C^\infty(\mathbb{R}^3), \quad \mathcal{Q}(\varphi) \leq C \|\varphi\|_{L^\infty(\mathbb{R}^3)}^4 \|\varphi\|_{L^2(\mathbb{R}^3)}^2 \|\omega\|_{L^2(\mathbb{R}^6)}^2.$$

Lemma 21. *Define B as in (108). For any $k \geq 0$, there exists $C_k > 0$ such that*

$$\forall (\lambda, N) \in [-1, 1] \times \mathbb{N} \setminus \{0\}, \quad \left\langle \Omega, e^{-\lambda B} \mathcal{N}^k e^{\lambda B} \Omega \right\rangle \leq C_k \left(1 + \frac{L^2}{N} \right)^k. \quad (110)$$

Choosing $L = \mathcal{O}(N^{1/2})$ gives the immediate corollary:

$$\forall k \geq 0, \exists C_k > 0, \forall N \in \mathbb{N} \setminus \{0\}, \quad \left\langle \Omega, U_N^* \mathcal{N}^k U_N \Omega \right\rangle \leq C_k. \quad (111)$$

Before proving these two lemmas, we first use them to prove the Corollary 19 of Theorem 18.

Proof of Corollary 19. For $\mu > 0$, denote

$$E^\mu(M, N) = \inf \sigma(H_{M,N}) - \mu(M + M^3/N^2),$$

where $H_{M,N}$ is defined in (106), and $E(M, N) := E^0(M, N)$.

We claim that there exists $\mu_0 > 0$ such that, for all $\mu > \mu_0$ and $N \in \mathbb{N}$, $\{E^\mu(M, N)\}_{M \in \mathbb{N}}$ is a non-increasing sequence. Indeed, let $\gamma_{M-1, N}$ be the density matrix of the ground state of $H_{M-1, N}$, then we have the inequalities

$$\begin{aligned} & E(M, N) \\ & \leq M \operatorname{Tr}(-\Delta + V_{\text{ext}}) \gamma_{M-1, N} + \binom{M}{3} \operatorname{Tr} V_N \gamma_{M-1, N} \\ & \leq E(M-1, N) \\ & \quad + \operatorname{Tr} \left\{ \left(\frac{M}{M-1} - 1 \right) (-\Delta + V_{\text{ext}}) + \left(\frac{M}{M-3} - 1 \right) \binom{M-1}{3} V_N \right\} \gamma_{M-1, N} \\ & \leq E(M-1, N) + \frac{C}{M} E(M-1, N) \\ & \leq E(M-1, N) + C \left(\|\varphi\|_{H^1(\mathbb{R}^3)}^6 + \left\| \sqrt{V_{\text{ext}}} \varphi \right\|_{L^2(\mathbb{R}^3)}^2 \right) \left(1 + \frac{M^2}{N^2} \right), \end{aligned}$$

where we bounded $E(M-1, N)$ by the energy of the ansatz $\varphi^{\otimes M-1}$ in the last inequality. Hence, it suffices to chose $\mu_0 = C(\|\varphi\|_{H^1(\mathbb{R}^3)}^6 + \|\sqrt{V_{\text{ext}}}\varphi\|_{L^2(\mathbb{R}^3)}^2)$.

Let $\tilde{N} = N - N^{2/3}$ and let us denote $\Psi_N := W(\sqrt{\tilde{N}}\varphi)U_N\Omega \in \mathcal{F}(L^2(\mathbb{R}^3))$. As a consequence of (109), (111) and

$$W(\sqrt{\tilde{N}}\varphi)^*\mathcal{N}W(\sqrt{\tilde{N}}\varphi) = \mathcal{N} + \sqrt{\tilde{N}}(a^*(\varphi) + a(\varphi)) + \tilde{N},$$

Ψ_N satisfies, for $k \geq 0$,

$$\langle \Psi_N, \mathcal{N}\Psi_N \rangle = N - N^{2/3} + \mathcal{O}(1), \quad (112)$$

$$\langle \Psi_N, (\mathcal{N} - \langle \Psi_N, \mathcal{N}\Psi_N \rangle)^2 \Psi_N \rangle = \mathcal{O}(N), \quad (113)$$

$$\langle \Psi_N, \mathcal{N}^3\Psi_N \rangle = N^3 + \mathcal{O}(N^{8/3}). \quad (114)$$

Let us first argue that

$$\langle \mathbb{H}_N \rangle_{\Psi_N} \leq N\mathcal{E}_{GP}(\varphi) + CN^{2/3}. \quad (115)$$

Indeed, following the proof of Theorem 18, we have

$$\begin{aligned} \langle \Omega, W(\sqrt{\tilde{N}}\varphi)^*\mathbb{H}_N W(\sqrt{\tilde{N}}\varphi) \rangle_{U_N\Omega} &= \langle \Omega, U_N^* W(\sqrt{N}\varphi)^*\mathbb{H}_N W(\sqrt{N}\varphi) \rangle_{U_N\Omega} \\ &\quad + \sum_{i=0}^6 ((\tilde{N}/N)^{(6-i)/2} - 1) \langle \mathcal{L}_i \rangle_{U_N\Omega} + (\tilde{N} - N) \langle \varphi, (-\Delta + V_{\text{ext}})\varphi \rangle, \end{aligned}$$

and using Theorem 18 and that $|\langle \mathcal{L}_i \rangle_{U_N\Omega}| \leq CN$ (see proof of Theorem 18) is enough to obtain (115).

Let us now denote $\Psi_{M,N} = \mathbb{1}_{\mathcal{N}=M}\Psi_N$, then for $\mu > \mu_0$ we have

$$\begin{aligned} E^\mu(N, N) &\leq \frac{1}{1 - \langle \mathbb{1}_{\mathcal{N}>N} \rangle_{\Psi_N}} \sum_{M \leq N} E^\mu(M, N) \|\Psi_{M,N}\|^2 \\ &\leq \frac{1}{1 - \langle \mathbb{1}_{\mathcal{N}>N} \rangle_{\Psi_N}} \left(\sum_{M=0}^{\infty} E^\mu(M, N) \|\Psi_{M,N}\|^2 - \sum_{M>N} E^\mu(M, N) \|\Psi_{M,N}\|^2 \right) \\ &\leq \frac{1}{1 - \langle \mathbb{1}_{\mathcal{N}>N} \rangle_{\Psi_N}} \left\langle \mathbb{H}_N - \mu \left(\mathcal{N} + \frac{\mathcal{N}^3}{N^2} \right) + \mu \mathbb{1}_{\mathcal{N}>N} \left(\mathcal{N} + \frac{\mathcal{N}^3}{N^2} \right) \right\rangle_{\Psi_N}, \end{aligned}$$

Now, using (112), (113), (114) and Chebyshev's inequality, we obtain that

$$\begin{aligned} \langle \mathbb{1}_{\mathcal{N}>N} \rangle_{\Psi_N} &= \langle \mathbb{1}_{\mathcal{N}>\langle \mathcal{N} \rangle_{\Psi_N} + N^{2/3}} \rangle_{\Psi_N} \leq N^{-4/3} \langle (\mathcal{N} - \langle \mathcal{N} \rangle_{\Psi_N})^2 \rangle_{\Psi_N} \leq CN^{-1/3}, \\ \langle \mathbb{1}_{\mathcal{N}>N} \mathcal{N} \rangle_{\Psi_N} &= \langle \mathbb{1}_{\mathcal{N}>N} (\mathcal{N} - \langle \mathcal{N} \rangle_{\Psi_N}) \rangle_{\Psi_N} + \langle \mathbb{1}_{\mathcal{N}>N} \mathcal{N} \rangle_{\Psi_N} \\ &\leq CN^{-2/3} \langle (\mathcal{N} - \langle \mathcal{N} \rangle_{\Psi_N})^2 \rangle_{\Psi_N} + CN^{-1/3} \langle \mathcal{N} \rangle_{\Psi_N} \leq CN^{2/3} \end{aligned}$$

and

$$\begin{aligned} &\langle \mathbb{1}_{\mathcal{N}>N} \frac{\mathcal{N}^3}{N^2} \rangle_{\Psi_N} \\ &= N^{-2} \langle \mathbb{1}_{\mathcal{N}>N} (\mathcal{N} - \langle \mathcal{N} \rangle_{\Psi_N})^3 \rangle_{\Psi_N} + 3N^{-2} \langle \mathbb{1}_{\mathcal{N}>N} (\mathcal{N} - \langle \mathcal{N} \rangle_{\Psi_N})^2 \rangle_{\Psi_N} \langle \mathcal{N} \rangle_{\Psi_N} \\ &\quad + 3N^{-2} \langle \mathbb{1}_{\mathcal{N}>N} (\mathcal{N} - \langle \mathcal{N} \rangle_{\Psi_N}) \rangle_{\Psi_N} \langle \mathcal{N} \rangle_{\Psi_N}^2 + N^{-2} \langle \mathbb{1}_{\mathcal{N}>N} \rangle_{\Psi_N} \langle \mathcal{N} \rangle_{\Psi_N}^3 \\ &\leq N^{-2} \langle (\mathcal{N} - \langle \mathcal{N} \rangle_{\Psi_N})^3 \rangle_{\Psi_N} + CN^{2/3} \leq CN^{2/3}. \end{aligned}$$

Hence we deduce from the above inequality that

$$E^\mu(N, N) \leq (1 + CN^{-1/3}) \langle \mathbb{H}_N \rangle_{\Psi_N} - 2\mu N + \mathcal{O}(N^{2/3}).$$

Using (115) for $\varphi = u_0$, the GP minimizer, finishes the proof. \square

We now prove the two previously stated lemmas.

Proof of Lemma 20. On the bosonic Fock space $\mathcal{F}(\mathfrak{H})$, we have

$$[a_x a_y a_z, a_{x'}^* a_{y'}^* a_{z'}^*] = 6\delta_{x=x'}\delta_{y=y'}\delta_{z=z'} + 18a_x^* a_{x'} \delta_{y=y'}\delta_{z=z'} + 9a_x^* a_y^* a_{x'} a_{y'} \delta_{z=z'}. \quad (116)$$

Hence,

$$\begin{aligned} [B_1, B_1^*] &= \frac{3\alpha^2}{2} \int_{\mathbb{R}^9} N^3 \omega_N(x, y, z)^2 |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz \\ &\quad + \frac{9\alpha^2}{2} \int_{\mathbb{R}^{12}} N^{3/2} \omega_N(x, y, z) \\ &\quad \quad \times N^{3/2} \omega_N(x', y, z) \varphi(x) \overline{\varphi(x')} |\varphi(y)|^2 |\varphi(z)|^2 a_x^* a_{x'} dx dx' dy dz \\ &\quad + \frac{9\alpha^2}{4} \int_{\mathbb{R}^{15}} N^{3/2} \omega_N(x, y, z) N^{3/2} \omega_N(x', y', z) \\ &\quad \quad \times \varphi(x) \overline{\varphi(x')} \varphi(y) \overline{\varphi(y')} |\varphi(z)|^2 a_x^* a_y^* a_{x'} a_{y'} dx dx' dy dy' dz \\ &= (\text{I}) + (\text{I}) + (\text{III}), \end{aligned}$$

which defines our terms (I), (I) and (III). We have

$$(\text{I}) \leq C \|\varphi\|_{L^\infty(\mathbb{R}^3)}^4 \|\varphi\|_{L^2(\mathbb{R}^3)}^2 \|N^{3/2} \omega_N\|_{L^2(\mathbb{R}^6)}^2 \leq C \|\varphi\|_{L^\infty(\mathbb{R}^3)}^4 \|\varphi\|_{L^2(\mathbb{R}^3)}^2 \|\omega\|_{L^2(\mathbb{R}^6)}^2.$$

Note that, contrarily to the two-body interaction case, here $\omega \in L^2(\mathbb{R}^6)$. Again, using the Cauchy-Schwarz inequality $a_x^* a_{x'} + h.c. \leq a_x^* a_x + a_{x'}^* a_{x'}$, we have that

$$\begin{aligned} (\text{II}) &\leq C \int_{\mathbb{R}^{12}} N^3 \omega_N(x, y, z) \omega_N(x', y, z) |\varphi(x) \varphi(x')| |\varphi(y)|^2 |\varphi(z)|^2 a_x^* a_x dx dx' dy dz \\ &\leq C \|\varphi\|_{L^\infty(\mathbb{R}^3)} \int_{\mathbb{R}^9} N \omega_N(x, y, z) |\varphi(x)| \frac{|\varphi(y)|^2 |\varphi(z)|^2}{|y-z|} a_x^* a_x dx dy dz \\ &\leq C \|N \omega_N\|_{L^2(\mathbb{R}^6)} \left\| \frac{|\varphi(y)|^2 |\varphi(z)|^2}{|y-z|} \right\|_{L^2(\mathbb{R}^6)} \|\varphi\|_{L^\infty(\mathbb{R}^3)}^2 \mathcal{N} \leq C \|\varphi\|_{H^2(\mathbb{R}^3)}^6 \frac{\mathcal{N}}{N^{1/2}}, \end{aligned}$$

where we used that

$$\int_{\mathbb{R}^3} N^2 \omega_N(x', y, z) |\varphi(x')| dx' \leq C \int_{\mathbb{R}^3} \frac{\|\varphi\|_\infty dx'}{(|x'-y|^2 + |y-z|^2)^2} \leq C \frac{\|\varphi\|_\infty}{|y-z|}. \quad (117)$$

We now estimate the last term (III), for this we define for $f \in L^2(\mathbb{R}^6)$ and $z \in \mathbb{R}^3$

$$T(f)(z) = \varphi(z) \int_{\mathbb{R}^6} N^{\frac{3}{2}} \omega_N(x, y, z) \varphi(x) \varphi(y) f(x, y) dx dy.$$

The term (III) is proportional to the second quantification of T^*T . On the other hand, using (105), we have that

$$\begin{aligned} &T(f)(z) \\ &\leq CN^{-\frac{1}{2}} |\varphi(z)| \int_{\mathbb{R}^3} |\varphi(y)| \left(\int_{\mathbb{R}^3} \frac{\varphi(x) f(x, y)}{(|x-y|^2 + |y-z|^2)^2} dx \right) dy \\ &\leq CN^{-\frac{1}{2}} |\varphi(z)| \int_{\mathbb{R}^3} |\varphi(y)| \left(\int_{\mathbb{R}^3} \frac{du}{(u^2 + |y-z|^2)^4} \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^3} |\varphi(x)|^2 |f(x, y)|^2 dx \right)^{\frac{1}{2}} dy \\ &\leq CN^{-\frac{1}{2}} \|\varphi\|_{L^\infty(\mathbb{R}^3)}^2 |\varphi(z)| \int_{\mathbb{R}^3} \frac{1}{|y-z|^{5/2}} \|f(\cdot, y)\|_{L^2(\mathbb{R}^3)} dy \end{aligned}$$

and hence by the Hardy–Littlewood–Sobolev inequality we obtain

$$\|T(f)\|_{L^2(\mathbb{R}^3)} \leq CN^{-\frac{1}{2}} \|\varphi\|_{L^\infty(\mathbb{R}^3)}^2 \|\varphi\|_{L^6(\mathbb{R}^3)} \|f\|_{L^2(\mathbb{R}^3)}.$$

From this we conclude that

$$(III) \leq C \|\varphi\|_{H^2(\mathbb{R}^3)}^6 \frac{\mathcal{N}^2}{N}. \quad \square$$

Proof of Lemma 21. Let $k \geq 1$ and $\xi(\mathcal{N}) = \mathcal{N}^k$, let us denote $\partial\xi = \xi(\cdot + 3) - \xi$. Then ξ is such that $\xi(0) = 0$ and $\partial^j \xi \geq 0$ for $j \geq 0$. We have

$$e^{-\lambda B} \xi(\mathcal{N}) e^{\lambda B} = \xi(\mathcal{N}) - \lambda [B, \xi(\mathcal{N})] + \lambda^2 \int_0^1 \int_0^s e^{-\lambda u B} [B, [B, \xi(\mathcal{N})]] e^{\lambda u B} du ds. \quad (118)$$

We are only interested in the expectation on the vacuum Ω , hence, since $\xi(0) = 0$, only the third term will give a non zero contribution. Using that $a_x \mathcal{N} = (\mathcal{N} + 1)a_x$, we obtain

$$-[B, \xi(\mathcal{N})] = [B_1 \chi^{\leq L}, \xi(\mathcal{N})] + h.c. = \partial\xi(\mathcal{N}) B_1 \chi^{\leq L} + h.c.$$

and

$$\begin{aligned} [B, [B, \xi(\mathcal{N})]] &= [B_1 \chi^{\leq L} - \chi^{\leq L} B_1^*, \partial\xi(\mathcal{N}) B_1 \chi^{\leq L}] + h.c. \\ &= \partial^2 \xi(\mathcal{N}) (B_1 \chi^{\leq L})^2 - \chi^{\leq L} B_1^* \partial^2 \xi(\mathcal{N}) B_1 \chi^{\leq L} \\ &\quad + \partial\xi(\mathcal{N}) [B_1 \chi^{\leq L}, \chi^{\leq L} B_1^*] + h.c. \end{aligned} \quad (119)$$

The last term, which is the most regular, is controlled using that $\partial\xi(\mathcal{N})$ is non-negative and commutes with $[B_1 \chi^{\leq L}, \chi^{\leq L} B_1^*]$, and that

$$\begin{aligned} [B_1 \chi^{\leq L}, \chi^{\leq L} B_1^*] &= \partial(\chi^{\leq L})^2 B_1^* B_1 + (\chi^{\leq L-3})^2 [B_1, B_1^*] \\ &\leq (\chi^{\leq L-3})^2 [B_1, B_1^*] \leq (\chi^{\leq L-3})^2 \left(\mathcal{Q}(\varphi) + C \frac{\mathcal{N}}{N^{1/2}} + C \frac{\mathcal{N}^2}{N} \right) \\ &\leq C (\chi^{\leq L-3})^2 \left(1 + \frac{L^2}{N} \right), \end{aligned} \quad (120)$$

where we used that χ and $B_1^* B_1$ commute and, for the first inequality, that $\partial(\chi^{\leq L})^2 \leq 0$, and Lemma 20 for the second inequality.

The first two terms in (119) are controlled using the Cauchy–Schwarz inequality, we have

$$\begin{aligned} &\partial^2 \xi(\mathcal{N}) (B_1 \chi^{\leq L})^2 - \chi^{\leq L} B_1^* \partial^2 \xi(\mathcal{N}) B_1 \chi^{\leq L} + h.c. \\ &= (B_1 \chi^{\leq L}) \partial^2 \xi(\mathcal{N} - 3) (B_1 \chi^{\leq L}) - 2 \chi^{\leq L} B_1^* \partial^2 \xi(\mathcal{N}) B_1 \chi^{\leq L} \\ &\leq (B_1 \chi^{\leq L}) \partial^2 \xi(\mathcal{N} - 3) (\chi^{\leq L} B_1^*) + (\chi^{\leq L} B_1^*) \partial^2 \xi(\mathcal{N} - 3) (\chi^{\leq L} B_1) \\ &\quad - 2 \chi^{\leq L} B_1^* \partial^2 \xi(\mathcal{N}) B_1 \chi^{\leq L} \\ &= \partial^2 \xi(\mathcal{N}) [B_1 \chi^{\leq L}, \chi^{\leq L} B_1^*] + (\partial^2 \xi(\mathcal{N}) + \partial^2 \xi(\mathcal{N} - 6)) \\ &\quad - 2 \partial^2 \xi(\mathcal{N} - 3) \chi^{\leq L} B_1^* B_1 \chi^{\leq L} \\ &= \partial^2 \xi(\mathcal{N}) [B_1 \chi^{\leq L}, \chi^{\leq L} B_1^*] + \partial^4 \xi(\mathcal{N} - 6) \chi^{\leq L} B_1^* B_1 \chi^{\leq L} \\ &\leq C (1 + L^2 N^{-1}) \partial^2 \xi(\mathcal{N}) + C \partial^4 \xi(\mathcal{N} - 6) (\mathcal{N} + 1)^3 \end{aligned}$$

Gathering the last three inequalities, we obtain

$$[B, [B, \xi(\mathcal{N})]] \leq C (\partial\xi(\mathcal{N}) + \partial^2 \xi(\mathcal{N})) (1 + L^2 N^{-1}) + C \partial^4 \xi(\mathcal{N} - 6) (\mathcal{N} + 1)^3.$$

Choosing $\xi(\mathcal{N}) = \mathcal{N}$, the above inequality shows that (110) holds for $k = 1$. For $k \geq 2$, taking $\xi(\mathcal{N}) = \mathcal{N}^k$, an immediate induction concludes the proof. \square

6.2. Computation of the upper bound: proof of Theorem 18. We first apply the Weyl operator to \mathbb{H}_N , we obtain

$$\begin{aligned} & W(\sqrt{N}\varphi)^* d\Gamma(-\Delta + V_{\text{ext}})W(\sqrt{N}\varphi) \\ &= \Gamma(-\Delta + V_{\text{ext}}) + \sqrt{N}a^*((-\Delta + V_{\text{ext}})\varphi) + \sqrt{N}a((-\Delta + V_{\text{ext}})\varphi) \\ & \quad + N\langle\varphi, (-\Delta + V_{\text{ext}})\varphi\rangle \end{aligned}$$

and

$$\frac{1}{6} \int_{\mathbb{R}^9} V_N(x-y, x-z) W(\sqrt{N}\varphi)^* a_x^* a_y^* a_z^* a_x a_y a_z W(\sqrt{N}\varphi) dx dy dz = \sum_{i=0}^6 \mathcal{L}_i,$$

where

$$\begin{aligned} \mathcal{L}_6 &= \frac{1}{6} \int_{\mathbb{R}^9} V_N(x-y, x-z) a_x^* a_y^* a_z^* a_x a_y a_z dx dy dz, \\ \mathcal{L}_5 &= \frac{N^{1/2}}{2} \int_{\mathbb{R}^9} V_N(x-y, x-z) a_x^* a_y^* a_z^* a_x a_y \varphi(z) dx dy dz + h.c., \\ \mathcal{L}_4 &= \frac{N}{2} \int_{\mathbb{R}^9} V_N(x-y, x-z) a_x^* a_y^* a_z^* a_x \varphi(y) \varphi(z) dx dy dz + h.c. \\ & \quad + \frac{N}{2} \int_{\mathbb{R}^9} V_N(x-y, x-z) a_x^* a_y^* a_x a_z \varphi(y) \overline{\varphi(z)} dx dy dz + h.c. \\ & \quad + \frac{N}{2} \int_{\mathbb{R}^9} V_N(x-y, x-z) a_x^* a_y^* a_x a_y |\varphi(z)|^2 dx dy dz \\ &=: \mathcal{L}_4^{(1)} + \mathcal{L}_4^{(2)} + \mathcal{L}_4^{(3)}, \\ \mathcal{L}_3 &= \frac{N^{3/2}}{6} \int_{\mathbb{R}^9} V_N(x-y, x-z) a_x^* a_y^* a_z^* \varphi(x) \varphi(y) \varphi(z) dx dy dz + h.c. \\ & \quad + \frac{N^{3/2}}{2} \int_{\mathbb{R}^9} V_N(x-y, x-z) a_x^* a_y^* a_x \varphi(y) |\varphi(z)|^2 dx dy dz + h.c. \\ & \quad + \frac{N^{3/2}}{2} \int_{\mathbb{R}^9} V_N(x-y, x-z) a_x^* a_y^* a_z \varphi(x) \varphi(y) \overline{\varphi(z)} dx dy dz + h.c. \\ &=: \mathcal{L}_3^{(1)} + \mathcal{L}_3^{(2)} + \mathcal{L}_3^{(3)}, \\ \mathcal{L}_2 &= \frac{N^2}{2} \int_{\mathbb{R}^9} V_N(x-y, x-z) a_x^* a_y^* \varphi(x) \varphi(y) |\varphi(z)|^2 dx dy dz + h.c. \\ & \quad + \frac{N^2}{2} \int_{\mathbb{R}^9} V_N(x-y, x-z) a_x^* a_x |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz \\ & \quad + N^2 \int_{\mathbb{R}^9} V_N(x-y, x-z) a_x^* a_y \varphi(x) \overline{\varphi(y)} |\varphi(z)|^2 dx dy dz \\ &=: \mathcal{L}_2^{(1)} + \mathcal{L}_2^{(2)} + \mathcal{L}_2^{(3)}, \\ \mathcal{L}_1 &= \frac{N^{5/2}}{2} \int_{\mathbb{R}^9} V_N(x-y, x-z) a_x^* \varphi(x) |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz + h.c. \end{aligned}$$

and

$$\mathcal{L}_0 = \frac{N}{6} \int_{\mathbb{R}^9} N^2 V_N(x-y, x-z) |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz.$$

Gathering up the constant terms, we obtain

$$N \left\{ \int_{\mathbb{R}^3} (|\nabla \varphi(x)|^2 + V_{\text{ext}} |\varphi(x)|^2) dx + \frac{1}{6} \int_{\mathbb{R}^9} N^3 V(N^{1/2}(x-z, y-z)) |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz \right\}$$

which is the mean-field energy associated to the particle φ . This quantity does not take into account the contribution from the scattering process. To obtain it, one must include contributions hidden in the terms $d\Gamma(-\Delta)$, $\mathcal{L}_3^{(1)}$ and \mathcal{L}_6 . Let us first prove that the other terms are negligible and then extract the main contribution from the previously mentioned terms.

6.2.1. *Controlling the error terms.* As in (109), we have

$$\langle \Omega, e^{-B} A e^B \Omega \rangle = 0, \quad \forall A \in \left\{ a^* ((-\Delta + V_{\text{ext}}) \varphi), \mathcal{L}_1, \mathcal{L}_2^{(1)}, \mathcal{L}_3^{(2)}, \mathcal{L}_3^{(3)}, \mathcal{L}_4^{(1)}, \mathcal{L}_5 \right\}$$

because they create or annihilate a number of particles which is coprime with 3. It remains to bound the following terms.

Lemma 22. *We have*

$$\langle \Omega, e^{-B} (\mathcal{L}_2^{(2)} + \mathcal{L}_2^{(3)}) e^B \Omega \rangle \leq C \quad \text{and} \quad \langle \Omega, e^{-B} (\mathcal{L}_4^{(2)} + \mathcal{L}_4^{(3)}) e^B \Omega \rangle \leq C N^{1/2}.$$

Proof. We will show that

$$\begin{aligned} \mathcal{L}_2^{(2)} + \mathcal{L}_2^{(3)} &\leq C \|V\|_{L^1(\mathbb{R}^6)} \|\varphi\|_{L^\infty(\mathbb{R}^3)}^2 \|\varphi\|_{L^2(\mathbb{R}^3)}^2 \mathcal{N}, \\ \mathcal{L}_4^{(2)} + \mathcal{L}_4^{(3)} &\leq C \sup_{x \in \mathbb{R}^3} \|V(\cdot, x)\|_{L^1(\mathbb{R}^3)} \|\varphi\|_{L^\infty(\mathbb{R}^3)}^2 N^{1/2} \mathcal{N}^2. \end{aligned}$$

Then applying (111) finishes the proof. The first bound comes from

$$\mathcal{L}_2^{(2)} = \frac{1}{2} \int_{\mathbb{R}^6} V |\varphi|^2 \otimes |\varphi|^2 \mathcal{N} \leq C \|V\|_{L^1(\mathbb{R}^6)} \|\varphi\|_{L^\infty(\mathbb{R}^3)}^4 \mathcal{N}$$

and, using the Cauchy–Schwarz inequality,

$$\begin{aligned} \mathcal{L}_2^{(3)} &\leq \int_{\mathbb{R}^9} N^3 V(N^{1/2}(x-y, x-z)) (a_x^* a_x |\varphi(y)|^2 + a_y^* a_y |\varphi(x)|^2) |\varphi(z)|^2 dx dy dz \\ &\leq C \|V\|_{L^1(\mathbb{R}^6)} \|\varphi\|_{L^\infty(\mathbb{R}^3)}^4 \mathcal{N}. \end{aligned}$$

For the second bound, on one hand we deduce from the pointwise inequality

$$\int_{\mathbb{R}^3} N^2 V(N^{1/2}(x-y, x-z)) |\varphi(z)|^2 dz \leq N^{1/2} \|\varphi\|_{L^\infty(\mathbb{R}^3)}^2 \sup_{u \in \mathbb{R}^3} \|V(u, \cdot)\|_{L^1(\mathbb{R}^3)}$$

that

$$\mathcal{L}_4^{(3)} \leq C \sup_{x \in \mathbb{R}^3} \|V(\cdot, x)\|_{L^1(\mathbb{R}^3)} \|\varphi\|_{L^\infty(\mathbb{R}^3)}^2 N^{1/2} \mathcal{N}^2,$$

and another hand by the Cauchy–Schwarz inequality we have

$$\mathcal{L}_4^{(2)} \leq \frac{N}{2} \int_{\mathbb{R}^9} V_N(x-y, x-z) (a_x^* a_y^* a_x a_y |\varphi(z)|^2 + a_x^* a_z^* a_x a_z |\varphi(y)|^2) dx dy dz$$

$$\leq C \sup_{x \in \mathbb{R}^3} \|V(\cdot, x)\|_{L^1(\mathbb{R}^3)} \|\varphi\|_{L^\infty(\mathbb{R}^3)}^2 N^{1/2} \mathcal{N}^2. \quad \square$$

6.2.2. Contribution from $d\Gamma(-\Delta)$.

Lemma 23. *We have*

$$\begin{aligned} & \langle \Omega, e^{-B} d\Gamma(-\Delta) e^B \Omega \rangle \\ & \leq 6\alpha^2 N^3 \int_{\mathbb{R}^9} (V_N f_N \omega_N)(x-z, y-z) |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz + \mathcal{O}(N^{1/2}). \end{aligned}$$

Proof. We have

$$e^{-B} d\Gamma(-\Delta) e^B = d\Gamma(-\Delta) - \int_0^1 e^{-sB} [B, d\Gamma(-\Delta)] e^{sB} ds. \quad (121)$$

Let us compute

$$\begin{aligned} & -[B, d\Gamma(-\Delta)] \quad (122) \\ & = [B_1 \chi^{\leq L}, d\Gamma(-\Delta)] + h.c. \\ & = \int_{\mathbb{R}^9} \int_{\mathbb{R}^3} \overline{b(x, y, z)} [a_x a_y a_z \chi^{\leq L}, \nabla_{x'} a_{x'}^* \nabla_{x'} a_{x'}] dx dy dz dx' + h.c. \\ & = 3 \int_{\mathbb{R}^9} -\Delta_1 b(x, y, z) a_x a_y a_z \chi^{\leq L} dx dy dz + h.c. \\ & = 3\alpha N^{3/2} \int_{\mathbb{R}^9} \left((-\Delta_1 \omega_N(x, y, z)) \varphi(x) \varphi(y) \varphi(z) - \omega_N(x, y, z) \Delta \varphi(x) \varphi(y) \varphi(z) \right. \\ & \quad \left. - 2 \nabla_1 \omega_N(x, y, z) \cdot \nabla \varphi(x) \varphi(y) \varphi(z) \right) a_x a_y a_z \chi^{\leq L} dx dy dz + h.c. \\ & =: \mathcal{K}_1 + \mathcal{K}_2 + \mathcal{K}_3. \quad (123) \end{aligned}$$

The main contribution comes from \mathcal{K}_1 . Indeed, using the Cauchy–Schwarz inequality with

$$\left\| N^{3/2} \nabla_1 \omega_N(x, y, z) \cdot \nabla \varphi(x) \varphi(y) \varphi(z) \right\|_{L^2(\mathbb{R}^9)} \leq CN^{1/2}$$

and

$$\|\omega_N(x, y, z) \Delta \varphi(x) \varphi(y) \varphi(z)\|_{L^2(\mathbb{R}^9)} \leq C,$$

we obtain that

$$\pm (\mathcal{K}_2 + \mathcal{K}_3) \leq C(1 + N^{1/2}) \mathcal{N}^{3/2}. \quad (124)$$

Noticing that

$$\begin{aligned} -\Delta_1 \omega_N(x, y, z) - \Delta_2 \omega_N(x, y, z) - \Delta_3 \omega_N(x, y, z) &= (\nabla_{\mathbb{R}^6} \cdot M \nabla_{\mathbb{R}^6} \omega)(x-z, y-z) \\ &= (V_N f_N)(x-z, y-z) \end{aligned}$$

and using symmetry, we can rewrite \mathcal{K}_1 as

$$\mathcal{K}_1 = \alpha N^{3/2} \int_{\mathbb{R}^9} (V_N f_N)(x-z, y-z) \varphi(x) \varphi(y) \varphi(z) a_x a_y a_z \chi^{\leq L} dx dy dz + h.c.$$

From (121)–(124), we obtain

$$e^{-B} d\Gamma(-\Delta) e^B = d\Gamma(-\Delta) + \int_0^1 e^{-sB} (\mathcal{K}_1 + \mathcal{K}_2 + \mathcal{K}_3) e^{sB} ds$$

$$\begin{aligned} &\leq d\Gamma(-\Delta) + \mathcal{K}_1 - \int_0^1 \int_0^s e^{-uB} [B, \mathcal{K}_1] e^{uB} du ds \\ &\quad + CN^{1/2} \int_0^1 e^{-sB} (1 + \mathcal{N})^{3/2} e^{sB} ds. \end{aligned}$$

Since we test against the vacuum, the two first terms will not contribute and the last one will give a contribution of order $N^{1/2}$. Let us then compute

$$\begin{aligned} &-[B, \mathcal{K}_1] \\ &= \alpha N^{3/2} \int_{\mathbb{R}^9} (V_N f_N)(x-z, y-z) \varphi(x) \varphi(y) \varphi(z) \\ &\quad \times [a_x a_y a_z \chi^{\leq L}, \chi^{\leq L} B_1^*] dx dy dz + h.c. \\ &= \alpha N^{3/2} \int_{\mathbb{R}^9} \int_{\mathbb{R}^9} (V_N f_N)(x-z, y-z) \varphi(x) \varphi(y) \varphi(z) b(x', y', z') \\ &\quad \times \chi^2(\cdot + 3) [a_x a_y a_z, a_{x'}^* a_{y'}^* a_{z'}^*] dx dy dz dx' dy' dz' + h.c. \\ &\quad + \alpha N^{3/2} \int_{\mathbb{R}^9} \int_{\mathbb{R}^9} (V_N f_N)(x-z, y-z) \varphi(x) \varphi(y) \varphi(z) b(x', y', z') \\ &\quad \times \partial \chi^2 a_{x'}^* a_{y'}^* a_{z'}^* a_x a_y a_z dx dy dz dx' dy' dz' + h.c. \\ &= \chi^2(\cdot + 3) \mathcal{K}_{1,1} + \partial \chi^2 \mathcal{K}_{1,2}. \end{aligned}$$

The main contribution comes from $\mathcal{K}_{1,1}$. Indeed, we have

$$\partial \chi^2 \mathcal{K}_{1,2} \leq C \partial \chi^2 \|N^{3/2} (V_N f_N) \varphi^{\otimes 3}\|_{L^2(\mathbb{R}^9)} \|b\|_{L^2(\mathbb{R}^9)} (\mathcal{N} + 1)^3 \leq C \frac{N}{L^k} \mathcal{N}^{k+3}.$$

Expanding the commutator in $\mathcal{K}_{1,1}$, we obtain

$$\begin{aligned} \mathcal{K}_{1,1} &= 12\alpha^2 N^3 \int_{\mathbb{R}^9} (V_N f_N \omega_N)(x-z, y-z) |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz \\ &\quad + 36\alpha^2 N^3 \int_{\mathbb{R}^{12}} (V_N f_N)(x, y, z) \omega_N(x', y, z) \varphi(x) \overline{\varphi(x')} |\varphi(y)|^2 |\varphi(z)|^2 \\ &\quad \quad \quad \times a_x^* a_{x'} dx dx' dy dz \\ &\quad + 18\alpha^2 N^3 \int_{\mathbb{R}^{15}} (V_N f_N)(x, y, z) \omega_N(x', y', z) \varphi(x) \varphi(y) \overline{\varphi(x') \varphi(y')} |\varphi(z)|^2 \\ &\quad \quad \quad \times a_x^* a_y^* a_{x'} a_{y'} dx dx' dy dy' dz \\ &=: \text{(I)} + \text{(II)} + \text{(III)}. \end{aligned}$$

Of course (I) is the main contribution. Let us bound the other terms. Using (117), we have

$$\begin{aligned} \text{(II)} &\leq CN^3 \int_{\mathbb{R}^{12}} (V_N f_N)(x, y, z) \omega_N(x', y, z) |\varphi(x) \overline{\varphi(x')}| |\varphi(y)|^2 |\varphi(z)|^2 \\ &\quad \quad \quad \times (a_x^* a_x + a_{x'}^* a_{x'}) dx dx' dy dz \\ &\leq C \|\varphi\|_\infty N \int_{\mathbb{R}^9} (V_N f_N)(x, y, z) |\varphi(x)| \frac{|\varphi(y)|^2 |\varphi(z)|^2}{|y-z|} a_x^* a_x dx dy dz \\ &\quad + CN^{\frac{5}{2}} \int_{\mathbb{R}^9} \omega_N(x', y, z) |\overline{\varphi(x')}| |\varphi(y)|^2 |\varphi(z)|^2 a_{x'}^* a_{x'} dx' dy dz \\ &\leq C \|\varphi\|_\infty^4 N^2 \left\| |\varphi|^2 * \frac{1}{|\cdot|} * \mathbf{1}_{|\cdot| \leq CN^{-1/2}} \right\|_{L^1} \mathcal{N} + N^{\frac{1}{2}} \left\| |\varphi|^2 * \frac{1}{|\cdot|} \right\|_{L^1} \|\varphi\|_\infty^4 \mathcal{N} \\ &\leq CN^{1/2} \mathcal{N}. \end{aligned}$$

Similar computations give

$$(III) \leq N^3 \int_{\mathbb{R}^{15}} (V_N f_N)(x, y, z) \omega_N(x', y', z) \varphi(x) \varphi(y) \overline{\varphi(x') \varphi(y')} |\varphi(z)|^2 \\ \times (a_x^* a_y^* a_x a_y + a_{x'}^* a_{y'}^* a_{x'} a_{y'}) dx dx' dy dy' dz \leq CN^{1/2} \mathcal{N}^2.$$

We therefore obtain

$$\mathcal{K}_{1,1} \leq 12\alpha^2 N^3 \int_{\mathbb{R}^9} (V_N f_N \omega_N)(x - z, y - z) |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz \\ + CN^{1/2} (1 + \mathcal{N}^2).$$

Noting that $\int_0^1 \int_0^s du ds = 1/2$, that $0 \leq 1 - \chi^2 \leq \mathcal{N}/L = \mathcal{N}/N^{1/2}$ and Lemma 21 concludes the proof of Lemma 23. \square

6.2.3. Contribution from \mathcal{L}_3 .

Lemma 24. *We have*

$$\langle \Omega, e^{-B} \mathcal{L}_3 e^B \Omega \rangle \\ \leq 2\alpha \int_{\mathbb{R}^9} N^3 (V_N \omega_N)(x, y, z) |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz + \mathcal{O}(N^{1/2}).$$

Proof. Let us first introduce the cut-off function χ in $\mathcal{L}_3 = \mathcal{L}_3^\dagger + \mathcal{L}_3^\circ$. We have

$$\mathcal{L}_3 \leq \chi^{\leq L} \mathcal{L}_3^\dagger + \mathcal{L}_3^\circ \chi^{\leq L} + C \frac{N}{L} \mathcal{N}^{5/2}.$$

Hence, we can look at

$$e^{-B} (\chi^{\leq L} \mathcal{L}_3^\dagger + \mathcal{L}_3^\circ \chi^{\leq L}) e^B = (\chi^{\leq L} \mathcal{L}_3^\dagger + \mathcal{L}_3^\circ \chi^{\leq L}) - \int_0^1 e^{sB} [B, \chi^L \mathcal{L}_3^\dagger + \mathcal{L}_3^\circ \chi^{\leq L}] e^{-sB} ds.$$

The first term vanishing when tested against Ω , let us compute the commutator. Using (116), we obtain

$$-[B, \chi^{\leq L} \mathcal{L}_3^\dagger + \mathcal{L}_3^\circ \chi^{\leq L}] = -[B, \mathcal{L}_3^\circ \chi^{\leq L}] + h.c. = [B_1 \chi^{\leq L} - \chi^{\leq L} B_1^*, \mathcal{L}_3^\circ \chi^{\leq L}] + h.c. \\ = [\mathcal{L}_3^\circ \chi^{\leq L}, \chi^{\leq L} B_1^*] + h.c. \\ = \chi^{\leq L-3} [\mathcal{L}_3^\circ, B_1^*] + \partial \chi^{\leq L} B_1^* \mathcal{L}_3^\circ + h.c. \\ \leq [\mathcal{L}_3^\circ, B_1^*] + h.c. + C \frac{N}{L} (\mathcal{N} + 1)^4.$$

Then, the same computations as in (125) gives

$$[\mathcal{L}_3^\circ, B_1^*] + h.c. \\ = \frac{\alpha}{6} \int_{\mathbb{R}^9 \times \mathbb{R}^9} N^3 V_N(x, y, z) \omega_N(x', y', z') \overline{\varphi(x) \varphi(y) \varphi(z) \varphi(x') \varphi(y') \varphi(z')} \\ \times [a_x a_y a_z, a_x^* a_y^* a_z^*] dx dy dz dx' dy' dz' + h.c. \\ \leq 2\alpha \int_{\mathbb{R}^9} N^3 (V_N \omega_N)(x, y, z) |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz + C(N^{1/2} \mathcal{N} + \mathcal{N}^2),$$

which concludes the proof. \square

6.2.4. Contribution from \mathcal{L}_6 .

Lemma 25. *We have*

$$\begin{aligned} & \langle \Omega, e^{-B} \mathcal{L}_6 \Omega \rangle \\ & \leq 6\alpha^2 N^3 \int_{\mathbb{R}^9} (V_N \omega_N^2)(x-z, y-z) |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz + \mathcal{O}(N^{1/2}). \end{aligned}$$

Proof. We have

$$e^{-B} \mathcal{L}_6 e^B = \mathcal{L}_6 - \int_0^1 e^{sB} [B, \mathcal{L}_6] e^{-sB} ds.$$

When estimating against Ω , only the last term will give a non zero contribution. Let us therefore focus on it. Using that $[\mathcal{A}\mathcal{B}, \mathcal{C}] = \mathcal{A}[\mathcal{B}, \mathcal{C}]$ when $[\mathcal{A}, \mathcal{C}] = 0$, we have

$$\begin{aligned} & - [B, \mathcal{L}_6] \\ & = \frac{\alpha}{6} \int_{\mathbb{R}^9} \int_{\mathbb{R}^9} V_N(x-y, x-z) [a_x a_y a_z, a_x^* a_{y'} a_{z'}^*] a_x a_y a_z \\ & \quad \times \chi^{\leq L} N^{3/2} \omega_N(x', y', z') \overline{\varphi(x') \varphi(y') \varphi(z')} dx dy dz dx' dy' dz' + h.c. \\ & = \alpha \int_{\mathbb{R}^9} N^{3/2} V_N(x-y, x-z) \omega_N(x, y, z) \overline{\varphi(x) \varphi(y) \varphi(z)} a_x a_y a_z \chi^{\leq L} dx dy dz + h.c. \\ & \quad + 12\alpha \int_{\mathbb{R}^{12}} N^{3/2} V_N(x-y, x-z) \omega_N(x', y, z) \overline{\varphi(x') \varphi(y) \varphi(z)} \\ & \quad \quad \quad \times a_x^* a_x a_y a_z a_{x'} \chi^{\leq L} dx dy dz dx' + h.c. \\ & \quad + \frac{3\alpha}{2} \int_{\mathbb{R}^{15}} N^{3/2} V_N(x-y, x-z) \omega_N(x', y', z) \overline{\varphi(x') \varphi(y') \varphi(z)} \\ & \quad \quad \quad \times a_x^* a_y^* a_x a_y a_{x'} a_{y'} a_z \chi^{\leq L} dx dy dz dx' dy' + h.c. \\ & =: \tilde{\mathcal{L}}_6^{(1)} + \tilde{\mathcal{L}}_6^{(2)} + \tilde{\mathcal{L}}_6^{(3)}, \end{aligned}$$

where we now prove that only the contribution of $\tilde{\mathcal{L}}_6^{(1)}$ is of order N .

On one hand, using the Cauchy–Schwarz inequality, we obtain for all $\eta > 0$ that

$$\begin{aligned} \tilde{\mathcal{L}}_6^{(2)} & \leq C\eta \int_{\mathbb{R}^{12}} \chi^{\leq L} a_x^* a_y^* a_z^* a_{x'}^* (\mathcal{N} + 1)^{-3/2} a_x a_y a_z a_{x'} dx dy dz dx' \\ & \quad + C\eta^{-1} \int_{\mathbb{R}^3} \chi^{\leq L-3} \left(\int_{\mathbb{R}^9} N^3 V_N(x-y, x-z)^2 \omega_N(x', y, z)^2 \right. \\ & \quad \quad \left. \times |\varphi(x')|^2 |\varphi(y)|^2 |\varphi(z)|^2 dy dz dx' \right) a_x^* (\mathcal{N} + 1)^{3/2} a_x dx \\ & \leq C \left(\eta + \eta^{-1} N^{1/2} \|\varphi\|_\infty^4 \left\| N^{3/2} \omega_N \right\|_2^2 \left\| |\varphi|^2 * N^{3/2} \mathbf{1}_{|\cdot| \leq CN^{-1/2}} \right\|_\infty \right) (\mathcal{N} + 1)^{5/2} \\ & \leq CN^{1/4} (\mathcal{N} + 1)^{5/2}, \end{aligned}$$

where we used that $V_N(x, y, z) \leq CN \mathbf{1}_{|x-z| \leq CN^{-1/2}}$ and optimized over η .

On another hand, and similarly, we have for all $\eta > 0$ that

$$\begin{aligned} \tilde{\mathcal{L}}_6^{(3)} & \leq C\eta \int_{\mathbb{R}^{15}} a_x^* a_y^* a_z^* a_{x'}^* a_{y'}^* (\mathcal{N} + 1)^{-3/2} a_x a_y a_z a_{x'} a_{y'} dx dy dz dx' dy' \\ & \quad + C\eta^{-1} \int_{\mathbb{R}^{15}} N^5 V(N^{1/2}(x-y, x-z))^2 \omega_N(x', y', z)^2 |\varphi(x')|^2 |\varphi(y')|^2 |\varphi(z)|^2 \\ & \quad \quad \quad \times a_x^* a_y^* (\mathcal{N} + 1)^{3/2} a_x a_y dx dy dz dx' dy' \end{aligned}$$

$$\begin{aligned} &\leq C \left(\eta + \eta^{-1} N^{1/2} \|\varphi\|_\infty^4 \left\| N^{3/2} \omega_N \right\|_2^2 \left\| |\varphi|^2 * N^{3/2} \mathbf{1}_{|\cdot| \leq CN^{-1/2}} \right\|_\infty \right) (\mathcal{N} + 1)^{7/2} \\ &\leq CN^{1/4} (\mathcal{N} + 1)^{7/2}, \end{aligned}$$

where we also optimized over η to obtain the last inequality.

Finally, to access the contribution of $\tilde{\mathcal{L}}_6^{(1)}$, we apply once more time the Duhamel formula:

$$\int_0^1 e^{-sB} \tilde{\mathcal{L}}_6^{(1)} e^{sB} ds = \tilde{\mathcal{L}}_6^{(1)} - \int_0^1 \int_0^s e^{-uB} [B, \tilde{\mathcal{L}}_6^{(1)}] e^{uB} du ds.$$

As before, the first term vanishes when tested against the vacuum state Ω and we focus therefore on estimating the second term. Let us decompose $\tilde{\mathcal{L}}_6^{(1)} =: \chi^{\leq L} \tilde{\mathcal{L}}_6^{(1,\dagger)} + \tilde{\mathcal{L}}_6^{(1,\circ)} \chi^{\leq L}$ and use that $[B_1 \chi^{\leq L}, \tilde{\mathcal{L}}_6^{(1,\circ)} \chi^{\leq L}] = 0$. We obtain

$$\begin{aligned} -[B, \tilde{\mathcal{L}}_6^{(1)}] &= [B_1 \chi^{\leq L} - B_1^* \chi^{\leq L}, \tilde{\mathcal{L}}_6^{(1,\circ)} \chi^{\leq L}] + h.c. = [\tilde{\mathcal{L}}_6^{(1,\circ)} \chi^{\leq L}, \chi^{\leq L} B_1^*] + h.c. \\ &= \chi^{\leq L-3} [\tilde{\mathcal{L}}_6^{(1,\circ)}, B_1^*] + \partial \chi^{\leq L} B_1^* \tilde{\mathcal{L}}_6^{(1,\circ)} + h.c. \\ &\leq [\tilde{\mathcal{L}}_6^{(1,\circ)}, B_1^*] + h.c. + C \frac{N}{L^k} \mathcal{N}^{3+k}, \end{aligned}$$

where we applied the Cauchy–Schwarz inequality and used

$$B_1^* B_1 \leq C (\mathcal{N} + 1)^3, \quad \tilde{\mathcal{L}}_6^{(1,\dagger)} \tilde{\mathcal{L}}_6^{(1,\circ)} \leq CN^2 (\mathcal{N} + 1)^3, \quad \text{and} \quad \partial \chi^{\leq L} \leq \frac{\mathcal{N}^k}{L^k},$$

for any $k \geq 0$. In view of (116), we have

$$\begin{aligned} [\tilde{\mathcal{L}}_6^{(1,\circ)}, B_1^*] &= 6\alpha^2 \int_{\mathbb{R}^9} N^3 (V_N \omega_N^2)(x, y, z) |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz \\ &+ 18\alpha^2 N^3 \int_{\mathbb{R}^{12}} (V_N \omega_N)(x, y, z) \omega(x, y, z') |\varphi(x)|^2 |\varphi(y)|^2 \varphi(z) \overline{\varphi(z')} a_z^* a_{z'} dx dy dz dz' \\ &+ 9\alpha^2 N^3 \int_{\mathbb{R}^{15}} (V_N \omega_N)(x, y, z) \omega(x, y', z') |\varphi(x)|^2 \varphi(y) \varphi(z) \overline{\varphi(y')} \overline{\varphi(z')} \\ &\quad \times a_y^* a_z^* a_{y'} a_{z'} dx dy dy' dz dz' \end{aligned}$$

hence

$$\begin{aligned} &[\tilde{\mathcal{L}}_6^{(1,\circ)}, B_1^*] \\ &\leq 6\alpha^2 \int_{\mathbb{R}^9} N^3 (V_N \omega_N^2)(x, y, z) |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz + C(N^{1/2} \mathcal{N} + \mathcal{N}^2), \end{aligned} \tag{125}$$

where we used that $V_N \leq \|V\|_{L^\infty(\mathbb{R}^3)} N$ and the computations done in the proof of Lemma 20. This concludes the proof of Lemma 25. \square

6.2.5. Concluding the proof of Theorem 18: summing up the leading contributions. In Lemmas 23, 24, and 25, we extracted the contributions of $d\Gamma(-\Gamma)$, \mathcal{L}_3 and \mathcal{L}_6 to the leading order, adding the contribution from \mathcal{L}_0 and controlling the remainder terms using Lemma 22, we obtain

$$\langle \Omega, e^{-B} W(\sqrt{N}\varphi)^* \mathbb{H}_N W(\sqrt{N}\varphi) e^B \Omega \rangle$$

$$\begin{aligned}
&\leq \frac{N}{6} \int_{\mathbb{R}^9} (N^2 V_N f_N)(x, y, z) |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz \\
&+ \int_{\mathbb{R}^9} \left(6\alpha^2 (f_N + \omega_N) + 2\alpha + \frac{1}{6} \right) (N^2 V_N \omega_N)(x, y, z) |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz \\
&\quad + N \int_{\mathbb{R}^3} (|\nabla \varphi(x)|^2 + V_{\text{ext}}(x) |\varphi(x)|^2) dx + \mathcal{O}(N^{1/2}) \\
&\leq N \mathcal{E}_{\text{GP}}(\varphi) + \mathcal{O}(N^{1/2}),
\end{aligned}$$

where we used that $f + \omega = 1$, that $\alpha = -1/6$ and that

$$\frac{1}{6} \int_{\mathbb{R}^9} (N^2 V_N f_N)(x, y, z) |\varphi(x)|^2 |\varphi(y)|^2 |\varphi(z)|^2 dx dy dz \leq \frac{b(V)}{6} \int_{\mathbb{R}^3} |\varphi(x)|^6 dx,$$

from $b(V) = \int V f = \|V f\|_{L^1}$ and the Hölder inequality. This concludes the proof of Theorem 18.

7. CONVERGENCE OF GROUND STATES

The convergence of states in Theorem 2 follows from a standard Hellmann–Feynman argument in [21, 22]. Let us quickly explain the proof for the reader's convenience. Let Ψ_N be a normalized state in \mathfrak{H}^N satisfying

$$\lim_{N \rightarrow \infty} \frac{\langle \Psi_N, H_N \Psi_N \rangle}{N} = e_{\text{GP}}. \quad (126)$$

Let us prove that $\text{Tr}(Q \gamma_{\Psi_N}^{(1)}) \rightarrow 0$ where $Q = 1 - |u_0\rangle \langle u_0|$ with u_0 the unique minimizer of e_{GP} . For every $\varepsilon > 0$ we introduce the perturbed Hamiltonian

$$H_N^\varepsilon := H_N - \varepsilon \sum_{i=1}^N Q_i \quad (127)$$

and the associated ground state energy

$$E^\varepsilon(N) := \inf_{\|\Psi\|_{L^2(\mathbb{R}^{3N})} = 1} \langle \Psi, H_N^\varepsilon \Psi \rangle.$$

By following the proof of the energy lower bound on E_N (Theorem 13), we obtain

$$\frac{E^\varepsilon(N)}{N} \geq e_{\text{GP}}^\varepsilon - C_\varepsilon o(1)_{N \rightarrow \infty},$$

where

$$e_{\text{GP}}^\varepsilon := \inf_{\|u\|_2=1} \mathcal{E}_{\text{GP}}^\varepsilon(u), \quad \mathcal{E}_{\text{GP}}^\varepsilon(u) := \mathcal{E}_{\text{GP}}(u) - \varepsilon \langle u, Qu \rangle. \quad (128)$$

Combining with the assumption (126) we get

$$\begin{aligned}
\text{Tr}[Q \gamma_{\Psi_N}^{(1)}] &= \frac{1}{N} \left\langle \Psi_N, \sum_{i=1}^N Q_i \Psi_N \right\rangle = \frac{1}{\varepsilon} \left[\frac{1}{N} \langle \Psi_N, H_N \Psi_N \rangle - \frac{1}{N} \langle \Psi_N, H_N^\varepsilon \Psi_N \rangle \right] \\
&\leq \frac{1}{\varepsilon} [e_{\text{GP}} - e_{\text{GP}}^\varepsilon + C_\varepsilon o(1)_{N \rightarrow \infty}],
\end{aligned}$$

namely

$$\limsup_{N \rightarrow +\infty} \text{Tr}[Q \gamma_{\Psi_N}^{(1)}] \leq \frac{e_{\text{GP}} - e_{\text{GP}}^\varepsilon}{\varepsilon}, \quad \forall \varepsilon > 0. \quad (129)$$

Now let us compute the right-hand side of (129) in the limit $\varepsilon \rightarrow 0^+$. By a standard compactness method (using $V_{\text{ext}}(x) \rightarrow \infty$ as $|x| \rightarrow \infty$), we can show that $e_{\text{GP}}^\varepsilon$ has a (L^2 -normalized) minimizer u_ε . Since

$$e_{\text{GP}}^\varepsilon \geq e_{\text{GP}}^\varepsilon = \mathcal{E}_{\text{GP}}(u_\varepsilon) - \varepsilon \langle u_\varepsilon, Qu_\varepsilon \rangle \geq \mathcal{E}_{\text{GP}}(u_\varepsilon) - \varepsilon \geq e_{\text{GP}} - \varepsilon$$

we find that $\{u_\varepsilon\}_{\varepsilon \rightarrow 0}$ is a minimizing sequence for e_{GP} . Consequently, u_ε converges to the unique minimizer u_0 of e_{GP} in $L^2(\mathbb{R}^3)$. Therefore, $\langle u_\varepsilon, Qu_\varepsilon \rangle \rightarrow 0$ when $\varepsilon \rightarrow 0$. Using again the variational principle

$$e_{\text{GP}}^\varepsilon = \mathcal{E}_{\text{GP}}(u_\varepsilon) - \varepsilon \langle u_\varepsilon, Qu_\varepsilon \rangle \geq e_{\text{GP}} - \varepsilon \langle u_\varepsilon, Qu_\varepsilon \rangle$$

we conclude from (129) that

$$\limsup_{N \rightarrow +\infty} \text{Tr}[Q\gamma_{\Psi_N}^{(1)}] \leq \frac{e_{\text{GP}} - e_{\text{GP}}^\varepsilon}{\varepsilon} \leq \langle u_\varepsilon, Qu_\varepsilon \rangle \xrightarrow{\varepsilon \rightarrow 0} 0.$$

Strictly speaking, $\text{Tr}(Q\gamma_{\Psi_N}^{(1)}) \rightarrow 0$ is equivalent to the fact that $\gamma_{\Psi_N}^{(1)} - |u_0\rangle\langle u_0| \rightarrow 0$ in the operator norm, while the statement in Theorem 1 is concerned the trace class convergence. However, by a well-known argument of Robert Seiringer, since $\gamma_{\Psi_N}^{(1)} - |u_0\rangle\langle u_0|$ has trace 0 and has at most one negative eigenvalue (as $\gamma_{\Psi_N}^{(1)} \geq 0$ and $|u_0\rangle\langle u_0|$ is rank-one), we find that its trace norm is equal to twice of its operator norm.

REFERENCES

- [1] M. H. ANDERSON, J. R. ENSHER, M. R. MATTHEWS, C. E. WIEMAN, AND E. A. CORNELL, *Observation of Bose–Einstein Condensation in a Dilute Atomic Vapor*, Science, 269 (1995), pp. 198–201.
- [2] N. BENEDIKTER, M. PORTA, AND B. SCHLEIN, *Effective evolution equations from quantum dynamics*, vol. 7 of SpringerBriefs in Mathematical Physics, Springer, Cham, 2016.
- [3] R. N. BISSET, AND P. B. BLAKIE, *Crystallization of a dilute atomic dipolar condensate*, Phys. Rev. A, 92 (2015), pp. 061603.
- [4] N. N. BOGOLIUBOV, *About the theory of superfluidity*, Izv. Akad. Nauk SSSR, 11 (1947), p. 77.
- [5] BOSE, *Plancks Gesetz und Lichtquantenhypothese*, Z. Physik, 26 (1924), pp. 178–181.
- [6] P. B. BLAKIE *Properties of a dipolar condensate with three-body interactions*, Phys. Rev. A, 93 (2016), pp. 033644.
- [7] H. P. BÜCHLER, A. MICHELI, AND P. ZOLLER, *Three-body interactions with cold polar molecules* Nature Physics, 3 (2007), pp. 726–731.
- [8] X. CHEN, *Second order corrections to mean field evolution for weakly interacting bosons in the case of three-body interactions*, Arch. Ration. Mech. Anal., 203 (2012), pp. 455–497.
- [9] X. CHEN AND J. HOLMER, *The derivation of the \mathbb{T}^3 energy-critical NLS from quantum many-body dynamics*, Invent. math., 217 (2019), pp. 433–547.
- [10] T. CHEN, AND N. PAVLOVIĆ, *The quintic NLS as the mean field limit of a Boson gas with three-body interactions*, J. Funct. Anal., 260 (2011), pp. 959–997.
- [11] K. B. DAVIS, M. O. MEWES, M. R. ANDREWS, N. J. VAN DRUTEN, D. S. DURFEE, D. M. KURN, AND W. KETTERLE, *Bose–Einstein Condensation in a Gas of Sodium Atoms*, Phys. Rev. Lett., 75 (1995), pp. 3969–3973.
- [12] J. DOLBEAULT, P. FELMER, M. LOSS, AND E. PATUREL, *Lieb–Thirring type inequalities and Gagliardo–Nirenberg inequalities for systems*, J. Funct. Anal., 238 (2006), pp. 193–220.
- [13] A. EINSTEIN, *Quantentheorie des einatomigen idealen Gases*, Sitzber. Kgl. Preuss. Akad. Wiss. (1924), pp. 261–267.
- [14] L. ERDŐS, B. SCHLEIN, AND H.-T. YAU, *Derivation of the nonlinear Schrödinger equation from a many body Coulomb system*, , Adv. Theor. Math. Phys., 5 (2001), pp. 1169–1205
- [15] L. LANDAU, *Theory of the Superfluidity of Helium II*, Phys. Rev., 60 (1941), pp. 356–358.
- [16] J. LEE, *Rate of convergence towards mean-field evolution for weakly interacting bosons with singular three-body interactions.*, Preprint arXiv:2006.13040, (2020).
- [17] M. LEWIN, P. T. NAM, AND N. ROUGERIE, *Derivation of Hartree’s theory for generic mean-field Bose systems*, Adv. Math., 254 (2014), pp. 570–621.

- [18] ———, *The mean-field approximation and the non-linear Schrödinger functional for trapped Bose gases*, Trans. Amer. Math. Soc., 368 (2016), pp. 6131–6157.
- [19] Y. LI, AND F. YAO, *Derivation of the nonlinear Schrödinger equation with a general nonlinearity and Gross–Pitaevskii hierarchy in one and two dimensions*, J. Math. Phys., 62 (2021), p. 021505.
- [20] E. H. LIEB AND M. LOSS, *Analysis*, vol. 14 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, second ed., 2001.
- [21] E. H. LIEB AND R. SEIRINGER, *Proof of Bose–Einstein Condensation for Dilute Trapped Gases*, Phys. Rev. Lett., 88 (2002), p. 170409.
- [22] ———, *Derivation of the Gross–Pitaevskii equation for rotating Bose gases*, Comm. Math. Phys., 264 (2006), pp. 505–537.
- [23] E. H. LIEB, R. SEIRINGER, AND J. P. SOLOVEJ, *Ground-state energy of the low-density Fermi gas*, Phys. Rev. A, 71 (2005), p. 053605.
- [24] E. H. LIEB, R. SEIRINGER, AND J. YNGVASON, *Bosons in a trap: A rigorous derivation of the Gross–Pitaevskii energy functional*, Phys. Rev. A, 61 (2000), p. 043602.
- [25] E. M. MAS, R. BUKOWSKI AND K. SZALEWICZ, , *Ab initio three-body interactions for water. II. Effects on structure and energetics of liquid*. J. Chem. Phys., 118 (2003), pp. 4404–4413.
- [26] R. D. MURPHY, AND J. A. BARKER, *Three-Body Interactions in Liquid and Solid Helium* Phys. Rev. A, 3 (1971), pp. 1037-1040.
- [27] P. T. NAM, M. NAPIÓRKOWSKI, J. RICAUD, AND A. TRIAY, *Optimal rate of condensation for trapped bosons in the Gross–Pitaevskii regime*, Anal. PDE, in press (2021).
- [28] P. T. NAM, N. ROUGERIE, AND R. SEIRINGER, *Ground states of large bosonic systems: the Gross–Pitaevskii limit revisited*, Anal. PDE, 9 (2016), pp. 459–485.
- [29] P. T. NAM AND R. SALZMANN, *Derivation of 3D Energy-Critical Nonlinear Schrödinger Equation and Bogoliubov Excitations for Bose Gases*, Commun. Math. Phys., 375 (2020), pp. 495–571.
- [30] P. T. NAM AND A. TRIAY, *Derivation of the 3D energy-critical nonlinear Schrödinger equation from dilute Bose gases*, Preprint 2021.
- [31] D. S. PETROV, *Three-body interacting bosons in free space*, Phys. Rev. Lett., 112 (2014), pp. 103201.
- [32] P. A. PIENIAZEK, C. J. TAINTER AND J. L. SKINNER *Surface of liquid water: Three-body interactions and vibrational sum-frequency spectroscopy*, J. Am. Chem. Soc. 133 (2011), pp. 10360-10363.
- [33] J. P. SOLOVEJ, *Upper bounds to the ground state energies of the one- and two-component charged Bose gases*, Comm. Math. Phys., 266 (2006), pp. 797–818.
- [34] S. UJEVIC AND S. A. VITIELLO, *Three-body interactions in the condensed phases of helium atom systems* J. Phys.: Condens. Matter 19 (2007) p. 116212
- [35] J. YUAN, *Derivation of the Quintic NLS from many-body quantum dynamics in \mathbb{T}^2* , Comm. Pure Appl. Anal., 14 (2015), p. 1941

DEPARTMENT OF MATHEMATICS, LMU MUNICH, THERESIENSTRASSE 39, 80333 MUNICH, AND MUNICH CENTER FOR QUANTUM SCIENCE AND TECHNOLOGY, SCHELLINGSTR. 4, 80799 MUNICH, GERMANY

Email address: `nam@math.lmu.de`

DEPARTMENT OF MATHEMATICS, LMU MUNICH, THERESIENSTRASSE 39, 80333 MUNICH, AND MUNICH CENTER FOR QUANTUM SCIENCE AND TECHNOLOGY, SCHELLINGSTR. 4, 80799 MUNICH, GERMANY

Email address: `julien.ricaud@polytechnique.edu`

DEPARTMENT OF MATHEMATICS, LMU MUNICH, THERESIENSTRASSE 39, 80333 MUNICH, AND MUNICH CENTER FOR QUANTUM SCIENCE AND TECHNOLOGY, SCHELLINGSTR. 4, 80799 MUNICH, GERMANY

Email address: `triay@math.lmu.de`