

Exact big Ramsey degrees via coding trees

Martin Balko¹, David Chodounský¹, Natasha Dobrinen², Jan Hubička¹,
Matěj Konečný¹, Lluís Vena³, and Andy Zucker⁴

¹*Department of Applied Mathematics (KAM), Charles University, Malostranské náměstí 25,
Praha 1, Czech Republic*

²*Department of Mathematics, University of Denver, C.M. Knudson Hall 302, Denver, USA*

³*Universitat Politècnica de Catalunya, Barcelona, Spain*

⁴*University of California San Diego, La Jolla, USA*

October 19, 2021

Abstract

We characterize the big Ramsey degrees of free amalgamation classes in finite binary languages defined by finitely many forbidden irreducible substructures, thus refining the recent upper bounds given by Zucker. Using this characterization, we show that the Fraïssé limit of each such class admits a big Ramsey structure. Consequently, the automorphism group of each such Fraïssé limit has a metrizable universal completion flow.

Contents

1	Introduction	3
2	Enumerated structures and Ramsey degrees	7
3	\mathcal{L}-structures and coding trees	9
4	Posets of classes of structures	11
4.1	Gluing	11
4.2	The poset $P(\rho)$	12
4.3	Operations on structures and classes	16
4.4	The spectrum of a gluing and a class	19
4.5	Consecutive age changes	21
4.6	Free relations	23
5	Diagonal substructures	26
5.1	Aged embeddings, envelopes, and closures	26
5.2	Weakly diagonal substructures	29
5.3	Controlled paths	32
5.4	Insertions	35
5.5	Controlled age-change levels	38
5.6	Controlled coding levels	41
5.7	Controlled splitting levels	46
5.8	Diagonal substructures	51
6	Diaries	53
6.1	Concrete diaries	53
6.2	Diagonal Diaries	56
6.3	Large subsets of coding nodes	62
6.4	The construction	66
7	Degenerate unary predicates	71
8	Big Ramsey structures	73

Overview

A central theme in the theory of infinite structures is the investigation of which structures have partition properties resembling the Infinite Ramsey Theorem [26]. Given a finite structure \mathbf{A} which embeds into an infinite structure \mathbf{K} , the program asks whether there is a positive integer ℓ so that whenever the set of embeddings of \mathbf{A} into \mathbf{K} is colored into finitely many colors, there is a subcopy of \mathbf{K} in which all embeddings take no more than ℓ colors. When such a bound exists, the least such integer is called the *big Ramsey degree* of \mathbf{A} in \mathbf{K} .

While finite structural Ramsey theory, which only looks for arbitrarily large, *finite* structures where the coloring takes one or few colors, has seen rapid progress over the past several decades, infinite structural Ramsey theory has seen a slower development. This has been due to the difficulty of preserving an isomorphic copy of an infinite structure while reducing the number of colors, and the lack of methods able to handle forbidden substructures. However, the recent development of new methods in [8], [9], and [32] have successively secured upper bounds for the Henson k -clique-free graphs and more generally for Fraïssé limits of the following free amalgamation classes:

Let \mathcal{L} be a finite language with relation symbols of arity at most two. Given a finite collection \mathcal{F} of finite irreducible \mathcal{L} -structures, let $\text{Forb}(\mathcal{F})$ denote the class of all finite \mathcal{L} -structures into which no member of \mathcal{F} embeds. Letting \mathbf{K} denote the Fraïssé limit of $\text{Forb}(\mathcal{F})$, and utilizing the upper bounds due to Zucker in [32], we characterize the big Ramsey degree of each structure $\mathbf{A} \in \text{Forb}(\mathcal{F})$ inside \mathbf{K} . Our characterization uses a notion of *diagonal diary*, from which one can construct an algorithm to calculate the exact big Ramsey degrees. Moreover, the diagonal diary encodes the additional structure necessary for a so-called big Ramsey structure (Definition 8.0.1), from which we obtain the following theorem:

Theorem. *With \mathcal{F} and \mathcal{K} as above, let $\mathbf{K} = \text{Flim}(\mathcal{K})$ be the Fraïssé limit of \mathcal{K} , and write $G = \text{Aut}(\mathbf{K})$. Then we exactly characterize the big Ramsey degrees of every structure in \mathcal{K} . As a consequence, \mathbf{K} admits a big Ramsey structure, and the topological group G has a metrizable universal completion flow, which is unique up to isomorphism.*

1 Introduction

The field of Ramsey theory was established by the following celebrated result.

Theorem 1.0.1 (Infinite Ramsey Theorem, [26]). *Given positive integers m and r , suppose the collection of all m -element subsets of \mathbb{N} is colored by r colors. Then there is an infinite set N of natural numbers such that all m -element subsets of N have the same color.*

From this, Ramsey deduced the following finite version.

Theorem 1.0.2 (Finite Ramsey Theorem, [26]). *Given positive integers m, n, r with $m \leq n$, there is an integer $p > n$ such that for any coloring of the m -element subsets of p into r colors, there is a subset $N \subseteq p$ of cardinality n such that all m -element subsets of N have the same color.*

Structural Ramsey theory investigates which classes of structures satisfy versions of Theorems 1.0.1 or 1.0.2. Given structures \mathbf{A} and \mathbf{B} , we write $\mathbf{A} \leq \mathbf{B}$ exactly when there is an embedding of \mathbf{A} into \mathbf{B} . A substructure \mathbf{A}' of \mathbf{B} is called a *copy* of \mathbf{A} if \mathbf{A}' is the image of some embedding of \mathbf{A} into \mathbf{B} . The collection of all copies of \mathbf{A} in \mathbf{B} is denoted by $\binom{\mathbf{B}}{\mathbf{A}}$. Given structures $\mathbf{A}, \mathbf{B}, \mathbf{C}$ with $\mathbf{A} \leq \mathbf{B} \leq \mathbf{C}$ and an integer $r \geq 1$, the Erdős–Rado partition arrow

$$\mathbf{C} \xrightarrow{c} (\mathbf{B})_r^{\mathbf{A}} \quad (1)$$

signifies that for each coloring $\gamma: \binom{\mathbf{C}}{\mathbf{A}} \rightarrow r$, there is a $\mathbf{B}' \in \binom{\mathbf{C}}{\mathbf{B}}$ for which γ takes only one color on $\binom{\mathbf{B}'}{\mathbf{A}}$. The c over the arrow indicates that we color the *copies* of \mathbf{A} in \mathbf{C} ; we will introduce several variants in Section 2. A class \mathcal{K} of structures is said to have the *Ramsey property* if given $\mathbf{A}, \mathbf{B} \in \mathcal{K}$ with $\mathbf{A} \leq \mathbf{B}$, for any positive integer r , there is some $\mathbf{C} \in \mathcal{K}$ for which $\mathbf{B} \leq \mathbf{C}$ and $\mathbf{C} \xrightarrow{c} (\mathbf{B})_r^{\mathbf{A}}$. An equivalent and illustrative form of the Ramsey property for Fraïssé classes is the following. Letting \mathbf{K} denote the Fraïssé limit of a Fraïssé class \mathcal{K} , \mathcal{K} has the *Ramsey property* if for any two structures $\mathbf{A} \leq \mathbf{B}$ in \mathcal{K} and any positive integer r ,

$$\mathbf{K} \xrightarrow{c} (\mathbf{B})_r^{\mathbf{A}} \quad (2)$$

The Ramsey theory of finite structures has a long and rich history. It was initiated in the 1970's when Nešetřil and Rödl, and independently Abramson and Harrington, proved that the class of all finite ordered graphs has the Ramsey property [1, 22]. In fact, the Nešetřil–Rödl theorem is much stronger, implying the Ramsey property for classes of all finite linear orderings of members of any relational free amalgamation class. In 2005, the Kechris–Pestov–Todorčević correspondence between the Ramsey property of \mathcal{K} and extreme amenability of the automorphism group of its Fraïssé limit [16] and Nešetřil's initiation of the classification program of Ramsey classes [21] boosted the development of the area which led to discoveries of many more classes with the Ramsey property culminating with the general theorem of Hubička and Nešetřil [15].

Generalizing further, while most classes of finite unordered structures do not have the Ramsey property, often equipping the class with an additional linear order or finitely many new relations produces the Ramsey property. When this is the case, the reducts of these expansions then retain a weaker property, called small Ramsey degrees. Here the term *small* refers to the fact that we are looking for copies of finite structures \mathbf{B} where the coloring attains one or few colors, while the term *big* is used to signify that we are interested in copies of the infinite structure \mathbf{K} where the coloring attains one or few colors. A Fraïssé class \mathcal{K} with Fraïssé limit \mathbf{K} has *finite small Ramsey degrees* if for each $\mathbf{A} \in \mathcal{K}$, there is a number ℓ such that for all $r \geq \ell$, we have

$$\mathbf{K} \xrightarrow{c} (\mathbf{B})_{r,\ell}^{\mathbf{A}} \quad (3)$$

For instance, the class of finite ordered graphs has the Ramsey property, while the class of finite graphs has finite small Ramsey degrees. One way to show that a Fraïssé class \mathcal{K} has finite small Ramsey degrees, developed in [16] and [24], is to show that \mathcal{K} admits a *pre-compact expansion class* \mathcal{K}^* which has the Ramsey property. For the class of finite graphs, this expansion class is the class of finite ordered graphs. So in a sense that can be made precise, it is because the class of finite

ordered graphs is Ramsey that the class of finite graphs has finite small Ramsey degrees. Zucker shows in [30] that in fact, every Fraïssé class \mathcal{K} with finite small Ramsey degrees admits a pre-compact expansion class with the Ramsey property.

When considering big Ramsey degrees instead of small, things become more complex. While many infinite structures have been shown to be indivisible, i.e. satisfying a pigeonhole principle for vertex partitions, it was known early on that exact analogues usually fail for colorings of structures of size two or more. Even in the simple case of the rationals, it was known from work of Sierpiński ([29]) that there is a coloring of pairs of rationals such that both colors persist in every subcopy of the rationals forming a dense linear order. However, it is sometimes possible to bound the number of colors which persist in some substructure isomorphic to the original infinite structure, leading to the following question.

Given an infinite structure \mathbf{K} and a finite substructure \mathbf{A} , is there a positive integer ℓ such that for any coloring of all copies of \mathbf{A} in \mathbf{K} into finitely many colors, there is a substructure of \mathbf{K} isomorphic to \mathbf{K} in which all copies of \mathbf{A} take no more than ℓ colors?

We say that \mathcal{K} has *finite big Ramsey degrees* if for each $\mathbf{A} \in \mathcal{K}$, there is a number ℓ such that for all $r \geq \ell$, we have

$$\mathbf{K} \xrightarrow{c} (\mathbf{K})_{r,\ell}^{\mathbf{A}}. \quad (4)$$

The least such number ℓ , when it exists, is called the *big Ramsey degree* (for copies) of \mathbf{A} in \mathbf{K} . It is clear from the formulations that finite big Ramsey degrees implies finite small Ramsey degrees.

Although work on big Ramsey degrees has been progressing sporadically since its inception by Sierpiński in 1933, the field has gained momentum beginning with the characterization of exact big Ramsey degrees of the Rado graph in [18] and more recently, with new techniques developed in [8] to handle triangle-free graphs. We refer to the paper [10] for comprehensive background on historical and recent results on big Ramsey degrees. This paper extends and concludes work in [8], [9], [11], [12], [13], [14], [17], [18], [25], [28], and [32] for free amalgamation classes with relations of arity at most two and finitely many forbidden irreducible substructures.

Let \mathcal{L} be a finite language with relation symbols of arity at most two. An \mathcal{L} -structure \mathbf{F} is *irreducible* if each pair of vertices in the universe of \mathbf{F} are in some relation of \mathbf{F} . Given a finite collection \mathcal{F} of finite irreducible \mathcal{L} -structures, let $\text{Forb}(\mathcal{F})$ denote the class of all finite \mathcal{L} -structures into which no member of \mathcal{F} embeds. Letting \mathbf{K} denote the Fraïssé limit of $\text{Forb}(\mathcal{F})$, Zucker proved in [32] that each structure in $\text{Forb}(\mathcal{F})$ has finite big Ramsey degree in \mathbf{K} .

The goal of this article is to characterize these degrees exactly, allowing a dynamical interpretation. Big Ramsey structures (see Definition 8.0.1) and completion flows were introduced in [31] as an attempt to connect big Ramsey degrees to dynamical phenomena, much as small Ramsey degrees are connected the behavior of minimal flows (see [16, 30]). Asserting that a Fraïssé limit $\mathbf{K} = \text{Flim}(\mathcal{K})$ admits a big Ramsey structure is stronger than asserting that the structures of \mathcal{K} have finite big Ramsey degrees. More-or-less, to say that \mathbf{K} admits a big Ramsey structure means that each $\mathbf{A} \in \mathcal{K}$ has finite big Ramsey degree *and* that the big Ramsey

behavior of every $\mathbf{A} \in \mathcal{K}$ can be described simultaneously. While for small Ramsey degrees, such a “simultaneous” description is an easy consequence of compactness, when working with big Ramsey degrees, we lose compactness. Nonetheless, in every known example where the big Ramsey degrees are finite and known exactly, the Fraïssé limit does admit a big Ramsey structure, and it is an open question whether finite big Ramsey degrees and admitting a big Ramsey structure are in fact equivalent.

The following is the main result of this paper.

Theorem 1.0.3. *Let \mathcal{L} be a finite binary language. Let \mathcal{F} be a finite set of finite, irreducible \mathcal{L} -structures, and set $\mathcal{K} = \text{Forb}(\mathcal{F})$. Let $\mathbf{K} = \text{Flim}(\mathcal{K})$ be the Fraïssé limit of \mathcal{K} , and write $G = \text{Aut}(\mathbf{K})$. Then \mathbf{K} admits a big Ramsey structure, and the topological group G has a metrizable universal completion flow, which is unique up to isomorphism.*

The proof of the theorem relies on a precise understanding of the big Ramsey degrees of every $\mathbf{A} \in \mathcal{K}$, which we develop in sections 5 and 6.

The main concepts behind our characterization are now presented, referring to sections where the ideas are precisely developed. After this, we will provide the characterization for the big Ramsey degrees of triangle-free graphs, as this example illustrates what is needed to work with forbidden substructures.

Exact big Ramsey degree results in [18] for the Rado graph utilized the tree of all finite binary sequences to represent a universal structure into which the Rado graph embeds. After applications of a Ramsey theorem for trees due to Milliken [20], an enumerated copy of the Rado graph represented by a so-called digonal antichain of nodes in the tree was extracted and shown to provide optimal results on big Ramsey degrees. Here, rather than working with universal structures, we work with enumerated structures (Section 2) and coding trees (Section 3) representing them. This approach is especially conducive to investigating Ramsey properties of structures which forbid certain substructures. Further, precisely because of forbidden substructures, it becomes necessary to keep track of those structures which are allowed to extend a given finite structure as coded in a tree. This leads to the notions of posets of classes of structures, gluings, and consecutive age changes, delineated in Section 4.

A key notion appearing so far in all results on big Ramsey degrees of binary relational structures is the notion of diagonal antichain, first seen in [7] for a characterization of the big Ramsey degrees of the rationals, and extended more generally in [28] for the Rado graph and similar structures. This notion forms the foundation of our characterization of big Ramsey degrees. However, in the presence of forbidden substructures, we need the more robust notion of diagonal substructure provided in Section 5. Diagonal substructures require the relationships between a substructure \mathbf{A} and an enumerated Fraïssé structure \mathbf{K} to change in a slow, controlled manner. Theorem 5.8.5 shows that each enumerated Fraïssé structure contains a diagonal substructure isomorphic to the whole Fraïssé limit. Paired with the main results from [32], this diagonal substructure yields upper bounds on the big Ramsey degrees, which are then shown to be exact in Section 6.

Diagonal diaries, defined in Section 6, are an abstraction of diagonal substructures which represent the essential relationships between an enumerated Fraïssé structure and a given substructure. Big Ramsey degrees are exactly characterized in Theorem 6.2.14 as the number of different diagonal diaries representing \mathbf{A} . From these diaries, one can derive an algorithm that calculates the degrees. Theorems 8.0.2 and 8.0.6 conclude the main results of this paper for big Ramsey structures.

While the main result of this paper requires much preamble to make precise, the special case of triangle-free graphs displays several aspects of the characterization necessary when forbidding substructures, and can be described in a relatively straightforward manner; for another description of the diagonal diaries of finite triangle-free graphs, see Example 6.2.10.

Let \mathcal{G}_3 denote the Fraïssé class of finite triangle-free graphs, and let \mathbf{G}_3 denote its Fraïssé limit with vertex set \mathbb{N} . For $I \subseteq \mathbb{N}$, let $\mathbf{G}_3 \upharpoonright I$ denote the induced subgraph of \mathbf{G}_3 on universe I , and given $x \in \mathbb{N}$, let $N(x)$ denote the \mathbf{G}_3 -neighbors of x . Given a finite induced subgraph $\mathbf{A} \subseteq \mathbf{G}_3$, we say that \mathbf{A} is *weakly diagonal* (Definition 5.2.2) if the following hold: For each pair $i < j$ in \mathbf{A} there is an $m(i, j) < i$ such that the set $\{m(i, j) : i < j \in \mathbf{A}\}$ is disjoint from \mathbf{A} , and $N(i) \cap m(i, j) = N(j) \cap m(i, j)$, while $N(i) \cap (m(i, j) + 1) \neq N(j) \cap (m(i, j) + 1)$. Furthermore, $\{m(i, j) : i < j \in \mathbf{A}\}$ has cardinality $2|\mathbf{A}| - 1$, and whenever we have $k \in \mathbf{A}$ with $k > m(i, j)$ and $k E m(i, j)$, then $N(k) \cap m(i, j) = N(i) \cap m(i, j)$. So far, this is the structural interpretation of the characterization of big Ramsey degrees in the Rado graph in [18]. For triangle-free graphs, there is one more requirement in the definition of a weakly diagonal substructure: For each $i \in \mathbf{A}$, we require that $i E m$ for some $m < i$; let m_i denote the least such m .

Diagonal substructures for triangle-free graphs involve two more requirements in addition to being weakly diagonal. First, for $i < j$ in \mathbf{A} , if $i \not E j$ and $m_j < i$, then there is some $n \in i$ such that $i E n$ and $j E n$, and the least such n , denoted $n(i, j)$ is not in $\mathbf{A} \cup \{m(i, j) : i, j \in \mathbf{A} \text{ and } i < j\}$. This is what we will call $i \in \mathbf{A}$ being a *controlled coding level* (Definition 5.6.5) of \mathbf{A} . Second, for each $i, j, k, \ell \in \mathbf{A}$ (not necessarily distinct) with $i < j$, $k < \ell$, $i \not E j$, $k \not E \ell$, $n_j < i$, $n_\ell < k$, and $n(i, j) = n(k, \ell) := n$, then we have $\{N(i) \cap n, N(j) \cap n\} = \{N(k) \cap n, N(\ell) \cap n\}$. This is equivalent to saying that \mathbf{A} has *controlled age-change levels* (Definition 5.5.1). Given a finite triangle-free graph \mathbf{A} , the big Ramsey degree of \mathbf{A} in \mathbf{G}_3 is the number of different diagonal substructures representing a copy of \mathbf{A} .

Notation

If $m < n < \omega$, we write $[m, n] = \{r < \omega : m \leq r \leq n\}$, and similarly for (m, n) , $(m, n]$, and $[m, n)$. We identify $m < \omega$ with the set $\{0, \dots, m-1\}$, though sometimes we will write the latter for emphasis. If I is a set, we sometimes write $\langle u_i : i \in I \rangle$ for the function σ with domain I and with $\sigma(i) = u_i$ for each $i \in I$.

2 Enumerated structures and Ramsey degrees

This section closely follows Section 1 of [32]. The key definition is the notion of *ordered Ramsey degree* appearing in Definition 2.0.2, where the order comes from

some choice of enumeration of the structures being considered. This is the form of big Ramsey degree which we work with in this paper. We also discuss how to translate between this notion of Ramsey degree and the more standard notions of big Ramsey degrees for embeddings and/or copies.

Definition 2.0.1. Suppose that \mathbf{K} is an infinite first-order structure, that \mathbf{A} is a finite structure with $\text{Emb}(\mathbf{A}, \mathbf{K}) \neq \emptyset$, and that $\ell < r < \omega$. We write

$$\mathbf{K} \rightarrow (\mathbf{K})_{r,\ell}^{\mathbf{A}}$$

if for any function $\chi: \text{Emb}(\mathbf{A}, \mathbf{K}) \rightarrow r$, there is $\eta \in \text{Emb}(\mathbf{K}, \mathbf{K})$ so that

$$|\text{Im}(\chi \cdot \eta)| = |\chi[\eta \cdot \text{Emb}(\mathbf{A}, \mathbf{K})]| \leq \ell.$$

The *Ramsey degree of \mathbf{A} in \mathbf{K}* , if it exists, is the least $\ell < \omega$ so that $\mathbf{K} \rightarrow (\mathbf{K})_{r,\ell}^{\mathbf{A}}$ holds for every $r > \ell$, or equivalently for $r = \ell + 1$.

When $\mathbf{K} = \text{Flim}(\mathcal{K})$ for some Fraïssé class \mathcal{K} , one says that $\mathbf{A} \in \mathcal{K}$ has *big Ramsey degree ℓ* exactly when \mathbf{A} has Ramsey degree ℓ in \mathbf{K} . We say that \mathcal{K} has *finite big Ramsey degrees* if every $\mathbf{A} \in \mathcal{K}$ has some finite big Ramsey degree.

Some authors formulate the above definitions with respect to *copies* rather than embeddings, where a copy is simply the image of an embedding. If \mathbf{A} has Ramsey degree ℓ in \mathbf{K} with respect to copies, then one can show that \mathbf{A} has Ramsey degree $\ell \cdot |\text{Aut}(\mathbf{A})|$ with respect to embeddings.

By considering *enumerated structures* and the *ordered embeddings* between them, we can minimize the distinction between copies and embeddings.

Definition 2.0.2. An *enumerated structure* is simply a structure \mathbf{K} whose underlying set is the cardinal $|\mathbf{K}|$, and we often identify \mathbf{K} and $|\mathbf{K}|$. In this paper, all enumerated structures will be countable. If $m < |\mathbf{K}|$, we write $\mathbf{K}_m := \mathbf{K} \cap m$.

We let $<$ denote the usual order on ω . If \mathbf{A} is another enumerated structure, then an *ordered embedding* of \mathbf{A} into \mathbf{K} is simply an embedding of $\langle \mathbf{A}, < \rangle$ into $\langle \mathbf{K}, < \rangle$. We denote the collection of ordered embeddings of \mathbf{A} into \mathbf{K} by $\text{OEmb}(\mathbf{A}, \mathbf{K})$.

The *ordered Ramsey degree* of \mathbf{A} in \mathbf{K} is just the Ramsey degree of $\langle \mathbf{A}, < \rangle$ in $\langle \mathbf{K}, < \rangle$.

The final goal of this section is to sharpen Proposition 1.3 from [32]. There, it is shown that given a Fraïssé class \mathcal{K} , then $\mathbf{A} \in \mathcal{K}$ has finite Ramsey degree iff every enumeration of \mathbf{A} has finite ordered Ramsey degree in an enumerated Fraïssé limit. Here, we wish to keep track of the exact degrees. The following notation will be helpful.

Notation 2.0.3. Suppose \mathbf{A} is a structure, S is a set, and $f: S \rightarrow \mathbf{A}$ is an injection. Then $\mathbf{A} \cdot f$ is the structure on S making f an embedding.

Proposition 2.0.4. *Let \mathcal{K} be a Fraïssé class, and suppose $\mathbf{K} = \text{Flim}(\mathcal{K})$ is enumerated. Fix $\mathbf{A} \in \mathcal{K}$, and write $B = \{\sigma_i : i < n\}$ for the set of bijections from $|\mathbf{A}|$ to \mathbf{A} . Suppose each enumerated structure $\mathbf{A} \cdot \sigma_i$ for $i < n$ has ordered Ramsey degree $\ell_i < \omega$ in \mathbf{K} . Then setting $\ell = \sum_{i < n} \ell_i$, we have that $\mathbf{A} \in \mathcal{K}$ has big Ramsey degree at most ℓ . If \mathcal{K} has strong amalgamation, then $\mathbf{A} \in \mathcal{K}$ has big Ramsey degree ℓ .*

Proof. We can write

$$\text{Emb}(\mathbf{A}, \mathbf{K}) = \bigsqcup_{i < n} \text{OEmb}(\mathbf{A} \cdot \sigma_i, \mathbf{K}).$$

Fix $r < \omega$ and an r -coloring $\chi: \text{Emb}(\mathbf{A}, \mathbf{K}) \rightarrow r$. Inductively on $i < n$, we can find $\eta_i \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ so that $\text{OEmb}(\mathbf{A} \cdot \sigma_i, \mathbf{K})$ sees at most ℓ_i colors for the coloring $\chi \cdot (\eta_0 \circ \dots \circ \eta_{i-1})$. Setting $\eta = \eta_0 \circ \dots \circ \eta_{n-1}$, then $|\chi \cdot \eta[\text{Emb}(\mathbf{A}, \mathbf{K})]| \leq \ell$. Hence \mathbf{A} has Ramsey degree at most ℓ in \mathbf{K} .

Now assume that \mathcal{K} has strong amalgamation. To show that \mathbf{A} has Ramsey degree exactly ℓ in \mathbf{K} , fix for each $i < n$ an *unavoidable coloring* $\chi_i: \text{OEmb}(\mathbf{A} \cdot \sigma_i, \mathbf{K})$, i.e., a coloring which witnesses that the ordered Ramsey degree of $\mathbf{A} \cdot \sigma_i$ is not less than ℓ_i . This gives rise to a coloring $\chi: \text{Emb}(\mathbf{A}, \mathbf{K}) \rightarrow \ell$. Now fix $\eta \in \text{Emb}(\mathbf{K}, \mathbf{K})$; we need to show that $|\chi \cdot \eta[\text{Emb}(\mathbf{A}, \mathbf{K})]| = \ell$. Because \mathcal{K} is a strong amalgamation class, we can find $\theta \in \text{Emb}(\mathbf{K}, \mathbf{K})$ so that $\eta \circ \theta \in \text{OEmb}(\mathbf{K}, \mathbf{K})$. It follows for each $i < n$ that $|\chi_i \cdot (\eta \circ \theta)[\text{OEmb}(\mathbf{A} \cdot \sigma_i, \mathbf{K})]| = \ell_i$. So we must have $|\chi \cdot \eta[\text{Emb}(\mathbf{A}, \mathbf{K})]| \geq |\chi \cdot (\eta \circ \theta)[\text{Emb}(\mathbf{A}, \mathbf{K})]| = \ell$. \square

3 \mathcal{L} -structures and coding trees

Throughout this article, we fix a finite binary relational language $\mathcal{L} = \{U_i : i < k^u\} \cup \{R_i : i < k\}$, where each U_i is unary and each R_i is binary. We put $\mathcal{L} = \mathcal{L}^u \cup \mathcal{L}^b$ for the unary and binary symbols, respectively. By adding extra relations if needed, we may assume that the following hold for every \mathcal{L} -structure \mathbf{A} we discuss.

1. For every $a \in \mathbf{A}$, there is a unique $i < k^u$ so that $U_i^{\mathbf{A}}(a)$ holds.
2. For any $a \in \mathbf{A}$ and any $i < k$, we never have $R_i^{\mathbf{A}}(a, a)$.
3. For any $a \neq b \in \mathbf{A}$, there is a unique $i < k$ so that $R_i^{\mathbf{A}}(a, b)$ holds.
4. There is an order 2 bijection $\text{Flip}: k \rightarrow k$ so that $R_i^{\mathbf{A}}(a, b)$ holds iff $R_{\text{Flip}(i)}^{\mathbf{A}}(b, a)$ holds. We demand that $\text{Flip}(0) = 0$.

By the above, we can treat $U^{\mathbf{A}}$ and $R^{\mathbf{A}}$ as functions, writing $U^{\mathbf{A}}(a) = i$ if $U_i^{\mathbf{A}}(a)$ holds, and similarly for $R^{\mathbf{A}}(a, b)$.

The relational symbol R_0 will play the role of “no relation,” and this is the sense in which we interpret notions such as “irreducible,” “free amalgam,” etc.

We now consider the k -branching tree with k^u -many roots $T := k^u \times k^{<\omega}$. If $k^u = 1$, we will often just think of T as the single-rooted tree $k^{<\omega}$. Given $t \in T$, we write $t = (t^u, t^b)$ for the *unary* and *binary* parts of t , and we write $\ell(t) = \text{dom}(t^b)$. If $i < k$, we write $t \frown i = (t^u, (t^b) \frown i)$. Often, in situations where t^u is understood and $m \in \text{dom}(t^b)$, we will simply write $t(m)$ instead of $t^b(m)$. If $i < k^u$, write $T_i := \{i\} \times k^{<\omega} = \{t \in T : t^u = i\}$, and we let $\emptyset_i \in T_i$ denote the root of the tree T_i . We write $T(n) := \{s \in T : \text{dom}(s^b) = n\}$ and $T_i(n) = T(n) \cap T_i$. We write $T(<n) := \bigcup_{m < n} T(m)$, and similarly for $T(>n)$, etc.

For now, let s, t , etc. range over elements of T . We write $\ell(s) := \text{dom}(s^b)$. We say that s is an *initial segment* of t and write $s \sqsubseteq t$ if $\ell(s) \leq \ell(t)$, $s^u = t^u$, and $t^b|_{\ell(s)} = s^b$. We write $s \perp t$ if $s \not\sqsubseteq t$ and $t \not\sqsubseteq s$. If $s^u = t^u = i$, we let $s \wedge t \in T_i$ denote the longest common initial segment of s and t . We write $\text{Succ}(s) := \{t \in T : t \sqsupseteq s\}$. The set of *immediate successors* of s is the set $\text{IS}(s) := \text{Succ}(s) \cap T(\ell(s) + 1)$; if

$S \subseteq T$, we write $\text{IS}(S) = \bigcup_{s \in S} \text{IS}(s)$. If $n \geq \ell(s)$, then $\text{Left}(s, n) \in T(n)$ denotes the successor of s where given $m < n$, we have $\text{Left}(s, n)^b(m) = s^b(m)$ if $m < \ell(s)$ or 0 otherwise.

We will frequently use the lexicographic order \preceq_{lex} on level subsets of T , where given $s, t \in T(n)$, we say $s \preceq_{lex} t$ if $s^u < t^u$ or if $s^u = t^u$ and $s^b \preceq_{lex} t^b$ in the ordinary sense.

We will think of $T(n)$ as the set of possible \mathcal{L} -types over an enumerated \mathcal{L} -structure of size n . We formalize this as follows. Given a countable, enumerated \mathcal{L} -structure \mathbf{A} , we form the *coding function* $c^{\mathbf{A}}: \mathbf{A} \rightarrow T$. For each $n \in \mathbf{A}$, we will have $c^{\mathbf{A}}(n) \in T(n)$, $c^{\mathbf{A}}(n)^u = U^{\mathbf{A}}(n)$, and given $m < n \in \mathbf{A}$, we set $c^{\mathbf{A}}(n)^b(m) = R^{\mathbf{A}}(n, m)$. So $c^{\mathbf{A}}(n)$ codes the 1-type of the vertex $n \in \mathbf{A}$ over \mathbf{A}_n . We refer to the set $c^{\mathbf{A}}[\mathbf{A}] \subseteq T$ as the *coding nodes* of \mathbf{A} . The *coding tree* $\text{CT}^{\mathbf{A}}$ is the set $\{t \in T : \exists a \in \mathbf{A} \text{ with } c^{\mathbf{A}}(a) \sqsupseteq t\}$ equipped with the coding function $c^{\mathbf{A}}$. If $m < \omega$ and $i < k^u$, we put $\text{CT}^{\mathbf{A}}(m) = \text{CT}^{\mathbf{A}} \cap T(m)$, $\text{CT}_i^{\mathbf{A}} = \text{CT}^{\mathbf{A}} \cap T_i$, and $\text{CT}_i^{\mathbf{A}}(m) = \text{CT}^{\mathbf{A}} \cap T_i(m)$.

We now fix a finite set \mathcal{F} of finite, irreducible \mathcal{L} -structures, and **we set** $\mathcal{K} = \text{Forb}(\mathcal{F})$. We can assume that the structures in \mathcal{F} are pairwise non-embeddable into each other. This implies that if $\mathbf{F} \in \mathcal{F}$ and $\mathbf{J} \subseteq \mathbf{F}$ is a proper substructure of \mathbf{F} , then $\mathbf{J} \in \mathcal{K}$. We may also assume that for each $i < k^u$, the singleton structure with unary predicate i is in \mathcal{K} , i.e., that $\min\{|\mathbf{F}| : \mathbf{F} \in \mathcal{F}\} \geq 2$.

Given an enumerated Fraïssé limit $\mathbf{K} \cong \text{Flim}(\mathcal{K})$, we say that \mathbf{K} is *left dense* if given any enumerated structure $\mathbf{A} \in \mathcal{K}$ with $|\mathbf{A}| = m + 1$ and $\mathbf{A}_m = \mathbf{K}_m$, there is $f \in \text{OEmb}(\mathbf{A}, \mathbf{K})$ which is the identity on \mathbf{A}_m and so that for any r with $m \leq r < f(m)$, we have $R^{\mathbf{K}}(f(m), r) = 0$. Equivalently, upon forming the coding function $c^{\mathbf{K}}$, we have that \mathbf{K} is left dense iff whenever $s \in \text{CT}^{\mathbf{K}}$, then there is $n > \ell(s)$ such that $\text{Left}(s, n) = c^{\mathbf{K}}(n)$. Note that if \mathbf{K} is left dense, then given \mathbf{A} as above, then there are in fact infinitely many such $f \in \text{OEmb}(\mathbf{A}, \mathbf{K})$; in particular, we can choose $f(m)$ to be as large as we want. **We now fix once and for all** a left-dense Fraïssé limit \mathbf{K} . When referring to this fixed \mathbf{K} , we will drop various superscripts, writing c, U, R, CT , etc.

We call a substructure $\mathbf{A} \subseteq \mathbf{K}$ *left-leaning* if whenever $a \in \mathbf{A}$, $x < a$, and $R(a, x) \neq 0$, then $x \in \mathbf{A}$. An embedding $f: \mathbf{A} \rightarrow \mathbf{K}$ is called *left-leaning* if $f[\mathbf{A}] \subseteq \mathbf{K}$ is left-leaning. Given $n < \omega$, we say that $f: \mathbf{A} \rightarrow \mathbf{K}_n$ is left-leaning iff $f: \mathbf{A} \rightarrow \mathbf{K}$ is left-leaning and $f[\mathbf{A}] \subseteq \mathbf{K}_n$. Write $\text{LEmb}(\mathbf{A}, \mathbf{K})$ or $\text{LEmb}(\mathbf{A}, \mathbf{K}_n)$ for the set of left-leaning embeddings of \mathbf{A} into \mathbf{K} or \mathbf{K}_n , respectively. If \mathbf{A} is an enumerated \mathcal{L} -structure, write $\text{OL}(\mathbf{A}, \mathbf{K})$ or $\text{OL}(\mathbf{A}, \mathbf{K}_n)$ for the set of ordered embeddings which are also left-leaning.

Lemma 3.0.1. *Let \mathbf{A} be a countable, enumerated \mathcal{L} -structure with $\text{Age}(\mathbf{A}) \subseteq \mathcal{K}$. Suppose $m < \omega$ and $f|_m \in \text{OL}(\mathbf{A}_m, \mathbf{K})$. Then we can extend $f|_m$ to $f \in \text{OL}(\mathbf{A}, \mathbf{K})$ while demanding $f(m)$ is as large as we want.*

Proof. By induction, it is enough to handle the case where $|\mathbf{A}| = m + 1$. Write $f(m-1) = N-1$; if $m = 0$, set $N = 0$. Let \mathbf{B} denote the free amalgam of $\mathbf{A}_m \hookrightarrow \mathbf{A}$ and $\mathbf{A}_m \xrightarrow{f|_m} \mathbf{K}_N$; we fix an enumeration of \mathbf{B} with $\mathbf{B}_N = \mathbf{K}_N$ and with $N \in \mathbf{B}$ corresponding to the point $m \in \mathbf{A}$. Note that if $m = 0$, then $\mathbf{B} = \mathbf{A}$. Since \mathbf{K} is left dense, we can find $g \in \text{OEmb}(\mathbf{B}, \mathbf{K})$ with g the identity on \mathbf{B}_N , with $g(N)$ as large

as we want, and so that for any r with $N \leq r < g(N)$, we have $R(g(N), r) = 0$. Set $f(m) = g(N)$; it follows that $f: \mathbf{A} \rightarrow \mathbf{K}$ is left-leaning. \square

4 Posets of classes of structures

4.1 Gluings

Definition 4.1.1. Let $\{V_n : n < \omega\}$ be new unary symbols not in \mathcal{L}^u . We set $\mathcal{L}_d = \mathcal{L}^b \cup \{V_0, \dots, V_{d-1}\}$. Similar to our conventions for \mathcal{L} -structures, when discussing \mathbf{B} an \mathcal{L}_d -structure on underlying set B , we demand that for every $x \in B$, there is a unique $i < d$ where $V_i^{\mathbf{B}}(x)$ holds; so this provides a partition of B into d -many (possibly empty) labelled pieces. We can then regard $V^{\mathbf{B}}: B \rightarrow d$ as a function with $V^{\mathbf{B}}(x) = i$ iff $V_i^{\mathbf{B}}(x)$ holds. Our conventions for the \mathcal{L}^b -part remain exactly the same. One mild difference is that we will almost never consider enumerated \mathcal{L}_d -structures; **in fact, unless stated otherwise, we will assume that a given \mathcal{L}_d -structure has underlying set disjoint from ω** . This is because we will be “gluing” \mathcal{L}_d -structures to enumerated \mathcal{L} -structures to form new \mathcal{L} -structures.

Definition 4.1.2. A *gluing* γ is a triple $\gamma = (\rho, \eta, \mathbf{I})$ where:

- $\rho: d \rightarrow k^u$ is a map for some $d < \omega$. We write $\rho = \text{Sort}(\gamma)$ and call ρ the *sort*, and the integer d is its *rank*.
- $\eta: d \times n \rightarrow k$ is a map for some $n < \omega$ (and d is the rank from above). We call η the *attachment map*, and the integer n is its *size*.
- $\mathbf{I} \in \mathcal{K}$ is an enumerated \mathcal{L} -structure with $|\mathbf{I}| = n$ (where n is the size from above).

When $k^u = 1$, then we omit ρ from the notation, and if $\mathbf{I} = \mathbf{K}_n$, then we omit \mathbf{I} from the notation. If both $k^u = 1$ and $\mathbf{I} = \mathbf{K}_n$, then we often just write η for the gluing.

Notice that the rank d is implicit as long as the sort is specified. We let $\text{Glue}(\rho, \mathbf{I}) = \{(\rho, \eta, \mathbf{I}) : (\rho, \eta, \mathbf{I}) \text{ is a gluing}\}$, and we let $\text{Glue}(d, \mathbf{I}) = \bigcup \{\text{Glue}(\rho, \mathbf{I}) : \text{dom}(\rho) = d\}$.

Definition 4.1.3. Let $\gamma = (\rho, \eta, \mathbf{I})$ be a gluing with rank d and size n . Suppose \mathbf{B} is an \mathcal{L}_d -structure with underlying set B . Then $\mathbf{B}[\gamma]$ denotes the \mathcal{L} -structure with underlying set $n \cup B$ with \mathbf{I} induced on n , the \mathcal{L}^b -part of \mathbf{B} induced on B , $U^{\mathbf{B}[\gamma]}(b) = \rho \circ V^{\mathbf{B}}(b)$ for each $b \in \mathbf{B}$, and $R^{\mathbf{B}[\gamma]}(b, x) = \eta(V^{\mathbf{B}}(b), x)$ whenever $b \in \mathbf{B}$ and $x \in \mathbf{I}$. Intuitively, we are gluing the structure \mathbf{B} to the structure \mathbf{I} using the instructions given by γ to determine the relations between vertices of \mathbf{B} and vertices of \mathbf{I} (see Figure 1).

In the case that $n = 0$ (i.e., \mathbf{I} is the empty structure and η is the empty map), we can identify the sort ρ with the corresponding gluing. So if $\mathbf{B} \in \mathcal{L}_d$, then $\mathbf{B}[\rho]$ is simply the \mathcal{L} -structure on underlying set B with the \mathcal{L}^b -part of \mathbf{B} induced on B and with $U^{\mathbf{B}[\rho]}(b) = \rho \circ V^{\mathbf{B}}(b)$.




Figure 1: Gluing $\mathbf{B}[\gamma]$.

Notation 4.1.4. A particular instance of Definition 4.1.2 that we will commonly use is as follows. Given $n < \omega$ and a function $\sigma: d \rightarrow T(n)$, we obtain maps $\sigma^u: d \rightarrow k^u$ and $\sigma^b: d \times n \rightarrow k$, where given $i < d$ and $m < n$, we have:

$$\begin{aligned}\sigma^u(i) &= \sigma(i)^u \\ \sigma^b(i, m) &= \sigma(i)^b(m)\end{aligned}$$

If $S = \{s_0 \preceq_{lex} \cdots \preceq_{lex} s_{d-1}\} \subseteq T(n)$, we define $\vec{S}: d \rightarrow T(n)$ by setting $\vec{S}(i) = s_i$ for each $i < d$. In this situation, we typically write S^u and S^b in place of $(\vec{S})^u$ and $(\vec{S})^b$. So in particular, $S^u(i) = j$ iff $s_i \in T_j$.

If \mathbf{A} is an enumerated \mathcal{L} -structure with $|\mathbf{A}| \geq n$, then $(\sigma^u, \sigma^b, \mathbf{A}_n)$ and (S^u, S^b, \mathbf{A}_n) are gluings. If \mathbf{B} is an \mathcal{L}_d -structure, we write $\mathbf{B}[\sigma, \mathbf{A}]$ in place of $\mathbf{B}[(\sigma^u, \sigma^b, \mathbf{A}_n)]$, where n is such that $T(n) = \text{ran}(\sigma)$, and similarly for $\mathbf{B}[S, \mathbf{A}]$. When $\mathbf{A} = \mathbf{K}_n$, we omit it from the notation. If $S = \{t\}$, we simply write $\mathbf{B}[t, \mathbf{A}]$, or $\mathbf{B}[t]$ if $\mathbf{A} = \mathbf{K}_n$.

4.2 The poset $P(\rho)$

Definition 4.2.1.

1. If $\rho: d \rightarrow k^u$ is a sort, we define the class of finite \mathcal{L}_d -structures

$$\mathcal{K}(\rho) = \{\mathbf{B} : \mathbf{B}[\rho] \in \mathcal{K}\}.$$

2. If γ is a rank d gluing, we define the *age* of γ to be the class of finite \mathcal{L}_d -structures

$$\mathcal{K}(\gamma) := \{\mathbf{B} : \mathbf{B}[\gamma] \in \mathcal{K}\}.$$

3. If $\sigma: d \rightarrow T(n)$ is a function, we similarly define the *age of σ* to be the class of finite \mathcal{L}_d -structures

$$\mathcal{K}(\sigma) := \{\mathbf{B} : \mathbf{B}[\sigma] \in \mathcal{K}\}.$$

If $S \subseteq T(n)$, we set $\mathcal{K}(S) = \mathcal{K}(\vec{S})$. If $\mathbf{A} \in \mathcal{K}$ is an enumerated structure with $|\mathbf{A}| \geq n$, we define the *age of σ relative to \mathbf{A}* to be the class of \mathcal{L}_d -structures

$$\mathcal{K}(\sigma, \mathbf{A}) = \{\mathbf{B} : \mathbf{B}[\sigma, \mathbf{A}] \in \mathcal{K}\},$$

and likewise for $\mathcal{K}(S, \mathbf{A})$.

Remark. Classes of the form $\mathcal{K}(S)$ for $S = \{s_0 \preceq_{lex} \cdots \preceq_{lex} s_{d-1}\} \subseteq T(n)$ have a somewhat more intuitive definition; recalling that $c: \mathbf{K} \rightarrow T$ denotes the coding function of our fixed enumerated Fraïssé limit \mathbf{K} , we have $\mathbf{B} \in \mathcal{K}(S)$ iff there is an $\mathcal{L}_{\mathbf{b}}$ -embedding $f: \mathbf{B} \rightarrow \mathbf{K}$ so that $c \circ f(b) \sqsupseteq_{s_{V_{\mathbf{B}}(b)}}$ for each $b \in \mathbf{B}$, i.e. so that the coding nodes corresponding to $\text{Im}(f)$ lie above the nodes of S as described by the unaries in \mathbf{B} . In particular, we have that $t \in \text{CT}$ iff $\mathcal{K}(t) \neq \emptyset$.

Definition 4.2.2. If $\rho: d \rightarrow k^u$ is a sort, we define the *poset of ages of ρ* via

$$P(\rho) := \{\mathcal{K}(S) : S \subseteq \text{CT}(n) \text{ for some } n \text{ and } S^u = \rho\}.$$

So each member of $P(\rho)$ is a class of \mathcal{L}_d -structures. We view $P(\rho)$ as a poset under inclusion. When $k^u = 1$, we simply write $P(d)$ for this poset.

In the $k^u = 1$ case, the poset $P(1)$ has been used by Sauer [27] to characterize when the Fraïssé limits of binary free amalgamation classes are *indivisible*, i.e. when the singleton structure has big Ramsey degree 1.

It turns out that the poset $P(\rho)$ is independent of the choice of left-dense Fraïssé limit \mathbf{K} that we chose; we will prove this later in Proposition 5.1.6. For now, we describe exactly which sorts ρ have $P(\rho) \neq \emptyset$ (Proposition 4.2.4), and we prove a strong closure property of $P(\rho)$, which implies that it is closed under intersections (Proposition 4.2.6).

Before stating Proposition 4.2.4, let us discuss a simplifying assumption that we will work with.

Definition 4.2.3. Fix $i < k^u$. We call i *non-degenerate* for \mathcal{K} if there is some structure of size 2 in \mathcal{K} where at least one of the vertices has unary predicate i and the two vertices are related by a non-zero binary relation. Otherwise, we call i *degenerate*.

Degenerate unaries will behave rather differently than non-degenerate ones. If $i < k^u$ is degenerate, then every single coding node in T_i is on the left branch. This would become annoying to deal with when we define diagonal substructures, where for non-degenerate unaries, we will ask that coding nodes form an antichain. Luckily, degenerate unaries will be easy to deal with at the very end. So **until Section 7, we will assume** that every $i < k^u$ is non-degenerate for \mathcal{K} . In section 7, we will indicate how to obtain exact big Ramsey degrees for classes which contain degenerate unary predicates.

Proposition 4.2.4. *Given $\rho: d \rightarrow k^u$, then $P(\rho) \neq \emptyset$ iff ρ is non-decreasing.*

Proof. From Definition 4.2.2, we must have that ρ is non-decreasing if $P(\rho) \neq \emptyset$. For the other direction, fix some $N < \omega$. It is enough to find some $n < \omega$ so that $|\text{CT}_i(n)| \geq N$ for each $i < k^u$. Because every $i < k^u$ is non-degenerate, we can find $j_i < k^u$ and a non-zero $q_i < k$ so that the two element enumerated structure \mathbf{I} with $U^{\mathbf{I}}(0) = j_i$, $U^{\mathbf{I}}(1) = i$, and $R^{\mathbf{I}}(1, 0) = q_i$ is in \mathcal{K} . Fix such choices j_i and q_i for now.

Let ℓ be chosen so that $N \leq 2^\ell$ for each $i < k^u$. Let \mathbf{A} be the enumerated structure of size $\ell \cdot k^u$ containing no non-trivial binary relations and so that for each

$i < k^u$ and $j < \ell$, we have $U^{\mathbf{A}}(i \cdot \ell + j) = j_i$. Fix $f \in \text{OL}(\mathbf{A}, \mathbf{K}_n)$ for some $n < \omega$. Now for each $i < k^u$ and $\sigma: \ell \rightarrow 2$, define a node $s_{i,\sigma} \in T(n)$ by setting $(s_{i,\sigma})^u = i$, and given $m < n$, we have

$$s_{i,\sigma}(m) = \begin{cases} q_i \cdot \sigma(j) & \text{if } m = f(i \cdot \ell + j) \\ 0 & \text{else.} \end{cases}$$

By construction $s_{i,\sigma} \in \text{CT}_i(n)$, so $|\text{CT}_i(m)| \geq 2^\ell \geq N$ as desired. \square

Remark. This proposition is false without the assumption that every unary is non-degenerate. In fact, if $i < k^u$ is degenerate and ρ is a sort with $P(\rho) \neq \emptyset$, then $|\rho^{-1}(\{i\})| \leq 1$.

From now on, whenever we mention a sort, we will always assume it is non-decreasing.

Definition 4.2.5. Suppose \mathcal{A} is a hereditary class of finite \mathcal{L}_d -structures. We call the class \mathcal{A} *reasonable* if \mathcal{A} contains every singleton \mathcal{L}_d -structure (one for each $i < d$). If $\rho: d \rightarrow k^u$ is a sort and $\gamma = (\rho, \eta, \mathbf{I})$ is a gluing, we call γ *reasonable* if the class $\mathcal{K}(\gamma)$ is reasonable.

Remark. If $S \subseteq \text{CT}(n)$, then $\mathcal{K}(S)$ is a reasonable class.

Proposition 4.2.6. Fix a sort $\rho: d \rightarrow k^u$. Let $\mathcal{A} \in P(\rho)$, and let $\gamma = (\rho, \eta, \mathbf{J})$ be a reasonable gluing. Then $\mathcal{A} \cap \mathcal{K}(\gamma) \in P(\rho)$.

Proof. Suppose $S = \{s_0 \preceq \dots \preceq s_{d-1}\} \subseteq \text{CT}(n)$ is such that $\mathcal{A} = \mathcal{K}(S)$. Fix some $h \in \text{OL}(\mathbf{J}, \mathbf{K})$ with $h(0) > n$, and choose some $N > \max\{h[\mathbf{J}]\}$. We now define for each $i < d$ a node $t_i \in T(N)$ with $t_i \supseteq s_i$:

$$t_i(m) = \begin{cases} s_i(m) & \text{if } m < n, \\ \eta(i, x) & \text{if } m = h(x), \\ 0 & \text{else.} \end{cases}$$

Set $X = \{t_0, \dots, t_{d-1}\}$. We will show that $\mathcal{A} \cap \mathcal{K}(\gamma) = \mathcal{K}(X)$. Note first that $\mathcal{K}(X) \subseteq \mathcal{K}(S) = \mathcal{A}$. Now fix $\mathbf{B} \in \mathcal{A}$.

Claim: The map $h_{\mathbf{B}}: \mathbf{B}[\gamma] \rightarrow \mathbf{B}[X]$ which is the identity on \mathbf{B} and with $h_{\mathbf{B}}(x) = h(x)$ for each $x \in \mathbf{J}$ is an embedding.

Proof of claim. Since h is an embedding and since γ and X have the same sort, we see that $h_{\mathbf{B}}$ respects unary relations and binary relations for those pairs (x, z) with both $x, z \in \mathbf{J}$ or $x, z \in \mathbf{B}$. Now suppose $x \in \mathbf{J}$ and $z \in \mathbf{B}$. We have:

$$\begin{aligned} R^{\mathbf{B}[X]}(h_{\mathbf{B}}(z), h_{\mathbf{B}}(x)) &= R^{\mathbf{B}[X]}(z, h(x)) \\ &= t_{V^{\mathbf{B}}(z)}(h(x)) \\ &= \eta(V^{\mathbf{B}}(z), x) \\ &= R^{\mathbf{B}[\gamma]}(z, x) \end{aligned} \quad \blacksquare$$

This shows that $\mathcal{K}(X) \subseteq \mathcal{K}(\gamma)$. In the other direction, suppose $\mathbf{B} \in \mathcal{A} \setminus \mathcal{K}(X)$, i.e., that $\mathbf{B}[X] \notin \mathcal{K}$. Since $\mathcal{K} = \text{Forb}(\mathcal{F})$, find $\mathbf{F} \in \mathcal{F}$ and an embedding $g: \mathbf{F} \rightarrow \mathbf{B}[X]$, and write $\mathbf{E} = g^{-1}(\mathbf{K}_N)$. Then there is a unique $\mathbf{I} \in \mathcal{I}$ and $f \in \text{OEmb}(\mathbf{I}, \mathbf{K}_N)$ with $f[\mathbf{I}] = g[\mathbf{E}]$. Since \mathbf{F} is irreducible, and by the construction of the t_i , we must have $f[\mathbf{I}] \subseteq \mathbf{K}_n \cup h[\mathbf{J}]$. Then, since h is left-leaning with $h(0) > n$, there are no non-trivial relations between \mathbf{K}_n and $h[\mathbf{J}]$. Since \mathbf{I} is irreducible, we must have either $f[\mathbf{I}] \subseteq \mathbf{K}_n$ or $f[\mathbf{I}] \subseteq h[\mathbf{J}]$. Since $\mathbf{B} \in \mathcal{A}$, the first case cannot hold, as $\mathbf{B}[S] \in \mathcal{K}$ is an induced substructure of $\mathbf{B}[X]$. So $f[\mathbf{I}] \subseteq h[\mathbf{J}]$, implying that $g[\mathbf{F}] \subseteq \text{Im}(h_{\mathbf{B}})$. So $\mathbf{B}[\gamma] \notin \mathcal{K}$, i.e., $\mathbf{B} \notin \mathcal{K}(\gamma)$.

To see that in fact $X \subseteq \text{CT}$, we note that $\mathcal{A} \cap \mathcal{K}(\gamma)$ is reasonable. So given $i < d$, the singleton \mathcal{L}_d -structure of unary i is in $\mathcal{K}(X)$, which means that there is $M < \omega$ with $c(M) \supseteq t_i$, i.e., that $t_i \in \text{CT}$. \square

Corollary 4.2.7. *$P(\rho)$ is closed under intersections.*

We end this subsection with some examples, describing the posets $P(\rho)$ in a few concrete cases.

Example 4.2.8. Let $\mathcal{K} = \mathcal{G}_3$ be the class of triangle-free graphs. Then $k^u = 1$ and $k = 2$. We think of \mathcal{L}_d -structures as graphs equipped with a labeled partition into d (possibly empty) pieces; in a mild abuse of terminology, we will refer to these pieces as V_0, \dots, V_{d-1} . Let $\mathbf{I} = \mathbf{K}_1$ be the graph on a single vertex, and let $\eta: 2 \times 1 \rightarrow 2$ be given by $\eta(0, 0) = \eta(1, 0) = 0$. Then $\mathcal{K}(\eta)$ is the class of all triangle-free graphs equipped with an arbitrary partition into pieces V_0 and V_1 . However, we have $\mathcal{K}(\eta) \notin P(2)$. This is because if $t \in \text{CT}$ satisfies that $\mathcal{K}(t)$ is the class of all triangle-free graphs with all points in V_0 , then we must have $t = 0^n$ for some $n < \omega$. In particular, if $S \subseteq \text{CT}(n)$ has $|S| = 2$, then for at least one $s \in S$, we have that $\mathcal{K}(s)$ is the class of edge-free graphs with all points in V_0 . In Subsection 4.6, we will see that this is all a consequence of the fact that \mathcal{G}_3 does not have a *free relation*.

For the class of triangle-free graphs, let us describe the posets $P(d)$ completely. The poset $P(1)$ contains two members – the class \mathcal{G}_3 and the class \mathcal{G}_2 of finite graphs with no edges. The poset $P(2)$ contains 3 members – the class of finite triangle-free graphs where V_1 is an anti-clique; the class of finite, bipartite graphs where V_0, V_1 is a bipartition, and the class of finite graphs with no edges with the vertex set split into V_0 and V_1 arbitrarily. For larger d , we will see that since we are omitting a structure of size 3, the posets $P(d)$ are entirely determined by what can happen on any pair V_i, V_j . If V is a set, write $\mathcal{G}(V)$ for the set of graphs on vertex set V . We can identify $P(d)$ with the set $\mathcal{G}(\{0, \dots, d-1\}) \cup \mathcal{G}(\{1, \dots, d-1\})$. Given a graph $H \in \mathcal{G}(\{0, \dots, d-1\})$, let \mathcal{A}_H denote the class of all finite, triangle-free graphs where each of the pieces V_0, \dots, V_{d-1} is an anti-clique, and edges are allowed between vertices in V_i and vertices in V_j iff $\{i, j\}$ is an edge of H . And if $H \in \mathcal{G}(\{1, \dots, d-1\})$, then the class \mathcal{A}_H contains all triangle-free graphs where each of the pieces V_1, \dots, V_{d-1} is an anti-clique and where for $1 \leq i < j \leq d-1$, edges are allowed between vertices in V_i and vertices in V_j iff $\{i, j\}$ is an edge of H . There are no restrictions (aside from triangle-free) on edges in V_0 nor on edges between V_0 and V_i for $1 \leq i \leq d-1$.




Figure 2: Structure \mathbf{B}^e .

Example 4.2.9. Let \mathcal{K} denote the class of directed graphs with no cyclic triangles. Here $k^u = 1$ and $k = 3$. The bijection $\text{Flip}: 3 \rightarrow 3$ is non-trivial, with $\text{Flip}(1) = 2$ and $\text{Flip}(2) = 1$. The cyclic triangle \mathbf{T} is the directed graph on three points $\{x, y, z\}$ with $R^{\mathbf{T}}(x, y) = R^{\mathbf{T}}(y, z) = R^{\mathbf{T}}(z, x) = 1$. The poset $P(1)$ contains only \mathcal{K} , while the poset $P(2)$ contains 4 members – the class of finite \mathbf{T} -free directed graphs with V_0, V_1 arbitrary; the class of finite \mathbf{T} -free directed graphs with no edges from V_0 to V_1 ; the class of finite \mathbf{T} -free directed graphs with no edges from V_1 to V_0 , and the class of finite \mathbf{T} -free directed graphs with no edges at all between V_0 and V_1 . Like in Example 4.2.8, because the class \mathcal{K} is described by forbidding substructures of size at most 3, we have that for $d \geq 3$, $P(d)$ is determined by what is allowed on any pair V_i, V_j . In this case, one can show that for any function $\sigma: \binom{d}{2} \rightarrow P(2)$, there is a class in $P(d)$ such that for $i < j < d$, the edges allowed between vertices in V_i and vertices in V_j are exactly described by the class $\sigma(i, j)$.

4.3 Operations on structures and classes

If $S \subseteq \text{CT}(n)$ and $S_0 \subseteq S$, we will want to describe the class $\mathcal{K}(S_0)$ using the class $\mathcal{K}(S)$. Indeed, this idea is implicit in Examples 4.2.8 and 4.2.9, when we described a member of $P(d)$ based on its “restriction” to V_i and V_j for some $i < j < d$. We now formalize this idea.

Definition 4.3.1.

1. Suppose \mathbf{B} is an \mathcal{L}_d -structure on underlying set B . If $d_0 \leq d$ and $e: d_0 \rightarrow d$ is an injection, we define the \mathcal{L}_{d_0} -structure \mathbf{B}^e , where the underlying set of \mathbf{B}^e is $\underline{B}^e := (V^{\mathbf{B}})^{-1}(\text{Im}(e)) \subseteq B$, the \mathcal{L}^b -part is induced from \mathbf{B} , and given $b \in \underline{B}^e$, we have $V^{\mathbf{B}^e}(b) = e^{-1}(V^{\mathbf{B}}(b))$ (see Figure 2).
2. Suppose $e: d_0 \rightarrow d$ is any function (not necessarily an injection) and \mathbf{B} is an \mathcal{L}_{d_0} -structure with underlying set B . Then we define $e \cdot \mathbf{B}$ to be the \mathcal{L}_d -structure on underlying set B with the \mathcal{L}^b -part induced from \mathbf{B} , and given $b \in B$, we have $V^{e \cdot \mathbf{B}}(b) = e \circ V^{\mathbf{B}}(b)$.
3. Let $e: d_0 \rightarrow d$ be any function. If \mathcal{A} is a hereditary class of finite \mathcal{L}_d -structures, we define the class of \mathcal{L}_{d_0} -structures $\mathcal{A}^e = \{\mathbf{B} : e \cdot \mathbf{B} \in \mathcal{A}\}$. If \mathcal{B} is a hereditary class of \mathcal{L}_{d_0} -structures, we define the class of \mathcal{L}_d -structures $e \cdot \mathcal{B} = \{e \cdot \mathbf{B} : \mathbf{B} \in \mathcal{B}\}$.

4. If \mathcal{A} is a hereditary class of \mathcal{L}_d -structures, \mathcal{B} is a hereditary class of \mathcal{L}_{d_0} -structures, and $e: d_0 \rightarrow d$ is an injection, we define $\mathcal{A}|_{(e,\mathcal{B})} = \{\mathbf{B} \in \mathcal{A} : \mathbf{B}^e \in \mathcal{B}\}$. One can think of this class as “ \mathcal{A} restricted at e to \mathcal{B} .”

Remark. If \mathcal{A} is a hereditary class of \mathcal{L}_d -structures and $e: d_0 \rightarrow d$ is an injection, we equivalently have $\mathcal{A}^e = \{\mathbf{B}^e : \mathbf{B} \in \mathcal{A}\}$; this holds because for any \mathcal{L}_d -structure \mathbf{B} , we have that $e \cdot (\mathbf{B}^e)$ is an induced substructure of \mathbf{B} .

Definition 4.3.2.

1. Let $\gamma = (\rho, \eta, \mathbf{I})$ be a gluing with rank d and size n . If $e: d_0 \rightarrow d$ is any function, then we write γ^e for the gluing $(\rho \circ e, \eta^e, \mathbf{I})$, where $\eta^e: d_0 \times n \rightarrow k$ is the map where given $i < d_0$ and $m < n$, we have $\eta^e(i, m) = \eta(e(i), m)$.
2. Suppose $e: d_0 \rightarrow d$ is an injection, and let $\zeta = (\sigma, \eta, \mathbf{I})$ be a gluing of rank d_0 and size n . Suppose $\rho: d \rightarrow k^u$ is a sort such that $\sigma = \rho \circ e$. Then we define the gluing $(\rho, e) \cdot \zeta = (\rho, e \cdot \eta, \mathbf{I})$, where $e \cdot \eta: d \times n \rightarrow k$ is the map where given $i < d$ and $m < n$, we have

$$(e \cdot \eta)(i, m) = \begin{cases} \eta(e^{-1}(i), m) & \text{if } i \in \text{Im}(e) \\ 0 & \text{else.} \end{cases}$$

Remark.

1. If $e: d_0 \rightarrow d$ is any function, \mathbf{B} is an \mathcal{L}_{d_0} -structure, and γ is a rank d gluing, then $(e \cdot \mathbf{B})[\gamma] = \mathbf{B}[\gamma^e]$. It follows that $(\mathcal{K}(\gamma))^e = \mathcal{K}(\gamma^e)$. If γ is reasonable (Definition 4.2.5), then so is γ^e .
2. If $e: d_0 \rightarrow d$ is an injection, \mathbf{B} is an \mathcal{L}_d -structure, $\zeta = (\sigma, \eta, \mathbf{I})$ is a rank d_0 -gluing, and $\rho: d \rightarrow k^u$ is a sort with $\sigma = \rho \circ e$, then $\mathbf{B}[(\rho, e) \cdot \zeta]$ is the free amalgam of $(\mathbf{B}^e)[\zeta]$ and $\mathbf{B}[\rho]$ along $(\mathbf{B}^e)[\sigma]$. It follows that $\mathcal{K}(\rho)|_{(e,\mathcal{K}(\zeta))} = \mathcal{K}((\rho, e) \cdot \zeta)$. The gluing $(\rho, e) \cdot \zeta$ is reasonable iff ζ is reasonable.

Proposition 4.3.3. *If $\rho: d \rightarrow k^u$ is a sort, $e: d_0 \rightarrow d$ is an increasing injection, $\mathcal{A} \in P(\rho)$, and $\mathcal{B} \in P(\rho \circ e)$, then we have $\mathcal{A}|_{(e,\mathcal{B})} \in P(\rho)$.*

Proof. First write $\mathcal{A}|_{(e,\mathcal{B})} = \mathcal{A} \cap \mathcal{K}(\rho)|_{(e,\mathcal{B})}$. Now let ζ be a gluing such that $\mathcal{B} = \mathcal{K}(\zeta)$. Then we have $\mathcal{K}(\rho)|_{(e,\mathcal{B})} = \mathcal{K}(\rho)|_{(e,\mathcal{K}(\zeta))} = \mathcal{K}((\rho, e) \cdot \zeta)$. Since $(\rho, e) \cdot \zeta$ is a reasonable gluing, we have $\mathcal{A}|_{(e,\mathcal{B})} \in P(\rho)$ by Proposition 4.2.6. \square

We end the subsection with a more complicated example along the lines of Examples 4.2.8 and 4.2.9.

Notation 4.3.4.

1. If $n < N \leq \omega$, we write $\iota_n: 1 \rightarrow N$ for the function given by $\iota_n(0) = n$. The range N is not explicit in the notation, but common choices of N are ω and k^u , and this is typically understood by context.
2. If $i < d$, then $\epsilon_{i,d}: (d-1) \rightarrow d$ is the increasing injection whose image skips i .

Example 4.3.5. Let $\mathcal{K} = \mathcal{G}_4$ be the class of K_4 -free graphs. Then $k^u = 1$ and $k = 2$. We describe the posets $P(1)$, $P(2)$, and $P(3)$, using Definition 4.3.1 to help with the description. The poset $P(1)$ contains 3 classes; the class \mathcal{G}_4 , the class \mathcal{G}_3 of triangle-free graphs, and the class \mathcal{G}_2 of finite graphs with no edges.

To describe the 10 members of $P(2)$, we categorize $\mathcal{A} \in P(2)$ based on the classes \mathcal{A}^{ι_0} and \mathcal{A}^{ι_1} (Notation 4.3.4). For instance, saying that $\mathcal{A}^{\iota_1} = \mathcal{G}_3$ means that in any $\mathbf{A} \in \mathcal{A}$, the induced subgraph on $V_1^{\mathbf{A}}$ must be triangle-free. For similar reasons as in Example 4.2.8, we cannot have $\mathcal{A}^{\iota_1} = \mathcal{G}_4$. The class $P(2)$ has the following 10 members:

- $\mathcal{A} \in P(2)$ with $(\mathcal{A}^{\iota_0}, \mathcal{A}^{\iota_1}) = (\mathcal{G}_4, \mathcal{G}_3)$:
 - $\mathcal{A} = \mathcal{G}_4^{4,3}$: Finite K_4 -free graphs where V_1 is triangle free,
- $\mathcal{A} \in P(2)$ with $(\mathcal{A}^{\iota_0}, \mathcal{A}^{\iota_1}) = (\mathcal{G}_4, \mathcal{G}_2)$:
 - $\mathcal{A} = \mathcal{G}_4^{4,2}$: Finite K_4 -free graphs where V_1 is an anti-clique,
- $\mathcal{A} \in P(2)$ with $(\mathcal{A}^{\iota_0}, \mathcal{A}^{\iota_1}) = (\mathcal{G}_3, \mathcal{G}_3)$:
 - $\mathcal{A} = \mathcal{G}_4^{3,3}$: Finite K_4 -free graphs where both V_0 and V_1 are triangle-free,
 - $\mathcal{A} = \mathcal{G}_3^{3,3}$: Finite triangle-free graphs with V_0, V_1 an arbitrary partition,
- $\mathcal{A} \in P(2)$ with $(\mathcal{A}^{\iota_0}, \mathcal{A}^{\iota_1}) = (\mathcal{G}_3, \mathcal{G}_2)$:
 - $\mathcal{A} = \mathcal{G}_4^{3,2}$: Finite K_4 -free graphs with V_0 triangle-free and V_1 an anticlique,
 - $\mathcal{A} = \mathcal{G}_3^{3,2}$: Finite triangle-free graphs with V_1 is an anticlique,
- $\mathcal{A} \in P(2)$ with $(\mathcal{A}^{\iota_0}, \mathcal{A}^{\iota_1}) = (\mathcal{G}_2, \mathcal{G}_3)$:
 - $\mathcal{A} = \mathcal{G}_4^{2,3}$: Finite K_4 -free graphs where V_0 is an anti-clique and V_1 is triangle-free,
 - $\mathcal{A} = \mathcal{G}_3^{2,3}$: Finite triangle-free graphs where V_0 is an anticlique,
- $\mathcal{A} \in P(2)$ with $(\mathcal{A}^{\iota_0}, \mathcal{A}^{\iota_1}) = (\mathcal{G}_2, \mathcal{G}_2)$:
 - $\mathcal{A} = \mathcal{G}_3^{2,2}$: Finite bipartite graphs where V_0, V_1 is a bipartition,
 - $\mathcal{A} = \mathcal{G}_2^{2,2}$: Finite graphs with no edges with V_0, V_1 an arbitrary partition.

To construct the members of $P(3)$, we can use the following procedure:

- First we decide what \mathcal{A}^{ι_0} , \mathcal{A}^{ι_1} , and $\mathcal{A}^{\iota_2} \in P(1)$ will be, with the restriction that if $\mathcal{A}^{\iota_i} = \mathcal{G}_4$, then $i = 0$.
- Now we decide what the classes $\mathcal{A}^{\epsilon_{0,3}}$, $\mathcal{A}^{\epsilon_{1,3}}$, and $\mathcal{A}^{\epsilon_{2,3}} \in P(2)$ will be (Notation 4.3.4). These choices need to be consistent with our choices for \mathcal{A}^{ι_j} ; for example, since $\epsilon_{0,3} \circ \iota_0 = \iota_1$, we must have $(\mathcal{A}^{\epsilon_{0,3}})^{\iota_0} = \mathcal{A}^{\iota_1}$.
- The last step is to decide whether structures in \mathcal{A} are allowed to have triangles. Sometimes, our previous choices will already dictate yes or no. If $\mathcal{A}^{\iota_0} = \mathcal{G}_4$, or if for any $i < 3$, $\mathcal{A}^{\epsilon_{i,3}}$ allows triangles, then \mathcal{A} must allow triangles. On the other hand, if all of the $\mathcal{A}^{\epsilon_{i,3}}$ forbid triangles and at least one of them forbids edges, then \mathcal{A} cannot allow triangles. Outside of these cases, we are allowed to choose whether or not structures in \mathcal{A} are allowed triangles; if so, any triangles must have one vertex each in V_0, V_1 , and V_2 .

Since \mathcal{G}_4 is defined by forbidding an irreducible structure of size 4, any $\mathcal{A} \in P(d)$ for $d \geq 4$ is completely determined by the classes \mathcal{A}^e where $e: 3 \rightarrow d$ ranges over increasing injections. Furthermore, letting $I(3, d)$ denote the set of increasing injections from 3 to d , one can show that if $w: I(3, d) \rightarrow P(3)$ is any “consistent” assignment of increasing injections to classes, where “consistent” just means that there are no obvious conflicts on overlapping pairs or singletons, then there is $\mathcal{A} \in P(d)$ with $\mathcal{A}^e = w(e)$ for every $e \in I(3, d)$.

Example 4.3.6. Generalizing Example 4.3.5, we can fix any $\ell \geq 3$ and consider the class $\mathcal{K} = \mathcal{G}_\ell$ of K_ℓ -free graphs. Given $d < \omega$, let $\binom{d}{<\ell}$ denote the collection of subsets of $\{0, \dots, d-1\}$ of size at most $\ell-1$. A typical member of $P(d)$ is described by a function $z: \binom{d}{<\ell} \rightarrow \{1, 2, \dots, \ell-1\}$ satisfying the following properties:

1. If $S \subseteq d$, $|S| = 1$, and $z(S) = \ell - 1$, then $S = \{0\}$,
2. z is monotone: If $S, T \in \binom{d}{<\ell}$ and $S \subseteq T$, then $z(S) \leq z(T)$,
3. z is consistent: For each $T \in \binom{d}{<\ell}$, there is a function $\sigma: T \rightarrow \omega$ with $\sum_{i \in T} \sigma(i) = z(T)$ and so that for any $S \subseteq T$, we have $\sum_{i \in S} \sigma(i) \leq z(S)$.

We call a function z satisfying the above 3 items a (d, ℓ) -graph-age function. Given a (d, ℓ) -graph-age function, we define the class \mathcal{G}_z of finite \mathcal{L}_d -structures via

$$\mathcal{G}_z := \left\{ \mathbf{B} : \forall T \in \binom{d}{<\ell} \left(\bigcup_{i \in T} V_i^{\mathbf{B}} \text{ contains no } (z(T) + 1)\text{-clique} \right) \right\}.$$

So the number $z(T)$ describes the largest clique which is allowed to appear among nodes in $\bigcup_{i \in T} V_i$. For instance, the class $\mathcal{G}_4^{3,2}$ from Example 4.3.5 is the class \mathcal{G}_z where $z(\{0\}) = 2$, $z(\{1\}) = 1$, and $z(\{0, 1\}) = 3$.

Unfortunately, checking whether a given function z satisfies the “consistency” of item 3 seems to be a challenging combinatorial problem. For example, this is the reason why in Example 4.3.5, in the last bullet describing the procedure to build a member of $P(3)$, \mathcal{A} was sometimes forced to not allow triangles.

4.4 The spectrum of a gluing and a class

This subsection begins a detailed analysis of gluings $\gamma = (\rho, \eta, \mathbf{I})$ and the resulting classes of structures $\mathcal{K}(\gamma)$. The idea is to pay special attention to irreducible substructures of \mathbf{I} and understand how the gluing restricted to an irreducible substructure of \mathbf{I} affects the class $\mathcal{K}(\gamma)$. In particular, Corollary 4.4.3 will tell us that the poset $P(\rho)$ is finite. We then turn our attention to classes in $P(\rho)$, showing that any $\mathcal{A} \in P(\rho)$ can be written as an intersection of classes $\mathcal{K}(\gamma)$, where γ ranges among a family of “basic” gluings.

Definition 4.4.1.

1. Recalling that $\mathcal{K} = \text{Forb}(\mathcal{F})$ for \mathcal{F} a finite set of finite irreducible \mathcal{L} -structures, we let \mathcal{I} denote the set of enumerated structures in \mathcal{K} which admit an embedding into some structure in \mathcal{F} . As \mathcal{F} is finite, so is \mathcal{I} .
2. Suppose $\gamma = (\rho, \eta, \mathbf{I})$ with rank d and size n . If \mathbf{J} is another enumerated \mathcal{L} -structure and $f: \mathbf{J} \rightarrow \mathbf{I}$ is an embedding, we write $\gamma \cdot f$ for the gluing $(\rho, \eta \cdot f, \mathbf{J})$, where given $i < d$ and $m < |\mathbf{J}|$, we have $(\eta \cdot f)(i, m) = \eta(i, f(m))$.

3. Suppose $\gamma = (\rho, \eta, \mathbf{I})$ is a gluing. The *spectrum* of γ is the following set of gluings:

$$\text{Spec}(\gamma) = \bigcup_{\mathbf{J} \in \mathcal{I}} \{\gamma \cdot f : f \in \text{OEmb}(\mathbf{J}, \mathbf{I})\}.$$

Since every gluing in the spectrum has sort ρ , and since \mathcal{I} is finite, we see that there are only finitely many possible spectra of gluings with sort ρ .

Lemma 4.4.2. *Fix a sort $\rho: d \rightarrow k^u$, and suppose $\gamma = (\rho, \eta_\gamma, \mathbf{I}_\gamma)$ and $\zeta = (\rho, \eta_\zeta, \mathbf{I}_\zeta)$ are gluings. If $\text{Spec}(\gamma) \subseteq \text{Spec}(\zeta)$, then $\mathcal{K}(\gamma) \supseteq \mathcal{K}(\zeta)$. In particular, if the spectra are equal, then the ages are equal.*

Proof. Suppose \mathbf{B} is a finite \mathcal{L}_d -structure so that $\mathbf{B} \notin \mathcal{K}(\gamma)$. Find $\mathbf{F} \in \mathcal{F}$ and an embedding $g_\gamma: \mathbf{F} \rightarrow \mathbf{B}[\gamma]$ which witnesses that $\mathbf{B} \notin \mathcal{K}(\gamma)$. There is a unique enumerated \mathcal{L} -structure \mathbf{J} and a unique $f_\gamma \in \text{OEmb}(\mathbf{J}, \mathbf{I}_\gamma)$ with $f_\gamma[\mathbf{J}] = g_\gamma[\mathbf{F}] \cap \mathbf{I}_\gamma$; since $\mathbf{I}_\gamma \in \mathcal{K}$, we have $\mathbf{J} \in \mathcal{I}$. So $\gamma \cdot f_\gamma \in \text{Spec}(\gamma) \subseteq \text{Spec}(\zeta)$. Find $f_\zeta: \mathbf{J} \rightarrow \mathbf{I}_\zeta$ with $\zeta \cdot f_\zeta = \gamma \cdot f_\gamma$. Write $h: \mathbf{J} \rightarrow \mathbf{F}$ for the embedding with $g_\gamma \circ h = f_\gamma$. Define $g_\zeta: \mathbf{F} \rightarrow \mathbf{B}[\zeta]$ via

$$g_\zeta(x) = \begin{cases} f_\zeta(y) & \text{if } x = h(y) \\ g_\gamma(x) & \text{if } x \notin h[\mathbf{J}] \end{cases}$$

Since f_ζ , h , and g_γ are embeddings, it's clear that $U^{\mathbf{B}[\zeta]}(g_\zeta(x)) = U^{\mathbf{F}}(x)$ for each $x \in \mathbf{F}$ and that $R^{\mathbf{B}[\zeta]}(g_\zeta(z), g_\zeta(x)) = R^{\mathbf{F}}(z, x)$ whenever both $x, z \in h[\mathbf{J}]$ or both $x, z \in \mathbf{F} \setminus h[\mathbf{J}]$. In the case that $x \in h[\mathbf{J}]$ and $z \in \mathbf{F} \setminus h[\mathbf{J}]$, set $x = h(y)$. Then:

$$\begin{aligned} R^{\mathbf{B}[\zeta]}(g_\zeta(z), g_\zeta(x)) &= R^{\mathbf{B}[\zeta]}(g_\gamma(z), f_\zeta(y)) \\ &= \eta_\zeta(V^{\mathbf{B}}(g_\gamma(z)), f_\zeta(y)) \\ &= \eta_\gamma(V^{\mathbf{B}}(g_\gamma(z)), f_\gamma(y)) \quad [\text{since } \zeta \cdot f_\zeta = \gamma \cdot f_\gamma] \\ &= \eta_\gamma(V^{\mathbf{B}}(g_\gamma(z)), g_\gamma(x)) \\ &= R^{\mathbf{B}[\gamma]}(g_\gamma(z), g_\gamma(x)) \\ &= R^{\mathbf{F}}(z, x) \end{aligned}$$

Since \mathbf{F} embeds into $\mathbf{B}[\zeta]$, we see that $\mathbf{B} \notin \mathcal{K}(\zeta)$. □

Corollary 4.4.3. *For any given sort $\rho: d \rightarrow k^u$, the poset $P(\rho)$ is finite.*

Definition 4.4.4. Fix a sort $\rho: d \rightarrow k^u$.

1. If $\mathbf{J} \in \mathcal{K}$ is an enumerated structure, we write

$$\text{Glue}(\rho, \mathbf{J}) = \{(\rho, \eta, \mathbf{J}) : (\rho, \eta, \mathbf{J}) \text{ is a gluing}\}.$$

2. Suppose $\mathcal{A} \in P(\rho)$. The *spectrum* of \mathcal{A} is the set

$$\text{Spec}(\mathcal{A}) := \bigcup_{\mathbf{J} \in \mathcal{I}} \{\gamma \in \text{Glue}(\rho, \mathbf{J}) : \mathcal{A} \subseteq \mathcal{K}(\gamma)\}.$$

Remark. Every $\gamma \in \text{Spec}(\mathcal{A})$ is reasonable since \mathcal{A} is reasonable (Definition 4.2.5).

Proposition 4.4.5. *In the setting of Definition 4.4.4, we have*

$$\mathcal{A} = \bigcap_{\gamma \in \text{Spec}(\mathcal{A})} \mathcal{K}(\gamma).$$

Proof. By definition, we have that \mathcal{A} is a subclass of the right hand side. So suppose $\mathbf{B} \notin \mathcal{A}$. Suppose $S \subseteq \text{CT}(n)$ is such that $\mathcal{K}(S) = \mathcal{A}$. Form the structure $\mathbf{B}[S] \notin \mathcal{K}$, and find some $\mathbf{F} \in \mathcal{F}$ and embedding $g: \mathbf{F} \rightarrow \mathbf{B}[S]$. Write $\mathbf{J} = g^{-1}(\mathbf{K}_n)$. Then there is a unique $\mathbf{I} \in \mathcal{I}$ and a unique $f \in \text{OEmb}(\mathbf{I}, \mathbf{K}_n)$ with $f[\mathbf{I}] = g[\mathbf{J}]$. Setting $\gamma = (S^u, S^b \cdot f, \mathbf{I})$, we see that $\gamma \in \text{Spec}(\mathcal{A})$ and $\mathbf{B} \notin \mathcal{K}(\gamma)$. \square

Corollary 4.4.6. *In the setting of Definition 4.4.4, suppose $\mathcal{A}, \mathcal{B} \in P(\rho)$ with $\mathcal{A} \supseteq \mathcal{B}$. Then $\text{Spec}(\mathcal{A}) \subsetneq \text{Spec}(\mathcal{B})$.*

4.5 Consecutive age changes

Since Corollary 4.4.3 tells us that the poset $P(\rho)$ is finite, it makes sense to study those pairs of classes which are consecutive in this poset.

Definition 4.5.1.

1. Fix a sort $\rho: d \rightarrow k^u$. A *consecutive age change* in $P(\rho)$ is a pair $(\mathcal{A}, \mathcal{B})$ with $\mathcal{A}, \mathcal{B} \in P(\rho)$, $\mathcal{B} \subsetneq \mathcal{A}$, and such that there is no $\mathcal{C} \in P(\rho)$ with $\mathcal{B} \subsetneq \mathcal{C} \subsetneq \mathcal{A}$. Write $\text{Con}(\rho)$ for the set of consecutive age changes in $P(\rho)$.
2. A consecutive age change $(\mathcal{A}, \mathcal{B})$ is called *essential* if for any increasing injection $e: d_0 \rightarrow d$ with $d_0 < d$, we have $\mathcal{A}^e = \mathcal{B}^e$. Write $\text{ECon}(\rho)$ for the set of essential consecutive age changes in $P(\rho)$.
3. Suppose $(\mathcal{A}, \mathcal{B}) \in \text{ECon}(\rho)$. Then by Corollary 4.4.6, we have $\text{Spec}(\mathcal{B}) \setminus \text{Spec}(\mathcal{A}) \neq \emptyset$. Fix some choice of $\gamma = (\rho, \eta, \mathbf{J}) \in \text{Spec}(\mathcal{B}) \setminus \text{Spec}(\mathcal{A})$ with \mathbf{J} of minimum possible cardinality. We set $\gamma = \text{Cause}(\mathcal{A}, \mathcal{B})$ and call γ the *cause* of the essential consecutive age change $(\mathcal{A}, \mathcal{B})$.

Remark. Suppose $(\mathcal{A}, \mathcal{B}) \in \text{ECon}(\rho)$, and write $\gamma = \text{Cause}(\mathcal{A}, \mathcal{B})$. Then $\mathcal{B} \subseteq \mathcal{A} \cap \mathcal{K}(\gamma) \subsetneq \mathcal{A}$, and by Proposition 4.2.6, we have $\mathcal{A} \cap \mathcal{K}(\gamma) \in P(\rho)$. It follows that $\mathcal{B} = \mathcal{A} \cap \mathcal{K}(\gamma)$.

Remark. One can show (see Lemma 8.0.3) that if $\text{ECon}(\rho) \neq \emptyset$ and $D = \max\{|\mathbf{F}| : \mathbf{F} \in \mathcal{F}\}$, then $\text{dom}(\rho) < D$.

The particular choice of “cause” that we made in Definition 4.5.1 is entirely arbitrary, but later, when constructing countable structures where we want to control exactly how age changes happen, we will create an age change by “causing” it using the specific gluing above. One could attempt to naïvely define the cause of any consecutive age change, not just essential consecutive age changes. However, we will soon see that given a consecutive age change $(\mathcal{A}, \mathcal{B}) \in \text{Con}(\rho)$, there is a

unique $d_0 \leq d$ and increasing injection $e: d_0 \rightarrow d$ with $(\mathcal{A}^e, \mathcal{B}^e) \in \text{ECon}(\rho \circ e)$. This will allow us to coherently extend the definition to any consecutive age change (see Definition 4.5.4).

Proposition 4.5.2. *Fix $\mathcal{A} \in P(\rho)$, an increasing injection $e: d_0 \rightarrow d$, and $\mathcal{B} \in P(\rho \circ e)$. Suppose $(\mathcal{A}^e, \mathcal{B}) \in \text{ECon}(\rho \circ e)$. Write $\mathcal{C} = \mathcal{A}|_{(e, \mathcal{B})}$ (Definition 4.3.1(4)). Then e is the unique increasing injection with range d such that $(\mathcal{A}^e, \mathcal{C}^e) \in \text{ECon}(\rho \circ e)$. More precisely, if $e_1: d_1 \rightarrow d$ is an increasing injection with $\text{Im}(e_1) \not\supseteq \text{Im}(e)$, then $\mathcal{A}^{e_1} = \mathcal{C}^{e_1}$.*

Remark. We may not have $(\mathcal{A}, \mathcal{A}|_{(e, \mathcal{B})}) \in \text{Con}(\rho)$. For a counterexample, consider the class $\mathcal{K} = \mathcal{G}_4$ of K_4 -free graphs (see Example 4.3.5). Consider the class $\mathcal{A} \in P(2)$ with $(\mathcal{A}^{\iota_0}, \mathcal{A}^{\iota_1}) = (\mathcal{G}_3, \mathcal{G}_2)$ and which allows triangles (so all triangles must have two vertices in V_0 and one in V_1). Consider $\mathcal{B} = \mathcal{G}_2 \in P(1)$ and $e = \iota_0$. Then $\mathcal{A}|_{(e, \mathcal{B})}$ is the class of bipartite graphs where V_0, V_1 is a bipartition; the class $\mathcal{D} \in P(2)$ with $(\mathcal{D}^{\iota_0}, \mathcal{D}^{\iota_1}) = (\mathcal{G}_3, \mathcal{G}_2)$ and where triangles are not allowed lies strictly between \mathcal{A} and $\mathcal{A}|_{(e, \mathcal{B})}$.

Proof. Fix $e_1: d_1 \rightarrow d$ an increasing injection different from e . First suppose $\text{Im}(e_1) \supseteq \text{Im}(e)$, and let $e_0: d_0 \rightarrow d_1$ be the increasing injection with $e_1 \circ e_0 = e$. Then $(\mathcal{A}^{e_1}, \mathcal{C}^{e_1})$ is not essential as $(\mathcal{A}^{e_1})^{e_0} = \mathcal{A}^e$ and $(\mathcal{C}^{e_1})^{e_0} = \mathcal{C}^e$. Now suppose that $\text{Im}(e_1) \not\supseteq \text{Im}(e)$. Then we argue that $\mathcal{C}^{e_1} = \mathcal{A}^{e_1}$. Certainly $\mathcal{C}^{e_1} \subseteq \mathcal{A}^{e_1}$. Now suppose $\mathbf{A} \in \mathcal{A}^{e_1}$, and form $e_1 \cdot \mathbf{A} \in \mathcal{A}$. To see that $e_1 \cdot \mathbf{A} \in \mathcal{C} := \mathcal{A}|_{(e, \mathcal{B})}$, suppose not, i.e., that $(e_1 \cdot \mathbf{A})^e \notin \mathcal{B}$. Let e_0 be the increasing enumeration of $e^{-1}(\text{Im}(e_1)) \subseteq d_0$. Then we have

$$e_0 \cdot ((e_1 \cdot \mathbf{A})^{e \circ e_0}) = (e_1 \cdot \mathbf{A})^e.$$

which implies that $(e_1 \cdot \mathbf{A})^{e \circ e_0} \notin \mathcal{B}^{e_0}$. But this is impossible as $(\mathcal{A}^e, \mathcal{B})$ is essential. \square

Corollary 4.5.3. *Let ρ be a sort, and fix $(\mathcal{A}, \mathcal{B}) \in \text{Con}(\rho)$.*

1. *For any increasing injection e with $\mathcal{B}^e \subsetneq \mathcal{A}^e$, we have $\mathcal{B} = \mathcal{A}|_{(e, \mathcal{B}^e)}$ and $(\mathcal{A}^e, \mathcal{B}^e) \in \text{Con}(\rho \circ e)$.*
2. *There is a unique increasing injection e with range d so that $(\mathcal{A}^e, \mathcal{B}^e) \in \text{ECon}(\rho \circ e)$.*

Proof. For item 1, note that by Proposition 4.3.3, we have $\mathcal{A}|_{(e, \mathcal{B}^e)} \in P(\rho)$. Since $\mathcal{B} \subseteq \mathcal{A}|_{(e, \mathcal{B}^e)} \subsetneq \mathcal{A}$, we must have $\mathcal{B} = \mathcal{A}|_{(e, \mathcal{B}^e)}$. Similarly, if there were some $\mathcal{C} \in P(\rho \circ e)$ with $\mathcal{B}^e \subsetneq \mathcal{C} \subsetneq \mathcal{A}^e$, then we would have $\mathcal{A}|_{(e, \mathcal{B}^e)} \subsetneq \mathcal{A}|_{(e, \mathcal{C})} \subsetneq \mathcal{A}$, a contradiction.

For item 2, simply pick any such e . By item 1, we have $\mathcal{B} = \mathcal{A}|_{(e, \mathcal{B}^e)}$, and then we appeal to Proposition 4.5.2. \square

Definition 4.5.4. Let $\rho: d \rightarrow k^u$ be a sort, and suppose $(\mathcal{A}, \mathcal{B}) \in \text{Con}(\rho)$. We extend the definition of $\text{Cause}(\mathcal{A}, \mathcal{B})$ to the case where $(\mathcal{A}, \mathcal{B})$ is possibly not essential as follows. By Corollary 4.5.3, let $e: d_0 \rightarrow d$ be the unique increasing injection with $(\mathcal{A}^e, \mathcal{B}^e) \in \text{ECon}(\rho \circ e)$. Writing $\gamma = \text{Cause}(\mathcal{A}^e, \mathcal{B}^e)$, we set $\text{Cause}(\mathcal{A}, \mathcal{B}) = (\rho, e) \cdot \gamma$.

4.6 Free relations

Notation 4.6.1. Given $\alpha < d < \omega$, we write $\delta_{\alpha,d}: (d+1) \rightarrow d$ for the non-decreasing surjection with $\delta_{\alpha,d}(\alpha) = \delta_{\alpha,d}(\alpha+1) = \alpha$.

We end this section with a more detailed analysis of how to compare members of $P(\rho)$ as we change the sort ρ . If $\text{dom}(\rho) = d$, we will be particularly interested in comparing members of $P(\rho)$ with members of $P(\rho \circ \delta_{\alpha,d})$. Suppose $S = \{s_0 \preceq_{lex} \dots \preceq_{lex} s_{d-1}\} \subseteq \text{CT}(m)$ has $S^u = \rho$, so that $\mathcal{K}(S) \in P(\rho)$. Then a common way of obtaining a member of $P(\rho \circ \delta_{\alpha,d})$ is to consider some $Q = \{t_0, \dots, t_d\} \subseteq \text{CT}(m+1)$ where for each $\beta < d$, we have $t_\beta \sqsupseteq s_{\delta_{\alpha,d}(\beta)}$. This will often be the case when s_α is a *splitting node* of some subset of CT that we are interested in. The main question we consider in this subsection is the following: how large can $\mathcal{K}(Q)$ be given $\mathcal{K}(S)$? An upper bound is given by the class $\mathcal{K}(S)^{\delta_{\alpha,d}}$. But we have already seen (see the first paragraph of Example 4.2.8) that sometimes $\mathcal{K}(S)^{\delta_{\alpha,d}} \notin P(\rho \circ \delta_{\alpha,d})$. It turns out that this phenomenon depends only on $s_\alpha^u < k^u$.

Notation 4.6.2. Given $i < k^u$, we write $P_i = P(\iota_i)$ (Notation 4.3.4). We remark that P_i has a maximum; this is the class $\mathcal{K}(\emptyset_i) = \mathcal{K}(\iota_i)$. We also remark that P_i has a minimum; by Corollary 4.2.7, P_i is closed under intersections, and by Corollary 4.4.3, P_i is finite. write $P_i^{\min} \in P_i$ for the minimum member of P_i , i.e., the intersection of all members of P_i .

Among the unary predicates $i < k^u$, some of them will admit a *free relation*, while others will not. To motivate the definition, consider the node $t = (i, 0^m) \in T_i$. Then $\mathcal{K}(t) \in P_i$ is the maximum member of P_i , since we have $\mathcal{K}(t) = \mathcal{K}(\emptyset_i)$. If $q \in [1, k]$, we want to understand when it is possible to have $\mathcal{K}(t \frown q) = \mathcal{K}(t) = \mathcal{K}(\emptyset_i)$.

Definition 4.6.3.

1. Fix $i, j < k^u$ and $q < k$. The gluing $\gamma\langle i, q, j \rangle = (\iota_i, \eta_q, \mathbf{J})$ is the gluing where \mathbf{J} is the singleton structure with unary predicate j , and $\eta_q: 1 \times 1 \rightarrow k$ is the map $\eta_q(0, 0) = q$. So given $\mathbf{B} \in \mathcal{L}_1$ with underlying set B , then $\mathbf{B}[\gamma\langle i, q, j \rangle]$ is the \mathcal{L} -structure on $B \cup \{0\}$ where $U^{\mathbf{B}[\gamma\langle i, q, j \rangle]}(b) = i$ for each $b \in B$, $U^{\mathbf{B}[\gamma\langle i, q, j \rangle]}(0) = j$, and we set $R^{\mathbf{B}[\gamma\langle i, q, j \rangle]}(b, 0) = q$ for each $b \in B$.
2. We say that $i < k^u$ has a *free relation* if there is some $(q, j) \in k \times k^u$ with $q > 0$ and $\mathcal{K}(\gamma\langle i, q, j \rangle) = \mathcal{K}(\emptyset_i)$; we call (q, j) a *free relation* for i . We let $\text{Free}: k^u \rightarrow k \times k^u$ be a function so that $\text{Free}(i) = (0, 0)$ if i does not have a free relation, and if i does have a free relation, then $\text{Free}(i)$ is an arbitrary choice of witness (q, j) . We write $\text{Free}(i)^u = j$ and $\text{Free}(i)^b = q$ for the unary and binary coordinates of the free relation function.

In a mild abuse of notation, we also write $\text{Free} \subseteq k^u$ for the set of $i < k^u$ which have a free relation.

Remark. Suppose $i \in k^u \setminus \text{Free}$, and suppose $s = (i, 0^a) \in T_i$. Then $\mathcal{K}(s) = \mathcal{K}(\emptyset_i)$, but for any $q < k$, we have $\mathcal{K}(s \frown q) \subsetneq \mathcal{K}(s)$. However, we may not have $(\mathcal{K}(s), \mathcal{K}(s \frown q)) \in \text{Con}(\iota_i)$. For an example of this, consider the language of finite

graphs where edges are either red or blue; so $k^u = 1$, $k = 3$, and we represent red by 1 and blue by 2. Now let \mathcal{K} be the class of such finite graphs with no blue K_4 and no red-red-blue triangle. Writing $s = 0^a$, we have $\mathcal{K}(s) \supseteq \mathcal{K}(s \frown 2) \supseteq \mathcal{K}(s \frown 1)$.

Example 4.6.4. The class $\mathcal{K} = \mathcal{G}_3$ of finite triangle-free graphs does not have a free relation. This is because $\mathcal{K}(\gamma\langle 0, 1, 0 \rangle) = \mathcal{G}_2$, the class of finite graphs with no edges. Similarly, the class $\mathcal{K} = \mathcal{G}_4$ of finite K_4 -free graphs also does not have a free relation.

Example 4.6.5. The class \mathcal{K} of finite directed graphs with no cyclic triangles (see Example 4.2.9) has a free relation; one can set $\text{Free}(0) = (1, 0)$. If $\mathbf{A} \in \mathcal{K}$, then viewing \mathbf{A} as an \mathcal{L}_1 -structure, we have $\mathbf{A}[\gamma\langle 0, 1, 0 \rangle] \in \mathcal{K}$, as adding a new point with directed edges emanating from the new point to all the old points will not add a cyclic triangle.

Example 4.6.6. Let $\mathcal{K} = \mathcal{K}_3^2$ denote the class of finite graphs whose nodes are colored in 2-colors and so that there are no monochromatic vertex triangles. In this example $k^u = 2$, and each of the unary predicates 0 and 1 has a free relation. For 0, the pair $(1, 1)$ will work, and for 1, the pair $(0, 1)$ will work.

Proposition 4.6.7. *Fix $i < k^u$. The following are equivalent:*

1. $i \in \text{Free}$,
2. Whenever $\rho: d \rightarrow k^u$ is a sort, $\alpha < d$ is such that $\rho(\alpha) = i$, and $\mathcal{A} \in P(\rho)$, then we have $\mathcal{A}^{\delta_{\alpha,d}} \in P(\rho \circ \delta_{\alpha,d})$.
3. Letting $\rho: 2 \rightarrow k^u$ be the sort with $\rho(0) = \rho(1) = i$, then we have $\mathcal{K}(\rho) \in P(\rho)$.

Proof. We first remark that item 3 is a particular instance of item 2. So we will show that 1 implies 2 and that $\neg 1$ implies $\neg 3$.

Assume that item 1 holds, and write $\text{Free}(i) = (q, j)$. Towards showing that item 2 holds, fix a sort $\rho: d \rightarrow k^u$, an $\alpha < d$ with $\rho(\alpha) = i$, and $\mathcal{A} \in P(\rho)$. Find $m < \omega$ and $S = \{s_0 \preceq_{lex} \cdots \preceq_{lex} s_{d-1}\} \subseteq \text{CT}(m)$ with $S^u = \rho$ and $\mathcal{K}(S) = \mathcal{A}$. Find $n \geq m$ with $c(n) = (j, 0^n)$, and define $Q \subseteq \text{CT}(n+1)$ via

$$Q = \{\text{Left}(s, n+1) : s \in S\} \cup \{\text{Left}(s_\alpha, n) \frown q\}.$$

We now check that $\mathcal{K}(Q) = \mathcal{A}^{\delta_{\alpha,d}}$; the left-to-right containment is clear, so fix $\mathbf{B} \in \mathcal{A}^{\delta_{\alpha,d}}$ with underlying set B . Then we have that $\delta_{\alpha,d} \cdot \mathbf{B} \in \mathcal{A}$, i.e., that $(\delta_{\alpha,d} \cdot \mathbf{B})[S] \in \mathcal{K}$. Now form $\mathbf{B}[Q]$. By our choice of n , $\mathbf{B}[Q]$ is the free amalgam of the induced substructures on the subsets $\{n\} \cup B$ and on $B \cup \{0, \dots, n-1\}$. The former is the free amalgam of the induced substructures on $\{n\} \cup (V^{\mathbf{B}})^{-1}(\{\alpha+1\})$ and B . And the former of these is in \mathcal{K} since $\text{Free}(i) = (q, j)$. As for the induced substructure on $B \cup \{0, \dots, n-1\}$, this is just the free amalgam of the induced substructures on $\{0, \dots, n-1\}$ (which is just \mathbf{K}_n) and $B \cup \{0, \dots, m-1\}$ (which is just $(\delta_{\alpha,d} \cdot \mathbf{B})[S]$).

Now assume that 1 does not hold. Fix $m < \omega$ and any $s \neq t \in \text{CT}_i(m)$. Without loss of generality, we have $t \neq (i, 0^m)$. Find $a < m$ with $t(a) = q \neq 0$, and write $U(a) = j < k^u$. Since (q, j) is not a free relation for $i < k^u$, we have $\mathcal{K}(\emptyset) \supseteq \mathcal{K}(\gamma\langle i, q, j \rangle) \supseteq \mathcal{K}(t)$. \square

When $i \in k^u \setminus \text{Free}$, we still want to prove a version of item 2 of Proposition 4.6.7. Even though we may not have $\mathcal{A}^{\delta_{\alpha,d}} \in P(\rho \circ \delta_{\alpha,d})$, we can in a meaningful sense get as close as possible.

Lemma 4.6.8. *Suppose $i \in k^u \setminus \text{Free}$. Then $|P(\iota_i)| \geq 2$.*

Proof. Since we are assuming $i < k^u$ is non-degenerate (see Definition 4.2.3), we can find $j_i < k^u$ and $q_i < k$ so that the two element structure $\mathbf{I} = \{x, y\}$ with $U^{\mathbf{I}}(x) = j_i$, $U^{\mathbf{I}}(y) = i$, and $R^{\mathbf{I}}(y, x) = q_i$ is in \mathcal{K} . Consider the gluing $\gamma := \gamma\langle i, q_i, j_i \rangle$. By our choice of q_i and j_i , γ is a reasonable gluing. However, since (q_i, j_i) is not a free relation for i , we have $\mathcal{K}(\gamma) \subsetneq \mathcal{K}(\emptyset_i)$. Since $\mathcal{K}(\emptyset_i) \cap \mathcal{K}(\gamma) = \mathcal{K}(\gamma)$, we can use Proposition 4.2.6 to conclude that $\mathcal{K}(\gamma) \in P_i$. \square

Lemma 4.6.9. *Suppose $i \in k^u \setminus \text{Free}$. Suppose $\mathcal{A} \in P_i$ has $(\mathcal{K}(\emptyset_i), \mathcal{A}) \in \text{Con}(\iota_i)$, and write $\gamma = (\iota_i, \eta, \mathbf{J}) = \text{Cause}(\mathcal{K}(\emptyset_i), \mathcal{A})$. Then we have $|\mathbf{J}| = 1$. In particular, there are $q < k$ and $j < k^u$ with $\gamma = \gamma\langle i, q, j \rangle$.*

Proof. By assumption, when defining the cause of an age change, \mathbf{J} is chosen with minimum possible cardinality. Suppose $|\mathbf{J}| = n$. Let \mathbf{I} be the enumerated \mathcal{L} -structure and $f \in \text{OEmb}(\mathbf{I}, \mathbf{J})$ the ordered embedding such that $f[\mathbf{I}] = \{m < |\mathbf{J}| : \eta(0, m) \neq 0\}$. By free amalgamation, we have $\mathcal{K}(\gamma \cdot f) = \mathcal{K}(\gamma)$ (Definition 4.4.1). By Definition 4.5.1, we therefore have $\mathbf{I} = \mathbf{J}$, and $\eta(0, m) \neq 0$ for every $m < n$. Set $q := \eta(0, 0) \neq 0$. Now if $U^{\mathbf{J}}(1) = j$, notice that $\gamma|_{\mathbf{J}_1} = \gamma\langle i, q, j \rangle$. Since $i \notin \text{Free}$, we have $\mathcal{K}(\gamma\langle i, q, j \rangle) \subsetneq \mathcal{K}(\emptyset_i)$. So we must have $\mathcal{K}(\gamma\langle i, q, j \rangle) = \mathcal{A}$, and the minimality of $|\mathbf{J}|$ in Definition 4.5.1 implies that $|\mathbf{J}| = 1$. \square

Lemma 4.6.10. *Suppose $i \in k^u \setminus \text{Free}$. Suppose $\rho: d \rightarrow k^u$ is a sort, and suppose $\alpha < d$ has $\rho(\alpha) = i$. Suppose $\mathcal{A} \in P(\rho)$ is such that $\mathcal{A}^{\iota_\alpha} = \mathcal{K}(\emptyset_i)$. Suppose $\mathcal{B} \in P(\rho)$ satisfies $\mathcal{B} \subsetneq \mathcal{A}$, and $\mathcal{B}^{\epsilon_{\alpha,d}} = \mathcal{A}^{\epsilon_{\alpha,d}}$. Then we have $\mathcal{B}^{\iota_\alpha} \subsetneq \mathcal{K}(\emptyset_i)$.*

In particular, given $\mathcal{D} \in P_i$ with $(\mathcal{K}(\emptyset_i), \mathcal{D}) \in \text{Con}(\iota_i)$, then if $\mathcal{A}|_{(\iota_\alpha, \mathcal{D})} \in P(\rho)$, then we have $(\mathcal{A}, \mathcal{A}|_{(\iota_\alpha, \mathcal{D})}) \in \text{Con}(\rho)$.

Proof. Enlarging \mathcal{B} if necessary, we may assume $(\mathcal{A}, \mathcal{B}) \in \text{Con}(\rho)$, so let $\gamma = (\rho, \eta, \mathbf{I}) = \text{Cause}(\mathcal{A}, \mathcal{B})$. Towards a contradiction, suppose $\mathcal{B}^{\iota_\alpha} = \mathcal{K}(\emptyset_i)$. Since $i \notin \text{Free}$, it follows that for every $x \in \mathbf{I}$, we must have $\eta(\alpha, x) = 0$. However, this in turn implies that $\mathcal{B}^{\epsilon_{\alpha,d}} \subsetneq \mathcal{A}^{\epsilon_{\alpha,d}}$, a contradiction.

For the ‘‘in particular,’’ the first part shows that if $\mathcal{B} \in P(\rho)$ satisfies $\mathcal{A}|_{(\iota_\alpha, \mathcal{D})} \subseteq \mathcal{B} \subsetneq \mathcal{A}$, then we must have $\mathcal{B}^{\iota_\alpha} = \mathcal{D}$. But then by definition of $\mathcal{A}|_{(\iota_\alpha, \mathcal{D})}$, we have $\mathcal{B} \subseteq \mathcal{A}|_{(\iota_\alpha, \mathcal{D})}$, so the two classes are equal. \square

Proposition 4.6.11. *Suppose $i \in k^u \setminus \text{Free}$. Fix a sort $\rho: d \rightarrow k^u$, a class $\mathcal{A} \in P(\rho)$, and $\alpha < d$ with $\rho(\alpha) = i$.*

1. *If $\mathcal{A}^{\iota_\alpha} \subsetneq \mathcal{K}(\emptyset_i)$, then $\mathcal{A}^{\delta_{\alpha,d}} \in P(\rho \circ \delta_{\alpha,d})$.*
2. *If $\mathcal{A}^{\iota_\alpha} = \mathcal{K}(\emptyset_i)$, then for any $\mathcal{D} \in P_i$ with $(\mathcal{K}(\emptyset_i), \mathcal{D}) \in \text{Con}(\iota_i)$, we have $(\mathcal{A}^{\delta_{\alpha,d}})|_{(\iota_{\alpha+1}, \mathcal{D})} \in P(\rho \circ \delta_{\alpha,d})$.*

Proof. Write $\mathcal{A} = \mathcal{K}(S)$, where $S = \{s_0 \preceq_{lex} \cdots \preceq_{lex} s_{d-1}\} \subseteq \text{CT}(m)$. First assume that $\mathcal{A}^{\iota_\alpha} \subsetneq \mathcal{K}(\emptyset_i)$. Then we must have $s_\alpha \neq (i, 0^m)$. Find $a < m$ with $s_\alpha(a) = q \neq 0$. Write $U(a) = j < k^u$. Find $n \geq m$ with $c(n) = (j, 0^n)$, and define $Q \subseteq \text{CT}(n+1)$ via

$$Q = \{\text{Left}(s, n+1) : s \in S\} \cup \{\text{Left}(s_\alpha, n) \frown q\}.$$

The verification that $\mathcal{K}(Q) = \mathcal{A}^{\delta_{\alpha,d}}$ is very similar to the one from the proof of Proposition 4.6.7. With notation as in that argument, the only difference is this: to conclude that the induced substructure on $\{n\} \cup (V^{\mathbf{B}})^{-1}(\{\alpha+1\})$ is in \mathcal{K} , we note that this is isomorphic to the induced substructure on $\{a\} \cup (V^{\mathbf{B}})^{-1}(\{\alpha+1\})$, and this is a substructure of $(\delta_{\alpha,d} \cdot \mathbf{B})[S]$.

Now assume that $\mathcal{A}^{\iota_\alpha} = \mathcal{K}(\emptyset_i)$, and fix $\mathcal{D} \in P_i$ with $(\mathcal{K}(\emptyset_i), \mathcal{D}) \in \text{Con}(\iota_i)$. Write $\text{Cause}(\mathcal{K}(\emptyset_i), \mathcal{D}) = \gamma$; by Lemma 4.6.9, there are $q < k$ and $j < k^u$ with $\gamma = \gamma\langle i, q, j \rangle$. Find $n \geq m$ with $c(n) = (j, 0^n)$, and define $Q \subseteq \text{CT}(n+1)$ via

$$Q = \{\text{Left}(s, n+1) : s \in S\} \cup \{\text{Left}(s_\alpha, n) \frown q\}.$$

To show that $\mathcal{K}(Q) = (\mathcal{A}^{\delta_{\alpha,d}})|_{(\iota_{\alpha+1}, \mathcal{D})}$, the left-to-right inclusion is clear, so fix \mathbf{B} in the right hand side, with underlying set B . Once again, the verification that $\mathbf{B}[Q] \in \mathcal{K}$ is very similar to the one from the proof of Proposition 4.6.7. This time, to show that the induced substructure on $\{n\} \cup (V^{\mathbf{B}})^{-1}(\{\alpha+1\})$ is in \mathcal{K} , we note that this structure is isomorphic to $\mathbf{B}^{\iota_{\alpha+1}}[\gamma]$. Since by assumption $\mathbf{B}^{\iota_{\alpha+1}} \in \mathcal{D} = \mathcal{K}(\gamma)$, we are done. \square

5 Diagonal substructures

5.1 Aged embeddings, envelopes, and closures

Definition 5.1.1. Let \mathbf{A}, \mathbf{B} be enumerated structures which embed into \mathbf{K} (finite or countably infinite). An *aged embedding* of $\text{CT}^{\mathbf{A}}$ into $\text{CT}^{\mathbf{B}}$ is a map $f: \text{CT}^{\mathbf{A}} \rightarrow \text{CT}^{\mathbf{B}}$ with the following properties:

- f preserves unaries: $f[\text{CT}_i^{\mathbf{A}}] \subseteq T_i$ for each $i < k^u$,
- f preserves meets, tree order, and relative levels; we write $\tilde{f}: |\mathbf{A}| \rightarrow |\mathbf{B}|$ for the induced map on levels,
- f preserves passing numbers: for each $s \in \text{CT}^{\mathbf{A}}$ and $m < \ell(s)$, we have $s(m) = f(s)(\tilde{f}(m))$,
- f preserves coding nodes: For each $m < |\mathbf{A}|$, we have $f(c^{\mathbf{A}}(m)) = c^{\mathbf{B}}(\tilde{f}(m))$,
- f preserves ages: for each $m < |\mathbf{A}|$, we have $\mathcal{K}(\text{CT}^{\mathbf{A}}(m), \mathbf{A}) = \mathcal{K}(f[\text{CT}^{\mathbf{A}}(m)], \mathbf{B})$.

We write $\text{AEmb}(\text{CT}^{\mathbf{A}}, \text{CT}^{\mathbf{B}})$ for the collection of aged embeddings of $\text{CT}^{\mathbf{A}}$ into $\text{CT}^{\mathbf{B}}$. Notice that if $f \in \text{AEmb}(\text{CT}^{\mathbf{A}}, \text{CT}^{\mathbf{B}})$ and $g \in \text{AEmb}(\text{CT}^{\mathbf{B}}, \text{CT}^{\mathbf{C}})$, then we have

$$g \circ f \in \text{AEmb}(\text{CT}^{\mathbf{A}}, \text{CT}^{\mathbf{C}}).$$

Notice that because f preserves unaries, coding nodes, and passing numbers, the map $\tilde{f}: \mathbf{A} \rightarrow \mathbf{B}$ is an ordered embedding. A key property of aged embeddings proven in [32] is that when $\mathbf{B} = \mathbf{K}$, any aged embedding f which has been defined up to $\text{CT}^{\mathbf{A}}(< m)$ for some $m < |\mathbf{A}|$ can be extended to all of $\text{CT}^{\mathbf{A}}$. The difficulty is finding coding nodes in CT where we need them; since f preserves ages and \mathbf{K} is left-dense, this is possible.

The following is Theorem 3.5 from [32].

Theorem 5.1.2 ([32]). *Fix $\mathbf{A} \in \mathcal{K}$ an enumerated structure, and let*

$$\chi: \text{AEmb}(\text{CT}^{\mathbf{A}}, \text{CT}) \rightarrow m$$

be a finite coloring. Then there is $h \in \text{AEmb}(\text{CT}, \text{CT})$ so that the coloring $\chi \cdot h: \text{AEmb}(\text{CT}^{\mathbf{A}}, \text{CT}) \rightarrow m$ given by $\chi \cdot h(f) = \chi(h \circ f)$ is constant.

Remark. Coding trees are defined slightly differently in [32]; instead of separating the unary predicates into the separate trees T_i at the very beginning, the coding trees in [32] equip the coding nodes with the information of which unary predicate is intended. However, the proof of Theorem 5.1.2 for the modified trees we use here is almost identical.

Theorem 5.1.2 means that members of $\text{OEmb}(\mathbf{A}, \mathbf{K})$ of the form \tilde{f} for some $f \in \text{AEmb}(\text{CT}^{\mathbf{A}}, \text{CT})$ are desirable to work with. The next definitions work towards the combinatorial characterization of such embeddings given in Proposition 5.1.4.

Definition 5.1.3. Fix $\mathbf{A} \subseteq \mathbf{K}$.

1. The *splitting node set* of \mathbf{A} is

$$\text{SpNode}(\mathbf{A}) = \{c(a) \wedge c(b) : a, b \in \mathbf{A}, U(a) = U(b), \text{ and } c(a) \perp c(b)\} \subseteq T.$$

The *splitting level set* of \mathbf{A} is $\text{Sp}(\mathbf{A}) = \{\ell(x) : x \in \text{SpNode}(\mathbf{A})\} \subseteq \mathbf{K}$.

2. If $m < \omega$, write $c[\mathbf{A}]|_m = \{c(a)|_m : a \in \mathbf{A} \setminus m\} \subseteq \text{CT}(m)$. The *age-change set* of \mathbf{A} is

$$\text{AC}(\mathbf{A}) = \{m \notin (\mathbf{A} \cup \text{Sp}(\mathbf{A})) : \mathcal{K}(c[\mathbf{A}]|_m) \neq \mathcal{K}(c[\mathbf{A}]|_{m+1})\}.$$

3. The *critical set* of \mathbf{A} is $\text{Crit}(\mathbf{A}) := \mathbf{A} \cup \text{Sp}(\mathbf{A}) \cup \text{AC}(\mathbf{A})$.
4. We call $\mathbf{A} \subseteq \mathbf{K}$ *closed* if $\text{Crit}(\mathbf{A}) = \mathbf{A}$, equivalently if $\text{Sp}(\mathbf{A}) \subseteq \mathbf{A}$ and $\text{AC}(\mathbf{A}) = \emptyset$, and we write $\overline{\mathbf{A}}$, the *closure* of \mathbf{A} , for the smallest closed set containing \mathbf{A} . We will soon discuss why this exists. If \mathbf{A} is an enumerated structure and $f \in \text{OEmb}(\mathbf{A}, \mathbf{K})$, we call f *closed* if $f[\mathbf{A}] \subseteq \mathbf{K}$ is closed.
5. If $\mathbf{A} \subseteq \mathbf{K}$ is finite, then the *interior* of \mathbf{A} , denoted $\text{Int}(\mathbf{A})$, is the smallest subset of \mathbf{A} with $\overline{\text{Int}(\mathbf{A})} = \overline{\mathbf{A}}$. We will soon discuss why this exists.
6. The *envelope* of \mathbf{A} , denoted $\tilde{\mathbf{A}}$, is the unique enumerated structure on underlying set $|\overline{\mathbf{A}}|$ such that the increasing bijection $\tilde{\mathbf{A}} \rightarrow \overline{\mathbf{A}}$ is an embedding.

Remark. There are some mild differences between the definitions and notation here compared to those given in [32]. Here, $\text{AC}(\mathbf{A})$ is defined to only pick up age changes occurring at levels not in \mathbf{A} . Also, the notion of envelope here is the enumerated structure $\tilde{\mathbf{A}}$ rather than the structure $\overline{\mathbf{A}}$.

As promised, we briefly discuss why closures and interiors exist by giving explicit constructions. For an infinite $\mathbf{A} \subseteq \mathbf{K}$, one can construct the closure of \mathbf{A} in countably many steps as follows. First set $\mathbf{A}^0 = \mathbf{A}$. If \mathbf{A}^n has been defined, set $\mathbf{A}^{n+1} = \mathbf{A}^n \cup \text{Sp}(\mathbf{A}^n) \cup \text{AC}(\mathbf{A}^n)$. Then $\overline{\mathbf{A}} = \bigcup_n \mathbf{A}^n$.

If we start with a finite $\mathbf{A} \subseteq \mathbf{K}$, we can compute $\overline{\mathbf{A}}$ using a finite, “top-down” procedure. Write $n = \max\{\mathbf{A}\}$, and set $\mathbf{A}^n = \{n\}$. If $m \in [0, n)$ and $\mathbf{A}^{m+1} \subseteq [m+1, n]$ has been determined, then set $\mathbf{A}^m = \mathbf{A}^{m+1} \cup \{m\}$ if $m \in \mathbf{A} \cup \text{Sp}(\mathbf{A}^{m+1}) \cup \text{AC}(\mathbf{A}^{m+1})$, and set $\mathbf{A}^m = \mathbf{A}^{m+1}$ otherwise. Then $\mathbf{A}^0 = \overline{\mathbf{A}}$. In particular, the closure of every finite substructure of \mathbf{K} is finite. However, the sizes of closures of substructures of size m could be unbounded.

Similarly, for finite $\mathbf{A} \subseteq \mathbf{K}$, we can also construct $\text{Int}(\mathbf{A})$ using a “top-down” procedure as follows: write $n = \max\{\mathbf{A}\}$, and set $\mathbf{A}^n = \{n\}$. If $m \in [0, n)$ and $\mathbf{A}^{m+1} \subseteq \mathbf{A} \setminus \{0, \dots, m\}$ has been determined, then set $\mathbf{A}^m = \mathbf{A}^{m+1} \cup \{m\}$ if $m \in \mathbf{A} \setminus \overline{\mathbf{A}^{m+1}}$, and set $\mathbf{A}^m = \mathbf{A}^{m+1}$ otherwise. Then $\mathbf{A}^0 = \text{Int}(\mathbf{A})$.

As promised, the notion of closed substructure/embedding is exactly what is needed to characterize which $f \in \text{OEmb}(\mathbf{A}, \mathbf{K})$ have the form $\tilde{\varphi}$ for some $\varphi \in \text{AEmb}(\text{CT}^{\mathbf{A}}, \text{CT}^{\mathbf{K}})$. The following is Proposition 4.2 from [32].

Proposition 5.1.4. *Let \mathbf{A} be a finite or countably infinite enumerated structure which embeds into \mathbf{K} . Then $f \in \text{OEmb}(\mathbf{A}, \mathbf{K})$ is closed iff there is $\varphi \in \text{AEmb}(\text{CT}^{\mathbf{A}}, \text{CT})$ with $f = \tilde{\varphi}$.*

An example of a closed set that we will often use is the following.

Proposition 5.1.5. *Suppose $\mathbf{B} \subseteq \mathbf{K}$ is left-leaning. Then \mathbf{B} is closed. In particular, if \mathbf{A} is an enumerated structure and $f \in \text{OL}(\mathbf{A}, \mathbf{K})$, then f is closed.*

Proof. Suppose $m \in \text{Sp}(\mathbf{A})$, and find $a, b \in \mathbf{A}$ with $c(a) \wedge c(b) \in T(m)$. If $a = m$ or $b = m$, then $m \in \mathbf{A}$, so assume $a, b > m$. Then for either a or b , say b , we have $c(b)(m) \neq 0$, i.e., $R(b, m) \neq 0$. Since $m < b$ and $\mathbf{A} \subseteq \mathbf{K}$ is left-leaning, we have $m \in \mathbf{A}$.

Now suppose $m \notin (\mathbf{A} \cup \text{Sp}(\mathbf{A}))$. If $a \in \mathbf{A}$ with $a > m$, we must have $R(a, m) = 0$. It follows that $t(m) = 0$ for every $t \in c[\mathbf{A}]|_{m+1}$, implying that $\mathcal{K}(c[\mathbf{A}]|_m) = \mathcal{K}(c[\mathbf{A}]|_{m+1})$. So $m \notin \text{AC}(\mathbf{A})$, and $\text{AC}(\mathbf{A}) = \emptyset$. \square

We can now fill in a gap remaining from Section 4.

Proposition 5.1.6. *Given a sort ρ , the poset $P(\rho)$ does not depend on the choice of left-dense Fraïssé limit \mathbf{K} .*

Proof. Let \mathbf{A} be another enumerated, left-dense Fraïssé limit, and write $P^{\mathbf{A}}(\rho)$ for the analog of $P(\rho)$ as defined using \mathbf{A} instead of \mathbf{K} . Fix $m < \omega$ and $S \subseteq \text{CT}^{\mathbf{A}}(m)$. Let $f \in \text{OL}(\mathbf{A}, \mathbf{K})$. By Proposition 5.1.5, f is closed, so find some $\varphi \in \text{AEmb}(\text{CT}^{\mathbf{A}}, \text{CT})$ with $f = \tilde{\varphi}$. Noting that φ is an aged embedding and in particular preserves lex-order, then $\mathcal{K}(\varphi[S]) = \mathcal{K}(S, \mathbf{A}_n)$. This shows that $P^{\mathbf{A}}(\rho) \subseteq P(\rho)$. By a symmetric argument, we must also have the reverse inclusion. \square

The study of envelopes is what allows us to compute the big Ramsey degrees of structures in \mathcal{K} . The following is a slight strengthening of Theorem 4.5 from [32].

Theorem 5.1.7. *Suppose there is some $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ with the following two properties:*

1. *For every $n < \omega$, there is $D_n < \omega$ so that any $\mathbf{B} \subseteq f[\mathbf{K}]$ with $|\mathbf{B}| \leq n$ has $|\overline{\mathbf{B}}| \leq D_n$,*
2. *For every finite $\mathbf{B} \subseteq f[\mathbf{K}]$, we have $\mathbf{B} = \text{Int}(\mathbf{B})$.*

Then given an enumerated $\mathbf{A} \in \mathcal{K}$, the ordered Ramsey degree of \mathbf{A} in \mathbf{K} is at most $|\{g[\mathbf{A}] : g \in \text{OEmb}(\mathbf{A}, f[\mathbf{K}])\}|$. In words, this is the number of envelopes of ordered copies of \mathbf{A} inside $f[\mathbf{K}]$.

The remainder of this section works towards the (rather technical) definition and construction of a *diagonal substructure* of \mathbf{K} (Definition 5.8.2 and Theorem 5.8.5). The idea is that given $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ with $f[\mathbf{K}]$ is diagonal, then for every enumerated $\mathbf{A} \in \mathcal{K}$, the number of envelopes of ordered copies of \mathbf{A} inside $f[\mathbf{K}]$ will be as small as possible, allowing us to use Theorem 5.1.7 to obtain upper bounds on big Ramsey degrees. In Section 6, we will give the corresponding lower bound arguments to show that these bounds are exact.

We will first define the notion of a *weakly diagonal* substructure. From there, the definition of a diagonal substructure will have four main parts; they will be weakly diagonal substructures of \mathbf{K} with *controlled paths*, *controlled age change levels*, *controlled splitting levels*, and *controlled coding levels*. One major difficulty in the construction is that in order to even make sense the notion of controlled paths, we first must consider a weaker notion of having *maximal paths*. So to build a diagonal substructure, we first build a substructure with maximal paths, then controlled paths, and then we control the age change, splitting, and coding levels. Somewhat circuitously, the most straightforward way of building a substructure with maximal paths is to build a substructure with controlled age-change, splitting, and coding levels, which will imply having maximal paths. Having done this, we perform a construction which produces a substructure with controlled paths, and then we have to redo the constructions which obtain controlled age-change, splitting, and coding levels while maintaining that we have controlled paths.

5.2 Weakly diagonal substructures

This subsection starts with the notion of a *weakly diagonal substructure*. Here, we will be mainly concerned with the “shape” of $c[\mathbf{A}]$ inside CT, without worrying yet about age structure. We also take the opportunity to discuss the notion of *start* levels.

Notation 5.2.1. Fix $t \in T$. If $t^b \neq 0^{\ell(t)}$, we let $\text{Start}(t) < \ell(t)$ be the least integer with $t^b(\text{Start}(t)) \neq 0$. If $t^b = 0^{\ell(t)}$, we say that $\text{Start}(t)$ does not exist. If $a \in \mathbf{K}$, we write $\text{Start}(a) = \text{Start}(c(a))$.

If $\mathbf{A} \subseteq \mathbf{K}$, we write $\text{Start}(\mathbf{A}) = \{\text{Start}(a) : a \in \mathbf{A}, c(a)^b \neq 0^a\}$, and if $i < k^u$, we write $\text{Start}(\mathbf{A}, i) = \{\text{Start}(a) : a \in \mathbf{A}, c(a)^b \neq 0^a, U(a) = i\}$. We write $\text{FreeStart}(\mathbf{A}) = \bigcup_{i \in \text{Free}} \text{Start}(\mathbf{A}, i)$. Write $\text{Crit}^+(\mathbf{A}) = \text{Crit}(\mathbf{A}) \cup \text{Start}(\mathbf{A})$.

Given $\mathbf{A} \subseteq \mathbf{K}$ and $a \in \mathbf{A}$, when do we have $\text{Start}(a) \in \text{Crit}(\mathbf{A})$? If $U(a) \notin \text{Free}$, the answer is always, since if $\text{Start}(a) \notin \mathbf{A} \cup \text{Sp}(\mathbf{A})$, the act of leaving the left branch results in an age change. If $U(a) \in \text{Free}$, the answer is sometimes. If $i \in \text{Free}$, then if $m \in \text{Start}(\mathbf{A}, i)$ is not the largest member of $\text{Start}(\mathbf{A}, i)$, then we will have $m \in \text{Sp}(\mathbf{A})$. However, if $m = \max(\text{Start}(\mathbf{A}, i))$, then m might not be in $\text{Crit}(\mathbf{A})$, and upon defining the notion of weakly diagonal structure, if $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal, then we will have $m \notin \text{Crit}(\mathbf{A})$. However, it is still possible to have $m \in \overline{\mathbf{A}}$, and we will need control over how this happens.

Definition 5.2.2. A substructure $\mathbf{A} \subseteq \mathbf{K}$ is *weakly diagonal* if the following all hold:

1. $c[\mathbf{A}] \subseteq \text{CT}$ is an anti-chain,
2. $\text{Sp}(\mathbf{A}) \cap \mathbf{A} = \emptyset$,
3. For each $m \in \text{Sp}(\mathbf{A})$, there is a unique $s \in \text{SpNode}(\mathbf{A})$ with $\ell(s) = a$. Furthermore, this s has exactly two successors in $c[\mathbf{A}]|_{m+1}$.
4. For each $m \in \text{Sp}(\mathbf{A})$ and $t \in c[\mathbf{A}]|_m$, we have $t \frown 0 \in c[\mathbf{A}]|_{m+1}$.
5. For every $a \in \mathbf{A}$, $\text{Start}(a)$ exists.
6. $\text{FreeStart}(\mathbf{A}) \subseteq \text{Crit}^+(\mathbf{A})$ forms an initial segment of $\text{Crit}^+(\mathbf{A})$.

If $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal and $m \in \text{Sp}(\mathbf{A})$, we write $\text{Sp}^{\mathbf{A}}(m)$ for the unique member of $\text{SpNode}(\mathbf{A}) \cap T(m)$. If $i < k^u$, we write $\text{Sp}_i^{\mathbf{A}}(\mathbf{A}) = \{m \in \text{Sp}(\mathbf{A}) : \text{Sp}^{\mathbf{A}}(m) \in T_i\}$.

Remark.

1. Notice that if $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal and $\mathbf{B} \subseteq \mathbf{A}$, then \mathbf{B} is also weakly diagonal. We have $\text{Sp}(\mathbf{B}) \subseteq \text{Sp}(\mathbf{A})$, but a level $m \in \text{AC}(\mathbf{B})$ could belong to any of \mathbf{A} , $\text{Sp}(\mathbf{A})$, or $\text{AC}(\mathbf{A})$.
2. If $m \in \mathbf{A}$, then since $m \notin \text{Sp}(\mathbf{A})$ and since $c[\mathbf{A}]$ is an antichain, we have $|c[\mathbf{A}]|_{m+1}| = |c[\mathbf{A}]|_m| - 1$. Similarly, if $m \in \text{Sp}(\mathbf{A})$, then as $m \notin \mathbf{A}$, we have $|c[\mathbf{A}]|_{m+1}| = |c[\mathbf{A}]|_m| + 1$. If $m \in \text{AC}(\mathbf{A})$, then $|c[\mathbf{A}]|_{m+1}| = |c[\mathbf{A}]|_m|$.

Our remaining goal for the subsection is to construct a weakly diagonal copy of \mathbf{K} inside \mathbf{K} .

Lemma 5.2.3. *Suppose $m < n \in \mathbf{K}$ have $U(m) = U(n)$, and let $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$. Then $\ell(c(f(m)) \wedge c(f(n))) \leq f(\ell(c(m) \wedge c(n)))$.*

Proof. If $c(m) \sqsubseteq c(n)$, then $\ell(c(m) \wedge c(n)) = m$, and certainly $\ell(c(f(m)) \wedge c(f(n))) \leq f(m)$. If $c(m) \not\sqsubseteq c(n)$, then $x := \ell(c(m) \wedge c(n)) < m$, and $c(m)(x) \neq c(n)(x)$, i.e., that $R(m, x) \neq R(n, x)$. As $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$, we have $R(f(m), f(x)) \neq R(f(n), f(x))$. So $\ell(c(f(m)) \wedge c(f(n))) \leq f(x) = f(\ell(c(m) \wedge c(n)))$. \square

Proposition 5.2.4. *There is $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ with $f[\mathbf{K}] \subseteq \mathbf{K}$ weakly diagonal.*

Proof. We are assuming that every $i < k^u$ is a non-degenerate unary predicate. This means that we can find $j_i < k^u$ and $q_i \in [1, k)$ so that the two element structure $\mathbf{J}_i = \{x, y\}$ with $U^{\mathbf{J}_i}(x) = j_i$, $U^{\mathbf{J}_i}(y) = i$, and $R^{\mathbf{J}_i}(y, x) = q_i$ is in \mathcal{K} . Fix such choices j_i and q_i .

We first create a countable \mathcal{L} -structure \mathbf{Y} which will contain \mathbf{K} as an induced substructure. For each $t \in \text{CT}(\geq 1)$, create a new vertex v_t . The underlying set of \mathbf{Y} will be $\omega \cup \{v_t : t \in \text{CT}(\geq 1)\} \cup \{u_i : i \in \text{Free}\}$. On ω , we simply reproduce \mathbf{K} . If $m < \omega$ and $t \in \text{CT}(m+1)$, there are two main cases regarding the new vertex v_t . If $t(m) \in [1, k)$, we set $U^{\mathbf{Y}}(v_t) = U(m)$, and if $n > m$ with $c(n) \supseteq t$, we set $R^{\mathbf{Y}}(n, v_t) = t(m)$. If $t(m) = 0$, then suppose that $U(m) = i$; we set $U^{\mathbf{Y}}(v_t) = j_i$, and we set $R(m, v_t) = q_i$. There are no other non-zero binary relations involving v_t . A helpful observation is that when $n > m$ and $c(n) \supseteq t$, the formula $R^{\mathbf{Y}}(n, v_t) = t(m)$ is valid in both cases. For the new vertices $\{u_i : i \in \text{Free}\}$, we set $U^{\mathbf{Y}}(u_i) = \text{Free}(i)^u$, and given $n \in \omega$ with $U(n) = i$, we set $R^{\mathbf{Y}}(n, u_i) = \text{Free}(i)^b$. There are no other non-zero binary relations involving u_i .

Claim. $\text{Age}(\mathbf{Y}) \subseteq \mathcal{K}$.

Proof of Claim. As there are no non-zero binary relations among the new vertices $\{v_t : t \in \text{CT}(\geq 1)\} \cup \{u_i : i \in \text{Free}\}$, it suffices to check that for any new vertex w , the induced substructure on $\mathbf{K} \cup \{w\}$ satisfies $\text{Age}(\mathbf{K} \cup \{w\}) \in \mathcal{K}$.

First, suppose $m < \omega$, $t \in T^*(m+1)$, and $w = v_t$. First assume $t(m) \neq 0$, and consider $S = \{n > m : c(n) \supseteq t\}$, viewed as an induced substructure of \mathbf{K} . Then $\mathbf{K} \cup \{v_t\}$ is the free amalgam of \mathbf{K} and $S \cup \{v_t\}$. Furthermore, we have $S \cup \{v_t\} \cong S \cup \{m\} \subseteq \mathbf{K}$. So $\text{Age}(\mathbf{Y}) \subseteq \mathcal{K}$ in this case. Now assume $t(m) = 0$, and suppose $U(m) = i$. Then $\mathbf{K} \cup \{v_t\}$ is the free amalgam of \mathbf{K} and $\{v_t, m\}$, and $\{v_t, m\} \cong \mathbf{J}_i \in \mathcal{K}$. So $\text{Age}(\mathbf{Y}) \subseteq \mathcal{K}$ in this case as well.

Now, suppose $i \in \text{Free}$ and $w = u_i$. Write $S = \{n < \omega : U(n) = i\}$. Then $\mathbf{K} \cup \{u_i\}$ is the free amalgam of \mathbf{K} and $S \cup \{u_i\}$, and we have $\text{Age}(\mathbf{K} \cup \{u_i\}) \subseteq \mathcal{K}$ by the definition of free relation. \blacksquare

We enumerate \mathbf{Y} via the function $g: \omega \rightarrow \mathbf{Y}$ which we now describe. First, g enumerates the set $\{u_i : i \in \text{Free}\}$. Then, given $m < \omega$, g first lists the set $\{v_t : t \in \text{CT}(m+1)\}$ in lex order, then lists m , doing this for each $m < \omega$ in order. With this enumeration, $\mathbf{Y} \cdot g$ is an enumerated structure (Notation 2.0.3) with $\text{Age}(\mathbf{Y} \cdot g) \subseteq \mathbf{K}$. Fix $h \in \text{OL}(\mathbf{Y} \cdot g, \mathbf{K})$, and write $f = h \circ g^{-1}$. So $f \in \text{Emb}(\mathbf{Y}, \mathbf{K})$, and restricting to \mathbf{K} yields the desired member of $\text{OEmb}(\mathbf{K}, \mathbf{K})$.

Claim. $f[\mathbf{K}] \subseteq \mathbf{K}$ is weakly diagonal.

Proof of claim. Towards showing items 1 and 2, fix $m < n < \omega$, and suppose $U(m) = U(n) := i$.

For item 1, suppose $c(n) \supseteq t \in \text{CT}(m+1)$. If $t(m) \neq 0$, then $R^{\mathbf{Y}}(n, v_t) = t(m)$ and $R^{\mathbf{Y}}(m, v_t) = 0$, and if $t(m) = 0$, then $R^{\mathbf{Y}}(m, v_t) = q_i \neq 0$ and $R^{\mathbf{Y}}(n, v_t) = 0$. Either way, since $f(v_t) < f(m) < f(n)$, we must have $\ell(c(f(m)) \wedge c(f(n))) \leq f(v_t)$. In particular, we cannot have $c(f(m)) \wedge c(f(n)) = c(f(m))$, so item 1 holds.

For item 2, suppose towards a contradiction that $\ell(c(f(m)) \wedge c(f(n))) = f(x)$ for some $x \in \mathbf{K}$. As item 1 holds, we have $x < m$. By Lemma 5.2.3, we must have

$x = \ell(c(m) \wedge c(n))$. In particular, at least one of $R(m, x)$ or $R(n, x)$ is non-zero; without loss of generality, suppose $R(n, x) \neq 0$. Setting $t = c(n)|_{x+1} \in \text{CT}(x+1)$, we have $R^{\mathbf{Y}}(n, v_t) = t(x) = R(n, x) \neq 0$ and $R^{\mathbf{Y}}(m, v_t) = 0$. This implies that $c(f(m))(f(v_t)) \neq c(f(n))(f(v_t))$, implying that $\ell(c(f(m)) \wedge c(f(n))) \leq f(v_t) < f(x)$. In particular, we cannot have $f(x) \in \text{Sp}(f[\mathbf{K}])$, so item 2 holds.

Let us also observe that $\ell(c(f(m)) \wedge c(f(n))) \neq f(u_j)$ for any $j \in \text{Free}$; to see this, we simply observe that $R^{\mathbf{Y}}(m, u_j) = R^{\mathbf{Y}}(n, u_j) = 0$ unless $i = j$, in which case $R^{\mathbf{Y}}(m, u_j) = R^{\mathbf{Y}}(n, u_j) = \text{Free}(i)^b$.

Once we know that every level in $\text{Sp}(f[\mathbf{K}])$ is of the form $f(v_t)$ for some $t \in \text{CT}$, the following argument is enough to show items 3 and 4. Suppose $m < n < \omega$, $t \in \text{CT}(x+1)$ for some $x \leq m$, and both $R^{\mathbf{Y}}(m, v_t)$ and $R^{\mathbf{Y}}(n, v_t)$ are non-zero. We will argue that

$$c(f(m))|_{f(v_t)+1} = c(f(n))|_{f(v_t)+1}.$$

First notice that we must have $x < m$; if $x = m$, then $R^{\mathbf{Y}}(m, v_t) \neq 0$ implies that $t(m) = 0$ and that $R^{\mathbf{Y}}(n, v_t) = 0$, a contradiction.

Now we must have $R^{\mathbf{Y}}(m, v_t) = t(x) = R^{\mathbf{Y}}(n, v_t)$; in particular, we have $c(m) \sqsupseteq t$ and $c(n) \sqsupseteq t$. Since f is left-leaning, it is enough to argue that $R(f(m), f(y)) = R(f(n), f(y))$ for any $y \in \mathbf{Y}$ with $f(y) \leq f(v_t)$. If $y \in \mathbf{K}$, then $y < x$, so since $c(m)|_{x+1} = c(n)|_{x+1}$, we are done. Now suppose $y = v_s$ for some $s \in \text{CT}(w+1)$ with $w \leq x$. Either we have $s \neq t|_{w+1}$, in which case $R(f(m), f(y)) = R(f(n), f(y)) = 0$, or we have $s = t|_{w+1}$, in which case we have $R(f(m), f(y)) = R(f(n), f(y)) = s(w)$.

Lastly, items 5 and 6 are straightforward. Item 5 holds since given $m < \omega$, then for any $t \in \text{CT}(m+1)$ with $t(m) = 0$, we have $R^{\mathbf{Y}}(m, v_t) = q_i \neq 0$ and $g(v_t) < g(m)$. Item 6 follows since $\text{FreeStart}(f[\mathbf{K}]) = \{f(u_i) : i \in \text{Free}\}$, $\text{Crit}^+(f[\mathbf{K}]) \subseteq f[\mathbf{Y}]$, and $\{f(u_i) : i \in \text{Free}\}$ is an initial segment of $f[\mathbf{Y}]$. Item 6 ■

□

5.3 Controlled paths

Definition 5.3.1. Given $i < k^u$, recall that we write $P_i = P(\iota_i)$ (Notation 4.6.2). A *path* in P_i is a linearly ordered subset of P_i . Write \vec{P}_i for the collection of paths in P_i . Given $t \in \text{CT}_i$, the *path* of t , denoted $\text{Path}(t)$, is the path $\{\mathcal{K}(t|_m) : m \leq \ell(t)\} \in \vec{P}_i$. If $a \in \mathbf{K}$, we sometimes write $\text{Path}(a) = \text{Path}(c(a))$.

For every $t \in \text{CT}_i$, $\text{Path}(t)$ is a linearly ordered subset of the poset P_i with maximum element $\mathcal{K}(\emptyset_i)$ and minimum element $\mathcal{K}(t)$. We will be particularly interested in the case where these subsets are as large as possible.

Definition 5.3.2. Given $i < k^u$, a *maximal path* in P_i is simply a path which is maximal under inclusion. Write $\max(\vec{P}_i)$ for the collection of maximal paths in P_i . We equip $\bigcup_{i < k^u} \max(\vec{P}_i)$ with an arbitrary linear order \leq_{path} with the property that whenever $i < j$, $\mathfrak{p} \in \max(\vec{P}_i)$, and $\mathfrak{q} \in \max(\vec{P}_j)$, then $\mathfrak{p} \leq_{\text{path}} \mathfrak{q}$. We say that a substructure $\mathbf{A} \subseteq \mathbf{K}$ has *maximal paths* if for every $a \in \mathbf{A}$, we have $\text{Path}(a) \in \max(\vec{P}_i)$.

Example 5.3.3. Let $\mathcal{K} = \mathcal{G}_3^{r,b}$ denote the class of finite graphs, where all edges are either red or blue, and where we forbid edge-monochromatic triangles. So $k^u = 1$ and $k = 3$, where 1 and 2 represent blue and red edges, respectively. Then $P(1)$ has four elements: $\mathcal{G}_3^{r,b}$, the class \mathcal{G}_3^r of finite, triangle-free graphs with all red edges, the class \mathcal{G}_3^b , defined analogously with all blue edges, and the class \mathcal{G}_2 of finite graphs with no edges. The set $\max(\vec{P}_0)$ has 2 elements, the path $\{\mathcal{G}_2, \mathcal{G}_3^r, \mathcal{G}_3^{r,b}\}$ and the path $\{\mathcal{G}_2, \mathcal{G}_3^b, \mathcal{G}_3^{r,b}\}$.

As we mentioned at the end of Subsection 5.1, we won't yet worry about building $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ where $f[\mathbf{K}] \subseteq \mathbf{K}$ has maximal paths. We will do this in the next few subsections. For now, we are interested in starting with a substructure $\mathbf{A} \subseteq \mathbf{K}$ with maximal paths and building $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ so that $f[\mathbf{A}] \subseteq \mathbf{K}$ has *controlled paths* (Definition 5.3.4). The definition will depend on which $i < k^u$ admit a free relation (Definition 4.6.3).

Definition 5.3.4. Fix a weakly diagonal substructure $\mathbf{A} \subseteq \mathbf{K}$. We say that $\mathbf{A} \subseteq \mathbf{K}$ has *controlled paths* if \mathbf{A} has maximal paths and the following all hold.

1. Suppose $i \in k^u \setminus \text{Free}$. Then given $a, b \in \mathbf{A}$ with $U(a) = U(b) = i$ and $\text{Start}(a) = \text{Start}(b)$, we demand that $\text{Path}(a) = \text{Path}(b)$.
2. Suppose $i \in \text{Free}$. Then given $a, b \in \mathbf{A}$ with $U(a) = U(b) = i$, we demand that $\text{Start}(a) \leq \text{Start}(b)$ iff $\text{Path}(a) \leq_{\text{path}} \text{Path}(b)$, that $U(\text{Start}(a)) = \text{Free}(i)^u$, that $c(a)(\text{Start}(a)) = \text{Free}(i)^b$, and if $\text{Path}(a) \neq \text{Path}(b)$, that $c(b)(\text{Start}(a)) = 0$.
3. Suppose $i < j \in \text{Free}$. Then for any $a, b \in \mathbf{A}$ with $U(a) = i$ and $U(b) = j$, we have $\text{Start}(a) < \text{Start}(b)$.

We note that if $\mathbf{A} \subseteq \mathbf{K}$ has controlled paths and $\mathbf{B} \subseteq \mathbf{A}$, then $\mathbf{B} \subseteq \mathbf{K}$ also has controlled paths.

A useful intuition is as follows. Since \mathcal{K} is a free amalgamation class, the left branch of each T_i , i.e., the set of nodes of the form $(i, 0^m)$, has special importance. Therefore, it is interesting to keep track of when a node in T_i first leaves the left branch. When a substructure $\mathbf{A} \subseteq \mathbf{K}$ has controlled paths, this means that given $a \in \mathbf{A}$ with $U(a) = i < k^u$, the moment the coding node $c(a)$ leaves the left branch, then $\text{Path}(a)$ is completely determined. When $i \in \text{Free}$, the free relation gives us even more control; we can consider coding nodes with the same path and introduce a new point related to all of them, ensuring that they all depart the left branch at the same time. Regarding items 2 and 3, the intuition is that all of the different possible maximal paths that can appear among the free unaries are separated in the tree CT in some arbitrary, but fixed fashion, and this occurs before any other critical events.

Proposition 5.3.5. *Suppose $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal and has maximal paths. Then there is $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ so that $f[\mathbf{A}] \subseteq \mathbf{K}$ is weakly diagonal and has controlled paths.*

Proof. The proof will have a similar outline to that of Proposition 5.2.4. We first build an \mathcal{L} -structure $\mathbf{Y} \supseteq \mathbf{K}$ with $\text{Age}(\mathbf{Y}) \subseteq \mathcal{K}$. Then we will build $f \in \text{Emb}(\mathbf{Y}, \mathbf{K})$ so that upon restricting to \mathbf{K} , we will have $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ as desired.

First, if $i \in \text{Free}$, write $\text{Free}(i) = (q, j)$. For each $\mathfrak{p} \in \max(\vec{P}_i)$, introduce a new vertex $u_{\mathfrak{p}}$. We set $U^{\mathbf{Y}}(u_{\mathfrak{p}}) = j$, and given $a \in \mathbf{A}$ with $U(a) = i$ and $\text{Path}(a) = \mathfrak{p}$, we set $R^{\mathbf{Y}}(a, u_{\mathfrak{p}}) = q$. The vertex $u_{\mathfrak{p}}$ participates in no other non-zero binary relations.

Now given $m < \omega$ and $i < k^u$, let us call (m, i) an *important pair* if we have at least one of the following:

1. $m \in \text{Sp}_i(\mathbf{A})$, in which case we write $s = \text{Sp}^{\mathbf{A}}(m)$,
2. $i \in k^u \setminus \text{Free}$, $s := (i, 0^m) \in c[\mathbf{A}]|_m$, and for some $q \in [1, k)$ we have $s \frown q \in c[\mathbf{A}]|_{m+1}$.

We remark that for a given $m < \omega$, there might be multiple $i < k^u$ with (m, i) an important pair; if so, then this must be due to item (2) for each such $i < k^u$. We also remark that (m, i) might satisfy both items above; if so, this is because $\text{Sp}^{\mathbf{A}}(m) = (i, 0^m)$.

For each important pair (m, i) and each $\mathfrak{p} \in \max(\vec{P}_i)$, introduce a new vertex $v_{m, \mathfrak{p}}$. We set $U^{\mathbf{Y}}(v_{m, \mathfrak{p}}) = U(m)$, and given $a \in \mathbf{A}$ with $c(a) \sqsupseteq s$ and $\text{Path}(a) = \mathfrak{p}$, we set $R^{\mathbf{Y}}(a, v_{m, \mathfrak{p}}) = R(a, m)$. The vertex $v_{m, \mathfrak{p}}$ participates in no other non-zero binary relations, and this concludes the construction of \mathbf{Y} .

To see that $\text{Age}(\mathbf{Y}) \subseteq \mathcal{K}$, the new vertices $u_{\mathfrak{p}}$ which we attach when $i < k^u$ has a free relation pose no problem precisely because of the free relation. Now suppose (m, i) is an important pair and that $\mathfrak{p} \in \max(\vec{P}_i)$. Suppose that $s \in c[\mathbf{A}]|_m$ is as in item (1) and/or (2) above, and suppose $\mathbf{B} \subseteq \mathbf{A}$ is a finite substructure with $c(b)|_m = s$ and $\text{Path}(b) = \mathfrak{p}$ for every $b \in \mathbf{B}$. Then the substructure of \mathbf{Y} induced on $\mathbf{B} \cup \{v_{m, \mathfrak{p}}\}$ is isomorphic to the substructure of \mathbf{K} induced on $\mathbf{B} \cup \{m\}$.

Let $g: \omega \rightarrow \mathbf{Y}$ be the following enumeration. First, g lists the new vertices $u_{\mathfrak{p}}$ in \leq_{path} -order for each $\max(\vec{P}_i)$, handling those $i < k^u$ with free relations in increasing order. Then g lists elements of \mathbf{K} in order, but with $g^{-1}(m-1) < g^{-1}(v_{m, \mathfrak{p}}) < g^{-1}(m)$ for every $m < \omega$ and new vertex $v_{m, \mathfrak{p}}$. For a fixed $m < \omega$, we do not care how g enumerates the new vertices of the form $v_{m, \mathfrak{p}}$, so long as these all appear between $g^{-1}(m-1)$ and $g^{-1}(m)$. We now pick any $h \in \text{OL}(\mathbf{Y} \cdot g, \mathbf{K})$ and define $f \in \text{Emb}(\mathbf{Y}, \mathbf{K})$ via $f = h \circ g^{-1}$.

Claim. $f[\mathbf{A}]$ is weakly diagonal.

Proof of claim. Items 1 and 2 of Definition 5.2.2 hold for any $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$. To check items 3 and 4, first notice that any $n \in \text{Sp}(f[\mathbf{A}])$ is of the form $f(u_{\mathfrak{p}})$ or $f(v_{m, \mathfrak{p}})$. Now suppose $a, b \in \mathbf{A}$ and $n \in \text{Sp}(f[\mathbf{A}])$ are such that both $R(f(a), n)$ and $R(f(b), n)$ are non-zero. We want to show that $c(f(a))|_{n+1} = c(f(b))|_{n+1}$. If $n = f(u_{\mathfrak{p}})$ for some new vertex of the form $u_{\mathfrak{p}}$, then this is clear. So suppose $n = f(v_{m, \mathfrak{p}})$, where $i < k^u$ is such that (m, i) is an important pair and with $\mathfrak{p} \in \max(\vec{P}_i)$. Then let $s \in c[\mathbf{A}]|_m$ be as in the definition of important pair. So we must have $\text{Path}(a) = \text{Path}(b) = \mathfrak{p}$, $c(a)|_m = c(b)|_m = s$, and $R(a, m) = R(b, m)$ both equal to the unique $q \in [1, k)$ with $s \frown q \in c[\mathbf{A}]|_{m+1}$. In particular, we have $c(a)|_{m+1} = c(b)|_{m+1}$. This implies that $R(f(a), f(\ell)) = R(f(b), f(\ell))$ for every $\ell \leq m$. And if $r \leq n$ with $r \in f[\mathbf{Y} \setminus \mathbf{K}]$ and either $R(f(a), r)$ or $R(f(b), r)$ are non-zero, then we must have $r = f(u_{\mathfrak{p}})$ (if $i < k^u$ has a free relation) or $r = f(v_{\ell, \mathfrak{p}})$ for some $\ell < m$. In either case, we have $R(f(a), r) = R(f(b), r)$. Hence $c(f(a))|_{n+1} = c(f(b))|_{n+1}$ as desired.

To check items 5 and 6, item 5 holds for $f[\mathbf{A}] \subseteq \mathbf{K}$ since it holds for $\mathbf{A} \subseteq \mathbf{K}$ and since $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$. Item 6 holds since $\text{FreeStart}(f[\mathbf{A}]) \subseteq \{f(u_{\mathfrak{p}}) : \mathfrak{p} \in \max(\vec{P}_i), i \in \text{Free}\}$, $\text{Crit}^+(f[\mathbf{A}]) \subseteq f[\mathbf{Y}]$, and $\{f(u_{\mathfrak{p}}) : \mathfrak{p} \in \max(\vec{P}_i), i \in \text{Free}\}$ is an initial segment of $f[\mathbf{Y}]$. \blacksquare

Claim. For every $a \in \mathbf{A}$, we have $\text{Path}(a) = \text{Path}(f(a))$.

Proof of claim. Fix $a \in \mathbf{A}$. It is enough to show that for every $m \leq a$, we have $\mathcal{K}(c(a)|_m) = \mathcal{K}(c(f(a))|_{f(m)})$. The right-to-left inclusion is clear. For the left-to-right inclusion, fix $\mathbf{B} \in \mathcal{K}(c(a)|_m)$ with underlying set B , and form $\mathbf{B}[c(f(a))|_{f(m)}]$. Upon deleting vertices from $\mathbf{K}_{f(m)} \setminus f[\mathbf{Y}]$ (which have no non-zero relations to members of \mathbf{B}), we obtain a structure which is the free amalgam of the substructures induced on $B \cup f[\{0, \dots, m-1\}] \cup \{x\}$, where x ranges over the new vertices of the form $u_{\mathfrak{p}}$ or $v_{\ell, \mathfrak{p}}$ for some $\ell \leq m$. So it is enough to show that each of these is in \mathcal{K} ; write \mathbf{C}_x for this induced substructure. First we note that $\mathbf{C}_x \setminus \{x\}$ is isomorphic to $\mathbf{B}[c(a)|_m]$, which is in \mathcal{K} by assumption. If x is of the form $u_{\mathfrak{p}}$, then $\mathbf{C} \in \mathcal{K}$ by the properties of a free relation. If $x = v_{\ell, \mathfrak{p}}$, then compare \mathbf{C} to the structure \mathbf{X} obtained by taking $\mathbf{B}[c(a)|_m]$ and duplicating ℓ with a new vertex y_{ℓ} , setting $R^{\mathbf{X}}(\ell, y_{\ell}) = 0$. Then $\mathbf{X} \in \mathcal{K}$, and \mathbf{C} is isomorphic to a structure obtained from \mathbf{X} by possibly deleting some non-zero binary relations. \blacksquare

Once we have that $\text{Path}(a) = \text{Path}(f(a))$ for each $a \in \mathbf{A}$, the rest of the argument that $f[\mathbf{A}] \subseteq \mathbf{K}$ has controlled paths is straightforward. Fix $a \in \mathbf{A}$ with $U(a) = i < k^u$. If i has a free relation, then $\text{Start}(c(f(a))) = f(u_{\text{Path}(a)})$. If i does not have a free relation, then $\text{Start}(c(f(a))) = f(v_{\text{Start}(a), \text{Path}(a)})$; we quickly remark that since \mathbf{A} is assumed to have maximal paths and by Lemma 4.6.8, we have that $\text{Start}(a)$ exists. Knowing this, the three items of Definition 5.3.4 are easy to verify. \square

5.4 Insertions

This subsection introduces the main technical tool we will use in our construction of “controlled levels.” We will start with a weakly diagonal $\mathbf{A} \subseteq \mathbf{K}$ and discuss how to modify elements of $\text{Crit}(\mathbf{A})$ one at a time. Particular details depend on whether the level we are modifying belongs to \mathbf{A} , $\text{Sp}(\mathbf{A})$, or $\text{AC}(\mathbf{A})$, and will be discussed in more detail in future subsections. For the remainder of the subsection, fix a weakly diagonal $\mathbf{A} \subseteq \mathbf{K}$.

Notation 5.4.1. Suppose we have $\mathbf{B} \subseteq \mathbf{K}$ and $M < \omega$ with $\min\{\mathbf{B}\} \geq M$. Suppose $S = \{s_0 \preceq_{lex} \dots \preceq_{lex} s_{d-1}\} \subseteq T(M)$ is such that $c[\mathbf{B}]|_M \subseteq S$. For each $b \in \mathbf{B}$, introduce a new vertex x_b . Then \mathbf{B}/S , said “ \mathbf{B} over S ,” is the \mathcal{L}_d -structure on underlying set $\{x_b : b \in \mathbf{B}\}$, with $R^{\mathbf{B}/S}(x_a, x_b) = R^{\mathbf{B}}(a, b)$ for any $a, b \in \mathbf{B}$, and such that for $b \in \mathbf{B}$, we have $V^{\mathbf{B}/S}(x_b) = j$ iff $c(b)|_M = s_j$.

The only reason we introduce the new vertices $\{x_b : b \in \mathbf{B}\}$ is to keep with our convention that the underlying sets of \mathcal{L}_d -structures should be disjoint from ω .

Definition 5.4.2. Let $\mathbf{A} \subseteq \mathbf{K}$ be weakly diagonal, and fix $m \leq M < \omega$. An *insertion for \mathbf{A} with parameters $m \leq M$* is a map $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ constructed using the following procedure:

1. Writing $S = c[\mathbf{A}]|_M$ and $\rho = S^u$, first choose a gluing $\gamma = (\rho, \eta, \mathbf{I})$ so that for every finite $\mathbf{B} \subseteq \mathbf{A}$ with $\min\{\mathbf{B}\} \geq M$, we have $(\mathbf{B}/S)[\gamma] \in \mathcal{K}$.
2. Create new vertices $\{v_x : x \in \mathbf{I}\}$. Form the \mathcal{L} -structure \mathbf{Y} on $\mathbf{K} \cup \{v_x : x \in \mathbf{I}\}$ with \mathbf{K} induced and with $\{v_x : x \in \mathbf{I}\} \cong \mathbf{I}$ in the obvious way. For the binary relations between \mathbf{K} and $\{v_x : x \in \mathbf{I}\}$, suppose $a \in \mathbf{A}$ with $a \geq M$ and $x \in \mathbf{I}$. Suppose $c[\mathbf{A}]|_M = \{s_0 \preceq_{lex} \cdots \preceq_{lex} s_{d-1}\}$ and that $c(a)|_M = s_j$. Then we set $R^{\mathbf{Y}}(a, v_x) = \eta(j, x)$. There are no other new non-zero binary relations.
3. Note that by item 1, we have $\text{Age}(\mathbf{Y}) = \mathcal{K}$. Let $g: \omega \rightarrow \mathbf{Y}$ be the enumeration of \mathbf{Y} which enumerates $\{0, \dots, m-1\}$ in order, then $\{v_x : x \in \mathbf{I}\}$ in order, then $\omega \setminus \{0, \dots, m-1\}$ in order. Form the enumerated structure $\mathbf{Y} \cdot g$ (Notation 2.0.3). Then using Lemma 3.0.1, choose $h \in \text{OL}(\mathbf{Y} \cdot g, \mathbf{K})$ with h the identity on $\{0, \dots, m-1\}$.
4. We now define an embedding $f: \mathbf{Y} \rightarrow \mathbf{K}$ via $f = h \circ g^{-1}$. Restricting to $\mathbf{K} \subseteq \mathbf{Y}$, we have $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$. Note that f is also the identity on $\{0, \dots, m-1\}$.

Remark. In practice, we will have $m \in \text{Crit}(\mathbf{A})$. We will set $M = m$ when $m \in \mathbf{A} \cup \text{AC}(\mathbf{A})$, and we will set $M = m + 1$ when $m \in \text{Sp}(\mathbf{A})$.

We have seen the idea of putting \mathbf{K} inside a superstructure \mathbf{Y} before, in the proofs of Propositions 5.2.4 and 5.3.5. What makes insertions special is that \mathbf{Y} has a very nice construction. Intuitively, an insertion builds \mathbf{Y} by “inserting” a copy of the structure \mathbf{I} just below vertex m using the instructions given by the gluing. The fact that we only glue to vertices from \mathbf{A} above M , and not every vertex from \mathbf{K} above M , gives us some flexibility to change the age of a given level.

Lemma 5.4.3. *Suppose $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ is an insertion for \mathbf{A} with parameters $m \leq M$. Suppose $a, b \in \mathbf{A}$ have $\ell(c(a) \wedge c(b)) < m$ or $\ell(c(a) \wedge c(b)) \geq M$. Then*

$$\ell(c(f(a)) \wedge c(f(b))) = f(\ell(c(a) \wedge c(b))).$$

In particular, if $m = M$, we have

$$\text{Sp}(f[\mathbf{A}]) = f[\text{Sp}(\mathbf{A})].$$

Remark. Lemma 5.2.3 yields $\ell(c(f(a)) \wedge c(f(b))) \leq f(\ell(c(a) \wedge c(b)))$, so the content of this lemma is the other inequality.

Proof. Given $L < m$, then for $\alpha \in \{a, b\}$, we have $R(\alpha, L) = R(f(\alpha), f(L)) = R(f(\alpha), L)$. The lemma in the case that $\ell(c(a) \wedge c(b)) < m$ follows. Now suppose that $\ell(c(a) \wedge c(b)) \geq M$. We need to show for any $p < f(\ell(c(a) \wedge c(b)))$ that $R(f(a), p) = R(f(b), p)$. If $p \notin f[\mathbf{Y}]$, then $R(f(a), p) = R(f(b), p) = 0$. If $p \in f[\mathbf{K}]$, then $p = f(n)$ for some $n < \ell(c(a) \wedge c(b))$, so we have $R(f(a), p) = R(a, n) = R(b, n) = R(f(b), p)$. And if $p = f(v_x)$ for some $x \in \mathbf{I}$, write $c[\mathbf{A}]|_M = \{s_0 \preceq_{lex} \cdots \preceq_{lex} s_{d-1}\}$, and suppose that $c(a)|_M = c(b)|_M = s_j$. Then $R(f(a), p) = R^{\mathbf{Y}}(a, v_x) = \eta(j, x) = R^{\mathbf{Y}}(b, v_x) = R(f(b), p)$.

For the “in particular,” we have just shown that $f[\text{Sp}(\mathbf{A}) \setminus (m, M)] \subseteq \text{Sp}(f[\mathbf{A}])$, and since $m = M$, we have $(m, M) = \emptyset$, giving the right-to-left inclusion. For

the left-to-right inclusion, suppose $r \in \text{Sp}(f[\mathbf{A}])$. Find $a, b \in \mathbf{A}$ with $\ell(c(f(a)) \wedge c(f(b))) = r$. But since $m = M$, we can apply the first part of the lemma to conclude that $r = f(\ell(c(a) \wedge c(b)))$, i.e., that $r \in f[\text{Sp}(\mathbf{A})]$. \square

Notation 5.4.4. Given $\mathbf{A} \subseteq \mathbf{K}$, we define the *body* of \mathbf{A} to be $\text{Body}(\mathbf{A}) = \bigcup_n c[\mathbf{A}]|_n$. So $\text{Body}(\mathbf{A})$ is just all nodes in CT which extend to a coding node from \mathbf{A} . As an example, we have $\text{Body}(\mathbf{K}) = \text{CT}$.

Definition 5.4.5. Suppose $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ is an insertion for \mathbf{A} with parameters $m \leq M$. We define the map

$$\hat{f}: \text{Body}(\mathbf{A}) \setminus \left(\bigcup_{n \in (m, M)} T(n) \right) \rightarrow \text{CT}.$$

where given $t \in \text{Body}(\mathbf{A})$ with $\ell(t) \notin (m, M)$, we choose some $a \in \mathbf{A}$ with $c(a) \sqsupseteq t$, and we set $\hat{f}(t) = c(f(a))|_{f(\ell(t))}$. By Lemma 5.4.3, this does not depend on the choice of $a \in \mathbf{A}$. Let us record some basic properties of the map \hat{f} :

- \hat{f} is the identity on $\text{Body}(\mathbf{A}) \cap T(< m)$.
- \hat{f} respects tree structure and relative levels.
- \hat{f} respects passing numbers; for any $t \in \text{Body}(\mathbf{A})$ with $\ell(t) \notin (m, M)$ and any $n < \ell(t)$, we have $t(n) = \hat{f}(t)(f(n))$.

Proposition 5.4.6. *Suppose $f: \text{OEmb}(\mathbf{K}, \mathbf{K})$ is an insertion for \mathbf{A} with equal parameters $m = M$ such that $m > \max(\text{FreeStart}(\mathbf{A}))$. Then $f[\mathbf{A}] \subseteq \mathbf{K}$ is weakly diagonal.*

Proof. This is immediate upon considering the map \hat{f} ; since $m = M$, the domain of \hat{f} is all of $\text{Body}(\mathbf{A})$. Since \hat{f} preserves tree structure, relative levels, and passing numbers, we see that the first five items of Definition 5.2.2 are satisfied for $f[\mathbf{A}]$. Then, since $\text{FreeStart}(f[\mathbf{A}]) = \text{FreeStart}(\mathbf{A}) \subseteq m$, item 6 is also satisfied. \square

While \hat{f} has many desirable properties, note that we may not have $\mathcal{K}(\hat{f}[c[\mathbf{A}]|_n]) = \mathcal{K}(c[\mathbf{A}]|_n)$. Indeed, the purpose of an insertion is often to change the age of a level set in $\text{Body}(\mathbf{A})$. However, there are several situations where this does hold.

Proposition 5.4.7. *Suppose $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ is an insertion for \mathbf{A} with parameters $m \leq M$ constructed with gluing $\gamma = (\rho, \eta, \mathbf{I})$.*

1. For any $n < m$, we have $\mathcal{K}(\hat{f}[c[\mathbf{A}]|_n]) = \mathcal{K}(c[\mathbf{A}]|_n)$.
2. For any $n < \omega$, we have $\mathcal{K}(\hat{f}[c[\mathbf{A}]|_n]) \subseteq \mathcal{K}(c[\mathbf{A}]|_n)$.
3. Suppose there is $r \geq M$ so that the following hold:
 - $\mathbf{A} \cap [M, r) = \emptyset$
 - Write $c[\mathbf{A}]|_r = \{t_0 \preceq_{\text{lex}} \cdots \preceq_{\text{lex}} t_{d_0-1}\}$, $c[\mathbf{A}]|_M = \{s_0 \preceq_{\text{lex}} \cdots \preceq_{\text{lex}} s_{d-1}\}$, and let $e: d_0 \rightarrow d$ be the function with $t_j|_M = s_{e(j)}$. Then $\mathcal{K}(c[\mathbf{A}]|_r) \subseteq \mathcal{K}(\gamma^e)$.

Then for any $n \geq r$, we have $\mathcal{K}(c[\mathbf{A}]|_n) = \mathcal{K}(\hat{f}[c[\mathbf{A}]|_n])$.

Remark. Keep in mind that by definition, $\hat{f}[c[\mathbf{A}]|_n] = c[f[\mathbf{A}]]|_{f(n)}$.

Proof. Item 1 is clear since \hat{f} is the identity on $\text{Body}(\mathbf{A}) \cap T(< m)$.

For item 2, if $\mathbf{B} \in \mathcal{K}(\hat{f}[c[\mathbf{A}]|_n])$, then $\mathbf{B}[\hat{f}[c[\mathbf{A}]|_n]] \in \mathcal{K}$, and $\mathbf{B}[c[\mathbf{A}]|_n]$ is isomorphic to the substructure of $\mathbf{B}[\hat{f}[c[\mathbf{A}]|_n]]$ induced on $\mathbf{B} \cup f[\{0, \dots, n-1\}]$.

For item 3, fix r and n as in the item statement, write $c[\mathbf{A}]|_n = \{u_0, \dots, u_{d_1-1}\}$, and let \mathbf{B} be an \mathcal{L}_{d_1} -structure with underlying set B and with $\mathbf{B} \in \mathcal{K}(c[\mathbf{A}]|_n)$. To show that $\mathbf{B} \in \mathcal{K}(\hat{f}[c[\mathbf{A}]|_n])$, the idea is to have \mathbf{B} “steal” the vertices $\mathbf{A} \cap [r, n-1]$, view the result as an \mathcal{L}_{d_0} -structure, and then use the assumption that $\mathcal{K}(c[\mathbf{A}]|_r) \subseteq \mathcal{K}(\gamma^e)$. More precisely, create new vertices $X := \{x_\alpha : \alpha \in \mathbf{A} \cap [r, n-1]\}$, and form the \mathcal{L}_{d_0} -structure \mathbf{C} on $B \cup X$ as follows:

- Given $b \in B$, we have $V^{\mathbf{C}}(b) = j$ iff $u_{V^{\mathbf{B}}(b)}|_r = t_j$.
- Given $\alpha \in \mathbf{A} \cap [r, n-1]$, we have $V^{\mathbf{C}}(x_\alpha) = j$ iff $c(\alpha)|_r = t_j$.
- The binary relations on B are those from \mathbf{B} , and given $x_\alpha, x_\beta \in X$, we have $R^{\mathbf{C}}(x_\alpha, x_\beta) = R(\alpha, \beta)$.
- Given $\alpha \in \mathbf{A} \cap [r, n-1]$ and $b \in B$, we have $R^{\mathbf{C}}(b, x_\alpha) = R^{\mathbf{B}[c[\mathbf{A}]|_n]}(b, \alpha) = u_{V^{\mathbf{B}}(b)}(\alpha)$.

By design, we have $\mathbf{C}[c[\mathbf{A}]|_r] \cong \mathbf{B}[c[\mathbf{A}]|_n] \setminus ([r, n-1] \setminus \mathbf{A})$, so $\mathbf{C} \in \mathcal{K}(c[\mathbf{A}]|_r) \subseteq \mathcal{K}(\gamma^e)$. Now consider $\mathbf{B}[\hat{f}[c[\mathbf{A}]|_n]]$; this is the free amalgam of the induced substructures on $\mathbf{K}_{f(n)}$ and on $B \cup f[\{0, \dots, n-1\}] \cup f[\{v_x : x \in \mathbf{I}\}]$. The latter is the free amalgam of the induced substructures on $B \cup f[\{0, \dots, n-1\}]$ (isomorphic to $\mathbf{B}[c[\mathbf{A}]|_n]$) and on $B \cup f[\{v_x : x \in \mathbf{I}\}] \cup f[\{0, \dots, r-1\}] \cup f[\mathbf{A} \cap [r, n-1]]$. The latter of these is the free amalgam of the substructures on $B \cup f[\mathbf{A} \cap [r, n-1]] \cup f[\{v_x : x \in \mathbf{I}\}]$ (isomorphic to $\mathbf{C}[\gamma^e]$) and $B \cup f[\mathbf{A} \cap [r, n-1]] \cup f[\{0, \dots, r-1\}]$ (isomorphic to $\mathbf{C}[c[\mathbf{A}]|_r]$). Hence $\mathbf{B} \in \mathcal{K}(\hat{f}[c[\mathbf{A}]|_n])$ as desired. \square

Remark. One can pull off the proof without the assumption that $\mathcal{K}(c[\mathbf{A}]|_r) \subseteq \mathcal{K}(\gamma^e)$; all that we needed was that $\mathbf{C} \in \mathcal{K}(\gamma^e)$. We will use this idea to prove the “furthermore” of Proposition 5.6.9.

Corollary 5.4.8. *Suppose $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal and has maximal paths. Let $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ be an insertion for \mathbf{A} with parameters m and M , where $M \leq m+1$. Suppose for any $n \geq m+1$ that we have $\mathcal{K}(c[\mathbf{A}]|_n) = \mathcal{K}(\hat{f}[c[\mathbf{A}]|_n])$. Then for any $a \in \mathbf{A}$, we have $\text{Path}(a) = \text{Path}(f(a))$.*

Proof. Whenever $\ell \leq m$, we have $c[\mathbf{A}]|_\ell = c[f[\mathbf{A}]]|_\ell = \hat{f}[c[\mathbf{A}]]|_\ell$ since f is the identity on $\{0, \dots, m-1\}$. It follows that for every $a \in \mathbf{A}$, we have $\text{Path}(a) \subseteq \text{Path}(f(a))$. The reverse inclusion follows because $\text{Path}(a)$ is maximal. \square

5.5 Controlled age-change levels

The main intuition behind the notion of a *controlled age-change level* is that any age change should take place as gradually as possible (Item (1) of Definition 5.5.1). However, the definition is a bit more complicated; the reason for this is that given

a substructure $\mathbf{A} \subseteq \mathbf{K}$ with a controlled age-change level $m \in \text{AC}(\mathbf{A})$, we want $\overline{\mathbf{A}}$ “near m ” to be entirely determined by the age change. We will make this idea precise in subsection 6.1.

Definition 5.5.1. Suppose $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal, and let $m \in \text{AC}(\mathbf{A})$. We call m a *controlled age change level* of \mathbf{A} if the following all hold:

1. Write $\rho = (c[\mathbf{A}]|_m)^u = (c[\mathbf{A}]|_{m+1})^u$ (Notation 4.1.4). Write $\mathcal{A} = \mathcal{K}(c[\mathbf{A}]|_m)$ and $\mathcal{B} = \mathcal{K}(c[\mathbf{A}]|_{m+1})$. Then we demand that $(\mathcal{A}, \mathcal{B}) \in \text{Con}(\rho)$.
2. Write $\gamma = (\rho, \eta, \mathbf{I}) = \text{Cause}(\mathcal{A}, \mathcal{B})$ (Definition 4.5.4). Then we demand that there is $\varphi \in \text{OL}(\mathbf{I}, \mathbf{K})$ with $\max\{\varphi[\mathbf{I}]\} = m$, and writing $c[\mathbf{A}]|_{m+1} = \{t_0, \dots, t_{d-1}\}$ we demand for each $j < d$ and $y \in [\min\{\varphi[\mathbf{I}]\}, \max\{\varphi[\mathbf{I}]\}]$ that

$$t_j(y) = \begin{cases} \eta(j, \varphi^{-1}(y)) & \text{if } y \in \varphi[\mathbf{I}] \\ 0 & \text{if } y \notin \varphi[\mathbf{I}]. \end{cases}$$

Recall that in Definition 4.5.1, \mathbf{I} is chosen with minimum possible cardinality. From this and the above formula for $t_j(y)$, we see that such a $\varphi \in \text{OL}(\mathbf{I}, \mathbf{K})$ is unique.

3. Write $i = \min(\varphi[\mathbf{I}])$. Then we demand that $\text{Crit}^+(\mathbf{A}) \cap [i, m] = \{m\}$.

We say that \mathbf{A} has *controlled age change levels* if every $m \in \text{AC}(\mathbf{A})$ is a controlled age change level.

Remark. In the setting of item 3, we can write $c[\mathbf{A}]|_i = \{s_0, \dots, s_{d-1}\}$ with $s_j \sqsubseteq t_j$ for each $j < d$, and we have $\mathcal{K}(c[\mathbf{A}]|_i) = \mathcal{K}(c[\mathbf{A}]|_m)$.

The most important part of Definition 5.5.1 is item 1. Items 2 and 3 are used to control the closure of $\mathbf{A} \subseteq \mathbf{K}$ near a controlled age-change level, but in a sense, the purpose of having this extra control is to allow us to forget the technicalities; upon defining the notion of a *diagonal diary*, which is an abstraction of the information contained in a diagonal substructure, the only part of a controlled age-change that we will record is the nature of the age-change taking place.

Lemma 5.5.2. *Suppose $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal, and let $m \in \text{AC}(\mathbf{A})$ be a controlled age change level. Suppose $\mathbf{B} \subseteq \mathbf{A}$. Then either $m \notin \text{AC}(\mathbf{B})$ or m is a controlled age change level of \mathbf{B} .*

Proof. We adopt all notation from Definition 5.5.1. Write $|c[\mathbf{B}]|_m| = |c[\mathbf{B}]|_{m+1}| = d_1$, and let $e_1: d_1 \rightarrow d$ be the increasing injection so that $c[\mathbf{B}]|_{m+1} = \{t_{e_1(j)} : j < d_1\}$. So in particular, we have $\mathcal{K}(c[\mathbf{B}]|_m) = \mathcal{A}^{e_1}$ and $\mathcal{K}(c[\mathbf{B}]|_{m+1}) = \mathcal{B}^{e_1}$. By Proposition 4.5.3, let $e: d_0 \rightarrow d$ be the unique increasing injection with $(\mathcal{A}^e, \mathcal{B}^e) \in \text{ECon}(\rho \circ e)$. If $\text{Im}(e) \not\subseteq \text{Im}(e_1)$, then Proposition 4.5.2 tells us that $\mathcal{A}^{e_1} = \mathcal{B}^{e_1}$, so in this case $m \notin \text{AC}(\mathbf{B})$. Now assume $\text{Im}(e) \subseteq \text{Im}(e_1)$; by Corollary 4.5.3, we have $(\mathcal{A}^{e_1}, \mathcal{B}^{e_1}) \in \text{Con}(\rho \circ e_1)$. Letting $e_0: d_0 \rightarrow d_1$ be the increasing injection with $e_1 \circ e_0 = e$, we have that e_0 is the unique increasing injection with $((\mathcal{A}^{e_1})^{e_0}, (\mathcal{B}^{e_1})^{e_0}) = (\mathcal{A}^e, \mathcal{B}^e)$ essential. It follows from Definition 4.5.4 that $\text{Cause}(\mathcal{A}^{e_1}, \mathcal{B}^{e_1}) = \gamma^{e_1} = (\rho \circ e_1, \eta^{e_1}, \mathbf{I})$ (Definition 4.3.2). Letting $\varphi \in \text{OL}(\mathbf{I}, \mathbf{K})$ be the unique map witnessing item 2 of Definition 5.5.1 for \mathbf{A} , we see that φ is as required with respect to \mathbf{B} for items 2 and 3 of Definition 5.5.1, and m is a controlled age-change level of \mathbf{B} . \square

Proposition 5.5.3. *Let $\mathbf{A} \subseteq \mathbf{K}$ be weakly diagonal, and suppose $m \in \text{AC}(\mathbf{A})$. Then there is $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ with the following properties:*

1. $f[\mathbf{A}] \subseteq \mathbf{K}$ weakly diagonal,
2. f is the identity on $\{0, \dots, m-1\}$,
3. Writing $\rho = c[\mathbf{A}]|_m^u$, $\mathcal{A} = \mathcal{K}(c[\mathbf{A}]|_m)$, and $\mathcal{B} = \mathcal{K}(c[f[\mathbf{A}]]|_{f(m)})$, we have $\mathcal{B} \supseteq \mathcal{K}(c[\mathbf{A}]|_{m+1})$ and $(\mathcal{A}, \mathcal{B}) \in \text{Con}(\rho)$,
4. $\text{Crit}^+(f[\mathbf{A}]) \cap [m, f(m))$ consists of a single controlled age-change level of $f[\mathbf{A}]$.
5. If $\mathbf{A} \subseteq \mathbf{K}$ has controlled paths, then so does $f[\mathbf{A}]$.

Proof. Write $c[\mathbf{A}]|_m = \{s_0 \preceq_{lex} \dots \preceq_{lex} s_{d-1}\}$. Setting $\mathcal{A} = \mathcal{K}(c[\mathbf{A}]|_m)$, pick any $\mathcal{B} \in P(\rho)$ with $\mathcal{B} \supseteq \mathcal{K}(c[\mathbf{A}]|_{m+1})$ and $(\mathcal{A}, \mathcal{B}) \in \text{Con}(\rho)$. Write $\gamma = (\rho, \eta, \mathbf{I}) = \text{Cause}(\mathcal{A}, \mathcal{B})$. Then $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ will be an insertion for \mathbf{A} with both parameters equal to m and using the gluing γ . Item 1 of Definition 5.4.2 holds since $\mathcal{B} \supseteq \mathcal{K}(c[\mathbf{A}]|_{m+1})$. So $f[\mathbf{A}] \subseteq \mathbf{K}$ is weakly diagonal, and f is the identity on $\{0, \dots, m-1\}$.

Define $\varphi: \mathbf{I} \rightarrow \mathbf{K}$ by setting $\varphi(x) = f(v_x)$, where $v_x \in \mathbf{Y}$ comes from the construction of the insertion f . Notice that $\varphi \in \text{OL}(\mathbf{I}, \mathbf{K})$ for each $i < n$.

Claim. $\mathcal{K}(c[f[\mathbf{A}]]|_{f(m)}) = \mathcal{B}$.

Proof. If $\mathbf{B} \in \mathcal{K}(c[f[\mathbf{A}]]|_{f(m)})$, then since $\mathbf{B}[\gamma]$ is isomorphic to the induced substructure of $\mathbf{B}[c[f[\mathbf{A}]]|_{f(m)}]$ on $\mathbf{B}[\rho] \cup \{f(v_x) : x \in \mathbf{I}\}$, we have $\mathbf{B} \in \mathcal{B}$. In the other direction, assume $\mathbf{B} \in \mathcal{B}$, and form $\mathbf{B}[c[f[\mathbf{A}]]|_{f(m)}]$. Then $\mathbf{B}[c[f[\mathbf{A}]]|_{f(m)}]$ is the free amalgam of the substructures induced on the subsets $\mathbf{B}[\rho] \cup \{f(v_x) : x \in \mathbf{I}\} \cup f[\{0, \dots, m-1\}]$ and $\mathbf{K}_{f(m)}$. The first of these is isomorphic to the free amalgam of $\mathbf{B}[\gamma]$ and $\mathbf{B}[c[\mathbf{A}]|_m]$ about $\mathbf{B}[\rho]$. It follows that $\mathbf{B}[c[f[\mathbf{A}]]|_{f(m)}] \in \mathcal{K}$, and $\mathbf{B} \in \mathcal{K}(c[f[\mathbf{A}]]|_{f(m)})$. \blacksquare

Claim. $\text{Crit}^+(f[\mathbf{A}]) \cap [m, f(m)) = \text{AC}(f[\mathbf{A}]) \cap [m, f(m)) = \{\max(\varphi[\mathbf{I}])\}$, and $\max(\varphi[\mathbf{I}])$ is a controlled age-change level of $f[\mathbf{A}]$.

Proof of claim. We first recall by Lemma 5.4.3 that $\text{Sp}(f[\mathbf{A}]) = f[\text{Sp}(\mathbf{A})]$, since the two parameters m and M used in constructing the insertion f are equal. From this, we see that the only possible members of $\text{Crit}(f[\mathbf{A}])$ in $[m, f(m))$ are age-change levels.

Now since f is the identity on $\{0, \dots, m-1\}$, we have $c[\mathbf{A}]|_m = c[f[\mathbf{A}]]|_m$, so in particular $\mathcal{K}(c[f[\mathbf{A}]]|_m) = \mathcal{A}$. To show that $\max(\varphi[\mathbf{I}])$ is the unique age-change level in the interval $[m, f(m))$, we will show given $y \in [m, f(m))$ that $\mathcal{K}(c[f[\mathbf{A}]]|_y) = \mathcal{A}$ if $y \leq \max(\varphi[\mathbf{I}])$, and $\mathcal{K}(c[f[\mathbf{A}]]|_y) = \mathcal{B}$ if $y > \max(\varphi[\mathbf{I}])$. The left-to-right containments are clear, and since we have already argued that $\mathcal{K}(c[f[\mathbf{A}]]|_{f(m)}) = \mathcal{B}$ in an earlier claim, we have the desired equality for $y > \max(\varphi[\mathbf{I}])$. For $y \leq \max(\varphi[\mathbf{I}])$, the right-to-left containment follows from the fact that \mathbf{I} is chosen with minimum possible cardinality in Definition 4.5.1.

Lastly, we note that $\text{Crit}^+(f[\mathbf{A}]) \setminus \text{Crit}(f[\mathbf{A}]) \subseteq \text{FreeStart}(f[\mathbf{A}])$. As \mathbf{A} is weakly diagonal, we have $\text{FreeStart}(\mathbf{A}) \subseteq m$, so it follows that $\text{FreeStart}(f[\mathbf{A}]) = \text{FreeStart}(\mathbf{A}) \subseteq m$. Hence $\text{Crit}^+(f[\mathbf{A}]) \cap [m, f(m)) = \{\max(\varphi[\mathbf{I}])\}$.

To see that $\max(\varphi[\mathbf{I}])$ is a controlled age-change level of $f[\mathbf{A}]$, items 1 and 3 of Definition 5.5.1 follow from the above paragraph, and item 2 is witnessed by the map $\varphi: \mathbf{I} \rightarrow \mathbf{K}$. \blacksquare

For item 5, assume that $\mathbf{A} \subseteq \mathbf{K}$ has controlled paths. We first argue for each $a \in \mathbf{A}$ that $\text{Path}(a) = \text{Path}(f(a))$. To do this, we use Corollary 5.4.8. This, in turn, follows from item 3 of Proposition 5.4.7 by setting $r = m + 1$, $d_0 = d$, and $e: d_0 \rightarrow d$ the identity map. Then $\gamma^e = \gamma$, and we have $\mathcal{K}(\gamma) \supseteq \mathcal{B} \supseteq \mathcal{K}(c[\mathbf{A}]|_{m+1})$.

Next, we note that if $a \in \mathbf{A}$ is such that $U(a) \in \text{Free}$, then $\text{Start}(c(a)) < m$, since $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal. This implies that $\text{Start}(c(f(a))) = \text{Start}(c(a)) < m$.

Now suppose $a \in \mathbf{A}$ is such that $U(a) \in k^u \setminus \text{Free}$. Observe that if $\text{Start}(c(f(a))) \notin [m, f(m))$, then $\text{Start}(c(f(a))) = f(\text{Start}(c(a)))$. Suppose $\text{Start}(c(f(a))) = f(v_x)$ for some $x \in \mathbf{I}$. We must have $\text{Start}(c(a)) \geq m$. However, we have $\mathcal{K}(c(f(a))|_{\max(\varphi[\mathbf{I}]}) \neq \mathcal{K}(\emptyset_{U(a)})$. This implies that $\mathcal{K}(c(a)|_{m+1}) \neq \mathcal{K}(\emptyset_{U(a)})$. So we must have $\text{Start}(c(a)) = m$.

These observations are enough to conclude that if $a, b \in \mathbf{A}$ satisfy $\text{Start}(f(a)) = \text{Start}(f(b))$, then $\text{Start}(a) = \text{Start}(b)$. This in turn is enough to conclude item 1 of Definition 5.3.4. The other items of Definition 5.3.4 are straightforward, as f is the identity on $\{0, \dots, m-1\}$. \square

5.6 Controlled coding levels

Given $\mathbf{A} \subseteq \mathbf{K}$, a *coding level* of \mathbf{A} is simply a member of \mathbf{A} ; these are the levels of CT which will contain a coding node $c(m)$ for some $m \in \mathbf{A}$.

Before giving the definition of a *controlled coding level* of \mathbf{A} , we first discuss some aspects of age classes in more detail, in particular, how the placement of the coding node in $\text{CT}(m)$ affects $\mathcal{K}(\text{CT}(m+1))$.

Definition 5.6.1. Suppose $\ell, d < \omega$, fix $j < d$, and let $\pi: \ell \rightarrow d$ and $\varphi: \ell \rightarrow k$ be maps.

1. Given \mathbf{B} an \mathcal{L}_ℓ -structure on underlying set B , we form an \mathcal{L}_d -structure $\mathbf{B}\langle j, \pi, \varphi \rangle := \mathbf{C}$ as follows. The underlying set of $\mathbf{B}\langle j, \pi, \varphi \rangle$ is $B \cup \{x\}$, where $x \notin B$ is some new point. We set $V^{\mathbf{C}}(x) = j$ and $V^{\mathbf{C}}(b) = \pi(V^{\mathbf{B}}(b))$ for each $b \in B$. The \mathcal{L}^b -part of \mathbf{C} on B is induced from \mathbf{B} . Lastly, given $b \in B$, we set $R^{\mathbf{C}}(b, x) = \varphi(V^{\mathbf{B}}(b))$.
2. If \mathcal{A} is a hereditary class of \mathcal{L}_d -structures, we define the class of \mathcal{L}_ℓ -structures $\mathcal{A}\langle j, \pi, \varphi \rangle^{-1} := \{\mathbf{B} : \mathbf{B}\langle j, \pi, \varphi \rangle \in \mathcal{A}\}$.

Proposition 5.6.2. Suppose $m < \omega$, $S_m = \{s_0 \preceq_{\text{lex}} \dots \preceq_{\text{lex}} s_{d-1}\} \subseteq \text{CT}(m)$, and suppose $j < d$ is such that $c(m) = s_j$. Suppose $S_{m+1} = \{t_0 \preceq_{\text{lex}} \dots \preceq_{\text{lex}} t_{\ell-1}\} \subseteq \text{CT}(m+1)$ is such that $S_{m+1}|_m \subseteq S_m$. Let $\pi: \ell \rightarrow d$ and $\varphi: \ell \rightarrow k$ be the maps so that $t_\alpha = (s_{\pi(\alpha)}) \frown \varphi(\alpha)$ for each $\alpha < \ell$. Then $\mathcal{K}(S_{m+1}) = \mathcal{K}(S_m)\langle i, \pi, \varphi \rangle^{-1}$.

Proof. For both containments, suppose \mathbf{B} is an \mathcal{L}_ℓ -structure, and write $\mathbf{B}\langle j, \pi, \varphi \rangle := \mathbf{C}$. Then considering the \mathcal{L} -structures $\mathbf{C}[S_m]$ and $\mathbf{B}[S_{m+1}]$, we observe that the map renaming the point x to m is an isomorphism between $\mathbf{C}[S_m]$ and $\mathbf{B}[S_{m+1}]$. \square

For the purposes of defining the notion of a controlled coding level, we will be interested in considering Definition 5.6.1 when $\ell = d - 1$ and $\pi = \epsilon_{j,d}$. So each $\alpha \in d \setminus \{j\}$, there is a unique extension of s_α to a member of $\{t_0, \dots, t_{d-2}\}$.

Definition 5.6.3. Fix a sort $\rho: d \rightarrow k^u$, $\mathcal{A} \in P(\rho)$, $j < d$, and a function $\varphi: (d - 1) \rightarrow k$. We call $(\mathcal{A}, j, \varphi)$ a *controlled coding triple* if all of the following hold:

1. Writing $i = \rho(j)$, we have $\mathcal{A}^{t_j} = P_i^{\min}$,
2. $\mathcal{A}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1} = \mathcal{A}^{\epsilon_{j,d}}$; call this class \mathcal{D} ,
3. For any $\mathcal{B} \in P(\rho)$ with $\mathcal{B} \subsetneq \mathcal{A}$, either $\mathcal{B}^{\epsilon_{j,d}} \subsetneq \mathcal{D}$ or $\mathcal{B}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1} \subsetneq \mathcal{D}$.

Call (\mathcal{A}, j) a *controlled coding pair* if there is some function $\varphi: (d - 1) \rightarrow k$ making $(\mathcal{A}, j, \varphi)$ a controlled coding triple.

Controlled coding triples are the main ingredient in the definition of controlled coding level (Definition 5.6.5). Even though the definition of a controlled coding triple looks rather strict, let us show that in fact, controlled coding triples are somewhat common.

Lemma 5.6.4. *Suppose $\rho: d \rightarrow k^u$ is a sort, and suppose $\mathcal{A} \in P(\rho)$, $j < d$, and $\varphi: (d - 1) \rightarrow k$ are such that $\mathcal{A}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1}$ is reasonable. Then there is $\mathcal{B} \in P(\rho)$ with $\mathcal{B} \subseteq \mathcal{A}$, $\mathcal{B}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1} = \mathcal{A}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1}$, and with $(\mathcal{B}, j, \varphi)$ a controlled coding triple.*

Proof. Write $\mathcal{D} = \mathcal{A}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1}$. Define

$$\mathcal{B} = \bigcap \{ \mathcal{C} \in P(\rho) : \mathcal{C}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1} = \mathcal{D} \}.$$

By Corollary 4.2.7, we have $\mathcal{B} \in P(\rho)$, and we also have $\mathcal{B}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1} = \mathcal{D}$. We also see that for any $\mathcal{B}' \in P(\rho)$ with $\mathcal{B}' \subsetneq \mathcal{B}$, we must have $\mathcal{B}'\langle j, \epsilon_{j,d}, \varphi \rangle^{-1} \subsetneq \mathcal{D}$, showing that item 3 of Definition 5.6.3 holds. To see that items 1 and 2 hold, write $i = \rho(j)$, and consider the classes $\mathcal{A}|_{(t_j, P_i^{\min})}$ and $\mathcal{A}|_{(\epsilon_{j,d}, \mathcal{D})}$. Letting \mathcal{C} denote either of these classes, we have that $\mathcal{C} \in P(\rho)$ (in the second case, this uses that \mathcal{D} is reasonable), and it is straightforward to show that $\mathcal{C}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1} = \mathcal{D}$. This implies that $\mathcal{B}^{t_j} = P_i^{\min}$ and $\mathcal{B}^{\epsilon_{j,d}} = \mathcal{D}$, showing that items 1 and 2 hold. \square

Definition 5.6.5. Suppose $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal, and let $m \in \mathbf{A}$. Write $c[\mathbf{A}]|_m = \{s_0 \preceq_{lex} \dots \preceq_{lex} s_{d-1}\}$, $c[\mathbf{A}]|_{m+1} = \{t_0 \preceq_{lex} \dots \preceq_{lex} t_{d-2}\}$, $\mathcal{A} = \mathcal{K}(c[\mathbf{A}]|_m)$, and let $j < d$ be such that $s_j = c(m)$. Let $\varphi: (d - 1) \rightarrow k$ be given by $\varphi(\alpha) = t_\alpha(m)$ for each $\alpha < d - 1$. We call m a *controlled coding level* of \mathbf{A} if $(\mathcal{A}, j, \varphi)$ is a controlled coding triple. We say that \mathbf{A} has *controlled coding levels* if every $m \in \mathbf{A}$ is a controlled coding level of \mathbf{A} .

Remark. Suppose $m \in \mathbf{A}$ a controlled coding level of \mathbf{A} , and suppose $\mathbf{B} \subseteq \mathbf{A}$ with $m \notin \mathbf{B}$. Then item 2 of Definition 5.6.3 and Proposition 5.6.2 imply that $m \notin \text{AC}(\mathbf{B})$.

Example 5.6.6. Let $\mathcal{K} = \mathcal{G}_3$ be the class of triangle-free graphs. It will be helpful to start thinking about age-sets for \mathcal{G}_3 using the same ideas we developed to talk about age-sets in \mathcal{G}_ℓ -free graphs. So fix $d < \omega$, z a $(d, 3)$ -graph-age function (Example 4.3.6), $j < d$, and $\varphi: (d-1) \rightarrow 2$, and suppose that $(\mathcal{G}_z, j, \varphi)$ is a controlled coding triple. Going through Definition 5.6.3, item 1 just says that $z(\{j\}) = 1$. In the setting of Definition 5.6.5, this just says that the coding node $c(m) = s_j$ cannot be on the left branch.

To understand the content of item 2, suppose $\alpha < \beta < d-1$ are such that $\varphi(\alpha) = \varphi(\beta) = 1$. In the terminology of [8], this is a “parallel 1.” Let us write $\mathcal{D} = \mathcal{G}_z \langle j, \epsilon_{j,d}, \varphi \rangle^{-1} = (\mathcal{G}_z)^{\epsilon_{j,d}}$. Letting $e: 2 \rightarrow (d-1)$ be the increasing injection with range $\{\alpha, \beta\}$, this implies that \mathcal{D}^e is the class $\mathcal{G}_2^{2,2}$ of finite graphs with no edges and an arbitrary V_0, V_1 partition. Item 2 demands that $z(\{\epsilon_{j,d}(\alpha), \epsilon_{j,d}(\beta)\}) = 1$.

Item 3 is perhaps the most subtle. Consider some $\alpha < d-1$ with $\varphi(\alpha) = 0$ and $z(\{\alpha\}) = 1$. Item 3 then requires that $z(\{j, \epsilon_{j,d}(\alpha)\}) = 1$. If this were not the case, i.e. if $z(\{j, \epsilon_{j,d}(\alpha)\}) = 2$, then one could form the $(d, 3)$ -graph-age function y which is identical to z except that $y(\{j, \epsilon_{j,d}(\alpha)\}) = 1$. Then $\mathcal{G}_y \subsetneq \mathcal{G}_z$, but $\mathcal{G}_y \langle j, \epsilon_{j,d}, \varphi \rangle^{-1} = (\mathcal{G}_y)^{\epsilon_{j,d}} = \mathcal{D}$, contradicting item 3.

It turns out that the restrictions pointed out above are the only restrictions for $(\mathcal{G}_z, j, \varphi)$ to be a controlled coding triple. A useful observation that we will take advantage of later, when precisely describing diagonal diaries for \mathcal{G}_3 (see Example 6.2.10) is that whenever (\mathcal{G}_z, j) is a controlled coding pair, then there is a unique φ such that $(\mathcal{G}_z, j, \varphi)$ is a controlled coding triple. We note that (\mathcal{G}_z, j) is a controlled coding pair exactly when given $\alpha \neq \beta \in d \setminus \{j\}$ with $z(\{\alpha\}) = z(\{\beta\}) = 1$ and $z(\{\alpha, j\}) = z(\{\beta, j\}) = 2$, then $z(\{\alpha, \beta\}) = 1$. The unique φ making $(\mathcal{G}_z, j, \varphi)$ a controlled coding triple is defined for $\alpha < (d-1)$ by setting $\varphi(\alpha) = 1$ iff $z(\{\epsilon_{j,d}(\alpha)\}) = 1$ and $z(\{\epsilon_{j,d}(\alpha), j\}) = 2$.

Contrast this example with the class \mathcal{G} of all finite graphs, where every triple is a controlled coding triple.

Example 5.6.7. Consider the class \mathcal{G}_ℓ of K_ℓ -free finite graphs. Fix $d < \omega$, z a (d, ℓ) -graph-age function (see Example 4.3.6), $j < d$ and $\varphi: (d-1) \rightarrow 2$. In order to understand when $(\mathcal{G}_z, j, \varphi)$ is a controlled coding triple, we first need to understand when $\mathcal{G}_z \langle j, \epsilon, \varphi \rangle^{-1} = (\mathcal{G}_z)^{\epsilon_{j,d}}$. Writing $X = \epsilon_{j,d}[\varphi^{-1}(\{1\})]$, one can show that this happens exactly when both of the following occur:

1. For every $T \in \binom{X}{<\ell}$, we have $z(T) < \ell - 1$,
2. For every $T \in \binom{X}{<(\ell-1)}$, we have $z(T \cup \{j\}) = z(T) + 1$.

Now suppose $(\mathcal{G}_z, j, \varphi)$ satisfies the above two items. When is it a controlled coding triple? First, we need $z(\{j\}) = 1$ so that item 1 of Definition 5.6.3 holds. Aside from this, we need to enforce item 3 by ensuring that z is “as small as possible.” Given $S \in \binom{d}{<\ell}$, call S *determined* if either $S \subseteq d \setminus \{j\}$ or $S \subseteq X \cup \{j\}$. We enforce item 3 by demanding for every $T \in \binom{d}{<\ell}$ that $z(T) = \max\{z(S) : S \subseteq T \text{ is determined}\}$.

As in the case of the class \mathcal{G}_3 , we have that whenever (\mathcal{G}_z, j) is a controlled coding pair, then there is a unique φ making $(\mathcal{G}_z, j, \varphi)$ a controlled coding triple. We summarize by noting that (\mathcal{G}_z, j) is a controlled coding pair iff all of the following hold:

1. $z(\{j\}) = 1$,
2. Writing $X = \{\alpha \in d \setminus \{j\} : z(\{\alpha, j\}) = z(\{\alpha\}) + 1\}$, then for every $T \in \binom{X}{<\ell}$, we have $z(T) < \ell - 1$, and for every $T \in \binom{X}{<(\ell-1)}$, we have $z(T \cup \{j\}) = z(T) + 1$,
3. With the same notion of “determined” as above, we have for every $T \in \binom{d}{<\ell}$ that $z(T) = \max\{z(S) : S \subseteq T \text{ is determined}\}$.

The unique φ making $(\mathcal{G}_z, j, \varphi)$ a controlled coding triple is defined for $\alpha < (d - 1)$ by setting $\varphi(\alpha) = 1$ iff $\epsilon_{j,d}(\alpha) \in X$.

The two remaining goals for this subsection are as follows. First, we show that the notion of controlled coding level behaves well when passing to a substructure of \mathbf{A} . Second, given $m \in \mathbf{A}$, we build $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ which fixes $\{0, \dots, m - 1\}$ and with $f(m)$ a controlled coding level of $f[\mathbf{A}]$.

Lemma 5.6.8. *Suppose $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal, and say $\mathbf{B} \subseteq \mathbf{A}$. If $m \in \mathbf{B}$ is a controlled coding level of \mathbf{A} , then m is a controlled coding level of \mathbf{B} . If $m \notin \mathbf{B}$, then $m \notin \text{AC}(\mathbf{B})$. In particular, if \mathbf{A} has controlled coding levels, then so does \mathbf{B} .*

Proof. The case where $m \notin \mathbf{B}$ follows from item 2 of Definition 5.6.3. We focus on the case where $m \in \mathbf{B}$.

We rephrase the lemma statement in terms of controlled coding triples. So fix $d < \omega$, a sort $\rho: d \rightarrow k^u$, a controlled coding triple $(\mathcal{A}, j, \varphi)$ with $\mathcal{A} \in P(\rho)$, and an increasing injection $e: r \rightarrow d$ with $j \in \text{Im}(e)$. Let $\ell < r$ be such that $e(\ell) = j$, and let $\sigma: (r - 1) \rightarrow (d - 1)$ be induced by e in the obvious way, i.e. $\sigma = \delta_{j,d-1} \circ e \circ \epsilon_{\ell,r}$. Our goal is to show that $(\mathcal{A}^e, \ell, \varphi \circ \sigma)$ is a controlled coding triple; we verify the three items of Definition 5.6.3.

For item 1, writing $i = \rho(j)$, we have $\mathcal{A}^{i,j} = P_i^{\min}$, so $(\mathcal{A}^e)^{\ell,\ell} = P_i^{\min}$ as well.

For item 2, we first observe that $e \circ \epsilon_{\ell,r} = \epsilon_{j,d} \circ \sigma$. Then we have:

$$\begin{aligned} (\mathcal{A}^e)^{\ell,r} &= (\mathcal{A}^{\epsilon_{j,d}})^{\sigma} \\ &= (\mathcal{A}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1})^{\sigma} \\ &= \mathcal{A}^e \langle \ell, \epsilon_{\ell,r}, \varphi \circ \sigma \rangle^{-1}. \end{aligned}$$

Intuitively, going back to thinking about a structure $\mathbf{A} \subseteq \mathcal{K}$ and a level m , the above equation just says that if \mathbf{A} has no age change at level m off of the coding node, then the same is true for \mathbf{B} .

For item 3, suppose $\mathcal{B} \subseteq \mathcal{A}^e$ is such that $\mathcal{B}^{\ell,r} = \mathcal{B}\langle \ell, \epsilon_{\ell,r}, \varphi \circ \sigma \rangle^{-1} = (\mathcal{A}^e)^{\ell,r} = \mathcal{A}^e \langle \ell, \epsilon_{\ell,r}, \varphi \circ \sigma \rangle^{-1}$. We need to show that $\mathcal{B} = \mathcal{A}^e$. To that end, form the class $\mathcal{C} := \mathcal{A}|_{(e,\mathcal{B})}$. It is routine to check that $\mathcal{C}^{\epsilon_{j,d}} = \mathcal{A}^{\epsilon_{j,d}}$. To check that $\mathcal{C}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1} = \mathcal{A}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1}$, the left-to-right inclusion is clear. For the right-to-left inclusion, fix $\mathbf{D} \in \mathcal{A}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1}$. This means that $\mathbf{D}\langle j, \epsilon_{j,d}, \varphi \rangle \in \mathcal{A}$, and we want to show that it is also in \mathcal{C} . To that end, we note that $(\mathbf{D}\langle j, \epsilon_{j,d}, \varphi \rangle)^e = \mathbf{D}^{\sigma} \langle \ell, \epsilon_{\ell,r}, \varphi \circ \sigma \rangle \in \mathcal{A}^e$. Then $\mathbf{D}^{\sigma} \in \mathcal{A}^e \langle \ell, \epsilon_{\ell,r}, \varphi \circ \sigma \rangle^{-1} = \mathcal{B}\langle \ell, \epsilon_{\ell,r}, \varphi \circ \sigma \rangle^{-1}$. So $\mathbf{D}^{\sigma} \langle \ell, \epsilon_{\ell,r}, \varphi \circ \sigma \rangle = (\mathbf{D}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1})^e \in \mathcal{B}$, which means that $\mathbf{D}\langle j, \epsilon_{j,d}, \varphi \rangle \in \mathcal{C}$. Because $(\mathcal{A}, j, \varphi)$ is a controlled coding triple, we must have $\mathcal{A} = \mathcal{C}$, implying that $\mathcal{A}^e = \mathcal{B}$. \square

Proposition 5.6.9. *Suppose $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal, and fix $m \in \mathbf{A}$ with $U(m) = i$. Then there is $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ which is the identity on $\{0, \dots, m-1\}$, with $f[\mathbf{A}] \subseteq \mathbf{K}$ weakly diagonal, and with $f(m)$ a controlled coding level of $f[\mathbf{A}]$.*

Furthermore, for any $n \geq m+1$, we have $\mathcal{K}(f[c[\mathbf{A}]|_n]) = \mathcal{K}(c[\mathbf{A}]|_n)$. Hence if $\mathbf{A} \subseteq \mathbf{K}$ has controlled paths, then so does $f[\mathbf{A}]$.

Proof. The desired f will be an insertion for \mathbf{A} with both parameters equal to m . To describe the gluing we use, write $c[\mathbf{A}]|_m = \{s_0 \preceq_{lex} \dots \preceq_{lex} s_{d-1}\}$ and $c[\mathbf{A}]|_{m+1} = \{t_0 \preceq_{lex} \dots \preceq_{lex} t_{d-2}\}$, and suppose $j < d$ is such that $s_j = c(m)$. Write $\mathcal{B} = \mathcal{K}(c[\mathbf{A}]|_{m+1})$, and let $\varphi: (d-1) \rightarrow k$ be the function so that $t_\alpha = (s_{\epsilon_{j,d}(\alpha)})^\frown \varphi(\alpha)$. So by Proposition 5.6.2, we have $\mathcal{B} = (\mathcal{K}(c[\mathbf{A}]|_m)) \langle j, \epsilon_{j,d}, \varphi \rangle^{-1}$.

Write $\rho = c[\mathbf{A}]|_m^u$, and use Lemma 5.6.4 to find $\mathcal{A} \in P(\rho)$ with $\mathcal{A} \subseteq \mathcal{K}(c[\mathbf{A}]|_m)$, $\mathcal{A} \langle j, \epsilon_{j,d}, \varphi \rangle^{-1} = \mathcal{B}$, and so that $(\mathcal{A}, j, \varphi)$ is a controlled coding triple. We can find some gluing $\gamma = (\rho, \eta, \mathbf{I})$ so that $\mathcal{A} = \mathcal{K}(\gamma)$. This is the gluing we use to form the insertion $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$. To check that this is indeed an insertion, we need to check that item 1 of Definition 5.4.2 holds. Let $\mathbf{B} \subseteq \mathbf{A}$ be finite with $\min\{\mathbf{B}\} \geq m$, and form the structure $(\mathbf{B}/c[\mathbf{A}]|_m)[\gamma]$. From the definition of γ , we have $(\mathbf{B}/c[\mathbf{A}]|_m)[\gamma] \in \mathcal{K}$ iff $\mathbf{B}/c[\mathbf{A}]|_m \in \mathcal{A}$. If $m \notin \mathbf{B}$, then $\mathbf{B}/c[\mathbf{A}]|_m \in \mathcal{A}$ since $\mathcal{A}^{\epsilon_{j,d}} = \mathcal{B}$. If $m \in \mathbf{B}$, then we can write $\mathbf{B}/c[\mathbf{A}]|_m \cong (\mathbf{B}/c[\mathbf{A}]|_m)^{\epsilon_{j,d}} \langle j, \epsilon_{j,d}, \varphi \rangle$, and this is in \mathcal{A} since $(\mathbf{B}/c[\mathbf{A}]|_m)^{\epsilon_{j,d}} \in \mathcal{B}$ and since $\mathcal{A} \langle j, \epsilon_{j,d}, \varphi \rangle^{-1} = \mathcal{B}$.

We have that f is the identity on $\{0, \dots, m-1\}$, and by Proposition 5.4.6, $f[\mathbf{A}] \subseteq \mathbf{K}$ is weakly diagonal. Towards showing that $f(m)$ is a controlled coding level of $f[\mathbf{A}]$, write $c[f[\mathbf{A}]]|_{f(m)} = \{\sigma_0 \preceq_{lex} \dots \preceq_{lex} \sigma_{d-1}\}$ and $c[f[\mathbf{A}]]|_{f(m)+1} = \{\tau_0 \preceq_{lex} \dots \preceq_{lex} \tau_{d-2}\}$. Since f is the identity on $\{0, \dots, m-1\}$, we have that $s_\ell \sqsubseteq \sigma_\ell$ for each $\ell < d$. Also notice that given $a \in \mathbf{A}$ with $a > m$, then if $\ell < d-1$ is such that $c(a)|_{m+1} = t_\ell$, then $c(f(a))|_{f(m)+1} = \tau_\ell$. In particular, we have $\tau_\ell(f(m)) = t_\ell(m)$ for any $\ell < d-1$. Hence it is enough to show that $\mathcal{K}(c[f[\mathbf{A}]]|_{f(m)}) = \mathcal{A}$. The left-to-right inclusion is clear. For the right-to-left inclusion, fix $\mathbf{B} \in \mathcal{A}$, and form $\mathbf{B}[c[f[\mathbf{A}]]|_{f(m)}]$. By the construction of f , this is the free amalgam of $\mathbf{B}[c[\mathbf{A}]|_m]$ and $\mathbf{B}[\gamma]$ along $\mathbf{B}[\rho]$. Since $\mathcal{A} \subseteq \mathcal{K}(c[\mathbf{A}]|_m)$, we see that $\mathbf{B}[c[\mathbf{A}]|_m] \in \mathcal{K}$, and by the definition of \mathbf{A} , we have $\mathbf{B}[\gamma] \in \mathcal{K}$. So $\mathbf{B} \in \mathcal{K}(c[f[\mathbf{A}]]|_{f(m)})$ as desired.

We move to the ‘‘furthermore’’ in the proposition statement, so fix any $n \geq m+1$. The left-to-right inclusion is item 2 of Proposition 5.4.7. For the right-to-left inclusion, we cannot directly use item 3 of Proposition 5.4.7 since $m \in \mathbf{A}$; however, the proof will follow a very similar idea. Write $c[\mathbf{A}]|_n = \{u_0 \preceq_{lex} \dots \preceq_{lex} u_{d_1-1}\}$. Fix $\mathbf{B} \in \mathcal{K}(c[\mathbf{A}]|_n)$ with underlying set B , and now \mathbf{B} steals the nodes from $\mathbf{A} \cap [m, n-1]$. We create new nodes $X := \{x_\alpha : \alpha \in \mathbf{A} \cap [m, n-1]\}$, and form an \mathcal{L}_d -structure \mathbf{X} on $B \cup X$ as follows:

- Given $b \in B$ and $p < d$, we have $V^{\mathbf{X}}(b) = p$ iff $u_{V^{\mathbf{B}}(b)}|_m = s_p$.
- Given $\alpha \in \mathbf{A} \cap [m, n-1]$, we have $V^{\mathbf{X}}(x_\alpha) = p$ iff $c(\alpha)|_m = s_p$.
- The binary relations on B are those from \mathbf{B} , and given $x_\alpha, x_\beta \in X$, we have $R^{\mathbf{X}}(x_\alpha, x_\beta) = R(\alpha, \beta)$.
- Given $\alpha \in \mathbf{A} \cap [m, n-1]$ and $b \in B$, we have $R^{\mathbf{X}}(b, x_\alpha) = u_{V^{\mathbf{B}}(b)}(\alpha)$.

By design, we have $\mathbf{X}[c[\mathbf{A}]|_m] \cong \mathbf{B}[c[\mathbf{A}]|_n] \setminus ([m, n-1] \cap \mathbf{A})$. The rest of the proof is exactly the same as that of Proposition 5.4.7, provided we can show that

$\mathbf{X} \in \mathcal{K}(\gamma) = \mathcal{A}$. To do this, consider forming from \mathbf{B} an \mathcal{L}_{d-1} -structure \mathbf{W} exactly as above, but only adding the new nodes $W := X \setminus \{x_m\}$, and in the four bullets above, replacing every instance of m with $m + 1$, and replacing every instance of s_i with t_i . Then $\mathbf{W} \in \mathcal{B}$ and $\mathbf{X} \cong \mathbf{W}\langle j, \epsilon_{j,d}, \varphi \rangle$, where x_m plays the role of the new vertex x from Definition 5.6.1. So by the definition of \mathcal{A} , we have $\mathbf{X} \in \mathcal{A}$. One can now duplicate the rest of the proof of Proposition 5.4.7 to conclude that $\mathbf{B} \in \mathcal{K}(\hat{f}[c[\mathbf{A}]|_n])$.

Now assume that $\mathbf{A} \subseteq \mathbf{K}$ has controlled paths. First note that by Corollary 5.4.8 that for any $a \in \mathbf{A}$, we have $\text{Path}(a) = \text{Path}(f(a))$. The rest of the argument is much like the end of the proof of Proposition 5.5.3. We note that if $a \in \mathbf{A}$ is such that $U(a) \in \text{Free}$, then $\text{Start}(c(a)) < m$, since $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal. This implies that $\text{Start}(c(f(a))) = \text{Start}(c(a)) < m$.

Now suppose $a \in \mathbf{A}$ is such that $U(a) \in k^u \setminus \text{Free}$. Observe that if $\text{Start}(c(f(a))) \notin [m, f(m))$, then $\text{Start}(c(f(a))) = f(\text{Start}(c(a)))$. Suppose $\text{Start}(c(f(a))) = f(v_x)$ for some $x \in \mathbf{I}$. Then $\mathcal{K}(c(f(a))|_{f(m)}) \neq \mathcal{K}(\emptyset_{U(a)})$, so also $\mathcal{K}(c(a)|_{m+1}) \neq \mathcal{K}(\emptyset_{U(a)})$. So we must have $\text{Start}(c(a)) = m$.

These observations are enough to conclude that if $a, b \in \mathbf{A}$ satisfy $\text{Start}(f(a)) = \text{Start}(f(b))$, then $\text{Start}(a) = \text{Start}(b)$. This in turn is enough to conclude item 1 of Definition 5.3.4. The other items of Definition 5.3.4 are straightforward, as f is the identity on $\{0, \dots, m-1\}$. \square

5.7 Controlled splitting levels

The notion of *controlled splitting level* is more complicated than our definitions of controlled age change and controlled coding level. Part of the reason why is the following observation: if $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal, $m \in \text{Sp}(\mathbf{A})$, and $\mathbf{B} \subseteq \mathbf{A}$, then it is possible to have $m \in \text{AC}(\mathbf{B})$. For a concrete example of this, suppose $\mathcal{K} = \mathcal{G}_3$ is the class of triangle-free graphs, $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal, and $0 \in \text{Sp}(\mathbf{A})$. If $\mathbf{B} = \{a \in \mathbf{A} : R(a, 0) = 1\}$, then $0 \in \text{AC}(\mathbf{B})$.

Our definition of controlled splitting node needs to take this into account by eliminating this possibility if at all possible, and if not, then ensuring that upon passing to a substructure in which the splitting level becomes an age-change level, that we will have a controlled age-change level of the substructure. It turns out that much will depend on the $i < k^u$ for which we have $m \in \text{Sp}_i(\mathbf{A})$. In particular, if i does not have a free relation and $\text{Sp}_i(m) = (i, 0^m)$, then we must worry about possible age changes.

Definition 5.7.1. Suppose $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal, and let $m \in \text{Sp}_i(\mathbf{A})$ for some $i < k^u$. Write $\rho = c[\mathbf{A}]|_m^u$, $c[\mathbf{A}]|_m := \{s_0 \preceq_{\text{lex}} \dots \preceq_{\text{lex}} s_{d-1}\}$ and $c[\mathbf{A}]|_{m+1} = \{t_0 \preceq_{\text{lex}} \dots \preceq_{\text{lex}} t_d\}$. Suppose $\alpha < d$ is such that $s_\alpha \in T_i$ is the splitting node; so t_α and $t_{\alpha+1}$ are the immediate successors of s_α .

Our definition of m being a *controlled splitting level* for \mathbf{A} will depend on if i has a free relation.

- If $i \in \text{Free}$, write $\text{Free}(i) = (j, q)$. Then we demand that $c(m) = (j, 0^m)$ and that $t_{\alpha+1} = s_\alpha \widehat{\ } q$.
- If $i \in k^u \setminus \text{Free}$, but we have $s_\alpha \neq (i, 0^m)$, write $q = s_\alpha(\text{Start}(s_\alpha))$ and $j = U(\text{Start}(s_\alpha))$. Then we demand that $c(m) = (j, 0^m)$ and that $t_{\alpha+1} = s_\alpha \widehat{\ } q$.

- The tricky situation is when $i \in k^u \setminus \text{Free}$ and $s_\alpha = (i, 0^m)$. We demand that the following all hold:
 1. Writing $\mathcal{A} = \mathcal{K}(c[\mathbf{A}]|_{m+1} \setminus \{t_\alpha\}) \in P(\rho)$, we demand that $(\mathcal{K}(c[\mathbf{A}]|_m), \mathcal{A}) \in \text{Con}(\rho)$. We note that $\mathcal{K}(c[\mathbf{A}]|_m)^{\iota_\alpha} = \mathcal{K}(s_\alpha) = \mathcal{K}(\emptyset_i)$.
 2. Suppose $q \in [1, k]$ is such that $s_\alpha \widehat{\ } q = t_{\alpha+1} \in c[\mathbf{A}]|_{m+1}$. Since $i \in k^u \setminus \text{Free}$, we must have $\mathcal{A}^{\iota_\alpha} = \mathcal{K}(s_\alpha \widehat{\ } q) \subsetneq \mathcal{K}(\emptyset_i)$, so by Proposition 4.5.2, we must have $(\mathcal{K}(s_\alpha), \mathcal{K}(s_\alpha \widehat{\ } q)) \in \text{Con}(\iota_i)$. By Lemma 4.6.9, $\text{Cause}(\mathcal{K}(s_\alpha), \mathcal{K}(s_\alpha \widehat{\ } q)) = \gamma\langle i, q', j \rangle$ for some $j < k^u$ and $q' < k$. We demand that $q' = q$ and $c(m) = (j, 0^m)$.

We say that $\mathbf{A} \subseteq \mathbf{K}$ has *controlled splitting levels* if every $m \in \text{Sp}(\mathbf{A})$ is a controlled splitting level for \mathbf{A} .

Given $m \in \text{Sp}_i(\mathbf{A})$ a controlled splitting level for \mathbf{A} , we will want to understand what this implies about $\mathcal{K}(c[\mathbf{A}]|_{m+1})$ in relation to $\mathcal{K}(c[\mathbf{A}]|_m)$.

Lemma 5.7.2. *Suppose $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal and $m \in \text{Sp}_i(\mathbf{A})$ is a controlled splitting level for \mathbf{A} with $m > \max(\text{FreeStart}(\mathbf{A}))$. Write $c[\mathbf{A}]|_m := \{s_0 \preceq_{\text{lex}} \cdots \preceq_{\text{lex}} s_{d-1}\}$ and $c[\mathbf{A}]|_{m+1} = \{t_0 \preceq_{\text{lex}} \cdots \preceq_{\text{lex}} t_d\}$. Suppose $\alpha < d$ is such that $s_\alpha \in T_i$ is the splitting node; so $t_\alpha = s_\alpha \widehat{\ } 0$ and $t_{\alpha+1} = s_\alpha \widehat{\ } q$ for some $q \in [1, k]$. Write $\mathcal{A} = \mathcal{K}(c[\mathbf{A}]|_m)^{\delta_{\alpha,d}}$.*

1. If $i \in \text{Free}$, then $\mathcal{K}(c[\mathbf{A}]|_{m+1}) = \mathcal{A}$.
2. If $i \in k^u \setminus \text{Free}$ and $s_\alpha \neq (i, 0^m)$, then $\mathcal{K}(c[\mathbf{A}]|_{m+1}) = \mathcal{A}$.
3. If $i \in k^u \setminus \text{Free}$ and $s_\alpha = (i, 0^m)$, then write $\mathcal{B} = \mathcal{K}(t_{\alpha+1})$. Then $\mathcal{K}(c[\mathbf{A}]|_{m+1}) = \mathcal{A}_{(\iota_{\alpha+1}, \mathcal{B})}$.

Proof. For all three items, the left-to-right inclusion is clear.

For item 1, suppose $\mathbf{B} \in \mathcal{A}$; this means that the structure $\mathbf{C} := (\delta_{\alpha,d} \cdot \mathbf{B})[c[\mathbf{A}]|_m]$ is in \mathcal{K} . We want to show that $\mathbf{D} := \mathbf{B}[c[\mathbf{A}]|_{m+1}]$ is in \mathcal{K} . We note that $\mathbf{C} \subseteq \mathbf{D}$ is an induced substructure with $\mathbf{D} \setminus \mathbf{C} = \{m\}$. Writing $\text{Free}(i) = (q, j)$, our assumption that m is a controlled splitting level for \mathbf{A} tells us that $U(m) = j$, that $R^{\mathbf{D}}(b, m) = q$ for $b \in \mathbf{B}$ with $V^{\mathbf{B}}(b) = \alpha + 1$, and that $R^{\mathbf{D}}(b, m) = 0$ for $b \in \mathbf{B}$ with $V^{\mathbf{B}}(b) \neq \alpha + 1$. This assumption also tells us that $R(m, \ell) = 0$ for every $\ell < m$. Since $\text{Free}(i) = (q, j)$, we know that the induced substructure of \mathbf{D} on $\mathbf{B} \cup \{m\}$ is in \mathcal{K} . So \mathbf{D} is a free amalgam of structures in \mathcal{K} .

For item 2, suppose $\mathbf{B} \in \mathcal{A}$. The structure of the argument is very similar to item 1; with \mathbf{C} and \mathbf{D} as above, we know for some $j < k^u$ and $q < k$ that $U(m) = j$, $R^{\mathbf{D}}(b, m) = q$ for $b \in \mathbf{B}$ with $V^{\mathbf{B}}(b) = \alpha + 1$, and otherwise $R^{\mathbf{D}}(b, m) = 0$. We also know that $R(m, \ell) = 0$ for every $\ell < m$. This time, we don't have that (q, j) is a free relation for i ; however, we do know that $\text{Start}(s_\alpha) < m$ has $U(\text{Start}(s_\alpha)) = j$ and $R^{\mathbf{C}}(b, \text{Start}(s_\alpha)) = q$ for every $b \in \mathbf{B}$. So $\{b \in \mathbf{B} : V^{\mathbf{B}}(b) = \alpha + 1\} \cup \{m\} \subseteq \mathbf{D}$ is isomorphic to a substructure of \mathbf{C} , and we once again conclude that \mathbf{D} is a free amalgam of structures in \mathbf{K} .

For item 3, suppose $\mathbf{B} \in \mathcal{A}_{(\iota_{\alpha+1}, \mathcal{B})}$. Arguing as in the first two items, this time we use the fact that writing $\text{Cause}(\mathcal{K}(\emptyset_i), \mathcal{B}) = \gamma\langle i, q, j \rangle$, then j and q will play the same roles that they do in each of the above proofs. The fact that $\mathbf{B}^{\iota_{\alpha+1}} \in \mathcal{B}$ tells

us that $\mathbf{B} \cup \{m\} \subseteq \mathbf{D}$ is in \mathcal{K} , and once again \mathbf{D} is a free amalgam of structures in \mathcal{K} . \square

Example 5.7.3. Let \mathcal{K} be the class of finite graphs where edges can either be red, green, or blue, and where we forbid the following irreducible structures:

- K_4 , no matter the colors of edges involved,
- monochromatic blue triangles,
- monochromatic red triangles.

So $k^u = 1$ and $k = 4$, where 1, 2, and 3 refer to green, red, and blue edges, respectively. The 6 members of $P(1)$ are as follows:

- \mathcal{K} , the maximum,
- \mathcal{A} , the class of all members of \mathcal{K} which are triangle-free,
- \mathcal{A}^r , the class of all members of \mathcal{K} which are triangle-free and have no red edges,
- \mathcal{A}^b , the class of all members of \mathcal{K} which are triangle-free and have no blue edges,
- $\mathcal{A}^{r,b}$, the class of all members of \mathcal{K} which are triangle-free and have only green edges,
- \mathcal{B} , the class of all members of \mathcal{K} with no edges.

Fix some $m < \omega$, and for each $q \in \{1, 2, 3\}$, set $t_q = (0^m) \frown q \in \text{CT}(m+1)$. We have

$$\mathcal{K}(t_q) = \begin{cases} \mathcal{A} & \text{if } q = 1 \\ \mathcal{A}^r & \text{if } q = 2 \\ \mathcal{A}^b & \text{if } q = 3 \end{cases}$$

It follows that if $\mathbf{A} \subseteq \mathbf{K}$, m is a controlled splitting level of \mathbf{A} , and $\text{Sp}^{\mathbf{A}}(m) = 0^m$, then we must have $t_1 \in c[\mathbf{A}]|_{m+1}$, as this is the only choice resulting in a consecutive age change. Conversely, Lemma 4.6.10 ensures that if $c[\mathbf{A}]|_m = \{s_0 \preceq_{\text{lex}} \cdots \preceq_{\text{lex}} s_{d-1}\}$, where $s_0 = 0^m$, and if $c[\mathbf{A}]|_m = \{s_i \frown 0 : i < d\} \cup \{t_1\}$, then m is a controlled splitting level.

Lemma 5.7.4. *Suppose $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal, and let $m \in \text{Sp}_i(\mathbf{A})$ be a controlled splitting level. Let other notation be as in the statement of Lemma 5.7.2. Fix $\mathbf{B} \subseteq \mathbf{A}$.*

1. *If t_α and $t_{\alpha+1}$ are in $c[\mathbf{B}]|_{m+1}$, then $m \in \text{Sp}(\mathbf{B})$ is a controlled splitting level of \mathbf{B} .*
2. *If $t_{\alpha+1} \notin c[\mathbf{B}]|_{m+1}$, then $m \notin \text{Crit}(\mathbf{B})$.*
3. *If $t_\alpha \notin c[\mathbf{B}]|_{m+1}$ and $t_{\alpha+1} \in c[\mathbf{B}]|_{m+1}$, then if i has a free relation, or if $s_\alpha \neq (i, 0^m)$, then $m \notin \text{Crit}(\mathbf{B})$. If i does not have a free relation and $s_\alpha = (i, 0^m)$, then $m \in \text{AC}(\mathbf{B})$ is a controlled age change level.*

Proof. Item 1 is obvious, except possibly in the case where i does not have a free relation and $s_\alpha = (i, 0^m)$, in which case it follows from Lemma 5.7.2 and Corollary 4.5.3.

For item 2, it is clear that $m \notin \text{Sp}(\mathbf{B}) \cup \mathbf{B}$. To see that $m \notin \text{AC}(\mathbf{B})$, we use Lemma 5.7.2.

Similarly, the first part of item 3 also follows from Lemma 5.7.2. Now suppose i does not have a free relation and $s_\alpha = (i, 0^m)$. Write $Q = c[\mathbf{A}]|_{m+1} \setminus \{t_\alpha\}$, and write $\mathcal{D} = \mathcal{K}(t_{\alpha+1})$. By Lemma 5.7.2, we have $\mathcal{K}(Q) = \mathcal{K}(c[\mathbf{A}]|_m)|_{(\iota_\alpha, \mathcal{D})}$. By Lemma 4.6.10, we have $(\mathcal{K}(c[\mathbf{A}]|_m), \mathcal{K}(Q)) \in \text{Con}(\rho)$. By Corollary 4.5.3, ι_α is the unique increasing injection so that we have $(\mathcal{K}(c[\mathbf{A}]|_m)^{\iota_\alpha}, \mathcal{K}(Q)^{\iota_\alpha}) = (\mathcal{K}(s_\alpha), \mathcal{K}(t_{\alpha+1})) \in \text{ECon}(\iota_i)$. By Lemma 4.6.9, we can write $\text{Cause}(\mathcal{K}(s_\alpha), \mathcal{K}(t_{\alpha+1})) = \gamma\langle i, q, j \rangle$ for some $q \in [1, k]$ and $j < k^u$. Since m is a controlled splitting level of \mathbf{A} we have $t_{\alpha+1} = s_\alpha \widehat{c} q$ and $c(m) = (j, 0^m)$. Write $\sigma = c[\mathbf{B}]|_m^u$; since $t_{\alpha+1} \in c[\mathbf{B}]|_{m+1} \subseteq Q$, Corollary 4.5.3 gives us that $(\mathcal{K}(c[\mathbf{B}]|_m), \mathcal{K}(c[\mathbf{B}]|_{m+1})) \in \text{Con}(\sigma)$. Letting $b < |c[\mathbf{B}]|_m|$ be chosen so that $\overrightarrow{c[\mathbf{B}]|_m}(b) = s_\alpha$ (Notation 4.1.4), then ι_b is the unique increasing injection with $(\mathcal{K}(c[\mathbf{B}]|_m)^{\iota_b}, \mathcal{K}(c[\mathbf{B}]|_{m+1})^{\iota_b}) = (\mathcal{K}(s_\alpha), \mathcal{K}(t_{\alpha+1}))$ essential, so by Definition 4.5.4 we have $\text{Cause}((\mathcal{K}(c[\mathbf{B}]|_m), \mathcal{K}(c[\mathbf{B}]|_{m+1}))) = (\sigma, \iota_b) \cdot \gamma\langle i, q, j \rangle$; see Definition 4.3.2 item 2. If \mathbf{J} denotes the singleton enumerated \mathcal{L} -structure with $U^{\mathbf{J}}(0) = j$, then since $c(m) = (j, 0^m)$, the map $f: \mathbf{J} \rightarrow \mathbf{K}$ given by $f(x) = m$ is in $\text{OL}(\mathbf{J}, \mathbf{K})$, and for $t \in c[\mathbf{B}]|_{m+1}$, we have $t(m) = q$ if $t = t_{\alpha+1}$ and $t(m) = 0$ otherwise. This verifies that m satisfies all of the criteria from Definition 5.5.1 to be a controlled age change level. \square

Corollary 5.7.5. *Suppose $\mathbf{A} \subseteq \mathbf{K}$ is a substructure with controlled age change levels and controlled splitting levels. Then if $\mathbf{B} \subseteq \mathbf{A}$, then \mathbf{B} also has controlled age change levels and controlled splitting levels.*

The remaining goal of this subsection is to construct $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ turning a splitting level of $\mathbf{A} \subseteq \mathbf{K}$ into a controlled splitting level of $f[\mathbf{A}]$.

Proposition 5.7.6. *Suppose $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal and $m \in \text{Sp}_i(\mathbf{A})$ with $m > \max(\text{FreeStart}(\mathbf{A}))$. Then there is $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ with the following properties:*

1. f is the identity on $\{0, \dots, m-1\}$,
2. $f[\mathbf{A}] \subseteq \mathbf{K}$ is weakly diagonal,
3. $\text{Crit}(f[\mathbf{A}]) \cap [m, f(m))$ consists of a single controlled splitting level of $f[\mathbf{A}]$.
4. If $\mathbf{A} \subseteq \mathbf{K}$ has controlled paths, then $f[\mathbf{A}]$ also has controlled paths.

Proof. We will construct f as an insertion for \mathbf{A} with parameters m and $m+1$. Write $c[\mathbf{A}]|_m = \{s_0 \preceq_{\text{lex}} \dots \preceq_{\text{lex}} s_{d-1}\}$, and write $c[\mathbf{A}]|_{m+1} = \{t_0 \preceq_{\text{lex}} \dots \preceq_{\text{lex}} t_d\}$. Let $\alpha < d$ be such that $\text{Sp}^{\mathbf{A}}(m) = s_\alpha$. Write $\sigma = c[\mathbf{A}]|_m^u$ and $\rho = c[\mathbf{A}]|_{m+1}^u$.

Our construction will need to deal with the three main cases, which we number for reference.

1. $i \in \text{Free}$,
2. $i \in k^u \setminus \text{Free}$ and $s_\alpha \neq (i, 0^m)$,
3. $i \in k^u \setminus \text{Free}$ and $s_\alpha = (i, 0^m)$.

The construction is similar in all three cases, so we will establish some common notation. In case 1, write $\text{Free}(i) = (q, j)$. In case 2, write $(s_\alpha(\text{Start}(s_\alpha)), U(\text{Start}(s_\alpha))) = (q, j)$. In case 3, write $\mathcal{A} = \mathcal{K}(c[\mathbf{A}]|_m)$ and fix any $\mathcal{B} \supseteq \mathcal{K}(c[\mathbf{A}]|_{m+1} \setminus \{t_\alpha\})$ with $(\mathcal{A}, \mathcal{B}) \in \text{Con}(\sigma)$. Then $\mathcal{A}^{\epsilon_{\alpha,d}} = \mathcal{B}^{\epsilon_{\alpha,d}}$, so Lemma 4.6.10 yields that $\mathcal{B}^{\iota_\alpha} \subsetneq \mathcal{K}(\emptyset_i)$. By Lemma 4.6.9, we can find (q, j) with $\text{Cause}(\mathcal{K}(\emptyset_i), \mathcal{B}^{\iota_\alpha}) = \gamma\langle i, q, j \rangle$. This concludes our definitions of $q < k$ and $j < k^u$ in all three cases.

Writing $\rho = c[\mathbf{A}]|_{m+1}^u$, we define our insertion in all three cases using the gluing $\gamma = (\rho, \eta, \mathbf{J}) := (\rho, \iota_{\alpha+1}) \cdot \gamma\langle i, q, j \rangle$. Explicitly, this is the gluing where \mathbf{J} is the singleton enumerated structure with $U^{\mathbf{J}}(0) = j$, and $\eta: 1 \times (d+1) \rightarrow k$ is the map where given $\ell < d+1$, we have $\eta(0, \ell) = 0$ unless $\ell = \alpha+1$, in which case $\eta(0, \ell) = q$.

We need to check item 1 of Definition 5.4.2 in all three cases. It holds in case 1 since (q, j) is a free relation for $i < k^u$. In cases 2 and 3, suppose $\mathbf{B} \subseteq \mathbf{A}$ has $\min\{\mathbf{B}\} \geq m+1$ and for every $b \in \mathbf{B}$ we have $c(b)|_{m+1} = t_{\alpha+1}$. Forming $\mathbf{C} := (\mathbf{B}/\{t_{\alpha+1}\})[\gamma\langle i, q, j \rangle]$, we have $\mathbf{C} \in \mathcal{K}$ since $\mathcal{K}(t_{\alpha+1}) \subseteq \mathcal{K}(\gamma\langle i, q, j \rangle)$; in case 2, we have $\mathcal{K}(\gamma\langle i, q, j \rangle) = \mathcal{K}(s_\alpha)$, and in case 3, we have $\mathcal{K}(\gamma\langle i, q, j \rangle) = \mathcal{B}^{\iota_\alpha}$.

So running the construction of f as in Definition 5.4.2, \mathbf{Y} is the \mathcal{L} -structure on $\mathbf{K} \cup \{v_0\}$ where $U^{\mathbf{Y}}(v_0) = j$ and $R^{\mathbf{Y}}(a, v_0) = q$ for any $a \in \mathbf{A}$ with $a \geq m+1$. We note that f , being an insertion with parameters m and $m+1$, is the identity on $\{0, \dots, m-1\}$.

We discuss how to extend Lemma 5.4.3, Definition 5.4.5, and Proposition 5.4.6 to deal with level m ; the discussion is the same in all three cases. The only situation not covered by Lemma 5.4.3 is if we have $a, b \in \mathbf{A}$ with $\ell(c(a) \wedge c(b)) = m$. The only way this can happen is if $c(a)|_{m+1} = t_\alpha$ and $c(b)|_{m+1} = t_{\alpha+1}$, or vice versa. The construction of \mathbf{Y} in this setting shows that we will have $\ell(c(f(a)) \wedge c(f(b))) = f(v_0)$. One can then extend the domain of \hat{f} to $c[\mathbf{A}]|_m$; given $t \in c[\mathbf{A}]|_m$, choose $a \in \mathbf{A}$ with $c(a)|_m = t$, and set $\hat{f}(t) = c(f(a))|_{f(v_0)}$. This does not depend on the choice of $a \in \mathbf{A}$. Then $\hat{f}: \text{Body}(\mathbf{A}) \rightarrow T$ respects relative levels, tree structure, and passing numbers except at level m (i.e., we might have $\hat{f}(t)(\ell(\hat{f}(u))) \neq t(\ell(u))$ when $\ell(u) = m$). To recover the result of Proposition 5.4.6 that $f[\mathbf{A}] \subseteq \mathbf{K}$ is weakly diagonal, items 1, 2, 5, and 6 are clear; we need to check items 3 and 4 of Definition 5.2.2 at level $f(v_0)$, but this follows from the choice of gluing γ .

Now consider some $y \in [m, f(m))$. If $y \neq f(v_0)$, then for every $a \in \mathbf{A}$ with $a > m$, we have $c(f(a))(y) = 0$. It follows that $\text{Crit}(f[\mathbf{A}]) \cap [m, f(m)) = \{f(v_0)\}$. Verifying that $f(v_0)$ is a controlled splitting level of $f[\mathbf{A}]$ is routine in cases 1 and 2. In case 3, we note that $\mathcal{K}(c[f[\mathbf{A}]]|_{f(v_0)}) = \mathcal{A}$ and $\mathcal{K}(c[f[\mathbf{A}]]|_{f(v_0)+1})^{\epsilon_{\alpha,d+1}} = \mathcal{B}$, and the remaining verifications in case 3 are straightforward.

Now assume that $\mathbf{A} \subseteq \mathbf{K}$ has controlled paths. First, we show that for any $n \geq m+1$, we have $\mathcal{K}(\hat{f}[c[\mathbf{A}]|_n]) = \mathcal{K}(c[\mathbf{A}]|_n)$. This follows from item 3 of Proposition 5.4.7. Set $r = m+1$, keeping in mind that $m+1 = M$ is the top parameter of the insertion f . Then $e: d+1 \rightarrow d+1$ is just the identity map ($d+1$ plays the same role here as d does in Proposition 5.4.7), and $\gamma^e = \gamma$. Then $\mathcal{K}(\gamma) \supseteq \mathcal{K}(c[\mathbf{A}]|_{m+1})$ in all three cases. Now we use Corollary 5.4.8 to conclude that for any $a \in \mathbf{A}$, we have $\text{Path}(a) = \text{Path}(f(a))$.

To finish the proof that $f[\mathbf{A}] \subseteq \mathbf{K}$ has controlled paths, suppose $a \in \mathbf{A}$. If $\text{Start}(f(a)) \neq f(v_0)$, then we have $\text{Start}(f(a)) = f(\text{Start}(a))$. If $\text{Start}(f(a)) = f(v_0)$, then we must have $\text{Start}(a) = m$. Lastly, if $\text{Start}(a) = m$, then we have $\text{Start}(c(a)) = f(v_0)$. These observations are enough to show that for any $a, b \in$

\mathbf{A} , we have $\text{Start}(a) = \text{Start}(b)$ iff $\text{Start}(f(a)) = \text{Start}(f(b))$. This is enough to conclude the three items of Definition 5.3.4. \square

5.8 Diagonal substructures

We are at last ready to define and construct a diagonal substructure. As a shorthand, let us say that $\mathbf{A} \subseteq \mathbf{K}$ *has controlled critical levels* if $\mathbf{A} \subseteq \mathbf{K}$ has controlled age-change levels, coding levels, and splitting levels (Definitions 5.5.1, 5.6.5, and 5.7.1). We point out explicitly the following properties of substructures with controlled critical levels.

Lemma 5.8.1. *Suppose $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal and has controlled critical levels.*

1. *Then $\mathbf{A} \subseteq \mathbf{K}$ has maximal paths.*
2. *If $\mathbf{B} \subseteq \mathbf{A}$, then $\mathbf{B} \subseteq \mathbf{K}$ is weakly diagonal and has controlled critical levels.*

Proof. The first item is immediate since $\mathbf{A} \subseteq \mathbf{K}$ has controlled coding nodes and controlled age-change levels. The second follows from Lemma 5.6.8 and Corollary 5.7.5. \square

Definition 5.8.2. A substructure $\mathbf{A} \subseteq \mathbf{K}$ is *diagonal* if it is weakly diagonal (Definition 5.2.2), has controlled paths (Definition 5.3.4), and has controlled critical levels.

We note that if $\mathbf{A} \subseteq \mathbf{K}$ is diagonal and $\mathbf{B} \subseteq \mathbf{A}$, then $\mathbf{B} \subseteq \mathbf{K}$ is also diagonal.

The following is the last major construction that we need, which will allow us to combine the major results of the previous sections.

Proposition 5.8.3. *Fix $\mathbf{A} \subseteq \mathbf{K}$ weakly diagonal. Then there is $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ so that $f[\mathbf{A}] \subseteq \mathbf{K}$ is weakly diagonal and has controlled critical levels.*

Furthermore, suppose $\mathbf{A} \subseteq \mathbf{K}$ has controlled paths and has every member of $\text{FreeStart}(\mathbf{A}) \cap \text{Crit}(\mathbf{A})$ controlled. Then there is $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ with $f[\mathbf{A}] \subseteq \mathbf{K}$ diagonal.

Proof. Set $\mathbf{A} = \mathbf{A}^0$. If $\mathbf{A}^0, \dots, \mathbf{A}^{n-1}$ and $f_1, \dots, f_{n-1} \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ have been defined, then let $m_{n-1} = \min\{m \in \text{Crit}(\mathbf{A}^{n-1}) : m \text{ is not controlled}\}$. Depending on whether m_{n-1} is in \mathbf{A}^{n-1} , $\text{Sp}(\mathbf{A}^{n-1})$, or $\text{AC}(\mathbf{A}^{n-1})$, apply Proposition 5.5.3, 5.6.9, or 5.7.6 to find $f_n \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ as guaranteed by the respective proposition, then set $\mathbf{A}^n = f_n[\mathbf{A}^{n-1}]$. Write $g_0 = \text{id}$ and $g_n = f_n \circ \dots \circ f_1$, and note that $g_n[\mathbf{A}] = \mathbf{A}^n$. We will set $f = \lim_n g_n$; if we can argue that this definition makes sense, then $f[\mathbf{A}] \subseteq \mathbf{K}$ will be weakly diagonal and have controlled critical levels. It is enough to argue that for $a := \min\{\mathbf{A}\}$, the sequence $\lim_n g_n(a)$ eventually stabilizes.

First we note that the sequence $(m_n)_n$ is non-decreasing, so if $n, r < \omega$ with $r < m_n$, then $f_N(r) = r$ for any $N > n$. So we want to show that for n large enough, we have $g_n(a) < m_n$. Towards a contradiction, suppose this was not the case.

Write $|\text{Sp}(\mathbf{A}) \cap [0, a]| = \alpha$; then for every $n < \omega$, we also have $|\text{Sp}(g_n[\mathbf{A}]) \cap [0, g_n(a)]| = \alpha$. We first argue that for some n , we have $|\text{Sp}(\mathbf{A}^n) \cap [m_n, g_n(a)]| = 0$.

We note that $|\mathrm{Sp}(\mathbf{A}^n) \cap [0, m_n]|$ is non-decreasing, and if $m_n \in \mathrm{Sp}(\mathbf{A}^n)$, then $|\mathrm{Sp}(\mathbf{A}^{n+1}) \cap [0, m_{n+1}]| = |\mathrm{Sp}(\mathbf{A}^n) \cap [0, m_n]| + 1$. So suppose for some n that $m_n \in \mathrm{AC}(\mathbf{A}^n)$, and suppose $r > m_n$ is the least member of $\mathrm{Sp}(\mathbf{A}^n)$ above. Towards a contradiction, suppose for every $N > n$ that $m_N \in \mathrm{AC}(\mathbf{A}^N)$ and $f_N \circ \cdots \circ f_{n+1}(r) > m_n$. By Lemma 5.4.3, $f_N \circ \cdots \circ f_{n+1}(r) \in \mathrm{Sp}(\mathbf{A}^{n+1})$ is the least splitting node above m_N . Writing $\rho = c[\mathbf{A}^n]_{m_n}^u$, we have for every $N \geq n$ that $\rho = c[\mathbf{A}^N]_{m_N}^u$. By Proposition 5.5.3, we have for every $N \geq n$ that $\mathcal{K}(c[\mathbf{A}^{N+1}]_{m_{N+1}}) \subsetneq \mathcal{K}(c[\mathbf{A}^N]_{m_N})$, contradicting that $P(\rho)$ is finite.

So now assume we have n with $|\mathrm{Sp}(\mathbf{A}^n) \cap [m_n, g_n(a)]| = 0$. Write $\rho = c[\mathbf{A}^n]_{m_n}$. Arguing as in the previous paragraph, we cannot have $m_N \in \mathrm{AC}(\mathbf{A}^N)$ for every $N \geq n$. So we can find $N \geq n$ with $m_N \in \mathrm{Sp}(\mathbf{A}^N)$, in which case $m_N > g_N(a)$ and we are done, or we have $m_N \in \mathbf{A}^N$, and we possibly have $m_N = g_N(a)$. Here we need to be a bit careful, since the application of Proposition 5.6.9 can introduce new age-change levels between $g_N(a) - 1$ and $g_{N+1}(a)$, and controlling these age-changes can destroy our control of the coding level. But if $m_N = g_N(a)$, $m_{N+1} \in \mathrm{AC}(\mathbf{A}^{N+1})$ and $m_{N+1} < g_{N+1}(a)$, we will still have $c[\mathbf{A}^{N+1}]_{m_{N+1}} = \rho$, and we have

$$\mathcal{K}(c[\mathbf{A}^N]_{m_N}) = \mathcal{K}(c[\mathbf{A}^{N+1}]_{m_N}) = \mathcal{K}(c[\mathbf{A}^{N+1}]_{m_{N+1}}).$$

Furthermore, if $N > n$, we will have $\mathcal{K}(c[\mathbf{A}^{N-1}]_{m_{N-1}}) \supsetneq \mathcal{K}(c[\mathbf{A}^N]_{m_N})$. So even though we may bounce between the cases $m_N \in \mathrm{AC}(\mathbf{A}^N)$ and $m_N = g_N(a)$ for a while, the finiteness of $P(\rho)$ prevents this from continuing indefinitely.

Now assume that $\mathbf{A} \subseteq \mathbf{K}$ has controlled paths and that every member of $\mathrm{FreeStart}(\mathbf{A}) \cap \mathrm{Crit}(\mathbf{A})$ is controlled. Then every application of Proposition 5.5.3, 5.6.9, or 5.7.6 preserves the fact that we have controlled paths, and the limit map will preserve this as well. \square

We need one last observation before completing our construction.

Lemma 5.8.4. *In the setting of Proposition 5.3.5, the $f \in \mathrm{OEmb}(\mathbf{K}, \mathbf{K})$ we construct has the property that every level in $\mathrm{FreeStart}(f[\mathbf{A}]) \cap \mathrm{Crit}(f[\mathbf{A}])$ is a controlled splitting level.*

Proof. Fix $m \in \mathrm{Crit}(f[\mathbf{A}]) \cap \mathrm{FreeStart}(f[\mathbf{A}])$. By the last paragraph of the proof of Proposition 5.3.5, we can find $i \in \mathrm{Free}$ and $\mathfrak{p} \in \max(\vec{P}_i)$ with $m = f(u_{\mathfrak{p}})$. Then writing $\mathrm{Free}(i) = (q, j)$, we have $U(m) = U(f(u_{\mathfrak{p}})) = U^{\mathbf{Y}}(u_{\mathfrak{p}}) = j$, and for any $y \in \mathbf{Y}$ with $R^{\mathbf{Y}}(y, u_{\mathfrak{p}}) \neq 0$, we have $R^{\mathbf{Y}}(y, u_{\mathfrak{p}}) = q$ and $R^{\mathbf{Y}}(y, u_{\mathfrak{q}}) = 0$ for any $u_{\mathfrak{q}} \in \mathbf{Y}$ previously enumerated by g . This shows that for any $y_0, y_1 \in \mathbf{Y}$ with $R^{\mathbf{Y}}(y_0, u_{\mathfrak{p}}) = R^{\mathbf{Y}}(y_1, u_{\mathfrak{p}}) = q$, then we have $y_0|_{f(u_{\mathfrak{p}})} = y_1|_{f(u_{\mathfrak{p}})}$. It follows that m is a controlled splitting level of $f[\mathbf{A}]$. \square

Theorem 5.8.5. *There is $f \in \mathrm{OEmb}(\mathbf{K}, \mathbf{K})$ with $f[\mathbf{K}] \subseteq \mathbf{K}$ diagonal.*

Proof. First use Proposition 5.2.4 to find $f_0 \in \mathrm{OEmb}(\mathbf{K}, \mathbf{K})$ with $f_0[\mathbf{K}]$ weakly diagonal. Then use Proposition 5.8.3 to find $f_1 \in \mathrm{OEmb}(\mathbf{K}, \mathbf{K})$ so that $f_1 \circ f_0[\mathbf{K}]$ is weakly diagonal and has controlled critical levels. By Lemma 5.8.1, $f_1 \circ f_0[\mathbf{K}]$ has maximal paths, so use Proposition 5.3.5 to find $f_2 \in \mathrm{OEmb}(\mathbf{K}, \mathbf{K})$ so that $f_2 \circ f_1 \circ f_0[\mathbf{K}]$ is weakly diagonal, has controlled paths, and satisfies the conclusion of Lemma 5.8.4. Then use Proposition 5.8.3 one more time to find $f_3 \in \mathrm{OEmb}(\mathbf{K}, \mathbf{K})$ with $f_3 \circ f_2 \circ f_1 \circ f_0[\mathbf{K}]$ diagonal. We then set $f = f_3 \circ f_2 \circ f_1 \circ f_0$. \square

6 Diaries

Having defined and constructed diagonal substructures in Section 5, we still need to show how to apply Theorem 5.1.7 to obtain finite upper bounds for the big Ramsey degrees of members of \mathcal{K} . In order to do this, given a diagonal $\mathbf{A} \subseteq \mathbf{K}$, we need a precise understanding of the closure $\overline{\mathbf{A}}$. In the process of doing this, we introduce the notion of a *diary*, a fragment of the information contained in $\overline{\mathbf{A}}$ which in the case that $\mathbf{A} \subseteq \mathbf{K}$ is diagonal, is enough to reconstruct all of $\overline{\mathbf{A}}$. We then turn to lower bounds, defining a notion of *diagonal diary*. We then show that given $\mathbf{A} \in \mathcal{K}$, every diagonal diary for \mathbf{A} appears in every diagonal copy of \mathbf{K} inside \mathbf{K} . Thus the ordered big Ramsey degree of an enumerated $\mathbf{A} \in \mathcal{K}$ is precisely the number of diagonal diaries which code \mathbf{A} .

6.1 Concrete diaries

We begin by defining *concrete diaries*, where given a weakly diagonal $\mathbf{A} \subseteq \mathbf{K}$, we create a function $\Delta^{\mathbf{A}}$ which codes certain limited information about $\overline{\mathbf{A}}$.

Definition 6.1.1. Let Label denote the set $\{\text{Cd}, \text{AC}, \text{Sp}\}$, and define

$$\Omega = \bigcup \left\{ \left(\text{Label} \times P(\rho) \times d \times k^{d-1} \right) : d < \omega, \rho: d \rightarrow k^u \text{ non-decreasing} \right\}.$$

Given a tuple $\alpha = (\mathbf{X}, \mathcal{A}, i, f) \in \Omega$, we write $\text{Lab}(\alpha) = \mathbf{X}$, $\text{Cl}(\alpha) = \mathcal{A}$, $\text{Pos}(\alpha) = i$, and $\text{Pass}(\alpha) = f$, the *label*, *class*, *position*, and *passing numbers* of α for the respective 4 coordinates, and we write $\text{Sort}(\alpha) = \rho$ and $\text{Size}(\alpha) = d$. Note that $\text{Size}(\alpha) = \text{dom}(\text{Sort}(\alpha))$.

Suppose $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal. The *diary* of \mathbf{A} is the function

$$\Delta^{\mathbf{A}}: |\text{Crit}(\mathbf{A})| \rightarrow \Omega$$

defined as follows. Suppose $g: |\text{Crit}(\mathbf{A})| \rightarrow \text{Crit}(\mathbf{A})$ is the increasing bijection, and fix $n < |\text{Crit}(\mathbf{A})|$ with $g(n) = m \in \text{Crit}(\mathbf{A})$.

- $\text{Lab}(\Delta^{\mathbf{A}}(n))$ indicates whether m is a coding, age-change, or splitting level of $\text{Crit}(\mathbf{A})$.
- We set $\text{Cl}(\Delta^{\mathbf{A}}(n)) = \mathcal{K}(c[\mathbf{A}]|_m)$. In particular, we have $\text{Sort}(\Delta^{\mathbf{A}}(n)) = c[\mathbf{A}]|_m^u$. For the remainder of the definition, write $\text{Sort}(\Delta^{\mathbf{A}}(n)) = \rho$ and $\text{dom}(\rho) = d$.
- If $\text{Lab}(\Delta^{\mathbf{A}}(n)) = \text{AC}$, we don't care about the position coordinate, and simply declare that $\text{Pos}(\Delta^{\mathbf{A}}(n)) = 0$. If $\text{Lab}(\Delta^{\mathbf{A}}(n)) \in \{\text{Cd}, \text{Sp}\}$, then $\text{Pos}(\Delta^{\mathbf{A}}(n)) < d$ indicates which node in $c[\mathbf{A}]|_m$ in lexicographic order is the coding or splitting node.
- If $\text{Lab}(\Delta^{\mathbf{A}}(n)) \in \{\text{AC}, \text{Sp}\}$, then we do not care about the passing numbers coordinate, and simply declare that $\text{Pass}(\Delta^{\mathbf{A}}(n))$ is the constant zero function. If $\text{Lab}(\Delta^{\mathbf{A}}(n)) = \text{Cd}$, then write $c[\mathbf{A}]|_{m+1} = \{t_0 \preceq_{\text{lex}} \cdots \preceq_{\text{lex}} t_{d-2}\}$. Then $\text{Pass}(\Delta^{\mathbf{A}}(n))$ is the function given by $\text{Pass}(\Delta^{\mathbf{A}}(n))(j) = t_j(m)$ for each $j < d - 1$.

One can think of the diary of $\mathbf{A} \subseteq \mathbf{K}$ as giving some very basic information about the envelope of \mathbf{A} . Certainly, given $\tilde{\mathbf{A}}$ (Definition 5.1.3), one can compute

$\Delta^{\mathbf{A}}$. The main result of this subsection is that if $\mathbf{A} \subseteq \mathbf{K}$ is diagonal, then we can in fact reconstruct $\tilde{\mathbf{A}}$ just from $\Delta^{\mathbf{A}}$. We state the main consequences of this construction in the next proposition.

Proposition 6.1.2. *Suppose $\mathbf{A}, \mathbf{B} \subseteq \mathbf{K}$ are diagonal substructures with $\Delta^{\mathbf{A}} = \Delta^{\mathbf{B}}$. Then we have $\tilde{\mathbf{A}} = \tilde{\mathbf{B}}$. Furthermore, we have $\mathbf{A} = \text{Int}(\overline{\mathbf{A}})$.*

Proof. The proposition will be proven by starting with a diagonal $\mathbf{A} \subseteq \mathbf{K}$ and constructing $\overline{\mathbf{A}}$ using only the information from $\Delta^{\mathbf{A}}$ along with the knowledge that $\mathbf{A} \subseteq \mathbf{K}$ is diagonal. The “furthermore” will follow from analyzing the construction.

The main difficulty is to decide what is needed beyond $\text{Crit}(\mathbf{A})$ to form the closure $\overline{\mathbf{A}}$. There are two main sources of members of $\overline{\mathbf{A}} \setminus \text{Crit}(\mathbf{A})$. First, whenever $m \in \text{AC}(\mathbf{A})$ is a controlled age change level, $\text{Cause}(\mathcal{K}(c[\mathbf{A}]|_m), \mathcal{K}(c[\mathbf{A}]|_{m+1})) = (\rho, \eta, \mathbf{I})$, and $\varphi \in \text{OL}(\mathbf{I}, \mathbf{K})$ is as in Definition 5.5.1, then we will show that $\varphi[\mathbf{I}] \subseteq \overline{\mathbf{A}}$. Second, there are some members of $\text{FreeStart}(\mathbf{A})$ in $\overline{\mathbf{A}} \setminus \text{Crit}(\mathbf{A})$. We now discuss this second source of new points in some detail.

Let us define

$$\text{Path}(\mathbf{A}, i) = \{\text{Path}(a) : a \in \mathbf{A} \text{ with } U(a) = i\} \subseteq \max(\vec{P}_i).$$

Note that $\text{Path}(\mathbf{A}, i) \neq \emptyset$ iff there is some $a \in \mathbf{A}$ with $U(a) = i$. For each $i \in \text{Free}$ and each $\mathfrak{p} \in \text{Path}(\mathbf{A}, i)$, let $m_{\mathfrak{p}} = \text{Start}(a)$ for some (equivalently any, since $\mathbf{A} \subseteq \mathbf{K}$ has controlled paths) $a \in \mathbf{A}$ with $\text{Path}(a) = \mathfrak{p}$. Also, for each $i \in \text{Free}$ with $\text{Path}(\mathbf{A}, i) \neq \emptyset$, let \mathfrak{p}_i denote the \leq_{path} -largest member of $\text{Path}(\mathbf{A}, i)$. We note that $m_{\mathfrak{p}_i} \notin \text{Crit}(\mathbf{A})$, and may or may not be a member of $\overline{\mathbf{A}}$. For every other $\mathfrak{p} \in \text{Path}(\mathbf{A}, i)$, we have that $m_{\mathfrak{p}} \in \text{Sp}(\mathbf{A})$. So we need to decide for which $i \in \text{Free}$ we will have $m_{\mathfrak{p}_i} \in \overline{\mathbf{A}}$.

To do this, we define the set $X^{\mathbf{A}} \subseteq \text{Free}$ via the following algorithm. The algorithm considers each $i \in \text{Free}$ in decreasing order, and adds i to $X^{\mathbf{A}}$ if at least one of the following holds

- There is some $m \in \text{Sp}(\mathbf{A})$ with $m > m_{\mathfrak{p}_i}$ and with $c(m) = (i, 0^m)$.
- There is some $m \in \text{AC}(\mathbf{A})$ such that, writing $\gamma = (\rho, \eta, \mathbf{I}) = \text{Cause}(\mathcal{K}(c[\mathbf{A}]|_m), \mathcal{K}(c[\mathbf{A}]|_{m+1}))$, there is some $x \in \mathbf{I}$ with $U^{\mathbf{I}}(x) = i$.
- There is some $j \in \text{Free}$ previously added to $X^{\mathbf{A}}$ by the algorithm with $\text{Free}(j)^u = i$

Since $\mathbf{A} \subseteq \mathbf{K}$ is diagonal, we can detect the above events using only the diary of \mathbf{A} . To rephrase the first event to avoid the mention of $m_{\mathfrak{p}_i}$, we note that the demand $m > m_{\mathfrak{p}_i}$ is equivalent to asking that either $\text{Sp}^{\mathbf{A}}(m) \in T_j$ for some $i < j < k^u$ with a free relation, or that $\text{Sp}^{\mathbf{A}}(m) \in T_j$ for any $j < k^u$ with no free relation, or that whenever $a, b \in \mathbf{A}$ have $c(a) \wedge c(b) = \text{Sp}^{\mathbf{A}}(m)$, we have $\text{Path}(a) = \text{Path}(b)$.

For each $m \in \text{AC}(\mathbf{A})$, let us write

$$\gamma^{(m)} = (\rho^{(m)}, \eta^{(m)}, \mathbf{I}^{(m)}) = \text{Cause}(\mathcal{K}(c[\mathbf{A}]|_m), \mathcal{K}(c[\mathbf{A}]|_{m+1})),$$

and let $\varphi^{(m)} \in \text{OL}(\mathbf{I}^{(m)}, \mathbf{K})$ be as in Definition 5.5.1. We now define

$$\mathbf{C} := \mathbf{A} \cup \text{Sp}(\mathbf{A}) \cup \{m_{\mathfrak{p}_i} : i \in X^{\mathbf{A}}\} \cup \bigcup \{\varphi^{(m)}[\mathbf{I}^{(m)}] : m \in \text{AC}(\mathbf{A})\}.$$

Note that $\text{AC}(\mathbf{A})$ is a subset of the last component. We will show that $\mathbf{C} = \overline{\mathbf{A}}$. Let us emphasize that an enumerated structure isomorphic to \mathbf{C} can be constructed by using only the information from $\Delta^{\mathbf{A}}$ and the fact that $\mathbf{A} \subseteq \mathbf{K}$ is diagonal.

First we argue that $\mathbf{C} \subseteq \mathbf{K}$ is closed. As a first observation, given $m \in \text{AC}(\mathbf{A})$ and $x \in \varphi^{(m)}[\mathbf{I}^{(m)}]$, then since $\varphi^{(m)} \in \text{OL}(\mathbf{I}^{(m)}, \mathbf{K})$, then either $x = (x^u, 0^{\ell(x)})$ or we have $\text{Start}(x) \in \varphi^{(m)}[\mathbf{I}^{(m)}]$. Towards showing that $\text{Sp}(\mathbf{C}) \subseteq \mathbf{C}$, let $a, b \in \mathbf{C}$ with $U(a) = U(b) = i < k^u$. We may assume throughout that $c(a) \perp c(b)$. There are several straightforward cases to discuss:

- Suppose $a \in \mathbf{A}$.
 - If $b \in \mathbf{A}$, then $\ell(c(a) \wedge c(b)) \in \text{Sp}(\mathbf{A})$.
 - If $b \in \text{Sp}(\mathbf{A})$, then $c(b) = (i, 0^b)$, so $\ell(c(a) \wedge c(b)) = \text{Start}(a)$. If $i \in k^u \setminus \text{Free}$, then we always have $\text{Start}(a) \in \text{Crit}(\mathbf{A})$ for any $a \in \mathbf{A}$. If $i \in \text{Free}$, then either $\text{Path}(a) < \mathfrak{p}_i$, in which case $\text{Start}(a) \in \text{Sp}(\mathbf{A})$, or $\text{Path}(a) = \mathfrak{p}_i$, in which case $\text{Start}(a) = m_{\mathfrak{p}_i}$, and $b > m_{\mathfrak{p}_i}$ will witness that $i \in X^{\mathbf{A}}$.
 - If $b = m_{\mathfrak{p}_j}$ for some $j \in X^{\mathbf{A}}$, then we must have $i = \text{Free}(j)^u$ and $c(b) = (i, 0^b)$. So $\ell(c(a) \wedge c(b)) = \text{Start}(a)$. Either $\text{Start}(a) \in \text{Crit}(a)$, or we have $\text{Start}(a) = m_{\mathfrak{p}_i}$. In this case, we must have $i < j$, so $j \in X^{\mathbf{A}}$ will witness that $i \in X^{\mathbf{A}}$.
 - If for some $m \in \text{AC}(\mathbf{A})$ we have $b \in \varphi^{(m)}[\mathbf{I}^{(m)}]$, let us first note that $\text{Start}(a) \neq \text{Start}(b)$, since $\text{Start}(b) \in \varphi^{(m)}[\mathbf{I}^{(m)}]$ and $\text{Start}(a) \in \text{Crit}^+(\mathbf{A})$. It follows that $\ell(c(a) \wedge c(b)) = \min\{\text{Start}(a), \text{Start}(b)\}$. If $\ell(c(a) \wedge c(b)) = \text{Start}(a)$, then either $\text{Start}(a) \in \text{Crit}(\mathbf{A})$ or $\text{Start}(a) = m_{\mathfrak{p}_i}$, and in this case $b \in \varphi^{(m)}[\mathbf{I}^{(m)}]$ will witness that $i \in X^{\mathbf{A}}$. If $\ell(c(a) \wedge c(b)) = \text{Start}(b)$, then $\text{Start}(b) \in \varphi^{(m)}[\mathbf{I}^{(m)}]$.
- Suppose $a \in \text{Sp}(\mathbf{A})$. Then we must have $c(a) = (i, 0^a)$. So whenever $b \in \mathbf{C}$ satisfies $c(a) \perp c(b)$, we must have $\ell(c(a) \wedge c(b)) = \text{Start}(b)$.
 - If $b \in \text{Sp}(\mathbf{A})$, then $c(b) = (i, 0^b)$, so $c(a) \not\perp c(b)$, contradicting our assumption.
 - If $b = m_{\mathfrak{p}_j}$ for some $j \in \text{FreeStart}(\mathbf{A})$, then $i = \text{Free}(j)^u$ and $c(b) = (i, 0^b)$, so $c(a) \not\perp c(b)$, contradicting our assumption.
 - If for some $m \in \text{AC}(\mathbf{A})$ we have $b \in \varphi^{(m)}[\mathbf{I}^{(m)}]$, then also $\text{Start}(b) \in \varphi^{(m)}[\mathbf{I}^{(m)}]$.
- Suppose $a = m_{\mathfrak{p}_j}$ for some $j \in X^{\mathbf{A}}$. Then we must have $i = \text{Free}(j)^u$ and $c(a) = (i, 0^a)$. So whenever $c(a) \perp c(b)$, we must have $\ell(c(a) \wedge c(b)) = \text{Start}(b)$.
 - If $b = m_{\mathfrak{p}_\ell}$ for some $\ell \in X^{\mathbf{A}}$, then we must have $i = \text{Free}(\ell)^u$ and $c(b) = (i, 0^b)$, so $c(a) \not\perp c(b)$, contradicting our assumption.
 - If for some $m \in \text{AC}(\mathbf{A})$ we have $b \in \varphi^{(m)}[\mathbf{I}^{(m)}]$, then also $\text{Start}(b) \in \varphi^{(m)}[\mathbf{I}^{(m)}]$.
- Suppose there are $m, n \in \text{AC}(\mathbf{A})$ with $a \in \varphi^{(m)}[\mathbf{I}^{(m)}]$ and $b \in \varphi^{(n)}[\mathbf{I}^{(n)}]$. If $m = n$, then $\ell(c(a) \wedge c(b)) \in \varphi^{(m)}[\mathbf{I}^{(m)}]$ since $\varphi^{(m)} \in \text{OL}(\mathbf{I}^{(m)}, \mathbf{K})$. If $m < n$, then $\ell(c(a) \wedge c(b)) = \text{Start}(a) \in \varphi^{(m)}[\mathbf{I}^{(m)}]$.

Now we need to argue that $\text{AC}(\mathbf{C}) = \emptyset$. So fix $m \notin \mathbf{C} \cup \text{Sp}(\mathbf{C}) = \mathbf{C}$. There are two main cases to consider.

- If there is no $n \in \text{AC}(\mathbf{A})$ with $m \in [\min(\varphi^{(n)}[\mathbf{I}^{(n)}]), \max(\varphi^{(n)}[\mathbf{I}^{(n)}])]$, then $c[\mathbf{C}]|_m \setminus c[\mathbf{A}]|_m \subseteq \{(i, 0^m) : i < k^u\}$ and $c[\mathbf{C}]|_{m+1} \setminus c[\mathbf{A}]|_{m+1} \subseteq \{(i, 0^{m+1}) : i < k^u\}$. Since $m \notin \text{AC}(\mathbf{A})$, it follows that $m \notin \text{AC}(\mathbf{C})$.
- If for some (unique) $n \in \text{AC}(\mathbf{A})$ we have $m \in [\min(\varphi^{(n)}[\mathbf{I}^{(n)}]), \max(\varphi^{(n)}[\mathbf{I}^{(n)}])]$, then also $c[\varphi^{(n)}[\mathbf{I}^{(n)}]]|_m \subseteq c[\mathbf{C}]|_m$. However, since $m \notin \mathbf{C}$, also $m \notin \varphi^{(n)}[\mathbf{I}^{(n)}]$. By the properties of $\varphi^{(n)}$ given in Definition 5.5.1, we have for any $t \in c[\varphi^{(n)}[\mathbf{I}^{(n)}]]|_{m+1}$ that $t(m) = 0$. Thus, since $m \notin \text{AC}(\mathbf{A})$, we also have that $m \notin \text{AC}(\mathbf{C})$.

Having argued that \mathbf{C} is closed, we now need to argue that \mathbf{C} is in fact the smallest closed substructure of \mathbf{K} containing \mathbf{A} . This will be clear once we argue that $\varphi^{(m)}[\mathbf{I}^{(m)}] \subseteq \overline{\mathbf{A}}$ whenever $m \in \text{AC}(\mathbf{A})$. Write $\mathbf{J} = (\varphi^{(m)})^{-1}(\overline{\mathbf{A}} \cap \varphi^{(m)}[\mathbf{I}^{(m)}])$. If $\mathbf{J} \subsetneq \mathbf{I}^{(m)}$, then upon restricting the gluing $\gamma^{(m)}$ to \mathbf{J} , we obtain a strictly smaller gluing which also results in the age change. As $\gamma^{(m)} = \text{Cause}(\mathcal{K}(c[\mathbf{A}]|_m), \mathcal{K}(c[\mathbf{A}]|_{m+1}))$ is chosen so that $\mathbf{I}^{(m)}$ has minimum possible cardinality, this is a contradiction, and we must have $\mathbf{J} = \mathbf{I}^{(m)}$.

We now briefly discuss the “furthermore” of Proposition 6.1.2. So suppose $\mathbf{A}' \subsetneq \mathbf{A}$, and fix some $a \in \mathbf{A} \setminus \mathbf{A}'$. Then $\mathbf{A}' \subseteq \mathbf{K}$ is also diagonal, so we can use the above construction to build $\overline{\mathbf{A}'}$, and we simply note that $a \notin \overline{\mathbf{A}'}$. \square

This concludes the construction of $\tilde{\mathbf{A}}$ using only $\Delta^{\mathbf{A}}$ and the knowledge that $\mathbf{A} \subseteq \mathbf{K}$ is diagonal. We are now in a position to apply Theorem 5.1.7.

Corollary 6.1.3. *Fix an enumerated $\mathbf{A} \in \mathcal{K}$, and fix $f \in \text{OEmb}(\mathbf{K}, \mathbf{K})$ with $f[\mathbf{K}] \subseteq \mathbf{K}$ diagonal. Then the ordered big Ramsey degree of \mathbf{A} in \mathcal{K} is at most $|\{\Delta^{g[\mathbf{A}]} : g \in \text{OEmb}(\mathbf{A}, f[\mathbf{K}])\}|$. In words, it is at most the number of diaries of ordered copies of \mathbf{A} inside $f[\mathbf{K}]$.*

6.2 Diagonal Diaries

While it is tempting to simply define a “diagonal diary” as a concrete diary of the form $\Delta^{\mathbf{A}}$ for some diagonal $\mathbf{A} \subseteq \mathbf{K}$, we will want a more constructive definition, so that in principle, one could write a computer program which takes as input a finite enumerated $\mathbf{A} \in \mathcal{K}$, then outputs the number of diagonal diaries which code \mathbf{A} . This leads to a rather abstract definition; every “concrete” diagonal diary will be a diagonal diary, but in principle there could be more “abstract” diagonal diaries than concrete ones. However, we will eventually show that every abstract diagonal diary is concrete, and that in any diagonal copy of \mathbf{K} inside \mathbf{K} , every diagonal diary appears as the diary of some further substructure. This will allow us to strengthen Corollary 6.1.3 into the exact result.

We now turn towards the definition of diagonal diary. The idea is to rephrase many of the properties of being a diagonal substructure in ways that can be detected by the diary.

Definition 6.2.1. Let Label , Ω , and other related notation have the same meanings as in Definition 6.1.1. Fix N with $1 \leq N \leq \omega$. A function $\Delta : N \rightarrow \Omega$ for some $1 \leq N$ is a *diary* if the following hold:

- $\text{Sort}(\Delta(0))$ is an injection.
- Suppose $n < \omega$ with $n + 1 < N$. Then:
 1. If $\text{Lab}(\Delta(n)) = \text{AC}$, we have $\text{Sort}(\Delta(n + 1)) = \text{Sort}(\Delta(n))$.
 2. If $\text{Lab}(\Delta(n)) = \text{Sp}$, then writing $j = \text{Pos}(\Delta(n))$, we have $\text{Sort}(\Delta(n + 1)) = \text{Sort}(\Delta(n)) \circ \delta_{j,d}$ (i.e., position j has split into two nodes).
 3. If $\text{Lab}(\Delta(n)) = \text{Cd}$, then writing $j = \text{Pos}(\Delta(n))$, we have $\text{Sort}(\Delta(n + 1)) = \text{Sort}(\Delta(n)) \circ \epsilon_{j,d}$ (i.e., position j is a terminal coding node).

Definition 6.2.2. Given a diary Δ of height $N \leq \omega$, the *tree of Δ* is a downwards closed $T^\Delta \subseteq T$ equipped with age-set structure and coding nodes which we will define. Given $n < \omega$ and $i < k^u$, we write $T^\Delta(n)$, T_i^Δ , etc. for the obvious subsets of T_Δ . We now define T^Δ inductively as follows:

- $T^\Delta(0) = \{\emptyset_i : i \in \text{Im}(\text{Sort}(\Delta(0)))\}$.
- For every $n < \omega$, we have $T^\Delta(n)^u = \text{Sort}(\Delta(n))$, and we assign $T^\Delta(n)$ the age $\text{Cl}(\Delta(n))$. If $\sigma : d \rightarrow T^\Delta(n)$ is any function, we set $\text{Cl}_\Delta(\sigma) = \text{Cl}(\Delta(n))^\sigma$. If $S \subseteq T^\Delta(n)$, we set $\text{Cl}_\Delta(S) := \text{Cl}_\Delta(\vec{S})$. If $S = \{t\}$, we simply write $\text{Cl}_\Delta(t)$ instead of $\text{Cl}_\Delta(\{t\})$.
- If $\text{Lab}(\Delta(n)) = \text{Sp}$, $\text{Size}(\Delta(n)) = d$, $\text{Pos}(\Delta(n)) = i$, and $T^\Delta(n) = \{t_0 \preceq_{lex} \cdots \preceq_{lex} t_{d-1}\}$, then $T^\Delta(n + 1) = \{t_j \widehat{\ } 0 : j < d\} \cup \{t_i \widehat{\ } 1\}$.
- If $\text{Lab}(\Delta(n)) = \text{AC}$, then $T^\Delta(n + 1) = \{t \widehat{\ } 0 : t \in T^\Delta\}$.
- If $\text{Lab}(\Delta(n)) = \text{Cd}$, $\text{Size}(\Delta(n)) = d$, $\text{Pos}(\Delta(n)) = i$, and $T^\Delta(n) = \{t_0 \preceq_{lex} \cdots \preceq_{lex} t_{d-1}\}$, then we declare that t_i is a *coding node* of T^Δ . We set

$$T^\Delta(n + 1) = \{(t_{\epsilon_{i,d}(j)}) \widehat{\ } \text{Pass}(\Delta(n))(j) : j < d - 1\}.$$

We write $\{c_\Delta(n) : n < |(\text{Lab} \circ \Delta)^{-1}(\text{Cd})|\}$ for the set of coding nodes of T^Δ ordered by increasing level, noting that unlike the coding function c , we will not have in general that $c_\Delta(n) \in T^\Delta(n)$. We write $\mathbf{Str}(\Delta)$ for the enumerated \mathcal{L} -structure coded by the coding nodes of T^Δ . More precisely, given $n < |\mathbf{Str}(\Delta)|$, we have $U^{\mathbf{Str}(\Delta)}(n) = c_\Delta(n)^u$, and given $m < n < |\mathbf{Str}(\Delta)|$, we have $R^{\mathbf{Str}(\Delta)}(n, m) = c_\Delta(n)(\ell(c_\Delta(m)))$.

If \mathbf{A} is an enumerated structure which embeds into \mathbf{K} , we say that a diary Δ *codes \mathbf{A}* if $\mathbf{Str}(\Delta) = \mathbf{A}$. Notice that if we start with a weakly diagonal $\mathbf{A} \subseteq \mathbf{K}$ and then form $\Delta^{\mathbf{A}}$ as in Definition 6.1.1, then $\mathbf{Str}(\Delta^{\mathbf{A}})$ is simply the enumerated structure which is order-isomorphic to \mathbf{A} .

Definition 6.2.3. A *diary with controlled critical levels* is a diary Δ which satisfies the following properties:

1. Coding nodes are dense in T^Δ .
2. For any $n < N$, if $\text{Lab}(\Delta(n)) = \text{AC}$, then $\text{Pos}(\Delta(n)) = 0$.
3. $\text{Pass}(\Delta(n))$ is the constant zero function unless $\text{Lab}(\Delta(n)) = \text{Cd}$.

4. (Controlled coding levels) Suppose we have $\text{Lab}(\Delta(n)) = \text{Cd}$, $\text{Sort}(\Delta(n)) = \rho$, $\text{Size}(\Delta(n)) = d$, $\text{Pos}(\Delta(n)) = j < d$, and $\rho(j) = i < k^u$. Write $\varphi := \text{Pass}(\Delta(n))$. We have that $\text{Cl}(\Delta(n+1)) = \text{Cl}(\Delta(n))\langle j, \epsilon_{j,d}, \varphi \rangle^{-1}$ and that $(\text{Cl}(\Delta(n)), j, \varphi)$ is a controlled coding triple.
5. (Controlled age-change levels) If $\text{Lab}(\Delta(n)) = \text{AC}$ and $\text{Sort}(\Delta(n)) = \rho$, then $(\text{Cl}(\Delta(n)), \text{Cl}(\Delta(n+1))) \in \text{Con}(\rho)$.
6. (Controlled splitting levels) Suppose we have $\text{Lab}(\Delta(n)) = \text{Sp}$, $\text{Sort}(\Delta(n)) = \rho$, $\text{Size}(\Delta(n)) = d$, $\text{Pos}(\Delta(n)) = \alpha$, and $\rho(\alpha) = i < k^u$. Then $\text{Cl}(\Delta(n+1)) = \text{Cl}(\Delta(n))^{\delta_{\alpha,d}}$ unless $i \in k^u \setminus \text{Free}$ and $\text{Cl}(\Delta(n))^{\iota_\alpha} = \mathcal{K}(\emptyset_i)$. In this case, we have $(\text{Cl}(\Delta(n)), \text{Cl}(\Delta(n+1))^{\epsilon_{\alpha,d+1}}) \in \text{Con}(\rho)$ and that $(\text{Cl}(\Delta(n))^{\iota_\alpha}, \text{Cl}(\Delta(n+1))^{\iota_{\alpha+1}}) \in \text{Con}(\iota_i)$.

Before defining diagonal diaries, let us take a few moments to establish some useful observations and definitions about diaries with controlled critical levels. Consider CT; the age-set structure of CT is a direct consequence of the location of the coding nodes in CT. By contrast, in a diary Δ , the age-set structure is completely arbitrary, and has no relation to where the coding nodes are and/or what structure they code. However, we will show that in a diary with controlled critical levels, we recover some correspondence between the locations of coding nodes and the age-set structure. In particular, we will show that the coding nodes at or below level m place an upper bound on what $\text{Cl}(\Delta(m))$ can be, while the coding nodes above level m place a lower bound on what $\text{Cl}(\Delta(m))$ can be.

Lemma 6.2.4. *Suppose $m \leq M < \omega$. Suppose $d < \omega$ and that $\sigma: d \rightarrow T^\Delta(M)$ is a function. Then we have $\text{Cl}_\Delta(\sigma) \subseteq \text{Cl}_\Delta(\langle \sigma(j)|_m : j < d \rangle)$.*

Proof. This is clear from the definition of controlled critical levels. \square

Proposition 6.2.5. *Suppose Δ is a diary with controlled critical levels. Suppose $1 \leq n \leq |\text{Str}(\Delta)|$, and fix any $m > L := \ell(c_\Delta(n-1))$. Let $X = \{x_0 \preceq_{\text{lex}} \cdots \preceq_{\text{lex}} x_{r-1}\} \subseteq T^\Delta(m)$. Let $\gamma = (X^u, \eta, \text{Str}(\Delta)|_n)$ be the gluing where $\eta: r \times n \rightarrow k$ is given by $\eta(\alpha, \ell) = x_\alpha(\ell(c_\Delta(\ell)))$ for each $\alpha < r$ and $\ell < n$. Then we have $\text{Cl}_\Delta(X) \subseteq \mathcal{K}(\gamma)$.*

Proof. We induct on n . By Lemma 6.2.4, we may assume that $m = L + 1$. In this case, it will be helpful to set $r = d - 1$, so that $X = \{x_0, \dots, x_{d-2}\}$. Write $X|_L \cup \{c_\Delta(n-1)\} := S = \{s_0 \preceq_{\text{lex}} \cdots \preceq_{\text{lex}} s_{d-1}\}$, and let $j < d$ be such that $c_\Delta(n-1) = s_j$. Let $\theta: d \times (n-1) \rightarrow k$ be given by $\theta(\alpha, \ell) = s_\alpha(\ell(c_\Delta(\ell)))$ for each $\alpha < d$ and $\ell < n-1$. Writing $\zeta = (S^u, \theta, \text{Str}(\Delta)|_{n-1})$, our inductive assumption gives us that $\text{Cl}_\Delta(S) \subseteq \mathcal{K}(\zeta)$. Write $\varphi: (d-1) \rightarrow k$ for the function given by $\varphi(\alpha) = y_\alpha(L)$. Then since L is a controlled coding level of Δ and by Lemma 6.2.4, we have that

$$\text{Cl}_\Delta(X) = \text{Cl}_\Delta(S)\langle j, \epsilon_{j,d}, \varphi \rangle^{-1} \subseteq \mathcal{K}(\zeta)\langle j, \epsilon_{j,d}, \varphi \rangle^{-1} \subseteq \mathcal{K}(\gamma).$$

To verify the last containment (which is actually an equality, but we only show what we need), suppose $\mathbf{B} \in \mathcal{K}(\zeta)\langle j, \epsilon_{j,d}, \varphi \rangle^{-1}$. This means that $\mathbf{B}\langle j, \epsilon_{j,d}, \varphi \rangle := \mathbf{C} \in \mathcal{K}(\zeta)$. This means that $\mathcal{C}[\zeta] \in \mathcal{K}$. Now we note that $\mathbf{C}[\zeta]$ and $\mathbf{B}[\gamma]$ are isomorphic; if we write $B \cup \{x\}$ for the underlying set of \mathbf{C} , where x is the new vertex, then the map sending x to $n-1$ is an isomorphism. \square

Proposition 6.2.6. *Suppose Δ is a diary with controlled critical levels. Fix $m < \omega$ and $X = \{x_0, \dots, x_{d-1}\} \subseteq T^\Delta(m)$. Suppose $\mathbf{A} \subseteq \mathbf{Str}(\Delta)$ is such that $\ell(c_\Delta(n)) \geq m$ for each $n \in \mathbf{A}$ and $c_\Delta[\mathbf{A}]|_m \subseteq X$. Let \mathbf{B} be the \mathcal{L}_d -structure on vertex set $\{v_n : n \in \mathbf{A}\}$ with \mathcal{L}^b part isomorphic to \mathbf{A} in the obvious way and so that $V^{\mathbf{B}}(v_n) = j$ iff $c_\Delta(n)|_m = x_j$. Then $\mathbf{B} \in \text{Cl}_\Delta(X)$.*

Proof. We induct on the size of \mathbf{A} . If $\mathbf{A} = \emptyset$, there is nothing to show, so assume $|\mathbf{A}| \geq 1$. We may assume that $c_\Delta[\mathbf{A}]|_m = X$, and by Lemma 6.2.4, we may suppose that $m = \ell(c_\Delta(\min(\mathbf{A})))$. Suppose $j < d$ is such that $x_j = c_\Delta(n)$. Write $Y = \{y_0, \dots, y_{d-2}\} \subseteq T^\Delta(m+1)$ for the set of immediate successors of members of X , and let $\varphi: (d-1) \rightarrow k$ be the function given by $\varphi(\alpha) = y_\alpha(m)$ for each $\alpha < d-1$. Write $\mathbf{A}' = \mathbf{A} \setminus \{\min(\mathbf{A})\}$, and define the \mathcal{L}_{d-1} -structure \mathbf{D} on vertex set $\{w_n : n \in \mathbf{A}'\}$ with \mathcal{L}^b -part isomorphic to \mathbf{A}' and with $V^{\mathbf{D}}(w_n) = \alpha$ iff $c_\Delta(n)|_{m+1} = y_\alpha$. By induction, we have $\mathbf{D} \in \text{Cl}_\Delta(Y)$. Since m is a controlled coding level of Δ , we have $\text{Cl}_\Delta(Y) = \text{Cl}_\Delta(X)\langle j, \epsilon_{j,d}, \varphi \rangle$. So $\mathbf{D}\langle j, \epsilon_{j,d}, \varphi \rangle \in \text{Cl}_\Delta(X)$, and $\mathbf{D}\langle j, \epsilon_{j,d}, \varphi \rangle \cong \mathbf{B}$. \square

We also take the opportunity to define an abstract notion of *sub-diary*. Theorem 6.2.14 will show that given a diagonal diary Δ with $\mathbf{Str}(\Delta)$ a Fraïssé limit of \mathcal{K} and an enumerated $\mathbf{A} \in \mathcal{K}$, the function sending $f \in \text{OEmb}(\mathbf{A}, \mathbf{K})$ to the sub-diary $\text{Sub}^\Delta(f[\mathbf{A}])$ exactly describes the ordered big Ramsey degree of \mathbf{A} in \mathbf{K} .

Definition 6.2.7. Fix a diary Δ with controlled critical levels, and fix $\mathbf{A} \subseteq \mathbf{Str}(\Delta)$.

1. We set $\text{Sp}^\Delta(\mathbf{A}) = \{\ell(c_\Delta(m) \wedge c_\Delta(n)) : m \neq n \in \mathbf{A}\}$.
2. We set $\text{AC}^\Delta(\mathbf{A}) = \{m < \omega : \text{Cl}_\Delta(c_\Delta[\mathbf{A}]|_{m+1}) \subsetneq \text{Cl}_\Delta(c_\Delta[\mathbf{A}]|_m)\}$.
3. We set $\mathbf{A}^\Delta = \{\ell(c_\Delta(n)) : n \in \mathbf{A}\}$.
4. We set $\text{Crit}^\Delta(\mathbf{A}) = \mathbf{A}^\Delta \cup \text{Sp}^\Delta(\mathbf{A}) \cup \text{AC}^\Delta(\mathbf{A})$.
5. The *sub-diary* $\text{Sub}^\Delta(\mathbf{A}) : |\text{Crit}^\Delta(\mathbf{A})| \rightarrow \Omega$ is now defined in the obvious way. Let $g : |\text{Crit}^\Delta(\mathbf{A})| \rightarrow \text{Crit}^\Delta(\mathbf{A})$, and fix $n < |\text{Crit}^\Delta(\mathbf{A})|$ with $g(n) = m \in \text{Crit}^\Delta(\mathbf{A})$.

- $\text{Lab}(\text{Sub}^\Delta(\mathbf{A})(n))$ indicates whether $m \in \mathbf{A}^\Delta$, $\text{Sp}^\Delta(\mathbf{A})$, or $\text{AC}^\Delta(\mathbf{A})$.
- We set $\text{Cl}(\text{Sub}^\Delta(\mathbf{A})(n)) = \text{Cl}_\Delta(c_\Delta[\mathbf{A}]|_m)$.
- If $\text{Lab}(\text{Sub}^\Delta(\mathbf{A})(n)) = \text{AC}$, we set $\text{Pos}(\text{Sub}^\Delta(\mathbf{A})(n)) = 0$. Otherwise, then writing $c_\Delta[\mathbf{A}]|_m = \{t_0 \preceq_{\text{lex}} \dots \preceq_{\text{lex}} t_{d-1}\}$, set $\text{Pos}(\text{Sub}^\Delta(\mathbf{A})(n)) = i$ iff t_i is the coding or splitting node.
- If $\text{Lab}(\text{Sub}^\Delta(\mathbf{A})(n)) \in \{\text{AC}, \text{Sp}\}$, we set $\text{Pass}(\text{Sub}^\Delta(\mathbf{A})(n))$ to be the constant zero function. If $\text{Lab}(\text{Sub}^\Delta(\mathbf{A})(n)) = \text{Cd}$, then writing $c_\Delta[\mathbf{A}]|_{m+1} = \{t_0 \preceq_{\text{lex}} \dots \preceq_{\text{lex}} t_{d-2}\}$, we set $\text{Pass}(\text{Sub}^\Delta(\mathbf{A})(n))(j) = t_j(m)$ for each $j < d-1$.

We remark that this notion of sub-diary behaves as expected in the concrete setting, i.e., if $\mathbf{A} \subseteq \mathbf{K}$ has controlled critical levels and $\mathbf{B} \subseteq \mathbf{A}$, then letting \mathbf{A}' denote the enumerated structure order isomorphic to \mathbf{A} , and letting $\mathbf{B}' \subseteq \mathbf{A}'$ be the substructure corresponding to $\mathbf{B} \subseteq \mathbf{A}$, then we have $\text{Sub}^{\Delta^{\mathbf{A}}}(\mathbf{B}') = \Delta^{\mathbf{B}}$.

A *diagonal diary* will be a diary with controlled critical levels as well as some analog of “controlled paths.” One small difference comes from the fact that when

$\mathbf{A} \subseteq \mathbf{K}$ is diagonal, we have $\text{FreeStart}(\mathbf{A}) \setminus \text{Crit}(\mathbf{A}) \neq \emptyset$. We won't be able to discuss these levels using the diary directly.

Fix Δ a diary with controlled critical levels. Given $t \in T^\Delta$, let $\text{Path}_\Delta(t) = \{\text{Cl}_\Delta(t|_m) : m < \ell(t)\}$, and if $n < |\mathbf{Str}(\Delta)|$, set $\text{Path}_\Delta(n) := \text{Path}_\Delta(c_\Delta(n))$. Since Δ has controlled critical levels, it follows that if $c_\Delta(n)^u = i$, then we have $\text{Path}_\Delta(n) \in \max(\vec{P}_i)$. Given $i < k^u$, we also write $\text{Path}(\Delta, i) = \{\text{Path}_\Delta(n) : n < |\mathbf{Str}(\Delta)|, c_\Delta(n)^u = i\}$.

Definition 6.2.8. A diary Δ is a *diagonal diary* if it has controlled critical levels and the following analog of ‘‘controlled paths:’’

1. For any $m, n < |\mathbf{Str}(\Delta)|$ with $U^{\mathbf{Str}(\Delta)}(m) = U^{\mathbf{Str}(\Delta)}(n) \notin \text{Free}$ and with $\text{Start}(c_\Delta(m)) = \text{Start}(c_\Delta(n))$, we have $\text{Path}_\Delta(m) = \text{Path}_\Delta(n)$.
2. Write $X = \{i \in \text{Free} : T_i^\Delta \neq \emptyset\}$. For each $i \in X$, let $\mathbf{p}_i^{\max} \in \text{Path}(\Delta, i)$ denote the \leq_{path} -largest member. Write $Y = (\bigcup_{i \in X} \text{Path}(\Delta, i)) \setminus \{\mathbf{p}_i^{\max} : i \in X\}$, and write $g : Y \rightarrow |Y|$ for the \leq_{path} -increasing bijection.

We demand that for every $L < |Y|$, we have $\text{Lab}(\Delta(L)) = \text{Sp}$ and $\text{Pos}(\Delta(L)) = 0$. We demand that whenever $n < |\mathbf{Str}(\Delta)|$ has $\text{Path}_\Delta(n) \in Y$, we have $\text{Start}(c_\Delta(n)) = g(\text{Path}_\Delta(n))$. We demand that whenever $n < |\mathbf{Str}(\Delta)|$ has $\text{Path}_\Delta \in \{\mathbf{p}_i^{\max} : i \in X\}$, then $\text{Start}(c_\Delta(n)) \geq |Y|$.

Regarding item 2 of Definition 6.2.8, the role is very similar to the roles of items 2 and 3 of Definition 5.3.4. The important thing is that the different possible maximal paths among the free unaries are separated in T^Δ in some arbitrary, but fixed fashion, and this occurs before any other critical events. So in some sense, all of the ‘‘non-trivial’’ content of the diagonal diary Δ occurs starting at level $|Y|$.

The prototypical example of a diagonal diary is a diary of the form $\Delta^{\mathbf{A}}$ for some diagonal $\mathbf{A} \subseteq \mathbf{K}$. In particular, Theorem 5.8.5 tells us that there is some diagonal diary Δ with $\mathbf{Str}(\Delta) = \mathbf{K}$.

A helpful way to think about item 2 of Definition 6.2.8 is as follows. Noting that $|\{t \in T^\Delta(0) : t^u \in \text{Free}\}| = |X|$, we have that $|\{t \in T^\Delta(|Y|) : t^u \in \text{Free}\}| = |X| + |Y| = |\bigcup_{i \in X} \text{Path}(\Delta, i)|$. We then have for every $n < |\mathbf{Str}(\Delta)|$ with $U^{\mathbf{Str}(\Delta)}(n) \in \text{Free}$ that $\text{Path}_\Delta(n)$ depends only on $c_\Delta(n)|_{|Y|}$. We also note that given $i \in \text{Free}$ and $n < |\mathbf{Str}(\Delta)|$, then $\text{Path}_\Delta(n) = \mathbf{p}_i^{\max}$ iff $c_\Delta(n)|_{|Y|} = (i, 0^{|Y|})$.

When considering diagonal diaries, the following definition will be useful.

Definition 6.2.9. Fix a diagonal diary Δ . We say that $v \in T^\Delta$ has a *defined path continuation* if whenever $m, n < |\mathbf{Str}(\Delta)|$ have $c_\Delta(m) \supseteq v$ and $c_\Delta(n) \supseteq v$, then $\text{Path}_\Delta(c_\Delta(m)) = \text{Path}_\Delta(c_\Delta(n))$.

So one way to paraphrase what it means for Δ to be diagonal is that ‘‘most’’ $v \in T^\Delta$ have a defined path. The only exceptions are if $v = (j, 0^{\ell(v)})$ for some $j \notin \text{Free}$, or if $v = (j, 0^m)$ for some $j \in \text{Free}$ and $m < |Y|$, where Y is as in item 2 of Definition 6.2.8.

Example 6.2.10. We describe all finite diagonal diaries for the class \mathcal{G}_3 of triangle free graphs. A helpful preliminary observation is to note that if $S \subseteq \text{CT}(m)$ and

$0^m \notin S$, then $\mathcal{K}(S)$ can be described as a graph on vertex set S . If $S = \{s_0 \preceq_{lex} \dots \preceq_{lex} s_{d-1}\}$ and z is the $(d, 3)$ -graph-age function with $\mathcal{G}_z = \mathcal{K}(S)$, then we say that s_i and s_j are adjacent iff $z(\{s_i, s_j\}) = 2$. Next, we note that $\mathcal{K}(\{0^m\} \cup S)$ can be recovered from just $\mathcal{K}(S)$. Thus, given a diagonal diary Δ , by paying attention to only the part of T^Δ off of the left branch, we obtain the following general procedure to produce a diagonal diary coding a finite enumerated triangle-free graph:

1. At stage 0, we start with the empty graph.
2. Suppose at stage $m < \omega$, we have produced a finite ordered graph G_m . If G_m is the empty graph, we can choose to stop.
3. If at stage m we choose to (or must) continue, we can choose exactly one of the following options for producing G_{m+1} :
 - (a) We can introduce a new least vertex which is adjacent to every vertex from G_m (i.e. the node on the left branch splits).
 - (b) If $G_m \neq \emptyset$, we can choose a vertex in G_m and duplicate it; these two new vertices will be order consecutive and graph non-adjacent in G_{m+1} , and will have the same neighbors in G_m as before (i.e. a node off of the left branch splits).
 - (c) If G_m has edges, we can choose an edge in G_m and delete it (i.e. an age change level).
 - (d) If G_m contains a vertex $v \in G_m$ whose set of neighbors is an anti-clique, we can delete it (i.e. a coding level).

Finite diagonal diaries for enumerated triangle-free graphs are in one-one correspondence with finite runs of this procedure. The size of the enumerated triangle-free graph which is coded by this procedure is exactly the number of vertex deletion events, and the edges of this triangle-free graph are determined by the edges in G_m at the time of the vertex deletion events. For instance, if we wish to code an enumerated anti-clique, this is the same as demanding that we only delete isolated vertices from G_m . Thus the ordered big Ramsey degree of the n -element anti-clique for $n = 1, 2, 3, 4$ are 1, 5, 161, 134397. It would be an interesting problem in enumerative combinatorics to obtain an exact formula for the ordered big Ramsey degree of the n -element anti-clique, or to even understand how this function grows asymptotically.

Example 6.2.11. Fixing $\ell \geq 3$, we describe all finite diagonal diaries for the class \mathcal{G}_ℓ of K_ℓ -free graphs. Much like in Example 6.2.10, for any $S \subseteq \text{CT}(m)$, we can recover $\mathcal{K}(\{0^m\} \cup S)$ from $\mathcal{K}(S)$. So given a diagonal diary Δ , we can give a complete description of it while only paying attention to what's going on off of the left branch. It will be helpful to generalize the notion of (d, ℓ) -graph-function by replacing the number d with any finite linear order X . We call a finite linear order X equipped with an (X, ℓ) -graph-age function an *ordered ℓ -graph-age structure*. If X is a finite linear order and $z: \binom{X}{<\ell}$ is a (X, ℓ) graph-age function, then by viewing z as an $(|X|, \ell)$ -graph-age function, we can write \mathcal{G}_z for the corresponding member of $P(|X|)$. We can now describe the procedure by which we produce a diagonal diary coding a finite enumerated K_ℓ -free graph:

1. At stage 0, we start with an empty ℓ -graph-age structure.
2. Suppose at stage $m < \omega$, we have produced a finite ordered ℓ -graph-age structure (X_m, z_m) . If X_m is empty, we can choose to stop.
3. If at stage m we choose to (or must) continue, we can choose exactly one of the following options for producing (X_{m+1}, z_{m+1}) :
 - (a) We can introduce a new least vertex x , set $z_{m+1}(\{x\}) = \ell - 2$, and for each non-empty $S \in \binom{X_m}{<\ell-1}$, we set $z_{m+1}(S \cup \{x\}) = \ell - 1$ (i.e. the node on the left branch splits).
 - (b) If $X_m \neq \emptyset$, we can choose a vertex v in X_m and duplicate it; these two new vertices x and y will be order consecutive, and letting $\pi: X_{m+1} \rightarrow X_m$ be the identity except on x and y , where $\pi(x) = \pi(y) = v$, then given $S \subseteq X_{m+1}$, we set $z_{m+1}(S) = z_m(\pi[S])$ (i.e. a node off off the left branch splits).
 - (c) If $X_m \neq \emptyset$ and there is $S \subseteq X_m$ with $z_m(S) \geq 2$, we can set $X_{m+1} = X_m$ and choose $z_{m+1} \leq z_m$ (pointwise as functions) so that $(\mathcal{G}_{z_m}, \mathcal{G}_{z_{m+1}}) \in \text{Con}(P(|X_m|))$ (i.e. an age change level).
 - (d) If $X_m \neq \emptyset$, say $X_m = \{x_0 \leq \dots \leq x_{d-1}\}$, and $j < d$ is such that (\mathcal{G}_{z_m}, j) is a controlled coding pair, we can delete the vertex x_j (i.e. a coding level).

Example 6.2.12. An example with non-trivial paths...

Definition 6.2.13. Let Δ and Θ be diaries. An *embedding* of Δ into Θ is simply an aged embedding of T^Δ into T^Θ , i.e., a map which preserves tree structure, relative levels, coding nodes, passing numbers, and age-sets.

Our main lower bound theorem for big Ramsey degrees is the following.

Theorem 6.2.14. *Let Δ be any diagonal diary so that $\mathbf{Str}(\Delta)$ is a Fraïssé limit of \mathcal{K} . Let Θ be any other diagonal diary. Then there is an embedding of Θ into Δ . In particular, given an enumerated $\mathbf{A} \in \mathcal{K}$, the ordered big Ramsey degree of \mathbf{A} in \mathbf{K} is exactly the number of diagonal diaries which code \mathbf{A} .*

The remainder of this section is spent proving this theorem, so for the time being, fix Δ and Θ as in the theorem statement.

6.3 Large subsets of coding nodes

Our proof of Theorem 6.2.14 is adapted from the characterization of the exact big Ramsey degrees for the class of finite graphs given by Laflamme, Sauer, and Vuksanovic [18]. In their proof, they more-or-less treat all nodes as coding nodes, and while the resulting graph is not the Rado graph, it does embed a copy of it (so is bi-embeddable with it). Upon defining a suitable notion of diagonal substructure, the authors then fix an embedding ψ from the universal graph coded by $2^{<\omega}$ and the diagonal copy. This leads to an important collection of pairs of nodes, namely those pairs (u, v) where $\psi^{-1}(\text{Succ}(v))$ is dense above u . Our strategy is similar, but

we require several notions of largeness which are more sophisticated than merely being dense above u , as we will need to keep track of the relevant age-set structure.

Notation 6.3.1. Fix $i < k^u$. Recall that \vec{P}_i denotes the collection of paths in P_i , i.e., those subsets of P_i which are linearly ordered. Also recall that $\max(\vec{P}_i)$ denotes those elements of P_i which are maximal under inclusion. We let $\text{Full}(\vec{P}_i) \subseteq \vec{P}_i$ denote those paths which are initial segments of members of $\max(\vec{P}_i)$.

Given $\mathfrak{p} \in \text{Full}(\vec{P}_i)$, we let \mathfrak{p}' denote the member of $\text{Full}(\vec{P}_i)$ obtained from \mathfrak{p} by deleting the largest element of \mathfrak{p} .

Definition 6.3.2. Suppose that $S \subseteq c[\mathbf{K}]$. Fix $i < k^u$, and let $\mathfrak{p} \in \text{Full}(\vec{P}_i)$. Let $u \in \text{CT}_i$ with $\mathcal{K}(u) = \max(\mathfrak{p})$. We define the property that S is (u, \mathfrak{p}) -large by induction on $|\mathfrak{p}|$.

- If $|\mathfrak{p}| = 1$, then S is (u, \mathfrak{p}) -large if it is dense in CT over u .
- Otherwise, we say that S is (u, \mathfrak{p}) -large if for every $v \sqsupseteq u$ with $\mathcal{K}(v) = \mathcal{K}(u) = \max(\mathfrak{p})$, there is $w \sqsupseteq v$ with $\mathcal{K}(w) = \max(\mathfrak{p}')$ so that S is (w, \mathfrak{p}') -large.

As a convention, if we refer to anything being (u, \mathfrak{p}) -large for some $\mathfrak{p} \in \text{Full}(\vec{P}_i)$, it is assumed that $u \in \text{CT}_i$ and $\mathcal{K}(u) = \max(\mathfrak{p})$.

Lemma 6.3.3. *Suppose $i < k^u$, $u \in \text{CT}_i$ and $S \subseteq c[\mathbf{K}]$ is dense in CT over u . Then S is (u, \mathfrak{p}) -large for any $\mathfrak{p} \in \text{Full}(\vec{P}_i)$ with $\max(\mathfrak{p}) = \mathcal{K}(u)$.*

Proof. We induct on $|\mathfrak{p}|$. When $|\mathfrak{p}| = 1$, then we have $\mathfrak{p} = \{P_i^{\min}\}$, and this is just the definition of (u, \mathfrak{p}) -large. Now suppose $|\mathfrak{p}| \geq 2$. Consider $v \sqsupseteq u$ with $\mathcal{K}(v) = \mathcal{K}(u) = \max(\mathfrak{p})$. Find $w \sqsupseteq v$ with $\mathcal{K}(w) = \max(\mathfrak{p}')$. Since S is dense in CT over w , our inductive assumption implies that S is (w, \mathfrak{p}') -large. Hence S is (u, \mathfrak{p}) -large as desired. \square

Lemma 6.3.4. *Suppose $i < k^u$, $\mathfrak{p} \in \text{Full}(\vec{P}_i)$, $u \in \text{CT}_i$, and that $S \subseteq c[\mathbf{K}]$ is (u, \mathfrak{p}) -large. If $n < \omega$ and we write $S = \bigcup_{j < n} S_j$, then for some $j < n$ and some $v \sqsupseteq u$ with $\mathcal{K}(v) = \mathcal{K}(u) = \max(\mathfrak{p})$, we have that S_j is (v, \mathfrak{p}) -large.*

Proof. First assume that $\mathfrak{p} = \{P_i^{\min}\}$. Towards a contradiction, suppose we could find $u \in \text{CT}_i$ with $\mathcal{K}(u) = P_i^{\min}$, $S \subseteq c[\mathbf{K}]$ dense in CT over u , and a partition $S = S_0 \cup \dots \cup S_{n-1}$ so that the conclusion of the lemma does not hold. Set $u = u_0$. Suppose $u_j \sqsupseteq \dots \sqsupseteq u_0$ have been defined with $\mathcal{K}(u_j) = P_i^{\min}$. As S_j is not dense in CT over u_j , find $u_{j+1} \sqsupseteq u_j$ with $\mathcal{K}(u_{j+1}) = P_i^{\min}$ and so that $S_j \cap \text{Succ}(u_{j+1}) = \emptyset$. Continue until u_n has been defined. Then S is dense in CT over u_n , but also $S_j \cap \text{Succ}(u_n) = \emptyset$ for every $j < n$. This is a contradiction.

For larger \mathfrak{p} , we proceed in a similar fashion. Set $u = u_0$. Suppose $u_j \sqsupseteq \dots \sqsupseteq u_0$ have been defined for some $j < n$ with $\mathcal{K}(u_j) = \max(\mathfrak{p})$. Since S_j is not (u_j, \mathfrak{p}) -large, we can find $u_{j+1} \sqsupseteq u_j$ with $\mathcal{K}(u_{j+1}) = \max(\mathfrak{p})$ so that for any $v \sqsupseteq u_{j+1}$ with $\mathcal{K}(v) = \max(\mathfrak{p}')$, we have that S_j is not (v, \mathfrak{p}') -large. However, since S is (u, \mathfrak{p}) -large, we can find $v \sqsupseteq u_n$ with $\mathcal{K}(v) = \max(\mathfrak{p}')$ and so that S is (v, \mathfrak{p}') -large. By our induction assumption, we can find $j < n$ and $w \sqsupseteq v$ with $\mathcal{K}(w) = \max(\mathfrak{p}')$ and so that S_j is (w, \mathfrak{p}') -large. But since $w \sqsupseteq u_{j+1}$, this is a contradiction. \square

The next proposition says that large sets of coding nodes code suitably rich substructures of \mathbf{K} . The proof is straightforward, but a bit long, and can be skipped on a first reading.

Proposition 6.3.5. *Let $m < \omega$, let $X = \{x_0, \dots, x_{d-1}\} \subseteq \text{CT}(m)$, and write $\rho = X^u$. For each $i < d$, let $\mathbf{p}_i \in \text{Full}(\vec{P}_{\rho(i)})$ and suppose that $S_i \subseteq c[\mathbf{K}] \cap \text{Succ}(x_i)$ is (x_i, \mathbf{p}_i) -large for each $i < d$. Setting $S = \bigcup_{i < d} S_i$ and viewing $c^{-1}(S)$ as an \mathcal{L}_d -structure, we have $\text{Age}(c^{-1}(S)) = \mathcal{K}(X)$.*

Proof. If $|\mathbf{p}_i| = 1$ for each $i < d$, the assumption on the S_i amounts to saying that S is dense in CT above each x_i . For each $\mathbf{B} \in \mathcal{K}(X)$, we show that $\mathbf{B} \in \text{Age}(c^{-1}(S))$ by induction on the size of \mathbf{B} (for all d simultaneously). If \mathbf{B} contains a single point b with $V^{\mathbf{B}}(b) = i$, the result is clear. Now suppose \mathbf{B} is an \mathcal{L}_d -structure with underlying set B and that $|B| \geq 2$. Pick some $b \in \mathbf{B}$, and write $V^{\mathbf{B}}(b) = i$. Let $w_i \sqsupseteq x_i$ be any member of S_i , and for each $j < d \setminus \{i\}$, we set $w_j = \text{Left}(x_j, \ell(v_i))$. Write $W = \{w_0 \preceq_{\text{lex}} \dots \preceq_{\text{lex}} w_{d-1}\}$, and write $Y = \text{CT} \cap \text{IS}(W) := \{y_0 \preceq_{\text{lex}} \dots \preceq_{\text{lex}} y_{\alpha-1}\}$. Note that since $\mathcal{K}(x_i) = \max(\mathbf{p}_i) = P_{\rho(i)}^{\min}$ (since $|\mathbf{p}_i| = 1$), we have $\mathcal{K}(X) = \mathcal{K}(W)$. Let \mathbf{C} be the \mathcal{L}_α -structure on underlying set $B \setminus \{b\}$ defined as follows. The \mathcal{L}^b -part of \mathbf{C} is induced from \mathbf{B} . For the unary part, fix $a \in B \setminus \{b\}$, and suppose that $V^{\mathbf{B}}(a) = j$ and $R^{\mathbf{B}}(a, b) = q$. Then $w_j \widehat{\ } q \in \text{CT}$ by Proposition 5.6.2. Now if $\beta < \alpha$ is such that $y_\beta = w_j \widehat{\ } q$, we set $V^{\mathbf{C}}(a) = \beta$. Since S is dense above each member of Y , our inductive hypothesis shows that viewing $c^{-1}(S \cap \text{Succ}(Y))$ as an \mathcal{L}_α -structure, we have $\mathbf{C} \in \text{Age}(c^{-1}(S \cap \text{Succ}(Y)))$. It follows that $\mathbf{B} \in \text{Age}(c^{-1}(S))$.

Now suppose we are given $\{\mathbf{p}_j : j < d\}$, where $|\mathbf{p}_j| \leq n$ for each $j < d$. Suppose we know the result (for all d) whenever the paths have size less than n . Write $I = \{i < d : |\mathbf{p}_i| = n\}$. Let $\mathbf{B} \in \mathcal{K}(X)$ have underlying set B , and write \mathbf{B}_i for the induced substructure on $\{b \in B : V^{\mathbf{B}}(b) = i\}$. Fix some $N \geq \max\{|\mathbf{B}_i| : i \in I\}$, and fix some $i \in I$. Since S_i is (x_i, \mathbf{p}_i) -large, there is $x'_{i,0} \sqsupseteq x_i$ so that S_i is $(x'_{i,0}, \mathbf{p}'_i)$ -large. If $x'_{i,r}$ has been defined, then since S_i is (x_i, \mathbf{p}_i) -large and $\mathcal{K}(\text{Left}(x_i, \ell(x'_{i,r}))) = \mathcal{K}(x_i) = \max(\mathbf{p}_i)$, there is $x'_{i,r+1} \sqsupseteq \text{Left}(x_i, \ell(x'_{i,r}))$ so that S_i is $(x'_{i,r+1}, \mathbf{p}'_i)$ -large. Continue until $x'_{i,N-1}$ is defined, and write $X_i = \{x'_{i,N-1} \preceq_{\text{lex}} \dots \preceq_{\text{lex}} x'_{i,0}\}$.

Do the same procedure for every $i \in I$, at each stage moving relevant points up leftmost far enough to be above anything previously considered, producing a set X_i as above. If $i \in d \setminus I$, set $X_i = \{x_i\}$. Pick some level L above everything we have considered, and set $Y = \bigcup_{i < d} \text{Left}(X_i, L) = \{y_0 \preceq_{\text{lex}} \dots \preceq_{\text{lex}} y_{\alpha-1}\}$. Let $\pi : \alpha \rightarrow d$ be the map so that $y_\beta \sqsupseteq x_{\pi(\beta)}$ for each $\beta < \alpha$. Then by the construction of Y , we have

$$\mathcal{K}(Y) = \mathcal{K}(X)^\pi \cap \bigcap_{i \in I} \bigcap_{\beta \in \pi^{-1}(\{i\})} \mathcal{K}(Y^u)|_{(\iota_\beta, \max(\mathbf{p}'_i))}.$$

From \mathbf{B} , we form an \mathcal{L}_α -structure \mathbf{C} on underlying set B as follows. The \mathcal{L}^b -part is the same as \mathbf{B} . For each $i \in d \setminus I$, let $\beta_i < \alpha$ be the unique member of $\pi^{-1}(\{i\})$, and set $V^{\mathbf{C}}(b) = \beta_i$ for each $b \in \mathbf{B}_i$. If $i \in I$, then for each $b \in \mathbf{B}_i$, choose a distinct $\beta_b \in \pi^{-1}(\{i\})$, and set $V^{\mathbf{C}}(b) = \beta_b$. Now $\mathbf{C} \in \mathcal{K}(Y)$, so by our inductive hypothesis, $\mathbf{C} \in \text{Age}(c^{-1}(S \cap \text{Succ}(Y)))$, with the latter viewed as an \mathcal{L}_α -structure. So also $\mathbf{B} \in \text{Age}(c^{-1}(S))$ as desired. \square

Returning to the setting of Theorem 6.2.14, fix $\psi \in \text{OEmb}(\mathbf{K}, \mathbf{Str}(\Delta))$. By an abuse of notation, let us also write ψ for the induced map on coding nodes,

i.e., $\psi(c(n)) := c_\Delta(\psi(n))$. This will remain fixed for the remainder of the proof of Theorem 6.2.14.

Definition 6.3.6. Suppose $\mathfrak{p} \in \text{Full}(\vec{P}_i)$. We say that a pair $(u, v) \in \text{CT} \times T^\Delta$ is \mathfrak{p} -compatible if $\text{Cl}_\Delta(v) = \max(\mathfrak{p})$ and $\psi^{-1}(\text{Succ}(v))$ is (u, \mathfrak{p}) -large. We write $\text{Pair}(\mathfrak{p})$ for the set of \mathfrak{p} -compatible pairs.

Notice that if $(u, v) \in \text{Pair}(\mathfrak{p})$ and $u_0 \sqsupseteq u$ satisfies $\mathcal{K}(u_0) = \mathcal{K}(u)$, then also $(u_0, v) \in \text{Pair}(\mathfrak{p})$. We also note that by Lemma 6.3.3, given any $i < k^u$ and any $\mathfrak{p} \in \max(\vec{P}_i)$, we have $(\emptyset_i, \emptyset_i) \in \text{Pair}(\mathfrak{p})$. The last results of this subsection collect some straightforward “pair-extension” properties and their corollaries. Before proving these, let us rephrase Proposition 6.2.5 in a way that is more suitable for our setting.

Notation 6.3.7. When $\sigma: d \rightarrow T(n)$ is a function, Definition 4.2.1 gives us a class $\mathcal{K}(\sigma)$ of \mathcal{L}_d -structures. It will be useful to generalize this to the case where $\sigma: d \rightarrow T$, i.e., where $\text{Im}(\sigma)$ is not a subset of just one level. To do this, pick any $N < \omega$ with $N \geq \max\{\ell(\sigma(i)) : i < d\}$, and let $\sigma^+: d \rightarrow T(N)$ be given by $\sigma^+(i) = \text{Left}(\sigma(i), N)$. We then set $\mathcal{K}(\sigma) = \mathcal{K}(\sigma^+)$.

Let us emphasize that we do not make this generalization for diaries; when referring to a diary Γ , we will only refer to $\text{Cl}_\Gamma(\sigma)$ for functions σ mapping to one level of T^Γ .

Lemma 6.3.8. Fix a diary Δ with controlled critical levels. Suppose $\sigma: d \rightarrow \text{CT}$ is a function, and suppose $N > \max\{\ell(c_\Delta(\psi(n))) : n < \max\{\ell(\sigma(j)) : j < d\}\}$. For each $j < d$, suppose $s_j \in c[\mathbf{K}] \cap \text{Succ}(\sigma(j))$ satisfies $\ell(\psi(s_j)) \geq N$. Then we have $\text{Cl}_\Delta(\langle \psi(s_j)|_N : j < d \rangle) \subseteq \mathcal{K}(\sigma)$.

Proof. Since $\psi \in \text{OEmb}(\mathbf{K}, \text{Str}(\Delta))$, we have for each $n < \max\{\ell(\sigma(j)) : j < d\}$ that

$$\psi(s_j)(\ell(c_\Delta(\psi(n)))) = s_j(n) = u_j(n).$$

The result now follows from Proposition 6.2.5. \square

Proposition 6.3.9. Suppose $i < k^u$, $\mathfrak{p} \in \text{Full}(\vec{P}_i)$, and $(u, v) \in \text{Pair}(\mathfrak{p})$.

1. Suppose $N > \ell(v)$. Then there is $v_0 \sqsupseteq v$ with $\ell(v_0) = N$ and $u_0 \sqsupseteq u$ with $(u_0, v_0) \in \text{Pair}(\mathfrak{p})$.
2. Suppose $|\mathfrak{p}| \geq 2$. Then there is $(u', v') \in \text{Pair}(\mathfrak{p}')$ with $u' \sqsupseteq u$ and $v' \sqsupseteq v$.

Proof.

1. Write $\text{Succ}(v) \cap T^\Delta(N) = \{t_0, \dots, t_{d-1}\}$. There is a finite set F for which we have

$$\psi^{-1}(\text{Succ}(v)) \setminus F = \bigcup_{j < d} \psi^{-1}(\text{Succ}(t_j)).$$

By Lemma 6.3.4, there is $u_0 \sqsupseteq u$ and $j < d$ so that $\psi^{-1}(\text{Succ}(t_j))$ is (u_0, \mathfrak{p}) -large. It follows by Proposition 6.3.5 that $t_j := v_0$ must have $\text{Cl}_\Delta(v_0) = \max(\mathfrak{p})$, i.e., $(u_0, v_0) \in \text{Pair}(\mathfrak{p})$.

2. Because $\psi^{-1}(\text{Succ}(v))$ is (u, \mathbf{p}) -large, find $u_0 \sqsupseteq u$ with $\mathcal{K}(u_0) = \max(\mathbf{p}')$ and so that $\psi^{-1}(\text{Succ}(v))$ is (u_0, \mathbf{p}') -large. Using Lemma 6.3.8, find N so that whenever $s \in \psi^{-1}(\text{Succ}(v)) \cap \text{Succ}(u_0)$ and $\ell(\psi(s)) > N$, we have $\text{Cl}_\Delta(\psi(s)|_N) \subseteq \max(\mathbf{p}')$. Let us write

$$S = \{s \in \psi^{-1}(\text{Succ}(v)) \cap \text{Succ}(u) : \ell(\psi(s)) > N\}.$$

Then S is a cofinite subset of $\psi^{-1}(\text{Succ}(v)) \cap \text{Succ}(u)$, a (u_0, \mathbf{p}') -large set; hence S is also (u_0, \mathbf{p}') -large. Writing $\psi[S]|_N = \{t_0, \dots, t_{d-1}\}$, let $S_j = \{s \in S : \psi(s)|_N = t_j\}$. We can find $u' \sqsupseteq u_0$ so that some S_j is (u', \mathbf{p}') -large. Set $v' = t_j$. It remains to check that $\text{Cl}_\Delta(v') = \max(\mathbf{p}')$. By choice of v' we must have $\text{Cl}_\Delta(v') \subseteq \max(\mathbf{p}')$. The reverse inclusion follows from Propositions 6.3.5 and 6.2.6. \square

Corollary 6.3.10. *Suppose $i \in k^u \setminus \text{Free}$. Then $(i, 0^n) \in T^\Delta$ for every $n < \omega$, and $((i, 0^m), (i, 0^n)) \in \text{Pair}(\mathbf{p})$ for every $m, n < \omega$ and $\mathbf{p} \in \max(\vec{P}_i)$.*

Proof. Start from the observation that $(\emptyset_i, \emptyset_i) \in \text{Pair}(\mathbf{p})$ and run Proposition 6.3.9; we obtain some $(u, v) \in \text{Pair}(\mathbf{p})$ with $\ell(v) = n$ and $u \in \text{CT}_i$. Since $\mathcal{K}(u) = \text{Cl}_\Delta(v) = \mathcal{K}(\emptyset_i)$, and since $i \notin \text{Free}$, we must have $u = (i, 0^m)$ for some $m < \omega$, and we must have $v = (i, 0^n)$. So $\psi^{-1}(\text{Succ}(v))$ is $((i, 0^m), \mathbf{p})$ -large, and it is enough to show that it is $(\emptyset_i, \mathbf{p})$ -large. But if $u \sqsupseteq \emptyset$ has $\mathcal{K}(u) = \mathcal{K}(\emptyset_i)$, then $u = 0^{m'}$ for some $m' < \omega$. Set $M = \max\{m, m'\}$; then because $\psi^{-1}(\text{Succ}(v))$ is $((i, 0^m), \mathbf{p})$ -large, we can find $w \sqsupseteq 0^M$ so that $\psi^{-1}(\text{Succ}(v))$ is (w, \mathbf{p}') -large. So this shows that $\psi^{-1}(\text{Succ}(v))$ is $(\emptyset_i, \mathbf{p})$ -large as desired, \square

Corollary 6.3.11. *Suppose $i < k^u$, $\mathbf{p} \in \text{Full}(\vec{P}_i)$, and $(u, v) \in \text{Pair}(\mathbf{p})$. Then there is $n < |\mathbf{Str}(\Delta)|$ with $c_\Delta(n) \sqsupseteq v$ and $\text{Path}_\Delta(n) = \mathbf{p} \cup \text{Path}_\Delta(v)$.*

Proof. First note that $\mathbf{p} \cap \text{Path}_\Delta(v) = \{\max(\mathbf{p})\}$. Our proof proceeds by induction on $|\mathbf{p}|$. When $|\mathbf{p}| = 1$, then $\text{Path}_\Delta(v) \in \max(\vec{P}_i)$, and the result is clear. Now suppose $|\mathbf{p}| \geq 2$. Use Proposition 6.3.9 to find $u' \sqsupseteq u$ and $v' \sqsupseteq v$ with $(u', v') \in \text{Pair}(\mathbf{p}')$. By our inductive assumption, find $n < |\mathbf{Str}(\Delta)|$ with $c_\Delta(n) \sqsupseteq v' \sqsupseteq v$ and $\text{Path}_\Delta(n) = \mathbf{p}' \cup \text{Path}_\Delta(v') = \mathbf{p} \cup \text{Path}_\Delta(v)$. So the result is true for \mathbf{p} as well. \square

6.4 The construction

We now begin the construction for the proof of Theorem 6.2.14. The idea is roughly as follows: given Θ , we will build our aged embedding $f: T^\Theta \rightarrow T^\Delta$ level by level. When constructing f , we will write $f(m)$ for $\ell(f(x))$ for some (any) $x \in T^\Theta(m)$. If f has been defined on $T_\Theta(m-1)$, then $f^+: T^\Theta(m) \rightarrow T^\Delta$ is defined, for $v \in T_\Theta(m-1)$ and $i < k$, via

$$f^+(v \frown i) = f(v) \frown i.$$

Let us also set $f^+(\emptyset_i) = \emptyset_i$ for each $i < k^u$.

To give the first few levels of the construction, let us establish some notation similar to item 2 of Definition 6.2.8. Write $X_\Delta = \{i \in \text{Free} : T_i^\Delta \neq \emptyset\}$; since

$\mathbf{Str}(\Delta) \cong \mathbf{K}$, we have $X_\Delta = \mathbf{Free}$, but we will adopt similar notations for Θ , where we might have $X_\Theta \subsetneq \mathbf{Free}$. For each $i \in X_\Delta$, let $\mathbf{p}_{i,\Delta}^{\max} \in \text{Path}(\Delta, i)$ denote the \leq_{path} -largest member (we will soon see that $\text{Path}(\Delta, i) = \max(\vec{P}_i)$). Write $Y_\Delta = \left(\bigcup_{i \in X_\Delta} \text{Path}(\Delta, i) \right) \setminus \{\mathbf{p}_{i,\Delta}^{\max} : i \in X_\Delta\}$, and write $g_\Delta: Y_\Delta \rightarrow |Y_\Delta|$ for the \leq_{path} -increasing bijection. As mentioned before, the versions of this notation using the diary Θ are defined analogously. Building on the remark after Definition 6.2.8, let us write $\bigcup_{i \in \mathbf{Free}} T_i^\Delta(|Y_\Delta|) := \{v_{\Delta, \mathbf{p}} : \mathbf{p} \in \bigcup_{i \in X_\Delta} \text{Path}(\Delta, i)\}$, where any $n < |\mathbf{Str}(\Delta)|$ with $\text{Path}_\Delta(n) = \mathbf{p} \in \bigcup_{i \in X_\Delta} \text{Path}(\Delta, i)$ has $c_\Delta(n)|_{Y_\Delta} = v_{\Delta, \mathbf{p}}$. Also define analogous notation for Θ .

First, we argue that since Δ codes the Fraïssé limit, we must have $\text{Path}(\Delta, i) = \max(\vec{P}_i)$. We state this below just for unaries in \mathbf{Free} , but we will eventually see that the result holds more generally.

Lemma 6.4.1. *Suppose $i \in \mathbf{Free}$ and $\mathbf{p} \in \max(\vec{P}_i)$. Then $\mathbf{p} \in \text{Path}(\Delta, i)$, and there is $u \in \text{CT}$ with $(u, v_{\Delta, \mathbf{p}}) \in \text{Pair}(\mathbf{p})$.*

Proof. We have that $(\emptyset_i, \emptyset_i) \in \text{Pair}(\mathbf{p})$. Using Proposition 6.3.9, find $u \in \text{CT}_i$ and $v \in T_i^\Delta(|Y_\Delta|)$ with $(u, v) \in \text{Pair}(\mathbf{p})$. By Corollary 6.3.11, we must have that $\mathbf{p} \in \text{Path}(\Delta, i)$ and that $v = v_{\Delta, \mathbf{p}}$. \square

This allows us to define the first few levels of f . Given $m < |Y_\Theta|$, suppose that $m = \text{Start}(v_{\Theta, \mathbf{p}})$ for some $\mathbf{p} \in \max(\vec{P}_i)$ with $i \in \mathbf{Free}$. We set $\tilde{f}(m) = \text{Start}(v_{\Delta, \mathbf{p}})$, noting that $v_{\Delta, \mathbf{p}}$ exists by Lemma 6.4.1. Now fix $t \in T^\Theta(m)$. If $t^u \notin \mathbf{Free}$, then $t = (t^u, 0^m)$, and we set $f(t) = (t^u, 0^{\tilde{f}(m)})$. If $t \in T^\Theta(m)$ is such that $t = v_{\Theta, \mathbf{q}}|_m$ for some $\mathbf{q} \in \bigcup_{j \in \mathbf{Free}} \max(\vec{P}_j)$, we set $f(t) = v_{\Delta, \mathbf{q}}|_{\tilde{f}(m)}$. Running this construction up to level $|Y_\Theta| - 1$, we note that for any $i \in \mathbf{Free}$ and $\mathbf{p} \in \text{Path}(\Theta, i)$ that $f^+(v_{\Theta, \mathbf{p}}) \sqsubseteq v_{\Delta, \mathbf{p}}$. This along with Lemma 6.4.1 imply that for each such \mathbf{p} , there is $u_{\mathbf{p}} \in \text{CT}$ with $(u_{\mathbf{p}}, f^+(v_{\Theta, \mathbf{p}})) \in \text{Pair}(\mathbf{p})$. We will use these $u_{\mathbf{p}}$ to continue the construction.

Before we continue the construction beyond the free start levels, let us streamline our notion of compatible pair while also elaborating on Definition 6.2.9.

Definition 6.4.2.

1. Suppose $v \in T_i^\Delta$ has a defined path continuation (Definition 6.2.9). The *path continuation* of v , denote $\text{PCon}(v)$, is the unique $\mathbf{p} \in \text{Full}(\vec{P}_i)$ so that $\text{Cl}_\Delta(v) = \max(\mathbf{p})$ and so that for any $n < |\mathbf{Str}(\Delta)|$ with $c_\Delta(n) \sqsupseteq v$, we have $\text{Path}_\Delta(c_\Delta(n)) = \text{Path}_\Delta(v) \cup \mathbf{p}$.
2. We say that $(u, v) \in \text{CT} \times T^\Delta$ is *compatible* if either one of the following happens.
 - If $v = (j, 0^{\ell(v)})$ for some $j \in k^u \setminus \mathbf{Free}$, then we simply ask that $u = (j, 0^{\ell(u)})$.
 - Otherwise, we demand that v has a defined path continuation and that $(u, v) \in \text{Pair}(\text{PCon}(v))$.

Given $d < \omega$, we say that functions $\sigma: d \rightarrow \text{CT}$ and $\xi: d \rightarrow T^\Delta$ are compatible if $(\sigma(i), \xi(i))$ is compatible for every $i < d$. We write $(u, v) \in \text{Pair}$, or in a mild abuse of notation, $(\sigma, \xi) \in \text{Pair}$.

Briefly returning to what we have constructed so far, write $f^+[T^\Theta(|Y_\Theta|)] = \{v_0 \preceq_{lex} \cdots \preceq_{lex} v_{d-1}\}$. We note that $\text{Cl}_\Theta(T^\Theta(|Y_\Theta|)) = \text{Cl}_\Delta(f^+[T^\Theta(|Y_\Theta|)]) = \mathcal{K}(T^\Theta(|Y_\Theta|)^u)$. Let $\xi: d \rightarrow T^\Delta$ be the function with $\xi(i) = v_i$ for each $i < d$. Then there is $\sigma: d \rightarrow \text{CT}$ with $(\sigma, \xi) \in \text{Pair}$. If v_i has the form $f^+(v_{\Theta, \mathfrak{p}})$ as discussed before Definition 6.4.2, we set $\sigma(i) = u_{\mathfrak{p}}$. Otherwise, we must have v_i on the left branch of T_j for some $j \notin \text{Free}$, and we can set $\sigma(i) = v_i$.

Now fix some $m \geq |Y_\Theta|$, and assume that f is defined on $T^\Theta(<m)$. Write $T^\Theta(m) = \{t_0 \preceq_{lex} \cdots \preceq_{lex} t_{d-1}\}$ and $f^+[T^\Theta(m)] = \{v_0 \preceq_{lex} \cdots \preceq_{lex} v_{d-1}\}$. We assume that our construction to this point satisfies all of the following:

1. $\text{Cl}_\Theta(T^\Theta(m)) = \text{Cl}_\Delta(f^+[T^\Theta(m)])$.
2. There are functions $\sigma: d \rightarrow \text{CT}$ and $\xi: d \rightarrow T^\Delta$ with $\xi(i) \sqsupseteq v_i$ for each $i < d$, with $\text{Im}(\xi)$ a level set, with $\text{Cl}_\Delta(f^+[T^\Theta(m)]) = \text{Cl}_\Delta(\xi)$, and with $(\sigma, \xi) \in \text{Pair}$.

We will repeatedly use the following strengthening of Proposition 6.3.9, which allows us to run the proposition on functions rather than just pairs, while preserving the ages of said functions.

Proposition 6.4.3. *Suppose $\sigma: d \rightarrow \text{CT}$ and $\xi: d \rightarrow T^\Delta$ are functions with $(\sigma, \xi) \in \text{Pair}$. Let*

$$N > \max(\{\ell(\xi(i)) : i < d\} \cup \{\ell(c_\Delta(\psi(n))) : n < \max\{\ell(\sigma(i)) : i < d\}\}).$$

Then there are functions $\sigma' \sqsupseteq \sigma$ and $\xi' \sqsupseteq \xi$ so that $\ell(\xi'(i)) = N$ for each $i < d$, $(\sigma', \xi') \in \text{Pair}$, and $\mathcal{K}(\sigma) = \mathcal{K}(\sigma') = \text{Cl}_\Delta(\xi')$.

If ξ maps to one level, say $\text{Im}(\xi) \subseteq T^\Delta(m)$, and $\mathcal{K}(\sigma) = \text{Cl}_\Delta(\xi)$, then N only needs to satisfy $N \geq m$.

Remark. Suppose $i < d$ and $\mathfrak{p} = \text{PCon}(\xi(i))$, so that $(\sigma(i), \xi(i)) \in \mathfrak{p}$. Then since $\xi'(i) \sqsupseteq \xi(i)$ and $\mathcal{K}(\sigma) = \text{Cl}_\Delta(\xi')$, we will also have $\text{PCon}(\xi'(i)) = \mathfrak{p}$.

Proof. Suppose $i < d$ and that we have chosen $\sigma'(j)$ and $\xi'(j)$ for every $j < i$. If $\xi(i) = (h, 0^r)$ for some $r < \omega$ and $h \in k^u \setminus \text{Free}$, we simply set $\xi'(i) = (h, 0^N)$ and $\sigma'(i) = \sigma(i)$. If $\xi(i)$ is not of this form, write $\mathfrak{p} = \text{PCon}(\xi(i))$, pick some $M > \max(\{\ell(\sigma'(j)) : j < i\} \cup \{\ell(\sigma_j) : j < d\})$ and run Proposition 6.3.9 on $(\text{Left}(\sigma(i), M), \xi(i)) \in \text{Pair}(\mathfrak{p}))$ to obtain $\sigma'(i) \sqsupseteq \text{Left}(\sigma(i), M)$ and $\xi'(i) \sqsupseteq \xi(i)$ with $\ell(\xi'(i)) = N$ and $(\sigma'(i), \xi'(i)) \in \text{Pair}(\mathfrak{p}_i)$. Then $\mathcal{K}(\sigma') = \mathcal{K}(\sigma)$ by construction, the key here being that at each stage, we moved up and left beyond anything built so far before applying Proposition 6.4.3. Proposition 6.3.5 gives us that $\text{Cl}_\Delta(\xi') \supseteq \mathcal{K}(\sigma)$. So in the case that $\text{Im}(\xi) \subseteq T^\Delta(m)$ and $\mathcal{K}(\xi) = \mathcal{K}(\sigma)$, we have $\mathcal{K}(\xi') = \mathcal{K}(\sigma)$. Otherwise, we use Lemma 6.3.8 to conclude that $\mathcal{K}(\xi') = \mathcal{K}(\sigma)$. \square

There are now three cases to consider depending on $\text{Lab}(\Theta(m))$. The splitting case is by far the most challenging, so we address it first.

Case 1: $\text{Lab}(\Theta(m)) = \text{Sp}$. Note that $\text{Size}(\Theta(m)) = d$. Write $\text{Pos}(\Theta(m)) = i < d$, and write $\text{Sort}(\Theta(m))(i) = h < k^u$. This case splits into three further sub-cases. Two of these cases are quite similar. In the first case, we have $h \in \text{Free}$. Write $\text{Free}(\text{Sort}(\Theta(m))(i)) = (q, j)$.

In the second case, we have $h \notin \text{Free}$, but with $t_i \neq (h, 0^m)$. We note that $\text{Cl}_\Theta(t_i) = \text{Cl}_\Delta(\xi(i)) = \mathcal{K}(\sigma(i)) \neq \mathcal{K}(\emptyset_h)$. It follows that $\sigma(i) \neq (h, 0^{\ell(\sigma(i))})$. Write $q = \sigma(i)(\text{Start}(\sigma(i)))$, and write $c(\text{Start}(\sigma(i)))^u = j$.

For both cases, the proof is now identical with these choices of $q < k$ and $j < k^u$. Find some $r > \max\{\ell(u) : u \in \text{Im}(\sigma)\}$ so that the following both hold:

- $c(r) = (j, 0^r)$,
- $\ell(\psi \circ c(r)) > \max\{\ell(\xi(\alpha)) : \alpha < d\}$.

It follows that we have

$$\mathcal{K}(\sigma(i)) = \mathcal{K}(\text{Left}(\sigma(i), r+1)) = \mathcal{K}(\text{Left}(\sigma(i), r) \frown q).$$

In particular, writing $\mathfrak{p} = \text{PCon}(\xi(i))$, we have that both $(\text{Left}(\sigma(i), r+1), \xi(i))$ and $(\text{Left}(\sigma(i), r) \frown q, \xi(i))$ are in $\text{Pair}(\mathfrak{p})$. Using Proposition 6.4.3, find $w_0 \sqsupseteq \text{Left}(\sigma(i), r+1)$, $w_q \sqsupseteq \text{Left}(\sigma(i), r) \frown q$, and $x_0, x_q \sqsupseteq \xi(i)$ with the following properties:

- $\ell(x_0) = \ell(x_q) = \ell(\psi \circ c(r)) + 1$,
- $(w_0, x_0), (w_q, x_q) \in \text{Pair}(\mathfrak{p})$.
- $\mathcal{K}(\langle w_0, w_q \rangle) = \max(\mathfrak{p})^{\delta_{0,1}}$.

Notice that since ψ is induced by an isomorphism of structures, we must have that $x_0(\ell(\psi \circ c(r))) = 0$ and $x_q(\ell(\psi \circ c(r))) = q$. In particular, $x_0 \neq x_q$. We set $\tilde{f}(m) = \ell(x_0 \wedge x_q)$ and $f(t_i) = x_0 \wedge x_q$. Put $y_i = f(t_i) \frown 0$ and $y_{i+1} = f(t_i) \frown 1$. We also rename w_0 to either w_i or w_{i+1} depending on if $x_0|_{\tilde{f}(m)+1} = y_i$ or y_{i+1} . We similarly rename w_q to the other of w_i or w_{i+1} . Define $\sigma' : (d+1) \rightarrow \text{CT}$ and $\xi' : (d+1) \rightarrow T^\Delta$ be given by

$$\sigma'(b) = \begin{cases} \sigma \circ \delta_{i,d}(b) & \text{if } b \notin \{i, i+1\} \\ w_b & \text{if } b \in \{i, i+1\} \end{cases}$$

$$\xi'(b) = \begin{cases} \xi \circ \delta_{i,d}(b) & \text{if } b \notin \{i, i+1\} \\ y_b & \text{if } b \in \{i, i+1\}. \end{cases}$$

We note that $\mathcal{K}(\sigma') = \mathcal{K}(\sigma)^{\delta_{i,d}} = \text{Cl}_\Theta(T^\Theta(m))^{\delta_{i,d}}$. Now use Proposition 6.4.3 on (σ', ξ') using a suitably large $N > \tilde{f}(m) + 1$. Doing this, we obtain functions $\sigma'' : (d+1) \rightarrow \text{CT}$ and $\xi'' : (d+1) \rightarrow T^\Delta(N)$ with $\sigma'' \sqsupseteq \sigma'$, $\xi'' \sqsupseteq \xi'$, $(\sigma'', \xi'') \in \text{Pair}$, and $\mathcal{K}(\sigma'') = \mathcal{K}(\sigma') = \text{Cl}_\Delta(\xi'')$. For each $b \in d \setminus \{i\}$, we now set

$$f(t_b) = (\xi'' \circ \epsilon_{i,d+1}(b))|_{\tilde{f}(m)}.$$

Note that if we write $f^+[T^\Theta(m+1)] = \{z_0 \preceq_{\text{lex}} \cdots \preceq_{\text{lex}} z_d\}$, we have $\xi''(b) \sqsupseteq z_b$ for each $b < d+1$. As σ'' and ξ'' will play the roles of σ and ξ when continuing the construction to level $m+1$, it only remains to verify that $\text{Cl}_\Delta(f^+[T^\Theta(m+1)]) = \text{Cl}_\Theta(T^\Theta(m+1))$. Looking at the right hand side, we have:

$$\begin{aligned} \text{Cl}_\Theta(T^\Theta(m+1)) &= \text{Cl}_\Theta(T^\Theta(m))^{\delta_{i,d}} \\ &= \mathcal{K}(\sigma') \\ &= \text{Cl}_\Delta(\xi'') \\ &\subseteq \text{Cl}_\Delta(f^+[T^\Theta(m+1)]). \end{aligned}$$

The reverse inequality follows from the observation that

$$\begin{aligned}\mathrm{Cl}_\Delta(f^+[T^\Theta(m+1)]) &\subseteq \mathrm{Cl}_\Delta(f[T^\Theta(m)])^{\delta_{i,d}} \\ &= \mathrm{Cl}_\Theta(T^\Theta(m))^{\delta_{i,d}} \\ &= \mathrm{Cl}_\Theta(T^\Theta(m+1)).\end{aligned}$$

In the third case, we have $h \notin \text{Free}$ and $t_i = (h, 0^m)$. Since we have $\mathrm{Cl}_\Delta(\xi) = \mathrm{Cl}_\Delta(f^+[T^\Theta(m)])$, we must also have $\xi(i)$ on the left branch of T_h . Since Θ has controlled paths, let $\mathbf{p} \in \max(\vec{P}_h)$ be the path so that every coding node of T^Θ with start level m has path \mathbf{p} . We first apply Proposition 6.3.9 to find $u \sqsupseteq \sigma(i)$ and $v \sqsupseteq \xi(i)$ with $(u, v) \in \text{Pair}(\mathbf{p}')$. We set $\tilde{f}(m) = \text{Start}(v)$. Form the functions $\sigma': (d+1) \rightarrow \text{CT}$ and $\xi': (d+1) \rightarrow T^\Delta$ as follows.

$$\begin{aligned}\sigma'(b) &= \begin{cases} \sigma \circ \delta_{i,d}(b) & \text{if } b \neq i+1 \\ u & \text{if } b = i+1 \end{cases} \\ \xi'(b) &= \begin{cases} \xi \circ \delta_{i,d}(b) & \text{if } b \neq i+1 \\ v & \text{if } b = i+1. \end{cases}\end{aligned}$$

Writing $\mathcal{D} = \max(\mathbf{p}')$, we note that $\mathcal{K}(\sigma') = (\mathcal{K}(\sigma)^{\delta_{i,d}})|_{(\iota_{i+1}, \mathcal{D})} = \mathrm{Cl}_\Theta(T^\Theta(m+1))$. Run Proposition 6.4.3 on (σ', ξ') using a suitably large $N > \tilde{f}(m) + 1$. Doing this, we obtain functions $\sigma'': (d+1) \rightarrow \text{CT}$ and $\xi'': (d+1) \rightarrow T^\Delta(N)$ with $\sigma'' \sqsupseteq \sigma'$, $\xi'' \sqsupseteq \xi'$, $(\sigma'', \xi'') \in \text{Pair}$, and $\mathcal{K}(\sigma') = \mathcal{K}(\sigma'') = \mathrm{Cl}_\Delta(\xi'')$. We note that by the equality of the age-sets, we must have $\xi''(i)$ on the left branch of T_h . For each $b \in d \setminus \{i\}$, we now set

$$f(t_b) = (\xi'' \circ \epsilon_{i,d+1}(b))|_{\tilde{f}(m)}.$$

As in the first two cases, it only remains to verify that $\mathrm{Cl}_\Delta(f^+[T^\Theta(m+1)]) = \mathrm{Cl}_\Theta(T^\Theta(m+1))$. Looking at the right hand side, we have:

$$\begin{aligned}\mathrm{Cl}_\Theta(T^\Theta(m+1)) &= (\mathrm{Cl}_\Theta(T^\Theta(m))^{\delta_{i,d}})|_{(\iota_{i+1}, d, \mathcal{D})} \\ &= \mathcal{K}(\sigma') \\ &= \mathrm{Cl}_\Delta(\xi'') \\ &\subseteq \mathrm{Cl}_\Delta(f^+[T^\Theta(m+1)]).\end{aligned}$$

For the reverse inequality, we first note that

$$\begin{aligned}\mathrm{Cl}_\Delta(f^+[T^\Theta(m+1)]) &\subseteq \mathrm{Cl}_\Delta(f[T^\Theta(m)])^{\delta_{i,d}} \\ &= \mathrm{Cl}_\Theta(T^\Theta(m))^{\delta_{i,d}}.\end{aligned}$$

Now we recall that since $h \notin \text{Free}$, we have $\mathrm{Cl}_\Theta(T^\Theta(m))^{\delta_{i,d}} \notin P(T^\Theta(m+1)^u)$. Finally, as no members of $P(T^\Theta(m+1)^u)$ lie strictly between $(\mathrm{Cl}_\Theta(T^\Theta(m))^{\delta_{i,d}})|_{(\iota_{i+1}, d, \mathcal{D})}$ and $\mathrm{Cl}_\Theta(T^\Theta(m))^{\delta_{i,d}}$, we must have $\mathrm{Cl}_\Delta(f^+[T^\Theta(m+1)]) = \mathrm{Cl}_\Theta(T^\Theta(m+1))$ as desired.

Case 2: $\text{Lab}(\Theta(m)) = \text{AC}$. Write $\rho = T^\Theta(m)^u$, and write $e: d_0 \rightarrow d$ for the unique increasing injection with $(\mathrm{Cl}_\Theta(T^\Theta(m))^e, \mathrm{Cl}_\Theta(T^\Theta(m+1))^e) \in \text{ECon}(\rho \circ e)$ (Corollary 4.5.3). There are two cases. If $d_0 > 1$, we first find a function $\sigma': d \rightarrow$

CT with $\sigma' \sqsupseteq \sigma$ so that $\mathcal{K}(\sigma') = \text{Cl}_\Theta(T^\Theta(m+1))$. In particular, we still have $(\sigma', \xi) \in \text{Pair}$, so run Proposition 6.4.3 on (σ', ξ) with some suitably large N , obtaining functions $\sigma'' : (d+1) \rightarrow \text{CT}$ and $\xi'' : (d+1) \rightarrow T^\Delta(N)$ with $\sigma'' \sqsupseteq \sigma'$, $\xi'' \sqsupseteq \xi$, $(\sigma'', \xi'') \in \text{Pair}$, and $\mathcal{K}(\sigma') = \mathcal{K}(\sigma'') = \text{Cl}_\Delta(\xi'')$. To determine f , we search between the levels of ξ and ξ'' to find the level of the age-change, and call this level $\tilde{f}(m)$, and then set $f(t_b) = \xi''(b)|_{\tilde{f}(m)}$.

If $d_0 = 1$, suppose $e = \nu_i$ for some $i < d$. Set $\mathfrak{p} = \text{PCon}(\xi(i))$. We use Proposition 6.3.9 to find $(u, v) \in \text{Pair}(\mathfrak{p}')$ with $u \sqsupseteq \sigma(i)$ and $v \sqsupseteq \xi(i)$. We then consider the functions σ' and ξ' , which are identical to σ and ξ except we set $\sigma'(i) = u$ and $\xi'(i) = v$, and then apply Proposition 6.4.3. The rest of the proof is similar to the $d > 1$ case.

Case 3: $\text{Lab}(\Theta(m)) = \text{Cd}$. Write $\text{Pos}(\Theta(m)) = j < d$ and $\text{Sort}(\Theta(m))(j) = i < k^u$. We set $\mathcal{A} = \text{Cl}_\Theta(T^\Theta(m))$, so also $\mathcal{K}(\sigma) = \text{Cl}_\Delta(\xi) = \mathcal{A}$. We also write $\varphi = \text{Pass}(\Theta(m))$. Saying that m is a controlled coding level of Θ amounts to saying that $(\mathcal{A}, j, \varphi)$ is a controlled coding triple. In particular, by item 1 of Definition 5.6.3, we have $\text{Cl}_\Theta(\xi(j)) = P_i^{\min}$, implying that $(\sigma(j), \xi(j)) \in \text{Pair}(P_i^{\min})$. So find r so that $c(r) \sqsupseteq \sigma(j)$ and $\psi \circ c(r) \sqsupseteq \xi(j)$. Define $\sigma' : (d-1) \rightarrow \text{CT}$ by setting $\sigma'(\alpha) = \text{Left}(\sigma \circ \epsilon_{j,d}(\alpha), r) \frown \varphi(\alpha)$. Note that $\mathcal{K}(\sigma') = \mathcal{A}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1}$, so by item 2 of Definition 5.6.3, we also have $\mathcal{K}(\sigma') = \mathcal{A}^{\epsilon_{j,d}}$.

We now run Proposition 6.4.3 on $(\sigma', \xi \circ \epsilon_{j,d})$ using some suitably large $N > r$. Doing this, we obtain functions $\sigma'' : (d-1) \rightarrow \text{CT}$ and $\xi'' : (d-1) \rightarrow T^\Delta(N)$ with $\sigma'' \sqsupseteq \sigma'$, $\xi'' \sqsupseteq \xi \circ \epsilon_{j,d}$, $(\sigma'', \xi'') \in \text{Pair}$, and $\mathcal{K}(\sigma') = \mathcal{K}(\sigma'') = \text{Cl}_\Delta(\xi'')$. We note that since ψ is an isomorphism of structures and since $\sigma''(\alpha) \sqsupseteq \sigma'(\alpha)$ for each $\alpha < d-1$, we must have $\xi''(\alpha)(r) = \varphi(\alpha)$. We now set $\tilde{f}(m) = \ell(\psi \circ c(r))$, $f(t_j) = \psi \circ c(r)$, and for $b \in d \setminus \{j\}$, we set $f(t_b) = \xi'' \circ \delta_{j,d-1}(b)|_{\tilde{f}(m)}$. Lastly, to show that $\text{Cl}_\Delta(f[T^\Theta(m)]) = \mathcal{A}$, write $\mathcal{B} = \text{Cl}_\Delta(f[T^\Theta(m)])$. Then $\mathcal{B} \subseteq \mathcal{A}$, $\mathcal{B}^{\epsilon_{j,d}} = \mathcal{A}^{\epsilon_{j,d}}$, and $\mathcal{B}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1} = \mathcal{A}\langle j, \epsilon_{j,d}, \varphi \rangle^{-1}$. As $(\mathcal{A}, j, \varphi)$ is a controlled coding triple, item 3 of Definition 5.6.3 gives us that $\mathcal{A} = \mathcal{B}$.

7 Degenerate unary predicates

Up until now, we have been assuming that every $i < k^u$ is a non-degenerate unary predicate, meaning that there is some $\mathbf{A} \in \mathcal{K}$ with underlying set 2 such that $U^{\mathbf{A}}(0) = i$ and $R^{\mathbf{A}}(0, 1) \neq 0$. In other words, a non-degenerate unary predicate is able to participate in some non-trivial binary relation. We now briefly discuss the modifications that need to be made when there are degenerate unary predicates; luckily, these modifications are all very straightforward. The idea is that for degenerate unaries, all the coding nodes will be on the left branch, and these coding levels can appear interspersed almost everywhere in a diagonal diary.

Notation 7.0.1. We now assume for some $k^d \leq k^u$ that every $i < k^d$ is degenerate. With \mathbf{K} denoting a fixed enumerated left-dense Fraïssé limit as before, we have for each $i < k^d$ that $\{a \in \mathbf{K} : U(a) = i\}$ is an infinite set of isolated points. Letting $\mathbf{K}_0 = \{a \in \mathbf{K} : k^d \leq U(a) < k^u\} \subseteq \mathbf{K}$ denote the induced substructure on the points with non-degenerate unary predicate, we have $\text{Aut}(\mathbf{K}) \cong \text{Aut}(\mathbf{K}_0) \times (S_\infty)^{k^d}$.

If $i < k^d$, then the set $\{c(a) : a \in \mathbf{K}, U(a) = i\}$ of coding nodes for points with unary i all lie on the left branch of T_i . In particular, it is impossible to ask for an anti-chain of coding nodes, as we do in Definition 5.2.2. However, this in many ways makes our work much easier, as the tree T_i will not meaningfully participate in splitting or age-change levels. We now collect all of the new conventions and definition modifications needed to define diagonal substructures and diaries in this slightly more general setting.

1. Proposition 4.2.4 must be modified as follows. Given a non-decreasing function $\rho: d \rightarrow k^u$, we have $P(\rho) \neq \emptyset$ iff for each $i < k^d$, we have $|\rho^{-1}(\{i\})| \leq 1$.
2. Degenerate unary predicates do not have a free relation. If $i < k^d$, then $P(\iota_i) = \{\mathcal{K}(\emptyset_i)\}$. All results from earlier which mention some $j \in k^u \setminus \mathbf{Free}$ should now be interpreted as $j \in k^u \setminus (\mathbf{Free} \cup k^d)$.
3. In the definition of a weakly diagonal substructure (Definition 5.2.2), we only demand items 1 and 5 for the non-degenerate vertices/coding nodes. We emphasize that if $\mathbf{A} \subseteq \mathbf{K}$ is weakly diagonal, then all levels of $\mathbf{FreeStart}(\mathbf{A})$ must come before any coding nodes, even degenerate ones.
4. The definitions of maximal/controlled paths (Definitions 5.3.2 and 5.3.4) remain the same.
5. The definition of a controlled age-change level (Definition 5.5.1) remains the same. In particular, these levels will never meaningfully involve any degenerate unary predicates.
6. Given $\mathbf{A} \subseteq \mathbf{K}$ and $a \in \mathbf{A}$ with $U(a) < k^d$, we simply declare that a is a controlled coding level of \mathbf{A} . We only use Definition 5.6.5 for $a \in \mathbf{A}$ with $a \notin k^d$.
7. The definition of a controlled splitting level (Definition 5.7.1) remains the same. In particular, these levels will never meaningfully involve any degenerate unary predicates.
8. With the modifications given above, the definition of a substructure with controlled critical levels and of a diagonal substructure (Definition 5.8.2) remain the same.
9. Given a weakly diagonal $\mathbf{A} \subseteq \mathbf{K}$, the definition of the diary of \mathbf{A} (Definition 6.1.1) remains the same. If $\mathbf{A} \subseteq \mathbf{K}$ is diagonal, then writing $\mathbf{A}_0 = \mathbf{A} \cap \mathbf{K}_0$, we have $\overline{\mathbf{A}} = \mathbf{A} \cup \overline{\mathbf{A}_0}$. In particular, the only extra information contained in $\overline{\mathbf{A}}$ which is not already in $\overline{\mathbf{A}_0}$ is the location of degenerate coding levels. As before, one can compute $\overline{\mathbf{A}}$ using only $\Delta^{\mathbf{A}}$ along with the knowledge that $\mathbf{A} \subseteq \mathbf{K}$ is diagonal.
10. In the definition of an (abstract) diary (Definition 6.2.1), we modify item 3 of the second bullet as follows. If $\mathbf{Lab}(\Delta(n)) = \mathbf{Cd}$ and $j = \mathbf{Pos}(\Delta(n))$, and $\mathbf{Sort}(\Delta(n))(j) = i < k^d$, then there are two cases. If $n < \mathbf{dom}(\Delta)$ is largest with $\mathbf{Lab}(\Delta(n)) = \mathbf{Cd}$ and $\mathbf{Sort}(\Delta(n))(\mathbf{Pos}(\Delta(n))) = i$, then we have $\mathbf{Sort}(\Delta(n+1)) = \mathbf{Sort}(\Delta(n)) \circ \epsilon_{j,d}$ (i.e., position j is a terminal coding node). If $n < \mathbf{dom}(\Delta)$ is not largest, then we have $\mathbf{Sort}(\Delta(n+1)) = \mathbf{Sort}(\Delta(n))$ (i.e., position j is a *non-terminal* degenerate coding node).

11. Given a diary Δ , the last bullet of the definition of T^Δ (Definition 6.2.2) is modified in the obvious way to include the case that the relevant coding node is a non-terminal degenerate coding node.
12. In the definition of a diary with controlled critical levels (Definition 6.2.3), item 4 must be modified to include the case that the relevant coding node is degenerate. In this case, we simply demand that there is no age-change and that the passing function is constant zero.
13. The definition of a sub-diary (Definition 6.2.7) remains almost the same, keeping in mind that when defining the passing function in the coding level case, the domain of $\text{Pass}(\text{Sub}^\Delta(\mathbf{A})(n))$ could be d instead of $d - 1$.

8 Big Ramsey structures

Our characterization of big Ramsey degrees given in Theorem 6.2.14 in fact yields an a priori stronger result. Not only does each $\mathbf{A} \in \mathcal{K}$ have finite big Ramsey degree, but there is a single expansion of \mathbf{K} which describes the exact big Ramsey degrees for every $\mathbf{A} \in \mathcal{K}$. The following definition is from [31].

Definition 8.0.1. We say that \mathbf{K} admits a big Ramsey structure if there is a language $\mathcal{L}' \supseteq \mathcal{L}$ and an \mathcal{L}' -structure \mathbf{K}' so that the following all hold.

1. $\mathbf{K}'|_{\mathcal{L}} = \mathbf{K}$.
2. Every $\mathbf{A} \in \mathcal{K}$ has finitely many \mathcal{L}' -expansions to a structure $\mathbf{A}' \in \text{Age}(\mathbf{K}')$; denote the set of expansions by $\mathbf{K}'(\mathbf{A})$.
3. Every $\mathbf{A} \in \mathcal{K}$ has big Ramsey degree $|\mathbf{K}'(\mathbf{A})|$.
4. For each $\mathbf{A} \in \mathcal{K}$, the function $\gamma_{\mathbf{A}} : \text{Emb}(\mathbf{A}, \mathbf{K}) \rightarrow \mathbf{K}'(\mathbf{A})$ given by $\gamma_{\mathbf{A}}(f) = \mathbf{K}' \cdot f$ witnesses the fact that the big Ramsey degree of \mathbf{A} is not less than $|\mathbf{K}'(\mathbf{A})|$.

Call a structure \mathbf{K}' satisfying (1)-(4) a *big Ramsey structure* for \mathbf{K} .

It is an open problem whether or not given a Fraïssé class, the property of having finite big Ramsey degrees implies that the Fraïssé limit admits a big Ramsey structure. The crucial difference is that a big Ramsey structure provides a coherent collection of “unavoidable” colorings of $\text{Emb}(\mathbf{A}, \mathbf{K})$ for every $\mathbf{A} \in \mathcal{K}$ simultaneously.

Theorem 8.0.2. *The Fraïssé structure $\mathbf{K} = \text{Flim}(\text{Forb}(\mathcal{F}))$ admits a big Ramsey structure.*

Proof. For now, fix an enumerated $\mathbf{A} \in \mathcal{K}$. Given $f \in \text{Emb}(\mathbf{A}, \mathbf{K})$, let $\sigma_f \in \text{Sym}(\mathbf{A})$ denote the permutation of \mathbf{A} such that $f \circ \sigma_f$ is an increasing injection.

Fix a diagonal diary Δ with $\text{Str}(\Delta) = \mathbf{K}$. For each enumerated $\mathbf{A} \in \mathcal{K}$ and each diagonal diary Θ which codes an enumerated structure of size n , we add an n -ary relational symbol R_Θ to \mathcal{L}' . To expand \mathbf{K} into an \mathcal{L}' -structure \mathbf{K}' , given an n -tuple $\{a_0, \dots, a_{n-1}\} \subseteq \mathbf{K}$, we declare that $R_\Theta^{\mathbf{K}'}(a_0, \dots, a_{n-1})$ holds iff $a_0 < \dots < a_{n-1}$ and $\text{Sub}^\Delta(\{a_0, \dots, a_{n-1}\}) = \Theta$.

The key observation is that the notion of sub-diary is coherent in the following sense: suppose $a_0 < \dots < a_{n-1} \in \mathbf{K}$ and $b_0 < \dots < b_{n-1} \in \mathbf{K}$ are such that $\text{Sub}^\Delta(\{a_0, \dots, a_{n-1}\}) = \text{Sub}^\Delta(\{b_0, \dots, b_{n-1}\})$. Then if $f: m \rightarrow n$ is an increasing injection, we have $\text{Sub}^\Delta(\{a_{f(0)}, \dots, a_{f(m-1)}\}) = \text{Sub}^\Delta(\{b_{f(0)}, \dots, b_{f(m-1)}\})$. It follows that $\mathbf{K}'|_{\{a_0, \dots, a_{n-1}\}} \cong \mathbf{K}'|_{\{b_0, \dots, b_{n-1}\}}$. From this, it follows that \mathbf{K}' is a big Ramsey structure from Theorem 6.2.14 and Proposition 2.0.4. \square

The language $\mathcal{L}' \supseteq \mathcal{L}$ we used in the proof of Theorem 8.0.2 is much larger than we need. Our next goal is to show that we can in fact take \mathcal{L}' to be a finite relational language. The maximum arity of the new relational symbols we need depends on the size of the largest member of \mathcal{F} . We need a lemma which could have appeared back in Section 4

Lemma 8.0.3. *Write $D = \max\{|\mathbf{F}| : \mathbf{F} \in \mathcal{F}\}$. Suppose $\rho: d \rightarrow k^u$ is a sort and that γ and ζ are gluings with sort ρ . If $\mathcal{K}(\gamma) \neq \mathcal{K}(\zeta)$, then there is some $d_0 < D$ and some increasing injection $e: d_0 \rightarrow d$ with $\mathcal{K}(\gamma^e) \neq \mathcal{K}(\zeta^e)$.*

In particular, if $\text{ECon}(\rho) \neq \emptyset$, then we have $d < D$.

Proof. Without loss of generality, suppose $\mathbf{B} \in \mathcal{K}(\zeta) \setminus \mathcal{K}(\gamma)$. Find $\mathbf{F} \in \mathcal{F}$ and $g: \mathbf{F} \rightarrow \mathbf{B}[\gamma]$ which witnesses that $\mathbf{B} \notin \mathcal{K}(\gamma)$. Write $\mathbf{J} = g^{-1}(\mathbf{B})$, which we view as an induced substructure of \mathbf{F} . Notice that since $\mathbf{B}[\zeta] \in \mathcal{K}$ and both ζ and γ have sort ρ , we have $\mathbf{J} \in \mathcal{K}$. In particular, $\mathbf{J} \subsetneq \mathbf{F}$, so $|\mathbf{J}| < D$. Set

$$X = \{i < d : \exists x \in g[\mathbf{J}] \text{ with } V^{\mathbf{B}}(x) = i\},$$

and set $d_0 = |X|$. Note that $d_0 \leq |g[\mathbf{J}]| < D$. Let $e: d_0 \rightarrow d$ be the increasing enumeration of X . Then considering the \mathcal{L}_d -structure \mathbf{B}_0 induced on $g[\mathbf{J}] \subseteq \mathbf{B}$, we have that $\mathbf{B}_0[\zeta^e] \subseteq \mathbf{B}[\zeta]$ and $\mathbf{B}_0[\gamma^e] \subseteq \mathbf{B}[\gamma]$ are induced substructures, and $g[\mathbf{F}] \subseteq \mathbf{B}_0[\gamma^e]$. This shows that $\mathbf{B}_0 \in \mathcal{K}(\zeta^e) \setminus \mathcal{K}(\gamma^e)$. \square

Theorem 8.0.4. *\mathbf{K} admits a big Ramsey structure \mathbf{K}' in a finite relational language \mathcal{L}' . Letting $D = \max\{|\mathbf{F}| : \mathbf{F} \in \mathcal{F}\}$, we can take the language \mathcal{L}' to have relational symbols of arity at most $\max\{2(D-1), 4\}$.*

Proof. Fix a diagonal diary Δ with $\text{Str}(\Delta) = \mathbf{K}$. The idea is to create a language \mathcal{L}' which can code enough information about T^Δ that we can in fact recover all of T^Δ . To do this, we add the following new relational symbols to \mathcal{L}' and describe how they are interpreted to form the expansion \mathbf{K}' .

- \preceq_{lex} , which we interpret in the usual sense, i.e., $a \preceq_{lex}^{\mathbf{K}'} b$ iff $c_\Delta(a) \preceq_{lex} c_\Delta(b)$.
- S , a 4-ary relational symbol which describes the relative levels of meets. More precisely, given $a, b, m, n \in \mathbf{K}$, we set $S^{\mathbf{K}'}(a, b, m, n)$ iff $\ell(c_\Delta(a) \wedge c_\Delta(b)) < \ell(c_\Delta(m) \wedge c_\Delta(n))$. By considering $a = b$ and/or $m = n$, we can compare the levels of meets with coding nodes or the levels of two coding nodes.
- Suppose $\rho: d \rightarrow k^u$ is a sort and that $(\mathcal{A}, \mathcal{B}) \in \text{ECon}(\rho)$. Then we introduce $X_{(\mathcal{A}, \mathcal{B})}$ a $(d+2)$ -ary relational symbol. Given $a_0, \dots, a_{d-1}, b_0, b_1 \in \mathbf{K}$, we declare that $X_{(\mathcal{A}, \mathcal{B})}^{\mathbf{K}'}(a_0, \dots, a_{d-1}, b_0, b_1)$ holds iff $a_0 \preceq_{lex} \dots \preceq_{lex} a_{d-1}$,

there is a (necessarily unique) level m with $\text{Cl}_\Delta(c_\Delta[\{a_0, \dots, a_{d-1}\}]|_m) = \mathcal{A}$ and $\text{Cl}_\Delta(c_\Delta[\{a_0, \dots, a_{d-1}\}]|_{m+1}) = \mathcal{B}$, and $m < \ell(c_\Delta(b_0) \wedge c_\Delta(b_1))$. By considering $b_0 = b_1$, we can compare the level of the age change with the level of a coding node.

- Suppose for each $i \in \{0, 1\}$ that $\rho_i: d_i \rightarrow k^u$ are sorts and that $(\mathcal{A}_i, \mathcal{B}_i) \in \text{ECon}(\rho_i)$. We introduce $Y_{(\mathcal{A}_0, \mathcal{B}_0), (\mathcal{A}_1, \mathcal{B}_1)}$ a $(d_0 + d_1)$ -ary relational symbol, which for now we shorten to Y . Given $a_0, \dots, a_{d_0-1}, b_0, \dots, b_{d_1-1} \in \mathcal{K}$, we declare that $Y^{\mathbf{K}'}(\vec{a}, \vec{b})$ holds iff $a_0 \preceq_{\text{lex}} \dots \preceq_{\text{lex}} a_{d_0-1}, b_0 \preceq_{\text{lex}} \dots \preceq_{\text{lex}} b_{d_1-1}$, there are (necessarily unique) levels m_0 and m_1 where $\text{Cl}_\Delta(c_\Delta[\vec{a}]|_{m_0}) = \mathcal{A}_0$, $\text{Cl}_\Delta(c_\Delta[\vec{b}]|_{m_0+1}) = \mathcal{B}_0$, $\text{Cl}_\Delta(c_\Delta[\vec{b}]|_{m_1}) = \mathcal{A}_1$, $\text{Cl}_\Delta(c_\Delta[\vec{b}]|_{m_1+1}) = \mathcal{B}_1$, and we have $m_0 < m_1$.

To see that \mathbf{K}' is a big Ramsey structure for \mathbf{K} , it suffices to note that \mathbf{K}' contains enough information about T^Δ to reconstruct T^Δ entirely. \square

The fact that \mathbf{K} admits a big Ramsey structure allows us to give a dynamical interpretation to the results of this paper (see [31] for more discussion). Given a topological group G , recall that the *left completion* of G , denoted \widehat{G} , is the completion of G with respect to its left uniform structure. One can endow \widehat{G} with the structure of a topological semigroup; for our purposes, all we need is that if $G = \text{Aut}(\mathbf{K})$, then $\widehat{G} = \text{Emb}(\mathbf{K}, \mathbf{K})$. Now if $X \times G \rightarrow X$ is a right G -flow, the action continuously extends to an action $X \times \widehat{G} \rightarrow X$. A point $x_0 \in X$ is called a *completion point* if for any $\eta \in \widehat{G}$, the point $x_0 \cdot \eta \in X$ has dense orbit. The flow X is called a *completion flow* if X contains a completion point. Theorem 1.6 from [31] now gives us the following.

Theorem 8.0.5. *The group $G = \text{Aut}(\mathbf{K})$ admits a metrizable universal completion flow, i.e., a completion flow which admits a surjective G -map to any other completion flow.*

This completion flow is simply the orbit closure of the big Ramsey structure \mathbf{K}' in the space of all \mathcal{L}' -structures with underlying set ω .

As a final remark, we observe that Theorem 6.2.14 actually gives us a somewhat stronger result. Given a G -flow X and $x_0 \in X$, let us call x_0 a *strong completion point* if x_0 has dense orbit and if for any $\eta \in \text{Emb}(\mathbf{K}, \mathbf{K})$, there is $\theta \in \text{Emb}(\mathbf{K}, \mathbf{K})$ with $x_0 \cdot \eta \cdot \theta = x_0$. Similarly, we can call X a *strong completion flow* if X contains some strong completion point. We now have the following.

Theorem 8.0.6. *The universal completion flow of the group $G = \text{Aut}(\mathbf{K})$ is a strong completion flow. In particular, every completion flow of G is a strong completion flow.*

Proof. This is more-or-less a rephrasing of Theorem 6.2.14. Start with a diagonal diary Δ with $\text{Str}(\Delta) = \mathbf{K}$ and form \mathbf{K}' as in Theorem 8.0.2 or 8.0.4. So \mathbf{K}' is a completion point of its orbit closure, and we show that it is a strong completion point. Let $\eta \in \text{Emb}(\mathbf{K}, \mathbf{K})$, and write $\mathbf{A} = \eta[\mathbf{K}]$. Then $\text{Sub}^\Delta(\mathbf{A})$ is another diary coding the Fraïssé limit of \mathcal{K} , so Theorem 6.2.14 yields an embedding of

Δ into $\text{Sub}^\Delta(\mathbf{A})$. This embedding gives rise to an embedding θ of $\mathbf{Str}(\Delta)$ into $\mathbf{Str}(\text{Sub}^\Delta(\mathbf{A}))$, i.e., an element of $\text{Emb}(\mathbf{K}, \mathbf{K})$. It follows that $\mathbf{K}' \cdot \eta \cdot \theta = \mathbf{K}'$ as desired. \square

In light of this result, we can add to the list of intriguing open questions about big Ramsey structures and universal completion flows. In particular, is there any topological group G which admits some completion flow which is not a strong completion flow?

Acknowledgement

J. H. and M. K. are supported by the project 21-10775S of the Czech Science Foundation (GAČR). This article is part of a project that has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 810115). M. B. and J. H. were supported by the Center for Foundations of Modern Computer Science (Charles University project UNCE/SCI/004). M. K. was supported by the Charles University Grant Agency (GA UK), project 378119. N. D. is supported by National Science Foundation grant DMS-1901753. A. Z. is supported by National Science Foundation grant DMS-2054302. L. V. is supported by Beatriu de Pinós BP2018, funded by the AGAUR (Government of Catalonia) and by the Horizon 2020 programme No 801370.

Index

- $(\rho, e) \cdot \zeta$, 17
- $P(\rho), P(d)$, 13
- T , 9
 - $T_i, T(n), T(<n)$, 9
 - $\text{Left}(s, n)$, 10
 - $\text{IS}(s)$, 9
 - $\text{Succ}(s)$, 9
 - $s \preceq_{\text{lex}} t$, 10
 - $s \sqsupseteq t$, 9
 - t^u, t^b , 9
- $\text{AEmb}(\text{CT}^{\mathbf{A}}, \text{CT}^{\mathbf{B}})$, 26
- $\mathbf{A} \cdot f$, 8
- \mathbf{K} , 10
 - R , 10
 - U , 10
 - CT , 10
 - c , 10
- $\text{Con}(\rho)$, 21
- $\text{CT}^{\mathbf{A}}$, 10
- \mathcal{F} , 10
- \mathcal{I} , 19
- \mathcal{K} , 10
- \mathcal{L} , 9
 - $R^{\mathbf{A}}$, 9
 - R_i , 9
 - $U^{\mathbf{A}}$, 9
 - U_i , 9
 - $\mathcal{L}^u, \mathcal{L}^b$, 9
 - Flip , 9
 - k , 9
 - k^u , 9
- \mathcal{L}_d , 11
 - $V^{\mathbf{B}}$, 11
 - V_n , 11
 - \mathbf{B}^e , 16
 - $\mathcal{A}|_{(e, \mathcal{B})}$, 17
 - \mathcal{A}^e , 16
 - $e \cdot \mathbf{B}$, 16
- γ^e , 17
- $\text{Cause}(\mathcal{A}, \mathcal{B})$, 21, 22
- $\text{Glue}(\rho, \mathbf{J})$, 20
- $\text{Spec}(\mathcal{A})$, 20
- $\text{Spec}(\gamma)$, 20
- $\text{OEmb}(\mathbf{A}, \mathbf{K})$, 8
- age of a gluing, 12
 - $\mathcal{K}(S)$, 12
 - $\mathcal{K}(\gamma)$, 12
 - $\mathcal{K}(\sigma)$, 12
 - $\mathcal{K}(\sigma, \mathbf{A})$, 12
- aged embedding, 26
 - $\text{AEmb}(\text{CT}^{\mathbf{A}}, \text{CT}^{\mathbf{B}})$, 26
- coding function, 10
 - $c^{\mathbf{A}}$, 10
- coding tree, 10
 - coding node, 10
- consecutive age change, 21
 - cause, 21
 - essential, 21
- enumerated structure, 8
- gluing, 11
- gluing map
 - $\gamma \cdot f, \gamma|_{\mathbf{J}}, \gamma_n, \gamma^e$, 19
- left dense Fraïssé limit, 10
- left-leaning embedding, 10
 - $\text{OL}(\mathbf{A}, \mathbf{K})$, 10
- non-degenerate, 13
- ordered embedding, 8
- Ramsey degree, 8
 - \mathcal{K} has finite big Ramsey degrees, 8
 - big Ramsey degree of $\mathbf{A} \in \mathcal{K}$, 8
 - of \mathbf{A} in \mathbf{K} , 8
 - ordered Ramsey degree, 8
- reasonable class, 14
- spectrum of a class, 20
- spectrum of a gluing, 20

References

- [1] Fred G. Abramson and Leo A. Harrington, Models without indiscernibles. *Journal of Symbolic Logic*, 43:572–600, 1978.
- [2] M. Balko, D. Chodounský, N. Dobrinen, J. Hubička, M. Konečný, L. Vena, and A. Zucker, Big Ramsey degrees of the generic partial order. In *Extended Abstracts EuroComb 2021*,
- [3] M. Balko, D. Chodounský, J. Hubička, M. Konečný, J. Nešetřil, and L. Vena, Big Ramsey degrees and forbidden cycles. *Extended abstract in EUROCOMB 2021* 2021, arXiv:2105.12184.
- [4] M. Balko, D. Chodounský, J. Hubička, M. Konečný, and L. Vena, Big Ramsey degrees of 3-uniform hypergraphs. *Acta Math. Univ. Com.* **LXXXVIII** (2019), no. 3, 415–422.
- [5] M. Balko, D. Chodounský, J. Hubička, M. Konečný, and L. Vena, Big Ramsey degrees of 3-uniform hypergraphs are finite. 2020, arXiv:2008.00268.
- [6] R. Coulson, N. Dobrinen, and R. Patel, Fraïssé classes with simply characterized big Ramsey structures. submitted (2020) arXiv:2010.02034
- [7] D. Devlin, *Some partition theorems for ultrafilters on ω* . Ph.D. thesis, Dartmouth College, 1979.
- [8] N. Dobrinen, The Ramsey theory of the universal homogeneous triangle-free graph. *Journal of Mathematical Logic*, **20** (2020) no. 2, 2050012, 75pp.
- [9] N. Dobrinen, The Ramsey theory of Henson graphs. submitted (2019), <https://arxiv.org/pdf/1901.06660.pdf>.
- [10] N. Dobrinen, Ramsey theory of homogeneous structures: Current trends and open problems. *Proceedings of the 2022 ICM*, to appear (2022), <https://arxiv.org/pdf/2110.00655.pdf>.
- [11] P. Erdős, A. Hajnal, and L. Pósa, Strong embeddings of graphs into colored graphs. In *Infinite and finite sets (Colloq., Keszthely, 1973; dedicated to P. Erdős on his 60th birthday*, vol. I, pp. 585–595, Colloq. Math. Soc. János Bolyai, 10, North-Holland, Amsterdam, 1975.
- [12] M. El-Zahar and N. Sauer, The indivisibility of the homogeneous K_n -free graphs. *Jour. Combin. Th., Ser. B* **47** (1989), no. 2, 162–170.
- [13] M. El-Zahar and N. Sauer, On the divisibility of homogeneous directed graphs. *Canadian Jour. Math.* **45** (1993), no. 2, 284–294.
- [14] Hubička, J.: Big Ramsey degrees using parameter spaces. *preprint*, arXiv:2009.00967 (2020)
- [15] J. Hubička and J. Nešetřil, All those Ramsey classes (Ramsey classes with closures and forbidden homomorphisms). *Adv. Math.* **356** (2019), 106791, 89 pp.
- [16] A. Kechris, V. Pestov, and S. Todorcević, Fraïssé limits, Ramsey theory, and topological dynamics of automorphism groups, *Geometric and Functional Analysis*, **15** (2005), 106–189.

- [17] P. Komjáth and V. Rödl, Coloring of universal graphs. *Graphs and Combin.* **2** (1986), no. 1, 55–60.
- [18] C. Laflamme, N. Sauer, and V. Vuksanovic, Canonical partitions of universal structures, *Combinatorica*, **26(2)** (2006), 183–205.
- [19] J. Larson, Counting canonical partitions in the random graph. *Combinatorica* **28** (2008), no. 6, 659–678.
- [20] K.R. Milliken, A Ramsey Theorem for Trees. *Jour. Combin. Th., Ser. A* **26** (1976), 215–237.
- [21] Jaroslav Nešetřil, Ramsey classes and homogeneous structures. *Combinatorics, probability and computing*, 14(1-2):171–189, 2005.
- [22] J. Nešetřil and V. Rödl, Partitions of finite relational and set systems. *Jour. Combin. Th. Ser. A* **22** (1977), no. 3, 289–312.
- [23] J. Nešetřil and V. Rödl, Ramsey classes of set systems. *Jour. Combin. Th. Ser. A* **34** (1983), no. 2, 183–201.
- [24] L. Nguyen Van Thé, More on the Kechris-Pestov-Todorčević Correspondence: Precompact Expansions, *Fund. Math.*, **222** (2013), 19–47.
- [25] M. Pouzet and N. Sauer, Edge partitions of the Rado graph. *Combinatorica* **16** (1996), no. 4, 505–520.
- [26] F.P. Ramsey, On a problem of formal logic. *Proc. London Math. Soc.* **30** (1929), no. 4, 264–296.
- [27] N. Sauer, Canonical vertex partitions, *Combinatorics, Probability, and Computing*, **12(6)** (2003), 671–704.
- [28] N. Sauer, Coloring subgraphs of the Rado graph. *Combinatorica* **26** (2006), no. 2, 231–253.
- [29] W. Sierpiński, Sur une problème de la théorie des relations. *Ann. Scuola Norm. Super. Pisa, Ser. 2* **2** (1933), 239–242.
- [30] A. Zucker, Topological dynamics of automorphism groups, ultrafilter combinatorics, and the generic point problem, *Trans. Amer. Math. Soc.*, **368(9)** (2016), 6715–6740.
- [31] A. Zucker, Big Ramsey degrees and topological dynamics, *Groups, Geom., Dyn.*, **13(1)** (2019), 235–276.
- [32] A. Zucker, A note on big Ramsey degrees, preprint, <https://arxiv.org/pdf/2004.13162.pdf>.