

UNIQUENESS OF THE SOLUTION OF SOME NONLINEAR SINGULAR PARTIAL DIFFERENTIAL EQUATIONS OF THE SECOND ORDER

HIDETOSHI TAHARA

ABSTRACT. In this paper, we consider a nonlinear singular second order partial differential equation of the form

$$\left(t \frac{\partial}{\partial t}\right)^2 u = F\left(t, x, \left\{\left(t \frac{\partial}{\partial t}\right)^i \left(\frac{\partial}{\partial x}\right)^\alpha u\right\}_{i+|\alpha| \leq 2, i < 2}\right)$$

in the complex domain. If $F(t, x, z)$ (with $z = \{z_{i,\alpha}\}_{i+|\alpha| \leq 2, i < 2}$) is a holomorphic function satisfying $F(0, x, 0) \equiv 0$ and $(\partial F / \partial z_{i,\alpha})(0, x, 0) \equiv 0$ (if $|\alpha| > 0$), then this equation is called a nonlinear Fuchsian type partial differential equation in t . Under a very weak assumption, we show the uniqueness of the solution. The result is applied to the problem of analytic continuation of local holomorphic solutions of this equation.

1. INTRODUCTION

To study the uniqueness of the solution is one of the most fundamental problems in the theory of partial differential equations. In this paper, we consider the case of nonlinear singular partial differential equations (1.1) given below.

Let $m \in \mathbb{N}^*$ ($= \{1, 2, \dots\}$). Let $(t, x) = (t, x_1, \dots, x_n) \in \mathbb{C}_t \times \mathbb{C}_x^n$ be a complex variable and let $z = \{z_{i,\alpha}\}_{i+|\alpha| \leq m, i < m} \in \mathbb{C}^N$ be also a complex variable, where we used the notations: $i \in \mathbb{N}$ ($= \{0, 1, 2, \dots\}$), $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$, $|\alpha| = \alpha_1 + \dots + \alpha_n$, and $N = \#\{(i, \alpha) \in \mathbb{N} \times \mathbb{N}^n; i + |\alpha| \leq m, i < m\}$. We write $(\partial / \partial x)^\alpha = (\partial / \partial x_1)^{\alpha_1} \dots (\partial / \partial x_n)^{\alpha_n}$.

Let $F(t, x, z)$ be a function in a neighborhood Δ of the origin of $\mathbb{C}_t \times \mathbb{C}_x^n \times \mathbb{C}_z^N$. Set $\Delta_0 = \Delta \cap \{t = 0, z = 0\}$. In this paper, we consider a nonlinear singular partial differential equation

$$(1.1) \quad \left(t \frac{\partial}{\partial t}\right)^m u = F\left(t, x, \left\{\left(t \frac{\partial}{\partial t}\right)^i \left(\frac{\partial}{\partial x}\right)^\alpha u\right\}_{i+|\alpha| \leq m, i < m}\right)$$

(in the germ sense at $(0, 0) \in \mathbb{C}_t \times \mathbb{C}_x^n$) under the following assumptions:

- A₁) $F(t, x, z)$ is a holomorphic function in Δ .
- A₂) $F(0, x, 0) \equiv 0$ in Δ_0 .
- A₃) $(\partial F / \partial z_{i,\alpha})(0, x, 0) \equiv 0$ in Δ_0 , if $|\alpha| > 0$.

Then, (1.1) is called a nonlinear Fuchsian type partial differential equation with respect to t , and the roots $\lambda_1(x), \dots, \lambda_m(x)$ of

$$\lambda^m - \sum_{i < m} (\partial F / \partial z_{i,0})(0, x, 0) \lambda^i = 0$$

2020 *Mathematics Subject Classification*. Primary 35A02; Secondary 35G20, 35B60.

Key words and phrases. Uniqueness of the solution, nonlinear partial differential equation, second order equation.

are called the characteristic exponents of (1.1).

Equations of this type were first studied by Gérard-Tahara [3, 4], and then by Tahara-Yamazawa [7]. The most fundamental result on (1.1) is:

Theorem 1 (Gérard-Tahara [3]). *Suppose that $\lambda_i(0) \notin \mathbb{N}^*$ holds for $i = 1, \dots, m$. Then, the equation (1.1) has a unique holomorphic solution $u_0(t, x)$ in a neighborhood of $(0, 0) \in \mathbb{C}_t \times \mathbb{C}_x^n$ satisfying $u_0(0, x) \equiv 0$.*

In this paper, we consider the uniqueness of the solution of (1.1) under the assumption

$$(1.2) \quad \operatorname{Re} \lambda_i(0) < 0, \quad i = 1, \dots, m.$$

We denote by $\mathcal{R}(\mathbb{C}_t \setminus \{0\})$ the universal covering space of $\mathbb{C}_t \setminus \{0\}$. For a nonempty open interval $I = (\theta_1, \theta_2)$ and $r > 0$ we write $S_I = \{t \in \mathcal{R}(\mathbb{C} \setminus \{0\}); \theta_1 < \arg t < \theta_2\}$ and $S_I(r) = \{t \in S_I; 0 < |t| < r\}$. For $R > 0$ we write $D_R = \{x \in \mathbb{C}^n; |x| < R\}$, where $|x| = \max_{1 \leq j \leq n} |x_j|$.

We know:

(1)(Gérard-Tahara [3]). If (1.2) is satisfied, and if $u(t, x)$ is a holomorphic solution of (1.1) on $S_I(r_0) \times D_{R_0}$ (for some nonempty open interval I , $r_0 > 0$ and $R_0 > 0$) satisfying

$$(1.3) \quad \sup_{x \in D_{R_0}} |u(t, x)| = O(|t|^a) \quad (\text{as } S_I \ni t \rightarrow 0)$$

for some $a > 0$, we have $u(t, x) = u_0(t, x)$ on $S_I(r) \times D_R$ for some $r > 0$ and $R > 0$.

(2) In Tahara [5], this condition (1.3) was weakened to the following one: there is an $\epsilon > 0$ such that

$$\sup_{x \in D_{R_0}} |u(t, x)| = O\left(\frac{1}{|\log t|^\epsilon}\right) \quad (\text{as } S_I \ni t \rightarrow 0).$$

(3) After that, the author has been trying to prove the following conjecture as a working hypothesis.

Conjecture. Suppose (1.2). If $u(t, x)$ is a holomorphic solution of (1.1) on $S_I(r_0) \times D_{R_0}$ (for some nonempty open interval I , $r_0 > 0$ and $R_0 > 0$) satisfying

$$\sup_{x \in D_{R_0}} |u(t, x)| = o(1) \quad (\text{as } S_I \ni t \rightarrow 0),$$

we have $u(t, x) = u_0(t, x)$ on $S_I(r) \times D_R$ for some $r > 0$ and $R > 0$.

The purpose of this paper is to show that the above conjecture is true in the case $m = 2$. The case $m = 1$ has already been proved in Tahara [6] by a method similar to the Cauchy's characteristics method. In this paper, we will modify its argument so that it works also in the case $m = 2$. Since the argument here works only in the case $m = 2$, the above conjecture is still open in the case $m \geq 3$.

2. MAIN RESULT

From now, we consider the case $m = 2$. Our equation is

$$(2.1) \quad \left(t \frac{\partial}{\partial t}\right)^2 u = F\left(t, x, \left\{\left(t \frac{\partial}{\partial t}\right)^i \left(\frac{\partial}{\partial x}\right)^\alpha u\right\}_{i+|\alpha| \leq 2, i < 2}\right).$$

Let $\lambda_1(x), \lambda_2(x)$ be the characteristic exponents of (2.1). We suppose the conditions $A_1), A_2), A_3)$ and

$$(2.2) \quad \operatorname{Re}\lambda_i(0) < 0, \quad i = 1, 2.$$

The following result is the main theorem of this paper.

Theorem 2. *Let $u(t, x)$ be a holomorphic solution of (2.1) on $S_I(r_0) \times D_{R_0}$ for some nonempty open interval I , $r_0 > 0$ and $R_0 > 0$. If $u(t, x)$ satisfies*

$$(2.3) \quad \overline{\lim}_{R \rightarrow +0} \left[\lim_{r \rightarrow +0} \left(\frac{1}{R^4} \sup_{S_I(r) \times D_R} |u(t, x)| \right) \right] = 0,$$

we have $u(t, x) = u_0(t, x)$ on $S_I(\sigma) \times D_\delta$ for some $\sigma > 0$ and $\delta > 0$, where $u_0(t, x)$ is the unique holomorphic solution in Theorem 1.

If

$$(2.4) \quad \sup_{x \in D_R} |u(t, x)| = o(1), \quad (\text{as } S_I \ni t \rightarrow 0)$$

holds for some $R > 0$ we have (2.3), and so we have

Corollary 1. *If $u(t, x)$ satisfies (2.4), we have $u(t, x) = u_0(t, x)$ on $S_I(\sigma) \times D_\delta$ for some $\sigma > 0$ and $\delta > 0$.*

This proves the conjecture posed in §1 in the case $m = 2$.

Remark. (1) If $\operatorname{Re}\lambda_1(0) > 0$ or $\operatorname{Re}\lambda_2(0) > 0$ holds, we have many solutions satisfying (2.4) and so the uniqueness of the solution is not valid. See [3, 7].

(2) In the case $\operatorname{Re}\lambda_1(0) = 0$ or $\operatorname{Re}\lambda_2(0) = 0$, we have the following counter example: the equation

$$\left(t \frac{\partial}{\partial t}\right)^2 u = -\left(t \frac{\partial}{\partial t}\right)u + \left(\frac{\partial u}{\partial x}\right)^2 + 8u \left(\frac{\partial^2 u}{\partial x^2}\right)^2$$

has a trivial solution $u(t, x) \equiv 0$ and a nontrivial solution

$$u(t, x) = \frac{-x^2}{4 \log t}$$

which satisfies (2.4). In this case, the characteristic exponents are 0 and -1 .

(3) The following example shows that the condition (2.3) will be reasonable: the equation

$$\left(t \frac{\partial}{\partial t}\right)^2 u = -3\left(t \frac{\partial}{\partial t}\right)u - 2u + \left(\frac{\partial^2 u}{\partial x^2}\right)^2$$

has a trivial solution $u(t, x) \equiv 0$ and a nontrivial solution $u = x^4/72$. We note that for $u = x^4/72$ we have

$$\overline{\lim}_{R \rightarrow +0} \left[\lim_{r \rightarrow +0} \left(\frac{1}{R^4} \sup_{S_I(r) \times D_R} |u(t, x)| \right) \right] = \frac{1}{72}.$$

In this case, the characteristic exponents are -1 and -2 .

3. SOME PREPARATORY DISCUSSION

Before the proof of Theorem 2, let us present some preparatory discussion. Let $u(t, x)$ be a holomorphic solution of (2.1) on $S_I(\sigma_0^*) \times D_{R_0^*}$ (for some nonempty open interval I , $\sigma_0^* > 0$ and $R_0^* > 0$) satisfying the condition (2.3). For simplicity, we set

$$\begin{aligned}\Lambda &= \{(i, \alpha) \in \mathbb{N} \times \mathbb{N}^n; i + |\alpha| \leq 2, i < 2\}, \\ N &= \#\Lambda \text{ (the cardinal of } \Lambda), \\ z &= \{z_{i,\alpha}\}_{(i,\alpha) \in \Lambda} \quad \text{and} \quad z' = \{z_{0,\alpha}\}_{|\alpha|=2}.\end{aligned}$$

For $\nu = \{\nu_{i,\alpha}\}_{(i,\alpha) \in \Lambda} \in \mathbb{N}^N$ we write

$$|\nu| = \sum_{(i,\alpha) \in \Lambda} \nu_{i,\alpha}, \quad z^\nu = \prod_{(i,\alpha) \in \Lambda} (z_{i,\alpha})^{\nu_{i,\alpha}}.$$

Suppose the conditions A₁), A₂), A₃): then $F(t, x, z)$ is a holomorphic function in a neighborhood of the origin of $\mathbb{C}_t \times \mathbb{C}_x^n \times \mathbb{C}_z^N$ having the expansion of the form

$$F(t, x, z) = a(x)t + \beta_0^*(x)z_{0,0} + \beta_1^*(x)z_{1,0} + R(t, x, z)$$

with

$$R(t, x, z) = \sum_{i+|\nu| \geq 2} \gamma_{i,\nu}^*(x) t^i z^\nu.$$

Step 1. We set $w(t, x) = u(t, x) - u_0(t, x)$. Then, we have

$$(3.1) \quad \overline{\lim}_{R \rightarrow +0} \left[\lim_{r \rightarrow +0} \left(\frac{1}{R^4} \sup_{S_I(r) \times D_R} |w(t, x)| \right) \right] = 0,$$

and $w(t, x)$ is a holomorphic solution of the equation

$$(3.2) \quad \left(t \frac{\partial}{\partial t} \right)^2 w = H \left(t, x, \left\{ \left(t \frac{\partial}{\partial t} \right)^i \left(\frac{\partial}{\partial x} \right)^\alpha w \right\}_{(i,\alpha) \in \Lambda} \right)$$

on $S_I(\sigma_0^*) \times D_{R_0^*}$, where

$$\begin{aligned}H(t, x, z) &= F(t, x, z + \{u_{0,i,\alpha}(t, x)\}_{(i,\alpha) \in \Lambda}) \\ &\quad - F(t, x, \{u_{0,i,\alpha}(t, x)\}_{(i,\alpha) \in \Lambda})\end{aligned}$$

under the notation $u_{0,i,\alpha}(t, x) = (t\partial/\partial t)^i (\partial/\partial x)^\alpha u_0(t, x)$ ($(i, \alpha) \in \Lambda$). It is easy to see that $H(t, x, z)$ can be expressed in the form

$$\begin{aligned}H(t, x, z) &= \beta_0^*(x)z_{0,0} + \beta_1^*(x)z_{1,0} + t \sum_{(i,\alpha) \in \Lambda} a_{i,\alpha}^*(t, x, z) z_{i,\alpha} \\ &\quad + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} b_{i,\alpha}^*(x, z) z_{i,\alpha} + \sum_{|\alpha|=|\beta|=2} c_{\alpha,\beta}^*(x, z') z_{0,\alpha} z_{0,\beta}\end{aligned}$$

for some holomorphic functions $\beta_i^*(x)$ ($i = 0, 1$), $a_{i,\alpha}^*(t, x, z)$ ($(i, \alpha) \in \Lambda$), $b_{i,\alpha}^*(x, z)$ ($(i, \alpha) \in \Lambda, |\alpha| \leq 1$), and $c_{\alpha,\beta}^*(x, z')$ ($|\alpha| = |\beta| = 2$) in a common neighborhood of $(0, 0, 0) \in \mathbb{C}_t \times \mathbb{C}_x^n \times \mathbb{C}_z^N$ satisfying $b_{i,\alpha}^*(x, 0) \equiv 0$ ($(i, \alpha) \in \Lambda, |\alpha| \leq 1$). In addition, we see that the roots of $\lambda^2 - \beta_1^*(x)\lambda - \beta_0^*(x) = 0$ are just the same as $\lambda_1(x), \lambda_2(x)$.

From now, we consider the equation (3.2). Our purpose is to show that $w(t, x) \equiv 0$ holds on $S_I(\sigma) \times D_\delta$ for some $\sigma > 0$ and $\delta > 0$.

Step 2. We set $\Theta_0 = 1$ and

$$\begin{aligned}\Theta_1 &= \left(t \frac{\partial}{\partial t} - \lambda_1(0)\right), \\ \Theta_2 &= \left(t \frac{\partial}{\partial t} - \lambda_2(0)\right) \left(t \frac{\partial}{\partial t} - \lambda_1(0)\right).\end{aligned}$$

We write $Dw = \{D_{i,\alpha}w\}_{(i,\alpha) \in \Lambda}$ and $D'w = \{D_{0,\alpha}w\}_{|\alpha|=2}$ with

$$D_{i,\alpha}w = \Theta_i \left(\frac{\partial}{\partial x}\right)^\alpha w, \quad (i, \alpha) \in \Lambda.$$

Then, (3.2) is expressed in the form

$$\begin{aligned}(3.3) \quad \Theta_2 w &= \beta_0(x)w + \beta_1(x)\Theta_1 w + t \sum_{(i,\alpha) \in \Lambda} a_{i,\alpha}(t, x, Dw) \Theta_i \left(\frac{\partial}{\partial x}\right)^\alpha w \\ &+ \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} b_{i,\alpha}(x, Dw) \Theta_i \left(\frac{\partial}{\partial x}\right)^\alpha w \\ &+ \sum_{|\alpha|=|\beta|=2} c_{\alpha,\beta}(x, D'w) \left[\left(\frac{\partial}{\partial x}\right)^\alpha w\right] \times \left[\left(\frac{\partial}{\partial x}\right)^\beta w\right]\end{aligned}$$

for some holomorphic functions $\beta_i(x)$ ($i = 0, 1$), $a_{i,\alpha}(t, x, z)$ ($(i, \alpha) \in \Lambda$), $b_{i,\alpha}(x, z)$ ($(i, \alpha) \in \Lambda, |\alpha| \leq 1$) and $c_{\alpha,\beta}(x, z')$ ($|\alpha| = |\beta| = 2$) in a common neighborhood of $(0, 0, 0) \in \mathbb{C}_t \times \mathbb{C}_x^n \times \mathbb{C}_z^N$ satisfying $\beta_i(0) = 0$ ($i = 0, 1$) and $b_{i,\alpha}(x, 0) \equiv 0$ ($(i, \alpha) \in \Lambda, |\alpha| \leq 1$).

Step 3. By a rotation, we may suppose that $0 \in I$ holds: then we have $(0, \sigma_0^*) \subset S_I(\sigma_0^*)$. From now, we consider the equation (3.3) only on $(0, \sigma_0^*) \times D_{R_0^*}$. By (3.1) and Nagumo type lemma in a sector (for example, see Lemma 4.2 in [1]) we have the condition

$$(3.4) \quad \overline{\lim}_{R \rightarrow +0} \left[\lim_{\sigma \rightarrow +0} \left(\frac{1}{R^4} \sup_{(0,\sigma] \times D_R} \left| \left(t \frac{\partial}{\partial t}\right)^i w(t, x) \right| \right) \right] = 0, \quad i = 0, 1, 2.$$

For a formal power series $f(t, x) = \sum_{|\alpha| \geq 0} f_\alpha(t) x^\alpha$ with coefficients in $C^0((0, T))$ we write

$$\|f(t)\|_\rho = \sum_{|\alpha| \geq 0} |f_\alpha(t)| \frac{\alpha!}{|\alpha|!} \rho^{|\alpha|}$$

(which is a formal power series in ρ with coefficients in $C^0((0, T))$). When $\|f(t)\|_\rho$ converges, we regard it as a function in (t, ρ) . We can easily see:

$$\begin{aligned}\|f(t)g(t)\|_\rho &\ll \|f(t)\|_\rho \|g(t)\|_\rho, \\ \|(\partial/\partial x_j)f(t)\|_\rho &\ll (\partial/\partial \rho)\|f(t)\|_\rho, \quad j = 1, \dots, n.\end{aligned}$$

where $\sum_{i \geq 0} a_i \rho^i \ll \sum_{i \geq 0} b_i \rho^i$ means that $|a_i| \leq b_i$ holds for all $i \geq 0$. For a holomorphic function $f(t, x, z) = \sum_{i+|\nu| \geq 0} f_{i,\nu}(x) t^i z^\nu$ we write

$$\|f\|_\rho(t, z) = \sum_{i+|\nu| \geq 0} \|f_{i,\nu}\|_\rho t^i z^\nu.$$

By (3.4) we have

Lemma 1. *Under the above situation, for $i = 0, 1, 2$ we have*

$$(3.5) \quad \lim_{\sigma \rightarrow +0} \left(\sup_{(0, \sigma] \times [0, R]} \left\| \left(t \frac{\partial}{\partial t} \right)^i w(t) \right\|_{\rho} \right) = o(R^4) \quad (\text{as } R \rightarrow +0).$$

Step 4. Set $J = \{(i, j) \in \mathbb{N}^2; i + j \leq 2, i < 2\}$: actually, we have

$$J = \{(0, 0), (1, 0), (0, 1), (1, 1), (0, 2)\}.$$

For $(i, j) \in J$ we set

$$\begin{aligned} \phi_{0,0}(t, \rho) &= \int_0^t \left(\frac{\tau}{t} \right)^{-\operatorname{Re}\lambda_1(0)} \|\Theta_1 w(\tau)\|_{\rho} \frac{d\tau}{\tau}, \\ \phi_{1,0}(t, \rho) &= \int_0^t \left(\frac{\tau}{t} \right)^{-\operatorname{Re}\lambda_2(0)} \|\Theta_2 w(\tau)\|_{\rho} \frac{d\tau}{\tau}, \\ \phi_{0,1}(t, \rho) &= \frac{\partial}{\partial \rho} \phi_{0,0}(t, \rho), \\ \phi_{1,1}(t, \rho) &= \frac{\partial}{\partial \rho} \phi_{1,0}(t, \rho), \\ \phi_{0,2}(t, \rho) &= \frac{\partial}{\partial \rho} \phi_{0,1}(t, \rho) = \left(\frac{\partial}{\partial \rho} \right)^2 \phi_{0,0}(t, \rho). \end{aligned}$$

By making $\sigma_0^* > 0$ and $R_0^* > 0$ smaller if necessary, we may suppose that $\phi_{i,j}(t, \rho)$ ($(i, j) \in J$) are convergent on $(0, \sigma_0^*] \times [0, R_0^*]$. By (3.5) we have

Lemma 2. (1) *For any $(i, j) \in J$ we have*

$$\lim_{\sigma \rightarrow +0} \left(\sup_{(0, \sigma] \times [0, R]} \phi_{i,j}(t, \rho) \right) = o(R^{4-j}) \quad (\text{as } R \rightarrow +0).$$

(2) *For any $\epsilon > 0$ we can find $\sigma > 0$ and $R > 0$ such that $|\phi_{i,j}(t, \rho)| \leq \epsilon$ holds on $(0, \sigma] \times [0, R]$ for any $(i, j) \in J$.*

Step 5. By (2.2) we can take an $h > 0$ such that

$$\operatorname{Re}\lambda_i(0) < -2h, \quad i = 1, 2.$$

Lemma 3. (1) *For any $(i, \alpha) \in \Lambda$ we have*

$$\|D_{i,\alpha} w(t)\|_{\rho} = \left\| \Theta_i \left(\frac{\partial}{\partial x} \right)^{\alpha} w(t) \right\|_{\rho} \ll \phi_{i,|\alpha|}(t, \rho) \quad \text{on } (0, \sigma_0^*].$$

(2) *We set $\Phi = \Phi(t, \rho) = (\phi_{i,|\alpha|}(t, \rho); (i, \alpha) \in \Lambda)$ and $\Phi' = \Phi'(t, \rho) = (\phi_{0,|\alpha|}(t, \rho); |\alpha| = 2)$: we have*

$$\begin{aligned} & \left(t \frac{\partial}{\partial t} + 2h \right) \phi_{0,0}(t, \rho) \ll \phi_{1,0}(t, \rho), \\ & \left(t \frac{\partial}{\partial t} + 2h \right) \phi_{1,0}(t, \rho) \\ & \ll \|\beta_0\|_{\rho} \phi_{0,0}(t, \rho) + \|\beta_1\|_{\rho} \phi_{1,0}(t, \rho) + t \sum_{(i,\alpha) \in \Lambda} \|a_{i,\alpha}\|_{\rho}(t, \Phi) \phi_{i,|\alpha|}(t, \rho) \\ & + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \|b_{i,\alpha}\|_{\rho}(\Phi) \phi_{i,|\alpha|}(t, \rho) + \sum_{|\alpha|=|\beta|=2} \|c_{\alpha,\beta}\|_{\rho}(\Phi') (\phi_{0,2}(t, \rho))^2 \end{aligned}$$

on $(0, \sigma_0^*]$.

(3) By taking $R_0^* > 0$ a smaller one if necessary, we may assume that $\|\beta_i\|_\rho$ ($i = 0, 1$), $\|a_{i,\alpha}\|_\rho(t, z)$ ($(i, \alpha) \in \Lambda$), $\|b_{i,\alpha}\|_\rho(z)$ ($(i, \alpha) \in \Lambda, |\alpha| \leq 1$) and $\|c_{\alpha,\beta}\|_\rho(z')$ are convergent on $(0, \sigma_0^*] \times [0, R_0^*] \times D_L$ (where $D_L = \{z \in \mathbb{C}^N; |z_{i,\alpha}| < L \text{ } ((i, \alpha) \in \Lambda)\}$) for some $L > 0$. In addition, we have

$$\begin{aligned} \|\beta_i\|_\rho &\leq H_i \rho \quad \text{on } [0, R_0^*], \quad i = 0, 1, \\ \|b_{i,\alpha}\|_\rho(z) &\leq B_{i,\alpha} |z| \quad \text{on } [0, R_0^*] \times D_L, \quad (i, \alpha) \in \Lambda, |\alpha| \leq 1 \end{aligned}$$

for some constants $H_i > 0$ ($i = 0, 1$) and $B_{i,\alpha} > 0$ ($(i, \alpha) \in \Lambda, |\alpha| \leq 1$), where $|z| = \sum_{(i,\alpha) \in \Lambda} |z_{i,\alpha}|$.

(4) For any $(i, \alpha) \in \Lambda$ with $|\alpha| \leq 1$ we have

$$\lim_{\sigma \rightarrow +0} \left(\sup_{(0,\sigma] \times [0,R]} \frac{\partial \|b_{i,\alpha}\|_\rho(\Phi)}{\partial \rho} \right) = o(R) \quad (\text{as } R \rightarrow +0).$$

Proof. Let us show (1). Since

$$\left(t \frac{\partial}{\partial t} - \lambda_{i+1}(0) \right) \Theta_i w = (\Theta_{i+1} w)(t, x), \quad i = 0, 1$$

hold, by integrating this we have

$$\Theta_i w(t, x) = \int_0^t \left(\frac{\tau}{t} \right)^{-\lambda_{i+1}(0)} (\Theta_{i+1} w)(\tau, x) \frac{d\tau}{\tau}$$

and so by taking the norm

$$\|\Theta_i w(t)\|_\rho \ll \int_0^t \left(\frac{\tau}{t} \right)^{-\text{Re} \lambda_{i+1}(0)} \|\Theta_{i+1} w(\tau)\|_\rho \frac{d\tau}{\tau} = \phi_{i,0}(t, \rho), \quad i = 0, 1.$$

Hence, for any $(i, \alpha) \in \Lambda$ we have

$$\left\| \Theta_i \left(\frac{\partial}{\partial x} \right)^\alpha w(t) \right\|_\rho \ll \left(\frac{\partial}{\partial \rho} \right)^{|\alpha|} \|\Theta_i w(t)\|_\rho \ll \left(\frac{\partial}{\partial \rho} \right)^{|\alpha|} \phi_{i,0}(t, \rho) = \phi_{i,|\alpha|}(t, \rho).$$

Let us show (2). We have

$$\begin{aligned} \left(t \frac{\partial}{\partial t} + 2h \right) \phi_{0,0}(t, \rho) &\ll \left(t \frac{\partial}{\partial t} + \text{Re} \lambda_1(0) \right) \phi_{0,0}(t, \rho) \\ &= \|\Theta_1 w(\tau)\|_\rho \ll \phi_{1,0}(t, \rho) \quad \text{on } (0, \sigma_0^*]. \end{aligned}$$

This proves the first inequality of (2). Since

$$\left(t \frac{\partial}{\partial t} + 2h \right) \phi_{1,0}(t, \rho) \ll \left(t \frac{\partial}{\partial t} + \text{Re} \lambda_2(0) \right) \phi_{1,0}(t, \rho) = \|\Theta_2 w(\tau)\|_\rho,$$

by applying (3.3) and by using (1) we can easily see the second inequality of (2). The condition (3) is clear. Since $\|b_{j,\alpha}\|_\rho(\Phi) \leq B_{j,\alpha} |\Phi|$ ($(j, \alpha) \in \Lambda$) hold, by (1) of Lemma 2 we have

$$\lim_{\sigma \rightarrow +0} \left(\sup_{(0,\sigma] \times [0,R]} \|b_{i,\alpha}\|_\rho(\Phi) \right) = o(R^2) \quad (\text{as } R \rightarrow +0).$$

This leads us to the condition (4). □

Step 6. Let $\varepsilon_{0,0} > 0$, $\varepsilon_{1,0} = 1$, $\varepsilon_{0,1} > 0$, $\varepsilon_{1,1} > 0$ and $0 < \kappa < 1$. We set

$$\begin{aligned} q(t, \rho) &= \varepsilon_{0,0} \phi_{0,0}(t, \rho) + \phi_{1,0}(t, \rho) + t^\kappa \phi_{0,2}(t, \rho) \\ &\quad + \varepsilon_{0,1} \phi_{0,1}(t, \rho) + \varepsilon_{1,1} \phi_{1,1}(t, \rho) + (\phi_{0,2}(t, \rho))^{3/2}. \end{aligned}$$

Lemma 4. *We have the following inequality:*

$$(3.6) \quad \left(t \frac{\partial}{\partial t} + 2h\right)q \leq A(t, \rho)q + B(t, \rho) \frac{\partial}{\partial \rho} q$$

on $(0, \sigma_0^*] \times [0, R_0^*]$, where

$$\begin{aligned} A(t, \rho) &= \varepsilon_{0,0} + \left(\frac{1}{\varepsilon_{0,0}} \|\beta_0\|_\rho + \|\beta_1\|_\rho\right) \\ &+ \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \frac{1}{\varepsilon_{i,|\alpha|}} \times t \|a_{i,\alpha}\|_\rho(t, \Phi) + \sum_{|\alpha|=2} t^{1-\kappa} \|a_{0,\alpha}\|_\rho(t, \Phi) \\ &+ \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \frac{1}{\varepsilon_{i,|\alpha|}} \times \|b_{i,\alpha}\|_\rho(\Phi) + \sum_{|\alpha|=|\beta|=2} \|c_{\alpha,\beta}\|_\rho(\Phi') (\phi_{0,2})^{1/2} \\ &+ \kappa + \frac{\varepsilon_{0,1}}{\varepsilon_{1,1}} + \varepsilon_{1,1} \left(\frac{1}{\varepsilon_{0,0}} \frac{\partial \|\beta_0\|_\rho}{\partial \rho} + \frac{\partial \|\beta_1\|_\rho}{\partial \rho}\right) \\ &+ \varepsilon_{1,1} \left(\frac{\|\beta_0\|_\rho}{\varepsilon_{0,1}} + \frac{\|\beta_1\|_\rho}{\varepsilon_{1,1}}\right) + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \frac{\varepsilon_{1,1}}{\varepsilon_{i,|\alpha|}} \times t \frac{\partial \|a_{i,\alpha}\|_\rho(t, \Phi)}{\partial \rho} \\ &+ \varepsilon_{1,1} \sum_{|\alpha|=2} t^{1-\kappa} \frac{\partial \|a_{0,\alpha}\|_\rho(t, \Phi)}{\partial \rho} + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \frac{\varepsilon_{1,1}}{\varepsilon_{i,|\alpha|}} \times \frac{\partial \|b_{i,\alpha}\|_\rho(\Phi)}{\partial \rho} \\ &+ \varepsilon_{1,1} \sum_{|\alpha|=|\beta|=2} \frac{\partial \|c_{\alpha,\beta}\|_\rho(\Phi')}{\partial \rho} (\phi_{0,2})^{1/2}, \\ B(t, \rho) &= \frac{t^\kappa}{\varepsilon_{1,1}} + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \frac{\varepsilon_{1,1}}{\varepsilon_{i,|\alpha|}} \times t \|a_{i,\alpha}\|_\rho(t, \Phi) \\ &+ \varepsilon_{1,1} \sum_{|\alpha|=2} t^{1-\kappa} \|a_{0,\alpha}\|_\rho(t, \Phi) + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \frac{\varepsilon_{1,1}}{\varepsilon_{i,|\alpha|}} \times \|b_{i,\alpha}\|_\rho(\Phi) \\ &+ \frac{4\varepsilon_{1,1}}{3} \sum_{|\alpha|=|\beta|=2} \|c_{\alpha,\beta}\|_\rho(\Phi') (\phi_{0,2})^{1/2} + \frac{3}{2\varepsilon_{1,1}} (\phi_{0,2})^{1/2}. \end{aligned}$$

Proof. By the definition of $q(t, \rho)$ we have

$$(3.7) \quad \phi_{i,j} \leq \frac{1}{\varepsilon_{i,j}} q \quad (i, j = 0, 1), \quad \phi_{0,2} \leq q^{2/3}, \quad t^\kappa \phi_{0,2} \leq q$$

on $(0, \sigma_0^*] \times [0, R_0^*]$. Since

$$\begin{aligned} \frac{\partial}{\partial \rho} q &= \varepsilon_{0,0} \frac{\partial}{\partial \rho} \phi_{0,0} + \frac{\partial}{\partial \rho} \phi_{1,0} + t^\kappa \frac{\partial}{\partial \rho} \phi_{0,2} \\ &+ \varepsilon_{0,1} \frac{\partial}{\partial \rho} \phi_{0,1} + \varepsilon_{1,1} \frac{\partial}{\partial \rho} \phi_{1,1} + \frac{3}{2} (\phi_{0,2})^{1/2} \frac{\partial}{\partial \rho} \phi_{0,2} \end{aligned}$$

we have also

$$(3.8) \quad \begin{aligned} \frac{\partial}{\partial \rho} \phi_{i,j} &\leq \frac{1}{\varepsilon_{i,j}} \frac{\partial}{\partial \rho} q \quad (i, j = 0, 1), \\ (\phi_{0,2})^{1/2} \frac{\partial}{\partial \rho} \phi_{0,2} &\leq \frac{2}{3} \frac{\partial}{\partial \rho} q, \quad t^\kappa \frac{\partial}{\partial \rho} \phi_{0,2} \leq \frac{\partial}{\partial \rho} q \end{aligned}$$

on $(0, \sigma_0^*] \times [0, R_0^*]$. By using these inequality, let us do a calculation.

1) About $\varepsilon_{0,0}\phi_{0,0}(t, \rho)$, by (2) of Lemma 3 and (3.7) we have

$$\left(t \frac{\partial}{\partial t} + 2h\right)(\varepsilon_{0,0}\phi_{0,0}) \ll \varepsilon_{0,0}\phi_{1,0} \leq \varepsilon_{0,0}q.$$

2) About $\phi_{1,0}$, by (2) of Lemma 3 and (3.7) we have

$$\begin{aligned} & \left(t \frac{\partial}{\partial t} + 2h\right)\phi_{1,0} \\ & \ll \|\beta_0\|_\rho \phi_{0,0} + \|\beta_1\|_\rho \phi_{1,0} + t \sum_{(i,\alpha) \in \Lambda} \|a_{i,\alpha}\|_\rho(t, \Phi) \phi_{i,|\alpha|} \\ & \quad + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \|b_{i,\alpha}\|_\rho(\Phi) \phi_{i,|\alpha|} + \sum_{|\alpha|=|\beta|=2} \|c_{\alpha,\beta}\|_\rho(\Phi')(\phi_{0,2})^2 \\ & \leq \left(\frac{1}{\varepsilon_{0,0}} \|\beta_0\|_\rho + \|\beta_1\|_\rho\right)q \\ & \quad + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \frac{1}{\varepsilon_{i,|\alpha|}} t \|a_{i,\alpha}\|_\rho(t, \Phi) q + t^{1-\kappa} \sum_{|\alpha|=2} \|a_{0,\alpha}\|_\rho(t, \Phi) q \\ & \quad + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \frac{1}{\varepsilon_{i,|\alpha|}} \|b_{i,\alpha}\|_\rho(\Phi) q + \sum_{|\alpha|=|\beta|=2} \|c_{\alpha,\beta}\|_\rho(\Phi')(\phi_{0,2})^{1/2} q. \end{aligned}$$

3) Let us consider $t^\kappa \phi_{0,2}(t, \rho)$. Since $(\partial/\partial\rho)\phi_{1,1} \ll (1/\varepsilon_{1,1})(\partial/\partial\rho)q$ holds, we have

$$\left(t \frac{\partial}{\partial t} + 2h\right)\phi_{0,2} = \left(\frac{\partial}{\partial\rho}\right)^2 \left(t \frac{\partial}{\partial t} + 2h\right)\phi_{0,0} \ll \left(\frac{\partial}{\partial\rho}\right)^2 \phi_{1,0} = \frac{\partial}{\partial\rho} \phi_{1,1} \leq \frac{1}{\varepsilon_{1,1}} \frac{\partial}{\partial\rho} q$$

and so

$$\left(t \frac{\partial}{\partial t} + 2h\right)t^\kappa \phi_{0,2} = t^\kappa \left(t \frac{\partial}{\partial t} + 2h\right)\phi_{0,2} + \kappa t^\kappa \phi_{0,2} \leq \frac{t^\kappa}{\varepsilon_{1,1}} \frac{\partial}{\partial\rho} q + \kappa q.$$

4) About $\varepsilon_{0,1}\phi_{0,1}(t, \rho)$, we have

$$\begin{aligned} \left(t \frac{\partial}{\partial t} + 2h\right)(\varepsilon_{0,1}\phi_{0,1}) & = \varepsilon_{0,1} \frac{\partial}{\partial\rho} \left(t \frac{\partial}{\partial t} + 2h\right)\phi_{0,0} \\ & \ll \varepsilon_{0,1} \frac{\partial}{\partial\rho} \phi_{1,0} = \varepsilon_{0,1}\phi_{1,1} \leq \frac{\varepsilon_{0,1}}{\varepsilon_{1,1}} q. \end{aligned}$$

5) About $\varepsilon_{1,1}\phi_{1,1}(t, \rho)$, by (2) of Lemma 3 we have

$$\begin{aligned} & \left(t \frac{\partial}{\partial t} + 2h\right)(\varepsilon_{1,1}\phi_{1,1}) = \varepsilon_{1,1} \frac{\partial}{\partial\rho} \left(t \frac{\partial}{\partial t} + 2h\right)\phi_{1,0} \\ & \ll \varepsilon_{1,1} \frac{\partial}{\partial\rho} \left[\|\beta_0\|_\rho \phi_{0,0} + \|\beta_1\|_\rho \phi_{1,0} + t \sum_{(i,\alpha) \in \Lambda} \|a_{i,\alpha}\|_\rho(t, \Phi) \phi_{i,|\alpha|} \right. \\ & \quad \left. + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \|b_{i,\alpha}\|_\rho(\Phi) \phi_{i,|\alpha|} + \sum_{|\alpha|=|\beta|=2} \|c_{\alpha,\beta}\|_\rho(\Phi')(\phi_{0,2})^2 \right]. \end{aligned}$$

By calculating the right side of the above formula and then by using (3.7) and (3.8) we have

$$\begin{aligned}
& \left(t \frac{\partial}{\partial t} + 2h\right)(\varepsilon_{1,1} \phi_{1,1}) \\
& \leq \varepsilon_{1,1} \left(\frac{1}{\varepsilon_{0,0}} \frac{\partial \|\beta_0\|_\rho}{\partial \rho} + \frac{\partial \|\beta_1\|_\rho}{\partial \rho} \right) q + \varepsilon_{1,1} \left(\frac{\|\beta_0\|_\rho}{\varepsilon_{0,1}} + \frac{\|\beta_1\|_\rho}{\varepsilon_{1,1}} \right) q \\
& + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \frac{\varepsilon_{1,1}}{\varepsilon_{i,|\alpha|}} \times t \frac{\partial \|a_{i,\alpha}\|_\rho(t, \Phi)}{\partial \rho} \times q \\
& + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \frac{\varepsilon_{1,1}}{\varepsilon_{i,|\alpha|}} \times t \|a_{i,\alpha}\|_\rho(t, \Phi) \frac{\partial}{\partial \rho} q \\
& + \varepsilon_{1,1} t^{1-\kappa} \sum_{|\alpha|=2} \frac{\partial \|a_{0,\alpha}\|_\rho(t, \Phi)}{\partial \rho} q + \varepsilon_{1,1} t^{1-\kappa} \sum_{|\alpha|=2} \|a_{0,\alpha}\|_\rho(t, \Phi) \frac{\partial}{\partial \rho} q \\
& + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \frac{\varepsilon_{1,1}}{\varepsilon_{i,|\alpha|}} \times \frac{\partial \|b_{i,\alpha}\|_\rho(\Phi)}{\partial \rho} q + \sum_{(i,\alpha) \in \Lambda, |\alpha| \leq 1} \frac{\varepsilon_{1,1}}{\varepsilon_{i,|\alpha|}} \times \|b_{i,\alpha}\|_\rho(\Phi) \frac{\partial}{\partial \rho} q \\
& + \varepsilon_{1,1} \sum_{|\alpha|=|\beta|=2} \frac{\partial \|c_{\alpha,\beta}\|_\rho(\Phi')}{\partial \rho} (\phi_{0,2})^{1/2} q \\
& + \frac{4\varepsilon_{1,1}}{3} \sum_{|\alpha|=|\beta|=2} \|c_{\alpha,\beta}\|_\rho(\Phi') (\phi_{0,2})^{1/2} \frac{\partial}{\partial \rho} q
\end{aligned}$$

6) Let us consider $(\phi_{0,2})^{3/2}$. Since

$$\left(t \frac{\partial}{\partial t} + 2h\right) \phi_{0,2} = \frac{\partial}{\partial \rho} \left(t \frac{\partial}{\partial t} + 2h\right) \phi_{0,1} \ll \frac{\partial}{\partial \rho} \phi_{1,1} \leq \frac{1}{\varepsilon_{1,1}} \frac{\partial}{\partial \rho} q$$

we have

$$\begin{aligned}
& \left(t \frac{\partial}{\partial t} + 2h\right) (\phi_{0,2})^{3/2} = \frac{3}{2} (\phi_{0,2})^{1/2} t \frac{\partial}{\partial t} \phi_{0,2} + 2h (\phi_{0,2})^{3/2} \\
& \leq \frac{3}{2} (\phi_{0,1})^{1/2} \left(t \frac{\partial}{\partial t} + 2h\right) \phi_{0,2} \leq \frac{3}{2\varepsilon_{1,1}} (\phi_{0,2})^{1/2} \frac{\partial}{\partial \rho} q.
\end{aligned}$$

7) Thus, by taking the summation from 1) to 6) we have the result (3.6). \square

Step 7. Let us estimate $A(t, \rho)$ and $B(t, \rho)$. We have

Lemma 5. *By taking $\varepsilon_{0,0} > 0$, $\varepsilon_{0,1} > 0$, $\varepsilon_{1,1} > 0$, $0 < \kappa < 1/2$, $\sigma_0 > 0$ and $R_0 > 0$ suitably, we have*

$$(3.9) \quad A(t, \rho) \leq h,$$

$$(3.10) \quad B(t, \rho) \leq C_1 t^\kappa + C_2 q + C_3 q^{2/3} + C_4 q^{1/3}$$

on $(0, \sigma_0] \times [0, R_0]$ for some $C_i > 0$ ($i = 1, 2, 3, 4$).

Proof. 1) First, we take $\varepsilon_{0,0}$ so that $0 < \varepsilon_{0,0} \leq h/4$.

2) Since $\varepsilon_{0,0}$ is a fixed constant, by taking $\varepsilon_{1,1} > 0$ sufficiently small we have the condition

$$\varepsilon_{1,1} \left(\frac{1}{\varepsilon_{0,0}} \frac{\partial \|\beta_0\|_\rho}{\partial \rho} + \frac{\partial \|\beta_1\|_\rho}{\partial \rho} \right) \leq \frac{h}{4} \quad \text{on } (0, \sigma_0^*] \times [0, R_0^*].$$

3) Thirdly, we take $\kappa > 0$ and $\varepsilon_{0,1} > 0$ so that $\kappa + \varepsilon_{0,1}/\varepsilon_{1,1} \leq h/4$.

4) Now, $\varepsilon_{i,|\alpha|} > 0$ ($(i, \alpha) \in \Lambda, |\alpha| \leq 1$) are fixed. Since $\|\beta_i\|_\rho \leq H_i \rho$ ($i = 0, 1$) and $\|b_{i,\alpha}\|_\rho(\Phi) \leq B_{i,\alpha}|\Phi|$ ($(i, \alpha) \in \Lambda, |\alpha| \leq 1$) are known (by (3) of Lemma 3), by the conditions (2) of Lemma 2 and (4) of Lemma 3 we can take $\sigma_0 > 0$ and $R_0 > 0$ suitably sufficiently small so that the condition (3.9) is valid on $(0, \sigma_0] \times [0, R_0]$.

5) Since $0 < \kappa < 1/2$ is supposed, we have $t = O(t^\kappa)$ and $t^{1-\kappa} = O(t^\kappa)$. Hence, by the condition $\|b_{i,\alpha}\|_\rho(\Phi) \leq B_{i,\alpha}|\Phi|$ we have

$$B(t, \rho) \leq K_1 t^\kappa + K_2 |\Phi| + K_3 (\phi_{0,2})^{1/2}$$

for some $K_i > 0$ ($i = 1, 2, 3$). By applying (3.7) to this estimate we have the condition (3.10). \square

4. PROOF OF THEOREM 2

In order to prove Theorem 2, it is enough to show

Proposition 1. *Under the situation in §3, there are $\sigma > 0$ and $\delta > 0$ such that $q(t, \rho) = 0$ holds on $(0, \sigma) \times [0, \delta]$.*

Proof. Let us show this, step by step.

Step 1. By Lemmas 4 and 5 we have

$$(4.1) \quad \left(t \frac{\partial}{\partial t} + h\right)q \leq B(t, \rho) \frac{\partial}{\partial \rho} q,$$

$$(4.2) \quad B(t, \rho) \leq C_1 t^\kappa + C_2 q + C_3 q^{2/3} + C_4 q^{1/3}$$

on $(0, \sigma_0] \times [0, R_0]$. For $\sigma > 0$ and $R > 0$ we set

$$r = \sup_{(0, \sigma] \times [0, R]} q(t, \rho).$$

By (1) of Lemma 2 and the definition of $q(t, \rho)$ we have

$$\lim_{\sigma \rightarrow +0} r = o(R^3) \quad (\text{as } R \rightarrow +0).$$

This shows

Lemma 6. *By taking $\sigma > 0$ and $R > 0$ sufficiently small, we have the condition*

$$\frac{C_1}{\kappa} \sigma^\kappa + \frac{C_2}{h} r + \frac{3C_3}{2h} r^{2/3} + \frac{3C_4}{h} r^{1/3} < \frac{R}{2} \quad \text{on } (0, \sigma] \times [0, R].$$

Step 2. Let $\sigma > 0$ and $R > 0$ be as in Lemma 6. Take any $t_0 \in (0, \sigma]$ and $\xi \in (0, R)$; for a while we fix them.

Let us consider the equation

$$(4.3) \quad \frac{d\rho}{dt} = -\frac{B(t, \rho)}{t}, \quad \rho(t_0) = \xi$$

in the region $(0, t_0] \times (0, R)$. Since $B(t, \rho)/t$ is a continuous function on $(0, t_0] \times (0, R)$ we have a local solution (not necessarily unique). We take a maximally extended solution $\rho(t)$ and we denote by $(t_\xi, t_0]$ the interval of existence of this maximally extended solution. Set

$$q^*(t) = q(t, \rho(t)), \quad t_\xi < t \leq t_0.$$

Lemma 7. *Under the above situation, we have the following inequality for any (t_1, τ) satisfying $t_\xi < t_1 < \tau \leq t_0$:*

$$(4.4) \quad q^*(\tau) \leq \left(\frac{t_1}{\tau}\right)^h q^*(t_1).$$

Proof. By (4.1) we have

$$\begin{aligned} \left(t \frac{d}{dt} + h\right) q^*(t) &= \left(t \frac{\partial}{\partial t} + h\right) q(t, \rho) \Big|_{\rho=\rho(t)} + \frac{\partial q}{\partial \rho}(t, \rho(t)) \times t \frac{d\rho(t)}{dt} \\ &\leq \left[B(t, \rho) \frac{\partial q}{\partial \rho} + \frac{\partial q}{\partial \rho}(-B(t, \rho)) \right] \Big|_{\rho=\rho(t)} = 0, \end{aligned}$$

that is,

$$\left(t \frac{d}{dt} + h\right) q^*(t) \leq 0 \quad \text{on } (t_\xi, t_0].$$

Since this is equivalent to

$$\frac{d}{dt}(t^h q^*(t)) \leq 0 \quad \text{on } (t_\xi, t_0],$$

by integrating this from t_1 to τ we have $\tau^h q^*(\tau) \leq t_1^h q^*(t_1)$. This proves (4.4). \square

Step 3. By (4.2) and Lemma 7 we have

Lemma 8. *Under the above situation, we have the following inequalities for any t_1 satisfying $t_\xi < t_1 < t_0$:*

$$(4.5) \quad \xi \leq \rho(t_1) \leq \xi + \frac{C_1}{\kappa}(t_0^\kappa - t_1^\kappa) + \frac{C_2}{h}r + \frac{3C_3}{2h}r^{2/3} + \frac{3C_4}{h}r^{1/3}.$$

Proof. Let $t_\xi < t_1 < t_0$. By (4.3) we have

$$\rho(t_1) = \xi + \int_{t_1}^{t_0} B(\tau, \rho(\tau)) \frac{d\tau}{\tau}.$$

Since $B(t, \rho) \geq 0$ we have $\rho(t_1) \geq \xi$. Since $q(\tau, \rho(\tau)) = q^*(\tau)$, by (4.2) and (4.4) we have

$$\begin{aligned} B(\tau, \rho(\tau)) &\leq C_1 \tau^\kappa + C_2 q^*(\tau) + C_3 (q^*(\tau))^{2/3} + C_4 (q^*(\tau))^{1/3} \\ &\leq C_1 \tau^\kappa + C_2 \left(\frac{t_1}{\tau}\right)^h q^*(t_1) + C_3 \left(\frac{t_1}{\tau}\right)^{2h/3} q^*(t_1)^{2/3} + C_4 \left(\frac{t_1}{\tau}\right)^{h/3} q^*(t_1)^{1/3} \\ &\leq C_1 \tau^\kappa + C_2 \left(\frac{t_1}{\tau}\right)^h r + C_3 \left(\frac{t_1}{\tau}\right)^{2h/3} r^{2/3} + C_4 \left(\frac{t_1}{\tau}\right)^{h/3} r^{1/3} \end{aligned}$$

for $t_1 < \tau \leq t_0$, and so

$$(4.6) \quad \begin{aligned} \rho(t_1) &\leq \xi + \int_{t_1}^{t_0} \left(C_1 \tau^\kappa + C_2 \left(\frac{t_1}{\tau}\right)^h r \right. \\ &\quad \left. + C_3 \left(\frac{t_1}{\tau}\right)^{2h/3} r^{2/3} + C_4 \left(\frac{t_1}{\tau}\right)^{h/3} r^{1/3} \right) \frac{d\tau}{\tau}. \end{aligned}$$

Since

$$\int_{t_1}^{t_0} \left(\frac{t_1}{\tau}\right)^a \frac{d\tau}{\tau} = \frac{1}{a} \left(1 - \frac{t_1^a}{t_0^a}\right) \leq \frac{1}{a}$$

holds for any $a > 0$, by applying this to (4.6) we have (4.5). \square

Corollary 2. *If $\xi \in (0, R/2)$ we have $t_\xi = 0$.*

Proof. Let $\xi \in (0, R/2)$. Let us show that the condition $t_\xi > 0$ leads us to a contradiction.

Suppose that $t_\xi > 0$ holds. Then, by Lemmas 6 and 8 we have

$$0 < \xi \leq \rho(t_1) \leq \xi + \frac{C_1}{\kappa} \sigma^\kappa + \frac{C_2}{h} r + \frac{3C_3}{2h} r^{2/3} + \frac{3C_4}{h} r^{1/3} = R_1 < R$$

for any t_1 satisfying $t_\xi < t_1 < t_0$. Since $K = \{\rho \in (0, R) ; \xi \leq \rho \leq R_1\}$ is a compact subset of $(0, R)$ and since $\rho(t_1) \in K$ holds for any $t_1 \in (t_\xi, t_0]$, by a general theory of ordinary differential equations (for example, by Theorem 4.1 in [2]) we can extend $\rho(t)$ to $(t_\xi - \delta, t_0]$ for some $\delta > 0$. This contradicts the assumption that $(t_\xi, t_0]$ is the interval of existence of a maximally extended solution $\rho(t)$. \square

Step 4. Since $t_\xi = 0$, by (4.4) with $\tau = t_0$ we have

$$0 \leq q^*(t_0) \leq \left(\frac{t_1}{t_0}\right)^h q^*(t_1) \leq \left(\frac{t_1}{t_0}\right)^h r$$

for any $0 < t_1 < t_0$. Since $r > 0$ is independent of t_1 , by letting $t_1 \rightarrow 0$ we have $q^*(t_0) = 0$. Since $q^*(t_0) = q(t_0, \xi)$ we have $q(t_0, \rho) = 0$ for any $\rho \in (0, R/2)$. By the continuity we have $q(t_0, \rho) = 0$ for any $\rho \in [0, R/2)$. Since $t_0 \in (0, \sigma]$ is taken arbitrarily we have $q(t, \rho) = 0$ on $(0, \sigma] \times [0, R/2)$.

This completes the proof of Proposition 1. \square

5. APPLICATION

As an application of Theorem 2, we have

Theorem 3 (Analytic continuation). *Let $u(t, x)$ be a holomorphic solution of (2.1) on $S_I(r_0) \times D_{R_0}$ for some nonempty open interval I , $r_0 > 0$ and $R_0 > 0$. If $u(t, x)$ satisfies (2.3), $u(t, x)$ can be continued holomorphically up to a neighborhood of $(0, 0) \in \mathbb{C}_t \times \mathbb{C}_x^n$.*

REFERENCES

- [1] D. B. Bacani and H. Tahara, Unique solvability of some nonlinear partial differential equations with Fuchsian and irregular singularities, *J. Math. Soc. Japan*, 66 (2014), no. 3, 1017-1042.
- [2] E. A. Coddington and N. Levinson, *Theory of ordinary differential equations*, McGraw-Hill, 1955.
- [3] R. Gérard and H. Tahara, Solutions holomorphes et singulières d'équations aux dérivées partielles singulières non linéaires, *Publ. Res. Inst. Math. Sci.*, 29 (1993), no. 1, 121-151.
- [4] R. Gérard and H. Tahara, Singular nonlinear partial differential equations, *Aspects of Mathematics*, Friedr. Vieweg & Sohn, Braunschweig, 1996.
- [5] H. Tahara, Uniqueness of the solution of non-linear singular partial differential equations, *J. Math. Soc. Japan*, 48 (1996), no. 4, 729-744.
- [6] H. Tahara, Uniqueness of the solution of nonlinear singular first order partial differential equations, *J. Math. Soc. Japan*, 72 (2020), no. 4, 1259-1282.
- [7] H. Tahara and H. Yamazawa, Structure of solutions of nonlinear partial differential equations of Gérard-Tahara type, *Publ. Res. Inst. Math. Sci.*, 41 (2005), no. 2, 339-373.

DEPARTMENT OF INFORMATION AND COMMUNICATION SCIENCES, SOPHIA UNIVERSITY, TOKYO 102-8554, JAPAN.

Email address: h-tahara@sophia.ac.jp