

# EXTENDED-CYCLE INTEGRALS OF MODULAR FUNCTIONS FOR BADLY APPROXIMABLE NUMBERS

YUYA MURAKAMI

ABSTRACT. Cycle integrals of modular functions are expected to play a role in real quadratic analogue of singular moduli. In this paper, we extend the definition of cycle integrals of modular functions from real quadratic numbers to badly approximable numbers. We also give explicit representations of values of extended-cycle integrals for some cases.

## CONTENTS

1.	Introduction	1
	Acknowledgement	5
2.	Fundamental properties of the extended val function	5
3.	The extended val function as the limit value along a geodesic	5
4.	The extended val function as the limit value along a continued fraction	9
5.	Elementary units for badly approximable numbers	13
6.	An explicit computation of values of extended val function	14
7.	The value of val function at the Thue-Morse word	16
	References	17

## 1. INTRODUCTION

The elliptic modular  $j$ -function  $j(z)$  is an  $\mathrm{SL}_2(\mathbb{Z})$ -invariant holomorphic function on the upper half-plane  $\mathbb{H} := \{z = x + y\sqrt{-1} \in \mathbb{C} \mid x, y \in \mathbb{R}, y > 0\}$ . It plays an essential role in the complex multiplication theory. Indeed, its special values at imaginary quadratic points generate the Hilbert class fields of imaginary quadratic fields. On the other hand, for real quadratic fields, any definitive construction of the Hilbert class fields or ray class fields is still unknown except for a few cases, that is [Shi72], [Shi78]. As he expected a candidate for that purpose, Kaneko [Kan09] introduced the “value” of  $j(z)$ , written  $\mathrm{val}(w)$  derived from Dedekind’s notation, at any real quadratic number  $w$  which is defined by

$$(1.1) \quad \mathrm{val}(w) := \frac{1}{2 \log \varepsilon_w} \int_{z_0}^{\gamma_w z_0} j(z) \left( \frac{1}{z - w'} - \frac{1}{z - w} \right) dz.$$

Here,  $z_0 \in \mathbb{H}$  is any point,  $w'$  is the non-trivial Galois conjugate of  $w$ ,  $\mathrm{SL}_2(\mathbb{Z})_w$  be the stabilizer of  $w$  in  $\mathrm{SL}_2(\mathbb{Z})$ , and  $\gamma_w = \begin{pmatrix} * & * \\ c & d \end{pmatrix}$  be the unique element of  $\mathrm{SL}_2(\mathbb{Z})_w$  such that  $\mathrm{SL}_2(\mathbb{Z})_w = \{\pm \gamma_w^n \mid n \in \mathbb{Z}\}$  and  $\varepsilon_w := cw + d > 1$ . The integral is independent of  $z_0$  since the integrand is holomorphic and invariant under  $\gamma_w$ . We remark that  $\varepsilon_w$  is the minimal unit greater than 1 with the norm 1 in the order  $\mathfrak{O}_w$  in the real quadratic field  $\mathbb{Q}(w)$  whose discriminant coincides with the discriminant of  $w$ .

This value has a representation as a cycle integral. Let

$$\widetilde{\mathrm{val}}(w) := \int_{z_0}^{\gamma_w z_0} j(z) \left( \frac{1}{z - w'} - \frac{1}{z - w} \right) dz, \quad \widetilde{\mathfrak{I}}(w) := \int_{z_0}^{\gamma_w z_0} \left( \frac{1}{z - w'} - \frac{1}{z - w} \right) dz,$$

---

*Date:* October 19, 2021.

The author is supported by JSPS KAKENHI Grant Number JP 20J20308.

and

$$\text{length}(\text{SL}_2(\mathbb{Z})_w \backslash S_w) := \int_{\text{SL}_2(\mathbb{Z})_w \backslash S_w} ds$$

where  $ds := y^{-1} \sqrt{dx^2 + dy^2}$  and  $S_w$  is the geodesic in  $\mathbb{H}$  from  $w'$  to  $w$ . Then, we have

$$\widetilde{\text{val}}(w) = \int_{\text{SL}_2(\mathbb{Z})_w \backslash S_w} j(z) ds, \quad \widetilde{1}(w) = 2 \log \varepsilon_w = \text{length}(\text{SL}_2(\mathbb{Z})_w \backslash S_w).$$

In the last decade, it turned out that the values of  $\widetilde{\text{val}}$  have interesting properties. They are related to a mock modular form of weight  $1/2$  for  $\Gamma_0(4)$  studied by Duke-Imamoglu-Tóth [DIT11], which is a real quadratic analogue of Zagier's work on traces of singular moduli in [Zag02]. They are also related to a locally harmonic Maass form of weight 2 studied by Matsusaka [Mat20, Theorem 1.1]. Bengoechea-Imamoglu [BI19, Theorem 1.1] showed some kind of continuity for the values of  $\text{val}$  at Markov quadratics which is conjectured by Kaneko [Kan09]. The author [Mur20, Theorem 1.1] generalized it to the values of  $\text{val}$  at real quadratic numbers whose periods of continued fraction expansions have long cyclic parts. It revealed that the continuity of  $\text{val}$  differs from Euclidean topology. However, the definition of the continuity of  $\text{val}$  is still missing. We would expect a topological space  $H$  containing all real quadratic numbers and a continuous function  $?: H \rightarrow \mathbb{C}$  such that the diagram

$$\begin{array}{ccc} H & & \\ \uparrow & \searrow ? & \\ \{\text{real quadratic numbers}\} & \xrightarrow{\text{val}} & \mathbb{C} \end{array}$$

commutes.

In this paper, we attempt to choose  $H$  as a set of badly approximable numbers and give some partial continuity of  $\text{val}$  on  $H$ .

To explain the main results, we introduce some notation. Each real number can be represented by the unique continued fraction

$$[k_1, k_2, k_3, \dots] := k_1 + \frac{1}{k_2 + \frac{1}{k_3 + \frac{1}{\ddots}}}$$

where  $k_1$  is an integer and  $k_2, k_3, \dots$  are positive integers. An irrational number  $x = [k_1, k_2, k_3, \dots]$  is called a badly approximable number if the set  $\{k_1, k_2, k_3, \dots\}$  is bounded. For example, real quadratic irrational numbers are badly approximable since they have periodic continued fraction expansions. We denote  $\mathbb{R}_{\text{ba}}$  as the set of badly approximable numbers. The group  $\text{SL}_2(\mathbb{Z})$  acts on  $\mathbb{R}_{\text{ba}}$  as a linear fractional transformation.

We denote  $\infty$  as the point at infinity. Let  $\mathbb{P}^1(\mathbb{R}) := \mathbb{R} \cup \{\infty\}$ . For two points  $x', x \in \mathbb{H} \cup \mathbb{P}^1(\mathbb{R})$ , we denote by  $S_{x', x}$  the geodesic in  $\mathbb{H}$  from  $x'$  to  $x$ . For an even number  $2r \in 2\mathbb{Z}_{\geq 0}$  and positive integers  $k_1, \dots, k_{2r}$ , we call  $W = (k_1, \dots, k_{2r})$  an even word. In the case when  $r = 0$ , we call it the empty word and denote it by  $\emptyset$ . For even words  $W_1 = (k_1^{(1)}, \dots, k_{2r_1}^{(1)}), \dots, W_n = (k_1^{(n)}, \dots, k_{2r_n}^{(n)})$ , define its product

$$W_1 \cdots W_n := (k_1^{(1)}, \dots, k_{2r_1}^{(1)}, \dots, k_1^{(n)}, \dots, k_{2r_n}^{(n)}).$$

For an infinite sequence of even words  $W_1 = (k_1^{(1)}, \dots, k_{2r_1}^{(1)}), \dots, W_n = (k_1^{(n)}, \dots, k_{2r_n}^{(n)}), \dots$ , define

$$[W_1 \cdots W_n \cdots] := [k_1^{(1)}, \dots, k_{2r_1}^{(1)}, \dots, k_1^{(n)}, \dots, k_{2r_n}^{(n)}, \dots].$$

For the empty word  $\emptyset$ , an even word  $V = (l_1, \dots, l_{2s})$ , and a non-empty even word  $W = (k_1, \dots, k_{2r})$ , define

$$\begin{aligned}\gamma_\emptyset &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}), \\ \gamma_W &:= \begin{pmatrix} k_1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} k_2 & 1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} k_{2r} & 1 \\ 1 & 0 \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z}), \\ |W| &:= r, \\ [V\overline{W}] &:= [l_1, \dots, l_{2s}, k_1, \dots, k_{2r}, k_1, \dots, k_{2r}, k_1, \dots, k_{2r}, \dots], \\ N(W) &:= \max\{n \in \mathbb{Z}_{>0} \mid \text{there exists an even word } W_1 \text{ such that } W = W_1^n\}.\end{aligned}$$

For the real quadratic number  $w = [V\overline{W}]$ , define

$$(1.2) \quad \widehat{\mathrm{val}}(w) = \frac{N(W)}{|W|} \widetilde{\mathrm{val}}(w), \quad \widehat{\mathbf{1}}(w) = \frac{N(W)}{|W|} \widetilde{\mathbf{1}}(w), \quad \widehat{\varepsilon}_w = \varepsilon_w^{N(W)/|W|}.$$

We now define  $\mathrm{val}(x)$  for a badly approximable number  $x$  as follows. Take  $x' \in \mathbb{P}^1(\mathbb{R}) \setminus \{x\}$  and  $z_0 \in \mathbb{H}$ . Let  $W_1, W_2, \dots$  be an infinite sequence of even words such that  $x = [W_1 W_2 \cdots]$ . The set  $\{W_1, W_2, \dots\}$  is finite since  $x$  is badly approximable. We remark that if  $x < 1$  then the first entry of  $W_1$  is not positive. Let  $c_n \in \mathbb{Z}_{>0}$  be the denominator of the rational number  $[W_1 \cdots W_n]$  and put

$$L := \lim_{n \rightarrow \infty} \frac{|W_1| + \cdots + |W_n|}{n}.$$

We define the limits  $\mathrm{val}(x)$ ,  $\widehat{\mathrm{val}}(x)$ ,  $\widehat{\mathbf{1}}(x)$ , and  $\widehat{\varepsilon}_x$  by

$$(1.3) \quad \mathrm{val}(x) := \lim_{\substack{z \in S_{z_0, x} \\ z \rightarrow x}} \frac{1}{\mathrm{length}(S_{z_0, z})} \int_{S_{z_0, z}} j ds,$$

$$(1.4) \quad \widehat{\mathrm{val}}(x) := \lim_{n \rightarrow \infty} \frac{1}{|W_1| + \cdots + |W_n|} \int_{z_0}^{\gamma_{W_1 \cdots W_n} z_0} j(z) \frac{-1}{z-x} dz,$$

$$(1.5) \quad \widehat{\mathbf{1}}(x) := \lim_{n \rightarrow \infty} \frac{1}{|W_1| + \cdots + |W_n|} \int_{z_0}^{\gamma_{W_1 \cdots W_n} z_0} \frac{-1}{z-x} dz,$$

$$(1.6) \quad \widehat{\varepsilon}_x := \lim_{n \rightarrow \infty} c_n^{1/Ln}.$$

We study the limit values on the right-hand side of each of (1.3) through (1.6). It will be proved that the limit values are bounded, but we do not know if the limit values are singleton. Hence, the right-hand sides of the above equations do converge for any badly approximable number. We can also prove that if  $x$  is a real quadratic number, then the limits  $\mathrm{val}(x)$ ,  $\widehat{\mathrm{val}}(x)$ ,  $\widehat{\mathbf{1}}(x)$ , and  $\widehat{\varepsilon}_x$  converge and coincide with the values defined in (1.1) and (1.2).

Our purpose in this paper is to give uncountable infinitely many badly approximable numbers  $x$  such that the limits  $\mathrm{val}(x)$ ,  $\widehat{\mathrm{val}}(x)$ ,  $\widehat{\mathbf{1}}(x)$ , and  $\varepsilon_x$  converge. We also explicitly represent values of extended val function at badly approximable numbers which have infinitely long cyclic parts in their continued fractions. Our first main result is as follows.

**Theorem 1.1.** *Let  $V_0, \dots, V_k$  be even words,  $W_1, \dots, W_k$  be non-empty even words, and  $\{a_{1,n}\}_{n=1}^\infty, \dots, \{a_{k,n}\}_{n=1}^\infty$  be sequences in  $\mathbb{Z}_{\geq 0}$  such that  $a_{i,n} \rightarrow \infty$  and  $2^{-n} a_{i,n} \rightarrow 0$  when  $n$  goes to infinity. Let  $w_i := [W_i]$ ,  $k' := \{0 \leq i \leq k \mid V_i \neq \emptyset\}$ ,*

$$A_n := k'n + \sum_{1 \leq i \leq k} \sum_{1 \leq j \leq n} a_{i,j}, \quad a_i := \lim_{n \rightarrow \infty} \frac{1}{A_n} \sum_{1 \leq j \leq n} a_{i,j},$$

and

$$U_n := V_1 W_1^{a_{1,n}} V_2 W_2^{a_{2,n}} \cdots V_k W_k^{a_{k,n}}, \quad x := [U_1 U_2 \cdots U_n \cdots].$$




FIGURE 1. The paths for the definitions of  $\text{val}(x)$ ,  $\widehat{\text{val}}(x)$ , and  $\widehat{1}(x)$  in (1.3), (1.4), and (1.5)

Then the limits  $\text{val}(x)$ ,  $\widehat{\text{val}}(x)$ ,  $\widehat{1}(x)$ , and  $\widehat{\varepsilon}_x$  converge and it holds  $\widehat{1}(x) = 2 \log \widehat{\varepsilon}_x > 0$  and  $\text{val}(x) = \widehat{\text{val}}(x)/\widehat{1}(x)$ . Moreover, it also holds

$$(1.7) \quad \text{val}(x) = \frac{a_1 N(W_1) \widehat{\text{val}}(w_1) + \cdots + a_k N(W_k) \widehat{\text{val}}(w_k)}{a_1 N(W_1) \widehat{1}(w_1) + \cdots + a_k N(W_k) \widehat{1}(w_k)}.$$

It is proved that the right-hand side in (1.7) equals to the limit of  $\text{val}([\overline{U}_n])$  in [Mur20, Theorem 1.1].

Next, we consider the value of  $\text{val}$  function at the Thue-Morse word. It is a typical word which is repetition-free. We fix two even words  $V$  and  $W$ . Let  $\{V, W\}^\omega$  be the set of even words generated by  $V, W$  which are of finite length. Let  $\{V, W\}^\omega$  be the set of even words generated by  $V, W$  which are of infinite length. We define the monoid homomorphism  $h: \{V, W\}^* \rightarrow \{V, W\}^*$  by  $h(V) := VW, h(W) := WV$ . In this setting, the word

$$V_h := \lim_{n \rightarrow \infty} h^n(V) = VWVWVWVW \cdots \in \{V, W\}^\omega$$

is called the Thue-Morse word. It is known that  $V_h$  is cubefree (for example, see [CK97, Section 3]), and thus it is not a word which satisfies the conditions in Theorem 1.1.

The value of  $\text{val}$  function at the Thue-Morse word is calculated as follows.

**Theorem 1.2.** *Let  $x := [V_h]$  and  $w_n := [h^n(V)]$ . Then  $\widehat{\text{val}}(x)$  converges if and only if  $\lim_{n \rightarrow \infty} \widehat{\text{val}}(w_{2n})/2^{2n}$  converges. Moreover, we have*

$$\widehat{\text{val}}(x) = \lim_{n \rightarrow \infty} \frac{1}{2^{2n}} \widehat{\text{val}}(w_{2n}), \quad \widehat{1}(x) = \lim_{n \rightarrow \infty} \frac{1}{2^{2n}} \widehat{1}(w_{2n}), \quad \text{val}(x) = \lim_{n \rightarrow \infty} \frac{1}{2^{2n}} \text{val}(w_{2n}).$$

The definitions of  $\widehat{\text{val}}(x)$  and  $\widehat{1}(x)$  are like the Birkoff average in ergodic theory. Khinchin-Lévy's theorem in ergodic theory asserts that for almost all  $x \in [0, 1]$  with respect to the measure

$$d\mu := \frac{1}{\log 2} \frac{dx}{1+x}$$

on  $[0, 1]$ , the limit defining  $\widehat{\varepsilon}_x$  converges and we have

$$\log \widehat{\varepsilon}_x = -2 \int_0^1 \log(x) d\mu = \frac{\pi^2}{6 \log 2}.$$

Although ergodic theory makes it possible to study values like the Birkoff average at almost all points, it does not tell us about a value at each point, for example, a badly approximable number. We can state that this paper studies a range which ergodic theory does not deal with.

This paper will be organized as follows. In Section 2, we state some fundamental properties of the limits  $\text{val}(x)$ ,  $\widehat{\text{val}}(x)$ ,  $\widehat{1}(x)$ , and  $\widehat{\varepsilon}_x$  for each badly approximable number  $x$ . In Section 3, we describe  $\text{val}(x)$  in terms of the limit value along a geodesic and study its properties. In Section 4, we also study basic properties of  $\widehat{\text{val}}(x)$  and  $\widehat{1}(x)$  which are defined using the continued fraction expansion of  $x$ . In Section 5, we study  $\varepsilon_x$  and a relation between it and  $\widehat{1}(x)$ . Finally we prove Theorem 1.1 in Section 6 and Theorem 1.2 in Section 7.

#### ACKNOWLEDGEMENT

I would like to show my greatest appreciation to Professor Takuya Yamauchi for giving many pieces of advice. I am deeply grateful to Dr. Toshiki Matsusaka for giving many comments. I would like to express my gratitude to Professor Shun'ichi Yokoyama and Dr. Toshihiro Suzuki for giving me constructive comments on computing  $\text{val}(x)$  numerically. It is a pleasure to extend my thanks to Professor Tatsuya Tate for teaching me ergodic theory. I also thank Dr. Daisuke Kazukawa, Dr. Hiroki Nakajima, and Dr. Shin'ichiro Kobayashi for teaching me geodesics, hyperbolic geometry, and metric spaces. I appreciate the technical assistance of Dr. Naruaki Kato for introducing me to how to write works by using GitHub.

#### 2. FUNDAMENTAL PROPERTIES OF THE EXTENDED $\text{val}$ FUNCTION

In this section, we state fundamental properties of the limits  $\text{val}(x)$ ,  $\widehat{\text{val}}(x)$ ,  $\widehat{1}(x)$ , and  $\widehat{\varepsilon}_x$ . In the later sections, we will prove the following.

**Theorem 2.1.** *Let  $x$  be a badly approximable number and  $W_1, W_2, \dots$  be an infinite sequence of even words such that  $x = [W_1 W_2 \dots]$  and the set  $\{W_1, W_2, \dots\}$  is finite. Take any points  $x' \in \mathbb{P}^1(\mathbb{R}) \setminus \{x\}$  and  $z_0 \in \mathbb{H}$ . Then the followings hold.*

- (i) *The limit values  $\text{val}(x)$ ,  $\widehat{\text{val}}(x)$ ,  $\widehat{1}(x)$ , and  $\widehat{\varepsilon}_x$  defined in each of (1.3), (1.4), (1.5), and (1.6) are bounded. Moreover, if they converge then they are independent of  $x', z_0$  and  $W_1, W_2, \dots$ . Further, they are  $\text{SL}_2(\mathbb{Z})$ -invariant.*
- (ii) *If  $x$  is a real quadratic number, then  $\text{val}(x)$ ,  $\widehat{\text{val}}(x)$ ,  $\widehat{1}(x)$ , and  $\widehat{\varepsilon}_x$  converge and coincide with the values defined in (1.1) and (1.2).*
- (iii) *If  $\widehat{\text{val}}(x)$  and  $\widehat{1}(x)$  converge, then for any point  $x' \in \mathbb{P}^1(\mathbb{R}) \setminus \{x\}$  and a sequence  $\{x_n\}_{n=1}^\infty$  of real numbers with  $z_n := \gamma_{W_1 \dots W_n}(x_n + \sqrt{-1}) \in S_{\infty, x}$ ,  $z_n \rightarrow x$ , we have*

$$\widehat{\text{val}}(x) = \lim_{n \rightarrow \infty} \frac{1}{|W_1| + \dots + |W_n|} \int_{z_0}^{z_n} j(z) \left( \frac{1}{z - x'} - \frac{1}{z - x} \right) dz,$$

$$\widehat{1}(x) = \lim_{n \rightarrow \infty} \frac{1}{|W_1| + \dots + |W_n|} \int_{z_0}^{z_n} \left( \frac{1}{z - x'} - \frac{1}{z - x} \right) dz.$$

Here we define  $1/(z - \infty) := 0$  if  $x' = \infty$ .

- (iv) *The limit  $\widehat{\varepsilon}_x$  converges if and only if  $\widehat{1}(x)$  converges. Moreover, we have  $\widehat{1}(x) = 2 \log \widehat{\varepsilon}_x$ .*
- (v) *It holds  $\widehat{\varepsilon}_x \geq (3 + \sqrt{5})/2$ . In particular,  $\widehat{1}(x) > 0$ .*
- (vi) *If  $\widehat{\text{val}}(x)$  and  $\widehat{1}(x)$  converge, then  $\text{val}(x)$  converges and we have  $\text{val}(x) = \widehat{\text{val}}(x)/\widehat{1}(x)$ .*

Here we remark that Theorem 2.1 (vi) follows from Theorem 2.1 (iv) and (v).

#### 3. THE EXTENDED $\text{val}$ FUNCTION AS THE LIMIT VALUE ALONG A GEODESIC

In this section, we prove Theorem 2.1 (i) and (ii) for  $\text{val}(x)$ . We start with the following properties of badly approximable numbers. Let  $\pi: \mathbb{H} \rightarrow \text{SL}_2(\mathbb{Z}) \backslash \mathbb{H}$  be the natural projection.

**Proposition 3.1** ([Dal11, Chapter VII, Theorem 3.4], [Dal11, Chapter VII, Lemma 3.5]). *For an irrational number  $x \in \mathbb{R} \setminus \mathbb{Q}$ , the followings are equivalent.*

- (i)  *$x$  is badly approximable;*




FIGURE 2. The path of integrals for the limit values in Theorem 2.1 (iii)

- (ii)  $L(x) := \sup \{L > 0 \mid \exists p_n, q_n \in \mathbb{Z} \text{ s.t. } (p_n, q_n) = 1, q_n \rightarrow \infty, |x - p_n/q_n| \leq 1/Lq_n^2\} < \infty;$
- (iii)  $h(x) := \sup \{t > 0 \mid \exists z_n \in S_{z_0, x} \text{ s.t. } z_n \rightarrow x, \pi(z_n) \in \pi(\{z \in \mathbb{H} \mid \text{Im}(z) = t\})\} < \infty;$
- (iv) For any point  $z_0 \in \mathbb{H}$ , the closure  $\overline{\pi(S_{z_0, x})} \subset \text{SL}_2(\mathbb{Z}) \backslash \mathbb{H}$  is compact.

The constant  $L(x)$  is called the Lagrange number of  $x$ . Usually, badly approximable numbers are defined as irrational numbers with finite Lagrange numbers.

The following is a key fact in this section.

**Theorem 3.2** (Theorem 2.1 (i) and (ii) for  $\text{val}(x)$ ). *Let  $x$  be a badly approximable number. Take any point  $z_0 \in \mathbb{H}$ . Then the followings hold.*

- (i) *The limit values*

$$\text{val}(x) := \lim_{\substack{z \in S_{z_0, x} \\ z \rightarrow x}} \frac{1}{\text{length}(S_{z_0, z})} \int_{S_{z_0, z}} j(z) ds$$

*is bounded.*

- (ii) *For any matrix  $\gamma \in \text{SL}_2(\mathbb{Z})$ , we have  $\text{val}(\gamma x) = \text{val}(x)$ .*
- (iii) *If the limit  $\text{val}(x)$  converges, then it is independent of  $z_0$ .*
- (iv) *For each a real quadratic number  $x$ ,  $\text{val}(x)$  converges and coincides with the value defined in (1.1).*

*Proof.* First we will prove (i). Take  $\sigma \in \text{SL}_2(\mathbb{R})$  and  $t \in \mathbb{R}_{>0}$  such that  $\sigma^{-1}x = \infty$  and  $\sigma^{-1}z_0 = t\sqrt{-1}$ . For a point  $z \in S_{z_0, x}$ , let  $d := \text{length}(S_{z_0, z})$ . Then we have

$$\int_{S_{z_0, z}} j(z) ds = \int_{t\sqrt{-1}}^{e^d t\sqrt{-1}} (j \circ \sigma) ds$$

since  $\sigma^{-1}z = e^d t\sqrt{-1}$ . Let  $a_d$  denote this integral. Since

$$\text{val}(x) = \lim_{d \rightarrow \infty} \frac{a_d}{d},$$

it is enough to show that the set  $\{a_d/d\}_{d \in \mathbb{R}_{>0}}$  is bounded.

Since  $x$  is badly approximable, the closure  $\overline{\pi(S_{x,z_0})} \subset \mathrm{SL}_2(\mathbb{Z}) \backslash \mathbb{H}$  is compact by Proposition 3.1. Thus, we can choose  $K > 0$  such that  $|j(z)| \leq K$  on  $S_{z_0,x}$ . Since

$$\left| \frac{a_d}{d} \right| = \left| \frac{1}{d} \int_{t\sqrt{-1}}^{e^{dt}\sqrt{-1}} (j \circ \sigma) ds \right| \leq \frac{K}{d} \int_{t\sqrt{-1}}^{e^{dt}\sqrt{-1}} \frac{dz}{z} = K,$$

$\{a_d/d\}$  is bounded.

The second claim (ii) follows from the fact that  $j(z)$  is  $\mathrm{SL}_2(\mathbb{Z})$ -invariant.

As for the third claim (iii), for each point  $z_0 \in \mathbb{H}$ , we consider

$$\mathrm{val}(x, z_0) := \lim_{\substack{z \in S_{z_0,x} \\ z \rightarrow x}} \frac{1}{\mathrm{length}(S_{z_0,z})} \int_{S_{z_0,z}} j(z) ds.$$

We will show that this value is independent of  $z_0$  by two steps.

**Step 1.** We will show that  $\mathrm{val}(x, z_0) = \mathrm{val}(x, z_1)$  for any point  $z_1 \in S_{z_0,x}$ . For  $i \in \{0, 1\}$ , let

$$a_i(z) := \int_{S_{z_i,z}} j(z) ds, \quad b_i(z) := \int_{S_{z_i,z}} ds.$$

Clearly,

$$\lim_{\substack{z \in S_{z_0,x} \\ z \rightarrow x}} b_i(z) = \infty.$$

Since

$$a_0(z) - a_1(z) = \int_{S_{z_0,z_1}} j(z) ds, \quad b_0(z) - b_1(z) = \int_{S_{z_0,z_1}} ds$$

is bounded, we have

$$\mathrm{val}(x, z_0) = \lim_{\substack{z \in S_{z_0,x} \\ z \rightarrow x}} \frac{a_0(z)}{b_0(z)} = \lim_{\substack{z \in S_{z_0,x} \\ z \rightarrow x}} \frac{a_1(z) + O(1)}{b_1(z) + O(1)} = \lim_{\substack{z \in S_{z_0,x} \\ z \rightarrow x}} \frac{a_1(z)}{b_1(z)} = \mathrm{val}(x, z_1).$$




FIGURE 3. The paths of integrals appearing Step 2 in the proof of Theorem 3.2 (iii)

**Step 2.** We will show that  $\mathrm{val}(x, z_0) = \mathrm{val}(x, z_1)$  for any points  $z_0, z_1 \in \mathbb{H}$ . This part is due to Matsusaka. Take a matrix  $\sigma \in \mathrm{SL}_2(\mathbb{R})$  such that  $\sigma^{-1}x = \infty$ . Let  $v_0$  be any positive number. By Step 1, we can replace  $z_i$  by  $\mathrm{Re}(z_i) + v_0\sqrt{-1}$ . Thus, we may assume  $v_0 = \mathrm{Im}(\sigma^{-1}z_0) =$

$\text{Im}(\sigma^{-1}z_1)$ . Let  $u_i := \text{Re}(\sigma^{-1}z_i)$  for  $i \in \{0, 1\}$ . Then  $\sigma^{-1}S_{z_i, x}$  is a subgeodesic of  $S_{u_i, \infty}$ . Since  $ds = y^{-1}\sqrt{dx^2 + dy^2} = y^{-1}dy$  on  $S_{u_i, \infty}$ , we have

$$\begin{aligned} \text{val}(x, z_i) &= \lim_{\substack{z \in S_{z_i, x} \\ z \rightarrow x}} \frac{1}{\text{length}(S_{z_i, z})} \int_{S_{z_i, z}} j(z) ds \\ &= \lim_{v \rightarrow \infty} \frac{1}{\text{length}(S_{\sigma(u_i + v\sqrt{-1}), \sigma(u_i + v_0\sqrt{-1})})} \int_{v_0}^v j(\sigma(u_i + y\sqrt{-1})) \frac{dy}{y}. \end{aligned}$$

Since

$$\text{length}(S_{\sigma(u_i + v\sqrt{-1}), \sigma(u_i + v_0\sqrt{-1})}) = \text{length}(S_{v\sqrt{-1}, v_0\sqrt{-1}}) = \int_{v_0}^v \frac{dy}{y} = \log \frac{v}{v_0},$$

we obtain

$$\text{val}(x, z_0) - \text{val}(x, z_1) = \lim_{v \rightarrow \infty} \frac{1}{\log v/v_0} \int_{v_0}^v (j(\sigma(u_0 + y\sqrt{-1})) - j(\sigma(u_1 + y\sqrt{-1}))) \frac{dy}{y}.$$

For two points  $z, z' \in \mathbb{H}$ , let

$$d_{\text{hyp}}(z, z') := \text{length}(S_{z, z'}), \quad d_{\text{SL}_2(\mathbb{Z}) \backslash \mathbb{H}}(\pi(z), \pi(z')) := \min \{d_{\text{hyp}}(\gamma z, z') \mid \gamma \in \text{SL}_2(\mathbb{Z})\}.$$

They define the hyperbolic distances on  $\mathbb{H}$  and  $\text{SL}_2(\mathbb{Z}) \backslash \mathbb{H}$  respectively. Since

$$\cosh d_{\text{hyp}}(\sigma(u_0 + y\sqrt{-1}), \sigma(u_1 + y\sqrt{-1})) = \cosh d_{\text{hyp}}(u_0 + y\sqrt{-1}, u_1 + y\sqrt{-1}) = 1 + \frac{|u_0 - u_1|}{2y^2}$$

by [Bea83, Theorem 7.2.1], we have

$$\lim_{y \rightarrow \infty} d_{\text{hyp}}(\sigma(u_0 + y\sqrt{-1}), \sigma(u_1 + y\sqrt{-1})) = 0.$$

Since  $x$  is badly approximable,  $\overline{\pi(S_{x, z_0})} \cup \overline{\pi(S_{x, z_1})} \subset \text{SL}_2(\mathbb{Z}) \backslash \mathbb{H}$  is compact by Proposition 3.1. Thus,  $j(z)$  is continuous on  $\overline{\pi(S_{x, z_0})} \cup \overline{\pi(S_{x, z_1})}$  with respect to the metric  $d_{\text{SL}_2(\mathbb{Z}) \backslash \mathbb{H}}$ . Hence for any  $\varepsilon > 0$ , there exists  $v_0 > 0$  such that

$$|j(\sigma(u_0 + y\sqrt{-1})) - j(\sigma(u_1 + y\sqrt{-1}))| < \varepsilon$$

for any  $y > v_0$ . Then we have

$$|\text{val}(x, z_0) - \text{val}(x, z_1)| \leq \lim_{v \rightarrow \infty} \frac{1}{\log v/v_0} \int_{v_0}^v \varepsilon \frac{dy}{y} = \varepsilon,$$

that is,  $\text{val}(x, z_0) = \text{val}(x, z_1)$ .

Finally, we prove (iv). Take any real quadratic irrational  $w$  and any point  $z_0 \in S_w$ . For a positive integer  $n$ , let  $z_n \in S_{z_0, w}$  be the unique point such that

$$\text{length}(S_{z_0, z_n}, d_{\text{hyp}}) = n \text{length}(\text{SL}_2(\mathbb{Z})_w \backslash S_w, d_{\text{SL}_2(\mathbb{Z}) \backslash \mathbb{H}}).$$

The value  $\text{val}(w)$  defined in (1.1) can be written as

$$\begin{aligned} \text{val}(w) &= \frac{1}{\text{length}(\text{SL}_2(\mathbb{Z})_w \backslash S_w)} \int_{\text{SL}_2(\mathbb{Z})_w \backslash S_w} j(z) ds \\ &= \lim_{n \rightarrow \infty} \frac{1}{\text{length}(S_{z_0, z_n})} \int_{S_{z_0, z_n}} j(z) ds \\ &= \lim_{\substack{z \in S_{z_0, w} \\ z \rightarrow w}} \frac{1}{\text{length}(S_{z_0, z})} \int_{S_{z_0, z}} j(z) ds. \end{aligned}$$

This value coincides with the definition of  $\text{val}(x)$  in (1.3).  $\square$



4. THE EXTENDED  $\text{val}$  FUNCTION AS THE LIMIT VALUE ALONG A CONTINUED FRACTION

In this section, we prove Theorem 2.1 (i), (ii), and (iii) for  $\widehat{\text{val}}(x)$  and  $\widehat{1}(x)$ . To begin with, we consider the case for real quadratic numbers. The following lemma proves Theorem 2.1 (iii).

**Lemma 4.1.** *For integers  $r \geq 0$  and  $s \geq 1$  and non-empty even words  $W_1, \dots, W_{r+s}$ , let  $w := [W_1 \cdots W_r \overline{W_{r+1} \cdots W_{r+s}}]$ . For integers  $0 \leq l < s$  and  $n \geq 1$ , let  $W_{r+l+ns} := W_{r+l}$ . For a positive integer  $i$ , let  $\gamma_i := \gamma_{W_i}$ . Let  $z_0 \in \mathbb{H}$  be a point. Then we have*

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \int_{z_0}^{\gamma_1 \cdots \gamma_n z_0} j(z) \left( \frac{1}{z-w'} - \frac{1}{z-w} \right) dz &= \frac{N(W_{r+1} \cdots W_{r+s})}{s} \widetilde{\text{val}}(w), \\ \lim_{n \rightarrow \infty} \frac{1}{n} \int_{z_0}^{\gamma_1 \cdots \gamma_n z_0} \left( \frac{1}{z-w'} - \frac{1}{z-w} \right) dz &= \frac{N(W_{r+1} \cdots W_{r+s})}{s} \widetilde{1}(w). \end{aligned}$$

*Proof.* Let  $N := N(W_{r+1} \cdots W_{r+s})$ . For a positive integer  $n$ , let

$$a_n := \int_{\gamma_1 \cdots \gamma_{n-1} z_0}^{\gamma_1 \cdots \gamma_n z_0} j(z) \left( \frac{1}{z-w'} - \frac{1}{z-w} \right) dz.$$

For  $n > r$ , it holds  $\gamma_{n+s} = \gamma_n$  and  $(\gamma_1 \cdots \gamma_{n+s-1})^{-1} w = (\gamma_1 \cdots \gamma_{n-1})^{-1} w$  since  $W_{n+r} = W_n$ . Thus, we have  $a_{n+s} = a_n$ . Since

$$\frac{1}{n} \int_{z_0}^{\gamma_1 \cdots \gamma_n z_0} j(z) \eta_w = \frac{a_1 + \cdots + a_n}{n},$$

it suffices to show that

$$\lim_{n \rightarrow \infty} \frac{a_1 + \cdots + a_{m+l+ns}}{m+l+ns} = \frac{N}{s} \widetilde{\text{val}}(w)$$

for a positive integer  $l$ .

Let  $z_1 := \gamma_1 \cdots \gamma_r z_0$ . Since  $\gamma_1 \cdots \gamma_r \gamma_{r+1} \cdots \gamma_{r+s} (\gamma_1 \cdots \gamma_r)^{-1} = \gamma_w^N$  by [Mur20, Lemma 2.3 (iv)], we have

$$\begin{aligned} N \widetilde{\text{val}}(w) &= \int_{z_1}^{\gamma_1 \cdots \gamma_r \gamma_{r+1} \cdots \gamma_{r+s} (\gamma_1 \cdots \gamma_r)^{-1} z_1} j(z) \left( \frac{1}{z-w'} - \frac{1}{z-w} \right) dz \\ &= \int_{\gamma_1 \cdots \gamma_r z_0}^{\gamma_1 \cdots \gamma_{r+s} z_0} j(z) \left( \frac{1}{z-w'} - \frac{1}{z-w} \right) dz \\ &= a_{r+1} + \cdots + a_{m+l+ns}. \end{aligned}$$

Thus, we have

$$\begin{aligned} &\lim_{n \rightarrow \infty} \frac{a_1 + \cdots + a_{m+l+ns}}{m+l+ns} \\ &= \lim_{n \rightarrow \infty} \left( \frac{a_1 + \cdots + a_m}{m+l+ns} + \frac{a_{m+1} + \cdots + a_{m+ns}}{m+l+ns} + \frac{a_{m+ns+1} + \cdots + a_{m+ns+l}}{m+l+ns} \right) \\ &= \lim_{n \rightarrow \infty} \left( \frac{n(a_{m+1} + \cdots + a_{m+s})}{m+l+ns} + \frac{a_{m+1} + \cdots + a_{m+l}}{m+l+ns} \right) \\ &= \frac{a_{m+1} + \cdots + a_{m+s}}{s} \\ &= \frac{N}{s} \widetilde{\text{val}}(w). \end{aligned}$$

The second equality is similarly proved.  $\square$

**Remark 4.2.** The first finite term of a continued fraction does not contribute to the left-hand side in Lemma 4.1. Thus, the right-hand side in Lemma 4.1 depends only on the period in the continued fraction.

For  $x', x \in \mathbb{P}^1(\mathbb{R}) = \mathbb{R} \cup \{\infty\}$ , define

$$\eta_{x',x}(z) := \left( \frac{1}{z-x'} - \frac{1}{z-x} \right) dz.$$

Here, let  $1/(z - \infty) := 0$  if  $x = \infty$  or  $x' = \infty$ . For any  $x, x' \in \mathbb{P}^1(\mathbb{R})$  and  $\gamma \in \mathrm{SL}_2(\mathbb{R})$ , we have  $\gamma^* \eta_{x',x} = \eta_{\gamma^{-1}x', \gamma^{-1}x}$  by direct calculation.

To prove Theorem 2.1 (ii), we prepare several lemmas.

**Lemma 4.3.** *For a badly approximable number  $x = [k_1, k_2, \dots]$ , let  $\gamma_n := \gamma_{(k_1, \dots, k_{2n})}$ . Then for any point  $x' \in \mathbb{P}^1(\mathbb{R})$ , the set  $\{\gamma_n^{-1}x' \mid n \in \mathbb{Z}_{>0}\}$  is bounded.*

*Proof.* Let  $h(x)$  be the number defined in Proposition 3.1. By Proposition 3.1, we have

$$\gamma_n^{-1}S_{x',x} \cap \{z \in \mathbb{H} \mid \mathrm{Im}(z) \geq h(x) + 1\} = \emptyset$$

for all sufficiently large  $n$ . Thus, the radius of  $\gamma_n^{-1}S_{x',x}$  is less than  $h(x) + 1$ . Since coefficients of the continued fraction expansion of  $x$  are bounded,  $\{\gamma_n^{-1}x \mid n \in \mathbb{Z}_{>0}\}$  is bounded. Thus,  $\{\gamma_n^{-1}x' \mid n \in \mathbb{Z}_{>0}\}$  is also bounded.  $\square$




FIGURE 4. Geodesics in the proof of Lemma 4.3

**Lemma 4.4.** *For an even word  $W$ , a point  $z_0 \in \mathbb{H}$ , and sequences  $\{x_n\}, \{x'_n\}, \{y_n\}, \{y'_n\}$ , we have*

$$\int_{z_0}^{\gamma_W z_0} j(z) (\eta_{x'_n, x_n} - \eta_{y'_n, y_n}) = O(|x_n - y_n| + |x'_n - y'_n|).$$

*Proof.* Since

$$\left| \frac{1}{z - x_n} - \frac{1}{z - y_n} \right| \leq \frac{|x_n - y_n|}{\min\{\mathrm{Im}(z_0), \mathrm{Im}(\gamma_W z_0)\}}$$

on the contour  $\{tz_0 + (1-t)\gamma_W z_0 \mid 0 \leq t \leq 1\}$ , it holds

$$\left| \int_{z_0}^{\gamma_W z_0} j(z) \left( \frac{1}{z - x_n} - \frac{1}{z - y_n} \right) \right| \leq \frac{|x_n - y_n|}{\min\{\mathrm{Im}(z_0), \mathrm{Im}(\gamma_W z_0)\}} \left| \int_{z_0}^{\gamma_W z_0} j(z) dz \right| = O(|x_n - y_n|).$$

Similar argument shows

$$\left| \int_{z_0}^{\gamma_W z_0} j(z) \left( \frac{1}{z - x'_n} - \frac{1}{z - y'_n} \right) \right| = O(|x'_n - y'_n|).$$

$\square$

**Lemma 4.5.** *For two points  $x, x' \in \mathbb{P}^1(\mathbb{R})$  with  $x \neq x'$ , we have  $ds = \eta_{x',x}$  on  $S_{x',x}$ .*

*Proof.* Let

$$\sigma := \begin{cases} \begin{pmatrix} x & x'/(x-x') \\ 1 & 1/(x-x') \end{pmatrix} & \text{if } x \neq \infty, \\ \begin{pmatrix} 1 & -x \\ 0 & 1 \end{pmatrix} & \text{if } x = \infty \end{cases}$$

be a matrix in  $\mathrm{SL}_2(\mathbb{R})$ . Since  $\sigma^* \eta_{x',x} = \eta_{0,\infty}$ ,  $\sigma^* ds = ds$ , it is enough to show when  $x = \infty$  and  $x' = 0$ . On the geodesic  $S_{0,\infty} = \{t\sqrt{-1} \mid t \in \mathbb{R}_{>0}\}$ , we have  $ds = dt/t = \eta_{0,\infty}$ .  $\square$

A key result in this section is the following.

**Theorem 4.6** (Theorem 2.1 (i) for  $\widehat{\mathrm{val}}(x)$  and  $\widehat{1}(x)$ ). *Let  $x$  be a badly approximable number and  $W_1, W_2, \dots$  be an infinite sequence of even words such that  $x = [W_1 W_2 \dots]$  and the set  $\{W_1, W_2, \dots\}$  is finite. Take a point  $z_0 \in \mathbb{H}$ . Then the followings hold.*

(i) *The sequences defining the limits*

$$\begin{aligned} \widehat{\mathrm{val}}(x) &:= \lim_{n \rightarrow \infty} \frac{1}{|W_1| + \dots + |W_n|} \int_{z_0}^{\gamma_{W_1 \dots W_n} z_0} j(z) \frac{-1}{z-x} dz, \\ \widehat{1}(x) &:= \lim_{n \rightarrow \infty} \frac{1}{|W_1| + \dots + |W_n|} \int_{z_0}^{\gamma_{W_1 \dots W_n} z_0} \frac{-1}{z-x} dz \end{aligned}$$

*is bounded.*

(ii) *The limits  $\widehat{\mathrm{val}}(x)$  and  $\widehat{1}(x)$  are independent of  $z_0$ .*

(iii) *The limits  $\widehat{\mathrm{val}}(x)$  and  $\widehat{1}(x)$  are independent of  $\{W_n \mid n \geq 1\}$ .*

(iv) *The limits  $\widehat{\mathrm{val}}(x)$  and  $\widehat{1}(x)$  are  $\mathrm{SL}_2(\mathbb{Z})$ -invariant.*

*Proof.* For the first claim (i), let

$$\begin{aligned} L &:= \lim_{n \rightarrow \infty} \frac{|W_1| + \dots + |W_n|}{2n}, \\ a_n(z_0) &:= \int_{\gamma_{W_1 \dots W_{n-1}} z_0}^{\gamma_{W_1 \dots W_n} z_0} j(z) \frac{-1}{z-x} dz, \\ K &:= \max \{ |j(z)| \mid |\mathrm{Re}(z)| \leq 1/2, |z| \leq 1, \mathrm{Im}(z) \leq h(x) \}, \\ R &:= \min \{ \mathrm{Im}(z_0), \mathrm{Im}(\gamma_{W_n} z_0) \mid n \in \mathbb{Z}_{>0} \}, \\ M &:= \sup \{ \gamma_{W_1 \dots W_n}^{-1} x, \gamma_{W_1 \dots W_n}^{-1} \infty \mid n \in \mathbb{Z}_{>0} \}. \end{aligned}$$

Here  $K$  and  $R$  are positive real numbers and  $L$  converges since the set  $\{W_1, W_2, \dots\}$  is finite. Since  $x$  is badly approximable,  $M$  is a positive real number by Lemma 4.3. We can write

$$\widehat{\mathrm{val}}(x) = \frac{1}{L} \lim_{n \rightarrow \infty} \frac{a_1(z_0) + \dots + a_n(z_0)}{n}.$$

For any positive integer  $n_0$ , we have

$$\begin{aligned} |a_1(z_0) + \dots + a_n(z_0)| &\leq \int_{z_0}^{\gamma_{W_1 \dots W_n} z_0} \left| j(z) \frac{-1}{z-x} \right| |dz| \\ &\leq \frac{2MK}{R^2} n \end{aligned}$$

and thus the sequence  $\{(a_1(z_0) + \dots + a_n(z_0))/n\}_{n \geq 1}$  is bounded.

For (ii), pick any other point  $z'_0 \in \mathbb{H}$ . Since

$$\begin{aligned} (a_1(z_0) + \dots + a_n(z_0)) - (a_1(z'_0) + \dots + a_n(z'_0)) &= \left( \int_{\gamma_{W_1 \dots W_n} z'_0}^{\gamma_{W_1 \dots W_n} z_0} - \int_{z'_0}^{z_0} \right) j(z) \frac{-1}{z-x} dz \\ &= \int_{z'_0}^{z_0} j(z) \left( \frac{1}{z - \gamma_{W_1 \dots W_n}^{-1} x} - \frac{1}{z-x} \right) dz \end{aligned}$$

is bounded by Lemma 4.3 and 4.4, we have

$$\lim_{n \rightarrow \infty} \frac{a_1(z_0) + \cdots + a_n(z_0)}{n} = \lim_{n \rightarrow \infty} \frac{a_1(z'_0) + \cdots + a_n(z'_0)}{n}.$$

As for (iii), let  $x = [k_1, k_2, \dots]$ ,  $V_n := (k_{2n-1}, k_{2n})$ , and  $L_n := |W_1| + \cdots + |W_n|$ . Since  $\gamma_{W_1 \cdots W_n} = \gamma_{V_1 \cdots V_{L_n}}$ , we have

$$\frac{1}{L_n} \int_{z_0}^{\gamma_{W_1 \cdots W_n} z_0} = \frac{1}{L_n} \int_{z_0}^{\gamma_{V_1 \cdots V_{L_n}} z_0}$$

and thus the values  $\widehat{\text{val}}(x)$  and  $\widehat{1}(x)$  are independent of  $\{W_n \mid n \geq 1\}$ .

Finally, we prove (iv). Since  $\text{SL}_2(\mathbb{Z})$ -equivalent real numbers have the same continued fraction expansions except for the first few terms, it suffices to show  $\widetilde{f}(x) = \widetilde{f}(\gamma^{-1}x)$  in the case when  $x = [W_1 W_2 \dots]$ ,  $\gamma = \gamma_{W_1} \cdots \gamma_{W_k}$ . This follows from the equality

$$\int_{\gamma_{W_1} \cdots \gamma_{W_m} z_0}^{\gamma_{W_1} \cdots \gamma_{W_{m+n+k}} z_0} j(z) \frac{-1}{z-x} dz = \int_{\gamma_{W_{k+1}} \cdots \gamma_{W_{m+n+k}} z_0}^{\gamma_{W_{k+1}} \cdots \gamma_{W_{m+n+k}} z_0} j(z) \frac{1}{z-\gamma^{-1}x}$$

for  $m > k$ . □

In Theorem 4.6 (i), we consider the differential form  $-dz/(z-x) = \eta_{\infty, x}$ . In fact, we can replace it by  $\eta_{x', x}$  for any point  $x' \in \mathbb{P}^1(\mathbb{R}) \setminus \{x\}$ . To prove it, we prepare the following lemma.

**Lemma 4.7.** *In the setting of Theorem 4.6, let  $\{x'_n\}_{n=1}^\infty \subset \mathbb{R}$  be a bounded sequence such that  $x$  is not an accumulation point. Then the sequence*

$$\int_{z_0}^{\gamma_{W_1 \cdots W_n} z_0} j(z) \frac{dz}{z-x'_n}$$

*is bounded.*

*Proof.* Let

$$K := \max \{ |j(z)| \mid |\text{Re}(z)| \leq 1/2, |z| \leq 1, \text{Im}(z) \leq h(x) \}.$$

Then

$$\begin{aligned} \left| \int_{z_0}^{\gamma_{W_1 \cdots W_n} z_0} j(z) \frac{dz}{z-x'_n} \right| &\leq K \int_{S_{z_0, \gamma_{W_1 \cdots W_n} z_0}} \frac{|dz|}{|z-x'_n|} \\ &\leq K \int_{S_{z_0, x}} \frac{|dz|}{|z-x'_n|}. \end{aligned}$$

□

By Lemma 4.7, we can reformulate the definition of  $\widehat{\text{val}}(x)$  in Theorem 4.6 (i) as follows.

**Proposition 4.8.** *In the setting of Theorem 4.6, let  $\{x'_n\}_{n=1}^\infty \subset \mathbb{R}$  be a bounded sequence such that  $x$  is not an accumulation point. Then*

$$\widehat{\text{val}}(x) = \lim_{n \rightarrow \infty} \frac{1}{|W_1| + \cdots + |W_n|} \int_{z_0}^{\gamma_{W_1 \cdots W_n} z_0} j \eta_{x'_n, x}$$

*holds.*

Finally, we prove Theorem 2.1 (iii).

**Theorem 4.9** (Theorem 2.1 (iii)). *In the setting of Theorem 4.6, for any point  $x' \in \mathbb{P}^1(\mathbb{R}) \setminus \{x\}$  and a sequence  $\{x_n\}_{n=1}^\infty$  of real numbers such that  $z_n := \gamma_{W_1 \cdots W_n}(x_n + \sqrt{-1}) \in S_{\infty, x}$  and  $z_n \rightarrow x$ , we have*

$$\begin{aligned} \widehat{\text{val}}(x) &= \lim_{n \rightarrow \infty} \frac{1}{|W_1| + \cdots + |W_n|} \int_{z_0}^{z_n} j \eta_{x', x}, \\ \widehat{1}(x) &= \lim_{n \rightarrow \infty} \frac{1}{|W_1| + \cdots + |W_n|} \int_{z_0}^{z_n} \eta_{x', x}. \end{aligned}$$

*Proof.* We prove only the first equality. Let  $\gamma_n := \gamma_{W_1 \dots W_n}$ . By Proposition 4.8, we have

$$\widehat{\text{val}}(x) = \lim_{n \rightarrow \infty} \frac{1}{|W_1| + \dots + |W_n|} \int_{\sqrt{-1}}^{\gamma_n \sqrt{-1}} j\eta_{x',x}.$$

Take a point  $z_0 \in S_{x',x}$  such that  $\pi(z_0) \in \pi(\{z \in \mathbb{H} \mid 1 \leq \text{Im}(z) \leq h(x)\})$ . Then we have

$$\widehat{\text{val}}(x) = \lim_{n \rightarrow \infty} \frac{1}{|W_1| + \dots + |W_n|} \int_{z_0}^{\gamma_n \sqrt{-1}} j\eta_{x',x}.$$

Since  $x_n + \sqrt{-1} \in \gamma_n^{-1}(S_{x',x})$ , we have  $|x_n| \leq \max\{|\gamma_n^{-1}x|, |\gamma_n^{-1}x'|\}$ . Thus,  $|x_n|$  is bounded by Lemma 4.3. Since

$$|d(z_n, z_0) - d(\gamma_n \sqrt{-1}, z_0)| \leq d(z_n, \gamma_n \sqrt{-1}) \leq d(x_n + \sqrt{-1}, \sqrt{-1}) = 1 + \frac{x_n^2}{2}$$

by the triangle inequality and [Bea83, Theorem 7.2.1],  $d(z_n, z_0) - d(\gamma_n \sqrt{-1}, z_0)$  is bounded. Let

$$K := \max\{|j(z)| \mid |\text{Re}(z)| \leq 1/2, |z| \leq 1, \text{Im}(z) \leq h(x)\}.$$

Then

$$\begin{aligned} \left( \int_{z_0}^{z_n} - \int_{z_0}^{\gamma_n \sqrt{-1}} \right) j\eta_{x',x} &= \int_{\sqrt{-1}}^{x_n + \sqrt{-1}} j\eta_{\gamma_n^{-1}x', \gamma_n^{-1}x} \\ &\leq K \int_{\sqrt{-1}}^{x_n + \sqrt{-1}} \frac{|\gamma_n^{-1}x - \gamma_n^{-1}x'|}{(\min\{\text{Im}(x_n + \sqrt{-1}), \text{Im}(\sqrt{-1})\})^2} dz \\ &\leq K |\gamma_n^{-1}x - \gamma_n^{-1}x'| |x_n| \end{aligned}$$

is bounded by Lemma 4.3. Thus, we obtain

$$\widehat{\text{val}}(x) = \lim_{n \rightarrow \infty} \frac{1}{|W_1| + \dots + |W_n|} \int_{z_0}^{z_n} j\eta_{x',x}.$$

□

## 5. ELEMENTARY UNITS FOR BADLY APPROXIMABLE NUMBERS

In this section, we consider an analog of elementary units for badly approximable numbers and prove Theorem 2.1 (iv) and (v).

**Theorem 5.1** (Theorem 2.1 (iv) and (v)). *Let  $x$  be a badly approximable number and  $W_1, W_2, \dots$  be an infinite sequence of even words such that  $x = [W_1 W_2 \dots]$  and the set  $\{W_1, W_2, \dots\}$  is finite. Let  $c_n \in \mathbb{Z}_{>0}$  be the denominator of a rational number  $[W_1 \dots W_n]$  and*

$$L := \lim_{n \rightarrow \infty} \frac{|W_1| + \dots + |W_n|}{n}.$$

*Then the limit*

$$\varepsilon_x := \lim_{n \rightarrow \infty} c_n^{1/Ln}$$

*converges if and only if  $\widehat{1}(x)$  converges. In that case, it holds  $\widehat{1}(x) = 2 \log \varepsilon_x$ .*

Here we call  $\varepsilon_x$  the elementary units for each badly approximable number  $x$ . Before giving a proof, we state a remark.

**Remark 5.2.** In the situation in Theorem 5.1, we have  $\varepsilon(W_1, W_2, \dots) \geq (3 + \sqrt{5})/2$ .

*Proof of Theorem 5.1.* Let

$$\gamma_n := \gamma_{W_1 \dots W_n} = \begin{pmatrix} * & * \\ c_n & d_n \end{pmatrix}.$$

Here  $c_n$  is a denominator of the rational number  $\gamma_n(\infty) = [W_1 \dots W_n]$ . Pick any point  $z_0 \in \mathbb{H}$ . Then we have

$$\widetilde{1}(x) = \lim_{n \rightarrow \infty} \frac{1}{n} \int_{z_0}^{\gamma_n z_0} \frac{-1}{z-x} dz = \lim_{n \rightarrow \infty} \frac{1}{n} \left[ -\log(z-x) \right]_{z_0}^{\gamma_n z_0}.$$

Since  $\gamma_n z_0$  converges to  $x$  as  $n \rightarrow \infty$  and the argument of  $\log$  is bounded, this limit value is equal to

$$\lim_{n \rightarrow \infty} \frac{-1}{n} \log |\gamma_n z_0 - x| = \lim_{n \rightarrow \infty} \frac{-1}{n} \log \left| \frac{z_0 - \gamma_n^{-1} x}{(c_n z_0 + d_n)(c_n \gamma_n^{-1} x + d_n)} \right|.$$

This is equal to

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log |(c_n z_0 + d_n)(c_n \gamma_n^{-1} x + d_n)|$$

since  $x$  is badly approximable and  $\{\gamma_n^{-1} x \mid n \in \mathbb{Z}_{>0}\}$  is bounded by Lemma 4.3. Since  $\gamma_n^{-1} \infty = -d_n/c_n$  is bounded by Lemma 4.3,

$$\left\{ \frac{c_n z + d_n}{c_n \gamma_n^{-1} x + d_n} \right\}_{n \in \mathbb{Z}_{>0}}$$

is bounded for any point  $z \in \mathbb{H} \cup \mathbb{R}$ . Thus, we obtain

$$\widehat{1}(x) = \lim_{n \rightarrow \infty} \frac{2}{n} \log |c_n z + d_n|.$$

By substituting  $z = 0$ , we have

$$\widehat{1}(x) = \lim_{n \rightarrow \infty} \frac{2}{n} \log d_n.$$

Since  $d_n/c_n$  is bounded, we obtain

$$\widehat{1}(x) = \lim_{n \rightarrow \infty} \frac{2}{n} \log c_n = 2 \log \varepsilon_x.$$

□

**Example 5.3.** For the case when  $x = \phi := (1 + \sqrt{5})/2 = [1, 1, \dots]$ , since

$$\underbrace{[1, \dots, 1]}_{2n} = \frac{F_{2n+1}}{F_{2n}}$$

where  $F_n$  denotes the  $n$ -th Fibonacci number, we have

$$\varepsilon_\phi = \lim_{n \rightarrow \infty} \sqrt[n]{F_{2n}} = \lim_{n \rightarrow \infty} \sqrt[n]{\frac{1}{\sqrt{5}} (\phi^{2n} - (-\phi)^{2n})} = \phi^2 = \frac{3 + \sqrt{5}}{2}.$$

## 6. AN EXPLICIT COMPUTATION OF VALUES OF EXTENDED val FUNCTION

In this section, we prove Theorem 1.1. The most important point of the proof below is the following lemma which is based on the proof of [BI19, Theorem 4.3] and is a generalization of [Mur20, Lemma 3.3].

**Lemma 6.1** (Repetition frequency estimation). *Let  $\{a_{1,n}\}_{n=1}^\infty, \dots, \{a_{k,n}\}_{n=1}^\infty$  be sequences in  $\mathbb{Z}_{\geq 0}$  such that  $a_{i,n} \rightarrow \infty$  and  $2^{-n} a_{i,n} \rightarrow 0$  as  $n \rightarrow \infty$ . Let  $V$  be a non-empty even word and put  $v := [V]$ . Let  $V_n$  and  $W_n$  be non-empty even words such that  $V_n$  and  $V_{n+1}$ ,  $W_n$  and  $W_{n+1}$  coincide in the first  $n$  terms respectively. Let  $\{x_n\}_{n=1}^\infty$  and  $\{x'_n\}_{n=1}^\infty$  be sequences of real numbers whose continued fraction expansions are*

$$x_n = [V^{a_n} V_n \dots], \quad -x'_n = [0, W_n \dots]$$

respectively. Then

$$I_n := \int_{z_0}^{\gamma_V^{a_n} z_0} j\eta_{x'_n, x_n} - a_n N(V) \widehat{\text{val}}(v)$$

is a Cauchy sequence. In particular, we have

$$\int_{z_0}^{\gamma_V^{a_n} z_0} j\eta_{x'_n, x_n} = a_n N(V) \widehat{\text{val}}(v) + O(1) \quad \text{as } n \rightarrow \infty.$$

*Proof.* Take positive integers  $m < n$  with  $a_m < a_n$ . We have

$$I_n - I_m = \int_{z_0}^{\gamma_V z_0} j \left( \sum_{0 \leq i < a_n} \eta_{\gamma_V^{i-a_n} x'_n, \gamma_V^{i-a_n} x_n} - \sum_{0 \leq i < a_m} \eta_{\gamma_V^{i-a_m} x'_m, \gamma_V^{i-a_m} x_m} + (a_n - a_m) \eta_{v', v} \right).$$

Let

$$\begin{aligned} S_1(m, n; z) &:= \sum_{0 \leq i < a_m} \left( \frac{1}{z - \gamma_V^{i-a_n} x_n} - \frac{1}{z - \gamma_V^{i-a_m} x_m} \right), \\ S'_1(m, n; z) &:= \sum_{0 \leq i < a_m} \left( \frac{1}{z - \gamma_V^{i-a_n} x'_n} - \frac{1}{z - \gamma_V^{i-a_m} x'_m} \right), \\ S_2(m, n; z) &:= \sum_{a_m \leq i < a_n} \left( \frac{1}{z - \gamma_V^{i-a_n} x_n} - \frac{1}{z - v} \right), \\ S'_2(m, n; z) &:= \sum_{a_m \leq i < a_n} \left( \frac{1}{z - \gamma_V^{i-a_n} x'_n} - \frac{1}{z - v'} \right). \end{aligned}$$

Then we can write

$$I_n - I_m = \int_{z_0}^{\gamma_V z_0} j (-S_1(m, n; z) + S'_1(m, n; z) + S_2(m, n; z) - S'_2(m, n; z)) dz.$$

For each  $0 \leq i < a_m$ , since

$$\begin{aligned} \gamma_V^{i-a_n} x_n &= [V^i V_n \cdots], & \gamma_V^{i-a_m} x_m &= [V^i V_m \cdots], \\ \gamma_V^{i-a_n} x'_n &= -[0, V^i W_n \cdots], & \gamma_V^{i-a_m} x'_m &= -[0, V^i W_m \cdots], \end{aligned}$$

we have

$$\left| \gamma_V^{i-a_n} x_n - \gamma_V^{i-a_m} x_m \right| \leq 2^{2-m}, \quad \left| \gamma_V^{i-a_n} x'_n - \gamma_V^{i-a_m} x'_m \right| \leq 2^{2-m}.$$

For each  $a_m \leq i < a_n$ , since

$$\gamma_V^{i-a_n} x_n = [V^i V_n \cdots], \quad \gamma_V^{i-a_n} x'_n = -[0, V^i W_n \cdots],$$

we have

$$\left| \gamma_V^{i-a_n} x_n - v \right| \leq 2^{2-a_m}, \quad \left| \gamma_V^{i-a_n} x'_n - v' \right| \leq 2^{2-a_m}.$$

By Lemma 4.4, we obtain

$$|I_n - I_m| = \sum_{0 \leq i < a_m} O(2^{-m}) + \sum_{a_m \leq i < a_n} O(2^{-a_m}) = O(a_m 2^{-m}) + O((a_n - a_m) 2^{-a_m}).$$

Since this converges to 0 as  $m \rightarrow \infty$  by the assumption,  $\{I_n\}$  is a Cauchy sequence.  $\square$

*Proof of Theorem 1.1.* Keep the notation in Theorem 1.1 and let

$$L := \lim_{n \rightarrow \infty} \frac{|U_1| + \cdots + |U_n|}{A_n}.$$

It suffices to show that

$$L \widehat{\text{val}}(x) = a_1 \widehat{\text{val}}(W_1) + \cdots + a_k \widehat{\text{val}}(W_k).$$

Take any real number  $x' < -1$  and for positive integers  $n$  and  $1 \leq i \leq k$ , let

$$\begin{aligned} x_n &:= \gamma_{U_1 \cdots U_{n-1}}^{-1} x = [U_n U_{n+1} \cdots], \\ x'_n &:= \gamma_{U_1 \cdots U_{n-1}}^{-1} x' = -[0, U'_1 \cdots U'_n, \delta_0 x'], \\ x_{i,n} &:= \gamma_{V_1 W_1^{a_1, n} V_2 W_2^{a_2, n} \cdots V_{i-1} W_{i-1}^{a_{i-1}, n}}^{-1} x_n = [V_i W_i^{a_{i,n}} \cdots V_k W_k^{a_{k,n}} U_{n+1} U_{n+2} \cdots], \\ x'_{i,n} &:= \gamma_{V_1 W_1^{a_1, n} V_2 W_2^{a_2, n} \cdots V_{i-1} W_{i-1}^{a_{i-1}, n}}^{-1} x'_n = -[0, W'^{a_{i-1}, n} V'_{i-1} \cdots W'^{a_{1,n}} V'_1 U'_{n-1} \cdots U'_1, \delta_0 x']. \end{aligned}$$

Since  $x' < -1$ , the far right-hand sides in the second and fourth rows are continued fraction expansions. Then we have

$$\begin{aligned} L\widehat{\text{val}}(x) &= \lim_{n \rightarrow \infty} \frac{1}{A_n} \sum_{1 \leq m \leq n} \int_{\gamma_{U_1 \cdots U_{m-1}}^{z_0}}^{\gamma_{U_1 \cdots U_m}^{z_0}} j\eta_{x', x} \\ &= \lim_{n \rightarrow \infty} \frac{1}{A_n} \sum_{1 \leq m \leq n} \int_{z_0}^{\gamma_{U_m}^{z_0}} j\eta_{x'_m, x_m} \\ &= \lim_{n \rightarrow \infty} \frac{1}{A_n} \sum_{1 \leq m \leq n} \sum_{1 \leq i \leq k} \left( \int_{z_0}^{\gamma_{V_i}^{z_0}} j\eta_{x'_{i,m}, x_{i,m}} + \int_{z_0}^{W_i^{a_{i,m}} z_0} j\eta_{\gamma_{V_i}^{-1} x'_{i,m}, \gamma_{V_i}^{-1} x_{i,m}} \right). \end{aligned}$$

Since we have

$$|x_{i,m} - \gamma_{V_i} w_i| \leq 2^{2-a_{i,m}}, \quad |x'_{i,m} - \gamma_{w'_{i-1}}| \leq 2^{2-a_{i-1,m}}$$

by continued fraction expansion, it follows from Lemma 4.4 that

$$\int_{z_0}^{\gamma_{V_i}^{z_0}} j \left( \eta_{x'_{i,m}, x_{i,m}} - \eta_{w'_{i-1}, \gamma_{V_i} w_i} \right) = O(2^{-a_{i,m}} + 2^{-a_{i-1,m}}).$$

Since

$$\begin{aligned} \gamma_{V_i}^{-1} x_{i,m} &= [W_i^{a_{i,m}} \cdots V_k W_k^{a_{k,m}} U_{n+1} U_{n+2} \cdots], \\ \gamma_{V_i}^{-1} x'_{i,m} &= -[0, V_i' W_i'^{a_{i-1,m}} V_{i-1}' \cdots W_1'^{a_{1,m}} V_1' U_{m-1}' \cdots U_1', \delta_0 x'], \end{aligned}$$

we have

$$\int_{z_0}^{W_i^{a_{i,m}} z_0} j\eta_{\gamma_{V_i}^{-1} x'_{i,m}, \gamma_{V_i}^{-1} x_{i,m}} = a_{i,m} N(W_i) \widehat{\text{val}}(w_i) + O(1)$$

by Lemma 6.1. Thus, we obtain

$$\begin{aligned} L\widehat{\text{val}}(x) &= \lim_{n \rightarrow \infty} \frac{1}{A_n} \sum_{1 \leq m \leq n} \sum_{1 \leq i \leq k} a_{i,m} N(W_i) \widehat{\text{val}}(w_i) \\ &= \sum_{1 \leq i \leq k} a_i N(W_i) \widehat{\text{val}}(w_i). \end{aligned}$$

□

## 7. THE VALUE OF val FUNCTION AT THE THUE-MORSE WORD

In this section, we calculate the value of val function at the Thue-Morse word and prove Theorem 1.2.

We fix two even words  $V$  and  $W$ . Let  $h: \{V, W\}^* \rightarrow \{V, W\}^*$  be a monoid homomorphism such that  $h(V) := VW$ ,  $h(W) := WV$  and  $V_h := \lim_{n \rightarrow \infty} h^n(V)$  be the Thue-Morse word. For a word  $U = U_1 \cdots U_n \in \{V, W\}^*$  with  $U_1, \dots, U_n \in \{V, W\}$ , let  $U' := U_n \cdots U_1$ .

To begin with, we give a simple representation of the Thue-Morse word.

**Lemma 7.1.** *Let  $\tau: \{V, W\}^* \rightarrow \{V, W\}^*$  be a monoid homomorphism such that  $\tau(V) := W$  and  $\tau(W) := V$ . Then the followings hold.*

- (i) *For any positive integer  $n$ , we have  $h^n(V) = h^{n-1}(V)\tau h^{n-1}(W)$ .*
- (ii)  *$h^{2n}(V)' = h^{2n}(V)$ .*

*Proof.* The first claim (i) follows from the facts that  $h \circ \tau = \tau \circ h$  and  $h^2(U) = h(U)\tau h(U)$  for a word  $U \in \{V, W\}^*$  such that  $h(U) = U\tau(U)$ . The last claim (ii) follows from the fact that  $h^2(U)' = h^2(U)$  for a word  $U \in \{V, W\}^*$  such that  $h(U) = U\tau(U)$  and  $U' = U$ . □

*Proof of Theorem 1.2.* Let  $V_n := h^n(V)$ . Since  $[V_n]' = -[0, V_n]$  by Lemma 7.1 (ii), we have

$$\widehat{\text{val}}([V_h]) = \lim_{n \rightarrow \infty} \frac{1}{2^{2n}} \int_{z_0}^{\gamma_{V_n}^{z_0}} j\eta_{-[0, V_h], V_h}, \quad \widehat{\text{val}}([V_n]) = \int_{z_0}^{\gamma_{V_n}^{z_0}} j\eta_{-[0, V_n], V_n}.$$

Here the first  $2^{2n}$  terms of  $V_h$  is  $V_n$  by Lemma 7.1 (i). Thus, we obtain the claim. □



## REFERENCES

- [Bea83] A. F. Beardon. *The geometry of discrete groups*, volume 91 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1983.
- [BI19] P. Bengoechea and Ö. Imamoglu. Cycle integrals of modular functions, Markov geodesics and a conjecture of Kaneko. *Algebra Number Theory*, 13(4):943—962, 2019.
- [CK97] C. Choffrut and J. Karhumäki. Combinatorics of words. In *Handbook of formal languages, Vol. 1*, pages 329–438. Springer, Berlin, 1997.
- [Dal11] F. Dal’Bo. *Geodesic and horocyclic trajectories*. Universitext. Springer-Verlag London, Ltd., London; EDP Sciences, Les Ulis, 2011. Translated from the 2007 French original.
- [DIT11] W. Duke, Ö. Imamoglu, and Á. Tóth. Cycle integrals of the  $j$ -function and mock modular forms. *Ann. of Math. (2)*, 173(2):947–981, 2011.
- [Kan09] M. Kaneko. Observations on the ‘values’ of the elliptic modular function  $j(\tau)$  at real quadratics. *Kyushu Journal of Mathematics*, 63(2):353–364, 2009.
- [Mat20] T. Matsusaka. A hyperbolic analogue of the rademacher symbol. 2020. arXiv:2003.12354.
- [Mur20] Y. Murakami. A continuity of cycle integrals of modular functions. *Ramanujan J.*, 2020.
- [Shi72] G. Shimura. Class fields over real quadratic fields and Hecke operators. *Ann. of Math. (2)*, 95:130–190, 1972.
- [Shi78] Takuro Shintani. On certain ray class invariants of real quadratic fields. *J. Math. Soc. Japan*, 30(1):139–167, 1978.
- [Zag02] D. Zagier. Traces of singular moduli. In *Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998)*, volume 3 of *Int. Press Lect. Ser.*, pages 211–244. Int. Press, Somerville, MA, 2002.

MATHEMATICAL INST. TOHOKU UNIV., 6-3, AOBA, ARAMAKI, AOBA-KU, SENDAI 980-8578, JAPAN  
Email address: yuya.murakami.s8@dc.tohoku.ac.jp