

DOUBLE SIDED TORUS ACTIONS AND COMPLEX GEOMETRY ON $SU(3)$

HIROAKI ISHIDA AND HISASHI KASUYA

ABSTRACT. We construct explicit complex structures and transversely Kähler holomorphic foliations on $SU(3)$ corresponding to variations of real quadratic equations on a complex quadric in \mathbb{C}^6 as generalizations of left-invariant complex structures on $SU(3)$ and an invariant Kähler structure on the flag variety $SU(3)/T$. Consequently, we obtain orbifold variants of the flag variety $SU(3)/T$ as quotients of double sided torus actions.

1. INTRODUCTION

Let G be a compact connected semi-simple Lie group and T a maximal torus. The homogeneous space G/T is called flag variety and it admits a G -invariant Kähler structure. Suppose G is even dimensional. Then, in [15] Samelson proved that there exist left-invariant complex structures on G . Moreover, for G equipped with a left-invariant complex structure, the principal bundle $T \rightarrow G \rightarrow G/T$ is holomorphic. We are interested in constructing variants of these geometric objects.

We consider the $T \times T$ -action (double sided action) on G such that for $t = (t_1, t_2) \in T \times T$, $g \in G$, the action is defined by $t \cdot g = t_1 g t_2^{-1}$. Now, we have $(S^1)^r \cong T$ where r is the rank of G . We consider two-sided variations of the right T -action. Let $\rho_L, \rho_R: (S^1)^r \rightarrow T$ be smooth homomorphisms. Consider the $(S^1)^r$ -action on G associated with $\rho_L \times \rho_R: (S^1)^r \rightarrow T \times T$ and the double sided action. Then, in general the orbit space $G/(\rho_L, \rho_R)((S^1)^r)$ may not be a smooth manifold. We recall the notion for the studying non-smooth $G/(\rho_L, \rho_R)((S^1)^r)$. Let M be a complex manifold with the complex structure J . A holomorphic foliation F on M is called transversely Kähler if there exists a closed $(1, 1)$ -form ω such that $\omega(X, JX) \geq 0$ for any $X \in TM$ and equality holds if and only if $X \in TF$. Such a $(1, 1)$ -form ω is called a transverse Kähler form. Transversely Kähler holomorphic foliations on complex manifolds can be seen as generalizations of holomorphic fiber bundles over Kähler manifolds. As an extension of the above argument, we desire a statement of the following type: “under certain conditions on $\rho_L, \rho_R: (S^1)^r \rightarrow T$, we can construct a (not necessarily left-invariant) complex structure on G so that $(\rho_L, \rho_R)((S^1)^r)$ -orbits form a transversely Kähler holomorphic foliation”.

In this paper, we focus on the case when $G = SU(3)$. We study the double sided torus action on $SU(3)$ and its quotient space from the point of view of moment maps. Let T be the maximal compact torus $\{g = \text{diag}(g_1, g_2, g_3) \mid g_1, g_2, g_3 \in S^1, g_1 g_2 g_3 = 1\}$ in $SU(3)$, where $\text{diag}(g_1, g_2, g_3)$ denote the diagonal matrix of size 3 whose (j, j) -entry is g_j . For $t = (t_1, t_2) \in (S^1)^2$ and $w = (w_1, w_2) \in \mathbb{Z}^2$, by t^w we mean $t_1^{w_1} t_2^{w_2}$. For s_1, \dots, s_k in \mathbb{R}^2 , by $\text{pos}(s_1, \dots, s_k)$ we mean the cone spanned by s_1, \dots, s_k . Namely, $\text{pos}(s_1, \dots, s_k) = \{\sum_j \lambda_j s_j \mid \lambda_j \geq 0\}$. The following is the main theorem of this paper:

Theorem 1.1. *Let $\rho_L, \rho_R: (S^1)^2 \rightarrow T$ be smooth homomorphisms given by*

$$\begin{aligned}\rho_L(t) &= \text{diag}(t^{w_1^L}, t^{w_2^L}, t^{w_3^L}), \\ \rho_R(t) &= \text{diag}(t^{w_1^R}, t^{w_2^R}, t^{w_3^R}),\end{aligned}$$

Date: October 19, 2021.

2020 Mathematics Subject Classification. 22C05, 32M25, 53D20, 57S25.

here $w_j^L, w_j^R \in \mathbb{Z}^2$. Put

$$A_j := w_j^L - w_1^R, \quad B_j := -w_j^L + w_3^R, \quad C := -w_1^R + w_3^R.$$

Assume that A_j, B_j for $j = 1, 2, 3$ and C satisfy

- (\star) $C \notin \text{pos}(A_i, A_j), \text{pos}(B_i, B_j)$ for all $i, j \in \{1, 2, 3\}$ and $C \in \text{pos}(A_i, B_j)$ for all $i, j \in \{1, 2, 3\}$.

Then the followings hold:

- (1) There exists a $T \times T$ -invariant complex structure on $SU(3)$ such that $(\rho_L, \rho_R)((S^1)^2)$ -orbits form a transversely Kähler holomorphic foliation.
- (2) There exists a $T \times T / (\rho_L, \rho_R)((S^1)^2)$ -invariant Kähler orbifold structure on $SU(3) / (\rho_L, \rho_R)((S^1)^2)$.

There is an advantage to our case $G = SU(3)$. Our construction is really explicit and may not be simply generalized to general compact Lie groups. We know that $SU(3)$ is presented by explicit real quadric equations on a complex quadric in \mathbb{C}^6 (see Section 3). The main idea is to vary such real quadric equations corresponding to (ρ_L, ρ_R) . A complex structure is defined by an explicit holomorphic foliation (Haefliger's trick as in [8]) and a transversely Kähler structure comes from the standard Kähler structure on \mathbb{C}^6 . These constructions are closely related to LVM manifolds [10, 11]. In [12], it is shown that certain non-Kähler manifolds (now known as LVM manifolds) admit structures of principal holomorphic Seifert fibering over toric varieties under some condition. Theorem 1.1 is a variant of this result by replacing toric varieties to "twisted" flag varieties.

Under the assumption (\star) in Theorem 1.1, we can check that the action of $(S^1)^2$ on $SU(3)$ given by (ρ_L, ρ_R) on $SU(3)$ is free if and only if ρ_L is trivial and ρ_R is an isomorphism (Proposition 5.3). That is to say, we obtain essentially new objects only if the action of $(S^1)^2$ is not free. On the other hand, it has been shown that there exists the 2 dimensional torus S_{12} in $SU(3) \times SU(3)$ acting freely on $SU(3)$ not contained in $1 \times SU(3)$ such that the quotient space $SU(3)/S_{12}$ admits a structure of Kähler "manifold" as well as the flag manifold, in [3] and [4] implicitly and [5] explicitly. However, it is also shown in [5, Theorem 3.3] that there is no Kähler structure on $SU(3)/S_{12}$ which is invariant under the action of 2 dimensional torus $T \times T/S_{12}$, not like the flag manifold. Thus, such example seem to be essentially different from our objects we obtain by Theorem 1.1.

We notice that if ρ_L is non-trivial, then a complex structure in Theorem 1.1 may not be left-invariant. Non-left-invariant complex structures on compact Lie groups are studied in [9] and [7].

This paper is organized as follows. In Section 2, we develop a key tool to construct a complex manifold equipped with a transverse Kähler structure via moment maps. In Section 3, we construct an embedding of $SU(3)$ into a complex quadric and see that the image coincides with a level set of a moment map for some torus action. In Section 4, we study the intersections of real quadrics in a complex quadric. This is an analogue of intersections of special real quadrics in \mathbb{C}^n developed in [12] and we use similar arguments. Using results obtained in previous sections, we show Theorem 1.1. In Section 5, we study the freeness of the action of $(S^1)^2$ on $SU(3)$ under the condition (\star). As a result, we see that the quotient $SU(3) / (\rho_L, \rho_R)((S^1)^2)$ has an orbifold singularity except $SU(3) / (\rho_L, \rho_R)((S^1)^2)$ is the ordinal flag manifold. We also give a nontrivial concrete example of (ρ_L, ρ_R) such that $SU(3) / (\rho_L, \rho_R)((S^1)^2)$ has a Kähler orbifold structure. In Section 6, we compute the cohomology associated with each structure in Theorem 1.1.

2. TRANSVERSE KÄHLER STRUCTURES ON LEVEL SETS OF MOMENT MAPS

In this section, we construct a transverse Kähler structure on a real submanifold in a Kähler manifold with a commuting Hamiltonian. For the notions about equivariant symplectic geometry, we mention the book [1] as an excellent reference.

Let (M, ω, J) be a Kähler manifold, that is, M is a smooth manifold, ω is a symplectic form on M and J is an integrable complex structure on M compatible with ω . Let $f_1, \dots, f_n, g_1, \dots, g_n$ be

smooth functions on M and $X_1, \dots, X_n, Y_1, \dots, Y_n$ the Hamiltonian vector fields of $f_1, \dots, f_n, g_1, \dots, g_n$, respectively. We assume that $X_1, \dots, X_n, Y_1, \dots, Y_n$ are Killing and commute with each other. Let $\alpha \in \mathbb{R}^{2n}$ be a regular value of $\Phi := (f_1, \dots, f_n, g_1, \dots, g_n): M \rightarrow \mathbb{R}^{2n}$ and assume that $\Phi^{-1}(\alpha) \neq \emptyset$.

First, we construct a holomorphic foliation \mathcal{F} on a neighborhood of $N := \Phi^{-1}(\alpha)$ that is transverse to N . For $i = 1, \dots, n$, we define vector fields $Z_i := X_i - JY_i$, $W_i := JX_i + Y_i$ on M . Let $p \in N$. We claim that $T_p M = T_p N \oplus \langle Z_1(p), \dots, Z_n(p), W_1(p), \dots, W_n(p) \rangle$. Let $a_1, \dots, a_n, b_1, \dots, b_n \in \mathbb{R}$. Assume that $\sum_i a_i Z_i(p) + b_i W_i(p) \in T_p N$. Then

$$\begin{aligned} 0 &= \omega\left(\sum_i b_i X_i(p) - a_i Y_i(p), \sum_i a_i Z_i(p) + b_i W_i(p)\right) \\ &= \omega\left(\sum_i b_i X_i(p) - a_i Y_i(p), J\left(\sum_i b_i X_i(p) - a_i Y_i(p)\right)\right) \end{aligned}$$

because $X_1, \dots, X_n, Y_1, \dots, Y_n$ commute and Hamiltonian. Since the bilinear form $\omega(-, J-)$ is positive definite, we have that $\sum_i b_i X_i(p) - a_i Y_i(p) = 0$. Since p is a regular point, we have that $X_1(p), \dots, X_n(p), Y_1(p), \dots, Y_n(p)$ are linearly independent. Therefore $a_1 = \dots = a_n = b_1 = \dots = b_n = 0$. This shows that

$$T_p N \cap \langle Z_1(p), \dots, Z_n(p), W_1(p), \dots, W_n(p) \rangle = 0$$

and $\dim \langle Z_1(p), \dots, Z_n(p), W_1(p), \dots, W_n(p) \rangle = 2n$. Thanks to the dimensions, we have

$$T_p M = T_p N \oplus \langle Z_1(p), \dots, Z_n(p), W_1(p), \dots, W_n(p) \rangle.$$

Since X_i and Y_i are Killing and preserve ω , they also preserve J . Hence $Z_1, \dots, Z_n, W_1, \dots, W_n$ commute with each other. Therefore $Z_1, \dots, Z_n, W_1, \dots, W_n$ generates a foliation $\tilde{\mathcal{F}}$ on a neighborhood of N that is transverse to N . Since $JZ_i = W_i$, the foliation $\tilde{\mathcal{F}}$ is a holomorphic foliation.

We construct the complex structure J_N on N using the foliation $\tilde{\mathcal{F}}$. Let $\{U_\alpha, \phi_\alpha: U_\alpha \rightarrow V_\alpha \times W_\alpha\}_\alpha$ be a holomorphic foliated atlas on the neighborhood of N such that, for any $z \in V_\alpha$ there uniquely exists $w \in W_\alpha$ such that $\phi_\alpha^{-1}(z, w) \in N$. Let $\phi_{\alpha 1}: U_\alpha \rightarrow V_\alpha$ be the first factor of ϕ_α . Since the atlas is holomorphic and foliated, $\phi_{\beta 1} \circ \phi_{\alpha 1}^{-1}$ is defined and holomorphic for any α and β . The family $\{U_\alpha \cap N, \phi_{\alpha 1}|_{U_\alpha \cap N}: U_\alpha \cap N \rightarrow V_\alpha\}_\alpha$ is a holomorphic atlas on N . In other words, the complex structure J_N on N is given as follows. Let $i: TN \rightarrow TM|_N/T\tilde{\mathcal{F}}_N$ be the isomorphism induced by the inclusion and $\pi: TM|_N \rightarrow TM|_N/T\tilde{\mathcal{F}}_N$ the projection. Then J_N is given by $J_N = i^{-1} \circ \pi \circ J$.

Since N is a level set of $f_1, \dots, f_n, g_1, \dots, g_n$ and $X_1, \dots, X_n, Y_1, \dots, Y_n$ commute with each other, we have that $X_1, \dots, X_n, Y_1, \dots, Y_n$ define the foliation \mathcal{F} on N . We shall see that \mathcal{F} is holomorphic. Since $J_N = i^{-1} \circ \pi \circ J$, we have $J_N X_i(p) \in JX_i(p) + T_p \tilde{\mathcal{F}}$. On the other hand, $Y_i(p) - JX_i(p) = W_i(p) \in T_p \tilde{\mathcal{F}}$. Therefore $J_N X_i(p) = Y_i(p)$. By the same argument we have that $J_N Y_i(p) = -X_i(p)$. This shows that \mathcal{F} is a holomorphic foliation on N .

Finally, we show that $\omega|_N$ is a transverse Kähler form on N with respect to \mathcal{F} . Let $T_p \mathcal{F}^\perp$ denote the orthogonal complement of $T_p \mathcal{F}$ in $T_p N$ with respect to the Riemann metric $\omega(-, J-)$. Since

$$\begin{aligned} T_p \mathcal{F}^\perp &= \{v \in T_p N \mid \omega(v, Ju) = 0 \text{ for all } u \in T_p \mathcal{F}\} \\ &= \{v \in T_p M \mid \omega(v, u) = \omega(v, Ju) = 0 \text{ for all } u \in T_p \mathcal{F}\}, \end{aligned}$$

we have that $T_p \mathcal{F}^\perp$ is invariant under the complex structure J on M . Therefore $J_N|_{T_p \mathcal{F}^\perp} = J|_{T_p \mathcal{F}^\perp}$. Let $X, Y \in T_p N$ and assume that $X = X' + X''$, $Y = Y' + Y''$, $X', Y' \in T_p \mathcal{F}^\perp$ and $X'', Y'' \in T_p \mathcal{F}$. Then

$$\omega(J_N X, J_N Y) = \omega(JX' + J_N X'', JY' + J_N Y'') = \omega(JX', JY')$$

because $J_N X'', J_N Y'' \in T_p \mathcal{F}$ and $\omega|_N(-, Z) = 0$ for any $Z \in T_p \mathcal{F}$. On the other hand,

$$\omega(X, Y) = \omega(X' + X'', Y' + Y'') = \omega(X', Y')$$

by the same reason. Thus $\omega(J_N X, J_N Y) = \omega(X, Y)$, showing that $\omega|_N$ is of type $(1, 1)$. Since

$$\omega(X, J_N X) = \omega(X' + X'', JX' + J_N X'') = \omega(X', JX')$$

and $\omega(-, J-)$ is positive definite, we have that $\omega(X, J_N X) \geq 0$ and the equality holds if and only if $X \in T_p \mathcal{F}$. This shows that $\omega|_N$ is a transverse Kähler form on N with respect to \mathcal{F} .

We shall state the conclusion in this section as a theorem for later use.

Theorem 2.1. *The level set N has the complex structure J_N such that the 2-form $\omega|_N$ is transverse Kähler form with respect to the foliation \mathcal{F} generated by commuting Hamiltonian vector fields. The foliation \mathcal{F} is holomorphic.*

We remark that we can obtain the *Kähler reduction* for even dimensional torus action as a corollary of Theorem 2.1:

Corollary 2.2. *In addition, if $X_1, \dots, X_n, Y_1, \dots, Y_n$ define an action of the torus T on M , then the quotient space N/T is a Kähler orbifold. Moreover the T -action restricted to N is free, then N/T is a Kähler manifold.*

Remark 2.3. For $A \in GL(2n, \mathbb{R})$, replacing $(X'_1, \dots, X'_n, Y'_1, \dots, Y'_n) = (X_1, \dots, X_n, Y_1, \dots, Y_n)A$, $(f'_1, \dots, f'_n, g'_1, \dots, g'_n) = (f_1, \dots, f_n, g_1, \dots, g_n)A$ and $\alpha' = \alpha A$, we obtain another complex structure J'_N on N by the same construction. Thus, in fact, we obtain the family of complex structures on N parametrized by $GL(2n, \mathbb{R})/GL(n, \mathbb{C})$. By the construction of the foliation \mathcal{F} , the complex structures on leaves of the holomorphic foliation \mathcal{F} depend on parameter but the structure of transversely Kähler C^∞ -foliation \mathcal{F} on N is independent. In particular, the Kähler orbifold structure on N/T is unique.

3. $SU(3)$ AS A LEVEL SET

In this section, we embed $SU(3)$ into a Kähler manifold as a level set of a smooth map. Let $SL(3, \mathbb{C})$ denote the complex special linear group of degree 3. Let B denote the subgroup of upper triangular matrices in $SL(3, \mathbb{C})$. Let U denote the subgroup of upper triangular matrices whose diagonal entries are 1. We will construct a Kähler form on $SL(3, \mathbb{C})/U$ explicitly. Let $V = \mathbb{C}^3$ and $\rho: SL(3, \mathbb{C}) \rightarrow GL(V)$ the natural representation. Let V^* be the dual of V and ρ^* the dual of ρ . Consider the (left) action of $SL(3, \mathbb{C})$ on $V \oplus V^*$ given by $\rho \oplus \rho^*$.

Let e_1, e_2, e_3 be the standard basis vectors of V and e_1^*, e_2^*, e_3^* the dual basis vectors. Suppose that $A = (a_{ij}) \in SL(3, \mathbb{C})$. Then, with respect to the basis e_1^*, e_2^*, e_3^* of V^* , the endmorphism $\rho^*(A)$ is represented as the cofactor matrix $\rho^*(A) = {}^t A^{-1}$.

Claim. *The subgroup of stabilizers of $SL(3, \mathbb{C})$ at $e_1 \oplus e_3^*$ coincides with U .*

Proof. Let $A = (a_{ij}) \in SL(3, \mathbb{C})$. Suppose that $(\rho \oplus \rho^*)(A)(e_1 \oplus e_3^*) = e_1 \oplus e_3^*$. Since $\rho(A)(e_1) = e_1$, we have $a_{11} = 1, a_{21} = 0$ and $a_{31} = 0$. Since $\rho^*(A)(e_3^*) = e_3^*$, we have $a_{21}a_{32} - a_{22}a_{31} = 0$, $a_{12}a_{31} - a_{11}a_{32} = 0$ and $a_{11}a_{22} - a_{12}a_{21} = 1$. It follows from $a_{11} = 1$ and $a_{12}a_{31} - a_{11}a_{32} = 0$ that $a_{32} = 0$. This together with $a_{21} = a_{31} = 0$ implies that $A \in B$. It follows from $a_{11} = 1, a_{21} = 0$ and $a_{11}a_{22} - a_{12}a_{21} = 1$ that $a_{22} = 1$. Since $a_{11}a_{22}a_{33} = 1$, we have $a_{33} = 1$. Therefore $A \in U$.

Conversely, let

$$A' = \begin{pmatrix} 1 & b_{12} & b_{13} \\ 0 & 1 & b_{23} \\ 0 & 0 & 1 \end{pmatrix} \in U.$$

Then $\rho(A')(e_1) = e_1$ is obvious. Since the representation matrix of $\rho^*(A')$ is

$$\begin{pmatrix} 1 & 0 & 0 \\ -b_{12} & 1 & 0 \\ b_{12}b_{23} - b_{13} & -b_{23} & 1 \end{pmatrix},$$

we have that $\rho^*(A')(e_3^*) = e_3^*$. Therefore elements in U stabilize $e_1 \oplus e_3^*$. \square

Thus the homogeneous space $SL(3, \mathbb{C})/U$ can be identified with the $SL(3, \mathbb{C})$ -orbit through $e_1 \oplus e_3^*$.

Claim. *The $SL(3, \mathbb{C})$ -orbit through $e_1 \oplus e_3^*$ is $\{v \oplus f \in V \oplus V^* \mid f(v) = 0, v \neq 0, f \neq 0\}$.*

Proof. Let $A \in SL(3, \mathbb{C})$. Put $v = \rho(A)(e_1)$ and $f = \rho^*(A)(e_3^*)$. Then $v \neq 0$, $f \neq 0$ and $f(v) = e_3^*(\rho(A)^{-1}\rho(A)(e_1)) = e_3^*(e_1) = 0$. Thus the $SL(3, \mathbb{C})$ -orbit through $e_1 \oplus e_3^*$ is contained in $\{v \oplus f \in V \oplus V^* \mid f(v) = 0, v \neq 0, f \neq 0\}$.

Let $v \oplus f \in V \oplus V^*$. Suppose that $f(v) = 0$, $v \neq 0$ and $f \neq 0$. Since $v \neq 0$, there exists $A_1 \in SL(3, \mathbb{C})$ such that $\rho(A_1)(e_1) = v$. Then $\rho^*(A_1)^{-1}(f)$ is a linear combination of e_2^* and e_3^* because $f(v) = 0$ and $\rho^*(A_1)^{-1}(f)(e_1) = f(\rho(A_1)(e_1))$. Suppose that $\rho^*(A_1)^{-1}(f) = ae_2^* + be_3^*$ for some $a, b \in \mathbb{C}$. Since $f \neq 0$, either a or b is nonzero. Put $\|f\| := \sqrt{|a|^2 + |b|^2}$ and

$$A_2 := \begin{pmatrix} 1 & 0 & 0 \\ 0 & \|f\|^{-2}\bar{b} & a \\ 0 & -\|f\|^{-2}\bar{a} & b \end{pmatrix}$$

Then $A_2 \in SL(3, \mathbb{C})$ and $\rho^*(A_2)^{-1}(e_3) = ae_2^* + be_3^*$. Thus $f = \rho^*(A_1A_2^{-1})(e_3^*)$.

Moreover, $\rho(A_1A_2^{-1})(e_1) = \rho(A_1)(e_1) = v$. Therefore we have that $v \oplus f = \rho(A_2A_1^{-1})(e_1) \oplus \rho^*(A_2A_1^{-1})(e_3^*)$, showing that $v \oplus f$ sits in the $SL(3, \mathbb{C})$ -orbit through $e_1 \oplus e_3^*$. \square

Let M be the quasi-affine variety in $\mathbb{C}^6 = \mathbb{C}^3 \times \mathbb{C}^3$ defined as

$$M := \{(z, w) \in \mathbb{C}^3 \times \mathbb{C}^3 \mid z \neq 0, w \neq 0, \sum_{j=1}^3 z_j w_j = 0\}$$

where $z = (z_1, z_2, z_3)$ and $w = (w_1, w_2, w_3)$. As we saw above, $SL(3, \mathbb{C})/U$ is isomorphic to M as varieties via the isomorphism given by $SL(3, \mathbb{C})/U \ni [A] \mapsto (z, w) \in M$, where

$$(3.1) \quad \begin{pmatrix} z_1 \\ z_2 \\ z_3 \\ w_1 \\ w_2 \\ w_3 \end{pmatrix} = \begin{pmatrix} A & O \\ O & {}^t A^{-1} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}.$$

Since M is quasi-affine, we have a Kähler form on M which is obtained by restricting the standard Kähler form on \mathbb{C}^6 . Moreover, $SL(3, \mathbb{C})/U$ admits $T \times T$ -action which we will introduce below. We will describe the inherited $T \times T$ -action on M and its moment map.

For $g, h \in T$ and $A \in SL(3, \mathbb{C})$, we define $(g, h) \cdot A := gAh^{-1}$. This is an action of $T \times T$ on $SL(3, \mathbb{C})$ and it descends to an action on $SL(3, \mathbb{C})/U$ because $hUh^{-1} = U$ for any $h \in T$. Suppose

that $g = \text{diag}(g_1, g_2, g_3)$ and $h = \text{diag}(h_1, h_2, h_3)$. Then

$$(3.2) \quad \begin{pmatrix} gAh^{-1} & O \\ O & {}^t(gAh^{-1})^{-1} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} g & O \\ O & g^{-1} \end{pmatrix} \begin{pmatrix} A & O \\ O & {}^tA^{-1} \end{pmatrix} \begin{pmatrix} h_1^{-1} \\ 0 \\ 0 \\ 0 \\ 0 \\ h_3 \end{pmatrix} \\ = \begin{pmatrix} \text{diag}(g_1h_1^{-1}, g_2h_1^{-1}, g_3h_1^{-1}) & O \\ O & \text{diag}(g_1^{-1}h_3, g_2^{-1}h_3, g_3^{-1}h_3) \end{pmatrix} \begin{pmatrix} A & O \\ O & {}^tA^{-1} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}.$$

It follows from (3.1) and (3.2) that the $T \times T$ action on M is obtained by restricting the standard $(S^1)^6$ -action on \mathbb{C}^6 . Thanks to the weights in (3.2), we have the following Proposition.

Proposition 3.1. *The function $f: M \rightarrow \mathbb{R}$ is a Hamiltonian for a fundamental vector field of the $T \times T$ -action on M if and only if it is of the form*

$$f(z, w) = \sum_j (a_j |z_j|^2 + b_j |w_j|^2) + c$$

for some real numbers $a_j, b_j, j = 1, 2, 3$ with $a_1 + b_1 = a_2 + b_2 = a_3 + b_3$ and c .

Remark 3.2. Let $f_1, f_2: M \rightarrow \mathbb{R}$ be the function defined by

$$f_1(z, w) := \sum_j |z_j|^2, \quad f_2(z, w) := \sum_j |w_j|^2.$$

The Hamiltonian vector field generated by a linear combination of f_1 and f_2 is a fundamental vector field of the $\{1\} \times T$ -action on M and vice versa. The preimage of $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ by $\begin{pmatrix} f_1 \\ f_2 \end{pmatrix}$ coincides with $SU(3)$ in $SL(3, \mathbb{C})/U$.

4. INTERSECTION OF REAL QUADRICS IN M

Theorem 4.1. *Suppose that $A_1, A_2, A_3, B_1, B_2, B_3, C \in \mathbb{R}^2$ satisfy*

(\star) $C \notin \text{pos}(A_i, A_j), \text{pos}(B_i, B_j)$ for all $i, j \in \{1, 2, 3\}$ and $C \in \text{pos}(A_i, B_j)$ for all $i, j \in \{1, 2, 3\}$.

Then,

$$X = \{(z, w) \in \mathbb{C}^3 \times \mathbb{C}^3 \mid \sum_{j=1}^3 z_j w_j = 0, \sum_{j=1}^3 (A_j |z_j|^2 + B_j |w_j|^2) = C\}$$

is $T \times T$ -equivariantly diffeomorphic to $SU(3)$.

We admit Proposition 4.2 below for a moment and then prove Theorem 4.1. After that we prove Proposition 4.2.

Proposition 4.2. *Let $A_j \in \mathbb{R}^2, j = 1, \dots, 6$ and $C \in \mathbb{R}^2$ satisfy*

(\star') *Suppose that $i < j$. $C \in \text{pos}(A_i, A_j)$ if and only if $i \leq 3$ and $4 \leq j$.*

Let $\Phi: \mathbb{C}^6 \rightarrow \mathbb{R}^4$ be the map given by

$$\begin{pmatrix} z_1 \\ z_2 \\ z_3 \\ z_4 \\ z_5 \\ z_6 \end{pmatrix} \mapsto \begin{pmatrix} \sum_{j=1}^6 A_j |z_j|^2 \\ \operatorname{Re} \sum_{j=1}^3 z_j z_{j+3} \\ \operatorname{Im} \sum_{j=1}^3 z_j z_{j+3} \end{pmatrix}.$$

Then $\Phi^{-1}\left(\begin{pmatrix} C \\ 0 \\ 0 \\ 0 \end{pmatrix}\right)$ is nonempty and $\begin{pmatrix} C \\ 0 \\ 0 \\ 0 \end{pmatrix}$ is a regular value of Φ .

Remark 4.3. Proposition 4.2 and Remark 3.2 yield that $SU(3)$ in $SL(3, \mathbb{C})/U$ is a regular level set for the moment map for the action of $T \times T$ restricted to $\{1\} \times T$.

Proof of Theorem 4.1. By (\star) , there uniquely exist $s_A, s_B > 0$ such that $s_A A_1 + s_B B_1 = C$. Put $A := s_A A_1$ and $B := s_B B_1$ so that $A + B = C$. For $t \in [0, 1]$, we define $A_j^t := A + t(A_j - A)$ and $B_j^t := B + t(B_j - B)$, $j = 1, 2, 3$. We show that $A_1^t, A_2^t, A_3^t, B_1^t, B_2^t, B_3^t, C \in \mathbb{R}^2$ satisfy (\star) first.

Assume that $C \in \operatorname{pos}(A_i^t, A_j^t)$ for some $i, j \in \{1, 2, 3\}$. Then $C \in \operatorname{pos}(A_1, A_2, A_3)$. By Carathéodory's theorem, there exist $i', j' \in \{1, 2, 3\}$ such that $C \in \operatorname{pos}(A_{i'}, A_{j'})$. This contradicts to (\star) . By the same argument we have that $C \notin \operatorname{pos}(B_i^t, B_j^t)$.

Since $C \in \operatorname{pos}(A_i, B_j)$ for $i, j \in \{1, 2, 3\}$, there exist $r_{ij}^A, r_{ij}^B > 0$ such that $C = r_{ij}^A A_i + r_{ij}^B B_j$. Since $C = r_{1j}^A A_1 + r_{1j}^B B_j$, we have

$$\begin{aligned} C &= sC + (1-s)C \\ &= s(r_{ij}^A A_i + r_{ij}^B B_j) + (1-s)(r_{1j}^A A_1 + r_{1j}^B B_j) \\ &= sr_{ij}^A A_i + (1-s)r_{1j}^A A_1 + (sr_{ij}^B + (1-s)r_{1j}^B)B_j \end{aligned}$$

for $s \in [0, 1]$. Choose s so that $sr_{ij}^A A_i + (1-s)r_{1j}^A A_1 \in \operatorname{pos}(A_i^t)$. Then we have that there exist $r_{ij}^{A'}, r_{ij}^{B'} > 0$ such that $C = r_{ij}^{A'} A_i^t + r_{ij}^{B'} B_j$. By the same argument as above, we have that there exist $r_{ij}^{A''}, r_{ij}^{B''} > 0$ such that $C = r_{ij}^{A''} A_i^t + r_{ij}^{B''} B_j^t$. Therefore (\star) is fulfilled.

Let $\Psi: \mathbb{C}^6 \times [0, 1] \rightarrow \mathbb{R}^4$ be the map defined by

$$\left(\begin{pmatrix} z_1 \\ z_2 \\ z_3 \\ w_1 \\ w_2 \\ w_3 \end{pmatrix}, t \right) \mapsto \begin{pmatrix} \sum_{j=1}^3 (A_j^t |z|^2 + B_j^t |w_j|^2) \\ \operatorname{Re} \sum_{j=1}^3 z_j w_j \\ \operatorname{Im} \sum_{j=1}^3 z_j w_j \end{pmatrix}.$$

By proposition 4.2, we have that $\Psi^{-1}\left(\begin{pmatrix} C \\ 0 \\ 0 \\ 0 \end{pmatrix}\right)$ is nonempty and $\begin{pmatrix} C \\ 0 \\ 0 \\ 0 \end{pmatrix}$ is a regular value of Ψ . Thus

$\Psi^{-1}\left(\begin{pmatrix} C \\ 0 \\ 0 \\ 0 \end{pmatrix}\right)$ is a manifold with boundary. Let $\pi: \Psi^{-1}\left(\begin{pmatrix} C \\ 0 \\ 0 \\ 0 \end{pmatrix}\right) \rightarrow [0, 1]$ be the second projection.

We show that $\pi^{-1}(t)$ is bounded. To show this, we show the existence of a linear map $\alpha: \mathbb{R}^2 \rightarrow \mathbb{R}$ such that $\alpha(A_j^t), \alpha(B_j^t) > 0$ for all $j = 1, 2, 3$. Since $A_1^t, A_2^t, A_3^t, B_1^t, B_2^t, B_3^t$ and C satisfy (\star) , we have that B_1^t and C form a basis of \mathbb{R}^2 . Let β_1, γ be the dual of B_1^t, C . Then $A_i^t = \beta_1(A_i^t)B_1^t + \gamma(A_i^t)C$. Since $C \in \operatorname{pos}(A_i^t, B_1^t)$, we have that $\beta_1(A_i^t) < 0$ and $\gamma(A_i^t) > 0$ for all i . Let $i_0 \in \{1, 2, 3\}$ and

assume that $\beta_1(A_{i_0}^t)^{-1}\gamma(A_{i_0}^t) \geq \beta_1(A_i^t)^{-1}\gamma(A_i^t)$ for $i = 1, 2, 3$. Since $A_i^t = \beta_1(A_i^t)B_1^t + \gamma(A_i^t)C$ and $B_1^t = \beta_1(A_{i_0}^t)^{-1}(A_{i_0}^t - \gamma(A_{i_0}^t)C)$, we have

$$\begin{aligned} A_i^t &= \beta_1(A_i^t)B_1^t + \gamma(A_i^t)C \\ &= \beta_1(A_i^t)(\beta_1(A_{i_0}^t)^{-1}(A_{i_0}^t - \gamma(A_{i_0}^t)C) + \gamma(A_{i_0}^t)C) \\ &= \beta_1(A_i^t)\beta_1(A_{i_0}^t)^{-1}A_{i_0}^t + (\gamma(A_i^t) - \beta_1(A_i^t)\beta_1(A_{i_0}^t)^{-1}\gamma(A_{i_0}^t))C, \end{aligned}$$

showing that $A_i^t \in \text{pos}(A_{i_0}^t, C)$. By the same argument, we may find $j_0 \in \{1, 2, 3\}$ such that $B_j^t \in \text{pos}(B_{j_0}^t, C)$. Since $C \in \text{pos}(A_{i_0}^t, B_{j_0}^t)$, we have that $A_1^t, A_2^t, A_3^t, B_1^t, B_2^t, B_3^t \in \text{pos}(A_{i_0}^t, B_{j_0}^t)$. Let $\alpha: \mathbb{R}^2 \rightarrow \mathbb{R}$ be the sum of the dual of $A_{i_0}^t, B_{j_0}^t$. Then $\alpha(A_j^t), \alpha(B_j^t) > 0$ for all $j = 1, 2, 3$. Let

$$\begin{pmatrix} z_1 \\ z_2 \\ z_3 \\ w_1 \\ w_2 \\ w_3 \end{pmatrix} \in \pi^{-1}(t). \text{ Since } \sum_{j=1}^3 (\alpha(A_j^t)|z_j|^2 + \alpha(B_j^t)|w_j|^2) = \alpha(C) \text{ and all coefficients are positive, we}$$

have that $\pi^{-1}(t)$ is bounded.

As we saw in Remark 3.2, the preimage $\pi^{-1}(0)$ is equivariantly diffeomorphic to $SU(3)$. π is an equivariant proper surjective submersion, here the $T \times T$ -action on $[0, 1]$ is trivial. By the equivariant version of Ehressmann's lemma, we have that $\pi^{-1}(0) = SU(3)$ and $\pi^{-1}(1) = X$ are equivariantly diffeomorphic. Theorem is proved. \square

Proof of Proposition 4.2. By (\star') , there exist $r_1, r_5 > 0$ such that $r_1A_1 + r_5A_5 = C$. Then

$$\Phi \left(\begin{pmatrix} \sqrt{r_1} \\ 0 \\ 0 \\ 0 \\ \sqrt{r_5} \\ 0 \end{pmatrix} \right) = \begin{pmatrix} C \\ 0 \\ 0 \end{pmatrix}.$$

Thus $\Phi^{-1} \left(\begin{pmatrix} C \\ 0 \\ 0 \end{pmatrix} \right)$ is not empty. Let

$$z = \begin{pmatrix} z_1 \\ z_2 \\ z_3 \\ z_4 \\ z_5 \\ z_6 \end{pmatrix} \in \Phi^{-1} \left(\begin{pmatrix} C \\ 0 \\ 0 \end{pmatrix} \right).$$

Put $J := \{j \in \{1, \dots, 6\} \mid z_j \neq 0\}$. Since $\sum_{j=1}^6 A_j |z_j|^2 = C$, it follows from **(C)** that there exist $i, j \in J$ with $i \leq 3$ and $4 \leq j$ such that $|z_i|^2 \neq 0$ and $|z_j|^2 \neq 0$. We may assume that $j \neq i+3$ because $\sum_{j=1}^3 z_j z_{j+3} = 0$. For simplicity, suppose that $z_1 \neq 0$ and $z_5 \neq 0$. We define the $(S^1)^4$ -action on \mathbb{C}^6

by

$$(g_1, g_2, h_1, h_3) \cdot \begin{pmatrix} z_1 \\ z_2 \\ z_3 \\ z_4 \\ z_5 \\ z_6 \end{pmatrix} = \begin{pmatrix} g_1 h_1^{-1} z_1 \\ g_2 h_1^{-1} z_2 \\ (g_1 g_2)^{-1} h_1^{-1} z_3 \\ g_1^{-1} h_3 z_4 \\ g_2^{-1} h_3 z_5 \\ g_1 g_2 h_3 z_6 \end{pmatrix}$$

for $(g_1, g_2, h_1, h_3) \in (S^1)^4$ (see also (3.2)). There exist $g_1, g_2, h_1, h_3 \in S^1$ such that

$$g_1 h_1^{-1} z_1, \quad g_2 h_1^{-1} z_2, \quad (g_1 g_2)^{-1} h_1^{-1} z_3, \quad g_2^{-1} h_3 z_5 \in \mathbb{R}_{\geq 0}.$$

Indeed, if

$$\begin{aligned} g_1 &= \exp\left(\frac{\sqrt{-1}}{3}(-2 \arg z_1 + \arg z_2 + \arg z_3)\right), \\ g_2 &= \exp\left(\frac{\sqrt{-1}}{3}(\arg z_1 - 2 \arg z_2 + \arg z_3)\right), \\ h_1 &= \exp\left(\frac{\sqrt{-1}}{3}(\arg z_1 + \arg z_2 + \arg z_3)\right), \\ h_3 &= \exp(-\sqrt{-1} \arg(z_5 g_2)) \end{aligned}$$

then $g_1 h_1^{-1} z_1 = |z_1|$, $g_2 h_1^{-1} z_2 = |z_2|$, $(g_1 g_2)^{-1} h_1^{-1} z_3 = |z_3|$ and $g_2 h_3 z_5 = |z_5|$. We define the linear action of $(S^1)^4$ on \mathbb{R}^4 by

$$(g_1, g_2, h_1, h_3) \cdot \begin{pmatrix} u_1 \\ u_2 \\ u_3 \\ u_4 \end{pmatrix} = \begin{pmatrix} u_1 \\ u_2 \\ \operatorname{Re}(h_1^{-1} h_3 (u_3 + \sqrt{-1} u_4)) \\ \operatorname{Im}(h_1^{-1} h_3 (u_3 + \sqrt{-1} u_4)) \end{pmatrix}.$$

Then Φ is $(S^1)^4$ -equivariant. Since the action is smooth and Φ is $(S^1)^4$ -equivariant, it is enough to consider the case when $z_1, z_2, z_3, z_5 \in \mathbb{R}_{\geq 0}$ because the diagram

$$\begin{array}{ccc} T_z \mathbb{C}^6 & \xrightarrow{d\Phi} & T_{\Phi(z)} \mathbb{R}^4 \\ \downarrow & & \downarrow \\ T_{(g_1, g_2, h_1, h_3) \cdot z} \mathbb{C}^6 & \xrightarrow{d\Phi} & T_{\Phi(z)} \mathbb{R}^4 \end{array}$$

is commutative, where $z \in \Phi^{-1}\left(\begin{pmatrix} C \\ 0 \\ 0 \end{pmatrix}\right)$ and the vertical arrows are the isomorphisms coming from

the action of $(S^1)^4$.

Let x_j and y_j be the real and imaginary parts of z_j , $j = 1, \dots, 6$. If $z_2 = 0$, then

$$\begin{aligned} \det \begin{pmatrix} \frac{\partial \Phi}{\partial x_1} & \frac{\partial \Phi}{\partial x_5} & \frac{\partial \Phi}{\partial x_2} & \frac{\partial \Phi}{\partial y_2} \end{pmatrix}_z &= \det \begin{pmatrix} 2x_1 A_1 & 2x_5 A_5 & 0 & 0 \\ x_4 & 0 & x_5 & 0 \\ y_4 & 0 & 0 & x_5 \end{pmatrix} \\ &= 4x_1 x_5^3 \det(A_1 \quad A_5). \end{aligned}$$

Since A_1 and A_5 are linearly independent and $x_1, x_5 \neq 0$, the minor above of the Jacobian matrix at z is not zero. Thus z is a regular point.

If $z_4 = 0$, then

$$\begin{aligned} \det \begin{pmatrix} \frac{\partial \Phi}{\partial x_1} & \frac{\partial \Phi}{\partial x_5} & \frac{\partial \Phi}{\partial x_4} & \frac{\partial \Phi}{\partial y_4} \end{pmatrix}_z &= \det \begin{pmatrix} 2x_1 A_1 & 2x_5 A_5 & 0 & 0 \\ 0 & x_2 & x_1 & 0 \\ 0 & 0 & 0 & x_1 \end{pmatrix} \\ &= 4x_1^3 x_5 \det(A_1 \ A_5). \end{aligned}$$

By the same reason as the case when $z_2 = 0$, we have that z is a regular point.

Suppose that $z_2, z_4 \neq 0$. Suppose that $z_4 \notin \mathbb{R}$. Then

$$\begin{aligned} \det \begin{pmatrix} \frac{\partial \Phi}{\partial x_1} & \frac{\partial \Phi}{\partial x_5} & \frac{\partial \Phi}{\partial y_1} & \frac{\partial \Phi}{\partial y_5} \end{pmatrix}_z &= \det \begin{pmatrix} 2x_1 A_1 & 2x_5 A_5 & 0 & 0 \\ x_4 & x_2 & -y_4 & 0 \\ y_4 & 0 & x_4 & x_2 \end{pmatrix} \\ &= -4x_1 x_2 y_4 x_5 \det(A_1 \ A_5). \end{aligned}$$

Since $y_4 \neq 0$, we have that z is a regular point.

Suppose that $z_4 \in \mathbb{R}_{>0}$. Then $z_3 z_6 \in \mathbb{R}_{<0}$ because $\sum_{j=1}^3 z_j z_{j+3} = 0$. Since $z_3 \in \mathbb{R}_{\geq 0}$, we have that $z_3 \in \mathbb{R}_{>0}$ and $z_6 \in \mathbb{R}_{<0}$. Put

$$x'_j := \begin{cases} x_{j+3} & j = 1, 2, 3 \\ x_{j-3} & j = 4, 5, 6. \end{cases}$$

and $A'_j := \frac{x_j}{x'_j} A_j$ for $j = 1, \dots, 6$. We will show that A'_1, A'_4, A'_3, A'_6 are not on the same line in \mathbb{R}^2 .

Suppose that A'_1, A'_4, A'_3, A'_6 are on the same line in \mathbb{R}^2 . Since $C \in \text{pos}(A_1, A_4) = \text{pos}(A'_1, A'_4)$, there exist $r_1, r_4 > 0$ such that $C = r_1 A_1 + r_4 A_4$. Since $C \in \text{pos}(A_3, A_6) = \text{pos}(-A'_3, -A'_6)$, there exist $r_3, r_6 > 0$ such that $-C = r_3 A'_3 + r_6 A'_6$. Then $0 = r_1 A'_1 + r_3 A'_3 + r_4 A'_4 + r_6 A'_6$. In particular, 0 sits in the convex hull of A'_1, A'_3, A'_4, A'_6 . Thus we have that 0 is also on the same line in \mathbb{R}^2 . It turns out that A'_1 and A'_4 are linearly dependent. Since A'_1 and A'_4 are linearly dependent, so are A_1 and A_4 . This contradicts to (\star') . Let $i, j, k \in \{1, 3, 4, 6\}$ such that A'_i, A'_j, A'_k are not on the same line in \mathbb{R}_2 . Then

$$\begin{aligned} \det \begin{pmatrix} \frac{\partial \Phi}{\partial x_i} & \frac{\partial \Phi}{\partial x_j} & \frac{\partial \Phi}{\partial x_k} & \frac{\partial \Phi}{\partial y_1} \end{pmatrix}_z &= \det \begin{pmatrix} 2x_i A_i & 2x_j A_j & 2x_k A_k & 0 \\ x'_i & x'_j & x'_k & 0 \\ 0 & 0 & 0 & x_4 \end{pmatrix} \\ &= 8x'_i x'_j x'_k x_4 \det \begin{pmatrix} A'_i & A'_j & A'_k \\ 1 & 1 & 1 \end{pmatrix}. \end{aligned}$$

Since A'_i, A'_j, A'_k are affinely independent, we have that $\begin{pmatrix} A'_i \\ 1 \end{pmatrix}, \begin{pmatrix} A'_j \\ 1 \end{pmatrix}, \begin{pmatrix} A'_k \\ 1 \end{pmatrix}$ are linearly independent. Therefore z is a regular point.

Finally, if $z_4 \in \mathbb{R}_{<0}$, then we have that A'_1, A'_2, A'_4, A'_5 are not on the same line in \mathbb{R}^2 by the same argument as above. Let $i, j, k \in \{1, 2, 4, 5\}$ such that A'_i, A'_j, A'_k are not on the same line in \mathbb{R}_2 . Then

$$\begin{aligned} \det \begin{pmatrix} \frac{\partial \Phi}{\partial x_i} & \frac{\partial \Phi}{\partial x_j} & \frac{\partial \Phi}{\partial x_k} & \frac{\partial \Phi}{\partial y_1} \end{pmatrix}_z &= \det \begin{pmatrix} 2x_i A_i & 2x_j A_j & 2x_k A_k & 0 \\ x'_i & x'_j & x'_k & 0 \\ 0 & 0 & 0 & x_4 \end{pmatrix} \\ &= 8x'_i x'_j x'_k x_4 \det \begin{pmatrix} A'_i & A'_j & A'_k \\ 1 & 1 & 1 \end{pmatrix}. \end{aligned}$$

Since A'_i, A'_j, A'_k are affinely independent, we have that $\begin{pmatrix} A'_i \\ 1 \end{pmatrix}, \begin{pmatrix} A'_j \\ 1 \end{pmatrix}, \begin{pmatrix} A'_k \\ 1 \end{pmatrix}$ are linearly independent. Therefore z is a regular point.

The proposition is proved. \square

Now we are in a position to show Theorem 1.1.

Proof of Theorem 1.1. Assume that A_j, B_j for $j = 1, 2, 3$ and C satisfy (\star) . Let M be the quasi-affine variety defined as

$$M := \{(z, w) \in \mathbb{C}^3 \times \mathbb{C}^3 \mid z \neq 0, w \neq 0, \sum_{j=1}^3 z_j w_j = 0\}.$$

It follows from Proposition 4.2 that C is a regular value of the map $\Phi: M \rightarrow \mathbb{R}^2$ given by $\Phi(z, w) := \sum_{j=1}^3 (A_j |z_j|^2 + B_j |w_j|^2)$ and $\Phi^{-1}(C)$ is not empty. It follows from Proposition 3.1 that Φ is a moment map for the $(\rho_L, \rho_R)(T)$ -action on M . It follows from Theorem 4.1 that $\Phi^{-1}(C)$ is $T \times T$ equivariantly diffeomorphic to $SU(3)$. Thus by Theorem 2.1 we have (1). By Corollary 2.2 we have (2). \square

Remark 4.4. As Remark 2.3, for each (ρ_L, ρ_R) under the condition (\star) , we obtain the family of complex structures on $SU(3)$ parametrized by $GL(2, \mathbb{R})/GL(1, \mathbb{C})$. The Kähler orbifold structure on $SU(3)/(\rho_L, \rho_R)((S^1)^2)$ is uniquely determined by (ρ_L, ρ_R) .

Remark 4.5. Theorem 1.1 (1) can be generalized to the double sided \mathbb{R}^2 -action on $SU(3)$. Let $w_1^L, w_2^L, w_3^L, w_1^R, w_2^R, w_3^R \in \mathbb{R}^2$ such that $\sum_{j=1}^3 w_j^L = \sum_{j=1}^3 w_j^R = 0$ and (\star) is fulfilled. We consider the action of \mathbb{R}^2 on $SU(3)$ defined by

$$v \cdot A := \text{diag}(e^{\sqrt{-1}\langle v, w_1^L \rangle}, e^{\sqrt{-1}\langle v, w_2^L \rangle}, e^{\sqrt{-1}\langle v, w_3^L \rangle}) \text{Adiag}(e^{\sqrt{-1}\langle v, w_1^R \rangle}, e^{\sqrt{-1}\langle v, w_2^R \rangle}, e^{\sqrt{-1}\langle v, w_3^R \rangle})^{-1}$$

for $v \in \mathbb{R}^2$ and $A \in SU(3)$, where $\langle -, - \rangle$ denote the standard inner product on \mathbb{R}^2 . Then there exists a $T \times T$ -invariant complex structure on $SU(3)$ such that the foliation whose leaves are \mathbb{R}^2 -orbits is holomorphic and transversely Kähler. All leaves are closed if and only if $A_j, B_j, j = 1, 2, 3$ generate a lattice of \mathbb{R}^2 .

Consider $A_j^t, B_j^t \in \mathbb{R}^2$ as in the Proof of Theorem 4.1. Corresponding to

$$w_1^L(t), w_2^L(t), w_3^L(t), w_1^R(t), w_2^R(t), w_3^R(t) \in \mathbb{R}^2$$

so that

$$A_j^t := w_j^L(t) - w_1^R(t), \quad B_j^t := -w_j^L(t) + w_3^R(t), \quad C := -w_1^R(t) + w_3^R(t),$$

by the above arguments, we also obtain a smooth family $\{J_t\}$ of $T \times T$ -invariant complex structures on $SU(3)$ parametrized by $t \in [0, 1]$. We can check that $\{J_0\}$ is a left-invariant complex structure on $SU(3)$ (see Remark 5.4). In [7], we compute the Kuranishi spaces of left-invariant complex structures on even-dimensional simply connected compact Lie groups. This allows us to describe all small deformations of left-invariant complex structures. However, we use Kodaira-Spencer theory in [7]. Therefore it is difficult to construct deformations in the large such as $\{J_t\}$ in this way.

5. FREENESS OF THE DOUBLE SIDED ACTION

In this section, we consider the case when $(\rho_L, \rho_R)((S^1)^2)$ -action is free under the condition (\star) .

Lemma 5.1. *Let $\rho_L, \rho_R: (S^1)^2 \rightarrow T$ be smooth homomorphisms and A_j, B_j as Theorem 1.1. The $(S^1)^2$ -action on $SU(3)$ given by ρ_L, ρ_R is free if and only if A_i, B_j form a \mathbb{Z} -basis of \mathbb{Z}^2 for all pair (i, j) with $i \neq j$.*

Proof. As we saw in Remark 3.2, $SU(3)$ is equivariantly diffeomorphic to $X_0 := \{(z, w) \in \mathbb{C}^3 \times \mathbb{C}^3 \mid \sum_{j=1}^3 |z_j|^2, \sum_{j=1}^3 |w_j|^2, \sum_{j=1}^3 z_j w_j = 0\} \subset M$. We shall see the isotropy subgroups of the T -action on X_0 given by (ρ_L, ρ_R) . Let $(z, w) \in X_0$. Put $I_z := \{i \mid z_i \neq 0\}$ and $J_w := \{j \mid w_j \neq 0\}$. Since the $(S^1)^2$ -action on X_0 given by (ρ_L, ρ_R) satisfies

$$t \cdot (z, w) = (t^{A_1} z_1, t^{A_2} z_2, t^{A_3} z_3, t^{B_1} w_1, t^{B_2} w_2, t^{B_3} w_3)$$

by (3.2), the isotropy subgroup at (z, w) is isomorphic to

$$\bigcap_{i \in I_z} \ker(t \mapsto t^{A_i}) \cap \bigcap_{j \in J_w} \ker(t \mapsto t^{B_j}).$$

Namely, the isotropy subgroup at (z, w) is trivial if and only if A_i for $i \in I_z$ and B_j for $j \in J_w$ span \mathbb{Z}^2 .

For “if” part, assume that A_i, B_j form a \mathbb{Z} -basis of \mathbb{Z}^2 for all pair (i, j) with $i \neq j$. Since $z \neq 0$ and $w \neq 0$, we have that $I_z \times J_w \neq \emptyset$. Since $\sum_{j=1}^3 z_j w_j = 0$, we have that there exists $(i, j) \in I_z \times J_w$ with $i \neq j$. Thus we have that the isotropy subgroup at (z, w) is trivial. Since $(z, w) \in X_0$ is arbitrary, we have that the T -action on X_0 given by (ρ_L, ρ_R) is free.

For “only if” part, assume that the $(S^1)^2$ -action on X_0 given by (ρ_L, ρ_R) is free. Let $(i_0, j_0) \in \{1, 2, 3\} \times \{1, 2, 3\}$ satisfies $i_0 \neq j_0$. Put

$$z_i = \begin{cases} 0, & i \neq i_0, \\ 1, & i = i_0, \end{cases} \quad w_j = \begin{cases} 0, & j \neq j_0, \\ 1, & j = j_0. \end{cases}$$

Then $(z, w) := (z_1, z_2, z_3, w_1, w_2, w_3) \in X_0$. Since the isotropy subgroup at (z, w) is trivial, we have that A_i and B_j span \mathbb{Z}^2 . Therefore A_i, B_j form a \mathbb{Z} -basis of \mathbb{Z}^2 . □

The lemma is proved. □

Lemma 5.2. *If $A_j, B_j \in \mathbb{Z}^2$ for $j = 1, 2, 3$ and $C \in \mathbb{Z}^2$ satisfy $A_j + B_j = C$ for all j , (\star) and A_i, B_j form a \mathbb{Z} -basis of \mathbb{Z}^2 , then $A_1 = A_2 = A_3$ and $B_1 = B_2 = B_3$.*

Proof. Since A_1, B_2 form a \mathbb{Z} -basis of \mathbb{Z}^2 , $C \in \text{pos}(A_1, B_2)$ and $C \notin \text{pos}(A_1, A_1), \text{pos}(B_2, B_2)$, we have that there exist $a, b \in \mathbb{Z}_{>0}$ such that $C = aA_1 + bB_2$. Also, since A_1, B_3 form a \mathbb{Z} -basis, we have that there exists $c \in \mathbb{Z}$ such that $B_2 = \pm B_3 + cA_1$. By substituting $B_2 = \pm B_3 + cA_1$ for $C = aA_1 + bB_2$, we have that $C = (a + bc)A_1 \pm bB_3$. Since $C = \text{pos}(A_1, B_3)$ and $b > 0$, we have that the sign \pm should be $+$ and $a + bc > 0$. Since $C = A_j + B_j$ for all j , we have that $B_1 = (a - 1)A_1 + bB_2$, $A_2 = aA_1 + (b - 1)B_2$ and $A_3 = (a + bc)A_1 + (b - 1)B_3 = (a + c)A_1 + (b - 1)B_2$. Since A_3, B_2 form a \mathbb{Z} -basis of \mathbb{Z}^2 , we have that $a + c = \pm 1$. By substituting $A_1 = \pm A_3 + (1 - b)B_2$ for $C = aA_1 + bB_2$, we have that $C = \pm aA_1 + (b \pm (a - ab))B_2$. Since $C \in \text{pos}(A_3, B_2)$, we have that $a + c = 1$. This together with $a > 0$ yields that $a = 1$ and $c = 0$. Therefore $B_1 = bB_2$. Since $b > 0$ and B_1 is primitive, we have that $b = 1$. These computations show that $A_1 = A_2 = A_3$ and $B_1 = B_2 = B_3$, proving the lemma. □

Proposition 5.3. *Let $\rho_L, \rho_R: T \rightarrow T$ be smooth homomorphisms and A_j, B_j, C as Theorem 1.1. Assume that A_j, B_j, C satisfy the condition (\star) . Then, the action of $(S^1)^2$ on $SU(3)$ given by (ρ_L, ρ_R) is free if and only if ρ_L is trivial and ρ_R is an isomorphism.*

Proof. The “only if” part is obvious. For the “if” part, assume that the action of $(\rho_L, \rho_R)((S^1)^2)$ on $SU(3)$ is free. By Lemmas 5.1 and 5.2, we have that $A_1 = A_2 = A_3$ and $B_1 = B_2 = B_3$. Thus $w_1^L = w_2^L = w_3^L$. Since $\sum_{j=1}^3 w_j^L = 0$, we have $w_j^L = 0$ for all j . This means that ρ_L is trivial. Since $w_1^R = -A_1 = -A_2 = -A_3$ and $w_3^R = B_1 = B_2 = B_3$ form a \mathbb{Z} -basis of \mathbb{Z}^2 , we have that ρ_R is an isomorphism. □

Remark 5.4. In case ρ_L is trivial (equivalently $w_1^R = -A_1 = -A_2 = -A_3$ and $w_3^R = B_1 = B_2 = B_3$), the set $\Phi^{-1}\left(\begin{pmatrix} C \\ 0 \\ 0 \end{pmatrix}\right)$ is an orbit of $SU(3)$ -action on $SL(3, \mathbb{C})/U$. In this case, since the complex structure, the Kähler structure and the fundamental vector fields of the $\{1\} \times T$ -action on $SL(3, \mathbb{C})/U$ are invariant under $SU(3)$ -action on $SL(3, \mathbb{C})/U$, we have that a complex structure and a transversely Kähler structure on $SU(3)$ given by Theorem 1.1 are left-invariant.

Proposition 5.3 yields that, under the condition (\star) , if the action of $(\rho_L, \rho_R)((S^1)^2)$ is free, then the quotient $SU(3)/(\rho_L, \rho_R)((S^1)^2)$ is nothing but a flag manifold with an invariant Kähler structure.

Finally, we shall see a nontrivial example of (ρ_L, ρ_R) which satisfies (\star) .


Example 5.5. We begin with the configuration of points $A_1, A_2, A_3, B_1, B_2, B_3, C \in \mathbb{R}^2$. Since $C \notin (A_i, A_j)$ for all $i, j = 1, 2, 3$, the 3 points A_1, A_2, A_3 should sit in the same half plane whose boundary is the line spanned by C . For example, choose A_1, A_2, A_3 and C as

$$A_1 = A_2 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad A_3 = \begin{pmatrix} 2 \\ -1 \end{pmatrix}, \quad C = \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

We choose B_1, B_2, B_3 as

$$B_1 = B_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad B_3 = \begin{pmatrix} -1 \\ 2 \end{pmatrix}$$

so that $A_j + B_j = C$. Then $A_1, A_2, A_3, B_1, B_2, B_3, C$ satisfy (\star) and $A_1 + B_1 = A_2 + B_2 = A_3 + B_3$ (see the figure below).



Now we solve the linear equations

$$A_j = w_j^L - w_1^R, \quad B_j = -w_j^L + w_3^R, \quad \sum_{j=1}^3 w_j^L = \sum_{j=1}^3 w_j^R = 0$$

about $w_j^L, w_j^R, j = 1, 2, 3$. Then we have

$$\begin{aligned} w_1^L &= -(A_1 + A_2 + A_3)/3 + A_1 = \begin{pmatrix} -1/3 \\ 1/3 \end{pmatrix}, \\ w_2^L &= -(A_1 + A_2 + A_3)/3 + A_2 = \begin{pmatrix} -1/3 \\ 1/3 \end{pmatrix}, \\ w_3^L &= -(A_1 + A_2 + A_3)/3 + A_3 = \begin{pmatrix} 2/3 \\ -2/3 \end{pmatrix}, \\ w_1^R &= -(A_1 + A_2 + A_3)/3 = \begin{pmatrix} -4/3 \\ 1/3 \end{pmatrix}, \\ w_2^R &= (A_1 + A_2 + A_3)/3 - (B_1 + B_2 + B_3)/3 = \begin{pmatrix} 5/3 \\ -5/3 \end{pmatrix}, \\ w_3^R &= (B_1 + B_2 + B_3)/3 = \begin{pmatrix} -1/3 \\ 4/3 \end{pmatrix}. \end{aligned}$$

By multiplying 3 so that every entries become integers, we get the homomorphisms

$$\begin{aligned} \rho_L(t) &= \text{diag}(t^{3w_1^L}, t^{3w_2^L}, t^{3w_3^L}) = \text{diag}(t_1^{-1}t_2, t_1^{-1}t_2, t_1^2t_2^{-2}), \\ \rho_R(t) &= \text{diag}(t^{3w_1^R}, t^{3w_2^R}, t^{3w_3^R}) = \text{diag}(t_1^{-4}t_2, t_1^5t_2^{-5}, t_1^{-1}t_2^4) \end{aligned}$$

such that the quotient space $SU(3)/(\rho_L, \rho_R)((S^1)^2)$ has a Kähler orbifold structure.

6. COHOMOLOGY

We consider $SU(3)$ equipped with the complex structure J and the transversely Kähler holomorphic foliation \mathcal{F} determined by $\rho_L, \rho_R : (S^1)^2 \rightarrow T$ under the assumption (\star) as in Theorem 1.1. In this case, we can apply [6, Theorem 4.13]. That is, for a real 2-dimensional vector space W , we have a differential graded algebra $(A^* = H_B^*(SU(3)) \otimes \wedge W, d)$ so that

- $H_B^*(SU(3))$ is the basic cohomology algebra of $SU(3)$ associated with the foliation \mathcal{F} ;
- W consists elements of degree 1;
- $dW \subset H_B^2(SU(3))$;
- d on $H_B^*(SU(3))$ is trivial;
- $H^*(A^*) \cong H^*(SU(3), \mathbb{R})$.

It is known that $H^*(SU(3), \mathbb{R}) \cong \wedge \langle a_3, a_5 \rangle$ with $\deg(a_3) = 3$ and $\deg(a_5) = 5$. Consequently, $H_B^1(SU(3)) = 0$ and $d : W \rightarrow H_B^2(SU(3))$ is bijective. Thus, for a basis w_1, w_2 of W , we have that $H_B^2(SU(3)) = \langle dw_1, dw_2 \rangle$. Since the foliation \mathcal{F} is transversely Kähler, we have that $H_B^6(SU(3)) = \langle [\omega]^3 \rangle$, where ω is a transverse Kähler form with respect to \mathcal{F} (see [6, Proposition 4.8]). By the hard Lefschetz property ([2]), we have that $H_B^4(SU(3)) = \langle [\omega] \wedge dw_1, [\omega] \wedge dw_2 \rangle$ and $H_B^5(SU(3)) = 0$. It follows from $A^3 = H_B^2(SU(3)) \otimes W$, $A^4 = H_B^3(SU(3)) \otimes W \oplus H_B^4(SU(3))$ and $H^4(SU(3)) = 0$ that $H_B^3(SU(3)) = 0$.

In summary, we have that

$$\dim H_B^k(SU(3)) = \begin{cases} 1, & k = 0, 6, \\ 2, & k = 2, 4, \\ 0 & \text{otherwise.} \end{cases}$$

Remark 6.1. The basic cohomology $H_B^*(SU(3))$ is canonically isomorphic to the singular cohomology $H^*(SU(3)/(\rho_L, \rho_R)((S^1)^2), \mathbb{R})$ of the topological space $SU(3)/(\rho_L, \rho_R)((S^1)^2)$ (see [13, Corollary 5.3.3]).

We consider the basic Dolbeault cohomology $H_B^{*,*}(SU(3))$ for holomorphic foliation \mathcal{F} . Then, we have the Hodge decomposition $H_B^r(SU(3)) \otimes \mathbb{C} = \bigoplus_{p+q=r} H_B^{p,q}(SU(3))$ ([2]). The transverse Kähler form defines the non-zero class $[\omega] \in H_B^{1,1}(SU(3))$. It follows from the symmetry of basic Hodge numbers that $H_B^2(SU(3)) \otimes \mathbb{C} = H_B^{1,1}(SU(3))$. Thus we also have $H_B^4(SU(3)) \otimes \mathbb{C} = H_B^{2,2}(SU(3))$.

We denote by $H^{*,*}(SU(3))$ the Dolbeault cohomology of $SU(3)$ equipped with the complex structure determined by $\rho_L, \rho_R : (S^1)^2 \rightarrow T$ under the assumption (\star) as in Theorem 1.1. By [6, Theorem 4.13], for an almost complex structure $W \otimes \mathbb{C} = W^{1,0} \oplus \overline{W^{1,0}}$, we have a differential bi-graded algebra $(B^{*,*} = H_B^{*,*}(SU(3)) \otimes \wedge(W^{1,0} \oplus \overline{W^{1,0}}), \bar{\partial})$ so that

- $W^{1,0}$ (respectively, $\overline{W^{1,0}}$) consists elements of degree $(1, 0)$ (respectively, $(0, 1)$);
- $\bar{\partial}W^{1,0} \subset H_B^{1,1}(SU(3))$ and $\bar{\partial}\overline{W^{1,0}} \subset H_B^{0,2}(SU(3))$;
- $\bar{\partial}$ on $H_B^{*,*}(SU(3))$ is trivial;
- $H^{*,*}(B^{*,*}) \cong H^{*,*}(SU(3))$.

Since $H_B^{0,2}(SU(3)) = 0$, we have that $H^{0,1}(SU(3)) \cong \overline{W^{1,0}}$. Since $H_B^{2,0}(SU(3)) = 0$, we have that $H_B^{1,1}(SU(3)) \ni \bar{\partial}w \neq 0$ for all $W^{1,0} \ni w \neq 0$ and hence $H^{1,0}(SU(3)) = 0$. Other Hodge numbers can be obtained by direct computation. The Hodge diamond is as below:

$$\begin{array}{ccccccc}
 & & & & & & 1 \\
 & & & & & & 0 & 1 \\
 & & & & & & 0 & 1 & 0 \\
 & & & & & & 0 & h^{2,1} & 1 & 0 \\
 & & & & & & 0 & 0 & h^{2,2} & 0 & 0 \\
 & & & & & & 0 & 1 & h^{2,3} & 0 \\
 & & & & & & 0 & 1 & 0 \\
 & & & & & & 1 & 0 \\
 & & & & & & 1
 \end{array}$$

The Hodge numbers $h^{2,1}, h^{2,2}, h^{2,3}$ are $(0, 0, 0)$ or $(1, 2, 1)$, depending on an almost complex structure $W \otimes \mathbb{C}$.

Remark 6.2. If ρ_L is non-trivial, then the complex structure on $SU(3)$ is not left-invariant. The computations of Dolbeault cohomology on compact Lie groups equipped with left-invariant complex structures (e.g. [14]) can not be applied.

Remark 6.3. We can describe W and $d: W \rightarrow H_B^2(SU(3))$ explicitly for the differential graded algebra $(A^* = H_B^*(SU(3)) \otimes \wedge W, d)$ see [6, Page 68]. We denote by $(-, -)$ the Cartan-Killing form on $SU(3)$. For a vector field X on $SU(3)$, we define the 1-form $w_X = (X, -)$. Let $\bar{\mathfrak{t}}$ be the vector space of fundamental vector fields on $SU(3)$ defined by the $(S^1)^2$ -action associated with $\rho_L, \rho_R : (S^1)^2 \rightarrow T$. Then, for any $X \in \bar{\mathfrak{t}}$, the 1-form w_X is $T \times T$ -invariant. We can take $W = \{w_X \mid X \in \bar{\mathfrak{t}}\}$ and $d: W \ni w \mapsto [dw] \in H_B^2(SU(3))$.

Remark 6.4. For the differential bi-graded algebra $(B^{*,*} = H_B^{*,*}(SU(3)) \otimes \wedge(W^{1,0} \oplus \overline{W^{1,0}}), \bar{\partial})$, we have to reconstruct W so that W is J -invariant see [6, Proposition 4.9]. Recall that we have a direct sum $TN = T\mathcal{F} \oplus T\mathcal{F}^\perp$ so that $T\mathcal{F}^\perp$ is J -invariant and invariant under the $(S^1)^2$ -action associated with $\rho_L, \rho_R : (S^1)^2 \rightarrow T$ see Section 2. Regarding $TN^* = T\mathcal{F}^* \oplus (T\mathcal{F}^\perp)^*$, we should replace W with the vector space of 1-forms which is generated by the locally free $(S^1)^2$ -action associated with $\rho_L, \rho_R : (S^1)^2 \rightarrow T$. An almost complex structure $W \otimes \mathbb{C} = W^{1,0} \oplus \overline{W^{1,0}}$ corresponds to parameter in $GL(2, \mathbb{R})/GL(1, \mathbb{C})$ as in Remark 4.4.

REFERENCES

- [1] Michèle Audin, *Torus actions on symplectic manifolds*, Second revised edition, Progress in Mathematics, vol. 93, Birkhäuser Verlag, Basel, 2004. MR2091310
- [2] Aziz El Kacimi-Alaoui, *Opérateurs transversalement elliptiques sur un feuilletage riemannien et applications*, *Compositio Math.* **73** (1990), no. 1, 57–106. MR1042454
- [3] J.-H. Eschenburg, *Inhomogeneous spaces of positive curvature*, *Differential Geom. Appl.* **2** (1992), no. 2, 123–132, DOI 10.1016/0926-2245(92)90029-M. MR1245552
- [4] Christine Escher and Wolfgang Ziller, *Topology of non-negatively curved manifolds*, *Ann. Global Anal. Geom.* **46** (2014), no. 1, 23–55, DOI 10.1007/s10455-013-9407-8. MR3205800
- [5] Oliver Goertsches, Panagiotis Konstantis, and Leopold Zoller, *Symplectic and Kähler structures on biquotients*, *J. Symplectic Geom.* **18** (2020), no. 3, 791–813, DOI 10.4310/JSG.2020.v18.n3.a6. MR4142487
- [6] Hiroaki Ishida and Hisashi Kasuya, *Transverse Kähler structures on central foliations of complex manifolds*, *Ann. Mat. Pura Appl. (4)* **198** (2019), no. 1, 61–81, DOI 10.1007/s10231-018-0762-8. MR3918619
- [7] ———, *Non-invariant deformations of left-invariant complex structures on compact Lie groups. preprint arXiv:2004.02523*, *Ann. Mat. Pura Appl. (4)* **1** (2020), 61–81, DOI 10.1007/s10231-018-0762-8. MR3918619
- [8] A. Haefliger, *Deformations of transversely holomorphic flows on spheres and deformations of Hopf manifolds*, *Compositio Math.* **55** (1985), no. 2, 241–251. MR795716
- [9] Jean-Jacques Loeb, Mónica Manjarín, and Marcel Nicolau, *Complex and CR-structures on compact Lie groups associated to abelian actions*, *Ann. Global Anal. Geom.* **32** (2007), no. 4, 361–378, DOI 10.1007/s10455-007-9067-7. MR2346223
- [10] Santiago López de Medrano and Alberto Verjovsky, *A new family of complex, compact, non-symplectic manifolds*, *Bol. Soc. Brasil. Mat. (N.S.)* **28** (1997), 253–269. MR1479504
- [11] Laurent Meersseman, *A new geometric construction of compact complex manifolds in any dimension*, *Math. Ann.* **317**(1) (2000), 79–115, DOI 10.1007/s002080050360. MR1760670
- [12] Laurent Meersseman and Alberto Verjovsky, *Holomorphic principal bundles over projective toric varieties*, *J. Reine Angew. Math.* **572** (2004), 57–96, DOI 10.1515/crll.2004.054. MR2076120
- [13] Markus J. Pflaum, *Analytic and geometric study of stratified spaces*, Second revised edition, Lecture Notes in Mathematics, vol. 1768, Springer-Verlag, Berlin, 2001. MR1869601
- [14] Harsh V. Pittie, *The Dolbeault-cohomology ring of a compact, even-dimensional Lie group.*, *Proc. Indian Acad. Sci. Math. Sci.* **98** (1988), 117–152, DOI 10.1007/BF02863632. MR0994129
- [15] H. Samelson, *A class of complex-analytic manifolds*, *Portugal. Math.* **12** (1953), 129–132. MR59287
- [16] Hsien-Chung Wang, *Closed manifolds with homogeneous complex structure*, *Amer. J. Math.* **76** (1954), 1–32, DOI 10.2307/2372397. MR66011

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, GRADUATE SCHOOL OF SCIENCE AND ENGINEERING,
KAGOSHIMA UNIVERSITY

Email address: ishida@sci.kagoshima-u.ac.jp

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, OSAKA, JAPAN.

Email address: kasuya@math.sci.osaka-u.ac.jp