

Enlargeable Foliations and the Monodromy Groupoid

Guangxiang Su and Zelin Yi

Abstract

Let M be a spin manifold, the Dirac operator with coefficient in the universal flat Hilbert $C^*\pi_1(M)$ -module determines a "Rosenberg index element" which, according to B.Hanke and T.Schick, subsumes the enlargeability obstruction of positive scalar curvature on M . In this note, we generalize this result to the case of spin foliation. More precisely, given a foliation (M, F) with F spin, we shall define a foliation version of "Rosenberg index element" and prove that it is nonzero at the presence of enlargeability of (M, F) .

1 Introduction

If M is an even dimensional spin manifold with the Dirac operator

$$D : C_c^\infty(M, S^+) \rightarrow C_c^\infty(M, S^-),$$

according to [MF79], the Dirac-type operator twisted by the canonical flat $C^*\pi_1(M)$ -bundle

$$(1.1) \quad \widetilde{M} \times_{\pi_1} C^*\pi_1$$

determines an element $[\alpha(M)]$ (we shall simple write $[\alpha]$ when there is no confusion) in $K_0(C^*\pi_1)$ which is usually called the Rosenberg index element. In fact, if M is of odd dimensional, by replacing M with $M \times S^1$, $[\alpha] \in K_1(C^*\pi_1)$ can also be defined. The main result of [HS06] is the following

1.1 Theorem ([HS06, Prop 4.2]). *If M is a compactly enlargeable spin manifold, then $[\alpha] \neq 0$ in $K_*(C^*\pi_1)$ where $*$ depends on the parity of the dimension of M .*

Let us summarize the main idea of [HS06] as follows: If M is compactly enlargeable, there exist a sequence of almost flat vector bundles E_i of dimension d_i over M (almost flat means the norms of curvatures of E_i 's converge to zero as $i \rightarrow \infty$). Moreover, the sequence E_i can be chosen such that all Chern classes vanish except the top degree part: $c_{\text{top}}(E_i) \neq 0$. Let P_i be the principal frame bundle of E_i , \mathcal{K} be the C^* -algebra of compact operators. Unitary matrices act on \mathcal{K} by the inclusion $U(d_i) \hookrightarrow \mathcal{K}$. The associated product

$$(1.2) \quad P_i \times_{U(d_i)} \mathcal{K}$$

is a Hilbert \mathcal{K} -module bundle.

1.2 Definition. Let A be the C^* -algebra of bounded sequence of compact operators. Namely

$$A = \left\{ (a_i) \in \prod_{\mathbb{N}} \mathcal{K} : \sup_{i \in \mathbb{N}} \|a_i\| < \infty \right\}.$$

Let $A_i \subset A$ be the subalgebra of sequences such that all but the i -th component vanish. It is clear that we have $A_i \cong \mathcal{K}$ for all $i \in \mathbb{N}$. Let $A' \subset A$ be the subalgebra consisting of sequences that converge to zero. In other words, A' is the closure of $\bigoplus \mathcal{K} \subset A$. Let Q be the quotient C^* -algebra A/A' .

Notice that A, A', Q are all nuclear C^* -algebras.

Thanks to the boundedness of curvature of E_i , the sequence of Hilbert module bundles $P_i \times_{U(d_i)} \mathcal{K}$ can be assembled into a Hilbert A -module bundle V . The almost flatness of E_i is reflected in the fact that the curvature of V is endomorphisms of A which take value in $\text{hom}(A, A')$. Therefore, V can be reduced into a genuinely flat Hilbert Q -module bundle $W = V/V \cdot A'$. Thanks to the flatness of W , there is a holonomy representation of the fundamental group $\pi_1(M)$ and correspondingly, a C^* -algebra homomorphism $C^*\pi_1 \rightarrow Q$. To detect the non-vanishing of $[\alpha]$, it is enough to show the non-vanishing of its image under the map

$$(1.3) \quad K_0(C^*\pi_1) \rightarrow K_0(Q).$$

It is known ([HS06, Prop 3.6]) that the K -theory of Q is explicitly computable as a quotient of $\prod \mathbb{Z}$, and the i -th argument of $[\alpha]$ in $K_0(Q)$ is computed as the index of the Dirac-type operator twisted by E_i . The non-vanishing of $[\alpha]$ then follows from the non-vanishing of the top degree Chern class and the Atiyah-Singer index theorem.

In this paper, we shall generalize the result to the case of compactly enlargeable foliations by following the same path. If (M, F) is a compactly enlargeable foliation with F spin. Let G_M and G_H be the monodromy groupoid and the holonomy groupoid of (M, F) respectively. The leafwise Dirac operator

$$D : C^\infty(M, S^+(F)) \rightarrow C^\infty(M, S^-(F))$$

defines a K -theory element $[\alpha(M, F)]$ (we shall simply write $[\alpha]$ when there is no confusion) in $K_0(C^*G_M)$. We shall prove in this paper that the enlargeability of M implies that $[\alpha] \neq 0$ in $K_0(C^*G_M)$.

The enlargeability condition gives a sequence of leafwise almost flat vector bundles $\{E_i\}$ of dimension d_i whose Chern classes vanish except the top degree part. Then the sequence of principal frame bundles and their associated product with compact operators can be defined in the same way as in (1.2). Again, the sequence $\{E_i\}$ can be assembled into a Hilbert A -module bundle V , and the almost leafwise flatness will be reflected in a genuinely leafwise flat Hilbert Q -module bundle $W = V/V \cdot A'$. However, to the best of the authors' knowledge, there is no characterization of leafwise flat vector bundle in the form of (1.1). To find the counterpart of (1.3), we shall make use of basic KK -theory.

The foliation counterpart of universal cover and fundamental group is the monodromy groupoid G_M . If W is a leafwise flat Hilbert Q -module bundle over M , the space of smooth compactly supported sections of the pull back bundle $r^*W \rightarrow G_M$ can be completed into a Hilbert $C^*G_M \otimes Q$ -module \mathcal{E}_Q (Thanks to the nuclearity of Q , there is a canonical choice of tensor product C^* -norm). It can be shown, due to the leafwise flatness, this module also has a left C^*G_M -action which, together with the zero operator, determines a KK -theory element in

$$(1.4) \quad KK(C^*G_M, C^*G_M \otimes Q)$$

which play the role of the map (1.3). Again, due to the nuclearity of A, A' and Q , the sequence of tensor product

$$0 \rightarrow C^*G_H \otimes A' \rightarrow C^*G_H \otimes A \rightarrow C^*G_H \otimes Q \rightarrow 0$$

is exact and induces the following exact sequence at the level of K-theory

$$(1.5) \quad K_0(C^*G_H \otimes A') \rightarrow K_0(C^*G_H \otimes A) \rightarrow K_0(C^*G_H \otimes Q).$$

The image of $[\alpha]$ under the composition

$$K_0(C^*G_M) \rightarrow K_0(C^*G_M \otimes Q) \rightarrow K_0(C^*G_H \otimes Q),$$

where the first arrow is given by Kasparov product with the KK-element (1.4) and the second map is given by the canonical map between full groupoid C*-algebra and the reduced groupoid C* algebra, is contained in the range of the second map of (1.5). This image can be realized as index of the Dirac type operators D_V twisted by the Hilbert A-module bundle V . It suffices to show that this index is not an element of $K_0(C^*G_H \otimes A')$. Indeed, Let p_i be the projection $A \rightarrow \mathcal{K}$ into the i -th component, it induces a map at the level of K-theory

$$K_0(C^*G_H \otimes A) \rightarrow K_0(C^*G_H).$$

Then the image of D_V under this map is precisely the longitudinal index of the leafwise Dirac type operator twisted by the vector bundle E_i . The non-vanishing of $[\alpha]$ is then a consequence of the non-vanishing of the top degree Chern class.

This paper is organized as follows: in section 2, we briefly recall the definition of monodromy groupoids and holonomy groupoids of a foliated manifold; in section 3, we review the notion of Haar system on Lie groupoids and the construction of full and reduced groupoid C*-algebras; in section 4, under the assumption that (M, F) is a foliation with F spin and even dimensional, we define the foliation counterpart of Rosenberg index $[\alpha] \in K_0(C^*G_M)$ and relate it to the longitudinal index element; in section 5, we construct the Hilbert module \mathcal{E}_Q at the presence of a leafwise flat Hilbert Q -module bundle. This Hilbert module \mathcal{E}_Q will later determines a KK-theory element which play the role of (1.3); in section 6, we write down the definition of the genuinely leafwise flat Hilbert Q -module bundle out of the enlargeability of (M, F) and prove the non-vanishing of $[\alpha]$; in the final section, we deal with odd dimensional F . We define $[\alpha] \in K_1(C^*G_M)$ and show how to reduce the non-vanishing problem to the even dimensional case.

2 Monodromy groupoids and holonomy groupoids

Let (M, F) be a compact foliation, we shall denote the monodromy groupoid by G_M and the holonomy groupoid by G_H . The unit space of G_H is the compact manifold M , the morphism space is the set of holonomy classes of curves along leaves of (M, F) . The range map and the source map $r, s : G_H \rightarrow M$ are given by sending curves to their terminal and initial points respectively. Groupoid multiplications are given by concatenation of curves.

2.1 Proposition ([MM03, Prop 5.6]). *The morphism space of holonomy groupoid G_H has a manifold structure.*

Proof. Let $\gamma \in G_H$ be some curves in a leaf of the foliation (M, F) . We shall construct an open neighborhood of γ which is homeomorphic to some Euclidean space.

Assume that $r(\gamma) = x$ and $s(\gamma) = y$. Pick local foliation charts $x \in U = T_1 \times L_1 \rightarrow \mathbb{R}^p \times \mathbb{R}^q$ with $x = (x_T, x_L) \in U$ and $y \in V = T_2 \times L_2 \rightarrow \mathbb{R}^p \times \mathbb{R}^q$ with $y = (y_T, y_L) \in V$. If we pick two foliation charts small enough, there is a smooth map

$$(2.1) \quad H : T_1 \times [0, 1] \rightarrow T_2$$

such that $H(x_T, t) = \gamma(t)$ and $H(*, t)$ is a curve within some leaves connecting $H(*, 0) \in T_1$ and $H(*, 1) \in T_2$. Now we can define a map

$$(2.2) \quad T_1 \times L_1 \times L_2 \rightarrow \text{Hol}(M, F)$$

which assign $(a, b, c) \in T_1 \times L_1 \times L_2$ the curve $\tau \circ H(a, t) \circ \eta$ where τ is any curve connecting (a, b) to (a, x_L) in U , H is the smooth map described in (2.1) with $H(a, 0) = (a, x_L)$ and η is any curve connecting $H(a, 1)$ with (y_T, c) in V . The map (2.2) is well-defined since the holonomy class of $\tau \circ H(a, t) \circ \eta$ is independent of the choice of τ, H, η . It is also clear that (2.2) is injective, and form a topological basis. In this way, we define the local Euclidean structure, and hence, a manifold structure of G_H . \square

2.2 Remark. As for the monodromy groupoid G_M , the unit space is given by M , the morphism space is the set of homotopy classes of curves along leaves of (M, F) , the manifold structure and the source and range maps are given in a similar way.

The source fibers of G_H and G_M over $x \in M$ is the holonomy cover and universal cover of the leaf passing through $x \in M$.

2.3 Example. If $F = TM$, the holonomy groupoid degenerates into the pair groupoid, namely $G_H = M \times M$. Under the same assumption $F = TM$, the morphism space of the monodromy groupoid is given by the space of homotopy classes of all curves in M . In this particular case, G_M is usually called fundamental groupoid and denoted by $\Pi(M) \rightrightarrows M$. It can be shown that the fundamental groupoid is Morita equivalent to fundamental group taken as groupoid over a single point. Hence, their corresponding groupoid C^* -algebras are Morita equivalent.

3 Groupoid C^* -algebras

Parallel to the notion of Haar measures on locally compact topological groups, there is a notion of Haar systems on Lie groupoids.

3.1 Definition. Let G be a Lie groupoid, a family of measures $\{\mu_x\}_{x \in G^0}$ is called a Haar system on G if

1. The measure μ_x is supported on the source fiber G_x ;
2. For any smooth compactly supported function f on G , the function on the unit space G^0 given by the assignment

$$x \mapsto \int_{G_x} f(\gamma) d\mu_x(\gamma)$$

is smooth;

3. Let $\eta \in G$, f be any smooth compactly supported function on G , then

$$\int_{G_{s(\eta)}} f(\gamma) d\mu_{s(\eta)}(\gamma) = \int_{G_{r(\eta)}} f(\gamma \circ \eta) d\mu_{r(\eta)}(\gamma).$$

The above family of measures is sometimes referred to as right invariant Haar system. Left invariant Haar system $\{\mu^x\}_{x \in G^0}$ can be defined in a similar way where we replace the source fibers G_x with the range fibers G^x .

3.2 Example. Recall that (M, F) is a compact foliation. Fix a metric on F , then there is an induced measure $\{\mu_x\}_{x \in M}$ on each leaf $L_x \subset M$. The leafwise measures, in turn, determine measures $\{\mu_x^H\}$ on their holonomy covers $G_{H,x}$ and measures $\{\mu_x^M\}$ on universal covers $G_{M,x}$ accordingly. It can be checked that these measures form Haar systems on G_H and G_M respectively.

At the presence of a Haar system $\{\mu_x\}_{x \in G^0}$, the space of compactly supported smooth functions on groupoid G can be made into an algebra. Let $f, g \in C_c^\infty(G)$, the multiplication $f * g$ is given by

$$f * g(\gamma) = \int_{\gamma_1 \in G_s(\gamma)} f(\gamma \circ \gamma_1^{-1}) g(\gamma_1) d\mu_s(\gamma)(\gamma_1).$$

3.3 Definition. Let $f \in C_c^\infty(G)$ and define

$$\|f\|_I = \sup_{x \in G^0} \left\{ \int_{G_x} |f(\gamma)| d\mu_x(\gamma), \int_{G_x} |f(\gamma^{-1})| d\mu_x(\gamma) \right\}.$$

It is easy to check that $\|\cdot\|_I$ is a norm. We shall say a representation $\varphi : C_c^\infty(G) \rightarrow \mathcal{B}(H_\varphi)$ is bounded if it satisfies

$$\|\varphi(f)\|_{\mathcal{B}(H_\varphi)} \leq \|f\|_I$$

for all $f \in C_c^\infty(G)$. The full groupoid C^* -algebra is the completion of $C_c^\infty(G)$ with respect to the norm

$$\sup_{\varphi} \|\varphi(f)\|_{\mathcal{B}(H_\varphi)}$$

where φ ranges over all bounded representations of $C_c^\infty(G)$ such that the span of $\{\varphi(f)\xi \mid f \in C_c^\infty(G), \xi \in \mathcal{H}_\varphi\}$ is a dense subspace of \mathcal{H}_φ . The full groupoid C^* -algebra is usually denoted by C^*G .

Analogous to the fact that the holonomy group at a fixed point is a quotient of the fundamental group of the leaf passing through the fixed point, there is a canonical quotient map $\pi : G_M \rightarrow G_H$ which sends the universal cover of a leaf to its holonomy cover. Therefore, G_M acts on G_H in a natural way. There is a left $C_c^\infty(G_M)$ -action on $C_c^\infty(G_H)$ which is given by the following formula

$$(3.1) \quad f.g(\gamma) = \int_{\gamma_1 \in G_{M,s(\gamma)}} f(\gamma \circ \gamma_1^{-1}) g_0(\gamma_1) d\mu_{s(\gamma)}(\gamma_1)$$

where f, g are compactly supported smooth functions on G_M and G_H respectively, $\gamma_1, \gamma_2 \in G_M$ and g_0 is the smooth function on G_M given by $g_0(\gamma) = g(\pi(\gamma))$.

3.4 Proposition. *The action (3.1) extends to a C^* -algebras homomorphism $C^*G_M \rightarrow C^*G_H$.*

Proof. It suffices to verify the inequality

$$\|f.g\|_{C^*G_H} \leq \|f\|_I \cdot \|g\|_{C^*G_H}.$$

Let $\Phi(f)$ be the function on G_H given by $\Phi(f)(\gamma) = \sum_{\pi(\eta)=\gamma} f(\eta)$. It is clear that $\|f.g\|_{C^*G_H} \leq \|\Phi(f)\|_{C^*G_H} \cdot \|g\|_{C^*G_H}$. It is then straightforward to check that $\|\Phi(f)\|_{C^*G_H} \leq \|\Phi(f)\|_I \leq \|f\|_I$ which completes the proof. \square

Within the set of bounded representations of groupoid algebra $C_c^\infty(G)$ there is a distinguished one called regular representation which is described as follows. Let $\{\mu_x\}$ be a right Haar system on the groupoid G . For any $x \in G^0$, the groupoid algebra $C_c^\infty(G)$ acts on the Hilbert space $L^2(G_x, \mu_x)$ as follows

$$\pi_x(f)\xi(\gamma) = \int_{\eta \in G_x} f(\gamma \circ \eta^{-1}) \xi(\eta) d\mu_x(\eta).$$

It is easy to check that this is a bounded representation. The completion of $C_c^\infty(G)$ with respect to the norm

$$\|f\| = \sup_{x \in G^0} \|\pi_x(f)\|$$

is denoted by C_r^*G and called the reduced groupoid C^* -algebra. By definition $\|\cdot\|_{C_r^*G} \leq \|\cdot\|_{C^*G}$, so there is a canonical map $C^*G \rightarrow C_r^*G$.

4 Rosenberg Index

From this section on, except the last section, we shall assume that $F \rightarrow M$ is a spin vector bundle of even dimensional with spinor given by $S = S^+(F) \oplus S^-(F)$. Let D be a leafwise Dirac operator acting on S , which means the following

- $D : C^\infty(M, S) \rightarrow C^\infty(M, S)$ is a usual differential operator;
- For any smooth section ξ of $S \rightarrow M$ and any leaf L of (M, F) , the restriction $D\xi|_L$ only depends on the restriction $\xi|_L$;
- For any leaf L , $D|_L : C_c^\infty(L, S) \rightarrow C_c^\infty(L, S)$ is the classical Dirac operator on L .

The notion of leafwise Dirac type operator or more generally, the notion of leafwise elliptic differential operator can be defined in a similar way(see [Con94, Ch 2, Sec 9]).

4.1 Proposition. *The space $C_c^\infty(G_M, r^*S)$ can be completed into a Hilbert C^*G_M -module which will be denoted by \mathcal{E} .*

Proof. Let $\varphi, \psi \in C_c^\infty(G_M, r^*S)$, $\gamma \in G_M$ the formula

$$\langle \varphi, \psi \rangle(\gamma) = \int_{\gamma_1 \in G_s(\gamma)} \langle \varphi(\gamma_1 \circ \gamma_1^{-1}), \psi(\gamma_1) \rangle d\mu_s(\gamma)(\gamma_1)$$

defines a C^*G_M -valued inner product on $C_c^\infty(G_M, r^*S)$. Let $f \in C_c^\infty(G_M)$, it acts on $C_c^\infty(G_M, r^*S)$ by

$$\varphi.f(\gamma) = \int_{\gamma_1 \in G_s(\gamma)} \varphi(\gamma \circ \gamma_1^{-1})f(\gamma_1) d\mu_s(\gamma)(\gamma_1).$$

It is easily checked that this inner product satisfies the pre-Hilbert module condition and $\langle \varphi, \varphi \rangle = 0$ implies $\varphi = 0$. The completion of $C_c^\infty(G_M, r^*S)$ under the norm

$$\|\varphi\|_{\mathcal{E}}^2 = \|\langle \varphi, \varphi \rangle\|_{C_{\max}^*G_M}$$

is a Hilbert C^*G_M -module. □

Recall that in [HR00, Chp 10], a continuous function $f : \mathbb{R} \rightarrow [-1, 1]$ is called normalizing if

- f is odd;
- $f(c) \geq 0$ if $c \geq 0$;
- $\lim_{c \rightarrow \pm\infty} f(c) \rightarrow \pm 1$.

According to the theory of [Vas06] and [Kuc02], the leafwise elliptic differential operator D defines a regular self-adjoint operator on Hilbert module \mathcal{E} . Moreover, if $f \in C_0(\mathbb{R})$ then $f(D)$ is a compact operator on \mathcal{E} . Therefore, $(\mathcal{E}, f(D))$ form a Kasparov module and determines an element $[\alpha]$ in $K_0(C^*G_M)$ for any normalizing function f . We shall call this element the Rosenberg index of D .

4.2 Proposition. *If F is spin and (M, F) admits leafwise positive scalar curvature, then $[\alpha] = 0$ as a K -theory element in $K_0(C^*G_M)$.*

Proof. By Lichnerowicz formula, if (M, F) has leafwise positive scalar curvature, the leafwise Dirac D is invertible. It has a spectrum gap around $0 \in \mathbb{R}$. We can choose the normalizing function f such that $f^2 = 1$ on spectrum of D . Under this circumstances, the Kasparov module $(\mathcal{E}, f(D))$ is degenerated. \square

Now, let B be a C^* -algebra, E be a Hilbert B -module bundle over M . Let D_E denote the leafwise Dirac type operator twisted by E . We have the following easy fact.

4.3 Proposition. *The space $C_c^\infty(G_M, r^*S \otimes r^*E)$ can be completed into a Hilbert $C^*G_M \otimes_{\min} B$ -module \mathcal{E}_B such that the pair $(\mathcal{E}_B, f(D_E))$ form a Kasparov module and determines an element in $K_0(C^*G_M \otimes_{\min} B)$. This element will be called twisted Rosenberg index. \square*

4.4 Remark. In [CS84], the authors define the longitudinal index as an element of $K_0(C_r^*G_H)$. Following their method, we set the Rosenberg index to live in $K_0(C^*G_M)$. In fact, under the map

$$(4.1) \quad C^*G_M \rightarrow C^*G_H \rightarrow C_r^*G_H$$

the Rosenberg index defined above is mapped to the longitudinal index.

5 The Hilbert module \mathcal{E}_Q

The basic theory of Hilbert C^* -module and Hilbert C^* -module bundle can be found in [Sch05]. Let W be a leafwise flat Hilbert Q -module bundle over M .

The space $C_c^\infty(G_M, r^*W)$ has a $C_c^\infty(G_M) \otimes Q \subset C^*G_M \otimes Q$ valued inner product

$$\langle \varphi, \psi \rangle(\gamma) = \int_{\gamma_1 \in G_s(\gamma)} \langle \varphi(\gamma_1 \circ \gamma^{-1}), \psi(\gamma_1) \rangle d\mu_s(\gamma)(\gamma_1).$$

Let \mathcal{E}_Q be the completion of $C_c^\infty(G_M, r^*W)$ under the norm $\|\varphi\| = \|\langle \varphi, \varphi \rangle\|^{1/2}$. The space $C_c^\infty(G_M, r^*W)$ has an obvious right $C_c^\infty(G_M) \otimes Q$ action which is given by

$$(5.1) \quad \varphi.f(\gamma) = \int_{\gamma_1 \in G_s(\gamma)} \varphi(\gamma \circ \gamma_1^{-1})f(\gamma_1) d\mu_s(\gamma)(\gamma_1)$$

where $\varphi \in C_c^\infty(G_M, r^*W)$ and $f \in C_c^\infty(G_M) \otimes Q$.

5.1 Proposition. *The action (5.1) extends to a right $C^*G_M \otimes Q$ action on \mathcal{E}_Q . As a consequence, \mathcal{E}_Q has a Hilbert $C^*G_M \otimes Q$ -module structure.*

Proof. It is straightforward to check that

$$\langle \varphi.f, \psi.g \rangle = f^* \langle \varphi, \psi \rangle g$$

for all $\varphi, \psi \in C_c^\infty(G_M, r^*W)$ and $f, g \in C_c^\infty(G_M) \otimes Q$. Therefore, we have

$$\|\varphi.f\|^2 = \|\langle \varphi.f, \varphi.f \rangle\| \leq \|f\|^2 \|\varphi\|^2$$

which proves the statement. \square

There is a left $C_0(M)$ -action on $C_c^\infty(G_M, r^*W)$ which is given by

$$(5.2) \quad \mathfrak{h} \cdot \varphi(\gamma) = \mathfrak{h}(r(\gamma)) \cdot \varphi(\gamma)$$

for all $\mathfrak{h} \in C_0(M)$ and $\varphi \in C_c^\infty(G_M, r^*W)$.

5.2 Proposition. *For any $\mathfrak{h} \in C_0(M)$ and $\varphi \in C_c^\infty(G_M, r^*W)$, we have*

$$\|\mathfrak{h} \cdot \varphi\| \leq \|\mathfrak{h}\| \cdot \|\varphi\|$$

where $\|\mathfrak{h}\|$ is the sup-norm of \mathfrak{h} in $C_0(M)$ and $\|\varphi\|$ is the norm in \mathcal{E}_Q .

Proof. We shall use the estimate in [Ren80, Lem 1.1.13]. Let $\mathfrak{k} \in C_0(M)$ be the function defined by

$$\mathfrak{k}(m) = (\|\mathfrak{h}\|^2 - |\mathfrak{h}(m)|^2)^{1/2}.$$

Then, it is easy to check the following

$$\begin{aligned} \|\mathfrak{h} \cdot \varphi\|^2 &= \|\langle \mathfrak{h} \cdot \varphi, \mathfrak{h} \cdot \varphi \rangle\| \\ &= \left| \|\mathfrak{h}\|^2 \langle \varphi, \varphi \rangle - \langle \mathfrak{k} \cdot \varphi, \mathfrak{k} \cdot \varphi \rangle \right| \\ &\leq \|\mathfrak{h}\|^2 \|\langle \varphi, \varphi \rangle\| \end{aligned}$$

which completes the proof. \square

5.3 Corollary. [Ren80, Prop 2.1.14] *The action (5.2) extends to a *-homomorphism $C_0(M) \rightarrow \mathcal{L}(\mathcal{E}_Q)$.*

Proof. Thanks to the above Proposition, the action extends to the Hilbert module \mathcal{E}_Q . It is a matter of direct calculation to check that it preserves the *-operation. \square

We shall show, in the rest of this section, the Hilbert module \mathcal{E}_Q determines a KK-theory element in $\text{KK}(C^*G_M, C^*G_M \otimes Q)$. There is a left $C_c^\infty(G_M)$ action on $C_c^\infty(G_M, r^*W)$ which is given by

$$(5.3) \quad \mathfrak{f} \cdot \varphi(\gamma) = \int_{\gamma_1 \in G_M, s(\gamma_1) = \gamma} \mathfrak{f}(\gamma \circ \gamma_1^{-1})(\gamma \circ \gamma_1^{-1}) \cdot \varphi(\gamma_1) d\mu_{s(\gamma)}(\gamma_1)$$

where $(\gamma \circ \gamma_1^{-1}) \cdot \varphi(\gamma_1)$ is the image of $\varphi(\gamma_1)$ under the parallel translation along the curve $\gamma \circ \gamma_1^{-1}$. Thanks to the flatness of W , this parallel translation is well defined. It is also convenient to have an alternative description of the above action.

Let $\mu = \{\mu_x\}$ be a right invariant Haar system on G_M . The inverse map $\iota : G_M \rightarrow G_M$ induces a left invariant Haar system which we denote by $\tilde{\mu}$. The space $C_c^\infty(G_M)$ can be completed into Hilbert $C_0(M)$ -module in two ways given by two inner products

$$\langle f, g \rangle_s(m) = \int_{s(\gamma)=m} \bar{f}(\gamma) \cdot g(\gamma) d\mu(\gamma)$$

and

$$\langle f, g \rangle_r(m) = \int_{r(\gamma)=m} \bar{f}(\gamma) \cdot g(\gamma) d\tilde{\mu}(\gamma)$$

where $f, g \in C_c^\infty(G_M)$. Following [BHM18], we shall denote the completions by $L^2(G_M, s, \mu)$ and $L^2(G_M, r, \tilde{\mu})$ respectively. According to Coro 5.3, we can form the inner tensor product $L^2(G_M, s, \mu) \otimes \mathcal{E}_Q$ and $L^2(G_M, r, \tilde{\mu}) \otimes \mathcal{E}_Q$. There is an isometry $U : L^2(G, s, \mu) \otimes \mathcal{E}_Q \rightarrow L^2(G, r, \tilde{\mu}) \otimes \mathcal{E}_Q$ given by

$$U(F)(\gamma_1, \gamma_2) = U_{\gamma_1} F(\gamma_1, \gamma_1^{-1} \circ \gamma_2)$$

when $F \in L^2(G, s, \mu) \otimes C_c^\infty(G_M, r^*E)$. Let $f \in C_c^\infty(G_M)$, let $T_f : \mathcal{E}_Q \rightarrow L^2(G, s, \mu) \otimes \mathcal{E}_Q$ denote the Hilbert module map $x \mapsto f \otimes x$ and $T_f^* : L^2(G_M, r, \tilde{\mu}) \otimes \mathcal{E}_Q \rightarrow \mathcal{E}_Q$ be the operator which sends $g \otimes x$ to $\langle f, g \rangle_{r,x}$. Choose $f_1, f_2 \in C_c^\infty(G_M)$ such that $f = \bar{f}_1 \cdot f_2$, where \cdot is the point-wise multiplication. Under this setting, the action (5.3) can be realized as $T_{f_1}^* \mathcal{U} T_{f_2}$.

5.4 Proposition. *The action (5.3) extends to a *-homomorphism $C^*G_M \rightarrow \mathcal{L}(\mathcal{E}_Q)$ whose image is contained in the algebra of compact operators on \mathcal{E}_Q .*

Proof. According to the above discussion,

$$(5.4) \quad \|f \cdot \varphi\| \leq \|T_{f_1}^*\| \cdot \|T_{f_2}\| \cdot \|\varphi\|$$

where we omit the norm of \mathcal{U} since it is an isometry. It is easy to check that $\|T_f\| = \|f\|_{L^2(G_M, s, \mu)}$ and $\|T_f^*\| = \|f\|_{L^2(G_M, r, \tilde{\mu})}$. According to the definition $\|f\|_I = \max\{\|T_f\|, \|T_f^*\|\}$. Therefore, the inequality (5.4) becomes

$$\|f \cdot \varphi\| \leq \|f\|_I \cdot \|\varphi\|$$

which completes the extension part of the proof.

As for the compactness, let $\varphi, \psi, \zeta \in C_c^\infty(G_M, r^*W)$, $\gamma \in G_M$, the rank one operator $\theta_{\varphi, \psi} \in C_c^\infty(G_M, s^*W^* \otimes r^*W)$ is given by

$$\begin{aligned} \theta_{\varphi, \psi}(\zeta)(\gamma) &= \varphi \langle \psi, \zeta \rangle(\gamma) \\ &= \int_{\eta \in G_s(\gamma)} \int_{\eta_1 \in G_s(\gamma)} \varphi(\gamma \circ \eta^{-1}) \langle \psi(\eta_1 \circ \eta^{-1}), \zeta(\eta_1) \rangle d\eta_1 d\eta \\ &= \int_{\eta_2 \in G_s(\gamma)} F(\gamma \circ \eta_2^{-1}) \zeta(\eta_2) d\eta_2 \end{aligned}$$

where

$$F(\gamma) = \int_{\eta \in G_s(\gamma)} \varphi(\gamma \circ \eta^{-1}) \psi^*(\eta^{-1}) d\eta.$$

Here we use $\psi^*(\gamma) \in W^*$ to denote the dual element of $\psi(\gamma) \in W$. It suffices to show the C^* -algebra $\mathcal{K}_{C^*G_M}(\mathcal{E})$ contains $C_c^\infty(G_M, s^*W^* \otimes r^*W)$. Let ψ_0^* be a section of $W^* \rightarrow M$, φ_0 be a section of $W \rightarrow M$, $f \in C_c^\infty(G)$. Elements in $C_c^\infty(G_M, s^*W^* \otimes r^*W)$ can be written as span of

$$(5.5) \quad \psi_0^* \otimes f \otimes \varphi_0$$

where $\psi_0^* \otimes f \otimes \varphi_0(\gamma) = \psi_0^*(s(\gamma))f(\gamma)\varphi_0(r(\gamma))$. It suffices to show elements of the form (5.5) is a compact operator.

If there are $f_1, f_2 \in C_c^\infty(G)$ such that $f_1 * f_2 = f$, we shall pick $\varphi(\gamma) = \varphi_0(r(\gamma))f_2(\gamma)$ and $\psi^*(\gamma^{-1}) = \psi_0^*(s(\gamma))f_1(\gamma)$, it is then easy to check that $\theta_{\varphi, \psi} = \psi_0^* \otimes f \otimes \varphi_0$. In general, f can be approximated by span of $f_1 * f_2$, the result follows. \square

5.5 Corollary. *The Hilbert module \mathcal{E}_Q together with the zero operator $(\mathcal{E}_Q, 0)$ form a Kasparov module which defines an element in $\text{KK}(C^*G_M, C^*G_M \otimes Q)$. \square*

6 Enlargeable foliation

6.1 Definition. A foliation (M, F) is compactly enlargeable if there is $C > 0$ such that for any $\varepsilon > 0$, there is a compact covering \widetilde{M} of M and a smooth map

$$f : \widetilde{M} \rightarrow S^n$$

with

- $|f_*X| \leq \varepsilon|X|$ for all $X \in C^\infty(\widetilde{M}, \widetilde{F})$, where \widetilde{F} is the lifting of F to \widetilde{M} ;
- $|f_*X| \leq C \cdot |X|$ for all $X \in C^\infty(\widetilde{M}, \widetilde{TM})$;
- f has nonzero degree.

6.2 Remark. Notice that our notion of enlargeable foliation is in between that of [Zha20] and [BH19], where [Zha20] only requires enlargeability in the leaf direction and [BH19] requires enlargeability in all directions.

Pick a complex vector bundle E_0 over sphere S^n such that all its Chern classes vanish except the top-degree one $c_n(E_0) \neq 0$. Let \widetilde{M}_i be the compact cover with covering group G and the constant $\varepsilon = 1/i$, the pull back bundle f^*E_0 can be extended to a G -equivariant bundle

$$\bigoplus_{g \in G} g^*(f^*E_0) \rightarrow \widetilde{M}_i$$

which can be reduced to a vector bundle E_i over M . As a result all Chern classes of E_i vanish except the top-degree one $c_n(E_i) \neq 0$. We shall denote by $P_i \rightarrow M$ the frame bundle of E_i which are by themselves principal $U(d_i)$ bundles. They are equipped with natural connections whose curvatures tend to zero as $i \rightarrow \infty$.

We shall consider the family of Hilbert \mathcal{K} -module bundles

$$(6.1) \quad V_i = P_i \times_{U(d_i)} \mathcal{K}.$$

where $U(d_i)$ acts on \mathcal{K} by matrix multiplications. Let q_i denote the image of $1 \in U(d_i)$ on \mathcal{K} . We shall briefly explain how they can be assembled into a leafwise flat Hilbert Q -module (see [HS06, Sec 2] for a detailed construction). Indeed, let $\{U_\alpha\}$ be an open cover of M over which each V_i is trivialisable and each U_α is homeomorphic to an unit open disc $[0, 1]^n$. We can choose local trivialisations

$$(6.2) \quad \psi_{\alpha,i} : V_i|_{U_\alpha} \rightarrow U_\alpha \times q_i \mathcal{K}$$

as in [HS06, Sec 2] such that

$$(6.3) \quad \nabla_{\frac{\partial}{\partial x_k}}^i s = 0$$

if s is a smooth section which is constant, under the trivialisations, in $[0, 1]^k \times \{0\} \times \dots \times \{0\}$ the first k variable of U_α . Here ∇^i is the connection on V_i .

The corresponding transition functions is denoted by

$$\varphi_{\alpha,\beta,i} : U_\alpha \cap U_\beta \rightarrow \text{End}(q_i \mathcal{K}) \cong q_i \mathcal{K} q_i$$

Since the norm of curvature of V_i is universally bounded with respect to $i \in \mathbb{N}$. According to [HS06, Lem 2.3, Lem 2.5 and Prop 2.6], $\varphi_{\alpha,\beta,i}$ is a Lipschitz function with Lipschitz constant independent of i . Therefore, the transition functions can be assembled into $\varphi_{\alpha,\beta} : U_\alpha \cap U_\beta \rightarrow qAq$ and determines a Lipschitz Hilbert A -module bundle over M which can be approximated by a smooth Hilbert A -module bundle V over M .

The properties of bundle A is summarized in the following

6.3 Proposition. *There is a Hilbert A -module bundle V over M such that*

- V_i , defined in (6.1), is isomorphic to $V \cdot A_i$ as Hilbert \mathcal{K} -module bundle;
- The connection of V preserves subbundle V_i ;
- The leafwise curvature take value in $\text{hom}(A, A')$.

Therefore the bundle $W = V/V \cdot A'$ is a leafwise flat Hilbert Q -module bundle which, according to Coro 5.5, determines an element in $\text{KK}(C^*G_M, C^*G_M \otimes Q)$. The KK-element induces $(\phi_1)_* : K_0(C^*G_M) \rightarrow K_0(C^*G_M \otimes Q)$. The above procedure can be replicated if we start with a sequence of trivial principal bundles $\{P'_i\}$ with $P'_i = M \times U(d_i)$. We shall get a new KK-theory element in $\text{KK}(C^*G_M, C^*G_M \otimes Q)$ and corresponding $(\phi_2)_* : K_0(C^*G_M) \rightarrow K_0(C^*G_M \otimes Q)$. Let

$$(6.4) \quad \phi_* = (\phi_1)_* - (\phi_2)_*.$$

6.4 Proposition. *Let $[D_W] \in K_0(C^*G_M \otimes Q)$ denote the image of $[\alpha] \in K_0(C^*G_M)$ under the map $(\phi_1)_* : K_0(C^*G_M) \rightarrow K_0(C^*G_M \otimes Q)$. Then $[D_W]$ coincides with the Rosenberg index twisted by the leafwise flat Hilbert Q -module bundle W .*

Proof. $[\alpha]$ is given by the Kasparov module $(\mathcal{E}, f(D))$ while the KK-theory element is given by the Kasparov module $(\mathcal{E}_Q, 0)$. Their Kasparov product is given by the pair $(\mathcal{E} \otimes_{C^*G_M} \mathcal{E}_Q, f(D) \otimes 1)$. It can be shown that the inner tensor product $\mathcal{E} \otimes_{C^*G_M} \mathcal{E}_Q$ is the completion of $C_c^\infty(G_M, r^*S \otimes r^*W)$ and the operator $f(D) \otimes 1$ is precisely $f(D_W)$. \square

By the same reasoning, the image of $[\alpha]$ under the map $(\phi_2)_*$ is the Rosenberg index $[D_{qQ}]$ twisted by the trivial bundle $M \times qQ$. Let $\pi : A \rightarrow Q$ be the canonical projection, it induces $\pi_* : K_0(C^*G_M \otimes A) \rightarrow K_0(C^*G_M \otimes Q)$. Let $[D_V], [D_{qA}] \in K_0(C^*G_M \otimes A)$ be the elements defined by the leafwise Dirac-type operators twisted by the non-flat bundle V and the trivial bundle $M \times qA$ respectively. Then it is straightforward to verify that we have $\pi_*[D_V] = [D_W]$ and $\pi_*[D_{qA}] = [D_{qQ}]$. (see also [HS06, Lem 3.1]).

Consider the following composition:

$$(6.5) \quad K_0(C^*G_M \otimes A) \rightarrow K_0(C_r^*G_H \otimes A) \rightarrow K_0(C_r^*G_H)$$

where the first arrow is given by (4.1) and the second arrow is given by the homomorphism sending A to its i -th component \mathcal{K} . Since K-theory of C^* -algebras is unchanged by tensoring with \mathcal{K} , the second arrow lands in $K_0(C_r^*G_H)$.

6.5 Proposition. *The image of $[D_V]$ under the map (6.5) is computed by the longitudinal index element corresponds to D_{E_i} .*

Proof. It is a consequence of Rmk 4.4. \square

Since the functor K_0 commutes with direct limit, we have

$$K_0(C_r^*G_H \otimes A') = \bigoplus K_0(C_r^*G_H).$$

6.6 Proposition. *Let $\phi_* : K_0(C^*G_M) \rightarrow K_0(C^*G_M \otimes Q)$ be defined as in (6.4). Then $\phi_*[\alpha] \neq 0$ in $K_0(C^*G_M \otimes Q)$.*

Proof. By Prop 6.4, the image of $[D_V] - [D_{qA}] \in K_0(C^*G_M \otimes A)$ under the map $\pi_* : K_0(C^*G_M \otimes A) \rightarrow K_0(C^*G_M \otimes Q)$ is precisely $\phi_*[\alpha]$. By the exact sequence (1.5), it suffices to show that $[D_V] - [D_{qA}] \in K_0(C^*G_M \otimes A)$ does not come from the image of $K_0(C^*G_M \otimes A')$.

Consider the following commutative diagram

$$\begin{array}{ccc} K_0(C^*G_M \otimes A') & \longrightarrow & K_0(C^*G_M \otimes A) \\ & \searrow & \downarrow \\ & & \prod K_0(C^*G_M) \end{array}$$

where the downward arrows are given by sending A and A' to A_i 's. It then suffices to show the image of $[D_V] - [D_{qA}]$ under the vertical downward arrow has infinitely many nonzero terms.

Indeed, according to Prop 6.5, under the map $K_0(C^*G_M) \rightarrow K_0(C^*_rG_H)$ the i -th component of $[D_V] - [D_{qA}]$ is given by the longitudinal index of the Dirac type operator twisted by the virtual bundle $E_i - \mathbb{C}^{d_i}$. According to Connes[Con86], there is a transverse fundamental class μ such that

$$\mu([D_{E_i - \mathbb{C}^{d_i}}]) = \langle \widehat{A}(F) \text{ch}(E_i - \mathbb{C}^{d_i}), [M] \rangle$$

where $[M]$ is a fundamental class of M . It is clear that the sequence $\mu([D_{E_i - \mathbb{C}^{d_i}}])$ does not converges to zero. This completes the proof. \square

The above Proposition directly implies our main theorem

6.7 Theorem. *If (M, F) is an enlargeable foliation in the sense of Def 6.1 with F spin and even dimensional, then $[\alpha] \neq 0$ in $K_0(C^*G_M)$.*

7 Reduction to the even dimensional case

If the foliation $F \rightarrow M$ is of odd dimensional, the exterior product of vector bundles $F \boxtimes TS^1 \rightarrow M \times S^1$ defines an even dimensional foliation. Moreover, the monodromy groupoid of $(M \times S^1, F \boxtimes TS^1)$ is the direct product of monodromy groupoid of (M, F) and the fundamental groupoid of S^1 . Accordingly, the corresponding maximal groupoid C^* -algebra is $C^*G_M \otimes \mathcal{K} \otimes C^*\mathbb{Z}$ whose K -theory is computed by the universal coefficient theorem:

$$K_*(C^*G_M \otimes \mathcal{K} \otimes C^*\mathbb{Z}) = K_*(C^*G_M) \otimes K_*(C^*\mathbb{Z}).$$

Here \mathcal{K} is the C^* -algebra of compact operators and $C^*\mathbb{Z}$ is the group C^* algebra of \mathbb{Z} . In particular, we have

$$K_0(C^*G_M \otimes \mathcal{K} \otimes C^*\mathbb{Z}) = K_0(C^*G_M) \otimes 1 \oplus K_1(C^*G_M) \otimes e$$

where 1 is the generator of $K_0(C^*\mathbb{Z})$ and e is the generator of $K_1(C^*\mathbb{Z})$. As in [HS06], $[\alpha(M, F)] \in K_1(C^*G_M)$ is defined by requiring

$$[\alpha(M, F)] \otimes e = [\alpha(M \times S^1, F \boxtimes TS^1)] \in K_0(C^*G_M \otimes \mathcal{K} \otimes C^*\mathbb{Z}).$$

7.1 Proposition. *The foliation $(M \times S^1, F \boxtimes TS^1)$ is enlargeable if (M, F) is enlargeable.*

Proof. Assume that (M, F) is enlargeable. Then for any $\varepsilon > 0$ there is compact covering space $\widetilde{M}_\varepsilon \rightarrow M$ and map $f_\varepsilon : \widetilde{M}_\varepsilon \rightarrow S^n$ with the properties given in Def 6.1. Since S^1 is also enlargeable, there is $g_\varepsilon : S^1 \rightarrow S^1$ with the properties of Def 6.1. It is straightforward to check that

$$(f_\varepsilon, g_\varepsilon) : \widetilde{M}_\varepsilon \times S^1 \rightarrow S^{n+1}$$

has the wanted property. \square

According to Thm 6.7 and Prop 7.1, $[\alpha(M \times S^1, F \boxtimes TS^1)]$ is nonzero, so $[\alpha(M, F)]$ is also nonzero.

Acknowledgment

The authors would like to thank Professor Weiping Zhang for kindly suggesting this problem. The authors also want to thank Professor Georges Skandalis for very helpful guidance on groupoid C^* -algebras, in particular for pointing out Prop 3.4.

References

- [BH19] Moulay-Tahar Benameur and James L. Heitsch. Enlargeability, foliations, and positive scalar curvature. *Invent. Math.*, 215(1):367–382, 2019.
- [BHM18] Alcides Buss, Rohit D. Holkar, and Ralf Meyer. A universal property for groupoid C^* -algebras. I. *Proc. Lond. Math. Soc. (3)*, 117(2):345–375, 2018.
- [Con86] A. Connes. Cyclic cohomology and the transverse fundamental class of a foliation. In *Geometric methods in operator algebras (Kyoto, 1983)*, volume 123 of *Pitman Res. Notes Math. Ser.*, pages 52–144. Longman Sci. Tech., Harlow, 1986.
- [Con94] Alain Connes. *Noncommutative geometry*. Academic Press, Inc., San Diego, CA, 1994.
- [CS84] A. Connes and G. Skandalis. The longitudinal index theorem for foliations. *Publ. Res. Inst. Math. Sci.*, 20(6):1139–1183, 1984.
- [HR00] Nigel Higson and John Roe. *Analytic K-homology*. Oxford Mathematical Monographs. Oxford University Press, Oxford, 2000. Oxford Science Publications.
- [HS06] B. Hanke and T. Schick. Enlargeability and index theory. *J. Differential Geom.*, 74(2):293–320, 2006.
- [Kuc02] Dan Kucerovsky. Functional calculus and representations of $C_0(\mathbb{C})$ on a Hilbert module. *Q. J. Math.*, 53(4):467–477, 2002.
- [MF79] A. S. Miščenko and A. T. Fomenko. The index of elliptic operators over C^* -algebras. *Izv. Akad. Nauk SSSR Ser. Mat.*, 43(4):831–859, 1979.
- [MM03] I. Moerdijk and J. Mrcun. *Introduction to foliations and Lie groupoids*, volume 91 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2003.
- [Ren80] Jean Renault. *A groupoid approach to C^* -algebras*, volume 793 of *Lecture Notes in Mathematics*. Springer, Berlin, 1980.
- [Sch05] Thomas Schick. L^2 -index theorems, KK-theory, and connections. *New York J. Math.*, 11:387–443, 2005.
- [Vas06] Stéphane Vassout. Unbounded pseudodifferential calculus on Lie groupoids. *J. Funct. Anal.*, 236(1):161–200, 2006.
- [Zha20] Weiping Zhang. Positive scalar curvature on foliations: the enlargeability. In *Geometric analysis*, volume 333 of *Progr. Math.*, pages 537–544. Birkhäuser/Springer, Cham, [2020] ©2020.

Chern Institute of Mathematics and LPMC, Nankai University, Tianjin 300071, P. R. China.

Email: guangxiangsu@nankai.edu.cn

School of Mathematical Sciences, Tongji University, Shanghai 200092, P. R. China.

Email: zelin@tongji.edu.cn