

CLOSED-FORM FORMULAS AND PROPERTIES OF COEFFICIENTS IN MACLAURIN'S SERIES EXPANSION OF WILF'S FUNCTION

FENG QI AND MARK DANIEL WARD

Dedicated to people facing and battling COVID-19

ABSTRACT. In the paper, with the aid of the Faà di Bruno formula, by virtue of several identities for the Bell polynomials of the second kind, with the help of two combinatorial identities, by means of the (logarithmically) complete monotonicity of generating functions of several integer sequences, and in the light of Wronski's theorem, the authors establish Taylor's series expansions of several functions involving the inverse (hyperbolic) tangent function, find out Maclaurin's series expansion of a complex function posed by Herbert S. Wilf, and analyze some properties, including generating functions, limits, positivity, monotonicity, and logarithmic convexity, of the coefficients in Maclaurin's series expansion of Wilf's function. These coefficients in Maclaurin's series expansion of Wilf's function are closed-form expressions in terms of the Stirling numbers of the second kind. The authors also derive a closed-form formula for a sequence of special values of Gauss' hypergeometric function, discover a closed-form formula for a sequence of special values of the Bell polynomials of the second kind, present several infinite series representations of the circular constant Pi and other sequences, recover an asymptotic rational approximation to the circular constant Pi, and connect several integer sequences by determinants.

CONTENTS

1.	Motivations	1
2.	Preparations and lemmas	3
3.	Taylor's series expansions of the inverse tangent function	6
4.	Maclaurin's series expansions of Wilf's function	13
5.	Series expansions of the inverse hyperbolic tangent function	15
6.	Analysis on coefficients in series expansions of Wilf's function	16
7.	Conclusions	24
8.	Declarations	24
	References	24

1. MOTIVATIONS

In this paper, we use the following notation:

$$\begin{aligned} \mathbb{N} &= \{1, 2, \dots\}, & \mathbb{N}_- &= \{-1, -2, \dots\}, & \mathbb{N}_0 &= \{0, 1, 2, \dots\}, \\ \mathbb{Z} &= \{0, \pm 1, \pm 2, \dots\}, & \mathbb{R} &= (-\infty, +\infty), & \mathbb{C} &= \{x + iy : x, y \in \mathbb{R}, i = \sqrt{-1}\}. \end{aligned}$$

Let

$$W(z) = \frac{\arctan \sqrt{2e^{-z} - 1}}{\sqrt{2e^{-z} - 1}}. \tag{1.1}$$

The third problem posed by Herbert S. Wilf (1931–2012) at the webpage <https://www2.math.upenn.edu/~wilf/website/UnsolvedProblems.pdf> states that, if $W(z) = \sum_{n=0}^{+\infty} a_n z^n$, find the first term of the asymptotic behaviour of the a_n 's.

2020 *Mathematics Subject Classification.* Primary 41A58; Secondary 11B73, 11B83, 26A24, 33B10.

Key words and phrases. Maclaurin's series expansion; Taylor's series expansion; inverse tangent function; inverse hyperbolic tangent function; coefficient; Faà di Bruno formula; Bell polynomial of the second kind; Stirling number of the second kind; combinatorial identity; special value; closed-form formula; Gauss' hypergeometric function; generating function; logarithmically complete monotonicity; positivity; logarithmic convexity; asymptotic rational approximation; Wronski's theorem.

This paper was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$.

In the conference paper [33], Mark Daniel Ward at Purdue University considered this problem.

We recite the texts of the review MR2735366 written by Xiao-He Tian at Illinois Wesleyan University for the paper [33] as follows.

The coefficient a_n can be written as $a_n = b_n\pi - c_n$, where b_n and c_n are nonnegative rational numbers. In fact, $\lim_{n \rightarrow +\infty} \frac{a_n}{b_n} = 0$, and the rational numbers of the form $\frac{c_n}{b_n}$ provide approximations to π . A complete expansion of the coefficients a_n is found by the author. It is probably the best that can be done, given the oscillatory nature of the terms.

We recite Wilf's comments on the paper [33] on 13 December 2010 as follows.

Mark Ward has found a complete expansion of these coefficients. It's not quite an asymptotic series in the usual sense, but it is probably the best that can be done, given the oscillatory nature of the terms.

For our own convenience, we call the function $W(z)$ defined by (1.1) Wilf's function.

In the theory of complex functions, the inverse tangent function $\text{Arctan } z$ is defined by

$$\text{Arctan } z = \frac{1}{2i} \text{Ln} \frac{1+iz}{1-iz} = \frac{1}{2i} \text{Ln} \frac{i-z}{i+z}$$

on the punctured complex plane $\mathbb{C} \setminus \{\pm i\}$ and its principal branch $\arctan z$ such that $\arctan 0 = 0$ is taken as

$$\arctan z = \frac{i}{2} [\ln(1-iz) - \ln(1+iz)]$$

on the cut complex plane

$$\mathcal{A} = \mathbb{C} \setminus (-i\infty, -i] \setminus [i, i\infty),$$

where $i = \sqrt{-1}$ is the imaginary unit, $\ln z = \ln |z| + i \arg z$ for $z \in \mathbb{C} \setminus (-\infty, 0]$ stands for the principal branch of the logarithm $\text{Ln } z$, and $\arg z \in (-\pi, \pi]$ represents the principal value of the argument $\text{Arg } z$. On the cut complex plane \mathcal{A} , the range of the principal branch $\arctan x$ for real numbers $x \in \mathbb{R}$ is determined as $(-\frac{\pi}{2}, \frac{\pi}{2})$. The principal branch $\arctan z$ has a unique zero $z = 0$ on \mathcal{A} .

The power function $f(z) = \sqrt{z}$ is defined on the whole complex plane \mathbb{C} . In polar coordinates, we can write $z = r e^{i\theta}$ for $r \geq 0$ and $\theta \in (-\pi, \pi]$. Then $f(z) = \sqrt{r} e^{i(\theta+2k\pi)/2}$ for $k = 0, 1$. On the cut complex plane $\mathbb{C} \setminus (-\infty, 0)$, the principal branch of $f(z)$ such that $f(1) = 1$ is $\sqrt{r} e^{i\theta/2}$ for $r \geq 0$ and $\theta \in (-\pi, \pi)$. The range of the principal branch of $f(z) = \sqrt{z}$ is the right half complex plane plus the origin point. This range is a subset of \mathcal{A} . Therefore, the principal branch of the function $\frac{\arctan \sqrt{z}}{\sqrt{z}}$ is defined on $\mathbb{C} \setminus (-\infty, 0)$.

The function $g(z) = 2e^{-z} - 1$ is analytic on the complex plane \mathbb{C} and its range is $\mathbb{C} \setminus \{-1\}$.

It is clear that Wilf's function $W(z)$ can be regarded as the composition of three functions

$$h(z) = \frac{\arctan z}{z}, \quad f(z) = \sqrt{z}, \quad g(z) = 2e^{-z} - 1.$$

When considering the composition of three principal branches of these functions $h(z)$, $f(z)$, and $g(z)$, Wilf's function $W(z)$ has no definition on points such that

$$2e^{-z} - 1 = 2e^{-x} \cos y - 1 - (2e^{-x} \sin y) i \in (-\infty, 0),$$

which is equivalent to

$$2e^{-x} \cos y - 1 < 0 \quad \text{and} \quad 2e^{-x} \sin y = 0.$$

These points constitute infinitely many parallel and horizontal rays

$$\{z = x + yi \in \mathbb{C} : x > \ln 2, y = 2k\pi, k \in \mathbb{Z}\}$$

and infinitely many parallel, horizontal, and straight lines

$$\{z = x + yi \in \mathbb{C} : x \in \mathbb{R}, y = (2k+1)\pi, k \in \mathbb{Z}\}$$

on the complex plane \mathbb{C} . In a word, Wilf's function $W(z)$ is defined on the region

$$\mathbb{C} \setminus \{z = x + yi \in \mathbb{C} : x > \ln 2, y = 2k\pi, k \in \mathbb{Z}\} \setminus \{z = x + yi \in \mathbb{C} : x \in \mathbb{R}, y = (2k+1)\pi, k \in \mathbb{Z}\}.$$

Consequently, on the open disc $|z| < \ln 2$, Wilf's function $W(z)$ is analytic.

By choosing an alternative branch of $\arctan z$, Mark Daniel Ward proved in [33, Lemma 2.2] that the maximum radius around the origin for which Wilf's function $W(z)$ has an analytic extension is

$$|\ln 2 + 2\pi i| = \sqrt{(\ln 2)^2 + 4\pi^2}.$$

In this paper, we will establish a closed-form formula involving the Stirling numbers of the second kind $S(n, k)$ for the coefficients a_n in Maclaurin's series expansion $W(z) = \sum_{n=0}^{+\infty} a_n z^n$ of Wilf's function $W(z)$ on the open disc $|z| < \ln 2$. From this, we will express a_n as the form $b_n \pi - c_n$ in terms of closed-form formulas for b_n and c_n .

For reaching the above aim, we need to prepare the Faà di Bruno formula, several identities for the Bell polynomials of the second kind, some basic knowledge of Gauss' hypergeometric function ${}_2F_1(\alpha, \beta; \gamma; z)$, two combinatorial identities, and other notations in Section 2 of this paper.

By virtue of some preparations in Section 2, we establish in Section 3 Taylor's or Maclaurin's series expansions of four functions $\arctan z$, $\frac{\arctan \sqrt{z}}{\sqrt{z}}$, $\frac{\arctan z}{z}$, and $\sqrt{2e^{-z}-1}$, and discover a closed-form formula for a sequence of special values of the Bell polynomials of the second kind.

In Section 4, by means of Taylor's series expansion of $\frac{\arctan \sqrt{z}}{\sqrt{z}}$ presented in Section 3, with the help of some preparations in Section 2, we find two forms for Maclaurin's series expansion $W(z) = \sum_{n=0}^{+\infty} a_n z^n$ of Wilf's function $W(z)$, the latter is a closed-form formula for coefficients a_n . We also obtain a closed-form formula of ${}_2F_1(n + \frac{1}{2}, n + 1; n + \frac{3}{2}; -1)$ for $n \in \mathbb{N}_0$.

In Section 5, from some results in Sections 3 and 4, with the aid of a relation between $\arctan z$ and $\operatorname{arctanh} z$, we derive series expansions of four functions

$$\operatorname{arctanh} z, \quad \frac{\operatorname{arctanh} \sqrt{z}}{\sqrt{z}}, \quad \frac{\operatorname{arctanh} z}{z}, \quad \frac{\operatorname{arctanh}(\sqrt{1-2e^{-z}})}{\sqrt{1-2e^{-z}}}.$$

In Section 6, we return to Maclaurin's series expansion

$$W(z) = \sum_{n=0}^{+\infty} a_n z^n = \sum_{n=0}^{+\infty} (b_n \pi - c_n) z^n, \quad |z| < \ln 2 \quad (1.2)$$

and analyze some properties of the sequences a_n , b_n , and c_n .

2. PREPARATIONS AND LEMMAS

For $\alpha \in \mathbb{C}$, the falling and rising factorials of α are respectively defined [15, p. 618, 26.1.1] by

$$\langle \alpha \rangle_n = \prod_{k=0}^{n-1} (\alpha - k) = \begin{cases} \alpha(\alpha - 1) \cdots (\alpha - n + 1), & n \in \mathbb{N}; \\ 1, & n = 0 \end{cases}$$

and

$$(\alpha)_n = \prod_{k=0}^{n-1} (\alpha + k) = \begin{cases} \alpha(\alpha + 1) \cdots (\alpha + n - 1), & n \in \mathbb{N}; \\ 1, & n = 0. \end{cases}$$

In particular, we note that $\langle 0 \rangle_0 = (0)_0 = 1$ and $\langle 0 \rangle_n = (0)_n = 0$ for $n \in \mathbb{N}$. The double factorial of negative odd integers $-(2k + 1)$ for $k \geq 0$ is defined by

$$[-(2k + 1)]!! = \frac{(-1)^k}{(2k - 1)!!} = (-1)^k \frac{(2k)!!}{(2k)!}. \quad (2.1)$$

For $\alpha, \beta \in \mathbb{C}$ and $\gamma \in \mathbb{C} \setminus \{0, -1, -2, \dots\}$, Gauss' hypergeometric function ${}_2F_1(\alpha, \beta; \gamma; z)$ can be defined by the series

$${}_2F_1(\alpha, \beta; \gamma; z) = \sum_{n=0}^{+\infty} \frac{(\alpha)_n (\beta)_n}{(\gamma)_n} \frac{z^n}{n!}. \quad (2.2)$$

The series in (2.2) absolutely and uniformly converges on the open unit disc $|z| < 1$. If $\Re(\alpha + \beta - \gamma) < 0$, the series in (2.2) absolutely converges over the unit circle $|z| = 1$. If $0 \leq \Re(\alpha + \beta - \gamma) < 1$, the series in (2.2) conditionally converges at all points of the unit circle $|z| = 1$ except $z = 1$. If $\Re(\alpha + \beta - \gamma) \geq 1$, the series in (2.2) diverges over the circle $|z| = 1$. There exists an analytic continuation of Gauss' hypergeometric function ${}_2F_1(\alpha, \beta; \gamma; z)$ to the region $\{z \in \mathbb{C} : |z| > 1\} \setminus (1, +\infty)$. Gauss' hypergeometric function ${}_2F_1(\alpha, \beta; \gamma; z)$ is a univalent analytic function on $\mathbb{C} \setminus (1, +\infty)$. If α or β is zero or a negative integer, the series in (2.2) terminates after a finite number of terms, and Gauss' hypergeometric function

${}_2F_1(\alpha, \beta; \gamma; z)$ is a polynomial in z . For details and more information, please refer to [12, pp. 64–66, Section 3.7] and [32, Chapter 5].

The classical Euler's gamma function $\Gamma(z)$ can be defined [32, Chapter 3] by

$$\Gamma(z) = \lim_{n \rightarrow +\infty} \frac{n!n^z}{\prod_{k=0}^n (z+k)}, \quad z \in \mathbb{C} \setminus \{0, -1, -2, \dots\}.$$

The extended binomial coefficient $\binom{z}{w}$ for $z, w \in \mathbb{C}$ is defined by

$$\binom{z}{w} = \begin{cases} \frac{\Gamma(z+1)}{\Gamma(w+1)\Gamma(z-w+1)}, & z \notin \mathbb{N}_-, \quad w, z-w \notin \mathbb{N}_-; \\ 0, & z \notin \mathbb{N}_-, \quad w \in \mathbb{N}_- \text{ or } z-w \in \mathbb{N}_-; \\ \frac{\langle z \rangle_w}{w!}, & z \in \mathbb{N}_-, \quad w \in \mathbb{N}_0; \\ \frac{\langle z \rangle_{z-w}}{(z-w)!}, & z, w \in \mathbb{N}_-, \quad z-w \in \mathbb{N}_0; \\ 0, & z, w \in \mathbb{N}_-, \quad z-w \in \mathbb{N}_-; \\ \infty, & z \in \mathbb{N}_-, \quad w \notin \mathbb{Z}. \end{cases}$$

The Stirling numbers of the second kind $S(n, k)$ for $n \geq k \geq 0$ can be analytically generated [6, p. 51] by

$$\frac{(e^x - 1)^k}{k!} = \sum_{n=k}^{+\infty} S(n, k) \frac{x^n}{n!} \quad (2.3)$$

and can be explicitly computed [6, p. 204, Theorem A] by

$$S(n, k) = \begin{cases} \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \ell^n, & n > k \in \mathbb{N}_0; \\ 1, & n = k \in \mathbb{N}_0. \end{cases} \quad (2.4)$$

By the way, we recommend the papers [17, 22] for several new results of the Stirling numbers of the second kind $S(n, k)$.

The Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ for $n \geq k \geq 0$ can be defined [6, p. 134, Theorem A] by

$$B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{1 \leq i \leq n-k+1 \\ \ell_i \in \{0\} \cup \mathbb{N} \\ \sum_{i=1}^{n-k+1} i \ell_i = n \\ \sum_{i=1}^{n-k+1} \ell_i = k}} \frac{n!}{\prod_{i=1}^{n-k+1} \ell_i!} \prod_{i=1}^{n-k+1} \left(\frac{x_i}{i!} \right)^{\ell_i}.$$

The Faà di Bruno formula can be described [6, p. 139, Theorem C] in terms of the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ by

$$\frac{d^n}{dt^n} f \circ h(t) = \sum_{k=0}^n f^{(k)}(h(t)) B_{n,k}(h'(t), h''(t), \dots, h^{(n-k+1)}(t)), \quad n \in \mathbb{N}_0. \quad (2.5)$$

The Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ satisfy the identities

$$B_{n,k}(\alpha\beta x_1, \alpha\beta^2 x_2, \dots, \alpha\beta^{n-k+1} x_{n-k+1}) = \alpha^k \beta^n B_{n,k}(x_1, x_2, \dots, x_{n-k+1}), \quad (2.6)$$

$$B_{n,k}(1, 1, \dots, 1) = S(n, k), \quad (2.7)$$

$$B_{n,k}(\alpha, 1, 0, \dots, 0) = \frac{(n-k)!}{2^{n-k}} \binom{n}{k} \binom{k}{n-k} \alpha^{2k-n}, \quad (2.8)$$

$$B_{n,k} \left(\left\langle \frac{1}{2} \right\rangle_1, \left\langle \frac{1}{2} \right\rangle_2, \dots, \left\langle \frac{1}{2} \right\rangle_{n-k+1} \right) = (-1)^{n+k} \frac{[2(n-k)-1]!!}{2^n} \binom{2n-k-1}{2(n-k)}, \quad (2.9)$$

and

$$\frac{1}{k!} \left(\sum_{m=1}^{+\infty} x_m \frac{t^m}{m!} \right)^k = \sum_{n=k}^{+\infty} B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) \frac{t^n}{n!} \quad (2.10)$$

for $n \geq k \geq 0$ and $\alpha, \beta \in \mathbb{C}$. See [5, p. 412], [6, pp. 133 and 135], [20, Theorem 4.1], and [23, p. 169, (3.6)], respectively. These identities can also be found in the review article [25].

Lemma 2.1. *For all $\ell, n \in \mathbb{Z}$, the combinatorial identity*

$$\sum_{k=\ell}^n \binom{2n-k-1}{n-1} 2^k k = \binom{2n-\ell}{n} 2^\ell n \quad (2.11)$$

is valid, where an empty sum is understood to be 0.

Proof. We will prove it by the descending induction mentioned in [11, p. 88].

It is trivial that, when $n - \ell = 0$, the identity (2.11) is valid.

Assume that the identity (2.11) is valid for some $n - \ell > 0$. Then it is easy to see that

$$\begin{aligned} \sum_{k=\ell-1}^n \binom{2n-k-1}{n-1} 2^k k &= \sum_{k=\ell}^n \binom{2n-k-1}{n-1} 2^k k + \binom{2n-\ell}{n-1} 2^{\ell-1} (\ell-1) \\ &= \binom{2n-\ell}{n} 2^\ell n + \binom{2n-\ell}{n-1} 2^{\ell-1} (\ell-1) \\ &= \binom{2n-\ell+1}{n} 2^{\ell-1} n. \end{aligned}$$

Inductively, we conclude that the identity (2.11) is valid for all $n \geq \ell$. The proof of Lemma 2.1 is complete. \square

Remark 2.1. The idea of the descending induction utilized in the proof of Lemma 2.1 comes from Dr. Darij Grinberg (Drexel University, Germany; darijgrinberg@gmail.com; <http://www.cip.ifi.lmu.de/~grinberg/>) at the site <https://mathoverflow.net/q/402822> online. The identity (2.11) in Lemma 2.1 has been posted at <https://mathoverflow.net/a/402833> and <https://mathoverflow.net/a/403278> online.

Lemma 2.2. *Let*

$$P(n, k) = k! [2(n-k) - 1]!! \binom{2n-k-1}{2(n-k)}, \quad n \geq k \geq 0. \quad (2.12)$$

Then

$$\sum_{m=0}^k (-1)^m \binom{k}{m} \binom{m/2}{\ell} = \begin{cases} 0, & k > \ell \in \mathbb{N}_0; \\ \frac{(-1)^\ell}{(2\ell)!!} P(\ell, k), & \ell \geq k \in \mathbb{N}_0. \end{cases} \quad (2.13)$$

Proof. In [28, p. 165, (12.1)], there is a relation

$$n! \binom{z}{n} = \sum_{\ell=0}^n s(n, \ell) z^\ell, \quad n \in \mathbb{N}_0, \quad z \in \mathbb{C}. \quad (2.14)$$

Making use of the relation 2.14 and utilizing the explicit formula (2.4), we find

$$\begin{aligned} \sum_{m=0}^k (-1)^m \binom{k}{m} \binom{m/2}{\ell} &= \frac{1}{\ell!} \sum_{m=0}^k (-1)^m \binom{k}{m} \left[\ell! \binom{m/2}{\ell} \right] \\ &= \frac{1}{\ell!} \sum_{m=0}^k (-1)^m \binom{k}{m} \sum_{q=0}^{\ell} s(\ell, q) \left(\frac{m}{2} \right)^q \\ &= \frac{1}{\ell!} \sum_{q=0}^{\ell} \frac{s(\ell, q)}{2^q} \sum_{m=0}^k (-1)^m \binom{k}{m} m^q \\ &= (-1)^k \frac{k!}{\ell!} \sum_{q=0}^{\ell} s(\ell, q) \left(\frac{1}{2} \right)^q S(q, k) \end{aligned} \quad (2.15)$$

for $k, \ell \in \mathbb{N}_0$, where 0^0 was regarded as 1. Hence, it is easy to see that

$$\sum_{m=0}^k (-1)^m \binom{k}{m} \binom{m/2}{\ell} = 0, \quad 0 \leq \ell < k.$$

In [23, Theorem 2.1], the identity

$$B_{n,k}(\langle \alpha \rangle_1, \langle \alpha \rangle_2, \dots, \langle \alpha \rangle_{n-k+1}) = \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \langle \alpha \ell \rangle_n \quad (2.16)$$

for $n \geq k \in \mathbb{N}_0$ and $\alpha \in \mathbb{C}$ was proved. Taking $\alpha = \frac{1}{2}$ in (2.16), applying the relation

$$\langle z \rangle_n = \sum_{\ell=0}^n s(n, \ell) z^\ell, \quad z \in \mathbb{C}, \quad n \in \mathbb{N}_0$$

in [32, p. 9, (1.26)], interchanging the order of sums, and employing (2.4) yield

$$\begin{aligned} B_{n,k} \left(\left\langle \frac{1}{2} \right\rangle_1, \left\langle \frac{1}{2} \right\rangle_2, \dots, \left\langle \frac{1}{2} \right\rangle_{n-k+1} \right) &= \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \left\langle \frac{\ell}{2} \right\rangle_n \\ &= \frac{(-1)^k}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \sum_{m=0}^n s(n, m) \left(\frac{\ell}{2} \right)^m \\ &= \frac{(-1)^k}{k!} \sum_{m=0}^n s(n, m) \left(\frac{1}{2} \right)^m \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \ell^m \\ &= \sum_{m=0}^n s(n, m) \left(\frac{1}{2} \right)^m S(m, k) \end{aligned}$$

for $n \geq k \geq 0$. Combining this with (2.9) results in

$$\sum_{m=k}^n s(n, m) \left(\frac{1}{2} \right)^m S(m, k) = (-1)^{n+k} \frac{[2(n-k)-1]!!}{2^n} \binom{2n-k-1}{2(n-k)} \quad (2.17)$$

for $n \geq k \geq 0$. Substituting the identity (2.17) into (2.15), we figure out

$$\sum_{m=0}^k (-1)^m \binom{k}{m} \binom{m/2}{\ell} = (-1)^\ell k! \frac{[2(\ell-k)-1]!!}{(2\ell)!!} \binom{2\ell-k-1}{2(\ell-k)} = \frac{(-1)^\ell}{(2\ell)!!} P(\ell, k)$$

for $\ell \geq k \in \mathbb{N}_0$. The proof of Lemma 2.2 is thus complete. \square

Remark 2.2. The identity (2.13) in Lemma 2.2 has been announced at <https://math.stackexchange.com/a/4268339> and <https://math.stackexchange.com/a/4268341> online.

Recall from [13, Chapter XIII] and [34, Chapter IV] that an infinitely differentiable function f is said to be completely (or absolutely, respectively) monotonic on an interval I if it has derivatives of all orders on I and satisfies $(-1)^n f^{(n)}(x) \geq 0$ (or $f^{(n)}(x) \geq 0$, respectively) for all $x \in I$ and $n \in \mathbb{N}_0$. Recall from [7, Definition 1], [19, Definition 1] and [30, Definition 5.10 and Comments 5.29] that an infinitely differentiable and positive function f is said to be logarithmically absolutely (or completely, respectively) monotonic on an interval I if $[\ln f(x)]^{(n)} \geq 0$ (or $(-1)^n [\ln f(x)]^{(n)} \geq 0$, respectively) for all $n \in \mathbb{N}$ and $x \in I$. In [3, Theorem 1.1], [7, Theorem 1], [19, Theorem 1], and [21, p. 627, (1.4)], the following two relations between these two pairs of functions were discovered or reviewed:

- (1) A logarithmically completely monotonic function on an interval I is also completely monotonic on I , but not conversely.
- (2) A logarithmically absolutely monotonic function on an interval I is also absolutely monotonic on I , but not conversely.

It is easy to see that, a function $f(x)$ is (logarithmically) completely monotonic on an interval I if and only if it is (logarithmically) absolutely monotonic on the interval $-I$ respectively. These concepts and conclusions will be employed in Section 6.

3. TAYLOR'S SERIES EXPANSIONS OF THE INVERSE TANGENT FUNCTION

In this section, by means of some preparations in Section 2, we present several Maclaurin's and Taylor's series expansions of the functions $\arctan z$, $\frac{\arctan \sqrt{z}}{\sqrt{z}}$, $\frac{\arctan z}{z}$, and $\sqrt{2e^{-z}-1}$.

Theorem 3.1. For $n \in \mathbb{N}$, the derivative of $\arctan z$ is

$$(\arctan z)^{(n)} = \frac{(n-1)!}{(2z)^{n-1}} \sum_{k=0}^{n-1} (-1)^k \binom{k}{n-k-1} \frac{(2z)^{2k}}{(1+z^2)^{k+1}}.$$

The function $\frac{\arctan z}{z}$ has Taylor's series expansion

$$\frac{\arctan z}{z} = \sum_{n=0}^{+\infty} (-1)^n \left[\frac{\pi}{4} + T(n) \right] (z-1)^n, \quad |z-1| < \sqrt{2}, \quad (3.1)$$

where

$$T(n) = \begin{cases} 0, & n = 0; \\ \sum_{k=1}^n \frac{(-1)^k}{2^{k/2} k} \sin \frac{3k\pi}{4}, & n \in \mathbb{N}. \end{cases} \quad (3.2)$$

Proof. For $n \in \mathbb{N}$, by virtue of the Faà di Bruno formula (2.5), we find

$$\begin{aligned} (\arctan z)^{(n)} &= \left(\frac{1}{1+z^2} \right)^{(n-1)} \\ &= \sum_{k=0}^{n-1} \frac{(-1)^k k!}{(1+z^2)^{k+1}} B_{n-1,k}(2z, 2, 0, \dots, 0) \\ &= \sum_{k=0}^{n-1} \frac{(-1)^k k!}{(1+z^2)^{k+1}} 2^k B_{n-1,k}(z, 1, 0, \dots, 0) \\ &= \sum_{k=0}^{n-1} \frac{(-1)^k k!}{(1+z^2)^{k+1}} 2^k \frac{(n-k-1)!}{2^{n-k-1}} \binom{n-1}{k} \binom{k}{n-k-1} z^{2k-n+1} \\ &= \frac{(n-1)!}{(2z)^{n-1}} \sum_{k=0}^{n-1} (-1)^k \binom{k}{n-k-1} \frac{(2z)^{2k}}{(1+z^2)^{k+1}}, \end{aligned}$$

where we also utilized the identities (2.6) and (2.8). Accordingly, for $n \in \mathbb{N}_0$, we have

$$\begin{aligned} \left(\frac{\arctan z}{z} \right)^{(n)} &= \sum_{k=0}^n \binom{n}{k} (\arctan z)^{(k)} \left(\frac{1}{z} \right)^{(n-k)} \\ &= \frac{(-1)^n n! \arctan z}{z^{n+1}} + \sum_{k=1}^n \binom{n}{k} \frac{(k-1)!}{(2z)^{k-1}} \sum_{\ell=0}^{k-1} (-1)^\ell \binom{\ell}{k-\ell-1} \frac{(2z)^{2\ell} (-1)^{n-k} (n-k)!}{(1+z^2)^{\ell+1} z^{n-k+1}} \\ &\rightarrow (-1)^n n! \left[\frac{\pi}{4} + \sum_{k=1}^n \frac{(-1)^k}{2^k k} \sum_{\ell=0}^{k-1} (-2)^\ell \binom{\ell}{k-\ell-1} \right] \\ &= (-1)^n n! \left[\frac{\pi}{4} + T(n) \right] \end{aligned}$$

as $z \rightarrow 1$, where we used the identity

$$\sum_{\ell=0}^k (-2)^\ell \binom{\ell}{k-\ell} = 2^{k/2} \left(\cos \frac{3k\pi}{4} - \sin \frac{3k\pi}{4} \right) = 2^{(k+1)/2} \sin \frac{3(k+1)\pi}{4} \quad (3.3)$$

for $k \geq 0$, which can be deduced from the identity

$$\sum_{k=0}^{n/2} \binom{n-k}{k} z^k = \frac{1}{\sqrt{1+4z}} \left[\left(\frac{1+\sqrt{1+4z}}{2} \right)^{n+1} - \left(\frac{1-\sqrt{1+4z}}{2} \right)^{n+1} \right]$$

collected as (1.70) and (1.71) on page 33 in [31]. Consequently, we obtain Taylor's series expansion (3.1).

The singular points of $\frac{\arctan z}{z}$ closest to the point $z = 1$ are $\pm i$. Therefore, the convergent open disc of Taylor's series expansion (3.1) is $|z-1| < \sqrt{2}$. The proof of Theorem 3.1 is complete. \square

Remark 3.1. The first few values of the sequence $T(n)$ for $0 \leq n \leq 11$ are given by

$$0, \quad -\frac{1}{2}, \quad -\frac{3}{4}, \quad -\frac{5}{6}, \quad -\frac{5}{6}, \quad -\frac{97}{120}, \quad -\frac{63}{80}, \quad -\frac{109}{140}, \quad -\frac{109}{140}, \quad -\frac{7883}{10080}, \quad -\frac{15829}{20160}.$$

These values reveal that $T(n) < 0$ and $T(4n-1) = T(4n)$ for $n \in \mathbb{N}$.

Taking the limit $z \rightarrow 0$ on both side of the series expansion (3.1) in Theorem 3.1 shows

$$\sum_{n=0}^{+\infty} \left[\frac{\pi}{4} + T(n) \right] = 1.$$

By the necessary condition for a series expansion to be convergent, we obtain

$$\lim_{n \rightarrow +\infty} T(n) = \sum_{k=1}^{+\infty} \frac{(-1)^k}{2^{k/2} k} \sin \frac{3k\pi}{4} = -\frac{\pi}{4}. \quad (3.4)$$

We can regard (3.4) as a series representation of the circular constant π .

Corollary 3.1. For $|z-1| < 1$, the function $\frac{\arctan z - \frac{\pi}{4}}{z}$ has Taylor's series expansion

$$\frac{\arctan z - \frac{\pi}{4}}{z} = \sum_{n=1}^{+\infty} (-1)^n T(n) (z-1)^n, \quad (3.5)$$

where $T(n)$ is defined by (3.2).

Proof. Taylor's series expansion (3.1) can be reformulated as

$$\frac{\arctan z}{z} = \frac{\pi}{4} \sum_{n=0}^{+\infty} (1-z)^n + \sum_{n=0}^{+\infty} T(n) (1-z)^n = \frac{\pi}{4z} + \sum_{n=1}^{+\infty} T(n) (1-z)^n$$

for $|z-1| < 1$. The proof of Corollary 3.1 is complete. \square

Remark 3.2. Letting $z = \sqrt{3}$ on both sides of Taylor's series expansion (3.5) deduces a series representation

$$\pi = 12\sqrt{3} \sum_{n=1}^{+\infty} (-1)^{n+1} T(n) (\sqrt{3}-1)^n,$$

where $T(n)$ is defined by (3.2).

Theorem 3.2. The function $\frac{\arctan \sqrt{z}}{\sqrt{z}}$ has Taylor's series expansion

$$\frac{\arctan \sqrt{z}}{\sqrt{z}} = \sum_{n=0}^{+\infty} \frac{(-1)^n}{2^n} \left[(2n-1)!! \frac{\pi}{4} + \frac{n!}{2^n} \sum_{k=1}^n (-1)^k \binom{2n-k}{n} \frac{2^{k/2}}{k} \sin \frac{3k\pi}{4} \right] \frac{(z-1)^n}{n!} \quad (3.6)$$

for $|z-1| < 1$, where $(-1)!! = 1$ is defined by (2.1).

Proof. Let $w = f(z) = \sqrt{z} \rightarrow 1$ as $z \rightarrow 1$. By virtue of the Faà di Bruno formula (2.5) and by means of the identities (2.6) and (2.9), we discover

$$\begin{aligned} \left(\frac{\arctan \sqrt{z}}{\sqrt{z}} \right)^{(n)} &= \sum_{k=0}^n \left(\frac{\arctan w}{w} \right)^{(k)} B_{n,k} \left(\left\langle \frac{1}{2} \right\rangle_1 \frac{1}{z^{1/2}}, \left\langle \frac{1}{2} \right\rangle_1 \frac{1}{z^{3/2}}, \dots, \left\langle \frac{1}{2} \right\rangle_{n-k+1} \frac{1}{z^{[2(n-k)+1]/2}} \right) \\ &= \sum_{k=0}^n \left(\frac{\arctan w}{w} \right)^{(k)} B_{n,k} \left(\left\langle \frac{1}{2} \right\rangle_1 \frac{z^{1/2}}{z}, \left\langle \frac{1}{2} \right\rangle_1 \frac{z^{1/2}}{z^2}, \dots, \left\langle \frac{1}{2} \right\rangle_{n-k+1} \frac{z^{1/2}}{z^{n-k+1}} \right) \\ &= \sum_{k=0}^n \left(\frac{\arctan w}{w} \right)^{(k)} z^{k/2-n} B_{n,k} \left(\left\langle \frac{1}{2} \right\rangle_1, \left\langle \frac{1}{2} \right\rangle_1, \dots, \left\langle \frac{1}{2} \right\rangle_{n-k+1} \right) \\ &= \sum_{k=0}^n \left(\frac{\arctan w}{w} \right)^{(k)} z^{k/2-n} (-1)^{n+k} \frac{[2(n-k)-1]!!}{2^n} \binom{2n-k-1}{2(n-k)} \\ &\rightarrow \frac{(-1)^n}{2^n} \sum_{k=0}^n P(n,k) \left[\frac{\pi}{4} + T(k) \right] \end{aligned}$$

as $z \rightarrow 1$ for $n \in \mathbb{N}_0$, where we used in the last step the series expansion (3.1) in Theorem 3.1 and used the notation $P(n, k)$ and $T(k)$ defined by (2.12) and (3.2) respectively. Consequently, we acquire Taylor's series expansion

$$\frac{\arctan \sqrt{z}}{\sqrt{z}} = \sum_{n=0}^{+\infty} (-1)^n \left(\sum_{k=0}^n P(n, k) \left[\frac{\pi}{4} + T(k) \right] \right) \frac{(z-1)^n}{(2n)!!}, \quad |z-1| < 1. \quad (3.7)$$

The singular point of the function $\frac{\arctan \sqrt{z}}{\sqrt{z}}$ closest to the point $z = 1$ is $z = 0$. Therefore, the convergent open disc of the series in (3.7) on the complex plane \mathbb{C} is $|z-1| < 1$.

Replacing \sqrt{z} by z on both sides of (3.7) leads to

$$\frac{\arctan z}{z} = \sum_{n=0}^{+\infty} (-1)^n \left(\sum_{k=0}^n P(n, k) \left[\frac{\pi}{4} + T(k) \right] \right) \frac{(z^2-1)^n}{(2n)!!}, \quad |z^2-1| < 1. \quad (3.8)$$

Comparing the series expansion (3.1) in Theorem 3.1 with the series expansion (3.8) shows

$$\sum_{n=0}^{+\infty} (-1)^n \left[\frac{\pi}{4} + T(n) \right] (z-1)^n = \sum_{n=0}^{+\infty} (-1)^n \left(\sum_{k=0}^n P(n, k) \left[\frac{\pi}{4} + T(k) \right] \right) \frac{(z^2-1)^n}{(2n)!!}.$$

Differentiating $q \in \mathbb{N}_0$ times with respect to z on both sides of this equation gives

$$\begin{aligned} \sum_{n=q}^{+\infty} (-1)^n \left[\frac{\pi}{4} + T(n) \right] \langle n \rangle_q (z-1)^{n-q} &= \sum_{n=0}^{+\infty} \frac{(-1)^n}{(2n)!!} \left(\sum_{k=0}^n P(n, k) \left[\frac{\pi}{4} + T(k) \right] \right) [(z^2-1)^n]^{(q)} \\ &= \sum_{n=0}^{+\infty} \frac{(-1)^n}{(2n)!!} \left(\sum_{k=0}^n P(n, k) \left[\frac{\pi}{4} + T(k) \right] \right) \sum_{p=0}^q (2z)^{2p-q} (q-p)! \binom{q}{p} \binom{p}{q-p} \langle n \rangle_p (z^2-1)^{n-p} \\ &= \sum_{p=0}^q (2z)^{2p-q} (q-p)! \binom{q}{p} \binom{p}{q-p} \sum_{n=0}^{+\infty} \frac{(-1)^n}{(2n)!!} \left(\sum_{k=0}^n P(n, k) \left[\frac{\pi}{4} + T(k) \right] \right) \langle n \rangle_p (z^2-1)^{n-p}, \end{aligned}$$

where we used the derivative formula

$$\begin{aligned} [(z^2-1)^n]^{(q)} &= \sum_{p=0}^q (u^n)^{(p)} B_{q,p}(2z, 2, 0, \dots, 0), \quad u = u(z) = z^2-1 \\ &= \sum_{p=0}^q \langle n \rangle_p (z^2-1)^{n-p} 2^p B_{q,p}(z, 1, 0, \dots, 0) \\ &= \sum_{p=0}^q (2z)^{2p-q} (q-p)! \binom{q}{p} \binom{p}{q-p} \langle n \rangle_p (z^2-1)^{n-p} \end{aligned}$$

for $q \in \mathbb{N}_0$, in which we employed the formulas (2.5), (2.6), and (2.8). Furthermore, letting $z = 1$ and simplifying reveal

$$\begin{aligned} (-1)^q q! \left[\frac{\pi}{4} + T(q) \right] &= \sum_{p=0}^q (-1)^p 2^{p-q} (q-p)! \binom{q}{p} \binom{p}{q-p} \sum_{k=0}^p P(p, k) \left[\frac{\pi}{4} + T(k) \right] \\ &= \sum_{k=0}^q \left[\sum_{p=k}^q (-1)^p 2^{p-q} (q-p)! \binom{q}{p} \binom{p}{q-p} P(p, k) \right] \left[\frac{\pi}{4} + T(k) \right] \\ &= \frac{\pi}{4} \sum_{k=0}^q \left[\sum_{p=k}^q (-1)^p 2^{p-q} (q-p)! \binom{q}{p} \binom{p}{q-p} P(p, k) \right] \\ &\quad + \sum_{k=0}^q \left[\sum_{p=k}^q (-1)^p 2^{p-q} (q-p)! \binom{q}{p} \binom{p}{q-p} P(p, k) \right] T(k) \end{aligned}$$

for $q \in \mathbb{N}_0$. Further equating the rational and irrational parts on both sides yields

$$\sum_{k=0}^q \sum_{p=k}^q (-1)^p 2^{p-q} (q-p)! \binom{q}{p} \binom{p}{q-p} P(p, k) = (-1)^q q!$$

and

$$\sum_{k=0}^q \left[\sum_{p=k}^q (-1)^p 2^{p-q} (q-p)! \binom{q}{p} \binom{p}{q-p} P(p, k) \right] T(k) = (-1)^q q! T(q)$$

for $q \in \mathbb{N}_0$. Replacing q by n and simplifying result in

$$\sum_{k=0}^n \sum_{\ell=k}^n \frac{(-2)^\ell}{\ell!} \binom{\ell}{n-\ell} P(\ell, k) = \sum_{\ell=0}^n \binom{\ell}{n-\ell} \frac{(-2)^\ell}{\ell!} \sum_{k=0}^{\ell} P(\ell, k) = (-2)^n \quad (3.9)$$

and

$$\sum_{k=0}^n \left[\sum_{\ell=k}^n \binom{\ell}{n-\ell} \frac{(-2)^\ell}{\ell!} P(\ell, k) \right] T(k) = \sum_{\ell=0}^n \binom{\ell}{n-\ell} \frac{(-2)^\ell}{\ell!} \sum_{k=0}^{\ell} P(\ell, k) T(k) = (-2)^n T(n) \quad (3.10)$$

for $n \in \mathbb{N}_0$.

For $k \in \mathbb{N}$, let s_k and S_k be two sequences independent of n such that $n \geq k \in \mathbb{N}$. The inversion theorem in [27, p. 528, Theorem 4.4] reads that

$$s_n = \sum_{k=1}^n \binom{k}{n-k} S_k \quad \text{if and only if} \quad (-1)^n n S_n = \sum_{k=1}^n \binom{2n-k-1}{n-1} (-1)^k k s_k. \quad (3.11)$$

Applying the inversion theorem in (3.11) to the second identities in (3.9) and (3.10) respectively, we acquire

$$\sum_{\ell=0}^n P(n, \ell) = \frac{(n-1)!}{2^n} \sum_{k=1}^n \binom{2n-k-1}{n-1} 2^k k = \frac{(n-1)!}{2^n} \binom{2n}{n} n = (2n-1)!!$$

and

$$(-1)^n n \frac{(-2)^n}{n!} \sum_{\ell=0}^n P(n, \ell) T(\ell) = \sum_{k=1}^n \binom{2n-k-1}{n-1} (-1)^k k (-2)^k T(k)$$

for $n \in \mathbb{N}$, where we used the identity (2.11) for $\ell = 0$ in Lemma 2.1. In a word, the identities

$$\sum_{k=0}^n P(n, k) = (2n-1)!! \quad (3.12)$$

and

$$\sum_{k=0}^n P(n, k) T(k) = \frac{(n-1)!}{2^n} \sum_{k=1}^n \binom{2n-k-1}{n-1} k 2^k T(k) \quad (3.13)$$

are valid for $n \in \mathbb{N}_0$. Applying (3.2) to (3.13) gives

$$\begin{aligned} \sum_{\ell=0}^n P(n, \ell) T(\ell) &= \frac{(n-1)!}{2^n} \sum_{\ell=1}^n \left[\sum_{k=\ell}^n \binom{2n-k-1}{n-1} 2^k k \right] \frac{(-1)^\ell}{\ell} \frac{1}{2^{\ell/2}} \sin \frac{3\ell\pi}{4} \\ &= \frac{(n-1)!}{2^n} \sum_{\ell=1}^n \binom{2n-\ell}{n} 2^\ell n \frac{(-1)^\ell}{\ell} \frac{1}{2^{\ell/2}} \sin \frac{3\ell\pi}{4} \\ &= \frac{n!}{2^n} \sum_{k=1}^n (-1)^k \binom{2n-k}{n} \frac{2^{k/2}}{k} \sin \frac{3k\pi}{4}, \end{aligned} \quad (3.14)$$

where we used the identity (2.11) for $\ell \geq 1$ in Lemma 2.1.

Substituting the identities (3.12) and (3.14) into (3.7) yields the series expansion (3.6). The proof of Theorem 3.2 is complete. \square

Remark 3.3. Setting $z = \frac{1}{3}$ on both sides of the series expansion (3.6) in Theorem 3.2 and simplifying lead to a series representation

$$\pi = 4\sqrt{3} \sum_{n=0}^{+\infty} \frac{1}{6^n} \sum_{k=1}^n (-1)^{k+1} \binom{2n-k}{n} \frac{2^{k/2}}{k} \sin \frac{3k\pi}{4}.$$

Remark 3.4. Let $s_n = (-2)^n$ and

$$S_n = (-1)^n 2 \binom{2n-1}{n}$$

for $n \in \mathbb{N}$. Applying these two sequences to the inversion theorem recited in (3.11) and considering the identity (2.11) for $\ell = 1$ lead to a combinatorial identity

$$\sum_{k=1}^n (-1)^k \binom{k}{n-k} \binom{2k-1}{k} = (-1)^n 2^{n-1}, \quad n \in \mathbb{N}.$$

This identity has been announced at <https://mathoverflow.net/a/405812> online.

Let $S_n = (-2)^n$ and

$$s_n = 2^{(n+1)/2} \sin \frac{3(n+1)\pi}{4}$$

for $n \in \mathbb{N}$. Applying these two sequence to the inversion theorem recited in (3.11) and utilizing the identity (3.3) reveal that

$$\sum_{k=1}^n (-1)^k \binom{2n-k-1}{n-1} 2^{(k+1)/2} k \sin \frac{3(k+1)\pi}{4} = 2^n n, \quad n \in \mathbb{N}.$$

Theorem 3.3. *The function $\sqrt{2e^{-z}-1}$ has Maclaurin's series expansion*

$$\sqrt{2e^{-z}-1} = \sum_{n=0}^{+\infty} (-1)^{n+1} \left[\sum_{k=0}^n (-1)^k S(n,k) (2k-3)!! \right] \frac{z^n}{n!}, \quad |z| < \ln 2, \quad (3.15)$$

where $S(n,k)$ denotes the Stirling numbers of the second kind in (2.4) and the double factorial of negative odd integers $-(2k+1)$ for $k \geq 0$ is defined by (2.1).

Proof. Let $w = g(z) = 2e^{-z} - 1$. Then

$$\sqrt{w} = \sqrt{g(z)} = \sqrt{2e^{-z}-1} \rightarrow 1, \quad z \rightarrow 0.$$

By the Faà di Bruno formula (2.5), we acquire

$$\begin{aligned} (\sqrt{2e^{-z}-1})^{(n)} &= \sum_{k=0}^n (\sqrt{w})^{(k)} B_{n,k}(g'(z), g''(z), \dots, g^{(n-k+1)}(z)) \\ &= \sum_{k=0}^n \left\langle \frac{1}{2} \right\rangle_k w^{1/2-k} B_{n,k}(-2e^{-z}, 2e^{-z}, \dots, (-1)^{n-k+1} 2e^{-z}) \\ &= \sum_{k=0}^n \left\langle \frac{1}{2} \right\rangle_k (2e^{-z}-1)^{1/2-k} (-1)^n (2e^{-z})^k B_{n,k}(1, 1, \dots, 1) \\ &= (-1)^n \sum_{k=0}^n S(n,k) (-1)^{k-1} (2k-3)!! (2e^{-z}-1)^{1/2-k} e^{-kz} \\ &\rightarrow (-1)^n \sum_{k=0}^n (-1)^{k-1} S(n,k) (2k-3)!! \end{aligned}$$

as $z \rightarrow 0$, where we employed the identities (2.6) and (2.7). The series expansion (3.15) is thus proved. The proof of Theorem 3.3 is complete. \square

Theorem 3.4. *For $n \geq k \geq 0$, the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ satisfy*

$$\begin{aligned} B_{n,k} \left(1, 0, 1, 3, 16, 105, \dots, (-1)^{n-k} \sum_{\ell=0}^{n-k+1} (-1)^{\ell-1} S(n-k+1, \ell) (2\ell-3)!! \right) \\ = \frac{(-1)^n}{k!} \sum_{\ell=k}^n (-1)^\ell S(n, \ell) P(\ell, k), \end{aligned} \quad (3.16)$$

where $P(\ell, k)$ is given by (2.12) and the Stirling numbers of the second kind $S(n, \ell)$ are generated by (2.3).

Proof. Making use of the formula (2.10) yields

$$B_{n+k,k}(x_1, x_2, \dots, x_{n+1}) = \binom{n+k}{k} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} \left[\sum_{m=0}^{+\infty} \frac{x_{m+1}}{(m+1)!} z^m \right]^k. \quad (3.17)$$

Setting $x_m = [\sqrt{g(z)}]^{(m)}|_{z=0} = (\sqrt{2e^{-z}-1})^{(m)}|_{z=0}$ for $m \in \mathbb{N}$ in (3.17) leads to

$$\begin{aligned} & B_{n+k,k} \left([\sqrt{g(z)}]'|_{z=0}, [\sqrt{g(z)}]''|_{z=0}, \dots, [\sqrt{g(z)}]^{(n+1)}|_{z=0} \right) \\ &= \binom{n+k}{k} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} \left[\sum_{m=0}^{+\infty} \frac{[\sqrt{g(z)}]^{(m+1)}|_{z=0}}{(m+1)!} z^m \right]^k \\ &= \binom{n+k}{k} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} \left[\frac{1}{z} \sum_{m=1}^{+\infty} \frac{[\sqrt{g(z)}]^{(m)}|_{z=0}}{m!} z^m \right]^k \\ &= \binom{n+k}{k} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} \left[\frac{\sqrt{g(z)} - 1}{z} \right]^k \\ &= \binom{n+k}{k} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} \left[\frac{1}{z^k} \sum_{\ell=0}^k (-1)^{k-\ell} \binom{k}{\ell} g^{\ell/2}(z) \right] \\ &= \binom{n+k}{k} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} \left[\frac{1}{z^k} \sum_{\ell=0}^k (-1)^{k-\ell} \binom{k}{\ell} (2e^{-z}-1)^{\ell/2} \right] \\ &= \binom{n+k}{k} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} \left[\frac{1}{z^k} \sum_{\ell=0}^k (-1)^{k-\ell} \binom{k}{\ell} \sum_{q=0}^{+\infty} \binom{\ell/2}{q} (2e^{-z}-2)^q \right] \\ &= \binom{n+k}{k} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} \left(\frac{1}{z^k} \sum_{\ell=0}^k (-1)^{k-\ell} \binom{k}{\ell} \sum_{q=0}^{+\infty} (2q)!! \binom{\ell/2}{q} \sum_{p=q}^{+\infty} (-1)^p S(p, q) \frac{z^p}{p!} \right) \\ &= \binom{n+k}{k} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} \sum_{\ell=0}^k (-1)^{k-\ell} \binom{k}{\ell} \sum_{p=0}^{+\infty} (-1)^p \left[\sum_{q=0}^p (2q)!! \binom{\ell/2}{q} S(p, q) \right] \frac{z^{p-k}}{p!} \\ &= \binom{n+k}{k} \sum_{\ell=0}^k (-1)^{k-\ell} \binom{k}{\ell} \sum_{p=0}^{+\infty} (-1)^p \left[\sum_{q=0}^p (2q)!! \binom{\ell/2}{q} S(p, q) \right] \langle p-k \rangle_n \lim_{z \rightarrow 0} \frac{z^{p-k-n}}{p!} \\ &= \binom{n+k}{k} (-1)^k \sum_{p=0}^{+\infty} (-1)^p \langle p-k \rangle_n \left[\sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \sum_{q=0}^p (2q)!! \binom{\ell/2}{q} S(p, q) \right] \lim_{z \rightarrow 0} \frac{z^{p-k-n}}{p!}, \end{aligned}$$

where we used the generating function (2.3). Therefore, we acquire

$$\begin{aligned} & B_{n+k,k} \left([\sqrt{g(z)}]'|_{z=0}, [\sqrt{g(z)}]''|_{z=0}, \dots, [\sqrt{g(z)}]^{(n+1)}|_{z=0} \right) \\ &= \frac{(-1)^n}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \left[\sum_{q=0}^{n+k} (2q)!! \binom{\ell/2}{q} S(n+k, q) \right] \end{aligned}$$

for $n \geq k \geq 0$. Replacing n by $n-k$ reduces to

$$\begin{aligned} & B_{n,k} \left([\sqrt{g(z)}]'|_{z=0}, [\sqrt{g(z)}]''|_{z=0}, \dots, [\sqrt{g(z)}]^{(n-k+1)}|_{z=0} \right) \\ &= \frac{(-1)^{n-k}}{k!} \sum_{\ell=0}^k (-1)^\ell \binom{k}{\ell} \left[\sum_{q=0}^n S(n, q) (2q)!! \binom{\ell/2}{q} \right]. \end{aligned}$$

The series expansion (3.15) in Theorem 3.3 implies that

$$[\sqrt{g(z)}]^{(n)}|_{z=0} = (-1)^n \sum_{k=0}^n (-1)^{k-1} S(n, k) (2k-3)!!, \quad n \in \mathbb{N}.$$

Consequently, it follows that

$$\begin{aligned} B_{n,k} & \left(-1, 0, -1, -3, -16, -105, \dots, (-1)^{n-k+1} \sum_{\ell=0}^{n-k+1} (-1)^{\ell-1} S(n-k+1, \ell) (2\ell-3)!! \right) \\ & = \frac{(-1)^{n-k}}{k!} \sum_{\ell=0}^n S(n, \ell) (2\ell)!! \sum_{m=0}^k (-1)^m \binom{k}{m} \binom{m/2}{\ell}. \end{aligned}$$

Employing the identity (2.6) results in the closed-form formula

$$\begin{aligned} B_{n,k} & \left(1, 0, 1, 3, 16, 105, \dots, (-1)^{n-k} \sum_{\ell=0}^{n-k+1} (-1)^{\ell-1} S(n-k+1, \ell) (2\ell-3)!! \right) \\ & = \frac{(-1)^n}{k!} \sum_{\ell=0}^n S(n, \ell) (2\ell)!! \sum_{m=0}^k (-1)^m \binom{k}{m} \binom{m/2}{\ell} \end{aligned} \quad (3.18)$$

for $n \geq k \geq 0$. Further considering the formula (2.13) in Lemma 2.2, we can conclude (3.16) from (3.18). The proof of Theorem 3.4 is complete. \square

Remark 3.5. The closed-form formula (3.16) in Theorem 3.4 can be applied to establish Maclaurin's series expansion of functions of the form $F(\sqrt{2e^{-z}-1})$, if all derivatives of the function $F(z)$ at $z = 1$ is computable.

4. MACLAURIN'S SERIES EXPANSIONS OF WILF'S FUNCTION

In this section, by means of Taylor's series expansions of $\frac{\arctan \sqrt{z}}{\sqrt{z}}$ and $\frac{\arctan z}{z}$ established in Section 3, and with the help of some preparations in Section 2, we find two forms for Maclaurin's series expansion $W(z) = \sum_{n=0}^{+\infty} a_n z^n$ of Wilf's function $W(z)$ defined by (1.1) on the open disc centered at $z = 0$ with a radius $\ln 2$. We also obtain a closed-form formula of ${}_2F_1\left(n + \frac{1}{2}, n + 1; n + \frac{3}{2}; -1\right)$ for $n \in \mathbb{N}_0$.

Theorem 4.1. For $|z| < \ln 2$, Wilf's function $W(z)$ has Maclaurin's series expansion

$$W(z) = \sum_{n=0}^{+\infty} (-1)^n \left[\sum_{k=0}^n (-1)^k S(n, k) \frac{(2k)!!}{2k+1} {}_2F_1\left(k + \frac{1}{2}, k + 1; k + \frac{3}{2}; -1\right) \right] \frac{z^n}{n!}, \quad (4.1)$$

where ${}_2F_1(\alpha, \beta; \gamma; z)$ is Gauss' hypergeometric function defined by (2.2) and $S(n, k)$ stands for the Stirling numbers of the second kind in (2.3).

Proof. It is well known [1, p. 81, 4.4.42] that

$$\arctan z = \sum_{n=0}^{+\infty} (-1)^n \frac{z^{2n+1}}{2n+1}, \quad |z| \leq 1, \quad z^2 \neq -1.$$

From this, it follows that

$$\frac{\arctan \sqrt{t}}{\sqrt{t}} = \sum_{n=0}^{+\infty} (-1)^n \frac{t^n}{2n+1}, \quad 0 \leq t \leq 1. \quad (4.2)$$

The series expansion (4.2) can be reformulated as

$$\begin{aligned} \frac{\arctan \sqrt{t}}{\sqrt{t}} & = \sum_{k=0}^{+\infty} (-1)^k \frac{[(t-1)+1]^k}{2k+1} \\ & = \sum_{k=0}^{+\infty} \frac{(-1)^k}{2k+1} \sum_{n=0}^k \binom{k}{n} (t-1)^n \\ & = \sum_{n=0}^{+\infty} \left[\sum_{k=n}^{+\infty} \frac{(-1)^k}{2k+1} \binom{k}{n} \right] (t-1)^n \end{aligned}$$

for $0 \leq t \leq 1$. This means that

$$\begin{aligned} \lim_{t \rightarrow 1^-} \frac{d^n}{dt^n} \left(\frac{\arctan \sqrt{t}}{\sqrt{t}} \right) &= n! \sum_{k=n}^{+\infty} \frac{(-1)^k}{2k+1} \binom{k}{n} \\ &= \frac{(-1)^n n!}{2n+1} {}_2F_1 \left(n + \frac{1}{2}, n+1; n + \frac{3}{2}; -1 \right) \end{aligned} \quad (4.3)$$

for $n \in \mathbb{N}_0$.

Let

$$H(z) = \frac{\arctan \sqrt{z}}{\sqrt{z}}, \quad \Re(z) > 0.$$

Then Wilf's function $W(z)$ can be regarded as the composition of $H(g(z))$, where $w = g(z) = 2e^{-z} - 1 \rightarrow 1$ as $z \rightarrow 0$. Accordingly, from (4.3), by the Faà di Bruno formula (2.5), we drive

$$\begin{aligned} W^{(n)}(z) &= [H(g(z))]^{(n)} \\ &= \sum_{k=1}^n H^{(k)}(w) B_{n,k}(g'(z), g''(z), \dots, g^{(n-k+1)}(z)) \\ &\rightarrow \sum_{k=1}^n \lim_{w \rightarrow 1} H^{(k)}(w) B_{n,k} \left(\lim_{z \rightarrow 0} g'(z), \lim_{z \rightarrow 0} g''(z), \dots, \lim_{z \rightarrow 0} g^{(n-k+1)}(z) \right) \\ &= \sum_{k=1}^n \frac{(-1)^k k!}{2k+1} {}_2F_1 \left(k + \frac{1}{2}, k+1; k + \frac{3}{2}; -1 \right) B_{n,k}(-2, 2, \dots, (-1)^{n-k+1} 2) \\ &= \sum_{k=1}^n \frac{(-1)^k k!}{2k+1} {}_2F_1 \left(k + \frac{1}{2}, k+1; k + \frac{3}{2}; -1 \right) (-1)^n 2^k B_{n,k}(1, 1, \dots, 1) \\ &= (-1)^n \sum_{k=1}^n (-1)^k S(n, k) \frac{(2k)!!}{2k+1} {}_2F_1 \left(k + \frac{1}{2}, k+1; k + \frac{3}{2}; -1 \right) \end{aligned}$$

as $z \rightarrow 0$ for $n \in \mathbb{N}$, where we used the identities (2.6) and (2.7). Consequently, the series expansion of Wilf's function $W(z)$ is

$$W(z) = \frac{\pi}{4} + \sum_{n=1}^{+\infty} (-1)^n \left[\sum_{k=1}^n (-1)^k S(n, k) \frac{(2k)!!}{2k+1} {}_2F_1 \left(k + \frac{1}{2}, k+1; k + \frac{3}{2}; -1 \right) \right] \frac{z^n}{n!}. \quad (4.4)$$

In [32, p. 109, Example 5.1], it is given that

$${}_2F_1 \left(\frac{1}{2}, 1; \frac{3}{2}; -z^2 \right) = \frac{\arctan z}{z}, \quad |z| < 1.$$

By virtue of Abel's limit theorem [2, p. 245, Theorem 9.31], we can take $z = 1$ and obtain

$${}_2F_1 \left(\frac{1}{2}, 1; \frac{3}{2}; -1 \right) = \frac{\pi}{4}.$$

Combining this with (4.4) brings out the series expansion (4.1). The proof of Theorem 4.1 is complete. \square

Corollary 4.1. For $n \in \mathbb{N}_0$, Gauss' hypergeometric function ${}_2F_1(\alpha, \beta; \gamma; z)$ satisfies

$${}_2F_1 \left(n + \frac{1}{2}, n+1; n + \frac{3}{2}; -1 \right) = \frac{(2n+1)!! \pi}{(2n)!!} \frac{\pi}{4} + \frac{2n+1}{2^{2n}} \sum_{k=1}^n (-1)^k \binom{2n-k}{n} \frac{2^{k/2}}{k} \sin \frac{3k\pi}{4}. \quad (4.5)$$

Proof. From the series expansion (3.6) in Theorem 3.2, it follows that

$$\lim_{z \rightarrow 1} \left(\frac{\arctan \sqrt{z}}{\sqrt{z}} \right)^{(n)} = \frac{(-1)^n}{2^n} \left[\frac{(2n-1)!! \pi}{4} + \frac{n!}{2^n} \sum_{k=1}^n (-1)^k \binom{2n-k}{n} \frac{2^{k/2}}{k} \sin \frac{3k\pi}{4} \right] \quad (4.6)$$

for $n \in \mathbb{N}_0$. Comparing (4.6) with (4.3) and simplifying lead to the explicit formula (4.5). The proof of Corollary 4.1 is complete. \square

Remark 4.1. The closed-form formula (4.5) in Corollary 4.1 can be reformulated as

$$\frac{\pi}{4} = \frac{(2n)!!}{(2n+1)!!} {}_2F_1\left(n + \frac{1}{2}, n + 1; n + \frac{3}{2}; -1\right) - \frac{(n!)^2}{(2n)!} \sum_{k=1}^n (-1)^k \binom{2n-k}{n} \frac{2^{k/2}}{k} \sin \frac{3k\pi}{4} \quad (4.7)$$

for all $n \in \mathbb{N}$. Since the quantities $2^{k/2} \sin \frac{3k\pi}{4}$ for $k \in \mathbb{N}$ are all integers and ${}_2F_1\left(n + \frac{1}{2}, n + 1; n + \frac{3}{2}; -1\right)$ is an infinite series, we can regard (4.7) as a series representation of the circular constant π .

Theorem 4.2. For $|z| < \ln 2$, Wilf's function $W(z)$ defined by (1.1) has Maclaurin's series expansion

$$W(z) = \sum_{n=0}^{+\infty} (-1)^n \left[\frac{\pi}{4} \sum_{k=0}^n (-1)^k S(n, k) (2k-1)!! + \sum_{k=1}^n (-1)^k S(n, k) \frac{k!}{2^k} \sum_{\ell=1}^k (-1)^\ell \binom{2k-\ell}{k} \frac{2^{\ell/2}}{\ell} \sin \frac{3\ell\pi}{4} \right] \frac{z^n}{n!}, \quad (4.8)$$

where $S(n, k)$ denotes the Stirling numbers of the second kind in (2.4).

Proof. The series expansion (4.8) comes from substituting the closed-form formula (4.5) in Corollary 4.1 into Maclaurin's series expansion (4.1) in Theorem 4.1 and simplifying. The proof of Theorem 4.2 is complete. \square

5. SERIES EXPANSIONS OF THE INVERSE HYPERBOLIC TANGENT FUNCTION

By means of the relation

$$\arctan z = -i \operatorname{arctanh}(iz), \quad z^2 \neq -1$$

in [1, p. 80, 4.4.22], from the series expansions in (3.1), (3.5), (3.6), and (4.8), we derive the following series expansions related to the inverse hyperbolic tangent function $\operatorname{arctanh} z$.

(1) For $|z - i| < \sqrt{2}$, we have

$$\frac{\operatorname{arctanh} z}{z} = \sum_{n=0}^{+\infty} \left[\frac{\pi}{4} + T(n) \right] (iz + 1)^n,$$

where $T(n)$ is defined by (3.2).

(2) For $|z - i| < 1$, we have

$$\frac{\operatorname{arctanh} z - \frac{\pi i}{4}}{z} = \sum_{n=1}^{+\infty} T(n) (iz + 1)^n.$$

(3) For $|z + 1| < 1$, we have

$$\frac{\operatorname{arctanh} \sqrt{z}}{\sqrt{z}} = \sum_{n=0}^{+\infty} \frac{1}{2^n} \left[(2n-1)!! \frac{\pi}{4} + \frac{n!}{2^n} \sum_{k=1}^n (-1)^k \binom{2n-k}{n} \frac{2^{k/2}}{k} \sin \frac{3k\pi}{4} \right] \frac{(z+1)^n}{n!}.$$

(4) For $|z| < \ln 2$, we have

$$\frac{\operatorname{arctanh} \sqrt{1 - 2e^{-z}}}{\sqrt{1 - 2e^{-z}}} = \sum_{n=0}^{+\infty} (-1)^n \left[\frac{\pi}{4} \sum_{k=0}^n (-1)^k S(n, k) (2k-1)!! + \sum_{k=1}^n (-1)^k S(n, k) \frac{k!}{2^k} \sum_{\ell=1}^k (-1)^\ell \binom{2k-\ell}{k} \frac{2^{\ell/2}}{\ell} \sin \frac{3\ell\pi}{4} \right] \frac{z^n}{n!},$$

where $S(n, k)$ denotes the Stirling numbers of the second kind in (2.4).

6. ANALYSIS ON COEFFICIENTS IN SERIES EXPANSIONS OF WILF'S FUNCTION

Considering the series expansion (4.8) in Theorem 4.2, we can see that the sequences b_n and c_n in (1.2) can be expressed as

$$b_n = \frac{1}{4} \frac{(-1)^n}{n!} \sum_{k=0}^n (-1)^k S(n, k) (2k-1)!! \quad (6.1)$$

and

$$c_n = \frac{(-1)^{n+1}}{n!} \sum_{k=1}^n (-1)^k S(n, k) \frac{k!}{2^k} \sum_{\ell=1}^k (-1)^\ell \binom{2k-\ell}{k} \frac{2^{\ell/2}}{\ell} \sin \frac{3\ell\pi}{4} \quad (6.2)$$

for $n \in \mathbb{N}_0$ respectively, where $S(n, k)$ denotes the Stirling numbers of the second kind generated in (2.3).

It is easy to see that the explicit formulas in (6.1) and (6.2) are closed-form expressions and that the sequence b_n is rational. Since $2^{\ell/2} \sin \frac{3\ell\pi}{4}$ for $\ell \in \mathbb{N}$ is an integer, the sequence c_n is also rational.

From closed-form formulas in (6.1) and (6.2), we can easily compute and list the first twelve values of the sequences b_n and c_n in Table 1. These twenty-four special values coincide with corresponding ones

TABLE 1. The first twelve values of b_n and c_n

n	0	1	2	3	4	5	6	7	8	9	10	11
b_n	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{7}{24}$	$\frac{35}{96}$	$\frac{113}{240}$	$\frac{1787}{2880}$	$\frac{16717}{20160}$	$\frac{2257}{2016}$	$\frac{315883}{207360}$	$\frac{4324721}{2073600}$	$\frac{447448}{155925}$
c_n	0	$\frac{1}{2}$	$\frac{3}{4}$	$\frac{11}{12}$	$\frac{55}{48}$	$\frac{71}{48}$	$\frac{2807}{1440}$	$\frac{8753}{3360}$	$\frac{94541}{26880}$	$\frac{694663}{145152}$	$\frac{47552791}{7257600}$	$\frac{719718067}{79833600}$

listed in the table on [33, p. 592].

Theorem 6.1. For $n \in \mathbb{N}_0$, the sequences b_n and c_n can be generated by

$$G(x) = \frac{1}{4\sqrt{2e^{-x}-1}} = \sum_{n=0}^{+\infty} b_n x^n, \quad |x| < \ln 2 \quad (6.3)$$

and

$$\frac{\frac{\pi}{4} - \arctan \sqrt{2e^{-x}-1}}{\sqrt{2e^{-x}-1}} = \sum_{n=0}^{+\infty} c_n x^n, \quad |x| < \ln 2. \quad (6.4)$$

Proof. In light of the Faà di Bruno formula (2.5) and the identities (2.6) and (2.7), we obtain

$$\begin{aligned} 4[G(x)]^{(n)} &= \sum_{k=0}^n (v^{-1/2})^{(k)} B_{n,k}(-2e^{-x}, 2e^{-x}, \dots, (-1)^{n-k+1} 2e^{-x}) \\ &= \sum_{k=0}^n \left\langle -\frac{1}{2} \right\rangle_k v^{-1/2-k} (-1)^n 2^k e^{-kx} B_{n,k}(1, 1, \dots, 1) \\ &= (-1)^n \sum_{k=0}^n (-1)^k (2k-1)!! (2e^{-x}-1)^{-1/2-k} e^{-kx} S(n, k) \\ &\rightarrow (-1)^n \sum_{k=0}^n (-1)^k S(n, k) (2k-1)!! \end{aligned}$$

as $x \rightarrow 0$ for $n \in \mathbb{N}_0$, where $v = v(x) = 2e^{-x} - 1$. This derivative implies that, via comparing with the explicit expression of b_n in (6.1), the function $G(x)$ defined on the interval $(-\infty, \ln 2)$ is a generating function of the sequence b_n for $n \in \mathbb{N}_0$.

From the series expansions (1.2) and (6.3), we derive

$$\sum_{n=0}^{+\infty} c_n x^n = \pi G(x) - W(x) = \frac{\frac{\pi}{4} - \arctan \sqrt{2e^{-x}-1}}{\sqrt{2e^{-x}-1}},$$

the series expansion (6.4) is thus proved. The proof of Theorem 6.1 is complete. \square

Remark 6.1. We can regard the generating function in (6.4) as a composite of the functions $\frac{\frac{\pi}{4} - \arctan z}{z}$ and $z = z(x) = \sqrt{g(x)} = \sqrt{2e^{-x} - 1}$. Therefore, by the aid of the Faà di Bruno formula (2.5), we acquire

$$\begin{aligned} \left(\frac{\frac{\pi}{4} - \arctan \sqrt{2e^{-x} - 1}}{\sqrt{2e^{-x} - 1}} \right)^{(n)} &= \sum_{k=0}^n \left(\frac{\frac{\pi}{4} - \arctan z}{z} \right)^{(k)} B_{n,k}(z'(x), z''(x), \dots, z^{(n-k+1)}(x)) \\ &\rightarrow \sum_{k=0}^n (-1)^{k+1} k! T(k) B_{n,k} \left(-1, 0, -1, -3, \dots, (-1)^{n-k+1} \sum_{\ell=0}^{n-k+1} (-1)^{\ell-1} S(n-k+1, \ell) (2\ell-3)!! \right) \\ &= - \sum_{k=0}^n k! T(k) B_{n,k} \left(1, 0, 1, 3, 16, 105, \dots, (-1)^{n-k} \sum_{\ell=0}^{n-k+1} (-1)^{\ell-1} S(n-k+1, \ell) (2\ell-3)!! \right) \\ &= (-1)^{n+1} \sum_{k=0}^n T(k) \sum_{\ell=k}^n (-1)^\ell S(n, \ell) P(\ell, k) \\ &= (-1)^{n+1} \sum_{\ell=0}^n (-1)^\ell S(n, \ell) \frac{\ell!}{2^\ell} \sum_{k=1}^{\ell} (-1)^k \binom{2\ell-k}{\ell} \frac{2^{k/2}}{k} \sin \frac{3k\pi}{4} \end{aligned}$$

as $x \rightarrow 0$ and $z \rightarrow 1$, where we used the explicit formula (3.16) in Theorem 3.4 and the identity (3.14). The closed-form expression (6.2) of the sequence c_n is thus recovered.

Remark 6.2. The generating function $G(x)$ for $|x| < \ln 2$ in (6.3) can also be rearranged as

$$\begin{aligned} G(x) &= \frac{e^{x/2}}{4\sqrt{2}\sqrt{1-e^x/2}} \\ &= \frac{1}{4\sqrt{2}} \sum_{j=0}^{+\infty} \binom{-1/2}{j} \left(-\frac{1}{2}\right)^j e^{(j+1/2)x} \\ &= \frac{1}{4\sqrt{2}} \sum_{j=0}^{+\infty} \binom{-1/2}{j} \left(-\frac{1}{2}\right)^j \sum_{n=0}^{+\infty} \frac{1}{n!} \left(j + \frac{1}{2}\right)^n x^n \\ &= \frac{1}{4\sqrt{2}} \sum_{n=0}^{+\infty} \frac{1}{n!} \left[\sum_{j=0}^{+\infty} \binom{-1/2}{j} \left(-\frac{1}{2}\right)^j \left(j + \frac{1}{2}\right)^n \right] x^n \\ &= \frac{1}{4\pi} \sum_{n=0}^{+\infty} \left[\sum_{j=0}^{+\infty} \left(j + \frac{1}{2}\right)^{n-1} \frac{2^{j+1/2}}{\binom{2j+1}{j+1/2}} \right] \frac{x^n}{n!}. \end{aligned}$$

Then the infinite series representation

$$b_n = \frac{1}{4\pi} \frac{1}{n!} \sum_{j=0}^{+\infty} \left(j + \frac{1}{2}\right)^{n-1} \frac{2^{j+1/2}}{\binom{2j+1}{j+1/2}}, \quad n \in \mathbb{N}_0, \quad (6.5)$$

which appears at the site <https://oeis.org/A014307>, is proved.

Theorem 6.2. *The following conclusions are valid:*

- (1) *the function $G(x)$ is logarithmically completely monotonic on $(-\infty, \ln 2)$;*
- (2) *the sequences*

$$4n!b_n = (-1)^n \sum_{k=0}^n (-1)^k S(n, k) (2k-1)!!, \quad n \in \mathbb{N}_0 \quad (6.6)$$

and

$$(-1)^n \sum_{k=1}^n (-1)^k S(n, k) (2k-2)!!, \quad n \in \mathbb{N} \quad (6.7)$$

are positive, increasing, and logarithmically convex.

Proof. For $n \in \mathbb{N}$ and $v = v(x) = 2e^{-x} - 1$, making use of the Faà di Bruno formula (2.5) and the identities (2.6) and (2.7), we acquire

$$[\ln G(x)]^{(n)} = \left(\ln \frac{1}{4\sqrt{2}e^{-x} - 1} \right)^{(n)}$$

$$\begin{aligned}
&= -\frac{1}{2} \sum_{k=1}^n (\ln v)^{(k)} B_{n,k}(-2e^{-x}, 2e^{-x}, \dots, (-1)^{n-k+1} 2e^{-x}) \\
&= -\frac{1}{2} \sum_{k=1}^n \frac{(-1)^{k-1} (k-1)!}{v^k} (-1)^n 2^k e^{-kx} B_{n,k}(1, 1, \dots, 1) \\
&= (-1)^n \sum_{k=1}^n (-1)^k \frac{(2k-2)!!}{(2e^{-x}-1)^k} e^{-kx} S(n, k) \\
&\rightarrow (-1)^n \sum_{k=1}^n (-1)^k S(n, k) (2k-2)!!
\end{aligned}$$

as $x \rightarrow 0$.

On the other hand, it is obvious that the first logarithmic derivative

$$[\ln G(x)]' = \left(\ln \frac{1}{4\sqrt{2e^{-x}-1}} \right)' = \frac{1}{2-e^x} = \frac{1}{2} \sum_{k=0}^{+\infty} \left(\frac{e^x}{2} \right)^k, \quad x < \ln 2$$

is absolutely monotonic on $(-\infty, \ln 2)$. Hence, the function $G(x)$ is logarithmically absolutely monotonic on $(-\ln 2, +\infty)$. As a result, the positivity of the sequence in (6.7) is valid.

Basing on [7, Definition 1 and Theorem 1] recited in Section 2, from the logarithmically absolute monotonicity of $G(x)$, we derive that the function $G(x)$ is absolutely monotonic on $(-\infty, \ln 2)$. Combining this result with the proof of Theorem 6.1 leads to the positivity of the sequence in (6.6).

Applying the positivity of the sequence in (6.6) to the second closed-form formula in (6.1) for b_n reveals that the sequence b_n for $n \in \mathbb{N}_0$ is positive.

In [13, p. 369], it was stated that, if f is a completely monotonic function on $(0, +\infty)$ such that $f^{(k)}(t) \neq 0$ for $k \geq 0$, then the sequence $(-1)^k f^{(k)}(t)$ for $k \geq 0$ is logarithmically convex. Applying this result to the functions $G(-x)$ and $\ln G(-x)$, both of which are completely monotonic on $(0, +\infty) \subset (-\ln 2, +\infty)$, yields the logarithmic convexity of the sequences in (6.6) and (6.7).

From the logarithmic convexity of the sequences in (6.6) and (6.7) and the increasing property of the first few values of the sequences in (6.6) and (6.7) respectively, we conclude that both of the sequences in (6.6) and (6.7) are increasing. The proof of Theorem 6.2 is complete. \square

Remark 6.3. A few of results of the integer sequence $4n!b_n$ for $n \in \mathbb{N}_0$ have been being collected at the site <https://oeis.org/A014307>. For example, the recursive relation

$$(n+1)!b_{n+1} = \frac{1}{4} + \sum_{j=1}^n \left[\binom{n+1}{j} - 1 \right] j!b_j, \quad n \in \mathbb{N}_0$$

is listed at the site, which implies the positivity and increasing property of the sequence $n!b_n$ for $n \in \mathbb{N}_0$.

From the complete monotonicity of the functions $G(-x)$ and $\ln G(-x)$ on the infinite interval $(0, +\infty) \subset (-\ln 2, +\infty)$ and a number of properties of completely monotonic functions collected in [13, Chapter XIII], [34, Chapter IV], and the monograph [30], we can deduce many properties, which do not appear at the site <https://oeis.org/A014307>, of the sequence $4n!b_n$ for $n \in \mathbb{N}_0$, as done in [16, 24] and closely related references therein. Due to the limitation of length, we do not write down them in details.

Remark 6.4. A lot of conclusions for the sequence $2n!c_n$ with $n \in \mathbb{N}$ have been listed at <http://oeis.org/A180875>. At the site, we do not see the generating function in (6.4) for the sequence c_n with $n \in \mathbb{N}_0$.

Theorem 6.3. *The limits*

$$\lim_{n \rightarrow +\infty} b_n = +\infty \tag{6.8}$$

and

$$\lim_{n \rightarrow +\infty} \frac{c_n}{b_n} = \pi \tag{6.9}$$

are valid. Consequently, we have

$$\lim_{n \rightarrow +\infty} c_n = +\infty \tag{6.10}$$

Proof. In [4, p. 2, (1.2)], the double inequality

$$\frac{1}{\sqrt{\pi(n + 4/\pi - 1)}} \leq \frac{(2n - 1)!!}{(2n)!!} < \frac{1}{\sqrt{\pi(n + 1/4)}}, \quad n \in \mathbb{N} \quad (6.11)$$

was recited, where the constants $\frac{4}{\pi} - 1$ and $\frac{1}{4}$ in (6.11) are the best possible in the sense that these two constants cannot be replaced by smaller and bigger ones, respectively. In [4, Theorem 1.1], the double inequality

$$\frac{\sqrt{\pi}}{2\sqrt{n + 9\pi/16 - 1}} \leq \frac{(2n)!!}{(2n + 1)!!} < \frac{\sqrt{\pi}}{2\sqrt{n + 3/4}}, \quad n \in \mathbb{N} \quad (6.12)$$

was established, where the constants $\frac{9\pi}{16} - 1$ and $\frac{3}{4}$ in (6.12) are the best possible in the sense that these two numbers cannot be replaced by smaller and bigger ones, respectively. See also [8, p. 658, Theorem 4].

The infinite series representation (6.5) can be reformulated as

$$b_n = \frac{1}{n!} \sum_{j=0}^{+\infty} \left(j + \frac{1}{2}\right)^{n-1} \frac{(2j+1)!!}{(2j)!!} \frac{1}{2^{j+7/2}}, \quad n \in \mathbb{N}_0. \quad (6.13)$$

Substituting (6.12) into (6.13) yields

$$\begin{aligned} b_n &> \frac{1}{n!} \left[\frac{1}{2^{n-5/2}} + \frac{1}{2^{5/2}\sqrt{\pi}} \sum_{j=1}^{+\infty} \left(j + \frac{1}{2}\right)^{n-1} \frac{\sqrt{j + 9\pi/16 - 1}}{2^j} \right] \\ &> \frac{1}{n!} \left[\frac{1}{2^{n-5/2}} + \frac{1}{2^{5/2}\sqrt{\pi}} \sum_{j=1}^{+\infty} \left(j + \frac{1}{2}\right)^{n-1/2} \frac{1}{2^j} \right] \\ &= \frac{1}{n!} \left[\frac{1}{2^{n-5/2}} + \frac{1}{2^{7/2}\sqrt{\pi}} \Phi\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}\right) \right] \end{aligned}$$

for $n \in \mathbb{N}_0$, where $\Phi(z, s, a)$ is the Lerch transcendent which is defined [15, p. 612, Section 25.14] by

$$\Phi(z, s, a) = \sum_{j=0}^{+\infty} \frac{z^j}{(j+a)^s}, \quad |z| < 1, \quad a \neq 0, -1, -2, \dots$$

Let $0 < |\lambda| < 1$ and $\lambda \notin (-1, 0)$. Corollary 4 in [14, p. 29] reads that the limit

$$\lim_{\Re s \rightarrow -\infty} \frac{(-\ln \lambda)^{1-s} \Phi(\lambda, s, z)}{\Gamma(1-s)} = \frac{1}{\lambda^z}$$

converges uniformly for z on compact subset of \mathbb{C} . As a result, we arrive at

$$\begin{aligned} \lim_{n \rightarrow +\infty} \left[\frac{1}{n!} \Phi\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}\right) \right] &= \lim_{n \rightarrow +\infty} \left[\frac{(\ln 2)^{1/2+n}}{\Gamma(1/2+n)} \Phi\left(\frac{1}{2}, \frac{1}{2} - n, \frac{3}{2}\right) \right] \lim_{n \rightarrow +\infty} \frac{\Gamma(1/2+n)}{n!(\ln 2)^{1/2+n}} \\ &= \frac{\sqrt{\pi}}{2^{3/2}(\ln 2)^{1/2}} \lim_{n \rightarrow +\infty} \frac{(2n-1)!!}{(2n)!!(\ln 2)^n} \\ &= +\infty, \end{aligned}$$

where we used in the last step the double inequality (6.11). Consequently, the limit (6.8) is proved.

By virtue of the necessity of convergence for power series expansions to be convergent, we see that the coefficients a_n in the series expansion (1.2) satisfy

$$a_n = b_n \pi - c_n = b_n \left(\pi - \frac{c_n}{b_n} \right) \rightarrow 0, \quad n \rightarrow +\infty.$$

Accordingly, from the limit (6.8), we see that it is mandatory that

$$\lim_{n \rightarrow +\infty} \left(\pi - \frac{c_n}{b_n} \right) = 0.$$

The limit (6.9) is thus verified.

From any one of the limits (6.8) and 6.9, we can trivially deduce the limit (6.10). The proof of Theorem 6.3 is complete. \square

Remark 6.5. Since b_n and c_n for all $n \in \mathbb{N}_0$ are rational numbers, we can regard the limit (6.9) as an asymptotic rational approximation to the circular constant π .

Remark 6.6. At the sites <https://oeis.org/A014307> and <http://oeis.org/A180875>, there exist two asymptotic formulas

$$4n!b_n \sim \frac{\sqrt{2}n^n}{e^n(\ln 2)^{n+1/2}} = \sqrt{\frac{2}{\ln 2}} \left(\frac{n}{e \ln 2}\right)^n, \quad n \rightarrow +\infty \quad (6.14)$$

and

$$2(n+1)!c_{n+1} \sim \pi \frac{n^{n+1}}{\sqrt{2}e^n(\ln 2)^{n+3/2}} = \frac{e\pi}{\sqrt{2}\ln 2} \left(\frac{n}{e \ln 2}\right)^{n+1}, \quad n \rightarrow +\infty, \quad (6.15)$$

respectively. On 4 September 2021, Dr. Professor Vaclav Kotesovec (kotesovec2@gmail.com, retired) told us that, using the Maple module equivalent, these two asymptotic formulas (6.14) and (6.15) can be proved as showed in Figures 1 and 2. Consequently, it is easy to see that

```
> march('open', "C:\maple\MLA\algotlib.mla");
      libname := "C:\Program Files\Maple 16\lib", ".", "C:\maple\MLA\algotlib.mla"
> simplify(convert(equivalent(sqrt(exp(x)/(2-exp(x))), x, n, 1), polynomial)) assuming n > 0;
      ln(2)^(1/2-n)
      -----
      sqrt(pi) sqrt(n)
>
> simplify((n^n/exp(n)) * sqrt(2*Pi*n) * convert(equivalent(sqrt(exp(x)/(2-exp(x))), x, n, 1), polynomial)) assuming n
> 0;
      sqrt(2) n^n ln(2)^(1/2-n) e^-n
```

FIGURE 1. Asymptotic formula of b_n

```
> march('open', "C:\maple\MLA\algotlib.mla");
      libname := "C:\Program Files\Maple 16\lib", ".", "C:\maple\MLA\algotlib.mla"
> |
>
> simplify((n^n/exp(n)) * sqrt(2*Pi*n) * convert(equivalent(2*((-1*log(1*(exp(x/2)/sqrt(2)) + sqrt(1-(exp(x/2)
/sqrt(2))^2))) - Pi/4) * sqrt(exp(x)/(2-exp(x))^3) + exp(x)/(2-exp(x))), x, n, 1), polynomial))
      assuming n > 0;
      1/2 sqrt(2) n^(n+1) pi ln(2)^(3/2-n) e^-n
```

FIGURE 2. Asymptotic formula of c_n

$$\frac{c_n}{b_n} \sim e\pi \left(1 - \frac{1}{n}\right)^n \rightarrow \pi, \quad n \rightarrow +\infty.$$

The limit (6.9) is thus proved again.

From asymptotic formulas (6.14) and (6.15), we derive the asymptotic formula

$$a_n = b_n\pi - c_n \sim \frac{\pi e}{2\sqrt{2}\ln 2} \frac{n^n}{n!} \frac{1}{(e \ln 2)^n} \left[\frac{1}{e} - \left(1 - \frac{1}{n}\right)^n \right], \quad n \rightarrow +\infty.$$

Remark 6.7. We conjecture that the sequences b_n and c_n are increasing and convex in $n \in \mathbb{N}_0$.

Theorem 6.4. *Let*

$$d_n = \frac{(-1)^n}{n!} \sum_{k=0}^n (-1)^k S(n, k) (2k-3)!!, \quad n \in \mathbb{N}_0. \quad (6.16)$$

For $n \in \mathbb{N}$, the sequence d_n is nonnegative.

Proof. Differentiating on both sides of the series expansion (3.15) in Theorems 3.3 and simplifying give

$$\frac{1}{\sqrt{e^x(2-e^x)}} = \sum_{n=0}^{+\infty} (n+1)d_{n+1}x^n, \quad |x| < \ln 2. \quad (6.17)$$

Taking the logarithm and differentiating result in

$$\left[\ln \frac{1}{\sqrt{e^x(2-e^x)}} \right]' = -\frac{1}{2}[x + \ln(2-e^x)]' = \frac{1}{2-e^x} - 1 = \frac{1}{2} \left[\sum_{k=1}^{+\infty} \left(\frac{e^x}{2} \right)^k - 1 \right]$$

which is absolutely monotonic on the interval $[0, \ln 2)$. This means that the function $\frac{1}{\sqrt{e^x(2-e^x)}}$ is (logarithmically) absolutely monotonic on $[0, \ln 2)$. Hence, the derivatives

$$\lim_{x \rightarrow 0^+} \left[\frac{1}{\sqrt{e^x(2-e^x)}} \right]^{(n)} = (n+1)!d_{n+1} \geq 0, \quad n \in \mathbb{N}_0.$$

The proof of Theorem 6.4 is complete. □

Remark 6.8. We can rewrite the generating function in (6.17) as

$$\begin{aligned} \frac{1}{\sqrt{e^x(2-e^x)}} &= \frac{e^{-x/2}}{\sqrt{2}\sqrt{1-e^x/2}} \\ &= \frac{1}{\sqrt{2}} \sum_{j=0}^{+\infty} \binom{-1/2}{j} \left(-\frac{1}{2}\right)^j e^{(j-1/2)x} \\ &= \frac{1}{\sqrt{2}} \sum_{j=0}^{+\infty} \binom{-1/2}{j} \left(-\frac{1}{2}\right)^j \sum_{n=0}^{+\infty} \frac{1}{n!} \left(j - \frac{1}{2}\right)^n x^n \\ &= \frac{1}{\sqrt{2}} \sum_{n=0}^{+\infty} \frac{1}{n!} \left[\sum_{j=0}^{+\infty} \binom{-1/2}{j} \left(-\frac{1}{2}\right)^j \left(j - \frac{1}{2}\right)^n \right] x^n \\ &= \frac{1}{\pi} \sum_{n=0}^{+\infty} \left[\sum_{j=0}^{+\infty} \frac{(j-1/2)^n}{j+1/2} \frac{2^{j+1/2}}{\binom{2j+1}{j+1/2}} \right] \frac{x^n}{n!}. \end{aligned}$$

Therefore, we acquire an infinite series representation

$$d_n = \frac{1}{\pi} \frac{1}{n!} \sum_{j=0}^{+\infty} \frac{(j-1/2)^n}{j+1/2} \frac{2^{j+1/2}}{\binom{2j+1}{j+1/2}}, \quad n \in \mathbb{N}. \quad (6.18)$$

By a similar approach used in the proof of the limit (6.8), as in the proof of Theorem 6.3, we can confirm that $\lim_{n \rightarrow +\infty} d_n = +\infty$.

Remark 6.9. The sequence $n!d_n$ is an integer sequence. The first few values of $n!d_n$ for $0 \leq n \leq 21$ are

$$\begin{aligned} &-1, \quad 1, \quad 0, \quad 1, \quad 3, \quad 16, \quad 105, \quad 841, \quad 7938, \quad 86311, \quad 1062435, \\ &14605306, \quad 221790723, \quad 3687263581, \quad 66609892440, \quad 1299237505021, \\ &27213601303983, \quad 609223983928576, \quad 14516520372130245, \\ &366820998284761861, \quad 9798039716677045218, \quad 275837214061454446171. \end{aligned}$$

When $n \in \mathbb{N}$, that is, when removing off the first term -1 , the sequence $n!d_n$ appears at the site <https://oeis.org/A014304>.

We guess that the sequences $n!d_n$ and d_n for $n \in \mathbb{N}$ are increasing and convex.

Theorem 6.5. *The sequences b_n and d_n are connected by the relations*

$$b_n = \frac{1}{4} \begin{vmatrix} 1 & -1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & \cdots & 0 & 0 & 0 \\ \frac{1}{6} & 0 & 1 & -1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ d_{n-2} & d_{n-3} & d_{n-4} & d_{n-5} & \cdots & 1 & -1 & 0 \\ d_{n-1} & d_{n-2} & d_{n-3} & d_{n-4} & \cdots & 0 & 1 & -1 \\ d_n & d_{n-1} & d_{n-2} & d_{n-3} & \cdots & \frac{1}{6} & 0 & 1 \end{vmatrix} \quad (6.19)$$

and

$$d_n = (-1)^{n-1} 4^n \begin{vmatrix} \frac{1}{4} & \frac{1}{4} & 0 & 0 & \cdots & 0 & 0 & 0 \\ \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & 0 & \cdots & 0 & 0 & 0 \\ \frac{7}{24} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ b_{n-2} & b_{n-3} & b_{n-4} & b_{n-5} & \cdots & \frac{1}{4} & \frac{1}{4} & 0 \\ b_{n-1} & b_{n-2} & b_{n-3} & b_{n-4} & \cdots & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \\ b_n & b_{n-1} & b_{n-2} & b_{n-3} & \cdots & \frac{7}{24} & \frac{1}{4} & \frac{1}{4} \end{vmatrix} \quad (6.20)$$

for $n \in \mathbb{N}$.

Proof. Wronski's theorem [9, p. 17, Theorem 1.3] reads that, if $\alpha_0 \neq 0$ and

$$P(x) = \alpha_0 + \alpha_1 x + \alpha_2 x^2 + \cdots$$

is a formal series, then the coefficients of the reciprocal series

$$\frac{1}{P(x)} = \beta_0 + \beta_1 x + \beta_2 x^2 + \cdots$$

are given by

$$\beta_n = \frac{(-1)^n}{\alpha_0^{n+1}} \begin{vmatrix} \alpha_1 & \alpha_0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ \alpha_2 & \alpha_1 & \alpha_0 & 0 & \cdots & 0 & 0 & 0 \\ \alpha_3 & \alpha_2 & \alpha_1 & \alpha_0 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ \alpha_{n-2} & \alpha_{n-3} & \alpha_{n-4} & \alpha_{n-5} & \cdots & \alpha_1 & \alpha_0 & 0 \\ \alpha_{n-1} & \alpha_{n-2} & \alpha_{n-3} & \alpha_{n-4} & \cdots & \alpha_2 & \alpha_1 & \alpha_0 \\ \alpha_n & \alpha_{n-1} & \alpha_{n-2} & \alpha_{n-3} & \cdots & \alpha_3 & \alpha_2 & \alpha_1 \end{vmatrix}, \quad n = 1, 2, \dots \quad (6.21)$$

The formula (6.21) can also be found in [10, p. 347], [18, Lemma 2.4], [26, Lemma 2.3], and [29, Section 2].

Using Wronski's theorem (6.21), from the series expansions (3.15) and (6.3) in Theorems 3.3 and 6.1 respectively, we arrive at the relations (6.19) and (6.20) immediately. The proof of Theorem 6.5 is thus complete. \square

Theorem 6.6. *For $n \in \mathbb{N}_0$, let*

$$e_n = -\frac{1}{(2n)!!} \sum_{k=0}^n \binom{n}{k} 2^k \sum_{\ell=0}^k S(k, \ell) \frac{(2\ell-3)!!}{2^\ell}.$$

Then Maclaurin's series expansion

$$\sqrt{e^x(2-e^x)} = \sum_{n=0}^{+\infty} e_n x^n, \quad |x| < \ln 2 \quad (6.22)$$

holds and the relations

$$d_{n+1} = \frac{(-1)^n}{n+1} \begin{vmatrix} 0 & 1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ -\frac{1}{2} & 0 & 1 & 0 & \cdots & 0 & 0 & 0 \\ -\frac{1}{2} & -\frac{1}{2} & 0 & 1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ e_{n-2} & e_{n-3} & e_{n-4} & e_{n-5} & \cdots & 0 & 1 & 0 \\ e_{n-1} & e_{n-2} & e_{n-3} & e_{n-4} & \cdots & -\frac{1}{2} & 0 & 1 \\ e_n & e_{n-1} & e_{n-2} & e_{n-3} & \cdots & -\frac{1}{2} & -\frac{1}{2} & 0 \end{vmatrix} \quad (6.23)$$

and

$$e_n = (-1)^n \begin{vmatrix} 0 & 1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ \frac{1}{2} & 0 & 1 & 0 & \cdots & 0 & 0 & 0 \\ \frac{1}{2} & \frac{1}{2} & 0 & 1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ (n-1)d_{n-1} & (n-2)d_{n-2} & (n-3)d_{n-3} & (n-4)d_{n-4} & \cdots & 0 & 1 & 0 \\ nd_n & (n-1)d_{n-1} & (n-2)d_{n-2} & (n-3)d_{n-3} & \cdots & \frac{1}{2} & 0 & 1 \\ (n+1)d_{n+1} & nd_n & (n-1)d_{n-1} & (n-2)d_{n-2} & \cdots & \frac{1}{2} & \frac{1}{2} & 0 \end{vmatrix} \quad (6.24)$$

ar valid for $n \in \mathbb{N}$, where d_n for $n \in \mathbb{N}_0$ are defined by (6.16).

Proof. It is standard that

$$\begin{aligned} [\sqrt{e^x(2-e^x)}]^{(n)} &= (e^{x/2} \sqrt{2-e^x})^{(n)} \\ &= \sum_{k=0}^n \binom{n}{k} (\sqrt{2-e^x})^{(k)} (e^{x/2})^{(n-k)} \\ &= \sum_{k=0}^n \binom{n}{k} \frac{e^{x/2}}{2^{n-k}} \sum_{\ell=0}^k \left\langle \frac{1}{2} \right\rangle_{\ell} (2-e^x)^{1/2-\ell} \mathbf{B}_{k,\ell}(-e^x, -e^x, \dots, -e^x) \\ &= -\frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} 2^k e^{x/2} \sum_{\ell=0}^k \frac{(2\ell-3)!!}{2^\ell} (2-e^x)^{1/2-\ell} e^{\ell x} S(k, \ell) \\ &\rightarrow -\frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} 2^k \sum_{\ell=0}^k \frac{(2\ell-3)!!}{2^\ell} S(k, \ell) \end{aligned}$$

as $x \rightarrow 0$, where we used the Faà di Bruno formula (2.5) and two identities (2.6) and (2.7).

Combining the series expansions (6.17) and (6.22) with Wronski's theorem in (6.21) recited in the proof of Theorem 6.5 leads to two relations (6.23) and (6.24) for $n \in \mathbb{N}$. The proof of Theorem 6.6 is complete. \square

Remark 6.10. The generating function in (6.22) can be alternatively expanded as

$$\begin{aligned} \sqrt{e^x(2-e^x)} &= \sqrt{2} e^{x/2} \sqrt{1 - \frac{e^x}{2}} \\ &= \sqrt{2} \sum_{j=0}^{+\infty} \binom{1/2}{j} \left(-\frac{1}{2}\right)^j e^{(j+1/2)x} \\ &= \sqrt{2} \sum_{j=0}^{+\infty} \binom{1/2}{j} \left(-\frac{1}{2}\right)^j \sum_{n=0}^{+\infty} \left(j + \frac{1}{2}\right)^n \frac{x^n}{n!} \\ &= \sqrt{2} \sum_{n=0}^{+\infty} \left[\sum_{j=0}^{+\infty} \binom{1/2}{j} \left(-\frac{1}{2}\right)^j \left(j + \frac{1}{2}\right)^n \right] \frac{x^n}{n!} \\ &= -\frac{1}{\pi} \sum_{n=0}^{+\infty} \left[\sum_{j=0}^{+\infty} \frac{(j+1/2)^{n-1}}{j-1/2} \frac{2^{j+1/2}}{\binom{2j+1}{j+1/2}} \right] \frac{x^n}{n!}. \end{aligned}$$

Accordingly, we obtain an infinite series representation

$$e_n = -\frac{1}{\pi} \frac{1}{n!} \sum_{j=0}^{+\infty} \frac{(j+1/2)^{n-1}}{j-1/2} \frac{2^{j+1/2}}{\binom{2j+1}{j+1/2}}, \quad n \in \mathbb{N}_0. \quad (6.25)$$

By a similar approach used in the proof of the limit (6.8), as in the proof of Theorem 6.3, we can confirm that $\lim_{n \rightarrow +\infty} e_n = -\infty$.

Remark 6.11. The sequence $n!e_n$ for $n \in \mathbb{N}_0$ is an integer sequence. Its first few values for $0 \leq n \leq 21$ are

$$\begin{aligned} &1, \quad 0, \quad -1, \quad -3, \quad -10, \quad -45, \quad -271, \quad -2058, \quad -18775, \\ &\quad -199335, \quad -2410516, \quad -32683563, \quad -490870315, \\ &\quad -8087188200, \quad -144994236661, \quad -2810079139143, \\ &\quad -58536519252130, \quad -1304198088413265, \quad -30946462816602331, \\ &\quad -779104979758256298, \quad -20742005411397108595, \quad -582214473250983046155. \end{aligned}$$

We do not find the integer sequence $n!e_n$ for $n \in \mathbb{N}_0$ or its subsequences at the site <https://oeis.org/>.

7. CONCLUSIONS

In this paper, with the aid of the Faà di Bruno formula (2.5), by virtue of several identities from (2.6) to (2.10) for the Bell polynomials of the second kind, with the help of two combinatorial identities (2.11) and (2.13), and by means of Wronski's theorem (6.21) and the (logarithmically) complete monotonicity of generating functions (6.3) and (6.4), we established Taylor's series expansions (3.1), (3.5), (3.6), and those in Section 5, found out Maclaurin's series expansions (4.1) and (4.8) for Wilf's function (1.1), and analyzed some properties, including two generating functions in Theorem 6.1 and including the positivity, monotonicity, and logarithmic convexity in Theorems 6.2 and 6.4, of the coefficients b_n and c_n in Maclaurin's series expansion (1.2) of Wilf's function (1.1). These coefficients are closed-form expressions in (6.1) and (6.2). We also derived a closed-form formula (4.5) for a sequence of special values ${}_2F_1(n + \frac{1}{2}, n + 1; n + \frac{3}{2}; -1)$ of Gauss' hypergeometric function ${}_2F_1(\alpha, \beta; \gamma; z)$ defined in (2.2), discovered a closed-form formula (3.16) in Theorem 3.4 for a sequence of special values $B_{n,k}(1, 0, 1, 3, 16, 105, \dots)$ of the Bell polynomials of the second kind $B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$, presented several infinite series representations (6.18) and (6.25), recovered an asymptotic rational approximation in Theorem 6.3 to the circular constant π , and connected several integer sequences by determinants in Theorems 6.5 and 6.6.

8. DECLARATIONS

Acknowledgements: The authors thank

- (1) Dr. Darij Grinberg (Drexel University, Germany; darijgrinberg@gmail.com; <http://www.cip.ifi.lmu.de/~grinberg/>) and others for their discussions and supplying the idea of the descending induction used in the proof of Lemma 2.1 at the site <https://mathoverflow.net/q/402822>;
- (2) Dr. Professor Vaclav Kotesovec (kotesovec2@gmail.com, retired) for his providing Figures 1 and 2 in this paper via e-mail;
- (3) Dr. Professor Hjalmar Rosengren (Chalmers University of Technology and University of Gothenburg, Sweden) and others for their discussions and an alternative proof of the infinite series representation (6.5) at the site <https://mathoverflow.net/q/403281>.

Availability of data and material: Data sharing is not applicable to this article as no new data were created or analyzed in this study.

Competing interests: The authors declare that they have no conflict of competing interests.

Funding: Not applicable.

Authors' contributions: All authors contributed equally to the manuscript and read and approved the final manuscript.

REFERENCES

- [1] ABRAMOWITZ, M. AND STEGUN, I. A. (eds.): *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Dover Publications, Inc., New York, 1992, reprint of the 1972 edition.

- [2] APOSTOL, T. M.: *Mathematical Analysis*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1974, 2nd edn.
- [3] BERG, C.: *Integral representation of some functions related to the gamma function*, *Mediterr. J. Math.*, **1** (2004), No. 4, 433–439.
URL <https://doi.org/10.1007/s00009-004-0022-6>
- [4] CAO, J., NIU, D.-W., AND QI, F.: *A Wallis type inequality and a double inequality for probability integral*, *Aust. J. Math. Anal. Appl.*, **4** (2007), No. 1, Art. 3, 6 pages.
URL <http://ajmaa.org/cgi-bin/paper.pl?string=v4n1/V4I1P3.tex>
- [5] CHARALAMBIDES, C. A.: *Enumerative Combinatorics*, CRC Press Series on Discrete Mathematics and its Applications, Chapman & Hall/CRC, Boca Raton, FL, 2002.
- [6] COMTET, L.: *Advanced Combinatorics*, D. Reidel Publishing Co., Dordrecht, 1974, enlarged edn., the art of finite and infinite expansions.
URL <https://doi.org/10.1007/978-94-010-2196-8>
- [7] GUO, B.-N. AND QI, F.: *A property of logarithmically absolutely monotonic functions and the logarithmically complete monotonicity of a power-exponential function*, *Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys.*, **72** (2010), No. 2, 21–30.
- [8] GUO, B.-N. AND QI, F.: *A class of completely monotonic functions involving divided differences of the psi and tri-gamma functions and some applications*, *J. Korean Math. Soc.*, **48** (2011), No. 3, 655–667.
URL <https://doi.org/10.4134/JKMS.2011.48.3.655>
- [9] HENRIK, P.: *Applied and Computational Complex Analysis. Vol. 1*, Wiley Classics Library, John Wiley & Sons, Inc., New York, 1988, reprint of the 1974 original, A Wiley-Interscience Publication.
- [10] INSELBERG, A.: *On determinants of Toeplitz-Hessenberg matrices arising in power series*, *J. Math. Anal. Appl.*, **63** (1978), No. 2, 347–353.
URL [https://doi.org/10.1016/0022-247X\(78\)90080-X](https://doi.org/10.1016/0022-247X(78)90080-X)
- [11] LANG, S.: *Topics in Cohomology of Groups*, vol. 1625 of *Lecture Notes in Mathematics*, Springer-Verlag, Berlin, 1996, translated from the 1967 French original by the author, Chapter X based on letters written by John Tate.
URL <https://doi.org/10.1007/BFb0092624>
- [12] LUKE, Y. L.: *The Special Functions and Their Approximations, Vol. I*, Mathematics in Science and Engineering, Vol. 53, Academic Press, New York-London, 1969.
- [13] MITRINOVIĆ, D. S., PEČARIĆ, J. E., AND FINK, A. M.: *Classical and New Inequalities in Analysis*, vol. 61 of *Mathematics and its Applications (East European Series)*, Kluwer Academic Publishers Group, Dordrecht, 1993.
URL <https://doi.org/10.1007/978-94-017-1043-5>
- [14] NAVAS, L. M., RUIZ, F. J., AND VARONA, J. L.: *Asymptotic behavior of the Lerch transcendent function*, *J. Approx. Theory*, **170** (2013), 21–31.
URL <https://doi.org/10.1016/j.jat.2012.03.006>
- [15] OLVER, F. W. J., LOZIER, D. W., BOISVERT, R. F., AND CLARK, C. W. (eds.): *NIST Handbook of Mathematical Functions*, U.S. Department of Commerce, National Institute of Standards and Technology, Washington, DC; Cambridge University Press, Cambridge, 2010, with 1 CD-ROM (Windows, Macintosh and UNIX).
URL <http://dlmf.nist.gov/>
- [16] QI, F.: *Integral representations and properties of Stirling numbers of the first kind*, *J. Number Theory*, **133** (2013), No. 7, 2307–2319.
URL <https://doi.org/10.1016/j.jnt.2012.12.015>
- [17] QI, F.: *Diagonal recurrence relations, inequalities, and monotonicity related to the Stirling numbers of the second kind*, *Math. Inequal. Appl.*, **19** (2016), No. 1, 313–323.
URL <https://doi.org/10.7153/mia-19-23>
- [18] QI, F. AND CHAPMAN, R. J.: *Two closed forms for the Bernoulli polynomials*, *J. Number Theory*, **159** (2016), 89–100.
URL <https://doi.org/10.1016/j.jnt.2015.07.021>
- [19] QI, F. AND CHEN, C.-P.: *A complete monotonicity property of the gamma function*, *J. Math. Anal. Appl.*, **296** (2004), No. 2, 603–607.
URL <https://doi.org/10.1016/j.jmaa.2004.04.026>
- [20] QI, F. AND GUO, B.-N.: *Explicit formulas for special values of the Bell polynomials of the second kind and for the Euler numbers and polynomials*, *Mediterr. J. Math.*, **14** (2017), No. 3, Paper No. 140, 14.
URL <https://doi.org/10.1007/s00009-017-0939-1>
- [21] QI, F. AND LI, W.-H.: *A logarithmically completely monotonic function involving the ratio of gamma functions*, *J. Appl. Anal. Comput.*, **5** (2015), No. 4, 626–634.
URL <https://doi.org/10.11948/2015049>
- [22] QI, F., LIM, D., AND GUO, B.-N.: *Some identities related to Eulerian polynomials and involving the Stirling numbers*, *Appl. Anal. Discrete Math.*, **12** (2018), No. 2, 467–480.
URL <https://doi.org/10.2298/AADM171008014Q>
- [23] QI, F., NIU, D.-W., LIM, D., AND GUO, B.-N.: *Closed formulas and identities for the Bell polynomials and falling factorials*, *Contrib. Discrete Math.*, **15** (2020), No. 1, 163–174.
URL <https://doi.org/10.11575/cdm.v15i1.68111>
- [24] QI, F., NIU, D.-W., LIM, D., AND GUO, B.-N.: *Some properties and an application of multivariate exponential polynomials*, *Math. Methods Appl. Sci.*, **43** (2020), No. 6, 2967–2983.
URL <https://doi.org/10.1002/mma.6095>

- [25] QI, F., NIU, D.-W., LIM, D., AND YAO, Y.-H.: *Special values of the Bell polynomials of the second kind for some sequences and functions*, J. Math. Anal. Appl., **491** (2020), No. 2, 124382, 31.
URL <https://doi.org/10.1016/j.jmaa.2020.124382>
- [26] QI, F., SHI, X.-T., AND GUO, B.-N.: *Some properties of the Schröder numbers*, Indian J. Pure Appl. Math., **47** (2016), No. 4, 717–732.
URL <https://doi.org/10.1007/s13226-016-0211-6>
- [27] QI, F., ZOU, Q., AND GUO, B.-N.: *The inverse of a triangular matrix and several identities of the Catalan numbers*, Appl. Anal. Discrete Math., **13** (2019), No. 2, 518–541.
URL <https://doi.org/10.2298/AADM190118018Q>
- [28] QUAINANCE, J. AND GOULD, H. W.: *Combinatorial Identities for Stirling Numbers*, World Scientific Publishing Co. Pte. Ltd., Singapore, 2016, the unpublished notes of H. W. Gould, With a foreword by George E. Andrews.
- [29] RUTISHAUSER, H.: *Eine formel von Wronski und ihre bedeutung für den quotienten-differenzen-algorithmus*, Z. Angew. Math. Phys., **7** (1956), 164–169.
URL <https://doi.org/10.1007/BF01600787>
- [30] SCHILLING, R. L., SONG, R., AND VONDRAČEK, Z.: *Bernstein Functions*, vol. 37 of *De Gruyter Studies in Mathematics*, Walter de Gruyter & Co., Berlin, 2012, 2nd edn., theory and applications.
URL <https://doi.org/10.1515/9783110269338>
- [31] SPRUGNOLI, R.: *Riordan Array Proofs of Identities in Gould’s Book*, Dipartimento di Sistemi e Informatica, University of Florence, Viale Morgagni, 65 - Firenze, Italy, February 7, 2006.
- [32] TEMME, N. M.: *Special Functions*, A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York, 1996, an introduction to the classical functions of mathematical physics.
URL <https://doi.org/10.1002/9781118032572>
- [33] WARD, M. D.: *Asymptotic rational approximation to Pi: solution of an “unsolved problem” posed by Herbert Wilf*, in: *21st International Meeting on Probabilistic, Combinatorial, and Asymptotic Methods in the Analysis of Algorithms (AofA’10)*, Discrete Math. Theor. Comput. Sci. Proc., AM, Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2010 pp. 591–601.
URL <https://hal.inria.fr/hal-01185575>
- [34] WIDDER, D. V.: *The Laplace Transform*, Princeton Mathematical Series, vol. 6, Princeton University Press, Princeton, N. J., 1941.

INSTITUTE OF MATHEMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO 454010, HENAN, CHINA
 SCHOOL OF MATHEMATICAL SCIENCES, TIANGONG UNIVERSITY, TIANJIN 300387, CHINA
 Email address: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com
 URL: <https://qifeng618.wordpress.com>, <https://orcid.org/0000-0001-6239-2968>

THE DATA MINE, PURDUE UNIVERSITY, WEST LAFAYETTE, INDIANA, USA
 DEPARTMENT OF STATISTICS, PURDUE UNIVERSITY, WEST LAFAYETTE, INDIANA, USA
 Email address: mdw@purdue.edu
 URL: <https://orcid.org/0000-0001-8795-746X>