

SUMS OF RANDOM POLYNOMIALS WITH DIFFERING DEGREES

ISABELLE KRAUS AND SEAN O'ROURKE

ABSTRACT. Let μ and ν be probability measures in the complex plane, and let p and q be independent random polynomials of degree n , whose roots are chosen independently from μ and ν , respectively. As $n \rightarrow \infty$, the limiting distribution for the zeros of the sum $p + q$ was computed by Reddy and the second author [J. Math. Anal. Appl. 495 (2021) 124719]. In this paper, we generalize this result to the case where p and q have different degrees. In this case, the logarithmic potential of the limiting distribution is given by the pointwise maximum of the logarithmic potentials of μ and ν , scaled by the limiting ratio of the degrees of p and q . Our methods also allow us to consider the sum of m independent random polynomials with differing degrees when m is fixed and the degree of the sum tends to infinity.

1. INTRODUCTION

Motivated by the results of Reddy and the second author [10], this paper focuses on a model of random polynomials with independent and identically distributed (iid) roots. Namely, we consider monic polynomials (in a single complex variable) of the form

$$p_n(z) := \prod_{i=1}^n (z - X_i),$$

where the roots X_1, \dots, X_n are iid random variables in the complex plane. Various properties of p_n have been studied by a number of authors, see [2, 3, 4, 6, 7, 11, 12, 13, 16, 20] and references therein.

The results in [10] do not focus on a single random polynomial, but instead describe zeros of sums of random polynomials. Specifically, let

$$p_n(z) := \prod_{i=1}^n (z - X_i), \quad q_n(z) := \prod_{j=1}^n (z - Y_j)$$

be two independent random polynomials of degree n , whose roots X_1, \dots, X_n and Y_1, \dots, Y_n are chosen independently from probability measures μ and ν in the complex plane, respectively. The main results of [10] describe the limiting distribution for the zeros of the sum $p + q$ as $n \rightarrow \infty$ in terms of the logarithmic potentials of μ and ν . More generally, the results in [10] apply to sums of m independent random polynomials when m is fixed and n tends to infinity. In order to state these results, we must first introduce some definitions and notation.

Let $\mathcal{P}(\mathbb{C})$ be the set of probability measures on \mathbb{C} with compact support. For a measure $\mu \in \mathcal{P}(\mathbb{C})$, we let $\text{supp}(\mu) \subset \mathbb{C}$ denote the support of μ .

S. O'Rourke has been supported in part by NSF grant DMS-1810500.

Definition 1.1. The *logarithmic potential* U_μ of $\mu \in \mathcal{P}(\mathbb{C})$ is the function $U_\mu : \mathbb{C} \rightarrow [-\infty, +\infty)$ defined for all $z \in \mathbb{C}$ by

$$U_\mu(z) := \int_{\mathbb{C}} \log |z - w| d\mu(w).$$

Definition 1.2 (Supported on a circle, [10]). Let $K \subset \mathbb{C}$ be nonempty. We say that $\mu \in \mathcal{P}(\mathbb{C})$ is *supported on a circle centered in K* if there exist $r \geq 0$ and $z \in K$ so that

$$\text{supp } \mu \subset \{w \in \mathbb{C} : |w - z| = r\}.$$

If not, we say μ is *not supported on a circle centered in K* . If μ is supported on a circle centered in \mathbb{C} , we simply say μ is *supported on a circle*, and when μ is not supported on a circle centered in \mathbb{C} , we say μ is *not supported on a circle*.

Let λ denote the Lebesgue measure on \mathbb{C} , and let $C_c^\infty(\mathbb{C})$ denote the set of all smooth functions $\varphi : \mathbb{C} \rightarrow \mathbb{C}$ with compact support.

Theorem 1.3 (Theorem 1.10 in [10]). *Let $m \geq 2$ be a fixed integer, and assume $\mu_1, \dots, \mu_m \in \mathcal{P}(\mathbb{C})$. Assume for each $1 \leq k \leq m - 1$, the measure μ_k is not supported on a circle. Let $\{X_{i,k} : 1 \leq k \leq m, i \geq 1\}$ be a collection of independent random variables so that $X_{i,k}$ has distribution μ_k for each $i \geq 1$. For each $n \geq 1$, define the degree n polynomials*

$$p_{n,k}(z) := \prod_{i=1}^n (z - X_{i,k}), \quad 1 \leq k \leq m.$$

Then there exists a (deterministic) probability measure ρ on \mathbb{C} so that, for any $\varphi \in C_c^\infty(\mathbb{C})$,

$$\frac{1}{n} \sum_{i=1}^n \varphi(z_i^{(n)}) \longrightarrow \int_{\mathbb{C}} \varphi d\rho$$

in probability as $n \rightarrow \infty$, where $z_1^{(n)}, \dots, z_n^{(n)}$ are the zeros of the sum $\sum_{k=1}^m p_{n,k}$. Here, ρ depends only on μ_1, \dots, μ_m and is uniquely defined by the condition that

$$\int_{\mathbb{C}} \varphi d\rho = \frac{1}{2\pi} \int_{\mathbb{C}} \Delta\varphi(z) \left(\max_{1 \leq k \leq m} U_{\mu_k}(z) \right) d\lambda(z) \quad \text{for all } \varphi \in C_c^\infty(\mathbb{C}).$$

Our main result generalizes Theorem 1.3 by allowing the polynomials $p_{n,1}, \dots, p_{n,m}$ to have different degrees. That is, we consider polynomials of the form

$$p_{n,k}(z) := \prod_{i=1}^{n_k} (z - X_{i,k}), \quad 1 \leq k \leq m,$$

where n_1, \dots, n_m are sequences of natural numbers, indexed by n (so that $n_k = n_k(n)$). Our main result focuses on the case where $n_1 := n \geq n_2 \geq \dots \geq n_m$ for all $n \in \mathbb{N}$, although our methods can handle slightly more general cases. In particular, our main result does not require that the sequences n_2, \dots, n_m tend to infinity; in fact, Example 1.5 below shows some interesting and unexpected behavior for the zeros of the sum when n_2, \dots, n_m do not tend to infinity.

Unlike the case when all the polynomials have the same degree (Theorem 1.3), the case when the polynomials have different degrees requires several new parameters.

These new parameters are $c_1, \dots, c_m \in [0, 1]$ and are defined by the limiting ratio of the degrees:

$$(1) \quad c_k := \lim_{n \rightarrow \infty} \frac{n_k}{n}, \quad 1 \leq k \leq m.$$

In particular, it is always the case that $c_1 = 1$. Our main result below shows that the limiting distribution for the zeros of the sum depends only on the measures μ_1, \dots, μ_m and the ratios c_1, \dots, c_m .

Theorem 1.4 (Main result). *Let $m \geq 2$ be a fixed integer, and assume $\mu_1, \dots, \mu_m \in \mathcal{P}(\mathbb{C})$. Assume for each $1 \leq k \leq m-1$, the measure μ_k is not supported on a circle. Let $\{X_{i,k} : 1 \leq k \leq m, i \geq 1\}$ be a collection of independent random variables so that $X_{i,k}$ has distribution μ_k for each $i \geq 1$. For each $n \geq 1$, define the polynomials*

$$(2) \quad p_{n,k}(z) := \prod_{i=1}^{n_k} (z - X_{i,k}), \quad 1 \leq k \leq m,$$

where n_1, \dots, n_m are sequences of natural numbers, indexed by n , so that $n_1 := n \geq n_2 \geq \dots \geq n_m$ for all $n \in \mathbb{N}$. In addition, assume the limits in (1) hold for some $c_1, \dots, c_m \in [0, 1]$. Then there exists a (deterministic) probability measure ρ on \mathbb{C} so that, for any $\varphi \in C_c^\infty(\mathbb{C})$,

$$\frac{1}{n} \sum_{i=1}^n \varphi(z_i^{(n)}) \longrightarrow \int_{\mathbb{C}} \varphi d\rho$$

in probability as $n \rightarrow \infty$, where $z_1^{(n)}, \dots, z_n^{(n)}$ are the zeros of the sum $\sum_{k=1}^m p_{n,k}$. Here, ρ depends only on μ_1, \dots, μ_m and c_1, \dots, c_m and is uniquely defined by the condition that

$$(3) \quad \int_{\mathbb{C}} \varphi d\rho = \frac{1}{2\pi} \int_{\mathbb{C}} \Delta\varphi(z) \left(\max_{1 \leq k \leq m} c_k U_{\mu_k}(z) \right) d\lambda(z) \quad \text{for all } \varphi \in C_c^\infty(\mathbb{C}),$$

where we use the convention that $c_k U_{\mu_k}(z) = 0$ for all $z \in \mathbb{C}$ if $c_k = 0$.

We make a few remarks concerning Theorem 1.4. First, in the case where $n_1 = n_2 = \dots = n_m = n$, it follows that $c_1 = c_2 = \dots = c_m = 1$, and we recover Theorem 1.3. Second, the defining relation for ρ given in (3) implies that the function $U(z) := \max_{1 \leq k \leq m} c_k U_{\mu_k}(z)$ is the logarithmic potential of ρ . In fact, we can write (3) as

$$\rho = \frac{1}{2\pi} \Delta U,$$

where the Laplacian is interpreted in the distributional sense (see Section 3.7 in [15]). Third, as noted in [10], the assumption in Theorem 1.3 that the measures μ_k , $1 \leq k \leq m-1$ are not supported on any circle can be replaced with an alternative assumption (see Assumption 1.9 in [10]). The same is true for Theorem 1.4, although the replacement assumption is much more technical to state in this case and requires specific growth conditions on the degrees n_2, \dots, n_m . For simplicity, we have decided to omit this alternative assumption from Theorem 1.4.

Example 1.5. Consider Theorem 1.4 in case where $m = 2$, μ_1 is the uniform probability measure on the unit disk in the complex plane centered at 0, and μ_2 is the uniform probability measure on the unit disk centered at 2. Let $n_1 := n$ for each $n \in \mathbb{N}$, and let n_2 be any sequence of natural numbers indexed by n chosen




FIGURE 1. A numerical simulation of Example 1.5. The blue circles represent the roots of $p_{n,1}$, which are uniform on the unit disk centered at the origin. In this simulation, $p_{n,1}$ has degree 300. The red circles represent the roots of $p_{n,2}$, which are uniform on the unit disk centered at 2. In this simulation, $p_{n,2}$ has degree 25. The green circles represent the roots of the sum $p_{n,1} + p_{n,2}$.

so that $\lim_{n \rightarrow \infty} \frac{n_2}{n} = 0$ (in particular, this includes the case where n_2 is constant). Define the polynomials $p_{n,1}$ and $p_{n,2}$ as in (2).

Since the degree of $p_{n,2}$ is significantly smaller than the degree of $p_{n,1}$, it might be natural to conjecture that the limiting distribution of the sum $p_{n,1} + p_{n,2}$ is given by μ_1 . However, this is not the case (even when $n_2 = 1$ for all n). In this case, the limiting distribution for the zeros of $p_{n,1} + p_{n,2}$ is given by the uniform probability measure on the unit circle centered at the origin. To see this, note that for every $z \in \mathbb{C}$,

$$U_{\mu_1}(z) = \begin{cases} \log |z|, & \text{if } |z| > 1, \\ \frac{1}{2}(|z|^2 - 1), & \text{if } |z| \leq 1, \end{cases}$$

see for instance [18], and

$$c_2 U_{\mu_2}(z) = 0$$

since $c_2 = 0$. It follows that

$$U(z) := \max\{U_{\mu}(z), 0\} = \begin{cases} \log |z|, & \text{if } |z| > 1, \\ 0, & \text{if } |z| \leq 1. \end{cases}$$

Using the mean value property for harmonic functions, U can be seen to be equal almost everywhere to the logarithmic potential of the uniform probability measure on the unit circle centered at the origin. Thus, by uniqueness of the logarithmic potential (see, for instance, Lemma 4.1 from [1]), it follows that the resultant measure ρ given in Theorem 1.4 is the uniform probability measure on the unit circle centered at the origin. A numerical simulation of this example is presented in Figure 1.

1.1. Notation. Unless otherwise noted, asymptotic notation (such as O, o, \ll) is used under the assumption that $n \rightarrow \infty$. For example, $X = O(Y)$ and $X \ll Y$ denote the bound $|X| \leq CY$ for some constant $C > 0$ and all $n > C$, where C is independent of n . Subscripts such as $X = O_k(Y)$ indicate that the constant C depends on some parameter k , but C may depend on the measures μ_1, \dots, μ_m without denoting this dependence. We use $X = o(Y)$ if $|X| \leq a_n Y$ for some sequence (a_n) such that $\lim_{n \rightarrow \infty} a_n = 0$.

We define the open disk of radius $r > 0$ centered at $z \in \mathbb{C}$ to be

$$B(z, r) := \{w \in \mathbb{C} : |z - w| < r\},$$

and set $B(r) := B(0, r)$.

Let $\mathcal{P}(\mathbb{C})$ be the set of probability measures on \mathbb{C} with compact support; we let $\text{supp}(\mu)$ denote the support of μ . The Lebesgue measure on \mathbb{C} is denoted by λ . Let $C_c^\infty(\mathbb{C})$ denote the set of all smooth functions $\varphi : \mathbb{C} \rightarrow \mathbb{C}$ with compact support; $\text{supp}(\varphi)$ will denote the support of φ .

We let $\log(z)$ be the natural logarithm and $\text{Log}(z)$ the principal branch of the complex logarithm. Let the real part of $z \in \mathbb{C}$ be denoted $\text{Re}(z)$ and the imaginary part $\text{Im}(z)$.

For a finite set S , we let $|S|$ denote its cardinality.

1.2. Overview and outline. The proof of Theorem 1.4 is similar to the proofs appearing in [10, 19]. While our method follows closely the approach taken in [10], the fact that $p_{n,1}, \dots, p_{n,m}$ have different degrees introduces new technical difficulties. For example, when n_2, \dots, n_m are small compared to n , the anti-concentration bounds from [10] no longer hold with high enough probability for the methods there to apply. To get around this issue, we treat all low-degree polynomials simultaneously.

To this end, define

$$(4) \quad j_n := \begin{cases} \min\{1 < k \leq m : n_k \leq \log^6 n\}, & \text{if } n_m \leq \log^6 n \\ m, & \text{if } n_m > \log^6 n \end{cases}$$

and

$$(5) \quad \tilde{p}_{n,k}(z) := \begin{cases} p_{n,k}(z), & \text{for } 1 \leq k < j_n \\ p_{n,j_n}(z) + \dots + p_{n,m}(z) & \text{for } k = j_n. \end{cases}$$

By definition, it is the case that $1 < j_n \leq m$ and

$$(6) \quad \sum_{k=1}^{j_n} \tilde{p}_{n,k} = \sum_{k=1}^m p_{n,k}.$$

The idea behind this construction is that $\tilde{p}_{n,k} = p_{n,k}$ when $k < j_n$, while p_{n,j_n} captures any of the “low-degree” polynomials in the sum (if there are none, then we simply take $\tilde{p}_{n,j_n} = p_{n,m}$). Here, we have used $\log^6 n$ as the cutoff value for what is considered a “low-degree” polynomial, although there are many other choices for the cutoff value that would also work with our method. Since the sequences n_2, \dots, n_m do not necessarily converge to infinity, it may be the case that j_n fluctuates; in particular, the sequence (j_n) is bounded but may not necessarily converge.

The representation for the sum given on the left-hand side of (6) will allow us to handle all of the low-degree polynomials simultaneously. However, this representation introduces its own difficulties: the polynomial \tilde{p}_{n,j_n} is not necessarily

monic and its roots are not necessarily independent random variables. Because of these issues, several technical results from [10] no longer apply and a more detailed analysis is required (see Section 2 for the details).

Our main result, Theorem 1.4, is proved using Theorem 1.6 below; the proof of Theorem 1.6 is presented in Section 2. The assumptions in Theorem 1.6 differ slightly from those in Theorem 1.4: Theorem 1.6 uses condition (7) instead of requiring that the measures $\mu_k, 1 \leq k \leq m-1$ not be supported on circles. In Section 3, we show that condition (7) holds whenever the measures $\mu_k, 1 \leq k \leq m-1$ are not supported on circles (see Lemma 3.1). Appendix A contains a number of auxiliary deterministic results.

Theorem 1.6. *Let $m \geq 2$ be a fixed integer, and assume $\mu_1, \dots, \mu_m \in \mathcal{P}(\mathbb{C})$. Let $\{X_{i,k} : 1 \leq k \leq m, i \geq 1\}$ be a collection of independent random variables so that $X_{i,k}$ has distribution μ_k for each $i \geq 1$. For each $n \geq 1$, define the polynomials $p_{n,1}, \dots, p_{n,m}$ as in (2), where n_1, \dots, n_m are sequences of natural numbers, indexed by n , so that $n_1 := n \geq n_2 \geq \dots \geq n_m$ for all $n \in \mathbb{N}$. In addition, assume the limits in (1) hold for some $c_1, \dots, c_m \in [0, 1]$. Define j_n as in (4) and $\tilde{p}_{n,k}$ as in (5) for $1 \leq k \leq j_n$. Let $\varphi \in C_c^\infty(\mathbb{C})$, and assume that*

$$(7) \quad \mathbb{P} \left(\bigcup_{k \neq l} \left\{ \frac{1}{2} \leq \left| \frac{\tilde{p}_{n,k}(Z)}{\tilde{p}_{n,l}(Z)} \right| \leq 2 \right\} \right) = O_\varphi \left(\frac{1}{\log^3 n} \right),$$

where Z is a random variable, uniformly distributed on the support of φ , independent of $X_{i,k}, 1 \leq k \leq m, i \geq 1$. Then

$$\frac{1}{n} \sum_{i=1}^n \varphi(z_i^{(n)}) \longrightarrow \frac{1}{2\pi} \int_{\mathbb{C}} \Delta\varphi(z) \left(\max_{1 \leq k \leq m} c_k U_{\mu_k}(z) \right) d\lambda(z)$$

in probability as $n \rightarrow \infty$, where $z_1^{(n)}, \dots, z_n^{(n)}$ are the zeros of the sum $\sum_{k=1}^m p_{n,k}$.

The limiting probability measure in Theorem 1.4 can be identified from Theorem 1.6 by using the following result.

Lemma 1.7. *Let $m \geq 1$ be a fixed integer, and take $\mu_1, \dots, \mu_m \in \mathcal{P}(\mathbb{C})$ and $c_1, \dots, c_m \in [0, 1]$ with $c_1 = 1$. Then there exists a unique (deterministic) probability measure ρ on \mathbb{C} so that*

$$(8) \quad \int_{\mathbb{C}} \varphi d\rho = \frac{1}{2\pi} \int_{\mathbb{C}} \Delta\varphi(z) \left(\max_{1 \leq k \leq m} c_k U_{\mu_k}(z) \right) d\lambda(z) \quad \text{for all } \varphi \in C_c^\infty(\mathbb{C}),$$

where we use the convention that $c_k U_{\mu_k}(z) = 0$ for all $z \in \mathbb{C}$ if $c_k = 0$. In addition, (8) uniquely defines ρ .

Proof. The lemma follows from standard results on subharmonic functions. By Theorems 2.2.3 and 3.1.2 in [15], $\max_{1 \leq i \leq m} c_i U_{\mu_i}$ is a subharmonic function on \mathbb{C} . Thus, from the results in Section 3.7 of [15], it follows that there exists a unique Radon measure ρ on \mathbb{C} such that (8) holds. It remains to show that ρ is a probability measure. By Theorem 3.1.2 in [15], for each $1 \leq k \leq m$,

$$U_{\mu_k}(z) = \log |z| + O(|z|^{-1}) \quad \text{as } z \rightarrow \infty.$$

In other words, there exists a constant $C_k > 0$ (depending on μ_k) such that

$$|U_{\mu_k}(z) - \log |z|| \leq \frac{C_k}{|z|}$$

for all $|z| \geq C_k$. For the constant $C = \max\{C_1, \dots, C_m\} > 0$, we have that

$$\left| \max_{1 \leq k \leq m} U_{\mu_k}(z) - \log |z| \right| \leq \frac{C}{|z|}$$

and

$$|U_{\mu_1}(z) - \log |z|| \leq \frac{C}{|z|}$$

for all $|z| \geq C$. Since $c_k \in [0, 1]$ and $c_1 = 1$,

$$-\frac{C}{|z|} \leq U_{\mu_1}(z) - \log |z| \leq \max_{1 \leq k \leq m} c_k U_{\mu_k}(z) - \log |z| \leq \max_{1 \leq k \leq m} U_{\mu_k}(z) - \log |z| \leq \frac{C}{|z|}$$

for $|z| \geq C$, and therefore

$$(9) \quad \left| \max_{1 \leq k \leq m} c_k U_{\mu_k}(z) - \log |z| \right| \leq \frac{C}{|z|}$$

for all $|z| \geq C$. With (9) established, the fact that ρ is a probability measure now follows from the arguments given at the end of the proof of Lemma 1.15 in [10]. \square

2. PROOF OF THEOREM 1.6

2.1. Preliminary bounds. The following section provides several bounds under the assumptions of Theorem 1.6 that are necessary for its proof. These bounds are similar to the auxiliary results found in Section 2.1 of [10].

Lemma 2.1. *(Lemma 2.1 from [10]) Suppose $\mu \in \mathcal{P}(\mathbb{C})$. Then for each compact set $K \subset \mathbb{C}$, there exists a constant $C > 0$ (depending only on μ and K) so that*

$$\int_K U_{\mu}^2(z) d\lambda(z) \leq \int_K \int_{\mathbb{C}} \log^2 |z - w| d\mu(w) d\lambda(z) \leq C.$$

Lemma 2.2. *Let $M > 1$, and let n and n' be natural numbers with $n' \leq n$. Let q be the (deterministic) polynomial*

$$q(z) := \alpha \prod_{i=1}^{n'} (z - \xi_i),$$

where the roots ξ_i lie in $B(M)$, and $\alpha > 0$. Then, for any compact set $K \subset \mathbb{C}$, there exists a constant $C > 0$ (depending only on K and α) so that

$$(10) \quad \frac{1}{n^2} \int_K \log^2 |q(z)| d\lambda(z) \leq C \frac{n'}{n} \log^2(2M).$$

Proof. By the Cauchy–Schwarz inequality,

$$\log^2 |q(z)| = \left(\sum_{i=1}^{n'} \log |z - \xi_i| + \log \alpha \right)^2 \leq (n' + 1) \left(\sum_{i=1}^{n'} \log^2 |z - \xi_i| + \log^2 \alpha \right).$$

Since $n' \leq n$, it follows that $\frac{n'+1}{n^2} \leq \frac{2}{n}$, and hence

$$(11) \quad \begin{aligned} \frac{1}{n^2} \int_K \log^2 |q(z)| d\lambda(z) &\leq \frac{2}{n} \sum_{i=1}^{n'} \int_K \log^2 |z - \xi_i| d\lambda(z) + \frac{2 \log^2 \alpha}{n} \int_K d\lambda(z) \\ &\leq \frac{2n'}{n} \sup_{w \in B(M)} \int_K \log^2 |z - w| d\lambda(z) + \frac{2\lambda(K) \log^2 \alpha}{n}. \end{aligned}$$

The integral on the right-hand side can be bounded using Lemma A.5 to obtain

$$(12) \quad \sup_{w \in B(M)} \int_K \log^2 |z - w| d\lambda(z) \leq C \log^2(2M),$$

where $C > 0$ is a constant depending only on K . Combining (12) with (11) yields (10). \square

Lemma 2.3. *Under the assumptions of Theorem 1.6 and for any compact set $K \subset \mathbb{C}$,*

$$(13) \quad \sup_{1 \leq k \leq m} \frac{1}{n^2} \int_K \log^2 |p_{n,k}(z)| d\lambda(z) = O_K(1)$$

and

$$(14) \quad \frac{1}{n^2} \int_K \log^2 \left(\max_{1 \leq k \leq m} |p_{n,k}(z)| \right) d\lambda(z) = O_{K,m}(1)$$

with probability 1.

Proof. We begin by establishing (13). By supposition, $\cup_{k=1}^m \text{supp}(\mu_k)$ is a compact set, and there exists $M > 1$ so that $\cup_{k=1}^m \text{supp}(\mu_k) \subset B(M)$. Thus, the random variables $X_{i,k}, 1 \leq k \leq m, i \geq 1$ lie in $B(M)$ with probability 1. In other words, the roots of $p_{n,1}, \dots, p_{n,m}$ lie in $B(M)$ with probability 1. Hence, (13) follows from Lemma 2.2.

The bound in (14) follows from (13) and the fact that

$$(15) \quad \log^2 \left(\max_{1 \leq k \leq m} a_k \right) \leq \sum_{k=1}^m \log^2 a_k$$

when $a_1, \dots, a_m > 0$. \square

Lemma 2.4. *Under the assumptions of Theorem 1.6, there exists a constant $C > 0$ (depending only on μ_1, \dots, μ_m) so that, with probability 1, all the roots of $\sum_{k=1}^m p_{n,k}$ are contained in the disk $B(Cn^{m-1})$.*

Proof. The claim follows immediately from Lemma A.4 since $\cup_{k=1}^m \text{supp}(\mu_k)$ is a compact set and therefore contained in $B(M)$ for some constant $M > 0$, which implies the random variables $X_{i,k}, 1 \leq k \leq m, i \geq 1$ (and hence the roots of the monic polynomials $p_{n,1}, \dots, p_{n,m}$) lie in $B(M)$ with probability 1. \square

Lemma 2.5. *Under the assumptions of Theorem 1.6, there exists a constant $C > 0$ (depending only on μ_1, \dots, μ_m) so that, with probability 1, all the roots of \tilde{p}_{n,j_n} are contained in the disk $B(C \log^{6(m-j_n)} n)$.*

Proof. Again we have that $\cup_{k=1}^m \text{supp}(\mu_k)$ is a compact set and therefore contained in $B(M)$ for some constant $M > 0$, which implies the random variables $X_{i,k}, 1 \leq k \leq m, i \geq 1$ (and hence roots of the monic polynomials $p_{n,1}, \dots, p_{n,m}$) lie in $B(M)$ with probability 1. We consider two cases. If $n_m > \log^6 n$, then $j_n = m$ and $\tilde{p}_{n,j_n} = p_{n,m}$. Since the roots of $p_{n,m}$ are contained in $B(M)$, the lemma holds with $C = M$. If $n_m \leq \log^6 n$, then by (4), $\tilde{p}_{n,j_n} = p_{n,j_n} + \dots + p_{n,m}$. Since each $p_{n,k}$ has degree at most $\log^6 n$ for $j_n \leq k \leq m$, the conclusion of the lemma follows from Lemma A.4 by noting that we are summing $m - j_n + 1$ polynomials. \square

Lemma 2.6. *Under the assumptions of Theorem 1.6 and for any $M > 0$,*

$$\frac{1}{n^2} \int_{B(M)} \log^2 \left| \sum_{k=1}^m p_{n,k}(z) \right| d\lambda(z) = O_{M,m}(\log^2 n)$$

with probability 1.

Proof. Recall that $z_1^{(n)}, \dots, z_n^{(n)}$ are the roots of $\sum_{k=1}^m p_{n,k}(z)$. Then

$$\left| \sum_{k=1}^m p_{n,k}(z) \right| = \alpha \prod_{i=1}^n |z - z_i^{(n)}|,$$

where α is the number of polynomials with degree equal to n . By Lemma 2.4, the roots $z_1^{(n)}, \dots, z_n^{(n)}$ are contained in $B(Cn^{m-1})$, so the result follows from applying Lemma 2.2 and noting that $\alpha \leq m$. \square

Lemma 2.7. *Under the assumptions of Theorem 1.6 and for any $M > 0$,*

$$(16) \quad \frac{1}{n^2} \int_{B(M)} \log^2 \left(\max_{1 \leq k \leq j_n} |\tilde{p}_{n,k}(z)| \right) d\lambda(z) = O_{M,m}(1)$$

with probability 1.

Proof. Since $j_n \leq m$ and

$$\frac{1}{n^2} \int_{B(M)} \log^2 \left(\max_{1 \leq k \leq j_n} |\tilde{p}_{n,k}(z)| \right) d\lambda(z) \leq \frac{1}{n^2} \sum_{k=1}^{j_n} \int_{B(M)} \log^2 |\tilde{p}_{n,k}(z)| d\lambda(z),$$

it suffices to bound

$$\sup_{1 \leq k \leq j_n} \int_{B(M)} \log^2 |\tilde{p}_{n,k}(z)| d\lambda(z).$$

If $\tilde{p}_{n,k} = p_{n,k}$, then Lemma 2.3 implies that

$$\frac{1}{n^2} \int_{B(M)} \log^2 |\tilde{p}_{n,k}(z)| d\lambda(z) = \frac{1}{n^2} \int_{B(M)} \log^2 |p_{n,k}(z)| d\lambda(z) = O_M(1)$$

with probability 1. If $n_m > \log^6 n$, then $\tilde{p}_{n,k} = p_{n,k}$ for all $1 \leq k \leq m$ and the result follows.

If $n_m \leq \log^6 n$, then $\tilde{p}_{n,k} = p_{n,k}$ for $1 \leq k < j_n$, and it only remains to consider \tilde{p}_{n,j_n} . In this case, \tilde{p}_{n,j_n} has degree at most $\log^6 n$. Thus, by Lemma 2.5, the roots of $\tilde{p}_{n,j_n}(z)$ are contained in $B(C \log^{6(m-j_n)} n)$ for some constant $C > 0$ depending on μ_1, \dots, μ_m . Therefore, by Lemma 2.2, we find

$$\frac{1}{n^2} \int_{B(M)} \log^2 |\tilde{p}_{n,j_n}(z)| d\lambda(z) = O_M \left(\frac{\log^6 n}{n} \log^2 (2 \log^{6(m-j_n)} n) \right) = O_M(1)$$

with probability 1. (In fact, the first bound above implies this term actually tends to zero.) \square

2.2. A reduction. In this section, we prove Lemma 2.8 (below) and exploit the result to simplify the proof of Theorem 1.6. The proofs in this section differ substantially from the proofs given in [10].

Lemma 2.8. *Under the assumptions of Theorem 1.6, if $\varphi \in C_c^\infty(\mathbb{C})$, then*

$$\frac{1}{n} \int_{\mathbb{C}} \Delta\varphi(z) \log \left(\max_{1 \leq k \leq j_n} |\tilde{p}_{n,k}(z)| \right) d\lambda(z) \longrightarrow \int_{\mathbb{C}} \Delta\varphi(z) \left(\max_{1 \leq k \leq m} c_k U_{\mu_k}(z) \right) d\lambda(z)$$

almost surely as $n \rightarrow \infty$.

Tao and Vu [21] provide the following result which we use to prove Lemma 2.8.

Lemma 2.9 (Dominated convergence; Lemma 3.1 from [21]). *Let (X, ρ) be a finite measure space. For integers $n \geq 1$, let $f_n : X \rightarrow \mathbb{R}$ be random functions which are jointly measurable with respect to X and the underlying probability space. Assume that:*

- (1) (uniform integrability) *there exists $\delta > 0$ such that $\int_X |f_n(x)|^{1+\delta} d\rho(x)$ is bounded in probability (resp., almost surely);*
- (2) (pointwise convergence) *for ρ -almost every $x \in X$, $f_n(x)$ converges in probability (resp., almost surely) to zero.*

Then $\int_X f_n(x) d\rho(x)$ converges in probability (resp., almost surely) to zero.

Proof of Lemma 2.8. Suppose the support of φ is contained in $B(M)$ for some $M > 0$. We will prove the lemma by applying Lemma 2.9 with

$$f_n(z) := \Delta\varphi(z) \left(\frac{1}{n} \log \left(\max_{1 \leq k \leq j_n} |\tilde{p}_{n,k}(z)| \right) - \max_{1 \leq k \leq j_n} c_k U_{\mu_k}(z) \right)$$

on the finite measure space $B(M)$ with the Lebesgue measure λ .

We begin by showing that $f_n(z)$ satisfies condition (2) of Lemma 2.9. To this end, we divide the natural numbers into two subsequences. Let $\{n^+\}$ denote the subsequence of all natural numbers n where $n_m > \log^6 n$ (recall that n_m is indexed by n), and let $\{n^-\}$ denote the subsequence of all natural numbers n where $n_m \leq \log^6 n$. We will show that, for almost every $z \in B(M)$, $f_n(z)$ converges to zero almost surely along both $\{n^+\}$ and $\{n^-\}$. If $\{n^+\}$ is finite, we may ignore these finitely many terms and consider the limit along $\{n^-\}$ instead. Similarly, if $\{n^-\}$ is finite, we only need to consider the limit along $\{n^+\}$. As such, our arguments below focus only on the cases where $\{n^+\}$ or $\{n^-\}$ are infinite.

If $n \in \{n^+\}$, then $n_m > \log^6 n$ and $j_n = m$. Let $1 \leq k \leq m$ be fixed and consider the polynomial $\tilde{p}_{n,k}$. Since $n = n_1 \geq n_2 \geq \dots \geq n_m \geq \log^6 n$, it follows that $\tilde{p}_{n,k}(z) = p_{n,k}(z)$ for all $n \in \{n^+\}$, and therefore,

$$\frac{1}{n} \log |\tilde{p}_{n,k}(z)| = \frac{1}{n} \log |p_{n,k}(z)| = \frac{1}{n} \sum_{i=1}^{n_k} \log |z - X_{i,k}| = \left(\frac{n_k}{n} \right) \left(\frac{1}{n_k} \sum_{i=1}^{n_k} \log |z - X_{i,k}| \right)$$

for all $n \in \{n^+\}$. If $\{n^+\}$ is infinite, by the law of large numbers, for Lebesgue almost every $z \in \mathbb{C}$,

$$(17) \quad \frac{1}{n} \log |\tilde{p}_{n,k}(z)| = \left(\frac{n_k}{n} \right) \left(\frac{1}{n_k} \sum_{i=1}^{n_k} \log |z - X_{i,k}| \right) \longrightarrow c_k U_{\mu_k}(z)$$

almost surely as $n \rightarrow \infty$ along $\{n^+\}$. Here, we used the fact that $\log |\cdot|$ is locally integrable on \mathbb{C} , and so Fubini's theorem implies that $c_1 U_{\mu_1}, c_2 U_{\mu_2}, \dots, c_m U_{\mu_m}$ are

finite Lebesgue almost everywhere. By continuity of the pointwise maximum, it follows that, for almost every $z \in B(M)$, $f_n(z)$ converges almost surely to zero along the subsequence $\{n^+\}$.

We now consider the subsequence $\{n^-\}$. If $n \in \{n^-\}$, then $n_m \leq \log^6 n$, so by definition $j_n := \min\{1 < k \leq m : n_k \leq \log^6 n\}$. We now divide $\{n^-\}$ into several further subsequences. For $2 \leq k \leq m$, we let $\{n_{=k}^-\}$ denote the subsequence of $\{n^-\}$ for which $j_n = k$. Since $j_n \geq 2$ for all n , it follows that

$$(18) \quad \bigcup_{2 \leq k \leq m} \{n_{=k}^-\} = \{n^-\},$$

and hence it suffices to show that, for almost every $z \in B(M)$, $f_n(z)$ converges almost surely to zero along $\{n_{=k}^-\}$ for each $2 \leq k \leq m$. As discussed above, we will ignore any of these subsequences that are finite.

Let $2 \leq k \leq m$ be fixed, and assume $\{n_{=k}^-\}$ is infinite. For any $1 \leq i < k$, by factoring and applying the law of large numbers to $\frac{1}{n} \log |\tilde{p}_{n,i}(z)|$ as in (17), we find that for Lebesgue almost every $z \in B(M)$, $\frac{1}{n} \log |\tilde{p}_{n,i}(z)| \rightarrow c_i U_{\mu_i}(z)$ as $n \rightarrow \infty$ along $\{n_{=k}^-\}$. We will now consider the case when $i = k = j_n$. In this case, $\tilde{p}_{n,j_n} = \sum_{l=j_n}^m p_{n,l}$. For $n \in \{n_{=k}^-\}$, we write

$$\frac{1}{n} \log |\tilde{p}_{n,j_n}(z)| = \frac{1}{n} \log \left| \alpha \prod_{i=1}^{n_{j_n}} (z - \xi_i) \right| = \frac{\log \alpha}{n} + \frac{1}{n} \sum_{i=1}^{n_{j_n}} \log |z - \xi_i|$$

where α is the number of polynomials of degree equal to n_{j_n} and $\xi_1, \dots, \xi_{n_{j_n}}$ denote the roots of \tilde{p}_{n,j_n} . Note that α depends on n , but we always have the bound $\alpha \leq m$.

Consider

$$\begin{aligned} & \sum_{n \in \{n_{=k}^-\}} \int_{B(M)} \left(\frac{1}{n} \log |\tilde{p}_{n,j_n}(z)| \right)^2 d\lambda(z) \\ &= \sum_{n \in \{n_{=k}^-\}} \int_{B(M)} \left(\frac{\log \alpha}{n} + \frac{1}{n} \sum_{i=1}^{n_{j_n}} \log |z - \xi_i| \right)^2 d\lambda(z) \\ &\ll \sum_{n \in \{n_{=k}^-\}} \int_{B(M)} \left(\frac{\log^2 \alpha}{n^2} + \frac{1}{n^2} \left(\sum_{i=1}^{n_{j_n}} \log |z - \xi_i| \right)^2 \right) d\lambda(z). \end{aligned}$$

For the first term on the right-hand side, clearly

$$\sum_{n \in \{n_{=k}^-\}} \int_{B(M)} \frac{\log^2 \alpha}{n^2} d\lambda(z) \ll_{M,m} \sum_{n=1}^{\infty} \frac{1}{n^2} < \infty.$$

We now bound the second term. In view of Lemma 2.5, the roots of $\tilde{p}_{n,j_n}(z)$ are contained in the disk $B(C \log^{6(m-j_n)} n)$ with probability 1 for some constant $C > 0$ (depending on μ_1, \dots, μ_m). Thus, by the Cauchy–Schwarz inequality, Lemma A.5,

and since $n_{j_n} \leq \log^6 n$, we have

$$\begin{aligned}
& \sum_{n \in \{n_{=k}^-\}} \int_{B(M)} \frac{1}{n^2} \left(\sum_{i=1}^{n_{j_n}} \log |z - \xi_i| \right)^2 d\lambda(z) \\
& \leq \sum_{n \in \{n_{=k}^-\}} \int_{B(M)} \frac{n_{j_n}}{n^2} \sum_{i=1}^{n_{j_n}} \log^2 |z - \xi_i| d\lambda(z) \\
& \leq \sum_{n \in \{n_{=k}^-\}} \frac{\log^{12}(n)}{n^2} \sup_{w \in B(C \log^{6(m-j_n)}(n))} \int_{B(M)} \log^2 |z - w| d\lambda(z) \\
& \ll_M \sum_{n \in \{n_{=k}^-\}} \left(\frac{\log^{12}(n)}{n^2} \right) \left(\log^2(2C \log^{6(m-j_n)}(n)) \right) \\
& \leq \sum_{n=1}^{\infty} \left(\frac{\log^{12}(n)}{n^2} \right) \left(\log^2(2C \log^{6(m-2)}(n)) \right) \\
& < \infty.
\end{aligned}$$

Putting together the bounds above, we conclude that

$$(19) \quad \sum_{n \in \{n_{=k}^-\}} \int_{B(M)} \left(\frac{1}{n} \log |\tilde{p}_{n,j_n}(z)| \right)^2 d\lambda(z) < \infty.$$

Let $\epsilon > 0$. By Markov's inequality,

$$\lambda \left\{ z \in B(M) : \frac{1}{n} |\log |\tilde{p}_{n,j_n}(z)|| > \epsilon \right\} \leq \frac{1}{\epsilon^2} \int_{B(M)} \left(\frac{1}{n} |\log |\tilde{p}_{n,j_n}(z)|| \right)^2 d\lambda(z).$$

Summing over all $n \in \{n_{=k}^-\}$, we see by (19) that

$$\sum_{n \in \{n_{=k}^-\}} \lambda \left\{ z \in B(M) : \frac{1}{n} |\log |\tilde{p}_{n,j_n}(z)|| > \epsilon \right\} < \infty$$

with probability 1. By the Borel–Cantelli lemma,

$$\lambda \left\{ z \in B(M) : \limsup_{n \rightarrow \infty} \frac{1}{n} |\log |\tilde{p}_{n,j_n}(z)|| > \epsilon \right\} = 0$$

with probability 1, where the limit superior is taken along the subsequence $\{n_{=k}^-\}$. Therefore, we conclude that for Lebesgue almost every $z \in B(M)$,

$$\frac{1}{n} \log |\tilde{p}_{n,j_n}(z)| \longrightarrow 0$$

almost surely along $\{n_{=k}^-\}$. Since $c_k U_{\mu_k}(z) = 0$ for all $z \in \mathbb{C}$ in this case, we have that, for almost every $z \in B(M)$,

$$\frac{1}{n} \log |\tilde{p}_{n,j_n}(z)| \longrightarrow c_k U_{\mu_k}(z)$$

almost surely along $\{n_{=k}^-\}$. By (18) and continuity of the pointwise maximum, it follows that, for almost every $z \in B(M)$, $f_n(z)$ converges almost surely to zero along the subsequence $\{n^-\}$. Combined with the same result for $\{n^+\}$ given above, we conclude that, condition (2) of Lemma 2.9 is satisfied.

In order to apply Lemma 2.9, we need to verify that

$$\frac{1}{n^2} \int_{B(M)} |\Delta\varphi(z)|^2 \log^2 \left(\max_{1 \leq k \leq j_n} |\tilde{p}_{n,k}(z)| \right) d\lambda(z)$$

and

$$\int_{B(M)} |\Delta\varphi(z)|^2 \left(\max_{1 \leq k \leq j_n} c_k^2 U_{\mu_k}^2(z) \right) d\lambda(z)$$

are bounded almost surely. By bounding $\Delta\varphi$ in L^∞ -norm, it suffices to show that

$$(20) \quad \frac{1}{n^2} \int_{B(M)} \log^2 \left(\max_{1 \leq k \leq j_n} |\tilde{p}_{n,k}(z)| \right) d\lambda(z)$$

and

$$(21) \quad \int_{B(M)} \max_{1 \leq k \leq j_n} c_k^2 U_{\mu_k}^2(z) d\lambda(z)$$

are bounded almost surely. A bound for (20) follows from Lemma 2.7. The bound for (21) follows from Lemma 2.1.

Applying Lemma 2.9, we find

$$(22) \quad \int_{\mathbb{C}} \Delta\varphi(z) \left(\frac{1}{n} \log \left(\max_{1 \leq k \leq j_n} |\tilde{p}_{n,k}(z)| \right) - \max_{1 \leq k \leq j_n} c_k U_{\mu_k}(z) \right) \rightarrow 0$$

almost surely.

It remains to show that

$$(23) \quad \max_{1 \leq k \leq j_n} c_k U_{\mu_k}(z) = \max_{1 \leq k \leq m} c_k U_{\mu_k}(z), \quad z \in B(M)$$

for all sufficiently large n . If $n \in \{n^+\}$, then $j_n = m$ and (23) follows. If $n \in \{n^-\}$ is sufficiently large, then $c_k U_{\mu_k}(z) = 0$ for all $k \geq j_n$ and all $z \in \mathbb{C}$. Thus, (23) also holds for all $n \in \{n^-\}$ sufficiently large. \square

With Lemma 2.8 in hand, the proof of Theorem 1.6 reduces to showing that

$$(24) \quad \frac{1}{n} \sum_{i=1}^n \varphi(z_i^{(n)}) - \frac{1}{2\pi n} \int_{\mathbb{C}} \Delta\varphi(z) \log \left(\max_{1 \leq k \leq j_n} |\tilde{p}_{n,k}(z)| \right) d\lambda(z) \rightarrow 0$$

in probability as $n \rightarrow \infty$, where $z_1^{(n)}, \dots, z_n^{(n)}$ are the zeros of the sum $\sum_{k=1}^m p_{n,k}$. For any smooth and compactly supported function $\varphi : \mathbb{C} \rightarrow \mathbb{C}$,

$$\begin{aligned} \sum_{i=1}^n \varphi(z_i^{(n)}) &= \frac{1}{2\pi} \int_{\mathbb{C}} \Delta\varphi(z) \log \left| \sum_{k=1}^m p_{n,k}(z) \right| d\lambda(z) \\ &= \frac{1}{2\pi} \int_{\mathbb{C}} \Delta\varphi(z) \log \left| \sum_{k=1}^{j_n} \tilde{p}_{n,k}(z) \right| d\lambda(z), \end{aligned}$$

where the formula in the first line is established in Section 2.4.1 of [5] and the second line follows from (6). Hence, the convergence in (24) is a consequence of the following lemma.

Lemma 2.10. *Under the assumptions of Theorem 1.6,*

$$\frac{1}{n} \int_{\mathbb{C}} \Delta\varphi(z) \log \left| \sum_{k=1}^{j_n} \tilde{p}_{n,k}(z) \right| d\lambda(z) - \frac{1}{n} \int_{\mathbb{C}} \Delta\varphi(z) \log \left(\max_{1 \leq k \leq j_n} |\tilde{p}_{n,k}(z)| \right) d\lambda(z) \rightarrow 0$$

in probability as $n \rightarrow \infty$.

2.3. Proof of Lemma 2.10. The proof of Lemma 2.10 is nearly identical to the proof of Lemma 2.7 given in [10]. We begin with the following lemma.

Lemma 2.11. *Suppose the conditions and assumptions of Theorem 1.6 hold. Let Z be a random variable, uniformly distributed on the support of φ , independent of $X_{i,k}$, $1 \leq k \leq m$, $i \geq 1$. Then*

$$\frac{1}{n} \log \left| \sum_{k=1}^{j_n} \tilde{p}_{n,k}(Z) \right| = \frac{1}{n} \log \left(\max_{1 \leq k \leq j_n} |\tilde{p}_{n,k}(Z)| \right) + O_m \left(\frac{1}{n} \right)$$

with probability $1 - O_\varphi((\log n)^{-3})$.

Proof. This proof of Lemma 2.11 is essentially the same as the proof of Lemma 2.8 in [10]. Only the following changes must be made:

- m needs to be replaced by j_n and $p_{n,1}, \dots, p_{n,m}$ need to be replaced with $\tilde{p}_{n,1}, \dots, \tilde{p}_{n,j_n}$, respectively.
- References to [10, Equation (4)] need to be replaced with (7).

□

Proof of Lemma 2.10. The proof of Lemma 2.10 repeats the proof of Lemma 2.7 given in [10] almost exactly. Only the following changes need to be made:

- $\sum_{k=1}^m p_{n,k}$ needs to be replaced by $\sum_{k=1}^{j_n} \tilde{p}_{n,k}$ and $\max_{1 \leq k \leq m} |p_{n,k}(z)|$ needs to be replaced by $\max_{1 \leq k \leq j_n} |\tilde{p}_{n,k}(z)|$.
- References to [10, Lemmas 2.2 and 2.4] need to be replaced by Lemmas 2.6 and 2.7.
- The use of [10, Lemma 2.8] needs to be replaced by the use of Lemma 2.11.

□

3. PROOF OF THEOREM 1.4

In this section, we prove Theorem 1.4 using Theorem 1.6. The following lemma justifies assumption (7) in Theorem 1.6. This is the only place in the proof where the assumption that the measures μ_k , $1 \leq k \leq m-1$ are not supported on circles is needed.

Lemma 3.1. *Let $K \subset \mathbb{C}$ be a compact set with positive Lebesgue measure, and let $m \geq 2$ be a fixed integer. Let $\mu_1, \dots, \mu_m \in \mathcal{P}(\mathbb{C})$ be such that, for each $1 \leq k \leq m-1$, the measure μ_k is not supported on a circle. Let $\{X_{i,k} : 1 \leq k \leq m, i \geq 1\}$ be a collection of independent random variables so that $X_{i,k}$ has distribution μ_k for each $i \geq 1$. Let $n_1 := n \geq n_2 \geq \dots \geq n_m$, and define the polynomials*

$$p_{n,k}(z) := \prod_{i=1}^{n_k} (z - X_{i,k}), \quad 1 \leq k \leq m.$$

Define j_n as in (4) and $\tilde{p}_{n,k}$ as in (5) for $1 \leq k \leq j_n$. Then

$$\max_{k \neq l} \mathbb{P} \left(\frac{1}{2} \leq \left| \frac{\tilde{p}_{n,k}(Z)}{\tilde{p}_{n,l}(Z)} \right| \leq 2 \right) = O_K \left(\frac{1}{\log^3 n} \right),$$

where Z is a random variable, uniformly distributed on K , independent of $X_{i,k}$, $1 \leq k \leq m$, $i \geq 1$.

Applying Theorem 1.6 and Lemma 1.7 together with Lemma 3.1 completes the proof of Theorem 1.4. The rest of this section is devoted to the proof of Lemma 3.1. Following [10], we will use an inequality due to Rogozin [17] (see also Theorem 2.15 in [14]) to bound the Lévy concentration function.

Definition 3.2 (Lévy concentration function). Let ξ be a real-valued random variable. The *Lévy concentration function* of ξ is defined as

$$\mathcal{L}(\xi, t) := \sup_{u \in \mathbb{R}} \mathbb{P}(|\xi - u| \leq t)$$

for all $t \geq 0$.

Theorem 3.3 (Kolmogorov–Rogozin inequality, [17]). *Let ξ_1, \dots, ξ_n be independent real-valued random variables, and let t_1, \dots, t_n be some positive real numbers. Then for any $t \geq \max_j t_j$ we have*

$$\mathcal{L}\left(\sum_{j=1}^n \xi_j, t\right) \leq Ct \left(\sum_{j=1}^n (1 - \mathcal{L}(\xi_j, t_j)) t_j^2\right)^{-1/2},$$

where $C > 0$ is an absolute constant.

The Kolmogorov–Rogozin inequality has previously been used by Kabluchko and Zaporozhets [8] to study the zeros of random polynomials. The last ingredient in the proof of Lemma 3.1 is the following.

Proposition 3.4. *Under the assumptions of Lemma 3.1,*

$$\max_{1 \leq k < l \leq j_n} \mathbb{P}(|\log |\tilde{p}_{n,k}(Z)| - \log |\tilde{p}_{n,l}(Z)|| \leq \log 2) = O_K\left(\frac{1}{\log^3 n}\right).$$

Remark 3.5. The choice of $\log 2$ in the bound above is simply for convenience. $\log 2$ can be replaced with any fixed constant and the conclusion would still hold.

Since

$$\mathbb{P}\left(\frac{1}{2} \leq \left|\frac{\tilde{p}_{n,k}(Z)}{\tilde{p}_{n,l}(Z)}\right| \leq 2\right) = \mathbb{P}(|\log |\tilde{p}_{n,k}(Z)| - \log |\tilde{p}_{n,l}(Z)|| \leq \log 2),$$

Lemma 3.1 follows immediately from Proposition 3.4. It remains to prove Proposition 3.4.

Proof of Proposition 3.4. Fix $1 \leq k < l \leq j_n$ and consider the event

$$\{|\log |\tilde{p}_{n,k}(Z)| - \log |\tilde{p}_{n,l}(Z)|| \leq \log 2\}.$$

Since Z is continuously distributed, with probability 1, $\tilde{p}_{n,k}(Z) \neq 0$ and $\tilde{p}_{n,l}(Z) \neq 0$, and so the event above is well-defined, except on events which hold with probability zero. For the remainder of the proof, we condition on $\tilde{p}_{n,l}$ and consider it a fixed polynomial. In addition, since $k < j_n$, it follows that $\tilde{p}_{n,k} = p_{n,k}$. In other words, for the remainder of the proof, we will work with the event

$$\Omega_k := \{|\log |p_{n,k}(Z)| - \log |q(Z)|| \leq \log 2\},$$

where q is some fixed realization of $\tilde{p}_{n,l}$; our bounds will be independent of the choice of q .

By conditioning on Z to avoid the atoms of μ_k (which are at most countable) and the roots of q (for which there are at most n), we find that

$$\mathbb{P}(\Omega_k) \leq \sup_{z \in K''} \mathbb{P}(|\log |p_{n,k}(z)| - \log |q(z)|| \leq \log 2) \leq \sup_{z \in K'} \mathcal{L}(\log |p_{n,k}(z)|, \log 2),$$

where $K' \subset \mathbb{C}$ is the set K with the atoms of μ_k removed and $K'' \subset \mathbb{C}$ is the set K' with the roots of q removed.

To bound $\sup_{z \in K'} \mathcal{L}(\log |p_{n,k}(z)|, \log 2)$, we will apply the Kolmogorov–Rogozin inequality since $\log |p_{n,k}(z)| = \sum_{i=1}^{n_k} \log |X_{i,k} - z|$ is the sum of iid random variables. To this end, we will show the existence of $\varepsilon, p \in (0, 1/4)$ so that

$$\mathcal{L}(\log |X_{1,k} - z|, \varepsilon) \leq 1 - p$$

for all $z \in K'$. Since for any $u \in \mathbb{R}$ and $z \in K'$,

$$\mathbb{P}(|\log |X_{1,k} - z| - u| \leq \varepsilon) = \mathbb{P}(e^{u-\varepsilon} \leq |X_{1,k} - z| \leq e^{u+\varepsilon}),$$

it suffices to show that there exist $\varepsilon, p \in (0, 1/4)$ so that

$$\sup_{r \geq 0} \mathbb{P}(re^{-\varepsilon} \leq |X_{1,k} - z| \leq re^{\varepsilon}) \leq 1 - p$$

for all $z \in K$. This bound follows Proposition 3.5 in [10], which is reproduced below as Proposition 3.6. Applying the Kolmogorov–Rogozin inequality, we conclude that

$$\sup_{z \in K'} \mathcal{L}(\log |p_{n,k}(z)|, \log 2) \leq \frac{C}{\varepsilon \sqrt{n_k p}} \leq \frac{C}{\varepsilon \sqrt{p} \log^3 n}$$

where $C > 0$ is an absolute constant and we used the fact that $n_k \geq \log^6 n$ since $k < j_n$. \square

Proposition 3.6. *(Proposition 3.5 from [10]) Let $K \subset \mathbb{C}$ be compact. Assume $\mu \in \mathcal{P}(\mathbb{C})$, and let X be a random variable with distribution μ . If μ is not supported on a circle centered in K , then there exist $p_0, \varepsilon_0 \in (0, 1/4)$ such that*

$$(25) \quad \sup_{r \geq 0} \mathbb{P}(re^{-\varepsilon_0} \leq |X - z| \leq re^{\varepsilon_0}) \leq 1 - p_0$$

for all $z \in K$.

APPENDIX A. DETERMINISTIC RESULTS

This appendix contains a few deterministic tools used in the course of the proofs. The first few results control the locations of the zeros for the sum of polynomials; they are based on a bound due to Walsh [22]. Recall that $B(r)$ denotes the disk of radius $r > 0$ centered at the origin in the complex plane.

Lemma A.1. *Let p and q be polynomials of degrees n and m respectively with $n > m$, and assume that p has leading coefficient $\lambda_1 \geq 1$ and q has leading coefficient $\lambda_2 \geq 1$. If all the zeros of p and q are contained in $B(C_1 n^i)$ and $B(C_2 m^j)$ respectively for some $C_1, C_2 > 0, i, j \geq 0$, then all the zeros of $p + q$ are contained in the disk centered at the origin of radius Cn^{i+j+1} , for the constant $C = 2 \max\{C_1, C_2, e^{\max\{\log(\lambda_1), \log(\lambda_2)\}+2}\}$ whenever $n \geq 1$.*

Proof. We write $p + q = (\lambda_1) \frac{1}{\lambda_1} p + (\lambda_2) \frac{1}{\lambda_2} q$, and note that $\frac{1}{\lambda_1} p$ and $\frac{1}{\lambda_2} q$ are monic polynomials, whose roots are contained in $B(C_1 n^i)$ and $B(C_2 m^j)$ by supposition.

By Theorem (17,1) from [9], the zeros of the linear combination $p + q$ lie in the locus of the roots of the equation

$$(26) \quad \lambda_1(z - \alpha)^n + \lambda_2(z - \beta)^m = 0$$

where α and β vary independently over the disks $B(C_1 n^i)$ and $B(C_2 m^j)$.

Suppose to the contrary that z is a root of the above equation and that $|z| > C n^{i+j+1}$, where $C = 2 \max\{C_1, C_2, e^{\max\{\log(\lambda_1), \log(\lambda_2)\}+2}\}$. Rearranging and then taking the absolute value of (26) gives

$$|z - \alpha|^n = \frac{\lambda_2}{\lambda_1} |z - \beta|^m,$$

which can be factored into

$$|z|^{n-m} \left|1 - \frac{\alpha}{z}\right|^n = \frac{\lambda_2}{\lambda_1} \left|1 - \frac{\beta}{z}\right|^m.$$

By assumption, $|\frac{\alpha}{z}| < \frac{C_1 n^i}{C n^{i+j+1}} = \frac{C_1}{C n^{j+1}} < \frac{1}{2}$ and $|\frac{\beta}{z}| < \frac{C_2 m^j}{C n^{i+j+1}} \leq \frac{C_2}{C n^{i+1}} < \frac{1}{2}$, so taking logarithms gives

$$(27) \quad (n - m) \log |z| + n \log \left|1 - \frac{\alpha}{z}\right| = \log \left(\frac{\lambda_2}{\lambda_1}\right) + m \log \left|1 - \frac{\beta}{z}\right|.$$

Applying the Taylor expansion for $\text{Log}(1 - z)$, where we recall that $\text{Log}(\cdot)$ is the principal branch of the complex logarithm, we have

$$n \log \left|1 - \frac{\alpha}{z}\right| = n \text{Re} \left(\text{Log} \left(1 - \frac{\alpha}{z}\right) \right) = \text{Re} \left(-n \sum_{k=1}^{\infty} \left(\frac{1}{k}\right) \left(\frac{\alpha}{z}\right)^k \right),$$

which gives the bound

$$\begin{aligned} \left| n \log \left|1 - \frac{\alpha}{z}\right| \right| &\leq \left| \text{Re} \left(-n \sum_{k=1}^{\infty} \left(\frac{1}{k}\right) \left(\frac{\alpha}{z}\right)^k \right) \right| \\ &< n \sum_{k=1}^{\infty} \left(\frac{1}{k}\right) \left(\frac{C_1}{C n^{j+1}}\right)^k \\ &\leq n \left(\frac{C_1}{C n^{j+1}}\right) \sum_{k=0}^{\infty} \left(\frac{C_1}{C n^{j+1}}\right)^k \\ &\leq \left(\frac{1}{n^j}\right) \left(\frac{1}{2}\right) \sum_{k=0}^{\infty} \left(\frac{1}{2}\right)^k \\ &= \frac{1}{n^j}. \end{aligned}$$

Similarly,

$$\left| m \log \left|1 - \frac{\beta}{z}\right| \right| < \frac{1}{n^i}.$$

Recall (27), which when rearranged gives

$$(28) \quad (n - m) \log |z| = \log \left(\frac{\lambda_2}{\lambda_1}\right) + m \log \left|1 - \frac{\beta}{z}\right| - n \log \left|1 - \frac{\alpha}{z}\right|.$$

Since $n - m \geq 1$, we have that

$$(n - m) \log |z| > \log(C n^{i+j+1}) \geq \log(C) > \max\{\log(\lambda_1), \log(\lambda_2)\} + 2.$$

In addition,

$$\begin{aligned} \left| \log\left(\frac{\lambda_2}{\lambda_1}\right) + m \log\left|1 - \frac{\beta}{z}\right| - n \log\left|1 - \frac{\alpha}{z}\right| \right| &\leq \left| \log\left(\frac{\lambda_2}{\lambda_1}\right) \right| + \left| m \log\left|1 - \frac{\beta}{z}\right| \right| \\ &\quad + \left| n \log\left|1 - \frac{\alpha}{z}\right| \right| \\ &\leq \max\{\log(\lambda_1), \log(\lambda_2)\} + \frac{1}{n^i} + \frac{1}{n^j} \\ &\leq \max\{\log(\lambda_1), \log(\lambda_2)\} + 2. \end{aligned}$$

This shows that the left-hand side of (28) is strictly larger than the right-hand side, and we reach a contradiction for any value of $n \geq 1$. This completes the proof of the lemma. \square

Lemma A.2. (Lemma A.3 from [10]) *Let p_1, \dots, p_m be degree n monic polynomials. If all the zeros of p_1, \dots, p_m are contained in $B(M)$ for some $M > 0$, then all the zeros of the sum $\sum_{k=1}^m p_k$ are contained in the disk centered at the origin of radius $\frac{2^{m-1}M}{(\sin(\pi/n))^{m-1}}$.*

Remark A.3. For $n \geq 2$, Lemma A.2 implies that there exists a constant $C > 0$ (depending only on M and m) so that all the roots of the sum $\sum_{k=1}^m p_k$ lie in the disk centered at the origin with radius Cn^{m-1} . In fact, this also holds in the case $n = 1$ (with $C = M$) since in this case, the root of the sum $\sum_{k=1}^m p_k$ is a convex combination of the roots of p_1, \dots, p_m .

Lemma A.4. *Let p_1, \dots, p_m be monic polynomials of degrees n_1, \dots, n_m respectively, where $n = n_1 \geq n_2 \geq \dots \geq n_m$. If all of the zeros of p_1, \dots, p_m are contained in $B(M)$ for some $M > 0$, then all of the zeros of the sum $\sum_{k=1}^m p_k$ are contained in the disk centered at the origin of radius Cn^{m-1} for some constant $C > 0$ (depending only on M and m).*

Proof. Let r denote the cardinality of the set $\{n_1, n_2, \dots, n_m\}$, and suppose that n_{i_1}, \dots, n_{i_r} are the distinct values so that $\{n_1, n_2, \dots, n_r\} = \{n_{i_1}, \dots, n_{i_r}\}$. For $1 \leq j \leq r$, we let l_j denote the number of polynomials that have degree n_{i_j} :

$$l_j := |\{1 \leq k \leq m : \deg(p_k) = n_{i_j}\}|,$$

where $|S|$ denotes the cardinality of the set S . In particular, it follows that $\sum_{j=1}^r l_j = m$. For $1 \leq j \leq r$, we similarly let q_j be the sum of all polynomials that have degree n_{i_j} :

$$q_j := \sum_{\substack{1 \leq k \leq m \\ \deg(p_k) = n_{i_j}}} p_k.$$

Thus, we can rewrite

$$(29) \quad \sum_{k=1}^m p_k = \sum_{j=1}^r q_j,$$

where each polynomial q_j has degree n_{i_j} .

Since $n_m \leq \dots \leq n_1 = n$, it follows from Lemma A.2 (in particular, see Remark A.3) that the roots of q_j are contained in $B(C'n^{l_j-1})$ for all $1 \leq j \leq r$, where $C' > 0$ is a constant depending only on M and m . Therefore, by inductively applying Lemma A.1 ($r-1$)-times, we conclude that the roots of $\sum_{j=1}^r q_j$ are

contained in $B(Cn^{(l_1-1)+\dots+(l_{r-1})+(r-1)}) = B(Cn^{m-1})$, where $C > 0$ is a constant depending only on M and m . In view of (29), this completes the proof. \square

Lemma A.5. *For any compact set $K \subset \mathbb{C}$, there exists a constant $C > 0$ (depending only on K) so that for any $M > 1$,*

$$\sup_{w \in B(M)} \int_K \log^2 |z - w| d\lambda(z) \leq C \log^2(2M).$$

Proof. Since K is a compact set, $K \subseteq B(k-1)$ for some $k > 2$ that depends only on K . We consider two cases.

If $|w| \leq k$, then

$$\int_K \log^2 |z - w| d\lambda(z) \leq \int_{B(2k)} \log^2 |z - w| d\lambda(z),$$

where we decompose

$$\int_{B(2k)} \log^2 |z - w| d\lambda(z) = \int_{B(w,1)} \log^2 |z - w| d\lambda(z) + \int_{B(2k) \setminus B(w,1)} \log^2 |z - w| d\lambda(z).$$

For the first term on the right-hand side, by converting to polar coordinates and integrating by parts, we have

$$\begin{aligned} \int_{B(w,1)} \log^2 |z - w| d\lambda(z) &= \int_{B(0,1)} \log^2 |z| d\lambda(z) \\ &= \int_0^{2\pi} \int_0^1 r \log^2 r dr d\theta \\ &= \pi/2. \end{aligned}$$

For the second term on the right-hand side,

$$1 \leq |z - w| \leq |z| + |w| \leq 2k + k = 3k,$$

and so

$$\int_{B(2k) \setminus B(w,1)} \log^2 |z - w| d\lambda(z) \leq 4\pi k^2 \log^2(3k).$$

Hence, we conclude that

$$(30) \quad \sup_{|w| \leq k} \int_{B(2k)} \log^2 |z - w| d\lambda(z) \leq C,$$

for a constant $C > 0$ depending only on K .

We now consider the case when $k < |w| < M$. For $z \in B(k-1)$, it follows that $1 < |z - w| < k + M < 2M$ and

$$\int_K \log^2 |z - w| d\lambda(z) \leq \lambda(K) \log^2(2M).$$

We conclude that

$$(31) \quad \sup_{k < |w| < M} \int_K \log^2 |z - w| d\lambda(z) \leq \lambda(K) \log^2(2M).$$

Combining (30) and (31) completes the proof of the lemma. \square

REFERENCES

- [1] C. Bordenave and D. Chafaï. Around the circular law. *Probability Surveys*, 9:1–89, 2012.
- [2] S.-S. Byun, J. Lee, and T. R. Reddy. Zeros of random polynomials and its higher derivatives. Available at arXiv:1801.08974, 2018.
- [3] P.-L. Cheung, T. W. Ng, J. Tsai, and S. C. P. Yam. Higher-order, polar and Sz.-Nagy's generalized derivatives of random polynomials with independent and identically distributed zeros on the unit circle. *Comput. Methods Funct. Theory*, 15(1):159–186, 2015.
- [4] B. Hanin. Pairing of zeros and critical points for random polynomials. *Ann. Inst. Henri Poincaré Probab. Stat.*, 53(3):1498–1511, 2017.
- [5] J. B. Hough, M. Krishnapur, Y. Peres, and B. Virág. *Zeros of Gaussian analytic functions and determinantal point processes*, volume 51 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2009.
- [6] Z. Kabluchko. Critical points of random polynomials with independent identically distributed roots. *Proc. Amer. Math. Soc.*, 143(2):695–702, 2015.
- [7] Z. Kabluchko and H. Seidel. Distances between zeroes and critical points for random polynomials with i.i.d. zeroes. *Electron. J. Probab.*, 24:Paper No. 34, 25, 2019.
- [8] Z. Kabluchko and D. Zaporozhets. Asymptotic distribution of complex zeros of random analytic functions. *The Annals of Probability*, 42(4):1374–1395, 2014.
- [9] M. Marden. *Geometry of polynomials*. Mathematical Surveys, No. 3. American Mathematical Society, Providence, R.I., second edition, 1966.
- [10] S. O'Rourke and T. R. Reddy. Sums of random polynomials with independent roots. *Journal of Mathematical Analysis and Applications*, 495(1):124719, 23, 2021.
- [11] S. O'Rourke and N. Williams. Pairing between zeros and critical points of random polynomials with independent roots. *Trans. Amer. Math. Soc.*, 371(4):2343–2381, 2019.
- [12] S. O'Rourke and N. Williams. On the local pairing behavior of critical points and roots of random polynomials. *Electron. J. Probab.*, 25:Paper No. 100, 68, 2020.
- [13] R. Pemantle and I. Rivin. The distribution of zeros of the derivative of a random polynomial. In *Advances in combinatorics*, pages 259–273. Springer, Heidelberg, 2013.
- [14] V. V. Petrov. *Limit theorems of probability theory*, volume 4 of *Oxford Studies in Probability*. The Clarendon Press, Oxford University Press, New York, 1995. Sequences of independent random variables, Oxford Science Publications.
- [15] T. Ransford. *Potential theory in the complex plane*, volume 28 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 1995.
- [16] T. R. Reddy. Limiting empirical distribution of zeros and critical points of random polynomials agree in general. *Electron. J. Probab.*, 22:Paper No. 74, 18, 2017.
- [17] B. A. Rogozin. On the increase of dispersion of sums of independent random variables. *Akademiya Nauk SSSR. Teoriya Veroyatnostei i ee Primeneniya*, 6:106–108, 1961.
- [18] E. B. Saff and V. Totik. *Logarithmic potentials with external fields*, volume 316 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1997. Appendix B by Thomas Bloom.
- [19] M. Sodin. Value distribution of sequences of rational functions. In *Entire and subharmonic functions*, volume 11 of *Adv. Soviet Math.*, pages 7–20. Amer. Math. Soc., Providence, RI, 1992.
- [20] S. D. Subramanian. On the distribution of critical points of a polynomial. *Electron. Commun. Probab.*, 17:no. 37, 9, 2012.
- [21] T. Tao and V. Vu. Random matrices: universality of ESDs and the circular law. *The Annals of Probability*, 38(5):2023–2065, 2010. With an appendix by Manjunath Krishnapur.
- [22] J. L. Walsh. On the location of the roots of certain types of polynomials. *Trans. Amer. Math. Soc.*, 24(3):163–180, 1922.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COLORADO, CAMPUS BOX 395, BOULDER, CO 80309-0395, USA

Email address: isabelle.kraus@colorado.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COLORADO, CAMPUS BOX 395, BOULDER, CO 80309-0395, USA

Email address: sean.d.orourke@colorado.edu