

Newton Polygons of Sums on Curves II: Variation in p -adic Families

Joe Kramer-Miller and James Upton

October 19, 2021

In this article we study the behavior of Newton polygons along \mathbb{Z}_p -towers of curves. Fix an ordinary curve X over a finite field \mathbb{F}_q of characteristic p . By a \mathbb{Z}_p -tower X_∞/X we mean a tower of covers $\cdots \rightarrow X_2 \rightarrow X_1 \rightarrow X$ with $\text{Gal}(X_n/X) \cong \mathbb{Z}/p^n\mathbb{Z}$. We show that if the ramification along the tower is sufficiently moderate, then the slopes of the Newton polygon of X_n are equidistributed in the interval $[0, 1]$ as n tends to ∞ . Under a stronger congruence assumption on the ramification invariants, we completely determine the slopes of the Newton polygon of each curve. This is the first result towards ‘regularity’ in Newton polygon behavior for \mathbb{Z}_p -towers over higher genus curves. We also obtain similar results for \mathbb{Z}_p -towers twisted by a generic tame character.

Contents

1	Introduction	2
1.1	Motivation	2
1.2	Main Results	3
1.2.1	Results on Newton Polygons	4
1.2.2	Towers Twisted by Tame Characters	5
1.2.3	Equicharacteristic Results	6
1.3	Outline of Proof	8
1.4	Future Work	9
1.5	Acknowledgments	10
2	\mathbb{Z}_p-Towers	10
2.1	Artin-Schreier-Witt Theory via Witt Vectors	10
2.2	Artin-Schreier-Witt Theory via Flat Liftings	11
2.3	Characters and Splitting Functions	13
3	\mathbb{Z}_p-Towers of Local Fields	14
3.1	Local Towers with Strictly Stable Monodromy	15

3.2	Local Newton Polygons	17
4	\mathbb{Z}_p-Towers of Curves	20
4.1	Results for Global Towers	20
4.2	Results for Twisted Towers	21

1 Introduction

1.1 Motivation

An important theme in modern number theory is that of p -variation—the idea of interpolating a family of mathematical objects or invariants by a p -adic parameter. For instance, Iwasawa constructed an “arithmetic” p -adic L -function, which interpolates the p -part of class groups of a \mathbb{Z}_p -tower over a number field. The existence (and finiteness) of this object allows one to deduce the remarkable Iwasawa class number formula (see e.g. [20]). A general principle is as follows: if M is a motive over a number field K and $\rho : G_K \rightarrow \mathbb{Z}_p$ is a additive character, we may consider a family of motives $M \otimes \rho_\chi$. Here ρ_χ denotes the composition $\chi \circ \rho$ and χ varies over the space $\text{Hom}(\mathbb{Z}_p, \mathbb{C}_p^\times)$ of continuous characters. One expects that p -adic families of invariants can be constructed that interpolate classical invariants. For example, we may consider a motive M_f corresponding to an eigenform f . The algebraic special values of $L(M_f \otimes \chi, s)^{\text{alg}}$, where χ is a Dirichlet character of p -power degree, can be parameterized by a p -adic L -function (see e.g. [16]). This is somewhat miraculous, as it tells us that $L(M_f \otimes \chi, 1)^{\text{alg}}$ is close to $L(M_f \otimes \chi', 1)^{\text{alg}}$ p -adically if χ and χ' are sufficiently close.

It is natural to ask for similar p -adic variational phenomena to occur over function fields. Let X be a smooth affine curve over a finite field \mathbb{F}_q with smooth compactification \overline{X} . Let M be a motive (we are working p -adically so it makes sense to realize M as an overconvergent F -isocrystal on X) and let $\rho : \pi_1(X) \rightarrow \mathbb{Z}_p$ be a continuous character. Unlike the number field case, it is relatively easy to construct a p -adic object that interpolates the L -functions $L(M \otimes \rho_\chi, s)$. This object will not be well behaved in general—for instance, the underlying Iwasawa module will no longer be finitely generated. However, we do expect there to be well-behaved variational properties when ρ is sufficiently “nice”. When M is trivial this expectation manifests in a program devised by Daqing Wan in [19]. Wan identifies three properties that one may expect for nice ρ :

1. (monodromy stability) We say that ρ is *monodromy stable* if there exist $a, b \in \mathbb{Q}$ such that: for $n \gg 0$ and χ of finite order p^n we have $\deg(L(\rho_\chi, s)) = a + bp^n$.
2. (slope uniformity) We say that χ is *slope uniform* if for χ of finite order p^n , the slopes of $\text{NP}_q L(\rho_\chi, s)$ are equidistributed in $[0, 1]$ as n tends to infinity.
3. (slope stability) We say that χ is *slope stable* if there exist $\alpha_1, \dots, \alpha_d \in (0, 1) \cap \mathbb{Q}$ and $k > 0$ such that for all $n > k$ and all χ of finite order p^n , the slopes of

$\mathrm{NP}_q L(\rho_\chi, s)$ are

$$K \sqcup \bigsqcup_{i=1}^d \left\{ \frac{\alpha_i}{p^{n-k}}, \frac{\alpha_i + 1}{p^{n-k}}, \dots, \frac{\alpha_i + p^{n-k} - 1}{p^{n-k}} \right\},$$

where K is a set of $2g$ elements in $[0, 1]$.

These properties first appear in work of Davis-Wan-Xiao [4], who prove that all three are satisfied when $X = \mathbb{A}_{\mathbb{F}_q}^1$ and ρ is a particularly simple representation. The representations where we expect (at least some of) these properties to hold can be broken into two classes. The first class are ρ that *come from geometry* (e.g. ρ arises from the relative p -adic étale cohomology of a fiber-wise ordinary smooth proper fibration $Y \rightarrow X$). In [19], Wan conjectures that all three properties hold for ρ coming from geometry. The only progress is due to the first author in [10], where geometric representations are proven to satisfy a slightly weaker version of monodromy stability. The second class of representations are those with *strictly stable monodromy* (see §1.2-1.4 below). By work of Kosters and Zhu (see [9]), we know that strictly stable towers over $\mathbb{A}_{\mathbb{F}_q}^1$ are slope uniform.

In this article we prove slope uniformity for strictly stable representations over any ordinary curve. We also prove slope stability for strictly stable representations that satisfy a stronger monodromy condition. To the best of our knowledge, these are the first examples of these phenomena for curves other than $\mathbb{A}_{\mathbb{F}_q}^1$ and \mathbb{G}_m . Our proof uses the local-to-global methods developed in [12], [11], and the previous article [13].

1.2 Main Results

Definition 1.1. Let F be a field. A \mathbb{Z}_p -tower of fields F_∞/F is a sequence of finite Galois extensions

$$\cdots \supseteq F_2 \supseteq F_1 \supseteq F_0 = F, \quad (1)$$

together with an identification $\mathrm{Gal}(F_n/F) \cong \mathbb{Z}/p^n\mathbb{Z}$ for all $n \geq 0$.

Suppose that F is a local field of characteristic p . Let F_∞/F be a \mathbb{Z}_p -tower of local fields. For each $n \geq 0$, let v_n denote the highest ramification break (in upper numbering) of the finite extension F_n/F . By local class field theory, we have a lower bound on the growth of this sequence:

$$v_{n+1} \geq pv_n. \quad (2)$$

Definition 1.2. We say that F_∞/F has *strictly stable monodromy* if (2) is an equality for all $n \gg 0$. Equivalently, F_∞/F has strictly stable monodromy if and only if there exists $\delta \in \mathbb{Z}[\frac{1}{p}]$ such that

$$v_n = \delta p^{n-1}$$

for all $n \gg 0$. In this case, we also say that F_∞/F has δ -stable monodromy.

Definition 1.3. Let X be a curve. A \mathbb{Z}_p -tower of curves X_∞/X is a sequence of finite étale covers

$$\cdots \rightarrow X_2 \rightarrow X_1 \rightarrow X_0 = X.$$

together with an identification $\mathrm{Gal}(X_n/X) = \mathbb{Z}/p^n\mathbb{Z}$ for all $n \geq 0$.

Let X/\mathbb{F}_q be a smooth affine curve with smooth compactification \overline{X} . Let S denote the complement of X in \overline{X} . For each $P \in S$, let F_P denote the local field of X at P . If X_∞/X is a \mathbb{Z}_p -tower of curves, then by restriction we obtain a \mathbb{Z}_p -tower of local fields $F_{P,\infty}/F_P$.

Definition 1.4. We say that X_∞/X has *strictly stable monodromy* if $F_{P,\infty}/F_P$ has strictly stable monodromy for all $P \in S$. Given a tuple $\boldsymbol{\delta} = (\delta_P)_{P \in S}$ of positive numbers in $\mathbb{Z}[\frac{1}{p}]$, we say that F_∞/F has $\boldsymbol{\delta}$ -*stable monodromy* if $F_{P,\infty}/F_P$ has δ_P -stable monodromy for each $P \in S$.

Let X_∞/X be a \mathbb{Z}_p -tower of curves, corresponding to a continuous and surjective map $\rho : \pi_1(X) \rightarrow \mathbb{Z}_p$. If $\chi \in \text{Hom}(\mathbb{Z}_p, \mathbb{C}_p^\times)$ is a continuous character, then we may form the composite

$$\rho_\chi : \pi_1(X) \rightarrow \mathbb{Z}_p \xrightarrow{\chi} \mathbb{C}_p^\times.$$

Thus we obtain a family of p -adic characters of $\pi_1(X)$ parameterized by the p -adic space $\text{Hom}(\mathbb{Z}_p, \mathbb{C}_p^\times)$. In this light, it is convenient to view X_∞/X as a family of characters of $\pi_1(X)$. We can now state our first theorem:

Theorem 1.5. *Assume that \overline{X} is ordinary. If X_∞/X has strictly stable monodromy, then X_∞/X is slope uniform.*

1.2.1 Results on Newton Polygons

We now state our results regarding the interaction of Newton and Hodge polygons of characters. Let X_∞/X be a \mathbb{Z}_p -tower of curves with $\boldsymbol{\delta}$ -stable monodromy.

Definition 1.6. The *Hodge polygon* $\text{HP}(X_\infty/X)$ is the convex polygon with slope set

$$\underbrace{\{0, \dots, 0\}}_{g-1+|S|} \sqcup \bigsqcup_{P \in S} \left\{ \frac{p-1}{\delta_P}, \frac{2(p-1)}{\delta_P}, \dots \right\}.$$

Let $\chi : \mathbb{Z}_p \rightarrow \mathbb{C}_p^\times$ be a finite non-trivial character of order p^{m_χ} . We attach to χ the parameter

$$\pi_\chi = \chi(1) - 1,$$

and regard χ as taking values in the ring $\mathbb{Z}_p[\pi_\chi]$. Thus, the Artin L -function $L(\rho_\chi, s)$ is a polynomial with coefficients in $\mathbb{Z}_q[\pi_\chi]$. In this article, it will be convenient for us to adopt the following normalization convention for Newton polygons:

Definition 1.7. Let $\pi_{q,\chi} = \pi_\chi^{v_p(q)}$. The *normalized Newton polygon* of ρ_χ is

$$\text{NP}(\rho_\chi) = \text{NP}_{\pi_{q,\chi}} L(\rho_\chi, s).$$

Let $e_\chi = v_{\pi_\chi}(p) = p^{m_\chi - 1}(p - 1)$ denote the absolute ramification index of $\mathbb{Z}_p[\pi_\chi]$. Let m_0 be the smallest non-negative integer such that for each $P \in S$ the highest ramification break (in upper numbering) of $F_{P,m_0+1}/F_P$ is $p^{m_0}\delta_P$. From (2) we see that the highest

ramification break of $F_{P,m}/F_P$ is $p^{m-1}\delta_P$ for all $m > m_0$. By work of the first author in [11], the Hodge polygon $\text{HP}(X_\infty/X)$ provides a common lower bound for $\text{NP}(\rho_\chi)$ whenever $m_\chi > m_0$:

$$\text{NP}^{<e_\chi}(\rho_\chi) \succeq \text{HP}^{<e_\chi}(X_\infty/X). \quad (3)$$

If \overline{X} is ordinary, then these polygons have the same terminal point. The next theorem greatly improves the estimate (3) when \overline{X} is ordinary. It shows that the Newton and Hodge polygons touch periodically:

Theorem 1.8. *Assume that \overline{X} is ordinary. Let $d = p^{m_0} \sum_{P \in S} \delta_P$. For all $m_\chi > m_0$ and all $1 \leq n < p^{m_\chi - m_0 - 1}$, the polygons $\text{NP}^{<e_\chi}(\rho_\chi)$ and $\text{HP}(X_\infty/X)$ agree on the interval*

$$[g - 1 + nd, g - 1 + |S| + nd]. \quad (4)$$

Theorem 1.5 is then an easy consequence of Theorem 1.8. If we impose additional congruence conditions on the δ_P , we can completely determine the Newton polygon of each finite character:

Theorem 1.9. *Let X_∞/X be a \mathbb{Z}_p -tower of curves with δ -stable monodromy. Then $\text{NP}^{<e_\chi}(\rho_\chi) = \text{HP}^{<e_\chi}(X_\infty/X)$ for every finite χ if and only if:*

1. \overline{X} is ordinary.
2. $m_0 = 0$, so that $\delta_P \in \mathbb{Z}$ for all $P \in S$.
3. $p \equiv 1 \pmod{\delta_P}$ for all $P \in S$.

As a consequence of Theorem 1.9, we deduce the full form of [13, Theorem 1.3]. This gives a necessary and sufficient condition for the Newton polygon of a single character to coincide with its Hodge bound. Theorem 1.9 also provides our first examples of slope stability over curves of higher genus:

Corollary 1.10. *Every \mathbb{Z}_p -tower of curves X_∞/X satisfying the conditions of Theorem 1.9 is slope stable.*

1.2.2 Towers Twisted by Tame Characters

Let us continue to assume that X_∞/X is a \mathbb{Z}_p -tower with δ -stable monodromy, corresponding to a continuous and surjective map $\rho : \pi_1(X) \rightarrow \mathbb{Z}_p$. Let $\psi : \pi_1(X) \rightarrow \mathbb{Z}_p^\times$ be a tame character. We will write $\psi \otimes \rho$ or $\psi \otimes X_\infty/X$ for the “family” of characters $\psi \otimes \rho_\chi$ parameterized by the space $\text{Hom}(\mathbb{Z}_p, \mathbb{C}_p^\times)$. By combining Theorem 1.8 and Theorem 1.9 with the results of [11], we are able to prove stability and uniformity results for this twisted family.

Let $f : X^{\text{tame}} \rightarrow X$ be the finite cyclic covering corresponding to $\ker(\psi)$. For each $P \in S$, let s_P denote the ramification index of f at P . If Q is a point of $\overline{X}^{\text{tame}}$ lying above P , then we define $\delta_Q^{\text{tame}} = s_P \delta_P$. Our results are as follows:

Theorem 1.11. *Assume $\overline{X}^{\text{tame}}$ is ordinary. Then $\psi \otimes X_\infty/X$ is slope uniform.*

Theorem 1.12. *Assume that $\overline{X}^{\text{tame}}$ is ordinary and that $p \equiv 1 \pmod{\delta_Q^{\text{tame}}}$ for Q lying above $P \in S$. Then $\psi \otimes X_\infty/X$ is slope stable.*

Remark 1.13. Our assumption that $\overline{X}^{\text{tame}}$ is ordinary is mild: A Theorem of Bouw [3] guarantees that a generic cover will be ordinary. This uses the fact that $\deg(f) \mid p-1$ in an essential way. If $\deg(f)$ does not divide $p-1$ the lower Hodge bound is obtained by ‘averaging’ the Hodge decomposition of each isotypical component of $H^1(\overline{X}^{\text{tame}})$ over the entire Frobenius orbit.

Remark 1.14. Our proof of Theorem 1.12 gives a much stronger result. We completely determine the slopes, which depend on certain monodromy invariants. See 4.2 for more details.

Example 1.15. Let $\overline{X} = \mathbb{P}_{\mathbb{F}_q}^1$ and let $S = P_1, \dots, P_{2n}$. Let ψ be a quadratic character ramified at each P_i . Let X_∞/X be a $(1)_{P \in S}$ -stable tower. Such a tower may be obtained using Artin-Schreier-Witt theory (§2) from a rational function with simple poles at P_1, \dots, P_{2n} . Then X^{tame} is a hyperelliptic curve of genus $n-1$, which we assume is ordinary. Without being too precise, we obtain the decomposition

$$\text{Ind}_X^{X^{\text{tame}}}(X_\infty/X) = X_\infty/X \oplus (\psi \otimes X_\infty/X).$$

Let $\chi : \mathbb{Z}_p \rightarrow \mathbb{C}_p^\times$ be a finite character and let ρ_χ^{tame} denote the pullback of ρ_χ along f . There is a factorization $L(\rho_\chi^{\text{tame}}, s) = L(\rho_\chi, s)L(\psi \otimes \rho_\chi, s)$. The slopes of the L -functions break up as follows:

$$\begin{aligned} \text{NP}(\rho_\chi^{\text{tame}}) &= \underbrace{\{0, \dots, 0\}}_{3n-2} \sqcup \underbrace{\{e_\chi, \dots, e_\chi\}}_{3n-2} \sqcup \bigsqcup_{i=0}^{2n} \left\{ \frac{p-1}{2}, 2\frac{p-1}{2}, \dots, (2p^{m_\chi-1} - 1)\frac{p-1}{2} \right\}, \\ \text{NP}(\rho_\chi) &= \underbrace{\{0, \dots, 0\}}_{2n-1} \sqcup \underbrace{\{e_\chi, \dots, e_\chi\}}_{2n-1} \sqcup \bigsqcup_{i=0}^{2n} \left\{ 2\frac{p-1}{2}, 4\frac{p-1}{2}, \dots, (2p^{m_\chi-1} - 2)\frac{p-1}{2} \right\}, \\ \text{NP}(\psi \otimes \rho_\chi) &= \underbrace{\{0, \dots, 0\}}_{n-1} \sqcup \underbrace{\{e_\chi, \dots, e_\chi\}}_{n-1} \sqcup \bigsqcup_{i=0}^{2n} \left\{ \frac{p-1}{2}, 3\frac{p-1}{2}, \dots, (2p^{m_\chi-1} - 1)\frac{p-1}{2} \right\}. \end{aligned}$$

Let us point out that this decomposition can be deduced from Theorem 1.9, without any reference to [11]. This is because Theorem 1.9 allows us to determine $\text{NP}_q(\rho_\chi^{\text{tame}})$ and $\text{NP}_q(\rho_\chi)$. The ‘leftover’ slopes must be the slopes of the quadratic twist $\text{NP}_q(\psi \otimes \rho_\chi)$.

1.2.3 Equicharacteristic Results

It is well known that the space $\text{Hom}(\mathbb{Z}_p, \mathbb{C}_p)$ may be identified with the open unit disk in \mathbb{C}_p (via the map $\chi \mapsto \pi_\chi$), or equivalently with the \mathbb{C}_p -valued points of $\text{Spec}(\mathbb{Z}_p[[T]])$. From this vantage point, it is natural to consider the ‘special’ point, which can be thought of as compatifying the open disc. The special point corresponds to an equicharacteristic character $\chi_0 : \mathbb{Z}_p \rightarrow \mathbb{F}_p[[T]]^\times$ defined by

$$\chi_0(a) = (1+T)^a \in 1 + T\mathbb{F}_p[[T]].$$

As before, we define the composite character.

$$\rho_0 : \pi_1(X) \rightarrow \mathbb{Z}_p \xrightarrow{\chi_0} \mathbb{F}_p[[T]]^\times.$$

We may think of ρ_0 as the ‘limit’ of characters tending towards the boundary of the open unit disc.

Definition 1.16. The *equicharacteristic L -function* of X_∞/X is the Artin L -function $L(\rho_0, s) \in 1 + s\mathbb{F}_q[[T]][[s]]$ associated to ρ_0 .

The equicharacteristic L -function describes the limiting behavior of the $L(\rho_\chi, s)$ for finite χ as $m_\chi \rightarrow \infty$. As an example of the well behaved nature of this L -function, we have:

Theorem 1.17. *Suppose that X_∞/X has strictly stable monodromy. Then $L(\rho_0, s)$ is a T -adic entire function on $\mathbb{F}_q((T))$.*

In light of Theorem 1.17, it is convenient for us to put the L -functions $L(\rho_0, s)$ and $L(\rho_\chi, s)$ for finite χ on equal footing. Let $T_q = T^{v_p(q)}$ and define the *normalized Newton polygon* of ρ_0 to be the infinite convex polygon

$$\text{NP}(\rho_0) = \text{NP}_{T_q} L(\rho_0, s).$$

We can now state the equicharacteristic analogues of our main theorems:

Theorem 1.18. *Assume that \overline{X} is ordinary. Let X_∞/X be a \mathbb{Z}_p -tower with δ -stable monodromy. Then:*

1. *The Newton polygon $\text{NP}(\rho_0)$ lies above $\text{HP}(X_\infty/X)$.*
2. *Let d be as in Theorem 1.8. For all $n > 0$, $\text{NP}(\rho_0)$ and $\text{HP}(X_\infty/X)$ coincide on the interval (4).*
3. *Suppose that $m_0 = 0$ and that $p \equiv 1 \pmod{\delta_P}$ for all $P \in S$. Then $\text{NP}(\rho_0) = \text{HP}(X_\infty/X)$.*

Corollary 1.19. *In the setting of Theorem 1.17, the L -function $L(\rho_0, s)$ factors over $\mathbb{F}_p((T))$ as a product of polynomials of degree $\leq d$.*

The Equicharacteristic Riemann hypothesis In light of Corollary 1.19, it seems natural to ask if there is a bound on the denominators of the Newton slopes of $L(\rho_0, s)$. This question is in the spirit of Goss’s (wide open) Riemann hypothesis for Drinfeld modules. Thus we are led to make the following definition:

Definition 1.20. We say that a \mathbb{Z}_p -tower X_∞/X satisfies the *equicharacteristic Riemann hypothesis* if $L(\rho_0, s)$ is a T -adic entire function whose zeros are all contained in a finite extension of $\mathbb{F}_p((T))$.

It is not clear to us which towers should satisfy the equicharacteristic Riemann hypothesis. A hopeful guess is that it holds for all towers with strictly stable monodromy. From Theorem 1.18 we can establish the Riemann hypothesis for certain towers.

Theorem 1.21. *Assume that \overline{X} is ordinary. Let X_∞/X be a \mathbb{Z}_p -tower with δ -stable monodromy. Assume one of the two conditions hold:*

1. $\delta_P \in \mathbb{Z}$ for all $P \in S$ and $\sum \delta_P < p$.
2. $\delta_P \in \mathbb{Z}$ for all $P \in S$ and $p \equiv 1 \pmod{\delta_P}$ for all $P \in S$.

Then the equicharacteristic Riemann hypothesis holds for X_∞/X .

To deduce the Riemann hypothesis from the first condition, use Corollary 1.19 in conjunction with the observation that there are finitely many extensions of $\mathbb{F}_q((T))$ with degree less than p . To deduce the Riemann hypothesis from the second condition, use the third part of Theorem 1.18.

1.3 Outline of Proof

Artin-Schreier-Witt Theory and Splitting Functions The purpose of §2 is to upgrade some classical constructions from the theory of exponential sums on tori to more general varieties. In §2.1-2.2 we recall the basic notions of Artin-Schreier-Witt theory, and give an alternative formulation in terms of the flat liftings used in [13]. Our reason for working with flat liftings is that they lead to a very general theory of Dwork splitting functions. Such splitting functions are an integral part of the classical theory of exponential sums (see [1] and [15]), and can be constructed quite explicitly from the Artin-Hasse exponential series (as in [6]). However, these classical splitting functions are inapplicable to the study of exponential sums on more general varieties. In §2.3 we construct splitting functions using a generalized Artin-Hasse exponential. This generalized Artin-Hasse exponential is “compatible” with a given lifting of Frobenius (as opposed to the classical Artin-Hasse exponential, which is compatible with the Frobenius $t \rightarrow t^p$). In particular, it allows us to explicitly define splitting functions over any variety. This approach to splitting functions appears to be genuinely new, and allows us to deduce, for example, that the F -crystal associated to *any* character of $\pi_1(X)$ (not necessarily finite) factoring through a \mathbb{Z}_p -tower is free. This result is essential in order to apply the main theorems of [13].

Local Towers and Global Extensions In §3.1 we study the growth conditions satisfied by splitting functions associated to a δ -stable tower F_∞/F of local fields. Every such tower extends in a natural way to a δ -stable tower $X_\infty^{\text{ext}}/\mathbb{A}_{\mathbb{F}_q}^1$ over the line, whose localization at ∞ is F_∞/F . Let $\rho^{\text{ext}} : \pi_q(\mathbb{A}_{\mathbb{F}_q}^1) \rightarrow \mathbb{Z}_p$ be the corresponding map. The purpose of §3.2 is to study the family of characters ρ_χ^{ext} . Using work of Kesters and Zhu in [9], we know that $\text{HP}(X_\infty^{\text{ext}}/\mathbb{A}_{\mathbb{F}_q}^1)$ and $\text{NP}(\rho_\chi^{\text{ext}})$ touch periodically when χ is a finite character of sufficiently large order. With a minimal amount of extra work, the theory

of Kusters-Zhu may be extended to include the equicharacteristic character χ_0 (see Theorem 3.10). This periodic touching between Newton and Hodge polygons is necessary to verify a technical hypotheses needed to apply results from [13] (see Corollary 3.12).

Applying Local-to-Global results We are now in a position to apply the main results from [13]. Assume that \overline{X} is ordinary and let X_∞/X be a \mathbb{Z}_p -tower with δ -stable monodromy, corresponding to $\rho : \pi_1(X) \rightarrow \mathbb{Z}_p$. Let χ be either a finite character or the equicharacteristic character. For each $P \in S$ we may localize to obtain a tower $F_{P,\infty}/F_P$, which extends as above to a \mathbb{Z}_p -tower tower $X_{P,\infty}^{\text{ext}}/\mathbb{A}_{\mathbb{F}_q}^1$ over the line.

The main result of [13] roughly states that $\text{NP}^{<e_\chi}(\rho_\chi)$ and $\text{HP}^{<e_\chi}(X_\infty/X)$ share a vertex if and only if $\text{NP}^{<e_\chi}(\rho_{P,\chi}^{\text{ext}})$ and $\text{HP}^{<e_\chi}(X_{P,\infty}^{\text{ext}}/\mathbb{A}_{\mathbb{F}_q}^1)$ share a corresponding vertex for each $P \in S$. Using the periodic touching between Newton and Hodge polygons from the previous paragraph, we obtain periodic touching between $\text{NP}^{<e_\chi}(\rho_\chi)$ and $\text{HP}^{<e_\chi}(X_\infty/X)$. This immediately allows us to deduce Theorem 1.8. To establish Theorem 1.9, we use a classical result on exponential sums (see e.g. [1]). This result says that $\text{NP}^{<e_\chi}(\rho_{P,\chi}^{\text{ext}})$ and $\text{HP}^{<e_\chi}(X_{P,\infty}^{\text{ext}}/\mathbb{A}_{\mathbb{F}_q}^1)$ are equal when χ has order p and the congruence conditions of Theorem 1.9 is satisfied. We then use the work of Kusters-Zhu (and our equicharacteristic modification) to see that $\text{NP}^{<e_\chi}(\rho_{P,\chi}^{\text{ext}})$ and $\text{HP}^{<e_\chi}(X_{P,\infty}^{\text{ext}}/\mathbb{A}_{\mathbb{F}_q}^1)$ agree for any finite χ or for χ_0 . Theorem 1.9 then follows from the main result of [13].

1.4 Future Work

It would be interesting to understand what happens when one twists a δ -stable tower X_∞/X by a motive M pure of weight k . Our expectation is that $M \otimes X_\infty/X$ exhibits slope uniformity in the interval $[0, k + 1]$. Perhaps a natural first step would be to consider M coming from an Artin representation. Indeed, let $Y \rightarrow X$ be a Galois cover and consider the pullback tower Y_∞/Y . Then Y_∞/Y will be δ' -stable for some $\delta' = (\delta'_p)$ that depends on δ and the ramification of $Y \rightarrow X$. In particular, Y_∞/Y is slope uniform by Theorem 1.5. We have a decomposition $\text{Ind}_X^Y(Y_\infty/Y) = \bigoplus \psi \otimes X_\infty/X$, where the sum is taken over the irreducible representations of $\text{Gal}(Y/X)$. We expect that each $\psi \otimes X_\infty/X$ is slope uniform as well.

Another interesting direction would be to explore other classes of \mathbb{Z}_p -towers. Our notion of δ -stable monodromy is natural condition from an analytic standpoint, but is somewhat ad-hoc from a geometrical point of view. Instead, one may consider geometric towers, i.e., towers coming from the relative p -adic étale cohomology of a smooth fibration $Z \rightarrow X$. This includes the Igusa tower, where the equicharacteristic L -function is closely related to the space of $\mathbb{F}_q[[T]]$ -valued modular forms. In general, geometric towers are not δ -stable. However, as demonstrated by the first author in [10], the monodromy of a geometric towers exhibits a similar type of stability. Daqing Wan has conjectured that geometric towers should be slope stable in the appropriate sense, but essentially nothing is known in this direction.

Finally, we believe that for a sufficiently well-behaved tower X_∞/X , geometric data should be encoded in the various L -functions under consideration. For example, we hope to establish a combinatorial formula for the a -numbers of each \overline{X}_n , depending on

the Newton polygon of the equicharacteristic L -function and the ramification breaks of the tower. This is an ongoing project with Jeremy Booher and Bryden Cais, motivated by computations and conjectures formulated by Booher and Cais in [2]. Our hope is that this formula holds for all δ -stable towers over any ordinary curve. It seems likely that the gluing methods developed in the prequel article [13] will allow one to bootstrap results of this form from the special case $X = \mathbb{A}_{\mathbb{F}_q}^1$. More generally, we hope that for well behaved towers the T -adic L -function defined Liu-Wan [14] should contain information on invariants of the Dieudonné module associated to $\text{Jac}(X_n)$.

1.5 Acknowledgments

We would like to thank Daqing Wan for many discussions about the nature of \mathbb{Z}_p -towers.

2 \mathbb{Z}_p -Towers

2.1 Artin-Schreier-Witt Theory via Witt Vectors

Let A be a ring and let $W(A)$ denote the ring of p -typical Witt vectors of A . Recall that as sets we have $W(A) = A^{\mathbb{N}}$. The ring structure on $W(A)$ is characterized uniquely by the requirement that, for each $i \geq 0$, the *ghost map* $w_i : W(A) \rightarrow A$ defined by

$$w_i(a_0, a_1, \dots) = \sum_{j=0}^n p^j a_j^{p^{i-j}}$$

is a functorial homomorphism of rings ([7], 15.3.10). Let X be an \mathbb{F}_p -scheme. Let $X_{\text{ét}}$ denote the small étale site of X . For each $n \geq 0$, the assignment $U \mapsto W_n(\mathcal{O}_U)$ defines a sheaf of rings on $X_{\text{ét}}$, which we denote by \tilde{W}_n . We define $\tilde{W} = \varprojlim_n \tilde{W}_n$. We equip \tilde{W} with the \mathbb{Z}_p -module endomorphism

$$\wp = F - \text{id},$$

where F denotes the Frobenius endomorphism on \tilde{W} .

Lemma 2.1. *Let X be an \mathbb{F}_p -scheme. For each $n \geq 0$, there is an exact sequence of étale sheaves of \mathbb{Z}_p -modules*

$$0 \rightarrow W_n(\mathbb{F}_p) \rightarrow \tilde{W}_n \xrightarrow{\wp} \tilde{W}_n \rightarrow 0. \quad (5)$$

Proof. This is well known for the case $n = 0$ (see e.g. [5]). The general case follows from an inductive lifting argument. \square

Lemma 2.2. *Let X be an affine scheme over \mathbb{F}_p . Then for all $n \geq 0$, $H^1(X_{\text{ét}}, \tilde{W}_n) = 0$.*

Proof. Noting that $\tilde{W}_0 = \mathbb{G}_a$, by Hilbert's theorem 90 [17, Remark 9.8] we have

$$H^1(X_{\text{ét}}, \tilde{W}_0) = H^1(X, \mathcal{O}_X) = 0,$$

since X is affine. The general case follows from an inductive lifting argument as above. \square

Theorem 2.3. *Let $X = \text{Spec}(\overline{A})$ be an affine scheme over \mathbb{F}_p . There is a natural isomorphism of \mathbb{Z}_p -modules*

$$W(\overline{A})/\wp W(\overline{A}) \xrightarrow{\sim} \text{Hom}(\pi_1(X), \mathbb{Z}_p).$$

Proof. By Lemma 2.1, for each $n \geq 0$ we have a long exact sequence

$$0 \rightarrow W_n(\mathbb{F}_p) \rightarrow W_n(\overline{A}) \xrightarrow{\wp} W_n(\overline{A}) \rightarrow H^1(X_{\text{ét}}, W_n(\mathbb{F}_p)) \rightarrow H^1(X_{\text{ét}}, \tilde{W}_n) \rightarrow \dots.$$

Note that $H^1(X_{\text{ét}}, W_n(\mathbb{F}_p)) = \text{Hom}(\pi_1(X), W_n(\mathbb{F}_p))$. By Lemma 2.2, the map

$$W_n(\overline{A})/\wp W_n(\overline{A}) \rightarrow \text{Hom}(\pi_1(X), W_n(\mathbb{F}_p))$$

is an isomorphism. The theorem follows by passing to the inverse limit. \square

2.2 Artin-Schreier-Witt Theory via Flat Liftings

Let $X = \text{Spec}(\overline{A})$ be an affine scheme of characteristic p . It will be convenient to replace the ring of Witt vectors $W(\overline{A})$ by a suitable lifting of \overline{A} to characteristic 0. Let R be a complete Noetherian local ring with maximal ideal \mathfrak{m} and residue field \mathbb{F}_p . By a *flat lifting* of X over R , we will mean a pair (A, σ) where A is a lifting of \overline{A} to a flat R -algebra, and $\sigma : A \rightarrow A$ is a lifting of the absolute Frobenius endomorphism of X .

Following our approach in [13], we will typically fix a flat lifting (A, σ) of X over \mathbb{Z}_p and then pass to general R by base change. Over \mathbb{Z}_p , the connection between flat liftings and Witt vectors is provided by the following:

Lemma 2.4 ([7, Lemma 17.6.9]). *There is a unique ring map $D_\sigma : A \rightarrow W(A)$ making the following diagram commute for each $i \geq 0$:*

$$\begin{array}{ccc} A & \xrightarrow{D_\sigma} & W(A) \\ & \searrow \sigma^i & \downarrow w_i \\ & & A \end{array}$$

By Lemma 2.4 and functoriality of the Witt construction, we obtain a *Frobenius-compatible* map

$$\overline{D}_\sigma : A \xrightarrow{D_\sigma} W(A) \rightarrow W(\overline{A}).$$

We define as before an additive endomorphism $\wp = \sigma - \text{id} : A \rightarrow A$. Let A^∞ denote the p -adic completion of A . The following justifies our use of flat liftings in the context of Artin-Schreier-Witt theory:

Theorem 2.5. *The natural map $A^\infty/\wp A^\infty \rightarrow W(\overline{A})/\wp W(\overline{A})$ is an isomorphism. In particular, there is a natural isomorphism of \mathbb{Z}_p -modules.*

$$A^\infty/\wp A^\infty \xrightarrow{\sim} \text{Hom}(\pi_1(X), \mathbb{Z}_p).$$

Proof. The reduction of this map modulo p is the identity map on $\overline{A}/\wp\overline{A}$. Since both modules are p -adically complete, the claim follows from [13, Lemma 2.3]. \square

Remark 2.6. The isomorphism of Theorem 2.5 is functorial in the following sense: Let $Y = \text{Spec}(\overline{B})$ be another \mathbb{F}_p -scheme, and let (B, τ) be a flat lifting of Y over \mathbb{Z}_p . If $h : A \rightarrow B$ is any *Frobenius-compatible* map of flat liftings, then there is a commutative diagram:

$$\begin{array}{ccc} A^\infty/\wp A^\infty & \xrightarrow{\sim} & \text{Hom}(\pi_1(X), \mathbb{Z}_p) \\ h \downarrow & & \downarrow h^* \\ B^\infty/\wp B^\infty & \xrightarrow{\sim} & \text{Hom}(\pi_1(Y), \mathbb{Z}_p) \end{array}$$

Definition 2.7. Let x be a closed point of X . The *Teichmüller lifting* of x is the Frobenius-compatible \mathbb{Z}_p -algebra map

$$\hat{x} : A \xrightarrow{\overline{D}_\sigma} W(\overline{A}) \rightarrow W(k(x)),$$

where $k(x)$ denotes the residue field at x .

The Teichmüller lifting \hat{x} is the unique Frobenius-compatible map lifting the natural map $\overline{A} \rightarrow k(x)$. Given an element $f \in A$, we write $f(\hat{x}) \in W(k(x))$ for the value of f at the Teichmüller point \hat{x} .

Proposition 2.8. Let $n \in \mathbb{N} \sqcup \{\infty\}$ and let $f \in A^\infty$. Let $\rho : \pi_1(X) \rightarrow \mathbb{Z}_p$ be the associated map from Artin-Schreier-Witt theory. Let x be a closed point of X . As elements of \mathbb{Z}_p , we have

$$\rho(\text{Frob}_x) = \text{Tr}_{k(x)/\mathbb{F}_p} f(\hat{x}).$$

Here, we identify the Galois group of $k(x)/\mathbb{F}_p$ with that of $W(k(x))/\mathbb{Z}_p$.

Proof. Choose a separable closure $k/k(x)$. By Lemma 2.1, there exists $b \in W(k)$ such that $\wp(b) = f(\hat{x})$. The isomorphism

$$W(k(x))/\wp W(k(x)) \rightarrow \text{Hom}(\text{Gal}(k/k(x)), \mathbb{Z}_p)$$

is explicitly given by $f(\hat{x}) \mapsto (g \mapsto gb - b)$. Let $d = [k(x) : \mathbb{F}_p]$. We see that

$$\rho(\text{Frob}_x) = F^d b - b = \sum_{j=0}^{d-1} F^j (Fb - b) = \text{Tr}_{k(x)/\mathbb{F}_p} f(\hat{x}).$$

\square

2.3 Characters and Splitting Functions

In his proof of the rationality of the zeta function, Dwork uses the classical Artin-Hasse exponential to construct a “splitting function” for the L -functions of additive character sums over the torus \mathbb{G}_m^d . We will now use our notion of a flat lifting to construct general splitting functions over a smooth affine \mathbb{F}_p -scheme X .

Let R be a complete Noetherian local ring with maximal ideal \mathfrak{m} and residue field \mathbb{F}_p . Let $\chi : \mathbb{Z}_p \rightarrow R^\times$ be a continuous character. We associate to χ the parameter

$$\pi_\chi = \chi(1) - 1 \in \mathfrak{m}.$$

Note that for any $a \in \mathbb{Z}_p$, we have $\chi(a) = (1 + \pi_\chi)^a$ and thus $\chi \mapsto \pi_\chi$ gives a one-to-one correspondence between the set of R -valued characters of \mathbb{Z}_p and \mathfrak{m} . The functor sending $R \mapsto \mathfrak{m}$ is representable, namely by the Iwasawa algebra $\Lambda = \mathbb{Z}_p[[T]]$. It follows that there is a universal T -adic character $\chi_T : \mathbb{Z}_p \rightarrow \Lambda^\times$, which corresponds to the topologically nilpotent element $T \in \Lambda$.

Let $X = \text{Spec}(\overline{A})$ be an \mathbb{F}_p -scheme, and let (A, σ) be a flat lifting of X over \mathbb{Z}_p . For every character $\chi : \mathbb{Z}_p \rightarrow R^\times$ as above, we define a flat lifting (A_{π_χ}, σ) over R , where

$$A_{\pi_\chi} = R \otimes_{\mathbb{Z}_p} A,$$

and $\sigma : A_{\pi_\chi} \rightarrow A_{\pi_\chi}$ is the R -linear lifting of Frobenius obtained by base change. For every closed point $x \in |X|$, let us write $R(x) = R \otimes_{\mathbb{Z}_p} W(k(x))$. The Teichmüller map \hat{x} induces a Frobenius-compatible map

$$\hat{x} : A_{\pi_\chi} \rightarrow R(x).$$

Given $E \in A_{\pi_\chi}^\infty$, we will write $E(\hat{x}) \in R(x)$ for the “value” of E at \hat{x} .

Definition 2.9. Let $f \in A^\infty$. A *splitting function* for f at χ is an element $E \in A_{\pi_\chi}^\infty$ satisfying

$$(1 + \pi_\chi)^{\text{Tr}_{k(x)/\mathbb{F}_p} f(\hat{x})} = N_{k(x)/\mathbb{F}_p} E(\hat{x})$$

for all $x \in |X|$. Here, we identify the Galois group of $R(x)/R$ with that of $k(x)/\mathbb{F}_p$.

Remark 2.10. Let $\rho : \pi_1(X) \rightarrow \mathbb{Z}_p$ be the map corresponding to f by Artin-Schreier-Witt theory. Suppose that R is a discrete valuation ring. In the terminology of [13, §5.1], a splitting function for f at χ is a Frobenius structure for the unit-root σ -module over $A_{\pi_\chi}^\infty$ corresponding to $\rho_\chi = \chi \circ \rho$. If f admits a splitting function at χ , then this σ -module is free of rank 1.

Let us write (A_T, σ) for the flat lifting corresponding to the T -adic character χ_T . We claim that every $f \in A^\infty$ admits a splitting function $E_f(T) \in A_T^\infty$ at χ_T . By specialization, we obtain a splitting function at every character $\chi : \mathbb{Z}_p \rightarrow R^\times$ as above. Recall that the *Artin-Hasse exponential series* is the power series

$$E(t) = \exp\left(\sum_{i=0}^{\infty} \frac{t^{p^i}}{p^i}\right) = \prod_{p \nmid n} (1 - t^n)^{-\mu(n)/n} \in 1 + t + t^2 \mathbb{Z}_p[[t]].$$

Here, μ denotes the Möbius function.

Lemma 2.11 ([9, Lemma 2.1]). *For each $j \geq 0$, $E(t)$ induces a bijection $T^j \Lambda \rightarrow 1 + T^j \Lambda$. Moreover, for all $r \in \mathbb{Z}_p$ and all $j \geq 0$ we have*

$$E(rT^j + T^{j+1}\Lambda) = 1 + rT^j + T^{j+1}\Lambda.$$

Definition 2.12. For each $j \geq 0$, we let $\tau_j = \tau_j(T) \in T\Lambda$ be the unique series satisfying $E(\tau_j(T)) = (1 + T)^{p^j}$.

We are now ready to define our T -adic splitting functions. Recall from [7, 17.2.7] that there is a functorial embedding of \mathbb{Z}_p -modules

$$\begin{aligned} W(A) &\rightarrow 1 + tA[[t]] \\ a &\mapsto \prod_{j=0}^{\infty} E(a_j t^{p^j}). \end{aligned}$$

Definition 2.13. For each $j \geq 0$, we define the j th *Artin-Hasse exponential map* to be the composite

$$p^j A \xrightarrow{1/p^j} A \xrightarrow{D_\sigma} W(A) \rightarrow 1 + tA[[t]] \xrightarrow{t \mapsto \tau_j} 1 + \tau_j A[[\tau_j]] \subseteq A_\Lambda^\infty.$$

Let us write $E_f^j = E_f^j(T)$ for the j th Artin-Hasse exponential of $f \in p^j A$. By construction, the assignment $f \mapsto E_f^j$ is \mathbb{Z}_p -linear and is functorial in the flat lifting (A, σ) .

Lemma 2.14 (Dwork's Splitting Lemma). *Let $f \in p^j A^\infty$. Then $E_f^j(T)$ is a splitting function for f at χ_T .*

Proof. By functoriality, we have that $E_f^j(\hat{x}) = E_{f(\hat{x})}^j \in \Lambda(x)$. Since $k(x)$ is perfect, we have a Teichmüller expansion

$$f(\hat{x}) = p^j([f_0] + [f_1]p + \cdots) \in W(k(x)).$$

By \mathbb{Z}_p -linearity of the Artin-Hasse exponential map, it suffices to prove the statement when $f(\hat{x}) = p^{j+k}[c]$. Note that since $\sigma([c]) = [c]^p$, we have $E_{p^j[c]}^j = E([c]\tau_j)$. But then

$$(1 + T)^{\mathrm{Tr}_{k(x)/\mathbb{F}_p} p^{j+k}[c]} = E(\tau_j)^{p^k \mathrm{Tr}_{k(x)/\mathbb{F}_p}[c]} = N_{k(x)/\mathbb{F}_p} E([c]\tau_j)^{p^k} = N_{k(x)/\mathbb{F}_p} E_{p^{j+k}[c]}^j.$$

□

3 \mathbb{Z}_p -Towers of Local Fields

In this section we let F be a local field of characteristic p with residue field \mathbb{F}_q . Let $t \in F$ be a uniformizer. Consider the flat lifting (\mathcal{A}, σ) , where $\mathcal{A} = \mathbb{Z}_q((t))$ and $\sigma : \mathcal{A} \rightarrow \mathcal{A}$ is the unique lifting of Frobenius such that $\sigma(t) = t^p$. Let G denote the absolute Galois group of F , and let $f \in \mathcal{A}^\infty$. By Artin-Schreier-Witt theory, f determines a map

$$\rho : G \rightarrow \mathbb{Z}_p. \tag{6}$$

For each $j \geq 0$, the open subgroup $p^j \mathbb{Z}_p \subseteq \mathbb{Z}_p$ corresponds to a finite extension of local fields F_j/F . We abbreviate the resulting tower by F_∞/F .

3.1 Local Towers with Strictly Stable Monodromy

Our first goal is to characterize the stable monodromy condition (1.2) in terms of a suitable representative $\tilde{f} \in \mathcal{A}^\infty$ of the tower F_∞/F .

Proposition 3.1. *Let $\beta \in \mathbb{F}_q$ such that $\mathrm{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(\beta) \neq 0$. There is a unique $\tilde{f} \in \mathcal{A}^\infty$ such that $f \equiv \tilde{f} \pmod{\wp \mathcal{A}^\infty}$ and*

$$\tilde{f} = c[\beta] + \sum_{k=1}^{\infty} c_k t^{-k},$$

where $c \in \mathbb{Z}_p$, $c_k \in \mathbb{Z}_q$ with $c_k = 0$ whenever $p|k$, and $c_k \rightarrow \infty$ as $k \rightarrow \infty$.

Proof. Let C be any basis for \mathbb{Z}_q as a \mathbb{Z}_p -module, and let

$$B = \{[\beta]\} \sqcup \{ct^{-k} : c \in C, p \nmid k\}.$$

The reduction of B modulo p forms a basis for $F/\wp F$ (see e.g. [8, Example 2.4]). Since $\mathcal{A}^\infty/\wp \mathcal{A}^\infty$ is p -torsion free, it follows from [13, Lemma 2.3] that the p -adic completion of the map

$$\begin{aligned} \bigoplus_{b \in B} \mathbb{Z}_p &\rightarrow \mathcal{A}/\wp \mathcal{A} \\ (c_b)_{b \in B} &\mapsto \sum_{b \in B} c_b b \end{aligned}$$

is an isomorphism. □

From now on we will fix a choice of β as in Proposition 3.1 and let $c_0 = c[\beta]$. Note that the p -adic valuation of c_0 is independent of the choice of β . The p -adic growth of the coefficients c_k is closely connected to the monodromy of F_∞/F :

Proposition 3.2 ([8, Proposition 3.3]). *For all $j \geq 0$, let v_j denote the highest ramification break (in upper numbering) of the finite extension F_j/F . Then*

$$v_j = \begin{cases} p^{j-1} \max\{kp^{-v_p(c_k)} : v_p(c_k) < j\} & \text{if } v_p(c_k) < j \text{ for some } k \\ 0 & \text{otherwise} \end{cases}.$$

It follows from Proposition 3.2 that F_∞/F has δ -stable monodromy for $\delta \in \mathbb{Z}[\frac{1}{p}]$ if and only if the sequence $kp^{-v_p(c_k)}$ attains its maximum at some k_0 , and the maximal value is equal to δ . We will now give another characterization which is more useful for making explicit estimates: For each $k \geq 0$, consider the Teichmüller expansion

$$c_k = \sum_{j=0}^{\infty} [c_{j,k}] p^j \in \mathbb{Z}_q. \tag{7}$$

The convergence condition on the c_k guarantees that we may write

$$\tilde{f} = \sum_{j=0}^{\infty} p^j F_j = \sum_{j=0}^{\infty} p^j \sum_{k=0}^{d_j} [c_{j,k}] t^{-k}, \tag{8}$$

where F_j is a polynomial of degree d_j in t^{-1} .

Lemma 3.3. *The following are equivalent:*

1. F_∞/F has δ -stable monodromy.
2. The sequence (d_j/p^j) has a maximum, with the maximal value equal to δ .

Proof. Observe that if $v_p(c_k) > j$ then we must have $d_j < k$. In other words, the sequence (d_j) is bounded above by the increasing sequence

$$q_j = \max\{k : v_p(c_k) \leq j\}.$$

Moreover, we have $d_j = q_j$ whenever j is the minimal index at which d_j attains this value. It follows that (d_j/p^j) has a maximum if and only if (q_j/p^j) has a maximum, and that the two maxima agree. On the other hand, we have an upper bound

$$\begin{aligned} q_j &= p^j \max\{kp^{-j} : v_p(c_k) < j+1\} \\ &\leq p^j \max\{kp^{-v_p(c_k)} : v_p(c_k) < j+1\} \\ &= v_{j+1}, \end{aligned}$$

with equality if and only if there exists some k such that $v_p(c_k) = j$.

First assume that F_∞/F has δ -stable monodromy. Then the sequence $kp^{-v_p(c_k)}$ has maximal value δ , so that $q_j \leq p^j \delta$ for all j . Let k_0 be the minimal value of k such that $kp^{-v_p(c_k)} = \delta$, and let $j_0 = v_p(c_{k_0})$. Then $q_{j_0} = \delta p^{j_0}$ so that q_j/p^j attains its maximal value δ when $j = j_0$.

Conversely, suppose that (d_j/p^j) has a maximum with maximal value δ . Let j_0 be the smallest value of j such that $q^j/p^j = \delta$, and let $k_0 = q_{j_0} = p^{j_0} \delta$. Then by the definition of (q_j) we have $j_0 = v_p(c_{k_0})$, so that $q_{j_0} = v_{j_0+1}$. To complete the proof, it suffices to show that for all $k > k_0$ the value $kp^{-v_p(c_k)}$ is bounded by δ . To see this, let $j = v_p(c_k)$. Since $k > k_0$, we have $j \geq j_0$ and since $v_p(c_k) < j+1$ we have

$$k \leq q_j \leq p^j \delta.$$

Finally, we see that

$$kp^{-v_p(c_k)} \leq p^j \delta p^{-v_p(c_k)} = \delta.$$

□

Our next goal is to construct a local splitting function with good π_χ -adic growth properties. As in 2.3, we first define a splitting function at the T -adic character χ_T and then specialize to obtain splitting functions at all other χ .

Definition 3.4. The *local splitting function* of the tower F_∞/F is defined to be

$$\tilde{E}(T) = \prod_{j=0}^{\infty} E_{F_j}^j(T) = \prod_{j=0}^{\infty} \prod_{k=0}^{d_j} E([c_{j,k}] \tau_j(T) t^{-k}) \quad (9)$$

If χ is any continuous character of \mathbb{Z}_p , then we let $\tilde{E}(\pi_\chi) \in R$ denote the specialization of $\tilde{E}(T)$ along $T \mapsto \pi_\chi$. To study the π_χ -adic convergence properties of $\tilde{E}(\pi_\chi)$, we recall our conventions on growth conditions from [13, §2.1]. For every real number $m > 0$, consider the subring

$$\mathcal{A}_{\pi_\chi}^m = \left\{ \sum_{k=\infty}^{\infty} a_k t^{-k} : v_{\pi_\chi}(a_k) \geq \frac{k}{m} \right\}.$$

Each $\mathcal{A}_{\pi_\chi}^m$ is a π_χ -adically complete subring of $\mathcal{A}_{\pi_\chi}^\infty$. We define $\mathcal{A}_{\pi_\chi}^\dagger$ to be the union of the $\mathcal{A}_{\pi_\chi}^m$. Note in particular that $E_{F_j}^j(\pi_\chi) \in \mathcal{A}_{\pi_\chi}^m$, where $m = d_j/v_{\pi_\chi}(\tau_j(\pi_\chi))$.

Lemma 3.5. *Suppose that χ is finite or that R has characteristic p . Then for all $j \geq 0$,*

$$v_{\pi_\chi}(\tau_j(\pi_\chi)) = \begin{cases} p^j & 0 \leq j < m_\chi \\ \infty & j \geq m_\chi \end{cases}.$$

Proof. By Lemma 2.4, we have

$$v_{\pi_\chi}(\tau_j(\pi_\chi)) = v_{\pi_\chi}(E(\tau_j(\pi_\chi)) - 1) = v_{\pi_\chi}((1 + \pi_\chi)^{p^j} - 1).$$

If R has characteristic p , then this is equal to $v_{\pi_\chi}(\pi_\chi^{p^j}) = p^j$ for all j . Otherwise, the result follows since $1 + \pi_\chi$ is a p^{m_χ} -root of unity. \square

As an immediate consequence of Lemma 3.3 and Lemma 3.5, we obtain an estimate for our local splitting function:

Theorem 3.6. *Suppose that F_∞/F has δ -stable monodromy. If χ is equicharacteristic or finite, then $\tilde{E}_{\pi_\chi} \in \mathcal{A}_{\pi_\chi}^\delta$.*

3.2 Local Newton Polygons

Let χ be a non-trivial R -valued character of \mathbb{Z}_p . Borrowing our terminology from [13, §5.3], we will say that ρ_χ is *overconvergent* if $\tilde{E}(\pi_\chi) \in \mathcal{A}_{\pi_\chi}^\dagger$. More specifically, given a rational number $\delta > 0$ we will say that ρ_χ is δ -*overconvergent* if $\tilde{E}(\pi_\chi) \in \mathcal{A}_{\pi_\chi}^\delta$. For example, Theorem 3.6 states that if F_∞/F has δ -stable monodromy, then ρ_χ is δ -overconvergent if χ is finite or if R has characteristic p .

Let $U_p : \mathcal{A} \rightarrow \mathcal{A}$ be the local U_p -operator of [13, §4.2]. If ρ_χ is overconvergent, then we define a local p -Dwork operator

$$\tilde{\Theta} = U_p \circ \tilde{E}(\pi_\chi) : \mathcal{A}_{\pi_\chi}^\dagger \rightarrow \mathcal{A}_{\pi_\chi}^\dagger,$$

The operator $\tilde{\Theta}$ is R -linear, and the iterate $\tilde{\Theta}_q = \tilde{\Theta}^{v_p(q)}$ is R_q -linear. The R -submodule

$$\mathcal{A}_{\pi_\chi}^{\dagger, \text{tr}} = t^{-1} R_q \langle t^{-1} \rangle \cap \mathcal{A}_{\pi_\chi}^\dagger$$

consisting of truncated overconvergent series in t is invariant under the action of $\tilde{\Theta}$. By [18, Theorem 2.1], the action of $\tilde{\Theta}_q$ on the K_q -vector space $\mathcal{V}_{\pi_\chi}^{\dagger, \text{tr}} = K \otimes_R \mathcal{A}_{\pi_\chi}^{\dagger, \text{tr}}$ is nuclear.

Definition 3.7. The *local normalized Newton polygon* of ρ_χ is defined to be

$$\text{NP}(\rho_\chi) = \text{NP}_{\pi_{q,\chi}}(\tilde{\Theta}_q | \mathcal{V}_{\pi_\chi}^{\dagger, \text{tr}}).$$

Definition 3.8. Let $\delta > 0$ be a rational number. We define the δ -*Hodge polygon* to be the infinite convex polygon with slope set

$$\left\{ \frac{p-1}{\delta}, \frac{2(p-1)}{\delta}, \dots \right\}.$$

Proposition 3.9. *If ρ_χ is δ -overconvergent, then $\text{NP}(\rho_\chi) \succeq \text{HP}(\delta)$.*

Theorem 3.10. *Suppose that F_∞/F is a \mathbb{Z}_p -tower of local fields with δ -stable monodromy. Let m_0 be the smallest non-negative integer such that $v_j = p^{j-1}\delta$ for all $j > m_0$. Let $r > 0$. The following are equivalent:*

1. *For some $\chi : \mathbb{Z}_p \rightarrow R^\times$ which is equicharacteristic or finite with $m_\chi > m_0$, the polygons $\text{NP}^{<r}(\rho_\chi)$ and $\text{HP}^{<r}(\delta)$ have the same terminal point.*
2. *For every $\chi : \mathbb{Z}_p \rightarrow R^\times$ which is equicharacteristic or finite with $m_\chi > m_0$, the polygons $\text{NP}^{<r}(\rho_\chi)$ and $\text{HP}^{<r}(\delta)$ have the same terminal point.*

Proof. This theorem more or less follows from the theory developed in [9]. However, [9] ignores equicharacteristic characters. Thus, we believe it is beneficial to outline their work and explain the extra observations needed to handle this case. In [9], Kusters and Zhu construct a “very generic” discrete valuation ring R_δ with valuation v_δ (note that they denote this ring by R). For each χ there is an evaluation map $\text{ev}_\chi : R_\delta \rightarrow R_\chi$. If χ is a finite character, then valuations “go up” under the evaluation map

$$v_{\pi_\chi}(\text{ev}_\chi(x)) \geq v_\delta(x) \tag{10}$$

for all $x \in R_\delta$. Furthermore, the following holds:

$$\begin{array}{ccc} \text{For some finite character} & & \text{For every finite character} \\ \chi \text{ with } m_\chi > m_0 \text{ such that} & \iff & \chi \text{ with } m_\chi > m_0 \text{ we have} \\ v_{\pi_\chi}(\text{ev}_\chi(x)) = v_\delta(x) & & v_{\pi_\chi}(\text{ev}_\chi(x)) = v_\delta(x) \end{array} \tag{11}$$

This is Lemma 6.1 in [9]. The proof of this Lemma extends without modification to include any character with values in a discrete valuation ring of characteristic p .

Kusters and Zhu then construct a generic “characteristic series” $C(\pi, s) = 1 + sR_\delta[[s]]$ which satisfies a certain entireness property with respect to v_δ , as well as the interpolation property

$$\text{ev}_\chi C(\pi, s) = C(\tilde{\Theta}_q | \mathcal{V}_{\pi_\chi}^{\text{tr}, \dagger}, s)$$

for any non-trivial χ . The entireness property of $C(\pi, s)$ allows us to define a Newton polygon $\text{NP } C(\pi, s)$ using the normalized valuation $v_p(q)v_\delta$. The main estimate of Kusters-Zhu [9, Proposition 5.7] states that

$$\text{NP } C(\pi, s) \succeq \text{HP}(\delta).$$

Now if χ is finite or R has characteristic p , then (10) guarantees that $\text{NP}(\rho_\chi) \succeq \text{NPC}(\pi, s)$. The theorem follows immediately from (11). \square

Corollary 3.11. *In the setting of Theorem 3.10, suppose that χ equicharacteristic or finite with $m_\chi > m_0$. Let $d = p^{m_0}\delta$. For all $n \geq 1$, the polygons $\text{NP}(\rho_\chi^{\text{ext}})$ and $\text{HP}(\delta)$ agree on the interval $[nd - 1, nd]$.*

Proof. By Theorem 3.10, it suffices to prove the statement for a finite character χ of order $m_\chi = m_0 + 1$. This is [13, Theorem 7.10]. \square

Corollary 3.12. *In the setting of Theorem 3.10, suppose that χ equicharacteristic or finite with $m_\chi > m_0$. Then ρ_χ is π_χ -adically δ -Hodge.*

Proof. The statement means that the δ -Hodge polygon $\text{HP}(\delta)$ agrees with a certain local Hodge polygon $\text{HP}(\rho_\chi)$ that we have defined in [13, §7.2]. Since the polygons $\text{NP}(\rho_\chi)$ and $\text{HP}(\delta)$ meet periodically by Corollary 3.11, the statement follows immediately from the criterion [13, Proposition 7.9] \square

Theorem 3.13. *In the setting of Theorem 3.10, suppose that χ equicharacteristic or finite with $m_\chi > 0$. Then $\text{NP}(\rho_\chi)$ and $\text{HP}(\delta)$ agree if and only if:*

1. $m_0 = 0$, so that $\delta \in \mathbb{Z}$.
2. $p \equiv 1 \pmod{\delta}$.

Proof. By Theorem 3.10, it suffices to prove the claim when χ is finite of order p . But we have shown this previously [13, Theorem 1.3]. \square

To conclude this section, let us briefly recall the notion of local-to-global extensions from [13, §7.2]. Consider the pair (A, σ) , where $A = \mathbb{Z}_q[t^{-1}]$ and $\sigma : A \rightarrow A$ is the lifting of Frobenius defined by $\sigma(t^{-1}) = t^{-p}$. Since $\tilde{f} \in A^\infty$, the tower F_∞/F extends in a natural way to a tower $X_\infty/\mathbb{A}_{\mathbb{F}_q}^1$, or equivalently a continuous map

$$\rho^{\text{ext}} : \pi_1(\mathbb{A}_{\mathbb{F}_q}^1) \rightarrow \mathbb{Z}_p.$$

We may then consider the family of characters $\rho_\chi^{\text{ext}} = \chi \circ \rho_\chi$, as χ varies through the continuous R -valued characters of \mathbb{Z}_p as above. For each such χ , let us write $L(\rho_\chi^{\text{ext}}, s)$ for the Artin L -function of ρ_χ^{ext} over $\mathbb{A}_{\mathbb{F}_q}^1$. If ρ_χ is overconvergent, then the Monsky trace formula guarantees that $L(\rho_\chi, s)$ is analytic in the disk $v_\pi(s) > -v_\pi(q)$. Moreover, we have the relation:

$$\text{NP}^{<e_\chi}(\rho_\chi^{\text{ext}}) = \text{NP}_{\pi_{q,\chi}}^{<e_\chi} L(\rho_\chi^{\text{ext}}, s) = \text{NP}^{<e_\chi}(\rho_\chi).$$

This shows that the truncated local polygon $\text{NP}^{<e_\chi}(\rho_\chi)$ does not depend on our particular choice of lifting (\mathcal{A}, σ) , or the choice of local splitting function $\tilde{E}(T)$.

4 \mathbb{Z}_p -Towers of Curves

In this final section we deduce our main theorems on \mathbb{Z}_p -towers of curves. Let X/\mathbb{F}_q be a smooth affine curve whose smooth compactification \overline{X} is ordinary. Let X_∞/X be a \mathbb{Z}_p -tower of curves with δ -stable monodromy, corresponding to a continuous and surjective map

$$\rho : \pi_1(X) \rightarrow \mathbb{Z}_p$$

4.1 Results for Global Towers

For each $P \in S$, let G_P denote the absolute Galois group of F_P . We obtain by restriction a \mathbb{Z}_p -tower of local fields $F_{P,\infty}/F_P$ with δ_P -stable monodromy. This local tower corresponds to a continuous and surjective map

$$\rho_P : G_P \rightarrow \mathbb{Z}_p.$$

As in §3.2, the local character ρ_P extends to $\rho_P^{\text{ext}} : \pi_1(\mathbb{A}_{\mathbb{F}_q}^1) \rightarrow \mathbb{Z}_p$. Let χ be an R -valued character of \mathbb{Z}_p . Assume χ is finite or that R has characteristic p . By Theorem 3.6, ρ_χ is π_χ -adically δ -Hodge in the sense that $\rho_{\chi,P}$ is π_χ -adically δ_P -Hodge for every $P \in S$. By [13, Proposition 4.11] and the Monsky trace formula [13, Theorem 4.5], the L -function $L(\rho_\chi, s)$ is entire.

For each $j \geq 0$, let $v_{P,j}$ denote the highest ramification break (in upper numbering) of the finite extension $F_{P,j}/F_P$. We define m_0 to be the *smallest* integer such that $v_{P,m_0+1} = p^{m_0}\delta_P$ for all P . Note that this means $v_{P,j} = p^{j-1}\delta_P$ for any $j > m_0$. We can now prove the combined form of Theorems 1.8 and 1.18.

Theorem 4.1. *Assume that \overline{X} is ordinary. Let $d = p^{m_0} \sum_{P \in S} \delta_P$. Let χ be a character that is either finite or equicharacteristic and assume $m_\chi > m_0$. Then for all $1 \leq n < p^{m_\chi - m_0 - 1}$, the polygons $\text{NP}^{<e_\chi}(\rho_\chi)$ and $\text{HP}(X_\infty/X)$ agree on the interval*

$$[g - 1 + nd, g - 1 + |S| + nd]. \quad (12)$$

Proof. Let $e_0 = p^{m_0}(p - 1)$, and let $\varepsilon < \min\{(p - 1)/\delta_P : P \in S\}$. By Corollary 3.11, the polygons $\text{NP}^{<r}(\rho_{\chi,P}^{\text{ext}})$ and $\text{HP}^{<r}(\delta_P)$ have the same terminal point for $r = ne_0$ or $ne_0 - \varepsilon$. By [13, Theorem 6.14], the polygons $\text{NP}^{<r}(\rho_\chi)$ and $\text{HP}^{<r}(X_\infty/X)$ have the same terminal point for such r as long as $r \leq e_\chi$. This condition is equivalent to $n < p^{m_\chi - m_0 - 1}$. The theorem follows since the restriction of $\text{HP}(X_\infty/X)$ to the interval (12) consists of $|S|$ segments of slope ne_0 . \square

Proof of Theorem 1.5. Assume $m_\chi > m_0$. By renormalizing, we see that $\text{NP}_q^{<1}(\rho_\chi)$ lies above $\frac{1}{e_\chi} \text{HP}^{<e_\chi}(\delta)$ and that both polygons meet periodically with period d . In particular, we see that $\text{NP}_q^{<1}(\rho_\chi) \rightarrow \frac{1}{e_\chi} \text{HP}^{<e_\chi}(\delta)$ as $m_\chi \rightarrow \infty$. The result follows since the slopes of $\frac{1}{e_\chi} \text{HP}^{<e_\chi}(\delta)$ are equidistributed in $[0, 1)$ as $m_\chi \rightarrow \infty$. \square

Theorem 4.2. *If χ is finite (resp. the equicharacteristic character), then $\text{NP}^{<e_\chi}(\rho_\chi^{\text{ext}})$ and $\text{HP}^{<e_\chi}(X_\infty/X)$ (resp. $\text{NP}(\rho_\chi^{\text{ext}})$ and $\text{HP}(X_\infty/X)$) agree if and only if:*

1. $m_0 = 0$, so that $\delta_P \in \mathbb{Z}$ for all $P \in S$.
2. $p \equiv 1 \pmod{\delta_P}$ for all $P \in S$.

Proof. Consider the finite character case. We know from [13, Remark 1.4] that the conditions are necessary. By [13, Theorem 6.14], it suffices to prove that $\text{NP}^{<e_X}(\rho_{X,P}^{\text{ext}})$ and $\text{HP}^{<e_X}(\delta_P)$ agree for all $P \in S$. This follows immediately from Theorem 3.13. The equicharacteristic case is almost identical. \square

Corollary 4.3. *Let $\rho : \pi_1(X) \rightarrow \mathbb{C}_p^\times$ be a character of finite order p^n . For each $P \in S$, let d_P denote the Swan conductor at P . Then $\text{NP}^{<e_X}(\rho) = \text{HP}^{<e_X}(\rho)$ if and only if*

1. \overline{X} is ordinary.
2. $\delta_P = d_P/p^{n-1} \in \mathbb{Z}$ for all $P \in S$.
3. $p \equiv 1 \pmod{\delta_P}$ for all $P \in S$.

Proof. Once again, we know from [13, Remark 1.4] that the conditions are necessary. Moreover, by [13, Theorem 1.1], it suffices to prove the claim when $X = \mathbb{A}_{\mathbb{F}_q}^1$. By Theorem 4.2, we need only show that ρ factors through a \mathbb{Z}_p -tower $X_\infty/\mathbb{A}_{\mathbb{F}_q}^1$ with δ_∞ -stable monodromy at ∞ . To see this, choose a factorization

$$\rho : \pi_1(X) \rightarrow \mathbb{Z}/p^n\mathbb{Z} \rightarrow \mathbb{C}_p^\times.$$

The first map can be obtained via Artin-Schreier-Witt theory from a polynomial $\overline{f} = \overline{c}_0 + \cdots + \overline{c}_d t^d \in \mathbb{Z}_q/p^n\mathbb{Z}_q[t]$ with $\overline{c}_k = 0$ whenever $p|k$. Choose a lifting $f = c_0 + \cdots + c_d t^d \in \mathbb{Z}_q[t]$ of \overline{f} such that $c_k = 0$ whenever $p|k$, and let $X_\infty/\mathbb{A}_{\mathbb{F}_q}^1$ be the corresponding tower. From Proposition 3.2, we see immediately that $X_\infty/\mathbb{A}_{\mathbb{F}_q}^1$ has δ_∞ -stable monodromy at ∞ . This completes the proof. \square

4.2 Results for Twisted Towers

We are now able to prove Theorem 1.11 and Theorem 1.12 by combining the results of §4.1 with the results of [11]. We begin by recalling the notation and results from [11]. Let $\psi : \pi_1(X) \rightarrow \mathbb{Z}_p^\times$ be a finite tame character of order c . Note that $c|(p-1)$. Let X^{tame} be the cover of X corresponding to $\ker(\psi)$ and let s_P be the ramification index over P . For each $P \in S$ we obtain a local representation $\psi_P : G_P \rightarrow \mathbb{Z}_p^\times$. There exists $e_P \in \frac{1}{p-1}\mathbb{Z}$ such that G_P acts on $u_P^{e_P}$ by ψ_P . Note that e_P is well defined up to addition by an integer. We define the *exponent* $\mathbf{e}_P \in \frac{1}{p-1}\mathbb{Z}/\mathbb{Z}$ of ψ at P to be the equivalence class of e_P modulo \mathbb{Z} . This is analogous to the notion of exponents in the theory of complex differential equations with regular singularities. We then define $0 \leq \epsilon_P \leq p-2$ to be the unique integer such that $\frac{\epsilon_P}{p-1}$ is in \mathbf{e}_P . Note that $\epsilon_P = 0$ if and only if ψ is unramified at P . Finally, we define a global invariant

$$\Omega_\psi = \frac{1}{p-1} \sum_{P \in S} \epsilon_P.$$

One may show that Ω_ψ is an integer (see [11, §5.3.2]).

Let X_∞/X be δ -stable tower. We define the “twisted” Hodge polygon as follows:

$$\text{HP}(\psi \otimes X_\infty/X) = \underbrace{\{0, \dots, 0\}}_{g-1+|S|-\Omega_\psi} \sqcup \bigsqcup_{P \in S} \left\{ \frac{1-\epsilon_P}{\delta_P}, \frac{2-\epsilon_P}{\delta_P}, \dots \right\}.$$

By [11, Theorem 1.1] we have

$$\text{NP}^{<e_\chi}(\psi \otimes \chi) \succeq \text{HP}^{<e_\chi}(\psi \otimes X_\infty/X) \quad (13)$$

for any finite character $\chi : \text{Gal}(X_\infty/X) \rightarrow \mathbb{C}_p^\times$.

Proof. (Proof of Theorem 1.11 and Theorem 1.12) Let $X_\infty^{\text{tame}}/X^{\text{tame}}$ be the pullback of the tower X_∞/X along $X^{\text{tame}} \rightarrow X$. Note that $X_\infty^{\text{tame}}/X^{\text{tame}}$ is δ^{tame} -stable, where $\delta_P^{\text{tame}} = s_P \delta_P$. We have a decomposition

$$\text{Ind}_X^{X^{\text{tame}}}(X_\infty^{\text{tame}}/X^{\text{tame}}) = \bigoplus_{i=0}^{c-1} \psi^{\otimes i} \otimes X_\infty/X.$$

This follows by considering the induction $\text{Ind}_X^{X^{\text{tame}}}(\chi^{\text{tame}})$ for any character $\chi^{\text{tame}} : \text{Gal}(X_\infty^{\text{tame}}/X^{\text{tame}}) \rightarrow \mathbb{C}_p^\times$. We obtain a corresponding decomposition of Hodge polygons

$$\text{HP}(X_\infty^{\text{tame}}/X^{\text{tame}}) = \bigsqcup_{i=0}^{c-1} \text{HP}(\psi^{\otimes i} \otimes X_\infty/X). \quad (14)$$

This decomposition essentially follows from the Riemann-Hurwitz theorem and by observing how the exponents of $\psi^{\otimes i}$ vary with i . Similarly, for any $\chi : \text{Gal}(X_\infty/X) \rightarrow \mathbb{C}_p^\times$ we have a decomposition

$$\text{NP}(\chi^{\text{tame}}) = \bigsqcup_{i=0}^{c-1} \text{NP}(\psi^{\otimes i} \otimes \chi). \quad (15)$$

By combining equations (14) and (15) with the Newton-over-Hodge result (13), we see that $\text{NP}^{<r}(\chi^{\text{tame}})$ and $\text{HP}^{<r}(X_\infty^{\text{tame}}/X^{\text{tame}})$ have the same endpoints if and only if $\text{NP}^{<r}(\psi^{\otimes i} \otimes \chi)$ and $\text{HP}^{<r}(\psi^{\otimes i} \otimes X_\infty/X)$ have the same endpoints for each $0 \leq i < c$. Slope uniformity then follows from Theorem 1.8 applied to $X_\infty^{\text{tame}}/X^{\text{tame}}$. The slope stability result follows from Theorem 1.9 applied to $X_\infty^{\text{tame}}/X^{\text{tame}}$. \square

References

- [1] Alan Adolphson and Steven Sperber. Exponential sums and Newton polyhedra: cohomology and estimates. *Ann. of Math. (2)*, 130(2):367–406, 1989.

- [2] Jeremy Booher and Bryden Cais. Iwasawa theory for p -torsion class group schemes in characteristic p , 2021. preprint, arXiv:2107.12555.
- [3] Irene I. Bouw. The p -rank of ramified covers of curves. *Compositio Math.*, 126(3):295–322, 2001.
- [4] Christopher Davis, Daqing Wan, and Liang Xiao. Newton slopes for Artin-Schreier-Witt towers. *Math. Ann.*, 364(3-4):1451–1468, 2016.
- [5] P. Deligne. *Cohomologie étale*, volume 569 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1977. Séminaire de géométrie algébrique du Bois-Marie SGA 4 $\frac{1}{2}$.
- [6] Bernard Dwork. On the zeta function of a hypersurface. *Inst. Hautes Études Sci. Publ. Math.*, (12):5–68, 1962.
- [7] Michiel Hazewinkel. *Formal Groups and Applications*. AMS Chelsea Publishing, 1978.
- [8] Michiel Kusters and Daqing Wan. Genus growth in \mathbb{Z}_p -towers of function fields. *Proc. Amer. Math. Soc.*, 146(4):1481–1494, 2018.
- [9] Michiel Kusters and Hui June Zhu. On slopes of L -functions of \mathbb{Z}_p -covers over the projective line. *Journal of Number Theory*, 187:430 – 452, 2018.
- [10] Joe Kramer-Miller. The monodromy of unit-root F -isocrystals with geometric origin. to appear in *Compositio Mathematica*.
- [11] Joe Kramer-Miller. p -adic estimates of abelian Artin L -functions on curves, 2020. to appear in *Forum of Mathematics*, σ .
- [12] Joe Kramer-Miller. p -adic estimates of exponential sums on curves. *Algebra & Number Theory*, 15(1):141–171, Mar 2021.
- [13] Joe Kramer-Miller and James Upton. Newton polygons of sums on curves I: Local-to-global theorems. Preprint, 2021.
- [14] Chunlei Liu and Daqing Wan. T -adic exponential sums over finite fields. *Algebra & Number Theory*, 3(5):489–509, 2009.
- [15] Chunlei Liu and Dasheng Wei. The L -functions of Witt coverings. *Math. Z.*, 255(1):95–115, 2007.
- [16] B. Mazur, J. Tate, and J. Teitelbaum. On p -adic analogues of the conjectures of Birch and Swinnerton-Dyer. *Invent. Math.*, 84(1):1–48, 1986.
- [17] J. S. Milne. *Étale Cohomology (PMS-33)*. Princeton University Press, 1980.
- [18] Paul Monsky. Formal cohomology III. *Annals of Mathematics*, 88(2):181–217, 1968.

- [19] Daqing Wan. Class numbers and p -ranks in \mathbb{Z}_p^d -towers. *J. Number Theory*, 203:139–154, 2019.
- [20] Lawrence C. Washington. *Introduction to cyclotomic fields*, volume 83 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1997.