

EXPONENTIAL SUMS AND RIGID COHOMOLOGY

PEIGEN LI

ABSTRACT. In this article, we prove a comparison theorem between the Dwork cohomology introduced by Adolphson and Sperber and the rigid cohomology. As a corollary, we can calculate the rigid cohomology of Dwork isocrystal on torus.

1. INTRODUCTION

Let p be a prime number, \mathbf{Q}_p the field of p -adic numbers, $k = \mathbf{F}_q$ the finite field with $q = p^a$ elements. Let k_i be the extension of k of degree i and let \bar{k} be the algebraic closure of k . Fix a primitive p -th root of unity ζ_p . Let K_0 be the unramified extension of $\mathbf{Q}_p(\zeta_p)$ of degree a . Let Ω be the completion of an algebraic closure of K_0 . Denote by “ord” the additive valuation on Ω normalized by $\text{ord}(p) = 1$. The norm on Ω is given by $|u| = p^{-\text{ord}(u)}$ for any $u \in \Omega$.

For a morphism $f : X \rightarrow \mathbf{A}_k^1$ with X being a k -scheme of finite type of dimension n , and the nontrivial additive character $\psi : k \rightarrow K_0^\times$ defined by $\psi(t) = \zeta_p^{\text{Tr}_{k/\mathbf{F}_p}(t)}$, define exponential sums

$$S_i(X, f) = \sum_{x \in X(k_i)} \psi(\text{Tr}_{k_i/k}(f(x))).$$

The L -function is defined by

$$L(X, f, t) = \exp\left(\sum_{i=1}^{\infty} S_i(X, f)t^i/i\right).$$

According to [9, Theorem 6.3], we have

$$L(X, f, t) = \prod_{i=0}^{2n} \det(I - tF^* | H_{c,\text{rig}}^i(X/K_0, f^* \mathcal{L}_\psi))^{(-1)^{i+1}},$$

where \mathcal{L}_ψ is the Dwork F -isocrystal defined over \mathbf{A}_k^1 associated to ψ and F^* is the Frobenius endomorphism on the space $H_{c,\text{rig}}^i(X/K_0, f^* \mathcal{L}_\psi)$.

In [2, section 4], Baldassarri and Berthelot compare the Dwork cohomology and the rigid cohomology for singular hypersurfaces. We prove a similar comparison theorem for the complex introduced by Adolphson and Sperber in [1, section 2] to study the exponential sums on the torus \mathbf{T}_k^n . Suppose that f is defined by a Laurent polynomial

$$f(x_1, \dots, x_n) = \sum_{j=1}^N a_j x^{w_j} \in k[x_1, x_1^{-1}, \dots, x_n, x_n^{-1}].$$

Let $\Delta(f)$ be the Newton polyhedron at ∞ of f which is defined to be the convex hull in \mathbf{R}^n of the set $\{w_j\}_{j=1}^N \cup \{(0, \dots, 0)\}$ and let δ be the convex cone generated by $\{w_j\}_{j=1}^N$ in \mathbf{R}^n . Let $\text{Vol}(\Delta(f))$ be the volume of $\Delta(f)$ with respect to Lebesgue measure on \mathbf{R}^n . We say f is *nondegenerate with*

I would like to thank Hao Zhang and Professor Lei Fu for helpful discussions.

respect to $\Delta(f)$ if for any face σ of $\Delta(f)$ not containing the origin, the Laurent polynomials $\frac{\partial f_\sigma}{\partial x_i}$, $i = 1, \dots, n$ have no common zero in $(\bar{k}^\times)^n$, where $f_\sigma = \sum_{w_j \in \sigma} a_j x^{w_j}$. Define a weight function on $\delta \cap \mathbf{Z}^n$ by

$$w(u) := \inf \{c : u \in c\Delta(f), c \geq 0\}.$$

Note that there exists some $M \in \mathbf{Z}_{>0}$ such that $w(\delta \cap \mathbf{Z}^n) \subset \frac{1}{M}\mathbf{Z}_{\geq 0}$.

Consider the Artin-Hasse exponential series:

$$E(t) = \exp\left(\sum_{i=0}^{\infty} \frac{t^{p^i}}{p^i}\right).$$

By [8, Lemma 4.1], the series $\sum_{i=0}^{\infty} \frac{t^{p^i}}{p^i}$ has a zero at $\gamma \in \mathbf{Q}_p(\zeta_p)$ such that $\text{ord } \gamma = 1/(p-1)$ and $\zeta_p \equiv 1 + \gamma \pmod{\gamma^2}$. Set

$$\theta(t) = E(\gamma t) = \sum_{i \geq 0} \lambda_i t^i.$$

The series $\theta(t)$ is a splitting function in Dwork's terminology [8, §4a]. In particular, we have $\text{ord } \lambda_i \geq i/(p-1)$ and $\theta(1) = \zeta_p$.

Fix an M -th root $\tilde{\gamma}$ of γ in Ω . Let $K = K_0(\tilde{\gamma})$, and \mathcal{O}_K the ring of integers of K . Let $\hat{a}_j \in K$ be the Teichmüller lifting of a_j and set

$$\hat{f}(x) = \sum_{j=1}^N \hat{a}_j x^{w_j} \in K[x_1, x_1^{-1}, \dots, x_n, x_n^{-1}].$$

For any $b > 0, c \in \mathbf{R}$, consider the following spaces :

$$L(b, c) = \left\{ \sum_{u \in \delta \cap \mathbf{Z}^n} a_u x^u : a_u \in K, \text{ord}(a_u) \geq bw(u) + c \right\},$$

$$L(b) = \bigcup_{c \in \mathbf{R}} L(b, c), \quad L_b^\dagger = \bigcup_{b > 0} L(b).$$

Set $\gamma_l = \sum_{i=0}^l \gamma^{p^i}/p^i$, $h(t) = \sum_{l=0}^{\infty} \gamma_l t^{p^l}$. Then we have

$$\theta(t) = \exp\left(\sum_{l=0}^{+\infty} \frac{\gamma^{p^l} t^{p^l}}{p^l}\right) = \exp\left(\gamma t + \sum_{l=1}^{+\infty} (\gamma_l - \gamma_{l-1}) t^{p^l}\right) = \exp(h(t) - h(t^p)).$$

Define

$$H(x) = \sum_{j=1}^N h(\hat{a}_j x^{w_j}), \quad F_0(x) = \exp(H(x) - H(x^q)).$$

The estimate $\text{ord}(\lambda_i) \geq \frac{i}{p-1}$ implies that $H(x)$ and $F_0(x)$ are well defined as formal Laurent series. In fact, we have

$$H(x) \in L\left(\frac{1}{p-1}, 0\right), \quad F_0(x) \in L\left(\frac{p}{q(p-1)}, 0\right).$$

Define an operator ψ_q on formal Laurent series by

$$\psi_q\left(\sum_{u \in \mathbf{Z}^n} a_u x^u\right) = \sum_{u \in \mathbf{Z}^n} a_{qu} x^u.$$

Let $\alpha = \psi_q \circ F_0$. Formally, we have

$$\alpha = \psi_q \circ \exp(H(x) - H(x^q)) = \exp(-H(x)) \circ \psi_q \circ \exp(H(x)).$$

For $i = 1, \dots, n$, define operators

$$E_i = x_i \partial / \partial x_i, \quad H_i = E_i H, \quad \hat{D}_i = E_i + H_i = E_i + E_i H.$$

Formally, we have

$$\hat{D}_i = \exp(-H(x)) \circ E_i \circ \exp(H(x))$$

for $i = 1, \dots, n$. Note that α and \hat{D}_i operate on L_δ^\dagger . One can show that \hat{D}_i commute with one another. Denote by $K.(L_\delta^\dagger, \hat{\underline{D}})$ the Koszul complex on L_δ^\dagger associated to $\hat{D}_1, \dots, \hat{D}_n$. We have $\alpha \circ \hat{D}_i = q \hat{D}_i \circ \alpha$ for all i . This implies that α induces a chain map on $K.(L_\delta^\dagger, \hat{\underline{D}})$. Now we can give the main theorem in the present paper.

Theorem 1.1. *We have an isomorphism*

$$H_{n-i}(K.(L_\delta^\dagger, \hat{\underline{D}})) \xrightarrow{\sim} H_{\text{rig}}^i(\mathbf{T}_k^n / K, f^* \mathcal{L}_{-\psi})$$

for each i . The endomorphism α on the left corresponds to the endomorphism $F_* = q^n (F^*)^{-1}$ on the rigid cohomology. Furthermore, α is bijective on each homology group of $K.(L_\delta^\dagger, \hat{\underline{D}})$.

As a corollary, we prove the following theorem, which eliminates the condition $p \neq 2$ in [6].

Theorem 1.2. *Suppose that $f : \mathbf{T}_k^n \rightarrow \mathbf{A}_k^1$ is nondegenerate with respect to $\Delta(f)$ and that $\dim \Delta(f) = n$. Then*

- (i) $H_{c, \text{rig}}^i(\mathbf{T}_k^n, f^* \mathcal{L}_\psi) = H_{\text{rig}}^i(\mathbf{T}_k^n, f^* \mathcal{L}_{-\psi}) = 0$ if $i \neq n$.
- (ii) $\dim H_{c, \text{rig}}^n(\mathbf{T}_k^n, f^* \mathcal{L}_\psi) = \dim H_{\text{rig}}^n(\mathbf{T}_k^n, f^* \mathcal{L}_{-\psi}) = n! \text{Vol}(\Delta(f))$.

2. COMPARISON THEOREM

By [8, Lemma 4.1], there exists an element $\pi \in \mathbf{Q}_p(\zeta_p)$ such that $\pi^{p-1} + p = 0$ and

$$\zeta_p \equiv 1 + \pi \pmod{\pi^2}.$$

Hence $\text{ord}(\pi - \gamma) \geq \frac{2}{p-1}$ as $\zeta_p - 1 \equiv \pi \equiv \gamma \pmod{(\zeta_p - 1)^2}$. By [8, Lemma 4.1] and the definition of the weight function, we have

$$G(x) := \exp(\pi(\hat{f}(x) - \hat{f}(x^q))) \in L\left(\frac{p-1}{pq}, 0\right).$$

For $i = 1, \dots, n$, define operators

$$D_i = \exp(-\pi \hat{f}(x)) \circ E_i \circ \exp(\pi \hat{f}(x)).$$

We have the Koszul complex $K.(L_\delta^\dagger, \underline{D})$ and its endomorphism induced by $\alpha_1 = \exp(-\pi \hat{f}(x)) \circ \psi_q \circ \exp(\pi \hat{f}(x))$. Let $R = \exp(H(x) - \pi \hat{f}(x))$. We have

$$(2.1) \quad \alpha_1 = R \circ \alpha \circ R^{-1}, \quad D_i = R \circ \hat{D}_i \circ R^{-1}$$

for all i .

Proposition 2.1. (i) $R, R^{-1} \in L(b_0)$ where $b_0 = \min\left\{\frac{1}{p-1}, \frac{p-1}{p}\right\}$.

(ii) *The multiplication by R defines an isomorphism of Koszul complexes $\beta : K.(L_\delta^\dagger, \hat{\underline{D}}) \xrightarrow{\cdot R} K.(L_\delta^\dagger, \underline{D})$ and $\alpha_1 \circ \beta = \beta \circ \alpha$.*

Proof. (ii) follows from (i) and (2.1). Let $h_1(t) = \sum_{l=1}^{+\infty} \gamma_l t^{p^l}$. Then

$$R = \exp((\gamma - \pi)\hat{f}(x)) \prod_{j=1}^N \exp(h_1(\hat{a}_j x^{w_j})).$$

The estimate $\text{ord}(\gamma - \pi) \geq \frac{2}{p-1}$ implies that

$$\exp((\gamma - \pi)\hat{f}(x)) = \prod_{j=1}^N \exp((\gamma - \pi)\hat{a}_j x^{w_j}) \in L\left(\frac{1}{p-1}, 0\right).$$

Note that $\exp(\gamma_l t^{p^l}) = \sum_{k=0}^{+\infty} \frac{\gamma_l^k}{k!} t^{k p^l}$ and

$$\text{ord}(\gamma_l) = \text{ord}\left(-\sum_{i=l+1}^{+\infty} \gamma^{p^i}/p^i\right) \geq \frac{p^{l+1}}{p-1} - l - 1.$$

Then

$$\text{ord}\left(\frac{\gamma_l^k}{k!}\right) \geq k\left(\frac{p^{l+1}}{p-1} - l - 1 - \frac{1}{p-1}\right) \geq \frac{p-1}{p} \cdot k p^l$$

for each $l \geq 1$. As the term $c_i t^i$ in the expansion for $\exp(h_1(t)) = \prod_{l=1}^{+\infty} \exp(\gamma_l t^{p^l})$ only depends on finitely many $\exp(\gamma_l t^{p^l})$, we have

$$\text{ord}(c_i) \geq \frac{p-1}{p}, \text{ and } \prod_{j=1}^N \exp(h_1(\hat{a}_j x^{w_j})) \in L\left(\frac{p-1}{p}, 0\right).$$

Hence $R \in L\left(\frac{1}{p-1}\right)L\left(\frac{p-1}{p}\right) \subset L(b_0)$. The same is true for R^{-1} . \square

Consider the twisted de Rham complex $C^\bullet(L_\delta^\dagger, \underline{D})$ defined by

$$C^k(L_\delta^\dagger, \underline{D}) = \bigoplus_{1 \leq i_1 < \dots < i_k \leq n} L_\delta^\dagger \frac{dx_{i_1}}{x_{i_1}} \wedge \dots \wedge \frac{dx_{i_k}}{x_{i_k}} \cong (L_\delta^\dagger)^{\binom{n}{k}}$$

with the differential $d : C^k(L_\delta^\dagger, \underline{D}) \rightarrow C^{k+1}(L_\delta^\dagger, \underline{D})$ given by

$$d(\omega) = \exp(-\pi\hat{f}) \circ d_x \circ \exp(\pi\hat{f})(\omega)$$

for any $\omega \in C^k(L_\delta^\dagger, \underline{D})$. Define an endomorphism $\alpha_1^{(k)}$ of $C^k(L_\delta^\dagger, \underline{D})$ by

$$\alpha_1^{(k)} = \bigoplus_{1 \leq i_1 < \dots < i_k \leq n} q^{n-k} \alpha_1.$$

The equalities $\alpha_1 \circ D_i = q D_i \circ \alpha_1$ imply that α_1 induces a chain map on $C^\bullet(L_\delta^\dagger, \underline{D})$. For each $0 \leq k \leq n$, we have a natural isomorphism

$$(2.2) \quad H^k(C^\bullet(L_\delta^\dagger, \underline{D})) \simeq H_{n-k}(K.(L_\delta^\dagger, \underline{D}))$$

which is compatible with their endomorphisms α_1 .

Let Σ be the fan of all faces of the dual cone of δ . Let X be the toric scheme over $R = \mathcal{O}_K$ associated to Σ . By the definition of Σ , we have $X = \text{Spec}(A)$ with $A = R[x^{\delta \cap \mathbf{Z}^n}]$. Set

$$\mathcal{S} = \left\{ \sum_{j=1}^N r_j w_j : 0 \leq r_1, \dots, r_N \leq 1, \sum_{j=1}^N r_j w_j \in \mathbf{Z}^n \right\}.$$

Since \mathcal{S} is discrete and bounded, it is finite. Set $\mathcal{S} = \{s_1, \dots, s_L\}$. Then A is generated by $\{x^{s_i} \mid s_i \in \mathcal{S}\}$ as an R -algebra. More precisely, we have $A \simeq R[y_1, y_2, \dots, y_L]/I$ by [7, Proposition 1.1.9], where

$$I = \left\{ \sum_{a,b} \left(y_1^{a_1} \cdots y_L^{a_L} - y_1^{b_1} \cdots y_L^{b_L} \right) : \sum_{i=1}^L a_i s_i = \sum_{i=1}^L b_i s_i, a_i, b_i \in \mathbf{Z}_{\geq 0} \right\}.$$

Consider the canonical immersions $X \rightarrow \mathbf{A}_R^L \rightarrow \mathbf{P}_R^L$. Let \overline{X} be the closure of X in \mathbf{P}_R^L . Let $\widehat{\overline{X}}$ be the formal completion of \overline{X} with respect to the p -adic topology. Denote by X_k, X_K the special fiber and the generic fiber of X over R , respectively. Denote by X_K^{an} the analytic space associated to X_K . By [4, 1.2.4 (ii)], X_K^{an} is a strict neighborhood of $]X_k[_{\widehat{\overline{X}}}$ in $]\overline{X}_k[_{\widehat{\overline{X}}}$. Let V be a strict neighborhood of $]X_k[_{\widehat{\overline{X}}}$ in $]\overline{X}_k[_{\widehat{\overline{X}}}$ and E a sheaf of \mathcal{O}_V -modules. Define

$$j^\dagger E := \varinjlim_{V'} j_{V'*} j_{V'}^* E,$$

where the inclusion $j_{V'} : V' \rightarrow V$ runs through the strict neighborhoods of $]X_k[_{\widehat{\overline{X}}}$ contained in V .

Lemma 2.2. *We have an isomorphism $L_\delta^\dagger \simeq \Gamma(X_K^{\text{an}}, j^\dagger \mathcal{O}_{X_K^{\text{an}}})$ and $H^i(X_K^{\text{an}}, j^\dagger(\mathcal{O}_{X_K^{\text{an}}})) = 0$ for all $i \geq 1$.*

Proof. By [4, Exemples 1.2.4(iii)], $V_\lambda := B^L(0, \lambda) \cap X_K^{\text{an}}$ form a cofinal system of strict neighborhoods of $]X_k[_{\widehat{\overline{X}}}$ in X_K^{an} when $\lambda \rightarrow 1^+$, where $B^L(0, \lambda)$ is the closed polydisc with radius λ of dimension L in $\mathbf{A}_K^{L, \text{an}}$. Let j_λ be the inclusion $V_\lambda \rightarrow X_K^{\text{an}}$ and fix a λ_0 . By [5, 1.2.5], we have

$$j^\dagger \mathcal{O}_{X_K^{\text{an}}} \simeq j_{\lambda_0*} j^\dagger \mathcal{O}_{V_{\lambda_0}} \simeq R j_{\lambda_0*} j^\dagger \mathcal{O}_{V_{\lambda_0}}.$$

Note that the inclusion $V_\lambda \rightarrow V_{\lambda_0}$ is affine and V_λ is affinoid for each $\lambda \leq \lambda_0$. We have

$$H^i(X_K^{\text{an}}, j^\dagger \mathcal{O}_{X_K^{\text{an}}}) \simeq H^i(V_{\lambda_0}, j^\dagger \mathcal{O}_{V_{\lambda_0}}) \simeq \varinjlim_{\lambda < \lambda_0} H^i(V_\lambda, \mathcal{O}_{V_\lambda}).$$

Hence the second assertion holds. Set

$$W_L = \bigcup_{\lambda > 1} \left\{ \sum_{v \in \mathbf{Z}_{\geq 0}^L} a_v y^v : a_v \in K, |a_v| \lambda^{|v|} \rightarrow 0 \text{ as } |v| \rightarrow +\infty \right\}.$$

We have $\Gamma(X_K^{\text{an}}, j^\dagger \mathcal{O}_{X_K^{\text{an}}}) = \varinjlim_{\lambda > 1} \Gamma(V_\lambda, \mathcal{O}_{V_\lambda}) \cong W_L / IW_L$. For any $u \in \mathbf{Z}^n \cap \delta$, denote the set of solutions $v_1 s_1 + \dots + v_L s_L = u$ in $\mathbf{Z}_{\geq 0}^L$ by S_u . Define a morphism $\chi : W_L \rightarrow L_\delta^\dagger$ by

$$\chi \left(\sum_{v \in \mathbf{Z}_{\geq 0}^L} a_v y^v \right) = \sum_{u \in \delta \cap \mathbf{Z}^n} \left(\sum_{v \in S_u} a_v \right) x^u.$$

Since the map $v \mapsto u = v_1 s_1 + \dots + v_L s_L$ is a linear continuous map, there exists some $B > 0$ such that $|u| \leq B|v|$. We have

$$\left| \sum_{v \in S_u} a_v \right| \lambda^{B^{-1}|u|} \leq \max_{v \in S_u} |a_v| \lambda^{|v|}$$

for any $\lambda > 1$. Thus χ is well defined and it induces a morphism $\bar{\chi} : W_L / IW_L \rightarrow L_\delta^\dagger$. It suffices to show $\bar{\chi}$ is an isomorphism.

Take $g(x) = \sum_{u \in \delta \cap \mathbf{Z}^n} b_u x^u \in L(b, c)$ for some $b > 0, c \in \mathbf{R}$. By the definition of S , for any nonzero element $u \in \delta \cap \mathbf{Z}^n$, we can find a subset $s(u) \subset \{1, \dots, L\}$ such that

$$u = \sum_{i \in s(u)} v_i s_i, \quad v_i \in \mathbf{Z}_{\geq 1}$$

and $w(s_i)^{-1} s_i, w(u)^{-1} u$ lie in the same codimension 1 face of $\Delta(f)$ for any $i \in s(u)$. By [1, Lemma 1.9 (c)] and the choice of M , we have

$$w(u) = \sum_{i \in s(u)} v_i w(s_i) \geq \frac{\sum_{i \in s(u)} |v_i|}{M}.$$

Set $y^{v(u)} = \prod_{i \in s(u)} y_i^{v_i}$ and $h(y) = \sum_u b_u y^{v(u)}$. We have

$$\text{ord}(b_u) \geq b w(u) + c \geq \frac{b}{M} |v(u)| + c.$$

So $h(y) \in W_L$ and $\chi(h(y)) = g(x)$. Hence $\bar{\chi}$ is surjective.

Take $f = \sum_v a_v y^v \in W_L$ such that $\text{ord}(a_v) \geq b|v| + c$ for some $b, c \in \mathbf{R}$ and $b > 0$. If $\chi(f) = 0$, we have

$$\sum_{v \in S_u} a_v = 0$$

for any u . For each $u \in \delta \cap \mathbf{Z}^n$, take some $v_u \in S_u$ such that $|v_u| = \min\{|v| | v \in S_u\}$. By Lemma 2.4 below, we can find $h_1, \dots, h_m \in I$ such that for any $v \in S_u$, there exist $g_{vu1}, \dots, g_{vum} \in R[y_1, \dots, y_L]$ such that

$$y^v - y^{v_u} = \sum_{j=1}^m g_{vu_j} h_j$$

and $\deg(g_{vu_j}) \leq |v|$ for all j . So

$$f = \sum_u \sum_{v \in S_u} a_v y^v = \sum_u \sum_{v \in S_u} a_v (y^v - y^{v_u}) = \sum_{j=1}^m h_j \left(\sum_u \sum_{v \in S_u} a_v g_{vu_j} \right).$$

Set $g_{vu_j} = \sum_k g_{vu_j}^k y^k$ with $|k| \leq |v|$ and $\text{ord}(g_{vu_j}^k) \geq 0$. Define

$$g_j := \sum_u \sum_{v \in S_u} a_v g_{vu_j} = \sum_k y^k \left(\sum_u \sum_{v \in S_u} a_v g_{vu_j}^k \right).$$

We have

$$\text{ord} \left(\sum_u \sum_{v \in S_u} a_v g_{vu_j}^k \right) \geq \min_{v \in S_u, u} \text{ord}(a_v g_{vu_j}^k) \geq \min_{v \in S_u, u} \text{ord}(a_v) \geq \min_{v \in S_u, u} b|v| + c \geq b|k| + c.$$

Hence $g_j \in W_L$ and $f \in IW_L$ which imply that $\bar{\chi}$ is injective. \square

Lemma 2.3. *For any subset J of $\mathbf{Z}_{\geq 0}^n$, there exists a finite subset J_0 of J such that $J \subset \bigcup_{v \in J_0} (v + \mathbf{Z}_{\geq 0}^n)$.*

Proof. We use induction on n . When $n = 1$, it is trivial. Suppose that our assertion holds for any subset of $\mathbf{Z}_{\geq 0}^{n-1}$. Let J be a subset of $\mathbf{Z}_{\geq 0}^n$. Take

$$J' = \{v \in \mathbf{Z}_{\geq 0}^n : (a, v) \in J \text{ for some } a \in \mathbf{Z}_{\geq 0}\}.$$

By the induction hypothesis, there exists a finite subset $J'_0 = \{v'_1, \dots, v'_r\}$ such that

$$J' \subset \bigcup_{i=1}^r (v'_i + \mathbf{Z}_{\geq 0}^n).$$

For each i , take some $a_i \in \mathbf{Z}_{\geq 0}$ such that $v_i = (a_i, v'_i) \in J$. For any $b \in \mathbf{Z}_{\geq 0}$, let

$$J_b = \{(c_1, \dots, c_{n+1}) \in J : c_1 = b\}.$$

Using the induction hypothesis again, there exists a finite subset $J_{b,0}$ such that

$$J_b \subset \bigcup_{v \in J_{b,0}} (v + \mathbf{Z}_{\geq 0}^{n+1}).$$

Set $a_0 = \max_{1 \leq i \leq r} \{a_i\}$. We have

$$\begin{aligned} J &\subset \left(\bigcup_{b < a_0} J_b \right) \cup \left(\bigcup_{i=1}^r (v_i + \mathbf{Z}_{\geq 0}^{n+1}) \right) \\ &\subset \left(\bigcup_{b < a_0} \bigcup_{v \in J_{b,0}} (v + \mathbf{Z}_{\geq 0}^{n+1}) \right) \cup \left(\bigcup_{i=1}^r (v_i + \mathbf{Z}_{\geq 0}^{n+1}) \right) \end{aligned}$$

We can take $J_0 = \left(\bigcup_{b < a_0} J_{b,0} \right) \cup \{v_1, \dots, v_r\}$. \square

Lemma 2.4. *There are finitely many elements $h_1, \dots, h_m \in I$ such that for any $y^a - y^b \in I$, there exist $g_1, \dots, g_m \in R[y_1, \dots, y_L]$ such that*

$$y^a - y^b = \sum_{i=1}^m g_i h_i$$

and $\deg(g_i) \leq \max\{|a|, |b|\}$.

Proof. Take $J = \{(a, b) : y^a - y^b \in I\} - \{(0, 0)\}$. By Lemma 2.3 above, there is a finite subset $J_0 = \{(a_1, b_1), \dots, (a_m, b_m)\} \subset J$ such that $J \subset \bigcup_{v \in J_0} (v + \mathbf{Z}_{\geq 0}^{2L})$. Take $h_i = y^{a_i} - y^{b_i}$. We prove the assertion by induction on $|a| + |b|$. When $|a| + |b| = 0$, the assertion is trivial. Take $(a, b) \in J$. Suppose that the assertion holds for any $y^{a'} - y^{b'} \in J$ with $|a'| + |b'| < |a| + |b|$. By the choice of J_0 , there is some i_0 such that $(a, b) \geq (a_{i_0}, b_{i_0})$. We have $|a - a_{i_0}| + |b - b_{i_0}| < |a| + |b|$. By induction hypothesis, there exist $g'_1, \dots, g'_m \in R[y_1, \dots, y_L]$ such that

$$y^{a-a_{i_0}} - y^{b-b_{i_0}} = \sum_{i=1}^m g'_i h_i$$

and $\deg(g'_i) \leq \max\{|a - a_{i_0}|, |b - b_{i_0}|\}$.

If $|a_{i_0}| \geq |b_{i_0}|$, we have

$$y^a - y^b = y^{a-a_{i_0}} (y^{a_{i_0}} - y^{b_{i_0}}) + y^{b_{i_0}} (y^{a-a_{i_0}} - y^{b-b_{i_0}}).$$

Take $g_{i_0} = y^{a-a_{i_0}} + y^{b_{i_0}} g'_{i_0}$ and $g_i = y^{b_{i_0}} g'_i$ for any $i \neq i_0$. We have

$$y^a - y^b = \sum_{i=1}^m g_i h_i$$

and $\deg(g_i) \leq \max\{|a - a_{i_0}|, |b_{i_0}| + \deg(g'_i)\} \leq \max\{|a|, |b|\}$.

If $|b_{i_0}| \geq |a_{i_0}|$, we have

$$y^a - y^b = y^{b-b_{i_0}} (y^{a_{i_0}} - y^{b_{i_0}}) + y^{a_{i_0}} (y^{a-a_{i_0}} - y^{b-b_{i_0}}).$$

Then we conclude as above. \square

Set

$$L_0^\dagger = \bigcup_{r>1} \left\{ \sum_{u \in \mathbf{Z}^n} a_u x^u : a_u \in K, |a_u| r^{|u|} \rightarrow 0 \text{ as } |u| \rightarrow +\infty \right\}.$$

Note that $L_\delta^\dagger \subset L_0^\dagger$. Similarly, we can define the complex $C^\cdot(L_0^\dagger, \underline{D})$ and the endomorphism α_1 . By [3, Proposition 3.4] or [6], we have

Proposition 2.5. *We have a canonical isomorphism*

$$R\Gamma_{\text{rig}}(\mathbf{T}_k^n/K, f^* \mathcal{L}_{-\psi}) \xrightarrow{\sim} C^\cdot(L_0^\dagger, \underline{D})$$

which is compatible with their endomorphisms $F_* = q^n(F^*)^{-1}$ and α_1 . Furthermore, α_1 is bijective on each cohomology group of $C^\cdot(L_0^\dagger, \underline{D})$.

Let Σ' be a regular refinement of Σ and let X' be the toric scheme over R associated to Σ' . Denote by X'_k, X'_K the special fiber and the generic fiber of X' over R , respectively. Let $\mathbf{T}_R^n \rightarrow X'$ be the immersion of the maximal open dense torus, and let $D' = X' - \mathbf{T}_R^n$. Denote by X'_K^{an} (resp. D'_K^{an}) the analytic space associated to X'_K (resp. D'_K). Note that X' is smooth and D' is a normal crossing divisor in X' . Let $\Omega_{X'}^k(\log D')$ be the sheaf of differential forms of degree k with logarithmic poles along D' . Then $\Omega_{X_K^{\text{an}}}^k(\log D_K^{\text{an}})$ is a free $\mathcal{O}_{X_K^{\text{an}}}$ module with basis $\frac{dx_{i_1}}{x_{i_1}} \wedge \cdots \wedge \frac{dx_{i_k}}{x_{i_k}}$, where $1 \leq i_1 < \cdots < i_k \leq n$ and x_1, \dots, x_n are coordinates of \mathbf{T}_K^{an} . Consider the complex

$$E := (\Omega_{X_K^{\text{an}}}^k(\log D_K^{\text{an}}), d)$$

where the differential is given by $d = d_x + \sum_{i=1}^n \pi E_i \hat{f} \frac{dx_i}{x_i}$. Note that

$$\hat{f} \in \Gamma(X_K^{\text{an}}, \mathcal{O}_{X_K^{\text{an}}}) \cong \Gamma(X'_K^{\text{an}}, \mathcal{O}_{X'_K^{\text{an}}}).$$

Hence the twist de Rham complex E is well defined. Let Σ'' be a regular fan containing Σ' such that $|\Sigma''| = \mathbf{R}^n$, let X'' be the toric scheme over R associated to Σ'' and let \widehat{X}'' be its formal completion. In [2, Corollary A.4], taking $\mathcal{X} = \widehat{X}''$, $Z = X'' - \mathbf{T}_R^n$, $V = X'_k$, $U = \mathbf{T}_k^n$ and $W = X'_K^{\text{an}}$, we get a canonical isomorphism

$$(2.3) \quad R\Gamma(X'_K^{\text{an}}, j^\dagger \Omega_{X'_K^{\text{an}}}(\log D_K^{\text{an}})) \cong R\Gamma_{\text{rig}}(\mathbf{T}_k^n, \mathcal{L}_{-\psi, f}).$$

Now we can give the proof of **Theorem 1.1**.

Proof. Take cones $\sigma_1, \dots, \sigma_m \in \Sigma'$ such that the open sets U_i defined by σ_i form an affine open covering of X' . Let $j_{\underline{i}} := j_{i_1 \dots i_k}$ be the inclusion $U_{\underline{i}} := U_{i_1} \cap \cdots \cap U_{i_k} \subset X'$. Note that $U_{\underline{i}}$ is the affine open set defined by the cone $\sigma_{\underline{i}} := \sigma_{i_1} \cap \cdots \cap \sigma_{i_k}$. By [4, Proposition 2.1.8], we have an exact sequence

$$0 \rightarrow j^\dagger E \rightarrow \prod_i j_i^\dagger E \rightarrow \prod_{i_1 < i_2} j_{i_1 i_2}^\dagger E \rightarrow \cdots \rightarrow j_{1 \dots m}^\dagger E \rightarrow 0,$$

where $j_{\underline{i}}^\dagger E = \varinjlim_V j_{V*} j_V^*(E)$ and $j_V : V \rightarrow X'_K^{\text{an}}$ runs through strict neighborhoods of $]U_{\underline{i}, k}[_{\widehat{X}''}$ contained in X'_K^{an} . We still denote by $j_{\underline{i}}$ the inclusion $U_{\underline{i}, K}^{\text{an}} \subset X'_K^{\text{an}}$. We have $j_{\underline{i}}^\dagger E = j_{\underline{i}*} j_{\underline{i}}^\dagger(E|_{U_{\underline{i}, K}^{\text{an}}})$ by [4, 1.2.4 (ii) and 2.1.1]. Let $\check{\sigma}_{\underline{i}}$ be the dual of $\sigma_{\underline{i}}$ and $\sigma = \check{\delta}$. Applying Lemma 2.2 to each $U_{\underline{i}, K}^{\text{an}}$, we know that $R\Gamma(X'_K^{\text{an}}, j^\dagger E)$ can be represented by the total complex of

$$(2.4) \quad 0 \rightarrow \prod_i \Gamma(U_{i, K}^{\text{an}}, j^\dagger((E|_{U_{i, K}^{\text{an}}})) \rightarrow \cdots \rightarrow \Gamma(U_{1 \dots n, K}^{\text{an}}, j^\dagger((E|_{U_{1 \dots n, K}^{\text{an}}})) \rightarrow 0$$

and $\Gamma(U_{\underline{i},K}^{\text{an}}, j^\dagger \mathcal{O}_{U_{\underline{i},K}^{\text{an}}}) \cong L_{\sigma_{\underline{i}}}^\dagger$, where

$$L_{\sigma_{\underline{i}}}^\dagger = \bigcup_{\lambda > 1} \left\{ \sum_{u \in \sigma_{\underline{i}} \cap \mathbf{Z}^n} A_u x^u : A_u \in K, |A_u| \lambda^{|u|} \xrightarrow{|u| \rightarrow +\infty} 0 \right\}$$

which can be regarded as a subalgebra of L_0^\dagger . Thus (2.4) can be represented by the bicomplex

$$(2.5) \quad 0 \rightrightarrows \prod_i C^\cdot(L_{\sigma_i}^\dagger) \rightrightarrows \prod_{i_1 < i_2} C^\cdot(L_{\sigma_{\underline{i}}}^\dagger) \rightrightarrows \cdots \rightrightarrows C^\cdot(L_{\sigma_{i_1 \dots m}}^\dagger) \rightrightarrows 0,$$

where all $C^\cdot(L_{\sigma_{\underline{i}}}^\dagger)$ are regarded as subcomplexes of $C^\cdot(L_0^\dagger, \underline{D})$. By Lemma 2.7 below and a spectral sequence argument, $C^\cdot(L_\delta^\dagger, \underline{D})$ can be represented by the total complex of (2.5). Hence

$$R\Gamma(X_K^{\text{an}}, j^\dagger E) \cong C^\cdot(L_\delta^\dagger, \underline{D}).$$

By Proposition 2.5, the isomorphism (2.3) can be represented by the inclusion

$$C^\cdot(L_\delta^\dagger, \underline{D}) \hookrightarrow C^\cdot(L_0^\dagger, \underline{D}).$$

By (2.2) and Proposition 2.1, we have

$$H_{n-i}(K, (L_\delta^\dagger, \hat{\underline{D}})) \xrightarrow{\sim} H_{\text{rig}}^i(\mathbf{T}_k^n/K, f^* \mathcal{L}_{-\psi})$$

for each i . The second assertion of Theorem 1.1 follows from (2.2) and Proposition 2.5. \square

Lemma 2.6. *Notation as above, we have an exact sequence*

$$(2.6) \quad 0 \rightrightarrows L_{\check{\sigma}, R} \rightrightarrows \prod_i L_{\check{\sigma}_i, R} \rightrightarrows \prod_{i_1 < i_2} L_{\check{\sigma}_{\underline{i}}, R} \rightrightarrows \cdots \rightrightarrows L_{\check{\sigma}_{i_1 \dots m}, R} \rightrightarrows 0,$$

where $L_{\check{\tau}, R} = R[x^{\check{\tau} \cap \mathbf{Z}^n}]$ is regarded as a subalgebra of $R[x^{\mathbf{Z}^n}]$, for $\tau = \sigma, \sigma_{\underline{i}}, \dots$. The boundary map $\prod_{i_1 < \dots < i_k} L_{\check{\sigma}_{\underline{i}}, R} \xrightarrow{d} \prod_{i_1 < \dots < i_{k+1}} L_{\check{\sigma}_{\underline{i}}, R}$ is given by

$$(dg)_{i_1 \dots i_{k+1}} = \sum_{s=1}^{k+1} (-1)^{s-1} g_{i_1 \dots \hat{i}_s \dots i_{k+1}}$$

for any $g \in \prod_{i_1 < \dots < i_k} L_{\check{\sigma}_{\underline{i}}, R}$.

Proof. Consider the Čech complex $C^\cdot(\mathfrak{U}, \mathcal{O}_{X'})$ defined by the affine open covering $\mathfrak{U} = \{U_{\sigma_i}\}_{i=1}^m$. We have $\check{H}^k(\mathfrak{U}, \mathcal{O}_{X'}) \cong H^k(X', \mathcal{O}_{X'})$ for any k . Since $\phi : X' \rightarrow X$ is proper and X is affine, we have $R^k \phi_* \mathcal{O}_{X'} = H^k(X', \mathcal{O}_{X'})^\sim$. By [7, Proposition 9.2.5], we have $R^k \phi_* \mathcal{O}_{X'} = 0$ for any $k \geq 1$. Thus (2.6) is exact except at the second position. Since $\sigma = \cup_{i=1}^m \sigma_i$, we have $\check{\sigma} = \cap_{i=1}^m \check{\sigma}_i$. Thus $L_{\check{\sigma}, R} = \text{Ker}(\prod_i L_{\check{\sigma}_i, R} \rightarrow \prod_{i_1 < i_2} L_{\check{\sigma}_{\underline{i}}, R})$ and then (2.6) is exact. \square

Lemma 2.7. *Notation as above, we have an exact sequence*

$$(2.7) \quad 0 \rightrightarrows L_\delta^\dagger \rightrightarrows \prod_i L_{\sigma_i}^\dagger \rightrightarrows \prod_{i_1 < i_2} L_{\sigma_{\underline{i}}}^\dagger \rightrightarrows \cdots \rightrightarrows L_{\sigma_{i_1 \dots m}}^\dagger \rightrightarrows 0,$$

where the boundary map is the same as (2.6).

Proof. By the definition of Σ' , we have $\sigma = \cup_{i=1}^m \sigma_i$. Thus $\delta = \check{\sigma} = \cap_{i=1}^m \check{\sigma}_i$ and $L_{\check{\sigma}}^\dagger = \text{Ker}(\prod_i L_{\check{\sigma}_i}^\dagger \rightarrow \prod_{i_1 < i_2} L_{\check{\sigma}_{i_1}}^\dagger)$. Now consider $\prod_{i_1 < \dots < i_k} L_{\check{\sigma}_{i_1}}^\dagger \xrightarrow{d} \prod_{i_1 < \dots < i_{k+1}} L_{\check{\sigma}_{i_1}}^\dagger$, take $g = (g_{\underline{i}}) \in \prod_{i_1 < \dots < i_k} L_{\check{\sigma}_{i_1}}^\dagger$. Suppose that for all \underline{i} , $g_{\underline{i}} \in L''(b, c)$ for some $b > 0$, $c \in \mathbf{R}$, where

$$L''(b, c) = \left\{ \sum_u A_u x^u : A_u \in K, \text{ord } A_u \geq b|u| + c \right\}.$$

Since we always can choose some element $a \in R$ such that $ag_i \in L''(b, 0)$ for all i , we treat the case $c = 0$. Now $g_{\underline{i}}$ can be written as

$$g_{\underline{i}} = \sum_{k=0}^{+\infty} \pi^k g_{\underline{i}, k}, \quad g_{\underline{i}, k} \in R[x^{\check{\sigma}_{\underline{i}} \cap \mathbf{Z}^n}],$$

where $g_{\underline{i}, k}$ is a sum of monomials of degree lying in $[\frac{k}{b(p-1)}, \frac{k+1}{b(p-1)})$. Note that the degree of polynomial defines a grading on the complex (2.6), and d preserves the grading. Write $g = \sum_{k=0}^{+\infty} \pi^k g_k$, where $g_k = (g_{\underline{i}, k})$. If $d(g) = 0$, we have $d(g_k) = 0$ for each k . By Lemma 2.6, there exists

$$h_k = (h_{\underline{i}, k})_{\underline{i}} \in \prod_{i_1 < \dots < i_{k-1}} L_{\check{\sigma}_{i_1}, R}$$

such that $d(h_k) = g_k$ for each k and we may assume that each $h_{\underline{i}, k}$ is a sum of monomials of degree lying in $[\frac{k}{b(p-1)}, \frac{k+1}{b(p-1)})$. Suppose that $h_{\underline{i}, k} = \sum_u A_{\underline{i}, k, u} x^u$, we have

$$\text{ord}(\pi^k A_{\underline{i}, k, u}) \geq \frac{k}{p-1} > \frac{k}{(p-1)\frac{k+1}{b(p-1)}}|u| \geq \frac{b}{2}|u|$$

for any $k \geq 1$. Thus $\sum_{k=1}^{+\infty} \pi^k h_{\underline{i}, k} \in L''(\frac{b}{2}, 0)$. Note that $h_{\underline{i}, 0}$ is a polynomial for each \underline{i} . We have

$$h := \sum_{k=0}^{+\infty} \pi^k h_k \in \prod_{i_1 < \dots < i_{k-1}} L_{\check{\sigma}_{i_1}}^\dagger$$

and $d(h) = g$. Hence (2.7) is exact. \square

3. APPLICATIONS

Define

$$B = \left\{ \sum_{u \in \delta \cap \mathbf{Z}^n} a_u \tilde{\gamma}^{Mw(u)} x^u : a_u \in K, a_u \rightarrow 0 \text{ as } w(u) \rightarrow +\infty \right\}.$$

We can define the Koszul complexes $K.(L(b), \hat{\underline{D}})$ ($0 < b \leq \frac{p}{p-1}$) and $K.(B, \hat{\underline{D}})$ with actions of α , respectively. They can be regarded as subcomplexes of $K.(L_\delta^\dagger, \hat{\underline{D}})$.

Proposition 3.1. *For each i , the inclusions $B \hookrightarrow L(\frac{1}{p-1}) \hookrightarrow L_\delta^\dagger$ induce a surjective homomorphism $H_i(K.(B, \hat{\underline{D}})) \rightarrow H_i(K.(L_\delta^\dagger, \hat{\underline{D}}))$.*

Proof. We use the same method as [6, Corollaire 1.3]. The inclusions $L(\frac{p}{p-1}) \hookrightarrow B \hookrightarrow L_\delta^\dagger$ imply that it suffices to prove the homomorphism $H_i(K.(L(\frac{p}{p-1}), \hat{\underline{D}})) \rightarrow H_i(K.(L_\delta^\dagger, \hat{\underline{D}}))$ is surjective. Note that $L_\delta^\dagger = \bigcup_{0 < b \leq \frac{p}{p-1}} L(b)$. By Theorem 1.1 and [5, 3.10], $H_i(K.(L_\delta^\dagger, \hat{\underline{D}}))$ is finite dimensional.

Thus there exists some $0 < b \leq \frac{p}{p-1}$ such that $H_i(K.(L(b), \hat{D})) \rightarrow H_i(K.(L_\delta^\dagger, \hat{D}))$ is surjective. Note that $\psi_q(L(b)) \subset L(qb)$. Consider the following composite

$$L(b) \xrightarrow{\cdot F_0} L\left(\min\left\{\frac{p}{q(p-1)}, b\right\}\right) \xrightarrow{\psi_q} L\left(\min\left\{\frac{p}{p-1}, qb\right\}\right).$$

Hence

$$\alpha(L(b)) \subset L\left(\min\left\{\frac{p}{p-1}, qb\right\}\right).$$

So there exists a positive integer m such that

$$\alpha^m(H_i(K.(L(b), \hat{D}))) \subset H_i\left(K.\left(L\left(\frac{p}{p-1}\right), \hat{D}\right)\right).$$

We have a commutative diagram

$$\begin{array}{ccc} H_i(K.(L(b), \hat{D})) & \longrightarrow & H_i(K.(L_\delta^\dagger, \hat{D})) \\ \downarrow \alpha^m & & \downarrow \alpha^m \\ H_i(K.(L(\frac{p}{p-1}), \hat{D})) & \longrightarrow & H_i(K.(L_\delta^\dagger, \hat{D})). \end{array}$$

The top horizontal arrow is surjective by the choice of b . The right vertical arrow α^m is bijective by Theorem 1.1. So the bottom horizontal arrow is surjective. \square

Let $X = \mathbf{T}_k^r \times \mathbf{A}_k^{n-r}$ and

$$f \in k[x_1, \dots, x_n, (x_1 \cdots x_r)^{-1}].$$

For any subset $A \subset S_r = \{r+1, \dots, n\}$, let

$$\mathbf{R}_A^n = \{(x_1, \dots, x_n) : x_i = 0 \text{ for } i \in A\}.$$

Let $V_A(f)$ be the volume of $\Delta(f) \cap \mathbf{R}_A^n$ with respect to Lebesgue measure on \mathbf{R}_A^n . Set

$$v_A(f) = \sum_{B \subset A} (-1)^{|B|} (n - |B|)! V_B(f)$$

and $\tilde{r} = \dim \Delta(f_{S_r})$. We say that f is *commode with respect to S_r* if for all subsets $A \subset S_r$, we have $\dim \Delta(f_A) = \tilde{r} + |S_r - A|$, where $f_A := \prod_{j \in A} \theta_j \circ f(x)$ and θ_j is the map such that $x_j \mapsto 0$.

Theorem 3.2. *Let $X = \mathbf{T}_k^r \times \mathbf{A}_k^{n-r}$, $f \in \Gamma(X, \mathcal{O}_X)$. Suppose that f is commode with respect to S_r , nondegenerate with respect to $\Delta(f)$ and $\dim \Delta(f) = n$. Then*

$$(i) H_{c, \text{rig}}^i(X, f^* \mathcal{L}_\psi) = H_{\text{rig}}^i(X, f^* \mathcal{L}_{-\psi}) = 0 \text{ if } i \neq n.$$

$$(ii) \dim H_{c, \text{rig}}^n(X, f^* \mathcal{L}_\psi) = \dim H_{\text{rig}}^n(X, f^* \mathcal{L}_{-\psi}) = v_{S_r}(f).$$

Proof. First we consider the case $r = n$. Since X is affine, smooth and pure of dimension n , by the Poincaré duality and the trace formula in the rigid cohomology theory, we have

$$(3.1) \quad L(\mathbf{T}_k^n, f, t) = \prod_{i=0}^n \det(I - tq^n(F^*)^{-1} | H_{\text{rig}}^i(\mathbf{T}_k^n/K, f^* \mathcal{L}_{-\psi})^{(-1)^{i+1}}).$$

When f is nondegenerate, we have $H_i(K.(B, \hat{D})) = 0$ for $i \neq 0$ and $\dim H_0(K.(B, \hat{D})) = n! \text{Vol}(\Delta(f))$ by [1, Theorem 2.18]. By Proposition 3.1, we have $H_i(K.(L_\delta^\dagger, \hat{D})) = 0$ for $i \neq 0$. By Theorem 1.1, we have

$$H_{\text{rig}}^i(\mathbf{T}_k^n, f^* \mathcal{L}_{-\psi}) \cong H_{n-i}(K.(L_\delta^\dagger, \hat{D})) = 0$$

for $i \neq n$. By [1, (2.5)] and (3.1), we have

$$\det(I - tq^n(F^*)^{-1}|H_{\text{rig}}^n(X/K, f^*\mathcal{L}_{-\psi})) = \det(1 - tH_0(\alpha)|H_0(K.(B, \hat{D}))).$$

So

$$\dim H_{\text{rig}}^n(X/K, f^*\mathcal{L}_{-\psi}) = \dim H_0(K.(B, \hat{D})) = n! \text{Vol}(\Delta(f)).$$

We then use induction on $n - r$. Suppose that Theorem 3.2 holds for $n - (r + 1)$. By [5, 2.3.1] and [10, Theorem 4.1.1], we have a distinguished triangle

$$R\Gamma_{\text{rig}}(X, \mathcal{L}_{-\psi, f}) \rightarrow R\Gamma_{\text{rig}}(U, \mathcal{L}_{-\psi, f}) \rightarrow R\Gamma_{\text{rig}}(Z, \mathcal{L}_{-\psi, f_{r+1}})[-1] \rightarrow$$

where $U = X - V(x_{r+1}) \cong \mathbf{T}_k^{r+1} \times \mathbf{A}_k^{n-r-1}$ and $Z = V(x_{r+1}) \cong \mathbf{T}_k^r \times \mathbf{A}_k^{n-r-1}$. One can check that f, f_{r+1} are still commode with respect to S_{r+1} , and f_{r+1} is nondegenerate with respect to $\Delta(f_{r+1})$. By the induction hypothesis, (i) holds and

$$\begin{aligned} \dim H_{\text{rig}}^n(X, \mathcal{L}_{-\psi, f}) &= \dim H_{\text{rig}}^n(U, \mathcal{L}_{-\psi, f}) - \dim H_{\text{rig}}^{n-1}(Z, \mathcal{L}_{-\psi, f_{r+1}}) \\ &= v_{S_{r+1}}(f) - v_{S_{r+1}}(f_{r+1}) = v_{S_r}(f). \end{aligned}$$

□

REFERENCES

- [1] A. Adolphson and S. Sperber, Exponential sums and Newton polyhedra: Cohomology and estimates, *Ann. of Math.* 130 (1989), 367-406.
- [2] F. Baldassarri, P. Berthelot, On Dwork cohomology for singular hypersurfaces. *Geometric Aspects of Dwork's Theory*, 177-244, Proceedings of the Dwork Trimester in Italy, May-July 2001, F. Baldassarri, et al. (eds), de Gruyter, 2004.
- [3] P. Berthelot, Cohomologie rigide et théorie de Dwork: le cas des sommes exponentielles, *Astérisque* 119 (1984), 17-49.
- [4] P. Berthelot, Cohomologie rigide et cohomologie rigide à supports propres. Première Partie, prépublication, Université de Rennes (1996).
- [5] P. Berthelot, Finitude et pureté cohomologique en cohomologie rigide avec un appendice par Aise Johan de Jong, *Invent math* 128 (1997), 329-377.
- [6] P. Bourgeois, Annulation et pureté des groupes de cohomologie rigide associés à des sommes exponentielles, *C. R. Acad. Sci. Paris* 328(1999), 681-686.
- [7] D. Cox, J. Little and H. Schenck, *Toric Varieties*, Graduate Study in Mathematics 124, Amer. Math. Soc., 2011.
- [8] B. Dwork, On the zeta function of a hypersurface, *Publ. Math. IHÉS* 12 (1962), 5-68.
- [9] J.-Y. Étesse and B. Le Stum, Fonctions L associées aux F-isocristaux surconvergents, *Math. Ann.* 296 (1993), 557-576.
- [10] N. Tsuzuki, On the Gysin isomorphism of rigid cohomology, *Hiroshima Math. J.* 29 (1999), 479-527.

YAU MATHEMATICAL SCIENCES CENTER, TSINGHUA UNIVERSITY, BEIJING 100084, P. R. CHINA
Email address: lpg16@mails.tsinghua.edu.cn