

STATISTICS FOR p -RANKS OF ARTIN-SCHREIER COVERS

ANWESH RAY

ABSTRACT. Given a prime p and q a power of p , we study the statistics of p -ranks of Artin-Schreier covers of given genus defined over \mathbb{F}_q , in the large q -limit. We refer to this problem as the *geometric problem*. We also study an *arithmetic* variation of this problem, and consider Artin-Schreier covers defined over \mathbb{F}_p , letting p go to infinity. Distribution of p -ranks has been previously studied for a Artin-Schreier covers over a fixed finite field as the genus is allowed to go to infinity. The method requires that we count isomorphism classes of covers that are unramified at ∞ .

1. INTRODUCTION

The study of curves over finite fields leads to many interesting problems in arithmetic statistics. Recently, statistical questions have been framed and studied for curves varying in certain naturally occurring ensembles, see for instance, [9, 7, 12, 2, 11, 4, 1, 3, 10]. Throughout, p will denote a prime number and q will be a power of p . The field with q elements is denoted \mathbb{F}_q . Given a curve \mathcal{C} of positive genus g over \mathbb{F}_q , its arithmetic is better understood by studying the structure of its Jacobian $\text{Jac}(\mathcal{C})$, which is a g -dimensional abelian variety. An important invariant associated to the curve is the p -rank of $\text{Jac}(\mathcal{C})(\bar{\mathbb{F}}_p)$, which we denote by τ . This is the number such that $\text{Jac}(\mathcal{C})(\bar{\mathbb{F}}_p) \otimes_{\mathbb{Z}} (\mathbb{Z}/p\mathbb{Z})$ is isomorphic to $(\mathbb{Z}/p\mathbb{Z})^{\tau}$. The p -rank lies in the range $0 \leq \tau \leq g$ and when $\tau = g$, the curve is *ordinary*. In [3, 10], statistical questions are studied for p -ranks of Artin-Schreier covers of \mathbb{P}^1 defined over a fixed finite field \mathbb{F}_q , as the genus goes to infinity. It is shown that when p is odd, the proportion of Artin-Schreier covers that are ordinary is zero as $g \rightarrow \infty$. When $p = 2$, this proportion is shown to be non-zero, see [10, Theorem 1.4].

We study similar questions about the distributions of p -ranks, however, the curves will no longer be defined over a fixed finite field. There are two independent questions we study in this paper. For the former, we fix the prime p and let $q \rightarrow \infty$. This is referred to as the *geometric problem* since the results may (in spirit) be interpreted in terms of the distribution of points in moduli spaces. Such moduli spaces for Artin-Schreier curves of genus $g = d \left(\frac{p-1}{2} \right)$ have been constructed by Pries and Zhu, see [8]. The various irreducible components in these moduli spaces are classified by partitions of $d + 2$. It is no surprise that the densities that are obtained in this paper are also expressed in terms of partitions. In fact, our results indicate that the points on these moduli spaces are asymptotically equally distributed on the various components of maximal dimension. The number of such components of maximal dimension with p -rank $\tau = r(p-1)$ can be expressed in terms of partitions of $d+2$ into $r+1$ numbers satisfying additional conditions, see [8, Theorem 1.1, Corollary 3.11]. Since the geometric structure of the various strata

2010 *Mathematics Subject Classification*. 11G20, 11T06, 11T55, 14G17, 14H25.

Key words and phrases. Arithmetic statistics, curves over finite fields, Artin-Schreier covers.

in these irreducible components is complicated, it does not seem easy to obtain such results directly from the moduli-space description. The method used to obtain results in this paper are combinatorial and are obtained independent of the results of Pries-Zhu.

We make certain simplifications to the problem in question:

- the curves are not classified up to isomorphism, however, up to *isomorphism of covers*, see Definition 2.1.
- We restrict to isomorphism classes of covers that are unramified at ∞ .

In studying the arithmetic variant, the prime p itself is allowed to vary. By a standard application of the Riemann-Hurwitz formula, the genus g of an Artin-Schreier cover is always divisible by $\binom{p-1}{2}$. Since we are interested in the case when the genus is positive it follows that $g \rightarrow \infty$ as $p \rightarrow \infty$. In this setting, we fix the *speed* at which the genus grows. Hence, we fix an integer $d > 0$ throughout, and let $g(d, p) := d \binom{p-1}{2}$. Likewise, fix $r \geq 0$ and let $\tau(r, p) = r(p-1)$. The p -rank will thus also grow at a fixed linear rate.

Thus, the two main questions studied are as follows.

- (1) *Geometric problem:* Given a prime p , what is the statistics for p -ranks of Artin-Schreier covers with fixed genus over \mathbb{F}_q , as $q \rightarrow \infty$?
- (2) *Arithmetic problem:* Suppose we are given an integer $d > 0$. Then, what is the statistics for p -ranks of Artin-Schreier covers over \mathbb{F}_p with genus $g(d, p)$, as $p \rightarrow \infty$.

In section 4, the results are illustrated through an explicit example.

Acknowledgements. The author participated in the workshop [RNT July 12-23, 2021](#), in which he was introduced to the broader theme of arithmetic statistics for families of curves over finite fields. He would like to thank the organizers Allecher Serrano López, Heidi Goodson and McKenzie West for the marvelous experience. He would also like to thank Soumya Sankar for helpful discussions.

2. PRELIMINARIES

Let p denote a prime number and $q = p^n$ be a power of p . We assume that p is odd. To a rational function $f(x) \in \mathbb{F}_q(x)$, we associate the Artin-Schreier cover \mathcal{C}_f of \mathbf{P}^1 , taken to be the projective closure of the affine curve defined by the equation $y^p - y = f(x)$. The map $\mathcal{C}_f \rightarrow \mathbf{P}^1$ sends $(x, y) \mapsto x$. Note that the Galois group of the cover is isomorphic to $\mathbb{Z}/p\mathbb{Z}$ and generated by $(x, y) \mapsto (x, y + 1)$.

We count isomorphism classes of covers that are unramified at ∞ .

Definition 2.1. Two covers \mathcal{C}_f and \mathcal{C}_g associated to $f, g \in \mathbb{F}_q(x)$ are isomorphic if there is an isomorphism $\varphi : \mathcal{C}_f \xrightarrow{\sim} \mathcal{C}_g$ which fits into a commuting diagram:

$$\begin{array}{ccc} \mathcal{C}_f & \xrightarrow{\varphi} & \mathcal{C}_g \\ \downarrow & & \downarrow \\ \mathbf{P}^1 & \xrightarrow{\text{id}} & \mathbf{P}^1. \end{array}$$

We express this relation by simply writing $\mathcal{C}_f \simeq_{\mathbf{P}^1} \mathcal{C}_g$, note that it is stronger than requiring that \mathcal{C}_f and \mathcal{C}_g are isomorphic as curves.

Let $B \subset \mathbf{P}^1(\overline{\mathbb{F}}_p)$ be the set of poles of $f(x)$. Note that the cover is ramified at precisely the points in B . Letting d_α be the order of the pole of $f(x)$ at α , we set $\mathcal{D}_f = \sum_{\alpha \in B} d_\alpha \alpha$ to be the ramification divisor. We shall explicitly work with partial fractions. In setting up notation, for $\alpha \in B$, define

$$x_\alpha := \begin{cases} \frac{1}{(x-\alpha)} & \text{if } \alpha \neq \infty, \\ x & \text{if } \alpha = \infty. \end{cases}$$

Then, using a partial fraction decomposition, we write

$$f(x) = \sum_{\alpha} f_\alpha(x_\alpha),$$

where $f_\alpha(x)$ is a polynomial of degree d_α . The order of the pole of f at α is d_α . We wish to reduce the question of counting equivalence classes of Artin-Schreier covers to the simpler question of counting rational functions with prescribed properties. Following [10], we introduce a class of rational functions that are suitable for counting, which we shall refer to as *admissible*.

Definition 2.2. A rational function $f(x)$ is *admissible* if its partial fractions decomposition satisfies the following conditions

- (1) At ∞ , the cover $\mathcal{C}_f \rightarrow \mathbf{P}^1$ is unramified, in other words, the polynomial $f_\infty(x)$ is a constant.
- (2) For $\alpha \neq \infty$, f_α has no constant term.
- (3) If $p|j$, then the coefficient of x^j in f_α is 0.

Let \mathcal{S} be the set of admissible functions, and $\mathcal{S}(\mathbb{F}_q)$ the subset with coefficients in \mathbb{F}_q .

Note that (1) implies that $f(x)$ can be expressed as a quotient $f(x) = g(x)/h(x)$, where $\deg g(x) \leq \deg h(x)$. Given rational functions f and g , the following gives an explicit criterion for there to be an isomorphism $\mathcal{C}_f \simeq_{\mathbf{P}^1} \mathcal{C}_g$.

Proposition 2.3. Let f and g be rational functions. Then, there is an isomorphism of curves $\mathcal{C}_f \simeq \mathcal{C}_g$ if and only if $f(x) = ug(\gamma x) + \delta^p - \delta$ for $u \in (\mathbb{Z}/p\mathbb{Z})^\times$, $\gamma \in \text{PSL}_2(\mathbb{F}_q)$ and $\delta \in \mathbb{F}_q(x)$. There is an isomorphism of covers $\mathcal{C}_f \simeq_{\mathbf{P}^1} \mathcal{C}_g$ if and only if $f(x) = ug(x) + \delta^p - \delta$ for $u \in (\mathbb{Z}/p\mathbb{Z})^\times$ and $\delta \in \mathbb{F}_q(x)$. Furthermore, if f and g are both admissible and $\mathcal{C}_f \simeq_{\mathbf{P}^1} \mathcal{C}_g$, then, $g(x) = uf(x)$ for $u \in (\mathbb{Z}/p\mathbb{Z})^\times$.

Proof. The first part of the statement follows from the proof of [10, Proposition 2.3]. The remaining assertions are left as an exercise (see [8, Remark 3.9]). \square

Henceforth, the word "isomorphism" shall refer to isomorphisms of covers in the sense of Definition 2.1.

Corollary 2.4. Any Artin-Schreier cover $\mathcal{C} \rightarrow \mathbf{P}^1$ that is unramified at ∞ is isomorphic to one that is of the form $\mathcal{C}_f \rightarrow \mathbf{P}^1$, with $f \in \mathcal{S}$. Moreover, if $\mathcal{C} \rightarrow \mathbf{P}^1$ is defined over \mathbb{F}_q , then, $f \in \mathcal{S}(\mathbb{F}_q)$.

Proof. Let g be such that $\mathcal{C} = \mathcal{C}_g$. Let δ be such that for $f(x) = g(x) + \delta^p - \delta$ condition (3) is satisfied. In greater detail, all monomials involving x^j where $p|j$ in $f_\alpha(x)$ can be removed by adding $(\delta^p - \delta)$. Furthermore, the other conditions are automatic since it is assumed that the cover is unramified at ∞ . Then, it follows that all conditions (1)-(3) are satisfied. \square

Lemma 2.5. Let $f \in \mathcal{S}$ and α and α' that are conjugate over \mathbb{F}_q , then, $d_\alpha = d_{\alpha'}$.

Proof. Let $\sigma \in \text{Gal}(\overline{\mathbb{F}_p}/\mathbb{F}_q)$ be such that $\alpha' = \sigma(\alpha)$. Note that since $f(x) \in \mathbb{F}_q(x)$ we have that $(\sigma f)(x) = f(x)$. Here, $(\sigma f)(x)$ is the rational function obtained by applying σ to the coefficients of $f(x)$. This implies that

$$\sigma(f_\alpha(x_\alpha)) = f_{\alpha'}(x_{\alpha'}).$$

On the other hand,

$$\sigma(f_\alpha(x_\alpha)) = (\sigma f_\alpha)(\sigma(x_\alpha)) = (\sigma f_\alpha)(x_{\alpha'}).$$

Setting $y = x_{\alpha'}$, we find that $f_{\alpha'}(y) = \sigma(f_\alpha)(y)$, and hence, $d_{\alpha'} = d_\alpha$. \square

We introduce some further notation:

- Let $\mathcal{AS}_g(\mathbb{F}_q)$ be the set of all isomorphism classes of Artin-Schreier covers $\mathcal{C} \rightarrow \mathbf{P}^1$ defined over \mathbb{F}_q that are of genus g and unramified at ∞ .
- Set $\mathcal{S}_g(\mathbb{F}_q)$ to be the set of all admissible rational functions $f(x) \in \mathbb{F}_q(x)$, such that $\mathcal{C}_f \in \mathcal{AS}_g(\mathbb{F}_q)$. The map $\mathcal{S}_g(\mathbb{F}_q) \rightarrow \mathcal{AS}_g(\mathbb{F}_q)$ is surjective by Corollary 2.4.
- Set $\mathcal{AS}_{g,\tau}(\mathbb{F}_q)$ (resp. $\mathcal{S}_{g,\tau}(\mathbb{F}_q)$) to be the subset of $\mathcal{AS}_g(\mathbb{F}_q)$ (resp. $\mathcal{S}_g(\mathbb{F}_q)$) for which the p -rank (resp. p -rank of the associated Artin-Schreier cover) is τ .

We shall compute the large q -limit:

$$(2.1) \quad \mathfrak{d}'(p, g, \tau) := \lim_{q \rightarrow \infty} \left(\frac{\#\mathcal{S}_{g,\tau}(\mathbb{F}_q)}{\#\mathcal{S}_g(\mathbb{F}_q)} \right),$$

where $g > 0$ and q ranges over all powers of p . From this, it is easy to calculate

$$(2.2) \quad \mathfrak{d}(p, g, \tau) := \lim_{q \rightarrow \infty} \left(\frac{\#\mathcal{AS}_{g,\tau}(\mathbb{F}_q)}{\#\mathcal{AS}_g(\mathbb{F}_q)} \right).$$

We end this section by reviewing some well-known facts, see [10, section 2.1] for a more comprehensive treatment. Let $f \in \mathcal{S}(\mathbb{F}_q)$ and $Q(x)$ be an irreducible polynomial in $\mathbb{F}_q[x]$ for which $d_\alpha > 0$ for each of the zeros of $Q(x)$. Since f has coefficients in \mathbb{F}_q , it follows from Lemma 2.5 that d_α is the same for all roots α of f . We shall denote this number by d_Q .

Let d denote the following sum

$$(2.3) \quad d := -2 + \sum_{\alpha} (d_\alpha + 1) = -2 + \sum_Q \deg(Q)(d_Q + 1),$$

where α runs over all poles of $f(x)$ and Q ranges over all irreducible monic polynomials $Q \in \mathbb{F}_q[x]$ such that $d_Q > 0$. In the formula above, d_α is set to be equal to -1 if there is no pole at α . By an application of the Riemann-Hurwitz formula, the genus of \mathcal{C}_f is given by

$$g = \left(\frac{p-1}{2} \right) (d + (d_\infty + 1)).$$

Since f is unramified at ∞ , we have by definition, $d_\infty = -1$, and the above formula simplifies to $g = d \left(\frac{p-1}{2} \right)$. Letting $r+1$ be the number of poles of $f(x)$, it follows from the Deuring-Shafarevich formula [5, Corollary 1.8] that the p -rank is given by $\tau = r(p-1)$.

3. MAIN RESULTS

We prove the main results of this paper. First, we treat the geometric problem, after which we study the arithmetic variant, and see that the densities match up. In the next section, we provide an example to illustrate some of the results.

3.1. Geometric problem: fixed prime, large q -limit. Fix a prime number p and let $g = d \left(\frac{p-1}{2} \right)$. Assume throughout that $d > 0$. In this section we study the large q -limits $\mathfrak{d}'(p, g, \tau)$ and $\mathfrak{d}(p, g, \tau)$ defined previously. We introduce some further notation. Let $f(x) \in \mathcal{S}$, note that this means that each polynomial $f_\alpha(x)$ satisfies the conditions of Definition 2.2. For $\alpha \neq \infty$, we write $f_\alpha(x) = \sum_{i=1}^{d_\alpha} c_{i,\alpha} x^i$, where the terms $c_{i,\alpha} = 0$ for all terms i that are divisible by p . Thus, it follows from condition (3) that $d_\alpha \not\equiv 0 \pmod{p}$. Note that it is assumed that $f_\infty(x)$ is a constant. For $j \in \mathbb{Z}_{\geq 1}$ with $j \not\equiv 0 \pmod{p}$, we set $g_j(x)$ to be the product

$$g_j(x) := \left(\prod_{d_\alpha=j} (x - \alpha) \right),$$

and set $g(x) := \prod_j g_j(x)^j$. Note that if α and α' are conjugate, then, by Lemma 2.5, $d_\alpha = d_{\alpha'}$. It follows that $g_j(x) \in \mathbb{F}_q[x]$.

We write $f(x) = \frac{h(x)}{g(x)}$, where $\deg h(x) \leq \deg g(x)$. Note that the polynomials $g_j(x)$ are squarefree polynomials. Set $\kappa_j := \deg g_j(x)$, note that

$$d + 2 = \sum_j (j + 1) \kappa_j,$$

hence, this gives rise to a partition of $d + 2$. Here, κ_j is the multiplicity of $j + 1$ as a term in the partition, and $j \not\equiv 0 \pmod{p}$. We have set $r + 1$ to be the total number of poles of $f(x)$, and this is given by

$$r + 1 = \sum_j \deg g_j(x) = \sum_j \kappa_j.$$

Recall from the previous section that the p -rank of \mathcal{C}_f is given by $\tau = r(p - 1)$. Set $\lambda := \#\{j | \kappa_j > 0\}$, note that $r + 1$ is the total number of terms in the partition of m and λ the number of distinct terms. In the case when $\kappa_j = 0$ for all $j > 1$, we have that d is even and $\kappa_1 = \frac{d+2}{2}$. In this case, $r = \frac{d}{2}$ and the p -rank equal to the genus. Thus, the curve is ordinary precisely when the degree at each pole is at most 1. There is only one partition that corresponds to this case when d is even, and no partitions when d is odd.

Definition 3.1. Let $g = d \left(\frac{p-1}{2} \right) > 0$ and $r > 0$. Denote by $\Omega_p(r + 1, d + 2)$ be the set of partitions of $d + 2$ into $r + 1$ numbers that are $\not\equiv 1 \pmod{p}$, and set $N_p(r + 1, d + 2) := \#\Omega_p(r + 1, d + 2)$.

Represent a partition in $\Omega_p(r + 1, d + 2)$ by the *multiplicity vector*, i.e., the tuple of integers $\kappa = (\kappa_1, \dots, \kappa_t)$, where (by (2.3))

$$\sum_{j=1}^t (j + 1) \kappa_j = d + 2.$$

Here, κ_j is the multiplicity of $(j+1)$ and $j \not\equiv 1 \pmod{p}$. The number of terms in the associated partition is

$$r+1 = \sum_j \kappa_j.$$

The multiplicity vector of \mathcal{C}_f is denoted $\kappa(f)$.

Definition 3.2. Let $\mathcal{S}_g^\kappa(\mathbb{F}_q)$ be the subset of $\mathcal{S}_g(\mathbb{F}_q)$ consisting of all $f(x)$ for which $\kappa(f) = \kappa$.

Lemma 3.3. Let $\kappa \in \Omega_p(r+1, d+2)$. We have the following assertions:

(1) Suppose that $\kappa_j = 0$ for all $j \geq p$. Then, we have that

$$\left(1 - \frac{1}{q}\right)^\lambda q^{d+3} - (d+2) \left(1 - \frac{1}{q}\right)^\lambda q^{d+2} \leq \#\mathcal{S}_g^\kappa(\mathbb{F}_q) \leq \left(1 - \frac{1}{q}\right)^\lambda q^{d+3},$$

in particular, as $q \rightarrow \infty$,

$$\#\mathcal{S}_g^\kappa(\mathbb{F}_q) \sim q^{d+3}.$$

(2) Suppose that $\kappa_j \neq 0$ for some $j \geq p$. Then, we have that

$$\#\mathcal{S}_g^\kappa(\mathbb{F}_q) = o(q^{d+3}).$$

Proof. Recall that we have expressed $f(x) = h(x)/g(x)$, where $g(x) = \prod_j g_j(x)^j$ and $\kappa_j = \deg g_j(x)$, where $g_j(x)$ is a squarefree polynomial with coefficients in \mathbb{F}_q . Assume without loss of generality that $g_j(x)$ is a monic polynomial for all j . Note that the degree of $g(x)$ is

$$\sum_j j\kappa_j = d+2 - \tau.$$

Since $f_\infty(x)$ is a constant, we have that $\deg h(x) \leq \deg g(x)$.

First, we consider the case when $\kappa_j = 0$ for all $j \geq p$. In this case, for every pole α of $f(x)$, we have that $d_\alpha < p$. Thus, the condition (3), which requires that the coefficient of x^j be zero in $f_\alpha(x)$ be zero whenever $p|j$, is automatically satisfied. Since $f(x)$ is unramified at ∞ , $\deg h(x) \leq \deg g(x)$. Furthermore, $h(x)$ is coprime to $g(x)$. Thus for each choice of $g(x)$, the number of choices of $h(x)$ is $\leq q^{\deg g(x)+1} = q^{d+3-\tau}$. The number of polynomials in $\mathbb{F}_q[x]$ of degree $\leq \deg g(x)$ that share a common factor with $g(x)$ is

$$\leq \deg g(x) q^{\deg g(x)} = (d+2-\tau)q^{d+2-\tau} = O(q^{d+2-\tau}).$$

To see this, note that the number of irreducible factors of $g(x)$ is $\leq \deg g(x)$, and for each irreducible factor, there are at most $q^{\deg g(x)}$ choices of $h(x)$ which it divides.

Hence, we find that for each choice of $g(x)$, the number of choices for $h(x)$ lies in between $q^{d+3-\tau} - (d+2)q^{d+2-\tau}$ and $q^{d+2-\tau}$. We count the number of choices of $g(x)$.

The number of monic squarefree polynomials of degree κ_j with coefficients in \mathbb{F}_q is $(1 - \frac{1}{q})q^{\kappa_j}$, see for instance [6, Theorem 1]. Therefore, the number of choices of $g_j(x)$ is $(1 - \frac{1}{q})q^{\kappa_j}$. Hence, when κ is prescribed, the number of choices of $g(x) = \prod_j g_j(x)^j$ is

$$\prod_j \left(\left(1 - \frac{1}{q}\right) q^{\kappa_j} \right) = \left(1 - \frac{1}{q}\right)^\lambda q^\tau.$$

Hence, the total number of choices for $f(x) = h(x)/g(x)$ is

$$\leq \left(\left(1 - \frac{1}{q}\right)^\lambda q^\tau \right) \times q^{d+3-\tau} = \left(1 - \frac{1}{q}\right)^\lambda q^{d+3}$$

and

$$\begin{aligned} &\geq \left(\left(1 - \frac{1}{q}\right)^\lambda q^\tau \right) \times \left(q^{d+3-\tau} - (d+2)q^{d+2-\tau} \right) \\ &= \left(1 - \frac{1}{q}\right)^\lambda q^{d+3} - (d+2) \left(1 - \frac{1}{q}\right)^\lambda q^{d+2}. \end{aligned}$$

This proves part (1).

Next, assume that for some $j > p$, we have that $\kappa_j \neq 0$. In this case, the same calculation goes through with the one difference being that there is an additional constraint requiring that the coefficient of x^j in $f_\alpha(x)$ is 0 for all $j > 0$ such that $p|j$. This condition was automatically satisfied in the previous case. In this case, this gives an additional condition on $h(x)$, which forces the number of choices to be

$$\leq \left(1 - \frac{1}{q}\right)^\lambda q^{d+2} = o(q^{d+3}).$$

Let us explain this in greater detail.

Since $\kappa_j \neq 0$ for some $j > p$, there exists a pole α of $f(x)$ such that $d_\alpha = j > p$. The coefficient of x^p in $f_\alpha(x)$ is required to be 0. Set $\eta(x) \in \mathbb{F}_q(x)$ to denote the function given by

$$\eta(x) := \sum_{\alpha'} \frac{1}{(x - \alpha')},$$

and note that $\eta(x)^p = \sum_{\alpha'} \frac{1}{(x - \alpha')^p}$. Here, α' ranges over the \mathbb{F}_q -conjugates of α . This function $\eta(x)^p$ has coefficients in \mathbb{F}_q and is expressed as $\eta(x)^p = \frac{\eta_1(x)}{\eta_2(x)}$, with

$$\deg \eta_1(x) \leq \deg \eta_2(x) \leq \kappa_j.$$

Moreover, note that $\eta_2(x)$ divides $\prod_{\alpha'} (x - \alpha')^p$, and hence divides $\prod_{\alpha'} (x - \alpha')^j$. Since this latter product divides $g(x)$, it follows that $\eta_2(x)$ divides $g(x)$. For any choice of $0 \neq c \in \mathbb{F}_q$, the function $f_c(x) := f(x) + c\eta(x)^p$ does not satisfy condition (3) since now the coefficient of x^p in $f_{c,\alpha}(x)$ is nonzero, and $p < j = d_\alpha$. We may express $f_c(x) = h_c(x)/g(x)$, where

$$\deg h_c(x) \leq \deg g(x).$$

The number of functions $f(x) = h(x)/g(x)$ with:

- (1) $\deg h(x) \leq \deg g(x)$,
- (2) $g(x)$ expressible as a product

$$g(x) = \prod_j g_j(x)^j,$$

where $g_j(x) \in \mathbb{F}_q[x]$ is a squarefree monic polynomial of degree κ_j ,

is by the previous calculation, $\leq \left(1 - \frac{1}{q}\right)^\lambda q^{d+3}$. On the other hand, from an admissible function $f(x) = \frac{h(x)}{g(x)}$, we have constructed q new functions $f_c(x)$, only one of which is admissible (i.e., when $c = 0$). It is easy to see that if $f(x)$ and $F(x)$ are distinct admissible functions, then $f_c(x) \neq F_{c'}(x)$ for all c, c' . Hence, the total number of admissible functions for the partition vector κ is

$$\leq \frac{1}{q} \left(\left(1 - \frac{1}{q}\right)^\lambda q^{d+3} \right) = O(q^{d+2}) = o(q^{d+3}).$$

□

Recall that the p -rank is $\tau = r(p - 1)$.

Definition 3.4. Let $M_p(r + 1, d + 2)$ be the number of partitions $\kappa \in \Omega_p(r + 1, d + 2)$ such that $\kappa_j = 0$ unless $1 \leq j < p$. Set

$$M_p(d + 2) := \sum_{r=0}^{\lfloor d/2 \rfloor} M_p(r + 1, d + 2).$$

Theorem 3.5. Fix a prime number p . With respect to notation above, we have that

$$\mathfrak{d}'(p, g, \tau) = \mathfrak{d}(p, g, \tau) = \frac{M_p(r + 1, d + 2)}{M_p(d + 2)}.$$

Proof. Note that

$$\#\mathcal{S}_{g,\tau}(\mathbb{F}_q) = \sum_{\kappa \in \Omega_p(r+1, d+2)} \#\mathcal{S}_g^\kappa(\mathbb{F}_q),$$

where we recall that $g = d \left(\frac{p-1}{2} \right)$ and $\tau = r(p - 1)$. It follows from Lemma 3.3 that $\#\mathcal{S}_{g,\tau}(\mathbb{F}_q) \sim M_p(r + 1, d + 2)q^{d+3}$. Likewise, we have that $\#\mathcal{S}_g(\mathbb{F}_q) \sim M_p(d + 2)q^{d+3}$, and it thus follows that

$$\mathfrak{d}'(p, g, \tau) = \frac{M_p(r + 1, d + 2)}{M_p(d + 2)}.$$

On the other hand, if f and g are both admissible, then, by Proposition 2.3,

$$\mathcal{C}_f \simeq_{\mathbf{P}^1} \mathcal{C}_g \Leftrightarrow f = ug \text{ for } u \in (\mathbb{Z}/p\mathbb{Z})^\times.$$

Hence, each isomorphism class of covers (unramified at ∞) consists of exactly $(p - 1)$ admissible functions. As a result,

$$\#\mathcal{AS}_{g,\tau}(\mathbb{F}_q) \sim \frac{M_p(r + 1, d + 2)}{p - 1} q^{d+3} \text{ and } \#\mathcal{AS}_g(\mathbb{F}_q) \sim \frac{M_p(d + 2)}{p - 1} q^{d+3},$$

and consequently, $\mathfrak{d}(p, g, \tau) = \mathfrak{d}'(p, g, \tau)$. □

Note that $\mathfrak{d}(p, g, \tau)$ depends on p .

Corollary 3.6. Suppose that $(r + 1)p < d + 2$, then, $\mathfrak{d}(p, g, \tau) = 0$. In particular, if $p < d + 2$, then, the proportion of isomorphism classes of Artin-Schreier covers of genus $g = d \left(\frac{p-1}{2} \right)$ and p -rank $\tau = 0$ is zero in the large q -limit.

Proof. Note that when $(r+1)p < d+2$, it is not possible to partition $d+2$ into $r+1$ numbers, all of which are $\leq p$. In this case, $M_p(r+1, d+2) = 0$, and the result follows from Theorem 3.5. \square

3.2. Arithmetic problem: large p -limit. We now study the *arithmetic* variant of the problem considered in the previous section. Many of the calculations are similar in spirit. Fix $d > 0$, $r \geq 0$, and let $g = g(d, p) = d \left(\frac{p-1}{2} \right)$, $\tau = \tau(r, p) := r(p-1)$. Note that g and τ depend on p and are increasing at fixed speeds as $p \rightarrow \infty$. This is suppressed in the notation.

Set $\mathfrak{d}'(d, r)$ to be the limit

$$\mathfrak{d}'(d, r) := \lim_{p \rightarrow \infty} \frac{\#\mathcal{S}_g^\tau(\mathbb{F}_p)}{\#\mathcal{S}_g(\mathbb{F}_p)},$$

and set

$$\mathfrak{d}(d, r) := \lim_{p \rightarrow \infty} \frac{\#\mathcal{AS}_g^\tau(\mathbb{F}_p)}{\#\mathcal{AS}_g(\mathbb{F}_p)},$$

with $g = g(d, p)$ and $\tau = \tau(r, p)$.

Definition 3.7. Let $\Theta(r+1, d+2)$ be the set of all partitions of $d+2$ into $r+1$ numbers that are all ≥ 2 , and set $T(r+1, d+2) := \#\Theta(r+1, d+2)$. Let $\Theta(d+2)$ be the set of all partitions of $d+2$ into numbers ≥ 2 .

A partition in $\Theta(r+1, d+2)$ is represented by a partition vector $\kappa = (\kappa_1, \dots, \kappa_t)$, where κ_j is the multiplicity of $(j+1)$ in the partition. There is no dependence on p in the definition of $T(r+1, d+2)$. Note that

$$T(d+2) = \sum_{r=0}^{\lfloor d/2 \rfloor} T(r+1, d+2).$$

Lemma 3.8. Let $\kappa \in \Theta(r+1, d+2)$. Suppose that $\kappa_j = 0$ for all $j \geq p$. Then, we have that

$$\left(1 - \frac{1}{p}\right)^\lambda p^{d+3} - (d+1) \left(1 - \frac{1}{p}\right)^\lambda p^{d+2} \leq \#\mathcal{S}_g^\kappa(\mathbb{F}_p) \leq \left(1 - \frac{1}{p}\right)^\lambda p^{d+3}.$$

Proof. The result follows from the first part of Lemma 3.3. \square

Corollary 3.9. Let $\kappa \in \Theta(r+1, d+2)$, and $g = g(d, p)$, $\tau = \tau(r, p)$. We have that

$$\#\mathcal{S}_g^\kappa(\mathbb{F}_p) \sim p^{d+3},$$

as $p \rightarrow \infty$.

Proof. The condition requiring $\kappa_j = 0$ for all $j \geq p$ is satisfied when p is large, and the result thus follows from Lemma 3.8. \square

Theorem 3.10. Fix an integer d . We have that

$$\mathfrak{d}(d, r) = \mathfrak{d}'(d, r) = \frac{T(r+1, d+2)}{T(d+2)}.$$

Proof. The proof is similar to that of Theorem 3.5. Assume that p is large enough so that $\kappa_j = 0$ for all $j \geq p$. Recall that by abuse of notation, $g := g(d, p)$, $\tau = \tau(r, p)$. Since d and r are fixed, g and τ increase linearly with p (but this is suppressed in our notation). We have that

$$\#\mathcal{S}_{g,\tau}(\mathbb{F}_p) = \sum_{\kappa \in \Theta(r+1, d+2)} \#\mathcal{S}_g^\kappa(\mathbb{F}_p).$$

It follows from Lemma 3.8 and Corollary 3.9 that as $p \rightarrow \infty$,

$$\#\mathcal{S}_{g,\tau}(\mathbb{F}_p) \sim T(r+1, d+2)p^{d+3} \text{ and } \#\mathcal{S}_{g,\tau}(\mathbb{F}_p) \sim T(d+2)p^{d+3}.$$

Putting everything together, we obtain that

$$\mathfrak{d}'(d, r) = \frac{T(r+1, d+2)}{T(d+2)}.$$

That $\mathfrak{d}(d, r) = \mathfrak{d}'(d, r)$ follows from the equalities:

$$\#\mathcal{AS}_{g,\tau}(\mathbb{F}_p) \sim \frac{1}{(p-1)} \#\mathcal{S}_{g,\tau}(\mathbb{F}_p) \text{ and } \#\mathcal{AS}_g(\mathbb{F}_p) \sim \frac{1}{(p-1)} \#\mathcal{S}_g(\mathbb{F}_p),$$

see the proof of Theorem 3.5. □

4. AN EXAMPLE

We illustrate results proved in the previous section through an example. First, let us pick the prime $p = 5$, and let $d = 8$. The genus is given by $g = d \left(\frac{p-1}{2} \right) = 8 \times 2 = 16$. The p -rank $\tau = r(p-1) = 4r$. The values of r range from $r = 0$ to $r = 4$. Note that when $r = 4$, $\tau = 16 = g$, and this is the case when the curve is ordinary.

We compute the proportion of isomorphism classes of Artin-Schreier covers with genus $g = 16$ and 5-rank $\tau = 4r$, in the large q -limit. There are a total of 42 partitions of 10, however, there are constraints on the partitions we consider. Recall that $M_p(r+1, d+2) = M(r+1, 10)$ is the number of partitions

$$10 = \sum_j (j+1)\kappa_j$$

for which $\kappa_j = 0$ for j outside the range $1 \leq j < 5$. Thus, we are to use only numbers 2, 3, 4, 5 in partitioning 10 into $r+1$ numbers. Let's list the partitions for each choice of $r = 0, \dots, 4$.

- (1) $r=0$: There is no partition, since $10 = 10$ is not allowed. Hence, $M(1, 10) = 0$. Thus, the proportion of curves with 5-rank 0 is 0 in the q -limit.
- (2) $r=1$: We write $10 = a + b$, where $a, b \in \{2, \dots, 5\}$. The only partition is

$$10 = 5 + 5,$$

hence, $M(2, 10) = 1$.

- (3) $r=2$: We write $10 = a + b + c$, where $a, b, c \in \{2, \dots, 5\}$. The partitions are

$$10 = 5 + 3 + 2 = 4 + 4 + 2 = 4 + 3 + 3,$$

hence, $M(3, 10) = 3$.

- (4) $r=3$: We write $10 = a + b + c + d$ this time. Since they are all ≥ 2 , there are only two choices:

$$10 = 2 + 2 + 2 + 4 = 2 + 2 + 3 + 3,$$

hence, $M(4, 10) = 2$.

- (5) $r=4$: We come to single ordinary case. The only partition is

$$10 = 2 + 2 + 2 + 2 + 2,$$

and $M(5, 10) = 1$.

Putting it all together, we find that

$$M(10) = \sum_{r=1}^4 M(r+1, 10) = 0 + 1 + 3 + 2 + 1 = 7.$$

Therefore, according to Theorem 3.5, the proportion of isomorphism classes of covers of genus $g = 16$ and p -rank $\tau = 4r$ is given by the following proportions:

$$\begin{aligned} \mathfrak{d}(5, 16, 0) &= \frac{0}{7}, \\ \mathfrak{d}(5, 16, 4) &= \frac{1}{7}, \\ \mathfrak{d}(5, 16, 8) &= \frac{3}{7}, \\ \mathfrak{d}(5, 16, 12) &= \frac{2}{7}, \\ \mathfrak{d}(5, 16, 16) &= \frac{1}{7}. \end{aligned}$$

Let's forget the prime p , and study the arithmetic problem. The for $d = 8$. The value of r ranges from 0 to 4 and specifies a given speed at which τ increases with p . In this setting, we no longer have the constraint that the partitions should involve terms from $\{2, \dots, 5\}$, but only that the terms be ≥ 2 . One again, we enumerate partitions for the five values of r .

- (1) $r=0$: This time, $10 = 10$ is allowed. Hence, $T(1, 10) = 1$.
 (2) $r=1$: We write $10 = a + b$, where $a, b \geq 2$. There are 4 partitions

$$10 = 2 + 8 = 3 + 7 = 4 + 6 = 5 + 5,$$

hence, $T(2, 10) = 4$.

- (3) $r=2$: We write $10 = a + b + c$. The partitions are

$$10 = 6 + 2 + 2 = 5 + 3 + 2 = 4 + 4 + 2 = 4 + 3 + 3,$$

hence, $T(3, 10) = 4$.

- (4) $r=3$: We write $10 = a + b + c + d$ this time. Since they are all ≥ 2 , there are only two choices:

$$10 = 2 + 2 + 2 + 4 = 2 + 2 + 3 + 3,$$

hence, $T(4, 10) = 2$.

- (5) $r=4$: We come to single ordinary case. The only partition is

$$10 = 2 + 2 + 2 + 2 + 2,$$

and $T(5, 10) = 1$.

Hence, we have that $T(10) = 12$. According to Theorem 3.10 the proportions are as follows:

$$\begin{aligned}\mathfrak{d}(d, 0) &= \frac{1}{12}, \\ \mathfrak{d}(d, 1) &= \frac{4}{12}, \\ \mathfrak{d}(d, 2) &= \frac{4}{12}, \\ \mathfrak{d}(d, 3) &= \frac{2}{12}, \\ \mathfrak{d}(d, 4) &= \frac{1}{12}.\end{aligned}$$

Thus, as $p \rightarrow \infty$, and g increases with speed $\frac{d}{2} = 4$, the p -rank grows with expected speed

$$0 \times \frac{1}{12} + 1 \times \frac{4}{12} + 2 \times \frac{4}{12} + 3 \times \frac{2}{12} + 4 \times \frac{1}{12} = \frac{19}{12} = 1.5833 \dots$$

REFERENCES

- [1] A. Bucur, C. David, B. Feigon, N. Kaplan, M. Lalin, E. Ozman, and M. M. Wood. The distribution of \mathbb{F}_q -points on cyclic ℓ -covers of genus g . *International Mathematics Research Notices*, 2016(14):4297–4340, 2016.
- [2] A. Bucur, C. David, B. Feigon, and M. Lalin. Biased statistics for traces of cyclic p -fold covers over finite fields., 2011.
- [3] A. Bucur, C. David, B. Feigon, and M. Lalin. Statistics for ordinary artin-schreier covers and other p -rank strata. *Transactions of the American Mathematical Society*, 368(4):2371–2413, 2016.
- [4] G. Cheong, M. Wood, and A. Zaman. The distribution of points on superelliptic curves over finite fields. *Proceedings of the American Mathematical Society*, 143(4):1365–1375, 2015.
- [5] R. M. Crew. Etale p -covers in characteristic p . *Compositio Mathematica*, 52(1):31–45, 1984.
- [6] A. Knopfmacher and J. Knopfmacher. Counting irreducible factors of polynomials over a finite field. *Discrete Mathematics*, 112(1-3):103–118, 1993.
- [7] P. Kurlberg and Z. Rudnick. The fluctuations in the number of points on a hyperelliptic curve over a finite field. *Journal of Number Theory*, 129(3):580–587, 2009.
- [8] R. Pries and H. J. Zhu. The p -rank stratification of artin-schreier curves. In *Annales de l'Institut Fourier*, volume 62, pages 707–726, 2012.
- [9] Z. Rudnick. Traces of high powers of the frobenius class in the hyperelliptic ensemble. *arXiv preprint arXiv:0811.3649*, 2008.
- [10] S. Sankar. Proportion of ordinarity in some families of curves over finite fields. *arXiv preprint arXiv:1904.12173*, 2019.
- [11] F. Thorne and M. Xiong. Distribution of zeta zeroes for cyclic trigonal curves over a finite field. *preprint*, 2014.
- [12] M. Xiong. The fluctuations in the number of points on a family of curves over a finite field. *Journal de théorie des nombres de Bordeaux*, 22(3):755–769, 2010.

(Ray) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER BC, CANADA V6T 1Z2

Email address: anweshray@math.ubc.ca