

# EULER CHARACTERISTIC AND SIGNATURE OF REAL SEMI-STABLE DEGENERATIONS

ERWAN BRUGALLÉ

**ABSTRACT.** We give a motivic proof of the fact that for non-singular real tropical complete intersections, the Euler characteristic of the real part is equal to the signature of the complex part. This has originally been proved by Itenberg in the case of surfaces in  $\mathbb{C}P^3$ , and has been successively generalized by Bertrand, and by Bihan and Bertrand. Our proof, different from the previous approaches, is an application of the motivic nearby fiber of semi-stable degenerations. In particular it extends the original result by Itenberg-Bertrand-Bihan to real analytic families admitting a  $\mathbb{Q}$ -non-singular tropical limit.

Given a real algebraic variety  $X$ , we denote by  $\mathbb{C}X$  the set of its complex points, and by  $\mathbb{R}X$  the set of its real points. Recall that the *signature*  $\sigma(M)$  of an even dimensional oriented compact smooth manifold  $M$  is defined to be 0 if  $\dim M = 4n + 2$ , and to be the signature on the (symmetric) intersection form on  $H_{2n}(M; \mathbb{Q})$  if  $\dim M = 4n$ . The Euler characteristic with closed support (or Borel-Moore Euler characteristic) of a topological space  $M$  is denoted by  $\chi^c(M)$ .

## 1. STATEMENT

**1.1. Context.** A real algebraic variety  $X$  is said to *satisfy*  $\chi = \sigma$  if

$$\chi^c(\mathbb{R}X) = \sigma(\mathbb{C}X).$$

This definition is motivated by the old empirical observation that an important proportion of known real algebraic varieties satisfy  $\chi = \sigma$ , in particular among compact non-singular  $M$ -varieties. By the Smith-Thom inequality (see for example [Man17, Section 3.2]), any real algebraic variety  $X$  satisfies

$$\sum_i \beta_i(\mathbb{R}X) \leq \sum_i \beta_i(\mathbb{C}X),$$

where  $\beta_i$  denotes the  $i$ -th Betti number with  $\mathbb{Z}/2\mathbb{Z}$ -coefficients. Furthermore the difference between the two hand-sides is even, and we say that  $X$  is an  $(M - i)$ -variety if

$$\sum_i \beta_i(\mathbb{C}X) - \sum_i \beta_i(\mathbb{R}X) = 2i.$$

If  $X$  is a compact and non-singular  $M$ -variety, Rokhlin Congruence [DK00, 2.7.1] asserts that

$$\chi(\mathbb{R}X) = \sigma(\mathbb{C}X) \pmod{16},$$

yet almost all known examples of non-singular compact  $M$ -varieties satisfy the stronger inequality  $\chi(\mathbb{R}X) = \sigma(\mathbb{C}X)$ . If many examples among  $M$ -surfaces not satisfying this equality are known (see for example [DK00, Man17]), all non-singular compact  $M$ -varieties of dimension at least 3 that I know<sup>1</sup> satisfy  $\chi = \sigma$ . A heuristic explanation of this fact is the following : a large class of known real algebraic varieties is obtained by gluing “basic real algebraic varieties” satisfying  $\chi = \sigma$ , a property

---

*2020 Mathematics Subject Classification.* Primary 14P25, 14T90.

*Key words and phrases.* Topology of real algebraic varieties, semi-stable degenerations, combinatorial patchworking, tropical geometry.

<sup>1</sup>I should precise that I know very few such high-dimensional  $M$ -varieties.

that is preserved under gluing. This latter statement is the content of Proposition 2.1 below, which is an elementary observation relying on deep results about motivic nearby fibers and limit mixed Hodge structures. By “basic real algebraic varieties”, we mean for example real projective spaces, or more generally real toric varieties.

As an application of this observation, we prove Theorem 1.2 below: real algebraic varieties constructed out of non-singular tropical varieties satisfy  $\chi = \sigma$ . This generalizes the case of complete intersections, originally proved by Itenberg [Ite97] for hypersurfaces of dimension 2, by Bertrand [Ber10] for hypersurfaces of higher dimensions, and then by Bihan and Bertrand [BB07] for any complete intersections. Theorem 1.2 is actually about potentially non-compact algebraic varieties, and we first briefly recall the extension of the signature to all complex algebraic varieties.

Given a field  $\mathbb{K}$ , the *Grothendieck group*  $K_0(\text{Var}_{\mathbb{K}})$  is the abelian group generated by isomorphism classes  $[X]$  of algebraic varieties over  $k$  modulo the *scissor relation*

$$[X] = [X \setminus Y] + [Y]$$

for any closed algebraic subvariety  $Y$  of  $X$ . Considering the product

$$[X] \times [Y] = [X \times Y]$$

turns  $K_0(\text{Var}_{\mathbb{K}})$  into a commutative ring with  $0 = [\emptyset]$  and  $1 = [pt]$ . It is not too difficult to show (see for example [Pet10, Lecture 1]) that when  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{C}$ , Euler characteristic with closed support provides a ring morphism

$$\begin{array}{ccc} \chi^c : K_0(\text{Var}_{\mathbb{K}}) & \longrightarrow & \mathbb{Z} \\ X & \longmapsto & \chi^c(\mathbb{K}X) \end{array} .$$

Such a ring (or group) morphism is called a *motivic invariant*. It is a much less obvious result that the signature of non-singular projective complex algebraic varieties extends to a motivic invariant

$$\sigma : K_0(\text{Var}_{\mathbb{C}}) \rightarrow \mathbb{Z}.$$

This is a consequence of the combination of the two following facts:

- the signature of a non-singular projective complex algebraic manifold is the evaluation at 1 of its Hirzebruch genus; this is the Hodge index Theorem, see for example [Voi02, Theorem 6.33];
- the Hirzebruch genus extends to a motivic invariant  $\chi_y : K_0(\text{Var}_{\mathbb{C}}) \rightarrow \mathbb{Z}[y]$ ; see for example [PS08, Remark 5.6], since  $\chi_y$  is in its turn a specialization of the Hodge-Euler polynomial.

Using the scissor relation, it is easy to check that, as mentioned above, basic real algebraic varieties satisfy  $\chi = \sigma$ .

**Example 1.1.** One computes easily that  $\chi^c(\mathbb{R}) = -1$ . On the other hand, one has

$$0 = \sigma(\mathbb{C}P^1) = \sigma(\mathbb{C}) + \sigma(pt) = \sigma(\mathbb{C}) + 1.$$

Hence we deduce that  $\chi^c(\mathbb{R}) = \sigma(\mathbb{C}) = -1$ , that is to say  $\mathbb{C}$  satisfies  $\chi = \sigma$ . Since both  $\chi^c$  and  $\sigma$  are ring morphisms, we deduce that  $\chi^c(\mathbb{R}^n) = \sigma(\mathbb{C}^n) = (-1)^n$ . In other words, real affine spaces satisfy  $\chi = \sigma$ . By the scissor relation again, this implies that any real algebraic variety with a stratification by real affine spaces satisfies  $\chi = \sigma$ . In particular, all Grassmannians, and more generally all real flag varieties, satisfy  $\chi = \sigma$ . The case of real toric varieties, i.e. equipped with the standard real structure induced by the complex conjugation on  $\mathbb{C}^*$ , can be handled similarly. One has

$$\chi^c(\mathbb{R}^*) = -2 = -1 - 1 = \sigma(\mathbb{C}^*) - \sigma(pt).$$

Hence real torus  $(\mathbb{C}^*)^n$  satisfy  $\chi = \sigma$ , and so does any real toric varieties since it admits a stratification by real tori.

**1.2. Real approximations of  $\mathbb{Q}$ -non-singular tropical varieties.** Given a fan  $\Delta \subset \mathbb{R}^n$ , we denote by  $Tor_{\mathbb{C}}(\Delta)$  the complex toric variety defined by  $\Delta$ , see [Ful93]. In this note we only consider fans  $\Delta$  for which  $Tor_{\mathbb{C}}(\Delta)$  is non-singular, in which case we can also consider the tropical toric variety  $Tor_{\mathbb{T}}(\Delta)$  defined by  $\Delta$ , see [BIMS15, MR]. In what follows, we use the notions defined in [IKMZ19] of  $\mathbb{Q}$ -non-singular tropical varieties of  $Tor_{\mathbb{T}}(\Delta)$ , and of tropical limits of non-singular analytic families  $X \subset \mathbb{C}P^N \times D^*$  of algebraic subvarieties of  $\mathbb{C}P^N$  over the punctured unit disk  $D^* \subset \mathbb{C}$ . For such a family, we denote by  $X_t$  the member of the family corresponding to  $t \in D^*$ .

**Theorem 1.2.** *Let  $\Delta$  be a fan defining a non-singular projective toric variety  $Tor_{\mathbb{C}}(\Delta)$ . Suppose that  $X \subset Tor_{\mathbb{C}}(\Delta) \times D^*$  is a non-singular real analytic family of algebraic subvarieties of  $Tor_{\mathbb{C}}(\Delta)$ , admitting a  $\mathbb{Q}$ -non-singular tropical limit in  $Tor_{\mathbb{T}}(\Delta)$ . Then for any subfan  $\delta$  of  $\Delta$  and for any  $t_0 \in \mathbb{R}D^*$  small enough, the real analytic variety  $X_{t_0} \cap Tor_{\mathbb{C}}(\delta)$  satisfies  $\chi = \sigma$ .*

As mentioned above the case of compact hypersurfaces has been earlier proved by Itenberg [Ite97] and Bertrand [Ber10]. This has been generalized to possibly non-compact complete intersections by Bertand and Bihan [BB07], where they used motivic aspects of  $\chi^c$  and  $\sigma$  to reduce to the case of hypersurfaces. The strategy in each of these three papers is then to separately compute  $\chi^c$  and  $\sigma$ , and to check that both numbers coincide. Alternative proofs using tropical homology were proposed in the case of complete intersections by Arnal [Arn17] (still by equating two separate computations), and by Renaudineau and Shaw [RS18]. In this latter work, the equality  $\chi^c(\mathbb{R}X_t) = \sigma(\mathbb{C}X_t)$  is a consequence of a much stronger result: it follows from the existence of a spectral sequence starting from the tropical homology of the tropical limit of  $X$  and converging to the homology of  $\mathbb{R}X$ .

Hence in a sense all previous partial proofs of Theorem 1.2 are based on separate computations of both  $\chi^c$  and  $\sigma$ . Our proof of Theorem 1.2 uses a different strategy : both quantities satisfy the same gluing relations under totally real semi-stable degenerations by Proposition 2.1, while the tropical non-singularity assumption ensures that all pieces involved in the gluing satisfy  $\chi = \sigma$ .

Note that in the case of hypersurfaces, Itenberg, Bertrand, and Bertand and Bihan do not work in the tropical geometry framework, but in the dual setup of unimodular subdivisions of polytopes. It is interesting that their proof also applies to non-regular (or non-convex) subdivisions, that is to say *real combinatorial hypersurfaces* (see [IS03] for a definition) also satisfy  $\chi = \sigma$ . Our proof of Theorem 1.2, as well as the partial proofs by Arnal, and Renaudineau and Shaw, do not seem to extend to real combinatorial hypersurfaces.

**Acknowledgment.** This work is partially supported by the grant TROPICOUNT of Région Pays de la Loire, and the ANR project ENUMGEOM NR-18-CE40-0009-02.

## 2. PROOF

**2.1. Real semi-stable degenerations.** Let  $f : X \rightarrow D$  be a proper analytic map from a non-singular complex algebraic manifold  $X$  to the unit disk  $D \subset \mathbb{C}$  such that

- $X_t$  is a non-singular algebraic manifold for all  $t \in D^*$ ;
- $X_0 = f^{-1}(0)$  is a reduced algebraic variety with non-singular components crossing normally.

Such a map is called a *semi-stable degeneration* (of  $X_t$  for  $t \neq 0$ ). Denoting by  $(E_i)_{i \in J}$  the irreducible components of  $X_0$ , we define for  $I \subset J$

$$E_I^\circ = \bigcap_{i \in I} E_i \setminus \bigcup_{j \notin I} E_j.$$

Suppose now that  $X$  is real, and that  $f$  is real when  $D$  is equipped with the standard complex conjugation. we say that  $f$  is *totally real* if each irreducible component of  $X_0$  is real.

**Proposition 2.1.** *Let  $f : X \rightarrow D$  be a totally real semi-stable degeneration. Then for any  $t \in \mathbb{R}D^*$*

$$\sigma(\mathbb{C}X_t) = \sum_{\emptyset \neq I \subset J} 2^{|I|-1} \sigma(\mathbb{C}E_I^o) \quad \text{and} \quad \chi^c(\mathbb{R}X_t) = \sum_{\emptyset \neq I \subset J} 2^{|I|-1} \chi^c(\mathbb{R}E_I^o).$$

*In particular  $X_t$  satisfies  $\chi = \sigma$  as soon as all  $E_I^o$ 's satisfy  $\chi = \sigma$ .*

*Proof.* The statement about the signature follows from deep results on motivic nearby fibers and limit mixed Hodge structures. We refer to [KS16, Section 1 and 3] for a concise exposition of what is needed here, and to [DL01, Section 3], [Bit05, Section 2], and [PS08, Section 11] for more details. Recall first that the Hirzebruch genus  $\chi_y(Y)$  of a projective non-singular complex algebraic variety is the polynomial in  $y$  defined by

$$\chi_y(Y) = \sum_{p \geq 0} \left( \sum_{q \geq 0} (-1)^q h^{p,q}(Y) \right) y^p,$$

and that it satisfies  $\chi_1(Y) = \sigma(Y)$  by the Hodge index Theorem. Hirzebruch genus turns out to be a motivic invariant, that is to say it extends to a ring morphism  $\chi_y : K_0(\text{Var}_{\mathbb{C}}) \rightarrow \mathbb{Z}[y]$ . In particular  $\sigma(Y) = \chi_1(Y)$  for any complex algebraic variety  $Y$ . The motivic nearby fiber of  $f$ , introduced by Denef and Loeser, is defined by

$$\psi_f = \sum_{\emptyset \neq I \subset J} [E_I^o](1 - [\mathbb{C}])^{|I|-1} \in K_0(\text{Var}_{\mathbb{C}}),$$

and satisfies the great property  $\chi_y(\psi_f) = \chi_y(X_t)$  for  $t \neq 0$ . So one has

$$\chi_y(X_t) = \sum_{\emptyset \neq I \subset J} \chi_y(E_I^o)(2 - \chi_y(\mathbb{C}P^1))^{|I|-1} = \sum_{\emptyset \neq I \subset J} (1 + y)^{|I|-1} \chi_y(E_I^o).$$

The first statement of the proposition is obtained by evaluating this identity at  $y = 1$ .

The statement about Euler characteristic follows from the observation that  $\mathbb{R}X_t$  is the disjoint union of  $2^{|I|-1}$  topological copies of  $\mathbb{R}E_I^o$ , with  $I$  ranging over all possible sets  $\emptyset \neq I \subset J$ . Indeed, by assumption one can locally express  $X$  in coordinates at a point of  $\mathbb{R}E_I^o$  as the solutions of the equation

$$x_1 x_2 \cdots x_{|I|} = \alpha \subset \mathbb{R}^n \times \mathbb{R},$$

where  $\alpha$  is the deformation parameter, with  $\alpha = 0$  corresponding to the central fiber  $X_0$ . Given a non zero  $\alpha$ , the corresponding smooth fiber of  $f$  is locally given in  $\mathbb{R}^n = \mathbb{R}^{|I|} \times \mathbb{R}^{n-|I|}$  by the solutions of the above equation. This set is homeomorphic to the disjoint union of  $2^{|I|-1}$  copies of  $\mathbb{R}_{\geq 0}^{|I|} \times \mathbb{R}^{n-|I|}$ , and the result follows.  $\square$

**2.2. Proof of Theorem 1.2.** Our proof combines Proposition 2.1 and [IKMZ19, Section 4] to reduce to an elementary computation for complement of real hyperplane arrangements.

Recall that  $\Delta \subset \mathbb{R}^n$  is a fan defining a non-singular projective toric variety  $\text{Tor}_{\mathbb{C}}(\Delta)$ , and that  $f : X \rightarrow D^*$  is a non-singular real analytic family of algebraic subvarieties of  $\text{Tor}_{\mathbb{C}}(\Delta)$  admitting a  $\mathbb{Q}$ -non-singular tropical limit  $V$  in  $\text{Tor}_{\mathbb{T}}(\Delta)$ . Given torus orbits  $\mathcal{O}_{\mathbb{C}}$  and  $\mathcal{O}_{\mathbb{T}}$  of  $\text{Tor}_{\mathbb{C}}(\Delta)$  and  $\text{Tor}_{\mathbb{T}}(\Delta)$  respectively corresponding to the same cone of  $\Delta$ , the intersection of  $V \cap \mathcal{O}_{\mathbb{T}}$  is the tropical limit of  $X \cap (\mathcal{O}_{\mathbb{C}} \times D^*)$ . Since both  $\sigma$  and  $\chi^c$  are motivic invariants, it is enough to prove Theorem 1.2 in the case when  $\delta = \{0\}$ , that is to say when  $\text{Tor}_{\mathbb{C}}(\delta) = (\mathbb{C}^*)^n$ . In what follows we define  $X^o = X \cap ((\mathbb{C}^*)^n \times D^*)$  and  $V^o = V \cap \mathbb{R}^n$ .

By [IKMZ19, Proposition 51] (see also [HK12, Proposition 2.3] and [KS12, Lemma 7.9] in a slightly different realm), after shrinking  $D$  if necessary, and after a base change  $t \mapsto \pm t^d$ , one can extend the real family  $X^o$  over 0 to a real semi-stable degeneration  $f : \overline{X}^o \rightarrow D$  such that any irreducible component of  $E_I^o$  of the central fiber  $\overline{X}_0^o$  is isomorphic to the complement of a hyperplane

arrangement in some  $\mathbb{C}P^m$ . When  $d$  is even, the sign in the base change is chosen so that the real fiber  $X_{t_0}$  lifts to a real fiber (i.e. it is the sign of  $t_0$ ). Since the complex conjugation on  $(\mathbb{C}^*)^n$  induces, via the tropicalization procedure, the identity map on  $\mathbb{R}^n$ , the semi-stable degeneration  $f : \overline{X}^\circ \rightarrow D$  is totally real. In particular, each irreducible component of  $E_I^\circ$  is the complement of a real hyperplane arrangement in some  $\mathbb{C}P^m$  equipped with the standard complex conjugation.

Hence, thanks to Proposition 2.1, the proof of Theorem 1.2 reduces to verify that these latter complements satisfy  $\chi = \sigma$ . This is easily done by a double induction on  $m$  and the number  $k$  of hyperplanes in the arrangement.

(1) The case  $m = 0$  holds trivially.

(2) Assume that this is true for  $m - 1$  and any  $k$ , and let us prove by induction on  $k$  that it is also true for  $m$  and any  $k$ .

(a) One checks easily that the case  $k = 0$  holds, see Example 1.1.

(b) By definition, the complement  $\mathcal{A}$  of  $k$  real hyperplanes in  $\mathbb{C}P^m$  is obtained by removing the complement  $\mathcal{A}''$  of  $k - 1$  real hyperplanes in  $\mathbb{C}P^{m-1}$  to the complement  $\mathcal{A}'$  of  $k - 1$  real hyperplanes in  $\mathbb{C}P^m$ . By induction and the scissor relation, we have

$$\chi^c(\mathbb{R}\mathcal{A}) = \chi^c(\mathbb{R}\mathcal{A}') - \chi^c(\mathbb{R}\mathcal{A}'') = \sigma^c(\mathbb{C}\mathcal{A}') - \sigma^c(\mathbb{C}\mathcal{A}'') = \sigma^c(\mathbb{C}\mathcal{A}).$$

Now the proof of Theorem 1.2 is complete.  $\square$

### 3. FURTHER COMMENTS

We end this note with a couple of remarks. First, dropping compactness or  $M$ -condition one constructs easily real algebraic varieties which do not satisfy  $\chi = \sigma$ . For example  $X = \mathbb{C}^*$  equipped with the real structure  $\tau(z) = \frac{1}{z}$  is an  $M$ -curve and satisfies

$$\chi^c(\mathbb{R}X) = \chi(S^1) = 0 \neq -2 = \sigma(\mathbb{C}^*),$$

and a quadric ellipsoid  $X$  in  $\mathbb{C}P^3$  satisfies

$$\chi(\mathbb{R}X) = 2 \neq 0 = \sigma(\mathbb{C}X).$$

More generally, Gudkov-Kharlamov-Krakhnov Congruence [DK00, 2.7.1] implies that a non-singular compact  $(M - 1)$ -variety<sup>2</sup> cannot satisfy  $\chi = \sigma$ .

Lastly, Viro Conjecture for  $M$ -surfaces states that such a projective and simply connected  $M$ -surface  $X$  satisfies

$$\beta_1(\mathbb{R}X) \leq h^{1,1}(\mathbb{C}X),$$

which is equivalent in this case to the inequality

$$\chi^c(\mathbb{R}X) \geq \sigma(\mathbb{C}X).$$

I do not know any  $M$ -variety<sup>3</sup>, projective or not, singular or not, which does not satisfy this latter inequality.

### REFERENCES

- [Arn17] C. Arnal. Patchwork combinatoire et topologie d'hypersurfaces algébriques réelles. Master Thesis, 2017.
- [BB07] B. Bertrand and F. Bihan. Euler characteristic of real nondegenerate tropical complete intersections. arXiv:0710.1222, 2007.
- [Ber10] B. Bertrand. Euler characteristic of primitive  $T$ -hypersurfaces and maximal surfaces. *J. Inst. Math. Jussieu*, 9(1):1–27, 2010.
- [BIMS15] E. Brugallé, I. Itenberg, G. Mikhalkin, and K. Shaw. Brief introduction to tropical geometry. In *Proceedings of the Gökova Geometry-Topology Conference 2014*, pages 1–75. Gökova Geometry/Topology Conference (GGT), Gökova, 2015.

<sup>2</sup>Unfortunately, I do not know much more  $(M - 1)$  than  $M$ -varieties.

<sup>3</sup>Go back to footnote 1.

- [Bit05] F. Bittner. On motivic zeta functions and the motivic nearby fiber. *Math. Z.*, 249(1):63–83, 2005.
- [DK00] A. I. Degtyarev and V. M. Kharlamov. Topological properties of real algebraic varieties: Rokhlin’s way. *Russian Math. Surveys*, 55(4):735–814, 2000.
- [DL01] J. Denef and F. Loeser. Geometry on arc spaces of algebraic varieties. In *European Congress of Mathematics, Vol. I (Barcelona, 2000)*, volume 201 of *Progr. Math.*, pages 327–348. Birkhäuser, Basel, 2001.
- [Ful93] W. Fulton. *Introduction to toric varieties*, volume 131 of *Ann. Math. Studies*. Princeton Univ. Press., 1993.
- [HK12] D. Helm and E. Katz. Monodromy filtrations and the topology of tropical varieties. *Canad. J. Math.*, 64(4):845–868, 2012.
- [IKMZ19] I. Itenberg, L. Katzarkov, G. Mikhalkin, and I. Zharkov. Tropical homology. *Math. Ann.*, 374(1-2):963–1006, 2019.
- [IS03] I. Itenberg and E. Shustin. Viro theorem and topology of real and complex combinatorial hypersurfaces. *Israel J. Math.*, 133:189–238, 2003.
- [Ite97] I. Itenberg. Topology of real algebraic  $T$ -surfaces. volume 10, pages 131–152. 1997. Real algebraic and analytic geometry (Segovia, 1995).
- [KS12] E. Katz and A. Stapledon. Tropical geometry and the motivic nearby fiber. *Compos. Math.*, 148(1):269–294, 2012.
- [KS16] E. Katz and A. Stapledon. Tropical geometry, the motivic nearby fiber, and limit mixed Hodge numbers of hypersurfaces. *Res. Math. Sci.*, 3:Paper No. 10, 36, 2016.
- [Man17] F. Mangolte. *Variétés algébriques réelles*, volume 24 of *Cours Spécialisés*. Société Mathématique de France, Paris, 2017.
- [MR] G. Mikhalkin and J. Rau. Tropical geometry. <https://math.uniandes.edu.co/~j.rau/downloads/main.pdf>.
- [Pet10] C. Peters. *Motivic aspects of Hodge theory*, volume 92. New Delhi: Narosa Publishing House/Published for the Tata Institute of Fundamental Research, 2010.
- [PS08] C. Peters and J. Steenbrink. *Mixed Hodge structures*, volume 52 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2008.
- [RS18] A. Renaudineau and K. Shaw. Bounding the betti numbers of real hypersurfaces near the tropical limit. arXiv:1805.02030, 2018.
- [Voi02] C. Voisin. *Théorie de Hodge et géométrie algébrique complexe*, volume 10 of *Cours Spécialisés [Specialized Courses]*. Société Mathématique de France, Paris, 2002.

ERWAN BRUGALLÉ, UNIVERSITÉ DE NANTES, LABORATOIRE DE MATHÉMATIQUES JEAN LERAY, 2 RUE DE LA HOUSSINIÈRE, F-44322 NANTES CEDEX 3, FRANCE

*E-mail address:* [erwan.brugalle@math.cnrs.fr](mailto:erwan.brugalle@math.cnrs.fr)