

Homogenization of a nonlinear drift-diffusion system for multiple charged species in a porous medium

Apratim Bhattacharya*, Markus Gahn** and Maria Neuss-Radu*

Abstract

We consider a nonlinear drift-diffusion system for multiple charged species in a porous medium in 2D and 3D with periodic microstructure. The system consists of a transport equation for the concentration of the species and Poisson's equation for the electric potential. The diffusion terms depend nonlinearly on the concentrations. We consider non-homogeneous Neumann boundary condition for the electric potential. The aim is the rigorous derivation of an effective (homogenized) model in the limit when the scale parameter ϵ tends to zero. This is based on uniform *a priori* estimates for the solutions of the microscopic model. The crucial result is the uniform L^∞ -estimate for the concentration in space and time. This result exploits the fact that the system admits a nonnegative energy functional which decreases in time along the solutions of the system. By using weak and strong (two-scale) convergence properties of the microscopic solutions, effective models are derived in the limit $\epsilon \rightarrow 0$ for different scalings of the microscopic model.

Keywords: Drift-diffusion model; nonlinear diffusion; multiple charged species; porous media; homogenization; two-scale convergence.

2020 Mathematics Subject Classification: 35B27, 35K59, 35Q92, 78A35

1 Introduction

The aim of this paper is the rigorous homogenization of the nonlinear drift-diffusion model (non-dimensional) (1.1)-(1.3), for a number of $P \in \mathbb{N}$ charged species with

*Department Mathematik, Friedrich-Alexander-Universität Erlangen-Nürnberg, Cauerstr. 11, 91058 Erlangen, Germany, apratim.bhattacharya@fau.de, maria.neuss-radu@math.fau.de

**Interdisciplinary Center for Scientific Computing, University of Heidelberg, Im Neuenheimer Feld 205, 69120 Heidelberg, Germany, markus.gahn@iwr.uni-heidelberg.de

concentrations $c_{i,\epsilon}$, $1 \leq i \leq P$, and the electric potential ϕ_ϵ , in a periodically perforated domain Ω_ϵ representing the fluid (pore) phase of a porous medium (see also Figure 1):

$$\partial_t c_{i,\epsilon}(t, x) + \nabla \cdot J_{i,\epsilon}(t, x) = 0 \quad \text{in } (0, T) \times \Omega_\epsilon, \quad (1.1a)$$

$$J_{i,\epsilon}(t, x) \cdot \nu_\epsilon = 0 \quad \text{on } (0, T) \times \partial\Omega_\epsilon, \quad (1.1b)$$

$$c_{i,\epsilon}(0, x) = c_i^0(x) \quad \text{in } \Omega_\epsilon. \quad (1.1c)$$

$$-\epsilon^\alpha \Delta \phi_\epsilon(t, x) = \sum_{i=1}^P z_i c_i(t, x) \quad \text{in } (0, T) \times \Omega_\epsilon, \quad (1.1d)$$

$$\epsilon^\alpha \nabla \phi_\epsilon(t, x) \cdot \nu_\epsilon = \xi_\epsilon(x) \quad \text{on } (0, T) \times \partial\Omega_\epsilon, \quad (1.1e)$$

where the total flux $J_{i,\epsilon}$ of the i -th charged species is given by

$$J_{i,\epsilon}(t, x) = - \left(D_i \nabla h_p(c_{i,\epsilon}) + \epsilon^\beta D_i z_i c_{i,\epsilon}(t, x) \nabla \phi_\epsilon(t, x) \right). \quad (1.2)$$

Here $(0, T)$ is a time interval, whereas $D_i > 0$ and $z_i \in \mathbb{Z}$ denote the (scaled) diffusivity and charge number of the i -th species respectively. The (scaled) permittivity of the medium and the (scaled) mobility of the i -th charged species are given by ϵ^α and $D_i \epsilon^\beta$ respectively, where $\alpha, \beta \in \mathbb{R}$ with $\alpha \leq \beta$. The scale parameter $\epsilon > 0$ describes the length of the period of the porous microstructure, and it is also proportional to the radius of the pores. ν_ϵ represents the outward unit normal vector to the boundary $\partial\Omega_\epsilon$. The function h_p is defined by

$$h_p(r) = r + \eta r^p, \quad r \geq 0, \quad \eta \in (0, \infty), \quad p \in [4, \infty). \quad (1.3)$$

Equation (1.1a) models the transport of the charged species due to nonlinear diffusion and electromigration, in a domain with an impermeable boundary modeled by the no-flux boundary condition (1.1b), and with initial concentrations given in (1.1c). The electric potential is induced by the charges of the species and is given as the solution of Poisson's equation (1.1d) subject to the non-homogeneous Neumann boundary condition (1.1e). Physically, (1.1e) models a charged boundary and ξ_ϵ represents the surface charge density. The parameters $\alpha, \beta \in \mathbb{R}$ (obtained from a non-dimensionalization procedure) allow to consider different scalings in our microscopic model corresponding to different settings and applications.

Drift-diffusion systems of the form (1.1)-(1.2) arise, e.g., in the mathematical modeling of semiconductors and of transport of charged particles (ions) in solutions (electrophoresis). In semiconductor modeling, see e.g. [17,20,30], the case of two species is relevant (the electrons with valence -1 and the holes with valence $+1$) while in electrophoresis multiple types of charged particles (like e.g. ions) with different charge numbers are transported in solution under the influence of an electric field, see e.g. [12,24]. In these applications, the function h_p in (1.2) can be linear or nonlinear, the linear case represents the classical drift-diffusion (Poisson-Nernst-Planck) system. For a discussion about different (nonlinear) shapes of h_p , see e.g. the introduction in [23]. The particular model with h_p given by (1.3) plays an important role in the approximation of the models used in applications. For example, in [5] (inspired by contributions from




Figure 1: The standard cell $Y = Y^f \cup \overline{Y^s}$ (left), and the porous medium Ω with the fluid (pore) part Ω_ϵ (right).

[18]) this model is used as a regularization of the classical PNP system in order to show existence of the latter for multiple species and space dimension three and higher. Furthermore, in [23], a similar model is used to approximate a drift-diffusion system with a (possible) degenerate nonlinear diffusion term by a nondegenerate system.

In many applications, e.g., from geosciences, biology or biomedicine, electrophoretic processes take place in porous media. Due to the complex microstructure of the medium numerical simulations of microscopic models at the fine (pore) scale are very expensive. Therefore, effective (homogenized) approximations of the solutions, obtained in the scale limit $\epsilon \rightarrow 0$, are highly demanded. Homogenization results for drift-diffusion models in porous media usually deal with the Poisson-Nernst-Planck (PNP) system or with models consisting of the PNP system coupled with the Stokes system (SPNP). Formal upscaling of PNP system using formal asymptotic expansions is given e.g. in [4,33]. Rigorous homogenization results have been obtained for the SPNP system in the case of two charged species with opposite valences e.g. in [38,39]. Furthermore, in [3] a stationary and linearized SPNP for multiple charged species was homogenized. In the recent paper [26], a generalized PNP problem in a two-phase medium, with transmission conditions at the microscopic interface has been homogenized. Here multiple charged species were considered, however under the constraint of *total mass balance*. Corrector results related to the model from [38] were considered in [25]. Homogenization results for the PNP system for multiple charged species without additional constraints on the solutions (e.g. *total mass balance*) are not available in the literature so far. This might be related to the fact that existence results in dimension three for such problems are only available in very weak function spaces, see e.g. [5]. In contrast, problem (1.1)-(1.3) has (for fixed values

of the parameters $\epsilon > 0$ and $\eta > 0$) weak solutions with good regularity properties, especially for the potential, see Proposition 2.1 below for details. Thus, this problem is more suitable (than the classical PNP problem) for the investigation of the behavior of the solutions in the limit $\epsilon \rightarrow 0$, and for the derivation of an effective (macroscopic) approximation in case of multiple charged species without imposing further constraints.

From the point of view of multiscale analysis and homogenization the problem (1.1)-(1.3) (for fixed values of $\eta > 0$) is of independent interest due to the nonlinearity in the diffusion term and the strong nonlinear coupling via the drift-term. In the literature there are few contributions dealing with the homogenization of quasi-linear problems and nonlinear diffusion, we mention here, e.g., [7,16,28]. In [7] the rigorous upscaling of a two phase flow in a perforated domain was performed while in [28] fluid flow in an unsaturated porous medium containing a fracture was upscaled. In [16] a reaction-diffusion problem with nonlinear diffusion was homogenized in a domain consisting of two bulk regions separated by a thin layer with periodic microstructure. Further, problems including monotone operators were treated in [2] for the stationary case, and in [11] for nonstationary problems. As usually in the homogenization of nonlinear problems, the main challenge of our study is to derive *a priori* estimates of the solutions uniformly with respect to the scale parameter ϵ , which allow to pass to the limit $\epsilon \rightarrow 0$, especially in the nonlinear terms. In particular, an L^∞ -estimate in time and space of the concentration vector $(c_{i,\epsilon})_{1 \leq i \leq P}$ is needed. To obtain this estimate, we make use of an energy functional associated to the system, which allows us to estimate the $L^\infty L^p$ -norm of the concentrations. The energy functional is inspired from [5], where the existence of solutions of problem (1.1)-(1.3) with Robin-type boundary conditions for the potential was shown. The estimate of the concentration vector then allows to prove an $L^\infty H^1$ -estimate of the potential ϕ_ϵ and eventually, to obtain the $L^\infty L^\infty$ -estimate for the concentrations by using a classical result from [27]. A key point in our proof is to avoid the use of higher (than H^1) norms for the potential. Based on the *a priori* estimates effective (homogenized) problems are derived. It turns out that different results are obtained for different values of the parameters α and β . Whereas for $\alpha = \beta$ a drift-diffusion problem with homogenized coefficients (similar to the microscopic problem) is obtained, for $\alpha < \beta$ the homogenized problem reduces to a system of diffusion equations (with homogenized nonlinear diffusion) for the species concentrations, one way coupled to the homogenized Poisson's equation. To our knowledge, the result obtained in this paper is the first one providing the rigorous homogenization of a three dimensional (nonlinear) electro-diffusion system for multiple charged species (avoiding further restrictions such as above).

This paper is organized as follows. In Section 2, the microscopic model is introduced and existence of solutions is proved. In Section 3, estimates for the microscopic solutions uniformly with respect to ϵ are derived. These are the basis for the (two-scale) convergence results for the microscopic solutions proved in Section 4. Also in this section the homogenized drift-diffusion model is derived. The paper is concluded with a discussion and outlook in Section 5 and the appendix A consisting of auxiliary results.

2 The microscopic model

We consider a porous medium occupying a bounded and connected domain $\Omega \subset \mathbb{R}^n$, $n = 2, 3$, with $\partial\Omega$ of class C^3 . The medium has a periodic microstructure, generated with the help of the scaled standard periodicity cell $Y = (0, 1)^n$ which consists of a solid part Y^s and a fluid or pore part Y^f . We assume that Y^s is an open set such that $\overline{Y^s} \subset Y$, and that the boundary $\Gamma := \partial Y^s$ is of class C^3 . Furthermore, let $Y^f = Y \setminus \overline{Y^s}$, see also Figure 1 (left). For $k \in \mathbb{Z}^n$, let $Y_k := Y + k$ and $\Gamma_k := \Gamma + k$. Furthermore, for $j = f, s$, set $Y_k^j := Y^j + k$.

For a given (small) scale parameter $\epsilon > 0$, let $I_\epsilon = \{k \in \mathbb{Z}^n \mid \epsilon Y_k \subset \Omega\}$. We define the microscopic domain Ω_ϵ representing the pore part of the porous medium by

$$\Omega_\epsilon = \Omega \setminus \bigcup_{k \in I_\epsilon} \epsilon \overline{Y_k^s},$$

see also Figure 1 (right). We remark that the boundary of Ω_ϵ consists of two disjoint parts

$$\partial\Omega_\epsilon = \Gamma_\epsilon \cup \partial\Omega,$$

where $\Gamma_\epsilon := \bigcup_{k \in I_\epsilon} \epsilon \Gamma_k$ denotes the boundary of the microscopic solid grains. We also note that the domain Ω_ϵ is connected with boundary $\partial\Omega_\epsilon$ of class C^3 .

The aim of the paper is the rigorous homogenization of the nonlinear drift-diffusion model (1.1)-(1.3), i.e., the derivation of a macroscopic model in the scale limit $\epsilon \rightarrow 0$.

2.1 Assumptions on the data

- (A1) For the diffusion coefficients we assume $D_i > 0$, $1 \leq i \leq P$. Furthermore, let $T > 0$ be a fixed time point.
(A2) The surface charge density is given by

$$\xi_\epsilon(x) = \begin{cases} \epsilon \xi_1(x, \frac{x}{\epsilon}) & \text{if } x \in \Gamma_\epsilon, \\ \xi_2(x) & \text{if } x \in \partial\Omega, \end{cases}$$

where $\xi_1 \in C^2(\overline{\Omega} \times \overline{\Gamma})$, $\xi_1(x, y)$ periodically extended with respect to y with period Y , and $\xi_2 \in C^2(\partial\Omega)$. Let us denote

$$\xi^* := \max \left\{ \max_{(x,y) \in (\overline{\Omega} \times \overline{\Gamma})} |\xi_1(x, y)|, \max_{x \in \partial\Omega} |\xi_2(x)| \right\}.$$

- (A3) For the initial concentrations we assume $c_i^0 \in C^2(\overline{\Omega})$ with $c_i^0 \geq 0$ for $i = 1, \dots, P$.
(A4) We assume the following compatibility condition:

$$\int_{\Omega_\epsilon} \sum_{i=1}^P z_i c_i^0(x) dx + \int_{\partial\Omega_\epsilon} \xi_\epsilon(x) dS(x) = 0. \quad (2.1)$$

2.2 Variational formulation of the microscopic problem

The variational formulation of the problem (1.1)-(1.3) is given as follows. Find non-negative functions $c_{i,\epsilon} \in L^\infty((0, T) \times \Omega_\epsilon) \cap L^2(0, T; H^1(\Omega_\epsilon))$ with $\partial_t c_{i,\epsilon} \in L^2(0, T; H^1(\Omega_\epsilon)')$ and $\phi_\epsilon \in L^\infty(0, T; W^{2,6}(\Omega_\epsilon))$ with $\int_{\Omega_\epsilon} \phi_\epsilon(t, x) dx = 0$ for all $t \in [0, T]$, satisfying

$$\langle \partial_t c_{i,\epsilon}, \psi \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} + \int_{\Omega_\epsilon} (D_i \nabla h_p(c_{i,\epsilon}) + \epsilon^\beta D_i z_i c_{i,\epsilon} \nabla \phi_\epsilon) \nabla \psi dx = 0, \quad (2.2)$$

for all $\psi \in H^1(\Omega_\epsilon)$ and almost every $t \in (0, T)$, together with the initial condition

$$c_{i,\epsilon}(0) = c_i^0 \quad \text{in } L^2(\Omega_\epsilon), \quad (2.3)$$

and

$$\epsilon^\alpha \int_{\Omega_\epsilon} \nabla \phi_\epsilon(t) \nabla v dx = \int_{\Omega_\epsilon} \sum_{i=1}^P z_i c_{i,\epsilon}(t) v dx + \int_{\partial \Omega_\epsilon} \xi_\epsilon v dS(x), \quad (2.4)$$

for all $v \in H^1(\Omega_\epsilon)$, and all $t \in [0, T]$.

Remark 1. We emphasize that due to the regularity properties of the variational solution, the Poisson's problem for the electric potential ϕ_ϵ holds even pointwise almost everywhere in x and t .

Remark 2. In order to keep the notation as clear as possible, we skip the parameters α, β, η and p in the labeling of the solution $(c_{i,\epsilon}, \phi_\epsilon)$.

2.3 Existence for the microscopic model

In this section we prove the existence of solutions for the microscopic model (1.1)-(1.3). The proof follows from similar arguments as in [5, Lemma 5], where a similar model was considered, however, with Poisson's equation for the potential subject to Robin-type boundary condition (instead of purely Neumann condition used in our model). Therefore, we mainly highlight the new arguments needed to prove the existence result for our setting. Note that in the proof we use energy estimates from Proposition 3.1 in the next section.

Proposition 2.1. *Suppose the assumptions (A1) – (A4) are satisfied. Then there exists a solution $(c_{i,\epsilon}, \phi_\epsilon), i \in \{1, \dots, P\}$ of the variational problem (2.2)-(2.4). If $n = 2$, additionally, we have $\phi_\epsilon \in L^\infty(0, T; W^{2,q}(\Omega_\epsilon))$, for any $q \in [1, \infty)$.*

Proof. The existence is obtained using Schaefer's fixed point theorem. Consider any $\phi_\epsilon \in L^\infty(0, T; W^{1,\infty}(\Omega_\epsilon))$. For such a ϕ_ϵ there exists a unique non-negative $c_{i,\epsilon} \in L^\infty((0, T) \times \Omega_\epsilon) \cap L^2(0, T; H^1(\Omega_\epsilon))$ with $\partial_t c_{i,\epsilon} \in L^2(0, T; H^1(\Omega_\epsilon)')$ satisfying (2.2) and (2.3), see [5]. Note that here the C^3 regularity of the domain Ω_ϵ is used. Again for these $c_{i,\epsilon}, 1 \leq i \leq P$, there exists a unique $\hat{\phi}_\epsilon \in L^\infty(0, T; H^1(\Omega_\epsilon))$ with

$\int_{\Omega_\epsilon} \hat{\phi}_\epsilon(t, x) dx = 0$, for all $t \in [0, T]$ such that $\hat{\phi}_\epsilon$ satisfies Poisson's equation (2.4), see [10, Theorem 4.22]. We note that the compatibility condition

$$\int_{\Omega_\epsilon} \sum_{i=1}^P z_i c_{i,\epsilon}(t, x) dx + \int_{\partial\Omega_\epsilon} \xi_\epsilon dS(x) = 0, \quad \text{for all } t \in [0, T] \quad (2.5)$$

required in the existence proof for $\hat{\phi}_\epsilon$ is satisfied due to the fact that testing equation (2.2) with $\psi = 1$, we obtain

$$\frac{d}{dt} \int_{\Omega_\epsilon} c_{i,\epsilon}(t, x) dx = 0, \quad \text{for a.e. } t \in (0, T),$$

which together with assumption (A4) yields (2.5).

Let us now consider the mapping

$$\tau : L^\infty(0, T; W^{1,\infty}(\Omega_\epsilon)) \rightarrow L^\infty(0, T; W^{1,\infty}(\Omega_\epsilon)), \quad \tau(\phi_\epsilon) = \hat{\phi}_\epsilon. \quad (2.6)$$

Next, we show that $\hat{\phi}_\epsilon \in L^\infty(0, T; W^{1,\infty}(\Omega_\epsilon))$ so that τ becomes well-defined. Here again, we need to argue differently from [5]. Since $c_{i,\epsilon} \in L^\infty(0, T; L^2(\Omega_\epsilon))$, elliptic regularity results imply that $\hat{\phi}_\epsilon \in L^\infty(0, T; H^2(\Omega_\epsilon))$, see [32, Theorem 4, p. 217]. Consequently, using $c_{i,\epsilon} \in L^\infty((0, T) \times \Omega_\epsilon)$ and [21, Lemma 2.4.1.4], we obtain for almost every $t \in (0, T)$ that $\hat{\phi}_\epsilon(t) \in W^{2,6}(\Omega_\epsilon)$, if $n = 3$, and $\hat{\phi}_\epsilon(t) \in W^{2,q}(\Omega_\epsilon)$ for any $q \in [1, \infty)$, if $n = 2$. The L^∞ -regularity with respect to time, namely $\hat{\phi}_\epsilon \in L^\infty(0, T; W^{2,6}(\Omega_\epsilon))$, if $n = 3$, and $\hat{\phi}_\epsilon \in L^\infty(0, T; W^{2,q}(\Omega_\epsilon))$ for any $q \in [1, \infty)$, if $n = 2$, is then obtained from [21, Theorem 2.4.1.3]. Thus, by Sobolev embedding theorem, we have $\hat{\phi}_\epsilon \in L^\infty(0, T; W^{1,\infty}(\Omega_\epsilon))$.

Note that, a solution of the microscopic problem (2.2)-(2.4) is a fixed point of τ . Thus, in the following, we prove the existence of such a fixed point by using Schaefer's fixed point theorem (see e.g., [13, Theorem 4, p. 504]). Since $W^{2,6}(\Omega_\epsilon)$ is compactly embedded in $W^{1,\infty}(\Omega_\epsilon)$, the continuity and compactness of τ follow by similar arguments as in [5, Lemma 5]. It remains to prove that the set

$$\{\phi_\epsilon \in L^\infty(0, T; W^{1,\infty}(\Omega_\epsilon)) : \phi_\epsilon = \lambda\tau(\phi_\epsilon) \text{ for some } \lambda \in [0, 1]\} \quad (2.7)$$

is bounded. Here we again need alternative arguments to [5]. Suppose, $\tau(\phi_\epsilon) = \frac{\phi_\epsilon}{\lambda}$ for some $\lambda \in (0, 1]$. Then ϕ_ϵ is a solution of the potential equation (3.3) in Proposition 3.1. Firstly, let us show that $\tau(\phi_\epsilon)$ is bounded in $L^\infty(0, T; W^{2,4}(\Omega_\epsilon))$ independently of λ . By Proposition 3.1 and Corollary 3.2, we can bound the $L^\infty(0, T; L^p(\Omega_\epsilon))$ norm of $c_{i,\epsilon}$ independently of λ . Since by assumption, see (1.3), we have $p \geq 4$, it follows that $c_{i,\epsilon}$ is bounded in $L^\infty(0, T; L^4(\Omega_\epsilon))$ independent of λ . Hence, there exists a constant $C_1 > 0$ independent of λ such that

$$\left\| \sum_{i=1}^P z_i c_{i,\epsilon} \right\|_{L^\infty(0, T; L^4(\Omega_\epsilon))} < C_1. \quad (2.8)$$

Testing with $\tau(\phi_\epsilon)(t)$ in the weak formulation satisfied by $\tau(\phi_\epsilon)(t)$, we get

$$\begin{aligned} \epsilon^\alpha \|\nabla \tau(\phi_\epsilon)(t)\|_{L^2(\Omega_\epsilon)}^2 &\leq \left\| \sum_{i=1}^P z_i c_{i,\epsilon}(t) \right\|_{L^2(\Omega_\epsilon)} \|\tau(\phi_\epsilon)(t)\|_{L^2(\Omega_\epsilon)} \\ &\quad + \|\xi_\epsilon\|_{L^2(\partial\Omega_\epsilon)} \|\tau(\phi_\epsilon)(t)\|_{L^2(\partial\Omega_\epsilon)} \\ &\leq C_2 \left(\left\| \sum_{i=1}^P z_i c_{i,\epsilon}(t) \right\|_{L^2(\Omega_\epsilon)} + \|\xi_\epsilon\|_{L^2(\partial\Omega_\epsilon)} \right) \|\nabla \tau(\phi_\epsilon)(t)\|_{L^2(\Omega_\epsilon)}. \end{aligned}$$

In the last inequality we have used $\int_{\Omega_\epsilon} \tau(\phi_\epsilon)(t) = 0$. Consequently, we have

$$\|\tau(\phi_\epsilon)(t)\|_{H^1(\Omega_\epsilon)} \leq C_3 \left(\left\| \sum_{i=1}^P z_i c_{i,\epsilon}(t) \right\|_{L^2(\Omega_\epsilon)} + \|\xi_\epsilon\|_{L^2(\partial\Omega_\epsilon)} \right). \quad (2.9)$$

Using (2.8) in (2.9) we obtain that, $\|\tau(\phi_\epsilon)\|_{L^\infty(0,T;H^1(\Omega_\epsilon))}$ is bounded independent of λ . Due to the embedding $H^1(\Omega_\epsilon) \subset L^4(\Omega_\epsilon)$, we have for some $C_4 > 0$ independent of λ :

$$\|\tau(\phi_\epsilon)\|_{L^\infty(0,T;L^4(\Omega_\epsilon))} < C_4. \quad (2.10)$$

Again, with the help of [21, Theorem 2.4.1.3], for all $t \in [0, T]$, we obtain

$$\|\tau(\phi_\epsilon)(t)\|_{W^{2,4}(\Omega_\epsilon)} \leq C_5 \left(\left\| \epsilon^{-\alpha} \sum_{i=1}^P z_i c_{i,\epsilon}(t) + \mu \tau(\phi_\epsilon)(t) \right\|_{L^4(\Omega_\epsilon)} + \epsilon^{-\alpha} \|\xi_\epsilon\|_{W^{1-\frac{1}{4},4}(\partial\Omega_\epsilon)} \right),$$

for some $C_5, \mu > 0$ independent of λ . This implies

$$\|\tau(\phi_\epsilon)(t)\|_{W^{2,4}(\Omega_\epsilon)} \leq C_6 \left(\left\| \sum_{i=1}^P z_i c_{i,\epsilon}(t) \right\|_{L^4(\Omega_\epsilon)} + \|\tau(\phi_\epsilon)(t)\|_{L^4(\Omega_\epsilon)} + \|\xi_\epsilon\|_{W^{1-\frac{1}{4},4}(\partial\Omega_\epsilon)} \right), \quad (2.11)$$

for some $C_6 > 0$ independent of λ . Finally using (2.8), (2.10) in (2.11) we obtain the desired result that $\tau(\phi_\epsilon)$ is bounded in $L^\infty(0, T; W^{2,4}(\Omega_\epsilon))$ independent of λ .

Consequently, the embedding $W^{2,4}(\Omega_\epsilon) \subset W^{1,\infty}(\Omega_\epsilon)$, implies that $\tau(\phi_\epsilon)$ is bounded in $L^\infty(0, T; W^{1,\infty}(\Omega_\epsilon))$ independent of λ . Since $\phi_\epsilon = \lambda \tau(\phi_\epsilon)$, for some $\lambda \in (0, 1]$, we conclude that ϕ_ϵ is bounded in $L^\infty(0, T; W^{1,\infty}(\Omega_\epsilon))$ independent of λ and the proof is complete. \square

3 Uniform estimates for the microscopic solutions

In this section, we prove the uniform estimates for the microscopic solutions. To obtain compactness results which allow to pass to the limit in the nonlinear terms, we prove an

L^∞ -estimate for the concentrations, uniformly with respect to ϵ . The first step towards this result is to show that the concentrations are uniformly bounded in $L^\infty((0, T), L^p(\Omega_\epsilon))$. (Note that in the whole paper $p \in [4, \infty)$ is a fixed index entering the definition (1.3) of the nonlinear diffusion function h_p .) For this we use the following energy functional associated to our system, see also [5,6]:

$$V_\epsilon(t) = \frac{1}{2\lambda} \epsilon^{\alpha+\beta} \int_{\Omega_\epsilon} |\nabla \phi_\epsilon|^2 dx + \sum_{i=1}^P \int_{\Omega_\epsilon} \Psi(c_{i,\epsilon}) dx, \quad \text{for a.e. } t \in (0, T), \quad (3.1)$$

where $\lambda \in (0, 1]$, and

$$\Psi(r) = \left(r \log r - r + 1 + \frac{\eta}{p-1} r^p \right), \quad \text{for } r \geq 0. \quad (3.2)$$

Proposition 3.1. *Let $\lambda \in (0, 1]$. Consider non-negative functions $c_{i,\epsilon} \in L^\infty((0, T) \times \Omega_\epsilon) \cap L^2(0, T; H^1(\Omega_\epsilon))$ with $\partial_t c_{i,\epsilon} \in L^2(0, T; H^1(\Omega_\epsilon)')$, and $\phi_\epsilon \in L^\infty(0, T; W^{2,6}(\Omega_\epsilon))$ with $\int_{\Omega_\epsilon} \phi_\epsilon(t, x) dx = 0$ for all $t \in [0, T]$ that satisfy (2.2)-(2.3) and the following equations:*

$$-\epsilon^\alpha \Delta \phi_\epsilon(t, x) = \lambda \sum_{i=1}^P z_i c_{i,\epsilon}(t, x) \quad \forall t \in [0, T] \text{ and a.e. } x \in \Omega_\epsilon, \quad (3.3)$$

$$\epsilon^\alpha \nabla \phi_\epsilon(t, x) \cdot \nu_\epsilon = \lambda \xi_\epsilon(x) \quad \forall t \in [0, T] \text{ and a.e. } x \in \partial\Omega_\epsilon. \quad (3.4)$$

Then

$$\frac{d}{dt} V_\epsilon(t) \leq 0, \quad \text{for a.e. } t \in (0, T). \quad (3.5)$$

In particular, we have

$$V_\epsilon(t) \leq C \left(1 + \epsilon^{-\alpha+\beta} \right), \quad \text{for all } t \in [0, T], \quad (3.6)$$

with some constant $C > 0$ independent of ϵ and λ .

Proof. Using $\langle \partial_t c_{i,\epsilon}, 1 \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} = 0$, we have

$$\frac{d}{dt} V_\epsilon(t) = \frac{d}{dt} \left(\frac{1}{2\lambda} \epsilon^{\alpha+\beta} \int_{\Omega_\epsilon} |\nabla \phi_\epsilon|^2 \right) + \frac{d}{dt} \left(\sum_{i=1}^P \int_{\Omega_\epsilon} c_{i,\epsilon} \log c_{i,\epsilon} \right) + \frac{d}{dt} \left(\sum_{i=1}^P \int_{\Omega_\epsilon} \frac{\eta}{p-1} c_{i,\epsilon}^p \right), \quad (3.7)$$

for almost every $t \in (0, T)$, provided that each of the three time derivatives given in the right hand side of (3.7) exists. Next we show that this is the case. Suppose δ_m is a sequence of positive numbers tending to zero as m tends to infinity. Then from Lemma A.2, we have for all $m \in \mathbb{N}$ and $t_1 \in [0, T]$,

$$\int_{\Omega_\epsilon} [c_{i,\epsilon}(t_1) + \delta_m] \log [c_{i,\epsilon}(t_1) + \delta_m] dx - \int_{\Omega_\epsilon} [c_{i,\epsilon}(0) + \delta_m] \log [c_{i,\epsilon}(0) + \delta_m] dx$$

$$= \int_0^{t_1} \langle \partial_t c_{i,\epsilon}, \log(c_{i,\epsilon} + \delta_m) \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} dt. \quad (3.8)$$

Testing (2.2) with $\log(c_{i,\epsilon} + \delta_m)$ we see that the right hand side of (3.8) is equal to

$$\begin{aligned} & \int_0^{t_1} \int_{\Omega_\epsilon} J_{i,\epsilon} \cdot \frac{\nabla c_{i,\epsilon}}{c_{i,\epsilon} + \delta_m} dx dt \\ &= \int_0^{t_1} \int_{\Omega_\epsilon} \frac{1 - h'_p(c_{i,\epsilon})}{c_{i,\epsilon} + \delta_m} \nabla c_{i,\epsilon} \cdot J_{i,\epsilon} + \frac{\nabla h_p(c_{i,\epsilon})}{c_{i,\epsilon} + \delta_m} \cdot J_{i,\epsilon} dx dt \\ &= \int_0^{t_1} \int_{\Omega_\epsilon} \frac{1 - h'_p(c_{i,\epsilon})}{c_{i,\epsilon} + \delta_m} \nabla c_{i,\epsilon} \cdot J_{i,\epsilon} + \frac{D_i \nabla h_p(c_{i,\epsilon}) + \epsilon^\beta D_i z_i c_{i,\epsilon} \nabla \phi_\epsilon}{D_i(c_{i,\epsilon} + \delta_m)} \cdot J_{i,\epsilon} dx dt \\ &\quad - \int_0^{t_1} \int_{\Omega_\epsilon} \frac{\epsilon^\beta z_i c_{i,\epsilon} \nabla \phi_\epsilon}{c_{i,\epsilon} + \delta_m} \cdot J_{i,\epsilon} dx dt \\ &= - \int_0^{t_1} \int_{\Omega_\epsilon} \frac{\eta p c_{i,\epsilon}^{p-1}}{c_{i,\epsilon} + \delta_m} \nabla c_{i,\epsilon} \cdot J_{i,\epsilon} - \frac{|J_{i,\epsilon}|^2}{D_i(c_{i,\epsilon} + \delta_m)} - \frac{\epsilon^\beta z_i c_{i,\epsilon} \nabla \phi_\epsilon}{c_{i,\epsilon} + \delta_m} \cdot J_{i,\epsilon} dx dt. \end{aligned}$$

Hence from (3.8), we get

$$\begin{aligned} & \int_{\Omega_\epsilon} [c_{i,\epsilon}(t_1) + \delta_m] \log[c_{i,\epsilon}(t_1) + \delta_m] - [c_{i,\epsilon}(0) + \delta_m] \log[c_{i,\epsilon}(0) + \delta_m] dx \\ &= - \int_0^{t_1} \int_{\Omega_\epsilon} \frac{\eta p c_{i,\epsilon}^{p-1}}{c_{i,\epsilon} + \delta_m} \nabla c_{i,\epsilon} \cdot J_{i,\epsilon} - \frac{|J_{i,\epsilon}|^2}{D_i(c_{i,\epsilon} + \delta_m)} - \frac{\epsilon^\beta z_i c_{i,\epsilon} \nabla \phi_\epsilon}{c_{i,\epsilon} + \delta_m} \cdot J_{i,\epsilon} dx dt. \quad (3.9) \end{aligned}$$

Again, using the monotone convergence theorem, we have

$$\lim_{m \rightarrow \infty} \int_0^{t_1} \int_{\Omega_\epsilon} \frac{|J_{i,\epsilon}|^2}{D_i(c_{i,\epsilon} + \delta_m)} dx dt = \int_0^{t_1} \int_{\Omega_\epsilon} \frac{|J_{i,\epsilon}|^2}{D_i c_{i,\epsilon}} dx dt. \quad (3.10)$$

However, this limit can be infinite. Next, we shall show that this is not the case.

The inequalities

$$x \log x \geq -1, \quad \forall x > 0 \text{ and};$$

$$x \log x \leq \begin{cases} 0 & \text{if } x \in (0, 1), \\ x^2 & \text{if } x \geq 1; \end{cases}$$

together give the following:

$$|(c_{i,\epsilon} + \delta_m) \log(c_{i,\epsilon} + \delta_m)| \leq 1 + (c_{i,\epsilon} + \delta_m)^2 \leq 1 + (c_{i,\epsilon} + 1)^2 \quad (\text{for large } m),$$

for all $t \in [0, T]$ and almost every $x \in \Omega_\epsilon$. For each t , we have $1 + (c_{i,\epsilon} + 1)^2 \in L^1(\Omega_\epsilon)$. This fact and the continuity of the function $x \mapsto x \log x$ for all $x \geq 0$ allow us to use the dominated convergence theorem to obtain the following:

$$\lim_{m \rightarrow \infty} \int_{\Omega_\epsilon} (c_{i,\epsilon} + \delta_m) \log(c_{i,\epsilon} + \delta_m) dx = \int_{\Omega_\epsilon} c_{i,\epsilon} \log c_{i,\epsilon} dx, \quad \forall t \in [0, T]. \quad (3.11)$$

Using (3.10), (3.11), the fact that

$$0 \leq \frac{c_{i,\epsilon}}{c_{i,\epsilon} + \delta_m} < 1; \text{ and as } m \rightarrow \infty, \frac{c_{i,\epsilon}}{c_{i,\epsilon} + \delta_m} \rightarrow \chi_{\{c_{i,\epsilon} \neq 0\}} \text{ a.e. in } (0, T) \times \Omega_\epsilon;$$

we pass to the limit in (3.9):

$$\begin{aligned} & \sum_{i=1}^P \int_{\Omega_\epsilon} c_{i,\epsilon}(t_1) \log c_{i,\epsilon}(t_1) - c_{i,\epsilon}(0) \log c_{i,\epsilon}(0) dx \\ &= \sum_{i=1}^P - \int_0^{t_1} \int_{\Omega_\epsilon} \eta p c_{i,\epsilon}^{p-2} \nabla c_{i,\epsilon} \cdot J_{i,\epsilon} - \frac{|J_{i,\epsilon}|^2}{D_i c_{i,\epsilon}} - \epsilon^\beta z_i \chi_{\{c_{i,\epsilon} \neq 0\}} \nabla \phi_\epsilon \cdot J_{i,\epsilon} dx dt, \\ & \qquad \qquad \qquad \forall t_1 \in [0, T]. \end{aligned} \quad (3.12)$$

We note from (3.12) that $\sum_{i=1}^P \frac{|J_{i,\epsilon}|^2}{D_i c_{i,\epsilon}} \in L^1((0, t_1) \times \Omega_\epsilon)$ and hence the limit in (3.10) is finite. Now from (3.12), we see that $\sum_{i=1}^P \int_{\Omega_\epsilon} c_{i,\epsilon} \log c_{i,\epsilon}$ is absolutely continuous on $[0, T]$ and

$$\begin{aligned} & \frac{d}{dt} \sum_{i=1}^P \int_{\Omega_\epsilon} c_{i,\epsilon} \log c_{i,\epsilon} dx \\ &= \sum_{i=1}^P - \int_{\Omega_\epsilon} \eta p c_{i,\epsilon}^{p-2} \nabla c_{i,\epsilon} \cdot J_{i,\epsilon} - \frac{|J_{i,\epsilon}|^2}{D_i c_{i,\epsilon}} - \epsilon^\beta z_i \chi_{\{c_{i,\epsilon} \neq 0\}} \nabla \phi_\epsilon \cdot J_{i,\epsilon} dx, \end{aligned} \quad (3.13)$$

for almost every $t \in (0, T)$.

Next, consider the first term of the right hand side of (3.13). Using (2.2) we have

$$\begin{aligned} & \sum_{i=1}^P \int_{\Omega_\epsilon} \eta p c_{i,\epsilon}^{p-2} \nabla c_{i,\epsilon} \cdot J_{i,\epsilon} dx = \sum_{i=1}^P \int_{\Omega_\epsilon} \nabla \left(\frac{\eta p}{p-1} c_{i,\epsilon}^{p-1} \right) \cdot J_{i,\epsilon} dx \\ &= \sum_{i=1}^P \left\langle \partial_t c_{i,\epsilon}, \frac{\eta p}{p-1} c_{i,\epsilon}^{p-1} \right\rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} = \sum_{i=1}^P \frac{d}{dt} \int_{\Omega_\epsilon} \frac{\eta}{p-1} c_{i,\epsilon}^p dx, \end{aligned} \quad (3.14)$$

where the last equality can be proven using approximation arguments similar to Lemma A.2.

Since $\nabla c_{i,\epsilon} = 0$ almost everywhere on the set $A := \{(t, x) \in (0, T) \times \Omega_\epsilon : c_{i,\epsilon}(t, x) = 0\}$, we see that $J_{i,\epsilon}$ vanishes on A . Using this the last term of the right hand side of (3.13) becomes

$$\begin{aligned} - \sum_{i=1}^P \int_{\Omega_\epsilon} \epsilon^\beta z_i \chi_{\{c_{i,\epsilon} \neq 0\}} \nabla \phi_\epsilon \cdot J_{i,\epsilon} dx &= - \sum_{i=1}^P \int_{\Omega_\epsilon} \epsilon^\beta z_i \nabla \phi_\epsilon \cdot J_{i,\epsilon} dx \\ &= - \epsilon^\beta \left\langle \sum_{i=1}^P z_i \partial_t c_{i,\epsilon}, \phi_\epsilon \right\rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} \end{aligned} \quad (3.15)$$

Differentiating (3.3), (3.4) with respect to t , we see that $\partial_t \phi_\epsilon \in H^1(\Omega_\epsilon)$ and the last term in (3.15) is equal to

$$-\epsilon^\beta \frac{1}{\lambda} \int_{\Omega_\epsilon} \epsilon^\alpha \nabla \partial_t \phi_\epsilon \nabla \phi_\epsilon dx = -\frac{1}{2\lambda} \epsilon^{\alpha+\beta} \frac{d}{dt} \int_{\Omega_\epsilon} |\nabla \phi_\epsilon|^2 dx, \quad (3.16)$$

where we used $\partial_t \xi_\epsilon = 0$, since ξ_ϵ is independent of t . From (3.13), (3.14), (3.16) we conclude that V_ϵ is differentiable with respect to time and (3.7) holds. Moreover, using (3.13) (3.14), (3.15), (3.16) in (3.7), we have

$$\frac{d}{dt} V_\epsilon(t) + \sum_{i=1}^P \int_{\Omega_\epsilon} \frac{|J_{i,\epsilon}|^2}{D_{i,\epsilon} c_{i,\epsilon}} dx = 0, \quad \text{for a.e. } t \in (0, T).$$

Hence,

$$\frac{d}{dt} V_\epsilon(t) \leq 0, \quad \text{for a.e. } t \in (0, T),$$

what implies

$$V_\epsilon(t) \leq V_\epsilon(0) \quad \text{for all } t \in [0, T].$$

It remains to obtain uniform estimate of $V_\epsilon(0)$.

$$\begin{aligned} V_\epsilon(0) &= \underbrace{\frac{1}{2\lambda} \epsilon^{\alpha+\beta} \int_{\Omega_\epsilon} |\nabla \phi_\epsilon(0)|^2 dx}_{:= I_1} \\ &+ \underbrace{\sum_{i=1}^P \int_{\Omega_\epsilon} \left(c_i^0 \log c_i^0 - c_i^0 + 1 + \frac{\eta}{p-1} (c_i^0)^p \right) dx}_{:= I_2}. \end{aligned} \quad (3.17)$$

Now,

$$|I_2| \leq |\Omega| \sum_{i=1}^P \max_{c_i^0 \in \bar{\Omega}} \left| c_i^0 \log c_i^0 - c_i^0 + 1 + \frac{\eta}{p-1} (c_i^0)^p \right| \leq C, \quad (3.18)$$

for some $C > 0$ independent of ϵ .

Now it remains to estimate I_1 . Testing with $\phi_\epsilon(0)$ the weak formulation of (3.3)-(3.4) at $t = 0$, we get

$$\begin{aligned} &\epsilon^\alpha \int_{\Omega_\epsilon} |\nabla \phi_\epsilon(0)|^2 dx \\ &\leq C\lambda \left[\int_{\Omega_\epsilon} |\phi_\epsilon(0)| dx + \epsilon \int_{\Gamma_\epsilon} |\phi_\epsilon(0)| dS(x) + \int_{\partial\Omega} |\phi_\epsilon(0)| dS(x) \right]. \end{aligned}$$

Since $|\Omega_\epsilon| = O(1)$ and $|\Gamma_\epsilon| = O\left(\frac{1}{\epsilon}\right)$, we have

$$\epsilon^\alpha \|\nabla \phi_\epsilon(0)\|_{L^2(\Omega_\epsilon)}^2 \leq C\lambda \left[\|\phi_\epsilon(0)\|_{L^2(\Omega_\epsilon)} + \sqrt{\epsilon} \|\phi_\epsilon(0)\|_{L^2(\Gamma_\epsilon)} + \|\phi_\epsilon(0)\|_{L^2(\partial\Omega)} \right]. \quad (3.19)$$

From the properties of the extension operator in Lemma A.3 given in the appendix and the usual trace-inequality we obtain

$$\|\phi_\epsilon(0)\|_{L^2(\partial\Omega)} = \|\tilde{\phi}_\epsilon(0)\|_{L^2(\partial\Omega)} \leq C(\Omega)\|\tilde{\phi}_\epsilon(0)\|_{H^1(\Omega)} \leq C_1\|\phi_\epsilon(0)\|_{H^1(\Omega_\epsilon)}, \quad (3.20)$$

together with the scaled trace-inequality from Lemma A.5 from the appendix we obtain from (3.19)

$$\epsilon^\alpha \|\nabla\phi_\epsilon(0)\|_{L^2(\Omega_\epsilon)}^2 \leq C\lambda\|\phi_\epsilon(0)\|_{H^1(\Omega_\epsilon)}.$$

Then (A.11) leads to

$$\epsilon^\alpha \|\nabla\phi_\epsilon(0)\|_{L^2(\Omega_\epsilon)} \leq C\lambda.$$

Since $\lambda \in (0, 1]$, we obtain

$$I_1 \leq C\epsilon^{-\alpha+\beta}. \quad (3.21)$$

(3.18) and (3.21) complete the proof. \square

Corollary 3.2. *Due to the fact that $r \log r - r + 1 \geq 0$, for $r \geq 0$, and taking into account that $\alpha \leq \beta$, Proposition 3.1 immediately implies*

$$\|c_{i,\epsilon}\|_{L^\infty(0,T;L^p(\Omega_\epsilon))} \leq C, \quad (3.22)$$

with $C > 0$ independent of λ and ϵ .

Based on estimate (3.22), in the following theorem we derive energy estimates for the microscopic solutions $(c_{i,\epsilon}, \phi_\epsilon)$.

Proposition 3.3. *There exists a constant $C > 0$ independent of ϵ such that the microscopic solutions $(c_{i,\epsilon}, \phi_\epsilon)$ of (2.2)-(2.4) satisfy the following estimates:*

$$\epsilon^\alpha \|\phi_\epsilon\|_{L^\infty(0,T;L^2(\Omega_\epsilon))} + \epsilon^\alpha \|\nabla\phi_\epsilon\|_{L^\infty(0,T;L^2(\Omega_\epsilon))} \leq C, \quad (3.23)$$

$$\|c_{i,\epsilon}\|_{L^\infty(0,T;L^p(\Omega_\epsilon))} + \|\nabla c_{i,\epsilon}\|_{L^2(0,T;L^2(\Omega_\epsilon))} \leq C. \quad (3.24)$$

Proof. Testing the weak formulation of Poisson's equation (2.4) with ϕ_ϵ , we get almost everywhere in $(0, T)$

$$\begin{aligned} & \epsilon^\alpha \|\nabla\phi_\epsilon\|_{L^2(\Omega_\epsilon)}^2 \\ & \leq \left\| \sum_{i=1}^P z_i c_{i,\epsilon} \right\|_{L^2(\Omega_\epsilon)} \|\phi_\epsilon\|_{L^2(\Omega_\epsilon)} + C\epsilon \int_{\Gamma_\epsilon} |\phi_\epsilon| dS(x) + C \int_{\partial\Omega} |\phi_\epsilon| dS(x) \\ & \leq C\|\phi_\epsilon\|_{L^2(\Omega_\epsilon)} + C\sqrt{\epsilon}\|\phi_\epsilon\|_{L^2(\Gamma_\epsilon)} + C\|\phi_\epsilon\|_{L^2(\partial\Omega)}; \end{aligned} \quad (3.25)$$

here in the last inequality we used that $\sum_{i=1}^P z_i c_{i,\epsilon}$ is bounded in $L^\infty(0, T; L^2(\Omega_\epsilon))$ uniformly with respect to ϵ (see Proposition 3.1 & Corollary 3.2) and $|\Gamma_\epsilon| = O\left(\frac{1}{\epsilon}\right)$. Comparing (3.25) to (3.19) with $\lambda = 1$, we conclude that the estimate $\|\nabla\phi_\epsilon(t)\|_{L^2(\Omega_\epsilon)}$

follows from similar arguments that we used to estimate for $\|\nabla\phi_\epsilon(0)\|_{L^2(\Omega_\epsilon)}$ in Proposition 3.1. Also, we have

$$\epsilon^\alpha \|\nabla\phi_\epsilon\|_{L^\infty(0,T;L^2(\Omega_\epsilon))} \leq C, \quad (3.26)$$

for some $C > 0$ independent of ϵ . Consequently, using (A.11) and (3.26), we get

$$\epsilon^\alpha \|\phi_\epsilon\|_{L^\infty(0,T;L^2(\Omega_\epsilon))} \leq C.$$

Next we prove (3.24). Due to Corollary 3.2 we only have to estimate the norm of the gradient. We test the equation (2.2) with $c_{i,\epsilon}$ and get

$$\begin{aligned} & \langle \partial_t c_{i,\epsilon}, c_{i,\epsilon} \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} + D_i \int_{\Omega_\epsilon} |\nabla c_{i,\epsilon}|^2 dx \\ & + D_i \eta p \int_{\Omega_\epsilon} c_{i,\epsilon}^{p-1} |\nabla c_{i,\epsilon}|^2 dx + \epsilon^\beta D_i z_i \int_{\Omega_\epsilon} c_{i,\epsilon} \nabla\phi_\epsilon \nabla c_{i,\epsilon} dx = 0. \end{aligned}$$

Since the third term of the above equation is non-negative, we have

$$\frac{1}{2} \frac{d}{dt} \|c_{i,\epsilon}\|_{L^2(\Omega_\epsilon)}^2 + D_i \int_{\Omega_\epsilon} |\nabla c_{i,\epsilon}|^2 dx + \frac{1}{2} \epsilon^\beta D_i z_i \int_{\Omega_\epsilon} \nabla\phi_\epsilon \nabla(c_{i,\epsilon}^2) dx \leq 0. \quad (3.27)$$

Since $c_{i,\epsilon}(t) \in H^1(\Omega_\epsilon) \cap L^\infty(\Omega_\epsilon)$, we have $c_{i,\epsilon}^2(t) \in H^1(\Omega_\epsilon)$. We test Poisson's equation (2.4) with $c_{i,\epsilon}^2(t)$ to obtain

$$\epsilon^\alpha \int_{\Omega_\epsilon} \nabla\phi_\epsilon \nabla(c_{i,\epsilon}^2) dx = \int_{\Omega_\epsilon} c_{i,\epsilon}^2 \sum_{i=1}^P z_i c_{i,\epsilon} dx + \int_{\partial\Omega_\epsilon} \xi_\epsilon c_{i,\epsilon}^2 dS(x). \quad (3.28)$$

Now using (3.28) in (3.27) we get

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|c_{i,\epsilon}\|_{L^2(\Omega_\epsilon)}^2 + D_i \int_{\Omega_\epsilon} |\nabla c_{i,\epsilon}|^2 dx \\ & \leq \frac{1}{2} \epsilon^{-\alpha+\beta} D_i \left| \int_{\Omega_\epsilon} c_{i,\epsilon}^2 \sum_{i=1}^P z_i c_{i,\epsilon} dx \right| + \frac{1}{2} \epsilon^{-\alpha+\beta} D_i \left| \int_{\partial\Omega_\epsilon} \xi_\epsilon c_{i,\epsilon}^2 dS(x) \right|. \end{aligned} \quad (3.29)$$

Let $C' = \max_{1 \leq i \leq P} |z_i|$. Then

$$\begin{aligned} & \left| \int_{\Omega_\epsilon} c_{i,\epsilon}^2 \sum_{i=1}^P z_i c_{i,\epsilon} dx \right| \leq C' \int_{\Omega_\epsilon} c_{i,\epsilon}^2 \sum_{i=1}^P c_{i,\epsilon} dx \leq C' \int_{\Omega_\epsilon} \left(\sum_{i=1}^P c_{i,\epsilon} \right)^3 dx \\ & \leq C' \left(\sum_{i=1}^P \|c_{i,\epsilon}\|_{L^3(\Omega_\epsilon)} \right)^3 \leq C, \end{aligned} \quad (3.30)$$

for some $C > 0$ independent of ϵ and t . The last inequality in (3.30) follows from (3.22).

So

$$\frac{1}{2} \epsilon^{-\alpha+\beta} D_i \left| \int_{\Omega_\epsilon} c_{i,\epsilon}^2 \sum_{i=1}^P z_i c_{i,\epsilon} dx \right| \leq \epsilon^{-\alpha+\beta} C. \quad (3.31)$$

Again,

$$\left| \int_{\partial\Omega_\epsilon} \xi_\epsilon c_{i,\epsilon}^2 dS(x) \right| \leq \xi^* \epsilon \int_{\Gamma_\epsilon} c_{i,\epsilon}^2 dS(x) + \xi^* \int_{\partial\Omega} c_{i,\epsilon}^2 dS(x). \quad (3.32)$$

Using the scaled trace-inequality from Lemma A.5 we get

$$\int_{\Gamma_\epsilon} c_{i,\epsilon}^2 dS(x) \leq C_1 \left(\frac{1}{\epsilon} \int_{\Omega_\epsilon} c_{i,\epsilon}^2 dx + \epsilon \int_{\Omega_\epsilon} |\nabla c_{i,\epsilon}|^2 dx \right). \quad (3.33)$$

Next, we estimate the last term of (3.32) using the extension operator from Lemma A.3 and the weighted trace inequality (see [19, p. 63, Excercise II.4.1]) to obtain

$$\begin{aligned} \int_{\partial\Omega} c_{i,\epsilon}^2 dS(x) &= \int_{\partial\Omega} \widetilde{c}_{i,\epsilon}^2 dS(x) \leq C_2(\Omega, \delta) \int_{\Omega} \widetilde{c}_{i,\epsilon}^2 dx + \delta \int_{\Omega} |\nabla \widetilde{c}_{i,\epsilon}|^2 dx \\ &\leq C_3 \int_{\Omega_\epsilon} c_{i,\epsilon}^2 dx + C_4 \delta \int_{\Omega_\epsilon} |\nabla c_{i,\epsilon}|^2 dx \end{aligned} \quad (3.34)$$

for any $\delta > 0$. Finally, utilizing (3.34), (3.33), (3.31) in (3.29), we get

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \|c_{i,\epsilon}\|_{L^2(\Omega_\epsilon)}^2 + D_i \int_{\Omega_\epsilon} |\nabla c_{i,\epsilon}|^2 dx \\ &\leq C \epsilon^{-\alpha+\beta} + \frac{1}{2} \epsilon^{-\alpha+\beta} D_i |z_i| \xi^* (C_1 \epsilon^2 + C_4 \delta) \int_{\Omega_\epsilon} |\nabla c_{i,\epsilon}|^2 dx \\ &+ \frac{1}{2} \epsilon^{-\alpha+\beta} D_i |z_i| \xi^* (C_1 + C_3) \int_{\Omega_\epsilon} c_{i,\epsilon}^2 dx. \end{aligned} \quad (3.35)$$

Because of (3.22), we have that the last term of (3.35) is bounded by $C \epsilon^{-\alpha+\beta}$, for some $C > 0$ independent of ϵ and t . Again, for $z_i \neq 0$, if we choose ϵ and δ is small enough so that

$$C_1 \epsilon^2 < \frac{1}{4|z_i| \xi^*}, \quad C_4 \delta < \frac{1}{4|z_i| \xi^*},$$

then we can absorb the gradient term by the left hand side. These arguments lead to

$$\frac{d}{dt} \|c_{i,\epsilon}\|_{L^2(\Omega_\epsilon)}^2 + D_i \int_{\Omega_\epsilon} |\nabla c_{i,\epsilon}|^2 dx \leq C \epsilon^{-\alpha+\beta}, \quad (3.36)$$

for some $C > 0$ independent of ϵ and t . Integration with respect to time gives (3.24). \square

Now, we are able to prove the uniform L^∞ -estimate for the microscopic solutions $c_{i,\epsilon}$ for $i = 1, \dots, P$. Our results are based on [27, Theorem 6.1 and Remark 6.2]. An important aspect in the proof is the estimate of the drift term, where refined arguments are required in order to avoid the occurrence of higher (than H^1) norms of the potential.

Theorem 3.4. *There exists a constant $C > 0$ which is independent of ϵ such that*

$$\|c_{i,\epsilon}\|_{L^\infty((0,T) \times \Omega_\epsilon)} \leq C.$$

Proof. For $u \in L^2(0, T; H^1(\Omega))$ with $\partial_t u \in L^2(0, T; H^1(\Omega)')$, where Ω is an arbitrary bounded open set, we define $u^+(t, x) := \max\{u(t, x), 0\}$. Then we have, see e.g. [40],

$$\langle \partial_t u(t), u^+(t) \rangle_{H^1(\Omega)', H^1(\Omega)} = \frac{1}{2} \frac{d}{dt} \|u^+(t)\|_{L^2(\Omega)}^2 \quad \text{for a.e. } t \in (0, T). \quad (3.37)$$

Now, let $\omega \in (0, \infty)$ and $k \in [0, \infty)$ be arbitrary and set $W := e^{-\omega t} c_{i,\epsilon}$ and $W_k := (W - k)^+$. We test the equation (2.2) with $e^{-\omega t} W_k$ to obtain

$$\begin{aligned} & \langle \partial_t c_{i,\epsilon}, e^{-\omega t} W_k \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} + D_i \int_{\Omega_\epsilon} h'_p(c_{i,\epsilon}) \nabla c_{i,\epsilon} e^{-\omega t} \nabla W_k \, dx \\ &= -\epsilon^\beta D_i \int_{\Omega_\epsilon} z_i c_{i,\epsilon} \nabla \phi_\epsilon e^{-\omega t} \nabla W_k \, dx. \end{aligned} \quad (3.38)$$

Due to

$$\partial_t (e^{-\omega t} c_{i,\epsilon} - k) = -\omega e^{-\omega t} c_{i,\epsilon} + e^{-\omega t} \partial_t c_{i,\epsilon}, \quad \text{for a.e. } t \in (0, T),$$

and (3.37), we have

$$\begin{aligned} & \langle \partial_t c_{i,\epsilon}, e^{-\omega t} W_k \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} \\ &= \langle \partial_t (e^{-\omega t} c_{i,\epsilon} - k), W_k \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} + \omega \int_{\Omega_\epsilon} e^{-\omega t} c_{i,\epsilon} W_k \, dx \\ &= \frac{1}{2} \frac{d}{dt} \int_{\Omega_\epsilon} W_k^2 \, dx + \omega \int_{\Omega_\epsilon} W W_k \, dx. \end{aligned} \quad (3.39)$$

Now utilizing (3.39), $h'_p \geq 1$ in (3.38), we get

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\Omega_\epsilon} W_k^2 \, dx + D_i \|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 \\ & \leq -\epsilon^\beta D_i z_i \int_{\Omega_\epsilon} W \nabla W_k \nabla \phi_\epsilon \, dx - \omega \int_{\Omega_\epsilon} W W_k \, dx \\ & = -\epsilon^\beta D_i z_i \int_{\Omega_\epsilon} W \nabla W_k \nabla \phi_\epsilon \, dx - \omega \int_{\{W(t) > k\}} W^2 \, dx + \omega \int_{\{W(t) > k\}} W k \, dx \\ & \leq -\epsilon^\beta D_i z_i \int_{\Omega_\epsilon} W \nabla W_k \nabla \phi_\epsilon \, dx - \frac{\omega}{2} \int_{\{W(t) > k\}} W^2 \, dx + \frac{\omega}{2} \int_{\{W(t) > k\}} k^2 \, dx, \end{aligned} \quad (3.40)$$

where $\{W(t) > k\} := \{x \in \Omega_\epsilon : W(t, x) > k\}$.

First, we consider the first term of the right hand side of (3.40).

$$\begin{aligned} & -\epsilon^\beta D_i z_i \int_{\Omega_\epsilon} W \nabla W_k \nabla \phi_\epsilon \, dx \\ &= -\epsilon^\beta D_i z_i \int_{\Omega_\epsilon} (W - k) \nabla W_k \nabla \phi_\epsilon \, dx - \epsilon^\beta D_i z_i k \int_{\Omega_\epsilon} \nabla W_k \nabla \phi_\epsilon \, dx \\ &= -\epsilon^\beta D_i z_i \int_{\Omega_\epsilon} W_k \nabla W_k \nabla \phi_\epsilon \, dx - \epsilon^\beta D_i z_i k \int_{\Omega_\epsilon} \nabla W_k \nabla \phi_\epsilon \, dx \end{aligned}$$

$$= \underbrace{-\frac{1}{2}\epsilon^\beta D_i z_i \int_{\Omega_\epsilon} \nabla(W_k^2) \nabla \phi_\epsilon dx}_{I_1} \quad \underbrace{-\epsilon^\beta D_i z_i k \int_{\Omega_\epsilon} \nabla W_k \nabla \phi_\epsilon dx}_{I_2}. \quad (3.41)$$

Using W_k^2 as a test-function in the weak formulation of Poisson's-equation (2.4), we obtain

$$I_1 = \underbrace{-\frac{1}{2}\epsilon^{\beta-\alpha} D_i z_i \int_{\Omega_\epsilon} W_k^2 \sum_{i=1}^P z_i c_{i,\epsilon} dx}_{A_1} \quad \underbrace{-\frac{1}{2}\epsilon^{\beta-\alpha} D_i z_i \int_{\partial\Omega_\epsilon} W_k^2 \xi_\epsilon dS(x)}_{A_2}.$$

The term A_1 can be estimated using Corollary 3.2 by

$$A_1 \leq \frac{1}{2}\epsilon^{\beta-\alpha} D_i |z_i| \|W_k\|_{L^4(\Omega_\epsilon)}^2 \left\| \sum_{i=1}^P z_i c_{i,\epsilon} \right\|_{L^2(\Omega_\epsilon)} \leq C\epsilon^{-\alpha+\beta} \|W_k\|_{L^4(\Omega_\epsilon)}^2.$$

Using the Gagliardo-Nirenberg-interpolation inequality (see for example [19, Exercise II.3.12] or [37]) and Lemma A.3, for any $\delta_1 > 0$, we obtain

$$\begin{aligned} A_1 &\leq C\epsilon^{-\alpha+\beta} \|\widetilde{W}_k\|_{L^4(\Omega)}^2 \\ &\leq C\epsilon^{-\alpha+\beta} \left(\delta_1 \|\nabla \widetilde{W}_k\|_{L^2(\Omega)}^2 + C(\delta_1, \Omega) \|\widetilde{W}_k\|_{L^2(\Omega)}^2 \right) \\ &\leq C\epsilon^{-\alpha+\beta} \left(C_1 \delta_1 \|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 + C_2(\delta_1) \|W_k\|_{L^2(\Omega_\epsilon)}^2 \right). \end{aligned} \quad (3.42)$$

Further, for the term A_2 we get

$$\begin{aligned} A_2 &\leq C\xi^* \epsilon^{-\alpha+\beta} \int_{\Gamma_\epsilon} \epsilon W_k^2 dS(x) + C\xi^* \epsilon^{-\alpha+\beta} \int_{\partial\Omega} W_k^2 dS(x) \\ &\leq C\epsilon^{-\alpha+\beta} \left(\int_{\Omega_\epsilon} W_k^2 dx + \epsilon^2 \|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 \right) + C\epsilon^{-\alpha+\beta} \int_{\partial\Omega} W_k^2; \end{aligned}$$

here in the last inequality we used again Lemma A.5. Consequently, by means of [19, p. 63, Exercise II.4.1] and Lemma A.3, for any $\delta_2 > 0$ we get

$$\begin{aligned} A_2 &\leq C\epsilon^{-\alpha+\beta} \left(\int_{\Omega_\epsilon} W_k^2 dx + \epsilon^2 \|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 \right) \\ &\quad + C\epsilon^{-\alpha+\beta} \left(\delta_2 \|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 + C_3(\delta_2) \int_{\Omega_\epsilon} W_k^2 dx \right). \end{aligned} \quad (3.43)$$

So (3.42), (3.43) imply

$$\begin{aligned} I_1 &\leq C\epsilon^{-\alpha+\beta} (\epsilon^2 + C_1 \delta_1 + \delta_2) \|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 \\ &\quad + C\epsilon^{-\alpha+\beta} (C_2(\delta_1) + C_3(\delta_2) + 1) \int_{\Omega_\epsilon} W_k^2 dx. \end{aligned}$$

Then, the inequality

$$\int_{\Omega_\epsilon} W_k^2 dx \leq 2 \left(\int_{\{W(t)>k\}} W^2 dx + k^2 \int_{\{W(t)>k\}} dx \right)$$

leads to

$$\begin{aligned} I_1 &\leq C\epsilon^{-\alpha+\beta}(\epsilon^2 + C_1\delta_1 + \delta_2)\|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 \\ &\quad + C\epsilon^{-\alpha+\beta}C_4(\delta_1, \delta_2) \left(\int_{\{W(t)>k\}} W^2 dx + k^2 \int_{\{W(t)>k\}} dx \right). \end{aligned} \quad (3.44)$$

To estimate I_2 from (3.41) we use kW_k as a test-function in Poisson's-equation (2.4) to obtain

$$I_2 = \underbrace{-\epsilon^{\beta-\alpha} D_i z_i \int_{\Omega_\epsilon} kW_k \sum_{i=1}^P z_i c_{i,\epsilon} dx}_{A_3} \quad \underbrace{-\epsilon^{\beta-\alpha} D_i z_i \int_{\partial\Omega_\epsilon} kW_k \xi_\epsilon dS(x)}_{A_4}. \quad (3.45)$$

Using again Corollary 3.2 (remember $p \geq 4$) we immediately get

$$A_3 \leq C\epsilon^{-\alpha+\beta} \|k\|_{L^2(\{W(t)>k\})} \|W_k\|_{L^4(\{W(t)>k\})}. \quad (3.46)$$

Now, using similar arguments used in the estimation of A_1 , we have from (3.46) for any $\delta_3 > 0$,

$$A_3 \leq C\epsilon^{-\alpha+\beta} \left(\int_{\{W(t)>k\}} k^2 dx + C_5\delta_3 \|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 + C_6(\delta_3) \int_{\{W(t)>k\}} W_k^2 dx \right).$$

This gives

$$\begin{aligned} A_3 &\leq C\epsilon^{-\alpha+\beta}C_5\delta_3\|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 \\ &\quad + C\epsilon^{-\alpha+\beta}C_7(\delta_3) \left(\int_{\{W(t)>k\}} W^2 dx + \int_{\{W(t)>k\}} k^2 dx \right). \end{aligned} \quad (3.47)$$

Now let us estimate A_4 from (3.45). We have

$$A_4 \leq C\xi^* \epsilon^{-\alpha+\beta} \int_{\Gamma_\epsilon} \epsilon k |W_k| dS(x) + C\xi^* \epsilon^{-\alpha+\beta} \int_{\partial\Omega} k |W_k| dS(x). \quad (3.48)$$

The scaled trace-inequality from Lemma A.5 yields

$$\int_{\Gamma_\epsilon} k |W_k| dS(x) \leq C \left(\frac{1}{\epsilon} \int_{\Omega_\epsilon} k |W_k| dx + \|k \nabla W_k\|_{L^1(\Omega_\epsilon)} \right).$$

Also, [19, p. 63, Exercise II.4.1] and Lemma A.3 lead to

$$\int_{\partial\Omega} k |W_k| dx \leq C_8(\delta_4) \int_{\Omega_\epsilon} k |W_k| dx + C_9\delta_4 \|k \nabla W_k\|_{L^1(\Omega_\epsilon)},$$

for any $\delta_4 > 0$. Making use of these two inequalities in (3.48), we obtain

$$A_4 \leq C\epsilon^{-\alpha+\beta}(\epsilon + C_9\delta_4)\|k\nabla W_k\|_{L^1(\Omega_\epsilon)} + C\epsilon^{-\alpha+\beta}(1 + C_8(\delta_4)) \int_{\Omega_\epsilon} k|W_k| dx.$$

Hence,

$$\begin{aligned} A_4 &\leq C\epsilon^{-\alpha+\beta}(\epsilon + C_9\delta_4)\|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 \\ &\quad + C\epsilon^{-\alpha+\beta}C_{10}(\delta_4) \left(\int_{\{W(t)>k\}} W^2 dx + \int_{\{W(t)>k\}} k^2 dx \right). \end{aligned} \quad (3.49)$$

Consequently, from (3.47), (3.49), we get

$$\begin{aligned} I_2 &\leq C\epsilon^{-\alpha+\beta}(\epsilon + C_5\delta_3 + C_9\delta_4)\|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 \\ &\quad + C\epsilon^{-\alpha+\beta}C_{11}(\delta_3, \delta_4) \left(\int_{\{W(t)>k\}} W^2 dx + \int_{\{W(t)>k\}} k^2 dx \right). \end{aligned} \quad (3.50)$$

Now using (3.50), (3.44) in (3.40), we get

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \int_{\Omega_\epsilon} W_k^2 dx + D_i \|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 \\ &\leq C\epsilon^{-\alpha+\beta}(\epsilon^2 + \epsilon + C_1\delta_1 + \delta_2 + C_5\delta_3 + C_9\delta_4)\|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 \\ &\quad + C\epsilon^{-\alpha+\beta}C_{12}(\delta_1, \delta_2, \delta_3, \delta_4) \int_{\{W(t)>k\}} W^2 dx \\ &\quad + \left[C\epsilon^{-\alpha+\beta}C_{13}(\delta_1, \delta_2, \delta_3, \delta_4) + \frac{\omega}{2} \right] \int_{\{W(t)>k\}} k^2 dx - \frac{\omega}{2} \int_{\{W(t)>k\}} W^2 dx. \end{aligned}$$

From the above equation we see that if we choose $\epsilon, \delta_1, \dots, \delta_4$ small enough so that $C\epsilon^{-\alpha+\beta}(\epsilon^2 + \epsilon + C_1\delta_1 + \delta_2 + C_5\delta_3 + C_9\delta_4) < \frac{D_i}{2}$, then we can absorb the gradient term in the right hand side by the gradient term in the left hand side. Note that, $\delta_1, \dots, \delta_4$ do not depend on ϵ , whenever $\epsilon < \epsilon_0$, for some $\epsilon_0 > 0$. Then, if we assume $\frac{\omega}{2} > C\epsilon^{-\alpha+\beta}C_{12}(\delta_1, \dots, \delta_4)$, we can remove the term $\int_{\{W(t)>k\}} W^2 dx$ from the right hand side. Therefore, we get

$$\frac{d}{dt} \int_{\Omega_\epsilon} W_k^2 dx + D_i \|\nabla W_k\|_{L^2(\Omega_\epsilon)}^2 \leq (C\epsilon^{-\alpha+\beta} + \omega) \int_{\{W(t)>k\}} k^2 dx. \quad (3.51)$$

Choose $k \geq \|c_i^0\|_{L^\infty(\Omega)} + 1$, which implies $W_k(0) = 0$. Now, we integrate (3.51) with respect to t to obtain

$$\max_{t \in [0, T]} \int_{\Omega_\epsilon} W_k^2(t) dx + D_i \|\nabla W_k\|_{L^2((0, T) \times \Omega_\epsilon)}^2 \leq (C\epsilon^{-\alpha+\beta} + \omega) k^2 \int_{\{W>k\}} dx dt,$$

where $\{W > k\} := \{(t, x) \in (0, T) \times \Omega_\epsilon : W(t, x) > k\}$. Now utilizing [27, p. 102, Theorem 6.1 and p. 103, Remark 6.2] with $r = q = \frac{2(n+2)}{n}$, $\kappa = \frac{2}{n}$, it follows that

$$\operatorname{ess\,sup}_{(t,x) \in (0, T) \times \Omega_\epsilon} W(t, x) \leq C_{14} \left[1 + C_{15}(C_{16}\epsilon^{-\alpha+\beta} + \omega)^{\frac{n+2}{4}} \right]. \quad (3.52)$$

We emphasize that the constants C_{14}, C_{15}, C_{16} are independent of ϵ . In fact, the proof of [27, Theorem 6.1] is based on the embedding

$$L^\infty((0, T), L^2(\Omega_\epsilon)) \cap L^2((0, T), H^1(\Omega_\epsilon)) \hookrightarrow L^r((0, T) \times \Omega_\epsilon).$$

The embedding constant is independent of ϵ , since due to the extension operator from Lemma A.3 we easily obtain for any $u_\epsilon \in L^\infty((0, T), L^2(\Omega_\epsilon)) \cap L^2((0, T), H^1(\Omega_\epsilon))$ that

$$\left(\int_0^T \int_{\Omega_\epsilon} |u_\epsilon|^r \right)^{\frac{1}{r}} \leq c \left[\operatorname{ess\,sup}_{t \in (0, T)} \left(\int_{\Omega_\epsilon} |u_\epsilon(t, x)|^2 dx \right)^{\frac{1}{2}} + \|\nabla u_\epsilon\|_{L^2((0, T) \times \Omega_\epsilon)} \right],$$

with a constant c independent of ϵ .

Consequently, from (3.52) we conclude that W is bounded from above by some positive constant C in $(0, T) \times \Omega_\epsilon$ uniformly with respect to ϵ . Since $c_{i, \epsilon}$ are non-negative, the proof is complete. \square

Proposition 3.5. *There exists a constant $C > 0$ independent of ϵ such that*

$$\|\partial_t c_{i, \epsilon}\|_{L^2(0, T; H^1(\Omega_\epsilon)')} \leq C. \quad (3.53)$$

Proof. Let $\psi \in L^2(0, T; H^1(\Omega_\epsilon))$. From (2.2) we have

$$\begin{aligned} & \int_0^T \langle \partial_t c_{i, \epsilon}, \psi \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} dt \\ & \leq D_i \int_0^T \int_{\Omega_\epsilon} (1 + \eta c_{i, \epsilon}^{p-1}) |\nabla c_{i, \epsilon}| |\nabla \psi| dx dt + \epsilon^\beta D_i |z_i| \int_0^T \int_{\Omega_\epsilon} c_{i, \epsilon} |\nabla \phi_\epsilon| |\nabla \psi| dx dt \\ & \leq C \|\nabla c_{i, \epsilon}\|_{L^2(0, T; L^2(\Omega_\epsilon))} \|\psi\|_{L^2(0, T; H^1(\Omega_\epsilon))} + C \epsilon^\beta \|\nabla \phi_\epsilon\|_{L^2(0, T; L^2(\Omega_\epsilon))} \|\psi\|_{L^2(0, T; H^1(\Omega_\epsilon))} \\ & \leq C(1 + \epsilon^{-\alpha+\beta}) \|\psi\|_{L^2(0, T; H^1(\Omega_\epsilon))}; \end{aligned}$$

here we used Proposition 3.3 and Theorem 3.4. This completes the proof. \square

4 Derivation of macroscopic model

In this section, we derive the macroscopic (homogenized) drift-diffusion model by means of two-scale convergence concepts introduced in [2, 35, 36]. The two-scale convergence results are based on the uniform *a priori* estimates for the microscopic solutions $c_{i, \epsilon}, i = 1, \dots, P$ and ϕ_ϵ from Section 3.

Definition 4.1 ([2]). A sequence of functions u_ϵ in $L^2((0, T) \times \Omega)$ is said to two-scale converge to a function $u_0 \in L^2((0, T) \times \Omega \times Y)$, if for all $\psi \in L^2((0, T) \times \Omega; C_{per}(\bar{Y}))$ the following holds:

$$\lim_{\epsilon \rightarrow 0} \int_0^T \int_{\Omega} u_\epsilon(t, x) \psi \left(t, x, \frac{x}{\epsilon} \right) dx dt = \int_0^T \int_{\Omega} \int_Y u_0(t, x, y) \psi(t, x, y) dy dx dt.$$

The following compactness result is well known: For every bounded sequence $u_\epsilon \in L^2((0, T), H^1(\Omega))$ there exists $u_0 \in L^2((0, T), H^1(\Omega))$ and $u_1 \in L^2((0, T) \times \Omega, H^1_{per}(Y)/\mathbb{R})$ such that up to a subsequence

$$\begin{aligned} u_\epsilon &\rightarrow u_0 && \text{in the two-scale sense,} \\ \nabla u_\epsilon &\rightarrow \nabla u_0 + \nabla_y u_1 && \text{in the two-scale sense.} \end{aligned}$$

A proof can be found in [2] (see also [36]) for the time-independent case, and can be easily generalized to time-dependent problems. Note, however, that in our problem the microscopic solutions are only defined on the perforated domain Ω_ϵ , and to apply the above two-scale compactness results we have to extend $c_{i,\epsilon}$ and ϕ_ϵ in a suitable way to the whole domain Ω . In a trivial way we can extend a function $f : \Omega_\epsilon \rightarrow \mathbb{R}$ by zero to Ω (this extension is denoted by $\chi_{\Omega_\epsilon} f$, and χ_{Ω_ϵ} denotes the characteristic function of Ω_ϵ). For the zero-extension, the regularity properties are in general not preserved. These are, however, the basis for obtaining strong compactness results, which allow to pass to the limit in the nonlinear terms. Hence, we extend the functions $c_{i,\epsilon}$ by using the extension operator $\widetilde{c}_{i,\epsilon}$ from Lemma A.3, pointwise in $t \in (0, T)$. It is well known, see, for example, [9] and [1], that $\widetilde{c}_{i,\epsilon}$ fulfills the same regularity results and uniform *a priori* estimates as $c_{i,\epsilon}$ in Proposition 3.3 and 3.4. In Lemma A.3, we additionally prove that $\widetilde{c}_{i,\epsilon}$ is also essentially bounded and non-negative. These properties are needed to pass to the limit in the microscopic model, in particular in the nonlinear term $h_p(c_{i,\epsilon})$ entering the diffusion coefficients. Finally, we emphasize that we have no information about $\partial_t \widetilde{c}_{i,\epsilon}$, but this is not necessary for the derivation of the macroscopic model.

In the following proposition, we prove convergence results for the microscopic solutions. Here, the characteristic function on Y^f is denoted by χ_{Y^f} and it is Y -periodically extended to the whole \mathbb{R}^n .

Proposition 4.2. (a) *There exist $c_{i,0} \in L^\infty((0, T) \times \Omega) \cap L^2(0, T; H^1(\Omega))$, $c_{i,0} \geq 0$ almost everywhere in $(0, T) \times \Omega$ with $\partial_t c_{i,0} \in L^2(0, T; H^1(\Omega)')$ and $c_{i,1} \in L^2((0, T) \times \Omega; H^1_{per}(Y^f)/\mathbb{R})$ such that for every $q \in [1, \infty)$, up to a subsequence*

$$\chi_{\Omega_\epsilon} c_{i,\epsilon} \rightarrow \chi_{Y^f} c_{i,0} \quad \text{in the two-scale sense,} \quad (4.1)$$

$$\chi_{\Omega_\epsilon} \nabla c_{i,\epsilon} \rightarrow \chi_{Y^f} (\nabla c_{i,0} + \nabla_y c_{i,1}) \quad \text{in the two-scale sense,} \quad (4.2)$$

$$\widetilde{c}_{i,\epsilon} \rightarrow c_{i,0} \quad \text{strongly in } L^q((0, T) \times \Omega), \quad (4.3)$$

$$h'_p(\widetilde{c}_{i,\epsilon}) \rightarrow h'_p(c_{i,0}) \quad \text{strongly in } L^q((0, T) \times \Omega). \quad (4.4)$$

(b) *There exists $\phi_0 \in L^2(0, T; H^1(\Omega))$ and $\phi_1 \in L^2((0, T) \times \Omega; H^1_{per}(Y^f)/\mathbb{R})$ such that, up to a subsequence*

$$\epsilon^\alpha \chi_{\Omega_\epsilon} \phi_\epsilon \rightarrow \chi_{Y^f} \phi_0 \quad \text{in the two-scale sense,} \quad (4.5)$$

$$\epsilon^\alpha \chi_{\Omega_\epsilon} \nabla \phi_\epsilon \rightarrow \chi_{Y^f} (\nabla \phi_0 + \nabla_y \phi_1) \quad \text{in the two-scale sense.} \quad (4.6)$$

Proof. The two-scale convergence results are quite standard (see for example [2] for more details) and follow from the *a priori* estimates in Proposition 3.3. It remains to prove the strong convergence results for the extension. Using again the *a priori* estimates

for $c_{i,\epsilon}$, we obtain from [15, Lemma 9 and Lemma 10] the strong convergence of $\widetilde{c_{i,\epsilon}}$ to $c_{i,0}$ in $L^2((0, T) \times \Omega)$ (up to a subsequence). This implies, again up to a subsequence, pointwise almost everywhere convergence in $(0, T) \times \Omega$. Together with the L^∞ -estimate from Theorem 3.4 and the dominated convergence theorem we obtain the convergence of $\widetilde{c_{i,\epsilon}}$ in $L^q((0, T) \times \Omega)$ for every $q \in [1, \infty)$. The lower semi-continuity of the norm implies $c_{i,0} \in L^\infty((0, T) \times \Omega)$, see also [8, Exercise 4.6]. For the convergence of $h'_p(\widetilde{c_{i,\epsilon}})$ we use the local Lipschitz continuity of h'_p and again the essential boundedness of $\widetilde{c_{i,\epsilon}}$, which implies together with Lemma A.3

$$\|h'_p(\widetilde{c_{i,\epsilon}}) - h'_p(c_{i,0})\|_{L^q((0,T) \times \Omega)} \leq C \|\widetilde{c_{i,\epsilon}} - c_{i,0}\|_{L^q((0,T) \times \Omega)} \rightarrow 0$$

for $\epsilon \rightarrow 0$. This finishes the proof. \square

We now have all ingredients to pass to the limit $\epsilon \rightarrow 0$ in the variational formulation (2.2)-(2.4) and to obtain the macroscopic models for different values of the parameters α and β .

Theorem 4.3. (a) *Let $\alpha = \beta$. The limit functions $c_{i,0}$ and ϕ_0 from Proposition 4.2 are weak solutions of the following homogenized model:*

$$\begin{aligned} \partial_t c_{i,0} - \nabla_x \cdot [D_i A_{hom} \nabla h_p(c_{i,0}) + D_i z_i c_{i,0} A_{hom} \nabla \phi_0] &= 0 && \text{in } (0, T) \times \Omega, \\ - [D_i A_{hom} \nabla h_p(c_{i,0}) + D_i z_i c_{i,0} A_{hom} \nabla \phi_0] \cdot \nu &= 0 && \text{on } (0, T) \times \partial\Omega, \\ c_{i,0}(0, x) &= c_i^0(x) && \text{in } \Omega, \end{aligned} \quad (4.7)$$

for all $i \in \{1, \dots, P\}$, and

$$\begin{aligned} -\nabla_x \cdot [A_{hom} \nabla \phi_0(t, x)] &= \sum_{k=1}^P z_k c_{k,0}(t, x) + \frac{1}{|Y^f|} \int_{\Gamma} \xi_1(x, y) dy && \text{in } (0, T) \times \Omega, \\ A_{hom} \nabla \phi_0(t, x) \cdot \nu &= \frac{1}{|Y^f|} \xi_2(x) && \text{on } (0, T) \times \partial\Omega. \end{aligned} \quad (4.8)$$

Here A_{hom} is an $n \times n$ matrix given by

$$A_{hom} e_k = \frac{1}{|Y^f|} \int_{Y^f} (\nabla_y w_k + e_k) dy, \quad \text{for } 1 \leq k \leq n, \quad (4.9)$$

where e_k denotes the standard basis vector of \mathbb{R}^n and $w_k \in H_{per}^1(Y^f)/\mathbb{R}$ is the unique weak solution of the so-called cell problem

$$\begin{aligned} -\nabla_y \cdot (\nabla_y w_k(y) + e_k) &= 0 && \text{in } Y^f, \\ (\nabla_y w_k(y) + e_k) \cdot \nu &= 0 && \text{on } \partial Y^f, \end{aligned} \quad (4.10)$$

$y \mapsto w_k(y)$ is Y -periodic.

(b) *If $\alpha < \beta$, then the homogenized Poisson's equation (4.8) remains the same, however the convective term $D_i z_i c_{i,0} A_{hom} \nabla \phi_0$ in the homogenized transport equation (4.7)*

does not appear. That is, in that case (4.7) becomes

$$\begin{aligned} \partial_t c_{i,0} - \nabla_x \cdot D_i A_{hom} \nabla h_p(c_{i,0}) &= 0 && \text{in } (0, T) \times \Omega, \\ -D_i A_{hom} \nabla h_p(c_{i,0}) \cdot \nu &= 0 && \text{on } (0, T) \times \partial\Omega, \\ c_{i,0}(0, x) &= c_i^0(x) && \text{in } \Omega. \end{aligned} \quad (4.11)$$

Proof. We start by proving statement (a). Let $\psi_0(t, x) \in C^\infty([0, T] \times \overline{\Omega})$ with $\psi_0(T, \cdot) = 0$ and $\psi_1(t, x, y) \in C_0^\infty((0, T) \times \Omega \times \overline{Y})$, which is Y -periodic in y . Let us consider a subsequence of ϵ , still denoted by ϵ , along which the convergence results given in Proposition 4.2 hold. Now considering $\psi_\epsilon(t, x) := \psi_0(t, x) + \epsilon \psi_1(t, x, \frac{x}{\epsilon})$ as a test function in (2.2), we get

$$\begin{aligned} & - \int_0^T \int_{\Omega_\epsilon} c_{i,\epsilon} \left[\partial_t \psi_0(t, x) + \epsilon \partial_t \psi_1 \left(t, x, \frac{x}{\epsilon} \right) \right] dx dt \\ & + D_i \int_0^T \int_{\Omega_\epsilon} h'_p(c_{i,\epsilon}) \nabla c_{i,\epsilon} \left[\nabla \psi_0(t, x) + \epsilon \nabla_x \psi_1 \left(t, x, \frac{x}{\epsilon} \right) + \nabla_y \psi_1 \left(t, x, \frac{x}{\epsilon} \right) \right] dx dt \\ & + \epsilon^\beta D_i \int_0^T \int_{\Omega_\epsilon} z_i c_{i,\epsilon} \nabla \phi_\epsilon \left[\nabla \psi_0(t, x) + \epsilon \nabla_x \psi_1 \left(t, x, \frac{x}{\epsilon} \right) + \nabla_y \psi_1 \left(t, x, \frac{x}{\epsilon} \right) \right] dx dt \\ & = \int_{\Omega_\epsilon} c_i^0(x) \psi_0(0, x) dx. \end{aligned} \quad (4.12)$$

Let us first pass to the limit in the second term in the left-hand side of (4.12). Passing to the limit in the other terms will follow from simpler arguments. We have,

$$\begin{aligned} D_i \int_0^T \int_{\Omega_\epsilon} h'_p(c_{i,\epsilon}) \nabla c_{i,\epsilon} \nabla \psi_\epsilon dx &= D_i \int_0^T \int_{\Omega_\epsilon} [h'_p(c_{i,\epsilon}) - h'_p(c_{i,0})] \nabla c_{i,\epsilon} \nabla \psi_\epsilon dx \\ &+ D_i \int_0^T \int_{\Omega_\epsilon} h'_p(c_{i,0}) \nabla c_{i,\epsilon} \nabla \psi_\epsilon dx =: A_\epsilon^1 + A_\epsilon^2 \end{aligned}$$

For the first term we use the strong convergence of $h'_p(\widetilde{c}_{i,\epsilon})$ from Proposition 4.2 and the boundedness of $\nabla c_{i,\epsilon}$ from Proposition 3.3 (we emphasize that $\nabla \psi_\epsilon$ is essentially bounded):

$$|A_\epsilon^1| \leq C \|h'_p(\widetilde{c}_{i,\epsilon}) - h'_p(c_{i,0})\|_{L^2((0,T) \times \Omega)} \xrightarrow{\epsilon \rightarrow 0} 0.$$

For the term A_ϵ^2 we notice that $h'_p(c_{i,0}) \nabla \psi_\epsilon$ is an admissible test-function in the two-scale sense. Hence, we can use the two-scale convergence result for $\chi_{\Omega_\epsilon} \nabla c_{i,\epsilon}$ from Proposition 4.2 to obtain

$$\lim_{\epsilon \rightarrow 0} A_\epsilon^2 = D_i \int_0^T \int_\Omega \int_{Y^f} h'_p(c_{i,0}) (\nabla c_{i,0} + \nabla_y c_{i,1}) (\nabla \psi_0(t, x) + \nabla_y \psi_1(t, x, y)) dy dx dt.$$

The other terms in (4.12) can be treated in a similar way and we obtain for $\epsilon \rightarrow 0$

$$-|Y^f| \int_0^T \int_\Omega c_{i,0} \partial_t \psi_0 dx dt \quad (4.13)$$

$$\begin{aligned}
 & +D_i \int_0^T \int_{\Omega} \int_{Y^f} h'_p(c_{i,0}) (\nabla c_{i,0} + \nabla_y c_{i,1}) (\nabla \psi_0 + \nabla_y \psi_1) dy dx dt \\
 & +D_i z_i \int_0^T \int_{\Omega} \int_{Y^f} c_{i,0} (\nabla \phi_0 + \nabla_y \phi_1) (\nabla \psi_0 + \nabla_y \psi_1) dy dx dt \\
 & = |Y^f| \int_{\Omega} c_i^0 \psi_0(0, x) dx. \tag{4.14}
 \end{aligned}$$

Choosing $\psi_1 = 0$ we obtain that $c_{i,0}(0) = c_i^0$. Integrating by parts in time in (4.13) and using a density argument, we obtain for all $\psi_0 \in L^2(0, T; H^1(\Omega))$ and $\psi_1 \in L^2((0, T) \times \Omega; H_{per}^1(Y^f)/\mathbb{R})$ that

$$\begin{aligned}
 & |Y^f| \int_0^T \langle \partial_t c_{i,0}, \psi_0 \rangle_{H^1(\Omega)', H^1(\Omega)} dt \\
 & + D_i \int_0^T \int_{\Omega} \int_{Y^f} h'_p(c_{i,0}) (\nabla c_{i,0} + \nabla_y c_{i,1}) (\nabla \psi_0 + \nabla_y \psi_1) dy dx dt \tag{4.15} \\
 & + D_i z_i \int_0^T \int_{\Omega} \int_{Y^f} c_{i,0} (\nabla \phi_0 + \nabla_y \phi_1) (\nabla \psi_0 + \nabla_y \psi_1) dy dx dt = 0.
 \end{aligned}$$

(4.15) is the so-called two-scale homogenized model for the transport equation.

Now considering $\psi_{\epsilon}(t, x) := \psi_0(t, x) + \epsilon \psi_1(t, x, \frac{x}{\epsilon})$ as a test function in (2.4), by similar arguments (see [35] for the convergence of the integral over the microscopic boundary Γ_{ϵ}), we obtain the two-scale homogenized Poisson's equation given as follows: For all $\psi_0 \in L^2((0, T); H^1(\Omega))$ and $\psi_1 \in L^2((0, T) \times \Omega; H_{per}^1(Y^f)/\mathbb{R})$ it holds that

$$\begin{aligned}
 & \int_0^T \int_{\Omega} \int_{Y^f} (\nabla \phi_0 + \nabla_y \phi_1) (\nabla \psi_0 + \nabla_y \psi_1) dy dx dt \\
 & = |Y^f| \int_0^T \int_{\Omega} \sum_{i=1}^P z_i c_{i,0} \psi_0 dx dt + \int_0^T \int_{\Omega} \int_{\Gamma} \xi_1(x, y) \psi_0(t, x) dS(y) dx dt \\
 & + \int_0^T \int_{\partial\Omega} \xi_2(x) \psi_0(t, x) dS(x) dt. \tag{4.16}
 \end{aligned}$$

Choosing $\psi_0 = 0$ in (4.16), we get

$$\int_0^T \int_{\Omega} \int_{Y^f} [\nabla \phi_0 + \nabla_y \phi_1] \cdot \nabla_y \psi_1 dy dx dt = 0. \tag{4.17}$$

It is well-known, see for example [2], that for a given $\phi_0 \in L^2((0, T), H^1(\Omega))$ this problem admits a unique weak solution $\phi_1 \in L^2((0, T) \times \Omega; H_{per}^1(Y^f)/\mathbb{R})$ which has the representation

$$\phi_1(t, x, y) = \sum_{k=1}^n \partial_{x_k} \phi_0(t, x) w_k(y), \tag{4.18}$$

where w_k is the unique weak solution of (4.10). Now, choosing $\psi_0 = 0$ in the two-scale homogenized transport equation (4.15) and using the representation for ϕ_1 , we obtain

with similar arguments as above (we emphasize that $h'_p(c_{i,0}) \geq 1$)

$$c_{i,1}(t, x, y) = \sum_{k=1}^n \partial_{x_k} c_{i,0}(t, x) w_k(y).$$

Choosing $\psi_1 = 0$ in (4.16), we obtain with a standard calculation

$$\begin{aligned} \int_0^T \int_{\Omega} A_{hom} \nabla \phi_0 \nabla \psi_0 \, dx \, dt &= \int_0^T \int_{\Omega} \sum_{i=1}^P z_i c_{i,0} \psi_0 \, dx \, dt \\ &+ \frac{1}{|Y^f|} \int_0^T \int_{\Omega} \int_{\Gamma} \xi_1(x, y) \psi_0(t, x) \, dS(y) \, dx \, dt + \frac{1}{|Y^f|} \int_0^T \int_{\partial\Omega} \xi_2(x) \psi_0(t, x) \, dS(x) \, dt. \end{aligned}$$

This is the weak formulation of (4.8). With similar arguments we obtain from (4.15) with $\psi_1 = 0$ the weak formulation for (4.7). This completes the proof of (a).

The proof of (b) follows easily by noting that the last term of the left hand side of (4.12) goes to 0 in the limit as $\epsilon \rightarrow 0$, for the case $\alpha < \beta$.

□

5 Discussion and outlook

We performed a rigorous homogenization of a drift-diffusion model for multiple species with nonlinear (non-degenerate) diffusion terms involving the nonlinear function $h_p(r) = r + \eta r^p$, with $\eta > 0$ and $p \in [4, \infty)$. This nonlinear problem raises difficulties for the homogenization procedure. In particular, for getting uniform estimates of the solutions with respect to the scale parameter ϵ , it is necessary to exploit the structure of the system, which admits a nonnegative energy functional decreasing in time along solutions of the model. Combining such arguments with more classical energy estimates, uniform *a priori* estimates of the microscopic solutions could be established, and a homogenized model could be derived for different scalings of the microscopic problem.

The microscopic model as well as the homogenized (macroscopic) model both depend on the parameter $\eta > 0$. In [5] it has been shown that fixing all other parameters of the model, in the limit $\eta \rightarrow 0$, the solutions of the nonlinear drift-diffusion model (1.1)-(1.3) converge to a solution of a Poisson-Nernst-Planck (PNP) system (formally obtained by setting $\eta = 0$), thus leading to the following diagram:

$$\begin{array}{ccc} u_{\epsilon, \eta} & \xrightarrow{\eta \rightarrow 0, [5]} & u_{\epsilon, 0} \\ \epsilon \rightarrow 0 \downarrow & & \downarrow \epsilon \rightarrow 0 (?) \\ u_{0, \eta} & \xrightarrow{\eta \rightarrow 0 (?) } & u_{0, 0} \end{array}$$

Here, $u_{\epsilon, \eta}$ represents the solution of the nonlinear drift-diffusion system (1.1)-(1.3), $u_{\epsilon, 0}$ represents the solution of a PNP system, and $u_{0, \eta}$ represents the solution to the

homogenized model for fixed parameter η derived in Theorem 4.3. If we consider the case $\alpha = \beta$, the homogenized system for $u_{0,\eta}$ is again of the form (1.1)-(1.3), however with matrix-valued constant coefficients. Thus, by similar arguments like in [5], we may assume that we can pass to the limit $\eta \rightarrow 0$ to obtain a PNP model with solution $u_{0,0}$. The question which now arises is whether this limit problem can also be obtained by passing to the limit $\epsilon \rightarrow 0$ in the PNP-problem for $u_{\epsilon,0}$ (see diagram). Unfortunately, the convergence for $\eta \rightarrow 0$ takes place in function spaces with weaker regularity, and thus, the uniform (with respect to ϵ) estimates of the solutions $u_{\epsilon,\eta}$ cannot be transferred to the limit $u_{\epsilon,0}$ and cannot be used for passing to the limit in the microscopic PNP model. An alternative approach could be to show uniform error estimates (with respect to suitable norms) for the other three convergences in the diagram, which is a highly demanding aim to be addressed in future investigations.

A Auxiliary results

In this section we prove some auxiliary results which have been used to obtain the uniform estimates for the microscopic solutions $(c_{i,\epsilon}, \phi_\epsilon)$ in Section 3, and for the derivation of the macroscopic model in Section 4. We start with an approximation result.

Lemma A.1. *Let Ω be a bounded open set in \mathbb{R}^n and let $u \in L^2(0, T; H^1(\Omega)) \cap L^\infty((0, T) \times \Omega)$, $u \geq 0$ almost everywhere in $(0, T) \times \Omega$ with $\partial_t u \in L^2(0, T; H^1(\Omega)')$. For each fixed $\delta > 0$, there exists a sequence ψ_m in $C^\infty([0, T] \times \overline{\Omega})$ which satisfies for all m*

$$\psi_m(t, x) \in \left[-\frac{\delta}{2}, \|u\|_{L^\infty((0, T) \times \Omega)} + \frac{\delta}{2} \right] \quad \text{for all } (t, x) \in [0, T] \times \overline{\Omega},$$

and such that

- (i) ψ_m converges to u in $L^2(0, T; H^1(\Omega))$,
- (ii) for every sequence of functions v_m bounded in $L^2((0, T), H^1(\Omega))$ and $L^\infty((0, T) \times \Omega)$ with $v_m \rightarrow v$, $\nabla v_m \rightarrow \nabla v$ almost everywhere in $(0, T) \times \Omega$ for some $v \in L^2((0, T), H^1(\Omega))$, it holds that

$$\lim_{m \rightarrow \infty} \int_0^T \langle \partial_t \psi_m, v_m \rangle_{H^1(\Omega)', H^1(\Omega)} dt = \int_0^T \langle \partial_t u, v \rangle_{H^1(\Omega)', H^1(\Omega)} dt.$$

Proof. First, we observe from [41, Proposition 23.23, (iii)], there exists a sequence ϕ_m in $C^\infty([0, T] \times \overline{\Omega})$ such that

$$\phi_m \rightarrow u \text{ in } L^2(0, T; H^1(\Omega)) \text{ and } \partial_t \phi_m \rightarrow \partial_t u \text{ in } L^2(0, T; H^1(\Omega)'). \quad (\text{A.1})$$

Now, let $\delta > 0$ be any fixed number, and let $\rho_\delta \in C^\infty(\mathbb{R})$ be such that $\rho_\delta, \rho_\delta', \rho_\delta''$ are bounded functions in \mathbb{R} with $\rho_\delta(x) \in \left[-\frac{\delta}{2}, \|u\|_{L^\infty((0,T)\times\Omega)} + \frac{\delta}{2}\right]$ for all $x \in \mathbb{R}$ and

$$\rho_\delta(x) = x, \text{ if } x \in \left[-\frac{\delta}{4}, \|u\|_{L^\infty((0,T)\times\Omega)} + \frac{\delta}{4}\right].$$

For example, such a function can be obtained by mollifying the following function:

$$\tilde{\rho}_\delta(x) = \begin{cases} \|u\|_{L^\infty((0,T)\times\Omega)} + \frac{\delta}{2} & \text{if } x > \|u\|_{L^\infty((0,T)\times\Omega)} + \frac{\delta}{2}, \\ x & \text{if } x \in \left[-\frac{\delta}{2}, \|u\|_{L^\infty((0,T)\times\Omega)} + \frac{\delta}{2}\right], \\ -\frac{\delta}{2} & \text{if } x < -\frac{\delta}{2}. \end{cases}$$

For the sake of clarity, in the remaining part of the proof we skip the index δ and denote ρ_δ simply by ρ . Note that, $\rho(u) = u$, $\rho'(u) = 1$, $\rho''(u) = 0$ for almost every $(t, x) \in (0, T) \times \Omega$. Let $\|\rho'\|_{L^\infty(\mathbb{R})} < M$, for some $M > 0$. Since up to a subsequence, which is still indexed by m , ϕ_m converges to u almost everywhere in $(0, T) \times \Omega$ and ρ is continuous, we have

$$\rho(\phi_m) \rightarrow \rho(u) = u \text{ a.e. in } (0, T) \times \Omega.$$

Again $|\rho(\phi_m)| < C$ in $[0, T] \times \bar{\Omega}$, where $C > 0$ is some constant independent of m . So by the dominated convergence theorem, we have

$$\rho(\phi_m) \rightarrow u \text{ in } L^2(0, T; L^2(\Omega)). \quad (\text{A.2})$$

Again, for $j \in \{1, \dots, n\}$, up to a subsequence, still indexed by m , we have that

$$\frac{\partial}{\partial x_j} \rho(\phi_m) = \rho'(\phi_m) \frac{\partial \phi_m}{\partial x_j}$$

converges almost everywhere in $(0, T) \times \Omega$ to

$$\rho'(u) \frac{\partial u}{\partial x_j} = \frac{\partial u}{\partial x_j}.$$

Furthermore, it holds

$$\left| \frac{\partial}{\partial x_j} \rho(\phi_m) \right| = \left| \rho'(\phi_m) \frac{\partial \phi_m}{\partial x_j} \right| \leq M \left| \frac{\partial \phi_m}{\partial x_j} \right| \text{ a.e. in } (0, T) \times \Omega.$$

Thus, by the generalized dominated convergence theorem (see [14, Exercises 20, 21, p. 59]), we get

$$\frac{\partial}{\partial x_j} \rho(\phi_m) \rightarrow \frac{\partial u}{\partial x_j} \text{ in } L^2(0, T; L^2(\Omega)). \quad (\text{A.3})$$

Finally, we choose a subsequence of $\rho(\phi_m)$, still indexed by m , along which the above convergence results hold and set $\psi_m := \rho(\phi_m)$. This proves (i). To prove (ii) we use

$$\int_0^T \langle \partial_t \psi_m, v_m \rangle_{H^1(\Omega)', H^1(\Omega)} dt = \int_0^T \langle \partial_t \phi_m, \rho'(\phi_m) v_m \rangle_{H^1(\Omega)', H^1(\Omega)} dt. \quad (\text{A.4})$$

Due to the strong convergence of $\partial_t \phi_m$ in $L^2((0, T), H^1(\Omega)')$, the claim follows if we show the weak convergence of $\rho'(\phi_m)v_m$ in $L^2((0, T), H^1(\Omega))$. From the assumptions on v_m and similar arguments as above we get

$$\|\rho'(\phi_m)v_m\|_{L^2((0, T), H^1(\Omega))} \leq C,$$

and

$$\rho'(\phi_m)v_m \rightharpoonup v, \quad \nabla(\rho'(\phi_m)v_m) \rightharpoonup \nabla v \quad \text{a.e. in } (0, T) \times \Omega.$$

The weak convergence of $\rho'(\phi_m)v_m$ to v now follows from [22, (13.44) Theorem]. \square

Lemma A.2. *Let $c_{i,\epsilon}$ be a weak solution to the microscopic model (2.2)-(2.3). Then for each $\delta > 0$, the following equality holds in the space $L^1(0, T; \mathbb{R})$:*

$$\frac{d}{dt} \int_{\Omega_\epsilon} (c_{i,\epsilon} + \delta) \log(c_{i,\epsilon} + \delta) dx = \langle \partial_t c_{i,\epsilon}, \log(c_{i,\epsilon} + \delta) \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)}.$$

Proof. In order to prove the lemma, we show that the followings hold:

- (i) $\int_{\Omega_\epsilon} (c_{i,\epsilon} + \delta) \log(c_{i,\epsilon} + \delta) dx \in L^1(0, T)$.
- (ii) $\langle \partial_t c_{i,\epsilon}, \log(c_{i,\epsilon} + \delta) \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} \in L^1(0, T)$.
- (iii) For all $\psi \in C_0^\infty(0, T)$, we have

$$\begin{aligned} & \int_0^T \int_{\Omega_\epsilon} (c_{i,\epsilon} + \delta) \log(c_{i,\epsilon} + \delta) \psi'(t) dx dt \\ &= - \int_0^T \langle \partial_t c_{i,\epsilon}, \log(c_{i,\epsilon} + \delta) \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} \psi(t) dt. \end{aligned}$$

Statement (i) follows immediately from the fact that $c_{i,\epsilon} \in L^\infty((0, T) \times \Omega_\epsilon)$ and is non-negative. (ii) is also obtained easily by noting that

$$\begin{aligned} & \int_0^T \left| \langle \partial_t c_{i,\epsilon}, \log(c_{i,\epsilon} + \delta) \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} \right| dt \\ & \leq \int_0^T \|\partial_t c_{i,\epsilon}\|_{H^1(\Omega_\epsilon)'} \|\log(c_{i,\epsilon} + \delta)\|_{H^1(\Omega_\epsilon)} dt \\ & \leq \|\partial_t c_{i,\epsilon}\|_{L^2(0, T; H^1(\Omega_\epsilon)')} \|\log(c_{i,\epsilon} + \delta)\|_{L^2(0, T; H^1(\Omega_\epsilon))} < \infty. \end{aligned}$$

Next, let us prove (iii). Lemma A.1 guarantees the existence of a sequence ψ_m in $C^\infty([0, T] \times \overline{\Omega_\epsilon})$ converging strongly to $c_{i,\epsilon}$ in $L^2((0, T), H^1(\Omega_\epsilon))$ and for all m , the range of $\psi_m \subseteq [-\frac{\delta}{2}, \|c_{i,\epsilon}\|_{L^\infty((0, T) \times \Omega_\epsilon)} + \frac{\delta}{2}]$. Then,

$$\int_0^T \int_{\Omega_\epsilon} (c_{i,\epsilon} + \delta) \log(c_{i,\epsilon} + \delta) \psi'(t) dx dt$$

$$= \lim_{m \rightarrow \infty} \int_0^T \int_{\Omega_\epsilon} [\psi_m + \delta] \log[\psi_m + \delta] \psi'(t) dx dt. \quad (\text{A.5})$$

Note that $\log[\psi_m + \delta]$ is well-defined, since $\psi_m \geq -\frac{\delta}{2}$. After an integration by parts in time, the right hand side of (A.5) becomes

$$\begin{aligned} & - \lim_{m \rightarrow \infty} \int_0^T \int_{\Omega_\epsilon} \partial_t \psi_m \log[\psi_m + \delta] \psi(t) dx dt \\ & - \lim_{m \rightarrow \infty} \int_0^T \int_{\Omega_\epsilon} \partial_t \psi_m \psi(t) dx dt =: -I - II. \end{aligned}$$

Choosing $v_m := \log(\psi_m + \delta)\psi$ in Lemma A.1 (ii), it is easy to check that by the convergence of ψ_m in $L^2((0, T), H^1(\Omega))$, the sequence v_m fulfills the assumptions of Lemma A.1 with $v = \log(c_{i,\epsilon} + \delta)\psi$. Hence, we obtain

$$I = \int_0^T \langle \partial_t c_{i,\epsilon}, \log(c_{i,\epsilon} + \delta) \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} \psi(t) dt.$$

In the same way we obtain

$$II = \int_0^T \langle \partial_t c_{i,\epsilon}, 1 \rangle_{H^1(\Omega_\epsilon)', H^1(\Omega_\epsilon)} \psi(t) dt = 0,$$

where the last equality follows from testing the equation (2.2) with 1. \square

Lemma A.3. *For $1 \leq q < \infty$ there exists an extension operator $\tilde{\cdot} : W^{1,q}(\Omega_\epsilon) \rightarrow W^{1,q}(\Omega)$ such that for all $u_\epsilon \in W^{1,q}(\Omega_\epsilon)$, we have*

$$\|\tilde{u}_\epsilon\|_{L^q(\Omega)} \leq C \|u_\epsilon\|_{L^q(\Omega_\epsilon)}, \quad \|\nabla \tilde{u}_\epsilon\|_{L^q(\Omega)} \leq C \|\nabla u_\epsilon\|_{L^q(\Omega_\epsilon)}, \quad (\text{A.6})$$

with a constant $C > 0$ independent of ϵ .

Furthermore, if $u_\epsilon \in W^{1,q}(\Omega_\epsilon) \cap L^\infty(\Omega_\epsilon)$, then $\tilde{u}_\epsilon \in L^\infty(\Omega)$ and

$$\|\tilde{u}_\epsilon\|_{L^\infty(\Omega)} \leq C \|u_\epsilon\|_{L^\infty(\Omega_\epsilon)},$$

with a constant $C > 0$ independent ϵ , and if $u_\epsilon \in W^{1,q}(\Omega_\epsilon)$ is a non-negative function, so is the extension \tilde{u}_ϵ .

Proof. The existence of an extension operator satisfying the estimates (A.6) is by now a standard result in homogenization theory, see e.g. [9] and [1], and is based on a similar result in the standard cell followed by a decomposition of the domain Ω_ϵ in ϵ -cells and a scaling argument. Since however, we need further properties of the extension operator (like non-negativity and essential boundedness), let us sketch the construction of the extension operator in the standard cell, from which the additional properties can be derived.

In a first step we extend a function $u \in W^{1,q}(Y^f)$ by a reflection method to the whole cell Y . More precisely, we define the extension operator $P : W^{1,q}(Y^f) \rightarrow W^{1,q}(Y)$ in

the following way (see [8, Theorem 9.7] for more details and [34] for more general settings and higher order derivatives): let U_1, \dots, U_k be an open covering of Γ such that $U_i \subset Y$ and assume $\theta_i \in C_0^\infty(U_i)$ for all $i = 1, \dots, k$; $\theta_0 \in C^\infty(\bar{Y})$, $\text{supp } \theta_0 \subset \bar{Y} \setminus \Gamma$, with $\sum_{i=0}^k \theta_i = 1$ and $0 \leq \theta_i \leq 1$ for $i = 0, \dots, k$. We define

$$\bar{u}_0(y) := \begin{cases} \theta_0(y)u(y) & \text{if } y \in Y^f, \\ 0 & \text{if } y \in Y \setminus Y^f; \end{cases} \quad \hat{u}_i(y) := \begin{cases} \theta_i(y)w_i(y) & \text{if } y \in U_i, \\ 0 & \text{if } y \in Y \setminus U_i, \end{cases} \quad (\text{A.7})$$

for $1 \leq i \leq k$, where the functions w_i represent the extension of $u|_{U_i \cap Y^f}$ to U_i by reflection. We emphasize that

$$\|w_i\|_{L^\infty(U_i)} \leq \|u\|_{L^\infty(U_i \cap Y^f)}. \quad (\text{A.8})$$

Then we define the extension operator

$$Pu = \bar{u}_0 + \sum_{i=1}^k \hat{u}_i.$$

From (A.7) and (A.8) we immediately obtain for $u \in W^{1,q}(Y^f) \cap L^\infty(Y^f)$ and non-negative that

$$0 \leq Pu(y) \leq \|u\|_{L^\infty(Y^f)} \quad \text{f.a.e. } y \in Y. \quad (\text{A.9})$$

The crucial point in the construction of the global extension operator on Ω_ε is that the norm of the gradient of the extension can be estimated by norm of the gradient of the function itself. However, in general this is not the case for the operator P . Hence, in the second step, we construct a local extension operator $S : W^{1,q}(Y^f) \rightarrow W^{1,q}(Y)$ such that

$$\|\nabla Su\|_{L^q(Y)} \leq C\|\nabla u\|_{L^q(Y^f)}. \quad (\text{A.10})$$

Since Y^f is connected, the existence of such an operator is well-known, see [9] or [1]. We only have to show that the operator S also preserves the non-negativity and essential boundedness of a function. We denote the mean-value on Y^f of a function u by $(u)_{Y^f}$. Now, we define as in [9, Lemma 3] the extension operator

$$Su := P(u - (u)_{Y^f}) + (u)_{Y^f},$$

which especially fulfills (A.10). Further, for $u \in W^{1,q}(Y^f) \cap L^\infty(Y^f)$ non-negative, we obtain from (A.9)

$$\|Su\|_{L^\infty(Y)} \leq \|u\|_{L^\infty(Y^f)} + 2|(u)_{Y^f}| \leq C\|u\|_{L^\infty(Y^f)},$$

and

$$Su = Pu - \underbrace{P(u)_{Y^f}}_{\geq -(u)_{Y^f}} + (u)_{Y^f} \geq 0.$$

This completes the proof. \square

Lemma A.4. For all $u_\epsilon \in H^1(\Omega_\epsilon)$ with $\frac{1}{|\Omega_\epsilon|} \int_{\Omega_\epsilon} u_\epsilon dx = 0$, we have

$$\|u_\epsilon\|_{L^2(\Omega_\epsilon)} \leq C \|\nabla u_\epsilon\|_{L^2(\Omega_\epsilon)}, \quad (\text{A.11})$$

with a constant $C > 0$ independent of ϵ .

Proof. The proof is similar to the proof of [31, Lemma 1.3]. Since it is rather short, we include it here for the sake of completeness. Let \tilde{u}_ϵ be the extension of u_ϵ to Ω given in Lemma A.3. Now, we use the Poincaré–Wirtinger inequality and the fact that due to the zero mean value of u_ϵ on Ω_ϵ we have $\int_\Omega \tilde{u}_\epsilon = \int_{\Omega \setminus \Omega_\epsilon} \tilde{u}_\epsilon$. We obtain

$$\begin{aligned} \|\tilde{u}_\epsilon\|_{L^2(\Omega)} &\leq \left\| \tilde{u}_\epsilon - \frac{1}{|\Omega|} \int_\Omega \tilde{u}_\epsilon \right\|_{L^2(\Omega)} + \frac{1}{|\Omega|} \left| \int_\Omega \tilde{u}_\epsilon \right| |\Omega|^{\frac{1}{2}} \\ &\leq C(\Omega) \|\nabla \tilde{u}_\epsilon\|_{L^2(\Omega)} + \frac{1}{|\Omega|^{\frac{1}{2}}} \int_{\Omega \setminus \Omega_\epsilon} |\tilde{u}_\epsilon| dx \\ &\leq C(\Omega) \|\nabla \tilde{u}_\epsilon\|_{L^2(\Omega)} + \frac{|\Omega \setminus \Omega_\epsilon|^{\frac{1}{2}}}{|\Omega|^{\frac{1}{2}}} \|\tilde{u}_\epsilon\|_{L^2(\Omega \setminus \Omega_\epsilon)} \\ &\leq C(\Omega) \|\nabla \tilde{u}_\epsilon\|_{L^2(\Omega)} + \frac{|\Omega \setminus \Omega_\epsilon|^{\frac{1}{2}}}{|\Omega|^{\frac{1}{2}}} \|\tilde{u}_\epsilon\|_{L^2(\Omega)}. \end{aligned}$$

Since $\frac{|\Omega \setminus \Omega_\epsilon|^{\frac{1}{2}}}{|\Omega|^{\frac{1}{2}}} < C_1 < 1$, with C_1 is independent of ϵ , we have

$$(1 - C_1) \|\tilde{u}_\epsilon\|_{L^2(\Omega)} \leq C(\Omega) \|\nabla \tilde{u}_\epsilon\|_{L^2(\Omega)}.$$

Now using the properties (A.6) of the extension \tilde{u}_ϵ , we conclude the proof. \square

The following Lemma gives a trace-estimate with an explicit dependence of ϵ and is well-known in the theory of homogenization, so we skip the proof, which is based on a standard decomposition argument and the trace-inequality in the reference element.

Lemma A.5. For every $u_\epsilon \in W^{1,q}(\Omega_\epsilon)$ with $q \in [1, \infty)$ it holds that

$$\epsilon \|u_\epsilon\|_{L^q(\Gamma_\epsilon)}^q \leq C \left(\|u_\epsilon\|_{L^q(\Omega_\epsilon)}^q + \epsilon^q \|\nabla u_\epsilon\|_{L^q(\Omega_\epsilon)}^q \right),$$

for a constant $C > 0$ independent of ϵ .

Acknowledgements

AB acknowledges the support by the RTG 2339 “Interfaces, Complex Structures, and Singular Limits” of the German Science Foundation (DFG). MG was supported by the SCIDATOS project, funded by the Klaus Tschira Foundation (grant 00.0277.2015).

References

- [1] E. Acerbi, V. Chiadò, G. D. Maso, and D. Percivale. An extension theorem from connected sets, and homogenization in general periodic domains. *Nonlinear Analysis, Theory, Methods & Applications*, 18(5):481–496, 1992.
- [2] G. Allaire. *Homogenization and two-scale convergence*. SIAM J. Math. Anal. 23 (1992), no. 6, 1482–1518.
- [3] G. Allaire, A. Mikelić, A. Piatnitski. *Homogenization of the linearized ionic transport equations in rigid periodic porous media*. J. Math. Phys. 51 (2010), no. 12, 123103, 18 pp.
- [4] J.-L. Auriault, J. Lewandowska. *On the cross-effects of coupled macroscopic transport equations in porous media*. Transp. Porous Media 16 (1994) 31–52.
- [5] D. Bothe, A. Fischer, M. Pierre and G. Rolland. *Global existence for diffusion-electromigration systems in space dimension three and higher*. Nonlinear Anal. 99:152–166, 2014.
- [6] D. Bothe, A. Fischer, and J.Saal. *Global well-posedness and stability of electrokinetic flows*. SIAM J. Math. Anal. 46:1263–1316 2014.
- [7] A. Bourgeat, S. Luckhaus, and A. Mikelić. *Convergence of the homogenization process for a double-porosity model of immiscible two-phase flow*. SIAM J. Math. Anal., 27:1520–1543, 1996.
- [8] H. Brezis. *Functional analysis, Sobolev spaces and partial differential equations*. Springer, New York, 2011.
- [9] D. Cioranescu and J. S. J. Paulin. *Homogenization in open sets with holes*. J. Math. Anal. Appl. no. 2, 590–607, 1979.
- [10] D. Cioranescu and P. Donato. *An introduction to homogenization*. Oxford University Press, 1999.
- [11] G. W. Clark and R. E. Showalter. Two-scale convergence of a model for flow in a partially fissured medium. *Electron. J. Differential Equations*, 1999(02):1–20, 1999.
- [12] Z. Deyl. *Electrophoresis: A Survey of Techniques and Applications*. Elsevier, Amsterdam, 1979.
- [13] L. C. Evans. *Partial differential equations*. Second edition. American Mathematical Society, 2010.
- [14] G. B. Folland. *Real analysis: modern techniques and their applications*. Second edition. John Wiley & Sons, 1999.
- [15] M. Gahn, M. Neuss-Radu and P. Knabner. *Homogenization of reaction-diffusion processes in a two-component porous medium with nonlinear flux conditions at the interface*. SIAM J. Appl. Math. 76 (2016), no. 5, 1819–1843.
- [16] M. Gahn. *Singular limit for reactive transport through a thin heterogeneous layer including a nonlinear diffusion coefficient*. Comm. Pure Appl. Anal. 2021.
- [17] H. Gajewski, K. Gröger. *On the Basic Equations for Carrier Transport in Semiconductors*. J. Math. Anal. Appl. 113 (1996) 109–130.
- [18] H. Gajewski, K. Gröger. *Reaction-diffusion processes of electrically charged species*. Math. Nachr. 177 (1996) 109–130.
- [19] G. P. Galdi. *An introduction to the mathematical theory of the Navier-Stokes equations. Steady-state problems*. Second edition. Springer, New York, 2011.

- [20] A. Glitzky, K. Gröger, R. Hünlich. *Existence, uniqueness and asymptotic behavior of solutions to equations modelling transport of dopants in semiconductors*. In: Special topics in semiconductor analysis (Edited by J. Frehse and H. Gajewski), Bonner Mathematische Schriften 258 (1994) 49–78.
- [21] P. Grisvard. *Elliptic problems in nonsmooth domains*. Pitman Advanced Publishing Program, Boston, 1985.
- [22] E. Hewitt and K. Stromberg. *Real and Abstract Analysis*. Springer-Verlag, New York, 1975.
- [23] A. Jüngel. *A nonlinear drift-diffusion system with electric convection arising in electrophoretic and semiconductor modeling*. Math. Nachr. 185 (1997), 85–110.
- [24] J. Keener, J. Sneyd. *Mathematical Physiology*. Springer Verlag, New York, Inc., 1998.
- [25] V. A. Khoa, A. Muntean. *Corrector homogenization estimates for a non-stationary Stokes-Nernst-Planck-Poisson system in perforated domains*. Commun. Math. Sci. 17 (2019), no. 3, 705–738.
- [26] V. A. Kovtunenکو, A. V. Zubkova. *Homogenization of the generalized Poisson–Nernst–Planck problem in a two-phase medium: correctors and estimates*. Appl. Anal. 100 (2021), 253–274
- [27] O. A. Ladyženskaja, V. A. Solonnikov and N. N. Ural’ceva. *Linear and quasilinear equations of parabolic type* (translated from the Russian by S. Smith). American Mathematical Society, 1988.
- [28] F. List, K. Kumar, I.S. Pop, and F.A. Radu. *Rigorous upscaling of unsaturated flow in fractured porous media*. SIAM. J. Math. Anal. 52:239–276, 2020.
- [29] D. Lukkassen, G. Nguetseng and P. Wall. *Two-scale convergence*. Int. J. Pure Appl. Math. 2 (2002), no. 1, 35–86.
- [30] P. A. Markowich, C.A. Ringhofer, C. Schmeiser. *Semiconductor Equations*. Springer Verlag, 1990.
- [31] A. Mikelić. *Homogenization theory and applications to filtration through porous media*. In Filtration in Porous Media and Industrial Application, Springer-Verlag Berlin Heidelberg, 127-214, 2000.
- [32] V. P. Mikhailov. *Partial differential equations* (translated from the Russian by P. C. Sinha). Mir Publishers, Moscow, 1978.
- [33] C. Moyne, M. A. Murad. *Electro-chemo-mechanical couplings in swelling clays derived from a micro/macro-homogenization procedure*. Internat. J. Solids Structures 39 (2006) 6159–6190.
- [34] J. Nečas. *Direct methods in the theory of elliptic equations* (translated by Gerard Tronel and Alois Kufner). Springer, Heidelberg, 2012.
- [35] M. Neuss-Radu. *Some extensions of two-scale convergence*. C. R. Acad. Sci. Paris Sér. I Math. 322 (1996), no. 9, 899–904.
- [36] G. Nguetseng. *A general convergence result for a functional related to the theory of homogenization*. SIAM J. Math. Anal. 20 (1989), no. 3, 608–623.
- [37] L. Nirenberg. *On elliptic partial differential equations*. Ann. Scuola Norm. Sup. Pisa Sci. Fis. Math., 13:116–162, 1959.
- [38] N. Ray, A. Muntean, P. Knabner. *Rigorous homogenization of a Stokes-Nernst-Planck-Poisson system*. J. Math. Anal. Appl. 390 (2012), no. 1, 374–393.
- [39] M. Schmuck. *Modelling and deriving porous media Stokes-Poisson-Nernst-Planck equations by a multi-scale approach*. Commun. Math. Sci. 9 (2011) 685–710.

- [40] D. Wachsmuth. *The regularity of the positive part of functions in $L^2(I; H^1(\Omega)) \cap H^1(I; H^1(\Omega)^*)$ with applications to parabolic equations*. *Comment. Math. Univ. Carolin.* 57 (2016), no. 3, 327–332.
- [41] E. Zeidler. *Nonlinear functional analysis and its applications. II/A: linear monotone operators* (translated by the author and L. F. Boron). Springer, 1990.