

# INVOLUTIONS, KNOTS, AND FLOER $K$ -THEORY

HOKUTO KONNO, JIN MIYAZAWA, AND MASAKI TANIGUCHI

ABSTRACT. We establish a version of Seiberg–Witten Floer  $K$ -theory for knots, as well as a version of Seiberg–Witten Floer  $K$ -theory for 3-manifolds with involutions. The main theorem is a  $10/8$ -type inequality for spin 4-manifolds with boundary and with involutions, and that yields numerous applications to knots, such as genus bounds and stabilizing numbers. We also give applications to get obstructions to extending involutions on 3-manifolds to spin 4-manifolds.

## CONTENTS

1. Introduction	2
1.1. Main theorem	2
1.2. A knot concordance invariant	3
1.3. Applications to H-sliceness	4
1.4. Applications to non-extendable actions	6
1.5. Floer homotopy type and Floer $K$ -theory	7
1.6. Structure of the paper	8
2. An involution on the configuration space	8
3. Seiberg–Witten Floer homotopy theory for involutions	10
3.1. Representations	10
3.2. Space of type $G$ -SWF	12
3.3. Doubling construction and spectrum classes	14
3.4. Doubled Seiberg–Witten Floer stable homotopy type for involutions	17
3.5. Seiberg–Witten Floer $K$ -theory for involutions	20
3.6. Cobordisms	21
3.7. Proof of Theorem 1.1	23
3.8. Calculations	25
3.9. Knot invariants	30
4. Applications to knot theory	34
4.1. Branched covers of punctured 4-manifolds	34
4.2. Genus bounds from Theorem 1.1	36
5. Non-extendable actions	42
6. Appendix	44
6.1. Manolescu’s relative $10/8$ -inequality	44
6.2. Topological obstructions	45
References	46

## 1. INTRODUCTION

The purpose of this paper is to establish a version of Seiberg–Witten Floer  $K$ -theory for knots, as well as a version of Seiberg–Witten Floer  $K$ -theory for 3-manifolds with involutions. Our construction is based on Manolescu’s Seiberg–Witten Floer stable homotopy type [53] and an involutive symmetry on the Seiberg–Witten equations introduced by Kato [39]. Notably, Kato’s involutive symmetry acts on the spinors as an anti-complex linear involution. The study of anti-linear involutions was started by Atiyah [4] and developed as  $KR$ -theory, and Kato’s work [39] may be regarded as a non-linear version of Atiyah’s argument in Seiberg–Witten theory for closed 4-manifolds. In this paper, we develop a 3-dimensional version of [39] and extend Kato’s work to 4-manifolds with boundary. Our main theorem is a 10/8-type inequality for spin 4-manifolds with boundary and with involutions, and that yields numerous interesting applications to knots, such as genus bounds and stabilizing numbers. While various researchers have studied effects of group actions in several types of Floer homology and their applications to knots, such as [2, 8, 16, 24–30, 36–38, 49–52], to the best of our knowledge, this paper is the first study of Floer  $K$ -theory defined for knots with a 10/8-type theorem. Like the original 10/8-inequality by Furuta [19] and its generalization to 4-manifolds with boundary by Manolescu [54], constraints on knots obtained from our 10/8-type inequality are much different from constraints obtained from usual (i.e. ordinary cohomological) equivariant Floer theory. Our 10/8-type inequality also yields applications to get obstructions to extending involutions on 3-manifolds to spin 4-manifolds.

**1.1. Main theorem.** Let  $(Y, \mathfrak{t})$  be a spin rational homology 3-sphere and  $\iota$  be an orientation-preserving smooth involution whose fixed point set is non-empty and of codimension 2. Suppose that  $\iota$  preserves the spin structure  $\mathfrak{t}$ . (If  $Y$  is a  $\mathbb{Z}_2$ -homology sphere, this is the case for all  $\iota$ .) We shall define a numerical invariant

$$\kappa(Y, \mathfrak{t}, \iota) \in \mathbb{Q},$$

which we call the  *$K$ -theoretic Frøyshov invariant of the spin 3-manifold with involution*  $(Y, \mathfrak{t}, \iota)$ . This is a version (taking effects of  $\iota$  into account) of the  $K$ -theoretic invariant  $\kappa$  introduced by Manolescu [54], which was used in [54] to establish a relative version of Furuta’s 10/8-inequality [19].

The main theorem of this paper is a 10/8-type inequality for spin 4-manifolds with involutions:

**Theorem 1.1.** *Let  $(Y_0, \mathfrak{t}_0)$ ,  $(Y_1, \mathfrak{t}_1)$  be spin rational homology 3-spheres. Let  $\iota_0, \iota_1$  be smooth involutions on  $Y_0, Y_1$ . Suppose that  $\iota_0, \iota_1$  preserve the given orientations and spin structures  $\mathfrak{t}_0, \mathfrak{t}_1$  on  $Y_0, Y_1$  respectively, and suppose that the invariant sets of  $\iota_0, \iota_1$  are non-empty and of codimension 2. Let  $(W, \mathfrak{s})$  be a smooth compact oriented spin cobordism with  $b_1(W) = 0$  from  $(Y_0, \mathfrak{t}_0)$  to  $(Y_1, \mathfrak{t}_1)$ . Suppose that there exists a smooth involution  $\iota$  on  $W$  such that  $\iota$  preserves the given orientation and spin structure  $\mathfrak{s}$  on  $W$ , and that the restriction of  $\iota$  to the boundary is given by  $\iota_0, \iota_1$ . Then we have*

$$(1) \quad -\frac{\sigma(W)}{16} + \kappa(Y_0, \mathfrak{t}_0, \iota_0) \leq b^+(W) - b_\iota^+(W) + \kappa(Y_1, \mathfrak{t}_1, \iota_1),$$

where  $b_\iota^+(W)$  denotes the maximal dimension of  $\iota$ -invariant positive-definite subspaces of  $H^2(W; \mathbb{R})$ .

*Remark 1.2.* When  $Y_0 = Y_1 = S^3(\subset \mathbb{C}^2)$  and  $\iota_0, \iota_1$  are the complex conjugations, we obtain

$$-\frac{\sigma(W)}{16} \leq b^+(W) - b_\iota^+(W),$$

which recovers Kato's 10/8-inequality [39, Theorem 2.3]. There are variants of 10/8-inequalities taking effects of finite groups actions into account: see, for example, [11, 18, 63].

To define the  $K$ -theoretic Frøyshov invariant for involutions and prove Theorem 1.1, we shall construct a version of Seiberg–Witten Floer homotopy type, which we denote by

$$DSWF_G(Y, \mathfrak{t}, \iota),$$

taking effects of  $\iota$  into account. Here  $G$  stands for  $G = \mathbb{Z}_4$ , which is the subgroup of  $Pin(2)$  generated by  $j \in Pin(2)$ , and  $D$  stands for a “doubling” construction introduced in Subsection 3.3. On the construction of  $DSWF_G(Y, \mathfrak{t}, \iota)$ , see Subsection 1.5 for more details.

Moreover, we can generalize Theorem 1.1 to fixed-point free odd involutions. See Theorem 3.25 and Theorem 3.27.

**1.2. A knot concordance invariant.** Applying equivariant Floer theory to the double branched covers, several knot (concordance) invariants are defined in Heegaard Floer homology, such as [2, 16, 24, 25, 27, 28, 38], and Seiberg–Witten Floer theory [8]. In a certain orbifold setting, several versions of knot instanton Floer homology are developed ([13, 15, 46]). We also provide a knot concordance invariant from our Floer  $K$ -theory.

For a given oriented knot  $K$  in  $S^3$ , we can associate an oriented rational homology 3-sphere  $\Sigma(K)$  called the branched covering space along  $K$ , which is equipped with an involution  $\iota_K$ . Since it is known that  $H^1(\Sigma(K); \mathbb{Z}/2) = 0$ , the spin structure on  $\Sigma(K)$  is unique. We denote by  $\mathfrak{t}$  the unique spin structure on  $\Sigma(K)$ . Define the  $K$ -theoretic Frøyshov invariant for the knot  $K$  by

$$\kappa(K) := \kappa(\Sigma(K), \mathfrak{t}, \iota_K) \in \mathbb{Q}.$$

As a corollary of Theorem 1.1, we prove the following 10/8-type inequality:

**Theorem 1.3.** *The invariant  $\kappa(K)$  satisfies the following properties:*

- (i) *The invariant  $\kappa(K)$  is a knot concordance invariant.*
- (ii) *Let  $X$  be a simply-connected compact oriented smooth 4-manifold  $X$  with boundary  $\partial = S^3$ . Let  $S$  be an oriented compact connected properly smoothly embedded surface in  $X$  bounded by  $K$  such that the homology class  $[S]$  of  $S$  is divisible by 2 and  $PD(w_2(X)) = [S]/2 \pmod{2}$ . Then, we have*

$$(2) \quad -\frac{\sigma(X)}{8} + \frac{9}{32}[S]^2 - \frac{9}{16}\sigma(K) \leq b^+(X) + g(S) + \kappa(K),$$

where  $\sigma(K)$  denotes the signature of  $K$  (with the sign convention  $\sigma(T(2, 3)) = -2$ ) and  $g(S)$  is the genus of  $S$ .

*Remark 1.4.* When  $K$  is the unknot and  $S$  is an embedded null-homologous disk, the inequality (2) implies

$$-\frac{\sigma(X)}{8} \leq b^+(X),$$

which recovers Furuta's original 10/8-inequality [19] except for adding 1 on the left-hand side. Note that Manolescu's relative 10/8-inequality [54] applied to the branched covering spaces also gives a similar inequality to (2). In Remark 4.7, we compare such known inequalities with (2). Also, the invariant  $\kappa$  can be extended to an invariant of a pair  $(Y, K)$  of an oriented homology 3-sphere  $Y$  and an oriented knot  $K$  in  $Y$ . For more details, see Remark 3.41.

For two bridge knots, we can compute  $\kappa$ :

**Theorem 1.5.** *For any coprime integers  $p, q$ , the two bridge knot  $K(p, q)$ , whose branched cover is the lens space  $L(p, q)$ , we have*

$$(3) \quad \kappa(K(p, q)) = -\frac{1}{16}\sigma(K(p, q)).$$

The equation (3) also holds for any finite connected sum of two bridge knots.

Further, we also compute  $\kappa$  for a certain class of torus knots:

**Theorem 1.6.** *For any positive integers  $n$  and  $m$ , we have*

$$\begin{aligned} \kappa(\#_m T(3, 12n - 5)) &= -m/2, & \kappa(\#_m T(3, 12n - 5)^*) &= m/2, \\ \kappa(\#_m T(3, 12n - 1)) &= 0, & \kappa(\#_m T(3, 12n - 1)^*) &= 0, \\ \kappa(\#_m T(3, 12n - 7)) &= m/2, & \kappa(\#_m T(3, 12n - 7)^*) &= -m/2, \\ \kappa(\#_m T(3, 12n + 1)) &= 0, & \kappa(\#_m T(3, 12n + 1)^*) &= 0, \end{aligned}$$

where  $K^*$  denotes the mirror image of  $K$ .

**1.3. Applications to H-sliceness.** It is one of the most important problems in low dimensional topology to study various notions of sliceness for knots in  $S^3$ . Recently, Manolescu, Marengon, Sarkar and Willis [57] defined the notion of H-slice. A given knot  $K$  in  $S^3$  is said to be *smoothly (resp. topologically) H-slice* in an oriented closed smooth 4-manifold  $X$  if  $K$  bounds a properly embedded smooth (resp. locally flat) null-homologous disk in  $X \setminus \text{Int } D^4$ . There are various known smooth obstructions to H-sliceness in both definite 4-manifolds (for examples, see [47, 57, 66]) and indefinite 4-manifolds [34, 61]. For topological obstructions, see [20, 42, 56, 67]. Some of these techniques can be used to give lower bounds on the following genus:

$$g_X(K) := \min \left\{ g(S) \mid \left. \begin{array}{l} S \text{ is a properly and smoothly embedded null-homologous} \\ \text{orientable surface in } X \setminus \text{int } D^4 \text{ such that } \partial S = K \end{array} \right\}.$$

On the other hand, for indefinite 4-manifolds with vanishing gauge-theoretic invariant, there are few ways to obstruct H-sliceness, except for an approach based on 10/8-inequality [56] and topological obstructions coming from the Arf invariant ([17, 67]) and the Tristram–Levine signature function ([14]). We focus on such a class of 4-manifolds, in particular the  $n$ -fold connected sums  $\#_n S^2 \times S^2$ ,  $\#_n K3$ , and  $\#_n \mathbb{C}P^2 \#_n (-\mathbb{C}P^2)$  of  $S^2 \times S^2$ ,  $K3$  and  $\mathbb{C}P^2 \# (-\mathbb{C}P^2)$  respectively.

First we consider H-sliceness in the  $n$ -fold connected sum of  $S^2 \times S^2$ . It is proven in [70] that, for any knot  $K$  in  $S^3$  whose Arf invariant  $\text{Arf}(K)$  is zero, there is a positive integer  $N$  such that  $K$  is smoothly H-slice in  $\#_N S^2 \times S^2$ . This result enables us to define invariants

$$sn(K) := \min\{N \mid K \text{ is smoothly H-slice in } \#_N S^2 \times S^2\}$$

and

$$sn^{Top}(K) := \min\{N \mid K \text{ is topologically H-slice in } \#_N S^2 \times S^2\}$$

when  $\text{Arf}(K)$  is zero. These quantities have been studied in the literature, and for examples in [14], the invariant  $sn(K)$  (resp.  $sn^{Top}(K)$ ) is called the *smooth (resp. topological) stabilizing number* of  $K$ .

Our invariant  $\kappa(K)$  can be used to give a lower bound on  $sn(K)$ :

**Theorem 1.7.** *For any knot  $K \subset S^3$  with  $\text{Arf}(K) = 0$ , we have*

$$-\frac{9}{16}\sigma(K) - \kappa(K) \leq sn(K).$$

It is a natural and basic question if the topological stabilizing number and smooth stabilizing number have essential difference. More concretely, it is asked in [14, Question 1.4] whether there exists a knot  $K$  such that

$$0 < sn^{Top}(K) < sn(K).$$

Using Theorems 1.6 and 1.7, we can give an affirmative answer to this question:

**Theorem 1.8.** *For a positive integer  $l$  and a positive even integer  $m$ , we have the following estimates:*

$$\begin{aligned} 9ml &\leq sn(\#_m T(3, 12l - 1)), \\ 9ml &\leq sn(\#_m T(3, 12l + 1)), \\ 9ml - 4m &\leq sn(\#_m T(3, 12l - 5)), \text{ and} \\ 9ml - 5m &\leq sn(\#_m T(3, 12l - 7)). \end{aligned}$$

On the other hand, we also have

$$\begin{aligned} sn^{Top}(\#_m T(3, 12l - 1)) &= 8ml, \\ 8ml &\leq sn^{Top}(\#_m T(3, 12l + 1)) \leq 8ml + m, \\ 8ml - 4m &\leq sn^{Top}(\#_m T(3, 12l - 5)) \leq 8ml - 3m, \text{ and} \\ sn^{Top}(\#_m T(3, 12l - 7)) &= 8ml - 4m. \end{aligned}$$

In particular, if  $K$  is one of the following:  $T(3, 12l - 1)$ ,  $T(3, 12(l+1)+1)$ ,  $T(3, 12(l+1) - 5)$ , and  $T(3, 12(l+1) - 7)$  for  $l > 0$ , and if we set  $K_n = \#_{2n} K$  for  $n > 0$ , then we have

$$0 < sn^{Top}(K_n) < sn(K_n)$$

for all  $n$  and the sequence  $\{K_n\}_{n=1}^{\infty}$  satisfies that

$$\lim_{n \rightarrow \infty} (sn(K_n) - sn^{Top}(K_n)) = +\infty.$$

The equalities for  $sn^{Top}$  in Theorem 1.8 follow from results by [7, Theorem 1], [14, Theorem 5.15], and the lower bounds on  $sn$  in Theorem 1.8 shall be deduced from Theorems 1.6 and 1.7.

Next we focus on  $\#_n K3$ . Our invariant  $\kappa(K)$  can be used to give a lower bound on  $g_{\#_n K3}$ :

**Theorem 1.9.** *Let  $n$  be a non-negative integer. Let  $K$  be a finite connected sum of two bridge knots with  $\text{Arf}(K) = 0$ . Then we have*

$$-\frac{1}{2}\sigma(K) - n \leq g_{\#_n K3}(K).$$

Also, the same inequality holds for any homotopy  $\#_n K3$ .

*Example 1.10.* Let  $n$  be a non-negative integer. Let  $K_m$  be the  $m$ -fold connected sum of  $5_2^*$ . The signature of  $5_2^*$  is  $\sigma(5_2^*) = -2$ . (See [1]. Note that our sign convention of the knot signature is opposite to The Knot Atlas. See Remark 3.45.) Thus we have

$$m - n \leq g_{\#_n K3}(K_m).$$

The same conclusion holds for any homotopy  $\#_n K3$ .

In the end of this subsection, we give relative genus bounds for  $\#_n \mathbb{C}P^2 \#_n (-\mathbb{C}P^2)$ .

*Example 1.11.* Let  $K_n$  be the  $n$ -fold connected sum of  $T(3, 7)$  for a positive integer  $n$  and  $X_n$  be  $\#_n \mathbb{C}P^2 \#_n (-\mathbb{C}P^2)$ . Then,

$$\frac{7}{4}n \leq g_{(X_n, x)}(K_n),$$

where  $x = (2, \dots, 2) \in H_2(\#_n \mathbb{C}P^2 \#_n (-\mathbb{C}P^2))$ .

*Remark 1.12.* Using the Tristram–Levine signature function, another inequality (Remark 4.7) holds for the topological (locally flat) embeddings. Also, Manolescu’s relative 10/8-inequality ([54]) gives a similar inequality. We compare our result with such known results in Remarks 4.7 and 4.13 and Example 4.8.

**1.4. Applications to non-extendable actions.** In Section 5, we use Theorem 1.1 to obstruct an extension of involutions on 3-manifolds to spin 4-manifolds. Regard Brieskorn homology spheres  $\Sigma(p, q, r)$  as a subset of  $\mathbb{C}^3$  followings the standard definition. Consider the involution  $\iota$  on  $\Sigma(2, q, r)$  defined by  $\iota(z_1, z_2, z_3) = (-z_1, z_2, z_3)$ . Recent result by Anvari and Hambleton [3, Theorem A] showed that the standard finite cyclic group actions on  $\Sigma(p, q, r)$  does not extend to any contractible smooth 4-manifold bounded by  $\Sigma(p, q, r)$  (if exists), as a smooth involution. In particular,  $\iota$  does not extend to any contractible smooth 4-manifold bounded by  $\Sigma(2, q, r)$  as a smooth involution. (See also recent work by Baraglia–Hekmati [8, Example 7.7].) As a complementary result, we obstruct to extend such  $\iota$  to non-contractible 4-manifolds for some class of Brieskorn homology spheres:

**Theorem 1.13** (Theorem 5.1). *Set  $Y = \pm\Sigma(2, q, r)$  and let  $\iota : Y \rightarrow Y$  be the involution defined by  $\iota(z_1, z_2, z_3) = (-z_1, z_2, z_3)$ . Let  $W$  be a compact smooth spin 4-manifold bounded by  $Y$  with  $b_1(W) = 0$ . (Note that  $\iota$  can extend to  $W$  as a homologically trivial diffeomorphism. See Theorem 5.1.)*

- (i) *If  $Y = \pm\Sigma(2, 3, 12k+1)$  or  $\pm\Sigma(2, 3, 12k-1)$  for  $k > 0$ , then  $\iota$  cannot extend to  $W$  as a smooth involution so that*

$$-\sigma(W)/16 > b^+(W) - b_l^+(W).$$

- (ii) *If  $Y = \Sigma(2, 3, 12k-5)$  or  $Y = -\Sigma(2, 3, 12k-7)$  for  $k > 0$ , then  $\iota$  cannot extend to  $W$  as a smooth involution so that*

$$-\sigma(W)/16 > b^+(W) - b_l^+(W) - 1/2.$$

- (iii) *If  $Y = -\Sigma(2, 3, 12k-5)$  or  $Y = \Sigma(2, 3, 12k-7)$  for  $k > 0$ , then  $\iota$  cannot extend to  $W$  as a smooth involution so that*

$$-\sigma(W)/16 > b^+(W) - b_l^+(W) + 1/2.$$

- (iv) *In particular, suppose that  $Y$  and  $W$  satisfy one of the following three conditions:*

- $Y = \pm\Sigma(2, 3, 12k + 1)$  or  $\pm\Sigma(2, 3, 12k - 1)$ , and  $\sigma(W) \neq 0$ .
- $Y = \Sigma(2, 3, 12k - 5)$  or  $-\Sigma(2, 3, 12k - 7)$ , and  $\sigma(W) \neq 8$ .
- $Y = -\Sigma(2, 3, 12k - 5)$  or  $\Sigma(2, 3, 12k - 7)$ , and  $\sigma(W) \neq -8$ .

Then  $\iota$  cannot extend to  $W$  as a homologically trivial smooth involution.

**1.5. Floer homotopy type and Floer  $K$ -theory.** The main ingredients to establish Theorem 1.1 are versions of Floer homotopy type and of Floer  $K$ -theory. Let us clarify the nature of our construction of these ingredients. Based upon Manolescu's construction of Seiberg–Witten Floer stable homotopy type [53], we shall construct versions of *Seiberg–Witten Floer stable homotopy type for involutions*

$$DSWF_G(Y, \mathfrak{t}, \iota)$$

and of *Seiberg–Witten Floer  $K$ -theory for involutions*

$$DSWFK_G(Y, \mathfrak{t}, \iota),$$

which are defined for a spin rational homology 3-sphere  $(Y, \mathfrak{t})$  with an involution  $\iota$  whose fixed point set is of codimension 2. Through the construction of the double branched cover, we obtain the *Seiberg–Witten Floer stable homotopy type of knot  $K$*

$$DSWF(K)$$

and the *Seiberg–Witten Floer  $K$ -theory of knot  $K$*

$$DSWFK(K).$$

These are closely related to the recent work by Baraglia and Hekmati [8], where they established an equivariant Seiberg–Witten Floer stable homotopy type for finite groups, and an equivariant Seiberg–Witten Floer cohomology, and defined knot invariants in a similar way. (Note also a recent combinatorial construction of knot Floer homotopy type by Manolescu and Sarkar [59].) However, even restricting our attention to involutions and the case of spin 3-manifolds, there are significant differences between the equivariant Floer homotopy type by Baraglia–Hekmati [8] and our Floer homotopy type for involutions.

The first major difference is that our construction is based on an involutive symmetry on the Seiberg–Witten equations studied by Kato [39]. Here we explain the key point of this. A basic strategy to extract information from group actions is to study the fixed point set. However, given involution  $\iota$  with codimension 2 fixed point set on a spin 3- or 4-manifold, the invariant part of (lifts of)  $\iota$  in the space of spinors turns out to be trivial. Therefore we cannot extract any useful information from the Seiberg–Witten equations restricted to this invariant part. To resolve this situation, Kato [39] defined an involution  $I$  using  $\iota$  on the configuration space for the Seiberg–Witten equations, and observed that the Seiberg–Witten equations are equivariant for this symmetry  $I$ . Notably, Kato's involutive symmetry acts on the spinors as an *anti-complex linear* involution, and therefore this is much different from usual equivariant theory. Recall that the Seiberg–Witten equations on spin manifolds are  $Pin(2)$ -equivariant. A key point is that, while the  $I$ -invariant part of the Seiberg–Witten equations are not  $Pin(2)$ -equivariant anymore, they are still equivariant for  $\mathbb{Z}_4$ , which emerges as the subgroup generated by  $j$  in  $Pin(2)$ . Kato proved a  $10/8$ -type inequality for closed spin 4-manifolds with involutions using this  $\mathbb{Z}_4$ -symmetry.

We consider a 3-dimensional version of this work by Kato, and construct  $G = \mathbb{Z}_4$ -equivariant Seiberg–Witten Floer homotopy type. To show Theorem 1.1, we have

to consider an analogous argument to Manolescu’s relative 10/8-inequality [54] in our setting, but there is still a problem to carry this out. The main problem occurs when we define the  $K$ -theoretic Frøyshov invariant  $\kappa$  for involutions. Once we restrict the Seiberg–Witten equations to the  $I$ -invariant part, the behavior of (an ingredient of)  $\kappa$  is not controllable for suspensions of some representations of  $G$ , not as in Manolescu’s original argument. (See Remark 3.6.) We resolve this problem by considering the “complexification” or “double” of the whole construction of the Floer homotopy type. See Subsection 3.3 for more details on this point. Not only for this problem, the doubling construction allows us to avoid many technical complications in the construction of the Floer homotopy type for involutions.

**1.6. Structure of the paper.** We finish off this introduction with an outline of the contents of this paper. In Section 2, following the Kato’s work [39], we introduce the involution  $I$  on the configuration space, which we mainly use in the construction of our invariant. Using such a symmetry, we apply equivariant Conley index theory to the formal gradient vector field of the Chern–Simons–Dirac functional. In Section 3, for a rational homology sphere with an involution, we construct the (doubled) Seiberg–Witten Floer stable homotopy type, Floer  $K$ -theory,  $K$ -theoretic Frøyshov invariant, and corresponding invariants for knots. The proof of Theorem 1.1 is given in Subsection 3.7. Also, we give several computations of our Seiberg–Witten Floer homotopy type and  $K$ -theoretic Frøyshov invariant in Subsection 3.8. In Section 4, we first review several fundamental calculations related to double branched covering spaces for properly embedded surfaces in punctured 4-manifolds. Using such calculations and Theorem 1.1, we prove Theorem 1.3. We discuss several examples including  $\#_n K3$ ,  $\#_n \mathbb{C}P^2 \#_m (-\mathbb{C}P^2)$  and  $\#_n S^2 \times S^2$ . In Section 5, we obtain results on non-extendable actions of certain Seifert homology 3-spheres using Theorem 1.1. In particular, we prove Theorem 1.13. Section 6 is devoted to explaining other methods to obtain genus bounds such as Manolescu’s relative 10/8-inequality and Tristram–Levine signature, which are related to our inequality (Theorem 1.3).

*Acknowledgement.* The authors would like to express their gratitude to David Baraglia, Motoo Tange, Nobuhiro Nakamura and Oğuz Şavk for helpful comments.

The first author was partially supported by JSPS KAKENHI Grant Numbers 17H06461, 19K23412, and 21K13785. The second author was supported by JSPS KAKENHI Grant Number 21J22979 and WINGS-FMSP program at the Graduate school of Mathematical Science, the University of Tokyo. The third author was supported by JSPS KAKENHI Grant Number 20K22319 and RIKEN iTHEMS Program.

## 2. AN INVOLUTION ON THE CONFIGURATION SPACE

Following Kato’s work [39] in dimensional 4, we shall define an involution on the configuration space for the Seiberg–Witten equations in the 3-dimensional setting. First let us recall a term on involutions [5, 12]. We call a smooth involution  $\iota$  on a smooth spin manifold a *spin involution of odd type* if  $\iota$  lifts to an automorphism of the spin structure as a  $\mathbb{Z}_4$ -action. If the fixed point set is non-empty, this is equivalent to that  $\iota$  lifts to a spin automorphism and the fixed point set is of codimension 2, as far as the spin manifold is of  $\dim \leq 4$  [5, Proposition 8.46].



Throughout this paper, all involutions on spin 3- or 4-manifolds we consider are of odd type.

Let  $(Y, \mathfrak{t})$  be a spin rational homology 3-sphere and  $\iota$  be an involution on  $Y$  preserving the isomorphism class of  $\mathfrak{t}$ . Take an  $\iota$ -invariant Riemannian metric  $g$  on  $Y$ . Let us denote by  $\mathbb{S}$  the spinor bundle of  $Y$ . Suppose that  $\iota$  is of odd type. Then we can take a lift  $\tilde{\iota}$  of  $\iota$  to an automorphism of  $\mathbb{S}$  satisfying

$$\tilde{\iota}^2 = -1.$$

This lift  $\tilde{\iota}$  yields a  $\mathbb{Z}_4$ -action on  $\mathbb{S}$ . There exists exactly one more lift of  $\iota$ , which is given by  $-\tilde{\iota}$ .

On the quaternionic structure of the spinor bundle, we adopt the convention that the quaternion scalars act on the right. Following Kato [39, Subsection 4.2], we define an involution

$$(4) \quad I : T^*Y \oplus \mathbb{S} \rightarrow T^*Y \oplus \mathbb{S}$$

by

$$I(a_x, \phi(x)) = (-\iota^* a_x, \tilde{\iota}(\phi(\iota(x))) \cdot j).$$

Here  $a_x \in T_x^*Y$  and  $\phi(x) \in \mathbb{S}_x$  for  $x \in Y$ . This involution  $I$  is a direct sum of involutions  $I : T^*Y \rightarrow T^*Y$  and  $I : \mathbb{S} \rightarrow \mathbb{S}$ , and the involution  $I$  on the spinors commutes with the right action of  $j \in \text{Pin}(2)$ , but anti-commutes with the right action of  $i$ .

*Remark 2.1.* This involution  $I$  is the composition of the action of  $(\iota^*, \tilde{\iota})$  and the right action of  $(-1, j)$ . The second action was considered by Furuta [19] in his proof of the 10/8-inequality. The action of  $-1$  to the form part is necessary to get the equivariance for the Seiberg–Witten equations. This is because, for any spinor  $\phi$ , we have  $(\phi j (\phi j)^*)_0 = -(\phi \phi^*)_0$ .

The above  $I$  induces an involution on the space of sections of  $TY \oplus \mathbb{S}$ , and restricting this, we obtain an involution

$$I : \text{Ker}(d^* : \Omega^1(Y) \rightarrow \Omega^0(Y)) \times \Gamma(\mathbb{S}) \rightarrow \text{Ker}(d^* : \Omega^1(Y) \rightarrow \Omega^0(Y)) \times \Gamma(\mathbb{S}).$$

Here we consider the Sobolev norms  $L_{k-\frac{1}{2}}^2$  for the spaces  $\text{Ker}(d^* : \Omega^1(Y) \rightarrow \Omega^0(Y))$  and  $\Gamma(\mathbb{S})$  defined using the  $\iota$ -invariant metric and  $\iota$ -invariant connections for a fixed integer  $k \geq 3$ . Henceforth,  $\text{Ker}(d^* : \Omega^1(Y) \rightarrow \Omega^0(Y))$  will be simply denoted by  $\text{Ker } d^*$ .

*Remark 2.2.* There are two choices of lift of  $\iota$ , namely,  $\tilde{\iota}$  and  $-\tilde{\iota}$ . Thus we have two involutions  $I, I'$  from the lift  $\tilde{\iota}$  and  $-\tilde{\iota}$  respectively. We have an isomorphism from the  $I$  fixed part to the  $I'$  fixed part which is given by  $(a, \phi) \mapsto (a, \phi \cdot \sqrt{-1})$ . This isomorphism preserves the formal gradient flow of the Chern–Simons–Dirac functional. Therefore, the Conley index Subsection 3.4, made from the  $I$ -fixed part and the doubled Conley index made from the  $I'$ -fixed part are same.

**Lemma 2.3.** *The formal gradient of CSD*

$$\sigma : \text{Ker } d^* \times \Gamma(\mathbb{S}) \rightarrow \text{Ker } d^* \times \Gamma(\mathbb{S})$$

*is equivariant with respect to the action  $I$ .*

*Proof.* The action  $I$  is the composition of the  $(\iota^*, \tilde{\iota})$  and the right  $(-1, j)$  action. It is obvious that the action of  $(-1, j)$  preserves the formal gradient flow of CSD because this is an element of the ordinary  $\text{Pin}^-(2)$  action to  $\text{Ker } d^* \times \Gamma(\mathbb{S})$ . Let  $P$  is

the principal Spin bundle which gives the spin structure  $\mathfrak{t}$ . The automorphism  $\tilde{\iota}$  on  $P$  is a lift of the involution on the  $SO(TY)$  given by  $\iota$ . The vector bundles  $TY$  and  $\mathbb{S}$  are all associated bundle of  $P$  and the Clifford multiplication  $TY \otimes \mathbb{S} \rightarrow \mathbb{S}$  and the map  $\mathbb{S} \ni \phi \mapsto (\phi\phi^*)_0 \in \sqrt{-1}\Lambda^2 T^*Y$  are given by a multiplication of elements Spin(4) resresentation space. Thus we have that the formal gradient of CSD is equivariant under the action of  $(\iota^*, \tilde{\iota})$ .  $\square$

Define

$$V(Y, \mathfrak{t}, \iota) := (\text{Ker } d^* \times \Gamma(\mathbb{S}))^I.$$

The vector field  $\sigma$  induces a vector field

$$(5) \quad \sigma' : V(Y, \mathfrak{t}, \iota) \rightarrow V(Y, \mathfrak{t}, \iota),$$

The vector field  $\sigma'$  decomposes into the linear part  $l$  and quadratic part  $c$ :  $\sigma' = l + c$ . Here  $l$  is given as  $(*d, D)$ , where  $D$  is the spin Dirac operator. For  $\lambda < 0 \leq \mu$ , define the subspace  $V_\mu^\lambda$  of  $V(Y, \mathfrak{t}, \iota)$  as the direct sum of eigenspaces for  $l$  whose eigenvalues lie in  $(\lambda, \mu]$ . Note that  $V_\mu^\lambda$  is finite-dimensional for any  $\lambda, \mu$ , and we think of  $V_\mu^\lambda$  as a finite-dimensional approximation of  $V(Y, \mathfrak{t}, \iota)$ .

Also, we may define an involution  $I$  on  $Pin(2)$  by

$$I \cdot g := jgj^{-1},$$

and it is easy to see that

$$Pin(2)^I = \{1, j, -1, -j\} \cong \mathbb{Z}_4.$$

Set

$$G := \{1, j, -1, -j\} \subset Pin(2).$$

It follows from a direct calculation that

$$Ig^*(a, \phi) = (I \cdot g)^*(a, \phi).$$

In summary, the flow (5) inherits  $G$ -equivariance from the original  $Pin(2)$ -symmetry of the Seiberg–Witten flow.

For this  $G$ -equivariant flow  $\sigma'$ , we may repeat the construction by Manolescu [53] of the equivariant Conley index associated with the Seiberg–Witten flow. Thus we have a  $G$ -equivariant Conley index

$$I_\lambda^\mu(Y, \mathfrak{t}, \iota, \tilde{\iota}, g)$$

for sufficiently large  $-\lambda, \mu$ .

*Remark 2.4.* While  $I_\lambda^\mu(Y, \mathfrak{t}, \iota, \tilde{\iota}, g)$  depends also on the choice of lift  $\tilde{\iota}$  of  $\iota$ , this does not affect our construction of stable Floer homotopy type in an essential way. See Lemma 3.15.

### 3. SEIBERG–WITTEN FLOER HOMOTOPY THEORY FOR INVOLUTIONS

**3.1. Representations.** Recall that we defined the group  $G$  to be the cyclic group of order 4 generated by  $j \in Pin(2)$ , i.e.

$$G = \{1, j, -1, -j\}.$$

Define a subgroup  $H$  of  $G$  by

$$H = \{1, -1\} \subset G.$$

Let  $\tilde{\mathbb{R}}$  be the 1-dimensional real representation space of  $G$  defined by the surjection  $G \rightarrow \mathbb{Z}_2 = \{1, -1\}$  and the scalar multiplication of  $\mathbb{Z}_2$  on  $\mathbb{R}$ .

Let  $\tilde{\mathbb{C}}$  be a 1-dimensional complex representation of  $G$  defined via the surjection  $G \rightarrow \mathbb{Z}_2$  and the scalar multiplication of  $\mathbb{Z}_2$  on  $\mathbb{C}$ . Note that, if  $s$  is even, say  $2t$ , then there is an isomorphism of real representations  $\tilde{\mathbb{R}}^{2s} \cong \tilde{\mathbb{C}}^t$ .

We introduce also  $G$ -representations

$$(6) \quad \mathbb{R}, \tilde{\mathbb{R}}, \mathbb{C}_+, \mathbb{C}_-,$$

where  $\mathbb{R}, \tilde{\mathbb{R}}$  are real 1-dimensional representations and  $\mathbb{C}_+, \mathbb{C}_-$  are complex 1-dimensional representations, defined by assigning to  $j \in G$  multiplication of  $1, -1, i, -i$  respectively. Note that we have the relation

$$\mathbb{C}_- = \tilde{\mathbb{C}} \cdot \mathbb{C}_+$$

in  $R(G)$ .

We can think of these four spaces (6) as  $H$ -representations through the inclusion  $H \subset G$ , and they correspond to

$$\mathbb{R}, \tilde{\mathbb{R}}, \tilde{\mathbb{C}}, \tilde{\mathbb{C}}$$

as  $H$ -representations, respectively.

It is straightforward to check that the representation ring  $R(G)$  is given by

$$R(G) = \mathbb{Z}[w, z]/(w^2 - 2w, w - 2z + z^2).$$

Here the generators  $w, z$  are given as the  $K$ -theoretic Euler classes of  $\tilde{\mathbb{C}}, \mathbb{C}_+$ , namely,

$$w = 1 - \tilde{\mathbb{C}}, \quad z = 1 - \mathbb{C}_+.$$

The augmentation map  $R(G) \rightarrow \mathbb{Z}$  is given by

$$w, z \mapsto 0,$$

so the augmentation ideal is given by  $(w, z) \subset R(G)$ .

Compared with the standard expression of  $R(G)$ , given by  $R(G) = \mathbb{Z}[t]/(t^4 - 1)$ , an isomorphism  $\mathbb{Z}[w, z]/(w^2 - 2w, w - 2z + z^2) \rightarrow \mathbb{Z}[t]/(t^4 - 1)$  is given by  $w \mapsto 1 - t^2$  and  $z \mapsto 1 - t$ .

It is also straightforward to check that

$$(7) \quad \text{Ker}(R(G) \rightarrow R(H)) = \{ cw \in R(G) \mid c \in \mathbb{Z} \}.$$

In a finite-dimensional approximation of the configuration space on which the (finite-dimensional approximation of)  $G$ -equivariant Seiberg–Witten flow acts, only three of the representations (6) may appear:  $\tilde{\mathbb{R}}, \mathbb{C}_+, \mathbb{C}_-$ . Here  $\mathbb{C}_+, \mathbb{C}_-$  appear in the following way. Originally a finite-dimensional approximation of the configuration space is of the form  $\tilde{\mathbb{R}}^N \oplus \mathbb{H}^{N'}$  as  $Pin(2)$ -representation for large  $N, N'$ . Restricting attention to the  $I$ -invariant part, the remaining symmetry on the configuration space is given by  $G$ , and as  $G$ -representation space,  $\mathbb{H}$  splits into  $\mathbb{C}_+ \oplus \mathbb{C}_-$ . More concretely, let us write the set of quaternions by

$$(8) \quad \mathbb{H} = \mathbb{C} \oplus j\mathbb{C} = \mathbb{R} \oplus i\mathbb{R} \oplus j\mathbb{R} \oplus (-k)\mathbb{R} \cong (\mathbb{R} \oplus j\mathbb{R}) \oplus i(\mathbb{R} \oplus (-j)\mathbb{R}).$$

Let us equip  $\mathbb{R} \oplus j\mathbb{R}$  with the complex structure by  $j$ , and  $\mathbb{R} \oplus (-j)\mathbb{R}$  with the complex structure by  $-j$ . Then the  $(\mathbb{R} \oplus j\mathbb{R})$ -component of (8) corresponds to  $\mathbb{C}_+$ , and the  $(\mathbb{R} \oplus (-j)\mathbb{R})$ -component corresponds to  $\mathbb{C}_-$ .

### 3.2. Space of type $G$ -SWF.

**Definition 3.1.** Let  $X$  be a pointed finite  $G$ -CW complex. We call  $X$  a *space of type  $G$ -SWF* if

- $X^H$  is  $G$ -homotopy equivalent to  $(\mathbb{R}^s)^+$  for some  $s \geq 0$ .
- $G$  acts freely on  $X \setminus X^H$ .

The natural number  $s$  is called the *level* of  $X$ .

Let  $X$  be a space of type  $G$ -SWF at even level  $s = 2t$ . Let  $\iota : X^H \rightarrow X$  denote the inclusion. The image of  $\iota^* : \tilde{K}_G(X) \rightarrow \tilde{K}_G(X^H)$  is expressed of the form  $\mathfrak{J}(X) \cdot b_{t\tilde{C}}$ , where  $\mathfrak{J}(X)$  is an ideal of  $R(G)$  and  $b_{t\tilde{C}}$  is the Bott element.

Repeating the proof of [54, Lemma 1] under replacing  $Pin(2)$  and  $S^1$  with  $G$  and  $H$  respectively, the  $G$ -equivariant localization theorem deduces the following lemma:

**Lemma 3.2.** *Let  $X$  be a space of type  $G$ -SWF at even level  $s = 2t$ . Then there exists  $k \geq 0$  such that  $w^k, z^k \in \mathfrak{J}(X)$ .*

**Definition 3.3.** For a space of type  $G$ -SWF, define

$$k(X) := \min \{ k \geq 0 \mid \exists x \in \mathfrak{J}(X), wx = 2^k w \}.$$

By Lemma 3.2, there exists  $k \geq 0$  such that  $wx = 2^k w$  for some  $x \in \mathfrak{J}(X)$ , and thus  $k(X)$  is a well-defined natural number.

The rest of this subsection is devoted to establishing basic properties of the quantity  $k(X)$ .

**Lemma 3.4.** *Let  $X$  and  $X'$  be spaces of type  $G$ -SWF at the same even level. Suppose that there exists a  $G$ -equivariant map  $f : X \rightarrow X'$  whose  $H$ -fixed point set map is a  $G$ -homotopy equivalence. Then we have*

$$k(X) \leq k(X').$$

*Proof.* We have a commutative diagram

$$\begin{array}{ccc} \tilde{K}_G(X') & \xrightarrow{f^*} & \tilde{K}_G(X) \\ \downarrow & & \downarrow \\ \tilde{K}_G((X')^H) & \xrightarrow{(f^H)^*} & \tilde{K}_G(X^H), \end{array}$$

where  $(f^H)^*$  is an isomorphism. This implies  $\mathfrak{J}(X') \subset \mathfrak{J}(X)$ , and hence  $k(X) \leq k(X')$ .  $\square$

We need to know the behavior of the ideal  $\mathfrak{J}(X)$  under suspension to define the  $K$ -theoretic Frøyshov invariant  $\kappa(Y, \iota)$ :

**Lemma 3.5.** *Let  $X$  be a space of type  $G$ -SWF at an even level. Then we have*

$$(9) \quad \mathfrak{J}(\Sigma^{\tilde{C}} X) = \mathfrak{J}(X), \quad \mathfrak{J}(\Sigma^{C^+} X) = z \cdot \mathfrak{J}(X), \quad \mathfrak{J}(\Sigma^{C^-} X) = (w + z - wz) \cdot \mathfrak{J}(X),$$

and

$$(10) \quad k(\Sigma^{\tilde{C}} X) = k(X), \quad k(\Sigma^{C^+ \oplus C^-} X) = k(X) + 1.$$

*Proof.* The proof is similar to the proof of [54, Lemma 2]. First we shall prove (9). The statement about  $\tilde{\mathbb{C}}$  follows from  $(\Sigma^{\tilde{\mathbb{C}}}X)^H = \Sigma^{\tilde{\mathbb{C}}}(X^H)$  and the naturality of the Bott element.

Next, to prove the statement about  $\mathbb{C}_+$ , consider the diagram induced from the inclusion  $i : X \rightarrow \Sigma^{\mathbb{C}_+}X$  and the inclusions from the  $H$ -invariant part:

$$(11) \quad \begin{array}{ccc} \tilde{K}_G(\Sigma^{\mathbb{C}_+}X) & \xrightarrow{i^*} & \tilde{K}_G(X) \\ \downarrow & & \downarrow \\ \tilde{K}_G((\Sigma^{\mathbb{C}_+}X)^H) & \xrightarrow{(i^H)^*} & \tilde{K}_G(X^H). \end{array}$$

Since  $(\Sigma^{\mathbb{C}_+}X)^H = X^H$ , the bottom map  $(i^H)^*$  is the identity. By the Bott isomorphism, the top map  $i^*$  is given by multiplication by the Euler class  $z$  associated to the representation  $\mathbb{C}_+$ . This implies that

$$\mathfrak{J}(\Sigma^{\mathbb{C}_+}X) = z \cdot \mathfrak{J}(X)$$

in  $R(G)$ .

Similarly, about the statement on  $\mathbb{C}_-$ , we have

$$\begin{array}{ccc} \tilde{K}_G(\Sigma^{\mathbb{C}_-}X) & \xrightarrow{i^*} & \tilde{K}_G(X) \\ \downarrow & & \downarrow \\ \tilde{K}_G((\Sigma^{\mathbb{C}_-}X)^H) & \xrightarrow{=} & \tilde{K}_G(X^H). \end{array}$$

The top map  $i^*$  is given by multiplication by the Euler class  $1 - \mathbb{C}_-$ , and by the relation  $\mathbb{C}_- = \tilde{\mathbb{C}} \cdot \mathbb{C}_-$ , this Euler class is given by

$$1 - \mathbb{C}_- = 1 - (1 - w)(1 - z) = w + z - wz.$$

Next we prove (10). The statement about  $\tilde{\mathbb{C}}$  is clear by (9). To show the statement about  $\mathbb{C}_+ \oplus \mathbb{C}_-$ , first note that we have

$$\mathfrak{J}(\Sigma^{\mathbb{C}_+ \oplus \mathbb{C}_-}X) = z(w + z - wz) \cdot \mathfrak{J}(X)$$

by (9). Using the relations  $w^2 - 2w = 0$ ,  $w - 2z + z^2 = 0$ , we deduce that

$$(12) \quad w(z(w + z - wz)) = zw^2 + wz^2 - w^2z^2 = 2zw - wz^2 = 2zw - 2zw + w^2 = 2w.$$

For  $x \in \mathfrak{J}(X)$  and  $k \geq 0$ , it follows from (12) that  $wx = 2^k w$  holds if and only if  $w(z(w + z - wz))x = 2^{k+1}w$  holds. Thus we have  $k(\Sigma^{\mathbb{C}_+ \oplus \mathbb{C}_-}X) = k(X) + 1$ .  $\square$

*Remark 3.6.* We do not state the behavior of  $k(X)$  under suspension by  $\mathbb{C}_+$  and by  $\mathbb{C}_-$  in Lemma 3.5. In fact,  $k(X)$  does not behave well under these suspensions, not as in Manolescu's original argument. (Compare this with [54, Lemma 2].) For  $\mathbb{C}_+$ , the reason why  $k(X)$  does not behave well is that the relation  $zw - 2w = 0$  does not hold in  $R(G)$ , not as in  $R(\text{Pin}(2))$ . For  $\mathbb{C}_-$ , the reason is that the behavior of  $\mathfrak{J}(X)$  is already complicated as seen in (9).

**Lemma 3.7.** *Let  $X$  and  $X'$  be spaces of type  $G$ -SWF at levels  $2t$  and  $2t'$ , respectively, such that  $t < t'$ . Suppose that there exists a  $G$ -equivariant map  $f : X \rightarrow X'$  whose  $G$ -fixed point set map is a homotopy equivalence. Then we have*

$$k(X) + t \leq k(X') + t'.$$

*Proof.* The proof is similar to the proof of [54, Lemma 5]. Let us start with the diagram

$$(13) \quad \begin{array}{ccc} \tilde{K}_G(X') & \xrightarrow{f^*} & \tilde{K}_G(X) \\ \downarrow & & \downarrow \\ \tilde{K}_G((X')^H) & \xrightarrow{(f^H)^*} & \tilde{K}_G(X^H) \\ \downarrow & & \downarrow \\ \tilde{K}_G((X')^G) & \xrightarrow{(f^G)^*} & \tilde{K}_G(X^G). \end{array}$$

Since  $\tilde{K}_G((X')^H)$ ,  $\tilde{K}_G(X^H)$ ,  $\tilde{K}_G((X')^G)$ ,  $\tilde{K}_G(X^G)$  are free  $R(G)$ -modules of rank 1, we may regard the four maps among them as multiplications by elements of  $R(G)$ . Note that  $(f^G)^*$  is just the identity map, since  $f$  is supposed so that  $f^G$  is a homotopy equivalence. The vertical maps  $\tilde{K}_G((X')^H) \rightarrow \tilde{K}_G((X')^G)$  and  $\tilde{K}_G(X^H) \rightarrow \tilde{K}_G(X^G)$  are given by multiplication with the  $K$ -theoretic Euler classes  $w^{t'}$ ,  $w^t$  respectively. Thus we have that  $(f^H)^*$  is given as multiplication by an element  $y \in R(G)$  satisfying that

$$(14) \quad w^t \cdot y = w^{t'}.$$

We claim that  $(f^H)^*$  in the diagram (13) is given by multiplication by  $2^{t'-t-1}w$ , in other words, we show that  $y = 2^{t'-t-1}w$ . First, since  $t < t'$ , the map

$$(f^H)^* : \tilde{K}_H((X')^H) \rightarrow \tilde{K}_H(X^H)$$

is the zero map. Hence the image of  $y$  under the restriction  $R(G) \rightarrow R(H)$  is zero. From this and (7) we deduce that  $y = cw$  for some  $c \in \mathbb{Z}$ . Then it follows from (14) and the relation  $w^2 = 2w$  that

$$2^t cw = cw^{t+1} = w^{t'} = 2^{t'-1}w,$$

and thus we have  $c = 2^{t'-t-1}$ , and  $y = 2^{t'-t-1}w$ , as claimed.

Take  $x \in \mathfrak{J}(X')$  such that  $wx = 2^{k'}w$ , where  $k' = k(X')$ . By the previous paragraph, we have

$$(f^H)^*(x) = 2^{t'-t-1}wx = 2^{k'+t'-t-1}w.$$

It follows from this that

$$w \cdot (f^H)^*(x) = 2^{k'+t'-t}w.$$

On the other hand, by the commutativity of the diagram (13),  $(f^H)^*(x)$  belongs to  $\mathfrak{J}(X)$ . Thus we obtain  $k' + t' - t \geq k(X)$ .  $\square$

**3.3. Doubling construction and spectrum classes.** Given a spin rational homology 3-sphere  $Y$  with a smooth odd involution  $\iota$ , in Subsection 3.5, we shall define the Seiberg–Witten Floer  $K$ -theory and the  $K$ -theoretic Frøyshov invariant. This is based on a space-valued 3-manifold invariant constructed in Subsection 3.4, which is a  $G$ -equivariant analogue of Manolescu’s stable homotopy type. In this subsection, we prepare a set in which the space-valued invariant takes value.

Fixing an  $\iota$ -invariant metric  $g$  on  $Y$ , we obtain a  $G$ -equivariant Conley index  $I_\lambda^\mu(Y, \iota, g)$  as explained. One option of the definition of the Seiberg–Witten Floer  $K$ -theory is just the  $K$ -theory of (a certain degree shift of)  $I_\lambda^\mu(Y, \iota, g)$ , but it is not

convenient to define the  $K$ -theoretic Frøyshov invariant. The reason is that both of representations  $\mathbb{C}_+$  and  $\mathbb{C}_-$  may appear in the  $I$ -invariant part of the space of spinors, and as we have seen in Lemma 3.5, the ideal  $\mathfrak{J}(X)$  of  $R(G)$  associated to a space of type  $G$ -SWF  $X$  does not behave neatly under the suspension by  $\mathbb{C}_-$ , and  $k(X)$  does not behave well for both  $\mathbb{C}_+$  and  $\mathbb{C}_-$ : see Remark 3.6.

However, under the suspension by the direct sum  $\mathbb{C}_+ \oplus \mathbb{C}_-$ , the quantity  $k(X)$  *does* behave neatly, as seen in Lemma 3.5. Moreover,  $\mathbb{C}_+ \oplus \mathbb{C}_-$  appears as the complexification of  $\mathbb{C}_+$ , and of  $\mathbb{C}_-$  as well. This observation leads us to consider the ‘complexification’ or ‘double’ of  $I_\lambda^\mu(Y, \iota, g)$  and of the relative Bauer–Furuta invariant among them.

To do this, let us define the double of a general space of type  $G$ -SWF. Define a group automorphism  $\alpha : G \rightarrow G$  by  $\alpha(j) = -j$ .

**Definition 3.8.** Let  $X$  be a space of type  $G$ -SWF at level  $t$ . Denote by  $X^\dagger$  the space of type  $G$ -SWF at level  $t$  defined as the same topological space with  $X$ , but the  $G$ -action on  $X$  is given by composing the original  $G$  action on  $X$  with  $\alpha$ . Then  $X \wedge X^\dagger$  is also a space of type  $G$ -SWF, at level  $2t$ . Define the space of type  $G$ -SWF  $D(X)$  by

$$D(X) = X \wedge X^\dagger,$$

which we call the *double* of  $X$ .

Similarly, for a real or complex representation  $V$  of  $G$ , define a representation  $V^\dagger$  by the same vector space with  $V$ , but with  $G$ -action obtained by composing the original  $G$  action on  $V$  with  $\alpha$ . Define a representation  $D(V)$  of  $G$  by

$$D(V) = V \oplus V^\dagger.$$

*Example 3.9.* Since the automorphism  $\alpha : G \rightarrow G$  does not affect the real representation  $\tilde{\mathbb{R}}$ , we have

$$D(\tilde{\mathbb{R}}^t) = \tilde{\mathbb{R}}^{2t} \cong \tilde{\mathbb{C}}^t$$

for  $t \geq 0$ .

On the other hand,  $\alpha$  swaps  $\mathbb{C}_+$  for  $\mathbb{C}_-$ :

$$\begin{aligned} (\mathbb{C}_+)^\dagger &= \mathbb{C}_-, \\ (\mathbb{C}_-)^\dagger &= \mathbb{C}_+. \end{aligned}$$

Thus we have  $G$ -equivariant homeomorphisms

$$D(\mathbb{C}_+) \cong D(\mathbb{C}_-) \cong \mathbb{C}_+ \oplus \mathbb{C}_- \cong (\mathbb{C}_+)_{\mathbb{C}} \cong (\mathbb{C}_-)_{\mathbb{C}},$$

where  $(\mathbb{C}_\pm)_{\mathbb{C}}$  denotes the complexification of  $\mathbb{C}_\pm$ . More generally, we have

$$D(\mathbb{C}_+^m \oplus \mathbb{C}_-^n) \cong (\mathbb{C}_+ \oplus \mathbb{C}_-)^{m+n}$$

for  $m, n \geq 0$ .

Following [54, Section 4], consider a triple  $(D, m, n)$ , where  $D$  is a space of type  $G$ -SWF at an even level, and  $m \in \mathbb{Z}$  and  $n \in \mathbb{Q}$ . We have in mind the case that  $D$  is given as  $D = D(X)$  for some  $X$ , not necessarily at an even level.

**Definition 3.10.** For such triples  $(D, m, n), (D', m', n')$ , we say that they are  $G$ -stably equivalent to each other if  $n - n' \in \mathbb{Z}$  and there exist  $M, N \geq 0$  and a  $G$ -homotopy equivalence

$$\Sigma^{(M-m)\tilde{\mathbb{C}}}\Sigma^{(N-n)(\mathbb{C}_+\oplus\mathbb{C}_-)}D \rightarrow \Sigma^{(M-m')\tilde{\mathbb{C}}}\Sigma^{(N-n')(\mathbb{C}_+\oplus\mathbb{C}_-)}D'.$$

Define  $\mathfrak{C}_G$  as the set of  $G$ -stable equivalence classes of triples  $(D, m, n)$ . An element of  $\mathfrak{C}_G$  is called a *spectrum class*.

Informally, we may think of the triple  $(D, m, n)$  as the formal desuspension of  $X$  by  $m\tilde{\mathbb{C}}$  and by  $n(\mathbb{C}_+ \oplus \mathbb{C}_-)$ , so symbolically one may write

$$(D, m, n) = \Sigma^{-m\tilde{\mathbb{C}}}\Sigma^{-n(\mathbb{C}_+ \oplus \mathbb{C}_-)}D,$$

while we need to keep in mind that  $n$  may not be an integer.

As well as the non-equivariant case, we can define the notion of local equivalence, which was introduced by Stoffregen [71], in our  $G$ -equivariant setting:

**Definition 3.11.** Let  $(D, m, n), (D', m', n')$  be triples as above. A  $G$ -stable map  $(D, m, n) \rightarrow (D', m', n')$  is a based  $G$ -map

$$\Sigma^{(M-m)\tilde{\mathbb{C}}}\Sigma^{(N-n)(\mathbb{C}_+ \oplus \mathbb{C}_-)}D \rightarrow \Sigma^{(M-m')\tilde{\mathbb{C}}}\Sigma^{(N-n')(\mathbb{C}_+ \oplus \mathbb{C}_-)}D'$$

for some  $M, N \geq 0$ . A  $G$ -stable map  $(D, m, n) \rightarrow (D', m', n')$  is called a  $G$ -local map if it induces a  $G$ -homotopy equivalence on the  $H$ -fixed point sets. We say that  $(D, m, n)$  and  $(D', m', n')$  are  $G$ -locally equivalent if there exist  $G$ -local maps  $(D, m, n) \rightarrow (D', m', n')$  and  $(D', m', n') \rightarrow (D, m, n)$ .

The  $G$ -local equivalence is evidently an equivalence relation, and we call an equivalence class for this relation a  $G$ -local equivalence class. The set of  $G$ -local equivalence classes is denoted by  $\mathcal{LE}_G$ . Evidently the  $G$ -stable equivalence implies the  $G$ -local equivalence, and we have a natural surjection  $\mathfrak{C}_G \rightarrow \mathcal{LE}_G$ .

For a triple  $(D, m, n)$  above, we define

$$\tilde{K}_G^*(D, m, n) := \tilde{K}^{*+m+2n}(D)$$

and

$$k(D, m, n) := k(D) - n.$$

The second statement of the following lemma is the main motivation that we work with  $\mathfrak{C}_G$ , allowing only suspension by  $\tilde{\mathbb{C}}$  and  $\mathbb{C}_+ \oplus \mathbb{C}_-$ :

**Lemma 3.12.** Let  $(D, m, n)$  be a triple as above. Then the followings are invariants of the equivalence class  $\mathcal{D} = [(D, m, n)] \in \mathfrak{C}_G$ :

- (1) The isomorphism class of equivariant  $K$ -cohomology,

$$\tilde{K}_G^*(\mathcal{D}) = [\tilde{K}_G^*(D, m, n)],$$

as a graded  $R(G)$ -module.

- (2) The rational number  $k(\mathcal{D}) = k(D, m, n) \in \mathbb{Q}$ .

*Proof.* The first statement immediately follows from the Bott periodicity about suspensions by complex representations. The second statement follows from Lemma 3.5.  $\square$

*Remark 3.13.* Instead of considering elements of  $\mathfrak{C}_G$ , one may define a ‘ $G$ -equivariant suspension spectrum’ by allowing suspension only by  $\mathbb{C}_+$  and only by  $\mathbb{C}_-$ , not necessarily by the pair  $\mathbb{C}_+ \oplus \mathbb{C}_-$ . However, then the statement corresponding to (2) in Lemma 3.12 cannot be obtained. See Remark 3.6.

The statement for (2) in Lemma 3.12 can be improved as follows:

**Lemma 3.14.** The rational number  $k(\mathcal{D})$  is a  $G$ -local equivalence invariant, i.e.  $k(\mathcal{D})$  depends only on the  $G$ -local equivalence class of  $\mathcal{D} \in \mathfrak{C}_G$ .



*Proof.* Given two triples  $(D, m, n), (D', m', n')$ , suppose that they are  $G$ -local equivariant. Let

$$\Sigma^{(M-m)\tilde{\mathbb{C}}\Sigma^{(N-n)(\mathbb{C}_+\oplus\mathbb{C}_-)} D \rightarrow \Sigma^{(M-m')\tilde{\mathbb{C}}\Sigma^{(N-n')(\mathbb{C}_+\oplus\mathbb{C}_-)} D'$$

be a  $G$ -local map from  $(D, m, n)$  to  $(D', m', n')$ , where  $M, N \geq 0$ . Applying Lemma 3.4 to this, we obtain

$$k(\Sigma^{(M-m)\tilde{\mathbb{C}}\Sigma^{(N-n)(\mathbb{C}_+\oplus\mathbb{C}_-)} D) \leq k(\Sigma^{(M-m')\tilde{\mathbb{C}}\Sigma^{(N-n')(\mathbb{C}_+\oplus\mathbb{C}_-)} D').$$

By Lemma 3.5, this is equivalent to that  $k(D, m, n) \leq k(D', m', n')$ . Similarly we obtain  $k(D', m', n') \leq k(D, m, n)$  from a  $G$ -local map  $(D', m', n') \rightarrow (D, m, n)$ , and thus have  $k(D, m, n) = k(D', m', n')$ .  $\square$

### 3.4. Doubled Seiberg–Witten Floer stable homotopy type for involutions.

Now we are ready to construct an invariant of 3-manifolds with involution. Let  $(Y, \mathfrak{t})$  be a spin rational homology 3-sphere and  $\iota$  be a smooth orientation-preserving involution on  $Y$ . Suppose that  $\iota$  also preserves the given spin structure  $\mathfrak{t}$  on  $Y$  and is of odd type. Fix an  $\iota$ -invariant metric  $g$  on  $Y$ , and choose a lift  $\tilde{\iota}$  of  $\iota$  of order 4 on the spin structure. Then we obtain a  $G$ -equivariant Conley index  $I_\lambda^\mu(Y, \iota, \tilde{\iota}, g)$ , once we fix  $\lambda \ll 0 \ll \mu$ . Recall that, the finite-dimensional approximation of the configuration space is decomposed into

$$V(\tilde{\mathbb{R}})_\lambda^\mu \oplus V(\mathbb{C}_+)_\lambda^\mu \oplus V(\mathbb{C}_-)_\lambda^\mu.$$

Here each direct summand is isomorphic to the direct sum of some copies of  $\tilde{\mathbb{R}}, \mathbb{C}_+, \mathbb{C}_-$  respectively. The  $H$ -invariant part of  $I_\lambda^\mu(Y, \iota, g)$  is given by  $V(\tilde{\mathbb{R}})_\lambda^\mu$ .

First, we see that the dependence on the choice of lift  $\tilde{\iota}$ . Recall that  $\iota$  has exactly two lifts, and once we pick a lift  $\tilde{\iota}$ , the other lift is  $-\tilde{\iota}$ .

**Lemma 3.15.** *The double  $D(I_\lambda^\mu(Y, \iota, \tilde{\iota}, g))$  is independent of the choice of  $\tilde{\iota}$ . Namely, there is a canonical  $G$ -equivariant homeomorphism*

$$(15) \quad D(I_\lambda^\mu(Y, \iota, \tilde{\iota}, g)) \cong D(I_\lambda^\mu(Y, \iota, -\tilde{\iota}, g)).$$

*Proof.* To record the dependence, denote the involution  $I$  introduced in (4) by  $I^+$  if it is defined using  $\tilde{\iota}$ , and by  $I^-$  if it is defined using  $-\tilde{\iota}$ . Then it is straightforward to check that

$$I^+(a, \phi i) = I^-(a, \phi) i$$

for all  $(a, \phi) \in \Omega^1(Y) \oplus \Gamma(\mathbb{S})$ .

Recall also how the representations  $\mathbb{C}_+, \mathbb{C}_-$  appear in the configuration space for the Seiberg–Witten Floer theory. Decompose  $\mathbb{H}$  so that

$$\mathbb{H} = \mathbb{C} \oplus j\mathbb{C} = \mathbb{R} \oplus i\mathbb{R} \oplus j\mathbb{R} \oplus (-k)\mathbb{R} \cong (\mathbb{R} \oplus j\mathbb{R}) \oplus i(\mathbb{R} \oplus (-j)\mathbb{R}).$$

Then  $(\mathbb{R} \oplus j\mathbb{R}, j)$  and  $(i(\mathbb{R} \oplus (-j)\mathbb{R}), -j)$  are isomorphic to  $\mathbb{C}_+$  and  $\mathbb{C}_-$  respectively. Note that the right multiplication of  $i$  gives complex linear isomorphisms

$$(\mathbb{R} \oplus j\mathbb{R}, j) \rightarrow (i(\mathbb{R} \oplus (-j)\mathbb{R}), -j), \quad (i(\mathbb{R} \oplus (-j)\mathbb{R}), -j) \rightarrow (\mathbb{R} \oplus j\mathbb{R}, j).$$

Denote a finite-dimensional approximation of the  $I^\pm$ -invariant part of the configuration space by

$$V(\tilde{\mathbb{R}})_\lambda^\mu \oplus V^\pm(\mathbb{C}_+)_\lambda^\mu \oplus V^\pm(\mathbb{C}_-)_\lambda^\mu.$$

(Note that the choice of lift of  $\iota$  does not affect the  $H$ -invariant part  $V(\tilde{\mathbb{R}})_\lambda^\mu$ .) By the above two paragraphs, we see that the right multiplication of  $i$  gives complex linear isomorphisms

$$V^+(\mathbb{C}_+)^\mu \rightarrow V^-(\mathbb{C}_-)^\mu, \quad V^+(\mathbb{C}_-)^\mu \rightarrow V^-(\mathbb{C}_+)^\mu.$$

Therefore the right multiplication of  $i$  induces a  $G$ -equivariant homeomorphism

$$I_\lambda^\mu(Y, \iota, \tilde{\iota}, g) \rightarrow I_\lambda^\mu(Y, \iota, -\tilde{\iota}, g)^\dagger,$$

and this gives rise to the desired  $G$ -equivariant homeomorphism (15).  $\square$

In view of Lemma 3.15, henceforth we drop from our notation the choice of lift of  $\iota$ .

We also need a kind of correction term to absorb the dependence of the invariant metric. Recall a correction term introduced in [53]:

$$n(Y, \mathfrak{t}, g) := \frac{1}{2}(\eta_{dir}(Y, \mathfrak{t}, g) - \dim_{\mathbb{C}} \text{Ker } D_{(Y, \mathfrak{t}, g)}) - \frac{1}{4}\eta_{sign}(Y, g).$$

Here  $\eta_{dir}(Y, \mathfrak{t}, g)$  and  $\eta_{sign}(Y, g)$  are the eta invariants of the Dirac operator and the signature operator respectively, and  $D_{(Y, \mathfrak{t}, g)}$  is the Dirac operator on  $(Y, \mathfrak{t}, g)$ . Alternatively, we can write  $n(Y, \mathfrak{t}, g)$  as

$$n(Y, \mathfrak{t}, g) = \text{ind}_{\mathbb{C}} D_W + \frac{\sigma(W)}{8},$$

where  $W$  is a compact spin Riemann 4-manifold bounded by  $(Y, g)$ .

In our  $G$ -equivariant setting, a direct analogue of Manolescu's Floer stable homotopy type is a triple

$$(16) \quad (\Sigma^{-V(\tilde{\mathbb{R}})_\lambda^0} \Sigma^{-V(\mathbb{C}_+)_\lambda^0} \Sigma^{-V(\mathbb{C}_-)_\lambda^0} I_\lambda^\mu(Y, \mathfrak{t}, \iota, g), 0, n(Y, \mathfrak{t}, g)/4),$$

where the division by 4 for the last factor will be explained in Remark 3.17. However, the triple (16) does not lie in  $\mathfrak{C}_G$ . So, instead, we consider the 'double' of this triple:

$$(17) \quad (\Sigma^{-D(V(\tilde{\mathbb{R}})_\lambda^0)} \Sigma^{-D(V(\mathbb{C}_+)_\lambda^0)} \Sigma^{-D(V(\mathbb{C}_-)_\lambda^0)} D(I_\lambda^\mu(Y, \mathfrak{t}, \iota, g)), 0, n(Y, \mathfrak{t}, g)/2).$$

Rewriting desuspensions to make (17) precise, in view of Example 3.9, we arrive at:

**Definition 3.16.** Given  $Y, \mathfrak{t}, \iota, g$  as above, define an element  $DSWF_G(Y, \mathfrak{t}, \iota) \in \mathfrak{C}_G$  by

$$\begin{aligned} &DSWF_G(Y, \mathfrak{t}, \iota) \\ &:= [(D(I_\lambda^\mu(Y, \mathfrak{t}, \iota, g)), \dim_{\mathbb{R}} V(\tilde{\mathbb{R}})_\lambda^0, \dim_{\mathbb{C}} V(\mathbb{C}_+)_\lambda^0 + \dim_{\mathbb{C}} V(\mathbb{C}_-)_\lambda^0 + n(Y, \mathfrak{t}, g)/2)]. \end{aligned}$$

We call  $DSWF_G(Y, \mathfrak{t}, \iota)$  the *doubled  $G$ -equivariant Seiberg-Witten Floer stable homotopy type* or *doubled Seiberg-Witten Floer  $G$ -spectrum class*.

Recall that, symbolically,  $DSWF_G(Y, \mathfrak{t}, \iota)$  can be thought of as

$$\Sigma^{-(\dim_{\mathbb{R}} V(\tilde{\mathbb{R}})_\lambda^0)} \tilde{\mathbb{C}} \Sigma^{-(\dim_{\mathbb{C}} V(\mathbb{C}_+)_\lambda^0 + \dim_{\mathbb{C}} V(\mathbb{C}_-)_\lambda^0 + n(Y, \mathfrak{t}, g)/2)} (\mathbb{C}_+ \oplus \mathbb{C}_-) D(I_\lambda^\mu(Y, \mathfrak{t}, \iota, g)).$$

*Remark 3.17.* The number  $n(Y, \mathfrak{t}, g)/4$  in (16) was chosen by the following observation. First, since  $I$  and  $i$  anti-commute, the dimension of  $I$ -invariant part of the kernel/cokernel of the Dirac operator is half of the dimension of the original kernel/cokernel. Second, passing to the double, the dimension turns into the double of the original one. Lastly, for suspensions corresponding to the third factor of triples,

we use the direct sum  $\mathbb{C}_+ \oplus \mathbb{C}_-$ , rather than a 1-dimensional complex vector space. So in total

$$\frac{1}{2} \cdot 2 \cdot \frac{1}{2} n(Y, \mathfrak{t}, g) = \frac{1}{2} n(Y, \mathfrak{t}, g)$$

should be put at the third factor of the triple (17). Adopting this number is necessary to prove the invariance of the doubled  $G$ -Floer homotopy type, shown in the following proposition.

We shall show the invariance of  $DSWF_G(Y, \mathfrak{t}, \iota)$ . Before that, we note a technicality about trivializations of representations. Since  $\tilde{\mathbb{C}}$  is a complex representation of  $G$  and  $GL(N, \mathbb{C})$  is connected for all  $N$ , for any triple  $(D, m, n)$  and for any complex  $N$ -dimensional representation  $V$  of  $G$  which is isomorphic to the direct sum of copies of  $\tilde{\mathbb{C}}$ , we have a  $G$ -stable equivalence

$$(V^+ \wedge D, N + m, n) \simeq (D, m, n),$$

and this  $G$ -stable equivalence is canonical up to homotopy. A similar remark applies also to the desuspension by  $\mathbb{C}_+ \oplus \mathbb{C}_-$  since  $\mathbb{C}_+ \oplus \mathbb{C}_-$  is also a complex representation.

**Proposition 3.18.** *The spectrum class  $DSWF_G(Y, \mathfrak{t}, \iota) \in \mathfrak{C}_G$  is an invariant of  $(Y, \mathfrak{t}, \iota)$ , independent of  $\lambda, \mu$ , and  $g$ .*

*Proof.* We basically follow the original argument by Manolescu [54, Proof of Theorem 1]. First we fix  $g$  and show the independence on  $\lambda, \mu$ . Recall the behavior of the Conley index under suspension. That is, for  $\lambda < \lambda' \ll 0 \ll \mu' < \mu$ ,

$$\begin{aligned} I_\lambda^\mu &= I_{\lambda'}^{\mu'}, \\ I_\lambda^\mu &\cong I_{\lambda'}^{\mu'} \wedge (V_{\lambda'}^{\lambda'})^+ = I_{\lambda'}^{\mu'} \wedge (V(\tilde{\mathbb{R}})_{\lambda'}^{\lambda'} \oplus V(\mathbb{C}_+)_{\lambda'}^{\lambda'} \oplus V(\mathbb{C}_-)_{\lambda'}^{\lambda'})^+. \end{aligned}$$

This combined with Example 3.9 implies that

$$\begin{aligned} D(I_\lambda^\mu) &= D(I_{\lambda'}^{\mu'}), \\ D(I_\lambda^\mu) &\cong D(I_{\lambda'}^{\mu'}) \wedge (D(V_{\lambda'}^{\lambda'}))^+ \\ &\cong D(I_{\lambda'}^{\mu'}) \wedge \left( \tilde{\mathbb{C}}^{\dim_{\mathbb{R}} V(\tilde{\mathbb{R}})_{\lambda'}^{\lambda'}} \oplus (\mathbb{C}_+ \oplus \mathbb{C}_-)^{\dim_{\mathbb{C}} V(\mathbb{C}_+)_{\lambda'}^{\lambda'} + \dim_{\mathbb{C}} V(\mathbb{C}_-)_{\lambda'}^{\lambda'}} \right)^+, \end{aligned}$$

and from this the independence on  $\lambda, \mu$  follows.

Next, we show the independence on  $g$ . Fix  $\lambda, \mu$ , and to record the choice of  $g$  used in the construction, we temporarily denote by  $DSWF_G(Y, \mathfrak{t}, \iota, g)$  the doubled  $G$ -spectrum class constructed being used  $\lambda, \mu$ . Take two  $\iota$ -invariant metrics  $g_0$  and  $g_1$ . Since the space of  $G$ -invariant metrics is contractible, we may take a path of  $\iota$ -invariant metrics  $\{g_t\}_{t \in [0,1]}$  between  $g_0$  and  $g_1$ . The assumption that  $b_1(Y) = 0$  tells us that the de Rham operator does not involve here, as in the non-equivariant case, and we focus on the family of Dirac operators  $D_t$  associated with this path of metrics.

Recall that the  $I$ -invariant part of each Dirac operator  $D_t^I$  is  $G$ -equivariant, and we obtain  $G$ -equivariant spectral flow

$$\text{sf}_G(\{D_t^I\}) \in R(G),$$

which can be written as a linear combination only of  $\mathbb{C}_+, \mathbb{C}_-$ . Under the natural map  $R(G) \rightarrow \mathbb{Z}$  induced from taking the dimension of the representation, the image of the equivariant spectral flow  $\text{sf}_G(\{D_t^I\})$  is given by the usual (non-equivariant) spectral flow  $\text{sf}(\{D_t^I\}) \in \mathbb{Z}$ . The difference between the doubled  $G$ -spectrum classes

$DSWF_G(Y, \mathfrak{t}, \iota, g_0)$  and  $DSWF_G(Y, \mathfrak{t}, \iota, g_1)$  is given by the ‘doubled’ equivariant spectral flow

$$D(\text{sf}_G(\{D_t^I\})) \in R(G),$$

which is given by

$$(18) \quad (\mathbb{C}_+ \oplus \mathbb{C}_-)^{\text{sf}(\{D_t^I\})}.$$

Recall that we have a formula

$$(19) \quad \text{sf}(\{D_t\}) = n(Y, \mathfrak{t}, g_1) - n(Y, \mathfrak{t}, g_0),$$

and thus have

$$\text{sf}(\{D_t^I\}) = n(Y, \mathfrak{t}, g_1)/2 - n(Y, \mathfrak{t}, g_0)/2.$$

Note that  $\text{sf}(\{D_t\})$  is an even integer because of the existence of the action  $I$ . Thus we finally get

$$D(\text{sf}_G(\{D_t^I\})) = (\mathbb{C}_+ \oplus \mathbb{C}_-)^{n(Y, \mathfrak{t}, g_1)/2 - n(Y, \mathfrak{t}, g_0)/2} \in R(G),$$

and this completes the proof.  $\square$

*Remark 3.19.* As noted in [8], it is subtle to split the equivariant spectral flow  $\text{sf}_G(\{D_t\})$  as in the non-equivariant case (19). But here we consider the doubled homotopy type, and the doubling construction makes the subtlety from the equivariance disappear: the difference between two doubled homotopy types corresponding to two choices of metrics is determined only by the non-equivariant spectral flow, as seen in (18).

**3.5. Seiberg–Witten Floer  $K$ -theory for involutions.** As in the last subsection, let  $(Y, \mathfrak{t})$  be a spin rational homology 3-sphere and  $\iota$  be a smooth orientation-preserving involution. Suppose that  $\iota$  also preserves the given spin structure  $\mathfrak{t}$  and is of odd type.

**Definition 3.20.** Define the *doubled Seiberg–Witten Floer  $K$ -cohomology* by

$$DSWFK_G(Y, \mathfrak{t}, \iota) := \tilde{K}_G(DSWF_G(Y, \mathfrak{t}, \iota)),$$

defined as the isomorphism class of an  $R(G)$ -graded module. We define also the  *$K$ -theoretic Frøyshov invariant* by

$$\kappa(Y, \mathfrak{t}, \iota) := k(DSWF_G(Y, \mathfrak{t}, \iota)) \in \mathbb{Q}.$$

**Lemma 3.21.** *The isomorphism class  $DSWFK_G(Y, \mathfrak{t}, \iota)$  and the rational number  $\kappa(Y, \mathfrak{t}, \iota)$  are invariant of  $(Y, \mathfrak{t}, \iota)$ .*

*Proof.* This is a direct consequence of Lemma 3.12 and Proposition 3.18.  $\square$

If we allow us to use additional non-topological data  $g, \lambda, \mu$ , the invariant  $\kappa(Y, \mathfrak{t}, \iota)$  is concretely described as

$$(20) \quad \kappa(Y, \mathfrak{t}, \iota) = k(D(I_\lambda^\mu(Y, \mathfrak{t}, \iota, g))) - \dim_{\mathbb{C}}(V(\mathbb{C}_+)_\lambda^0) - \dim_{\mathbb{C}}(V(\mathbb{C}_-)_\lambda^0) - n(Y, \mathfrak{t}, g)/2.$$

*Remark 3.22.* In this paper, we apply  $K$ -theory to the doubled or ‘complexification’ of the  $I$ -fixed point part of the Seiberg–Witten Floer stable homotopy type to define the invariant  $\kappa$ . In our setting, it could be able to apply  $KR$ -theory, which was introduced by Atiyah in [4], to the whole space of Seiberg–Witten Floer stable homotopy type with involution  $I$ , and perhaps such formulation could be natural.

**3.6. Cobordisms.** Let  $(Y_0, \mathfrak{t}_0, \iota_0)$  and  $(Y_1, \mathfrak{t}_1, \iota_1)$  be spin closed 3-manifold with  $b_1(Y) = 0$ . We do not assume that  $Y_0$  and  $Y_1$  are connected. Suppose that we have an involution  $\iota_i$  on each of  $Y_i$ . Let  $(W, \mathfrak{s})$  be a smooth spin 4-dimensional oriented cobordism with  $b_1(W) = 0$ . We assume that there is a involution  $\iota$  on  $W$  such that  $\iota|_{Y_i} = \iota_i$  for  $i = 0, 1$ , and suppose that  $\iota$  preserves  $\mathfrak{s}$  and of odd type. We may take an  $\iota$ -invariant Riemannian metric  $g$  on  $W$  so that  $g$  is a cylindrical metric near  $\partial W$ . Then the metrics defined by  $g_i = g|_{Y_i}$  on  $Y_i$  are  $\iota_i$ -invariant metrics. Then  $\iota$  lifts to some  $\tilde{\iota}$ , which is a  $\mathbb{Z}_4$ -lift of  $\iota$  to the spinor bundle of  $\mathfrak{s}$ . As well as the case of dimension 3, following [39], we may define the involutions

$$\begin{aligned} I : \Omega^*(W) &\rightarrow \Omega^*(W), \\ I : \Gamma(\mathbb{S}^\pm) &\rightarrow \Gamma(\mathbb{S}^\pm) \end{aligned}$$

by

$$\begin{aligned} I(a_x) &= -(\iota^* a)_x, \\ I(\phi(x)) &= \tilde{\iota}(\phi(\iota(x))) \cdot j, \end{aligned}$$

where  $\mathbb{S}^\pm$  are positive and negative spinor bundles. Here we consider the Sobolev norms  $L_k^2$  for the spaces  $\Omega^*(W)$  and  $\Gamma(\mathbb{S}^\pm)$  obtained from  $\iota$ -invariant metrics and  $\iota$ -invariant connections for a fixed integer  $k \geq 3$ . The relative Bauer–Furuta invariant of  $W$  introduced by Manolescu [53] gives a map between the Seiberg–Witten Floer stable homotopy types of  $Y_0$  and  $Y_1$ . This is obtained from the Seiberg–Witten map on  $W$ , which is given as a finite-dimensional approximation of a map

$$(21) \quad SW : \Omega_{CC}^1(W) \times \Gamma(\mathbb{S}^+) \rightarrow \Omega^+(W) \times \Gamma(\mathbb{S}^-) \times \hat{V}_{-\infty}^\mu(-Y_0) \times \hat{V}_{-\infty}^\mu(Y_1),$$

for large  $\mu$ . Here

$$\Omega_{CC}^1(W) = \left\{ a \in \Omega^1(W) \mid d^* a = 0, d^* \mathfrak{t}_i a = 0, \int_{Y_i} \mathfrak{t}_i * a = 0. \right\}$$

is the space of all 1-forms satisfying the double coulomb condition. This is introduced by T. Khandhawit in [40]. Here for a general rational spin homology sphere  $Y$ ,  $\hat{V}(Y, \mathfrak{t})_{-\infty}^\mu$  is a subspace of

$$\hat{V}(Y, \mathfrak{t}) := \text{Ker } d^* \times \Gamma(\mathbb{S}),$$

which is defined as the direct sum of eigenspaces whose eigenvalues are less than  $\mu$ . The  $\Omega^+(W) \times \Gamma(\mathbb{S}^-)$ -factor of the map  $SW$  is given as the Seiberg–Witten equations, and the  $V_{-\infty}^\mu(-Y_0) \times V_{-\infty}^\mu(Y_1)$ -factor is given, roughly, as the restriction of 4-dimensional configurations to 3-dimensional ones. Taking the  $I$ -invariant part of (21), we obtain a  $G$ -equivariant map, and a finite-dimensional approximation of this gives us a  $G$ -equivariant map of the form

$$(22) \quad f : \Sigma^{m_0 \tilde{\mathbb{R}}} \Sigma^{n_0^+ \mathbb{C}_+} \Sigma^{n_0^- \mathbb{C}_+} I_{-\mu}^{-\lambda}(Y_0) \rightarrow \Sigma^{m_1 \tilde{\mathbb{R}}} \Sigma^{n_1^+ \mathbb{C}_+} \Sigma^{n_1^- \mathbb{C}_+} I_\lambda^\mu(Y_1),$$

where  $I_\lambda^\mu(Y_i) = I_\lambda^\mu(Y_i, \mathfrak{t}_i, \iota_i, g_i)$ , and  $m_i, n_i^\pm \geq 0$  and  $-\lambda, \mu$  are sufficiently large. Taking the double of  $f$ , we obtain the ‘doubled cobordism map’

$$(23) \quad D(f) : \Sigma^{m_0 \tilde{\mathbb{C}}} \Sigma^{n_0(\mathbb{C}_+ \oplus \mathbb{C}_-)} D(I_{-\mu}^{-\lambda}(Y_0)) \rightarrow \Sigma^{m_1 \tilde{\mathbb{C}}} \Sigma^{n_1(\mathbb{C}_+ \oplus \mathbb{C}_-)} D(I_\lambda^\mu(Y_1)),$$

where  $n_i = n_i^+ + n_i^-$ . Denote by  $V_0(\tilde{\mathbb{R}})_\lambda^\mu$  is the vector space  $V(\tilde{\mathbb{R}})_\lambda^\mu$  for  $Y_0$ , and let us use similar notations also for other representaions and for  $Y_1$ .

**Lemma 3.23.** *We have*

$$(24) \quad m_0 - m_1 = \dim_{\mathbb{R}}(V_1(\tilde{\mathbb{R}})_{\lambda}^0) - \dim_{\mathbb{R}}(V_0(\tilde{\mathbb{R}})_{-\mu}^0) - b^+(W) + b_t^+(W),$$

$$(25) \quad n_0 - n_1 = \dim_{\mathbb{C}}(V_1(\mathbb{C}_+)_{\lambda}^0) + \dim_{\mathbb{C}}(V_1(\mathbb{C}_-)_{\lambda}^0) \\ - \dim_{\mathbb{C}}(V_0(\mathbb{C}_+)_{-\mu}^0) - \dim_{\mathbb{C}}(V_0(\mathbb{C}_-)_{-\mu}^0) \\ - \frac{\sigma(W)}{16} + \frac{n(Y_1, \mathfrak{t}_1, g_1)}{2} - \frac{n(Y_0, \mathfrak{t}_0, g_0)}{2}.$$

*Proof.* It is sufficient to show the equality when  $\mu$  and  $\lambda$  are not eigenvalue. The proof of this lemma is parallel to the proof of [40, Proposition 2]. We will denote by  $U_i$  the space  $\text{Im}(d)_{Y_i}^I$ . Let  $r_i$  be the restriction to  $Y_i$  and  $\Pi^-$  be the projection from  $\Omega_1(Y_i)^I \times \Gamma(\mathbb{S})^I$  to  $\hat{V}_{-\infty}^0(Y_i)^I$  and let  $\Pi_2$  be the projection from  $\Omega_1(Y_i)^I \times \Gamma(\mathbb{S})^I$  to  $U_i$ . Let  $\mathfrak{t}a$  be the tangent component of the restriction of a 1-form  $a$  to the boundary. When  $\mu = 0$ , the index of the linearization of the  $I$ -invariant part of the map (21) coincides with the index of the map

$$(26) \quad F: \Omega^1(W)^I \times \Gamma(\mathbb{S}^+)^I \rightarrow \Omega^+(W)^I \times \Gamma(\mathbb{S}^-)^I \times \Omega^0(W)^I \times \hat{V}_{-\infty}^0(-Y_0)^I \times \hat{V}_{-\infty}^0(Y_1)^I \times U_0 \times U_1$$

given by

$$F(a, \phi) = (d^+a, D\phi, d^*a, \Pi^- \circ r_0(a, \phi), \Pi^- \circ r_1(a, \phi), \Pi_2 \circ r_0(a, \phi), \Pi_2 \circ r_1(a, \phi))$$

because the kernel of the map  $d^* \oplus \Pi_2 \circ r_0 \oplus \Pi_2 \circ r_1$  coincides with  $\Omega_{CC}^1(W)^I \times \Gamma(\mathbb{S})^I$  and the cokernel of  $d^* \oplus \Pi_2 \circ r_0 \oplus \Pi_2 \circ r_1$  is 0 because constant functions on  $W$  are not fixed by  $-\iota^*$ . Note that  $\int_{Y_i} * \mathfrak{t}a = 0$  for all  $a \in \Omega^1(W)^I$ . Let  $L_1$  be the operator acting on  $\text{Im}(d)_{Y_i}^I \times \Omega_0(Y_i)^I$  by

$$L_1 = \begin{pmatrix} 0 & -d \\ -d^* & 0 \end{pmatrix}.$$

Let  $\Pi_1^-$  be the projection to the non-negative eigenspace of the operator  $L_1$ . The kernel of the projection  $\Pi_2|_{\Omega_1(Y_i)^I}$  is  $\text{Ker}(d^*)_{Y_i}^I$  and the image of the  $\Pi_1^-$  is

$$\{(b, d^*(dd^*)^{-1/2}b) \mid b \in \Omega_1(Y_i)^I\}.$$

The kernel of  $(\Pi_2|_{\Omega_1(Y_i)^I}) \circ r_i$  is complementary to the non-positive eigenspace of  $L_1$  in the space  $\Omega_1(Y_i)^I \times \Omega_0(Y_i)^I$ . By the [48, Proposition 17.2.6], we have the operator  $F$  is Fredholm and the index coincides with the index of the following operator:

$$(27) \quad F': \Omega^1(W)^I \times \Gamma(\mathbb{S}^+)^I \rightarrow \Omega^+(W)^I \times \Gamma(\mathbb{S}^-)^I \times \Omega^0(W)^I \times \hat{U}_{-\infty}^0(-Y_0) \times \hat{U}_{-\infty}^0(Y_1)$$

Here  $\hat{U}_{-\infty}^0(-Y_0)$  and  $\hat{U}_{-\infty}^0(Y_1)$  are the sum of the non-positive eigenspaces in  $\Omega_1(Y_0)^I \times \Omega_0(Y_0)^I \times \Gamma(Y_0, \mathbb{S})^I$  and  $\Omega_1(Y_1)^I \times \Omega_0(Y_1)^I \times \Gamma(Y_1, \mathbb{S})^I$  of  $(L_1, D)^I$  respectively. We have the index of the spinor part of  $F'$  is the half of the Atiyah-Patodi-Singer index of the  $\text{Spin}^c$  Dirac operator because  $I$  is anti-commutes with  $i$ . We see that the index of the form part of  $F'$  is given by  $-b^+(W) + b_t^+(W)$  as follows: Elements in the kernel and the cokernel of  $F'$  are able to be extended to a  $L^2$  harmonic form of  $\hat{W} = (-\infty, 0] \times Y_0 \cup_{Y_0} W \cup_{Y_1} Y_1 \times [0, \infty)$  because  $Y_i$  are rational homology spheres. We see that the  $L^2$  harmonic form on  $\hat{W}$  coincides with the de Rham cohomology of  $\hat{W}$  from in [6, Proposition 4.9] and the kernel and cokernel

of  $F'$  coincides with the  $-\iota^*$ -invariant part of the space of  $L^2$  harmonic forms. Thus we have the index of the form part of  $F'$ .

In the case  $\mu > 0$ , we have the index of the  $\text{Spin}^c$  Dirac part of  $F'$  is  $\text{ind}_{\mathbb{C}}(D^+) - \dim_{\mathbb{C}}(\hat{V}_0^\mu(-Y_0)^I \cap \Gamma(\mathbb{S})^I) - \dim_{\mathbb{C}}(\hat{V}_0^\mu(Y_1)^I \cap \Gamma(\mathbb{S})^I)$  and the index of the form part of  $F'$  is  $-b^+(W) + b_i^+(W) - \dim(\hat{V}_0^\mu(-Y_0)^I \cap \text{Ker}(d^*)^I) - \dim(\hat{V}_0^\mu(Y_1)^I \cap \text{Ker}(d^*)^I)$ . Recall that the map (22) is given the same way in [55, Section 3.6] and that  $\hat{V}_\lambda^\mu(-Y_0) = \hat{V}_{-\mu}^{-\lambda}(Y_0)$ . From the construction of the map (39), we easily see that

$$\begin{aligned} & (\text{the index of the form part of } F') + \dim(V_0(\tilde{\mathbb{R}})_{-\mu}^{-\lambda}) - \dim(V_1(\tilde{\mathbb{R}})_0^\mu) - \dim(V_0(\tilde{\mathbb{R}})_{-\mu}^0) \\ &= (\text{the index of the form part of } F') + \dim(V_0(\tilde{\mathbb{R}})_0^{-\lambda}) - \dim(V_1(\tilde{\mathbb{R}})_0^\mu) \\ &= m_0 + \dim(V_0(\tilde{\mathbb{R}})_{-\mu}^{-\lambda}) - m_1 - \dim(V_1(\tilde{\mathbb{R}})_\lambda^\mu), \\ & (\text{the index of the } \text{Spin}^c \text{ Dirac part of } F') + \dim_{\mathbb{C}}(V_0(\mathbb{C}_+)_{-\mu}^{-\lambda}) + \dim_{\mathbb{C}}(V_0(\mathbb{C}_-)_{-\mu}^{-\lambda}) \\ & \quad - \dim_{\mathbb{C}}(V_0(\mathbb{C}_+)_{-\mu}^0) - \dim_{\mathbb{C}}(V_1(\mathbb{C}_+)_0^\mu) \\ & \quad - \dim_{\mathbb{C}}(V_0(\mathbb{C}_-)_{-\mu}^0) - \dim_{\mathbb{C}}(V_1(\mathbb{C}_-)_0^\mu) \\ &= (\text{the index of the } \text{Spin}^c \text{ Dirac part of } F') + \dim_{\mathbb{C}}(V_0(\mathbb{C}_+)_{-\mu}^{-\lambda}) - \dim_{\mathbb{C}}(V_1(\mathbb{C}_+)_{-\mu}^\mu) \\ & \quad + \dim_{\mathbb{C}}(V_0(\mathbb{C}_-)_{-\mu}^{-\lambda}) - \dim_{\mathbb{C}}(V_1(\mathbb{C}_-)_{-\mu}^\mu) \\ &= n_0 + \dim_{\mathbb{C}}(V_0(\mathbb{C}_+)_{-\mu}^{-\lambda}) + \dim_{\mathbb{C}}(V_0(\mathbb{C}_-)_{-\mu}^{-\lambda}) - n_1 - \dim_{\mathbb{C}}(V_1(\mathbb{C}_+)_{-\mu}^\mu) - \dim_{\mathbb{C}}(V_1(\mathbb{C}_-)_{-\mu}^\mu) \end{aligned}$$

respectively. Hence we have the formula (24) and (25).  $\square$

*Remark 3.24.* To calculate the differences  $n_0^+ - n_1^+, n_0^- - n_1^-$ , we need to use an additional equivariant index theorem. But for our purpose, it is enough to know only the difference  $n_0 - n_1$ .

**3.7. Proof of Theorem 1.1.** First, we state and prove the main result of this paper in the most general form:

**Theorem 3.25.** *Let  $(Y_0, \mathfrak{t}_0), (Y_1, \mathfrak{t}_1)$  be spin rational homology 3-spheres. Let  $\iota_0, \iota_1$  be smooth involutions on  $Y_0, Y_1$ . Suppose that  $\iota_0, \iota_1$  preserve the given orientations and spin structures  $\mathfrak{t}_0, \mathfrak{t}_1$  on  $Y_0, Y_1$  respectively, and suppose that  $\iota_0, \iota_1$  are of odd type. Let  $(W, \mathfrak{s})$  be a smooth compact oriented spin cobordism with  $b_1(W) = 0$  from  $(Y_0, \mathfrak{t}_0)$  to  $(Y_1, \mathfrak{t}_1)$ . Suppose that there exists a smooth involution  $\iota$  on  $W$  such that  $\iota$  preserves the given orientation and spin structure  $\mathfrak{s}$  on  $W$ , and that the restriction of  $\iota$  to the boundary is given by  $\iota_0, \iota_1$ . Then we have*

$$(28) \quad -\frac{\sigma(W)}{16} + \kappa(Y_0, \mathfrak{t}_0, \iota_0) \leq b^+(W) - b_i^+(W) + \kappa(Y_1, \mathfrak{t}_1, \iota_1),$$

where  $b_i^+(W)$  denotes the maximal dimension of  $\iota$ -invariant positive-definite subspaces of  $H^2(W; \mathbb{R})$ .

*Proof.* As in Subsection 3.6, take an  $\iota$ -invariant metric  $g$  on  $W$  so that  $g$  is a cylindrical metric near  $\partial W$ . The involution  $\iota$  lifts to the spin structure  $\mathfrak{s}$  as  $\mathbb{Z}_4$ -action, and we have the doubled cobordism map (39). In this proof, we take  $\mu = \nu$  and  $\lambda = -\nu$ . Recall that the  $H$ -invariant part of  $I_{-\nu}^\nu(Y_i)$  is of (real) dimension  $\dim_{\mathbb{R}}(V_i(\tilde{\mathbb{R}})_{-\nu}^\nu)$ , so the  $H$ -invariant part of the double  $D(I_{-\nu}^\nu(Y_i))$  is of dimension  $2 \dim_{\mathbb{R}}(V_i(\tilde{\mathbb{R}})_{-\nu}^\nu)$  and there is a canonical homotopy which collapses  $(V_i(\tilde{\mathbb{R}})_{-\nu}^\nu)^2$  part in  $D(I_{-\nu}^\nu(Y_i))$  to the base point. Hence the level of  $\Sigma^{m_i} \tilde{\mathbb{C}} \Sigma^{n_i} (\mathbb{C}_+ \oplus \mathbb{C}_-) D(I_{-\nu}^\nu(Y_i))$  is given by  $2(m_i + \dim_{\mathbb{R}}(V_i(\tilde{\mathbb{R}})_{-\nu}^0))$ .

First, consider the case that  $b^+(W) - b_l^+(W) > 0$ . Then, by (24), the level of the domain of the doubled cobordism map (39) is smaller than that of the codomain of (39). Therefore we can apply Lemma 3.7 to (39), and obtain that

$$\begin{aligned} & k(\Sigma^{m_0} \tilde{\mathbb{C}} \Sigma^{n_0}(\mathbb{C}_+ \oplus \mathbb{C}_-) D(I_{-\nu}^\nu(Y_0))) + m_0 + \dim_{\mathbb{R}}(V_0(\tilde{\mathbb{R}})_{-\nu}^0) \\ & \leq k(\Sigma^{m_1} \tilde{\mathbb{C}} \Sigma^{n_1}(\mathbb{C}_+ \oplus \mathbb{C}_-) D(I_{-\nu}^\nu(Y_1))) + m_1 + \dim_{\mathbb{R}}(V_1(\tilde{\mathbb{R}})_{-\nu}^0). \end{aligned}$$

By Lemma 3.5, this is equivalent to

$$\begin{aligned} & k(D(I_{-\nu}^\nu(Y_0))) + n_0 + m_0 + \dim_{\mathbb{R}}(V_0(\tilde{\mathbb{R}})_{-\nu}^0) \\ & \leq k(D(I_{-\nu}^\nu(Y_1))) + n_1 + m_1 + \dim_{\mathbb{R}}(V_1(\tilde{\mathbb{R}})_{-\nu}^0). \end{aligned}$$

From this combined with (20), (24), (25), we obtain the desired inequality (1).

Next, consider the case that  $b^+(W) - b_l^+(W) = 0$ . Then, by (24), the level of the domain of the doubled cobordism map (39) is the same as that of the codomain of (39). Then we can apply Lemma 3.4 to (39), instead of Lemma 3.7, and obtain the desired inequality (1) by the same argument for the case that  $b^+(W) - b_l^+(W) > 0$  above.  $\square$

*Proof of Theorem 1.1.* In general, for a spin manifold of  $\dim \leq 4$  and an involution  $\iota$  on this manifold preserving the spin structure with non-empty fixed point set, the condition that  $\iota$  is of odd type is equivalent to that the fixed point set of  $\iota$  is of codimension 2 [5, Proposition 8.46]. Thus Theorem 1.1 immediately follows from Theorem 3.25.  $\square$

*Remark 3.26.* In this paper, we only consider  $\mathbb{Z}_2$ -branched covering spaces. On the other hand,  $\mathbb{Z}_p$ -branched covering spaces have been used in gauge theory (see for examples [3, 8, 35]). However, it seems not to be straightforward to extend the  $\mathbb{Z}/p$  action to the settings in this paper.

Note that for a statement similar to Theorem 3.25 for a spin 4-manifold  $W$  with one boundary component follows from Theorem 3.25 by removing a ball in  $W$  near a fixed point, as far as the involution has non-empty fixed point set. But, for free involutions, it seems that the case of one boundary component is not deduced from the two component case Theorem 3.25. However, we can carry out the proof of Theorem 3.25 also for the one boundary component case without any essential change. We record this as a statement as follows. We wish to thank David Baraglia for pointing this out. See also Nakamura's work [63] in his  $Pin^-(2)$ -monopole setting for free involutions.

**Theorem 3.27.** *Let  $(Y, \mathfrak{t})$  be a spin rational homology 3-sphere. Let  $\iota$  be a smooth involution on  $Y$ . Suppose that  $\iota_Y$  preserves the given orientation and spin structure  $\mathfrak{t}$  respectively, and suppose that  $\iota_Y$  is of odd type. Let  $(W, \mathfrak{s})$  be a smooth compact oriented spin 4-manifold  $b_1(W) = 0$  bounded by  $(Y, \mathfrak{t})$ . Suppose that there exists a smooth involution  $\iota$  on  $W$  such that  $\iota$  preserves the given orientation and spin structure  $\mathfrak{s}$  on  $W$ , and that the restriction of  $\iota$  to the boundary is given by  $\iota_Y$ . Then we have*

$$(29) \quad -\frac{\sigma(W)}{16} \leq b^+(W) - b_l^+(W) + \kappa(Y, \mathfrak{t}, \iota_Y),$$

where  $b_l^+(W)$  denotes the maximal dimension of  $\iota$ -invariant positive-definite subspaces of  $H^2(W; \mathbb{R})$ .



**3.8. Calculations.** We carry out calculations of the doubled Seiberg-Witten Floer stable homotopy type  $DSWF_G(Y, \mathfrak{t}, \iota)$  and equivariant  $K$ -theoretic Frøyshov invariant  $\kappa(Y, \mathfrak{t}, \iota)$  for some  $(Y, \mathfrak{t}, \iota)$ .

**Proposition 3.28.** *Let  $(Y, \mathfrak{t})$  be a spin rational homology 3-sphere equipped with an involution  $\iota$  preserving  $\mathfrak{t}$ . Assume that  $Y$  admits an  $\iota$ -invariant positive scalar curvature metric  $g$ . Construct  $DSWF_G(Y, \mathfrak{t}, \iota)$  using  $g$ , as well as non-equivariant case [53, Subsection 5.1]. Then we have that*

$$DSWF_G(Y, \mathfrak{t}, \iota) = [(S^0, 0, n(Y, \mathfrak{t}, g)/2)]$$

and

$$\kappa(Y, \mathfrak{t}, \iota) = -n(Y, \mathfrak{t}, g)/2.$$

Moreover, we have

$$\kappa(-Y, \mathfrak{t}, \iota) = -\kappa(Y, \mathfrak{t}, \iota).$$

*Proof.* In general, we have

$$(30) \quad n(Y, \mathfrak{t}, g) + n(-Y, \mathfrak{t}, g) = \dim_{\mathbb{C}} \text{Ker } D_{(Y, \mathfrak{t}, g)},$$

where  $D_{(Y, \mathfrak{t}, g)}$  is the 3-dimensional Dirac operator on  $(Y, \mathfrak{t}, g)$  (see [55, Page 167]), and now we have  $\text{Ker } D_{(Y, \mathfrak{t}, g)} = 0$  since  $g$  is a positive scalar curvature metric.  $\square$

*Remark 3.29.* In the non-equivariant case [53, Subsection 5.1], the  $K$ -theoretic Frøyshov invariant  $\kappa(Y, \mathfrak{t})$  is given by

$$\kappa(Y, \mathfrak{t}) = -n(Y, \mathfrak{t}, g)$$

whenever  $Y$  admits a positive scalar metric  $g$ . Proposition 3.28 says that the equivariant  $\kappa$  is the half of the non-equivariant one provided the existence of  $\iota$ -invariant positive scalar curvature metric. Intuitively, this difference arises from the fact that the  $I$ -invariant part of the space of spinors is the ‘half’ of the whole space of spinors.

A large class of examples are obtained from Seifert homology spheres. We prepare some generality before calculating our invariants for Seifert homology spheres.

**Lemma 3.30.** *Let  $(Y, \mathfrak{t})$  be a spin rational homology sphere. Let  $\iota$  be a smooth involution preserving  $\mathfrak{t}$ . Let  $g$  be an  $\iota$ -invariant metric on  $Y$ . Suppose that  $\iota$  is isotopic to the identity of  $Y$  through the isometry group of  $(Y, g)$ . Then a path from  $\iota$  to  $\text{id}_Y$  in the isometry group of  $(Y, g)$  lifts to a path from  $\tilde{\iota}$  to the identity  $\text{id}$  or  $-\text{id}$  on the spin structure in the automorphism group of the spin structure.*

*Proof.* Let  $P_{SO}$  be the oriented orthogonal frame bundle of  $Y$  and  $P_{Spin}$  is the principal  $Spin(3)$  bundle of the spin structure  $\mathfrak{t}$ . We denote by the induced action on  $P_{SO}$  from  $\iota$  by the same notation. Note that  $P_{Spin} \times [0, 1]$  is a fiberwise double covering of  $P_{SO} \times [0, 1]$ , and the path from  $\iota$  to the identity of  $P_{SO}$  induces an automorphism  $f$  of the fiber bundle  $P_{SO} \times [0, 1]$  and the automorphism of the fundamental group of  $P_{SO} \times [0, 1]$  induced by  $f$  is the identity. Thus  $f$  lifts to an automorphism  $\tilde{f}$  on  $P_{Spin} \times [0, 1]$ . If necessary, we change that map by composing covering transformation, the lift coincides with the automorphism which is given by the path from  $\tilde{\iota}$  to  $\text{id}$  or  $-\text{id}$ . It is easy to see that  $\tilde{f}$  is an isomorphism of the principal  $Spin(3)$  bundle, since it is a lift of the isomorphism of the principal  $SO(3)$  bundle on  $P_{SO} \times [0, 1]$  and  $\tilde{\iota} \times \text{id}_{[0, 1]}$  is an isomorphism of principal  $Spin(3)$  bundle.  $\square$

Denote by  $Pin(2)_+$  the disjoint union of  $Pin(2)$  with the base point.

**Lemma 3.31.** *Let  $(Y, \mathfrak{t})$  be a spin rational homology sphere. Let  $\iota$  be a smooth odd involution preserving  $\mathfrak{t}$ . Let  $g$  be an  $\iota$ -invariant metric on  $Y$ . Suppose that the moduli space of irreducible solutions to the Seiberg–Witten equations on  $(Y, \mathfrak{t}, g)$  is discrete. Suppose also that  $\iota$  is isotopic to the identity of  $Y$  through the isometry group of  $(Y, g)$ . Let  $(a, \phi)$  be an irreducible solution to the Seiberg–Witten equations on  $(Y, \mathfrak{t}, g)$ . Then we have*

$$((a, \phi) \cdot Pin(2)_+)^I = *,$$

where  $*$  is the base point. Moreover, for any representation  $V$  of  $G$  and  $I$  (regarded as a  $\mathbb{Z}/2$ -action), we have

$$(\Sigma^V((a, \phi) \cdot Pin(2)_+))^I = *.$$

*Proof.* Denote by  $\text{Aut}(Y, \mathfrak{t}, g)$  the automorphism group of the spin 3-manifold  $(Y, \mathfrak{t}, g)$ . Note that all automorphisms of  $(Y, \mathfrak{t}, g)$  preserve the set of irreducible solutions  $\mathcal{S}^*(Y, \mathfrak{t}, g)$ , and its moduli space  $\mathcal{M}^*(Y, \mathfrak{t}, g) = \mathcal{S}^*(Y, \mathfrak{t}, g)/\mathcal{G}$ , where  $\mathcal{G} = \text{Map}(Y, S^1)$ . Let  $\mathcal{G}_0$  be a subgroup of  $\mathcal{G}$  with an exact sequence (such as the base point fibration)

$$1 \rightarrow \mathcal{G}_0 \rightarrow \mathcal{G} \rightarrow S^1 \rightarrow 1.$$

Let  $\tilde{\mathcal{M}}^*(Y, \mathfrak{t}, g) = \mathcal{S}^*(Y, \mathfrak{t}, g)/\mathcal{G}_0$ , on which the group  $Pin(2)$  acts. The action of  $Pin(2)$  descends to a free  $\mathbb{Z}_2$  action on  $\mathcal{M}^*(Y, \mathfrak{t}, g)$  via the exact sequence

$$1 \rightarrow S^1 \rightarrow Pin(2) \rightarrow \mathbb{Z}_2 \rightarrow 1.$$

Thus the image of  $(a, \phi) \cdot Pin(2)$  in  $\mathcal{M}^*(Y, \mathfrak{t}, g)$  consists of exactly two points,  $[(a, \phi)], [(a, \phi) \cdot j]$ .

By assumption, the moduli space  $\mathcal{M}^*(Y, \mathfrak{t}, g)$  is discrete, and hence the action of  $\text{Aut}(Y, \mathfrak{t}, g)$  on  $\mathcal{M}^*(Y, \mathfrak{t}, g)$  descends to an action of  $\pi_0(\text{Aut}(Y, \mathfrak{t}, g))$  on  $\mathcal{M}^*(Y, \mathfrak{t}, g)$ . By Lemma 3.30, the image of the pair  $(\iota, \tilde{\iota}) \in \text{Aut}(Y, \mathfrak{t}, g)$  under the natural map  $\text{Aut}(Y, \mathfrak{t}, g) \rightarrow \pi_0(\text{Aut}(Y, \mathfrak{t}, g))$  is either the identity element  $[(\text{id}_Y, \text{id}_{\mathfrak{t}})]$  or  $[(\text{id}_Y, -\text{id}_{\mathfrak{t}})]$ . Note that  $(\text{id}_Y, -\text{id}_{\mathfrak{t}})$  also acts trivially on  $\mathcal{M}^*(Y, \mathfrak{t}, g)$ , since the minus sign of  $-\text{id}_{\mathfrak{t}}$  corresponds to  $-1 \in S^1 \subset Pin(2)$ . Thus  $(\iota, \tilde{\iota})$  acts trivially on the two point set  $\{[(a, \phi)], [(a, \phi) \cdot j]\}$ , which is a subset of  $\mathcal{M}^*(Y, \mathfrak{t}, g)$ .

Recall that the involution  $I : \Gamma(\mathbb{S}) \rightarrow \Gamma(\mathbb{S})$  is given as the composition of  $(\iota, \tilde{\iota})$  and the right multiplication by  $j$ . Since we have seen that  $(\iota, \tilde{\iota})$  acts trivially on the two point set  $\{[(a, \phi)], [(a, \phi) \cdot j]\}$  and  $j$  acts non-trivially, we have that  $I$  acts non-trivially. This means that  $I$  swaps the connected component  $(a, \phi) \cdot S^1$  for  $(a, \phi) \cdot jS^1$ , and thus we have  $((a, \phi) \cdot Pin(2))^I = \emptyset$ . The assertion of the lemma immediately follows from this.  $\square$

Let  $p, q, r$  be coprime natural numbers. A Seifert metric  $g$  on  $\Sigma(p, q, r)$  is a Riemannian metric given by

$$(31) \quad \eta^2 + g_{S^2(p, q, r)}$$

where  $g_{S^2(p, q, r)}$  is the orbifold metric on  $S^2$  of type  $(p, q, r)$  and  $i\eta$  is a connection on the circle bundle  $Y \rightarrow S^2(p, q, r)$ . Henceforth we consider this metric on  $\Sigma(p, q, r)$ .

Recall that, except for  $S^3$  and  $\Sigma(2, 3, 5)$ , the mapping class group  $\Sigma(p, q, r)$  is isomorphic to  $\mathbb{Z}_2$ :

$$\pi_0(\text{Diff}^+(\Sigma(p, q, r))) \cong \mathbb{Z}_2.$$

(See, for example, [10, 60].) Therefore, the “half” of the orientation-preserving diffeomorphisms of  $\Sigma(p, q, r)$  are isotopic to the identity.

**Lemma 3.32.** *Let  $\iota : \Sigma(2, q, r) \rightarrow \Sigma(2, q, r)$  be the involution defined by*

$$(32) \quad \iota(z_1, z_2, z_3) = (-z_1, z_2, z_3).$$

*Then  $\iota$  is isotopic to the identity through the isometry group of the Seifert metric on  $\Sigma(2, q, r)$ .*

*Proof.* Recall that the standard  $S^1$ -action on  $\Sigma(2, q, r)$  is given by

$$u \cdot (z_1, z_2, z_3) := (u^{qr} z_1, u^{2r} z_2, u^{2q} z_3)$$

for  $u \in S^1$  and  $(z_1, z_2, z_3) \in \Sigma(2, q, r)$ . Since  $q, r$  are supposed to be odd, a path  $\{e^{i\theta}\}_{0 \leq \theta \leq \pi}$  in  $S^1$  gives rise to the desired isotopy between  $\text{id}_{\Sigma(2, q, r)}$  and  $\iota$ .  $\square$

*Remark 3.33.* In what follows, we state our results on calculations for the involution  $\iota$  on  $\Sigma(2, q, r)$  given by (32). But the same results explained below hold for all odd involution on  $\Sigma(2, q, r)$  which is isotopic to the identity, without any essential change.

**Lemma 3.34.** *For any  $(p, q, r)$ , we have*

$$n(\Sigma(p, q, r), \mathfrak{t}, g) = -n(-\Sigma(p, q, r), \mathfrak{t}, g)$$

*for the unique spin structure  $\mathfrak{t}$  and the Seifert metric  $g$ .*

*Proof.* Nicolaescu [64, Section 2.3] proved that the Dirac operator has zero kernel for general Seifert homology spheres with respect to the Seifert metrics. The assertion follows from this and (30).  $\square$

Henceforth, by abuse of notation, the unique spin structure on  $\Sigma(2, q, r)$  and the involution (32) are denoted by  $\mathfrak{t}, \iota$  respectively for all  $(2, q, r)$ .

**Proposition 3.35.** *Let  $Y_k = \Sigma(2, 3, 12k + 1)$  for a positive integer  $k$ . Then, there exists an integer  $b_k$  such that*

$$DSWF_G(Y_k, \mathfrak{t}, \iota) = [(S^0, b_k, 0)]$$

*for any positive integer  $k$ . In particular,*

$$\kappa(Y_k, \mathfrak{t}, \iota) = 0 \text{ and } \kappa(-Y_k, \mathfrak{t}, \iota) = 0$$

*hold for any positive integer  $k$ .*

*Proof.* First, we focus on the case  $k = 1$ . Let us recall Manolescu's calculation in [54] about the non-equivariant Seiberg–Witten Floer stable homotopy type of  $Y$  with respect to the unique spin structure  $\mathfrak{t}$  and the Seifert metric  $g$ . From [54, Equation (21)] and a sentence below it, it follows that

$$(33) \quad SWF(Y, \mathfrak{t}, g) = [(S^0 \vee \Sigma^{-1} Pin(2)_+, 0, 0)] = [(\mathbb{H}^+ \vee \Sigma^3 Pin(2)_+, 0, 1)],$$

where the convention of the (de)suspension in [54] is

$$(\mathbb{H}^+ \wedge \Sigma^3 Pin(2)_+, 0, 1) = \Sigma^{-\mathbb{H}} \Sigma^{\mathbb{R}^3} \mathbb{H}^+ \vee Pin(2)_+.$$

Here  $\mathbb{R}$  denotes the trivial  $Pin(2)$  representation. In the expression of the right-hand side of the first equality of (33), the third component 0 corresponds to  $n(Y, \mathfrak{t}, g) = 0$ . See Nicolaescu [65, Proposition 3.3].

In order to calculate  $DSWF_G$ , we replace  $[(\mathbb{H}^+ \vee \Sigma^3 Pin(2)_+, 0, 1)]$  with

$$[(\Sigma^{a\tilde{\mathbb{C}} \oplus (2b-2)(\mathbb{C}_+ \oplus \mathbb{C}_-)} D(\mathbb{H}^+ \vee \Sigma^3 Pin(2)_+), -a, 4 - 2b)]$$

so that  $\Sigma^{a\tilde{\mathbb{C}}\oplus(2b-2)(\mathbb{C}_+\oplus\mathbb{C}_-)}(\mathbb{H}^+ \vee \Sigma^3 Pin(2)_+)$  can be regarded as usual equivariant spaces.

We determine the  $I$ -invariant part of  $\Sigma^{a\tilde{\mathbb{C}}\oplus(2b-2)(\mathbb{C}_+\oplus\mathbb{C}_-)}D(\mathbb{H}^+ \vee \Sigma^3 Pin(2)_+)$ . This is  $(\Sigma^{a\tilde{\mathbb{C}}\oplus(2b-2)(\mathbb{C}_+\oplus\mathbb{C}_-)}D(\mathbb{H}^+ \vee \Sigma^3 Pin(2)_+))^I$ . The  $(\Sigma^{a\tilde{\mathbb{C}}\oplus(2b-2)(\mathbb{C}_+\oplus\mathbb{C}_-)}D(\mathbb{H}^+ \vee \Sigma^3 Pin(2)_+))^I$  is  $\Sigma^{a'\tilde{\mathbb{C}}\oplus(b-1)(\mathbb{C}_+\oplus\mathbb{C}_-)}$  for some  $a' \leq a$  because the  $I$  invariant part of the suspension  $\mathbb{H}^n$  to the space  $\mathbb{H}^+ \vee \Sigma^3 Pin(2)_+$  is  $\Sigma^{n^+\mathbb{C}_+ \vee \sigma^{n^-\mathbb{C}_-}$  and the double of this part is  $\Sigma^{(2b-2)(\mathbb{C}_+\oplus\mathbb{C}_-)}$  with  $n^+ + n^- = 2b - 2$ . It is easy to check that

$$D(\mathbb{H} \vee \Sigma^3 Pin(2)_+) = D(\mathbb{H}) \vee D(\Sigma^3 Pin(2)_+) \vee ((\mathbb{H}^\dagger)^+ \wedge \Sigma^3 Pin(2)_+) \vee (\mathbb{H}^+ \wedge (\Sigma^3 Pin(2)_+)^{\dagger}).$$

The  $I$ -invariant part of  $\mathbb{H}^+$  is  $\mathbb{C}_+$  or  $\mathbb{C}_-$  because  $I$  commutes the right action of  $i$ . Thus the  $I$ -invariant part of the  $D(\mathbb{H})$  is  $\mathbb{C}_+ \oplus \mathbb{C}_-$ . From the Lemma 3.31, the  $I$ -invariant parts of  $D(\Sigma^3 Pin(2)_+)$  and  $\Sigma^3 Pin(2)_+$  are  $*$ . Therefore, we have

$$D(\mathbb{H} \vee \Sigma^3 Pin(2)_+)^I = (\mathbb{C}_+ \oplus \mathbb{C}_-)^+$$

and

$$DSWF_G(Y, \mathfrak{t}, \iota) = [((\mathbb{C}_+ \oplus \mathbb{C}_-)^+, b_1, 1)]$$

for some integer  $b_1$ . When general  $k > 1$ , from [54], we have

$$(34) \quad SWF(Y, \mathfrak{t}, g) = [(S^0 \vee \underbrace{\Sigma^{-1} Pin(2)_+ \vee \cdots \vee \Sigma^{-1} Pin(2)_+}_k, 0, 0)]$$

$$(35) \quad = [(\mathbb{H}^+ \vee \underbrace{\Sigma^3 Pin(2)_+ \vee \cdots \vee \Sigma^3 Pin(2)_+}_k, 0, 1)].$$

A slightly change of the proof above actually shows that

$$DSWF_G(Y_k, \mathfrak{t}, \iota) = [(S^0, b_k, 0)]$$

for some integer  $b_k$  and

$$\kappa(Y_k, \mathfrak{t}, \iota) = 0$$

for all  $k$ . □

From the similar calculation of Proposition 3.35, we have the following result.

**Proposition 3.36.** *Let  $Y_k = \Sigma(2, 3, 12k + 5)$  for a non-negative integer  $k$ . Then, there exists an integer  $b_k$  such that*

$$DSWF_G(Y_k, \mathfrak{t}, \iota) = [(S^0, b_k, -1/2)]$$

for any positive integer  $k$ . Moreover,

$$\kappa(Y_k, \mathfrak{t}, \iota) = \frac{1}{2} \text{ and } \kappa(-Y_k, \mathfrak{t}, \iota) = -\frac{1}{2}$$

hold for any positive integer  $k$ .

*Proof.* From [54], we have

$$(36) \quad SWF(Y_k, \mathfrak{t}, g) = [(S^0 \vee \underbrace{\Sigma^{-1} Pin(2)_+ \vee \cdots \vee \Sigma^{-1} Pin(2)_+}_{k-1}, 0, -1/2)]$$

$$(37) \quad = [(\mathbb{H}^+ \vee \underbrace{\Sigma^3 Pin(2)_+ \vee \cdots \vee \Sigma^3 Pin(2)_+}_{k-1}, 0, 1/2)].$$

and this is only the  $-1/2$  shift of the third component of  $SWF(\Sigma(2, 3, 12k+1), \mathfrak{t}, g)$ . □

**Proposition 3.37.** *Let  $Y_k = \Sigma(2, 3, 12k - 5)$ , where  $k$  is a positive integer. Then, there exists an integer  $b_k$  such that*

$$DSWF_G(Y_k, \mathfrak{t}, \iota) = [(S^0, b_k, 1/2)]$$

for any positive integer  $k$ . In particular,

$$\kappa(Y_k, \mathfrak{t}, \iota) = -\frac{1}{2} \text{ and } \kappa(-Y_k, \mathfrak{t}, \iota) = \frac{1}{2}.$$

hold for any positive integer  $k$ .

*Proof.* Fix  $k$  and set  $Y = Y_k$ . Let  $g$  be a Seifert metric on  $Y$ . The Seiberg–Witten Floer stable homotopy type of  $Y$  is given by

$$SWF(Y, \mathfrak{t}, g) = \left[ (\widetilde{Pin}(2) \vee \vee^{k-1} \Sigma Pin(2)_+, 0, 1/2) \right],$$

where  $\widetilde{Pin}(2)$  denotes the unreduced suspension of  $Pin(2)$ . See [54, Section 5.2]. The third component  $1/2$  corresponds to  $n(Y, \mathfrak{t}, g)/2 = 1/2$ . See Nicolaescu [65, Proposition 3.8]. As explained in [54, Section 5.2],  $\widetilde{Pin}(2)$  and  $(k-1)$  copies of  $\Sigma Pin(2)_+$  correspond to  $2k$  irreducible solutions. So they arise from  $k$  copies of  $Pin(2)$ , which are the set of irreducible solutions (before dividing by  $S^1$ ).

As in the proof of Proposition 3.35, we calculate

$$\left( \Sigma^{a\tilde{\mathcal{C}} \oplus 2b(\mathbb{C}_+ \oplus \mathbb{C}_-)} (D(\widetilde{Pin}(2) \vee \vee^{k-1} \Sigma Pin(2)_+))^I \right).$$

Let  $\iota : Y \rightarrow Y$  for the involution on  $Y$  given by (32). Then, just like the argument in Proposition 3.35, one sees that  $(\iota, \tilde{\iota})$  acts trivially on the set of connected components of the  $k$  copies of  $Pin(2)$  since  $(\iota, \tilde{\iota})$  is isotopic to  $(\text{id}_Y, \text{id}_{P_{Spin}})$ , and thus we have that

$$\left( D(\widetilde{Pin}(2) \vee \vee^{k-1} \Sigma Pin(2)_+)^I = D(\widetilde{Pin}(2))^I \vee \vee^{k-1} \Sigma Pin(2)_+^I = D(S^0) \right)$$

from Lemma 3.31. This implies

$$\begin{aligned} & \left( \Sigma^{a\tilde{\mathcal{C}} \oplus 2b(\mathbb{C}_+ \oplus \mathbb{C}_-)} (\widetilde{Pin}(2) \vee \vee^{k-1} \Sigma Pin(2)_+)^I \right) \\ &= \Sigma^{a'\tilde{\mathcal{C}} \oplus b(\mathbb{C}_+ \oplus \mathbb{C}_-)} (D(S^0)). \end{aligned}$$

This implies

$$DSWF_G(Y, \mathfrak{t}, \iota) = [(D(S^0), a', n(Y, \mathfrak{t}, g)/2)]$$

for any positive integer  $k$ . This implies

$$\kappa(Y, \mathfrak{t}, \iota) = -n(Y, \mathfrak{t}, g)/2, \quad \kappa(-Y, \mathfrak{t}, \iota) = -n(-Y, \mathfrak{t}, g)/2.$$

As mentioned, Nicolaescu [65, Proposition 3.8] showed that  $n(Y, \mathfrak{t}, g) = 1$ . Moreover, Nicolaescu [64, Section 2.3] proved that the Dirac operator has zero kernel for general Seifert homology spheres with respect to the Seifert metrics. It follows from these facts and (30) that

$$\kappa(Y, \mathfrak{t}, \iota) = -1/2.$$

□

**Proposition 3.38.** *Let  $Y_k = \Sigma(2, 3, 12k - 1)$ , where  $k$  is a positive integer. Then, there exists a positive integer  $b_k$  such that*

$$DSWF_G(Y_k, \mathfrak{t}, \iota) = [(S^0, b_k, 0)]$$

for any positive integer  $k$ . In particular,

$$\kappa(Y_k, \mathfrak{t}, \iota) = \kappa(-Y_k, \mathfrak{t}, \iota) = 0.$$

*Proof.* The Seiberg–Witten Floer stable homotopy type of  $Y$  is given by

$$SWF(Y, \mathfrak{t}, g) = \left[ (\widetilde{Pin}(2) \vee \vee_{k-1} \Sigma Pin(2)_+, 0, 0) \right].$$

Then, by the similar calculation of Proposition 3.35,

$$DSWF_G(Y, \mathfrak{t}, \iota) = [(S^0, b_k, 0)].$$

□

For convenience, we summarize the value of  $\kappa$  for the Brieskorn spheres we considered above:

**Corollary 3.39.** *Let  $\iota : \Sigma(2, q, r) \rightarrow \Sigma(2, q, r)$  be the involution defined by  $\iota(z_1, z_2, z_3) = (-z_1, z_2, z_3)$ . Denote by  $\mathfrak{t}$  the unique spin structure on  $\Sigma(2, q, r)$ . Then we have:*

$$\begin{aligned} \kappa(\Sigma(2, 3, 12n - 5), \mathfrak{t}, \iota) &= -1/2, & \kappa(-\Sigma(2, 3, 12n - 5), \mathfrak{t}, \iota) &= 1/2, \\ \kappa(\Sigma(2, 3, 12n - 1), \mathfrak{t}, \iota) &= 0, & \kappa(-\Sigma(2, 3, 12n - 1), \mathfrak{t}, \iota) &= 0, \\ \kappa(\Sigma(2, 3, 12n + 5), \mathfrak{t}, \iota) &= 1/2, & \kappa(-\Sigma(2, 3, 12n + 5), \mathfrak{t}, \iota) &= -1/2, \\ \kappa(\Sigma(2, 3, 12n + 1), \mathfrak{t}, \iota) &= 0, & \kappa(-\Sigma(2, 3, 12n + 1), \mathfrak{t}, \iota) &= 0. \end{aligned}$$

*Proof.* This is just a collection of Propositions 3.35 to 3.38. □

**3.9. Knot invariants.** For any oriented knot  $K$  in  $S^3$ , we can uniquely associate a double branched cover

$$\Sigma(K) \rightarrow S^3$$

and an involution  $\iota$  on  $\Sigma(K)$ . It is proven that  $H^1(\Sigma(K); \mathbb{Z}_2) = \{0\}$ . So, the isomorphism class of the spin structure  $\mathfrak{t}$  on  $\Sigma(K)$  is automatically preserved by  $\iota$ .

**Definition 3.40.** For any knot  $K$  in  $S^3$ , we define the *Seiberg–Witten Floer stable homotopy type* of  $K$  by

$$DSWF(K) := DSWF_G(\Sigma(K), \mathfrak{t}, \iota).$$

We also define the *Seiberg–Witten Floer  $K$ -theory* of  $K$  by

$$DSWFK(K) := DSWFK_G(\Sigma(K), \mathfrak{t}, \iota),$$

and the  *$K$ -theoretic Frøyshov invariant* of  $K$  by

$$\kappa(K) := k(DSWF(K)) \in \mathbb{Q}.$$

*Remark 3.41.* Since the observation above (beginning of Subsection 3.9) only use the homological information of  $S^3$ , the same observation can be applied to the case of a pair  $(Y, K)$  of an oriented homology 3-sphere  $Y$  and an oriented knot  $Y$ . Thus, we can define the Seiberg–Witten Floer stable homotopy type  $DSWF(Y, K)$ , the Seiberg–Witten Floer  $K$ -theory  $DSWFK(Y, K)$  and the  $K$ -theoretic Frøyshov invariant  $\kappa(Y, K)$ . In Lemma 3.44, we will see that  $\kappa(K)$  is a concordance invariant.

For the generalization  $\kappa(Y, K)$ , we can see that  $\kappa(Y, K)$  is invariant under *homology concordance*, which is defined by the following way: if there are a homology cobordism  $W$  from  $Y_0$  to  $Y_1$  and an embedded concordance (annulus) in  $W$  from  $K_0$  to  $K_1$ , then we call  $(Y_0, K_0)$  and  $(Y_1, K_1)$  are homology concordance.

*Remark 3.42.* Before proving several properties of  $\kappa$ , we explain how to compute  $b_i^+$  for double branched covers. Let  $X$  be a simply-connected smooth closed 4-manifold  $X$  and an embedded oriented surface  $S$  with  $[S]$  is divisible by 2, we have a double branched cover

$$\Sigma(S) \rightarrow X.$$

It is proven in [31, Page 254 Lemma] that

$$(38) \quad b_i^+(\Sigma(S)) = b^+(X).$$

Also, we will use (38) for 4-manifolds  $X$  with several  $S^3$ -boundaries  $S_1^3 \cup \cdots \cup S_m^3$ . Suppose  $S$  is properly smoothly embedded surface in  $X$  such that  $S \cap S_i^3$  is a knot for any  $i$ . We can easily see the proof in [31] can also be applied to this case.

Henceforth, if we say a surface, we always assume that it is connected.

When  $K$  and  $K'$  are isotopic, then there is an orientation preserving diffeomorphism

$$\phi : \Sigma(K) \rightarrow \Sigma(K')$$

which is  $\mathbb{Z}_2$ -equivariant. This enables us to prove the following:

**Lemma 3.43.**  *$DSWF(K)$  and  $\kappa(K)$  are isotopy invariants of knots.*

The invariant  $\kappa(K)$  is actually invariant under knot concordance. Let  $\mathcal{C}$  be the knot concordance group.

**Lemma 3.44.** *The correspondence*

$$\mathcal{C} \rightarrow \mathcal{LE}_G, \quad K \mapsto [DSWF_G(\Sigma(K), \mathfrak{t}, \iota)]_{loc}$$

*is a well-defined map. Moreover, since  $\kappa(K)$  can be recovered from the local equivalence class*

$$[DSWF_G(\Sigma(K), \mathfrak{t}, \iota)]_{loc},$$

*$\kappa(K)$  is a concordance invariant.*

*Proof.* Let  $S$  be a concordance from  $K$  to  $K'$  in  $I \times S^3$ . Then the double branched cover  $\Sigma(S)$  along  $S$  gives a  $\mathbb{Z}_2$ -equivariant  $\mathbb{Z}_2$ -homology cobordism from  $\Sigma(K)$  to  $\Sigma(K')$ . Note that the isomorphism class of the spin structure on  $\Sigma(S)$  is preserved under the involution.

We have an associate doubled cobordism map

$$(39) \quad D(f) : \Sigma^{m_0} \tilde{\mathcal{C}} \Sigma^{n_0(\mathbb{C}_+ \oplus \mathbb{C}_-)} D(I_{-\mu}^{-\lambda}(\Sigma(K))) \rightarrow \Sigma^{m_1} \tilde{\mathcal{C}} \Sigma^{n_1(\mathbb{C}_+ \oplus \mathbb{C}_-)} D(I_{\lambda}^{\mu}(\Sigma(K'))).$$

(24) implies

$$\begin{aligned} & m_0 - m_1 \\ &= \dim_{\mathbb{R}}(V_1(\tilde{\mathbb{R}}_{\lambda}^0)) - \dim_{\mathbb{R}}(V_0(\tilde{\mathbb{R}}_{-\mu}^0)) - b^+(\Sigma(S)) + b_i^+(\Sigma(S)) \\ &= \dim_{\mathbb{R}}(V_1(\tilde{\mathbb{R}}_{\lambda}^0)) - \dim_{\mathbb{R}}(V_0(\tilde{\mathbb{R}}_{-\mu}^0)). \end{aligned}$$

Here we used Remark 3.42 to calculate  $b_t^+$ . This means  $D(f)$  is a local map. By considering the same discussion for  $-S$ , we see

$$[DSWF_G(\Sigma(K), \mathfrak{t}, \iota)]_{\text{loc}} = [DSWF_G(\Sigma(K'), \mathfrak{t}, \iota)]_{\text{loc}}.$$

By Lemma 3.14, we saw  $\kappa(K)$  is recovered from  $[DSWF_G(\Sigma(K), \mathfrak{t}, \iota)]_{\text{loc}}$ . This completes the proof.  $\square$

Let us give calculations of  $\kappa$  for two bridge knots. Before that, we remark on the sign convention of the knot signature.

*Remark 3.45.* We use the same convention of the signature as in [9, 69]. Namely, the signature of  $T(2, 3)$  is given by  $\sigma(T(2, 3)) = -2$ . On the other hand, the sign convention of the signature in the Knot Atlas [1] is opposite to ours.

**Lemma 3.46.** *For any two bridge knot  $K(p, q)$  whose branched cover is  $L(p, q)$ , one has*

$$(40) \quad DSWF(K) = [(S^0, 0, \frac{1}{16}\sigma(K(p, q)))]$$

and

$$(41) \quad \kappa(K(p, q)) = -\frac{1}{16}\sigma(K(p, q)).$$

*Proof.* In Proposition 3.28, we showed that

$$DSWF_G(Y, \mathfrak{t}, \iota) = [(S^0, 0, n(Y, \mathfrak{t}, g)/2)]$$

and

$$\kappa(Y, \mathfrak{t}, \iota) = -n(Y, \mathfrak{t}, g)/2$$

for any spin rational homology 3-sphere  $(Y, \mathfrak{t})$  equipped with an involution  $\iota$  preserving  $\mathfrak{t}$  such that  $Y$  admits an  $\iota$ -invariant positive scalar curvature metric  $g$ . Since the two bridge knot  $K(p, q)$  is regarded as the fixed point set of the complex conjugation of the lens space  $L(p, q)$ . Note that the standard positive scalar curvature metric  $g$  on  $L(p, q)$  is preserved by the conjugation. So, one has

$$(42) \quad \kappa(K(p, q)) = \kappa(L(p, q), \mathfrak{t}, \iota) = -n(L(p, q), \mathfrak{t}, g)/2.$$

On the other hand, since  $g$  is a positive scalar curvature metric again, we also have

$$(43) \quad \delta(L(p, q), \mathfrak{t}) = -n(L(p, q), \mathfrak{t}, g).$$

(See [55].) Also, it follows from [58, Theorem 1.2] that

$$(44) \quad 2d(\Sigma(K(p, q))) = -\frac{1}{2}\sigma(K(p, q)).$$

In general, for a general homology sphere  $Y$ , the  $d$  invariant and the  $\delta$  invariant are related by

$$(45) \quad \delta(Y) = d(Y)/2.$$

(See, for example, [51, Remark 1.1].) Now it follows from (42), (43), (44), (45) that

$$\kappa(K(p, q)) = -\frac{1}{16}\sigma(K(p, q)) = -n(L(p, q), \mathfrak{t}, g)/2,$$

and we have (40) and (41).  $\square$

Now we give a proof of Theorem 1.5.



*Proof of Theorem 1.5.* The proof is written by induction on the number of connected sum components. From Lemma 3.46, when  $K = K(p, q)$ , the quality

$$DSWF(K) = [(S^0, 0, \frac{1}{16}\sigma(K(p, q)))]$$

holds. In the next step, we consider  $K = K(p, q)\#K(p', q')$ . We will show

$$(46) \quad [DSWF(K)]_{\text{loc}} = [(S^0, 0, \frac{1}{16}(\sigma(K(p, q) + \sigma(K(p', q'))))].$$

In order to prove (46), we consider two knot cobordisms  $(W, S)$  and  $(W^*, S^*)$ :

- The knot cobordism  $(W, S)$  is a cobordism from  $K$  to  $K(p, q) \cup K(p', q')$ . The cobordism  $W$  is a 3-handle cobordism attached to  $I \times S^3$ . The cobordism  $S$  inside  $W$  is a 1-handle cobordism attached to  $K$  corresponding to the connected sum decomposition of  $K = K(p, q)\#K(p', q')$ .
- The knot cobordism  $(W^*, S^*)$  is a cobordism from  $K(p, q) \cup K(p', q')$  to  $K$ . The cobordism  $W^*$  is a 1-handle cobordism attached to  $I \times S^3$ . The cobordism  $S$  inside  $W$  is a 1-handle cobordism attached to  $K(p, q) \cup K(p', q')$  corresponding to the connected sum decomposition of  $K = K(p, q)\#K(p', q')$ .

Then, we consider the double branched covering spaces  $\Sigma(S)$  and  $\Sigma(S')$  of  $W$  and  $W^*$  along  $S$  and  $S'$ . It is easy to see  $b^+(\Sigma(S)) = b^+(\Sigma(S^*)) = b_1(\Sigma(S)) = b_1(\Sigma(S^*)) = 0$ . Also,  $H^1(\Sigma(S); \mathbb{Z}_2) = H^1(\Sigma(S^*); \mathbb{Z}_2) = 0$ . Associated to these  $\mathbb{Z}_2$ -equivariant cobordism  $\Sigma(S)$  and  $\Sigma(S')$ , we have two maps

$$\begin{aligned} D(f) : \Sigma^{m_0 \tilde{C}} \Sigma^{n_0(C_+ \oplus C_-)} D(I_{-\mu}^{-\lambda}(\Sigma(K))) &\rightarrow \\ \Sigma^{m_1 \tilde{C}} \Sigma^{n_1(C_+ \oplus C_-)} D(I_{\lambda}^{\mu}(\Sigma(K(p, q))) \wedge D(I_{\lambda}^{\mu}(\Sigma(K(p', q')))) & \end{aligned}$$

and

$$\begin{aligned} D(f^*) : \Sigma^{m_0^* \tilde{C}} \Sigma^{n_0^*(C_+ \oplus C_-)} D(I_{\lambda}^{\mu}(\Sigma(K(p, q))) \wedge D(I_{\lambda}^{\mu}(\Sigma(K(p', q')))) &\rightarrow \\ \Sigma^{m_1^* \tilde{C}} \Sigma^{n_1^*(C_+ \oplus C_-)} D(I_{-\mu}^{-\lambda}(\Sigma(K))) & \end{aligned}$$

Because of (24) and Remark 3.42, we see that these two maps  $D(f)$  and  $D(f^*)$  are local maps. So, one concludes

$$\begin{aligned} [DSWF(K)]_{\text{loc}} &= [DSWF(K(p, q) \wedge DSWF(K(p', q')))]_{\text{loc}} \\ &= [(S^0, 0, \frac{1}{16}(\sigma(K(p, q) + \sigma(K(p', q'))))]. \end{aligned}$$

When  $K = m_1 K(p_1, q_1)\#\dots\#m_n K(p_n, q_n)$ , an inductive proof can be applied. As a conclusion, we have

$$[DSWF(K)]_{\text{loc}} = [(S^0, 0, \frac{1}{16} \sum_{i=1}^n m_i \sigma(K(p_i, q_i))].$$

So, the invariant  $\kappa$  can be computed as

$$\kappa(K) = -\frac{1}{16} \sum_{i=1}^n m_i \sigma(K(p_i, q_i)) = -\frac{1}{16} \sigma(K),$$

since the signature is a homomorphism.  $\square$

We also provide calculations of  $\kappa(Y)$  for a certain class of torus knots giving a proof of Theorem 1.6.

*Proof of Theorem 1.6.* This is a corollary of Proposition 3.38, Proposition 3.37, Proposition 3.36, and Proposition 3.35. Indeed,  $T(3, 12k - 5)$  can be regarded as the branch locus of the branched cover

$$\Sigma(2, 3, 12k - 5) \rightarrow S^3$$

whose involution is given by (32). Next, we consider the  $n$ -fold connected sum  $K_n$  of  $T(3, 7)$ . It is proven that

$$DSWF(K_1) = [(S^0, 0, 1/2)].$$

So, by the same proof in Theorem 1.5, we obtain local maps

$$DSWF(K_1) \wedge DSWF(K_1) \rightarrow DSWF(K_1 \# K_1)$$

and

$$DSWF(K_1 \# K_1) \rightarrow DSWF(K_1) \wedge DSWF(K_1)$$

as in the proof of (3). This implies  $\kappa(K_2) = 1$  for  $K_2 = K_1 \# K_1$ . The remaining part is completely the same as in the proof of Theorem 1.5. Also, for  $T(3, 12k - 5)$ , since it is proven in (32) that

$$DSWF(K_1) = [(S^0, 0, 1/2)],$$

the same argument works. For the other torus knots, the same argument also works. This completes the proof.  $\square$

#### 4. APPLICATIONS TO KNOT THEORY

**4.1. Branched covers of punctured 4-manifolds.** For a simply-connected smooth closed 4-manifold  $X$  and an embedded oriented surface  $S$  with  $[S]$  is divisible by 2, we have a double branched cover

$$\Sigma(S) \rightarrow X.$$

When  $H^1(X; \mathbb{Z}_2) = 0$ , one can see the double branched cover is unique. The following calculations are proven in [32, 33]:

- (i)  $\sigma(\Sigma(S)) = 2\sigma(X) - \frac{1}{2}[S]^2$
- (ii)  $b^+(\Sigma(S)) = 2b^+(X) + g(S) - \frac{1}{4}[S]^2$ ,  $b^-(\Sigma(S)) = 2b^-(X) + g(S) + \frac{1}{4}[S]^2$  and
- (iii)  $b_1(\Sigma(S)) = 0$ .

Also, in [62, Theorem 1.1], it is proven that  $\Sigma(S)$  has a spin structure if and only if  $PD(w_2(X)) = \frac{1}{2}[S] \pmod{2}$ . We suppose  $PD(w_2(X)) = \frac{1}{2}[S] \pmod{2}$ .

**Lemma 4.1.** *Under above assumptions,  $H^1(\Sigma(S); \mathbb{Z}_2) = \{0\}$  holds. In particular, the spin structure on  $\Sigma(S)$  is unique up to isomorphism.*

*Proof.* In [45, Proposition 2.1], it is proven that there is no 2-torsion in  $H_*(\Sigma(S); \mathbb{Z})$ . Thus, the universal coefficient theorem implies

$$0 \rightarrow \text{Ext}_{\mathbb{Z}}^1(H_0(\Sigma(S); \mathbb{Z}), \mathbb{Z}_2) \rightarrow H^1(\Sigma(S); \mathbb{Z}_2) \rightarrow \text{Hom}(H_1(\Sigma(S); \mathbb{Z}), \mathbb{Z}_2) \rightarrow 0.$$

is exact. This shows  $H^1(\Sigma(S); \mathbb{Z}_2) = \{0\}$ .  $\square$

Let  $K$  be an oriented knot in  $S^3$ . Let  $X$  be a simply-connected 4-manifold  $X$  bounded by  $S^3$  and  $S$  be an oriented compact connected surface  $S$  bounded by  $K$  satisfying  $[S]$  is divisible by 2. Then one can associate the double branched cover

$$\Sigma(S) \rightarrow X$$

of 4-manifolds with boundary.

The following provides a sufficient condition so that the double branched cover has the unique spin structure.

**Lemma 4.2.** *We suppose  $PD(w_2(X)) = [S] \bmod 2$ . The double branched cover  $\Sigma(S)$  along  $S$  has the unique spin structure. Moreover, the following equalities*

$$(47) \quad \sigma(\Sigma(S)) = 2\sigma(X) - \frac{1}{2}[S]^2 + \sigma(K),$$

$$(48) \quad b^+(\Sigma(S)) = 2b^+(X) + g(S) - \frac{1}{4}[S]^2 + \frac{1}{2}\sigma(K),$$

$$(49) \quad b^-(\Sigma(S)) = 2b^-(X) + g(S) + \frac{1}{4}[S]^2 - \frac{1}{2}\sigma(K), \text{ and}$$

$$(50) \quad b_1(\Sigma(S)) = 0$$

hold.

Recall that, as mentioned in Remark 3.45, our sign convention of the knot signature is the same as in [9, 69].

*Proof.* When  $X$  is a closed 4-manifold, the existence condition of a spin structure on the double branched cover is completely determined in [62, Theorem 1.1]. Define  $X' := X \cup_{S^3} D^4$ . Take a properly embedded oriented surface  $S'$  in  $D^4$  bounded by  $-K^*$ . Consider the union  $S^\# := S \cup S' \subset X'$ . Then we just apply [62, Theorem 1.1] to  $(X', S^\#)$  and obtain a spin structure on  $\Sigma(S^\#)$ . As a restriction, we obtain a spin structure on  $\Sigma(S)$ . Next, we see the spin structure is unique up to isomorphism. It is sufficient to say  $H^1(\Sigma(S); \mathbb{Z}_2) = \{0\}$ . The Mayer-Vietoris exact sequence for  $(\Sigma(S), \Sigma(S'))$  implies

$$0 \rightarrow H^1(\Sigma(S^\#); \mathbb{Z}_2) \rightarrow H^1(\Sigma(S); \mathbb{Z}_2) \oplus H^1(\Sigma(S'); \mathbb{Z}_2) \rightarrow H^1(\Sigma(K); \mathbb{Z}_2) \rightarrow \dots$$

One can verify  $H^1(\Sigma(K); \mathbb{Z}_2) = 0$ . By Lemma 4.1, we see  $H^1(\Sigma(S); \mathbb{Z}_2) = \{0\}$ .

Next, we see the equations above. The double branched cover along  $S^\#$  has the following decomposition:

$$\Sigma(S^\#) = \Sigma(S) \cup \Sigma(S').$$

Thus, one has

$$\sigma(\Sigma(S^\#)) = \sigma(\Sigma(S)) + \sigma(\Sigma(S'))$$

and

$$b^+(\Sigma(S^\#)) = b^+(\Sigma(S)) + b^+(\Sigma(S')).$$

On the other hand, we have

$$\sigma(\Sigma(S^\#)) = 2\sigma(X) - \frac{1}{2}[S]^2$$

and

$$b^+(\Sigma(S^\#)) = 2b^+(X) + g(S^\#) - \frac{1}{4}[S^\#]^2.$$

It is proven in [22, Theorem 6] that

$$\sigma(\Sigma(S')) = \sigma(-K^*) = -\sigma(K).$$

Also, we can verify

$$b^+(\Sigma(S')) = g(S') + \frac{1}{2}\sigma(-K^*) = g(S') - \frac{1}{2}\sigma(K).$$

These equations imply

$$\sigma(\Sigma(S)) = 2\sigma(X) - \frac{1}{2}[S]^2 - \sigma(-K^*) = 2\sigma(X) - \frac{1}{2}[S]^2 + \sigma(K)$$

and

$$b^+(\Sigma(S)) = 2b^+(X) + g(S) - \frac{1}{4}[S]^2 + \frac{1}{2}\sigma(K).$$

□

**4.2. Genus bounds from Theorem 1.1.** In this subsection, we provide a genus bound in 4-manifolds. Let  $X$  be a simply-connected smooth 4-manifold.

**Definition 4.3.** For a fixed homology class  $x \in H_2(X; \mathbb{Z})$  and an oriented knot  $K$  in  $S^3$ , we define the  $X$ -genus of  $K$  is defined by

$$g_{X,x}(K) := \min\{g(\Sigma) \mid \Sigma \text{ is an oriented properly embedded connected surface in } X \setminus D^4 \text{ bounded by } K, x = [\Sigma] \in H_2(X, \partial X; \mathbb{Z})\}$$

The  $X$ -genus has been studied in various situations. Also, by considering locally flat embedding, we define the topological version  $g_{X,x}^{Top}(K)$  of  $g_{X,x}(K)$ . In general, these two invariants  $g_{X,x}^{Top}(K)$  and  $g_{X,x}(K)$  are different. When  $X = S^4$  and  $x = 0$ ,  $g_{X,x}(K)$  is called the *smooth slice genus* of  $K$ , and denoted by  $g_4(K)$ . Obviously, we have

$$g_{X,x}(K) \leq g_4(K).$$

Also, when we take  $K = U$  (the unknot),  $g_{X,x}(U)$  is called the *minimal genus* of  $(X, x)$ .

We now prove Theorem 1.3.

*Proof of Theorem 1.3.* We consider the double branched cover  $\Sigma(S)$  along  $S$ . By Lemma 4.2,  $\Sigma(S)$  has the unique spin structure. In particular, the branched involution preserves the isomorphism class of the spin structure. Now we apply Theorem 1.1 and obtain

$$-\frac{\sigma(\Sigma(S))}{16} \leq b^+(\Sigma(S)) - b^+_i(\Sigma(S)) + \kappa(K).$$

Moreover, using Lemma 4.2 and Remark 3.42, we obtain the equalities

$$(51) \quad \sigma(\Sigma(S)) = 2\sigma(X) - \frac{1}{2}[S]^2 + \sigma(K)$$

$$(52) \quad b^+(\Sigma(S)) = 2b^+(X) + g(S) - \frac{1}{4}[S]^2 - \frac{1}{2}\sigma(K) \text{ and}$$

$$(53) \quad b^+_i(\Sigma(S)) = b^+(X).$$

Thus, we obtain

$$-\left(\frac{2\sigma(X) - \frac{1}{2}[S]^2 + \sigma(K)}{16}\right) \leq b^+(X) + g(S) - \frac{1}{4}[S]^2 + \frac{1}{2}\sigma(K) + \kappa(K).$$

This completes the proof. □

*Remark 4.4.* When  $X = D^4$ , then Theorem 1.3 implies

$$(54) \quad -\frac{9}{16}\sigma(K) \leq g_4(K) + \kappa(K).$$

For a finite connected sum of two bridge knots, (54) shows

$$-\frac{1}{2}\sigma(K) \leq g_4(K),$$

which is equivalent to a constraint from the signature given in Theorem 6.7. Also, for torus knots,  $g_4(T(p, q)) = \frac{(p-1)(q-1)}{2}$  is known as Milnor's conjecture. On the other hand, for example, our inequalities for  $\#_n T(3, 7)$  imply the following weak inequality:

$$\frac{9}{4}n + \frac{1}{2}n = \frac{11}{4}n \leq g_4(\#_n T(3, 7)) = 3n.$$

So, our inequality can be also used to detect the 4-genus of  $\#_n T(3, 7)$ .

*Remark 4.5.* When  $K = U$  (the unknot), Theorem 1.3 implies a version of 10/8-inequality corresponding to the inequality given in [39]. In this case, (2) gives

$$-\frac{\sigma(X)}{8} + \frac{9}{32}[S]^2 \leq b^+(X) + g(S)$$

for a simply-connected closed oriented smooth 4-manifold  $X$  and an oriented compact properly smoothly embedded surface  $S$  in  $X$  such that the homology class  $[S]$  of  $S$  is divisible by 2 and  $PD(w_2(X)) = [S]/2 \pmod{2}$ . For example, for a positive integer  $n$  and any class  $x \in H_2(\#_n K3)$  satisfying that  $x$  is divisible by 2 and  $0 = x/2 \pmod{2}$ , we have the following genus bound:

$$\frac{9}{32}[S]^2 \leq n + g_{\#_n K3, x}(U).$$

*Remark 4.6.* As it is remarked in Remark 3.41, we can generalize the invariant  $\kappa(K)$  to an invariant  $\kappa(Y, K)$  for a pair of an oriented homology 3-sphere and an oriented knot  $K$ . We can also generalize Theorem 1.3 to a theorem for such an invariant.

When  $g_{X, H=0}(K) = 0$ , a knot  $K$  is called (*smoothly*) *H-slice in a closed 4-manifold*  $X$ .

4.2.1. *Results on  $\#_n K3$ .* We first give a proof of Theorem 1.9.

*Proof of Theorem 1.9.* We just apply Theorem 1.3 to the case  $X = \#_n K3$ ,  $[S] = 0 \in H_2(\#_n K3)$  and obtain the inequality:

$$2n - \frac{1}{16}\sigma(K) \leq 3n + \frac{1}{2}\sigma(K) + \kappa(K) + g(S).$$

By combining this with (41), we obtain

$$-\frac{1}{2}\sigma(K) \leq n + g(S).$$

This completes the proof.  $\square$

*Remark 4.7* (Comparison with other methods). We compare Theorem 1.3 with Maloescu's relative 10/8-inequality, obstructions from Arf invariant and signature function. We review these methods in Section 6.

- (1) Manolescu's relative 10/8 inequality [54] enable us to prove that for any finite connected sum of two bridge knots  $K$  in  $S^3$ , we have

$$(55) \quad -\frac{1}{2}\sigma(K) \leq 2n + g(S).$$

For more details, see Corollary 6.3.

- (2) From an obstruction via Arf invariant (For more details see Theorem 6.6), we have

$$(56) \quad \text{Arf}(K) \equiv 0 \pmod{2}.$$

- (3) From an obstruction via the Tristram-Levine signature  $\sigma_K(t)$  (For more details see Theorem 6.7), we have

$$(57) \quad \max_m \max_{r \in \{1, \dots, m-1\}} \frac{1}{2} |\sigma_K(e^{2\pi r i m}) - 16n| \leq 11n + g.$$

*Example 4.8* (Example 1.10). Let  $K_m$  be the  $m$ -fold connected sum of  $5_2$  for  $m \in \mathbb{Z}_{>0}$ . Note that  $5_2$  is two bridge knot  $K(7, 3)$  and  $\text{Arf}(5_2) = 0$ . Then  $\sigma(K_m) = -2m$ . Suppose  $S$  is a properly embedded oriented surface in  $\#_n K3 \setminus \text{int } D^4$  bounded by  $K$  such that  $[S] = 0$ .

Then so our inequality Theorem 1.9 implies

$$m \leq n + g(S)$$

holds. On the other hand, (55) implies

$$m \leq 2n + g(S).$$

Also (57) implies

$$\max\{8n, |-\frac{1}{2}m - 8n|, |-m - 8n|\} \leq 11n + g(S).$$

Note that a family of topologically H-slice but not smoothly H-slice knots in the punctured  $\#_3 K3$  are given in [34]. However, the Bauer-Furuta type invariant used in [34] vanishes for  $\#_n K3$  when  $n \geq 4$ . On the other hand, our invariant  $\kappa(K)$  may be used to give such examples.

**Problem 4.9.** *Let  $n$  be a positive integer with  $n \geq 4$ . Is there a topologically H-slice but not smoothly H-slice knot in the punctured  $\#_n K3$  ?*

4.2.2. *Results on  $\#_n \mathbb{C}P^2 \#_m \mathbb{C}P^2$ .* We also consider  $X := \#_n \mathbb{C}P^2 \#_m (-\mathbb{C}P^2)$  for positive integers  $n$  and  $m$ . When  $\min\{m, n\} \geq 2$ , since known gauge theoretic invariant of  $X$  vanishes, there is no way to obtain adjunction type inequality.

**Corollary 4.10.** *Let  $X$  be  $\#_n \mathbb{C}P^2 \#_m (-\mathbb{C}P^2)$  for a pair  $(m, n)$  of non-negative integers. Let  $H$  be an element in  $H_2(X; \mathbb{Z})$  such that*

$$H = \sum_{1 \leq i \leq n} a_i e_i^+ + \sum_{1 \leq j \leq m} b_j e_j^-$$

for integers  $a_i, b_j$  with  $a_i \equiv 2 \pmod{4}$  and  $b_j \equiv 2 \pmod{4}$ , where  $\{e_i^+\}$  and  $\{e_j^-\}$  are generators corresponding to  $H_2(\mathbb{C}P^2)$  and  $H_2(-\mathbb{C}P^2)$  for each summand. Then for any knot  $K$  in  $S^3$ , we have

$$(58) \quad \frac{1}{32} \left( -36n + 4m + 9 \left( \sum_{i=1}^n a_i^2 - \sum_{j=1}^m b_j^2 \right) - 18\sigma(K) - 32\kappa(K) \right) \leq g_{X,H}(K).$$

In particular, when  $K$  is a finite connected sum of two bridge knots, one has

$$\frac{1}{32} \left( -36n + 4m + 9 \left( \sum_{i=1}^n a_i^2 - \sum_{j=1}^m b_j^2 \right) - 16\sigma(K) \right) \leq g_{X,H}(K).$$

This is just a corollary of Theorem 1.3.

*Remark 4.11.* Under the same assumptions of Corollary 4.10, the following facts hold:

- (1) Manolescu's relative 10/8 inequality [54](see Theorem 6.1) enable us to prove that implies

$$\frac{1}{16} \left( -36n + 4m + 5 \left( \sum_{i=1}^n a_i^2 - \sum_{j=1}^m b_j^2 \right) - 10\sigma(K) - 16\kappa^*(K) \right) \leq g_{X,H}(S).$$

In particular, when  $K$  is a finite connected sum of two bridge knots, one has

$$\frac{1}{16} \left( -36n + 4m + 5 \left( \sum_{i=1}^n a_i^2 - \sum_{j=1}^m b_j^2 \right) - 8\sigma(K) \right) \leq g_{X,H}(S).$$

- (2) From an obstruction via Arf invariant (For more details see Theorem 6.6), we have

$$0 = \text{Arf}(K) + \text{Arf}(X, S).$$

- (3) Since  $H$  is divisible by 2, by Theorem 6.7, one has

$$(59) \quad \frac{1}{2} \left( \left| \sigma(K) + n - m + \sum_{i=1}^n a_i^2 - \sum_{j=1}^m b_j^2 \right| - n - m \right) \leq g_{X,H}^{\text{Top}}(K)$$

**Theorem 4.12** (Example 1.11). *Let  $K_m$  be the  $m$ -fold connected sum of  $T(3, 7)$  for a positive integer  $m$  and  $X$  be  $\#_m \mathbb{C}P^2 \#_m (-\mathbb{C}P^2)$ . Then,*

$$\frac{7}{4}m \leq g_{(X,x)}(K_m),$$

where  $x = (2, \dots, 2) \in H_2(\#_m \mathbb{C}P^2 \#_m (-\mathbb{C}P^2))$ .

*Proof.* We apply (58) to  $X$ ,  $K_m$ , and  $[S] = x$  and obtain

$$\begin{aligned} \frac{1}{32} \left( -36m + 4m - 18(-4m) - 32\left(-\frac{1}{2}m\right) \right) &\leq g_{X,x}(K_m) \\ \frac{7}{4}m &\leq g_{X,x}(K_m). \end{aligned}$$

□

*Remark 4.13.* We again compare Theorem 4.12 with the topological obstruction coming from the knot signature. Under the same assumptions in Theorem 4.12, the inequality (4.13) implies

$$\frac{1}{2} (|-4m + m - m + 0| - 2m) = m \leq g_{X,x}^{\text{Top}}(K_m).$$

4.2.3. *Results on  $\#_N S^2 \times S^2$ .* It is proven in [70] that for any knot  $K$  in  $S^3$  whose Arf invariant  $\text{Arf}(K)$  is zero, there is a positive integer  $N$  such that  $K$  is smoothly H-slice in  $\#_N S^2 \times S^2$ . This result enables us to define knot concordance invariants

$$sn(K) := \min\{N \mid K \text{ is smoothly H-slice in } \#_N S^2 \times S^2\}$$

and

$$sn^{\text{Top}}(K) := \min\{N \mid K \text{ is topologically H-slice in } \#_N S^2 \times S^2\}$$

when  $\text{Arf}(K)$  is zero. In [14],  $sn(K)$  is defined and called *stabilizing number*. As it is asked in [14, Question 1.4], the problem whether there exists a knot  $K$  such that

$$0 < sn^{Top}(K) < sn(K)$$

or not was open. We will give an answer. Our invariant  $\kappa(K)$  can be used to give a lower bound on  $sn(K)$ .

**Corollary 4.14** (Theorem 1.7). *For any knot  $K \subset S^3$  with  $\text{Arf}(K) = 0$ , we have*

$$-\frac{9}{16}\sigma(K) - \kappa(K) \leq sn(K).$$

*Proof of Corollary 4.14.* We just apply Theorem 1.3 to the case  $X = \#_N S^2 \times S^2$ ,  $[S] = 0 \in H_2(S^2 \times S^2)$  and  $g(S) = 0$  and obtain the inequality:

$$-\frac{1}{16}\sigma(K) \leq N + \frac{1}{2}\sigma(K) + \kappa(K).$$

This completes the proof.  $\square$

**Proposition 4.15.** *For an even integer  $m$  and a positive integer  $l$ ,*

- $9ml \leq sn(\#_m T(3, 12l - 1))$ ,
- $9ml \leq sn(\#_m T(3, 12l + 1))$ ,
- $9ml - 5m \leq sn(\#_m T(3, 12l - 7))$ , and
- $9ml - 4m \leq sn(\#_m T(3, 12l - 5))$ .

*Proof.* It is well-known that

$$(60) \quad \sigma(\#_m T(3, 6n - 1)) = -8nm \text{ and } \sigma(\#_m T(3, 6n + 1)) = -8nm$$

for positive integers  $m$  and  $n$ . (For example, see [58].) Putting  $n = 2l$ , it follows from (60) that

$$\sigma(\#_m T(3, 12l - 1)) = \sigma(\#_m T(3, 12l + 1)) = -16ml.$$

So our inequality in Theorem 1.7 and Theorem 1.6 implies that

$$9ml \leq sn(\#_m T(3, 12l - 1)) \text{ and } 9ml \leq sn(\#_m T(3, 12l + 1)).$$

We next put  $n = 2l - 1$ . Then it follows from (60) and Theorem 1.7 that

$$\sigma(\#_m T(3, 12l - 7)) = \sigma(\#_m T(3, 12l - 5)) = -16ml + 8m.$$

So Theorem 1.3 and Theorem 1.6 imply that

$$9ml - \frac{9}{2}m - \frac{1}{2}m = 9ml - 5m \leq sn(\#_m T(3, 12l - 7))$$

and

$$9ml - \frac{9}{2}m + \frac{1}{2}m = 9ml - 4m \leq sn(\#_m T(3, 12l - 5)).$$

$\square$

*Proof of Theorem 1.8.* From Proposition 4.15, we obtain the desired lower bounds on  $sn$ , and in what follows we give bounds on  $sn^{Top}$ . On the other hand, in [14, Theorem 5.15], we have

$$sn^{Top}(K) \leq g_4^{Top}(K)$$



for any knot  $K$  with  $\text{Arf}(K) = 0$ . Since it is proven in [7, Theorem 1] that, for a positive integer  $m$  coprime to 3,

$$g_4^{Top}(T(3, m)) = \left\lceil \frac{2m}{3} \right\rceil,$$

we have

$$g_4^{Top}(T(3, 6n - 1)) = 4n \text{ and } g_4^{Top}(T(3, 6n + 1)) = 4n + 1,$$

where  $g_4^{Top}$  is the topological slice genus. This proves

$$g_4^{Top}(\#_m T(3, 6n - 1)) \leq 4nm, \quad g_4^{Top}(\#_m T(3, 6n + 1)) \leq 4nm + m.$$

In particular,

$$g_4^{Top}(\#_m T(3, 12l - 1)) \leq 8lm, \quad g_4^{Top}(\#_m T(3, 12l + 1)) \leq 8lm + m$$

and

$$g_4^{Top}(\#_m T(3, 12l - 7)) \leq 8lm - 4m, \quad g_4^{Top}(\#_m T(3, 12l - 5)) \leq 8lm - 3m.$$

Also, from Theorem 6.7 and, we have

$$sn^{Top}(\#_m T(3, 12l - 1)) \geq 8lm, \quad sn^{Top}(\#_m T(3, 12l + 1)) \geq 8lm$$

and

$$sn^{Top}(\#_m T(3, 12l - 7)) \geq 8lm - 4m, \quad sn^{Top}(\#_m T(3, 12l - 5)) \geq 8lm - 4m.$$

This completes the proof.  $\square$

*Remark 4.16* (Comparison with other methods). We compare Theorem 1.7 with Maloescu's relative 10/8-inequality, obstructions from Arf invariant and signature function.

- (1) Manolescu's relative 10/8 inequality [54, Theorem 1.1] enable us to prove that for any knot  $K$ ,

$$(61) \quad -\frac{5}{8}\sigma(K) - \kappa(\Sigma(K), \mathfrak{t}) \leq 2sn(K).$$

For a finite connected sum of two bridge knots  $K$  in  $S^3$ , we have

$$(62) \quad -\frac{1}{2}\sigma(K) \leq 2sn(K),$$

where  $\kappa^*(K)$  is the Manolescu's kappa invariant  $\kappa(\Sigma(K), \mathfrak{t})$  for the double branched cover. For more details, see Corollary 6.3.

- (2) From an obstruction via Arf invariant (For more details see Theorem 6.6), we have

$$(63) \quad \text{Arf}(K) \equiv 0 \pmod{2}.$$

- (3) From an obstruction via the Tristram-Levine signature  $\sigma_K(t)$  (For more details see Theorem 6.7), we have

$$\frac{1}{2} \max_m \max_{r \in \{1, \dots, m-1\}} |\sigma_K(e^{2\pi r i m})| \leq sn(K)$$

## 5. NON-EXTENDABLE ACTIONS

We apply Theorem 1.1 to obtain obstructions to an extension of involutions on 3-manifolds to spin 4-manifolds. Regard Brieskorn homology spheres  $\Sigma(p, q, r)$  as a subset of  $\mathbb{C}^3$  following the standard definition. Consider the involution  $\iota$  on  $\Sigma(2, q, r)$  defined by  $\iota(z_1, z_2, z_3) = (-z_1, z_2, z_3)$ . Recent result by Anvari and Hambleton [3, Theorem A] showed that the standard finite cyclic group actions on  $\Sigma(p, q, r)$  does not extend to any contractible smooth 4-manifold bounded by  $\Sigma(p, q, r)$  (if exists), as a smooth involution. In particular,  $\iota$  does not extend to any contractible smooth 4-manifold bounded by  $\Sigma(2, q, r)$  as a smooth involution. (See also recent work by Baraglia–Hekmati [8, Example 7.7].) As a complementary result, we obstruct to extend such  $\iota$  to 4-manifolds typically with  $\sigma \neq 0$ :

**Theorem 5.1.** *Set  $Y = \pm\Sigma(2, q, r)$  and let  $\iota : Y \rightarrow Y$  be the involution defined by  $\iota(z_1, z_2, z_3) = (-z_1, z_2, z_3)$ . Let  $W$  be a compact smooth spin 4-manifold bounded by  $Y$  with  $b_1(W) = 0$ . Then  $\iota$  extends to  $W$  as a homologically trivial diffeomorphism, and we have the followings:*

- (i) *If  $Y = \pm\Sigma(2, 3, 12k+1)$  or  $\pm\Sigma(2, 3, 12k-1)$  for  $k > 0$ , then  $\iota$  cannot extend to  $W$  as a smooth involution so that*

$$-\sigma(W)/16 > b^+(W) - b_\iota^+(W).$$

- (ii) *If  $Y = \Sigma(2, 3, 12k-5)$  for  $k > 0$  or  $Y = -\Sigma(2, 3, 12k+5)$  for  $k \geq 0$ , then  $\iota$  cannot extend to  $W$  as a smooth involution so that*

$$-\sigma(W)/16 > b^+(W) - b_\iota^+(W) - 1/2.$$

- (iii) *If  $Y = -\Sigma(2, 3, 12k-5)$  for  $k > 0$  or  $Y = \Sigma(2, 3, 12k+5)$  for  $k \geq 0$ , then  $\iota$  cannot extend to  $W$  as a smooth involution so that*

$$-\sigma(W)/16 > b^+(W) - b_\iota^+(W) + 1/2.$$

- (iv) *In particular, suppose that  $Y$  and  $W$  satisfy one of the following conditions:*

- *$Y = \pm\Sigma(2, 3, 12k+1)$  or  $\pm\Sigma(2, 3, 12k-1)$ , and  $\sigma(W) \neq 0$ .*
- *$Y = \Sigma(2, 3, 12k-5)$  or  $-\Sigma(2, 3, 12k+5)$ , and  $\sigma(W) \neq 8$ .*
- *$Y = -\Sigma(2, 3, 12k-5)$  or  $\Sigma(2, 3, 12k+5)$ , and  $\sigma(W) \neq -8$ .*

*Then  $\iota$  cannot extend to  $W$  as a homologically trivial smooth involution.*

*Proof.* Since  $\iota$  on  $Y$  is isotopic to the identity as a diffeomorphism of  $Y$  (Lemma 3.32), it is clear that  $\iota$  can be extended to  $W$  as a homologically trivial diffeomorphism.

Consider the case (i):  $Y = \pm\Sigma(2, 3, 12k+1)$  or  $\pm\Sigma(2, 3, 12k-1)$  for  $k > 0$ . As seen in Propositions 3.35 and 3.38, we have  $\kappa(Y, \mathfrak{t}, \iota) = 0$  for the unique spin structure  $\mathfrak{t}$ . Suppose that  $\iota$  extends to  $W$  as a smooth involution so that  $-\sigma(W)/16 > b^+(W) - b_\iota^+(W)$ . Regarding  $W$  as a cobordism from  $S^3$  to  $Y$  and applying Theorem 1.1 to this cobordism, we obtain that

$$-\sigma(W)/16 \leq b^+(W) - b_\iota^+(W),$$

but this is a contradiction.

The cases (ii), (iii) are deduced from a totally same argument of the above proof of the case (i): apply Theorem 1.1 combined with the calculations Propositions 3.36 and 3.37 to  $W$ .

The ‘‘in particular’’ part (iv) follows from these non-extending results (i)-(iii) applied to  $W$  if  $\sigma(W) \leq 0$ , and from the results (i)-(iii) applied to  $-W$  if  $\sigma(W) > 0$ .  $\square$

*Remark 5.2.* One can deduce constraints on locally linear topological involutions by the  $G$ -signature theorem. It is summarized in Corollary 6.8. Note that this topological constraint involves the self-intersection number and genus of the surface  $S$  obtained as the fixed point, also involves the signature of the knot  $K = S \cap Y$ . But the constraint on smooth involutions obtained in Theorem 5.1 is free from these data of fixed point sets.

*Remark 5.3.* As mentioned in Remark 3.33, for all odd involutions on  $\Sigma(2, 3, 6l \pm 1)$  which are isotopic to the identity, the kappa invariants are just the same as that of the above  $\iota$ . Therefore the statement of Theorem 5.1 holds for all odd involutions on  $\Sigma(2, 3, 6l \pm 1)$  which are isotopic to the identity.

We note a consequence of Theorem 5.1 and Remark 5.3 for (relatively) small 4-manifolds. A preliminary result is:

**Proposition 5.4.** *For  $k > 0$ , let  $W$  be a compact spin smooth 4-manifold bounded by one of  $\Sigma(2, 3, 12k + 1)$ ,  $-\Sigma(2, 3, 12k + 1)$ , and  $-\Sigma(2, 3, 12k - 1)$  with the intersection form isomorphic to that of  $K3$ . Let  $f : W \rightarrow W$  be an orientation-preserving diffeomorphism whose restriction to the boundary is isotopic to the identity. Suppose that  $f$  lifts to a spin automorphism. (If  $W$  is simply-connected, this is the case for all  $f$ .) Then  $f$  preserves the orientation of  $H^+(W)$ .*

*Proof.* Set  $Y = \Sigma(2, 3, 13)$ . Let  $f'$  be a diffeomorphism of  $W$  which is isotopic to  $f$  and is the identity on  $Y$ . Since  $f$  lifts to a spin automorphism, so does  $f'$ . Let  $\tilde{f}'$  be a lift of  $f'$  on the spin structure. Then, as the mapping torus of  $W$  by  $(f', \tilde{f}')$ , we obtain a fiber bundle  $E_W \rightarrow S^1$  over a circle with fiber  $W$  equipped with a fiberwise spin structure. Associated to this fiber bundle, we have a real vector bundle  $H^+(E_W) \rightarrow S^1$ , whose fiber is a positive-definite subspace of the real second cohomology group of the fiber. If  $f$  reverses the orientation of  $H^+(W)$ , this implies that  $w_1(H^+(E_W)) \neq 0$ .

Recall the computation of Manolescu's  $\alpha$  invariant of  $Y$  (see [55, Proposition 3.8, Page 172]):

$$\begin{aligned}\alpha(\Sigma(2, 3, 12k + 1)) &= 0, \\ \alpha(-\Sigma(2, 3, 12k + 1)) &= -\gamma(\alpha(\Sigma(2, 3, 12k + 1))) = 0, \\ \alpha(-\Sigma(2, 3, 12k - 1)) &= -\gamma(\alpha(\Sigma(2, 3, 12k - 1))) = 0.\end{aligned}$$

Thus in any case we have  $\alpha(Y) = 0$ . Then it follows from [44, Theorem 1.2] that

$$-\sigma(W)/8 \leq \alpha(Y) = 0,$$

but since  $\sigma(W) = -16$ , this is a contradiction.  $\square$

Bryan proved in [12, Theorem 1.8] that, for a spin odd involution  $\iota$  on a spin rational cohomology  $K3$  surface  $X$ , we have  $b_\iota^+(X) = 1$ . If  $f$  in Proposition 5.4 is an involution  $\iota$ , then Proposition 5.4 implies that  $b_\iota^+(W) = 1$  or  $b_\iota^+(W) = 3$ . But if  $\iota$  is of odd type, there is an additional constraint similar to the result by Bryan:

**Corollary 5.5.** *let  $W$  be a compact spin smooth 4-manifold bounded by one of  $\Sigma(2, 3, 12k + 1)$ ,  $-\Sigma(2, 3, 12k + 1)$ , and  $-\Sigma(2, 3, 12k - 1)$  with the intersection form isomorphic to that of  $K3$ . Let  $\iota : W \rightarrow W$  be a smooth spin involution of odd type whose restriction to the boundary is isotopic to the identity. Then we have  $b_\iota^+(W) = 1$ .*

*Proof.* By Proposition 5.4, we have  $b_\ell^+(W) = 1$  or  $b_\ell^+(W) = 3$ . But the possibility that  $b_\ell^+(W) = 3$  is excluded by Theorem 5.1 and Remark 5.3.  $\square$

## 6. APPENDIX

In this section, we review several known genus bounds related to our result.

**6.1. Manolescu's relative 10/8-inequality.** First, we compare our main result and 10/8-inequality without involutions. A similar discussion for closed 4-manifolds can be founded in [23]. We consider Manolescu's relative 10/8-inequality [54]. The following theorem is just a corollary of usual relative 10/8-inequality combined with double branched covers.

In [54], Manolescu introduced a rational valued spin rational homology cobordism invariant

$$(Y, \mathfrak{s}) \mapsto \kappa(Y, \mathfrak{t}) \in \mathbb{Q}.$$

This enables us to define a knot concordance invariant by

$$K \mapsto \kappa^*(K) := \kappa(\Sigma(K), \mathfrak{t}) \in \mathbb{Q},$$

where  $\mathfrak{t}$  is the unique spin structure on the double branched cover  $\Sigma(K)$ . The following is an application of the relative 10/8-inequality proven in [54] and Lemma 4.2.

**Theorem 6.1** ([54]). *The invariant  $\kappa^*(K)$  is a knot concordance invariant satisfying the following property: For any simply-connected compact 4-manifold  $X$  bounded by  $S^3$  and any oriented compact surface  $S$  bounded by  $K$  such that  $PD(w_2(X)) = [S]/2 \pmod{2}$ , we have*

$$(64) \quad \frac{1}{16} (-4\sigma(X) + 5[S]^2 - 10\sigma(K) - 16\kappa^*(K)) \leq 2b^+(X) + g(S)$$

holds, where  $[S]/2$  denotes the element in  $H_2(S; \mathbb{Z})$  such that  $2([S]/2) = [S]$ .

*Remark 6.2.* When  $K$  is the unknot and  $S$  is an embedded null-homologous disk, (64) implies

$$-\frac{\sigma(X)}{8} \leq b^+(X),$$

which recovers Furuta's original 10/8-inequality ([19]) except for adding 1 or 2 on the left hand side as in the case of Theorem 1.3.

*Proof of Theorem 6.1.* Consider the double branched cover  $\Sigma(S)$  and apply [54, Theorem 1.1 and Remark 4.6] to  $\Sigma(S)$ . Then, we have the following inequality

$$(65) \quad -\frac{\sigma(\Sigma(S))}{8} \leq b^+(\Sigma(S)) + \kappa^*(K).$$

Moreover, using Lemma 4.2, we obtain the equalities

$$\begin{aligned} \sigma(\Sigma(S)) &= 2\sigma(X) - \frac{1}{2}[S]^2 + \sigma(K) \quad \text{and} \\ b^+(\Sigma(S)) &= 2b^+(X) + g(S) - \frac{1}{4}[S]^2 + \frac{1}{2}\sigma(K). \end{aligned}$$

Combining these two equations with (65), we complete the proof.  $\square$

Also, for spin rational homology 3-spheres with positive scalar curvature metrics,

$$\kappa(Y, \mathfrak{t}) = \delta(Y, \mathfrak{t})$$

holds ([54, Subsection 5.1]). In [58, Theorem 1.2], it is proved that for any finite connected sum of two bridge knots,

$$4\delta(\Sigma(K), \mathfrak{t}) = 2d(\Sigma(K), \mathfrak{t}) = \delta(K) = \frac{1}{2}\sigma(K)$$

holds, where  $d(\Sigma(K), \mathfrak{t})$  is the Heegaard Floer correction term. Since

$$\kappa(\Sigma(K), \mathfrak{t}) = \delta(\Sigma(K), \mathfrak{t}),$$

for a finite connected sum of two bridge knots, we have

$$\kappa^*(K) = \kappa(\Sigma(K), \mathfrak{t}) = \frac{1}{8}\sigma(K).$$

As a corollary of Theorem 6.1, we see

**Corollary 6.3.** *Let  $K$  be a finite connected sum of two bridge knots. For any compact simply-connected 4-manifold  $X$  bounded by  $S^3$  and any oriented compact surface  $S$  bounded by  $K$  such that  $[S]$  is divisible by 2 and  $PD(w_2(X)) = [S]/2 \pmod{2}$ , we have*

$$-\frac{1}{4}\sigma(X) + \frac{5}{16}[S]^2 - \frac{1}{2}\sigma(K) \leq 2b^+(X) + g(S).$$

*Remark 6.4.* Note that for a finite connected sum  $K$  of two bridge knots, we have

$$\kappa(K) = \frac{1}{2}\kappa^*(K).$$

It is natural to ask whether there exists a knot  $K$  such that  $\kappa(K) \neq \frac{1}{2}\kappa^*(K)$  or not. For examples, we consider  $K_n := T(3, 12n - 5)$  for positive integer  $n$ . Then, we saw  $\kappa(K_n) = -\frac{1}{2}$ . On the other hand, it is proven in [54, Theorem 1.2] that

$$\kappa^*(K_n) = \kappa(\Sigma(2, 3, 12n - 5)) = 1.$$

**6.2. Topological obstructions.** We review several topological obstructions to sliceness of knots related to our results. For more details, see [56, Section 3].

The following is a relative version of the Rochlin's result [68].

**Theorem 6.5** ([41, 43, 73]). *Let  $X$  be a smooth, closed, connected, oriented 4-manifold. If  $S \subset X^\circ$  is a properly embedded, locally flat characteristic surface with boundary a knot  $K$ , then*

$$\frac{\sigma(X) - [S]^2}{8} = \text{Arf}(K) + \text{Arf}(X, S).$$

When  $X$  is spin and  $[\Sigma] = 0$ , Theorem 6.5 implies the following:

**Theorem 6.6** ([67]). *If a knot  $K$  is topologically H-slice in a spin smooth 4-manifold, then  $\text{Arf}(K) = 0$ .*

Also, we review a genus bound coming from the Tristram-Levine signature. For a knot  $K \subset S^3$  and a value  $t \in S^1$ , the Tristram-Levine signature  $\sigma_K(t)$  is defined as the signature of

$$(1 - t)M + (1 - \bar{t})M^t,$$

where  $M$  is a Seifert matrix for  $K$ .

**Theorem 6.7** ([14, 21, 72]). *Let  $X$  be a topological closed oriented 4-manifold with  $H_1(X; \mathbb{Z}) = 0$ . Let  $S \subset X^\circ$  be a locally flat, properly embedded surface of genus  $g$ , with boundary a knot  $K \subset S^3$ . If the homology class  $[S] \in H_2(X)$  is divisible by 2, then*

$$|\sigma(K) + \sigma(X) + [S]^2| \leq b_2(X) + 2g(S)$$

*holds. Moreover, if the homology class  $[S] \in H_2(X)$  is divisible by a prime power  $m = p^k$ , then*

$$|\sigma_K(e^{2\pi rim}) + \sigma(X) - \frac{2r(m-r)}{m^2}[S]^2| \leq b_2(X) + 2g$$

*holds for  $r \in \{1, \dots, m-1\}$ . In particular, if  $K$  is topologically  $H$ -slice, then*

$$\max_{r \in \{1, \dots, m-1\}} |\sigma_K(e^{2\pi rim}) + \sigma(X)| \leq b_2(X) + 2g(S)$$

*for any prime power  $m = p^k$ .*

**Corollary 6.8.** *Let  $W$  be an oriented topological compact 4-manifold with boundary. Suppose that a locally linear involution  $\iota$  on  $W$  is given and that the fixed point set of  $\iota$  is of codimension 2. Let  $S$  be the fixed point set of  $\iota$  and set  $K = S \cap \partial W$ . Assume that  $S$  is connected. Then we have*

$$\frac{1}{2} |\sigma(W) + [S]^2 + \sigma(K)| \leq \frac{b_2(W)}{2} + g(S).$$

*Proof.* This follows from Theorem 6.7 and the computation of  $b^+$ ,  $b^-$  of the branched cover in Lemma 4.2. (Note that, in the proof of Lemma 4.2, the smoothness and spinness of the manifold and involution were not used to compute these quantities.)  $\square$

## REFERENCES

- [1] *The Knot Atlas*, available at <http://katlas.org/>.
- [2] Antonio Alfieri, Sungkyung Kang, and András I. Stipsicz, *Connected Floer homology of covering involutions*, Math. Ann. **377** (2020), no. 3-4, 1427–1452. MR4126897
- [3] Nima Anvari and Ian Hambleton, *Cyclic branched coverings of Brieskorn spheres bounding acyclic 4-manifolds*, Glasg. Math. J. **63** (2021), no. 2, 400–413. MR4244205
- [4] M. F. Atiyah, *K-theory and reality*, Quart. J. Math. Oxford Ser. (2) **17** (1966), 367–386. MR206940
- [5] M. F. Atiyah and R. Bott, *A Lefschetz fixed point formula for elliptic complexes. II. Applications*, Ann. of Math. (2) **88** (1968), 451–491. MR232406
- [6] M. F. Atiyah, V. K. Patodi, and I. M. Singer, *Spectral asymmetry and Riemannian geometry. I*, Math. Proc. Cambridge Philos. Soc. **77** (1975), 43–69. MR397797
- [7] S. Baader, I. Banfield, and L. Lewark, *Untwisting 3-strand torus knots*, Bull. Lond. Math. Soc. **52** (2020), no. 3, 429–436. MR4171377
- [8] David Baraglia and Pedram Hekmati, *Equivariant Seiberg-Witten-Floer cohomology*, arXiv preprint arXiv:2108.06855 (2021).
- [9] Joshua Batson, *Nonorientable slice genus can be arbitrarily large*, Math. Res. Lett. **21** (2014), no. 3, 423–436. MR3272020
- [10] Michel Boileau and Jean-Pierre Otal, *Scindements de Heegaard et groupe des homéotopies des petites variétés de Seifert*, Invent. Math. **106** (1991), no. 1, 85–107. MR1123375
- [11] Jim Bryan, *Seiberg-Witten theory and  $\mathbf{Z}/2^p$  actions on spin 4-manifolds*, Math. Res. Lett. **5** (1998), no. 1-2, 165–183. MR1617929
- [12] ———, *Seiberg-Witten theory and  $\mathbf{Z}/2^p$  actions on spin 4-manifolds*, Math. Res. Lett. **5** (1998), no. 1-2, 165–183. MR1617929
- [13] Olivier Collin and Brian Steer, *Instanton Floer homology for knots via 3-orbifolds*, J. Differential Geom. **51** (1999), no. 1, 149–202. MR1703606

- [14] Anthony Conway and Matthias Nagel, *Stably slice disks of links*, J. Topol. **13** (2020), no. 3, 1261–1301. MR4125756
- [15] Aliakbar Daemi and Christopher Scaduto, *Equivariant aspects of singular instanton floer homology* (2019), available at [arXiv:1912.08982](#).
- [16] Irving Dai, Matthew Hedden, and Abhishek Mallick, *Corks, involutions, and Heegaard Floer homology* (2020), available at [arXiv:2002.02326](#).
- [17] Michael Freedman and Robion Kirby, *A geometric proof of Rochlin's theorem*, Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 2, 1978, pp. 85–97. MR520525
- [18] Y. Fukumoto and M. Furuta, *Homology 3-spheres bounding acyclic 4-manifolds*, Math. Res. Lett. **7** (2000), no. 5-6, 757–766. MR1809299
- [19] M. Furuta, *Monopole equation and the  $\frac{11}{8}$ -conjecture*, Math. Res. Lett. **8** (2001), no. 3, 279–291. MR1839478
- [20] Patrick M. Gilmer, *Configurations of surfaces in 4-manifolds*, Trans. Amer. Math. Soc. **264** (1981), no. 2, 353–380. MR603768
- [21] ———, *Configurations of surfaces in 4-manifolds*, Trans. Amer. Math. Soc. **264** (1981), no. 2, 353–380. MR603768
- [22] C. McA. Gordon and R. A. Litherland, *On the signature of a link*, Invent. Math. **47** (1978), no. 1, 53–69. MR500905
- [23] M. J. D. Hamilton, *Homology classes of negative square and embedded surfaces in 4-manifolds*, Bull. Lond. Math. Soc. **45** (2013), no. 6, 1221–1226. MR3138489
- [24] Kristen Hendricks, *A rank inequality for the knot Floer homology of double branched covers*, Algebr. Geom. Topol. **12** (2012), no. 4, 2127–2178. MR3020203
- [25] Kristen Hendricks, Tye Lidman, and Robert Lipshitz, *Rank inequalities for the Heegaard Floer homology of branched covers* (2020), available at [arXiv:2007.03023](#).
- [26] Kristen Hendricks and Robert Lipshitz, *Involutive bordered Floer homology*, Trans. Amer. Math. Soc. **372** (2019), no. 1, 389–424. MR3968773
- [27] Kristen Hendricks, Robert Lipshitz, and Sucharit Sarkar, *A flexible construction of equivariant Floer homology and applications*, J. Topol. **9** (2016), no. 4, 1153–1236. MR3620455
- [28] ———, *Corrigendum: A flexible construction of equivariant Floer homology and applications*, J. Topol. **13** (2020), no. 3, 1317–1331. MR4125758
- [29] ———, *A simplicial construction of G-equivariant Floer homology*, Proc. Lond. Math. Soc. (3) **121** (2020), no. 6, 1798–1866. MR4201124
- [30] Kristen Hendricks and Ciprian Manolescu, *Involutive Heegaard Floer homology*, Duke Math. J. **166** (2017), no. 7, 1211–1299. MR3649355
- [31] F. Hirzebruch, *The signature of ramified coverings*, Global Analysis (Papers in Honor of K. Kodaira), 1969, pp. 253–265. MR0258060
- [32] ———, *The signature of ramified coverings*, Global Analysis (Papers in Honor of K. Kodaira), 1969, pp. 253–265. MR0258060
- [33] W. C. Hsiang and R. H. Szczarba, *On embedding surfaces in four-manifolds*, Algebraic topology (Proc. Sympos. Pure Math., Vol. XXII, Univ. Wisconsin, Madison, Wis., 1970), 1971, pp. 97–103. MR0339239
- [34] Nobuo Iida, Anubhav Mukherjee, and Masaki Taniguchi, *An adjunction inequality for the bauer-furuta type invariants, with applications to sliceness and 4-manifold topology* (2021), available at [arXiv:2102.02076](#).
- [35] Stanislav Jabuka, *Concordance invariants from higher order covers*, Topology Appl. **159** (2012), no. 10-11, 2694–2710. MR2923439
- [36] Sungkyung Kang,  *$\mathbb{Z}_2$ -equivariant Heegaard Floer cohomology of knots in  $S^3$  as a strong Heegaard invariant* (2018), available at [arXiv:1810.01919](#).
- [37] ———, *A transverse knot invariant from  $\mathbb{Z}_2$ -equivariant Heegaard Floer cohomology* (2018), available at [arXiv:1802.00351](#).
- [38] Çağrı Karakurt and Tye Lidman, *Rank inequalities for the Heegaard Floer homology of Seifert homology spheres*, Trans. Amer. Math. Soc. **367** (2015), no. 10, 7291–7322. MR3378830
- [39] Yuya Kato, *Nonsmoothable actions of  $\mathbb{Z}^2$  on spin four-manifolds*, [arXiv:1708.08030](#) (2017).
- [40] Tirasan Khandhawit, *A new gauge slice for the relative Bauer-Furuta invariants*, Geom. Topol. **19** (2015), no. 3, 1631–1655. MR3352245

- [41] Robion C. Kirby, *The topology of 4-manifolds*, Lecture Notes in Mathematics, vol. 1374, Springer-Verlag, Berlin, 1989. MR1001966
- [42] Michael Klug and Benjamin Ruppik, *Deep and shallow slice knots in 4-manifolds* (2020), available at [arXiv:2009.03053](https://arxiv.org/abs/2009.03053).
- [43] Michael R. Klug, *A relative version of Rochlin's theorem* (2020), available at [arXiv:2011.12418](https://arxiv.org/abs/2011.12418).
- [44] Hokuto Konno and Masaki Taniguchi, *The groups of diffeomorphisms and homeomorphisms of 4-manifolds with boundary*, [arXiv:2010.00340](https://arxiv.org/abs/2010.00340) (2020).
- [45] D. Kotschick and G. Matic, *Embedded surfaces in four-manifolds, branched covers, and SO(3)-invariants*, Math. Proc. Cambridge Philos. Soc. **117** (1995), no. 2, 275–286. MR1307081
- [46] P. B. Kronheimer and T. S. Mrowka, *Knot homology groups from instantons*, J. Topol. **4** (2011), no. 4, 835–918. MR2860345
- [47] ———, *Gauge theory and Rasmussen's invariant*, J. Topol. **6** (2013), no. 3, 659–674. MR3100886
- [48] Peter Kronheimer and Tomasz Mrowka, *Monopoles and three-manifolds*, New Mathematical Monographs, vol. 10, Cambridge University Press, Cambridge, 2007. MR2388043
- [49] Tim Large, *Equivariant Floer theory and double covers of three-manifolds* (2019), available at [arXiv:1910.12119](https://arxiv.org/abs/1910.12119).
- [50] Tye Lidman and Ciprian Manolescu, *Floer homology and covering spaces*, Geom. Topol. **22** (2018), no. 5, 2817–2838. MR3811772
- [51] Jianfeng Lin, Daniel Ruberman, and Nikolai Saveliev, *On the Frøyshov invariant and monopole Lefschetz number*, [arXiv:1802.07704](https://arxiv.org/abs/1802.07704) (2018).
- [52] ———, *On the monopole Lefschetz number of finite order diffeomorphisms*, [arXiv:2004.05497](https://arxiv.org/abs/2004.05497) (2020).
- [53] Ciprian Manolescu, *Seiberg-Witten-Floer stable homotopy type of three-manifolds with  $b_1 = 0$* , Geom. Topol. **7** (2003), 889–932. MR2026550
- [54] ———, *On the intersection forms of spin four-manifolds with boundary*, Math. Ann. **359** (2014), no. 3-4, 695–728. MR3231012
- [55] ———, *Pin(2)-equivariant Seiberg-Witten Floer homology and the triangulation conjecture*, J. Amer. Math. Soc. **29** (2016), no. 1, 147–176. MR3402697
- [56] Ciprian Manolescu, Marco Marengon, and Lisa Piccirillo, *Relative genus bounds in indefinite four-manifolds* (2020), available at [2012.12270](https://arxiv.org/abs/2012.12270).
- [57] Ciprian Manolescu, Marco Marengon, Sucharit Sarkar, and Michael Willis, *A generalization of Rasmussen's invariant, with applications to surfaces in some four-manifolds* (2019), available at [arXiv:1910.08195](https://arxiv.org/abs/1910.08195).
- [58] Ciprian Manolescu and Brendan Owens, *A concordance invariant from the Floer homology of double branched covers*, Int. Math. Res. Not. IMRN **20** (2007), Art. ID rnm077, 21. MR2363303
- [59] Ciprian Manolescu and Sucharit Sarkar, *A knot Floer stable homotopy type* (2021), available at [arXiv:2108.13566](https://arxiv.org/abs/2108.13566).
- [60] Darryl McCullough and Teruhiko Soma, *The Smale conjecture for Seifert fibered spaces with hyperbolic base orbifold*, J. Differential Geom. **93** (2013), no. 2, 327–353. MR3024309
- [61] Tomasz Mrowka and Yann Rollin, *Legendrian knots and monopoles*, Algebr. Geom. Topol. **6** (2006), 1–69. MR2199446
- [62] Seiji Nagami, *Existence of Spin structures on double branched covering spaces over four-manifolds*, Osaka J. Math. **37** (2000), no. 2, 425–440. MR1772842
- [63] Nobuhiro Nakamura, *Pin<sup>-</sup>(2)-monopole equations and intersection forms with local coefficients of four-manifolds*, Math. Ann. **357** (2013), no. 3, 915–939. MR3118618
- [64] Liviu I. Nicolaescu, *Eta invariants of Dirac operators on circle bundles over Riemann surfaces and virtual dimensions of finite energy Seiberg-Witten moduli spaces*, Israel J. Math. **114** (1999), 61–123. MR1738674
- [65] ———, *Finite energy Seiberg-Witten moduli spaces on 4-manifolds bounding Seifert fibrations*, Comm. Anal. Geom. **8** (2000), no. 5, 1027–1096. MR1846125
- [66] Peter Ozsváth and Zoltán Szabó, *Knot Floer homology and the four-ball genus*, Geom. Topol. **7** (2003), 615–639. MR2026543
- [67] Raymond A. Robertello, *An invariant of knot cobordism*, Comm. Pure Appl. Math. **18** (1965), 543–555. MR182965



- [68] V. A. Rohlin, *Two-dimensional submanifolds of four-dimensional manifolds*, Funkcional. Anal. i Priložen. **5** (1971), no. 1, 48–60. MR0298684
- [69] Kouki Sato, *On eigenvalues of double branched covers*, Proc. Amer. Math. Soc. **147** (2019), no. 6, 2707–2722. MR3951444
- [70] Rob Schneiderman, *Stable concordance of knots in 3-manifolds*, Algebr. Geom. Topol. **10** (2010), no. 1, 373–432. MR2602841
- [71] Matthew Stoffregen,  *$Pin(2)$ -equivariant Seiberg-Witten Floer homology of Seifert fibrations*, Compos. Math. **156** (2020), no. 2, 199–250. MR4044465
- [72] OY Viro, *Placements in codimension 2 and boundary*, Uspekhi Mat. Nauk **30** (1975), no. 1, 231–232.
- [73] Akira Yasuhara, *Connecting lemmas and representing homology classes of simply connected 4-manifolds*, Tokyo J. Math. **19** (1996), no. 1, 245–261. MR1391941

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1 KOMABA,  
MEGURO, TOKYO 153-8914, JAPAN

*Email address:* `konno@ms.u-tokyo.ac.jp`

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1 KOMABA,  
MEGURO, TOKYO 153-8914, JAPAN

*Email address:* `miyazawa@ms.u-tokyo.ac.jp`

2-1 HIROSAWA, WAKO, SAITAMA 351-0198, JAPAN

*Email address:* `masaki.taniguchi@riken.jp`