

COMPOSITION OPERATORS ON SOBOLEV SPACES, CAPACITY AND WEIGHTED SOBOLEV INEQUALITIES

ALEXANDER MENOVSCHIKOV AND ALEXANDER UKHLOV

ABSTRACT. In this paper we give the detailed study of set functions associated with composition operator norms by using capacity representations. On this way we establish connections between mappings which generate bounded composition operators on Sobolev spaces and Q -mappings. As an application weighted Sobolev inequalities with special weights are obtained.

1. INTRODUCTION

Composition operators on Sobolev spaces arise in [29, 36] and represent the significant part of the geometric analysis of Sobolev spaces due its applications to the Sobolev embedding theory, that includes embedding theorems, extension theorems and traces theorems [4, 9, 10, 14, 22] and to the spectral theory of elliptic operators, see, for example, [13, 15]. In the composition operators theory the basic role represent set functions associated with the operator norms which was introduced in [38]. The result of [38] states: *Let $\varphi : \Omega \rightarrow \tilde{\Omega}$ be a homeomorphism, where Ω and $\tilde{\Omega}$ are domains in \mathbb{R}^n . Suppose there exists a collection pairwise disjoint open sets $\Omega_k \subset \Omega$, $\tilde{\Omega}_k := \varphi(\Omega_k)$, $k = 1, 2, \dots$, such that composition operators $\varphi_{\Omega_k}^*(f) := f \circ \varphi$,*

$$\varphi_{\Omega_k}^* : L_p^1(\tilde{\Omega}_k) \rightarrow L_q^1(\Omega_k), \quad 1 \leq q < p \leq \infty,$$

are bounded. Then

$$\|\varphi_{\cup_{k=1}^{\infty} \Omega_k}^*\|_{\frac{1}{q}-\frac{1}{p}} = \sum_{k=1}^{\infty} \|\varphi_{\Omega_k}^*\|_{\frac{1}{q}-\frac{1}{p}}.$$

Hence [38], if $\varphi : \Omega \rightarrow \tilde{\Omega}$ generates the bounded operator

$$\varphi^* : L_p^1(\tilde{\Omega}) \rightarrow L_q^1(\Omega), \quad 1 \leq q < p \leq \infty,$$

by the composition rule $\varphi^*(f) = f \circ \varphi$, then the set function

$$\tilde{\Phi}_{p,q}(\tilde{A}) = \sup_{f \in L_p^1(\tilde{A}) \cap C_0(\tilde{A})} \left(\frac{\|\varphi^*(f) \mid L_q^1(\Omega)\|}{\|f \mid L_p^1(\tilde{A})\|} \right)^{1/q-1/p},$$

is a bounded monotone countably additive set function (an outer measure) defined on open bounded subsets $\tilde{A} \subset \tilde{\Omega}$. The case $p = \infty$ was considered in [45].

In the present work we give characterizations of set functions in capacity terms. This approach allows to define set functions as variation of capacity condensers. The

⁰**Key words and phrases:** Sobolev spaces, Quasiconformal mappings

⁰**2000 Mathematics Subject Classification:** 46E35, 30C65.

⁰The first named author was supported by the grant GAČR 20-19018Y.

geometric theory of composition operators on Sobolev spaces [38, 46] is closely connected with various generalizations of quasiconformal mappings which are defined in capacity (moduli) terms and were studied by many authors, see, for example, [21, 23, 24, 28, 34, 35, 37]. In [32], we consider *BMO*-quasiconformal mappings in connection with composition operators on Hardy–Sobolev spaces. In the present work we consider connections of composition operators on Sobolev spaces with Q -homeomorphisms $\varphi : \Omega \rightarrow \tilde{\Omega}$, $Q \in L_1(\Omega)$.

The notion of Q -mappings was introduced in [26] (see also [27]–[28]) as a generalization of quasiconformal mappings. The main idea behind such generalization is a connection between quasiconformal mappings theory and Beltrami equations. Recall that a homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$ of domains $\Omega, \tilde{\Omega} \subset \mathbb{R}^n$ is called a Q -homeomorphism, with a non-negative measurable function Q , if

$$M(\varphi\Gamma) \leq \int_{\Omega} Q(x) \cdot \rho^n(x) dx$$

for every family Γ of rectifiable paths in Ω and every admissible function ρ for Γ .

Q -mappings with a function Q belongs to the A_n -Muckenhoupt class are mappings which inverse to homeomorphisms generating bounded composition operators on the weighted Sobolev spaces [39] (see, also [42]).

In the present article we prove that in the case $n - 1 = q < p = n$ mappings generated bounded composition operators on Sobolev spaces are Q -homeomorphisms with $Q \in L_1(\Omega)$ and inverse, Q -homeomorphisms with $Q \in L_1(\Omega)$ generate bounded composition operators

$$\varphi^* : L_n^1(\tilde{\Omega}) \rightarrow L_1^1(\Omega).$$

In the case $n = 2$ we have: *Let Ω and $\tilde{\Omega}$ are bounded domains in \mathbb{R}^2 . Then $\varphi : \Omega \rightarrow \tilde{\Omega}$ is a Q -homeomorphism with $Q \in L_1(\Omega)$, if and only if φ generates by the composition rule $\varphi^*(f) = f \circ \varphi$ a bounded operator*

$$\varphi^* : L_2^1(\tilde{\Omega}) \rightarrow L_1^1(\Omega).$$

In particular, as a consequence, on the base of method suggested in [4, 13], we give applications of Q -homeomorphisms to the weighted Sobolev type embedding theorems: *Let $\tilde{\Omega} \subset \mathbb{R}^2$ be a simply connected bounded domain. Then the weighted Sobolev inequality*

$$\inf_{c \in \mathbb{R}} \left(\int_{\tilde{\Omega}} |f(x) - c|^2 w(x) dx \right)^{\frac{1}{2}} \leq B_{2,2}^w(\tilde{\Omega}) \left(\int_{\tilde{\Omega}} |\nabla f(x)|^2 dx \right)^{\frac{1}{2}}$$

holds for any function $f \in W_2^1(\tilde{\Omega})$ with the weight $w(x) = J_{\varphi^{-1}}(x)$, where $J_{\varphi^{-1}}(x)$ is a volume derivative of the mapping inverse to Q -homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$, $Q \in L_1(\Omega)$, of a bounded Lipschitz domain $\Omega \subset \mathbb{R}^2$ onto $\tilde{\Omega} \subset \mathbb{R}^2$.

Note, that weighted Sobolev inequalities with quasiconformal weights were considered, for example, in [20, 12].

Composition operators on Sobolev spaces have applications to the spectral theory of elliptic operators, see, for example, [13, 15, 6]. In some cases the composition operators method allows to obtain better estimates than the classical L. E. Payne and H. F. Weinberger estimates in convex domains [33]. In the last section of the

present work, as an example of applications Q -mappings, we give spectral estimates of the Neumann-Laplace operator in Hölder singular domains.

This paper is organized as follows. Section 2 contains definitions and preliminary results. In Section 3, we give the description of set functions in capacity terms. In Section 4, we prove a connection between composition operators on Sobolev spaces and Q -homeomorphisms. Section 5 contains the definition of a new p -capacity metric of the hyperbolic type and measure properties of weak (p, q) -quasiconformal mappings. Section 6 is devoted to the weighted Sobolev inequalities and in the final section, Section 7, we consider applications of Q -homeomorphisms to the spectral theory of elliptic operators.

2. SOBOLEV SPACES AND COMPOSITION OPERATORS

2.1. Sobolev spaces. Let us recall the basic notions of the Sobolev spaces and the change of variable formula.

Let Ω be an open subset of \mathbb{R}^n . The Sobolev space $W_p^1(\Omega)$, $1 \leq p \leq \infty$, is defined [30] as a Banach space of locally integrable weakly differentiable functions $f : \Omega \rightarrow \mathbb{R}$ equipped with the following norm:

$$\|f\|_{W_p^1(\Omega)} = \|f\|_{L_p(\Omega)} + \|\nabla f\|_{L_p(\Omega)},$$

where ∇f is the weak gradient of the function f , i. e. $\nabla f = (\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})$.

The homogeneous seminormed Sobolev space $L_p^1(\Omega)$, $1 \leq p \leq \infty$, is defined as a space of locally integrable weakly differentiable functions $f : \Omega \rightarrow \mathbb{R}$ equipped with the following seminorm:

$$\|f\|_{L_p^1(\Omega)} = \|\nabla f\|_{L_p(\Omega)}.$$

In the Sobolev spaces theory, a crucial role is played by capacity as an outer measure associated with Sobolev spaces [30]. In accordance to this approach, elements of Sobolev spaces $W_p^1(\Omega)$ are equivalence classes up to a set of p -capacity zero [31]. The exact definition of capacity will be given below.

The mapping $\varphi : \Omega \rightarrow \mathbb{R}^n$ belongs to the Sobolev space $W_{1,\text{loc}}^1(\Omega, \mathbb{R}^n)$, if its coordinate functions belongs to $W_{1,\text{loc}}^1(\Omega)$. In this case, the formal Jacobi matrix $D\varphi(x)$ and its determinant (Jacobian) $J(x, \varphi)$ are well defined at almost all points $x \in \Omega$. The norm $|D\varphi(x)|$ is the operator norm of $D\varphi(x)$,

Let us recall the change of variable formula in the Lebesgue integral [1, 17]. Suppose a homeomorphism $\varphi : \Omega \rightarrow \mathbb{R}^n$ be such that there exists a collection of closed sets $A_k \subset A_{k+1} \subset \Omega$, $k = 1, 2, \dots$, for which restrictions $\varphi|_{A_k}$ are Lipschitz mappings on the sets A_k and

$$\left| \Omega \setminus \sum_{k=1}^{\infty} A_k \right| = 0.$$

Then there exists a measurable set $S \subset \Omega$, $|S| = 0$, such that the homeomorphism $\varphi : \Omega \setminus S \rightarrow \mathbb{R}^n$ has the Luzin N -property (the image of a set of measure zero has measure zero) and the change of variable formula

$$(2.1) \quad \int_E f \circ \varphi(x) |J(x, \varphi)| dx = \int_{\mathbb{R}^n \setminus \varphi(S)} f(y) dy$$

holds for every measurable set $E \subset \Omega$ and every non-negative measurable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$.

Note, that Sobolev homeomorphisms of the class $W_{1,\text{loc}}^1(\Omega)$ satisfy the conditions of the change of variable formula [17] and, therefore, for Sobolev homeomorphisms the change of variable formula (2.1) holds.

If the mapping φ possesses the Luzin N -property, then $|\varphi(S)| = 0$ and the second integral can be rewritten as the integral on \mathbb{R}^n . Note, that Sobolev homeomorphisms of the class $L_p^1(\Omega)$, $p \geq n$, possess the Luzin N -property [43].

The following formula can be found, for example, in [18]. Let $\varphi : \Omega \rightarrow \tilde{\Omega}$ be a homeomorphism. We define a volume derivative of the inverse mapping as

$$J_{\varphi^{-1}}(y) := \lim_{r \rightarrow 0} \frac{|\varphi^{-1}(B(y, r))|}{|B(y, r)|}.$$

If $\varphi : \Omega \rightarrow \tilde{\Omega}$ possesses the Luzin N^{-1} -property (the preimage of a set of measure zero has measure zero), then the following change of variables formula holds:

$$(2.2) \quad \int_{\Omega} f \circ \varphi(x) dx = \int_{\tilde{\Omega}} f(y) J_{\varphi^{-1}}(y) dy.$$

2.2. Composition operators and set functions. Let Ω and $\tilde{\Omega}$ be domains in the Euclidean space \mathbb{R}^n . Then a homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$ generates a bounded composition operator

$$\varphi^* : L_p^1(\tilde{\Omega}) \rightarrow L_q^1(\Omega), \quad 1 \leq q \leq p \leq \infty,$$

by the composition rule $\varphi^*(f) = f \circ \varphi$, if for any function $f \in L_p^1(\tilde{\Omega})$, the composition $\varphi^*(f) \in L_q^1(\Omega)$ is defined quasi-everywhere in Ω and there exists a constant $K_{p,q}(\varphi; \Omega) < \infty$ such that

$$\|\varphi^*(f) \mid L_q^1(\Omega)\| \leq K_{p,q}(\varphi; \Omega) \|f \mid L_p^1(\tilde{\Omega})\|.$$

Recall that the p -dilatation [3] of a Sobolev mapping $\varphi : \Omega \rightarrow \tilde{\Omega}$ at a point $x \in \Omega$ is defined as

$$K_p(x) = \inf \{k(x) : |D\varphi(x)| \leq k(x) |J(x, \varphi)|^{\frac{1}{p}}\}.$$

The following theorem gives the characterization of composition operators in terms of integral characteristics of mappings of finite distortion. Recall that a weakly differentiable mapping $\varphi : \Omega \rightarrow \mathbb{R}^n$ is the mapping of finite distortion if $D\varphi(x) = 0$ for almost all x from $Z = \{x \in \Omega : J(x, \varphi) = 0\}$ [43].

Theorem 2.1. *A homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$ between two domains Ω and $\tilde{\Omega}$ generates a bounded composition operator*

$$\varphi^* : L_p^1(\tilde{\Omega}) \rightarrow L_q^1(\Omega), \quad 1 \leq q \leq p \leq \infty,$$

if and only if $\varphi \in W_{q,\text{loc}}^1(\Omega)$, has finite distortion, and

$$K_{p,q}(\varphi; \Omega) := \|K_p \mid L_\kappa(\Omega)\| < \infty, \quad 1/q - 1/p = 1/\kappa \quad (\kappa = \infty, \text{ if } p = q).$$

The norm of the operator φ^ is estimated as $\|\varphi^*\| \leq K_{p,q}(\varphi; \Omega)$.*

This theorem in the case $p = q$ was proved in [41] (see, also [5]), the case $q < p$ was proved in [38] and the limit case $p = \infty$ was considered in [11].

Let us recall, that homeomorphisms $\varphi : \Omega \rightarrow \tilde{\Omega}$ which satisfy conditions of Theorem 2.1 are called as weak (p, q) -quasiconformal mappings [5, 44].

In the case of weak (p, q) -quasiconformal mappings, the following composition duality theorem holds [38] (the detailed proof can be found in [16]):

Theorem 2.2. *Let a homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$ between two domains Ω and $\tilde{\Omega}$ generates a bounded composition operator*

$$\varphi^* : L_p^1(\tilde{\Omega}) \rightarrow L_q^1(\Omega), \quad n-1 < q \leq p < \infty,$$

and has finite distortion in the case $p = \infty$, then the inverse mapping $\varphi^{-1} : \tilde{\Omega} \rightarrow \Omega$ generates a bounded composition operator

$$(\varphi^{-1})^* : L_{q'}^1(\Omega) \rightarrow L_{p'}^1(\tilde{\Omega}),$$

where $p' = p/(p-n+1)$, $q' = q/(q-n+1)$. In the case $n = 2$ we have $1 \leq q \leq p < \infty$, the inverse assertion is also correct and $p'' = p$.

The geometric theory of composition operators on Sobolev spaces is based on the measure property of composition operators introduced in [38] (see, also [45]). This measure property use the additive set function that forms the basis of our results.

Let us define a set function $\tilde{\Phi}_{p,q}(\tilde{A})$, defined on open bounded subsets $\tilde{A} \subset \tilde{\Omega}$ and associated with a composition operator $\varphi^* : L_p^1(\tilde{\Omega}) \rightarrow L_q^1(\Omega)$, $1 \leq q < p < \infty$,

$$(2.3) \quad \tilde{\Phi}_{p,q}(\tilde{A}) = \sup_{f \in L_p^1(\tilde{A}) \cap C_0(\tilde{A})} \left(\frac{\|\varphi^*(f) \mid L_q^1(\Omega)\|}{\|f \mid L_p^1(\tilde{A})\|} \right)^\kappa, \quad 1/\kappa = 1/q - 1/p.$$

Theorem 2.3. *Let a homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$ between two domains Ω and $\tilde{\Omega}$ generates a bounded composition operator*

$$\varphi^* : L_p^1(\tilde{\Omega}) \rightarrow L_q^1(\Omega), \quad 1 \leq q < p \leq \infty.$$

Then function $\tilde{\Phi}_{p,q}(\tilde{A})$, defined by (2.3), is a bounded monotone countably additive set function defined on open bounded subsets $\tilde{A} \subset \tilde{\Omega}$.

3. CAPACITARY DESCRIPTION AND SET FUNCTIONS

In this section we study capacity properties of the set functions associated with norms of composition operators. First of all, we recall the notion of a variational p -capacity [8, 43].

A condenser in a domain $\Omega \subset \mathbb{R}^n$ is the pair (F_0, F_1) of connected disjoint closed relatively to Ω sets $F_0, F_1 \subset \Omega$. Then a continuous function $f \in L_p^1(\Omega)$ is called an admissible function for the condenser (F_0, F_1) , if the set $F_i \cap \Omega$ is contained in some connected component of the set $\text{Int}\{x \mid f(x) = i\}$, $i = 0, 1$.

The p -capacity of the condenser (F_0, F_1) relatively to domain Ω is the following quantity:

$$\text{cap}_p(F_0, F_1; \Omega) = \inf \|f \mid L_p^1(\Omega)\|^p.$$

Here the greatest lower bound is taken over all functions admissible for the condenser $(F_0, F_1) \subset \Omega$. If the condenser has no admissible functions we put the capacity equal to infinity.

The next two theorems provide a capacity description of composition operators.

Theorem 3.1. [5] *Let $1 < p < \infty$. A homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$ generates a bounded composition operator*

$$\varphi^* : L_p^1(\tilde{\Omega}) \rightarrow L_p^1(\Omega)$$

if and only if for every condenser $(F_0, F_1) \subset \tilde{\Omega}$ the inequality

$$\text{cap}_p^{1/p}(\varphi^{-1}(F_0), \varphi^{-1}(F_1); \Omega) \leq K_{p,p}(\varphi; \Omega) \text{cap}_p^{1/p}(F_0, F_1; \tilde{\Omega})$$

holds.

Theorem 3.2. [38] *Let $1 < q < p < \infty$. A homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$ generates a bounded composition operator*

$$\varphi^* : L_p^1(\tilde{\Omega}) \rightarrow L_q^1(\Omega)$$

if and only if there exists a bounded monotone countable-additive set function $\tilde{\Phi}_{p,q}$ defined on open subsets of $\tilde{\Omega}$ such that for every condenser $(F_0, F_1) \subset \tilde{\Omega}$ the inequality

$$\text{cap}_q^{1/q}(\varphi^{-1}(F_0), \varphi^{-1}(F_1); \Omega) \leq \tilde{\Phi}_{p,q}(\tilde{\Omega} \setminus F_0)^{\frac{p-q}{pq}} \text{cap}_p^{1/p}(F_0, F_1; \tilde{\Omega})$$

holds.

Further, one consider a distortion property of weak (p, q) -quasiconformal mappings for ring condensers. The ring condenser R (see, for example, [3, 25]) in a domain $\Omega \subset \mathbb{R}^n$ is the pair $R = (F; G)$ of sets $F \subset G \subset \Omega$, where F is a connected compact set and G is a connected open set. The p -capacity of the ring $R = (F; G)$ is defined by

$$\text{cap}_p(F; G) = \inf \|f\|_{L_p^1(\Omega)}^p,$$

where the greatest lower bound is taken over all continuous functions $f \in L_p^1(\Omega)$ such that $f \geq 1$ on F and $f = 0$ on $\Omega \setminus G$.

Next we show that in the case $n - 1 < q \leq p < \infty$ we can consider ring condensers only. Let us recall the geometric characterization of homeomorphisms which generate composition operators on Sobolev spaces. The homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$ is called (strong) (p, q) -quasiconformal, $1 < q \leq p < \infty$, if there exists a bounded monotone countable-additive absolutely continuous function Θ defined on open subsets of Ω such that

$$\limsup_{r \rightarrow 0} \frac{L_\varphi^p(x, r)r^{n-p}}{|\varphi(B(x, 2r))|} \bigg/ \left(\frac{\Theta(B(x, r))}{|B(x, r)|} \right)^{\frac{n-q}{q}} \leq H < \infty,$$

where $L_\varphi(x, r) = \max_{|x-y|=r} |\varphi(x) - \varphi(y)|$, for all points $x \in \Omega$.

This definition was introduced in [5] in the case $p = q$ and in [44] in the case $q < p$ for the case of Carnot group \mathbb{G} . Recall that the Euclidean space \mathbb{R}^n , $n \geq 2$, is an example of the abelian Carnot group \mathbb{G} .

Theorem 3.3. *Let $n - 1 < q \leq p < \infty$ ($1 \leq q \leq p < \infty$ if $n = 2$) and $\varphi : \Omega \rightarrow \tilde{\Omega}$ be a homeomorphism. Then the following conditions are equivalent:*

- (1) *The composition operator*

$$\varphi^* : L_p^1(\tilde{\Omega}) \rightarrow L_q^1(\Omega),$$

defined by the rule $\varphi^(f) = f \circ \varphi$ is bounded.*

- (2) *The homeomorphism $\varphi \in W_{q,\text{loc}}^1(\Omega)$, has finite distortion, and*

$$K_{p,q}(\varphi; \Omega) := \|K_p \mid L_\kappa(\Omega)\| < \infty, \quad 1/q - 1/p = 1/\kappa \quad (\kappa = \infty, \text{ if } p = q).$$

- (3) *There exists a bounded monotone countable-additive set function $\tilde{\Phi}_{p,q}$ defined on open subsets of $\tilde{\Omega}$ such that for every ring condenser $(F, \tilde{A}) \subset \tilde{\Omega}$ the inequality*

$$\text{cap}_q^{1/q}(\varphi^{-1}(F), \varphi^{-1}(\tilde{A})) \leq C_0 \tilde{\Phi}_{p,q}(\tilde{A})^{\frac{p-q}{pq}} \text{cap}_p^{1/p}(F, \tilde{A})$$

holds.

- (4) *There exists a bounded monotone countable-additive absolutely continuous function Θ defined on open subsets of Ω such that*

$$\limsup_{r \rightarrow 0} \frac{L_\varphi^p(x, r) r^{n-p}}{|\varphi(B(x, 2r))|} \bigg/ \left(\frac{\Theta(B(x, r))}{|B(x, r)|} \right)^{\frac{p-q}{q}} \leq H < \infty,$$

for all points $x \in \Omega$.

Proof. This theorem was proved, but not formulated in [5] in the case $p = q$ and in [44] in the case $q < p$. In the proof of corresponding assertions in [5, 44] were used ring condensers only and Teichmüller type capacity estimates. \square

To give characterizations of the set function $\tilde{\Phi}_{p,q}$ in capacity terms, we have to define another set function $\tilde{\Psi}$ and consider its variation. Let $1 < q < p < \infty$ and $\varphi : \Omega \rightarrow \tilde{\Omega}$ be a homeomorphism. We define the set function on open bounded subsets $\tilde{A} \subset \tilde{\Omega}$ by the rule

$$\tilde{\Psi}_{p,q}(\tilde{A}) = \sup_{(F;G) \subset \tilde{A}} \left(\frac{\text{cap}_q^{\frac{1}{q}}(\varphi^{-1}(F); \varphi^{-1}(G))}{\text{cap}_p^{\frac{1}{p}}(F; G)} \right)^{\frac{pq}{p-q}},$$

where the supremum is taken over all ring condensers $(F; G) \subset \tilde{A}$ such that $\text{cap}_p(F; G) \neq 0$.

For function $\tilde{\Psi}_{p,q}$, we consider its variation. Variation of a set function is a quasiadditive set function

$$V(\tilde{\Psi}_{p,q}, \tilde{\Omega}) = \sup_{\{\tilde{A}_k\}} \sum_{k=1}^{\infty} \tilde{\Psi}_{p,q}(\tilde{A}_k),$$

where the supremum is taken over all families of disjoint sets $\{\tilde{A}_k\}_{k \in \mathbb{N}}$, $\tilde{A}_k \subset \tilde{\Omega}$.

Proposition 3.4. *Let $n - 1 < q < p < \infty$, then the set function $\tilde{\Psi}_{p,q}$ is positive and non-decreasing.*

Proof. Let $\tilde{A}_1 \subset \tilde{A}_2 \subset \tilde{\Omega}$. Then

$$\begin{aligned} \tilde{\Psi}_{p,q}(\tilde{A}_1) &= \sup_{(F_1; G_1) \subset \tilde{A}_1} \left(\frac{\text{cap}_q^{\frac{1}{q}}(\varphi^{-1}(F_1); \varphi^{-1}(G_1))}{\text{cap}_p^{\frac{1}{p}}(F_1; G_1)} \right)^{\frac{pq}{p-q}} \\ &= \sup_{(F_1; G_1) \subset \tilde{A}_2} \left(\frac{\text{cap}_q^{\frac{1}{q}}(\varphi^{-1}(F_1); \varphi^{-1}(G_1))}{\text{cap}_p^{\frac{1}{p}}(F_1; G_1)} \right)^{\frac{pq}{p-q}} \\ &\leq \sup_{(F_2; G_2) \subset \tilde{A}_2} \left(\frac{\text{cap}_q^{\frac{1}{q}}(\varphi^{-1}(F_2); \varphi^{-1}(G_2))}{\text{cap}_p^{\frac{1}{p}}(F_2; G_2)} \right)^{\frac{pq}{p-q}} = \tilde{\Psi}_{p,q}(\tilde{A}_2), \end{aligned}$$

because any condenser $(F_1; G_1) \subset \tilde{A}_2$ coincides with some condenser $(F_2; G_2) \subset \tilde{A}_2$.

Let us proof, that for arbitrary open set $\tilde{A} \subset \mathbb{R}^n$ the function $\tilde{\Psi}_{p,q}(\tilde{A})$ is positive. Fix a point $x \in \tilde{A}$ and choose $r > 0$ such that ring condenser $(\overline{B}(x, r); B(x, 2r))$ is a subset of \tilde{A} . Then, by [8],

$$\text{cap}_p(\overline{B}(x, r); B(x, 2r)) \geq c_1 > 0.$$

The preimage of $\overline{B}(x, r)$ is a continuum set. Therefore, because in the case $q > n - 1$ the capacity of a continuum set is positive [8], we have

$$\text{cap}_p(\varphi^{-1}(\overline{B}(x, r)); \varphi^{-1}(B(x, 2r))) \geq c_2 > 0.$$

Hence, the set function $\tilde{\Psi}_{p,q}(\tilde{A}) > 0$ for an arbitrary open set $\tilde{A} \subset \mathbb{R}^n$. \square

Remark 3.5. *In the case $n = 2$ Proposition 3.4 is correct for all $1 \leq q < p < \infty$.*

The following statement establishes a correspondence between set functions $\tilde{\Phi}_{p,q}$ and $\tilde{\Psi}_{p,q}$.

Theorem 3.6. *Fix a constant $M < \infty$ and let $\varphi : \Omega \rightarrow \tilde{\Omega}$ be a homeomorphism. Then the following conditions are equivalent:*

- (1) *there exists a bounded monotone countable-quasiadditive set function $\tilde{\Phi}_{p,q}$ defined on open subsets of $\tilde{\Omega}$ such that for every ring condenser $(F; \tilde{A}) \subset \tilde{\Omega}$ the inequality*

$$\text{cap}_q^{1/q}(\varphi^{-1}(F); \varphi^{-1}(\tilde{A})) \leq \tilde{\Phi}_{p,q}(\tilde{A})^{\frac{p-q}{pq}} \text{cap}_p^{1/p}(F; \tilde{A})$$

holds and $\tilde{\Phi}_{p,q}(\tilde{\Omega}) \leq M$;

- (2) *function $\tilde{\Psi}_{p,q}$ has a bounded variation $V(\tilde{\Psi}_{p,q}, \tilde{\Omega}) \leq M$;*

- (3) *for every family of disjoint ring condensers $\{(F_k; \tilde{A}_k)\}_{k \in \mathbb{N}}$ in $\tilde{\Omega}$ with $\text{cap}_p^{1/p}(F_k; \tilde{A}_k) \neq 0$, the following estimate is true*

$$(3.1) \quad \sum_{k=1}^{\infty} \left(\frac{\text{cap}_q^{1/q}(\varphi^{-1}(F_k); \varphi^{-1}(\tilde{A}_k))}{\text{cap}_p^{1/p}(F_k; \tilde{A}_k)} \right)^{\frac{pq}{p-q}} \leq M.$$

Proof. First we show, that the second and the third condition are equivalent. If $V(\tilde{\Psi}_{p,q}, \tilde{\Omega}) \leq M$ then, by virtue of definitions of $\tilde{\Psi}$ and $V(\tilde{\Psi}_{p,q})$, we have

$$\sup_{\{\tilde{A}_k\}} \sum_{k=1}^{\infty} \sup_{(F_k; G_k) \subset \tilde{A}_k} \left(\frac{\text{cap}_q^{1/q}(\varphi^{-1}(F_k); \varphi^{-1}(G_k))}{\text{cap}_p^{1/p}(F_k; G_k)} \right)^{\frac{pq}{p-q}} \leq M,$$

where the first supremum is taken over all families of disjoint sets \tilde{A}_k , and the second is taken over all ring condensers in \tilde{A}_k with non-zero capacity. Hence, for an arbitrary family $\{\tilde{A}_k\}_{k \in \mathbb{N}}$ and condensers $(F_k; \tilde{A}_k)$, the sum (3.1) is bounded by the same constant M .

Now we assume that condition 3 is fulfill. For an arbitrary family of disjoint sets $\{(F_k; \tilde{A}_k)\}_{k \in \mathbb{N}}$ in $\tilde{\Omega}$ and an arbitrary $\varepsilon > 0$, we choose condensers $(F_k; G_k)$ such that

$$\tilde{\Psi}_{p,q}(\tilde{A}_k) < \left(\frac{\text{cap}_q^{1/q}(\varphi^{-1}(F_k); \varphi^{-1}(G_k))}{\text{cap}_p^{1/p}(F_k; G_k)} \right)^{\frac{pq}{p-q}} + \frac{\varepsilon}{2^k}.$$

Then $\sum_{k=1}^{\infty} \tilde{\Psi}_{p,q}(\tilde{A}_k) < M + \varepsilon$ and, therefore, $V(\tilde{\Psi}_{p,q}, \tilde{\Omega}) \leq M$ and condition 2 is fulfill.

Next we show that condition 1 and 2 are equivalent. If condition 1 is true, then, due to the capacity inequality for $\tilde{\Phi}_{p,q}$, $\tilde{\Psi}_{p,q}(\tilde{A}) \leq \sup \tilde{\Phi}_{p,q}(G)$, where the supremum is taken over all $G \subset \tilde{A}$. Hence, by monotonicity of $\tilde{\Phi}_{p,q}$, $\tilde{\Psi}_{p,q}(\tilde{A}) \leq \tilde{\Phi}_{p,q}(\tilde{A})$ for every $\tilde{A} \subset \tilde{\Omega}$. Further, by the definition of the variation,

$$V(\tilde{\Psi}_{p,q}, \tilde{\Omega}) = \sup_{\{\tilde{A}_k\}} \sum_{k=1}^{\infty} \tilde{\Psi}_{p,q}(\tilde{A}_k) \leq \sup_{\{\tilde{A}_k\}} \sum_{k=1}^{\infty} \tilde{\Phi}_{p,q}(\tilde{A}_k).$$

Function $\tilde{\Phi}_{p,q}$ is countable-quasiadditive and monotone, which implies

$$V(\tilde{\Psi}_{p,q}, \tilde{\Omega}) \leq \sup \tilde{\Phi}_{p,q}(\bigcup_{k=1}^{\infty} \tilde{A}_k) \leq \tilde{\Phi}_{p,q}(\tilde{\Omega})$$

and condition 2 holds.

If condition 2 holds, then for an arbitrary condenser (F, \tilde{A}) with $\text{cap}_p(F, \tilde{A}) \neq 0$ by the definition of $\tilde{\Psi}_{p,q}$ and $V(\tilde{\Psi}_{p,q})$ we obtain the inequality

$$\text{cap}_q^{1/q}(\varphi^{-1}(F); \varphi^{-1}(\tilde{A})) \leq V(\tilde{\Psi}_{p,q}, \tilde{A}) \text{cap}_p^{1/p}(F; \tilde{A}).$$

Function $V(\tilde{\Psi}_{p,q})$ is quasiadditive and for finishing the proof we have to show that the above inequality holds also for the case $\text{cap}_p(F; \tilde{A}) = 0$. For an arbitrary condenser $(F; \tilde{A})$ with $\text{cap}_p(F; \tilde{A}) = 0$, let us consider a decreasing sequence of compact sets $\{F_k\}_{k=1}^{\infty}$, $F_k \subset \tilde{A}$, $F = \bigcap_{k=1}^{\infty} F_k$, with $\text{cap}_p(F_k; \tilde{A}) > 0$ (for example, $F_k = \{x : \text{dist}(x, F) \leq 1/k\}$). For such condensers

$$\text{cap}_q^{1/q}(\varphi^{-1}(F_k); \varphi^{-1}(\tilde{A})) \leq V(\tilde{\Psi}_{p,q}, \tilde{A}) \text{cap}_p^{1/p}(F_k; \tilde{A}).$$

Due to the continuity of the capacity we can pass to the limit and obtain the above inequality for condensers with zero capacity. \square

Note, that in [23] the similar statement was proven in the particular case $p = n$.

4. COMPOSITION OPERATORS AND Q -HOMEOMORPHISMS

4.1. Modulus and capacity. The theory of Q -quasiconformal mappings has been extensively developed in recent decades, see, for example, [28]. Let us give the basic definitions.

The linear integral is denoted by

$$\int_{\gamma} \rho \, ds = \sup_{\gamma'} \int \rho \, ds = \sup \int_0^{l(\gamma')} \rho(\gamma'(s)) \, ds$$

where the supremum is taken over all closed parts γ' of γ and $l(\gamma')$ is the length of γ' . Let Γ be a family of curves in \mathbb{R}^n . Denote by $\text{adm}(\Gamma)$ the set of Borel functions (admissible functions) $\rho : \mathbb{R}^n \rightarrow [0, \infty]$ such that the inequality

$$\int_{\gamma} \rho \, ds \geq 1$$

holds for locally rectifiable curves $\gamma \in \Gamma$.

Let Γ be a family of curves in $\overline{\mathbb{R}^n}$, where $\overline{\mathbb{R}^n}$ is a one point compactification of the Euclidean space \mathbb{R}^n . The quantity

$$M(\Gamma) = \inf_{\mathbb{R}^n} \int \rho^n dx$$

is called the (conformal) module of the family of curves Γ [28]. The infimum is taken over all admissible functions $\rho \in \text{adm}(\Gamma)$.

Let Ω be a bounded domain in \mathbb{R}^n and F_0, F_1 be disjoint non-empty compact sets in the closure of Ω . Let $M(\Gamma(F_0, F_1; \Omega))$ stand for the module of a family of curves which connect F_0 and F_1 in Ω . Then [28]

$$(4.1) \quad M(\Gamma(F_0, F_1; \Omega)) = \text{cap}_n(F_0, F_1; \Omega),$$

where $\text{cap}_n(F_0, F_1; \Omega)$ is a conformal capacity of the condenser $(F_0, F_1; \Omega)$ [30].

Recall that a homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$ of domains $\Omega, \tilde{\Omega} \subset \mathbb{R}^n$ is called a Q -homeomorphism [28] with a non-negative measurable function Q if

$$M(\varphi\Gamma) \leq \int_{\Omega} Q(x) \cdot \rho^n(x) dx$$

for every family Γ of rectifiable paths in Ω and every admissible function ρ for Γ .

4.2. Mappings of finite distortion. Let us define two distortion functions for Sobolev mappings of finite distortion $\varphi : \Omega \rightarrow \tilde{\Omega}$.

The outer (quasi)conformal dilatation

$$K^O(x, \varphi) = \begin{cases} \frac{|D\varphi(x)|^n}{|J(x, \varphi)|}, & J(x, \varphi) \neq 0, \\ 0, & J(x, \varphi) = 0. \end{cases}$$

The inner (quasi)conformal dilatation

$$K^I(x, \varphi) = \begin{cases} \frac{|J(x, \varphi)|}{l(D\varphi(x))^n}, & J(x, \varphi) \neq 0, \\ 0, & J(x, \varphi) = 0. \end{cases}$$

Note that $K^I(x) \leq (K^O(x))^{n-1}$ and $K^O(x) \leq (K^I(x))^{n-1}$.

In this section we consider Q -homeomorphisms with $Q \in L_s(\Omega)$, $s \geq 1$. In [28], it was proven that Q -homeomorphisms $\varphi : \Omega \rightarrow \tilde{\Omega}$ with $Q \in L_{1,\text{loc}}(\Omega)$ belong to the Sobolev space $W_{1,\text{loc}}^1(\Omega)$, differentiable a.e., has a finite distortion and possesses the Luzin- N^{-1} property.

Theorem 4.1. *Let $\varphi : \Omega \rightarrow \tilde{\Omega}$ be a Q -homeomorphism with $Q \in L_s(\Omega)$. Then φ generates the bounded composition operator*

$$\varphi^* : L_n^1(\tilde{\Omega}) \rightarrow L_q^1(\Omega),$$

where $q = sn/(s+n-1)$. In particular, if $Q \in L_1(\Omega)$, then φ generates the bounded composition operator

$$\varphi^* : L_n^1(\tilde{\Omega}) \rightarrow L_1^1(\Omega).$$

Proof. Because φ is a Q -homeomorphism with $Q \in L_s(\Omega)$, then $Q \in L_{1,\text{loc}}(\Omega)$. Hence φ belongs to the Sobolev space $W_{1,\text{loc}}^1(\Omega)$ and has a finite distortion [28].

In [28] it was proved that for $Q \in L_{1,\text{loc}}(\Omega)$ the inequality

$$(4.2) \quad |D\varphi(x)| \leq C(n)|J(x, \varphi)|^{\frac{1}{n}} Q^{\frac{n-1}{n}}(x)$$

holds for almost all $x \in \Omega$. Hence

$$\left(\frac{|D\varphi(x)|^n}{|J(x, \varphi)|} \right)^{\frac{1}{n-1}} \leq Q(x) \text{ for almost all } x \in \Omega.$$

Since $Q \in L_s(\Omega)$, we have

$$\int_{\Omega} \left(\frac{|D\varphi(x)|^n}{|J(x, \varphi)|} \right)^{\frac{s}{n-1}} dx \leq \int_{\Omega} Q^s(x) dx < \infty.$$

Then, by Theorem 2.1 the mapping φ generates the bounded composition operator $\varphi^* : L_n^1(\tilde{\Omega}) \rightarrow L_q^1(\Omega)$, $q = sn/(s + n - 1)$. \square

Using the composition duality property in the case of planar domains $\Omega, \tilde{\Omega} \subset \mathbb{R}^2$ we obtain:

Theorem 4.2. *Let $\varphi : \Omega \rightarrow \tilde{\Omega}$, $\Omega, \tilde{\Omega} \subset \mathbb{R}^2$, be a Q -homeomorphism with $Q \in L_1(\Omega)$. Then the inverse mapping $\varphi^{-1} : \tilde{\Omega} \rightarrow \Omega$ generates the bounded composition operator*

$$(\varphi^{-1})^* : L_{\infty}^1(\Omega) \rightarrow L_2^1(\tilde{\Omega}).$$

In particular, $\varphi \in L_2^1(\Omega)$.

Now we prove the inverse property.

Theorem 4.3. *Let a homeomorphism $\varphi : \Omega \rightarrow \tilde{\Omega}$ generates the bounded composition operator*

$$\varphi^* : L_n^1(\tilde{\Omega}) \rightarrow L_{n-1}^1(\Omega).$$

Then φ is a Q -homeomorphism with $Q(x) = K^I(x, \varphi)$.

Proof. This statement was proved in [28] in a different formulation and under the additional assumption that the mapping satisfies Luzin N -property. The proof below completely repeats the proof of [28, Theorem 4.1] and is presented for the convenience of the reader and to avoid confusion.

By the definition of the Q -homeomorphism, we have to show, that for every family Γ of paths in Ω and every $\rho \in adm(\Gamma)$

$$M(\varphi\Gamma) \leq \int_{\Omega} K^I(x, \varphi) \rho^n(x) dx.$$

First, note, that by theorem 2.1, $\varphi \in W_{n-1,loc}^1(\Omega)$. Also, $\varphi^{-1} \in W_{n,loc}^1(\tilde{\Omega})$ by theorem 2.2. It implies, that $\varphi^{-1} \in ACL_{loc}^n(\tilde{\Omega})$, is differentiable a.e. and satisfies Luzin N -property

By Fuglede's theorem ([40], p. 95), if $\tilde{\Gamma}$ is the family of all paths $\gamma \in \varphi\Gamma$ for which φ^{-1} is absolutely continuous on all closed subpaths of γ , then $M(\varphi\Gamma) = M(\tilde{\Gamma})$. Then, for given $\rho \in adm(\Gamma)$, one consider

$$\tilde{\rho}(y) = \begin{cases} \rho(\varphi^{-1}(y)) |D\varphi^{-1}(y)|, & y \in \tilde{\Omega}, \\ 0, & \text{otherwise.} \end{cases}$$

Then, for $\tilde{\gamma} \in \tilde{\Gamma}$

$$\int_{\tilde{\gamma}} \tilde{\rho} ds \geq \int_{\varphi^{-1} \circ \tilde{\gamma}} \rho ds \geq 1,$$

and consequently $\tilde{\rho} \in \text{adm}(\tilde{\Gamma})$.

By change of variable formula, we obtain

$$\begin{aligned} M(\varphi\Gamma) &= M(\tilde{\Gamma}) \leq \int_{\tilde{\Omega}} \tilde{\rho}^n dy \\ &= \int_{\tilde{\Omega}} \rho^n(\varphi^{-1}(y)) |D\varphi^{-1}(y)|^n dy = \int_{\tilde{\Omega}} \frac{\rho^n(\varphi^{-1}(y))}{l(D\varphi(\varphi^{-1}(y)))^n} dy \\ &= \int_{\tilde{\Omega}} \rho^n(\varphi^{-1}(y)) K^I(\varphi^{-1}(y), \varphi) J_{\varphi^{-1}}(y) dy \leq \int_{\Omega} K^I(x, \varphi) \rho^n(x) dx \end{aligned}$$

which completes the proof. \square

Hence we obtain the following equivalence theorem in \mathbb{R}^2 .

Theorem 4.4. *Let $\varphi : \Omega \rightarrow \tilde{\Omega}$ be a homeomorphism of planar domains $\Omega, \tilde{\Omega} \subset \mathbb{R}^2$. Then φ be a Q -homeomorphism with $Q \in L_1(\Omega)$ if and only if φ generate a bounded composition operator*

$$\varphi^* : L_2^1(\tilde{\Omega}) \rightarrow L_1^1(\Omega).$$

5. THE MEASURE DISTORTION PROPERTIES AND THE CAPACITARY METRIC DISTORTION PROPERTIES

5.1. The measure distortion properties of weak (p, q) -quasiconformal mappings. The first our results, based on theorem 2.3, is the following measure distortion properties, which refines results of [45], in the case $1 < q \leq p < n$ the similar result was obtained in [45] by using another set function.

Theorem 5.1. *Let $\varphi : \Omega \rightarrow \tilde{\Omega}$ be a weak (p, q) -quasiconformal mapping, $1 < q \leq p < n$. Then for any measurable subset $\tilde{A} \subset \tilde{\Omega}$ the following inequality hold:*

$$|\varphi^{-1}(\tilde{A})|^{\frac{n-q}{nq}} \leq C \tilde{\Phi}_{p,q}(\tilde{A})^{\frac{p-q}{pq}} |\tilde{A}|^{\frac{n-p}{np}}, \quad 1 < q \leq p < n.$$

Proof. In the case $q < p$ by Theorem 2.3, the following inequality

$$\|\varphi^*(f) | L_q^1(\Omega)\| \leq \tilde{\Phi}_{p,q}(B(y_0, 2r))^{\frac{p-q}{pq}} \|f | L_p^1(\tilde{\Omega} \cap B(y_0, 2r))\|,$$

holds for any function $f \in L_p^1(\tilde{\Omega}) \cap C_0(B(y_0, 2r))$ and for any ball $B(y_0, 2r)$ with the center at fixed point y_0 and with radius $2r$ such that $0 < 2r < \text{dist}(y_0, \partial\tilde{\Omega})$. Fix a cut-off function $\eta \in C_0^\infty(\mathbb{R}^n)$ such that $\eta = 1$ on the ball $B(0, 1)$ and $\eta = 0$ on the set $\mathbb{R}^n \setminus B(0, 2)$. Substituting the test functions

$$f(y) = \eta\left(\frac{|y - y_0|}{r}\right), \quad 0 < 2r < \text{dist}(y_0, \partial\tilde{\Omega}),$$

into the above inequality, we obtain

$$\begin{aligned} \|\varphi^*(f) | L_q^1(\varphi^{-1}(B(y_0, r)))\| &\leq \|\varphi^*(f) | L_q^1(\Omega)\| \\ &\leq \tilde{\Phi}_{p,q}(B(y_0, 2r))^{\frac{p-q}{pq}} \|f | L_p^1(\tilde{\Omega} \cap B(y_0, 2r))\| \\ &\leq C \tilde{\Phi}_{p,q}(B(y_0, 2r))^{\frac{p-q}{pq}} |B(y_0, r)|^{\frac{n-p}{np}}. \end{aligned}$$

Now, because functions $\varphi^*(f)$ are equal to one on the set $\varphi^{-1}(B(y_0, r))$, by using the embedding of the of functions with compact supports of the Sobolev space

$L_q^1(\varphi^{-1}(B(y_0, r)))$ into the Lebesgue space $L_{\frac{n-q}{n-q}}(\varphi^{-1}(B(y_0, r)))$, we have the following inequality for an arbitrary ball $B(y_0, r)$, $0 < 2r < \text{dist}(y_0, \partial\tilde{\Omega})$:

$$|\varphi^{-1}(B(y_0, r))|^{\frac{n-q}{n-q}} \leq C\tilde{\Phi}_{p,q}(B(y_0, 2r))^{\frac{p-q}{pq}} |B(y_0, r)|^{\frac{n-p}{np}}.$$

It is known (see, e.g. [46]), that for an arbitrary open set $\tilde{A} \subset \tilde{\Omega}$ there exists a sequence of balls $\{B(y_i, r_i)\}$, such that families $\{B(y_i, r_i)\}$ and $\{B(y_i, 2r_i)\}$ are finite covers of \tilde{A} and $\sum_{i=1}^{\infty} \tilde{\Phi}(B(y_i, 2r_i)) \leq C(n)\tilde{\Phi}(\tilde{A})$. Taking such sequences, we rise both sides of the above inequality to the power $\frac{np}{n-p}$ and sum over this families:

$$\sum_{i=1}^{\infty} |\varphi^{-1}(B(y_0, r))| \leq C \sum_{i=1}^{\infty} \tilde{\Phi}_{p,q}(B(y_0, 2r))^{\frac{n(p-q)}{p(n-q)}} |B(y_0, r)|^{\frac{q(n-p)}{p(n-q)}}.$$

Due to the choice of families of balls and the Hölder inequality, we obtain

$$|\varphi^{-1}(\tilde{A})| \leq C\tilde{\Phi}_{p,q}(\tilde{A})^{\frac{n(p-q)}{p(n-q)}} |\tilde{A}|^{\frac{q(n-p)}{p(n-q)}}.$$

Raising to the power of $(n-q)/nq$ give us the claimed result.

In the case $p = q$ we have that the inequality

$$\|\varphi^*(f) | L_q^1(\Omega)\| \leq C\|f | L_p^1(\tilde{\Omega} \cap B(y_0, 2r))\|,$$

holds for any function $f \in L_p^1(\tilde{\Omega}) \cap C_0(B(y_0, 2r))$ and for any ball $B(y_0, 2r)$ with the center at fixed point y_0 and with radius $2r$ such that $0 < 2r < \text{dist}(y_0, \partial\tilde{\Omega})$. Hence, by same reasons, like in the case $q < p$, we have the required result. \square

Theorem 5.2. *Let $\varphi : \Omega \rightarrow \tilde{\Omega}$ be a weak (p, q) -quasiconformal mapping, $n < q \leq p < \infty$, then for any measurable subset $A \subset \Omega$ the following inequality hold:*

$$|\varphi(A)|^{\frac{(p-n)(n-1)}{np}} \leq C\Phi_{q',p'}(A)^{\frac{(p-q)(n-1)}{pq}} |A|^{\frac{(q-n)(n-1)}{nq}}, \quad n < q \leq p < \infty,$$

where, by (2.3), $\Phi_{q',p'}(A)$ is a set function, defined on open sets $A \subset \Omega$ and associated with operator

$$(\varphi^{-1})^* : L_{q'}^1(\Omega) \rightarrow L_{p'}^1(\tilde{\Omega}), \quad p' = \frac{p}{p-n+1}, \quad q' = \frac{q}{q-n+1}.$$

Proof. Since $\varphi : \Omega \rightarrow \tilde{\Omega}$ is a weak (p, q) -quasiconformal mapping, $n < q \leq p < \infty$, then by the duality theorem (Theorem 2.2) the mapping φ^{-1} is the weak (q', p') -quasiconformal mapping with $1 < p' \leq q' < n$.

Hence, by the previous theorem

$$|\varphi(A)|^{\frac{n-p'}{np'}} \leq C\Phi_{q',p'}(A)^{\frac{q'-p'}{q'p'}} |A|^{\frac{n-q'}{nq'}}, \quad 1 < p' \leq q' < n,$$

for any measurable subset $A \subset \Omega$. Taking into account that $p' = p/(p-n+1)$, $q' = q/(q-n+1)$ we obtain

$$|\varphi(A)|^{\frac{(p-n)(n-1)}{np}} \leq C\Phi_{q',p'}(A)^{\frac{(p-q)(n-1)}{pq}} |A|^{\frac{(q-n)(n-1)}{nq}}.$$

\square

As a consequence of the above theorem we obtain the following result for integrability of the Jacobian (the case $1 < q \leq p < n$ was proved in [45]) by another method.

Theorem 5.3. *Let $\varphi : \Omega \rightarrow \tilde{\Omega}$ be a weak (p, q) -quasiconformal mapping, $1 < q \leq p < n$. Then*

$$J(\cdot, \varphi^{-1}) = J_{\varphi^{-1}}(\cdot) \in L_{\frac{p(n-q)}{n(p-q)}}(\tilde{\Omega}).$$

If $\varphi : \Omega \rightarrow \tilde{\Omega}$ is a weak (p, q) -quasiconformal mapping with $n < q \leq p < \infty$, then

$$J(\cdot, \varphi) = J_{\varphi}(\cdot) \in L_{\frac{q(p-n)}{n(p-q)}}(\Omega).$$

Proof. Let $1 \leq q \leq p < n$. Fix a point y_0 and a ball $B(y_0, r) \subset \tilde{\Omega}$. By theorem 5.1,

$$|\varphi^{-1}(B(y_0, r))|^{\frac{n-q}{nq}} \leq C \tilde{\Phi}_{p,q}(B(y_0, r))^{\frac{p-q}{pq}} |B(y_0, r)|^{\frac{n-p}{np}}.$$

Dividing both sides on $|B(y_0, r)|^{\frac{n-q}{nq}}$

$$\left(\frac{|\varphi^{-1}(B(y_0, r))|}{|B(y_0, r)|} \right)^{\frac{n-q}{nq}} \leq C \left(\frac{\tilde{\Phi}_{p,q}(B(y_0, r))}{|B(y_0, r)|} \right)^{\frac{p-q}{pq}}.$$

Rising to the power $\frac{pq}{p-q}$ and taking the limit $r \rightarrow 0$, we obtain $J_{\varphi^{-1}}(y_0)$ in the right hand side and a derivative $\tilde{\Phi}'_{p,q}$ of the set function $\tilde{\Phi}_{p,q}$. In [46], it was shown, that for quasiadditive set function $\tilde{\Phi}_{p,q}$ its derivative $\tilde{\Phi}'_{p,q}$ is integrable and

$$\int_{\tilde{\Omega}} \tilde{\Phi}'_{p,q}(y) dy \leq C \tilde{\Phi}_{p,q}(\tilde{\Omega}).$$

Combining these estimates and integrate over $\tilde{\Omega}$, we infer

$$\int_{\tilde{\Omega}} (J_{\varphi^{-1}}(y))^{\frac{p(n-q)}{n(p-q)}} dy \leq C \int_{\tilde{\Omega}} \tilde{\Phi}'_{p,q}(y) dy \leq C \tilde{\Phi}_{p,q}(\tilde{\Omega}) < \infty.$$

Hence,

$$J(\cdot, \varphi^{-1}) = J_{\varphi^{-1}}(\cdot) \in L_{\frac{p(n-q)}{n(p-q)}}(\tilde{\Omega}).$$

The case $n < q \leq p < \infty$ now follows from the duality theorem (Theorem 2.2). \square

5.2. Capacitary metric. In this subsection we introduce capacity metric of the hyperbolic type and prove, that (p, q) -quasiconformal mappings are Lipschitz mappings in correspondence capacity metrics.

Let $n-1 < p \leq n$ ($1 \leq p \leq 2$ if $n = 2$) and let a domain $\Omega \subset \mathbb{R}^n$ have a boundary of non-zero p -capacity.

Let us define a capacity metric

$$d_p(x, y) = \inf_{\gamma} \text{cap}_p^{\frac{1}{p}}(\gamma; \Omega),$$

where the infimum is taken over all continuous curves γ , joining points x and y in Ω . In the case $p = n$ metrics of such type was considered in [2] in a connection with quasiconformal mappings.

Theorem 5.4. *Let $n-1 < p \leq n$ ($1 \leq p \leq 2$ if $n = 2$) and $\text{cap}_p(\partial\Omega) > 0$ then*

$$d_p(x, y) = \inf_{\gamma} \text{cap}_p^{\frac{1}{p}}(\gamma; \Omega)$$

is a metric.

Proof. The restriction on p arises because for $p > n$ only empty set has zero capacity and for $p \leq n - 1$ even a continuum has zero capacity. If boundary $\partial\Omega$ has zero capacity, then, by the definition of $\text{cap}_p^{\frac{1}{p}}(\gamma; \Omega)$ will be equal to zero for any continuum γ .

1. Identity:

If $x = y$ then the infimum is reached on one-point set, which capacity is zero, hence, $d_p(x, y) = 0$. Now, let $d_p(x, y) = 0$. Then there exists a sequence γ_k , such that $\text{cap}_p(\gamma_k; \Omega) \rightarrow 0$ for $k \rightarrow \infty$. The sequence of extremal functions u_k for capacitors $(\gamma_k; \Omega)$ converges to 0 in $L_p^1(\Omega)$ and a.e. except the set of zero capacity. Since any continuum has a non-zero capacity, $x = y$.

2. Symmetry: is a direct consequence of definition $d_p(x, y)$.

2. Triangle inequality: follows from the triangle inequality for L_p^1 -norm in the definition of the capacity. \square

As a consequence of theorem 3.2 we obtain the following statement.

Theorem 5.5. *Let $\varphi : \Omega \rightarrow \tilde{\Omega}$ be a (p, q) -quasiconformal mapping, $n - 1 < q \leq p \leq n$ ($1 \leq q \leq p \leq 2$ if $n = 2$). Then, for any two points $x, y \in \tilde{\Omega}$ the following inequality*

$$d_q(\varphi^{-1}(x), \varphi^{-1}(y)) \leq K_{p,q}(\varphi; \Omega) d_p(x, y),$$

holds.

Proof. Because $\varphi : \Omega \rightarrow \tilde{\Omega}$ is a (p, q) -quasiconformal mapping, then there exists a bounded monotone countable-additive set function $\tilde{\Phi}$ defined on open subsets of $\tilde{\Omega}$ such that for every condenser $(F_0, F_1) \subset \tilde{\Omega}$ the inequality

$$\begin{aligned} \text{cap}_q^{1/q}(\varphi^{-1}(F_0), \varphi^{-1}(F_1); \Omega) &\leq \tilde{\Phi}_{p,q}(\tilde{\Omega} \setminus F_0)^{\frac{p-q}{pq}} \text{cap}_p^{1/p}(F_0, F_1; \tilde{\Omega}) \\ &\leq K_{p,q}(\varphi; \Omega) \text{cap}_p^{1/p}(F_0, F_1; \tilde{\Omega}) \end{aligned}$$

holds.

Let $\tilde{\gamma} \subset \tilde{\Omega}$ be a continuous curve which join x and y in $\tilde{\Omega}$. Because φ is a homeomorphism, then $\varphi^{-1}(\tilde{\gamma})$ is a continuous curve which join $\varphi^{-1}(x)$ and $\varphi^{-1}(y)$ in Ω . Hence we obtain the following inequalities:

$$d_q(\varphi^{-1}(x), \varphi^{-1}(y)) \leq \text{cap}_q^{\frac{1}{q}}(\varphi^{-1}(\tilde{\gamma}); \Omega) \leq K_{p,q}(\varphi; \Omega) \text{cap}_p^{\frac{1}{p}}(\tilde{\gamma}; \tilde{\Omega}).$$

Now since $\tilde{\gamma}$ is an arbitrary continuous curve which join x and y , then, taking the infimum over all such $\tilde{\gamma}$, we obtain

$$d_q(\varphi^{-1}(x), \varphi^{-1}(y)) \leq \inf K_{p,q}(\varphi; \Omega) \text{cap}_p^{\frac{1}{p}}(\tilde{\gamma}; \tilde{\Omega}) = K_{p,q}(\varphi; \Omega) d_p(x, y). \quad \square$$

Using the composition duality theorem (Theorem!2.2) for composition operators, we immediately obtain the following result:

Theorem 5.6. *Let $\varphi : \Omega \rightarrow \tilde{\Omega}$ be a (p, q) -quasiconformal mapping, $n < q \leq p < \infty$. Then, for any two points $x, y \in \Omega$ the following inequality*

$$d_{p'}(\varphi(x), \varphi(y)) \leq K_{q',p'}(\varphi^{-1}; \tilde{\Omega}) d_{q'}(x, y), \quad p' = \frac{p}{p-n+1}, \quad q' = \frac{q}{q-n+1}$$

holds.

6. POINCARÉ INEQUALITIES

In this section we prove the weighted Sobolev–Poincaré inequalities and on this base we provide estimates of the exact constants in Sobolev–Poincaré inequalities.

Let us recall that a bounded domain $\Omega \subset \mathbb{R}^n$ is said to be (s, q) -Sobolev–Poincaré domain (see, for example, [10, 13]), $1 \leq q, s \leq \infty$, if the following Sobolev–Poincaré inequality

$$\inf_{c \in \mathbb{R}} \|g - c\|_{L_s(\Omega)} \leq B_{s,q}(\Omega) \|g\|_{L_q^1(\Omega)}$$

holds for any $g \in L_q^1(\Omega)$ with the constant $B_{s,q}(\Omega) < \infty$.

Theorem 6.1. *Let a bounded domain $\tilde{\Omega} \subset \mathbb{R}^n$ be such that there exists a weak (p, q) -quasiconformal mapping $\varphi : \Omega \rightarrow \tilde{\Omega}$, $1 \leq q \leq p < \infty$, of a bounded (s, q) -Sobolev–Poincaré domain $\Omega \subset \mathbb{R}^n$ onto $\tilde{\Omega}$, $q < s$. Then the weighted Poincaré inequality*

$$\inf_{c \in \mathbb{R}} \left(\int_{\tilde{\Omega}} |f(x) - c|^s w(x) dx \right)^{\frac{1}{s}} \leq B_{s,p}^w(\tilde{\Omega}) \left(\int_{\tilde{\Omega}} |\nabla f(x)|^p dx \right)^{\frac{1}{p}}$$

holds for any function $f \in W_p^1(\tilde{\Omega})$ with the weight $w(x) = J_{\varphi^{-1}}(x)$, where $J_{\varphi^{-1}}(x)$ is a volume derivative of the inverse mapping to $\varphi : \Omega \rightarrow \tilde{\Omega}$.

Proof. Since the mapping $\varphi : \Omega \rightarrow \tilde{\Omega}$ is a weak (p, q) -quasiconformal, then the composition operator $\varphi^* : L_p^1(\tilde{\Omega}) \rightarrow L_q^1(\Omega)$ is bounded and, for a function $f \in L_p^1(\tilde{\Omega})$, the composition $f \circ \varphi$ belongs to $L_q^1(\Omega)$.

By the change of variable formula and since Ω is a bounded (s, q) -Sobolev–Poincaré domain, we have

$$\begin{aligned} \inf_{c \in \mathbb{R}} \left(\int_{\tilde{\Omega}} |f(y) - c|^s J_{\varphi^{-1}}(y) dy \right)^{\frac{1}{s}} &= \inf_{c \in \mathbb{R}} \left(\int_{\Omega} |f(\varphi(x)) - c|^s dx \right)^{\frac{1}{s}} \\ &\leq B_{s,q}(\Omega) \left(\int_{\Omega} |\nabla f(\varphi(x))|^q dx \right)^{\frac{1}{q}}, \end{aligned}$$

where $B_{s,q}(\Omega)$ is a best constant in the (s, q) -Sobolev–Poincaré inequality.

By Theorem 2.1

$$\|\varphi^*(f)\|_{L_q^1(\Omega)} \leq K_{p,q}(\Omega; \varphi) \|f\|_{L_p^1(\tilde{\Omega})}.$$

Hence

$$\begin{aligned} \inf_{c \in \mathbb{R}} \left(\int_{\tilde{\Omega}} |f(y) - c|^s J_{\varphi^{-1}}(y) dy \right)^{\frac{1}{s}} &\leq B_{s,q}(\Omega) \left(\int_{\Omega} |\nabla f(\varphi(x))|^q dx \right)^{\frac{1}{q}} \\ &\leq B_{s,q}(\Omega) K_{p,q}(\Omega; \varphi) \left(\int_{\tilde{\Omega}} |\nabla f(y)|^p dy \right)^{\frac{1}{p}} = B_{s,p}^w(\tilde{\Omega}) \left(\int_{\tilde{\Omega}} |\nabla f(x)|^p dx \right)^{\frac{1}{p}}. \end{aligned}$$

□

Applications of Q -homeomorphisms to the spectral theory of two-dimensional elliptic operators are based on estimates of the constants in non-weighted case $B_{s,p}(\Omega)$ for $\Omega \subset \mathbb{R}^2$. In [7], for $\Omega \subset \mathbb{R}^2$ there was proved the next version of Poincaré inequality with the upper estimates of the sharp constant:

Theorem 6.2 ([7]). *Let $f \in W_1^1(\Omega)$, $\Omega \subset \mathbb{R}^2$. Then for any $r > 0$ and any $z_0 \in \Omega$, such that $\text{dist}(z_0, \partial\Omega) > 2r$, the following inequality holds:*

$$(6.1) \quad \left(\iint_{D(z_0, r)} |f(z) - f_{D(z_0, r)}|^2 dx dy \right)^{\frac{1}{2}} \leq \frac{3\sqrt{\pi^3}}{4} \iint_{D(z_0, r)} |\nabla f(z)| dx dy.$$

Using the bi-Lipschitz change of variables, we can extend this result from a disk to a general domain with Lipschitz boundary and obtain an estimates for the constant $B_{2,2}(\Omega)$.

Theorem 6.3. *Let there exist a bi-Lipschitz mapping $\varphi : D(0, r) \rightarrow \Omega$ and $f \in W_1^1(\Omega)$. Then the following inequality holds:*

$$(6.2) \quad \left(\iint_{\Omega} |f(x, y) - f_{\Omega}|^2 dx dy \right)^{\frac{1}{2}} \leq \frac{3\sqrt{L^5 \pi^3}}{4} \iint_{\Omega} |\nabla f(x, y)| dx dy,$$

where L is a Lipschitz constant for mapping φ .

Proof. For the simplicity, we prove the inequality for the case $f_{\Omega} = 0$. Since the function f belongs to $L_1^1(\Omega)$, composition $f \circ \varphi$ with bi-Lipschitz mapping belongs L_1^1 and we can apply the change of variable formula.

$$\begin{aligned} \left(\iint_{\Omega} |f(u, v)|^2 dudv \right)^{\frac{1}{2}} &= \left(\iint_{D(0, r)} |f(\varphi(x, y))|^2 |J(x, y, \varphi)| dx dy \right)^{\frac{1}{2}} \\ &\leq \text{ess sup}_{(x, y) \in D(0, r)} |J(x, y, \varphi)|^{\frac{1}{2}} \left(\iint_{D(0, r)} |f(\varphi(x, y))|^2 dx dy \right)^{\frac{1}{2}} \end{aligned}$$

By theorem 6.2, we obtain the following estimate:

$$\left(\iint_{\Omega} |f(u, v)|^2 dudv \right)^{\frac{1}{2}} \leq \frac{3\sqrt{\pi^3}}{4} \text{ess sup}_{(x, y) \in D(0, r)} |J(x, y, \varphi)|^{\frac{1}{2}} \iint_{D(0, r)} |\nabla f(\varphi(x, y))| dx dy.$$

Using the chain rule and the change of variable formula the second time, we infer

$$\left(\iint_{\Omega} |f(u, v)|^2 dudv \right)^{\frac{1}{2}} \leq \frac{3\sqrt{\pi^3}}{4} \operatorname{ess\,sup}_{(x,y) \in D(0,r)} |J(x, y, \varphi)|^{\frac{1}{2}} \operatorname{ess\,sup}_{(x,y) \in D(0,r)} \frac{|D\varphi(x, y)|}{|J(x, y, \varphi)|} \iint_{\Omega} |\nabla f(u, v)| dudv.$$

By the Hadamar inequality, we can choose a constant L such that $0 < L^{-1} \leq |J_{\varphi}(x, y)|^{1/2} \leq |D\varphi(x, y)| \leq L$. Then

$$\left(\iint_{\Omega} |f(u, v)|^2 dudv \right)^{\frac{1}{2}} \leq \frac{3\sqrt{L^5\pi^3}}{4} \iint_{\Omega} |\nabla f(u, v)| dudv.$$

□

7. SPECTRAL ESTIMATES OF ELLIPTIC OPERATORS

In this section we consider applications of Q -homeomorphisms to spectral estimates of the Laplace operator in Hölder cusp domains. The detailed description of the spectral theory of non-linear elliptic operators and applications of geometric theory of composition operators see, for example, [6, 13, 15]. Let us consider the Neumann spectral problem

$$\begin{cases} -\Delta u = \mu u & \text{in } \tilde{\Omega}, \\ \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\tilde{\Omega}. \end{cases}$$

By the Min-Max Principle the first non-trivial Neumann eigenvalues in the domains $\tilde{\Omega}$ can be characterized as

$$\mu_1(\tilde{\Omega}) = \left(B_{2,2}(\tilde{\Omega}) \right)^{-2}.$$

With the help of Q -homeomorphism we can estimate the first eigenvalue of the Neumann-Laplacian in the case of cusp domain. The estimate is based on the following anti-commutative diagram [4, 10]:

$$\begin{array}{ccc} L_2^1(\tilde{\Omega}) & \xrightarrow{\varphi^*} & L_1^1(\Omega) \\ \downarrow & & \downarrow \\ L_2(\tilde{\Omega}) & \xleftarrow{(\varphi^{-1})^*} & L_2(\Omega) \end{array}$$

By this diagram, $B_{2,2}(\tilde{\Omega}) \leq \|\varphi^*\| B_{2,1}(\Omega) \|(\varphi^{-1})^*\|$

Let us consider the Hölder cusp domain $\Omega_{\alpha} \subset \mathbb{R}^2$ which is the image of the square $\Omega = \{(x, y) : |x| + |y| \leq 1\}$ under Q -homeomorphism $\varphi : \Omega \rightarrow \Omega_{\alpha}$ of the form

$$\varphi(x, y) = \begin{cases} (x, y^{\alpha}), & y \geq 0, \\ (x, -y^{\alpha}), & y < 0. \end{cases}$$

Theorem 7.1. *Let $\Omega_{\alpha} \subset \mathbb{R}^2$ be the Hölder cusp domain. Then*

$$\mu_1(\Omega_{\alpha}) \geq \frac{(\alpha + 1)\alpha(2 - \alpha)(3 - \alpha)}{9\pi^3\alpha(\alpha + 1 + \alpha(2 - \alpha)(3 - \alpha))}.$$

In particular, if $\alpha = 4$, then $\mu_1(\Omega_\alpha) \geq (11, 7\pi^3)^{-1}$.

Proof. First of all, we calculate the constant in inequality (6.2) for the square $\Omega = \{(x, y) : |x| + |y| \leq 1\}$. As a bi-Lipschitz mapping, let us consider the radial transformation $\psi : D(0, 1) \rightarrow \Omega$, acting by the rule

$$\psi(x, y) = (l(x, y)x, l(x, y)y), \quad l(x, y) = \frac{\sqrt{x^2 + y^2}}{|x| + |y|}.$$

By simple calculation, we obtain

$$|D\psi(x, y)| = \left(\frac{2(x^2 + y^2)^2}{(|x| + |y|)^4} + \frac{x^2 + y^2}{(|x| + |y|)^2} \right)^{\frac{1}{2}}$$

and

$$J(x, y, \psi) = \frac{x^2 + y^2}{(x + y)^2}.$$

Then, for $(x, y) \in D(0, 1)$,

$$\sup_{(x, y)} |J(x, y, \psi)|^{\frac{1}{2}} = 1, \quad \sup_{(x, y)} \frac{|D\psi(x, y)|}{J(x, y, \psi)} = 2.$$

Substituting these values to inequality (6.2), we obtain

$$(7.1) \quad \left(\iint_{\Omega} |f(x, y) - f_{\Omega}|^2 dx dy \right)^{\frac{1}{2}} \leq \frac{3\sqrt{\pi^3}}{2} \iint_{\Omega} |\nabla f(x, y)| dx dy,$$

The next step is to calculate the norms of composition operators

$$\varphi^* : L_2^1(\Omega_\alpha) \rightarrow L_1^1(\Omega) \quad \text{and} \quad (\varphi^{-1})^* : L_2(\Omega) \rightarrow L_2(\Omega_\alpha).$$

By theorem 2.1 the norm

$$\|\varphi^*\| \leq K_{2,2}(\Omega; \varphi) = \left\| \frac{|D\varphi(\cdot)|}{|J(\cdot, \varphi)|^{\frac{1}{2}}} \Big| L_2(\Omega) \right\|.$$

In the [46], it was proven, that the norm $(\varphi^{-1})^*$ can be calculate as

$$\|(\varphi^{-1})^*\| = M_2(\Omega) = \sup_{(x, y) \in \Omega} |J(x, y, \varphi)|^{\frac{1}{2}}.$$

For the Q -homeomorphism $\varphi : \Omega_\alpha \rightarrow \Omega$,

$$|D\varphi(x, y)| = \sqrt{1 + \alpha^2 y^{2(\alpha-1)}}, \quad J(x, y, \varphi) = \alpha y^{\alpha-1};$$

$$K_{2,2}(\Omega) = \left(4 \int_0^1 dx \int_0^x \frac{1 + \alpha^2 y^{2(\alpha-1)}}{\alpha y^{\alpha-1}} dy \right)^{\frac{1}{2}} = \left(\frac{4}{\alpha(2-\alpha)(3-\alpha)} + \frac{4}{\alpha+1} \right)^{\frac{1}{2}};$$

$$M_2(\Omega) = \sup_{(x, y) \in \Omega} |J(x, y, \varphi)|^{\frac{1}{2}} = \sqrt{\alpha}.$$

Combine all the estimates, we infer that

$$\mu_1(\Omega_\alpha) \geq \frac{(\alpha+1)\alpha(2-\alpha)(3-\alpha)}{9\pi^3\alpha(\alpha+1+\alpha(2-\alpha)(3-\alpha))}.$$

□

REFERENCES

- [1] H. Federer, Geometric measure theory, Springer Verlag, Berlin, 1969.
- [2] J. Ferrand, G. J. Martin, M. K. Vuorinen, Lipschitz conditions in conformally invariant metrics, *J. Analyse Math.*, 56 (1991), 187–210.
- [3] F. W. Gehring, Lipschitz mappings and the p -capacity of rings in n -space, *Advances in the theory of Riemann surfaces (Proc. Conf., Stony Brook, N.Y., 1969)*, 175–193. *Ann. of Math. Studies*, No. 66. Princeton Univ. Press, Princeton, N.J., 1971.
- [4] V. Gol'dshtein, L. Gurov, Applications of change of variables operators for exact embedding theorems, *Integral Equations Operator Theory* 19 (1994), 1–24.
- [5] V. Gol'dshtein, L. Gurov, A. Romanov, Homeomorphisms that induce monomorphisms of Sobolev spaces, *Israel J. Math.*, 91 (1995), 31–60.
- [6] V. Gol'dshtein, V. Pchelintsev, A. Ukhlov, On the First Eigenvalue of the Degenerate p -Laplace Operator in Non-convex Domains. *Integral Equations Operator Theory* 90 (2018), 90:43.
- [7] V. Gol'dshtein, V. Pchelintsev, A. Ukhlov, Integral estimates of conformal derivatives and spectral properties of the Neumann-Laplacian. *Journal of Mathematical Analysis and Applications*, 463 (2018), 19:39.
- [8] V. M. Gol'dshtein, Yu. G. Reshetnyak, Quasiconformal mappings and Sobolev spaces, Dordrecht, Boston, London: Kluwer Academic Publishers, 1990.
- [9] V. M. Gol'dshtein, V. N. Sitnikov, Continuation of functions of the class W_p^1 across Hölder boundaries, Imbedding theorems and their applications, *Trudy Sem. S. L. Soboleva*, 1 (1982), 31–43.
- [10] V. Gol'dshtein, A. Ukhlov, Weighted Sobolev spaces and embedding theorems, *Trans. Amer. Math. Soc.*, 361, (2009), 3829–3850.
- [11] V. Gol'dshtein, A. Ukhlov, About homeomorphisms that induce composition operators on Sobolev spaces, *Complex Var. Elliptic Equ.*, 55 (2010), 833–845.
- [12] V. Gol'dshtein, A. Ukhlov, Brennan's Conjecture and universal Sobolev inequalities, *Bull. Sci. Math.*, 138 (2014), 253–269.
- [13] V. Gol'dshtein, A. Ukhlov, On the first Eigenvalues of Free Vibrating Membranes in Conformal Regular Domains, *Arch. Rational Mech. Anal.*, 221 (2016), no. 2, 893–915.
- [14] V. Gol'dshtein, A. Ukhlov, Traces of function of L^1_2 Dirichlet spaces on the Caratheodory boundary, *Studia Math.* 235 (2016), 209–224.
- [15] V. Gol'dshtein, A. Ukhlov, The spectral estimates for the Neumann-Laplace operator in space domains, *Adv. in Math.*, 315 (2017), 166–193.
- [16] V. Gol'dshtein, A. Ukhlov, On functional properties of weak (p, q) -quasiconformal homeomorphisms, *Ukrainian Math. Bull.*, 16 (2019), 329–344.
- [17] P. Hajlasz, Change of variables formula under minimal assumptions, *Colloq. Math.*, 64 (1993), 93–101.
- [18] P. Halmos, Measure Theory, Springer-Verlag New York, 1950.
- [19] J. Heinonen, T. Kilpelinen, O. Martio, *Nonlinear Potential Theory of Degenerate Elliptic Equations*. Clarendon Press. Oxford, New York, Tokio, 1993.
- [20] J. Heinonen, P. Koskela, Weighted Sobolev and Poincaré inequalities and quasiregular mappings of polynomial type, *Math. Scand.* 77 (1995), 251–271.
- [21] S. Hencl, P. Koskela, *Lectures on Mappings of Finite Distortion*, Springer, Berlin/Heidelberg, 2014.
- [22] P. Koskela, Zh. Zhu, Sobolev extensions via reflections, arXiv:1812.09037.
- [23] V. I. Kruglikov, Capacities of condensers and quasiconformal in the mean mappings in space. *Mat. Sb. (N.S.)* 130 (172) (1986), 185–206.
- [24] S. L. Krushkal, On mean quasiconformal mappings, *Dokl. Akad. Nauk SSSR*, 157 (1964), 517–519.
- [25] O. Martio, S. Rickman, J. Väisälä, Definitions for quasiregular mappings, *Ann. Acad. Sei. Fenn., Ser. AI*, 448 (1969), 1–40.
- [26] O. Martio, V. Ryazanov, U. Srebro, E. Yakubov, To the theory of Q -homeomorphisms, *Dokl. Akad. Nauk Rossii*, 381 (2001), 20–22.
- [27] O. Martio, V. Ryazanov, U. Srebro and E. Yakubov, Mappings with finite length distortion, *J. d'Anal. Math.*, 93 (2004), 215–236.

- [28] O. Martio, V. Ryazanov, U. Srebro, E. Yakubov, Moduli in modern mapping theory. Springer Monographs in Mathematics. Springer, New York, 2009.
- [29] V. G. Maz'ya, Weak solutions of the Dirichlet and Neumann problems, Trudy Moskov. Mat. Ob-va., 20 (1969), 137–172 (1969)
- [30] V. Maz'ya, Sobolev spaces: with applications to elliptic partial differential equations, Springer, Berlin/Heidelberg, 2010.
- [31] V. G. Maz'ya, V. P. Havin, Non-linear potential theory, Russian Math. Surveys, **27** (1972), 71–148.
- [32] A. Menovschikov, A. Ukhlov, Composition operators on Hardy-Sobolev spaces and BMO-quasiconformal mappings, Ukrainian Math. Bull., (2021)
- [33] L. E. Payne, H. F. Weinberger, An optimal Poincaré inequality for convex domains, Arch. Rat. Mech. Anal., 5 (1960), 286–292.
- [34] I. N. Pesin, Mappings that are quasiconformal in the mean, Soviet Math. Dokl., 10 (1969), 939–941.
- [35] V. I. Ryazanov, On mean quasiconformal mappings, Siberian Math. J., 37 (1996), 325–334.
- [36] S. L. Sobolev, On some groups of transformations of the n -dimensional space, Dokl. AN SSSR, 32 (1941), 380–382.
- [37] G. D. Suvorov, The generalized "length and area principle" in mapping theory, Naukova Dumka, Kiev, 1985.
- [38] A. Ukhlov, On mappings, which induce embeddings of Sobolev spaces, Siberian Math. J., 34 (1993), 185–192.
- [39] A. Ukhlov, S. K. Vodop'yanov, Mappings associated with weighted Sobolev spaces. Complex analysis and dynamical systems III, Contemp. Math., Amer. Math. Soc., Providence, RI. 455 (2008), 369–382.
- [40] J. Väisälä, Lectures on n -dimensional quasiconformal mappings. Lecture Notes in Math. 229, Springer Verlag, Berlin, 1971.
- [41] S. K. Vodop'yanov, Taylor Formula and Function Spaces, Novosibirsk Univ. Press., 1988.
- [42] S. K. Vodop'yanov, Composition Operators on Weighted Sobolev Spaces and the Theory of Q_p -Homeomorphisms, Doklady Mathematics, 102 (2020), 371–375.
- [43] S. K. Vodop'yanov, V. M. Gol'dshtein, Yu. G. Reshetnyak, On geometric properties of functions with generalized first derivatives, Uspekhi Mat. Nauk 34 (1979), 17–65.
- [44] S. K. Vodop'yanov, A. D. Ukhlov, Sobolev spaces and (p, q) -quasiconformal mappings of Carnot groups. Siberian Math. J., 39 (1998), 776–795.
- [45] S. K. Vodop'yanov, A. D. Ukhlov, Superposition operators in Sobolev spaces, Russian Mathematics (Izvestiya VUZ) 46 (2002), no. 4, 11–33.
- [46] S. K. Vodop'yanov, A. D. Ukhlov, Set Functions and Their Applications in the Theory of Lebesgue and Sobolev Spaces. I, Siberian Adv. Math., 14:4 (2004), 78–125.

Alexander Menovschikov; Department of Mathematics, University of Hradec Králové, Rokitanského 62, 500 03 Hradec Králové, Czech Republic. Faculty of Economics, University of South Bohemia, Studentská 13, 370 05 České Budějovice, Czech Republic.

E-mail address: alexander.menovschikov@uhk.cz

Alexander Ukhlov; Department of Mathematics, Ben-Gurion University of the Negev, P.O.Box 653, Beer Sheva, 8410501, Israel

E-mail address: ukhlov@math.bgu.ac.il