

# NICHOLS ALGEBRAS OF DIMENSION $n^m$ AND MULTINOMIAL EXPANSION

YUXING SHI

**ABSTRACT.** In this paper, we discover Nichols algebras of dimension  $n^m$  and Gelfand-Kirillov dimension  $m$  via their relations with multinomial expansion. This result is a generalization of the work in arXiv:2103.06489, which built a connection between the Nichols algebras of square dimension and Pascal's triangle.

## 1. INTRODUCTION

Nichols algebras first appeared in a work of Nichols [Nic78], for construction of certain pointed Hopf algebras. They also arose independently in Woronowicz, Lusztig, and Rosso's works [Wor89][Lus93][Ros98]. Finite dimensional Nichols algebras are important ingredients for the classification of finite dimensional non-semisimple Hopf algebras with dual Chevalley property [AS02]. One central task for the study of Nichols algebras is to classify all finite dimensional Nichols algebras. The classification for finite dimensional Nichols algebras of group type has archived great success. For examples, Nichols algebras of diagonal type were classified completely by Heckenberger [Hec09] based on the theory of reflections [HS20] and Weyl groupoid [Hec06]; the classification of Nichols algebras of non-simple semisimple Yetter-Drinfeld modules over non-abelian groups were almost finished by Heckenberger and Vendramin [HV17b] [HV17a].

The study for Nichols algebras of non-group type is rare. One possible approach to obtain finite dimensional Nichols algebras of non-group type is to investigate Yetter-Drinfeld categories of Hopf algebras which are not categorically Morita-equivalent to group algebras [Shi20a]. The other way is to study the Nichols algebras directly from braidings [AG18] [GS21], since a Nichols algebra is completely determined by its braiding. We are only interested in rigid braidings for they have realizations in Yetter-Drinfeld categories of Hopf algebras [Sch92] [Tak00]. In this paper, we define a class of rigid braidings arising from the actions of symmetric groups on sets. We prove the existence of Nichols algebras of dimension

---

2010 *Mathematics Subject Classification.* 16T05, 16T25, 17B37.

*Keywords:* Nichols algebra; Multinomial expansion; Symmetric group; Gelfand-Kirillov dimension; Hopf algebra.

$n^m$ , via its connections with the multinomial expansion  $(x_1 + \cdots + x_m)^n$ . The result generalizes a previous work of the author in [Shi20b], which associated the Nichols algebras of square dimension with Pascal's triangle. Those Nichols algebras  $\mathfrak{B}(W_{m,n_0})$  ( $n_0 \neq 0$ ) obtained in the paper are not of group type in general.

The paper is organized as follows. In the section 2, we introduce the Nichols algebra, braidings arising from actions of symmetric group on sets, and some notations. In the section 3, we define a class of actions of  $\mathbb{S}_n$  on  $\mathbb{Z}_m^n$ , and reveal a connection between the decomposition of  $\mathbb{Z}_m^n$  into orbits and multinomial expansion. We obtain Nichols algebras of dimension  $n^m$  and Gelfand–Kirillov dimension  $m$ , according to their connections with multinomial expansion.

## 2. PRELIMINARIES

**2.1. Nichols algebra.** Let  $\mathbb{k}$  be an algebraically closed field of characteristic 0, and  $\mathbb{k}^\times = \mathbb{k} - \{0\}$ . Here we give a brief introduction to the Nichols algebra. For more details, please refer to Heckenberger and Schneider's monograph [HS20].

**Definition 2.1.** [AS02, Definition 2.1] Let  $H$  be a Hopf algebra and  $V \in {}^H_H\mathcal{YD}$ . A braided  $\mathbb{N}$ -graded Hopf algebra  $R = \bigoplus_{n \geq 0} R(n) \in {}^H_H\mathcal{YD}$  is called the *Nichols algebra* of  $V$  if

- (i)  $\mathbb{k} \simeq R(0)$ ,  $V \simeq R(1) \in {}^H_H\mathcal{YD}$ .
- (ii)  $R(1) = \mathcal{P}(R) = \{r \in R \mid \Delta_R(r) = r \otimes 1 + 1 \otimes r\}$ .
- (iii)  $R$  is generated as an algebra by  $R(1)$ .

In this case,  $R$  is denoted by  $\mathfrak{B}(V) = \bigoplus_{n \geq 0} \mathfrak{B}^n(V)$ .

*Remark 2.2.* Let  $(V, c)$  be a braided vector space, then the Nichols algebra  $\mathfrak{B}(V)$  is completely determined by the braiding  $c$ . More precisely, as proved in [Sch96] and noted in [AS02],

$$\mathfrak{B}(V) = \mathbb{k} \oplus V \oplus \bigoplus_{n=2}^{\infty} V^{\otimes n} / \ker \mathfrak{S}_n = T(V) / \ker \mathfrak{S},$$

where  $\mathfrak{S}_{n,1} \in \text{End}_{\mathbb{k}}(V^{\otimes(n+1)})$ ,  $\mathfrak{S}_n \in \text{End}_{\mathbb{k}}(V^{\otimes n})$ ,

$$c_i := \text{id}^{\otimes(i-1)} \otimes c \otimes \text{id}^{\otimes(n-i-1)} \in \text{End}_{\mathbb{k}}(V^{\otimes n}),$$

$$\mathfrak{S}_{n,1} := \text{id} + c_n + c_{n-1}c_n + \cdots + c_1 \cdots c_{n-1}c_n = \text{id} + \mathfrak{S}_{n-1,1}c_n,$$

$$\mathfrak{S}_1 := \text{id}, \quad \mathfrak{S}_2 := \text{id} + c, \quad \mathfrak{S}_n := \mathfrak{S}_{n-1,1}(\mathfrak{S}_{n-1} \otimes \text{id}).$$

**2.2. Braidings arising from actions of symmetric group on sets.** Let  $V$  be a non-empty set,  $\mathbb{S}_n$  be the symmetric group on  $n$  letters. The symmetric group  $\mathbb{S}_n$  is generated by transpositions  $s_1, \cdots, s_{n-1}$ . Denote  $V^n = \{i_1 i_2 \cdots i_n \mid i_j \in V, 1 \leq j \leq n\}$ . Let  $\mathcal{O}(x) = \mathbb{S}_n \cdot x$  be the orbit of  $x \in V^n$ . Denote  $\mathfrak{G}(x, y) = \{\sigma \in \mathbb{S}_n \mid \sigma \cdot x = y\}$  for  $x, y \in V$ . Suppose that there is an action of  $\mathbb{S}_n$  on  $V^n$  as

$$s_k \cdot (i_1 \cdots i_{k-1} p q i_{k+2} \cdots i_n) = i_1 \cdots i_{k-1} p' q' i_{k+2} \cdots i_n,$$

then we can define a braided vector space  $(W, c)$  such that  $W = \sum_{i \in V} \mathbb{k}w_i$ ,  $c(w_p \otimes w_q) = R_{pq}^{p'q'} w_{p'} \otimes w_{q'}$  for suitable  $R_{pq}^{p'q'} \in \mathbb{k}^\times$ . For  $x = i_1 i_2 \cdots i_n \in V^n$ , we denote  $w_x = w_{i_1} w_{i_2} \cdots w_{i_n}$  for abbreviation of  $w_{i_1} \otimes w_{i_2} \otimes \cdots \otimes w_{i_n}$ . Define  $\mathcal{T} : \mathbb{S}_n \rightarrow \text{End}_k(W^{\otimes n})$  via  $s_i \mapsto c_i = \mathcal{T}_{s_i}$ , then  $\mathfrak{S}_n(w_x) = \sum_{s \in \mathbb{S}_n} \mathcal{T}_s(w_x)$  for  $x \in V^n$ .

For  $k_1 + \cdots + k_r = n$ , the set of  $(k_1, \cdots, k_r)$ -shuffles, i.e. the set of permutations  $w$  such that  $w(1) < w(2) < \cdots < w(k_1)$ ,  $w(k_1 + 1) < w(k_1 + 2) < \cdots < w(k_1 + k_2)$ ,  $\cdots$ ,  $w(k_1 + \cdots + k_{r-1} + 1) < \cdots < w(n)$ , is denoted by  $\text{shuffle}(k_1, \cdots, k_r)$ . Let  $\mathbb{G}_n$  be the set of  $n$ -th primitive roots of unity. Denote  $(n)_q = 1 + q + \cdots + q^{n-1}$ ,  $(n)_q! = \prod_{k=1}^n (k)_q$ .

### 3. MULTINOMIAL EXPANSION AND NICHOLS ALGEBRAS OF DIMENSION $n^m$

**3.1. Multinomial expansion and actions of  $\mathbb{S}_n$  on  $\mathbb{Z}_m^n$ .** Let  $m \in \mathbb{Z}$ ,  $m \geq 2$ ,  $\mathbb{Z}_m^n = \{i_1 i_2 \cdots i_n \mid i_j \in \mathbb{Z}_m, 1 \leq j \leq n\}$ . We define an action of  $\mathbb{S}_n$  on  $\mathbb{Z}_m^n$  as follows:

$$s_k \cdot (i_1 \cdots i_{k-1} p q i_{k+2} \cdots i_n) = i_1 \cdots i_{k-1} p' q' i_{k+2} \cdots i_n,$$

where  $p' \equiv q - n_0 \pmod{m}$ ,  $q' \equiv p + n_0 \pmod{m}$  for a fixed integer  $n_0 \in \mathbb{Z}_m$ .

Denote  $\mathcal{P}(n, m)$  the set of partitions  $\lambda = (\lambda_1, \lambda_2, \cdots, \lambda_m)$  such that  $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_m \geq 0$ ,  $\lambda_1 + \lambda_2 + \cdots + \lambda_m = n$ . In general, we omit the zero parts of  $\lambda$ . The number of permutations of  $\lambda = (\lambda_1, \lambda_2, \cdots, \lambda_m) \in \mathcal{P}(n, m)$  is denoted as  $\text{Perm}(\lambda)$ . For example, if  $\lambda = (5, 5, 0) \in \mathcal{P}(10, 3)$ , then  $\text{Perm}(\lambda) = 3$ .

Denote  $\chi_{a,b}^n = i_1 i_2 \cdots i_n$  such that  $i_{j+1} \equiv i_j + n_0 \pmod{m}$  for  $j = 1, \cdots, n-1$ ,  $i_1 = a$ ,  $i_n = b$ .

**Definition 3.1.** If  $x = \chi_{a_1, b_1}^{\lambda_1} \chi_{a_2, b_2}^{\lambda_2} \cdots \chi_{a_k, b_k}^{\lambda_k}$  such that  $\lambda = (\lambda_1, \cdots, \lambda_k) \in \mathcal{P}(n, m)$ ,

$$(3.1) \quad a_{i+1} \not\equiv b_i + n_0 = a_i + \lambda_i n_0 \pmod{m},$$

$$(3.2) \quad a_{i+1} \not\equiv \lambda_i n_0 + b_{i-1} + n_0 = a_{i-1} + (\lambda_i + \lambda_{i-1}) n_0 \pmod{m},$$

$$(3.3) \quad \cdots \cdots \cdots,$$

$$(3.4) \quad a_{i+1} \not\equiv a_1 + (\lambda_1 + \lambda_2 + \cdots + \lambda_i) n_0 \pmod{m}, \quad 1 \leq i \leq k-1,$$

then we say  $x$  is a  $\lambda$ -element and the orbit  $\mathcal{O}(x)$  is corresponding to the partition  $\lambda$ . And we let  $\mathcal{B}(\lambda)$  be the union of all orbits corresponding to the partition  $\lambda$ .

*Remark 3.2.* From the definition, we see there are  $m - i$  choices for  $a_{i+1}$ . So the number of  $\lambda$ -elements is  $\binom{m}{k}$ .

**Lemma 3.3.** If  $x = \chi_{a_1, b_1}^{\lambda_1} \chi_{a_2, b_2}^{\lambda_2} \cdots \chi_{a_k, b_k}^{\lambda_k}$  is a  $\lambda$ -element for the partition  $\lambda = (\lambda_1, \cdots, \lambda_k) \in \mathcal{P}(n, m)$ , then the number of  $\lambda$ -elements in  $\mathcal{O}(x)$  is  $k!$ .

*Proof.* If  $\lambda = (\lambda_1) \in \mathcal{P}(n, m)$ , then there is exactly one  $\lambda$ -element in  $\mathcal{O}(x)$ . If  $\lambda = (\lambda_1, \lambda_1) \in \mathcal{P}(n, m)$ , then there are exactly two  $\lambda$ -elements in  $\mathcal{O}(x)$ , see the figure 1. By induction, we only need to prove the lemma holds for  $k+1$  under the condition that the lemma holds for  $k$ .




FIGURE 1.  $\mathcal{O}(\chi_{a_1, b_1}^{\lambda_1} \chi_{a_2, b_2}^{\lambda_2}) = \mathcal{O}(\chi_{a_2 - \lambda_1 n_0, b_2 - \lambda_1 n_0}^{\lambda_2} \chi_{a_1 + \lambda_2 n_0, b_1 + \lambda_2 n_0}^{\lambda_1})$

Let  $x_1 = \chi_{a_1, b_1}^{\lambda_1} \chi_{a_2, b_2}^{\lambda_1} \cdots \chi_{a_k, b_k}^{\lambda_1} \chi_{a_{k+1}, b_{k+1}}^{\lambda_1}$  be a  $\lambda$ -element for the partition  $\lambda = (\lambda_1, \dots, \lambda_1) \in \mathcal{P}(n, m)$ . Then  $a_{i+1} \not\equiv a_{i-j} + j\lambda_1 n_0 \in \mathbb{Z}_m$  for  $i = 1, \dots, k$  and  $j = 1, \dots, i$ . Similarly to the figure 1, if we move the term  $\chi_{a_l, b_l}^{\lambda_1}$  to the head of  $x_1$  for  $l = 2, \dots, k+1$ , then we obtain  $x_l = \chi_{a_l - (l-1)\lambda_1 n_0, b_l - (l-1)\lambda_1 n_0}^{\lambda_1} \chi_{a_1 - \lambda_1 n_0, b_1 - \lambda_1 n_0}^{\lambda_1} \cdots \in \mathcal{O}(x_1)$ . Notice that the leading terms of  $x_l$  for  $l = 1, \dots, k+1$  are different in  $\mathbb{Z}_m$ , the lemma is true for the case  $k+1$  by induction.  $\square$

A composition is a sequence  $\mu = (\mu_1, \dots, \mu_k)$  such that  $\mu_i \in \mathbb{Z}^+$  for  $i = 1, \dots, k$ . The length of a composition is the number of its parts, denoted as  $l(\mu)$ . For any  $y \in \mathbb{Z}_m^n$ , we can write  $y$  in the form  $\chi_{a_1, b_1}^{\mu_1} \cdots \chi_{a_k, b_k}^{\mu_k}$  for a composition  $\mu = (\mu_1, \dots, \mu_k)$  such that  $a_j \not\equiv b_{j-1} + n_0 \in \mathbb{Z}_m$ ,  $j = 2, \dots, k$ ,  $\mu_1 + \dots + \mu_k = n$ . As for convenience, we say  $y$  is a composition  $\mu$ -element. Now we give two operations on  $y$ .

- (1) If there exist  $i, j$  such that  $1 \leq i < j \leq k$ ,  $a_j \equiv a_i + (\mu_{j-1} + \dots + \mu_{i+1} + \mu_i)n_0 \in \mathbb{Z}_m$ , then we can move  $\chi_{a_j, b_j}^{\mu_j}$  forward to join with  $\chi_{a_i, b_i}^{\mu_i}$ . In other words, there exists a composition  $\mu'$ -element  $y' \in \mathcal{O}(y)$ ,  $l(\mu') < l(\mu)$ .
- (2) If there exists  $i$  such that  $\mu_i < \mu_{i+1}$  for  $1 \leq i \leq k-1$ , then we can exchange the parts  $\chi_{a_i, b_i}^{\mu_i}, \chi_{a_{i+1}, b_{i+1}}^{\mu_{i+1}}$  as what does in the figure 1.

We can perform the above two operations repeatedly. In final, we obtain a  $\lambda$ -element  $\bar{y} \in \mathcal{O}(y)$  for some partition  $\lambda$  of  $n$ . The remark 3.2 tells us that  $\lambda \in \mathcal{P}(n, m)$ . Now we arrive at the following lemma.

**Lemma 3.4.**  $\mathbb{Z}_m^n = \bigcup_{\lambda \in \mathcal{P}(n, m)} \mathcal{B}(\lambda)$ .

**Example 3.5.** As for  $0121212020102 \in \mathbb{Z}_3^{13}$ ,  $n_0 = 1$ , we have

$$\begin{aligned} \mathcal{O}(0121212020102) &= \mathcal{O}(012\ 12\ 120\ 201\ 02) = \mathcal{O}(012\ 120\ 201\ 012\ 2) \\ &= \mathcal{O}(012\ 120\ 2012\ 120) = \mathcal{O}(2012\ 120\ 201\ 120) \\ &= \mathcal{O}(2012\ 120120\ 201) = \mathcal{O}(012012\ 2012\ 201). \end{aligned}$$

012012 2012 201 is a (6, 4, 3)-element.

**Theorem 3.6.** *Let  $\lambda = (\lambda_1, \dots, \lambda_k) \in \mathcal{P}(n, m)$ . The number of orbits corresponding to the partition  $\lambda$  is  $\text{Perm}(\lambda)$ . If  $x$  is a  $\lambda$ -element, then  $\mathfrak{G}(x, x) = \mathbb{S}_1 \times \dots \times \mathbb{S}_k$ ,  $|\mathcal{O}(x)| = \frac{n!}{\lambda_1! \lambda_2! \dots \lambda_k!}$ .*

*Proof.* According to the lemma 3.3 and remark 3.2, the number of orbits corresponding to the partition  $\lambda$  is  $\text{Perm}(\lambda)$ .

Since  $x$  is a  $\lambda$ -element,  $(\mathbb{S}_{\lambda_1} \times \dots \times \mathbb{S}_{\lambda_k}) \cdot x = x$ . We have  $|\mathcal{O}(x)| \leq \frac{n!}{\lambda_1! \lambda_2! \dots \lambda_m!}$  according to Lagrange's theorem. The multinomial expansion

$$(x_1 + x_2 + \dots + x_m)^n = \sum_{a_1 + a_2 + \dots + a_m = n} \frac{n!}{\lambda_1! \lambda_2! \dots \lambda_m!} x_1^{a_1} x_2^{a_2} \dots x_m^{a_m}$$

implies that  $m^n = \sum_{a_1 + a_2 + \dots + a_m = n} \frac{n!}{\lambda_1! \lambda_2! \dots \lambda_m!} = \sum_{\lambda \in \mathcal{P}(n, m)} \frac{n!}{\lambda_1! \lambda_2! \dots \lambda_m!} \cdot \text{Perm}(\lambda)$ . So  $|\mathcal{O}(x)| = \frac{n!}{\lambda_1! \lambda_2! \dots \lambda_k!}$  follows from the result  $\mathbb{Z}_m^n = \bigcup_{\lambda \in \mathcal{P}(n, m)} \mathcal{B}(\lambda)$ . According to Lagrange's theorem,  $\mathfrak{G}(x, x) = \mathbb{S}_1 \times \dots \times \mathbb{S}_k$ .  $\square$

*Remark 3.7.*  $\mathcal{O}(x) \cong \text{Ind}_{\mathbb{S}_{\lambda_1} \times \dots \times \mathbb{S}_{\lambda_k}}^{\mathbb{S}_n} \mathbb{k} \cong M^\lambda$ , where  $M^\lambda$  is the permutation module of  $\mathbb{S}_n$  on  $\lambda$ -tabloids, see [Ful97].

**Example 3.8.** Set  $n_0 = 1$ , then  $\mathbb{Z}_3^4$  is the union of the following orbits.

$$\begin{aligned} \mathcal{O}(0120) &= \{0120\} \cong M^{(4)}, \\ \mathcal{O}(1201) &= \{1201\} \cong M^{(4)}, \quad \mathcal{O}(2012) = \{2012\} \cong M^{(4)}, \\ \mathcal{O}(0122) &= \{0122, 0110, 0020, 2120\} \cong M^{(3,1)}, \\ \mathcal{O}(0121) &= \{0121, 0100, 0220, 1120\} \cong M^{(3,1)}, \\ \mathcal{O}(1200) &\cong M^{(3,1)}, \quad \mathcal{O}(1202) \cong M^{(3,1)}, \quad \mathcal{O}(2011) \cong M^{(3,1)}, \quad \mathcal{O}(2010) \cong M^{(3,1)}, \\ \mathcal{O}(0101) &= \{0101, 0221, 1121, 0200, 1100, 1220\} = \mathcal{O}(1220) \cong M^{(2,2)}, \\ \mathcal{O}(0112) &= \{0112, 0022, 2122, 0010, 2110, 2020\} = \mathcal{O}(2020) \cong M^{(2,2)}, \\ \mathcal{O}(1212) &= \{1212, 1002, 2202, 1011, 2211, 2001\} = \mathcal{O}(2001) \cong M^{(2,2)}, \\ \mathcal{O}(0102) &= \{0210, 1110, 0000, 0222, 1020, 1122, 2100, 0021, 0102, 2220, \\ &\quad 2121, 0111\} = \mathcal{O}(0111) \cong M^{(2,1,1)}, \\ \mathcal{O}(2000) &= \{2102, 0002, 2222, 2111, 0212, 0011, 1022, 2210, 2021, 1112, \\ &\quad 1010, 2000\} = \mathcal{O}(2021) \cong M^{(2,1,1)}, \\ \mathcal{O}(1210) &= \{1021, 2221, 1111, 1000, 2101, 2200, 0211, 1102, 1210, 0001, \\ &\quad 0202, 1222\} = \mathcal{O}(1222) \cong M^{(2,1,1)}. \end{aligned}$$

The number of orbits is  $3 + 6 + 3 + 3 = \binom{4+3-1}{3-1}$ .

**3.2. The Nichols algebras of dimension  $n^m$ .** In this section, we prove the existence of Nichols algebras of dimension  $n^m$  for any  $m, n \geq 2$ . Firstly, we define a braided vector space  $W_{m,n_0}$  arising from the action of  $\mathbb{S}_n$  on  $\mathbb{Z}_m^n$ .

**Definition 3.9.** Let  $m \in \mathbb{Z}^{\geq 2}$  and  $n_0$  be a fixed integer in  $\mathbb{Z}_m$ .  $W_{m,n_0} = \bigoplus_{i \in \mathbb{Z}_m} \mathbb{k}w_i$  is a braided vector space with the braiding given by  $c(w_i \otimes w_j) = R_{ij}^{j-n_0, i+n_0} w_{j-n_0} \otimes w_{j+n_0}$ , where  $R_{i, i+n_0}^{i+i_0} = q$ ,  $R_{ij}^{j-n_0, i+n_0} \neq 0$  for all  $i, j \in \mathbb{Z}_m$ .

*Remark 3.10.* According to the braided equation, it's easy to see that All  $R_{i, i+n_0}^{i+i_0}$  equal for  $i \in \mathbb{Z}_m$ . The braiding of  $W_{m,n_0}$  is rigid according to [Sch92, Lemma 3.1.3].

**Theorem 3.11.** If  $q \in \mathbb{G}_n$  for  $n \geq 2$ , and  $R_{ij}^{j-n_0, i+n_0} R_{j-n_0, i+n_0}^{i, j} = 1$  for any  $i, j \in \mathbb{Z}_m$  with  $j \not\equiv i + n_0 \pmod{m}$ , then  $\dim \mathfrak{B}(W_{m,n_0}) = n^m$ .

*Proof.* Given a partition  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k) \in \mathcal{P}(n', m)$ , we take  $x = \chi_{a_1, b_1}^{\lambda_1} \cdots \chi_{a_k, b_k}^{\lambda_k}$  satisfying the conditions of formulas (3.1)-(3.4). Then  $\mathcal{O}(x)$  is corresponding to the partition  $\lambda$  and  $\mathfrak{S}_{n'}(w_x) = \mathfrak{S}_{n'}(w_y)$  for any  $y \in \mathcal{O}(x)$ . There exists a unique element  $\sigma_y \in \text{shuffle}(\lambda_1, \dots, \lambda_k)$  for any  $y \in \mathcal{O}(x)$  such that  $\sigma_y \cdot x = y$ . This implies that there exists a parameter  $\xi_y \in \mathbb{k}^\times$  such that  $\mathcal{T}_{\sigma_y}(w_x) = \xi_y w_y$ . We have

$$\mathfrak{S}_{n'}(w_x) = (\lambda_1)_q! (\lambda_2)_q! \cdots (\lambda_k)_q! \sum_{y \in \mathcal{O}_x} \xi_y w_y.$$

So  $w_x = 0$  iff  $\lambda_1 \geq n$ . In other words, every orbit contributes at most one dimension to the Nichols algebra and those vanish for corresponding to the partition  $\lambda = (\lambda_1, \dots)$  with  $\lambda_1 \geq n$ . In case  $n_0 = 0$ ,  $\dim \mathfrak{B}(W_{m,n_0}) = n^m$  since it is of Cartan type  $A_1 \times \cdots \times A_1$  ( $m$  copies). Notice that the dimension of the Nichols algebras does not rely on the parameter  $n_0$ , we finish the proof.  $\square$

*Remark 3.12.* It is obvious that  $\mathfrak{B}(W_{m,n_0})$  with  $n_0 \not\equiv 0$  is not of group type in general. The theorem generalizes a result in [Shi20b], which associated the Nichols algebras of square dimension with Pascal's triangle. The Nichols algebras of square dimension were appeared first in Andruskiewitsch and Giralaldi's work [AG18]. The Nichols algebras  $\mathfrak{B}(W_{2,1})$  have realizations in Yetter-Drinfeld category of the Suzuki Hopf algebra  $A_{Nn}^{\mu\lambda}$ , see [Shi20a] [Shi21] [Shi19].

**Corollary 3.13.** If  $q \notin \mathbb{G}_n$  for all  $n \geq 2$ , and  $R_{ij}^{j-n_0, i+n_0} R_{j-n_0, i+n_0}^{i, j} = 1$  for any  $i, j \in \mathbb{Z}_m$  with  $j \not\equiv i + n_0 \pmod{m}$ , then the Gelfand–Kirillov dimension (abbreviated GKdim) of  $\mathfrak{B}(W_{m,n_0})$  is  $m$ .

*Remark 3.14.* The result  $\text{GKdim} \mathfrak{B}(W_{2,1}) = 2$  was obtained first by Andruskiewitsch and Giralaldi [AG18].

*Proof.* Since  $w_x = w_y$  for any  $y \in \mathcal{O}(x) \subseteq \mathbb{Z}_m^n$ , we have

$$\mathrm{GKdim} \mathfrak{B}(W_{m,n_0}) = \lim_{n \rightarrow \infty} \log_n \sum_{k=0}^n \binom{k+m-1}{m-1} = m.$$

□

**Problem 3.15.** For those Nichols algebras of dimension  $n^m$ , is it possible to obtain  $\mathfrak{B}(W_{m,n_0})$  with  $n_0 \neq 0 \in \mathbb{Z}_m$  from  $\mathfrak{B}(W_{m,0})$  by some Hopf 2-cocycle deformation or Drinfeld twist?

*Remark 3.16.* As for the case of infinite dimension, Masuoka provided an example, see [Shi20a] [Mas00]. A negative information is that the Suzuki Hopf algebras  $A_{Nn}^{\mu\lambda}$  are not categorically Morita-equivalent to group algebras in general, for example  $A_{12}^{+-}$  [NN08, Section 5.2].

**Example 3.17.** [AG18] [Shi20a] [Shi21]  $W_{2,1} = \bigoplus_{i \in \mathbb{Z}_2} \mathbb{k}w_i$  is a braided vector space defined as

$$\begin{aligned} c(w_0 \otimes w_0) &= aw_1 \otimes w_1, & c(w_0 \otimes w_1) &= qw_0 \otimes w_1, \\ c(w_1 \otimes w_0) &= qw_1 \otimes w_0, & c(w_1 \otimes w_1) &= ew_0 \otimes w_0, \end{aligned}$$

where  $aqe \in \mathbb{k}^\times$ . If  $ae = q^2$ , then  $\mathfrak{B}(W_{2,1})$  is of diagonal type and

$$\dim \mathfrak{B}(W_{2,1}) = \begin{cases} 4, & q = -1, \quad (\text{Cartan type } A_1 \times A_1), \\ 27, & q^3 = 1 \neq q, \quad (\text{Cartan type } A_2), \\ \infty, & \text{otherwise.} \end{cases}$$

If  $q^2 \neq ae$ ,  $\mathfrak{B}(W_{2,1})$  is obviously not of group type,

$$(3.5) \quad \dim \mathfrak{B}(W_{2,1}) = \begin{cases} 4n, & q = -1, ae \in \mathbb{G}_n, \\ n^2, & ae = 1, q \in \mathbb{G}_n \text{ for } n \geq 2, \\ \infty, & q^2 = (ae)^{-1}, q \in \mathbb{G}_n \text{ for } n \geq 3, \\ \infty, & q \notin \mathbb{G}_n \text{ for } n \geq 2, \\ \text{unknown,} & \text{otherwise.} \end{cases}$$

**Example 3.18.**  $W_{3,1} = \bigoplus_{i \in \mathbb{Z}_3} \mathbb{k}w_i$  is a braided vector space defined as

$$\begin{aligned} c(w_0 \otimes w_0) &= aw_2 \otimes w_1, & c(w_0 \otimes w_1) &= qw_0 \otimes w_1, & c(w_0 \otimes w_2) &= dw_1 \otimes w_1, \\ c(w_1 \otimes w_0) &= ew_2 \otimes w_2, & c(w_1 \otimes w_1) &= fw_0 \otimes w_2, & c(w_1 \otimes w_2) &= qw_1 \otimes w_2, \\ c(w_2 \otimes w_0) &= qw_2 \otimes w_0, & c(w_2 \otimes w_1) &= \frac{df}{a}w_0 \otimes w_0, & c(w_2 \otimes w_2) &= \frac{df}{e}w_1 \otimes w_0, \end{aligned}$$

where  $adefq \in \mathbb{k}^\times$ . If  $df = 1$ ,  $q \in \mathbb{G}_n$  for  $n \geq 2$ , then

$$\begin{aligned} w_0^2 - aw_2w_1 &= 0, & w_0w_2 - dw_1^2 &= 0, & w_1w_0 - ew_2^2 &= 0, \\ w_iw_{i+1} \cdots w_{i+n-1} &= 0, & i &\in \mathbb{Z}_3. \end{aligned}$$

In this case,  $\dim \mathfrak{B}(W_{3,1}) = n^3$ .

**Example 3.19.**  $W_{4,1} = \bigoplus_{i \in \mathbb{Z}_4} \mathbb{k}w_i$  is a braided vector space defined as

$$\begin{aligned}
c(w_0 \otimes w_0) &= x_1 w_3 \otimes w_1, & c(w_0 \otimes w_1) &= q w_0 \otimes w_1, \\
c(w_0 \otimes w_2) &= x_2 w_1 \otimes w_1, & c(w_0 \otimes w_3) &= x_3 w_2 \otimes w_1, \\
c(w_1 \otimes w_0) &= x_4 w_3 \otimes w_2, & c(w_1 \otimes w_1) &= x_5 w_0 \otimes w_2, \\
c(w_1 \otimes w_2) &= q w_1 \otimes w_2, & c(w_1 \otimes w_3) &= x_6 w_2 \otimes w_2, \\
c(w_2 \otimes w_0) &= \frac{x_2 x_4}{x_3} w_3 \otimes w_3, & c(w_2 \otimes w_1) &= \frac{x_2 x_4 x_5}{x_1 x_6} w_0 \otimes w_3, \\
c(w_2 \otimes w_2) &= \frac{x_2 x_5}{x_6} w_1 \otimes w_3, & c(w_2 \otimes w_3) &= q w_2 \otimes w_3, \\
c(w_3 \otimes w_0) &= q w_3 \otimes w_0, & c(w_3 \otimes w_1) &= \frac{x_2 x_5}{x_1} w_0 \otimes w_0, \\
c(w_3 \otimes w_2) &= \frac{x_2 x_3 x_5}{x_1 x_6} w_1 \otimes w_0, & c(w_3 \otimes w_3) &= \frac{x_3 x_5}{x_4} w_2 \otimes w_0,
\end{aligned}$$

where  $q x_1 x_2 x_3 x_4 x_5 x_6 \in \mathbb{k}^\times$ . If  $x_2 x_5 = 1$ ,  $x_1 x_6 = x_2 x_3 x_4 x_5$ ,  $q \in \mathbb{G}_n$  for  $n \geq 2$ , then

$$\begin{aligned}
w_0^2 - x_1 w_3 w_1 &= 0, & w_0 w_2 - x_2 w_1^2 &= 0, & w_0 w_3 - x_3 w_2 w_1 &= 0, \\
w_1 w_0 - x_4 w_3 w_2 &= 0, & w_1 w_3 - x_6 w_2^2 &= 0, & w_2 w_0 - \frac{x_2 x_4}{x_3} w_3^2 &= 0, \\
w_i w_{i+1} \cdots w_{i+n-1} &= 0, & i \in \mathbb{Z}_4.
\end{aligned}$$

In this case,  $\dim \mathfrak{B}(W_{4,1}) = n^4$ .

**Example 3.20.**  $W_{4,2} = \bigoplus_{i \in \mathbb{Z}_4} \mathbb{k}w_i$  is a braided vector space defined as

$$\begin{aligned}
c(w_0 \otimes w_0) &= x_1 w_2 \otimes w_2, & c(w_0 \otimes w_1) &= x_2 w_3 \otimes w_2, \\
c(w_0 \otimes w_2) &= q w_0 \otimes w_2, & c(w_0 \otimes w_3) &= x_3 w_1 \otimes w_2, \\
c(w_1 \otimes w_0) &= x_4 w_2 \otimes w_3, & c(w_1 \otimes w_1) &= x_5 w_3 \otimes w_3, \\
c(w_1 \otimes w_2) &= x_6 w_0 \otimes w_3, & c(w_1 \otimes w_3) &= q w_1 \otimes w_3, \\
c(w_2 \otimes w_0) &= q w_2 \otimes w_0, & c(w_2 \otimes w_1) &= x_7 w_3 \otimes w_0, \\
c(w_2 \otimes w_2) &= \frac{x_1 x_7 x_9}{x_2 x_3} w_0 \otimes w_0, & c(w_2 \otimes w_3) &= x_9 w_1 \otimes w_0, \\
c(w_3 \otimes w_0) &= \frac{x_3 x_6}{x_7} w_2 \otimes w_1, & c(w_3 \otimes w_1) &= q w_3 \otimes w_1, \\
c(w_3 \otimes w_2) &= \frac{x_4 x_9}{x_2} w_0 \otimes w_1, & c(w_3 \otimes w_3) &= \frac{x_3 x_5 x_9}{x_2 x_7} w_1 \otimes w_1,
\end{aligned}$$

where  $q x_1 x_2 x_3 x_4 x_5 x_6 x_7 x_9 \in \mathbb{k}^\times$ . If  $x_1^2 x_7 x_9 = x_2 x_3$ ,  $x_4 x_9 = 1$ ,  $x_3 x_6 = 1$ ,  $x_2 x_7 = x_3 x_5^2 x_9$ ,  $q \in \mathbb{G}_n$  for  $n \geq 2$ , then

$$\begin{aligned}
w_0^2 - x_1 w_2^2 &= 0, & w_0 w_1 - x_2 w_3 w_2 &= 0, & w_0 w_3 - x_3 w_1 w_2 &= 0, \\
w_1 w_0 - x_4 w_2 w_3 &= 0, & w_2 w_1 - x_7 w_3 w_0 &= 0, & w_1^2 - x_5 w_3^2 &= 0,
\end{aligned}$$



$$w_i w_{i+2} \cdots w_{i+2(n-1)} = 0, \quad i \in \mathbb{Z}_4.$$

In this case,  $\dim \mathfrak{B}(W_{4,2}) = n^4$ .

#### ACKNOWLEDGEMENTS

This work was partially supported by Foundation of Jiangxi Educational Committee (No.12020447).

#### REFERENCES

- [AG18] N. Andruskiewitsch and J. M. J. Giraldi, *Nichols algebras that are quantum planes*, Linear and Multilinear Algebra **66** (2018), no. 5, 961–991. MR 3775317
- [AS02] N. Andruskiewitsch and H.-J. Schneider, *Pointed Hopf algebras*, New directions in Hopf algebras, Math. Sci. Res. Inst. Publ., vol. 43, Cambridge Univ. Press, Cambridge, 2002, pp. 1–68. MR 1913436
- [Ful97] W. Fulton, *Young tableaux*, London Mathematical Society Student Texts, vol. 35, Cambridge University Press, Cambridge, 1997, With applications to representation theory and geometry. MR 1464693
- [GS21] J. M. J. Giraldi and L. D. Silva, *Examples of Nichols algebras associated to upper triangular solutions of the Yang-Baxter equation in rank 3*, Comm. Algebra **49** (2021), no. 5, 2209–2233. MR 4232494
- [Hec06] I. Heckenberger, *The Weyl groupoid of a Nichols algebra of diagonal type*, Invent. Math. **164** (2006), no. 1, 175–188. MR 2207786
- [Hec09] ———, *Classification of arithmetic root systems*, Adv. Math. **220** (2009), no. 1, 59–124. MR 2462836
- [HS20] I. Heckenberger and H.-J. Schneider, *Hopf algebras and root systems*, Mathematical Surveys and Monographs, vol. 247, American Mathematical Society, Providence, RI, [2020] ©2020. MR 4164719
- [HV17a] I. Heckenberger and L. Vendramin, *A classification of Nichols algebras of semisimple Yetter-Drinfeld modules over non-abelian groups*, J. Eur. Math. Soc. (JEMS) **19** (2017), no. 2, 299–356. MR 3605018
- [HV17b] ———, *The classification of Nichols algebras over groups with finite root system of rank two*, J. Eur. Math. Soc. (JEMS) **19** (2017), no. 7, 1977–2017. MR 3656477
- [Lus93] G. Lusztig, *Introduction to quantum groups*, Progress in Mathematics, vol. 110, Birkhäuser Boston, Inc., Boston, MA, 1993. MR 1227098
- [Mas00] A. Masuoka, *Cocycle deformations and Galois objects for some cosemisimple Hopf algebras of finite dimension*, New trends in Hopf algebra theory (La Falda, 1999), Contemp. Math., vol. 267, Amer. Math. Soc., Providence, RI, 2000, pp. 195–214. MR 1800713
- [Nic78] W. D. Nichols, *Bialgebras of type one*, Comm. Algebra **6** (1978), no. 15, 1521–1552. MR 506406
- [NN08] D. Naidu and D. Nikshych, *Lagrangian subcategories and braided tensor equivalences of twisted quantum doubles of finite groups*, Comm. Math. Phys. **279** (2008), no. 3, 845–872. MR 2386730
- [Ros98] M. Rosso, *Quantum groups and quantum shuffles*, Invent. Math. **133** (1998), no. 2, 399–416. MR 1632802
- [Sch92] P. Schauenburg, *On coquasitriangular Hopf algebras and the quantum Yang-Baxter equation*, Algebra Berichte [Algebra Reports], vol. 67, Verlag Reinhard Fischer, Munich, 1992. MR 1623636

- [Sch96] ———, *A characterization of the Borel-like subalgebras of quantum enveloping algebras*, *Comm. Algebra* **24** (1996), no. 9, 2811–2823. MR 1396857
- [Shi19] Y.-X. Shi, *Finite-dimensional Hopf algebras over the Kac-Paljutkin algebra  $H_8$* , *Rev. Un. Mat. Argentina* **60** (2019), no. 1, 265–298. MR 3981570
- [Shi20a] ———, *Finite dimensional Nichols algebras over the Suzuki algebra I: simple Yetter-Drinfeld modules of  $A_{N2n}^{\mu\lambda}$* , arXiv:2011.14274 (2020).
- [Shi20b] ———, *The nichols algebras  $\mathfrak{B}(v_{abe})$  and a class of combinatorial numbers*, arXiv:2103.06489 (2020).
- [Shi21] ———, *Finite dimensional Nichols algebras over the Suzuki algebra II: simple Yetter-Drinfeld modules of  $A_{N2n+1}^{\mu\lambda}$* , arXiv:2103.06475 (2021).
- [Tak00] M. Takeuchi, *Survey of braided Hopf algebras*, *New trends in Hopf algebra theory* (La Falda, 1999), *Contemp. Math.*, vol. 267, Amer. Math. Soc., Providence, RI, 2000, pp. 301–323. MR 1800719
- [Wor89] S. L. Woronowicz, *Differential calculus on compact matrix pseudogroups (quantum groups)*, *Comm. Math. Phys.* **122** (1989), no. 1, 125–170. MR 994499 (90g:58010)

SCHOOL OF MATHEMATICS AND STATISTICS, JIANGXI NORMAL UNIVERSITY, NANCHANG 330022, P. R. CHINA

*Email address:* yxshi@jxnu.edu.cn