

WEAK AMENABILITY OF FREE PRODUCTS OF HYPERBOLIC AND AMENABLE GROUPS

IGNACIO VERGARA

ABSTRACT. We show that, if G is an amenable group and H is a hyperbolic group, then the free product $G * H$ is weakly amenable. A key ingredient in the proof is the fact that $G * H$ is orbit equivalent to $\mathbb{Z} * H$.

1. INTRODUCTION

Let G be a countable group. We say that G is weakly amenable if there exists a sequence of finitely supported functions $\varphi_n : G \rightarrow \mathbb{C}$ converging pointwise to 1, and a constant $C \geq 1$ such that

$$\sup_{n \in \mathbb{N}} \|\varphi_n\|_{B_2(G)} \leq C.$$

Here $B_2(G)$ stands for the space of Herz–Schur multipliers on G . For more details, see [2, Appendix D]. The Cowling–Haagerup constant $\Lambda(G)$ is the infimum of all $C \geq 1$ such that the condition above holds. If G is not weakly amenable, we set $\Lambda(G) = \infty$.

It is not known if the free product of weakly amenable groups is weakly amenable, but some particular cases have been studied. Bożejko and Picardello [1] showed that the free product of amenable groups is weakly amenable with Cowling–Haagerup constant 1. This result was improved by Ricard and Xu [11], who proved that, if $\Lambda(G) = \Lambda(H) = 1$, then $\Lambda(G * H) = 1$. In [5], Guentner, Reckwerdt and Tessera studied weak amenability for relatively hyperbolic groups. A particular case of their result says that, if H is hyperbolic and G has polynomial growth, then $G * H$ is weakly amenable. This result builds on Ozawa’s work [10], where he shows that all hyperbolic groups are weakly amenable.

Here we prove the following.

Proposition 1.1. *Let G be an amenable group and let H be a hyperbolic group. Then the free product $G * H$ is weakly amenable.*

The proof relies on the notion of orbit equivalence of group actions.

2. ORBIT EQUIVALENCE AND WEAK AMENABILITY

Let G and H be countable groups, and let $G \curvearrowright (X, \mu)$, $H \curvearrowright (Y, \nu)$ be probability measure preserving (p.m.p.) actions. We say that these actions are orbit equivalent if there exists a measure space isomorphism $T : (X, \mu) \rightarrow (Y, \nu)$ sending G -orbits onto H -orbits. We call T an orbit equivalence between $G \curvearrowright (X, \mu)$ and $H \curvearrowright (Y, \nu)$. We will write $G \sim_{OE} H$ if such actions and such an orbit equivalence exist. We refer the reader to [4, §2.2] for a detailed treatment of this subject.

Cowling and Zimmer [3] showed that, if $G \sim_{OE} H$, then $\Lambda(G) = \Lambda(H)$. This was later extended by Jolissaint [8] to measure equivalent groups. Recently, Ishan [6] extended this even further to the context of von Neumann equivalence of groups, which was introduced in [7].

A very important result concerning orbit equivalence of amenable groups was proved by Ornstein and Weiss [9].

Theorem 2.1 (Ornstein–Weiss). *Any two ergodic probability measure preserving actions of any two infinite countable, amenable groups are orbit equivalent.*

In particular, every countable, amenable group G satisfies $G \sim_{OE} \mathbb{Z}$.

Proof of Proposition 1.1. Since G is amenable, we have $G \sim_{OE} \mathbb{Z}$. This implies that $G * H \sim_{OE} \mathbb{Z} * H$; see [4, §3]. Hence

$$\Lambda(G * H) = \Lambda(\mathbb{Z} * H).$$

On the other hand, since H and \mathbb{Z} are both hyperbolic, so is $\mathbb{Z} * H$. Therefore, by [10], $\mathbb{Z} * H$ is weakly amenable, i.e., $\Lambda(\mathbb{Z} * H) < \infty$. This shows that

$$\Lambda(G * H) < \infty,$$

which finishes the proof. □

REFERENCES

- [1] Marek Bożejko and Massimo A. Picardello. Weakly amenable groups and amalgamated products. *Proc. Amer. Math. Soc.*, 117(4):1039–1046, 1993.
- [2] Nathaniel P. Brown and Narutaka Ozawa. *C*-algebras and finite-dimensional approximations*, volume 88 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2008.
- [3] Michael Cowling and Robert J. Zimmer. Actions of lattices in $\mathrm{Sp}(1, n)$. *Ergodic Theory Dynam. Systems*, 9(2):221–237, 1989.
- [4] Alex Furman. A survey of measured group theory. In *Geometry, rigidity, and group actions*, Chicago Lectures in Math., pages 296–374. Univ. Chicago Press, Chicago, IL, 2011.
- [5] Erik Guentner, Eric Reckwerdt, and Romain Tessera. Proper actions and weak amenability for classical relatively hyperbolic groups. *Preprint*.
- [6] Ishan Ishan. Von neumann equivalence and group approximation properties. *arXiv preprint arXiv:2107.11335*, 2021.
- [7] Ishan Ishan, Jesse Peterson, and Lauren Ruth. Von neumann equivalence and properly proximal groups. *arXiv preprint arXiv:1910.08682*, 2019.
- [8] Paul Jolissaint. Approximation properties for measure equivalent groups. *Preprint*, 2001.
- [9] Donald S. Ornstein and Benjamin Weiss. Ergodic theory of amenable group actions. I. The Rohlin lemma. *Bull. Amer. Math. Soc. (N.S.)*, 2(1):161–164, 1980.
- [10] Narutaka Ozawa. Weak amenability of hyperbolic groups. *Groups Geom. Dyn.*, 2(2):271–280, 2008.
- [11] Éric Ricard and Quanhua Xu. Khintchine type inequalities for reduced free products and applications. *J. Reine Angew. Math.*, 599:27–59, 2006.