

INSCRIBED TVERBERG-TYPE PARTITIONS FOR ORBIT POLYTOPES

STEVEN SIMON AND TOBIAS TIMOFEYEV

ABSTRACT. Tverberg’s theorem states that any set of $T(r, d) = (r - 1)(d + 1) + 1$ points in \mathbb{R}^d can be partitioned into r subsets whose convex hulls have non-empty r -fold intersection. Moreover, any generic collection of fewer points cannot be so divided. Extending earlier work of the first author, we show that in many such circumstances one can still guarantee inscribed “polytopal partitions” with specified symmetry conditions. Namely, for any faithful and full-dimensional d -dimensional orthogonal G -representation of a given group G of order r , we show that a generic set of $T(r, d) - d$ points in \mathbb{R}^d can be partitioned into r subsets so that there are r points, one from each of the resulting convex hulls, which are the vertices of a convex d -polytope whose isometry group contains G via a regular action afforded by the representation. As with Tverberg’s theorem, the number of points is tight for this. At one extreme, this gives polytopal partitions for all regular r -gons in the plane as well as for three of the six regular 4-polytopes in \mathbb{R}^4 . On the other hand, one has polytopal partitions for d -polytopes on r vertices with isometry group equal to G whenever G is the isometry group of a vertex-transitive d -polytope.

1. INTRODUCTION AND STATEMENT OF MAIN RESULTS

Tverberg’s theorem [28] establishes that any set of $T(r, d) = (r - 1)(d + 1) + 1$ points in \mathbb{R}^d can be partitioned into r subsets whose convex hulls have non-empty r -fold intersection. The result is of fundamental importance in discrete and convex geometry, with a panoply of rich generalizations and variants in geometric combinatorics and beyond. We refer the reader to the surveys [3, 5, 10] for the history of Tverberg’s theorem and its relatives, as well as a sampling of its many applications.

The Tverberg number $T(r, d)$ is “generically tight”, in that almost any collection of fewer points cannot be so partitioned. In lieu of a full Tverberg partition, it was asked in [22] whether one can nonetheless ensure “inscribed” polytopal variants in such circumstances. Namely, given a prescribed convex d -polytope $P(r, d)$ on r vertices, we say that a set of N points in \mathbb{R}^d can be $P(r, d)$ -partitioned if there exists a partition of the set into r subsets so that there are r points, one from each of the resulting convex hulls, which form the vertex set of $P(r, d)$.

Question 1. *What is the minimum number $N_{P(r, d)}$ for which almost any set of N points in \mathbb{R}^d can be $P(r, d)$ -partitioned by a polytope similar to $P(r, d)$?*

The main result of [22] gave tight results for all regular polygonal partitions in the plane.

Theorem 1.1. *Let $r \geq 3$ and let $P_r = P(r, 2)$ be a regular polygon. Then $N_{P_r} = 3r - 4$.*

1.1. Prescribed symmetries and orbit polytopes. The aim of the present paper is to extend Theorem 1.1 to d -polytopes with prescribed symmetry conditions. Observing that the cyclic group \mathbb{Z}_r acts regularly (freely and transitively) on the vertex set of a regular r -gon, our central result Theorem 1.2 affords polytopal partitions more generally for d -polytopes whose isometry group

contains an arbitrarily prescribed subgroup, and in fact one which acts regularly on the vertex set via a prescribed d -dimensional orthogonal representation.

Let $\rho: G \rightarrow O(d)$ be a faithful orthogonal representation of a finite group. An *orbit polytope*

$$P(G, u) = \text{Conv}(G \cdot u)$$

for ρ is simply the convex hull of the orbit of some $u \in \mathbb{R}^d$ under the action afforded by the representation. These polytopes are very well-studied (see, e.g. [1, 9, 14, 15, 24, 25]) particularly so for finite reflection groups [4, 6, 17, 18], and have origins dating back to the classical Wythoff construction for uniform polytopes and the study of Coxeter groups [8]. It is not difficult to see that the vertex set of each $P(G, u)$ is precisely the orbit $G \cdot u$, so that any orbit polytope is in particular vertex-transitive. Clearly, $G \leq \text{Iso}(P(G, u))$ via the action of the representation.

For our purposes, we shall consider those representations which are *full-dimensional*, that is, those for which there exists some orbit $G \cdot u$ whose affine hull is all of \mathbb{R}^d . Letting $\mathbb{R}[G] = \{\sum_{g \in G} r_g g \mid r_g \in \mathbb{R}\}$ denote the (left) regular representation and taking the module-theoretic perspective, it can be shown that a representation is full-dimensional precisely when it is isomorphic to a subrepresentation of $\mathbb{R}[G]$ which does not contain the trivial subrepresentation [14, Theorem 4.3]. Lastly, we shall say that $P(G, u)$ is a *free orbit polytope* if u has trivial stabilizer, so that the vertex set is identifiable with G under the action of the representation.

Theorem 1.2. *Let $\rho: G \rightarrow O(d)$ be a faithful and full-dimensional representation of a finite group G of order $r \geq 3$. Then almost any collection of $T(r, d) - d$ points in \mathbb{R}^d can be $P(r, d)$ -partitioned by the translate of a free orbit polytope for ρ . Moreover, such a partition fails for almost any collection of fewer points.*

1.2. Summary of Examples. We provide some special cases of Theorem 1.2, which in particular gives the upper bounds $N_{P(r,d)} \leq T(r, d) - d$ for a number of classical vertex-transitive d -polytopes. A detailed discussion is given in Section 4.

First, Theorem 1.1 is recovered by considering the standard representation of finite cyclic groups as rotations in the plane. These groups are the unique finite subgroups of the unit complex numbers; a similar analysis for the non-cyclic finite subgroups of the unit quaternions yields polytopal partitions for three of the six regular 4-polytopes. As discussed in Section 4.1, we expect each of the following estimates to exceed the exact value by 3.

Theorem 1.3. *Let $P_{\{3,3,4\}}$, $P_{\{3,4,3\}}$, and $P_{\{3,3,5\}}$ denote the regular 4-dimensional cross-polytope, regular 24-cell, and regular 600-cell, respectively. Then*

- (a) $N_{\{3,3,4\}} \leq 32$,
- (b) $N_{\{3,4,3\}} \leq 112$, and
- (c) $N_{\{3,3,5\}} \leq 592$.

In the examples above, the isometry group of the orbit polytope is larger than the group prescribed. Whether or not this is true in general depends on the group G as well as the given representation. For instance, $\text{Iso}(P(G, u)) > G$ for any free orbit polytope of any G -representation when G is (i) a finite abelian group with exponent greater than 2 or (ii) generalized dicyclic (see, e.g., [1, 23]). Moreover, a theorem of Babai [1] shows that a group is realizable as the isometry group of a vertex-transitive d -polytope precisely when it is not of the form (i) or (ii). Using results of [13, 14, 15], we are guaranteed that $\text{Iso}(P(G, u)) = G$ in Theorem 1.2 in these later cases.

Theorem 1.4. *Let $d \geq 2$. If G is the isometry group of a vertex transitive d -polytope, then almost any set of $T(|G|, d) - d$ points in \mathbb{R}^d can be $P(|G|, d)$ -partitioned by a vertex transitive d -polytope with $\text{Iso}(P(|G|, d)) = G$.*

Notable examples of Theorem 1.4 include polytopal partitions for all Coxeter permutahedra, that is, the free orbit polytopes determined by the finite reflection groups (Corollary 4.1). Considering dimension 3 along with other finite subgroups of the orthogonal group $O(3)$, we obtain polyhedral partitions for all right regular prisms and antiprisms, as well as for almost uniform copies of some Platonic and Archimedean solids (see Section 4.3 for a complete description).

Theorem 1.5. *Let D_n denote the Dihedral group and let S_n and A_n denote the symmetric and alternating groups on n elements, respectively.*

(a) $N_{Pr(n)} \leq 8n - 6$ for all $n \geq 3$, where $Pr(n)$ is a right regular prism with regular n -gon base; $\text{Iso}(Pr(n)) \cong D_n \times \mathbb{Z}_2$.

(b) $N_{APr(n)} \leq 8n - 6$ for all $n \geq 2$, where $APr(n)$ is a right regular antiprism with regular n -gon base; $\text{Iso}(APr(n)) \cong D_{2n}$.

(c) $N_{tO} \leq 90$, where tO is a truncated octahedron; $\text{Iso}(tO) \cong S_4$.

(d) $N_{tCO} \leq 186$, where tCO is a truncated cuboctahedron; $\text{Iso}(tCO) \cong S_4 \times \mathbb{Z}_2$.

(e) $N_{tID} \leq 474$, where tID is a truncated icosidodecahedron; $\text{Iso}(tID) \cong A_5 \times \mathbb{Z}_2$.

(f) $N_I \leq 38$, where I is an icosahedron; $\text{Iso}(I) \cong A_4$.

(g) $N_{sC} \leq 90$, where sC is a snub cube; $\text{Iso}(sC) \cong S_4$.

(h) $N_{sD} \leq 234$, where sD is a snub dodecahedron; $\text{Iso}(sD) \cong A_5$.

We note that $APr(2)$ is a tetrahedron whose faces are congruent isosceles triangles, while $APr(3)$ is an octahedron. For (d)–(e) and (g)–(h), the isometry groups match those given by their *uniform* versions [20, Chapter 11.5]. As discussed in Section 4.3, we expect the estimates of (a)–(b) and those of (c)–(h) to exceed the true values by 2 and 3, respectively.

Acknowledgements. The authors are grateful to Frieder Ladisch for very useful discussions, as well as to Florian Frick for helpful comments which improved the exposition of the manuscript.

2. AFFINE ORBIT PARTITIONS

Theorem 1.2 is a special case of a more general result on “affine orbit partitions” which holds for all representations that do not contain the trivial representation. The existence of such partitions and their optimality is given as Theorem 2.1, which by Lemmas 3.1 and 3.2 of the next section are shown to satisfy certain genericity conditions. Theorem 1.2 follows by considering faithful and full-dimensional representations. Theorem 4.3 below gives an example in the non-faithful setting.

In Tverberg-type theory, it is often convenient to recast the desired partition-type for a set of N points in \mathbb{R}^d as an equivalent statement in terms of affine linear maps $f: \Delta_{N-1} \rightarrow \mathbb{R}^d$ from an $(N - 1)$ -simplex to \mathbb{R}^d , that is, those which send any convex sum of vertices to the convex sums of their images. With this viewpoint, any partition of N points in \mathbb{R}^d into r subsets corresponds to r pairwise faces of the simplex, while the convex hulls arising from the partition corresponds to the images of the r faces under the mapping. For instance, a non-empty r -fold intersection of convex hulls as in Tverberg’s theorem corresponds to a non-empty r -fold intersection of the images of r pairwise disjoint faces.

A collection $\{x_1, \dots, x_r\}$ of r points from Δ_{N-1} are said to have pairwise disjoint support if there exist r vertex-disjoint faces $\sigma_1, \dots, \sigma_r$ of Δ_{N-1} with $x_j \in \sigma_j$ for all $1 \leq j \leq r$. If G is a group of order r , we parametrize each set $\{x_g\}_{g \in G} \subset \Delta_{N-1}$ of r points with pairwise disjoint support by the group G .

Theorem 2.1. *Let G be a finite group of order $r \geq 3$, let $\rho: G \rightarrow O(d)$ be an orthogonal representation which does not contain the trivial representation, and let $N = T(r, d) - d$. Then for any affine linear map $f: \Delta_{N-1} \rightarrow \mathbb{R}^d$, there exists a collection $\{x_g\}_{g \in G} \subset \Delta_{N-1}$ of r points with pairwise disjoint support and points a and u in \mathbb{R}^d such that*

$$f(x_g) = a + g \cdot u \quad (2.1)$$

for all $g \in G$. Moreover, such a partition fails for almost any affine linear map if $N < T(r, d) - d$.

Our proof of existence in Theorem 2.1 follows the usual Configuration–Space/Test–Map Paradigm of topological combinatorics, whereby the existence of the desired partition is equivalent to the existence of a zero of a naturally associated G -equivariant map. The domain of this mapping is the deleted r -fold join $(\Delta_{N-1})_{\Delta}^{*r}$ commonly used in Tverberg-type theory, while the codomain is a certain sub-representation of the $(d + 1)$ -fold product of the regular representation. That such a map must vanish follows from Sarkaria’s “Linear Borsuk–Ulam” Theorem [26] (given as Theorem 2.5 below), or equivalently the colorful Carathéodory Theorem of Bárány [2].

2.1. Configuration space. Let $N \geq 1$. Recall that the r -fold join Δ_{N-1}^{*r} consists of all formal convex sums $\lambda x := \lambda_1 x_1 + \dots + \lambda_r x_r \in \sigma_1 * \dots * \sigma_r$ from the join of any r faces of Δ_{N-1} , including the possibility of empty faces (here $\sum_{j=1}^r \lambda_j = 1$ and $\lambda_j \geq 0$ for all $1 \leq j \leq r$). The deleted r -fold join

$$(\Delta_{N-1})_{\Delta}^{*r} = \{\lambda x \in \sigma_1 * \dots * \sigma_r \mid x_i \in \sigma_i \text{ and } \sigma_i \cap \sigma_j = \emptyset \forall i \neq j\}$$

is the subcomplex of Δ_{N-1}^{*r} consisting of all points from the joins of pairwise disjoint faces. Given a group G of order r , we follow Sarkaria [26] and parametrize each r -tuple of disjoint faces by G and denote the resulting complex by $(\Delta_{N-1})_{\Delta}^{*G}$.

We shall consider the *right* group action on $(\Delta_{N-1})_{\Delta}^{*G}$ induced by right multiplication in G . Thus $\lambda x \cdot h = \sum_g \lambda_{gh} x_{gh}$ for each $\lambda x = \sum_g \lambda_g x_g \in (\Delta_{N-1})_{\Delta}^{*G}$. This action is free because the faces are pairwise disjoint. Parametrizing r disjoint copies of the vertex set $\{v_j\}_{j=1}^N$ of Δ_{N-1} by G , this simplicial complex can be identified with the N -fold join

$$G^{*N} = \left\{ \sum_{j=1}^N t_j v_j^{g_j} \mid g_j \in G \text{ and } \sum_{j=1}^N t_j = 1, t_j \geq 0 \forall 1 \leq j \leq N \right\}.$$

For this complex, we consider the *left* G -action given by $h \cdot \sum_{j=1}^N t_j v_j^{g_j} = \sum_{j=1}^N t_j v_j^{g_j h^{-1}}$ arising from right multiplication in G by inverses. Again, this action is free. An equivariant isomorphism $\iota: G^{*N} \cong (\Delta_{N-1})_{\Delta}^{*G}$ between the complexes is given by grouping, which, as in [22], has the following explicit description: For each $v = \sum_{j=1}^N t_j v_j^{g_j} \in G^{*N}$ and each $g \in G$, let

$$J_g(v) = \{j \mid g_j = g \text{ and } t_j > 0\}.$$

One then defines $\iota(v) = \sum_{g \in G} \lambda_g x_g$, where

$$\lambda_g = \sum_{j \in J_g(v)} t_j \quad \text{and} \quad x_g = \sum_{j \in J_g(v)} \frac{t_j}{\lambda_g} v_j$$

if $J_g \neq \emptyset$, while one sets $\lambda_g(v) = 0$ otherwise. In particular, observe that $\iota(v_j^{g_j}) = v_j$, where v_j occurs in the g_j -th coordinate of the deleted join. As $g = g_j h^{-1}$ if and only if $g_j = gh$, it is easily verified that ι preserves the respective G -actions.

2.2. Target space. For a finite group G , let $\mathbb{R}[G] = \{\sum r_g g \mid r_g \in \mathbb{R}\}$ denote the *right* regular representation, that is, with *left* G -action arising from multiplication on the right by inverses. We let $\mathbb{R}^d[G]$ denote the d -fold sum of $\mathbb{R}[G]$ equipped with the diagonal action. For convenience, we denote each $(\sum_{g \in G} r_1^g g, \dots, \sum_{g \in G} r_d^g g)$ in $\mathbb{R}^d[G]$ by $\sum_{g \in G} (r_1^g, \dots, r_d^g) \mathbf{e}_g$, so that

$$\mathbb{R}^d[G] = \left\{ \sum_g v_g \mathbf{e}_g \mid v_g \in \mathbb{R}^d \right\}.$$

As before, $h \cdot \mathbf{e}_g = \mathbf{e}_{gh^{-1}}$ for each $h, g \in G$, so that each $h \in G$ acts as a right permutation of elements of $\mathbb{R}^d[G]$, that is $h \cdot \sum_g v_g \mathbf{e}_g = \sum_g v_{gh} \mathbf{e}_g$.

Now let $\rho: G \rightarrow \mathbb{R}^d$ be any orthogonal representation. For notational purposes, we denote the representation space by V and let \cdot denote the resulting left action. In what follows, we will need to establish the dimension of a certain subrepresentation W of $\mathbb{R}[G]$. To that end, define $V_\rho = \{\sum_g (g \cdot v) \mathbf{e}_g \mid v \in \mathbb{R}^d\}$ and let

$$W_\rho = \left\{ \sum_g w_g \mathbf{e}_g \mid \sum_g g^{-1} \cdot w_g = 0 \right\}.$$

Proposition 2.2. V_ρ and W_ρ are complementary subrepresentations of $\mathbb{R}^d[G]$. Moreover, $V_\rho \cong V$ and so $\dim W_\rho = d(|G| - 1)$.

Proof. Clearly, both V_ρ and W_ρ are linear subspaces of $\mathbb{R}^d[G]$. To see that they are subrepresentations, first consider V_ρ . For any $h \in G$ have $h \cdot \sum_g (g \cdot v) \mathbf{e}_g = \sum_g (g \cdot v) \mathbf{e}_{gh^{-1}} = \sum_g (gh \cdot v) \mathbf{e}_g = \sum_g g \cdot (h \cdot v) \mathbf{e}_g$. Thus V_ρ is a subrepresentation of $\mathbb{R}^d[G]$ and in fact is isomorphic to V . As for W_ρ , we have $h \cdot \sum_g w_g \mathbf{e}_g = \sum_g w_{gh} \mathbf{e}_g$. Then $\sum_g g^{-1} \cdot w_{gh} = h \cdot \sum_g (gh)^{-1} \cdot w_{gh} = h \cdot \sum_g g^{-1} \cdot w_g = 0$, so again W_ρ is a subrepresentation of $\mathbb{R}^d[G]$.

Now we show that V_ρ and W_ρ are complementary. First, note that $\sum_g (g \cdot v) \mathbf{e}_g$ lies in W_ρ if and only if $|G|v = 0$, so $V_\rho \cap W_\rho = \{0\}$. On the other hand, suppose that $\sum_g v_g \mathbf{e}_g \in \mathbb{R}^d[G]$. Letting $a = \frac{1}{|G|} \sum_g g^{-1} \cdot v_g$, we have that $b := \sum_g (g \cdot a) \mathbf{e}_g$ lies in V_ρ . On the other hand, $\sum_g v_g \mathbf{e}_g - b$ lies in W_ρ because $\sum_g g^{-1} \cdot (v_g - g \cdot a) = \sum_g (g^{-1} \cdot v_g - a) = 0$. \square

Let $V_0 = \{\sum_{g \in G} v \mathbf{e}_g \mid v \in \mathbb{R}^d\}$ denote the d -fold sum of the trivial representation inside $\mathbb{R}^d[G]$ and let

$$W_0 = \left\{ \sum_{g \in G} w_g \mathbf{e}_g \mid \sum_g w_g = 0 \right\}$$

denote its complementary subrepresentation.

Proposition 2.3. *Suppose that V does not contain the trivial representation. Then*

- (i) $V_0 \cap V_\rho = \{0\}$, and
- (ii) $V' := V_0 + V_\rho$ and $W := W_0 \cap W_\rho$ are complementary subrepresentations of $\mathbb{R}^d[G]$. In particular, $\dim W = d(|G| - 2)$.

Proof. To see (i), suppose that there exist v_0 and v_1 in \mathbb{R}^d such that $g \cdot v_0 = v_1$ for all $g \in G$. Certainly $v_1 = v_0$, and so v_0 is G -invariant. Thus $v_0 = 0$, since V does not contain the trivial representation.

For (ii), we first show that the intersection of V' and W is trivial. Suppose that $v_0, v_1 \in V$ and let $w := \sum_g (v_0 + g \cdot v_1) \mathbf{e}_g \in V'$. We have $\sum_g g^{-1} \cdot (v_0 + g \cdot v_1) = \sum_g g^{-1} \cdot v_0 + |G|v_1 = |G|v_1$ because $\sum_g g^{-1} \cdot v_0$ is G -invariant. Thus $v_1 = 0$ if w lies in W_ρ . Similar reasoning shows that $\sum_g (v_0 + g \cdot v_1) = |G|v_0$, so that $v_0 = 0$ if $w \in W_0$. To conclude that $\mathbb{R}^d[G] = V' \oplus W$, we show that $\dim(V') + \dim W \geq d|G|$. But $\dim V' = 2d$, while $\dim W = \dim W_0 + \dim W_\rho - \dim(W_0 + W_\rho) \geq d(|G| - 1) + d(|G| - 1) - d|G| = d(|G| - 2)$ by Proposition 2.2. \square

2.3. Test map. As motivation for our test map, we first make a simple observation. Let $\rho: G \rightarrow O(d)$ be an orthogonal representation. Fixing an ordering of G , let $(\Delta_{N-1})_\Delta^{\times G}$ denote the $|G|$ -fold *deleted product* of Δ_{N-1} consisting of all $|G|$ -tuples $x = (x_g)_{g \in G}$ of points from Δ_{N-1} with pairwise disjoint support. For any $x \in (\Delta_{N-1})_\Delta^{\times G}$, we let

$$c_\rho(x) = \frac{1}{|G|} \sum_{g \in G} g^{-1} \cdot f(x_g).$$

In particular, c_ρ is simply the average

$$c_0(x) = \frac{1}{|G|} \sum_{g \in G} f(x_g)$$

when ρ is the trivial representation.

Proposition 2.4. *Suppose that $\rho: G \rightarrow O(d)$ does not contain the trivial representation and let $f: \Delta_{N-1} \rightarrow O(d)$ be an affine linear map. If there exist $\{x_g\}_{g \in G} \subset \Delta_{N-1}$ with pairwise disjoint support which satisfy Equation 2.1, then $a = c_0(x)$ and $u = c_\rho(x)$.*

Proof. Assuming Equation 2.1 holds, we have $|G|c_0(x) = a|G| + \sum_{g \in G} g \cdot u$. But $\sum_{g \in G} g \cdot u = 0$ by G -invariance, so $a = c_0(x)$. Likewise, $\sum_g g^{-1} \cdot f(x_g) = \sum_{g \in G} g^{-1} \cdot a + |G|u$. Again $\sum_{g \in G} g^{-1} \cdot a$ is G -invariant so $u = c_\rho(x)$. \square

Let $f: \Delta_{N-1} \rightarrow \mathbb{R}^d$ be an affine linear map and suppose that $\rho: G \rightarrow O(d)$ does not contain the trivial representation. For each $\lambda x = \sum_{g \in G} \lambda_g x_g \in (\Delta_{N-1})_\Delta^{*G}$, we let

$$c_\rho(\lambda x) = \frac{1}{|G|} \sum_{g \in G} \lambda_g g^{-1} \cdot f(x_g) \tag{2.2}$$

and

$$c_0(\lambda x) = \frac{1}{|G|} \sum_{g \in G} \lambda_g f(x_g).$$

Observe that the deleted product $(\Delta_{N-1})_\Delta^{\times G}$ can be identified as the subcomplex of the deleted join $(\Delta_{N-1})_\Delta^{*G}$ consisting of all λx with $\lambda_g = \frac{1}{|G|}$ for all g . In that case, we have $c_\rho(x) = |G|c_\rho(\lambda x)$ and $c_0(x) = |G|c_0(\lambda x)$ as well.

Define $F: (\Delta_{N-1})_{\Delta}^{*G} \rightarrow \mathbb{R}^d[G]$ by $F(\lambda x) = \sum_g F_g(\lambda x) \mathbf{e}_g$, where

$$F_g(\lambda x) = \lambda_g f(x_g) - c_0(\lambda x) - g \cdot c_\rho(\lambda x).$$

It follows that $F(\lambda x \cdot h) = \sum_g F_{gh}(\lambda x) \mathbf{e}_g = \sum_g F_g(\lambda x) \mathbf{e}_{gh^{-1}} = h \cdot F(\lambda x)$ for all $h \in G$, and therefore F is equivariant with respect to the given group actions.

It is straightforward to show that image of F actually lies in the representation $W = W_0 \cap W_\rho$ of Proposition 2.3. Namely, $\sum_{g \in G} g \cdot c_\rho(\lambda x) = 0$ because the sum is G -invariant, and hence $\sum_g F_g(\lambda x) = \sum_g \lambda_g f_g(x) - \sum_g c_0(\lambda x) - \sum_g g \cdot c_\rho(\lambda x) = |G|c_0(\lambda x) - |G|c_0(\lambda x) - 0 = 0$. Thus $F(\lambda x) \in W_0$. Likewise, $\sum_g g^{-1} \cdot F_g(\lambda x) = \sum_g \lambda_g g^{-1} \cdot f_g(x) - \sum_g g^{-1} \cdot c_0(\lambda x) - \sum_g c_\rho(\lambda x) = |G|c_\rho(\lambda x) - 0 - |G|c_\rho(\lambda x) = 0$ because $\sum_g g^{-1} \cdot c_0(\lambda x)$ is G -invariant. Thus $F(\lambda x) \in W_\rho$.

Finally, let $\mathbb{R}^\perp[G] = \{\sum_{g \in G} r_g g \mid \sum_{g \in G} r_g = 0\}$ denote the orthogonal complement of the right regular representation $\mathbb{R}[G]$. To preclude empty faces, we define $R: (\Delta_{N-1})_{\Delta}^{*G} \rightarrow \mathbb{R}[G]^\perp$ by

$$R(\lambda x) = \sum_{g \in G} \left(\lambda_g - \frac{1}{|G|} \right) g.$$

This map is also G -equivariant, so that

$$F \oplus R: (\Delta_{N-1})_{\Delta}^{*G} \rightarrow W \oplus \mathbb{R}^\perp[G]$$

is as well.

The existence of some $\{x_g\}_{g \in G} \subset \Delta_{N-1}$ with disjoint support that satisfies Equation 2.1 is equivalent to a zero of $F \oplus R$. Indeed, the vanishing of $F \oplus R$ means that $F_g(\lambda x) = 0$ and $\lambda_g = \frac{1}{|G|}$ for all $g \in G$, or equivalently that there exists some $x = (x_g)_{g \in G}$ in the deleted product $(\Delta_{N-1})_{\Delta}^{\times G}$ such that $f(x_g) = c_0(x) + g \cdot c_\rho(x)$ for all $g \in G$. Conversely, any $x \in (\Delta_{N-1})_{\Delta}^{\times G}$ satisfying Equation 2.1 corresponds to a zero of $F \oplus R$ by Proposition 2.4.

Let $\iota: G^{*N} \rightarrow (\Delta_{N-1})_{\Delta}^{*G}$ be the identification isomorphism above. A straightforward argument below will show that the composition

$$L := (F \oplus R) \circ \iota: G^{*N} \rightarrow W \oplus \mathbb{R}^\perp[G] \tag{2.3}$$

is affine linear, that is, that $L(\sum_{j=1}^N t_j v_j^{g_j}) = \sum_{j=1}^N t_j L(v_j^{g_j})$ for any $v = \sum_{j=1}^N t_j v_j^{g_j}$ in G^{*N} . Clearly, L is G -equivariant with respect to the actions considered. The representation $W \oplus \mathbb{R}^\perp[G]$ does not contain the trivial representation, and has dimension $T(|G|, d) - d - 1$ by Proposition 2.3. Letting $N = T(|G|, d) - d$, that our map vanishes is a consequence of the ‘‘Linear Borsuk–Ulam’’ Theorem [26, Theorem 2.4] of Sarkaria:

Theorem 2.5. *Let U be a real $(N - 1)$ -dimensional linear representation of G which does not contain the trivial representation. If $L: G^{*N} \rightarrow U$ is an affine linear G -equivariant map, then there exists some $v \in G^{*N}$ such that $L(v) = 0$.*

To prove that the mapping L of Equation 2.3 is affine linear, we show that both $F \circ \iota$ and $R \circ \iota$ are. The argument for the later was given in the proof of Theorem 3.1 from [22]. For the former, it suffices to prove the affine linearity of $F_1 \circ \iota$ and $F_2 \circ \iota$, where $F_1(\lambda x) = \sum_g \lambda_g f(x_g) \mathbf{e}_g$ and $F_2(\lambda x) = \sum_g \lambda_g g^{-1} \cdot f(x_g) \mathbf{e}_g$. We show the proof for F_1 ; the proof for F_2 is nearly identical. Consider $v = \sum_{j=1}^N t_j v_j^{g_j} \in G^{*N}$. We have $(F_1 \circ \iota)(v_j^{g_j}) = f(v_j) \mathbf{e}_{g_j}$, so $(F_1 \circ \iota)(v) = \sum_g \lambda_g f(x_g) \mathbf{e}_g = \sum_{g \in G} \lambda_g [\sum_{j \in J_g(v)} \frac{t_j}{\lambda_g} f(v_j)] \mathbf{e}_g = \sum_{j=1}^N t_j f(v_j) \mathbf{e}_{g_j} = \sum_{j=1}^N t_j (F_1 \circ \iota)(v_j^{g_j})$ by affine linearity of f .

2.4. Optimality. To complete the proof of Theorem 2.1, we show that an affine orbit partition fails for almost any affine linear map $f: \Delta_{N-1} \rightarrow \mathbb{R}^d$ if $N < T(|G|, d) - d$. This follows from a standard (co-)dimension counting argument.

Let $N = T(r, d) - d - 1$. Given any affine linear map $f: \Delta_{N-1} \rightarrow \mathbb{R}^d$, we have seen that the existence of some collection $\{x_g\}_{g \in G} \subset \Delta_{N-1}$ with pairwise support which satisfies Equation 2.1 is equivalent to a zero of the map $L: G^{*N} \rightarrow W \oplus \mathbb{R}^\perp[G]$ of Equation 2.3. Let $\Sigma_{N-1} = \{s = (s_1, \dots, s_N) \in \mathbb{R}^N \mid \sum_{j=1}^N s_j = 1 \text{ and } s_j \geq 0 \text{ for all } j\}$ denote the standard $(N-1)$ -simplex in \mathbb{R}^N . If the images $f(v_j)$ of the vertices of Δ_{N-1} are generic, we show that any $v = \sum_{j=1}^N t_j v_j^{g_j}$ satisfying $L(v) = 0$ forces $t = (t_1, \dots, t_N)$ in Σ_{N-1} to satisfy N linearly independent conditions, a contradiction.

Consider $(g_1, \dots, g_N) \in G^{\times N}$ for $v = \sum_{j=1}^N t_j v_j^{g_j}$ above. It follows that $J_g := \{j \mid g_j = j\} \neq \emptyset$ for each $g \in G$. That $(R \circ i)(v) = 0$ means that $\sum_{j \in J_g} t_j = \frac{1}{|G|}$ for all $g \in G$, thereby giving $(|G| - 1)$ linearly independent conditions on t . Secondly, $(F \circ \iota)(v) = 0$ means that t is zero of each of the linear maps

$$M_g(s) = \sum_{j \in J_g} s_j f(v_j) - \sum_{j=1}^N \frac{s_j}{|G|} f(v_j) - g \cdot \sum_{j=1}^N \frac{s_j}{|G|} g_j^{-1} \cdot f(v_j)$$

defined on Σ_{N-1} . As $|G| \geq 3$, $N-1 = (|G|-2)(d+1) > d$, and so for any particular $g \in G$ one has d linearly independent conditions provided the $f(v_j)$ are chosen generically. On the other hand, the image of $M(s) = \sum_{g \in G} M_g(s) e_g$ lies in the subspace W of $\mathbb{R}^d[G]$ of Proposition 2.3. For generic $f(v_j)$, it follows from Proposition 2.3 that $M(t) = 0$ determines $d(|G| - 2)$ linearly independent conditions. Moreover, these conditions are independent from those given by $(R \circ \iota)(v) = 0$, again provided the $f(v_j)$ are generic. One therefore has N independent conditions on t in total.

3. GENERICITY CONDITIONS

We now show that the affine orbits $a + G \cdot u$ guaranteed by Theorem 2.1 satisfy two generic conditions (see, e.g., [15, Lemma 2.11] and [14, Lemma 4.2]), namely that u has minimal stabilizer and that $\text{Aff}(G \cdot u)$ has maximum dimension.

Lemma 3.1. *Let G be a finite group of order $r \geq 3$ and let $\rho: G \rightarrow O(d)$ be an orthogonal representation which does not contain the trivial representation. Let $K = \ker \rho$ and let $m = \max\{\dim \text{Aff}(G \cdot u) \mid u \in \mathbb{R}^d\}$. If $N = T(r, d) - d$, then the following is true for almost any affine linear map $f: \Delta_{N-1} \rightarrow \mathbb{R}^d$: If $\{x_g\}_{g \in G} \subset \Delta_{N-1}$ are points with pairwise disjoint support that satisfy Equation 2.1, then (i) $\text{Stab}(u) = K$ and (ii) $\dim \text{Aff}(G \cdot u) = m$.*

As with the proof of optimality in Theorem 2.1, Lemma 3.1 follows from a degrees of freedom argument. Note that (i) of Lemma 3.1 means that $|G \cdot u| = |G|$ if the representation is faithful, so that $P(G, u) = \text{Conv}(G \cdot u)$ is a free orbit polytope. On the other hand, (ii) $m = d$ if the representation is full-dimensional. Thus Theorem 2.1 together with Lemma 3.1 completes the proof of Theorem 1.2.

Proof of Lemma 3.1. Let $f: \Delta_{N-1} \rightarrow \mathbb{R}^d$ be affine linear and let $L: G^{*N} \rightarrow W \oplus \mathbb{R}^\perp[G]$ be the associated map given by Equation 2.3. It follows from Proposition 2.4, Equation 2.2, and the affine linearity of f that any u satisfying Equation 2.1 is of the form $u = |G|c_\rho(\iota(v)) = \sum_{j=1}^N t_j g_j^{-1} \cdot f(v_j)$,

where $v = \sum_{j=1}^N t_j v_j^{g_j}$ satisfies $L(v) = 0$. If f is generic, we have seen in the proof of Lemma 3.1 that corresponding to this $(g_1, \dots, g_N) \in G^{\times N}$ one has $(N - 1)$ linearly independent conditions on $t = (t_1, \dots, t_N)$ in Σ_{N-1} .

For each $s \in \Sigma_{N-1}$, let $u(s) = \sum_{j=1}^N s_j g_j^{-1} \cdot f(v_j)$. For any $h \in G - K$, consider the linear map $L_h(s) = u(s) - h \cdot u(s)$ defined on Σ_{N-1} . If the $g_j^{-1} \cdot f(v_j)$ are chosen to be in general position, then L_h is not the zero map because h is not in K . Moreover, so long as the $f(v_j)$ are generic, the condition that $L_h(t) = 0$ is independent from those determined by $L(v) = 0$. Thus u is not fixed by any $h \in G - K$, as otherwise t satisfies N independent conditions.

For maximum dimensionality of $\text{Aff}(G \cdot u)$, let m be as in the statement of Lemma 3.1. As in [14, Lemma 4.2], let $|G| = r$, fix an ordering g_1, \dots, g_r of G , and consider the $(d + 1) \times r$ matrix

$$A(s) = \begin{pmatrix} g_1 \cdot u(s) & \cdots & g_r \cdot u(s) \\ 1 & \cdots & 1 \end{pmatrix}$$

defined on Σ_{N-1} . Each $m \times m$ sub-determinant of $A(s)$ determines a polynomial $P(s)$. For generic $f(v_j)$, each of these polynomials is non-zero and the condition that $P(t) = 0$ is independent from the linear conditions determined by $L(v) = 0$. Thus $\dim \text{Aff}(G \cdot u) = m$, as otherwise $P(t) = 0$ for each of these polynomials and t would satisfy at least N independent conditions. \square

3.1. Generic Isometries. For the proof of Theorem 1.4 and the calculation of some of the isometry groups in Theorem 1.5, we will need to show that the translated orbit polytopes of Theorem 1.2 satisfy a minimal symmetry condition. To state this formally, we follow [14, 15], which introduced the notion of a generic linear symmetry group of a linear representation. An analogous group in the orthogonal setting was given in [13, Chapter 7].

Suppose that $\rho: G \rightarrow O(d)$ is a full-dimensional orthogonal representation. By definition, the “generating set”

$$\text{Gens}(G) := \{u \in \mathbb{R}^d \mid \text{Aff}(G \cdot u) = \mathbb{R}^d\}$$

is non-empty. Reasoning as in the proof of (ii) of Lemma 3.1, it can be shown that those u which do not lie in $\text{Gens}(G)$ form a proper algebraic subset of \mathbb{R}^d . In particular, $\text{Gens}(G)$ is a dense open set [14, Corollary 4.5].

Let S_G denote the set of all permutations of G . For any $u \in \mathbb{R}^d$, let

$$\text{OSym}(G, u) = \{\pi \in S_G \mid \exists A \in O(d) \text{ such that } A(g \cdot u) = \pi(g) \cdot u \text{ for all } g \in G\}.$$

If $u \in \text{Gens}(G)$, it is easily seen that any $A \in O(d)$ above is uniquely determined by the permutation π , and so that $\text{OSym}(G, u)$ can be identified with $\text{Iso}(P(G, u))$. Note that for each $g \in G$, the permutation π_g given by left multiplication by g corresponds to ρ_g under this identification. The *generic isometry group* is defined by

$$\text{OSym}(G) = \bigcap_{u \in \text{Gens}(G)} \text{OSym}(G, u).$$

For $u \in \text{Gens}(G)$, to say that $\text{OSym}(G, u) = \text{OSym}(G)$ therefore means that the d -dimensional orbit polytope $P(G, u)$ has minimal isometry group.

Considering the standard inner product on \mathbb{R}^d , an easy observation communicated to the authors by Frieder Ladisch shows that $\pi \in \text{OSym}(G, u)$ if and only if u is a zero of the polynomial

$$P_{g,h}^\pi(u) = \langle g \cdot u, h \cdot u \rangle - \langle \pi(g) \cdot u, \pi(h) \cdot u \rangle$$

for all distinct $g, h \in G$. If $\pi \in \text{OSym}(G)$, then each $P_{g,h}^\pi$ vanishes on $\text{Gens}(G)$, a non-empty open set in \mathbb{R}^d . Therefore $P_{g,h}^\pi(u)$ is the zero polynomial and $\text{OSym}(G) \leq \text{OSym}(G, u)$ for any $u \in \mathbb{R}^d$. If $\pi \in S_G - \text{OSym}(G)$, on the other hand, then by definition there is some $u \in \text{Gens}(G)$ with $\pi \notin \text{OSym}(G, u)$ and so at least one of the $P_{g,h}^\pi(u)$ is non-zero. In particular, the set of all $u \in \mathbb{R}^d$ with $\text{OSym}(G, u) > \text{OSym}(G)$ is a proper algebraic subset of \mathbb{R}^d .

Lemma 3.2. *Let G be a finite group of order $r \geq 3$, let $\rho: G \rightarrow O(d)$ be a faithful and full-dimensional representation, and let $N = T(r, d) - d$. Then the following holds for almost any affine linear map $f: \Delta_{N-1} \rightarrow \mathbb{R}^d$: If $\{x_g\}_{g \in G} \subset \Delta_{N-1}$ are points with pairwise disjoint support that satisfy Equation 2.1, then $\text{OSym}(G, u) = \text{OSym}(G)$.*

Proof of Lemma 3.2. For an affine linear map $f: \Delta_{N-1} \rightarrow \mathbb{R}^d$, again let $L: G^{*N} \rightarrow W \oplus \mathbb{R}^1[G]$ be the linear map given by Equation 2.3 and consider $u = \sum_{j=1}^N t_j g_j^{-1} \cdot f(v_j)$ where $v = \sum_{j=1}^N t_j v_j^{g_j}$ satisfies $L(v) = 0$. As before, let $u(s) = \sum_{j=1}^N s_j g_j^{-1} \cdot f(v_j)$ for $s \in \Sigma_{N-1}$. For each $\pi \in S_G$ and any distinct $g, h \in G$, consider the polynomial $P_{g,h}(s) = P_{g,h}^\pi(u(s))$. If the $g_j^{-1} \cdot f(v_j)$ are chosen to be in general position, then $P_{g,h}(s)$ is the zero polynomial exactly when $P_{g,h}^\pi(u)$ is. By the observations above, there is some $P_{g,h}^\pi$ which is not the zero polynomial precisely when $\pi \in S_G - \text{OSym}(G)$. For such $P_{g,h}^\pi$ and generic $f(v_j)$, the condition determined by $P_{g,h}(t) = 0$ is independent from the linear conditions determined by $L(v) = 0$. Thus $\pi \notin \text{OSym}(G, u)$ for any $\pi \notin \text{OSym}(G)$, as otherwise $t \in \Sigma_{N-1}$ satisfies N independent conditions. \square

To prove Theorem 1.4, we now only need to appeal to a result of [14]. As there, we shall say that a faithful and full-dimensional representation $\rho: G \rightarrow O(d)$ is (orthogonally) *closed* if $\text{OSym}(G)$ consists only of those permutations determined by left multiplication in G . Thus a closed representation means precisely that $\text{Iso}(P(G, u)) = G$ for all $u \in \text{Gens}(G)$.

By replacing orthogonal representations with arbitrary linear ones and the orthogonal group $O(d)$ with the general linear group $GL(\mathbb{R}^d)$, one has the groups $\text{LinSym}(G, u)$ and $\text{LinSym}(G)$ of [14] analogous to $\text{OSym}(G, u)$ and $\text{OSym}(G)$ above. One likewise has an analogous definition of linear closure. Theorem A of [14] shows that the inclusion representation $G \hookrightarrow GL(\mathbb{R}^d)$ is linearly closed whenever $G \leq GL(\mathbb{R}^d)$ is the affine linear symmetry group of a d -dimensional vertex transitive polytope. Of course, linear closure of an orthogonal representation implies orthogonal closure of the representation. Thus if $G < O(d)$ is the isometry group of a vertex-transitive d -polytope, the inclusion representation is automatically (orthogonally) closed. The proof of Theorem 1.4 follows immediately.

Proof of Theorem 1.4. Let $P(r, d) = a + \text{Conv}(G \cdot u)$ be the d -dimensional translated free orbit polytope guaranteed by Theorem 1.2. By Lemma 3.2, we have $\text{OSym}(G, u) = \text{OSym}(G)$. As the inclusion $G \hookrightarrow O(d)$ is closed, $\text{OSym}(G)$ is given by left multiplication and therefore $\text{Iso}(P(G, u)) = G$. \square

Another class of linearly closed representations $\rho: G \rightarrow GL(\mathbb{R}^d)$ are those which are *absolutely irreducible* [14, Theorem 5.5], that is, irreducible when viewed as a representation over the complex numbers. In particular, any absolutely irreducible orthogonal representation $\rho: G \rightarrow O(d)$ must be (orthogonally) closed. Any real irreducible representation is full-dimensional, and in fact $\text{Gens}(G) = \mathbb{R}^d - \{0\}$ because the linear subspace spanned by any orbit is a G -invariant. Note also that $|G| \geq 3$ for any (real) irreducible representation $\rho: G \rightarrow O(d)$ if $d \geq 2$.

Corollary 3.3. *Let $d \geq 2$ and let $\rho: G \rightarrow O(d)$ be a faithful and absolutely irreducible representation. Then almost any set of $T(|G|, d) - d$ points in \mathbb{R}^d can be $P(|G|, d)$ -partitioned by a vertex-transitive polytope with $\text{Iso}(P(|G|, d)) = G$.*

4. EXAMPLES

4.1. Finite Subgroups of Spheres. Let \mathbb{F} be either the field \mathbb{C} of complex numbers or the skew field \mathbb{H} of quaternions. The corresponding unit spheres $S(\mathbb{F})$ form groups under \mathbb{F} -multiplication. Left multiplication by any $u \in S(\mathbb{F})$ defines an \mathbb{F} -isometry, so that $S(\mathbb{C}) = S^1 = U(1)$ and $S(\mathbb{H}) = S^3$ are the 1-dimensional unitary and symplectic groups, respectively.

For any finite subgroup G of $S(\mathbb{F})$, left multiplication of \mathbb{F} by G defines a faithful representation, and in fact $\text{Stab}(u)$ is trivial for all non-zero $u \in \mathbb{F}$. The corresponding affine free orbit polytopes are thus the translates of $\text{Conv}(G)u$ for any $u \neq 0$, or equivalently the translated and scaled \mathbb{F} -unitary copies of $\text{Conv}(G)$. We therefore expect the upper bound $N_{P(G)} \leq (|G| - 2)(d + 1) + 2$ given by Theorem 1.2 for $P(G) = \text{Conv}(G)$ to exceed the exact value by $\dim(O(d)) - \dim S(\mathbb{F})$. This will be confirmed below when $\mathbb{F} = \mathbb{C}$ (in which case the difference is zero) and is left as a conjecture when $\mathbb{F} = \mathbb{H}$ (in which case the difference is 3).

4.1.1. Regular Polygons. The non-trivial finite subgroups $G < S^1$ are precisely the cyclic groups \mathbb{Z}_r realized as the r -th roots of unity. Letting $\omega_r = e^{2\pi i/r}$ denote the standard r -th root of unity, \mathbb{Z}_r is the vertex set $\{\omega_r^k \mid 0 \leq k < r\}$ of the regular r -gon $\text{Conv}(\mathbb{Z}_r)$ inscribed in the unit circle when $r \geq 3$. In particular, the inclusion $\mathbb{Z}_r \hookrightarrow U(1) \cong SO(2)$ is full-dimensional when $r \geq 3$ (and in fact irreducible). It is easily observed that any regular r -gon in the plane is the translate of some $P(\mathbb{Z}_r, u)$ with $u \neq 0$, so that Theorem 1.2 recovers Theorem 1.1 when $r \geq 3$.

4.1.2. Regular 4-polytopes. Proceeding analogously, we consider the finite subgroups of S^3 . It is a classical fact that these are the binary dihedral groups D_r^* for all $r \geq 2$ and the binary polyhedral groups T^*, O^* , and I^* , the pullbacks under the canonical 2-fold covering homomorphism $\phi: SU(2) \rightarrow SO(3)$ of the Dihedral groups D_r (viewing D_2 as $\mathbb{Z}_2 \times \mathbb{Z}_2$) and the rotation groups T, O and I of the regular tetrahedron, octahedron, and icosahedron, respectively. Each inclusion $G \hookrightarrow Sp(1) \cong SU(2) \subset SO(4)$ is irreducible and so is in particular full-dimensional.

Viewing \mathbb{Z}_r as the r -th roots of unity in \mathbb{C} as before, one has the explicit decomposition $D_r^* = \mathbb{Z}_{2r} \cup \mathbb{Z}_{2r}j \subset \mathbb{H}$. Thus $\text{Conv}(D_r^*) = P_{2r} \oplus P_{2r}j$ is the join of two congruent regular $2r$ -gons lying in orthogonal planes in \mathbb{R}^4 which intersect at the origin. In particular, one has the Quaternion group $Q_8 = \{\pm 1, \pm i, \pm j, \pm k\}$ when $r = 2$. Thus $\text{Conv}(Q_8)$ is the regular 4-dimensional cross-polytope, which gives part (a) of Theorem 1.3. When $r \geq 3$, the $P(D_r^*)$ are vertex and facet transitive “fusils” [20] with $\text{Iso}(P(D_r^*)) \cong D_{2r} \times D_{2r} \times \mathbb{Z}_2$, whose tetrahedral facets are composed of congruent isosceles triangle faces. For parts (b) and (c) of Theorem 1.3, we note that the T^* consists of the 24 units of the Hurwitz Quaternion ring, and these are the vertices of the standard regular 24-cell. Likewise, the 120 vertices of I^* are the vertex set of a regular 600-cell (see, e.g. [11, Chapter 4]).

4.2. Coxeter Permutahedra and Alternahedra. Recall that a d -dimensional finite reflection group is a finite subgroup of $O(d)$ which is generated by reflections across linear hyperplanes. A finite reflection group is called *irreducible* if the inclusion representation is. The irreducible finite reflection groups are precisely the isometry groups of certain uniform polytopes, including those

of all regular polytopes, while all other finite reflection groups are the products of these (see, e.g., [19, Chapter 2]).

It is easily seen that an orbit polytope $P(G, u)$ of a finite reflection group is free precisely when u is not contained in any linear hyperplane determined by a reflection of G . The free orbit polytopes are commonly called Coxeter permutahedra, with the classical permutahedra (see, e.g., [29]) arising from the standard representation of the symmetric group. All Coxeter permutahedra for a given reflection group are combinatorially equivalent (see, e.g., [17, Proposition 2.1], which need not be true for free orbit polytopes of a given representation in general (see, e.g., [24]).

Owing to the classification of finite reflection groups, it follows quickly that $\text{Iso}(P(G, u)) = G$ for the the Coxeter permutahedra guaranteed by Theorem 1.2. For irreducible groups, this is an immediate consequence of Theorem 1.4. For a direct product $G = \prod_{j=1}^k G_j$ of irreducible d_j -dimensional reflection groups, one has $P(G, u) = \prod_{j=1}^k P(G_j, u_j)$, for any $u = (u_1, \dots, u_k) \in \bigoplus_{j=1}^k \mathbb{R}^{d_j}$. If each u_j is non-zero, then each $P(G_j, u_j)$ is full dimensional and therefore so is $P(G, u)$. On the other hand, $\text{Iso}(P(G, u)) = \prod_{j=1}^k \text{Iso}(P(G_j, u_j)) = \prod_{j=1}^k G_j = G$ for a generic choice of u . Thus the result again follows from Theorem 1.4.

Corollary 4.1. *Let $d \geq 2$ and let $G < O(d)$ be a finite reflection group. Then almost any set of $T(|G|, d) - d$ points in \mathbb{R}^d can be $P(|G|, d)$ -partitioned by a Coxeter permutahedra with isometry group G .*

For irreducible d -dimensional reflection groups, we note that the isometry groups of the Coxeter permutahedra of Corollary 4.1 matches those given by *any* permutahedra, even uniform ones, so long as $d \geq 3$ and G is neither the isometry group S_d of the $(d - 1)$ -simplex nor the exceptional groups F_4 or E_6 [20, Chapter 11.6].

Given a finite reflection group G , we also consider its index two rotational subgroup G^+ , whose corresponding free orbit polytopes we shall call Coxeter alternahedra (see [9] for when $G = S_n$ is the symmetric group). Provided one takes $u \in \mathbb{R}^d$ to be trivially stabilized by the full reflection group G , the resulting alternahedron is obtained by “alternation” [7, Chapter 8.6] of the Coxeter permutahedron, that is, by taking the convex hull of those vertices lying on the permutahedron which are indexed by the subgroup G^+ . An easy adaption of the the proof of Lemma 3.1 shows that one can ensure that u has trivial stabilizer with respect to all the full group G and not just G^+ , so that the alternahedra guaranteed by Theorem 1.2 are always alternated permutahedra.

The isometry groups in (f)–(h) of Theorem 1.5 are given by the following proposition:

Proposition 4.2. *Let $d \geq 3$ and let $G^+ < SO(d)$ be the rotational subgroup of an irreducible finite reflection group $G < O(d)$. Then almost any set of $T(|G^+|, d) - d$ points in \mathbb{R}^d can be $P(|G^+|, d)$ -partitioned by a Coxeter alternahedra with with isometry group G^+ .*

We note that Proposition 4.2 is false when $d = 2$, as follows by considering the Dihedral group D_r when $r \geq 3$, in which case $D_r^+ = \mathbb{Z}_r$ and the corresponding alternahedra are regular r -gons.

Proof of Proposition 4.2. Let $\rho: G \hookrightarrow O(d)$ be the inclusion and let $\rho^+: G^+ \hookrightarrow SO(d)$ be its restriction to G^+ . We show that ρ^+ is absolutely irreducible and appeal to Proposition 3.3. The proof is a basic character theory argument along the lines of [16, Proposition 5.1].

Viewing ρ and ρ^+ as complex representations, denote their characters by χ and χ^+ , respectively. As shown in [12], ρ is in fact complex irreducible, so that $\langle \chi, \chi \rangle = |G|$. Thus $2|G|^+ =$

$\langle \chi, \chi \rangle = \sum_{g \in G} |\chi(g)|^2 = \sum_{r \in G^+} |\chi^+(r)|^2 + \sum_{s \notin G^+} |\chi(s)|^2$. Expressing $\chi^+ = \sum_{j=1}^k d_j \chi_j^+$ as the sum of its irreducible characters, linearity and orthogonality of characters gives $\sum_{r \in G^+} |\chi^+(r)|^2 = \langle \chi^+, \chi^+ \rangle = (\sum_{j=1}^k d_j^2) |G^+|$. Thus we either have $\langle \chi^+, \chi^+ \rangle = |G^+|$, in which case ρ^+ is complex irreducible, or else that $\langle \chi^+, \chi^+ \rangle = 2|G^+|$, in which case $\chi(s) = 0$ for each reflection $s \in G - G^+$. As $d \geq 3$ and $\chi(s) = d - 2$ for any reflection, we conclude that ρ^+ is complex irreducible. \square

4.3. Polyhedral Examples. We now consider the polyhedral partitions of Theorem 1.5.

4.3.1. Regular Prisms and Antiprisms. For part (a), let $n \geq 3$ and let $\rho: \mathbb{Z}_n \oplus \mathbb{Z}_2 \rightarrow O(3)$ be the product of the inclusions $\mathbb{Z}_n \hookrightarrow U(1) \cong SO(2)$ and $\mathbb{Z}_2 = \{\pm 1\} = O(1)$. For $u = (u_1, u_2) \in \mathbb{C} \times \mathbb{R}$, the orbit polytope is free provided $u_1, u_2 \neq 0$, in which case $P(\mathbb{Z}_n \oplus \mathbb{Z}_2, u) = P(\mathbb{Z}_n, u_1) \times P(\mathbb{Z}_2, u_2)$ is a regular prism $Pr(n)$ with regular n -gon base. This is a special case of the ‘‘multiprism’’ partitions of [22, Theorem 1.3].

For part (b), let $n \geq 2$ and let $\rho: \mathbb{Z}_{2n} \rightarrow O(3)$ be the sum of inclusion representation $\mathbb{Z}_{2n} \hookrightarrow U(1)$ and the representation $\chi: \mathbb{Z}_{2n} \rightarrow O(1)$ defined by $\chi(\omega_{2n}^k) = (-1)^k$ for all $0 \leq k < 2n$. As before, one has a free orbit polytope provided each u_i of $u = (u_1, u_2) \in \mathbb{C} \times \mathbb{R}$ is non-zero. For $n \geq 3$, the translation of an orbit $\mathbb{Z}_{2n} \cdot u$ consists of the vertices of a regular n -gon and those of a π/n -rotated copy lying directly above it in a parallel plane. The convex hull is thus a regular antiprism $APr(n)$, whose faces consist of the two congruent regular n -gons and $2n$ congruent isosceles triangles. In particular, $APr(3)$ is an octahedron. When $n = 2$, the regular n -gons degenerate to perpendicular congruent edges, so that $APr(2)$ is a tetrahedron with congruent isosceles triangle faces.

The regular n -gon bases for $Pr(n)$ and $APr(n)$ guaranteed by Theorem 1.2 are precisely those which are parallel to the coordinate plane $\mathbb{R}^2 \times 0$, in which case the upper bounds given in parts (a) and (b) of Theorem 1.5 are tight. As there are $\mathbb{R}P^2$ choices of linear planes in \mathbb{R}^3 , we therefore expect the upper bounds given there to exceed the true value by 2. Owing to the extra condition on edge length, we conjecture that $N_{Pr(n)} = N_{APr(n)} = 8n - 7$ in the uniform setting.

4.3.2. Omnitruncated and Snub Polyhedra. The polyhedra of parts (c)-(h) of Theorem 1.5 are the Coxeter permutahedra and their alternahedra arising from the irreducible 3-dimensional finite reflection groups. These are the isometry groups \mathbf{T} , \mathbf{O} , and \mathbf{I} of the Platonic solids, with orders 24, 48, and 120, respectively. The resulting permutahedra are the (not necessarily uniformly) omnitruncated polyhedra [21] – the truncated icosahedron tO , truncated cuboctahedron tCO , and truncated icosidodecahedron tID , respectively (see, e.g., [17, Example 2.7]). To describe their faces, we note that all d -dimensional Coxeter permutahedra are d -simple [17, Theorem 2.3] and that each j -face is itself a j -dimensional Coxeter permutahedra [17, Proposition 2.1]. The 2-dimensional finite reflection groups are precisely the Dihedral groups D_r , $r \geq 2$ (where again we view D_2 as $\mathbb{Z}_2 \times \mathbb{Z}_2$), and it is easily seen that the Coxeter permutahedra are equiangular $2r$ -gons with alternating congruent edges. The facets for parts (c)–(e) are therefore as follows: a single set of congruent rectangles and two sets of congruent hexagons for tO ; a set each of congruent rectangles, hexagons, and octagons for tCO ; and a set each of congruent rectangles, hexagons, and decagons for tID . Alternation of these permutahedra gives the (chiral) snub polyhedra: the icosahedron sT realized as a snub tetrahedron, the snub cube sC , and the snub dodecahedron sD , respectively, whose faces can be described as follows: in addition to a single class of congruent triangles, one has two sets of congruent equilateral triangles for sT ; a set of congruent equilateral

triangles and a set of congruent squares for sC ; and a set of congruent equilateral triangles and a set of congruent regular pentagons for sI .

Considering conjugation of the above groups by orthogonal matrices, we expect the estimates of parts (c)–(h) to exceed the exact value by $\dim(O(3)) = 3$ in each case. As there are three distinct edge lengths, two additional conditions are required to ensure uniform polyhedra, and so we again expect the exact values in the uniform setting to be one fewer than the upper bounds given there.

4.4. Non-Faithful Representations. As an example in the non-faithful setting, we provide a generalization of Theorem 1.1 with intersecting convex hulls. For partitions with a somewhat similar spirit, see [27].

Theorem 4.3. *Let $r = r_1 r_2$, where $r_1 \geq 3$ and $r_2 \geq 1$. Then almost any set of $3r - 4$ points in \mathbb{R}^2 can be partitioned into a collection of r subsets $\{A_i^j\}_{1 \leq i \leq r_1, 1 \leq j \leq r_2}$ such that*

- (a) $I_i := \bigcap_{j=1}^{r_2} \text{Conv}(A_i^j) \neq \emptyset$ for all $1 \leq i \leq r_1$, and
 - (b) there exist points x_1, \dots, x_{r_1} , one from each I_i , which form the vertex set of a regular r_1 -gon.
- Moreover, such a partition fails for almost any set of $N < 3r - 4$ points.

For instance, a generic set of 32 points in the plane can be partitioned into 12 subsets whose convex hulls intersect in 4 groups of 3 in such a way that there is a point from each of the four intersections which form a square. One may also divide the set into 12 subsets whose convex hulls intersect in 3 groups of 4 so that there is a point from each of the 3 intersections which form an equilateral triangle.

Proof of Theorem 4.3. Consider the representation $\rho: \mathbb{Z}_r \rightarrow U(1)$ given by $\rho(\omega_r^k) = \omega_r^{kr_2} = \omega_{r_1}^k$. Stated in terms of affine mappings $f: \Delta_{N-1} \rightarrow \mathbb{C}$, the existence of a partition satisfying (a) and (b) of Theorem 4.3 is equivalent to the existence of a collection of points $\{x_k\}_{k \in \mathbb{Z}_r}$ from the simplex with pairwise disjoint support for which there exist points a and $u \neq 0$ in the plane for which $f(x_k) = a + \omega_r^k \cdot u$ for all $k \in \mathbb{Z}_r$. Thus we are finished by Theorem 2.1 and Lemma 3.1. \square

4.5. A Conjecture for Vertex Transitive Polytopes. We conclude with a conjecture. While we have exclusively considered free orbit polytopes, any vertex-transitive polytope is the affine translate of the (not necessarily-free) orbit polytope determined by its isometry group. For example, going back to [8], many famous uniform polytopes (including all regular ones) can be realized as the orbit polytopes $P(G, u)$ where G is a finite reflection group and u has non-trivial stabilizer. We expect the upper bound for free orbit polytopes arising from Theorem 1.2 to carry over to the non-free setting.

Conjecture 1. $N_{P(r,d)} \leq T(r, d) - d$ for any vertex-transitive d -polytope $P(r, d)$ on r vertices.

REFERENCES

- [1] L. Babai. Symmetry groups of vertex-transitive polytopes. *Geom. Dedicata*, Vol. 6 (1977) 331–337.
- [2] I. Bárány. A generalization of Carathéodory’s theorem. *Discr. Math.*, Vol. 40, No. 2–3 (1982) 141–152.
- [3] I. Bárány and P. Soberón. Tverberg’s theorem is 50 years old: a survey. *Bull. Amer. Math. Soc.*, Vol. 55, No. 4 (2018) 459–492.
- [4] A. I. Barvinok and A. M. Vershik. Convex hulls of orbits of representations of finite groups, and combinatorial optimization. *Func. Anal. Appl.*, Vol. 22, No. 3 (1988) 224–225.

- [5] P. Blagojević and G.M. Ziegler. Beyond the Borsuk–Ulam theorem: The topological Tverberg story, *A journey through discrete mathematics*. Springer, Cham (2017) 273–341.
- [6] A. V. Borovik and A. Borovik. *Mirrors and Reflections: The Geometry of Finite Reflection Groups*. Universitext. Springer, New York (2010).
- [7] H. S. M. Coxeter. *Regular Polytopes*. Dover Publications, United Kingdom (1973).
- [8] H. S. M. Coxeter. Wythoff’s construction for uniform polytopes. *Proc. Lond. Math. Soc.*, Vol. 38 , No. 2 (1934) 327–339.
- [9] J. Cruickshank and S. Kell. Rearrangement Inequalities and the Alternahedron. *Discrete Comput. Geom.*, Vol. 35 (2006) 241–254.
- [10] J. A. De Loera, X. Goaoc, F. Meunier, and N. H. Mustafa. The discrete yet ubiquitous theorems of Carathéodory, Helly, Tucker, and Tverberg. *Bull. Amer. Math. Soc.* Vol. 56 (2019), 415–511.
- [11] P. Du Val. *Homographies, Quaternions, and Rotations*. Clarendon Press: Oxford University Press (1964).
- [12] W. G. Dwyer and C. W. Wilkerson. Centers and Coxeter elements. *Homotopy Methods in Algebraic Topology*. Contemp. Math. Vol. 271, American Mathematical Society (2001) 53–76.
- [13] E. Friese. *Generic Symmetries of Group Representations*, Doctoral Dissertation. Universität Rostock (2018).
- [14] E. Friese and F. Ladisch. Affine Symmetries of orbit polytopes. *Adv. Math.*, Vol. 288 (2016) 386–425.
- [15] E. Friese and F. Ladisch. Classification of affine symmetries of orbit polytopes. *J. Algebraic Combin.*, Vol. 48, No. 3 (2018), pp. 481–509.
- [16] W. Fulton and J. Harris. *Representation Theory: A First Course. Graduate Texts in Mathematics*. Springer, New York (2004).
- [17] C. Hohlweg. Permutahedra and associahedra: generalized associahedra from the geometry of finite reflection groups. *Associahedra, Tamari lattices and related structures*. Prog. Math. Phys. 299. Birkhäuser/Springer, Basel (2012) 129–159.
- [18] C. Hohlweg, C. E. M. C. Lange, and H. Thomas. Permutahedra and generalized associahedra. *Adv. Math.*, Vol. 226, No.1 (2011) 608–640.
- [19] J.E. Humphries. *Reflection Groups and Coxeter Groups*. Cambridge University Press (1992).
- [20] N. W. Johnson. *Geometries and Transformations*. Cambridge University Press (2018).
- [21] N.W. Johnson. *The Theory of Uniform Polytopes and Honeycombs*. Doctoral Dissertation. University of Toronto (1966).
- [22] L. Leiner and S. Simon. Regular polygonal partitions of a Tverberg type. *Discrete Comp. Geom.*, Vol. 66 (2021) 1053–1071.
- [23] J. Morris, P. Spiga, and G. Verret. Automorphisms of Cayley graphs on generalised dicyclic groups. *Eur. J. Comb.*, Vol. 43 (2015) 68–81.
- [24] S. Onn. Geometry, complexity, and combinatorics of permutation polytopes. *J. Combin. Theory Ser. A*, Vol. 64, No.1 (1993) 31–49.
- [25] R. Sanyal, F. Sottile, and B. Sturmfels. Orbitopes. *Mathematika*, Vol. 57, No. 2 (2011) 275–314.
- [26] K.S. Sarkaria. Tverberg partitions and Borsuk–Ulam theorems. *Pacific J. Math.*, Vol. 196, No. 1 (2000) 231–241.
- [27] S. Simon. Average-value Tverberg partitions via finite Fourier analysis. *Isr. J. Math.*, Vol. 216, No. 2 (2016) 891–904.
- [28] H. Tverberg. A generalization of Radon’s theorem. *J. London Math. Soc.*, Vol. 41 (1966) 123–128.
- [29] G. M. Ziegler. *Lectures on Polytopes*. Grad. Texts Math. 152, Springer-Verlag, New York (1995).

(SS) BARD COLLEGE, MATHEMATICS DEPARTMENT
 Email address: ssimon@bard.edu

(TT) BARD COLLEGE, MATHEMATICS DEPARTMENT
 Email address: tt6421@bard.edu