

DEPINNING IN THE INTEGER-VALUED GAUSSIAN FIELD AND THE BKT PHASE OF THE 2D VILLAIN MODEL

MICHAEL AIZENMAN, MATAN HAREL, RON PELED, AND JACOB SHAPIRO

To Barry Simon, on the occasion of his 75th birthday

ABSTRACT. It is shown that the Villain model of two-component spins over two dimensional lattices exhibits slow, non-summable, decay of correlations at any temperature at which the dual integer-valued Gaussian field exhibits depinning. For the latter, we extend the recent proof by P. Lammers of the existence of a depinning transition in the integer-valued Gaussian field in two-dimensional graphs of degree three to all doubly-periodic graphs, in particular to \mathbb{Z}^2 . Taken together these two statements yield a new perspective on the Berezinskii-Kosterlitz-Thouless phase transition in the Villain model, and complete a new proof of depinning in two-dimensional integer-valued height functions.

CONTENTS

1. Introduction	2
2. Depinning in the integer-valued Gaussian field	3
3. The BKT phase of two-dimensional spin models	10
4. Duality relations between the two models	12
5. New results connecting the Villain model to the ZGF.	14
6. Proof of the ZGF bound on the spin correlation function	18
7. Positivity of the ZGF's stiffness modulus	22
8. The relation of the spin correlation function with height fluctuations	27
Appendix A. The Regev Stephens-Davidowitz Monotonicity Theory	30
Appendix B. Perspectives on the Villain model	33
References	35

Date: October 22, 2021.

1. INTRODUCTION

1.1. A brief outline.

We discuss here a pair of phase transitions linked through a duality relation, related to different phenomena each of which is special to two dimensions. One is the depinning transition in the integer-valued Gaussian field (ZGF). The other concerns the Berezinskii-Kosterlitz-Thouless (BKT) phase transition in the $O(2)$ invariant Villain spin model on the dual lattice.

The configurations of the ZGF on $\mathbb{Z}^d \cap [-L, L]^d$ form a random surface of a discretized height, tied at the domain's boundary, whose fluctuations are controlled by elastic restoring forces. In its localized, or pinned, phase the fluctuations of the surface's height at any given point remain tight. Depinning means that in the limit $L \rightarrow \infty$ they diverge, and the surface fluctuates away. The corresponding phase transition is special to two dimensions: in one dimension the surface depins at any temperature, with Brownian bridge type fluctuations, whereas in $d > 2$ dimensions its fluctuations are tight at all temperatures. In two dimensions the ZGF exhibits a temperature driven phase transition.

The Berezinskii-Kosterlitz-Thouless (BKT) phase is a phenomenon encountered in two dimensional ferromagnetic systems with an $O(2)$ rotational symmetry. At low enough temperatures the spin-spin correlations in such systems transition from exponential to *slow decay*, or varying power rate, without exhibiting long range order (which for $d = 2$ is ruled out by the Mermin-Wagner theorem).

As is often the case, the duality relation under which the two systems are related also inverts the relevant temperature-like parameter. However this in itself does not explain the transitions discussed here since the less trivial phases of the two models coincide under duality. More explicitly: the more challenging behaviors to prove are the *depinning* of the ZGF which characterizes its *high* temperature phase and the slow decay of correlations which characterizes the spin system's *low* temperature BKT phase.

The analyses of the two phenomena have so far proceeded through their relation with yet another system, which is a lattice Coulomb gas—a system of integer charges with the two dimensional Coulomb interaction (i.e. a logarithmic potential). Its relevant transition is between a plasma state and a gas of neutral dipoles [15]. The rigorous proof of this behavior and its implications for the two transitions outlined above was presented in the groundbreaking work by J. Fröhlich and T. Spencer [10], the crux of which is establishing the condensation of the 2D Coulomb system at low temperatures into a gas of dipoles. The multi-scale analysis that was employed there has inspired a number of other applications, and it continues to draw refinements [13, 7]. Yet there may still be potential interest in alternative approaches to the two transitions highlighted above.

The main results presented below include:

- 1) A proof of the existence of a depinning transition in planar doubly periodic \mathbb{Z} GF models, which is intrinsic to these systems in the sense that it does not proceed through the phase transition of the dual Coulomb gas.

Such a proof was recently presented for cubic graphs (graphs of degree 3) in a breakthrough work by P. Lammers [18]. Here we present a construction which, with the help of recent inequalities of Regev and Stephens-Davidowitz [24], allows us to remove the degree constraint.

- 2) A direct link between depinning in the \mathbb{Z} GF and slow decay of correlations in the Villain $O(2)$ spin model.

More explicitly, it is shown that depinning in the dual \mathbb{Z} GF model implies that at the corresponding (inverted) temperature

$$\sum_{x \in \mathbb{Z}^2} \left(\langle \sigma_0 \cdot \sigma_x \rangle_\beta^{\text{Vill}} \right)^{1-\varepsilon} = \infty \quad \forall \varepsilon > 0. \quad (1.1)$$

which is in contrast to the exponential decay that characterizes high-temperature phases [8, 26, 1, 10].

Taken together, the two results yield a new perspective on the BKT phase of the Villain spin model.

Since the depinning transition is of independent interest, we discuss it first. After presenting the new result on the depinning transition, we introduce the spin model, and turn to the duality relations by which the two are linked, some known and some new.

2. DEPINNING IN THE INTEGER-VALUED GAUSSIAN FIELD

2.1. The \mathbb{Z} GF model and its two phases

The integer-valued Gaussian field (\mathbb{Z} GF) over a connected planar graph \mathcal{G} , of vertex set \mathcal{V} and edge set \mathcal{E} (with a locally-finite embedding in \mathbb{R}^2), has as its basic variables the values of the random function $n : \mathcal{V} \rightarrow \mathbb{Z}$. The \mathbb{Z} GF finite-volume partition function in $\mathcal{V}_L = \mathcal{V} \cap [-L, L]^2$, which by default is taken here with the Dirichlet boundary conditions, is

$$Z_{\lambda, L}^{\mathbb{Z}\text{GF}, \mathcal{G}} \equiv \sum_{n: \mathcal{V}_L \rightarrow \mathbb{Z}: n|_{\partial \mathcal{V}_L} = 0} \exp \left(-\frac{\lambda}{2} \sum_{\{u, v\} \in \mathcal{E}} (n_u - n_v)^2 \right). \quad (2.1)$$

At any given λ the probability of n is given by the configuration's normalized contribution to the above sum.

Quite generally, Gaussian domination holds in the sense that the fluctuations of the \mathbb{Z} GF are upper bounded by those of the corresponding Gaussian free field (GFF) [12] (cf. [16]).

Explicitly, for every $x, y \in \mathcal{V}_L$,

$$\mathbb{E}_{\lambda, L}^{\text{ZGF}, \mathcal{G}} [(n_x - n_y)^2] \leq \mathbb{E}_{\lambda, L}^{\text{GFF}, \mathcal{G}} [(\phi_x - \phi_y)^2] \quad (2.2)$$

and furthermore, for every $f : \mathcal{V}_L \rightarrow \mathbb{R}$,

$$\mathbb{E}_{\lambda, L}^{\text{ZGF}, \mathcal{G}} [\exp(\langle f, n \rangle)] \leq \mathbb{E}_{\lambda, L}^{\text{GFF}, \mathcal{G}} [\exp(\langle f, \phi \rangle)]. \quad (2.3)$$

In two dimensions such models exhibit a *localization–delocalization*, or *de-pinning*, phase transition, which can also be presented in terms of *symmetry breaking*.

A shift of the boundary conditions to $n_u = k$ at all $u \in \partial\mathcal{V}_L$ at other values of $k \in \mathbb{Z}$ does not affect the partition function, but the corresponding Gibbs state will be shifted accordingly. Uniform shifts therefore represent potential symmetry of the system's infinite-volume Gibbs states.

Definition 2.1. The ZGF is said to be *pinned*, or exhibit *localization* at a specified λ iff for every $x \in \mathcal{V}$,

$$\lim_{L \rightarrow \infty} \mathbb{E}_{\lambda, L}^{\text{ZGF}, \mathcal{G}} [n_x^2] < \infty. \quad (2.4)$$

Otherwise it is said to *depin*, or be *delocalized*.

The limit in (2.4) exists by the model's absolute-value FKG property, which was established by Lammers–Ott [19]. By (2.2), if (2.4) holds for a single x then it holds for all x . And, by the log-concavity of the marginal distributions of (n_x) , which was proven by Sheffield [25, Section 8.2], in the delocalized phase, for every $x \in \mathcal{V}$,

$$\lim_{L \rightarrow \infty} \mathbb{E}_{\lambda, L}^{\text{ZGF}, \mathcal{G}} [|n_x|] = \infty \quad (2.5)$$

and furthermore, for any $t < \infty$,

$$\lim_{L \rightarrow \infty} \mathbb{P}_{\lambda, L}^{\text{ZGF}, \mathcal{G}} \{|n_x| \leq t\} = 0. \quad (2.6)$$

In the pinned phase the system admits an infinite collection of infinite-volume Gibbs equilibrium states which are limits of the finite states with boundary values $k \in \mathbb{Z}$, and hence differ by simple shifts. In the doubly-periodic case, which is the case of interest here, the limiting states will be ergodic.

In contrast, in the de-pinned phase the distribution of $|n_x|$ drifts to ∞ as $L \rightarrow \infty$. However, it is still the case that the distribution of the gradients of n , or equivalently of each of the difference variables $\{n_x - n_y\}_{x, y \in \mathcal{V}}$ remains tight (though not uniformly in $\|x - y\|$).

In the latter case, the system behaves similarly to the GFF—the classical 2D Gaussian free field defined on the lattice, i.e., without the \mathbb{Z} -valued constraint. The GFF does not undergo a phase transition since, by its Hamiltonian's homogeneity, the field's distributions at different temperatures are related via simple rescaling by $\sqrt{\lambda}$. In contrast, for the ZGF there are two natural scales: that of the discretization step (here 1), and that of the typical

local fluctuations, $1/\sqrt{\lambda}$. Alternatively stated: there is an energy gap. This sets a relevant scale for λ , opening the possibility for a phase transition.

Our main result in this section is

Theorem 2.2. *For the ZGF on any graph \mathcal{G} as above, there exists $\lambda_c(\mathcal{G}) \in [0, \infty]$ such that the ZGF is pinned for $\lambda > \lambda_c(\mathcal{G})$ and depinned for $\lambda < \lambda_c(\mathcal{G})$.*

When \mathcal{G} is doubly periodic $0 < \lambda_c(\mathcal{G}) < \infty$. Furthermore, for the ZGF on \mathbb{Z}^2 , $\lambda_c(\mathbb{Z}^2) \geq \frac{2}{3} \ln 2$, and at $\lambda = \frac{2}{3} \ln 2$ the model is delocalized.

For the sake of simplicity of presentation we concentrate on \mathbb{Z}^2 but the argument presented here extends to the more general cases (albeit with the bounds on the critical coupling increased according to the maximal degree).

The delocalized phase for the ZGF on planar doubly periodic graphs was first established in [10] using the dual Coulomb gas perspective. Recently a new proof was presented by Lammers [18] for shift-invariant planar graphs of maximum degree 3.

The degree limitation played an important role in Lammers' new argument. Here we extend its applicability by presenting a series of surgeries on the model through which it is transformed into the ZGF on a cubic graph, albeit one with stronger couplings. This transformation is rather natural, but the key point is the proof that in each step the height fluctuations can only *decrease*. This is done here by adapting the question to the recently presented Regev Stephens-Davidowitz [24] monotonicity theory. The bound on the critical coupling constant stated in [Theorem 2.2](#) builds on Lammers proof that on shift invariant cubic graphs, the ZGF is delocalized at $\lambda \leq 2 \ln 2$.

2.2. An outline of Lammers' proof of delocalization on cubic planar graphs

Following is a sketch of Lammers' proof of delocalization, aimed at explaining the origin of the argument's limitation to degree three. It starts with the known observation that a necessary condition for the ZGF to be pinned at a given λ is the almost sure percolation of the set $\{x \in \mathcal{V} \mid n_x \geq 0\}$ as well as that of its flipped image $\{x \in \mathcal{V} \mid n_x \leq 0\}$. Key steps for that are:

- 1) by the shift invariance and ergodicity of the putative limiting state, for each of these sets the probability that it includes an infinite connected cluster is either 1 or 0,
- 2) in the absence of percolation, with probability one each finite volume is encircled by a *-connected path of sites along which $n_x < 0$, and also by another *-connected path of sites with $n_x > 0$,
- 3) by conditioning (initially over the outermost such paths in finite but large volumes) one may deduce that $\lim_{L \rightarrow \infty} \mathbb{E}_{\lambda, L}^{\text{ZGF}, \mathcal{G}} [n_0] \leq -1$, and $\lim_{L \rightarrow \infty} \mathbb{E}_{\lambda, L}^{\text{ZGF}, \mathcal{G}} [n_0] \geq +1$, which would be a contradiction.



FIGURE 1. The significance of the vertex degree for edge percolation. Through a vertex of degree 3 in any configuration only one of the \pm signs may edge percolate. That is not the case at vertices of higher degree. An essential part of Lammers’ argument is the reduction of influence percolation at small λ to an edge percolation process.

Continuing with known arguments, Lammers notes that on doubly-periodic planar graphs two incompatible percolation processes cannot occur simultaneously [25] (using the uniqueness argument of [5]). That does not yet apply to the case presented above since the two sets share the set of sites where $n = 0$. And while it seems reasonable to expect the simultaneous percolation to be inhibited when the density of the sets’ shared part is sufficiently small, as it is for $\lambda \ll 1$, a simple direct proof of this is not readily available.

The above discussion may be viewed as concerning the influence percolation process in which the information about the level to which the surface is pinning at the boundary propagates through a site percolation process. The new step in Lammers’ argument is to replace the above by an edge percolation.

For that purpose, Lammers shows that at λ sufficiently small it is possible to eliminate the neutral option $\{x \in \mathcal{V} \mid n_x = 0\}$, through the addition of an ad hoc field associated with the graph’s edges by which they are declared of either positive or negative sign (a related idea was employed in [2]). Through this device the influence propagation is presented as occurring through edge percolation, in a system where the edges carry a sign. On graphs of degree 3, paths of differently signed edges cannot cross (cf. Figure 1). By the non-coexistence principle, which now does apply, the above argument can be made to work.

Lammers’ theorem proves depinning for the ZGF on the hexagonal graph (whose dual is the triangular lattice). However, it obviously does not apply to \mathbb{Z}^2 .

2.3. A more general proof of depinning

Towards the proof of Theorem 2.2 we transform the given ZGF model through a sequence of local steps. The first is explained by the observation that by the divisibility of its distribution, a Gaussian variable of variance $\frac{1}{\lambda}$ can be presented as the sum of two independent Gaussian variables, of variance $\frac{2}{3\lambda}$ and $\frac{1}{3\lambda}$. Thus the ZGF distribution on any graph




FIGURE 2. Local surgery in which a vertex of degree 4 is replaced by two of degree 3. It proceeds through the addition of mediating vertices (at doubled coupling strength) in the midst of a pair of adjacent edges, with variables which are initially \mathbb{R} and then converted to \mathbb{Z} -valued. Subsequently the sites are merged through infinite coupling, or equivalently conditioning on having equal values. An essential element in our analysis is the proof that in each step the model's fluctuations can only decrease.

remains unaffected if the coupling across any of its edges is replaced by coupling through an intermediate site added in the midst of the edge. We proceed as follows.

- 1) Enhance the model by adding *real-valued* variables in the midpoints of the edges of \mathbb{Z}^2 , repealing the original interaction by one mediated by mid-edge spins with coupling at strength $3\lambda/2$ across the lower and left parts of the split edges, and 3λ across the upper and the right parts of the split edges. (This does affect the distribution of the sum along each of the original edges).
- 2) Next, constrain the mid-edge variables to take values in \mathbb{Z} instead of \mathbb{R} . It need not be initially obvious, but is proven below, that this change can only reduce the fluctuations of n .
- 3) Collapse the variables pairwise by forcing the pairs of variables north and east from the original lattice sites to assume equal value (see [Figure 2](#)). This again only reduces fluctuations, and results in a \mathbb{Z} GF model on a cubic planar graph with uniform coupling constants of 3λ .

Applied to the \mathbb{Z} GF model on \mathbb{Z}^2 this construction results in \mathbb{Z} GF on the hexagonal graph at λ increased to 3λ .

We next prove the monotonicity of fluctuations in these processes, applying for this purpose the Regev Stephens-Davidowitz monotonicity theory [24] which concerns Gaussian probability measures on lattices. Before presenting it, we alert the reader that in the following terminology, the relevant lattice \mathcal{L} is the collection of configurations of the \mathbb{Z} GF over the given finite graph. Its dimension k is the number of graph vertices.

A lattice \mathcal{L} in \mathbb{R}^k is the set $\mathcal{L} := LZ^k$ for some \mathbb{R} -valued $k \times k$ matrix $L \in \text{Mat}_k(\mathbb{R})$. Given a lattice \mathcal{L} and a positive $k \times k$ matrix A , a random field $\psi \in \mathcal{L}$ associated with (A, \mathcal{L}) is defined via the partition function

$$Z_{A, \mathcal{L}} := \sum_{\psi \in \mathcal{L}} \exp\left(-\frac{1}{2}\langle \psi, A\psi \rangle\right). \quad (2.7)$$

For example, the ZGF on a graph with the set of vertices Λ and coupling λ is obtained as the special case of this family of models by choosing $k = |\Lambda|$ and letting $A = -\lambda\Delta$ where Δ is the discrete Laplacian on $\ell^2(\Lambda)$ with Dirichlet boundary conditions.

Since, for any $v \in \mathbb{R}^k$, monotonicity of the moment-generating function implies monotonicity of the second moment of $\langle v, \psi \rangle$, let us focus on the former, which we denote:

$$\mathbb{M}_{A, \mathcal{L}}[v] := \mathbb{E}_{A, \mathcal{L}}[\exp(\langle v, \psi \rangle)] \quad (v \in \mathbb{R}^k). \quad (2.8)$$

In the application we have in mind, v is proportional to $\delta_x - \delta_y$ (y possibly being at the boundary, with $n(y) = 0$ fixed by the boundary conditions).

Then the Regev Stephens-Davidowitz theory provides us with monotonicity in the lattice:

Proposition 2.3. *If $\mathcal{M} \subseteq \mathcal{L}$ is a sub-lattice then*

$$\mathbb{M}_{A, \mathcal{M}}[v] \leq \mathbb{M}_{A, \mathcal{L}}[v] \quad (v \in \mathbb{R}^k, A > 0).$$

Additionally, it also says that $A \mapsto \mathbb{M}_{A, \mathcal{L}}[v]$ is matrix-monotone:

Proposition 2.4. *For any lattice \mathcal{L} and any two matrices A, B such that $A \geq B > 0$ (in the quadratic form sense), we have*

$$\mathbb{M}_{A, \mathcal{L}}[v] \leq \mathbb{M}_{B, \mathcal{L}}[v] \quad (v \in \mathbb{R}^k).$$

For the reader's convenience, a self-contained presentation of the relevant statements is presented in [Appendix A](#) below. These two propositions will next be applied in the proof of the results stated above.

Proof of [Theorem 2.2](#). The first assertion is a direct consequence of the monotonicity in λ which follows from [Proposition 2.4](#).

That $\lambda_c(\mathbb{Z}^2) < \infty$ is a known fact, provable by a Peierls-type argument. To prove the positivity of $\lambda_c(\mathbb{Z}^2)$, we follow the above construction, justifying here that in each step the fluctuations, such as $\mathbb{E}[n_x^2]$, can only decrease. In the following discussion $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ refers to the finite-volume restrictions of \mathbb{Z}^2 .

Step 1: As outlined above, using the identity

$$\exp\left(-\frac{1}{2}\lambda(n_x - n_y)^2\right) = \sqrt{\frac{4\lambda}{9\pi}} \int_{n_{xy} \in \mathbb{R}} \exp\left\{-\lambda\left[3(n_x - n_{xy})^2/2 + 3(n_y - n_{xy})^2\right]\right\} dn_{xy} \quad (2.9)$$

we may associate to each edge of \mathcal{G} an additional *real-valued* degree of freedom $n_{x,y}$, so that altogether the field has elastic energy which is associated with the discrete Laplacian on the union graph $\tilde{\mathcal{G}}$ with vertex set $\tilde{\mathcal{V}} := \mathcal{V} \cup \mathcal{E}$ and the field is a map $n \in \mathbb{Z}^{\mathcal{V}} \times \mathbb{R}^{\mathcal{E}}$, with the inhomogeneous coupling constants set at $3\lambda/2$ and 3λ as explained above. We denote the associated probability distribution $\mathbb{P}_{\mathbb{Z}^{\mathcal{V}} \times \mathbb{R}^{\mathcal{E}}, 2\lambda}^{\tilde{\mathcal{G}}}$. One should note that at this point it is of mixed \mathbb{R} -valued and \mathbb{Z} -valued variables.

Step 2: The \mathbb{R} -valued field can be viewed as a $\varepsilon \rightarrow 0$ limit of a $\varepsilon\mathbb{Z}$ -valued one, that is,

$$\lim_{\varepsilon \rightarrow 0} \mathbb{M}_{\mathbb{Z}^{\mathcal{V}} \times (\varepsilon\mathbb{Z})^{\mathcal{E}}, \bar{\lambda}}^{\tilde{\mathcal{G}}} [v] = \mathbb{M}_{\mathbb{Z}^{\mathcal{V}} \times \mathbb{R}^{\mathcal{E}}, \bar{\lambda}}^{\tilde{\mathcal{G}}} [v] \quad (v \in \mathbb{R}^{\mathcal{V}}), \quad (2.10)$$

where $\bar{\lambda}$ indicates the the inhomogeneous coupling constants described above. Applying **Proposition 2.3** to the sequence $\varepsilon_k = 2^{-k}$, in which case the corresponding lattices are nested ($(2\varepsilon\mathbb{Z})^{\mathcal{E}}$ being a sub-lattice of $(\varepsilon\mathbb{Z})^{\mathcal{E}}$), we conclude that

$$\mathbb{M}_{\mathbb{ZGF}, \bar{\lambda}}^{\tilde{\mathcal{G}}} [v] \leq \mathbb{M}_{\mathbb{Z}^{\mathcal{V}} \times (\mathbb{R})^{\mathcal{E}}, \bar{\lambda}}^{\tilde{\mathcal{G}}} [v] \quad (v \in \mathbb{R}^{\tilde{\mathcal{V}}}). \quad (2.11)$$

It follows that the conversion of the mixed \mathbb{R} -valued and \mathbb{Z} -valued Gaussian field to a pure ZGF on $\tilde{\mathcal{G}}$ with the same coupling constant can only lower the variance of $n(0)$.

Now, taking the limit along appropriate subsequences so that, e.g., $(2\varepsilon\mathbb{Z})^{\mathcal{E}}$ is a sub-lattice of $(\varepsilon\mathbb{Z})^{\mathcal{E}}$, we see from **Proposition 2.3** that

$$[0, 1] \ni \varepsilon \mapsto \mathbb{M}_{\mathbb{Z}^{\mathcal{V}} \times (\varepsilon\mathbb{Z})^{\mathcal{E}}, \bar{\lambda}}^{\tilde{\mathcal{G}}} [v]$$

is monotone decreasing for any vector $v \in \mathbb{R}^{\tilde{\mathcal{V}}}$. This implies that

$$\mathbb{M}_{\mathbb{ZGF}, \bar{\lambda}}^{\tilde{\mathcal{G}}} [v] \leq \mathbb{M}_{\mathbb{Z}^{\mathcal{V}} \times (\mathbb{R})^{\mathcal{E}}, \bar{\lambda}}^{\tilde{\mathcal{G}}} [v] \quad (v \in \mathbb{R}^{\tilde{\mathcal{V}}}), \quad (2.12)$$

i.e., converting the mixed \mathbb{R} -valued and \mathbb{Z} -valued Gaussian field to a pure ZGF on $\tilde{\mathcal{G}}$ with the same coupling constant only lowers fluctuations.

Remark 2.5. We note in passing that the above gives another proof of the Gaussian domination assertions stated in (2.2) and (2.3). Explicitly,

$$\mathbb{M}_{\lambda}^{\mathbb{ZGF}, \mathcal{G}} [v] \equiv \mathbb{E}_{\lambda}^{\mathbb{ZGF}, \mathcal{G}} \left[e^{\langle v, n \rangle} \right] \leq \exp \left(+ \frac{1}{2\lambda} \langle v, -\Delta^{-1} v \rangle \right) \quad (v \in \ker(-\Delta)^{\perp}). \quad (2.13)$$

Step 3: In a repeated application of the monotonicity principle expressed in **Proposition 2.3** we deduce that also in the third step of the construction, presented as the imposition of the constraint to a sub lattice of configurations in which the new edge variables are pairwise equal, the fluctuations can only decrease.

The assertion that the ZGF is depinned at $\lambda \leq \frac{2}{3} \ln(2)$ follows by applying Lammers' theorem to the hexagonal graph which is obtained by the above construction. \square




FIGURE 3. The degree-3 graph on which the fluctuations of n_0 , at adjusted couplings, provide a lower bound for those on \mathbb{Z}^2 . The (thick) positively-sloped lines result from the binding of two edges, at the construction's last step. The spitting ratio of the couplings at the first step is selected so as to equilibrate the resulting model's coupling. These end up at 3λ , in terms of the ZGF's coupling on the original graph \mathbb{Z}^2 .

3. THE BKT PHASE OF TWO-DIMENSIONAL SPIN MODELS

3.1. The Villain model

Prototypical examples of 2D ferromagnetic systems with $O(2)$ symmetry are two-dimensional arrays of two-component unit spins, attached to the sites \mathcal{V} of a two dimensional homogeneous graph \mathcal{G} with, e.g., the XY ferromagnetic nearest neighbor interaction

$$H^{XY}(\sigma) = - \sum_{\{u,v\}: \|u-v\|=1} \sigma_u \cdot \sigma_v = - \sum_{\{u,v\}: \|u-v\|=1} \cos(\theta_u - \theta_v). \quad (3.1)$$

The spins are presented here in two equivalent ways: as two-component unit vectors $\sigma_x \in \mathbb{R}^2$, or as complex unitary numbers $e^{i\theta_x} \in \mathbb{S}^1 \subseteq \mathbb{C}$.

The graphs \mathcal{G} considered here are planar and doubly periodic. Examples of interest include the easily drawn square lattice \mathbb{Z}^2 , and the triangular lattice whose dual (the hexagonal graph) is everywhere of degree 3. The graphs may be finite or infinite, the latter case conveniently viewed as the infinite-volume limit of its finite restrictions. Thus \mathbb{Z}^2 is studied as a limit ($L \rightarrow \infty$) of its truncated version $\mathcal{V}_L = [-L, L]^2 \cap \mathbb{Z}^2$. By default we take the model with free boundary conditions.

The simplest model to write down, though not the most convenient for analysis, is the basic $O(2)$ spin model whose partition function with free boundary conditions is

$$Z_\beta^{XY, \mathcal{G}} = \int_{\theta \in [-\pi, \pi]^{\mathcal{V}}} \prod_{\{u,v\} \in \mathcal{E}} e^{\beta \cos(\theta_u - \theta_v)} d\theta. \quad (3.2)$$

The corresponding Gibbs equilibrium state is the probability distribution which is obtained by normalizing (i.e. dividing by $Z_\beta^{XY, \mathcal{G}}$) the measure over which the configurations are integrated in (3.2); here \mathcal{E} denotes the graph's edges.

We find it convenient to discuss the transition to slow decay of correlations in the context of the Villain version of the $O(2)$ model [27]. In it the Gibbs factors in (3.2) are replaced through the following substitution

$$e^{\beta \cos(\theta_u - \theta_v)} \Rightarrow \sum_{m_{uv} \in \mathbb{Z}} e^{-\frac{\beta}{2}(\theta_u - \theta_v + 2\pi m_{uv})^2}. \quad (3.3)$$

An intrinsic motivation for considering this model is presented in [Appendix B](#). However, skipping that one may note that in both versions of the $O(2)$ spin models: i) the systems' symmetries include global spin rotations by a common angle, ii) the weights assigned to the spin configurations are of the same periodicity in θ and have the same maxima, in the vicinity of which they agree up to the fourth order (after a trivial correction by factors of e^β).

Of particular interest is the behavior of the correlation function

$$\langle \sigma_x \cdot \sigma_y \rangle_\beta = \lim_{L \rightarrow \infty} \langle \sigma_x \cdot \sigma_y \rangle_{\beta, L} \quad (3.4)$$

where $\sigma_x \cdot \sigma_y = \cos(\theta_x - \theta_y)$ and $\langle \dots \rangle_{\beta, L}$ denotes the expectation value with respect to the finite-volume Gibbs equilibrium state in $\Lambda(L) = [-L, L]^2$ at the inverse temperature $\beta = (k_B T)^{-1}$ with free boundary conditions. The limit's existence for both the basic XY model and its Villain version follows by the Ginibre correlation inequalities [14, 12]. These imply that $\langle \sigma_x \cdot \sigma_y \rangle_{\beta, L}$ is monotone increasing in L , and more generally in the volume.

By general arguments [8, 26], or more model-specific bounds [1], at high enough temperatures the spin-spin correlations decay exponentially fast. That is, for all $\beta < \beta_c$, at some $\beta_c > 0$,

$$\langle \sigma_x \cdot \sigma_y \rangle_\beta \leq A(\beta) e^{-m(\beta)\|x-y\|} \quad \text{with } m(\beta) > 0, A(\beta) < \infty. \quad (3.5)$$

In dimensions $d > 2$ such models exhibit also a low temperature phase with continuous symmetry breaking, where $\lim_{|y| \rightarrow \infty} \langle \sigma_x \cdot \sigma_y \rangle_\beta > 0$ [11]. However, by the Mermin–Wagner theorem, that does not occur in two dimensions [9, 22, 20].

Nevertheless, as was pointed out by Berezinskii [4] and Kosterlitz and Thouless [17], and proven rigorously in [10], two-component spin systems with a rotation-invariant interaction do exhibit a low temperature phase at which (3.5) does not hold. At that phase the spin-spin correlations decay slowly, at a temperature-dependent power law, i.e., there is some $\beta_{\text{BKT}} > 0$ such that

$$\langle \sigma_x \cdot \sigma_y \rangle_\beta \approx \frac{\text{Const.}}{\|x-y\|^{\eta(\beta)}} \quad (\text{for all } \beta > \beta_{\text{BKT}}). \quad (3.6)$$

It is expected that such behavior does not occur for systems in which the $O(2)$ spin-rotation symmetry is part of a larger non-Abelian $O(N)$ symmetry at $N > 2$, e.g., in the

three-component classical Heisenberg model [23]. However, in contrast to the case $N = 2$, still unresolved challenges were raised to the arguments on which this prediction rests [21].

As mentioned above, the BKT phenomenon has previously been analyzed through a dual system of integral charges coupled through the two-dimensional Coulomb interaction [15, 10, 13]. We next present a different approach, which is based on the system's direct relation with the integer-valued Gaussian field.

4. DUALITY RELATIONS BETWEEN THE TWO MODELS

4.1. Equality of the partition functions

As outlined in (3.3), the Villain model's partition function, on a finite graph \mathcal{G} and with the free boundary conditions, is

$$Z_\beta^{\text{Vill},\mathcal{G}} = \int_{\theta \in [-\pi, \pi]^\mathcal{V}} \prod_{\{u,v\} \in \mathcal{E}} \left[\sum_{m_{uv} \in \mathbb{Z}} e^{-\frac{\beta}{2}(\theta_u - \theta_v + 2\pi m_{uv})^2} \right] d\theta, \quad (4.1)$$

and the corresponding Gibbs equilibrium expectation values of local functions of $\{\theta_x\}_{x \in \mathcal{V}}$ are given by

$$\langle f \rangle_\beta^{\text{Vill},\mathcal{G}} = \frac{1}{Z_\beta^{\text{Vill},\mathcal{G}}} \int_{\theta \in [-\pi, \pi]^\mathcal{V}} f(\theta) \prod_{\{u,v\} \in \mathcal{E}} \left[\sum_{m_{uv} \in \mathbb{Z}} e^{-\frac{\beta}{2}(\theta_u - \theta_v + 2\pi m_{uv})^2} \right] d\theta. \quad (4.2)$$

In a known duality relation (c.f. [10, Appendix A]) for planar \mathcal{G} the Villain model's partition function equals that of the ZGF on the dual graph \mathcal{G}^* at $\lambda = 1/\beta$, and with the Dirichlet boundary conditions ($n_u = 0 \forall u \in \partial\mathcal{V}^*$):


$$(2\pi)^{|\mathcal{V}^*|} Z_\beta^{\text{Vill},\mathcal{G}} = \sum_{\substack{n \in \mathbb{Z}^{\mathcal{V}^*} \\ n_{\partial\mathcal{G}^*} \equiv 0}} \exp\left(-\frac{1}{2\beta} \|\nabla n\|^2\right) \equiv Z_{\lambda=\frac{1}{\beta}}^{\text{ZGF},\mathcal{G}^*} \quad (4.3)$$

where ∇n (the gradient of n) is a function defined over the edges of \mathcal{G}^* , associating to the oriented edge (x, y) the difference $\nabla n(x, y) = n(y) - n(x)$, and

$$\|\nabla n\|^2 \equiv \sum_{\{x,y\} \in \mathcal{E}^*} (n_x - n_y)^2.$$

This relation, and a further perspective on the model, are discussed in greater detail in [Appendix B](#).

To clarify the notation when discussing the dual models on the dual pair of graphs we adapt the following convention: The vertices of \mathcal{G} will continue to be called vertices, while the vertices of \mathcal{G}^* , on which the ZGF model is defined, will be referred to as faces.

FIGURE 4. A depiction of γ_{yx} and Γ_{yx} .

4.2. The dual expression for the spin correlation function

Under the above correspondence the spin-spin correlation function is mapped onto what may be viewed as the expectation value of a dislocation-favoring operator. In presenting it we shall employ the following notation: given a pair of sites $\{x, y\} \subset \mathcal{V}$ and an oriented path γ_{yx} from x to y along the edges of \mathcal{E} , we denote by Γ_{yx} the function defined over the oriented edges of \mathcal{G}^* by

$$\Gamma_{yx}(u, v) := \begin{cases} +1 \text{ (resp. } -1) & (u, v) \text{ is a counter-clockwise (resp. clockwise)} \\ & \text{rotation of an edge traversed by } \gamma_{yx} . \\ 0 & \text{otherwise} \end{cases}$$

The following expression represents a dislocation-favoring deformation of the elastic energy, which favors a jump discontinuity across γ_{yx} :

$$\|\nabla n + \Gamma_{yx}\|^2 \equiv \sum_{\{u,v\} \in \mathcal{E}^*} (n_u - n_v + \Gamma_{yx}(u, v))^2. \quad (4.4)$$

The duality transformation leading to (4.3) yields also the following relation [10], which holds for any choice of the \pm sign:

$$\begin{aligned} \langle \sigma_x \cdot \sigma_y \rangle_\beta^{\text{Vill}, \mathcal{G}} &\equiv \langle e^{\pm i(\theta_y - \theta_x)} \rangle_\beta^{\text{Vill}, \mathcal{G}} = \frac{\sum_{n \in \mathbb{Z}^{\mathcal{V}^*}} \mathbb{1}_{n_{\partial \mathcal{G}^*} \equiv 0} \exp(-\frac{1}{2\beta} \|\nabla n \pm \Gamma_{yx}\|^2)}{Z_{1/\beta}^{\text{ZGF}, \mathcal{G}^*}} \\ &= \mathbb{E}_{1/\beta}^{\text{ZGF}, \mathcal{G}^*} [T_{\gamma_{yx}}^+] = \mathbb{E}_{1/\beta}^{\text{ZGF}, \mathcal{G}^*} [T_{\gamma_{yx}}^-], \end{aligned} \quad (4.5)$$

where $T_{\gamma_{yx}}^+$ and $T_{\gamma_{yx}}^-$ are the two functions

$$T_{\gamma_{yx}}^\pm(n) = \exp\left(\frac{1}{2\beta} \left(\|\nabla n\|^2 - \|\nabla n \pm \Gamma_{yx}\|^2\right)\right). \quad (4.6)$$

One may note the homotopy invariance of the expression on the right in [Section 4.2](#): as a function of γ , its value is equal for any pair of paths with coinciding end points. That is obviously true for the LHS of [Section 4.2](#). For the RHS this can be explained through a “gauge transformation” which consists of an increase of each n_u by the winding number around u by the oriented loop formed by concatenating one of the paths with the inverse of the other.

5. NEW RESULTS CONNECTING THE VILLAIN MODEL TO THE ZGF.

The duality relation [Section 4.2](#) will be used below for a key lower bound on the spin correlation function in terms of the probability that the two sites lie on a common level line of the ZGF. We start by defining the relevant concepts.

5.1. Level lines of the ZGF

Viewing the ZGF process as describing a random height function on the graph \mathcal{G}^* it is natural to think of its realization in terms of the corresponding contour maps. Adapting to the fact that the values of $\{n_u\}_u$ may change discontinuously, we base the contour description on oriented contours of heights in $\mathbb{Z} + 1/2$.

More specifically: with each realization n of the ZGF on a planar graph \mathcal{G}^* , embedded in \mathbb{R}^2 , we associate the family of non-crossing contours, each either a loop or an infinite line, drawn in \mathbb{R}^2 through the following rules:

- i) an oriented contour line at level $q \in \mathbb{Z} + \frac{1}{2}$ passes through an edge $\{u, v\} \in \mathcal{E}$ if and only if $n < q$ on the face to its right and $n > q$ at the face to its left (thus the number of contour lines crossing the edge $\{u, v\}$ is exactly $|n_u - n_v|$, one for each value of q meeting these conditions).
- ii) the right-hand rule: at each vertex of \mathcal{G} each incoming q -line exits through the first outgoing q -line to its right in the counterclockwise order of the edges exiting the given vertex, as depicted in [Figure 5](#).

The construction is enabled by the fact that at each vertex the orientations of the incoming and outgoing q lines alternate in the circular ordering of the edges linked to the vertex.

Segment of an oriented q contour would be referred to as q -paths. And, for close contours we adapt the standard terminology for calling counterclockwise (resp. clockwise) loops as positively (resp. negatively) oriented.

For any face $f \in \mathcal{V}^*$ and $\eta = \pm$, the number of loops of orientation η that surround f will be denoted $\mathcal{N}_q^\eta(f)$, and with $r = \pm$ we further denote by $\mathcal{N}^{\eta,r}(f)$ the number of η -oriented loops with $\text{sgn } q = r$, i.e.

$$\mathcal{N}^{\eta,r}(f) = \sum_{\substack{q \in \frac{1}{2} + \mathbb{Z} \\ \text{sgn } q = r}} \mathcal{N}_q^\eta(f) \quad (5.1)$$




FIGURE 5. Example of level loops resolution through the right-hand rule.

From the definition of the level lines it follows that in any configuration

$$\mathcal{N}^{+,+}(f) + \mathcal{N}^{-,-}(f) \geq |n_f|. \quad (5.2)$$

5.2. Lower bounds on the spin correlation function

Our first set of new results relate the Villain model's BKT phase to depinning in the ZGF. A key intermediary step involves the event described below.

Definition 5.1. Let γ_{yx} be an arbitrary oriented, simple path in \mathcal{G} that begins at y and ends at x , and $e = (x, x')$ an oriented edge such that x' is not in γ_{yx} . For any $q \in \mathbb{Z} + 1/2$, we define as $A_{\gamma_{yx}, e}^q$ the event that there is q -path κ_{xy}^q which begins with the oriented edge e and terminates at y such that $\gamma_{yx} \circ \kappa_{xy}^q$ is a simple loop with orientation $\text{sgn}(q)$.

A particular example of the occurrence of $A_{\gamma_{yx}, e}^q$ where γ_{yx} is the 'L'-shaped path and q is positive is depicted in **Figure 6**. We also need to demarcate the two faces that border the oriented edge e that begins the path κ_{xy}^q , as they will play a special role in our analysis. Given an oriented edge e , we set x^+ and x^- to be the faces to the left and right of the edge, respectively.

We now show that the two point function of the Villain model can be bounded by the probability of the event described above. It should be emphasized here that, while the path γ_{yx} is arbitrary, for the following bound it must be preselected (i.e. not adjusted dynamically to the realization of the field n).

Theorem 5.2. *Given a dual pair of the Villain model on a finite planar graph \mathcal{G} and the corresponding ZGF on \mathcal{G}^* , for any path γ_{yx} and an oriented edge $e = (x, x')$ which does not intersect it, at any $q \in \mathbb{Z} + \frac{1}{2}$,*

$$\langle \sigma_x \cdot \sigma_y \rangle_\beta^{\text{Vill}, \mathcal{G}} \geq \mathbb{P}_{1/\beta}^{\text{ZGF}, \mathcal{G}^*} \left[A_{\gamma_{yx}, e}^q \right]. \quad (5.3)$$

Furthermore, (5.3) holds also for q replaced by $\tilde{q}(n)$ which is determined by the values of n on the pair of faces sharing the edge e (i.e. by $\{n(x^+), n(x^-)\}$).




FIGURE 6. A schematic depiction of the event A_{xy}^q for a positive value of q . The dots indicate the points on which the field n is defined, and the marks H, L the sites whose n values are uncovered in the κ_{xy}^q -exploration process.

The bound stated in (1.1) is obtained through a summation of the above, in a method which applies quite generally to doubly-periodic planar graphs. To convey the argument in a relatively simple way we present it in the context of \mathbb{Z}^2 .

By (5.3) fast decay of spin correlations implies that in the corresponding ZGF long level lines occur only rarely. For a more quantitative statement, we next combine this with the observation that any loop which encircles a face $f \in (\mathbb{Z}^2)^*$, at distance R from it, implies that the conditions for the event $A_{xy,e}^q$ in (5.3) are met for at least four distinct pairs $\{x, y\}$, with $\|x - y\| \geq R$, for which the difference vectors $y - x$ lie each in a different quadrant of \mathbb{Z}^2 .

For this purpose, consider the standard quadrants of \mathbb{Z}^2 shifted so that their intersection is the face $f_0 \in (\mathbb{Z}^2)^*$ centered at $(1/2, 1/2)$ and their corners form the four vertices of f_0 . Let Q_j be the collection of pairs of sites of $\{x, y\}$ one on the vertical and the other on the horizontal part of the boundary of Q_j . As depicted in Figure 7, each q loop encircling f , of orientation $\text{sgn } q$ implies the existence of at least one pair of sites in each Q_j ($j = 1, \dots, 4$) which are linked by a path meeting the conditions for κ_{xy}^q , whose first step is into the corresponding quadrant. This observation leads to the following statement.

6. PROOF OF THE ZGF BOUND ON THE SPIN CORRELATION FUNCTION

For the rest of this section, we will fix an oriented, simple path γ_{yx} and an oriented edge $e = (x, x')$ such that x' is disjoint from γ_{yx} .

6.1. Reduction to a conditional expectation

In this section we prove [Theorem 5.2](#) taken as granted the positivity of what we term the ZGF's stiffness modulus. The proof of this general property of the ZGF on finite graphs is presented to the next section.

In preparation for the proof, let us note that from the duality relation [Section 4.2](#) one may conclude the following

$$\begin{aligned} \langle \sigma_x \cdot \sigma_y \rangle_\beta^{\text{Vill}, \mathcal{G}} &= \mathbb{E}_{1/\beta}^{\text{ZGF}, \mathcal{G}^*} \left[T_{\gamma_{yx}}^+ \right] \geq \mathbb{E}_{1/\beta}^{\text{ZGF}, \mathcal{G}^*} \left[T_{\gamma_{yx}}^+ \chi_{A_{\gamma_{yx}, e}^{\tilde{q}}} \right] = \\ &= \mathbb{P}_{1/\beta}^{\text{ZGF}, \mathcal{G}^*} \left[A_{xy}^q \right] \mathbb{E}_{1/\beta}^{\text{ZGF}, \mathcal{G}^*} \left[T_{\gamma_{yx}}^+ | A_{\gamma_{yx}, e}^{\tilde{q}} \right] \end{aligned} \quad (6.1)$$

where $T_{\gamma_{yx}}^+$ is the function defined in [\(4.6\)](#), χ_A is the indicator function of the event A and the last factor is the conditional expectation of $T_{\gamma_{yx}}^+$ conditioned on the event $A_{\gamma_{yx}, e}^{\tilde{q}}$. The inequality remains true when $T_{\gamma_{yx}}^+$ is replaced by $T_{\gamma_{yx}}^-$.

Thus, to prove [Theorem 5.2](#) it is shown below that for any \tilde{q} which depends only on the values of n on the two faces whose boundaries include the edge e :

$$\mathbb{E}_{1/\beta}^{\text{ZGF}, \mathcal{G}^*} \left[T_{\gamma_{yx}}^{\text{sgn}(\tilde{q})} | A_{\gamma_{yx}, e}^{\tilde{q}} \right] \geq 1. \quad (6.2)$$

6.2. The Exploration Process

To study the conditional expectation conditioned on the event $A_{\gamma_{yx}, e}^{\tilde{q}}$, it is of help to employ an exploration process, in which the variables $n(u)$ are revealed along a dynamically defined sequence of steps which by design uncover the q -path κ_{xy}^q from x to y meeting the conditions of the event $A_{\gamma_{yx}, e}^{\tilde{q}}$ as in [Definition 5.1](#).

The exploration is constructed so that if $A_{\gamma_{yx}, e}^{\tilde{q}}$ fails that the exploration will stop in a bounded number of steps and reveal this fact. And if the path exists, the exploration will locate it, in a manner which allows a relatively simple description of the conditional distribution of n conditioned on the information revealed by the natural ‘‘stopping time’’.

Crucially, the process outputs two sets of faces: \mathcal{L} a collection of faces on which $n < q$ (lower), and \mathcal{H} a collection of faces on which $n > q$ (higher), see [Figure 6](#), such that κ_{xy}^q will be measurable with respect to the values of n on $\mathcal{L} \cup \mathcal{H}$.

We now outline the steps of the process, starting with the case of preselected value of q :

- (1) Start by revealing the values of n on $\mathcal{H}_1 := x^+$ and $\mathcal{L}_1 := x^-$, the pair of faces along the initial edge step e . In case $n_{x^-} > q$ or $n_{x^+} < q$, we learned that $A_{\gamma_{yx},e}^{\tilde{q}}$ fails, and the exploration stops. Otherwise we move to the next step.
- (2) To define the $(k+1)$ th step given all k steps before, expose the faces surrounding w_k in a counter-clockwise fashion, starting from the one to the right of the edge (w_{k-1}, w_k) , until an is reached which has $n < q$ on its right and $n > q$ on its left. For reasons explained above, such an edge will be reached. If the added edge does not terminate on γ_{yx} , the process is repeated. Otherwise the exploration stops.
- (3) From the thus constructed path one determines whether the event $A_{\gamma_{yx},e}^q$ occurred or not. It does iff the path terminates at y and complements γ_{xy} into a loop of the relevant orientation, in which case the constructed path is the path κ_{xy}^q by which $A_{\gamma_{yx},e}^q$ is defined.

It is of relevance to note that in case of success all faces that are revealed in this process have $n < q$ to the right of the path and $n > q$ on the left. Their respective unions are the sets \mathcal{L} and \mathcal{H} respectively.

Though we shall not use it in the argument, let us note that, in the standard percolation terminology, the \mathcal{L} set is connected while the \mathcal{H} set is $*$ -connected.

For any $q \in \mathbb{Z} + 1/2$, define \mathcal{F}_{xy}^q to be the σ -algebra generated by the values of the field n on $\mathcal{L} \cup \mathcal{H}$, i.e. the faces exposed by the exploration process for q .

The above construction extends simply to the more general class of exploration processes for paths $\kappa_{xy}^{\tilde{q}}$ whose q is given by a function \tilde{q} which depends only on the first two bits of information, i.e. n_{x^+} and n_{x^-} . Naturally, the corresponding σ algebra is denoted $\mathcal{F}_{xy}^{\tilde{q}}$.

The observations made above suffice to conclude the following statement.

Lemma 6.1. *For any function $\tilde{q} : (n_{x^+}, n_{x^-}) \mapsto \mathbb{Z} + \frac{1}{2}$, the event $A_{\gamma_{yx},e}^{\tilde{q}}$ is $\mathcal{F}_{xy}^{\tilde{q}}$ -measurable. In addition, if $\tilde{q} > 0$, the interior of $\kappa_{xy}^{\tilde{q}} \cup \gamma_{yx}$ includes \mathcal{H} and is disjoint from \mathcal{L} . If $\tilde{q} < 0$, then \mathcal{H} is disjoint from the interior of $\kappa_{xy}^{\tilde{q}} \cup \gamma_{yx}$ and \mathcal{L} is included in it.*

6.3. Proof of Theorem 5.2

We start with the simpler statement, proving (5.3) for any preselected value of $q = \mathbb{Z} + 1/2$.

First consider $q > 0$. The first stipulation of Lemma 6.1 can be restated as

$$\mathbb{E} \left[\chi_{A_{xy}^q} \mid \mathcal{F}_{xy}^q \right] (n) = \chi_{A_{xy}^q}(n). \quad (6.3)$$

For each specified n let

$$A(n) := \mathcal{H}(n), \quad B(n) := \mathcal{L}(n) \cup \partial\mathcal{V}^* \quad (6.4)$$

and let $F_A : A \rightarrow \mathbb{Z}$ and $F_B : B \rightarrow \mathbb{Z}$ denote the constrained values of n on these two sets (with $F_B|_{\partial\mathcal{V}^*} = 0$ under the Dirichlet boundary conditions).

It is now important to note that due to the nature of the exploration process, under the event A_{xy}^q the uncovered values of n satisfy

$$\min_{u \in A} F_A(u) - 1 \geq \max_{u \in B} F_B(u). \quad (6.5)$$

Summing over the values which n may assume in the complement of $A \cup B$ we find that for any $n \in A_{xy}^q$:

$$\begin{aligned} \frac{\mathbb{E} \left[T_{\gamma_{yx}}^+ \chi_{A_{xy}^q} \mid \mathcal{F}_{xy}^q \right] (n)}{\mathbb{E} \left[\chi_{A_{xy}^q} \mid \mathcal{F}_{xy}^q \right] (n)} &= \frac{\sum_{n: \mathcal{V}^* \rightarrow \mathbb{Z}} \exp \left(-\frac{1}{2} \lambda \|\nabla n + \Gamma_{x,y}\|^2 \right) \chi_{\{n|_A=F_A\}} \chi_{\{n|_B=F_B\}}}{\sum_{n: \mathcal{V}^* \rightarrow \mathbb{Z}} \exp \left(-\frac{1}{2} \lambda \|\nabla n\|^2 \right) \chi_{\{n|_A=F_A\}} \chi_{\{n|_B=F_B\}}} \\ &= \frac{\sum_{n: \mathcal{V}^* \rightarrow \mathbb{Z}} \exp \left(-\frac{1}{2} \lambda \|\nabla n\|^2 \right) \chi_{\{n|_A=F_A-1\}} \chi_{\{n|_B=F_B\}}}{\sum_{n: \mathcal{V}^* \rightarrow \mathbb{Z}} \exp \left(-\frac{1}{2} \lambda \|\nabla n\|^2 \right) \chi_{\{n|_A=F_A\}} \chi_{\{n|_B=F_B\}}} \end{aligned} \quad (6.6)$$

where for the last ratio we employed the change of variables in which n is increased over the set enclosed by the loop $\kappa_{xy}^q \circ \gamma_{yx}$, i.e.

$$n \mapsto n - \chi_{\text{Int}(\kappa_{xy}^q \circ \gamma_{yx})}, \quad (6.7)$$

which, in the terminology used above, is a gauge transformation which erases the shift operator $T_{\gamma_{yx}}^+$ while increasing the boundary values along the set A . By the second stipulation of [Lemma 6.1](#) and the fact that the interior must be disjoint from the boundary, this transformation leaves B unaltered.

For a more succinct expression let us denote

$$Z_{A,B;F_A,F_B}(n) := \sum_{n: \mathcal{V}^* \rightarrow \mathbb{Z}} \exp \left(-\frac{1}{2} \lambda \|\nabla n\|^2 \right) \chi_{\{n|_A=F_A\}} \chi_{\{n|_B=F_B\}}, \quad (6.8)$$

where the n dependence of A, B and F_A, F_B is omitted on the RHS.

In this notation, the above can be summarized by saying that for each $n \in A_{xy}^q$:

$$\frac{\mathbb{E} \left[T_{\gamma_{yx}}^+ \chi_{A_{xy}^q} \mid \mathcal{F}_{xy}^q \right] (n)}{\mathbb{E} \left[\chi_{A_{xy}^q} \mid \mathcal{F}_{xy}^q \right] (n)} = \frac{Z_{A,B;F_A-\chi_A,F_B}(n)}{Z_{A,B;F_A,F_B}(n)}. \quad (6.9)$$

Under the condition (6.5) the partition functions which appear in (6.9) are of a system which is stressed by the imposed boundary conditions, and the stress is higher for the term in the denominator.

This suggests that it may be the case that under the constraint (6.5),

$$\frac{Z_{\lambda; F_A - \chi_A, F_B}^{\text{ZGF}}}{Z_{\lambda; F_A, F_B}^{\text{ZGF}}} \geq 1. \quad (6.10)$$

The suggestive argument should be taken with a grain of salt, since it ignores the possibility that the integrality constraints could introduce some unexpected effects in the propagation of strain due to the increase in the stress introduced through the alteration of the boundary conditions.

Nevertheless, as we show in the next section, under the stated conditions (6.10) holds true. We refer to this assertion as an expression of the positivity of the model's stiffness modulus (τ) since, with a stretch of imagination, the quantity

$$\tau = -\frac{1}{\lambda|A|} \log \frac{Z_{F_A, F_B}}{Z_{F_A - \chi_A, F_B}} \quad (6.11)$$

can be viewed as the pressure needed to be applied along the set A , in order to increase the strain there by 1 (starting from with initial value $F_A - \chi_A$, for which (6.5) holds).

Combined with (6.9), (6.10) allows to deduce the claimed relation (5.3) – for the case q is set at a preselected, positive value.

For $q < 0$ the above is to be repeated with the following changes:

- 1) $T_{\gamma_{xy}}^+$ replaced by $T_{\gamma_{xy}}^-$
- 2) in the condition for successful exploration the loop $\gamma_{yx} \circ \kappa_{xy}^q$ is to be oriented negatively (i.e clockwise) rather than positively
- 3) in the definition of the sets $A(n)$ $B(n)$ the boundary included in the former, i.e. (6.4) replaced by

$$A(n) := \mathcal{H}(n) \cup \partial\mathcal{V}^*, \quad B(n) := \mathcal{L}(n). \quad (6.12)$$

- 4) the expression (6.9) replaced by

$$\frac{\mathbb{E} \left[T_{\gamma_{yx}}^- \chi_{A_{xy}}^q \mid \mathcal{F}_{xy}^q \right] (n)}{\mathbb{E} \left[\chi_{A_{xy}}^q \mid \mathcal{F}_{xy}^q \right] (n)} = \frac{Z_{A, B; F_A, F_B + \chi_B}(n)}{Z_{A, B; F_A, F_B}(n)}. \quad (6.13)$$

The stiffness condition (6.5) still holds, though now it may be written as $\min_{u \in A} F_A(u) \geq \max_{u \in B} F_B(u) + 1$. Under this correspondence, the ratio in (6.13) still represents the reduction of stress, and by the positivity of the stiffness modulus the relation (6.2) holds also in this case.

Finally, to establish the stronger statement that is made in [Theorem 5.2](#) we note that the above argument readily extends to the case where q instead of being constant is selected as a function of n_{x^+}, n_{x^-} (the first variables uncovered by the path exploration process).

Strictly speaking the above derivation of the lower bound on the spin-spin correlation which is expressed in [Theorem 5.2](#) is conditioned on the validity of the principle expressed above. We next turn to its proof.

7. POSITIVITY OF THE $\mathbb{Z}GF$ 'S STIFFNESS MODULUS

For the $\mathbb{Z}GF$ on a finite graph of vertex set \mathcal{V} let $Z_{\lambda;F_A,F_B}^{\mathbb{Z}GF,\mathcal{V}}$ be the restricted partition function of the system whose configurations are constrained to pre-specified values over a pair of disjoint subsets $A, B \subset \mathcal{V}$, over which the field n is given by a pair of functions $F_A : A \rightarrow \mathbb{Z}$ and $F_B : B \rightarrow \mathbb{Z}$ such that

$$F_A(u) \leq 0 \leq F_B(v) \quad \text{for all } u \in A \text{ and } v \in B. \quad (7.1)$$

For brevity we shall refer to the constraints as *boundary values* (though A and B may be arbitrary subsets of \mathcal{V}).

Under the above conditions the change

$$F_B \mapsto F_B + 1, \quad (7.2)$$

(to the function whose values are $[F_B(u) + 1]$) only increases the differences on n within each of the pairs of the constrained values. Thus it can be viewed as an increase of tensile strain. One could expect that under such circumstances the constrained partition function can only decrease (or, in physical terms, the system's free energy can only increase). Our goal in this section is to prove that this is indeed the case.

We prove the statement in a slightly greater generality, in the settings of a nearest-neighbor integer-valued gradient model on a finite graph $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ with the *interaction potential*, which for $\mathbb{Z}GF$ is $U(k) = \frac{1}{2}\lambda k^2$ generalized to a function which is symmetric, $U(k) = U(-k)$, and satisfies the *monotonicity assumption*

$$U(k+1) \geq U(k) \quad \text{for integer } k \geq 0. \quad (7.3)$$

Thus, the model's configuration space is

$$\Omega := \{n : \mathcal{V} \rightarrow \mathbb{Z}\} \quad (7.4)$$

and the weight of configurations $n \in \Omega$ is given by

$$w(n) := \prod_{v \sim w} \exp(-U(n_v - n_w)) \quad (7.5)$$

where $v \sim w$ means that $\{v, w\} \in \mathcal{E}$. The restricted configuration space is

$$\Omega_{F_A, F_B} := \{n \in \Omega : n|_A = F_A, n|_B = F_B\} \quad (7.6)$$

and the corresponding restricted partition function is

$$Z_{F_A, F_B} := \sum_{n \in \Omega_{F_A, F_B}} w(n). \quad (7.7)$$

Theorem 7.1. *In the above setup, for any pair of sets and functions $\{A, B; F_A, F_B\}$ satisfying (7.1),*

$$Z_{F_A, F_B} \geq Z_{F_A, F_B + 1}. \quad (7.8)$$

Let us first outline the proof strategy. Consider the *metric graph*, in which the edges in \mathcal{E} are metrized as segments of unit length. Suppose that the function n , initially defined on the vertex set \mathcal{V} , admits an extension to a *continuous* function on the metric graph, in such a way that the domain Markov property is preserved (for instance, the ZGF model permits such an extension by interpolating the values at vertices via independent Brownian bridges). Then one may perform the following procedure to transform a function n on the metric graph with the boundary values $F_A, F_B + 1$ to a function $T(n)$ on the metric graph with boundary values F_A, F_B :

- (1) Let C^0 be the union of the connected components of $\{s: n(s) \leq 0\}$ which intersect A (here s stands for points of the metric graph).
Similarly let $C^{1/2}$ be the union of the connected components of $\{s: n(s) \geq 1/2\}$ which intersect B .
Importantly, the two sets are disjoint due to our assumption that $F_A \leq 0$ and $F_B \geq 0$.
- (2) Define $T(n)$ by reflecting the values of n around 0 on C^0 and reflecting the values of n around $1/2$ on $C^{1/2}$. Precisely,

$$T(n)_s := \begin{cases} -n_s & s \in C^0, \\ 1 - n_s & s \in C^{1/2}, \\ n_s & s \notin C^0 \cup C^{1/2}. \end{cases} \quad (7.9)$$

- (3) Apply a global sign change to all values.

The domain Markov property and the fact that U is an even function implies that the above procedure is a measure-preserving map from the set of functions on the metric graph with boundary values $F_A, F_B + 1$ to the set of functions with boundary values F_A, F_B (the idea that a reflection of this type is measure preserving is commonly used in the famous *reflection principle* for Brownian motion). The existence of such a measure-preserving mapping implies the inequality (7.8).

The above description is informal. In particular, the idea of extending the model from the vertex set to a continuous function on the metric graph while preserving the domain Markov property needs to be explained. While this is possible for the ZGF model (via Brownian bridges), it is not clear how to perform such an extension for the more general interaction potentials U .

Thus, our proof follows a variant of the above description, in which we avoid the extension of the model to the metric graph. Instead, we attach to each edge a pair of boolean variables which serve as a replacement for the indicator functions of whether the levels 0 and $1/2$

were crossed by the continuous extension on that edge. Such ideas were used before in [6] and references within.

We proceed to give a formal proof of (7.8). Throughout we fix $F_A : A \rightarrow \mathbb{Z}$ and $F_B : B \rightarrow \mathbb{Z}$ satisfying (7.1).

For a set of edges $E \subset \mathcal{E}$, we write $\mathcal{P}(E) := \{\omega : E \rightarrow \{0, 1\}\}$ and identify a function $\omega \in \mathcal{P}(E)$ with the set of edges $\{e \in E : \omega_e = 1\}$ that it characterizes.

Given $n \in \Omega$, an edge $\{v, w\} \in \mathcal{E}$ and $m \in \frac{1}{2}\mathbb{Z}$, define

$$p_{n, \{v, w\}}^m := \max\{1 - e^{U(n_v - n_w) - U(2m - n_v - n_w)}, 0\}, \quad (7.10)$$

where one should interpret $p_{n, \{v, w\}}^m$ as the probability that, given n_v and n_w , the ‘‘continuous extension of n to $\{v, w\}$ ’’ *does not* cross the level m . The relevant properties of these probabilities are the following

$$p_{n, \{v, w\}}^m \text{ depends only on the } \textit{unordered} \text{ pair } \{n_v, n_w\}, \quad (7.11)$$

$$\text{If } \min\{n_v, n_w\} \leq m \text{ and } \max\{n_v, n_w\} \geq m \text{ then } p_{n, \{v, w\}}^m = 0, \quad (7.12)$$

using in both cases the fact that U is an even function and using the monotonicity assumption (7.3) to prove (7.12). In addition,

$$\text{if } n' \in \Omega \text{ satisfies } \{n'_v, n'_w\} = \{2m - n_v, 2m - n_w\} \text{ then } p_{n', \{v, w\}}^m = p_{n, \{v, w\}}^m, \quad (7.13)$$

and if instead either $(n'_v, n'_w) = (2m - n_v, n_w)$ or $(n'_v, n'_w) = (n_v, 2m - n_w)$ then

$$\exp(-U(n'_v - n'_w))(1 - p_{n', \{v, w\}}^m) = \exp(-U(n_v - n_w))(1 - p_{n, \{v, w\}}^m), \quad (7.14)$$

which is verified by separating to two cases according to whether $U(2m - n_v - n_w) \geq U(n_v - n_w)$ or not.

Given $n : \mathcal{V} \rightarrow \mathbb{Z}$, a set of edges $E \subset \mathcal{E}$, $m \in \frac{1}{2}\mathbb{Z}$ and an edge configuration $\omega \in \mathcal{P}(E)$, let

$$p_{n, E}^m(\omega) := \prod_{e \in E} (p_{n, e}^m)^{\omega_e} (1 - p_{n, e}^m)^{1 - \omega_e}, \quad (7.15)$$

with 0^0 defined to be 1. This should be interpreted as the probability, given n , of sampling the edge configuration ω on the edges in E using the probabilities (7.10). Recalling (7.7), it is clear that we have

$$Z_{G_A, G_B} = \sum_{\substack{n \in \Omega_{G_A, G_B} \\ \omega^0, \omega^{1/2} \in \mathcal{P}(E)}} w(n) p_{n, E}^0(\omega^0) p_{n, E}^{1/2}(\omega^{1/2}) \quad (7.16)$$

for any $\emptyset \neq E \subset \mathcal{E}$, $G_A : A \rightarrow \mathbb{Z}$ and $G_B : B \rightarrow \mathbb{Z}$.

Given $\omega^0, \omega^{1/2} \in \mathcal{P}(E)$, write $C^0(\omega^0)$ for the union of the connected components of A in the graph (\mathcal{V}, ω^0) . Similarly write $C^{1/2}(\omega^{1/2})$ for the union of the connected components of B in the graph $(\mathcal{V}, \omega^{1/2})$. We make the following observation.

Claim 1: Let $n \in \Omega_{F_A, F_B+1}$ and let $\omega^0, \omega^{1/2} \in \mathcal{P}(\mathcal{E})$ be such that $p_{n, \mathcal{E}}^0(\omega^0) > 0$ and $p_{n, \mathcal{E}}^{1/2}(\omega^{1/2}) > 0$. Then $n_v \leq 0$ for all $v \in C^0(\omega^0)$ and $n_v \geq 1$ for all $v \in C^{1/2}(\omega^{1/2})$. In particular, the sets $C^0(\omega^0)$ and $C^{1/2}(\omega^{1/2})$ are disjoint.

Proof. For both $m = 0$ and $m = 1/2$, property (7.12) shows that $p_{n, \{v, w\}}^m = 0$ when $\min\{n_v, n_w\} \leq 0$ and $\max\{n_v, n_w\} \geq 1$. Then, also $\omega_{\{v, w\}}^0 = \omega_{\{v, w\}}^{1/2} = 0$ by our assumptions that $p_{n, \mathcal{E}}^0(\omega^0) > 0$ and $p_{n, \mathcal{E}}^{1/2}(\omega^{1/2}) > 0$. The claim now follows from the fact that $n \in \Omega_{F_A, F_B+1}$ and the assumption that $F_A \leq 0$ and $F_B \geq 0$. \square

We wish to extend the formula (7.16) to a version in which only the parts of $\omega^0, \omega^{1/2}$ which intersect $C^0(\omega^0), C^{1/2}(\omega^{1/2})$, respectively, are enumerated upon. To this end, we introduce the following additional notation. For a set $S \subset \mathcal{V}$, write

$$E(S) := \{e \in \mathcal{E} : e \cap S \neq \emptyset\} \quad (7.17)$$

for the set of edges with at least one endpoint in S , and let

$$O^0(S) := \{\omega \in \mathcal{P}(E(S)) : C^0(\omega) = S\}, \quad (7.18)$$

$$O^{1/2}(S) := \{\omega \in \mathcal{P}(E(S)) : C^{1/2}(\omega) = S\}. \quad (7.19)$$

Note that $O^0(S) = \emptyset$ if $A \not\subset S$ and similarly $O^{1/2}(S) = \emptyset$ if $B \not\subset S$. An element $\omega \in O^0(S)$ can be thought of as the result of exploring the edges reachable from A in some edge set ω^0 with $C^0(\omega^0) = S$. A similar description applies to $O^{1/2}(S)$. With this in mind, one easily verifies the following extension of (7.16),

$$Z_{G_A, G_B} = \sum_{\substack{n \in \Omega_{G_A, G_B} \\ C^0, C^{1/2} \subset \mathcal{V}}} \sum_{\substack{\omega^0 \in O^0(C^0) \\ \omega^{1/2} \in O^{1/2}(C^{1/2})}} w(n) p_{n, E(C^0)}^0(\omega^0) p_{n, E(C^{1/2})}^{1/2}(\omega^{1/2}) \quad (7.20)$$

for any $G_A : A \rightarrow \mathbb{Z}$ and $G_B : B \rightarrow \mathbb{Z}$.

Lastly, given disjoint $C^0, C^{1/2} \subset \mathcal{V}$ with $A \subset C^0$ and $B \subset C^{1/2}$ we define a mapping $T_{C^0, C^{1/2}} : \Omega_{F_A, F_B+1} \rightarrow \Omega_{-F_A, -F_B}$ by

$$T_{C^0, C^{1/2}}(n) := \begin{cases} -n_s & s \in C^0, \\ 1 - n_s & s \in C^{1/2}, \\ n_s & s \notin C^0 \cup C^{1/2}. \end{cases} \quad (7.21)$$

The mapping $T_{C^0, C^{1/2}}$ is easily seen to be one-to-one (for fixed values of $C^0, C^{1/2}$).

To establish the inequality (7.8), we proceed to compare the partition functions Z_{F_A, F_B} and Z_{F_A, F_B+1} using the identity (7.20) and the mapping (7.21). On the one hand,

$$Z_{F_A, F_B+1} = \sum_{\substack{n \in \Omega_{F_A, F_B+1} \\ C^0, C^{1/2} \subset \mathcal{V}}} \sum_{\substack{\omega^0 \in O^0(C^0) \\ \omega^{1/2} \in O^{1/2}(C^{1/2})}} w(n) p_{n, E(C^0)}^0(\omega^0) p_{n, E(C^{1/2})}^{1/2}(\omega^{1/2}) \quad (7.22)$$

while on the other hand,

$$\begin{aligned}
Z_{F_A, F_B} &= Z_{-F_A, -F_B} \\
&= \sum_{\substack{\bar{n} \in \Omega_{-F_A, -F_B} \\ C^0, C^{1/2} \subset \mathcal{V}}} \sum_{\substack{\omega^0 \in O^0(C^0) \\ \omega^{1/2} \in O^{1/2}(C^{1/2})}} w(\bar{n}) p_{\bar{n}, E(C^0)}^0(\omega^0) p_{\bar{n}, E(C^{1/2})}^{1/2}(\omega^{1/2}) \\
&\geq \sum_{\substack{\bar{n} \in \Omega_{-F_A, -F_B} \\ C^0, C^{1/2} \subset \mathcal{V} \text{ disjoint}}} \sum_{\substack{\omega^0 \in O^0(C^0) \\ \omega^{1/2} \in O^{1/2}(C^{1/2})}} w(\bar{n}) p_{\bar{n}, E(C^0)}^0(\omega^0) p_{\bar{n}, E(C^{1/2})}^{1/2}(\omega^{1/2}) 1(\bar{n}, C^0, C^{1/2}) \\
&= \sum_{\substack{n \in \Omega_{F_A, F_B+1} \\ C^0, C^{1/2} \subset \mathcal{V} \text{ disjoint}}} \sum_{\substack{\omega^0 \in O^0(C^0) \\ \omega^{1/2} \in O^{1/2}(C^{1/2})}} w(T(n)) p_{T(n), E(C^0)}^0(\omega^0) p_{T(n), E(C^{1/2})}^{1/2}(\omega^{1/2})
\end{aligned} \tag{7.23}$$

where in the first equality we used the definition (7.7) and the fact that U is an even function, in the inequality the symbol $1(\bar{n}, C^0, C^{1/2})$ is defined to equal 1 if $\bar{n} = T_{C^0, C^{1/2}}(n)$ for some $n' \in \Omega_{F_A, F_B+1}$ and defined to equal 0 otherwise, and where in the last equality we abbreviated $T_{C^0, C^{1/2}}(n)$ to $T(n)$ for typographical reasons. The last equality uses the fact that $T_{C^0, C^{1/2}}$ is one-to-one. Comparing (7.22) with (7.23) and recalling Claim 1, we conclude that the requisite inequality (7.8) follows from the following statement.

Claim 2: Let $n \in \Omega_{F_A, F_B+1}$. Let $C^0, C^{1/2} \subset \mathcal{V}$ and let $\omega^0 \in O^0(C^0)$ and $\omega^{1/2} \in O^{1/2}(C^{1/2})$. Suppose that $p_{n, E(C^0)}^0(\omega^0) > 0$ and $p_{n, E(C^{1/2})}^{1/2}(\omega^{1/2}) > 0$. Then

$$w(T(n)) p_{T(n), E(C^0)}^0(\omega^0) p_{T(n), E(C^{1/2})}^{1/2}(\omega^{1/2}) \geq w(n) p_{n, E(C^0)}^0(\omega^0) p_{n, E(C^{1/2})}^{1/2}(\omega^{1/2}) \tag{7.24}$$

where we again abbreviate $T_{C^0, C^{1/2}}(n)$ to $T(n)$.

Proof. Unwrapping the definitions, the right-hand side of (7.24) equals

$$\prod_{v \sim w} \exp(-U(n_v - n_w)) \prod_{e \in E(C^0)} (p_{n, e}^0)^{\omega_e^0} (1 - p_{n, e}^0)^{1 - \omega_e^0} \prod_{e \in E(C^{1/2})} (p_{n, e}^{1/2})^{\omega_e^{1/2}} (1 - p_{n, e}^{1/2})^{1 - \omega_e^{1/2}}. \tag{7.25}$$

Similarly, the left-hand side of (7.24) equals

$$\prod_{v \sim w} \exp(-U(T(n)_v - T(n)_w)) \prod_{e \in E(C^0)} (p_{T(n), e}^0)^{\omega_e^0} (1 - p_{T(n), e}^0)^{1 - \omega_e^0} \prod_{e \in E(C^{1/2})} (p_{T(n), e}^{1/2})^{\omega_e^{1/2}} (1 - p_{T(n), e}^{1/2})^{1 - \omega_e^{1/2}}. \tag{7.26}$$

We prove the claim by proving that the terms corresponding to each edge $\{v, w\} \in \mathcal{E}$ in (7.26) are at least as large as the terms corresponding to the same edge in (7.25) (in fact, in all but one case equality holds). Fix an edge $\{v, w\} \in \mathcal{E}$ and consider the following cases.

Case 1: $v, w \in \mathcal{V} \setminus (C^0 \cup C^{1/2})$. We need to show that

$$\exp(-U(T(n)_v - T(n)_w)) \geq \exp(-U(n_v - n_w)) \quad (7.27)$$

and in this case equality holds since $T(n)_v = n_v$ and $T(n)_w = n_w$ by (7.21).

Case 2: $v, w \in C^0$. We need to show that

$$\begin{aligned} \exp(-U(T(n)_v - T(n)_w)) (p_{T(n),\{v,w\}}^0)^{\omega_{\{v,w\}}^0} (1 - p_{T(n),\{v,w\}}^0)^{1-\omega_{\{v,w\}}^0} \\ \geq \exp(-U(n_v - n_w)) (p_{n,\{v,w\}}^0)^{\omega_{\{v,w\}}^0} (1 - p_{n,\{v,w\}}^0)^{1-\omega_{\{v,w\}}^0}. \end{aligned} \quad (7.28)$$

In this case $T(n)_v = -n_v$ and $T(n)_w = -n_w$ (by (7.21)). Thus, in fact, equality holds, as $U(T(n)_v - T(n)_w) = U(n_v - n_w)$ since U is an even function and $p_{T(n),\{v,w\}}^0 = p_{n,\{v,w\}}^0$ by (7.13).

A similar analysis applies to the case $v, w \in C^{1/2}$.

Case 3: $v \in C^0$, $w \in \mathcal{V} \setminus (C^0 \cup C^{1/2})$. In this case we necessarily have $\omega^0(\{v, w\}) = 0$, so that we need to show

$$\exp(-U(T(n)_v - T(n)_w)) (1 - p_{\{v,w\}}^0(T(n))) \geq \exp(-U(n_v - n_w)). \quad (7.29)$$

This time $T(n)_v = -n_v$ and $T(n)_w = n_w$ and equality holds due to (7.14).

A similar analysis applies to the case $v \in C^{1/2}$, $w \in \mathcal{V} \setminus (C^0 \cup C^{1/2})$.

Case 4: $v \in C^0$, $w \in C^{1/2}$. In this case we necessarily have $n_v \leq 0$, $n_w \geq 1$ and $\omega^0(\{v, w\}) = \omega^{1/2}(\{v, w\}) = 0$ by Claim 1. We need to show that

$$\exp(-U(T(n)_v - T(n)_w)) (1 - p_{T(n),\{v,w\}}^0) (1 - p_{T(n),\{v,w\}}^{1/2}) \geq \exp(-U(n_v - n_w)). \quad (7.30)$$

Suppose first that $n_v = 0$. In this case $T(n)_v = n_v$ so the same analysis as the case $v \in \mathcal{V} \setminus (C^0 \cup C^{1/2})$, $w \in C^{1/2}$ applies.

Suppose next that $n_v < 0$. In this case $T(n)_v = -n_v \geq 1$ and $T(n)_w = 1 - n_w \leq 0$ so that $p_{T(n),\{v,w\}}^0 = p_{T(n),\{v,w\}}^{1/2} = 0$ by (7.12). Thus we need only show that

$$\exp(-U(T(n)_v - T(n)_w)) \geq \exp(-U(n_v - n_w)). \quad (7.31)$$

This follows from the fact that $|T(n)_v - T(n)_w| = |-n_v - (1 - n_w)| = |n_v - n_w| - 1$ using our monotonicity assumption (7.3). \square

8. THE RELATION OF THE SPIN CORRELATION FUNCTION WITH HEIGHT FLUCTUATIONS

As the last preparatory step towards the proof of [Theorem 5.3](#), we note the following deterministic relation. To state it, we recall that Q_j are the four collections of pairs of sites on the boundaries of short translates of the standard quadrants of \mathbb{Z}^2 (cf. [Figure 7](#)). For any $x, y \in Q_j$, we also set γ_{yx}^j to be the oriented path on Q_j from y to x , starting with the oriented edge e directed towards the interior of the corresponding quadrant.

Lemma 8.1. *For any $q \in \mathbb{Z} + \frac{1}{2}$,*

$$4 \cdot \mathcal{N}_q^{\text{sgn}(q)}(f_0) \leq \sum_{j=1}^4 \sum_{(x,y) \in Q_j} \chi_{A_{\gamma_{yx}^j, e}^q}$$

with χ_A the characteristic function of the event A (of [Definition 5.1](#)).

Proof. We begin by proving that for $q > 0$

$$\mathcal{N}_q^+(f_0) \leq \sum_{(x,y) \in Q_1} \chi_{A_{\gamma_{yx}^1, e}^q}.$$

Let κ be a simple, counter-clockwise oriented q -loop surrounding f_0 . For topological reasons, this loop must include the oriented edge e for some x in the horizontal part of Q_1 . Let x be the leftmost such edge. Similarly, κ must intersect the vertical portion of Q_1 ; we define y to be the first such intersection in counter-clockwise order, beginning at x . Then, the portion of the q -loop that begins at x and ends at y satisfies the requirements of the event $A_{\gamma_{yx}^1, e}^q$. Thus, every such q -loop implies the existence of a pair of points in Q_1 , and the desired inequality follows.

For negative values of q , we consider clockwise oriented q -loops, and define x to be the bottommost point on the vertical portion of Q_1 that includes the appropriate oriented edge. We set y to be the first intersection with the horizontal portion, in clockwise order beginning at x . The argument then follows. For different values of j , the construction is identical, up to rotations. \square

Proof of [Theorem 5.3](#). Fix x and y in \mathcal{G} . Using [Theorem 5.2](#), we learn that for any function $\tilde{q}(n)$ with values in $\mathbb{Z} + \frac{1}{2}$ which depends only on $\{n_{x^+}, n_{x^-}\}$

$$\langle \sigma_x \cdot \sigma_y \rangle_\beta^{\text{Vill}, \mathcal{G}} \geq \mathbb{P}_\lambda^{\text{ZGF}, \mathcal{G}^*} \left[A_{\gamma_{yx}^j, e}^{\tilde{q}} \right] = \mathbb{E}_\lambda^{\text{ZGF}, \mathcal{G}^*} \left[\mathbb{P}_\lambda^{\text{ZGF}, \mathcal{G}^*} \left[A_{\gamma_{yx}^j, e}^{\tilde{q}} \mid n_{x^+}, n_{x^-} \right] \right]. \quad (8.1)$$

We apply that to the function \tilde{q} which returns the value of q maximizing the conditional probability of A_{xy}^q to occur, i.e.

$$\tilde{q}(n) := \operatorname{argmax}_{q \in \{n_{x^-} + \frac{1}{2}, \dots, n_{x^+} - \frac{1}{2}\}} \left\{ \mathbb{P}_\lambda^{\text{ZGF}, \mathcal{G}^*} \left[A_{\gamma_{yx}^j, e}^{\tilde{q}} \mid n_{x^+}, n_{x^-} \right] \right\}.$$

The maximum is attained since the specified values of n leave only $|n_{x^+} - n_{x^-}|$ options for q at which the probability does not vanish.

Recalling the definitions

$$\mathcal{N}^{+,+}(f_0) = \sum_{q \in \mathbb{Z} + \frac{1}{2}; q > 0} \mathcal{N}_q^+(f_0), \quad \mathcal{N}^{-,-}(f_0) = \sum_{q \in \mathbb{Z} + \frac{1}{2}; q < 0} \mathcal{N}_q^-(f_0),$$

and summing the inequality in [Lemma 8.1](#) over all $q \in \mathbb{Z} + 1/2$, we may conclude that

$$4 \cdot \mathbb{E}[\mathcal{N}^{+,+}(f_0) + \mathcal{N}^{-,-}(f_0)] \leq \sum_{q \in \mathbb{Z} + \frac{1}{2}} \sum_{j=1}^4 \sum_{(x,y) \in Q_j} \mathbb{P}[A_{\gamma_{yx},e}^q] = \sum_{j=1}^4 \sum_{(x,y) \in Q_j} \mathbb{E} \left[\sum_{q \in \mathbb{Z} + \frac{1}{2}} \chi_{A_{\gamma_{yx},e}^q} \right]. \quad (8.2)$$

Once the sum over q is placed within the expectation, its convergence becomes clear since

$$\begin{aligned} \mathbb{E} \left[\sum_{q \in \mathbb{Z} + \frac{1}{2}} \chi_{A_{\gamma_{yx},e}^q} \right] &\leq \mathbb{E} \left[|n_{x^+} - n_{x^-}| \chi_{A_{\gamma_{yx},e}^{\tilde{q}}} \right] \\ &\leq \mathbb{E} \left[|n_{x^+} - n_{x^-}|^{1/\varepsilon} \right]^\varepsilon \mathbb{P} \left[A_{\gamma_{yx},e}^{\tilde{q}} \right]^{1-\varepsilon} \quad (\text{H\"older, for any } \varepsilon > 0) \\ &\leq C_\varepsilon \sqrt{\beta} (\langle \sigma_x \cdot \sigma_y \rangle)^{1-\varepsilon}. \quad (\text{Using (8.1) and defining } C_\varepsilon) \end{aligned} \quad (8.3)$$

where the first inequality is deduced through the intermediate conditioning on the values of n at x^+ and x^- , by noticing that under this information the number of relevant values of q is at most $|n_{x^+} - n_{x^-}|$, and the one whose contribution to the sum is maximal is, by definition $\tilde{q}(n)$.

The finiteness of C_ε is a special case of the Gaussian domination principle which is recalled below, applied to $v = \delta_{x^+} - \delta_{x^-}$. This, combined with the last two inequalities, yields the claimed (5.4). \square

Lemma 8.2. *For the ZGF on any finite graph, $v \in \ker(-\Delta)^\perp$ and $\varepsilon > 0$*

$$\mathbb{E} \left[|\langle v, n \rangle|^{1/\varepsilon} \right]^\varepsilon \leq D_\varepsilon \sqrt{\frac{2\langle v, -\Delta^{-1}v \rangle}{\lambda}} \quad (8.4)$$

where $D_\varepsilon := (\Gamma(1 + \frac{1}{2\varepsilon}))^\varepsilon < \infty$.

Proof. By the well-known Gaussian domination bound (see Remark 2.5 above),

$$\mathbb{E}_\lambda^{\text{ZGF}, \mathcal{G}^*} \left[e^{\langle n, v \rangle} \right] \leq \exp \left(+\frac{1}{2\lambda} \langle v, -\Delta^{-1}v \rangle \right) \quad (v \perp \ker -\Delta). \quad (8.5)$$

This, through the exponential Chebyshev stratagem yields the estimate

$$\mathbb{P} \left[\{ |\langle v, n \rangle| \geq t \} \right] \leq \exp \left(-\frac{\lambda}{2} t^2 / \langle v, -\Delta^{-1}v \rangle \right). \quad (8.6)$$

The claim then follows through the expectation value's layer-cake representation $\mathbb{E}[X] = \int_0^\infty \mathbb{P} \{ X \geq t \} dt$. \square

APPENDIX A. THE REGEV STEPHENS-DAVIDOWITZ MONOTONICITY THEORY

For the completeness of presentation, we summarize below the key points of the Regev Stephens-Davidowitz monotonicity theory [24], and state their relation to the case of interest here.

A lattice \mathcal{L} in \mathbb{R}^k is a set of the form $\mathcal{L} := L\mathbb{Z}^k$ for some $n \in \mathbb{N}$ and $L \in \text{Mat}_k(\mathbb{R})$. Given a positive-definite matrix $A : \mathbb{R}^k \rightarrow \mathbb{R}^k$, a random field $\psi \in \mathcal{L}$ associated with (A, \mathcal{L}) has as its partition function

$$Z_{A, \mathcal{L}} := \sum_{\psi \in \mathcal{L}} \exp\left(-\frac{1}{2}\langle \psi, A\psi \rangle\right), \quad (\text{A.1})$$

and the probability distribution of its configurations is given by their normalized contributions to the sum.

The corresponding moment generating function is

$$\mathbb{M}_{A, \mathcal{L}}[v] := \mathbb{E}_{A, \mathcal{L}}[\exp(\langle v, \psi \rangle)] \quad (v \in \mathbb{R}^k). \quad (\text{A.2})$$

The following result is an important tool for the derivation of monotonicity estimates.

Theorem A.1. *For any lattice \mathcal{L} and a positive matrix $A \geq 0$*

$$\mathbb{M}_{A, \mathcal{L}}[u] \mathbb{M}_{A, \mathcal{L}}[v] \leq \sqrt{\mathbb{M}_{A, \mathcal{L}}[u+v] \mathbb{M}_{A, \mathcal{L}}[u-v]} \quad (u, v \in \mathbb{R}^k). \quad (\text{A.3})$$

One may note that in the Gaussian case, where the moment generating function is the exponential of a quadratic form, the relation holds as equality (due to the Pythagorean theorem).

Proof. Rewriting the LHS in term of the Cartesian product of lattices $\mathcal{L}^2 = \mathcal{L} \oplus \mathcal{L}$:

$$\mathbb{M}_{A, \mathcal{L}}[u] \mathbb{M}_{A, \mathcal{L}}[v] = Z_{A, \mathcal{L}}^{-2} \sum_{\Psi \in \mathcal{L}^2} \exp\left(-\frac{1}{2}\langle \Psi, A \oplus A \Psi \rangle + \left\langle \begin{bmatrix} u \\ v \end{bmatrix}, \Psi \right\rangle\right). \quad (\text{A.4})$$

Under the linear transformation by $T = \begin{bmatrix} \mathbf{1}_k & \mathbf{1}_k \\ \mathbf{1}_k & -\mathbf{1}_k \end{bmatrix}$, the lattice $T\mathcal{L}^2$ is the disjoint union of translates of $(2\mathcal{L})^2$:

$$T\mathcal{L}^2 = \bigsqcup_{w \in \mathcal{L}/2\mathcal{L}} (2\mathcal{L} + w)^2. \quad (\text{A.5})$$

The matrix $\frac{1}{\sqrt{2}}T$ is unitary and $[T, M \oplus M] = 0$ for any $n \times n$ matrix M and. Hence for any $\psi \in \mathcal{L}^2$

$$-\frac{1}{2}\langle \Psi, A \oplus A \Psi \rangle + \left\langle \begin{bmatrix} u \\ v \end{bmatrix}, \Psi \right\rangle = -\frac{1}{4}\langle T\Psi, A \oplus AT\Psi \rangle + \frac{1}{2}\left\langle \begin{bmatrix} u+v \\ u-v \end{bmatrix}, T\Psi \right\rangle. \quad (\text{A.6})$$

Substituting this in (A.4) one gets

$$\begin{aligned} \mathbb{M}_{A,\mathcal{L}}[u] \mathbb{M}_{A,\mathcal{L}}[v] &= Z_{A,\mathcal{L}}^{-2} \sum_{w \in \mathcal{L}/2\mathcal{L}} \left(\sum_{\psi \in \mathcal{L}+w} \exp \left(-\frac{1}{4} \langle \psi, A\psi \rangle + \frac{1}{2} \langle u+v, \psi \rangle \right) \right) \times \\ &\quad \times \left(\sum_{\psi \in \mathcal{L}+w} \exp \left(-\frac{1}{4} \langle \psi, A\psi \rangle + \frac{1}{2} \langle u-v, \psi \rangle \right) \right). \end{aligned} \quad (\text{A.7})$$

In particular, with $v = 0$ the last equation yields

$$\mathbb{M}_{A,\mathcal{L}}[u] = Z_{A,\mathcal{L}}^{-2} \sum_{w \in \mathcal{L}/2\mathcal{L}} \left(\sum_{\psi \in \mathcal{L}+w} \exp \left(-\frac{1}{4} \langle \psi, A\psi \rangle + \frac{1}{2} \langle u, \psi \rangle \right) \right)^2. \quad (\text{A.8})$$

Applying now the Cauchy-Schwarz inequality on the RHS of (A.7) w.r.t. the sum $\sum_{w \in \mathcal{L}/2\mathcal{L}}$ and using (A.8) to re-identify $\mathbb{M}_{A,\mathcal{L}}[u+v]$, $\mathbb{M}_{A,\mathcal{L}}[u-v]$ yields the statement. \square

As was pointed out in [24], from the inequality (A.3) one may deduce *sub-lattice monotonicity* of the moment generating function that was stated above as:

Proposition 2.3.

$$\mathbb{M}_{A,\mathcal{M}}[v] \leq \mathbb{M}_{A,\mathcal{L}}[v] \quad (v \in \mathbb{R}^k, A \geq 0).$$

for any sub-lattice $\mathcal{M} \subseteq \mathcal{L}$.

Proof. Since $\mathcal{M} \subseteq \mathcal{L}$ is a sub-lattice, we have the disjoint decomposition $\mathcal{L} = \bigsqcup_{w \in \mathcal{L}/\mathcal{M}} \mathcal{M} + w$. Hence

$$\begin{aligned} Z_{A,\mathcal{L}} Z_{A,\mathcal{M}} \mathbb{M}_{A,\mathcal{M}}[v] &= \left(\sum_{w \in \mathcal{L}/\mathcal{M}} \sum_{\psi \in \mathcal{M}+w} \exp \left(-\frac{1}{2} \langle \psi, A\psi \rangle \right) \right) Z_{A,\mathcal{M}} \mathbb{M}_{A,\mathcal{M}}[v] \\ &= \sum_{w \in \mathcal{L}/\mathcal{M}} \exp \left(-\frac{1}{2} \langle w, Aw \rangle \right) Z_{A,\mathcal{M}}^2 \mathbb{M}_{A,\mathcal{M}}[-Aw] \mathbb{M}_{A,\mathcal{M}}[v]. \end{aligned}$$

Applying now (A.3) together with $\sqrt{ab} \leq \frac{1}{2}(a+b)$ on the last two factors of \mathbb{M} above yields

$$\begin{aligned} Z_{A,\mathcal{L}} Z_{A,\mathcal{M}} \mathbb{M}_{A,\mathcal{M}}[v] &\leq \sum_{w \in \mathcal{L}/\mathcal{M}} \exp \left(-\frac{1}{2} \langle w, Aw \rangle \right) Z_{A,\mathcal{M}}^2 \frac{1}{2} (\mathbb{M}_{A,\mathcal{M}}[-Aw+v] + \mathbb{M}_{A,\mathcal{M}}[-Aw-v]) \\ &= \sum_{w \in \mathcal{L}/\mathcal{M}} \exp \left(-\frac{1}{2} \langle w, Aw \rangle \right) Z_{A,\mathcal{M}}^2 \mathbb{M}_{A,\mathcal{M}}[-Aw+v] \\ &\quad (\mathcal{M} \mapsto -\mathcal{M} \text{ symmetry}) \\ &= Z_{A,\mathcal{L}} Z_{A,\mathcal{M}} \mathbb{M}_{A,\mathcal{L}}[v]. \quad (\text{Use } \mathcal{L} = \bigsqcup_{w \in \mathcal{L}/\mathcal{M}} \mathcal{M} + w \text{ again}) \end{aligned}$$

\square

While it is not used here, it is of interest to note also the following result of [24] (Theorem 5.1 there).

Theorem A.2. *If $\mathcal{N}, \mathcal{M} \subseteq \mathcal{L}$ are sub-lattices then*

$$Z_{A,\mathcal{M}}Z_{A,\mathcal{N}} \leq Z_{A,\mathcal{L}}Z_{A,\mathcal{M} \cap \mathcal{N}} \quad (A \geq 0).$$

In our context this yields new correlation inequalities for general ZGF models, for instance:

$$\mathbb{P}_\lambda^{\text{ZGF}} [\{n_x = n_y\} \cap \{n_u = n_v\}] \geq \mathbb{P}_\lambda^{\text{ZGF}} [\{n_x = n_y\}] \mathbb{P}_\lambda^{\text{ZGF}} [\{n_u = n_v\}]. \quad (\text{A.9})$$

The other property of interest is the operator monotonicity of the moment generating function, i.e. the monotonicity of $\mathbb{M}_{A,\mathcal{L}}[v]$ in the matrix A which specifies the interaction. Its derivation starts with a lower bound on the Hessian of the moment generating function, $(\mathbb{H}\mathbb{M}[u])_{ij} \equiv \partial_{u_i} \partial_{u_j} \mathbb{M}[u]$.

Lemma A.3. *For any \mathcal{L} and $A \geq 0$, as above, the moment generating function $\mathbb{M}[v] = \mathbb{M}_{A,\mathcal{L}}[v]$ satisfies*

$$\frac{1}{\mathbb{M}[u]} \mathbb{H}\mathbb{M}[u] \geq (\mathbb{H}\mathbb{M}[u])|_{u=0} + \frac{1}{\mathbb{M}[u]^2} \nabla \mathbb{M}[u] \otimes (\nabla \mathbb{M}[u]) \quad (u \in \mathbb{R}^k). \quad (\text{A.10})$$

In particular, since

$$\mathbb{H}\mathbb{M}[u] = \mathbb{E} [\exp(\langle u, \psi \rangle) \psi \otimes \psi]; \quad \nabla \mathbb{M}[u] = \mathbb{E} [\exp(\langle u, \psi \rangle) \psi] \quad (\text{A.11})$$

and $\text{tr}(MN) \geq 0$ for any $M, N \geq 0$, (A.10) implies

$$\mathbb{E} [\exp(\langle u, \psi \rangle) \langle \psi, M\psi \rangle] \geq \mathbb{M}[u] \mathbb{E} [\langle \psi, M\psi \rangle] \quad (M \in \text{Mat}_k(\mathbb{R}) : M \geq 0, u \in \mathbb{R}^k). \quad (\text{A.12})$$

Proof. Consider the function F of $u, v \in \mathbb{R}^k$,

$$F(u, v) := \mathbb{M}[u+v] \mathbb{M}[u-v] - \mathbb{M}[u]^2 \mathbb{M}[v]^2 \geq 0$$

where positivity is equivalent to (A.3). For any fixed u , $F(u, 0) = 0$ and hence $v = 0$ is a local minimum for each u . Hence, $(\mathbb{H}F)(u, 0) \geq 0$ (the Hessian is w.r.t. v). A calculation in which it is also used that $\mathbb{E}[\psi] = 0$ due to the symmetry $\mathcal{L} \mapsto -\mathcal{L}$, shows that

$$(\mathbb{H}F)(u, 0) = 2\mathbb{M}[u] \mathbb{E} [\exp(\langle u, \psi \rangle) \psi \otimes \psi] - 2\mathbb{E} [\exp(\langle u, \psi \rangle) \psi] \otimes \mathbb{E} [\exp(\langle u, \psi \rangle) \psi] - 2\mathbb{M}[u]^2 \mathbb{E} [\psi \otimes \psi].$$

The result then follows using (A.11). \square

This lemma is used in the proof of the matrix-monotonicity which was stated above as:

Proposition 2.4. For any lattice \mathcal{L} , and pair of matrices A, B such that $A \geq B \geq 0$

$$\mathbb{M}_{A,\mathcal{L}}[v] \leq \mathbb{M}_{B,\mathcal{L}}[v] \quad (v \in \mathbb{R}^k).$$

Proof. Define the homotopy of covariance matrices

$$[0, 1] \ni t \mapsto tA + (1 - t)B =: A_t.$$

Suffice then to show that $t \mapsto \mathbb{M}_{A_t, \mathcal{L}}[v]$ is monotone decreasing, hence suffice to show $\partial_t \mathbb{M}_{A_t, \mathcal{L}}[v] \leq 0$. But

$$2\partial_t \mathbb{M}_{A_t, \mathcal{L}}[v] = \mathbb{E}_{A_t, \mathcal{L}}[\langle \psi, (A - B)\psi \rangle] \mathbb{M}_{A_t, \mathcal{L}}[v] - \mathbb{E}_{A_t, \mathcal{L}}[\langle \psi, (A - B)\psi \exp(\langle u, \psi \rangle)] \leq 0$$

the last inequality is (A.12) with $M = A - B \geq 0$. \square

Corollary A.4. *For the ZGF on any graph, the variances*

$$\mathbb{E}_{A, \lambda}^{\text{ZGF}}[(n_x - n_y)^2]$$

are monotone decreasing in the coupling matrix A , and in the overall coupling constant λ .

APPENDIX B. PERSPECTIVES ON THE VILLAIN MODEL

B.1. An associated vector field

A convenient, and natural, choice of the Villain model's basic variables is the collection of edge variables

$$h_{uv} := \theta_u - \theta_v + 2\pi n_{uv}, \quad (\text{B.1})$$

associated with the edges $(u, v) \in \mathcal{E}$, plus a single additional angle variable θ_{x_0} of an arbitrarily preselected site $x_0 \in \mathcal{V}$.

Let $\mu_\beta^{\text{Vill}, \mathcal{G}}(dh)$ denote the (un-normalized) measure induced, as ‘‘push forward’’, on $\mathbf{h} = \{h_{uv}\}_{(u,v) \in \mathcal{E}}$ from the Villain measure

$$\prod_{\{u,v\} \in \mathcal{E}} \left[\sum_{n_{uv} \in \mathbb{Z}} e^{-\frac{\beta}{2}(\theta_u - \theta_v + 2\pi n_{uv})^2} \right] \prod_{x \in \mathcal{V}} d\theta_x. \quad (\text{B.2})$$

In this notation, the spin-spin correlation function of the Villain model takes the form

$$\langle \sigma_v \cdot \sigma_u \rangle_\beta^{\text{Vill}, \mathcal{G}} \equiv \langle e^{i(\theta_u - \theta_v)} \rangle_\beta^{\text{Vill}, \mathcal{G}} = \frac{1}{Z_\beta^{\text{Vill}, \mathcal{G}}} \int e^{i \sum_{b \in \gamma_{uv}^*} h(b)} d\mu_\beta^{\text{Vill}, \mathcal{G}}(\mathbf{h}). \quad (\text{B.3})$$

Here γ_{uv} is any path in \mathcal{V} from u to v .

For a slightly different perspective, it is natural to regard \mathbf{h} as a vector field. Inspired by that, the curl of any given realization \mathbf{h} is defined as

$$\text{curl } \mathbf{h}(q) = \sum_{b \in \partial q} h_b \quad (\text{B.4})$$

where q is a plaquette of \mathcal{G} (i.e. a point in $\mathcal{V}^* \equiv \mathcal{F}$) and ∂q is the (clockwise-oriented) edge boundary of the plaquette.

An alternative presentation of the induced measure which appears in (4.1) is

$$d\mu_\beta^{\text{Vill},\mathcal{G}}(\mathbf{h}) = \left(\prod_{q \in \mathcal{F}} \delta \left(e^{2\pi i \text{curl} \mathbf{h}(q)} - 1 \right) \right) \exp \left[-\frac{\beta}{2} (2\pi)^2 \sum_{b \in \mathcal{E}} h_b^2 \right] \prod_{b \in \mathcal{E}} dh_b \quad (\text{B.5})$$

While the δ function is a restriction to a set of zero measure, the integrals of functions with respect to it have a well known and standard interpretation. In the present case, the δ function's calculation is facilitated by the Helmholtz-Hodge decomposition of \mathbf{h} with prescribed values of $\text{curl} \mathbf{h} = \alpha_q$ into $|\mathcal{F}| + 1$ linearly independent parts:

$$\mathbf{h}_b = \left[\sum_{q \in \mathcal{F}} n_q \tilde{h}_b^q \right] + [\nabla^* \Psi_{b^*}]^* \quad (\text{B.6})$$

where \tilde{h}^q are the $\|\mathbf{h}\|_2$ minimizing solutions of

$$\text{curl} \tilde{h}_{b'}^q = \delta_{b',b} \quad (\text{B.7})$$

and Ψ is an arbitrary function defined over the dual lattice \mathcal{V}^* (whose points correspond to the faces of \mathcal{G}).

The bracketed sum in (B.6) is fully determined by $\text{curl} \mathbf{h} = \mathbf{n}$. The Ψ variables vary independently from $\{n_q\}$, and as functions in $\ell^2(\mathcal{E})$ the two parts on the right in (B.6) are orthogonal.

To answer a natural question it may be added that the total mass of $\mu_\beta^{\text{Vill},\mathcal{G}}$ on configurations with prescribed $\text{curl} \mathbf{h}$ is

$$\mu_\beta^{\text{Vill},\mathcal{G}}([\forall q \in \mathcal{F} : \text{curl} \mathbf{h}(q) = \alpha(q)]) = C'(\beta) \exp \left\{ -\frac{1}{2\beta} (2\pi)^2 \langle \alpha, (-\Delta)^{-1} \alpha \rangle \right\} \quad (\text{B.8})$$

(a fact which is not used in our discussion). This observation is related to the model's Coulomb gas representation which underlines the analysis of [15, 10].

B.2. The Villain model's duality relation with the ZGF

For a dual representation of the Villain model's Gibbs state, let us recall the distributional relation (in the dual space to Schwartz test functions)

$$\sum_{k \in \mathbb{Z}} \delta(q - 2\pi k) = \frac{1}{2\pi} \sum_{n \in \mathbb{Z}} e^{inq}. \quad (\text{B.9})$$

Inserting this relation for each plaquette q in (B.5) - (B.3), and then integrating over \mathbf{h} one gets

$$\begin{aligned} (2\pi)^{|\mathcal{V}^*|} Z_\beta^{\text{Vill}, \mathcal{G}} &= \int \exp \left\{ -\frac{\beta}{2} \sum_b h(b)^2 \right\} \prod_P \left[\sum_{n(P) \in \mathbb{Z}} e^{i \text{curl}[\mathbf{h}](q) \cdot \mathbf{n}(q)} \right] \prod_{b \in \mathcal{E}} dh(b) \\ &= \sum_{n: \mathcal{V}^* \rightarrow \mathbb{Z}} \int \exp \left\{ -\frac{\beta}{2} \sum_b [h(b)^2 + i h(b) \nabla_{b^*}^* [n]] \right\} \prod_{b' \in \mathcal{E}} dh(b') \end{aligned} \quad (\text{B.10})$$

where b^* , for an edge $b \in \mathcal{E}$, represents the dual edge, and $\nabla_{b^*}^* [n]$ is, in plain language, the difference between the values of $n(q)$ between the two plaquettes which are separated by b . The Gaussian integration over \mathbf{h} yields

$$\begin{aligned} (2\pi)^{|\mathcal{V}^*|} Z_\beta^{\text{Vill}, \mathcal{G}} &= \sum_{n: \mathcal{V}^* \rightarrow \mathbb{Z}} \exp \left\{ -\frac{\lambda}{2} \sum_{\{q, q'\} \in \mathcal{E}^*} (n_q - n_{q'})^2 \right\} \\ &\equiv \sum_{n: \mathcal{V}^* \rightarrow \mathbb{Z}} \exp \left\{ -\frac{1}{2\beta} \sum_{b \in \mathcal{E}} |\nabla_{b^*}^* [n]|^2 \right\} =: Z_{1/\beta}^{\mathbb{Z}\text{GF}, \mathcal{G}^*} \end{aligned} \quad (\text{B.11})$$

with the free boundary conditions of the Villain model corresponding to $n = 0$ along $\partial \mathcal{V}^*$. The superscripts in $Z_\beta^{\mathbb{Z}\text{GF}, \mathcal{G}^*}$ are there to convey that this is the partition function of the integer-valued Gaussian field formulated over the dual graph, whose vertices are the original graph's faces.

Under the joint probability distribution with the probability of each configuration $\{n(q)\}_{q \in \mathcal{E}}$ proportional to the weights which are summed over in (B.11) forms the “ \mathbb{Z} -valued Gaussian field” to which we refer as $\mathbb{Z}\text{GF}$.

Acknowledgements: Work on this project was supported in parts by the following grants: M.H. was partially supported by the European Research Council starting grant 678520 (LocalOrder). The research of R.P. is supported by the Israel Science Foundation grant 1971/19 and by the European Research Council starting grant 678520 (LocalOrder). J.S. was supported by the Swiss National Science Foundation (grant number P2EZIP2_184228), and the Princeton-Geneva Univ. collaborative travel funds. M.A. gratefully acknowledges the support of the Weston Visiting Professorship at The Weizmann Institute of Science (Rehovot, Israel), and the warm hospitality of Drs. Enrique and Maria Rodriguez Boulan in Quogue NY.

REFERENCES

- [1] M. Aizenman and B. Simon. A comparison of plane rotor and Ising models. *Phys. Lett. A*, 76(3):281–282, 1980.

- [2] Michael Aizenman, Matan Harel, and Ron Peled. Exponential Decay of Correlations in the 2D Random Field Ising Model. *J. Stat. Phys.*, 180(1):304–331, Sep 2020.
- [3] Michael Aizenman and Bruno Nachtergaele. Geometric aspects of quantum spin states. *Comm. Math. Phys.*, 164(1):17 – 63, 1994.
- [4] V. L. Berezinsky. Destruction of long range order in one-dimensional and two-dimensional systems having a continuous symmetry group. I. Classical systems. *Sov. Phys. JETP*, 32:493–500, 1971.
- [5] R. M. Burton and M. Keane. Density and uniqueness in percolation. *Comm. Math. Phys.*, 121(3):501 – 505, 1989.
- [6] Omri Cohen-Alloro and Ron Peled. Rarity of extremal edges in random surfaces and other theoretical applications of cluster algorithms. *The Annals of Applied Probability*, 30(5):2439 – 2464, 2020.
- [7] Paul Dario and Wei Wu. Massless phases for the Villain model in $d \geq 3$, 2020.
- [8] P. L. Dabrushin. The Description of a Random Field by Means of Conditional Probabilities and Conditions of its Regularity. *Theory of Probab. & Its Applic.*, 13(2):197–224, 1968.
- [9] R. L. Dabrushin and S. B. Shlosman. Absence of breakdown of continuous symmetry in two-dimensional models of statistical physics. *Comm. Math. Phys.*, 42(1):31 – 40, 1975.
- [10] Jürg Fröhlich and Thomas Spencer. The Kosterlitz-Thouless transition in two-dimensional abelian spin systems and the Coulomb gas. *Comm. Math. Phys.*, 81(4):527 – 602, 1981.
- [11] J. Fröhlich, B. Simon, and T. Spencer. Infrared bounds, phase transitions and continuous symmetry breaking. *Comm. Math. Phys.*, 50(1):79 – 95, 1976.
- [12] Jürg Fröhlich and Yong Moon Park. Correlation inequalities and the thermodynamic limit for classical and quantum continuous systems. *Comm. Math. Phys.*, 59(3):235–266, 1978.
- [13] Christophe Garban and Avelio Sepúlveda. Quantitative bounds on vortex fluctuations in $2d$ Coulomb gas and maximum of the integer-valued Gaussian free field.
- [14] J. Ginibre. General formulation of Griffiths’ inequalities. *Comm. Math. Phys.*, 16(4):310 – 328, 1970.
- [15] Jorge V. José, Leo P. Kadanoff, Scott Kirkpatrick, and David R. Nelson. Renormalization, vortices, and symmetry-breaking perturbations in the two-dimensional planar model. *Phys. Rev. B*, 16:1217–1241, Aug 1977.
- [16] Vital Kharash and Ron Peled. The Fröhlich-Spencer Proof of the Berezinskii-Kosterlitz-Thouless Transition. *arXiv preprint arXiv:1711.04720*, 2017.
- [17] J M Kosterlitz and D J Thouless. Ordering, metastability and phase transitions in two-dimensional systems. *Journal of Physics C: Solid State Physics*, 6(7):1181–1203, apr 1973.
- [18] Piet Lammers. Height function delocalisation on cubic planar graphs. *Probability Theory and Related Fields*, Sep 2021.
- [19] Piet Lammers and Sébastien Ott. Delocalisation and absolute-value-FKG in the solid-on-solid model, 2021.
- [20] Oliver A. McBryan and Thomas Spencer. On the decay of correlations in $SO(n)$ -symmetric ferromagnets. *Comm. Math. Phys.*, 53(3):299 – 302, 1977.
- [21] A. Patrascioiu and E. Seiler. Phase structure of two-dimensional spin models and percolation. *J. Stat. Phys.*, 69(3):573–595, Nov 1992.
- [22] Charles Edouard Pfister. On the symmetry of the Gibbs states in two-dimensional lattice systems. *Comm. Math. Phys.*, 79(2):181 – 188, 1981.
- [23] A.M. Polyakov. Interaction of goldstone particles in two dimensions. Applications to ferromagnets and massive Yang-Mills fields. *Physics Letters B*, 59(1):79 – 81, 1975.
- [24] Oded Regev and Noah Stephens-Davidowitz. An Inequality for Gaussians on Lattices. *SIAM Journal on Discrete Mathematics*, 31(2):749–757, 2017.
- [25] Scott Sheffield. *Random surfaces*. Number 304 in Astérisque. Société mathématique de France, 2005.
- [26] B. Simon. A remark on Dabrushin’s uniqueness theorem. *Comm. Math. Phys.*, 68:183–185, 1979.
- [27] J. Villain. Theory of one- and two-dimensional magnets with an easy magnetization plane. II. the planar, classical, two-dimensional magnet. *Journal de Physique*, 36(6):581–590, 1975.

MA: DEPARTMENTS OF PHYSICS AND MATHEMATICS, PRINCETON UNIVERSITY, USA
WESTON VISITING PROFESSOR AT THE WEIZMANN INSTITUTE OF SCIENCE, ISRAEL

MH: DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIV., BOSTON, USA

RP: SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, ISRAEL

JS: PHYSICS DEPARTMENT, PRINCETON UNIVERSITY, PRINCETON, NJ 08544, USA