

Eternal Domination and Clique Covering

Gary MacGillivray ^{*†} C. M. Mynhardt ^{*†} Virg lot Virgile [†]

October 20, 2021

Abstract

We study the relationship between the eternal domination number of a graph and its clique covering number. Using computational methods, we show that the smallest graph having its eternal domination number less than its clique covering number has 10 vertices. This answers a question of Klostermeyer and Mynhardt [Protecting a graph with mobile guards, *Appl. Anal. Discrete Math.* 10 (2016), no. 1, 1 – 29]. We also determine the complete set of 10-vertex and 11-vertex graphs having eternal domination numbers less than their clique covering numbers. In addition, we study the problem on triangle-free graphs, circulant graphs, planar graphs and cubic graphs. Our computations show that all triangle-free graphs and all circulant graphs of order 12 or less have eternal domination numbers equal to their clique covering numbers, and exhibit 13 triangle-free graphs and 2 circulant graphs of order 13 which do not have this property. Using these graphs, we describe a method to generate an infinite family of triangle-free graphs and an infinite family of circulant graphs with eternal domination numbers less than their clique covering numbers. Our computations also show that all planar graphs of order 11 or less, all 3-connected planar graphs of order 13 or less and all cubic graphs of order less than 18 have eternal domination numbers equal to their clique covering numbers. Finally, we show that for any integer $k \geq 2$ there exist infinitely many graphs having domination number and eternal domination number equal to k containing dominating sets which are not eternal dominating sets. This answers another question of Klostermeyer and Mynhardt [Eternal and Secure Domination in Graphs, *Topics in domination in graphs*, *Dev. Math.* 64 (2020), 445–478, Springer, Cham].

Keywords: Dominating sets, eternal dominating sets, independent sets, clique covering, combinatorial game, graph protection.

AMS MSC 2020: 05C69

1 Introduction

The eternal domination game played on graphs is a combinatorial game that was introduced by Burger, Cockayne, Grundlingh, Mynhardt, Van Vuuren and Winterbach [5]. The game is played between two players that alternate turns: a defender who controls a set of guards, and an attacker. To start the game, the defender chooses a subset of vertices D_0 on which to place the guards (at most one guard on each vertex). At each time $t = 1, 2, 3, \dots$ the attacker selects a vertex v on which there is no guard; we say the attacker *attacks* v . The defender responds by moving a guard on a neighbour of v to v ; we say the defender *defends* v . The guards (or the defenders) win if they are able to respond to the sequence of attacks; otherwise the attacker wins, that is, if at some

*Funded by a Discovery Grant from the Natural Sciences and Engineering Research Council of Canada.

[†]Department of Mathematics and Statistics, University of Victoria, Victoria BC, Canada.

E-mail addresses: gmacgill@uvic.ca, kiek@uvic.ca, virgilev@uvic.ca

time t there is no guard in the neighbourhood of the vertex selected by the attacker. The eternal domination number of a graph G , denoted by $\gamma^\infty(G)$, is the minimum number of guards necessary to respond to any sequence of attacks on G . For a survey on the eternal domination game and its variants, see [17] or [18].

For graphs belonging to certain families, the eternal domination number is closely related to another well-studied parameter which is the clique covering number. The *clique covering number* of a graph G , denoted by $\theta(G)$, is the minimum cardinality of a clique covering of G , i.e., a partition $\{V_1, V_2, \dots, V_k\}$ of $V(G)$ such that each V_i induces a clique. Observe that, as stated in [5], if we consider a minimum clique covering of G , place a guard in each clique of the covering and play the game independently in each clique, each guard is able to respond to any sequence of attacks on its respective clique. This strategy shows that the clique covering number of a graph is an upper bound on its eternal domination number. In fact, $\gamma^\infty(G) = \theta(G)$ if G is a perfect graph [5], an outerplanar graph [2], or a graph with $\theta(G) \leq 3$ [5]; a complete characterization of the graphs G with $\gamma^\infty(G) = \theta(G)$ is yet to be found. Goddard, Hedetniemi and Hedetniemi [10] showed that there exist graphs with $\gamma^\infty < \theta$; they gave an 11-vertex graph as example. Klostermeyer and Mynhardt [17] asked whether that 11-vertex graph is the smallest graph, in terms of the number of vertices, with this property. They further asked whether there exist graphs with $\gamma^\infty < \theta$ in some particular graph classes, for example planar graphs.

This paper is organised as follows. In Section 2, we introduce a few other graph parameters that are related to the eternal domination number of a graph. We also define a new graph product which we call the bow tie product and finally present some graph classes on which we study the problem. In Section 3, we introduce some preliminary results on the topic. Our main results lie in Section 4. To begin with, we show in Section 4.1 that the smallest graph with eternal domination number less than its clique covering number has 10 vertices. We also determine the complete set of 10-vertex and 11-vertex graphs having eternal domination numbers less than their clique covering numbers. In Section 4.2, we restrict our attention to certain graph classes, namely triangle-free graphs, circulant graphs, planar graphs and cubic graphs. In particular, we show that the smallest triangle-free graph with eternal domination number less than its clique covering number has 13 vertices. We also show that the smallest circulant graph with eternal domination number less than its clique covering number has 13 vertices. Our computations show that all planar graphs on fewer than 12 vertices, all 3-connected planar graphs on fewer than 14 vertices and all cubic graphs on fewer than 18 vertices have eternal domination numbers equal to their clique covering numbers. Finally, in Section 4.3, we consider a question and a conjecture of Klostermeyer and Mynhardt. We show that the answer to the question is no and verify the conjecture for all graphs up to order 11.

2 Definitions

The *domination number* of a graph G , denoted by $\gamma(G)$, is the minimum cardinality of a dominating set of G , that is, a set $D \subseteq V(G)$ such that any vertex $v \in V(G) \setminus D$ has a neighbour in D . This means that, in the eternal domination game, the guards must always be located on the vertices of a dominating set of G in order to defend a sequence of attack on G . A dominating set from which the guards can defend any sequence of attack is known as an *eternal dominating set*.

The *independence number* of G , denoted by $\alpha(G)$, is the maximum cardinality of an independent set of G , that is, a set $S \subseteq V(G)$ such that for any $u, v \in S$, $uv \notin E(G)$. The *clique number* of G , denoted by $\omega(G)$, is the maximum cardinality of a clique of G , where a clique is the complement of an independent set. This implies that $\alpha(G) = \omega(\overline{G})$ for any graph G and its complement \overline{G} .

The *chromatic number* of G , denoted by $\chi(G)$, is the minimum number of colours required to

colour the vertices of G so that no two adjacent vertices have the same colour. Observe that such a colouring with k colours is obtained by partitioning $V(G)$ into $\{V_1, V_2, \dots, V_k\}$ where each V_i induces an independent set. For this reason, the clique covering number of a graph is equal to the chromatic number of its complement. When the graph G is clear from context, we use $n, \alpha, \gamma^\infty, \theta$ to denote respectively $|V(G)|, \alpha(G), \gamma^\infty(G), \theta(G)$.

A graph G is said to be *vertex critical* with respect to θ if $\theta(G - \{v\}) = \theta(G) - 1$ for any vertex $v \in V(G)$. A graph G is said to be *edge critical* with respect to θ if $\theta(G + \{uv\}) = \theta(G) - 1$ for any edge $uv \notin E(G)$, where $G + \{uv\}$ is the graph obtained from G by adding the missing edge uv . Since we only consider criticality with respect to θ , we simply refer to G being vertex or edge critical without explicitly referring to θ . We say that a graph G is *critical* if G is vertex critical and edge critical.

The *circulant graph* $C_n[k_1, k_2, \dots, k_l]$, where $\{k_1, k_2, \dots, k_l\} \subseteq \mathbb{Z}^+$ and $1 \leq k_1 < k_2 < \dots < k_l \leq \lfloor \frac{n}{2} \rfloor$, is the graph with vertex set $\{v_0, v_1, v_2, \dots, v_{n-1}\}$ such that two vertices v_i and v_j are adjacent if and only if $i - j \equiv \pm k_p \pmod{n}$ for some $p \in \{1, 2, \dots, l\}$. A *perfect graph* is a graph such that the chromatic number of any of its induced subgraphs is equal to the clique number of that subgraph.

The *bow tie product* of a graph G with a graph H , denoted by $G \bowtie H$, is the graph with vertex set $\{(v_i, v_j) : v_i \in V(G), v_j \in V(H)\}$, where two vertices (v_i, v_j) and (v'_i, v'_j) are adjacent if and only if one of the following conditions holds: $v_i v'_i \in E(G)$ and $v_j = v'_j$, or $v_i v'_i \in E(G)$ and $v_j v'_j \in E(H)$ (see Figure 1). As shown in Figure 1, $G \bowtie H \not\cong H \bowtie G$ in general.




Figure 1: Bow tie product of two graphs.

We refer to a graph G as being a *smallest graph having some property \mathcal{P}* if no graph H on fewer than $|V(G)|$ vertices has property \mathcal{P} .

3 Preliminary results

It is straightforward to see that the eternal domination number of a graph is the sum of the eternal domination number of its components; therefore, we restrict our attention to connected graphs.

We begin with the following observation that we may obtain upper and lower bounds on $\gamma^\infty(G)$ by considering the problem on the induced and spanning subgraphs of G .

Fact 1. *Let G be a graph and let H be a subgraph of G .*

- * [15] *If H is an induced subgraph of G , then $\gamma^\infty(G) \geq \gamma^\infty(H)$.*
- * [2] *If H is a spanning subgraph of G , then $\gamma^\infty(G) \leq \gamma^\infty(H)$.*

The reader can easily check that $\gamma^\infty(K_n) = 1$ and $\gamma^\infty(\overline{K_n}) = n$. Hence, by considering a minimum clique covering of G and playing the game independently on each clique we obtain an upper bound on $\gamma^\infty(G)$. Likewise, if we consider a maximum independent set S of G and play the

game on the subgraph induced by S , we obtain a lower bound on $\gamma^\infty(G)$. In consequence, we have the following fact of Burger *et al.*

Fact 2 ([5]). *For any graph G , $\alpha(G) \leq \gamma^\infty(G) \leq \theta(G)$.*

It is well known [19, 20] that the complement of a perfect graph is perfect. As a direct consequence we have the following corollary.

Corollary 3 ([5]). *For any perfect graph G , $\alpha(G) = \gamma^\infty(G) = \theta(G)$.*

The inequalities in Fact 2 can both be strict. Observe that the 5-cycle is a graph with $\alpha = 2 < \gamma^\infty = 3$ and is the smallest such graph. As for the second inequality, the first proof of the existence of a graph with $\gamma^\infty < \theta$ is due to Goddard, Hedetniemi and Hedetniemi [10] and follows from Theorem 4 and its generalisation, Theorem 5, which shows that the eternal domination number of a graph is bounded above by a function of the independence number of the graph.

Theorem 4 ([10]). *If G is a graph such that $\alpha(G) = 2$, then $\gamma^\infty(G) \leq 3$.*

Theorem 5 ([14]). *For any graph G , $\gamma^\infty(G) \leq \binom{\alpha(G)+1}{2}$.*

It is known that for any integer $k \geq 2$ there exists a triangle-free graph G with chromatic number k . The first known construction of a family containing such graphs with arbitrarily large chromatic numbers is due to Blanche Descartes [7, 8]. Mycielski [22] described the construction of a family $\mathcal{F} = \{M_2, M_3, M_4, \dots\}$ of triangle-free graphs starting with $M_2 = K_2$, where, for each $k \geq 3$, the graph M_k is obtained from the graph M_{k-1} and has chromatic number k (see M_4 in Figure 2a).



Figure 2: The Grötzsch graph (a) and its complement (b).

Corollary 6 ([10]). *For any integer $k \geq 4$, $\gamma^\infty(\overline{M_k}) < \theta(\overline{M_k})$.*

Our primary purpose in this paper is to study the relationship between the eternal domination number and the clique covering number for some specific graphs; however, Theorem 5 implies a nice result on the problem regarding general random graphs that is worth mentioning.

Theorem 7 ([1], Theorem 3.1). *For almost all graphs G , $\alpha(G) \leq (2 + o(1)) \log_2 n$ and $\theta(G) \geq (1 + o(1)) \frac{n}{2 \log_2 n}$.*

Corollary 8. *For almost all graphs G , $\gamma^\infty(G) < \theta(G)$.*

Proof. We know from Theorem 7 that $\alpha(G) \leq (2 + o(1)) \log_2 n$ for almost all graphs G . Together with Theorem 5, this implies that $\gamma^\infty(G) \leq (2 + o(1)) (\log_2 n)^2$ for almost all graphs G . Since $\lim_{n \rightarrow \infty} \frac{2(\log_2 n)^2}{n/(2 \log_2 n)} = 0$, we conclude that $\gamma^\infty(G) < \theta(G)$ for almost all graphs G . \square

Theorem 9 ([6]). *The Grötzsch graph (Figure 2a) is the unique smallest triangle-free graph with chromatic number at least 4.*

Corollary 10. *The complement of the Grötzsch graph is the unique smallest graph with independence number 2 and clique covering number at least 4.*

Klostermeyer and Mynhardt [17, 18] posed the following questions.

Question 11 ([17]). *Is the complement of the Grötzsch graph the smallest graph with eternal domination number less than its clique covering number?*

Question 12 ([18]). *Let G be a graph with $\gamma(G) = \gamma^\infty(G)$. Is any minimum dominating set of G an eternal dominating set of G ?*

We answer Question 11 in Section 4.1 by finding two graphs of order 10 with this property. We also show that the answer to Question 12 is no by constructing an infinite family of counterexamples.

3.1 Algorithms and complexity

Consider the following decision problem.

ETERNAL DOMINATION
INSTANCE: A graph G and an integer $k > 0$.
QUESTION: Is $\gamma^\infty(G) \leq k$?

Although the precise complexity of ETERNAL DOMINATION is not known for general graphs, Klostermeyer and MacGillivray [13] showed that the problem can be solved in time exponential in n . Despite the fact that their algorithm is described for a variant of the eternal domination game, Algorithm 1 is adapted to the version of the game considered in this paper. Klostermeyer also studied the complexity of the problem under narrow assumptions on the sequence of attacks [12].

The precise complexity of ETERNAL DOMINATION is known for graphs belonging to certain graph classes, for example perfect graphs. We know from Corollary 3 that the eternal domination number of a perfect graph is equal to its independence number and hence its clique covering number. It is well known [11] that the independence number of a perfect graph can be found in polynomial time in n . As a result, ETERNAL DOMINATION is in P for perfect graphs.

Algorithm 1 ([13]). Determine whether $\gamma^\infty(G) \leq k$.

Given G , construct an arc-coloured digraph D as follows.

1. The vertex set of D is the set of k -vertex dominating sets of G . The set of colours is $V(G)$. There is an arc from X to Y of colour v when $Y - X = \{v\}$, and v is adjacent in G to the unique vertex $w \in X - Y$.
2. Delete any vertex X which is not the origin of an arc coloured x for some $x \in V(G) \setminus X$.
3. Repeat Step 2 until no further vertices can be deleted.

If D is the null graph, then $\gamma^\infty(G) > k$; otherwise, $\gamma^\infty(G) \leq k$.

4 Main results

4.1 General graphs

In this section, we show that the graphs G_1 and G_2 depicted in Figure 3 are the smallest graphs having their eternal domination numbers less than their clique covering numbers. Observe that G_2 is obtained from G_1 by adding the edge $(6,7)$.



Figure 3: Smallest graphs with $\gamma^\infty < \theta$.



Figure 4: Complements of the graphs in Figure 3.

We first show that these graphs have eternal domination numbers less than their clique covering numbers. To this end, we begin with the following facts; the proofs are easy and left to the reader.

Fact 13. For the graphs $G_1, G_2, \overline{G_1}$ and $\overline{G_2}$ depicted in Figures 3 and 4,

- * $\alpha(G_1) = \alpha(G_2) = \omega(\overline{G_1}) = \omega(\overline{G_2}) = 3$.
- * $\theta(G_1) = \theta(G_2) = \chi(\overline{G_1}) = \chi(\overline{G_2}) = 4$.

Fact 14. The graph $\overline{G_1}$ contains exactly six triangles, namely $\{2, 3, 4\}$, $\{5, 6, 7\}$, $\{0, 2, 8\}$, $\{0, 1, 7\}$, $\{1, 3, 9\}$ and $\{6, 8, 9\}$, any two of which share at most one vertex.

Fact 15. The subgraph induced by the vertices in the open neighbourhood of each vertex of $\overline{G_1}$ is isomorphic to either $2K_2$ or to $K_1 \cup K_2$.

The following result can be verified by computer, but we include a proof as well.

Proposition 16. For the graphs G_1 and G_2 depicted in Figure 3, $\gamma^\infty(G_1) = \gamma^\infty(G_2) < \theta(G_1) = \theta(G_2)$.

Proof. Since G_1 is a spanning subgraph of G_2 , it suffices to show that $\gamma^\infty(G_1) = 3$. Fact 1 will then imply that $\gamma^\infty(G_2) = 3$. We do this by contradiction and assume that $\gamma^\infty(G_1) \geq 4$. We may also assume without loss of generality that three guards are initially located on the vertices of an independent set of G_1 . This is because the attacker may sequentially attack all the vertices of a maximum independent set of G_1 and force a guard to be located on each of them. Let k be the smallest integer such that there exists a sequence of attacks of length k that the three guards cannot defend. Suppose the guards respond optimally to that sequence of attacks. At time $t = k - 1$, they will be located on the vertices b, c and d , none of which is adjacent to vertex a (see Figure 5 where a thick black edge corresponds to an edge in G_1 and a thin blue edge corresponds to an edge in $\overline{G_1}$).




Figure 5: Configuration of the guards at time $t = k - 1$.

Following Fact 15, we may assume that vertex c is not adjacent to vertex d and vertex b is adjacent to both of c and d . So, we colour the edge cd blue and the edges bc and bd black (see Figure 6).




Figure 6: Configuration of the guards at time $t = k - 1$.

Note that in the previous time ($t = k - 2$), the guards dominated all the vertices of G , in particular vertex a was dominated. We now consider two cases depending on the location of the guards at time $t = k - 2$.

- Case 1: In the previous turn, a guard moved from a vertex e to vertex c (see Figure 7b). In this case, e is adjacent to a because a was dominated at time $t = k - 2$ but is undominated at time $t = k - 1$, and b has a neighbour f which is non-adjacent to all of c, d and e (otherwise the guards would still be on a dominating set if the guard on b moved to c instead; see Figure 7c). This contradicts Fact 14 since the thin blue triangles $\{a, c, d\}$ and $\{f, c, d\}$ share two vertices (c and d ; see Figure 7c).




Figure 7: Configuration of the guards in Case 1 of the proof of Proposition 16.

- Case 2: In the previous turn, a guard moved from vertex e to vertex b (see Figure 8b). In this case, c has a neighbour f which is non-adjacent to b, d and e (otherwise the guards would still be on a dominating set if the guard on c moved to b instead). For the same reason, d

has a neighbour g which is not adjacent to b, c and e , as shown in Figure 8c. Following Fact 14, f must be adjacent to g , otherwise $\{f, b, g\}$ and $\{f, e, g\}$ would be two thin blue triangles sharing two vertices. By the same fact, a must be adjacent to f as $\{a, c, d\}$ and $\{a, f, d\}$ would share two vertices otherwise. For a similar reason, a must be adjacent to g as $\{a, c, g\}$ and $\{a, c, d\}$ would be two thin blue triangles sharing two vertices otherwise (see Figure 8d). Now, the edges ce and de cannot be both thin blue, otherwise the thin blue triangles $\{a, c, d\}$ and $\{e, c, d\}$ would share two vertices. We assume without loss of generality that the edge ce is black and we consider the two choices for the colour of the edge de .




Figure 8: Configuration of the guards in Case 2 of the proof of Proposition 16.

So, we obtain two blue/black colourings of the edges of the complete graph on 7 vertices. One of the thin blue subgraphs obtained (Figure 9) must be an induced subgraph of $\overline{G_1}$.




Figure 9: Induced subgraphs of $\overline{G_1}$.

The reader can check that $\overline{G_1}$ does not contain any of those graphs as an induced subgraph. Hence, we obtain a contradiction and conclude that $\gamma^\infty(G_1) = 3$. \square

In the remaining part of this section, we verify that any graph with fewer than 10 vertices has eternal domination number equal to its clique covering number.

Proposition 17. *For any graph G of order 9 or less, $\gamma^\infty(G) = \theta(G)$.*

Proposition 17 was verified by computer; the search can be described as follows. Suppose G is a smallest graph such that $\gamma^\infty(G) < \theta(G)$ with the maximum number of edges. Fact 2 implies that $\alpha(G) < \theta(G)$; moreover, by the choice of G and Fact 1, deleting a vertex from the graph or adding a missing edge to the graph decreases its clique covering number by 1. Consequently, G is a critical graph.

Table 1 shows the number of critical graphs of order 9 or less and each of these graphs is either drawn in Figures 10 or 11, or listed in Graph6 format in Table 8 in the appendix. To find these graphs, we used NAUTY [21] to generate the set of connected graphs on fewer than 10 vertices. Then, using some basic graph algorithms, we computed the independence number and the clique covering number of each of these graphs. All the computations were done in Python. We

found the independence number of each graph by computing the clique number of the complement graph using the built-in functions `complement()` and `graph_clique_number()` from the NetworkX package. We found the clique covering number of each graph by first finding the set of maximal cliques of the graph using the built-in function `find_cliques()` from the NetworkX package, then, using two different approaches (a naive method and an integer programming method), we found the size of the smallest subset of the set of maximal cliques which covers all the vertices of the graph. Our naive method is on average faster than the integer programming method on the set of graphs of order 10 or less. For those graphs with independence numbers less than their clique covering numbers, we checked whether they are critical graphs before computing their eternal domination numbers using Algorithm 1.

Using a similar technique, without considering only critical graphs, we found the complete set of 10-vertex and 11-vertex graphs with $\gamma^\infty < \theta$. Those graphs are listed in Table 9 in the appendix in Graph6 format.

The graph C_5 is the only critical graph of order 5 with $\alpha < \theta$. We display the critical graphs of order 7 and 8 in figures 10 and 11, and list the critical graphs of order 9 with $\alpha < \theta$ in Graph6 format in the appendix (Table 8).

Table 1: Number of critical graphs on n vertices with $\gamma^\infty < \theta$.

n	Total	$\alpha < \theta$	Vertex-Critical & $\alpha < \theta$	Critical & $\alpha < \theta$	Critical & $\gamma^\infty < \theta$
5	21	1	1	1	0
6	112	3	0	0	0
7	853	33	8	3	0
8	11117	498	7	4	0
9	261080	16539	353	38	0
10	11716571	975676	5159	290	1




Figure 10: Critical graphs of order 7 with $\alpha < \theta$.




Figure 11: Critical graphs of order 8 with $\alpha < \theta$.

4.2 Graph Classes

In this section, we focus on different classes of graphs, starting with triangle-free graphs in Sections 4.2.1 and 4.2.2, before moving on to circulant graphs in Section 4.2.3 and planar graphs in Section 4.2.4. We briefly mention claw-free graphs in Section 4.2.5 and cubic graphs in Section 4.2.6.

4.2.1 Triangle-free graphs

Erdős, Kleitman and Rothschild [9] showed that almost all triangle-free graphs are bipartite, therefore almost all triangle-free graphs are perfect and satisfy $\gamma^\infty = \theta$. However, Goddard, Hedetniemi and Hedetniemi [10] showed that there exist triangle-free graphs with $\gamma^\infty < \theta$. They gave the circulant graphs $C_{18}[1, 3, 8]$ and $C_{21}[1, 3, 8]$, which they found using computer assistance, as examples. Using Algorithm 1, we checked that any triangle-free graph of order 12 or less has eternal domination number equal to its clique covering number. To compute the clique covering number of a triangle-free graph, we first find the size of a maximum matching in the graph using the NetworkX's function `max_weight_matching()` and then we subtract this number from the order of the graph. We found 13 triangle-free graphs of order 13 with $\gamma^\infty < \theta$ (see Table 10). Previously the smallest known triangle-free graph with this property has 17 vertices and is a subgraph of the circulant graph $C_{18}[1, 3, 8]$. The following corollary describes a way to generate an infinite family of triangle-free graphs with $\gamma^\infty < \theta$.

Fact 18. *Let G be a triangle-free graph. Then $G \boxtimes K_2$ is triangle-free.*

Corollary 19. *Let G be a triangle-free graph such that $\gamma^\infty(G) < \theta(G) = \lceil \frac{n}{2} \rceil$. Then, $\gamma^\infty(G \boxtimes K_2) < \theta(G \boxtimes K_2)$.*

Proof. Let G be a triangle-free graph such that $\gamma^\infty(G) < \theta(G) = \lceil \frac{n}{2} \rceil$. Fact 18 implies that $G \boxtimes K_2$ is triangle-free; as a result, $\theta(G \boxtimes K_2) \geq n$. Observe that the vertices of $G \boxtimes K_2$ can be partitioned into two subsets, each of which induces a subgraph isomorphic to G . This means that $\gamma^\infty(G \boxtimes K_2) \leq 2\gamma^\infty(G) < 2\lceil \frac{n}{2} \rceil$; consequently, $\gamma^\infty(G \boxtimes K_2) \leq n - 1$. \square

Klostermeyer and Mynhardt [16] posed the following question.

Question 20 ([16]). *Does there exist a triangle-free graph G such that $\alpha(G) = \gamma^\infty(G) < \theta(G)$?*

Note that the 13 triangle-free graphs on 13 vertices with $\gamma^\infty < \theta$ we found also satisfy $\alpha < \gamma^\infty < \theta$. In Proposition 21, we describe a property of a smallest such a graph (if it exists), then we show by using computer assistance that no graph of order 14 or less satisfies this property.

Proposition 21. *Suppose there exist triangle-free graphs with $\alpha = \gamma^\infty < \theta$ and let G be such a graph with minimum order. Let X be a maximum independent set of G and let $Y = V(G) - X$. Then $|X| = |Y| - 1$.*

Proof. It is clear that $|X| \geq |Y| - 1$, otherwise, by Fact 1, $\alpha(G - \{v\}) = \gamma^\infty(G - \{v\}) < \theta(G - \{v\})$ for any $v \in Y$, and $G - \{v\}$ would be a smaller triangle-free graph with $\alpha = \gamma^\infty < \theta$. So, it remains to show that $|X| \leq |Y| - 1$. Suppose this is false, in other words $|X| \geq |Y|$. Let G' be the spanning bipartite subgraph of G obtained by deleting all edges having both endpoints in Y . Observe that the graph G' does not contain a matching that covers all of the vertices in Y , otherwise, G would contain a matching that matches each vertex in Y to a vertex in X , and this would imply that $\alpha(G) = \theta(G)$ (contradiction). Consequently, by Hall's matching condition, Y contains a subset S such that $|N_{G'}(S)| < |S|$. Now, consider the subgraph H of G induced by $S \cup N_{G'}(S)$. Since there is no edge between a vertex in H and a vertex in $X - V(H)$, we have $\alpha(H) = |N_{G'}(S)|$, otherwise

G contains an independent set of size at least $|X| + 1$. Moreover, since $\gamma^\infty(G) = |X|$ we must have $\gamma^\infty(H) = |N_{G'}(S)|$: this is true because the attacker may force the $|X|$ guards to be located in X and from there only attacks the vertices in H . In this case, only the $|N_{G'}(S)|$ guards are able to respond to that sequence of attacks on H . Hence, H would be a smaller triangle-free graph with $\alpha = \gamma^\infty < \theta$. \square




Figure 12: Cases considered in the proof of Proposition 21.

Using the property stated in Proposition 21, to find the smallest triangle-free graph G with $\alpha = \gamma^\infty < \theta$ we only need to check the triangle-free graphs of order $2k + 1$ with $\alpha = k$ and $\theta = k + 1$. Again, using NAUTY, we generated the set of triangle-free graphs of odd orders on fewer than 14 vertices; then, we computed the independence number and the clique covering number of each of these graphs using the algorithms described above. If $\alpha = \frac{n-1}{2}$ and $\theta = \frac{n+1}{2}$, then we computed the eternal domination number of the graph. We performed the computation listed in Table 2 on a cluster which required approximately 218 CPU days.

Observation 22. *There are no triangle-free graphs on $n \leq 14$ vertices with $\alpha = \gamma^\infty < \theta$.*

Table 2: Number of connected triangle-free graphs on n vertices with $\alpha = \lfloor \frac{n}{2} \rfloor$ and $\theta = \lceil \frac{n}{2} \rceil$.

n	Total	$\alpha = \lfloor \frac{n}{2} \rfloor$	$\alpha = \lfloor \frac{n}{2} \rfloor$ & $\theta = \lceil \frac{n}{2} \rceil$	$\alpha = \gamma^\infty = \lfloor \frac{n}{2} \rfloor$ & $\theta = \lceil \frac{n}{2} \rceil$
5	6	1	1	0
7	59	8	8	0
9	1380	276	276	0
11	90842	29660	29660	0
13	19425052	9606337	9606334	0

4.2.2 Maximal triangle-free graphs

A graph G is said to be *maximal triangle-free* if G is triangle-free and the insertion of any missing edge in G creates a triangle. The maximal triangle-free graphs are considered to be an interesting family of graphs since several problems on triangle-free graphs can be studied by restricting attention to maximal triangle-free graphs. Observe that if G is a triangle-free graph with $\lfloor \frac{n}{2} \rfloor = \alpha = \gamma^\infty < \theta = \lceil \frac{n}{2} \rceil$, then G is a subgraph of a maximal triangle-free graph with $\gamma^\infty < \theta = \lceil \frac{n}{2} \rceil$. This follows since the clique covering number of a triangle-free graph G is at least

$\lceil \frac{n}{2} \rceil$ and adding any missing edge to G that does not create a triangle increases neither its clique covering number nor its eternal domination number.

Proposition 23. *Suppose there exist maximal triangle-free graphs with $\alpha = \gamma^\infty < \theta$ and let G be such a graph with minimum order. Let X be a maximum independent set of G and $Y = V(G) - X$. Then $|X| = |Y| - 1$.*

Proof. Same proof as Proposition 21. □

Observation 24. *There are no maximal triangle-free graphs on $n \leq 16$ vertices with $\alpha = \gamma^\infty < \theta$.*

Observation 24 was verified using a similar computer search as in Observation 22.

Table 3: Number of maximal triangle-free graphs on n vertices with $\alpha = \lfloor \frac{n}{2} \rfloor$ and $\theta = \lceil \frac{n}{2} \rceil$.

n	Total	$\alpha = \lfloor \frac{n}{2} \rfloor$	$\alpha = \lfloor \frac{n}{2} \rfloor$ & $\theta = \lceil \frac{n}{2} \rceil$	$\alpha = \gamma^\infty = \lfloor \frac{n}{2} \rfloor$ & $\theta = \lceil \frac{n}{2} \rceil$
5	3	1	1	0
7	6	1	1	0
9	16	5	5	0
11	61	23	23	0
13	392	172	172	0
15	5036	1837	1837	0
17	164796	38606	38606	?

4.2.3 Circulant graphs

Circulant graphs are another interesting family of graphs on which we study the problem. Using Algorithm 1, we found all the circulant graphs of order 20 or less with $\gamma^\infty < \theta$ (see Table 4).

Table 4: List of small circulant graphs with $\gamma^\infty < \theta$.

n	List of graphs
13	$C_{13}[1, 3, 4], C_{13}[1, 2, 3, 5]$.
14	None
15	$C_{15}[1, 3, 4]$.
16	$C_{16}[1, 2, 4, 5], C_{16}[1, 2, 3, 4, 6]$.
17	$C_{17}[1, 2, 4, 8], C_{17}[1, 2, 3, 5, 6], C_{17}[1, 2, 3, 5, 8]$.
18	$C_{18}[1, 3, 8], C_{18}[1, 2, 4, 5, 6], C_{18}[1, 2, 4, 5, 6, 9]$.
19	$C_{19}[1, 4, 6], C_{19}[1, 3, 5, 6], C_{19}[1, 2, 3, 4, 5, 7], C_{19}[1, 2, 3, 5, 7, 8]$.
20	$C_{20}[1, 5, 8], C_{20}[2, 5, 6], C_{20}[1, 6, 8, 9], C_{20}[1, 2, 4, 5, 6], C_{20}[1, 2, 4, 5, 7]$ $C_{20}[1, 2, 5, 7, 8], C_{20}[1, 2, 3, 4, 5, 7, 8], C_{20}[1, 2, 3, 4, 6, 7, 10], C_{20}[1, 3, 4, 7, 8, 9, 10]$.

Proposition 25. *For any integer n , $C_n[k_1, k_2, \dots, k_l] \bowtie K_2 \cong C_{2n}[2k_1, 2k_2, \dots, 2k_l, 2k_1 + 1, 2k_2 + 1, \dots, 2k_l + 1]$.*

Proof. Let $G = C_n[k_1, k_2, \dots, k_l]$ be a circulant graph and let $H = C_n[k_1, k_2, \dots, k_l] \bowtie K_2$. Suppose $V(G) = \{v_1, v_2, \dots, v_n\}$ and $V(K_2) = \{0, 1\}$. For each $i \in \{1, 2, \dots, n\}$, let $u_{2i} = (v_i, 0)$ and $u_{2i+1} = (v_i, 1)$. The definition of the bow tie product implies the following statements:

- The vertices u_{2i} and u_{2j} are adjacent in H if and only if the vertices v_i and v_j are adjacent in G ; that is, if and only if $i - j \equiv \pm k \pmod{n}$ for some $k \in \{k_1, k_2, \dots, k_p\}$. Equivalently, u_{2i} is adjacent to u_{2j} if and only if $2i - 2j \equiv \pm k \pmod{2n}$ for some $k \in \{2k_1, 2k_2, \dots, 2k_l\}$.
- The vertices u_{2i+1} and u_{2j+1} are adjacent in H if and only if the vertices v_i and v_j are adjacent in G . Therefore, u_{2i+1} is adjacent to u_{2j+1} if and only if $2i - 2j \equiv \pm k \pmod{2n}$ for some $k \in \{2k_1, 2k_2, \dots, 2k_l\}$.
- The vertices u_{2i+1} and u_{2j} are adjacent in H if and only if the vertices v_i and v_j are adjacent in G ; that is, if and only if $i - j \equiv \pm k \pmod{n}$ for some $k \in \{k_1, k_2, \dots, k_p\}$. Equivalently, u_{2i+1} is adjacent to u_{2j} if and only if $(2i + 1) - 2j \equiv \pm k \pmod{2n}$ for some $k \in \{2k_1 + 1, 2k_2 + 1, \dots, 2k_l + 1\}$.

Thus, for each $i, j \in \{0, 1, 2, \dots, 2n - 1\}$, the vertices u_i and u_j are adjacent if and only if $i - j \equiv \pm k \pmod{2n}$ for some $k \in \{2k_1, 2k_2, \dots, 2k_l, 2k_1 + 1, 2k_2 + 1, \dots, 2k_l + 1\}$. \square

Corollary 26. *There exist infinitely many circulant graphs with $\gamma^\infty < \theta$.*

Proof. This follows from Fact 18, Corollary 19 and Proposition 25. \square

4.2.4 Planar graphs

Anderson, Barrientos, Brigham, Carrington, Vitray and Yellen [2] showed that any outerplanar graph has eternal domination number equal to its clique covering number. Moreover, no planar graph where $\gamma^\infty < \theta$ is known. This suggest that it might be true for general planar graphs and motivates the following question of Klostermeyer and Mynhardt.

Question 27 ([17]). *Is it true that $\gamma^\infty(G) = \theta(G)$ if G is planar?*

We first describe some properties of a smallest planar graph with $\gamma^\infty < \theta$, if it exists.

Proposition 28. *Let \mathcal{F} be a family of graphs satisfying a hereditary property \mathcal{P} ; i.e. if $G \in \mathcal{F}$ and H is a subgraph of G , then $H \in \mathcal{F}$. Suppose \mathcal{F} contains graphs G such that $\gamma^\infty(G) < \theta(G)$. Then, the smallest such graph G is a 2-connected vertex critical graph.*

Proof. It is clear that G is a vertex-critical graph; otherwise, by Fact 1, G contains a proper induced subgraph which is also in \mathcal{F} with $\gamma^\infty < \theta$ (contradiction). Suppose G is not 2-connected and let v be a cut vertex of G , that is a vertex such that $G - \{v\}$ has k components $\{G_1, G_2, \dots, G_k\}$ where $k \geq 2$. By the minimality of G , $\gamma^\infty(G_i) = \theta(G_i)$ for each $i \in \{1, 2, \dots, k\}$. Since $\gamma^\infty(G) \geq \sum_{i=1}^k \gamma^\infty(G_i)$, $\theta(G) \leq \sum_{i=1}^k \theta(G_i) + 1$ and $\gamma^\infty(G) < \theta(G)$, we conclude that $\gamma^\infty(G) = \sum_{i=1}^k \gamma^\infty(G_i)$ and $\theta(G) = \sum_{i=1}^k \theta(G_i) + 1$. This implies that there exists $i \in \{1, 2, \dots, k\}$ such that $\gamma^\infty(G_i) = \gamma^\infty(G_i + \{v\})$ and $\theta(G_i + \{v\}) = \theta(G_i) + 1$. In this case, $G_i + \{v\}$ is a smaller graph in \mathcal{F} with $\gamma^\infty < \theta$, which is a contradiction. \square

Since planarity is a hereditary property, the smallest planar graph with $\gamma^\infty < \theta$, if it exists, is a 2-connected vertex-critical graph. Observe that none of the graphs on 11 vertices or less satisfying $\gamma^\infty < \theta$ we found (Table 9) are planar, hence we have the following observation.

Observation 29. *There are no planar graphs on 11 vertices or less with $\gamma^\infty < \theta$.*

We also considered planar graphs of higher orders (12 and 13) in our search and we used plantri [4, 3] (version 5.2) to generate them. Due to the limitations of plantri, we only considered 3-connected planar graphs and obtained the following observation.

Observation 30. *There are no 3-connected planar graphs on 13 vertices or less with $\gamma^\infty < \theta$.*

Table 5: Number of 3-connected planar graphs on n vertices.

n	Total	$\alpha < \theta$	Vertex-Critical & $\alpha < \theta$	Vertex-Critical & $\gamma^\infty < \theta$
10	32300	2773	14	0
11	440564	25771	74	0
12	6384634	745440	878	0
13	96262938	6774391	2475	0

4.2.5 Claw-free graphs

A *claw-free graph* is a graph that does not contain a claw as an induced subgraph. In particular, any graph with independence number 2 is claw-free. The complements of the Mycielski graphs M_k described in Section 3 have independence number 2, eternal domination number 3 and clique covering number k for $k \geq 4$. Consequently, there are infinitely many claw-free graphs with $\gamma^\infty < \theta$.

4.2.6 Cubic graphs

A *cubic graph* is a graph in which all vertices have degree 3. Using NAUTY along with Algorithm 1, we generated the set of cubic graphs on fewer than 18 vertices and obtained the following observation.

Observation 31. *There are no cubic graph on $n \leq 16$ vertices with $\gamma^\infty < \theta$.*

Table 6: Number of connected cubic graphs on n vertices.

n	Total	$\alpha < \theta$	$\gamma^\infty < \theta$
4	1	0	0
6	2	0	0
8	5	2	0
10	19	9	0
12	85	46	0
14	509	320	0
16	4060	2888	0

4.3 Domination, eternal domination and clique covering


As we have seen above, the guards must always be located on the vertices of a dominating set in a graph G in order to defend a sequence of attacks on G . With this in mind, many researchers showed interest in characterizing graphs for which the domination number is equal to the eternal domination number. Klostermeyer and Mynhardt [18] posed the following question.

Question 32 ([18]). *Let G be a graph with $\gamma(G) = \gamma^\infty(G)$. Is any minimum dominating set of G an eternal dominating set of G ?*


The answer to Question 32 is no. We state this in the following proposition.

Proposition 33. *For any integer $k \geq 2$, there exists a graph G such that $\gamma(G) = \gamma^\infty(G) = k$ having minimum dominating sets which are not eternal dominating sets.*

Proof. Consider the graph G_1 in Figure 13a. It can be easily checked that $\gamma(G_1) = \gamma^\infty(G_1) = \theta(G_1) = 2$. Moreover, the set $\{u_1, u_4\}$ is a dominating set of G ; however, if the guards are located on the vertices u_1 and u_4 , then any response to an attack on the vertex u_0 leaves either the vertex u_2 or the vertex u_3 undominated on the next turn. Figure 13b shows a generalisation of the graph in Figure 13a where $\gamma(G_2) = \gamma^\infty(G_2) = k$. The set $\{u_1, u_4\} \cup \{v_0, v_1, \dots, v_{k-3}\}$ is a dominating set of size k , but any response to an attack on vertex u_0 leaves one of the vertices u_2, u_3 or w_0 undominated on the next turn. \square



(a) A graph G_1 with $\gamma(G_1) = \gamma^\infty(G_1) = 2$ having more minimum dominating sets than eternal dominating sets.



(b) A graph G_2 with $\gamma(G_2) = \gamma^\infty(G_2) = k$ having more minimum dominating sets than eternal dominating sets.

Figure 13: Two graphs having more minimum dominating sets than eternal dominating sets.

A question of Klostermeyer and MacGillivray, which was later stated as a conjecture by Klostermeyer and Mynhardt, is the following.

Conjecture 34 ($\gamma - \theta$ Conjecture [17, 18]). *For any graph G , $\gamma(G) = \gamma^\infty(G) \iff \gamma(G) = \theta(G)$.*

A computer search yields the following observation.

Observation 35. *There are no counterexample to the $\gamma - \theta$ conjecture of order $n \leq 11$.*

Table 7: Number of connected graphs on n vertices with $\gamma = \alpha$, $\gamma = \gamma^\infty$ and $\gamma = \theta$.

n	Total	$\gamma = \alpha$	$\gamma = \gamma^\infty$	$\gamma = \gamma^\infty = \theta$
5	21	6	5	5
6	112	24	22	22
7	853	88	67	67
8	11117	524	358	358
9	261080	4515	2265	2265
10	11716571	73515	23394	23394
11	1006700565	2324209	396755	396755

The computation in Table 7 was performed on a cluster which required approximately 85 CPU days. We found 54 graphs (listed in Table 9 in the appendix) with $\gamma^\infty < \theta$ among which there are 53 graphs with $\alpha = \gamma^\infty < \theta$ and none with $\gamma = \gamma^\infty < \theta$.

5 Acknowledgements

All computations in this paper were run on a PowerEdge R7425 server equipped with two AMD EPYC processors and totalling 64 threaded CPU cores. This machine was purchased using NSERC funding by Wendy Myrvold. We thank Wendy for her help with the computations.

We acknowledge the support of the Natural Sciences and Engineering Research Council of Canada (NSERC), PIN 04459, 253271.

Cette recherche a été financée par le Conseil de recherches en sciences naturelles et en génie du Canada (CRSNG), PIN 04459, 253271.



References

- [1] N. ALON AND J. H. SPENCER. The probabilistic method. With an appendix by Paul Erdős. *Wiley-Interscience Series in Discrete Mathematics and Optimization*, John Wiley & Sons, (1992).
- [2] M. ANDERSON, C. BARRIENTOS, R. C. BRIGHAM, J. R. CARRINGTON, R. P. VITRAY AND J. YELLEN. Maximum-demand graphs for eternal security. *J. Combin. Math. Combin. Comput.* 61 (2007), 111–127.
- [3] G. BRINKMANN. AND B. D. MCKAY. The program **plantri**. Available at <https://users.cecs.anu.edu.au/~bdm/plantri/>
- [4] G. BRINKMANN AND B. D. MCKAY. Fast generation of planar graphs. *MATCH Commun. Math. Comput. Chem.* 58 (2007), no. 2, 323–357.
- [5] A. P. BURGER, E. J. COCKAYNE, W. R. GRUNDLINGH, C. M. MYNHARDT, J. H. VAN VUUREN AND W. WINTERBACH. Infinite order domination in graphs. *J. Combin. Math. Combin. Comput.* 50 (2004), 179–194.
- [6] V. CHVÁTAL. The minimality of the Mycielski graph. *Graphs and Combinatorics (Proc. Capital Conf., George Washington Univ., Washington, D.C., 1973)*, *Lecture Notes in Math.* 406 (1974), 243–246, Springer, Berlin.
- [7] B. DESCARTES. A three colour problem. *Eureka* 9 (1947), 21.
- [8] B. DESCARTES. Solution to advanced problem no. 4526. *Amer. Math. Monthly* 61 (1954), 352.
- [9] P. ERDŐS, D. J. KLEITMAN AND B. L. ROTHSCHILD. Asymptotic enumeration of K_n -free graphs. *Colloquio Internazionale sulle Teorie Combinatorie (Rome, 1973)*, 2 (1976), 19–27.
- [10] W. GODDARD, S. M. HEDETNIEMI AND S. T. HEDETNIEMI. Eternal security in graphs. *J. Combin. Math. Combin. Comput.* 52 (2005), 169–180.
- [11] M. GRÖTSCHEL, L. LOVÁSZ AND A. SCHRIJVER. Geometric algorithms and combinatorial optimization. Springer-Verlag, (1988).
- [12] W. F. KLOSTERMEYER. Complexity of eternal security. *J. Combin. Math. Combin. Comput.* 61 (2007), 135–140.
- [13] W. F. KLOSTERMEYER, M. LAWRENCE AND G. MACGILLIVRAY. Dynamic dominating sets: the eviction model for external domination. *J. Combin. Math. Combin. Comput.* 97 (2016), 247–269.
- [14] W. F. KLOSTERMEYER AND G. MACGILLIVRAY. Eternal security in graphs of fixed independence number. *J. Combin. Math. Combin. Comput.* 63 (2007), 97–101.
- [15] W. F. KLOSTERMEYER AND G. MACGILLIVRAY. Eternal dominating sets in graphs. *J. Combin. Math. Combin. Comput.* 68 (2009), 97–111.
- [16] W. F. KLOSTERMEYER AND C. M. MYNHARDT. Domination, eternal domination and clique covering. *Discuss. Math. Graph Theory* 35 (2015), no. 2, 283–300.

- [17] W. F. KLOSTERMEYER AND C. M. MYNHARDT. Protecting a graph with mobile guards. *Appl. Anal. Discrete Math.* 10 (2016), no. 1, 1–29.
- [18] W. F. KLOSTERMEYER AND C. M. MYNHARDT. Eternal and secure domination in graphs. *Topics in domination in graphs, Dev. Math.* 64 (2020), 445–478, Springer, Cham.
- [19] L. LOVÁSZ. A characterization of perfect graphs. *J. Combin. Theory Ser. B* 13 (1972), 95–98.
- [20] L. LOVÁSZ. Normal hypergraphs and the perfect graph conjecture. *Discrete Math.* 2 (1972), no. 3, 253–267.
- [21] B. D. MCKAY AND A. PIPERNO. Practical graph isomorphism, II. *J. Symbolic Comput.* 60 (2014), 94–112.
- [22] J. MYCIELSKI. Sur le coloriage des graphes. *Colloq. Math.* 3 (1955), 161–162.

Appendix

All the graphs in this appendix are listed in Graph6 format.

Table 8: List of critical graphs with $\alpha < \theta$.

n	List of graphs
5	DUW
7	FCptO FCxv? FUzro
8	GCrbuW GCpveg GCxvcw GEhuSw
9	H?bBVbQ H?bBTjQ H?bBThY H?bBTiU H?bDJqU H?bF'xw H?bDjpw H?bfBqU H?bbVaU H?bbV_] H?bvbro H?rF'zo H?q'qjo H?o~fbo HCQ'faw HCQeJaL HCrfRym HCrbvW} HCrUrze HCpvfim HCpvexy HCpvVW} HCpvUzq HCpvUZU HCpvRym HCPunrM HCPunp] HCrJvi] HCzbvZe HCxvfri HCxvfpy HCxvez[HCxvezM HCvdjrm HEjfaxu HEhuTxm HEhvTy{ HUZvvx}

Table 9: List of connected graphs with $\gamma^\infty < \theta$.

n	List of graphs
10	IEhbtj{ro IEhbtn{ro
11	JQyurj yt ? JEhbtj{rvu? JEhbtj{rvx? JEhbtj{rvf? JEhbtj{rvv? JEhbtj{rvT_ JEhbtj{rtt_ JEhbtj{rrt_ JEhbtj{rv}? JEhbtj{rv ? JEhbtj{rvv? JEhbtj{rv^? JEhbtj{rvx_ JEhbtj{rvt_ JEhbtj{rvl_ JEhbtj{rv_ JEhbtj{rvf_ JEhbtj{rtv_ JEhbtj{rv~? JEhbtj{rv _ JEhbtj{rvv_ JEhbtj{rv~_ JEhbtnm~E ? JEhbtnm~FZ? JEhbtnm~D}_ JEhbtnm~@}_ JEhbtnN~Fu? JEhbtnN~Bv? JEhbtnN~Fw_ JEhbtnN~Fe_ JEhbtnN~Eu_ JEhbtnN~Bu_ JEhbtnN~Ef_ JEhbtnN~F}? JEhbtnN~Fv? JEhbtnN~F{_ JEhbtnN~Fu_ JEhbtnN~F}_ JEhbtnN~Ff_ JEhbtnN~Ev_ JEhbtnN~Bv_ JEhbtnN~F~? JEhbtnN~F}_ JEhbtnN~Fv_ JEhbtnN~F~_ JEhvUtn~D{_ JEhutz{xr_ JEhru^u~Fj? JEhru v~Et_ JEjbtnN~Dm_ JEnfbz\zvb? JCXetqu uz? JCXetq}vVm? JCXetq} uz?

Table 10: List of triangle-free graphs with $\gamma^\infty < \theta$.

n	List of graphs
13	L?'@F?M]DgYOFo L?'DAboU'w@{hS L?'@?boNAsLGBw L?'@?boNAsOwYO L?'@?boNAs@{os L?'@Cbod'w@{YS L?BDB?{{AsRGS L?'@C'w1?{DgQs L?BDB?{Ucq^?Fo L?'@F@w1EcBoBo L?'@F@w1EcBoFo L?'@F@w1EcBgFo L?'@F?kQ_]YS1C

Table 11: List of maximal triangle-free graphs with $\gamma^\infty < \theta$.

n	List of graphs
13	L?'DABOu'w@{hS L?BDB?{{AsRGBs L?BDB?{Ucq^?Fo L?'@F?kQ_}YS1C L?'@?boNAsOwYO
14	M?'@F?kQckBwsglC? M?'@?boNAs@{oshQ?
15	N??EFBOK_)DsrC~^?_? N??EFBOK_ZbwJgrC^?_? N?AAF@SBo}TSM_{CN_? N?AA@aoqTsVO^?BwGl_ N?AADB_HdsN_VOE[B{? N?AADB_HeobMm_VOVB{? N?AADB_HaqbM^?m_B{? N?ABB?WwGNN_eoFs^@? N?ABAAqFbgTG}?JWPX? N??CBBoK@HqfO^_@}? N?'@F?kQckBwsglCN@? N??ED?qw@{JgRW^?By? N??ED?qwAaHkjA^?By? N??ED?qwAaHkJa^?By? N??ED?qwAaPdRW^?By? N??ED?qwAyJgRW^?By? N??ED?qwEXHkN_~?By? N??ED?qwEXHkVOFs^?_? N??ED?qwAZHkN_~?By? N??CB@eOmCuiI^?No? N?AAD@WUDD_)N_VQ^?_ N??CFBOBw}Ds^~?No? N??EDBo@{@{JgFoB{^?_? N?AA@BOHn_N_m_uS@ ? N??CFBOK_)BwJgrCNo? N??CB@OKCyTcrC^_@}? N??ED@_MdoRC^?FwGj_ N??EDbgl?}EsLg^?HIo N?AA@B_@zwVouOFw^A? N?AAF?e{EpN_m_Fo@y? N?AA@Bo@{JgeoB{@}? N??ED@_nfo^?VOFG? _ N??ED@_NayRc^?Fw^?_? N??ED@_NayRcN_~?B{? N?AA@Boy?^FoVOeoAy? N?AADb_ro^Bs^?m_UQ? N?AADb_RtC^?N_BwBs? N?AA@b_HaYPWFguCN_? N??CB?Xs?}cufOxGNo? N?AA@BoZ?^QYm_VOFO? N??CABoyBKRHBwDsVg? N?AA@?O{BwVOuOyG@}? N?AA@?O{BwRWigRW@}? N?AA@?O{@kJgZGaw@N? N?AA@?O{@kJgigB{XL? N?AA@?O{@{JgeoyG@}? N?AA@?O{@{JgZGaw@}? N?AA@?O{@{XKVOig@}? N?AA@?O{?}DsRwaw@}? N??E@b_sEYBsJc^?@}? N?AAD@O{Ai^?N_ii@}? N?AAD@O{AiN_N_B{TT? N?AAD@O{AyN_m_ig@}? N??ED?WHiJJIMaxG^?_? N?ABA_oebgTGqGBwP\? N?ABA_oeDQXC^?BwGf? N?AA@b_qQYAu^?m_@{? N?AA@b_qO^Bs^?m_Eq? N?AA@b_qO1^?{^?m_??? N?AA@b_qO1^?{^?m_N_? N??EDaKIaYMCN_tCET? N?AA@BoJ_~YI^?eoVO? N?AA@?WwGNN_m_uON_? N?AA@?WwGNN_m_uOBy? N?AA@?WwGNN_m_uO^@? N?AA@b_RRgBsN_aquQ? N?AA@b_RSUEoFg}?XG_ N?AA@b_RSUUQ\AFg^?? N??CFBOKewBwJg^~^?_? N??CFBOKew'mFoJgNo? N??ED?qN_)HkrG^?^?_? N??ED?qN_)HkrGVO^?_? N??ED?qSROQ'FobK^?_? N??ED?qSO}HkJc^?^?_? N??ED?qSO}HkJcRg^?_? N??ED?qSO}HkJcK^?_? N??ED@OAw^FoVojG}a? N?AA@B_{BwVOuOFw@}? N?AA@B_{@{JgeoB{B{? N??EDaKRR'}?N_Br^?_? N??CB@OyDPRgBwDuNo? N?AAF@Sy?mtSm_FoN_? N?AA@AOwFoRWJgrWBy? N?AA@AOwDsZGZGFs@}? N?AA@ao@{@kGkqQB{VO_ N?AAD@Wgno^?N_VS? _ N??CEBoy?^Ay^?NoNo? N??CBBOyBKRHDsJg^?_? N?AADBOQRwDsm_FoEq? N?AAD?oqN_VoLcVO@ ? N?AAD?oqJgVOBwVOPT? N??CFBOQ_)BwJgLC^?_? N??CFBOQ_)Ds1C^~^?_? N??ED?qEqyHkxCN_~^?_? N??ED?qEo}HkxC^?^?_? N??ED?qEo}WeRW^?^?_? N??ED?qEqZHkN_xC^?_? N?AA@B_)BwFoVOeoB{? N?AA@B_)@{FoVOeoB{? N?AA@B_)@{JgeoFwB{? N?AA@AO{BwRWJgrW@}? N?AA@AO{DsZGFoZG@}? N?AA@AO{CuFoVOZG@}? N?AA@AO{BHVOVOeq@}? N?AA@AO{BHFOVOB{RX? N?AA@BO{FoJgeoJg@}? N?AA@BO{@{JgeoJg@}? N??CFBO{?}DsB^?No? N?'@?boNAs[G'oBwBo_ N?'@?boNAsLG'o'o@{? N?'@?boNAsLG'o'oN?W N?'@?boNAsSgooBo?Fw N?AA@B_)BwVOuOA{B{? N?AA@B_)BwVOasuOB{? N?AA@B_)DSPYtQuOB{? N?AA@B_)@[YH]?asB{? N?AA@B_)@[YHasVOB{? N?AA@B_)CU^?N_VOVB{? N?AA@B_)CUvON_VOVB{? N?AA@B_)CUvOuON_B{? N?AA@B_)CUvOuOFwB{? N?AA@B_)CUfoVOeoB{? N??ED@o}@sJGBw^?Of_ N?AA@_owBwVOFfcqW@}? N?AA@BO}FoJgeoJg?^? N?AA@_gow]As}??^?VO? N?AA@_gow]As^?m_N_? N?AA@_gow]As^?m_Fq? N?AA@_gow]As^?m_]@? N?AA@_gow]As^?m_YD? N?AA@_gow]As^?m_]@_ N?AA@b@BpLZ@^?m_Fs? N?AA@b@Bo\N_m_{EFs? N?AA@b@Bv_n_m_NgFs? N?AA@BOX@]VO}?RY? N?AAF@ScfoTSZCFoBw? N?AAF@Sc'kBwigZC@N? N??ED?ZDp{[cVOyC^?_? N?AA@_o{FoFoRWaw@}? N?AA@_o{@kGkB{qYVO_ N?AA@_o{AJN_qWB{VP? N??ED?ZFp{[cVO^?^?_? N??FEaK[N_BoJO^?@z? N?AA@?WeF@Cueo^?Gm? N??CB@OeRcCuFobY^?_? N?AADB_)FoVOVOA{B{? N?AADB_)BwVOVOA{B{? N?AA@B_HbwVOVOFwZI? N?AA@B_HbwVOFwFwZI? N?AA@B_HbwVOE[FwZI? N?AA@B_HaI^?uSmaB{? N?AA@B_HaIN_FwuSVP? N?AA@B_HaIN_EMeoN_? N?AA@B_HaIN_EMeoRW? N?AA@B_HaIN_EMeoB{? N?AA@B_HaIOW}?FwBF_ N?AA@B_HaIRWFwuSVP? N?AA@B_HaIB{uSmaB{? N?AA@B_HaITP}?uSB{? N?AA@B_HaITP^?FwZI? N?AA@B_HaITPm_FwZI? N?AA@B_HayRW}?FwZI? N?AA@B_HayRW^?FwZI? N?AA@B_H_rjgm_FwZI? N?AADa@{AJFoVOB{VP? N??ED@_@{FoVOFG@}? N??ED@_@{JgfgB{^?_? N??ED@_@{JgfgB{B{? N?AA@_o}@{^?RWaw?}_ N??CFBOQdguafOJg^?_? N?AADB_kdsN_RSfwb{? N?AADB_k_NHi^?Fw^@? N?ABAAQT?N_}?iSDM?