

SCHWARZ SYMMETRIZATIONS IN PARABOLIC EQUATIONS ON COMPLETE MANIFOLDS

HAIQING CHENG, TENGFEI MA, AND KUI WANG

ABSTRACT. In this article, we prove a sharp estimate for the solutions to parabolic equations on manifolds. Precisely, using symmetrization techniques and isoperimetric inequalities on Riemannian manifold, we obtain a Bandle's comparison on complete noncompact manifolds with nonnegative Ricci curvature and compact manifolds with positive Ricci curvature respectively. Our results generalize Bandle's result [6] to Riemannian setting, and Talenti's comparison for elliptic equation on manifolds by Colladay-Langford-McDonald [12] and Chen-Li [9] to parabolic equations.

1. INTRODUCTION

Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with smooth boundary, $\Omega^\# \subset \mathbb{R}^n$ be a round ball with the same volume as Ω , $f(x)$ and $g(x)$ be nonnegative functions on Ω , and $f^\#, g^\#$ be the Schwarz rearrangement of f and g , see Definition 2.1 below. Let $u(x)$ and $v(x)$ be solutions to

$$\begin{cases} u_t(x, t) - \Delta u(x, t) = f(x), & (x, t) \in \Omega \times (0, \infty), \\ u(x, 0) = g(x), & x \in \Omega, \\ u(x, t) = 0, & (x, t) \in \partial\Omega \times (0, \infty), \end{cases}$$

and

$$\begin{cases} v_t(x, t) - \Delta v(x, t) = f^\#(x), & (x, t) \in \Omega^\# \times (0, \infty), \\ v(x, 0) = g^\#(x), & x \in \Omega^\#, \\ v(x, t) = 0, & (x, t) \in \partial\Omega^\# \times (0, \infty). \end{cases}$$

Then for all $(a, t) \in (0, |\Omega|) \times [0, +\infty)$, it holds

$$(1.1) \quad \int_0^a u^*(s, t) ds \leq \int_0^a v^*(s, t) ds,$$

where u^* is the decreasing rearrangement of u , see (2.1) below. Inequality (1.1) was proved by Bandle [6] for strong solutions, by Vázquez [24] and Mossino-Rakotoson [20] for weak solutions. The elliptic version of (1.1) is known as Talenti's comparison [22, 23]. The proof of (1.1) is mainly based on the isoperimetric inequality and symmetrization techniques. Talent and Bandle's comparisons play an important role in mathematical studies, since there provide sharp estimates for solutions to elliptic and parabolic equations, Faber-Krahn type inequality for eigenvalues [22, 3], and the bounds on the exit times of Brownian motion[8, 12]. Bandle's comparison (1.1) was generalized to nonlinear parabolic equations with Dirichlet

2020 *Mathematics Subject Classification.* 53C21, 53C42.

Key words and phrases. Bandle's comparison, Parabolic equations, Manifolds.

The research of the third author is supported by NSFC No.11601359.

boundary condition (see for instance [4, 5] and references therein), and to parabolic equations with Neumann boundary condition [14] as well. We also refer the reader to the excellent books [16, 17] for related topics.

Recently, Talenti's comparison for Dirichlet boundary was generalized to the compact manifolds with positive Ricci curvature by Colladay, Langford and McDonald [12], and to complete noncompact manifold with nonnegative Ricci curvature and positive asymptotic volume ratio by Chen and Li [9]. Also Talenti's comparison for Robin boundary was proved in both Euclidean space [3, 2] and complete manifolds [10, 11]. To the best of our knowledge, Bandle's comparison, however, did not receive as much attention as Talenti's comparison on manifolds. The purpose of the present paper is to study the Bandle's comparison for solutions to parabolic equation on manifolds. In particular, we will establish a sharp comparison between the solutions to (1.3) and (1.4) on complete and noncompact manifolds with nonnegative Ricci curvature and positive asymptotic volume ratio, and compact manifolds with positive Ricci curvature, see Theorem 1.1 below.

Let (M, g) be a complete Riemannian manifold of dimension n , which is either compact with $\text{Ric} \geq (n-1)\kappa$ for $\kappa > 0$ or noncompact with nonnegative Ricci curvature and positive asymptotic volume ratio. Denote by

$$(1.2) \quad \theta := \begin{cases} \lim_{r \rightarrow \infty} \frac{|B_p(r)|}{\omega_n r^n}, & \kappa = 0, \\ \frac{|M|}{|M_\kappa|}, & \kappa > 0, \end{cases}$$

where $B_p(r)$ is the geodesic ball centered at p with radius r in M , M_κ is the n dimensional space form of constant sectional curvature κ , ω_n is the volume of the unit ball in \mathbb{R}^n , and $|M|$ denotes the volume of M . It follows from the volume comparison that $\theta \leq 1$. Let $\Omega \subset M$ be a bounded domain with smooth boundary, $f(x)$ and $g(x)$ be smooth and nonnegative function not identically zero on Ω , we consider the following Cauchy problem

$$(1.3) \quad \begin{cases} u_t(x, t) - \Delta u(x, t) = f(x), & (x, t) \in \Omega \times (0, \infty), \\ u(x, t) = 0, & (x, t) \in \partial\Omega \times (0, \infty), \\ u(x, 0) = g(x), & x \in \Omega. \end{cases}$$

Let Ω^\sharp be a geodesic ball in M_κ satisfying $\theta|\Omega^\sharp| = |\Omega|$, f^\sharp and g^\sharp defined on Ω^\sharp are the Schwarz rearrangement of f and g respectively. Denote by $v(x)$ the solution to the following Schwarz rearrangement system of (1.3)

$$(1.4) \quad \begin{cases} v_t(x, t) - \Delta v(x, t) = f^\sharp(x), & (x, t) \in \Omega^\sharp \times (0, \infty), \\ v(x, t) = 0, & (x, t) \in \partial\Omega^\sharp \times (0, \infty), \\ v(x, 0) = g^\sharp(x), & x \in \Omega^\sharp. \end{cases}$$

The main result of this paper is the following theorem, which gives a sharp Bandle's comparison on Riemannian manifolds.

Theorem 1.1. *Let $u(x)$ and $v(x)$ be the solutions to (1.3) and (1.4) respectively. Then*

$$(1.5) \quad U(a, t) \leq V(a, t), \quad (a, t) \in (0, |\Omega^\sharp|] \times (0, \infty).$$

Where

$$(1.6) \quad U(a, t) = \frac{1}{\theta} \int_0^{\theta a} u^*(s, t) ds, \quad V(a, t) = \int_0^a v^*(s, t) ds.$$

Moreover if inequality (1.5) holds as an equality for all $(s, t) \in (0, |\Omega^\sharp|] \times (0, \infty)$, then M is isometric to M_κ , Ω is isometric to Ω^\sharp , and $u(x, t) = v(x, t)$ on $\Omega^\sharp \times (0, \infty)$.

Remark 1.2. *In this paper, we mainly focus on Bandle's comparison (1.1) on manifolds and we then set up Theorem 1.1 in smooth case. In fact, Theorem 1.1 remains valid for weak solutions to (1.3), see [20] and [4].*

The following corollary is a simple consequence of (1.5), which gives a sharp L^p norm estimate of u by v on space variables for all t .

Corollary 1.3. *Let $u(x)$ and $v(x)$ be the solutions to (1.3) and (1.4) respectively. Then we have*

$$(1.7) \quad \left(\frac{1}{\theta} \int_{\Omega} u^p(x, t) dx \right)^{1/p} \leq \left(\int_{\Omega^\sharp} v^p(x, t) dx \right)^{1/p}$$

for all $t > 0$ and $p \in [1, \infty]$.

We mention finally that Talenti's comparison for Poisson equation with Robin boundary has recently been confirmed by Alvino, Nitsch, and Trombetti in dimension two, as well as an integral estimate for higher dimensions [3]. It is a natural question to extend Bandle's comparison to heat equation with Robin boundary, and we leave this to a future study.

This paper is organized as follows. In Section 2 we recall the Schwarz rearrangements and isoperimetric inequalities on manifolds. In Section 3 we prove Theorem 1.1.

2. PRELIMINARIES

Let Ω be a bounded smooth domain in M and Ω^\sharp be a geodesic ball in M_κ (i.e. \mathbb{R}^n or \mathbb{S}^n) with volume $|\Omega|/\theta$, where θ is a constant defined by (1.2). We recall the definitions and properties of the Schwarz rearrangement of nonnegative functions on manifolds, see also [9, Section 2] and [12, Section 2].

Definition 2.1. *Let $h(x)$ be a nonnegative measurable function on Ω . Denote by $\Omega_{h,s} = \{x \in \Omega : h(x) > s\}$ and $\mu_h(s) = |\Omega_{h,s}|$, the decreasing rearrangement h^* of h is defined by*

$$(2.1) \quad h^*(s) = \begin{cases} \text{ess sup}_{x \in \Omega} h(x), & s = 0, \\ \inf\{t \geq 0 : \mu_h(t) < s\}, & s > 0, \end{cases}$$

for $s \in [0, |\Omega|]$. The Schwarz rearrangement of h is defined by

$$(2.2) \quad h^\sharp(x) = h^*(\theta \omega_n r^n(x)), \quad x \in \Omega^\sharp,$$

where $r(x)$ is the distance function from the center of Ω^\sharp in M_κ , and ω_n is the volume of unit ball in \mathbb{R}^n .

It follows directly from (2.1) and (2.2) that

$$(2.3) \quad \mu_h(s) = \theta \mu_{h^\sharp}(s)$$

for $s \geq 0$. Meanwhile, the Fubini's theorem gives

$$(2.4) \quad \int_{\Omega} h^p(x) dx = \int_0^{|\Omega|} (h^*)^p(s) ds = \theta \int_{\Omega^\sharp} (h^\sharp)^p(x) dx,$$

for $h \in L^p(\Omega)$, $p \geq 1$. Moreover for any nonnegative functions $f(x)$ and $g(x)$, the following inequality, known as Hardy-Littlewood inequality, holds true

$$(2.5) \quad \int_{\Omega} f(x)g(x) dx \leq \int_0^{|\Omega|} f^*(s)g^*(s) ds.$$

Taking $g(x)$ as the characteristic function of $\Omega_{h,s}$ in above inequality yields

$$(2.6) \quad \int_{\Omega_{h,s}} f(x) dx \leq \int_0^{\mu_h(s)} f^*(\eta) d\eta.$$

To prove Theorem 1.1, we require the following isoperimetric inequality on manifolds with Ricci curvature bounded from below.

Theorem 2.2. *With M , θ , Ω and Ω^\sharp as above, there holds*

$$(2.7) \quad |\partial\Omega| \geq \theta |\partial\Omega^\sharp|,$$

where $|\partial\Omega|$ denotes the $(n-1)$ -dimensional area of $\partial\Omega$. The equality holds if only if Ω is isometric to Ω^\sharp .

When $\text{Ric} \geq n-1$, inequality (2.7) was shown by Lévy and Gromov [15], see also Theorem 2.1 of [21]. When M is noncompact and $\text{Ric} \geq 0$, inequality (2.7) was shown in dimension three by Agostiniani, Fogagnolo and Mazzieri [1] and in all dimensions by Brendle [7]. Meanwhile, Brendle proved in [7] that (2.7) also holds true when Ω is a compact minimal submanifold of M of dimension $n+2$ with nonnegative sectional curvature.

3. PROOF OF THEOREM 1.1

In this section, we will prove the main theorem. For simplicity, we write $\Omega_u(a, t)$ and $\Omega_u^\sharp(a, t)$ as $\Omega_{a,t}$ and $\Omega_{a,t}^\sharp$ respectively for short. Let

$$\Phi(s) = |\partial B_s|$$

for $s \in (0, |\Omega^\sharp|]$, where B_s is a round geodesic ball in M_κ with volume s . It can be easily checked that $\Phi(s) = n\omega_n^{1/n} s^{(n-1)/n}$ if $\kappa = 0$, and $\Phi(s)$ is monotone increasing in s .

Lemma 3.1. *Under the hypotheses of Theorem 1.1, we have*

$$(3.1) \quad \theta^2 \Phi^2\left(\frac{\mu_u(s, t)}{\theta}\right) \leq -\partial_s \mu_u(s, t) \int_{\Omega_{s,t}} f(x) - u_t(x, t) dx,$$

and

$$(3.2) \quad \Phi^2(\mu_v(s, t)) = -\partial_s \mu_v(s, t) \int_{\Omega_{s,t}^\sharp} f^\sharp(x) - v_t(x, t) dx,$$

for a.e. $t > 0$.

Proof. By the Sard's theorem, for each $t > 0$ we have

$$(3.3) \quad \partial\Omega_{s,t} = \{x \in \Omega : u(x, t) = s\}.$$

for almost every $s \geq 0$. Observing from isoperimetric inequality (2.7) that

$$(3.4) \quad \theta \Phi\left(\frac{|\Omega_{s,t}|}{\theta}\right) \leq |\partial\Omega_{s,t}|,$$

and from the Hölder inequality that

$$(3.5) \quad |\partial\Omega_{s,t}|^2 \leq \int_{\partial\Omega_{s,t}} \frac{1}{|\nabla u|} dA \int_{\partial\Omega_{s,t}} |\nabla u| dA,$$

then we get

$$(3.6) \quad \theta^2 \Phi^2\left(\frac{|\Omega_{s,t}|}{\theta}\right) \leq \int_{\partial\Omega_{s,t}} \frac{1}{|\nabla u|} dA \int_{\partial\Omega_{s,t}} |\nabla u| dA,$$

here and thereafter dA denotes the induced measure on $(n-1)$ dimensional surface in Ω . Because $u(x, t)$ is vanishing on $\partial\Omega$, so $\Omega_{s,t}$ is a compact set in Ω , therefore by the coarea formula we have

$$\partial_s \mu_u(s, t) = - \int_{\partial\Omega_{s,t}} \frac{1}{|\nabla u|} dA$$

then (3.6) becomes to

$$(3.7) \quad \theta^2 \Phi^2\left(\frac{|\Omega_{s,t}|}{\theta}\right) \leq \partial_s \mu_u(s, t) \int_{\partial\Omega_{s,t}} |\nabla u| dA.$$

For $t > 0$, noticing from Definition 2.1 and equality (3.3) that

$$|\nabla u| = -\frac{\partial u}{\partial \nu}$$

on $\partial\Omega_{s,t}$ for a.e. $s > 0$, where ν is the unit outer normal to $\partial\Omega_{s,t}$, we then get

$$\int_{\partial\Omega_{s,t}} |\nabla u| dA = \int_{\Omega_{s,t}} -\Delta u(x) dx = \int_{\Omega_{s,t}} f(x) - u_t(x, t) dx.$$

So inequality (3.1) follows from the above equality and inequality (3.7).

If $v(x, t)$ is the solution to system (1.4), $v(x, t)$ is radial and decreasing along the radial direction on Ω^\sharp for all t , hence $\Omega_{s,t}^\sharp$ is a round ball. Therefore both (3.4) and (3.5) hold as equalities with $\theta = 1$ if we replace u by v , hence equality (3.2) holds true. \square

Now we turn to prove the main theorem.

Proof of Theorem 1.1. Recall

$$U(a, t) = \frac{1}{\theta} \int_0^{\theta a} u^*(s, t) ds$$

for $a \in [0, |\Omega^\sharp|)$, then the first derivative of a gives

$$U'(a, t) = u^*(\theta a, t),$$

and from Lemma 1.1 of [6] it holds

$$(3.8) \quad U_t(a, t) = \frac{\partial}{\partial t} \left(\frac{1}{\theta} \int_{\Omega_{u^*,t}} u(x, t) dx \right) = \frac{1}{\theta} \int_{\Omega_{u^*,t}} u_t(x, t) dx,$$

where u^* is evaluated at $(\theta a, t)$ and $U' = \partial_a U$. Observing that

$$\mu_u(u^*(\theta a, t)) = \theta a,$$

for a.e. $a > 0$ (c.f. [20, page 66]), and differentiating above identity in a yields

$$\mu'_u(u^*(\theta a, t)) U''(a, t) = \theta,$$

then

$$U''(a, t) = \frac{\theta}{\mu'_u(u^*)} \geq -\frac{\int_{\Omega_{u^*,t}} f(x) - u_t(x, t) dx}{\theta \Phi^2(a)},$$

where we used (3.1) in the inequality. Recall from (2.6) that

$$(3.9) \quad \int_{\Omega_{u^*,t}} f(x) dx \leq \int_0^{a\theta} f^*(s) ds = \theta \int_0^a (f^\sharp)^*(s) ds,$$

then we deduce that

$$(3.10) \quad U_t(a, t) - \Phi^2(a)U''(a, t) - \int_0^a (f^\sharp)^*(s) ds \leq 0$$

for $(a, t) \in (0, |\Omega^\sharp|) \times (0, \infty)$ in the viscosity sense, where we used equality (3.8).

Note that $v(x, t)$ is radial function decreasing along the radial direction, and f^\sharp is radial as well, then the inequality (3.9) holds as equality if we replace u by v , Ω by Ω^\sharp and $f(x)$ by f^\sharp according to equality case of Hardy-Littlewood inequality, i.e.

$$(3.11) \quad \int_{\Omega_{v^*,t}^\sharp} f(x) dx = \int_0^a (f^\sharp)^*(s) ds.$$

Combining equality (3.2) and equality (3.11), we find similarly as (3.10) that

$$(3.12) \quad V_t(a, t) - \Phi^2(a)V''(a, t) - \int_0^a (f^\sharp)^*(s) ds = 0,$$

for $(a, t) \in (0, |\Omega^\sharp|) \times (0, \infty)$. It then follows from inequality (3.10) and equality (3.12) that

$$(3.13) \quad (U - V)_t(a, t) - \Phi^2(a)(U - V)''(a, t) \leq 0.$$

On the other hand, by the initial condition of u and v we have

$$(3.14) \quad \begin{aligned} U(a, 0) - V(a, 0) &= \frac{1}{\theta} \int_0^{\theta a} u^*(s, 0) ds - \int_0^a v^*(s, 0) ds \\ &= \frac{1}{\theta} \int_0^{\theta a} g^*(s) ds - \int_0^a (g^\sharp)^*(s) ds \\ &= 0. \end{aligned}$$

Moreover, by direct calculations we have

$$(3.15) \quad U(0, t) = V(0, t) = 0,$$

and

$$(3.16) \quad U'(|\Omega^\sharp|, t) - V'(|\Omega^\sharp|, t) = u^*(|\Omega|, t) - v^*(|\Omega^\sharp|, t) = 0.$$

We then conclude that differential inequality (3.13) holds with initial condition (3.14) and boundary conditions (3.15) and (3.16). Therefore (1.5) follows by the standard maximum principle for viscosity solutions to parabolic equation, see [13, Theorem 3.3] and [18, Section 2].

Now we consider the equality case of (1.5). If $U(a, t) = V(a, t)$, then $u^*(\theta a, t) = v^*(a, t)$, hence $\mu_u(u^*, t) = \theta \mu_v(v^*, t)$ for all $a > 0$. Then we estimate that

$$\begin{aligned}
 |\partial \Omega_{u^*, t}|^2 &\leq -\mu'_u(u^*, t) \left(\int_{\Omega_{u^*, t}} f(x, t) - u_t(x, t) \, dx \right) \\
 &\leq -\mu'_u(u^*, t) \left(\int_0^{\theta a} f^*(s, t) \, ds - \theta U_t(a, t) \right) \\
 &= -\theta^2 \mu'_v(v^*, t) \left(\int_0^a (f^\#)^*(s, t) \, ds - V_t(a, t) \right) \\
 &= \theta^2 |\partial \Omega_{v^*, t}^\#|^2.
 \end{aligned}
 \tag{3.17}$$

where in the first step we used inequality (3.1), in the second inequality we used (3.8) and (3.9). On the other hand, the isoperimetric inequality (2.7) gives

$$|\partial \Omega_{u^*, t}|^2 \geq \theta^2 |\partial \Omega_{v^*, t}^\#|^2,$$

so inequality (3.17) holds as equality. Thus we conclude from the equality case in (2.7) that $\Omega_{u^*, t}$ is a geodesic ball in M_κ , $\theta = 1$ and $u(x, t) = v(x, t)$ on $\Omega^\# \times [0, \infty)$, hence $f(x) = f^\#(x)$ and $g(x) = g^\#(x)$. \square

Proof of Corollary 1.3. The proof of this lemma is in the same spirit as in Theorem 3.2 of [6]. For the reader's convenience, we give the details. For $p \geq 1$, the Fubini's theorem gives

$$\frac{1}{\theta} \int_{\Omega} u^p(x, t) \, dx = \frac{1}{\theta} \int_0^{|\Omega|} (u^*)^p(s, t) \, ds = \int_0^{|\Omega^\#|} (u^*)^p(\theta s, t) \, ds$$

and

$$\int_{\Omega^\#} v^p(x, t) \, dx = \int_0^{|\Omega^\#|} (v^*)^p(s, t) \, ds.$$

Then we estimate by integration by parts that

$$\begin{aligned}
 &\frac{1}{\theta} \int_{\Omega} u^p(x, t) \, dx - \int_{\Omega^\#} v^p(x, t) \, dx \\
 &= \int_0^{|\Omega^\#|} (u^*)^p(\theta s, t) - (v^*)^p(s, t) \, ds \\
 &\leq \int_0^{|\Omega^\#|} p(u^*)^{p-1}(u^* - v^*) \, ds \\
 &= (U - V)p(u^*)^{p-1} \Big|_0^{|\Omega^\#|} - \int_0^{|\Omega^\#|} (U - V)p(p-1)(u^*)^{p-2} \partial_s u^* \, ds,
 \end{aligned}
 \tag{3.18}$$

where in the inequality we used the elementary inequality $x^p - y^p \leq px^{p-1}(x - y)$ for $x, y \in \mathbb{R}$ when $p \geq 1$. By (3.15), (1.5) and $\partial_s u^* \leq 0$, we see

$$(U - V)p(u^*)^{p-1} \Big|_0^{|\Omega^\#|} = (U - V)p(u^*)^{p-1} \Big|_{s=|\Omega^\#|} \leq 0,$$

and

$$(U - V)p(p-1)(u^*)^{p-2} \partial_s u^* \geq 0,$$

therefore it follows from (3.18) that

$$\frac{1}{\theta} \int_{\Omega} u^p(x, t) dx - \int_{\Omega^{\sharp}} v^p(x, t) dx \leq 0$$

for all $p \geq 1$, hence for $p = \infty$. We complete the proof of the corollary. \square

In Euclidean space, the argument in the proof of (1.5) can be adapted to prove Bandle's comparison for the heat kernels. This is outlined in Theorem 2.2 of [6]. It's not hard to adapt our argument to prove Bandle's comparison for the heat kernels on compact manifolds with positive Ricci curvature, and complete and noncompact manifolds with nonnegative Ricci curvature and positive asymptotic volume ratio. Hence one can easily obtain the Faber-Krahn inequality for the first Dirichlet eigenvalue on such manifolds via heat kernel comparison, which has already been proved by Mattia-Lorenzo [19] and Chen-Li [9] respectively.

REFERENCES

- [1] Virginia Agostiniani, Mattia Fogagnolo, and Lorenzo Mazzieri. Sharp geometric inequalities for closed hypersurfaces in manifolds with nonnegative Ricci curvature. *Invent. Math.*, 222(3):1033–1101, 2020.
- [2] A. Alvino, F. Chiacchio, C. Nitsch, and C. Trombetti. Sharp estimates for solutions to elliptic problems with mixed boundary conditions. *J. Math. Pures Appl. (9)*, 152:251–261, 2021.
- [3] A. Alvino, C. Nitsch, and C. Trombetti. A talenti comparison result for solutions to elliptic problems with robin boundary conditions. *to appear on Comm. Pure Appl. Math.*, *arXiv:1909.11950*, 2019.
- [4] A. Alvino, G. Trombetti, and P.-L. Lions. Comparison results for elliptic and parabolic equations via Schwarz symmetrization. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 7(2):37–65, 1990.
- [5] Angelo Alvino, Roberta Volpicelli, and Bruno Volzone. Comparison results for solutions of nonlinear parabolic equations. *Complex Var. Elliptic Equ.*, 55(5-6):431–443, 2010.
- [6] Catherine Bandle. On symmetrizations in parabolic equations. *J. Analyse Math.*, 30:98–112, 1976.
- [7] Simon Brendle. Sobolev inequalities in manifolds with nonnegative curvature. *to appear on Comm. Pure Appl. Math.*, *arXiv:2009.13717*, 2020.
- [8] A. Burchard and M. Schmuckenschläger. Comparison theorems for exit times. *Geom. Funct. Anal.*, 11(4):651–692, 2001.
- [9] Daguang Chen and Haizhong Li. Talenti's comparison theorem for poisson equation and applications on riemannian manifold with nonnegative ricci curvature. *arXiv:2104.05568*, 2021.
- [10] Daguang Chen, Haizhong Li, and Yilun Wei. Comparison results for solutions of poisson equations with robin boundary on complete riemannian manifolds. *arXiv:2110.0605*, 2021.
- [11] Haiqing Cheng, Tengfei Ma, and Wang Kui. Comparison results for poisson equation with mixed boundary condition on manifolds. *arXiv:2110.06814*, 2021.
- [12] Don Colladay, Jeffrey J. Langford, and Patrick McDonald. Comparison results, exit time moments, and eigenvalues on Riemannian manifolds with a lower Ricci curvature bound. *J. Geom. Anal.*, 28(4):3906–3927, 2018.
- [13] Michael G. Crandall, Hitoshi Ishii, and Pierre-Louis Lions. User's guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc. (N.S.)*, 27(1):1–67, 1992.
- [14] V. Ferone and A. Mercaldo. Neumann problems and Steiner symmetrization. *Comm. Partial Differential Equations*, 30(10-12):1537–1553, 2005.
- [15] Misha Gromov. *Metric structures for Riemannian and non-Riemannian spaces*, volume 152 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, 1999. Based on the 1981 French original [MR0682063 (85e:53051)], With appendices by M. Katz, P. Pansu and S. Semmes, Translated from the French by Sean Michael Bates.
- [16] Bernhard Kawohl. *Rearrangements and convexity of level sets in PDE*, volume 1150 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1985.
- [17] S. Kesavan. *Symmetrization & applications*, volume 3 of *Series in Analysis*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2006.
- [18] Xiaolong Li and Kui Wang. Moduli of continuity for viscosity solutions on manifolds. *J. Geom. Anal.*, 27(1):557–576, 2017.

- [19] Fogagnolo Mattia and Mazzieri Lorenzo. Minimising hulls, p-capacity and isoperimetric inequality on complete riemannian manifolds. *arXiv:2012.09490*, 2020.
- [20] J. Mossino and J.-M. Rakotoson. Isoperimetric inequalities in parabolic equations. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 13(1):51–73, 1986.
- [21] Lei Ni and Kui Wang. Isoperimetric comparisons via viscosity. *J. Geom. Anal.*, 26(4):2831–2841, 2016.
- [22] Giorgio Talenti. Elliptic equations and rearrangements. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 3(4):697–718, 1976.
- [23] Giorgio Talenti. Nonlinear elliptic equations, rearrangements of functions and Orlicz spaces. *Ann. Mat. Pura Appl. (4)*, 120:160–184, 1979.
- [24] Juan Luis Vázquez. Symétrisation pour $u_1 = \Delta\varphi(u)$ et applications. *C. R. Acad. Sci. Paris Sér. I Math.*, 295(2):71–74, 1982.

SCHOOL OF MATHEMATICAL SCIENCES, SOOCHOW UNIVERSITY, SUZHOU, 215006, CHINA

Email address: `chq4523@163.com`

SCHOOL OF MATHEMATICAL SCIENCES, SOOCHOW UNIVERSITY, SUZHOU, 215006, CHINA

Email address: `1829401184@stu.suda.edu.cn`

SCHOOL OF MATHEMATICAL SCIENCES, SOOCHOW UNIVERSITY, SUZHOU, 215006, CHINA

Email address: `kuiwang@suda.edu.cn`