

## Cyclic and Quasi-Cyclic DNA Codes

Adel Alahmadi · Alaa Altassan · Amani  
Alyoubi · Manish K. Gupta · Hatoon  
Shoab

Received: date / Accepted: date

**Abstract** In this paper, we discuss DNA codes that are cyclic or quasi-cyclic over  $\mathbb{Z}_4 + \omega\mathbb{Z}_4$ , where  $\omega^2 = 2 + 2\omega$  along with methods to construct these with combinatorial constraints. We also generalize results obtained for the ring  $\mathbb{Z}_4 + \omega\mathbb{Z}_4$ , where  $\omega^2 = 2 + 2\omega$ , and some other rings to the sixteen rings  $R_\theta = \mathbb{Z}_4 + \omega\mathbb{Z}_4$ , where  $\omega^2 = \theta \in \mathbb{Z}_4 + \omega\mathbb{Z}_4$ , using the generalized Gau map and Gau distance in [3]. We determine a relationship between the Gau distance and Hamming distance for linear codes over the sixteen rings  $R_\theta$  which enables us to attain an upper boundary for the Gau distance of free codes that are self-dual over the rings  $R_\theta$ .

**Keywords** Cyclic DNA codes · Quasi-cyclic DNA codes · Gau map · Self-dual codes · Codes over  $\mathbb{Z}_4 + \omega\mathbb{Z}_4$  etc

---

A. Alahmadi  
Saudi Arabia  
E-mail: adelnife2@yahoo.com

A. Altassan  
Saudi Arabia  
E-mail: aaltassan@kau.edu.sa

A. Alyoubi  
Saudi Arabia  
E-mail: moon61343@gmail.com

M. Gupta  
India  
E-mail: mankg@computer.org

H. Shoab  
Saudi Arabia  
E-mail: hashoab@kau.edu.sa

## 1 Introduction

The interdisciplinary field of DNA computing has developed into a captivating subject of research during the last twenty-six years. The first successful ingress into DNA computing was performed by Adleman in 1994. DNA is a double-stranded helix made up of chains of four chemicals (nucleotides), Adenine, Cytosine, Guanine and Thymine, which we denote by  $A$ ,  $C$ ,  $G$  and  $T$  respectively. Each of these is paired with another in the other strand, its complementary base (called the Watson-Cricks complement), the two kinds of complements given by  $A^c = T$  and  $C^c = G$ . Thus a DNA string can be represented as a finite sequence of the DNA letters  $A, C, G, T$ . Any set of DNA strings having this set of letters is a DNA code. For a such a string, the complement comes from changing each letter  $X$  to  $X^c$ . Similarly, the reverse string reflects it into reverse order, while the reverse complement is the combination of both actions (in either order). A DNA code is called reversible if it contains the reverse of each of its element. It is called complement if it contains the complement of each of its element. And if the reverse complement of each element of a DNA code belongs to this code, then we call it a reversible complement DNA code. Processing in DNA computing is mostly via DNA hybridization. However, this leads to the main source of errors. To avoid these errors, DNA codes have to satisfy many constraints, especially combinatorial and thermodynamic. Over the past sixteen years many different approaches have been tried to construct these DNA strings: combinatorial, algebraic, computational. In particular, many researchers used algebraic constructions over rings and fields for the DNA codes. However, those methods often gave DNA codes with non-optimal distances. In 2018, Limbachyia et al [1] discovered the Gau isometry map from the ring  $R_{2+2\omega} = \mathbb{Z}_4 + \omega\mathbb{Z}_4$ , where  $\omega^2 = 2 + 2\omega$ , to DNA strings of length two utilizing the Gau distance on codes over  $R_{2+2\omega}$  and produced several DNA codes with better parameters with some optimal ones. In 2019 [3], the Gau isometry map has been extended to cover all the sixteen rings  $R_\theta = \mathbb{Z}_4 + \omega\mathbb{Z}_4$ , where  $\omega^2 = \theta \in \mathbb{Z}_4 + \omega\mathbb{Z}_4$ , and a generalized definition of Gau distance was given for codes over  $R_\theta$ . Cyclic DNA codes have been intensively studied for many element sets such as fields and rings. In particular in [2], cyclic DNA codes were investigated using the Gray map over the ring  $\mathbb{Z}_4 + \omega\mathbb{Z}_4$  where  $\omega^2 = 2$ . It is still an interesting problem to study this type of code using a different map. Besides, quasi-cyclic DNA codes have not been studied substantially.

The ring  $\mathbb{Z}_4 + \omega\mathbb{Z}_4 = \{0, 1, 2, 3, \omega, 1 + \omega, 2 + \omega, 3 + \omega, 2\omega, 1 + 2\omega, 2 + 2\omega, 3 + 2\omega, 3\omega, 1 + 3\omega, 2 + 3\omega, 3 + 3\omega\}$  is an extension of  $\mathbb{Z}_4$ . Using the Gau map defined in [1], we have studied DNA codes that are cyclic or quasi-cyclic over  $\mathbb{Z}_4 + \omega\mathbb{Z}_4$ , where  $\omega^2 = 2 + 2\omega$ , and we have found necessary and sufficient conditions to obtain such codes satisfying the reverse, complement and reverse complement constraints. Furthermore, we have generalized many results obtained for the ring  $R_{2+2\omega} = \mathbb{Z}_4 + \omega\mathbb{Z}_4$  ( $\omega^2 = 2 + 2\omega$ ) and some other rings to all the sixteen rings  $R_\theta$  using the generalized Gau map defined in [3]. In addition, we have improved the maximum possible Hamming distances for self-dual codes that

are free over  $\mathbb{Z}_4 + \omega\mathbb{Z}_4$ , where  $\omega^2 \in \{1, 2\omega\}$  in [4], to the ring  $R_\theta$  and we have found a relationship between the Gau and Hamming distances over the ring  $R_\theta$ , leading us to find, for a self-dual code that is free over  $R_\theta$ , an upper limit to its Gau distance.

This paper is set out as follows: Section 2 gives some examples of cyclic and 2-quasi-cyclic DNA codes over the ring  $R_2 = \mathbb{Z}_4 + \omega\mathbb{Z}_4$  ( $\omega^2 = 2$ ) using the Gau map. In §3 cyclic DNA codes over the ring  $R_{2+2\omega}$  are looked at, along with sufficient conditions for the construction of such codes satisfying the reverse, complement or reverse complement constraints. Similarly, in §4 we find quasi-cyclic codes over the ring  $R_{2+2\omega}$  with sufficient conditions for reverse, complement and reverse complement DNA quasi-cyclic codes. Finally, §5 includes the generalization of results obtained for the ring  $R_{2+2\omega}$  to the general ring  $R_\theta$ , where  $\theta \in \mathbb{Z}_4 + \omega\mathbb{Z}_4$ , and as well contains a generalization of results achieved for self-dual codes over the ring  $\mathbb{Z}_4 + \omega\mathbb{Z}_4$ , where  $\omega^2 = 2\omega$  in [4], to the rings  $R_j = \mathbb{Z}_4 + \omega\mathbb{Z}_4$ , where  $j \in \{2\omega, 2\}$ . Also, it displays a relationship between the Hamming and Gau distances over the ring  $R_\theta$  and upper bounds for the Hamming and Gau distances of free self-dual codes over the ring  $R_\theta$ .

## 2 DNA Codes over $R_2 = \mathbb{Z}_4 + \omega\mathbb{Z}_4$ ( $\omega^2 = 2$ )

In [2] cyclic DNA codes have been studied over the ring  $R_2$  using the Gray image. Here, we shall use the Gau map which defined in [3] from the ring  $R_2$  to the set of DNA strings of length 2, where  $\lambda = 2$ , relating it to the Gau distance to give some examples of cyclic and quasi-cyclic DNA codes. The zero divisors  $Z$  and the units  $U$  of the ring  $R_2$  have been given in [2]. Clearly  $U = Z + 1$ . From basic ring theory,  $(U, \cdot)$  is a multiplicative group, while  $(Z, +)$  is an additive group, and in this case both have order eight. It is unusual that both these abelian groups are isomorphic to  $\mathbb{Z}_4 \times \mathbb{Z}_2$ . Indeed using this isomorphism we find that the multiplicative and additive orders of elements in the ring are related by the formula  $|x + 1| \cdot = |x|_+$ , for all  $x \in Z$ .

The Gau map that we will use has the following properties:

- There are four elements that have additive order 1 or 2: 0, 2,  $2\omega$  and  $2 + 2\omega$ . Their images  $AA$ ,  $TT$ ,  $CC$  and  $GG$  are self-reversible.
- There are four elements which have multiplicative order 1 or 2: 1, 3,  $1 + 2\omega$  and  $3 + 2\omega$ . Their images  $AT$ ,  $TA$ ,  $GC$  and  $CG$  are self-reversible-complement.

### 2.1 Examples

A cyclic code  $\mathcal{C}$  is a linear code such that whenever  $x_0x_1 \dots x_{n-1} \in \mathcal{C}$ , we have also  $x_{n-1}x_0 \dots x_{n-2} \in \mathcal{C}$  while the quasi-cyclic code  $Q$  of index  $l$  is a linear code such that for any  $x_0x_1 \dots x_{n-1} \in Q$ , we have  $x_{n-l}x_{n+1-l} \dots x_0 \dots x_{n-1-l} \in Q$ . We can get the 2-quasi-cyclic DNA code with parameters  $(2n = 8, M =$

32,  $d_H = 4$ ) by shifting the last two coordinates of the vector  $(1 + 3\omega, 1 + \omega, 1 + \omega, 1 + 3\omega)$ . These 32 DNA codewords are as follows:

*AAAAAAAA CCCCCCCC CCGGGGCC CTTCTCCT*  
*TAGCGCTA GCTATAGC ATCGCGAT ACTGTGAC*  
*TTTTTTTT GGGGGGGG TACGCGTA TTAAAATT*  
*ATGCGCAT CGATATCG GCATATGC AATTTTAA*  
*CACACACA ACACACAC GGCCCCGG TCCTCTTC*  
*TCGAGATC GATCTCGA CGTATACG GTCACAGT*  
*GTGTGTGT TGTGTGTG GAAGAGGA AGGAGAAG*  
*AGCTCTAG CTAGAGCT TGACACTG CAGTGTCA*

In similar ways, we have found many examples of cyclic and 2–quasi-cyclic DNA codes given in table 1 and table 2. For the cyclic codes, we will shift only the last coordinate of the vector.

**Table 1** Cyclic DNA Codes over  $R_2$ 

First row vector	$(2n, M, d_H)$
$(2 + \omega, 3, 1)$	$(6, 2048, 1)$
$(3, 1, 2 + 2\omega)$	$(6, 1024, 2)$
$(2 + \omega, 1, 3, 2 + \omega)$	$(8, 4096, 2)$
$(1, 1, 1)$	$(6, 16, 3)$

**Table 2** 2-Quasi-Cyclic DNA Codes over  $R_2$ 

First row vector	$(2n, M, d_H)$
$(2\omega, 1, 1, 2\omega)$	$(8, 256, 2)$
$(1 + 3\omega, 1 + \omega, 1 + \omega, 1 + 3\omega)$	$(8, 32, 4)$
$(3 + \omega, 1, 1, 3 + \omega)$	$(8, 64, 4)$
$(2, 1, \omega, \omega, 1, 2)$	$(12, 4096, 4)$
$(1 + 2\omega, 1, 2, 2\omega, 2\omega, 2, 1, 1 + 2\omega)$	$(16, 4096, 4)$

In both Table 1 and Table 2, all codes represented are reversible and reversible complement. Furthermore, Table 2 includes DNA codes with good parameters.

### 3 Cyclic Codes over $R_{2+2\omega} = \mathbb{Z}_4 + \omega\mathbb{Z}_4$ ( $\omega^2 = 2 + 2\omega$ )

In this section we consider cyclic DNA codes.

**Definition 1** Let  $g(x) = g_0 + g_1x + \dots + g_{n-1}x^{n-1}$  be a polynomial of degree  $n - 1$  with leading coefficient not equal to zero and  $g_i \in R_{2+2\omega}$  for all  $i \in \{0, 1, \dots, n - 1\}$ . Then the reciprocal of  $g$  denoted as  $g^*$  is defined by  $g^*(x) = g_0x^{n-1} + g_1x^{n-2} + \dots + g_{n-1}$ .

A self-reciprocal polynomial  $g(x)$  is one for which  $\exists v \in R_{2+2\omega}$  such that  $g^* = vg$ .

**Definition 2** Let  $x = x_0x_1\dots x_{n-1} \in R_{2+2\omega}^n$  be any string of length  $n$ . The reverse of  $x$  is  $x^r = x_{n-1}x_{n-2}\dots x_0$ . The complement of  $x$  is  $x^c = x_0^c x_1^c \dots x_{n-1}^c$ . The reverse complement of  $x$  is  $x^{rc} = x_{n-1}^c x_{n-2}^c \dots x_0^c$ . Let  $\mathbb{C}$  be any linear code over  $R_{2+2\omega}$ . The code  $\mathbb{C}$  is called reversible if for every  $x \in \mathbb{C}$ , we have  $x^r \in \mathbb{C}$ . The code is called complement if for every  $x \in \mathbb{C}$ , we have  $x^c \in \mathbb{C}$ . The code  $\mathbb{C}$  is called reversible complement if for every  $x \in \mathbb{C}$ , we have  $x^{rc} \in \mathbb{C}$ .

**Proposition 1** For any linear reversible complement code  $\mathbb{C}$  over  $R_{2+2\omega}$ , the constant string  $2 + 2\omega \in \mathbb{C}$ .

**Proof.** Assume that  $\mathbb{C}$  is reversible complement linear code over  $R_{2+2\omega}$ . Since  $\mathbb{C}$  is linear, then  $00\dots 0 \in \mathbb{C}$ .  $\phi(00\dots 0) = AA\dots A$  and  $\phi(00\dots 0)^r =$

$AA \dots A = \phi(00 \dots 0)$ . So,  $\phi^{-1}(\phi(00 \dots 0)^{rc}) = \phi^{-1}(\phi(00 \dots 0)^c) = 2+2\omega \dots 2+2\omega = \mathbf{2} + \mathbf{2}\omega \in \mathbb{C}$ .

**Theorem 2** For a cyclic code  $\mathcal{C}$  over  $R_{2+2\omega}[x]/\langle x^n - 1 \rangle$ ,  $\exists a(x), b(x)$  such that  $b(x)|a(x)|x^n - 1$  and  $\mathcal{C} = \langle a(x), \omega b(x) \rangle$ .

**Proof.** See [7] for the proof.

Using [2, Lemma 4], we obtain the following:

**Lemma 3** Let  $a, b \in R_{2+2\omega}[x]$  and assuming that  $\deg(a(x)) - \deg(b(x)) = t$  then:

- $(a(x)b(x))^* = a^*(x)b^*(x)$ .
- $(a(x) + b(x))^* = a^*(x) + x^t b^*(x)$ .

**Theorem 4** If  $\mathcal{C} = \langle a(x), \omega b(x) \rangle$  is a cyclic code over the ring  $R_{2+2\omega}$ , then a necessary and sufficient condition that  $\mathcal{C}$  is reversible is that  $a$  and  $b$  are self-reciprocal.

**Proof.** ( $\implies$ ) Assume that  $\mathcal{C}$  is reversible.

Let  $a(x) = a_0 + a_1x + \dots + a_sx^s, s \leq n - 1$ . Then if  $a(x) \in \mathcal{C}$ ,

$$a(x)^r = 0 + 0x + \dots + 0x^{n-s-2} + a_sx^{n-s-1} + \dots + a_0x^{n-1} \in \mathcal{C}.$$

Then,  $a_sx^{n-s-1} + \dots + a_0x^{n-1} \in \mathcal{C}$ , which implies,

$$x^{s+1-n}(a_sx^{n-s-1} + \dots + a_0x^{n-1}) \in \mathcal{C}.$$

Then,  $a_s + \dots + a_0x^s \in \mathcal{C}$  which means  $a^*(x) \in \mathcal{C}$ . Hence,

$$a^*(x) = a(x)c(x) + \omega b(x)d(x),$$

where  $c(x), d(x) \in Z_4[x]$ . We have  $a_i \in Z_4$  for  $i \in \{0, 1, \dots, s\}$ , so  $d(x) = 0$ . Then  $a^*(x) = a(x)c(x)$ . Hence,  $\deg(a^*(x)) \geq \deg(a(x))$  but  $\deg(a^*(x)) \leq \deg(a(x))$ , and so  $\deg(a^*(x)) = \deg(a(x))$ . Hence,  $c(x) = m \in Z_4$  where  $m \neq 0$ . Thus,  $a$  must be self-reciprocal.

Similarly, let  $b(x) = b_0 + b_1x + \dots + b_tx^t, t \leq n - 1$  then,

$$\omega b(x) = \omega b_0 + \omega b_1x + \dots + \omega b_tx^t$$

$$(\omega b(x))^r = \omega(b(x))^r = 0 + 0x + \dots + 0x^{n-t-2} + \omega h_tx^{n-t-1} + \dots + \omega h_0x^{n-1} \in \mathcal{C}$$

Then,

$$\omega(b_tx^{n-t-1} + \dots + b_0x^{n-1}) \in \mathcal{C}.$$

Then,

$$\omega x^{t+1-n}(b_tx^{n-t-1} + \dots + b_0x^{n-1}) \in \mathcal{C},$$

which means  $\omega(b_t + \dots + b_0x^t) \in \mathcal{C}$ . Then,  $\omega b^*(x) \in \mathcal{C}$ . Hence,

$$\omega b^*(x) = a(x)c'(x) + \omega b(x)d'(x),$$

where  $c'(x), d'(x) \in Z_4[x]$ . Since every term of  $a(x)$  does not contain  $\omega$ , then  $c'(x) = 0$ . Hence,  $b^*(x) = b(x)d'(x)$  which implies  $\deg(b^*(x)) = \deg(b(x))$ . Then,  $d'(x) = m' \in Z_4$  where  $m' \neq 0$ . Thus,  $b$  must be self-reciprocal.

( $\Leftarrow$ ) Assume that  $a(x)$  and  $b(x)$  are self-reciprocal.

Let  $h(x) = h_0 + h_1x + \cdots + h_u x^u \in \mathcal{C}$ ,  $u \leq n-1$ . Then,

$$h(x) = a(x)e(x) + \omega b(x)f(x),$$

where  $e(x), f(x) \in Z_4[x]$ .

By lemma 3,  $h^*(x) = (a(x)e(x) + \omega b(x)f(x))^* = [a(x)e(x)]^* + x^t[\omega b(x)f(x)]^* = a^*(x)e^*(x) + x^t[\omega h^*(x)f^*(x)] = a^*(x)e^*(x) + \omega b^*(x)(x^t f^*(x)) = ma(x) + \omega(m b(x))(x^t f^*(x))$ .

Hence,  $h^*(x) \in \mathcal{C}$ . So,

$$h^*(x) = h_u + h_{u-1}x + \cdots + h_0 x^u \in \mathcal{C}.$$

Since  $\mathcal{C}$  is cyclic, then  $x^{n-u-1}h^*(x) \in \mathcal{C}$ . Since  $\mathcal{C}$  is linear, then

$$[0 + 0x + \cdots + 0x^{n-1}] + x^{n-u-1}h^*(x) \in \mathcal{C}.$$

Therefore,  $h(x)^r \in \mathcal{C}$ . Thus,  $\mathcal{C}$  must be reversible.

**Lemma 5** For any  $x \in R_{2+2\omega}$ ;  $\omega x = (\omega x)^c - (2 + 2\omega)$ .

**Proof.** It is obvious from Table I in [1].

**Theorem 6** If  $\mathcal{C} = \langle a(x), \omega b(x) \rangle$  is a reversible complement odd length cyclic code over  $R_{2+2\omega}$ , then both  $a$  and  $b$  must be self-reciprocal.

**Proof.** Let  $a(x) = a_0 + a_1x + \cdots + a_s x^s \in \mathcal{C}$ ,  $s \leq n-1$ .

Because  $\mathcal{C}$  is reversible complement, we have

$$a(x)^{rc} = (2+2\omega) + (2+2\omega)x + \cdots + (2+2\omega)x^{n-s-2} + a_s^c x^{n-s-1} + \cdots + a_1^c x^{n-2} + a_0^c x^{n-1} \in \mathcal{C}.$$

Using linearity of  $\mathcal{C}$  and proposition 1, we get

$$(2 + 2\omega)(1 + x + \cdots + x^{n-1}) + a(x)^{rc} \in \mathcal{C}.$$

Then,

$$[(2 + 2\omega) + a_s^c]x^{n-s-1} + [(2 + 2\omega) + a_{s-1}^c]x^{n-s} + \cdots + [(2 + 2\omega) + a_0^c]x^{n-1} \in \mathcal{C}.$$

Since  $(2 + 2\omega) + x^c = x$  for any  $x \in R_{2+2\omega}$ , then

$$x^{n-s-1}[a_s + a_{s-1}x + \cdots + a_0 x^s] \in \mathcal{C}.$$

Then,  $x^{n-s-1}a^*(x) \in \mathcal{C}$  which implies  $a^*(x) \in \mathcal{C}$ .

Since  $\mathcal{C} = \langle a(x), \omega b(x) \rangle$ , then

$$a^*(x) = a(x)c(x) + \omega b(x)d(x)$$

where  $c(x), d(x) \in Z_4[x]$ . We have  $a_i \in Z_4$  for  $i \in \{0, 1, \dots, s\}$ , so  $d(x) = 0$ . Then  $a^*(x) = a(x)c(x)$ . Then,  $\deg(a^*(x)) = \deg(a(x))$ . Hence,  $c(x) = m \in Z_4$

where  $m \neq 0$ . Thus,  $a$  must be self-reciprocal.

Similarly, let  $b(x) = b_0 + b_1x + \dots + b_tx^t$ ,  $t \leq n-1$ . Then,

$$\omega b(x) = \omega b_0 + \omega b_1x + \dots + \omega b_tx^t.$$

$$(\omega b(x))^{rc} = (2+2\omega) + (2+2\omega)x + \dots + (2+2\omega)x^{n-t-2} + (\omega b_t)^c x^{n-t-1} + \dots + (\omega b_0)^c x^{n-1} \in \mathcal{C}.$$

Using linearity of  $\mathcal{C}$  and Proposition 1, we get

$$(\omega b(x))^{rc} - (2+2\omega)(1+x+\dots+x^{n-1}) \in \mathcal{C}$$

Then,

$$[(\omega b_t)^c - (2+2\omega)]x^{n-t-1} + \dots + [(\omega b_1)^c - (2+2\omega)]x^{n-2} + [(\omega b_0)^c - (2+2\omega)]x^{n-1} \in \mathcal{C}.$$

Then,

$$x^{n-t-1}[(\omega b_t)^c - (2+2\omega)] + \dots + ((\omega b_1)^c - (2+2\omega))x^{t-1} + ((\omega b_0)^c - (2+2\omega))x^t \in \mathcal{C}.$$

Using Lemma 5,

$$x^{n-t-1}[\omega b_t + \dots + \omega b_1x^{t-1} + \omega b_0x^t] \in \mathcal{C}.$$

Then,  $x^{n-t-1}\omega b^*(x) \in \mathcal{C}$  which implies  $\omega b^*(x) \in \mathcal{C}$ . So,

$$\omega b^*(x) = g(x)c'(x) + \omega b(x)d'(x)$$

where  $c'(x), d'(x) \in Z_4[x]$ . Since every term of  $a(x)$  does not contain  $\omega$ , then  $c'(x) = 0$ . Hence,  $b^*(x) = b(x)d'(x)$ , which implies  $\deg(b^*(x)) = \deg(b(x))$  then,  $d'(x) = m' \in Z_4$ , where  $m' \neq 0$ . Thus,  $b(x)$  is self-reciprocal.

We found the sufficient conditions for showing the converse of Theorem 6.

**Theorem 7** *If  $\mathcal{C} = \langle a(x), \omega b(x) \rangle$  is a cyclic code of odd length over the ring  $R_{2+2\omega}$ , and if  $(2+2\omega)(1+x+\dots+x^{n-1}) \in \mathcal{C}$  and  $a$  and  $b$  are self reciprocal, then  $\mathcal{C}$  must be reversible complement.*

**Proof.** Let  $h(x) \in \mathcal{C}$ . We want to show that  $h(x)^{rc} \in \mathcal{C}$ .

Let  $h(x) = h_0 + h_1x + \dots + h_tx^t \in \mathcal{C}$ ,  $t \leq n-1$ . Since  $h(x) \in \mathcal{C}$ , then

$$h(x) = a(x)c(x) + \omega b(x)d(x).$$

Then,

$$\begin{aligned} h^*(x) &= (a(x)c(x) + \omega b(x)d(x))^* = [a(x)c(x)]^* + x^s[\omega b(x)d(x)]^* \\ &= a^*(x)c^*(x) + x^s[\omega b^*(x)d^*(x)]. \end{aligned}$$

Since  $a^*(x) = ma(x)$  and  $b^*(x) = m'b(x)$ , then

$$h^*(x) = ma(x)c^*(x) + \omega b(x)[m'x^s d^*(x)].$$



Hence,  $h^*(x) \in \mathcal{C}$ .

Since  $\mathcal{C}$  is cyclic, then

$$x^{n-t-1}h(x) = h_0x^{n-t-1} + h_1x^{n-t} + \cdots + h_tx^{n-1} \in \mathcal{C}.$$

Since  $(2 + 2\omega)(1 + x + \cdots + x^{n-1}) \in \mathcal{C}$  and  $\mathcal{C}$  is linear, then

$$(2 + 2\omega)(1 + x + \cdots + x^{n-1}) + x^{n-t-1}h(x) \in \mathcal{C}.$$

Then,

$$(2+2\omega) + (2+2\omega)x + \cdots + (2+2\omega)x^{n-1} + [h_0x^{n-t-1} + h_1x^{n-t} + \cdots + h_tx^{n-1}] \in \mathcal{C},$$

which implies  $(2+2\omega) + (2+2\omega)x + \cdots + (2+2\omega)x^{n-t-2} + [(2+2\omega) + h_0]x^{n-t-1} + [(2+2\omega) + h_1]x^{n-t} + \cdots + [(2+2\omega) + h_t]x^{n-1} \in \mathcal{C}$ . Since,  $(2+2\omega) + x = x^c$ , then

$$(2+2\omega) + (2+2\omega)x + \cdots + (2+2\omega)x^{n-t-2} + h_0^c x^{n-t-1} + h_1^c x^{n-t} + \cdots + h_t^c x^{n-1} \in \mathcal{C}.$$

Hence,  $(h^*(x))^{rc} \in \mathcal{C}$ . We have  $h^*(x) \in \mathcal{C}$  for any polynomial  $h(x) \in \mathcal{C}$ . Then,

$$((h^*(x))^{rc})^* \in \mathcal{C}.$$

$$((h^*(x))^{rc})^* = h_t^c + h_{t-1}^c x + \cdots + h_0^c x^t + (2+2\omega)x^{t+1} + \cdots + (2+2\omega)x^{n-1} \in \mathcal{C}.$$

Note that  $((h^*(x))^{rc})^* = (x^{n-t-1}h(x))^{rc} \in \mathcal{C} \implies h(x)^{rc} \in \mathcal{C}$ . Thus,  $\mathcal{C}$  is reversible complement.

Using [1, Lemmas 2 and 3], we obtain the following:

**Lemma 8** *A necessary and sufficient condition that the DNA cyclic code  $\mathcal{C}$  over  $R_{2+2\omega}$  be reversible is that  $x^s \phi^{-1}(\phi(g(x)^r)) \in \mathcal{C}$  for all  $s$  where  $s \in \{0, 1, \dots, k-1\}$  and  $g(x)$  is the first row in the generator matrix of  $\mathcal{C}$ .*

**Lemma 9** *The DNA cyclic code  $\mathcal{C}$  over  $R_{2+2\omega}$  is complementary if  $(2 + 2\omega)(1 + x + \cdots + x^{n-1}) \in \mathcal{C}$  and conversely.*

Combining Lemmas 8 and 9 we obtain the following;

**Proposition 10** *A necessary and sufficient condition that the DNA cyclic code  $\mathcal{C}$  over  $R_{2+2\omega}$  be reversible complement is that  $x^s \phi^{-1}(\phi(g(x)^r)) \in \mathcal{C}$  and  $(2 + 2\omega)(1 + x + \cdots + x^{n-1}) \in \mathcal{C}$*

#### 4 Quasi-Cyclic Codes over $R_{2+2\omega}$

Every  $l$ -quasi-cyclic code has a generator matrix

$$G = \begin{bmatrix} g(x) \\ x^l g(x) \\ x^{2l} g(x) \\ \vdots \\ x^{(k-1)l} g(x) \end{bmatrix}$$

A quasi-cyclic code has length  $n = l \cdot m$ , where  $l$  is the index and  $m$  is the co-index, and the number of rows  $k \leq m$ .

Similarly, using [1, Lemmas 2 and 3], we obtain the following:

**Lemma 11** *A necessary and sufficient condition that the DNA quasi-cyclic code over  $R_{2+2\omega}$  be reversible is that  $x^{sl}\phi^{-1}(\phi(g(x)^r)) \in Q$  for all  $s$ , where  $s \in \{0, 1, \dots, k-1\}$  and  $g(x)$  is any row in the generator matrix of  $Q$ .*

**Lemma 12** *The DNA quasi-cyclic code  $Q$  over  $R_{2+2\omega}$  is complement if and only if  $(2+2\omega)(1+x+\dots+x^{n-1}) \in Q$ .*

From Lemmas 11 and 12 this holds:

**Proposition 13** *A DNA code which is quasi-cyclic over  $R_{2+2\omega}$  has all three reversible complement, reversible and complement conditions if and only if  $x^{sl}\phi^{-1}(\phi(g(x)^r)) \in Q$  and  $(2+2\omega)(1+x+\dots+x^{n-1}) \in Q$ .*

The following conjecture presents sufficient conditions to obtain reversible and reversible complement DNA quasi-cyclic codes.

*Conjecture 1* A DNA code which is quasi-cyclic with length  $2n = 2(l \cdot m)$  where  $l > 1$  is reversible, also reversible complement if and only if  $l \leq m$  and there exists a row  $g(x) = \sum_{i=0}^{n-1} g_i x^i$  in the generator matrix  $G$  such that  $g_i = g_{n-i-1}$  for all  $i \in \{0, 1, \dots, n-1\}$ .

Note that in the above conjecture, it is sufficient to know any matrix that generates the quasi-cyclic code to determine whether the corresponding DNA code meets or does not meet the combinatorial constraints.

**Example 14** *We can get the 2-quasi-cyclic DNA code with parameters  $(2n = 16, M = 256, d_H = 8)$  by shifting the last two coordinates of the vector  $(0, 3, 1, 2, 2, 1, 3, 0)$ .*

These 256 DNA codewords are as follows:

AAAAAAAAAAAAAAAAAA AGGGGGAGGAAAAAGA GGAAAAGGGGAAAAGG CTAAACTTCGGGGTC  
 GAGGGGGAAGAAAAAG TGCCCTGGTAAAAGT TATTTTAAATAAAAAAT TTGGGGTTTTGGGGTT  
 CACCCCAACAAAAAC CGTTTTCGGCAAAAGC CCAAAAACCCAAAAACC TCAAAATCCTGGGGCT  
 CTGGGGCTTCAAAATC TTA AAAATTTTAAAATT TCGGGGTCTAAAAC TTTTGTGGGGGTG  
 GTCCCGTTGAAAATG GCTTTTGCCGAAAACG ATTTTATTA AAAAATA GCCCCCGCCGGGGGGC  
 GGAGGAAAAAGAAAGG GAGAAGAGGAGAAGAG AAAGGAGGGGGAAGAA ACTTTTACCAGGGCA  
 AGGAAGGAAGGAAGGA CACTTCTGGTGAAGAC CGTCTTAATGAAGGC ATCCCTATTAGGGTA  
 TGCTTCAACGAAGGT TATCCTCGGCGAAGAT TTAGGACCCCGAAGTT GGGAAAGAAAAAGGAGG  
 TCGAAGCTTCGAAGCT GAAAAAGAAGGGGGAG TGTTTTTGGTGGGGCT GAAGGAAGGAAGGAAG  
 CTGAAGTCTGAAGTC ACCTTCGTTGGAAGCA ATTCTGCCGGAAGTA AAGAAGGGGGAGGAAA  
 GTCTTACCAGAAGTG GCTCCTATTAGAAGCG AAGGGGAAAAAGGGGAA AGAGGAGAAGAGGAGA  
 AGAAAAAGGAGGGGGA GGGGGGGGGGGGGGG CCGGGGGCCCGGGGCC CATCCTTGGTAGGAAC  
 TACCCCTAATGGGGAT CATTTC AACGGGGAC CGCCCCCGGGGGGGC CGCTTCTAATAGGAGC  
 TACTTCCGGCAGGAAT TTGAAGCCCCAGGATT TCAGGACTTCAGGACT TGTCTCAACAGGAGT  
 CCGAAGTTTTAGGACC CTAGGATCCTAGGATC ACTCCTGTTGAGGACA ATCTTCCCGAGGATA  
 GTTCTACCAAGGATG GCCTTCAATTAAGGACG CCTGGTAAAAGTTGCC CTC AACAGGAGTTGTC  
 TTTGGTGGGGGTTGTT TCCAACGAAGGTTGCT GTGTTGTGGTGTGTTG GCACCAT AATGTTGCG  
 ACCTTGC AACGTTGCA ATACCACGGC GTTGTGA GACA ACTCCTGTTGAG ATGGTCCCGTTGAA  
 GGTGTTTTTGTGGG GACTTCTGTTGAG TGGTTGGTGGTTGGT TAACAGCCGGTTGAT  
 CAGTTGACCAGTTGAC CGACCAATTAGTTGGC TTTAATAAAAAATTATT TCCGGCAGGAATTACT  
 CCTAATGGGGATTACC CTGCGGAAGATTATC ATACCACGGC GTTGTGA ATATTATAATATTAT  
 GTGCCGCAACATTATG GCATTACGGCATTACG GGTAATCCCCATTAGG GACGGCCTTCATTAAG  
 AATAATTTTTATTAAA AGCGGCTCCTATTAGA CAGCCGGTTGATTAA CCGATTAGCCGATTAGC  
 TGGCCGACCAATTAGT TAATTAATTAATTAAT CCAACAAAAAACACC ACCCTTGGTAGGAACCATC  
 TTCAACGGGGACCATT TCTGGTGAAGACC ACT GTACCATGGTACCATG GCGTTGT AATACCACG  
 ACACCACAACACCACA ATGTTGCGGCACCATA AACACCCCCACC AAA AGTGGTCTTCACCAGA  
 GGC AACTTTTACCAGG GATGGTCTTACCAAG TTGACAGTTGACCAGT TAGTTGGCCATCCAAAT  
 CAACCAACCAACCAAC CGGTTGATTAACCAGC TTCGGCAAAAAGCCGTT TCTAATAGGAGCCGCT  
 CCCGGCGGGGGCCGCG CTTAATGAAGGCCGTC ACATTATGGTGGCCGA ATGCCGTAATGCCGTA  
 GTATTACAACGTTGCA GCGCGCGCGCGCGCGG GCGGGCCCCCGCCGGG GATAAATCTTCGGCAG  
 AACGGCTTTTACCGAA AGTAATTCCTGCCGGA CAATTAGTTGGCCGAC CCGCCGGCCGGCCGGC  
 TGATTAACGACCCGTT TAGCCGATTAGCCGTA AACCCAAAAACCCCAA AGTTTTAGGACCCCGA  
 GGCCCCGGGGCCCCGG GATTTTGAAGCCCCAG TGA AAAATGGTCCCCGT TAGGGGT AATCCCCAT  
 CAAAAAC AACCCCCAC TCGGGGGCGCCCCCGC CCCCCCCCCCCCCCCC CTTTTTCTTCCCCCTC  
 TTCCCTTTTCCCTT ACAAAAACCCACCCCA GTTTTTCCTCCCTTGT GTAAAAGTTGCCCCCTG  
 AACTTCGGGGTCTTAA ATG GGGGATTACCCCTA ATGGGTTAAGCCCCAG GCGGGGGCGCCCCCG  
 TGAGGACAACCTCTGT TAGCCTGAAGTCTGTA TACCTTCCCTCCCTT GATAAATCTTCAG  
 CCCTTCTTTTCTCTC CTTCCCTTCCCTCCCTC CTTCCCTCCCTCCCTT TTCTCCCTCTCTCT  
 GTAGGAACCATCCTTG GCGAAGATTATCCTCG GCAAGGATGGTCTTAC CAAGGATGGTCTTAC  
 CAGTTTGGGTTTGG GACCCCGAAGTTTTAG TTCTTCCCTCCCTTT ACAGGAGTTGTCTTCA  
 CAGGGCAACTTTTAC CGAAAAACGGCTTTTTC CTTTTCCTTCCCTTCC AATTTTAAAATTTTAA  
 TTTT TTTT TTTT TTTT TTTT TTTT TTTT TTTT TTTT TTTT TTTT TTTT TTTT TTTT TTTT  
 CAGGGGACCATTTTCA ATAAAAATTATTTTAA TG GGGGTGGTTTTTGT TAAAAATAATTTTTAT  
 AATCCTGGGGCTTCAA AGCTTCAAGCTTTCGA TTTTCTCCCTTCTT CTTCCCTTCTTTTTC  
 TGGAAGCAACCTTCGT TAAGGACGGCCTTCA CTCTTCTCTCTTCTC CCGTTGAAAATGGTCC  
 CCTCCTTTTCTTCCC GCAGGAATTACTTCCG TTTTCTCCCTTCTT TTTTCTCCCTTCTT  
 GTGAAGACC ACTTCTG TCACCAAGAGTGGTCT AAGTTGCCCCTGGTAA AAGTTGCCCCTGGTAA  
 ACTGGTCAACTGGTCA GGGTTGTTTTTGGTGG GAACCATCCTTGGTAG TGTGGTGTGTTGGTGT  
 CCAGGATTTTGAAGCC CATGGTACCATGGTAC CGCAGGCTGTTGTTG CCGCAACATTATGGTGC  
 TACAACCGCGTGGTAT ATCAACCGGCTGGTTA AAGTTGCCCCTGGTAA AAGTTGCCCCTGGTAA  
 CCGCGCGCGCGCGCGC GTTAATCAACCGGCTG GCGCGCGCGCGCGCGC GCGCGCGCGCGCGCGC  
 GAATTACTTCCGGCAG AAGCCGTTTTTCGGCAA AGATTATCCTCGGCGA ATATTAGAAGCGGCTC  
 CGCGCGCGCGCGCGCGC TGTAATACCACGGCGT TACGGCATTACGGCAT TCGTTGGAAGCAACCT  
 CTGTTGAGGACAACCT TTACCAGGGGCAACTT TCGTTGGAAGCAACCT TCGTTGGAAGCAACCT  
 GCTGGTTAATCAACCG GACCAACCAACCAACCA AAGTTGCCCCTGGTAA AAGTTGCCCCTGGTAA  
 AGTTGCTTCCAACGA GGACCATTTTCAACCG GAGTTGCCCCTGGTAA AAGTTGCCCCTGGTAA  
 TATGGTGGCGCAACAT CACAACACCACAACAC CGTGGTATTACAACGC CGTGGTATTACAACGC  
 TCGCCGAGGATAATCT CCATTAGGGGTAATTC CTGCCGGAAGTAATTC CTGCCGGAAGTAATTC  
 ATTAATTAATTAATTA GTCGGCAACTAATTC GCTAATCGGCTAATTC GCTAATCGGCTAATTC  
 GAGCCGCTTCTAATAG AAATTATTTTTAATAA AGGCCGCTTCTAATGA AGGCCGCTTCTAATGA  
 CGTAATGCCGTAATGC TGCGGCACCATAATGT ACCCCACC AAAAAC A TATAATATTAT AATAT

**Table 3** Quasi-Cyclic DNA Code of index 2

First row vector	$(2n, M, d_H)$
$(1, 2, 2, 1)$	$(8, 256, 2)$
$(1, 1, 2\omega + 3, 2, 2, 2\omega + 3)$	$(12, 1024, 4)$
$(\omega, 1, 1, \omega, 3, 3)$	$(12, 2048, 4)$
$(0, 3, 1, 2, 2, 1, 3, 0)$	$(16, 256, 8)$

**Table 4** Quasi-Cyclic DNA Code of index 3

First row vector	$(2n, M, d_H)$
$(3, 3, 3, 1, 1, 1, 1, 1, 1)$	$(18, 256, 6)$
$(3 + \omega, 3 + \omega, 3 + \omega, 1, 1, 1, 1, 1, 3 + \omega, 3 + \omega, 3 + \omega, )$	$(24, 512, 6)$

In similar ways, we have found many examples of quasi-cyclic DNA codes of indices 2 and 3 which are reversible and reversible complement given in table 3 and table 4. For the quasi-cyclic codes of index 3, we will shift the last 3 coordinates of the vector. newpage

Using the Sphere Packing Bound defined in [3], we found that Table 3 and Table 4 contain good codes.

## 5 General Results

The structure of the sixteen rings  $R_\theta = \mathbb{Z}_4 + \omega\mathbb{Z}_4$  ( $\omega^2 = \theta$ ) where  $\theta \in \mathbb{Z}_4 + \omega\mathbb{Z}_4$  has been studied in [3]. We found in addition, the following general properties:

- There are special zero divisors which are self-invertible under addition;  $0, 2, 2\omega$  and  $2 + 2\omega$  will be the same in any ring  $R_\theta$ .
- There are special units,  $1, 3, 1 + 2\omega$  and  $3 + 2\omega$ , which are self-invertible under multiplication, that behave the same for any ring  $R_\theta$ .
- The ring  $R_\theta$  is a Frobenius ring for  $\theta \in \{0, 1, 2, 3, \omega, 1 + \omega, 2 + \omega, 3 + \omega, 2\omega, 1 + 2\omega, 2 + 2\omega, 3\omega, 1 + 3\omega, 2 + 3\omega, 3 + 3\omega\}$ .

We use the generalized Gau map which has been defined in [3] to generalize some of the results to the ring  $R_\theta$ .

**Proposition 15** *For any reversible complement linear code  $\mathbb{C}$  over the ring  $R_\theta$ , the string  $\lambda \in \mathbb{C}$  where  $\lambda$  is a string with each coordinte  $\lambda$  where  $\lambda \in \{2, 2\omega, 2 + 2\omega\}$ .*

**Theorem 16** *Suppose that  $\mathcal{C}$  is cyclic in  $R_\theta[x]/\langle x^n - 1 \rangle$ . Then there exists polynomials  $a, b$  in  $R_\theta[x]/\langle x^n - 1 \rangle$  such that  $b(x)|a(x)|x^n - 1$  and  $\mathcal{C} = \langle a(x), \omega b(x) \rangle$ .*

**Lemma 17** *Let  $a, b$  be two polynomials in  $R_\theta[x]$  with the property that  $\deg(a(x)) - \deg(b(x)) = t$ , then:*

- $(a(x)b(x))^* = a^*(x)b^*(x)$ .
- $(a(x) + b(x))^* = a^*(x) + x^t b^*(x)$ .

**Theorem 18** Let  $\mathcal{C} = \langle a(x), \omega b(x) \rangle$  be a cyclic code over the ring  $R_\theta$ . It follows that the necessary and sufficient condition for  $\mathcal{C}$  to be reversible is that both  $a$  and  $b$  be self-reciprocal.

**Lemma 19** For any  $x \in R_\theta$ ;  $\omega x = (\omega x)^c - (\lambda)$  where  $\lambda \in \{2, 2\omega, 2 + 2\omega\}$ .

**Theorem 20** If  $\mathcal{C} = \langle a(x), \omega b(x) \rangle$  is a reversible complement cyclic code over  $R_\theta$  having odd length, then both  $a$  and  $b$  are self-reciprocal.

**Theorem 21** Let  $\mathcal{C} = \langle a(x), \omega b(x) \rangle$  be a cyclic code of odd length over the ring  $R_\theta$ . If  $\lambda(1 + x + \dots + x^{n-1}) \in \mathcal{C}$  where  $\lambda \in \{2, 2\omega, 2 + 2\omega\}$  and  $a(x)$  and  $b(x)$  are self reciprocal, then  $\mathcal{C}$  is reversible complement.

Using [3, Lemma 10 and Remark 12], we obtain:

**Lemma 22** A necessary and sufficient condition that a DNA cyclic code  $\mathcal{C}$  over  $R_\theta$  be reversible is that  $x^s \phi^{-1}(\phi(g(x)^r)) \in \mathcal{C}$  for all  $s$ , where  $s \in \{0, 1, \dots, k-1\}$  and  $g(x)$  is the first row in a generator matrix of  $\mathcal{C}$ .

**Lemma 23** The DNA cyclic code  $\mathcal{C}$  over  $R_\theta$  is complement if and only if  $\lambda(1 + x + \dots + x^{n-1}) \in \mathcal{C}$  where  $\lambda \in \{2, 2\omega, 2 + 2\omega\}$ .

Combining Lemmas 22 and 23, we obtain the following;

**Proposition 24** The DNA cyclic code  $\mathcal{C}$  over the ring  $R_\theta$  is reversible complement, reversible and complement if and only if  $x^s \phi^{-1}(\phi(g(x)^r)) \in \mathcal{C}$  and  $\lambda(1 + x + \dots + x^{n-1}) \in \mathcal{C}$  where  $\lambda \in \{2, 2\omega, 2 + 2\omega\}$ .

Similarly, using [3, Lemma 10 and Remark 12], we obtain the following:

**Lemma 25** The DNA quasi-cyclic code  $Q$  over  $R_\theta$  is reversible if and only if  $x^{sl} \phi^{-1}(\phi(g(x)^r)) \in Q$  for all  $s$ , where  $s \in \{0, 1, \dots, k-1\}$  and  $g(x)$  is any row in the generator matrix of  $Q$ .

**Lemma 26** The DNA quasi-cyclic code  $Q$  over  $R_\theta$  is complement if and only if  $\lambda(1 + x + \dots + x^{n-1}) \in Q$  where  $\lambda \in \{2, 2\omega, 2 + 2\omega\}$ .

Combining Lemmas 25 and 26 we obtain:

**Proposition 27** The DNA quasi-cyclic code  $Q$  over the ring  $R_\theta$  is reversible complement, reversible and complement if and only if  $x^{sl} \phi^{-1}(\phi(g(x)^r)) \in Q$  and  $\lambda(1 + x + \dots + x^{n-1}) \in Q$  where  $\lambda \in \{2, 2\omega, 2 + 2\omega\}$ .

Self-dual codes over the ring  $R_i = \mathbb{Z}_4 + \omega \mathbb{Z}_4$ , where  $\omega^2 = i$  for  $i \in \{1, 2\omega\}$ , have been studied in [4]. We generalize some results obtained in that paper using the Gray map and the generalized Gau map.

In this paper, we consider the usual inner product regarding to the duality concepts.

**Definition 3** The dual code  $\mathbb{C}^\perp$  of a code  $\mathbb{C}$  over a ring  $R$  is defined as  $\mathbb{C}^\perp = \{c \in R^n \mid c \cdot c' = 0, \forall c' \in \mathbb{C}\}$ , where  $c \cdot c'$  denotes the usual inner product

**Definition 4** A code  $\mathbb{C}$  over a ring is called self-dual if  $\mathbb{C} = \mathbb{C}^\perp$ .

The Gray map  $\psi_i$  is defined in [4] for the ring  $R_i$  where  $i \in \{1, 2\omega\}$ . We extend the definition of Gray map for the ring  $R_\theta$ , so the Gray map  $\psi_\theta : R_\theta \rightarrow \mathbb{Z}_4^2$  is defined as:  $\psi_\theta(a + \omega b) = (b, a + b)$ . This is extended component wise to  $\psi_\theta : R_\theta^n \rightarrow \mathbb{Z}_4^{2n}$ .

**Theorem 28** Let  $\mathbb{C}$  be a linear code over the ring  $R_j$  where  $j \in \{2, 2\omega\}$ , Then  $\psi_j(\mathbb{C}^\perp) = \psi_j(\mathbb{C})^\perp$ .

**Proof.** The proof of this theorem for  $j = 2\omega$  is given in [4, Theorem 2.1]. Consider  $j = 2$  and assume that  $\phi(c) \in \phi(\mathbb{C}^\perp)$ , where  $c = a + \omega b = (a_0, a_1, \dots, a_{n-1}) + \omega(b_0, b_1, \dots, b_{n-1}) \in \mathbb{C}^\perp$ , it follows that  $c \cdot c' = 0$  for all  $c' = a' + \omega b' \in \mathbb{C}$  which implies  $a \cdot a' + 2(b \cdot b') = 0$  and  $a \cdot b' + b \cdot a' = 0$ . Then  $\psi_2(c) \cdot \psi_2(c') = (b, a + b) \cdot (b', a' + b') = a \cdot a' + 2(b \cdot b') + (a \cdot b' + a' \cdot b) = 0$ . Therefore,  $\psi_2(\mathbb{C}^\perp) \subseteq \psi_2(\mathbb{C})^\perp$ .

The second part of the proof is by similar technique as proving [4, Theorem 2.1].

**Corollary 29** Whenever  $\mathbb{C}$  is self-dual code over the ring  $R_j$  where  $j \in \{2, 2\omega\}$ , Then  $\psi_j(\mathbb{C})$  must be also self-dual over  $\mathbb{Z}_4$

**Proof.** Let  $\mathbb{C} = \mathbb{C}^\perp$ , then  $\psi_j(\mathbb{C}) = \psi_j(\mathbb{C}^\perp)$  and  $\psi_j(\mathbb{C}^\perp) = \psi_j(\mathbb{C})^\perp$ .

Now we consider the generalized Gau map  $\phi$  for the following results.

**Definition 5** The dual code of a DNA code  $\phi(\mathbb{C})$  where  $\mathbb{C}$  is a code over  $R_\theta$  is defined as  $\phi(\mathbb{C})^\perp = \{\phi(c) \mid c \cdot c' = 0, \forall c' \in \mathbb{C}\}$ .

We define the self-dual DNA code for a DNA strings of even length

**Definition 6** A DNA code  $\phi(\mathbb{C})$  is self-dual if  $\phi(\mathbb{C}) = \phi(\mathbb{C})^\perp$

one can easily show the following remark which is clear by definition 5

**Remark 30**  $\phi(\mathbb{C})^\perp = \phi(\mathbb{C}^\perp)$  for all linear codes  $\mathbb{C}$  over the ring  $R_\theta$ .

**Corollary 31**  $\mathbb{C}$  is a self-dual code over the ring  $R_\theta$  if and only if  $\phi(\mathbb{C})$  is self-dual.

**Proof.**  $\mathbb{C} = \mathbb{C}^\perp \Leftrightarrow \phi(\mathbb{C}) = \phi(\mathbb{C}^\perp) = \phi(\mathbb{C})^\perp$ .

**Definition 7** A self-dual code that is free is just one with no non-zero code-word made up of only zero divisors.

**Example 32** Let  $\mathbb{C}$  be a linear code over the ring  $R_2$  generated by

$$G = \begin{bmatrix} 1 & 0 & 0 & 0 & 1 & 1 + 2\omega & 1 + 2\omega & 0 \\ 0 & 1 & 0 & 0 & 3 + 2\omega & 1 & 0 & 1 + 2\omega \\ 0 & 0 & 1 & 0 & 1 + 2\omega & 0 & 3 & 1 + 2\omega \\ 0 & 0 & 0 & 1 & 0 & 3 + 2\omega & 1 + 2\omega & 1 \end{bmatrix}$$

This is an  $(8, 16^4)$ -free self-dual code over  $R_2$ .

**Theorem 33**  $\frac{n}{2} + 1$  is the largest possible value of minimum Hamming distance of a free self-dual code of length  $n$  over  $R_\theta$ .

**Proof.** The proof has the same technique [4, Theorem 3.8].

We have also found the upper bound of the Gau distance  $d_{G(\theta)}$  of a free self-dual code over  $R_\theta$  using the following relation between the Gau and Hamming distances.

**Lemma 34**  $d_{G(\theta)}(x, y) = d_H(x, y) + r$  for all  $x, y \in \mathbb{C}$ , where  $\mathbb{C}$  is a code over the ring  $R_\theta$  and  $r$  is how many coordinates  $i$  with  $d_{G(\theta)}(x_i, y_i) = 2$ , where  $i \in \{0, \dots, n-1\}$ .

**Proof.** Let  $x, y$  be arbitrary codewords in  $\mathbb{C}$  of length  $n$ ,  
 $x = x_0 \dots x_{n-1}$  and  $y = y_0 \dots y_{n-1}$ .

Case 1: If  $d_{G(\theta)}(x_i, y_i) \neq 2$  for all  $i \in \{0, 1, \dots, n-1\}$ ,  
then,  $r = 0$  and  $d_H(x_i, y_i) = d_{G(\theta)}(x_i, y_i)$  for all  $i \in \{0, 1, \dots, n-1\}$ .

$$d_{G(\theta)}(x, y) = \sum_{i=0}^{n-1} d_{G(\theta)}(x_i, y_i) = \sum_{i=0}^{n-1} d_H(x_i, y_i).$$

Thus,  $d_{G(\theta)}(x, y) = d_H(x, y) + r$ .

Case 2: If there exists  $i \in \{0, 1, \dots, n-1\}$  such that  $d_{G(\theta)}(x_i, y_i) = 2$ .

Let

$$S = \{i : d_{G(\theta)}(x_i, y_i) \neq 2\}$$

and let

$$T = \{j : d_{G(\theta)}(x_j, y_j) = 2\}.$$

Hence,  $d_H(x_i, y_i) = d_{G(\theta)}(x_i, y_i)$  for all  $i \in S$  and  $d_H(x_j, y_j) = 1$  for all  $j \in T$ .

$$d_{G(\theta)}(x, y) = \sum_{i \in S} d_{G(\theta)}(x_i, y_i) + \sum_{j \in T} d_{G(\theta)}(x_j, y_j), \dots\dots (*)$$

where

$$\sum_{i \in S} d_{G(\theta)}(x_i, y_i) = \sum_{i \in S} d_H(x_i, y_i),$$

and

$$\sum_{j \in T} d_{G(\theta)}(x_j, y_j) = 2 \sum_{j \in T} d_H(x_j, y_j) = \sum_{j \in T} d_H(x_j, y_j) + r.$$

By substituting in (\*), we obtain

$$d_{G(\theta)}(x, y) = \sum_{i \in S} d_H(x_i, y_i) + \sum_{j \in T} d_H(x_j, y_j) + r.$$

Thus,  $d_{G(\theta)}(x, y) = d_H(x, y) + r$ .

**Theorem 35**  $\frac{n}{2} + r + 1$  is an upper bound for the Gau distance of a free self-dual code  $\mathbb{C}$  over  $R_\theta$ .

**Proof.** If  $\mathbb{C}$  is a code over  $R_\theta$  and  $x$  and  $y$  are arbitrary codewords in  $\mathbb{C}$ , then by Theorem 33,  $d_H(x, y) \leq \frac{n}{2} + 1$  and by Lemma 34,  $d_{G(\theta)}(x, y) = d_H(x, y) + r \leq \frac{n}{2} + 1 + r$ . Thus,  $d_{G(\theta)}(x, y) \leq \frac{n}{2} + r + 1$ .

## 6 Conclusion

This paper investigated cyclic and quasi-cyclic DNA codes. Conditions for DNA codes that are cyclic or quasi-cyclic have having certain combinatorial constraints have been explored. Many examples of good DNA quasi-cyclic codes have been provided. it would be nice to study twisted and quasi-twisted DNA codes over  $R_\theta$ . Moreover, one can ascertain whether Conjecture 1 is true.

## References

1. D. Limbachiya, K. G. Benerjee, B. Rao, and M. K. Gupta, On DNA codes using the ring  $\mathbb{Z}_4 + \omega\mathbb{Z}_4$ , IEEE Int. Symp. Inf. Theory-Proc., pp 2401-2405 (2018)
2. A. Dertli, Y. Cengellenmis, and O. Mayis, On the cyclic DNA codes over the finite rings  $\mathbb{Z}_4 + \omega\mathbb{Z}_4$  and  $\mathbb{Z}_4 + \omega\mathbb{Z}_4 + v\mathbb{Z}_4 + \omega v\mathbb{Z}_4$ , BIOMATH, vol. 6, no.2 (2017)
3. A. Alahmadi, K. G. Benerjee, S. Deb, and M. K. Gupta, DNA codes over the ring  $\mathbb{Z}_4 + \omega\mathbb{Z}_4$ , submitted (2019) Available at <https://arxiv.org/abs/2110.09089>
4. R. Bandi, and M. Bhaintwal, self-dual codes over  $\mathbb{Z}_4 + \omega\mathbb{Z}_4$ , Discret.Math.Algorithms Appl., vol. 7, no.2, pp.1-10 (2015)
5. J. Watada, DNA computing and its application, Studies in Computational Intelligence, pp.1065-1089, Springer, Berlin (2008)
6. J. Li, Q. Zhang, R. Li, and S. Zhou, Optimization of DNA encoding based on combinatorial constraints, ICIC Express Letters, vol.2, no.1, pp 81-88, (2008)
7. N. Aydin, A. Dertli, and Y. Cengellenmis, Cyclic and constacyclic codes over  $\mathbb{Z}_4 + \omega\mathbb{Z}_4$ , preprint.
8. D. Limbachiya, B. Rao, and M. K. Gupta, The art of DNA strings: Sixteen years of DNA coding theory, arXiv:1607.00266, pp.1-19 (2016)
9. M. Bhaintwal, and S.K. Wasan, On quasi-cyclic codes over  $\mathbb{Z}_q$ , AAECC, vol.20, 459-480 (2009)
10. H. Q. Dinh, A. K. Singh, S. Pattanayak and S. Sriboonchitta, Construction of cyclic DNA codes over the ring  $\mathbb{Z}_4[u]/\langle u^2 - 1 \rangle$  based on the deletion distance, Theoretical Computer Science, vol. 773, pp 27-42 (2019)
11. S. Ling and P. Solé, On the Algebraic Structure of Quasi-cyclic Codes II:Chain Rings, Designs, Codes and Cryptography, vol. 30, pp. 113-130 (2003)
12. A. Dertli, and Y. Cengellenmis, On the Codes Over the Rings  $\mathbb{Z}_4 + u\mathbb{Z}_4 + v\mathbb{Z}_4$  Cyclic, Constacyclic, Quasi-Cyclic Codes, Thier Skew Codes, Cyclic DNA and Skew Cyclic DNA Codes, Prespacetime, vol. 10, no 2, pp.196-213 (2019)