

A note on morphisms to wreath products

Anthony Genevois and Romain Tessera

October 20, 2021

Abstract

Given a morphism $\varphi : G \rightarrow A \wr B$ from a finitely presented group G to a wreath product $A \wr B$, we show that, if the image of φ is a sufficiently large subgroup, then $\ker(\varphi)$ contains a non-abelian free subgroup and φ factors through an acylindrically hyperbolic quotient of G . As direct applications, we classify the finitely presented subgroups in $A \wr B$ up to isomorphism and we deduce that a group having a wreath product (non-trivial) \wr (infinite) as a quotient must be SQ-universal (extending theorems of Baumslag and Cornuier-Kar). Finally, we exploit our theorem in order to describe the structure of the automorphism groups of several families of wreath products, highlighting an interesting connection with the Kaplansky conjecture on units in group rings.

Contents

1	Introduction	1
2	A few words about graph products of groups	5
3	Proof of the main theorem	8
4	Automorphisms	11
A	A word about approximating infinitely presented groups	16
	References	19

1 Introduction

The starting point of the article is the following surprising observation, which can be found in Baumslag's book [Bau93, Theorem IV.7]: a finitely presented group G that admits a lamplighter group $\mathbb{Z}/n\mathbb{Z} \wr \mathbb{Z}$ ($n \geq 2$) as a quotient is necessarily *large*, i.e. it contains a finite-index subgroup that surjects onto a free group of rank two. The result is surprising, at least at first glance, because, under the seemingly innocent assumption that our group G is finitely presented, we deduce from the existence of a specific solvable quotient the fact that G is very far from being solvable. Unfortunately, the proof given in [Bau93] does not really clarify this phenomenon. A well-known alternative argument, which, in our opinion, is more transparent, is the following. It turns out that, even though $\mathbb{Z}/n\mathbb{Z} \wr \mathbb{Z}$ is solvable, it can be approximated by virtually free groups: by truncating the infinite presentation

$$\langle t, a_i \ (i \in \mathbb{Z}) \mid a_i^n = 1, ta_it^{-1} = a_{i+1} \ (i \in \mathbb{Z}), [a_i, a_j] = 1 \ (i, j \in \mathbb{Z}) \rangle$$

2010 *Mathematics subject classification*. Primary 20E22, 20F65. Secondary 20E36, 20F28, 20F67.
Key words and phrases. Wreath products, Automorphism groups, SQ-universal groups.

of $\mathbb{Z}/n\mathbb{Z} \wr \mathbb{Z}$ as

$$\langle t, a_i \ (i \in \mathbb{Z}) \mid a_i^n = 1, ta_it^{-1} = a_{i+1} \ (i \in \mathbb{Z}), [a_i, a_j] = 1 \ (|i - j| \leq k) \rangle$$

for some $k \geq 1$, one recognises the HNN extension

$$\left\langle t, a_0, \dots, a_k \left| \begin{array}{l} a_i^n = 1 \ (0 \leq i \leq k) \\ [a_i, a_j] = 1 \ (0 \leq i, j \leq k) \\ ta_it^{-1} = a_{i+1} \ (0 \leq i \leq k-1) \end{array} \right. \right\rangle = \langle a_0, \dots, a_k \rangle_{\langle a_0, \dots, a_{k-1} \rangle}^*$$

of $(\mathbb{Z}/n\mathbb{Z})^{k+1}$, which is virtually free non-abelian. Then, it is not difficult to deduce from the fact that G is finitely presented that a quotient map $G \twoheadrightarrow \mathbb{Z}/n\mathbb{Z} \wr \mathbb{Z}$ factors through one of these truncations (see Fact A.1), proving that G is indeed large.

This point of view was developed in [CK11] for arbitrary wreath products $A \wr B := \bigoplus_B A \rtimes B$, by observing that truncations of a well-chosen (infinite) presentation have a very specific algebraic structure: they decompose as a semidirect product between a graph product of copies of A with B . In particular, this observation allowed the authors of [CK11] to generalise Baumslag's observation (see Theorem 1.4 below).

In our article, we combine this point of view on wreath products together with the geometric perspective on graph products introduced by the first-named author in [Gen17]. As a result, we are able to deduce severe restrictions on morphisms from finitely presented groups to wreath products. The next statement, which is the main result of the article, summarises the restrictions we find:

Theorem 1.1. *Let A, B, G be three groups. Assume that G is finitely presented. Every morphism $\varphi : G \rightarrow A \wr B$ factors through a quotient \overline{G} of G such that:*

- *if $\varphi(G)$ has an infinite projection on B and intersects non-trivially $\bigoplus_B A \leq A \wr B$, then $\ker(\varphi)$ contains a non-abelian free subgroup and \overline{G} is acylindrically hyperbolic;*
- *if $\varphi(G)$ has infinite projection on B and is not contained in a conjugate of B , then \overline{G} acts on a finite-dimensional CAT(0) cube complex with unbounded orbits and with hyperplane-stabilisers virtually embedded in finite powers of A .*

In particular, in the second item, \overline{G} semisplits over a subgroup that virtually embeds in a power of A ; so, if A is finite, then \overline{G} is multi-ended.

Loosely speaking, this theorem states that either $\varphi : G \rightarrow A \wr B$ is uninteresting, in the sense that it essentially comes from a morphism to B or a power of A ; or it has a large kernel and it factors through a negatively curved quotient of G that acts non-trivially on a finite-dimensional CAT(0) cube complex.

Theorem 1.1 might be difficult to digest, so let us illustrate a few specific aspects of it.

Nearly injective morphisms. As a direct consequence of Theorem 1.1, a morphism $\varphi : G \rightarrow A \wr B$ from a finitely presented group G to a wreath product $A \wr B$ that has a *small* kernel (i.e. that does not contain a non-abelian free subgroup) must be quite specific: either the image $\varphi(G)$ of G projects isomorphically into B , or it lies in $\bigoplus_B A \rtimes F \leq A \wr B$ for some finite subgroup $F \leq B$. Applying this observation to injective morphisms allows us to classify the finitely presented subgroups in $A \wr B$.

Corollary 1.2. *Let A, B, G be three groups. Assume that G is finitely presented and B infinite. Then G is isomorphic to a subgroup of $A \wr B$ if and only if G is isomorphic to a subgroup of B or if there exists an $n \geq 1$ such that G is a (subgroup in A^n)-by-(finite subgroup in B).*

In other words, the finitely presented subgroups of a wreath product $A \wr B$ essentially comes from the finitely presented subgroups of B and the powers of A . The wreath product structure does not create new finitely presented subgroups.

Proof of Corollary 1.2. Assume that G is isomorphic to a subgroup of $A \wr B$. By applying Theorem 1.1 to an injective morphism $\varphi : G \hookrightarrow A \wr B$, it follows that either G is isomorphic to a finitely presented subgroup of B or it embeds in $\bigoplus_B A \rtimes F$ for some finite subgroup $F \leq B$. In the latter case, because G is finitely generated, only finitely many copies of A contains non-trivial coordinates of elements in the image of G , so G must be a (subgroup in A^n)-by-(finite subgroup in B) for some $n \geq 1$. Conversely, it is clear that finitely presented subgroups of B are finitely presented subgroups of $A \wr B$. According to [KK51], a group G that is a (subgroup in A^n)-by-(finite subgroup in B) embeds into $A^n \wr F$ for some finite subgroup $F \leq B$. But $A^n \wr F$ embeds into $A \wr B$ because B is infinite, so G necessarily embeds into $A \wr B$, as desired. \square

For instance, for all $n, m \geq 1$, the finitely presented subgroups of $\mathbb{Z}^n \wr \mathbb{Z}^m$ are free abelian; and the finitely presented subgroups of $\mathbb{Z}/n\mathbb{Z} \wr \mathbb{Z}^m$ are free abelian or sums of cyclic groups whose orders divide n .

Nearly surjective morphisms. In the opposite direction, if a finitely presented group G admits a morphism $\varphi : G \rightarrow A \wr B$ to a wreath product $A \wr B$ that has a “sufficiently large” image (i.e. such that $\varphi(G)$ has a non-trivial intersection with $\bigoplus_B A$ and an infinite projection on B) then Theorem 1.1 implies that G must have an acylindrically hyperbolic quotient. As a consequence, G must be *SQ-universal* (i.e. every countable group is isomorphic to a subgroup in a quotient of G). Thus, one obtains the following statement, in the same spirit as Baumslag’s observation:

Corollary 1.3. *Let A, B be two groups and G a finitely presented groups. Assume that A is non-trivial and that B is infinite. If G admits $A \wr B$ as a quotient, then it must be SQ-universal.*

Proof. The corollary follows from Theorem 1.1 and from the fact that acylindrically hyperbolic groups are SQ-universal [DGO17, Theorem 2.33]. \square

Observe that, compared to Baumslag’s result, we weakened the conclusion: instead of being large, our group is only SQ-universal. In fact, deducing that the group has to be large can be done for a vast class of wreath products:

Theorem 1.4 ([CK11]). *Let A be a group admitting a non-trivial finite quotient and B an infinite residually finite group. Every finitely presented group admitting $A \wr B$ as a quotient must be large.*

Nevertheless, one easily sees that it is not true that a finitely presented group admitting a wreath product (non-trivial) \wr (infinite) as a quotient is automatically large, as shown by the next examples.

Example 1.5. If A, B are two finitely presented groups with no proper finite-index subgroup, then the free product $A * B$ defines a finitely presented group that admits the wreath product $A \wr B$ as a quotient but that is not large. Indeed, every finite-index subgroup of $A * B$ has to contain both A and B , and so cannot be proper since A and B generate $A * B$. In other words, $A * B$ also has no proper finite-index subgroup. On the other hand, every morphism $A * B \rightarrow \mathbb{F}_2$ must contain A and B in its kernel, so its image must be trivial. In conclusion, $A * B$ is far from being large.

As a consequence, the free product $\mathbb{Z}/2\mathbb{Z} * A$, which admits $\mathbb{Z}/2\mathbb{Z} \wr A$ as a quotient, cannot be large either. Indeed, the kernel of the projection $\mathbb{Z}/2\mathbb{Z} * A \rightarrow \mathbb{Z}/2\mathbb{Z}$ is $A * zAz^{-1}$, where

z denotes the non-trivial element of the factor $\mathbb{Z}/2\mathbb{Z}$. But our previous observation shows that $A * zAz^{-1}$ cannot be large, which implies that $\mathbb{Z}/2\mathbb{Z} * A$ is not large, as desired.

One the simplest finitely presented group with no proper finite-index subgroup is the so-called Higman group $H := \langle x_i \ (i \in \mathbb{Z}/4\mathbb{Z}) \mid x_{i+1}x_i x_{i+1}^{-1} = x_i^2 \ (i \in \mathbb{Z}/4\mathbb{Z}) \rangle$ [Hig51]. Other possible examples come from infinite finitely presented simple groups, such as Thompson's groups T and V [CFP96].

Example 1.5 justifies that the property of being large has been replaced with the property of being SQ-universal.

As a by-product, observe that Corollary 1.3, combined with Pichot's observation from [Osa18], can be used in order to show that some (free) Burnside groups are not finitely presented. Even though this is known for the Burnside groups for which we know that they are infinite (see for instance the proof of [Cou14, Theorem 7.7]), up to our knowledge there is no proof that they cannot be finitely presented as soon as they are infinite.

Proposition 1.6. *Let $k, m, n \geq 1$ be three integers. If the Burnside group $B(m, n)$ is infinite, then $B(m + 1, kn)$ is not finitely presented.*

Recall that $B(m, n)$ is defined as $\langle x_1, \dots, x_m \mid w(x_1, \dots, x_m)^n = 1 \text{ for every word } w \rangle$.

Proof of Proposition 1.6. The wreath product $\mathbb{Z}/k\mathbb{Z} \wr B(m, n)$ admits a generating set of cardinality $m + 1$ and all its elements have exponent kn . Therefore, there exists a surjective morphism $B(m + 1, kn) \twoheadrightarrow \mathbb{Z}/k\mathbb{Z} \wr B(m, n)$. If $B(m, n)$ is infinite, then it follows from Corollary 1.3 that $B(m + 1, kn)$ cannot be finitely presented. (Observe that the latter assertion also follows from Corollary 1.7 below.) \square

Fixed-point property. Finally, it follows from Theorem 1.1 that finitely presented groups satisfying strong fixed-point properties do not have many morphisms to wreath products. More precisely:

Corollary 1.7. *Let A, B, G be three groups and $\varphi : G \rightarrow A \wr B$ a morphism. If G is finitely presented and satisfies the property (FW_{fin}) , then $\varphi(G)$ has a finite projection on B or it lies in a conjugate of B .*

We say that a group satisfies the property (FW_{fin}) if it cannot act on a finite-dimensional CAT(0) cube complex with unbounded orbits, or, equivalently, without a global fixed point. Because Kazhdan's property (T) can be formulated as a fixed-point property on (complete) median spaces [CDH10] and that median graphs, a discrete counterpart of median spaces, define canonically the same objects as CAT(0) cube complexes [Che00, Rol98], one can think of the property (FW_{fin}) as being a discrete and finite-dimensional version of Kazhdan's property (T). Examples include groups satisfying (T), such as $SL(n, \mathbb{Z})$ for $n \geq 3$ and uniform lattices in quaternionic hyperbolic spaces, but also finitely generated torsion groups [Sag95] and Thompson's groups T, V [Gen19b, Kat19].

Automorphisms. However, our main application of Theorem 1.1 deals with automorphisms of wreath products. It seems that automorphism groups of (finitely generated) wreath products have not been extensively studied in the literature. Here, we observe that, if B from our wreath product $A \wr B$ is finitely presented and one-ended, then Theorem 1.1 immediately implies that every automorphism of $A \wr B$ sends B to one of its conjugates. We deduce from this rigidity the following structure result:

Corollary 1.8. *Let A be a finite abelian group, thought of as the additive group of a commutative ring, and B a one-ended finitely presented group. Then*

$$\text{Aut}(A \wr B) = \left(\bigoplus_B A \oplus A[B]^\times \right) \rtimes \text{Aut}(B).$$

Here, $\bigoplus_B A$ is identified with its image under the canonical morphism $A \wr B \rightarrow \text{Inn}(A \wr B)$ (which is injective here) to the subgroup of inner automorphisms, $\text{Aut}(B)$ is the subgroup of $\text{Aut}(A \wr B)$ corresponding to the automorphisms of B that we naturally extend to $A \wr B$, and $A[B]^\times$ corresponds to a subgroup of $\text{Aut}(A \wr B)$ isomorphic to the units of the group ring $A[B]$. We refer to Section 4 for more details.

As a particular case, $\text{Out}(A \wr \mathbb{Z}^n)$ contains a finite-index subgroup isomorphic to $\text{GL}(n, A)$ for every $n \geq 2$. This contrasts with the fact that $\text{Out}(A \wr \mathbb{Z})$ is virtually isomorphic to $\bigoplus_{\mathbb{Z}} A$, and so is not finitely generated. (See Fact 4.6.)

Interestingly, Corollary 1.8 exhibits a connection between automorphism groups of wreath products and units in group rings. Understanding the structure of group rings, or even units in group rings, is notably difficult. As an illustration, recall that Kaplansky's unit conjecture - which claims that the units in the group ring $F[G]$ of a torsion-free group G over a field F are all of the form $k \cdot g$ for some $k \in K^*$, $g \in G$ - has been disproved only recently [Gar21]. By extension, fully understanding the structure of the automorphism groups of specific wreath products might be quite difficult.

As an immediate consequence of Corollary 1.8:

Corollary 1.9. *Let A be a finite abelian group, thought of as the additive group of a commutative ring, and B a one-ended finitely presented group. If $\text{Aut}(B)$ is finitely generated and if B satisfies the Kaplansky unit conjecture over A , then $\text{Out}(A \wr B)$ is finitely generated.*

Here, we say that B satisfies the Kaplansky unit conjecture over A if every unit in $A[B]$ is trivial, i.e. of the form $a \cdot b$ for some $a \in A^\times$, $b \in B$. For instance, when A is a field, it is known that B satisfies the Kaplansky unit conjecture over A if it satisfies the so-called *unique product property*, which includes in particular orderable [Pas85] and locally indicable groups [BH72].

Section 4 contains a discussion about automorphism groups of wreath products, including a generalisation of Corollary 1.8 (see Proposition 4.3) and some hints suggesting that the structure of automorphism groups of arbitrary wreath products can be quite tricky.

Acknowledgments. We are grateful to Rémi Coulon and Simon André for instructive discussions related to Propositions 1.6 and A.4; and to Yves Cornulier for his comments that allow us to improve the presentation of the article.

2 A few words about graph products of groups

As shown in the next section, wreath products turn out to be tightly related to *graph products of groups*. This section is dedicated to basic definitions and preliminary statements related to these products.

Given a simplicial graph Γ and a collection of groups $\mathcal{G} = \{G_u \mid u \in V(\Gamma)\}$ indexed by the vertices of Γ , the *graph product* $\Gamma\mathcal{G}$ is defined by relative presentation

$$\langle G_u \ (u \in V(\Gamma)) \mid [G_u, G_v] = 1 \ (\{u, v\} \in E(\Gamma)) \rangle$$

where $E(\Gamma)$ denotes the edge-set of Γ and where $[G_u, G_v] = 1$ is a shortcut for $[x, y] = 1$ for all $x \in G_u, y \in G_v$. The groups in \mathcal{G} are referred to as *vertex-groups*. If all the vertex-groups are isomorphic to a single group A , we also denote $\Gamma\mathcal{G}$ by ΓA .

Normal form. Fix a simplicial graph Γ and a collection of groups \mathcal{G} indexed by $V(\Gamma)$. A *word* in $\Gamma\mathcal{G}$ is a product $g_1 \cdots g_n$ where $n \geq 0$ and where each g_i belongs to a vertex-group; the g_i 's are the *syllables* of the word, and n is the *length* of the word. Clearly, the following operations on a word do not modify the element of $\Gamma\mathcal{G}$ it represents:

- (O1) delete the syllable $g_i = 1$;
- (O2) if $g_i, g_{i+1} \in G$ for some $G \in \mathcal{G}$, replace the two syllables g_i and g_{i+1} by the single syllable $g_i g_{i+1} \in G$;
- (O3) if g_i and g_{i+1} belong to two adjacent vertex-groups, switch them.

A word is *graphically reduced* if its length cannot be shortened by applying these elementary moves. Given a word $g_1 \cdots g_n$ and some $1 \leq i < n$, if the vertex-group associated to g_i is adjacent to each of the vertex-groups of g_{i+1}, \dots, g_n , then the words $g_1 \cdots g_n$ and $g_1 \cdots g_{i-1} \cdot g_{i+1} \cdots g_n \cdot g_i$ represent the same element of $\Gamma\mathcal{G}$; we say that g_i *shuffles to the right*. Analogously, one can define the notion of a syllable shuffling to the left. If $g = g_1 \cdots g_n$ is a graphically reduced word and h a syllable, then a graphical reduction of the product gh is given by

- $g_1 \cdots g_n$ if $h = 1$;
- $g_1 \cdots g_{i-1} \cdot g_{i+1} \cdots g_n$ if $h \neq 1$, g_i shuffles to the right and $g_i = h^{-1}$;
- $g_1 \cdots g_{i-1} \cdot g_{i+1} \cdots g_n \cdot (g_i h)$ if $h \neq 1$, g_i and h belong to the same vertex-group, g_i shuffles to the right, and $g_i \neq h^{-1}$, where $(g_i h)$ is thought of as a single syllable.

As a consequence, every element g of $\Gamma\mathcal{G}$ can be represented by a graphically reduced word, and this word is unique up to applying the operation (O3). Often, elements of $\Gamma\mathcal{G}$ and graphically reduced words are identified. We refer to [Gre90] for more details (see also [Gen19c] for a more geometric approach).

In graph products, there is also a notion of cyclic reduction. A word $g_1 \cdots g_n$ is *graphically cyclically reduced* if it is graphically reduced and if there do not exist two indices $1 \leq i < j \leq n$ such that g_i and g_j belong to the same vertex-group, g_i shuffles to the left, and g_j shuffles to the right. Notice that, as proved in [Gre90], every element $g \in \Gamma\mathcal{G}$ can be uniquely written as a graphically reduced product xyx^{-1} where y is graphically cyclically reduced. The subgraph of Γ generated by the vertices corresponding to the vertex-groups used to write the syllables of y is the *support* of g , denoted by $\text{supp}(g)$. In other words, the support of an element g is the smallest subgraph $\Lambda \subset \Gamma$ such that g belongs to a conjugate of the subgroup generated by the vertex-groups indexing the vertices of Λ .

Quasi-median geometry. Let Γ be a simplicial graph and \mathcal{G} a collection of groups labelled by $V(\Gamma)$. The geometry of the graph product $\Gamma\mathcal{G}$ turns out to be encoded in the following Cayley graph:

$$\text{QM}(\Gamma, \mathcal{G}) := \text{Cayl} \left(\Gamma\mathcal{G}, \bigcup_{u \in V(\Gamma)} G_u \setminus \{1\} \right),$$

i.e. the graph whose vertices are the elements of the group $\Gamma\mathcal{G}$ and whose edges link two distinct vertices $x, y \in \Gamma\mathcal{G}$ if $y^{-1}x$ is a non-trivial element of some vertex-group. Like




Figure 1: Example of a graph $\text{QM}(\Gamma, \mathcal{G})$.




Figure 2: Four hyperplanes in a quasi-median graph, coloured in red, blue, green, and orange. The orange hyperplane is transverse to the blue and red hyperplanes.

in any Cayley graph, edges of $\text{QM}(\Gamma, \mathcal{G})$ are labelled by generators, namely by elements of vertex-groups. By extension, paths in $\text{QM}(\Gamma, \mathcal{G})$ are naturally labelled by words of generators. In particular, geodesics in $\text{QM}(\Gamma, \mathcal{G})$ correspond to words of minimal length. More precisely:

Lemma 2.1. *Let Γ be a simplicial graph and \mathcal{G} a collection of groups indexed by $V(\Gamma)$. Fix two vertices $g, h \in \text{QM}(\Gamma, \mathcal{G})$. If $s_1 \cdots s_n$ is a graphically reduced word representing $g^{-1}h$, then*

$$g, gs_1, gs_1s_2, \dots, gs_1 \cdots s_{n-1}, gs_1 \cdots s_{n-1}s_n = h$$

defines a geodesic in $\text{QM}(\Gamma, \mathcal{G})$ from g to h . Conversely, if s_1, \dots, s_n is the sequence of elements of $\Gamma\mathcal{G}$ labelling the edges of a geodesic in $\text{QM}(\Gamma, \mathcal{G})$ from g to h , then $s_1 \cdots s_n$ is a graphically reduced word representing $g^{-1}h$.

In [Gen17], it is shown that $\text{QM}(\Gamma, \mathcal{G})$ is a *quasi-median graph* and a general geometry of such graphs is developed in analogy with $\text{CAT}(0)$ cube complexes. The central idea is that hyperplanes can be defined in quasi-median graphs in such a way that the geometry of the quasi-median graph reduces to the combinatorics of its hyperplanes.

Definition 2.2. Let X be a quasi-median graph. A *hyperplane* is a class of edges with respect to the transitive closure of the relation claiming that two edges that are opposite in a square or that belong to a common triangle are equivalent. The *carrier* of a hyperplane J , denoted by $N(J)$, is the subgraph of $\text{QM}(\Gamma, \mathcal{G})$ generated by J . Two hyperplanes J_1 and J_2 are *transverse* if they intersect a square along two distinct pairs of opposite edges.

We refer to Figure 2 for examples of hyperplanes. The main difference between quasi-median graphs and $\text{CAT}(0)$ cube complexes is that hyperplanes separate the complex into exactly two connected components in the latter case when they separate the graph in at least two (and possibly infinitely many) connected components in the former case.

Theorem 2.3. [Gen17, Proposition 2.15] *Let X be a quasi-median graph and J a hyperplane. The graph obtained from X by removing the interiors of all the edges in J contains at least two connected components, referred to as sectors.*

Negative curvature. Typically, graph products are acylindrically hyperbolic. More precisely, given a finite simplicial graph Γ and a collection of non-trivial groups \mathcal{G} indexed by $V(\Gamma)$, the graph product $\Gamma\mathcal{G}$ is acylindrically hyperbolic as soon as the graph Γ contains at least two vertices and is not a join [MO15]. (Recall that Γ is a join if there exists a non-trivial partition $V(\Gamma) = A \sqcup B$ such that every vertex in A is adjacent to every vertex in B .) This negative curvature comes from the *irreducible elements* in $\Gamma\mathcal{G}$, namely the elements whose support is neither a single vertex nor a join. Such elements act on the quasi-median graph $\text{QM}(\Gamma, \mathcal{G})$ like rank-one isometries, as justified by the following statement (which is a consequence of [Gen18, Proposition 4.24]).

Lemma 2.4. *Let Γ be a simplicial graph and \mathcal{G} a collection of groups indexed by $V(\Gamma)$. If $g \in \Gamma\mathcal{G}$ is an irreducible element, then it skewers a pair of strongly separated hyperplanes in $\text{QM}(\Gamma, \mathcal{G})$.*

Recall that, given a quasi-median graph X , an isometry $g \in \text{Isom}(X)$ skewers a pair of hyperplanes $\{J_1, J_2\}$ if there exist a power $n \in \mathbb{Z}$ and two sectors J_1^+, J_2^+ respectively delimited by J_1, J_2 such that $g^n \cdot J_1^+ \subset J_2^+$. The hyperplanes J_1 and J_2 are *strongly separated* if no hyperplane in X is transverse to both J_1 and J_2 .

Our proof of Theorem 1.1 will be based on the following criterion:

Proposition 2.5. *Let G be a group acting on a quasi-median graph X . If G contains an element that skewers a pair of strongly separated hyperplanes $\{J_1, J_2\}$ such that $\text{stab}(J_1) \cap \text{stab}(J_2)$ is finite, then G must be acylindrically hyperbolic or virtually cyclic.*

Proof. Consider the collection of walls

$$\mathcal{SW} := \{\{S, S^c\} \mid S \subset X \text{ sector}\}.$$

According to [Gen17, Lemma 4.17], any two vertices in X are separated by only finitely many walls in \mathcal{SW} , i.e. (X, \mathcal{SW}) is a wallspace. If G contains an element that skewers a pair $\{J_1, J_2\}$ of strongly separated hyperplanes in X , then by construction such an element also skewers a pair $\{J'_1, J'_2\}$ of strongly separated hyperplanes in the CAT(0) cube complex $C(X, \mathcal{SW})$ constructed by cubulating the wallspace (X, \mathcal{SW}) , where J'_1 (resp. J'_2) is the hyperplane in $C(X, \mathcal{SW})$ associated to the wall $\{S_1, S_1^c\}$ (resp. $\{S_2, S_2^c\}$) defined by the sector S_1 (resp. S_2) delimited by J_1 (resp. J_2) that contains J_2 (resp. J_1). As $\text{stab}(J_1) \cap \text{stab}(J_2) = \text{stab}(J'_1) \cap \text{stab}(J'_2)$, the desired conclusion follows from [Gen19a, Theorem 2]. \square

3 Proof of the main theorem

Fix two groups A and B . The wreath product $A \wr B$ admits the following relative presentation:

$$\langle A_b (b \in B), B \mid bA_1b^{-1} = A_b (b \in B), [A_1, A_b] = 1 (b \in B) \rangle$$

where $[A_1, A_b] = 1$ is a shortcut for $[x, y] = 1$ for all $x \in A_1, y \in A_b$. Fixing a subset $S \subset B$, let $A \square_S B$ denote the group defined by the truncated presentation

$$\langle A_b (b \in B), B \mid bA_1b^{-1} = A_b (b \in B), [A_1, A_b] = 1 (b \in S) \rangle,$$

or equivalently by

$$\langle A_b \ (b \in B), B \mid bA_1b^{-1} = A_b \ (b \in B), [A_{b_1}, A_{b_2}] = 1 \ (b_1, b_2 \in B, b_1^{-1}b_2 \in S) \rangle.$$

We denote by π_S the canonical quotient map $A\Box_S B \rightarrow A \wr B$. From the latter presentation, it follows that $A\Box_S B$ decomposes as the semidirect product $\Gamma_S A \rtimes B$ where $\Gamma_S := \text{Cayl}(B, S)$ and where B acts on $\Gamma_S A$ by permuting the vertex-groups according to its action on the Cayley graph Γ_S by left-multiplication.

The key idea, first expressed in [Cor06], is that the wreath product $A \wr B$ can be approximated by $A\Box_S B = \Gamma_S A \rtimes B$ by taking a sufficiently large subset $S \subset B$. As an illustration of this idea:

Lemma 3.1. *Let A, B be two groups. For every finitely presented group G and every morphism $\varphi : G \rightarrow A \wr B$, there exists a finite subset $S \subset B$, and a morphism $\psi : G \rightarrow A\Box_S B$ such that $\varphi = \pi_S \circ \psi$.*

Proof. Fix a generating set $\{s_1, \dots, s_k\}$ of G . For every $1 \leq i \leq k$, let w_i denote a word written over $A^{\pm 1} \cup B^{\pm 1}$ such that $\varphi(s_i) = w_i$ in $A \wr B$. Such a word exists because $A \cup B$ generates $A \wr B$. Thought of as words written over $\{w_1, \dots, w_k\}^{\pm 1}$, the relations of a finite presentation of G can be deduced from finitely many relations of the presentation of $A \wr B$ written above. Consequently, by taking a sufficiently large finite subset $S \subset B$, we can define a morphism $\psi : G \rightarrow A\Box_S B$ satisfying $\varphi = \pi_S \circ \psi$ by setting $\psi(s_i) = w_i$ in $A\Box_S B$. \square

We are now ready to prove the main theorem of the article.

Proof of Theorem 1.1. Let $\varphi : G \hookrightarrow A \wr B$ be a morphism. Because G is finitely presented, we know from Lemma 3.1 that there exist a finite subset $S \subset B$ and a morphism $\psi : G \rightarrow A\Box_S B$ such that $\varphi = \pi_S \circ \psi$. We claim that $\psi(G)$ is the quotient of G we are looking for.

In order to prove the first item in our theorem, we assume that $\varphi(G)$ has an infinite projection on B and a non-trivial intersection with $\bigoplus_B A \leq A \wr B$. As a consequence, $\psi(G)$ has an infinite projection on B and a non-trivial intersection with $\Gamma_S A$. We want to prove that $\ker(\varphi)$ contains a non-abelian free subgroup and that $\psi(G)$ is acylindrically hyperbolic.

Fix a non-trivial element $g \in \Gamma_S A \cap \psi(G)$. Because B acts metrically properly on Γ_S and because $\psi(G)$ has an infinite projection on B , there must exist some $h := (a, b) \in \psi(G)$ such that the distance in Γ_S between the supports of g and bgb^{-1} is at least two. Notice that the same statement holds for g and hgh^{-1} . As a consequence, $\langle g, h \rangle$ is non-abelian free subgroup in $\psi(G)$. Moreover, the derived subgroup of $\langle g, h \rangle$ has trivial image under π_S , so it must lie in $\ker(\psi)$ and a fortiori in $\ker(\varphi)$. Thus, we have proved our first assertion.

Next, consider the action of $\Gamma_S A$ on its quasi-median graph QM. This action clearly extends to $\Gamma_S A \rtimes B$ since the action of B on $\Gamma_S A$ by conjugations permutes the vertex-groups (and so stabilises the generating set used to define QM). To be precise, for all $g \in \Gamma_S$ and $b \in B$, we have

$$gb : x \mapsto gbx b^{-1}, \text{ for every vertex } x \in \text{QM}.$$

Our goal is to deduce from its action on QM that $\psi(G)$ is acylindrically hyperbolic, which will prove our second assertion.

Claim 3.2. *The group $\Gamma_S A \rtimes B$ acts on QM with trivial edge-stabilisers.*

Indeed, if $e = (x, xs)$ is an edge of QM and if $h \in \Gamma_S A \rtimes B$ stabilises e , then, writing $h = gb$ for some $g \in \Gamma_S A$ and $b \in B$, we have

$$\begin{cases} gbx b^{-1} = x \\ gbx s b^{-1} = xs \end{cases} .$$

From

$$x \cdot b s b^{-1} = g b x b^{-1} \cdot b s b^{-1} = g b x s b^{-1} = xs,$$

we deduce that $b s b^{-1} = s$. But B acts freely on the vertices of Γ_S , so necessarily $b = 1$. It then follows that $g = 1$ from the equality $g b x b^{-1} = x$.

Claim 3.3. *The intersection $\psi(G) \cap \Gamma_S A$ contains an irreducible element.*

Fix an enumeration $\{g_1, g_2, \dots\}$ of $\psi(G) \cap \Gamma_S A$. For every $n \geq 1$, set $G_n := \langle g_1, \dots, g_n \rangle$ and let Λ_n denote the smallest subgraph in Γ_S such that G_n lies in a conjugate of the subgroup $\langle \Lambda_n \rangle$ generated by the vertex-groups indexed by the vertices in Λ_n (such a subgraph exists according to [AM15, Proposition 3.10] and it is finite). Because B acts metrically properly on Γ_S and because $\psi(G)$ has an infinite projection on B , the sequence $\Lambda_1 \subset \Lambda_2 \subset \dots$ does not stabilise. As a consequence, since Γ_S is locally finite, there exists an index $n \geq 1$ such that Λ_n contains at least two vertices and is not a join. According to [MO15, Theorem 6.16], G_n contains an element whose support is Λ_n . Such an element is irreducible by definition, concluding the proof of our claim.

By combining Claim 3.3 and Lemma 2.4, it follows that $\psi(G)$ contains an element that skewers a pair $\{J_1, J_2\}$ of strongly separated hyperplanes in QM. Let $x_1 \in N(J_1)$ and $x_2 \in N(J_2)$ be two vertices minimising the distance between the carriers of J_1 and J_2 . Because J_1 and J_2 are strongly separated, it follows from [GM19, Corollary 2.16] that x_1 and x_2 are the only such vertices. Consequently, $\text{stab}(J_1) \cap \text{stab}(J_2) \subset \text{stab}(x_1) \cap \text{stab}(x_2)$. Because there exist only finitely many geodesics between x_1 and x_2 (see Lemma 2.1), we deduce from Claim 3.2 that $\text{stab}(x_1) \cap \text{stab}(x_2)$, and a fortiori $\text{stab}(J_1) \cap \text{stab}(J_2)$, is finite. As a consequence of Proposition 2.5, $\psi(G)$ is either acylindrically hyperbolic or virtually cyclic. But we saw in the second paragraph of this proof that $\psi(G)$ contains a non-abelian free subgroup, so the latter case cannot happen.

Finally, in order to justify the second item in our theorem, we need to show that, if $\varphi(G)$ has infinite projection on B and is not contained in a conjugate of B , then $\psi(G)$ acts non-trivially on a finite-dimensional CAT(0) cube complex with hyperplane-stabilisers that virtually embeds in finite powers of A . Because QM is canonically quasi-isometric to a CAT(0) cube complex of dimension $\text{clique}(\Gamma_S)$ according to [Gen17, Proposition 4.16 and Corollary 8.7], it suffices to show that, under our assumptions, $\psi(G)$ has unbounded orbits in QM and that hyperplane-stabilisers in $A \square_S B$ virtually embeds into a finite power of A . The latter assertion is clear thanks to [Gen17, Corollary 8.10] (see also [GM19, Theorem 2.10]). If $\psi(G)$ instead has finite orbits, then it has to stabilise a *prism* of QM, namely a coset $g \langle \Lambda \rangle \subset \text{QM}$ for some $g \in \Gamma_S A$ and some complete subgraph $\Lambda \subset \Gamma_S$ [Gen17, Theorem 2.115 and Corollary 8.7]. Up to conjugating $\psi(G)$ with an element in $\bigoplus_A B$, we assume that $g = 1$. Observe that, if Λ is empty, then $\langle \Lambda \rangle$ is reduced to the single vertex 1 and its stabiliser in $A \square_S B$ is B . If so,

$$\varphi(G) = \pi_S(\psi(G)) \subset \pi_S(B) = B,$$

proving that $\varphi(G)$ is contained in B (up to conjugacy). Otherwise, if Λ is non-empty, then the stabiliser of $\langle \Lambda \rangle$ in $\Gamma_S A$ has finite index in the stabiliser of $\langle \Lambda \rangle$ in $A \square_S B$, hence

$$\varphi(G) = \pi_S(\psi(G)) \underset{\text{virtually}}{\subset} \pi_S(\Gamma_S A) = \bigoplus_B A,$$

proving that $\varphi(G)$ has finite projection on B . This concludes the proof of our theorem. \square

4 Automorphisms

We fix two groups F, H .

Automorphisms of H . Given an automorphism $\varphi \in \text{Aut}(H)$, the map

$$\bar{\varphi} : \begin{cases} F \wr H & \rightarrow & F \wr H \\ (c, p) & \mapsto & (c \circ \varphi^{-1}, \varphi(p)) \end{cases}$$

defines an automorphism of $F \wr H$. Moreover, the map $\varphi \mapsto \bar{\varphi}$ defines a monomorphism $\text{Aut}(H) \hookrightarrow \text{Aut}(F \wr H)$. In the sequel, we will identify $\text{Aut}(H)$ with its image in $\text{Aut}(F \wr H)$ when this does not cause ambiguity.

H -equivariant automorphisms of $\bigoplus_H F$. For every automorphism $\varsigma \in \text{Aut}(\bigoplus_H F)$ that is equivariant with respect to the action of H on $\bigoplus_H F$ by coordinate-shifting, i.e. $\varsigma(h \cdot c) = h \cdot \varsigma(c)$ for all $h \in H$ and $c \in \bigoplus_H F$, the map

$$\bar{\varsigma} : \begin{cases} F \wr H & \rightarrow & F \wr H \\ (c, p) & \mapsto & (\varsigma(c), p) \end{cases}$$

defines an automorphism of $F \wr H$. We denote by $\text{Aut}_H(\bigoplus_H F)$ the subgroup of $\text{Aut}(F \wr H)$ consisting of all these automorphisms.

Verifying that these maps define automorphisms is a direct and straightforward computation. Notice that, for every automorphism $\varphi \in \text{Aut}(F)$,

$$\hat{\varphi} : (c, h) \mapsto (\varphi \circ c, h), \quad (c, h) \in F \wr H$$

yields an example of an H -equivariant automorphism of $\bigoplus_H F$. When there is no possible confusion, we identify $\text{Aut}(F)$ with its image in $\text{Aut}(F \wr H)$.

Our first observation is that these automorphisms, together with the inner automorphisms, are sufficient to generate the automorphism group in many cases.

Proposition 4.1. *Let F be a finite group and H a one-ended finitely presented group. Then*

$$\text{Aut}(F \wr H) = \text{Inn}(F \wr H) \cdot \left(\text{Aut}_H \left(\bigoplus_H F \right) \rtimes \text{Aut}(H) \right).$$

In particular, $\text{Aut}(F \wr H)$ is generated by the inner automorphisms, the automorphisms of H , and the H -equivariant automorphisms of $\bigoplus_H F$.

Proof. Let $\varphi \in \text{Aut}(F \wr H)$ be an automorphism. As a consequence of Theorem 1.1, $\varphi(H)$ lies in a conjugate of H . Up to composing φ with an inner automorphism, we assume that $\varphi(H) \subset H$. Similarly, $\varphi^{-1}(H)$ lies in a conjugate of H , say gHg^{-1} . Observe that $H = \varphi^{-1}(\varphi(H)) \subset \varphi^{-1}(H) \subset gHg^{-1}$. But H is almost malnormal (see for instance [GT21, Lemma 7.11]), so we must have $H = gHg^{-1}$. Thus, $\varphi(H) = H$. Up to composing φ with an automorphism of H , we assume that φ fixes H pointwise. In other words, there exists an automorphism $\varsigma : \bigoplus_H F \rightarrow \bigoplus_H F$ such that $\varphi(f, p) = (\varsigma(f), p)$ for every $(f, p) \in F \wr H$. Necessarily, ς must be H -equivariant, i.e. φ is induced by an H -equivariant automorphism of $\bigoplus_H F$.

Thus, we have proved the decomposition

$$\text{Aut}(F \wr H) = \text{Inn}(F \wr H) \cdot \text{Aut}(H) \cdot \text{Aut}_H \left(\bigoplus_H F \right).$$

As $\text{Aut}(H)$ intersects trivially and normalises $\text{Aut}_H(\bigoplus_H F)$, the desired conclusion follows. \square

It is worth noticing that the proof above shows that the conclusion of Proposition 4.1 holds more generally when F is an arbitrary group and when H is a finitely presented group that does not semisplit over a subgroup of $\bigoplus_H F$. For instance, this applies for wreath products of the form $\mathbb{Z} \wr$ (rigid hyperbolic group). (A hyperbolic group is *rigid* if it is a non-elementary and if it does not split over a virtually cyclic subgroup.)

In view of Proposition 4.1, a natural problem is to investigate the structure of the group of H -equivariant automorphisms of $\bigoplus_H F$. In the sequel, we focus on the case when F can be thought of as the underlying additive group of a ring, i.e. when F is abelian. Even restricting to this case, we shall see that understanding H -equivariant automorphisms may be quite tricky.

Units of $F[H]$. If F is the additive group of a commutative ring, identifying $\bigoplus_H F$ with the group ring $F[H]$ allows us to endow $\bigoplus_H F$ with a product \cdot such that $(\bigoplus_H F, +, \cdot)$ is a ring. For every unit $u \in \bigoplus_H F$, the map

$$\bar{u} : \begin{cases} F \wr H & \rightarrow & F \wr H \\ (c, p) & \mapsto & (c \cdot u, p) \end{cases}$$

defines an automorphism of $F \wr H$. Observe that two distinct units induce two distinct automorphisms of $F \wr H$. We denote by $F[H]^\times$ the subgroup of $\text{Aut}(F \wr H)$ consisting of all these automorphisms.

Verifying that \bar{u} defines an automorphism of $F \wr H$ is a straightforward computation once we have observed that the action of H on $\bigoplus_H F$ by coordinate-shifting coincides with the action of H by left multiplication in the ring $F[H]$.

Lemma 4.2. *Assume that F is the additive group of a commutative ring. The H -equivariant automorphisms of $\bigoplus_H F$ are induced by the units of $F[H]$.*

Proof. Let $\sigma \in \text{Aut}(\bigoplus_H F)$ be an H -equivariant automorphism. Being H -equivariant amounts to saying that $\sigma(h \cdot c) = h \cdot \sigma(c)$ for all $h \in H$ and $c \in \bigoplus_H F$. By linearity, it follows that $\sigma(c_1 \cdot c_2) = c_1 \cdot \sigma(c_2)$ for all $c_1, c_2 \in \bigoplus_H F$. In other words, σ is morphism of $(\bigoplus_H F)$ -module. Consequently,

$$\sigma(c) = \sigma(c \cdot 1) = c \cdot \sigma(1) \text{ for every } c \in \bigoplus_H F,$$

where $\sigma(1)$ is a unit (indeed, if $c \in \bigoplus_H F$ is such that $\sigma(c) = 1$, then one has $c \cdot \sigma(1) = \sigma(c) = 1$). \square

As a consequence of Lemma 4.2, understanding the H -equivariant automorphisms of $F \wr H$ is as hard as understanding the units of the group ring $F[H]$. The latter problem is well-known to be quite difficult. Indeed, the so-called Kaplansky unit conjecture, which states that the group ring $F[H]$ has only trivial units when F is a field and H torsion-free, remained an open problem for a long time and has been disproved only recently [Gar21]. Therefore, Lemma 4.2 suggests that the structure of $\text{Aut}_H(\bigoplus_H F)$, and a fortiori of $\text{Aut}(F \wr H)$, may be highly non-trivial and goes beyond the scope of our current understanding in full generality.

Proof of Corollary 1.8. It follows from Proposition 4.1 and Lemma 4.2 that

$$\text{Aut}(A \wr B) = \text{Inn}(A \wr B) \cdot A[B]^\times \cdot \text{Aut}(B).$$

Because a conjugation by an element of $A \wr B$ decomposes as a product of a conjugation by an element of $\bigoplus_B A$ with a conjugation of an element of B , and that the latter also belongs to $A[B]^\times$, we have

$$\text{Aut}(A \wr B) = \bigoplus_B A \cdot A[B]^\times \cdot \text{Aut}(B).$$

Clearly, $\bigoplus_B A$ and $A[B]^\times$ intersect trivially and commute, so $\bigoplus_B A \cdot A[B]^\times = \bigoplus_B A \oplus A[B]^\times$. Also, observe that $\text{Aut}(B)$ intersects trivially $\bigoplus_B A \cdot A[B]^\times$ and normalises both $\bigoplus_B A$ and $A[B]^\times$. The desired decomposition of $\text{Aut}(A \wr B)$ follows. \square

Outside the one-ended case, the elementary automorphisms introduced so far are not sufficient to generate the automorphism group of the corresponding lamplighter group. It seems difficult to exhibit a structure that is both completely general and really informative. Nevertheless, one can push further.

Partial conjugations. Assume that F is an abelian group and that H splits as a free product $H_1 * H_2$. Given an $a \in \bigoplus_H F$, the map

$$\kappa : s \mapsto \begin{cases} s & \text{if } s \in F \cup H_1 \\ asa^{-1} & \text{if } s \in H_2 \end{cases}$$

induces an automorphism of $F \wr H$.

Here, κ is only defined on generators of $F \wr H$, and it is naturally extended to words of generators. In order to justify that κ induces a well-defined morphism $F \wr H \rightarrow F \wr H$, it suffices to show that a relator of F, H_1, H_2 or from the relative presentation

$$\langle F, H \mid [hfh^{-1}, f] = 1 \text{ where } f \in F, h \in H \setminus \{1\} \rangle$$

has always trivial image under κ . Clearly, κ sends a relator of F, H_1, H_2 to the identity element, so the desired conclusion follows from the observation that κ fixes $\bigoplus_H F$ pointwise. Indeed, let $f \in F$ and $h \in H$ be two elements. Decompose h as an alternating product $a_1 b_1 \cdots a_n b_n$ where $a_1, \dots, a_n \in H_1$ and $b_1, \dots, b_n \in H_2$. Observe that

$$\begin{aligned} \kappa(a_1 b_1 \cdot f \cdot b_1^{-1} a_1^{-1}) &= a_1 a b_1 a^{-1} \cdot f \cdot a b_1^{-1} a^{-1} a_1^{-1} \\ &= a_1 a b_1 \cdot f \cdot b_1^{-1} a^{-1} a_1^{-1} = a_1 b_1 \cdot f \cdot b_1^{-1} a_1, \end{aligned}$$

where the penultimate equality is justified by the fact that F is abelian and where the last equality is justified by the fact that $b_1 f b_1^{-1}$ commutes with a since $\bigoplus_H F$ is abelian. By iterating, one shows that $\kappa(hfh^{-1}) = hfh^{-1}$, as desired. That the kernel of κ is trivial is a direct consequence of the facts that κ induces the identity on H and restricts to the identity on $\bigoplus_H F$. Finally, the surjectivity of κ follows from the easy observation that F, H_1, H_2 all lie in the image of κ .

Transvections. Assume that F is abelian and that H splits as a free product $H' * Z$ where Z is cyclic, say generated by $z \in Z$ of order $r \in \mathbb{N} \cup \{\infty\}$. Given an $a \in \bigoplus_H F$ such that $(az)^r = 1$ if $r < \infty$, the map

$$\kappa : s \mapsto \begin{cases} s & \text{if } s \in F \cup H' \\ as & \text{if } s = z \end{cases}$$

induces an automorphism of $F \wr H$.

Again, κ is only defined on generators and is naturally extended to words of generators. In order to justify that κ induces a well-defined automorphism of $F \wr H$, it suffices to reproduce the previous argument almost word for word, starting from the relative presentation

$$\langle F, H', z \mid z^r = 1, [hfh^{-1}, f] = 1 \text{ where } f \in F, h \in \langle H', z \rangle \setminus \{1\} \rangle.$$

When r is finite, the elements $a \in \bigoplus_H F$ satisfying $(az)^r = 1$ are obtained in the following way. Decompose $\text{supp}(a)$ as a disjoint union $S_1 \sqcup \dots \sqcup S_k$ of $\langle z \rangle$ -orbits. Write a as $a_1 \dots a_k$ such that $\text{supp}(a_i) = S_i$ for every $1 \leq i \leq k$. Observe that

$$\begin{aligned} (az)^r &= a \cdot z a z^{-1} \dots z^{r-1} a z^{-r+1} \\ &= (a_1 \cdot z a_1 z^{-1} \dots z^{r-1} a_1 z^{-r+1}) \dots (a_k \cdot z a_k z^{-1} \dots z^{r-1} a_k z^{-r+1}). \end{aligned}$$

Because z has order r , each S_i has size r and z acts on it through a cyclic shifting, hence

$$a_i \cdot z a_i z^{-1} \dots z^{r-1} a_i z^{-r+1} = \prod_{s \in S_i} s.$$

Thus, in order to construct a , we took a collection of $\langle z \rangle$ -invariant subsets $S_1, \dots, S_k \subset H$, and, for each $1 \leq i \leq k$, we coloured the lamps in S_i such that the product of all the colours is trivial.

We emphasize that the free product decompositions used in order to define partial conjugations and transvections may be trivial. For instance, $\mathbb{Z} \wr \mathbb{Z}$ has transvections (which turns out to generate the outer automorphism group up to finite index).

Thanks to these new families of automorphisms, we are now able to generalise Proposition 4.1 to some multi-ended groups.

Proposition 4.3. *Let F be a finite abelian group, thought of as the additive group of a commutative ring, and H a finitely presented group that decomposes as a free product of one-ended and cyclic groups. Then*

$$\text{Aut}(F \wr H) = (\text{PConj}(F \wr H) \cdot \text{Trans}(F \wr H) \cdot F[H]^\times) \rtimes \text{Aut}(H).$$

We emphasize that, in this statement, the partial conjugations and the transvections are taken with respect to a fixed free product decomposition.

Proof of Proposition 4.3. Decompose H as a free product $H_1 * \dots * H_n$ of one-ended and cyclic groups, and let $\varphi \in \text{Aut}(F \wr H)$ be an automorphism. Because $\bigoplus_H F$ is characteristic, up to composing φ with an automorphism of H we assume that φ induces the identity on H .

Assume that there exists some $1 \leq i \leq n$ such that H_i is one-ended. As a consequence of Theorem 1.1, $\varphi(H_i)$ must lie in a conjugate of H . So there exist $1 \leq j \leq n$ and $g \in F \wr H$ such that $\varphi(H_i) \subset g H_j g^{-1}$. Since we know that the projection of $\varphi(H_i)$ on H is H_i , necessarily $j = i$ and $g \in (\bigoplus_H F) H_i$. Therefore, $\varphi(H_i) \subset a H_i a^{-1}$ for some $a \in \bigoplus_H F$. By composing φ with a partial conjugation, we can assume that $\varphi(H_i) \subset H_i$. Because φ induces the identity on H , it follows that φ fixes H_i pointwise.

Next, assume that there exists some $1 \leq i \leq n$ such that H_i is cyclic. Fix a generator $h_i \in H_i$. Because φ induces the identity on H , there must exist some $a \in \bigoplus_H F$ such that $\varphi(h_i) = a h_i$. Of course, h_i and $a h_i$ have the same order, since φ is an automorphism, so we can assume that φ fixes h_i up to composing it with a transvection.

Thus, we have proved that, up to composing φ with an automorphism of H followed by a product of pairwise commuting partial conjugations and transvections, we can assume that φ fixes H pointwise. Then φ is induced by an H -equivariant automorphism of $\bigoplus_H F$, and the desired conclusion follows from Lemma 4.2.

In other words, we have proved that

$$\text{Aut}(F \wr H) = \text{Aut}(H) \cdot \text{PConj}(F \wr H) \cdot \text{Trans}(F \wr H) \cdot F[H]^\times.$$

Because $\text{Aut}(H)$ normalises $\text{PConj}(F \wr H)$, $\text{Trans}(F \wr H)$, and $F[H]^\times$, and because it intersects trivially $\text{PConj}(F \wr H) \cdot \text{Trans}(F \wr H) \cdot F[H]^\times$, the desired decomposition follows. \square

Proposition 4.3 describes the automorphism groups of many interesting lamplighter groups, but we are still far away from a global picture. Let us conclude this section with a few open questions.

First, it is worth noticing that lamplighters over groups that are not finitely presented are not considered at all in the article. This is justified by the fact that admitting a finite presentation is fundamental in Theorem 1.1. However, it is natural to ask what happens for infinitely presented groups. For instance:

Question 4.4. Let F be a non-trivial finite group. Describe the automorphism group of the iterated wreath product $F \wr (F \wr (\dots (F \wr \mathbb{Z}) \dots))$.

In the article, we mainly focused on lamps with only finitely many colours. Even though we saw that the proof of Proposition 4.1 also provides interesting information outside this situation, many cases of interest are not covered. As a particular case:

Question 4.5. Describe the automorphism group of $\mathbb{F}_n \wr \mathbb{Z}$, $n \geq 2$.

We excluded the case $n = 1$ from the question because it can be easily solved:

Fact 4.6. Let Z be a cyclic group (finite or infinite). Then

$$\text{Aut}(Z \wr \mathbb{Z}) = \text{Inn}(Z \wr \mathbb{Z}) \cdot \text{Aut}(Z) \cdot \text{Trans}(Z \wr \mathbb{Z}) \cdot \langle \iota \rangle$$

where ι denotes the automorphism that inverts the \mathbb{Z} -factor. As a consequence, $\text{Out}(Z \wr \mathbb{Z})$ contains a finite-index subgroup isomorphic to $\bigoplus_{\mathbb{Z}} Z$.

Proof. Let φ be an automorphism of $Z \wr \mathbb{Z}$. Because $\bigoplus_{\mathbb{Z}} Z$ is characteristic, φ induces an isomorphism $\mathbb{Z} \rightarrow \mathbb{Z}$, which we can assume to be trivial up to composing φ with ι . Consequently, there exists some $a \in \bigoplus_{\mathbb{Z}} Z$ such that $\varphi(z) = az$ where z denotes a generator of the \mathbb{Z} -factor. Up to composing φ with a transvection, we assume that $a = 1$. So φ fixes pointwise the \mathbb{Z} -factor. In other words, φ has to be a \mathbb{Z} -equivariant automorphism of $\bigoplus_{\mathbb{Z}} Z$, which amounts to saying that φ belongs to $Z[\mathbb{Z}]^\times$ according to Lemma 4.2. However, because \mathbb{Z} is abelian, $Z[\mathbb{Z}]^\times$ lies in $\text{Inn}(Z \wr \mathbb{Z}) \cdot \text{Aut}(Z)$, so the desired conclusion follows. \square

For wreath products of the form (finite group) \wr (multi-ended group), we only considered the case the finite group is abelian, because otherwise the partial conjugations and transvections may not be well-defined. This suggests the following problem.

Question 4.7. Let F be a non-abelian finite group and $n \geq 2$ an integer. Describe the automorphism group of $F \wr \mathbb{F}_n$.

The difficulty in extending Proposition 4.3 to lamplighters over arbitrary multi-ended groups comes from the torsion, which may introduce sufficient flexibility in order to define new types of automorphisms.

Generalised partial conjugations. Assume that F is abelian and that H splits as a graph of groups with underlying graph Γ . Fix a marked vertex $v_0 \in V(\Gamma)$, an orientation on Γ , and, for every edge $e \in E(\Gamma)$, an element $a_e \in C(H_e) \cap \bigoplus_H F$. Every path in Γ is naturally labelled by the element obtained by reading the oriented edges it passes through. Assume that the label of every cycle is trivial. For every vertex $v \in V(\Gamma)$, let $a_v \in \bigoplus_H F$ denote the label of a path from v_0 to v . (As a consequence of our previous assumption, this label does not depend on the path we chose.) Fix a maximal subtree $T \subset \Gamma$, let e_1, \dots, e_k denote the oriented edges outside T , and let t_1, \dots, t_k denote the corresponding stable letters. The map

$$\kappa : s \mapsto \begin{cases} s & \text{if } s \in F \\ a_v s a_v^{-1} & \text{if } s \in H_v \\ a_{o(e_i)} t_i a_{o(e_i)}^{-1} & \text{if } s = t_i \end{cases}$$

induces an automorphism of $F \wr H$.

The proof is in the same spirit as before, but we use Bass-Serre theory in order to find a relative presentation of H . We do not include the details here. Similarly, one can define generalised transvections.

Understanding such automorphisms in full generality turns out to be tricky since they seem to depend on the structures of the finite groups involved and on how they interact together. As a test question, we propose:

Question 4.8. Let A, B be two non-trivial finite groups and $C \leq B$ a subgroup. Describe the automorphism group of $A \wr (B *_C B)$.

A A word about approximating infinitely presented groups

The main idea underlying the article is that a group H admitting a presentation of the form $\langle X \mid R, r_1, r_2, \dots \rangle$, where R is a set of relations and where r_1, r_2, \dots is an infinite sequence of relations, can be “approximated” by the groups H_i defined by the truncated presentations $\langle X \mid R, r_1, \dots, r_i \rangle$. The proof of Theorem 1.1 is an algebraic illustration of this idea applied to wreath products. This appendix is a general discussion dedicated to this strategy from different points of view and for other infinitely presented groups.

Algebraic approximation. The fact that the H_i algebraically approximate, in a sense, the group H is illustrated by the following (well-known) observation:

Fact A.1. *For every finitely presented group G and every morphism $\varphi : G \rightarrow H$, there exist an $i \geq 1$ and a morphism $\psi : G \rightarrow H_i$ such that $\varphi = \pi_i \circ \psi$.*

The proof follows the same lines as Lemma 3.1, which is a particular case. Interestingly, the H_i may satisfy properties that are quite different from the properties satisfied by H . For instance, applying the approximation given in Section 3 to the lamplighter group $H = \mathbb{Z}/2\mathbb{Z} \wr \mathbb{Z}$ yields virtually non-abelian free groups H_i , even though H is solvable. This implies that every finitely presented group admitting $\mathbb{Z}/2\mathbb{Z} \wr \mathbb{Z}$ as a quotient must virtually surject onto \mathbb{F}_2 , as already mentioned in the introduction. Similar statements can be obtained for infinitely presented groups that are not wreath products, as shown below.

Geometric approximation. Fact A.1 admits a geometric version, justifying that the H_i also provide geometric approximations of H . More precisely:

Fact A.2. *Let X be a coarsely 1-connected graph. For every coarse (resp. quasi-isometric) embedding $\rho : X \rightarrow H$, there exist an $i \geq 1$ and a coarse (resp. quasi-isometric) embedding $\eta : X \rightarrow H_i$ such that $\rho = \pi_i \circ \eta$.*

Recall that a graph X is *coarsely 1-connected* if there exists an $R \geq 0$ such that the 2-complex obtained from X by filling in with discs all the cycle of length $\leq R$ is simply connected. If X is a Cayley graph of a finitely generated group G (constructed from a finite generating set), then X is coarsely 1-connected precisely when G is finitely presented.

Fact A.2 is central in the proof of the embedding theorem obtained in [GT21, Theorem 5.1]; see [GT21, Lemma 2.2].

Logic approximation. One can also observe that, in the space of marked groups, the (H_n, X) converge to (H, X) . Because convergence in the space of marked groups is connected to first-order properties, one obtains the following observation:

Fact A.3. *Let G be a finitely presented group and $S \subset G$ a finite generating set. If $\text{Th}_\forall(G) \supset \text{Th}_\forall(H)$, then (G, S) is a limit of marked groups (K_n, S_n) where each K_n is both a subgroup of H_n and a quotient of G .*

We refer to [CG05] for details about marked groups and universal theories of groups.

Proof of Fact A.3. According to [CG05, Proposition 5.3], the inclusion $\text{Th}_\forall(G) \supset \text{Th}_\forall(H)$ implies that (G, S) is a limit of marked subgroups of H in the space of marked groups. But we also know that H (endowed with an arbitrary marking) is the limit of the H_i (for some marking). Consequently, (G, S) is a limit of marked subgroups $K_n \leq H_n$. Since we know that some neighbourhood of (G, S) contains only marked quotients of G [CG05, Lemma 2.3], we can suppose without loss of generality that H_n is also a quotient of G for every $n \geq 0$. \square

As an illustration, let us show that no finitely presented group have the same universal theory as, for instance, the lamplighter group $\mathbb{Z} \wr \mathbb{Z}$.

Proposition A.4. *Let A, B two free abelian groups of finite rank ≥ 1 , and let G be a finitely generated group. If G and $A \wr B$ have the same universal theory, then G is not finitely presented.*

Proof. Notice that like $A \wr B$, G is metabelian but not abelian, since these properties can be expressed as universal or existential formulae. According to Fact A.3, if G is finitely presented then (G, S) can be described as a limit of marked groups (K_n, S_n) , where $S \subset G$ is an arbitrary finite generating set and where each K_n is both a quotient of G and a subgroup of $A \square_{R_n} B$ for some $R_n \subset B$. We distinguish two cases.

Case 1: The projection of K_n on B is trivial for infinitely many n . For such an n , K_n lies in $\Gamma_{R_n} A$ (which is a right-angled Artin group). Because K_n cannot contain a non-abelian free subgroup, necessarily K_n is abelian. So G has to be abelian, which is not possible.

Case 2: The intersection between K_n and $\Gamma_{R_n} A$ is trivial for infinitely many n . Then, for such an n , K_n embeds into B and so is abelian. Again, one gets a contradiction.

The remaining case: for all $n \geq 0$, K_n has an infinite projection on B and a non-trivial intersection with $\Gamma_{R_n} A$. By applying Theorem 1.1 to the morphism

$$G \twoheadrightarrow K_n \hookrightarrow A \square_{S_n} B \twoheadrightarrow A \wr B$$

for a sufficiently large n , it follows that G has to be SQ-universal, and a fortiori it contains a non-abelian free subgroup, a contradiction with the fact that G is metabelian. So G cannot be finitely presented. \square

Other infinitely presented groups. At first glance, it might be surprising that a finitely presented group that surjects onto a wreath product (non-trivial)(infinite) must be SQ-universal, even though the wreath product itself may be very far from being SQ-universal (e.g. it may be solvable). As described in Section 3, the reason comes from the fact that wreath products can be approximated by groups satisfying properties of negative curvature, and the SQ-universality is deduced from such properties.

Interestingly, such a phenomenon is not specific to wreath products and seems to be rather common. For instance, [BGdlH13] exhibits many similar examples among metabelian groups and branch groups. In the particular case of the Grigorchuk group \mathfrak{G} , the structure of the approximating groups is completely explicit. Recall from [Lys85, Lyu84] that \mathfrak{G} admits as a presentation

$$\langle a, b, c, d \mid a^2 = b^2 = c^2 = d^2 = bcd = 1, u_n = v_n = 1 \ (n \geq 0) \rangle$$

where the words u_n, v_n are defined inductively as

$$\begin{cases} u_0 = (ad)^4 \text{ and } v_0 = (adacac)^4 \\ u_n = \sigma^n(u_0) \text{ and } v_n = \sigma^n(v_0) \text{ for all } n \geq 0 \end{cases}$$

from the substitution σ defined by

$$\sigma(a) = aca, \ \sigma(b) = d, \ \sigma(c) = b, \ \sigma(d) = c.$$

Then, for every $n \geq 0$, the group \mathfrak{G}_n defined by the truncated presentation

$$\left\langle a, b, c, d \mid \begin{array}{l} a^2 = b^2 = c^2 = d^2 = bcd = 1 \\ u_0 = \cdots = u_n = v_0 = \cdots = v_{n-1} = 1 \end{array} \right\rangle$$

turns out to be virtually a product of free groups (see [BGdlH13, Theorem 5.3] for a more precise statement). Let us mention two applications of this point of view:

Proposition A.5. *For every finitely presented group G and every morphism $\varphi : G \rightarrow \mathfrak{G}$, if the image of φ is not virtually abelian, then G has to be large.*

Proof. As a consequence of Fact A.1, φ factors through a morphism $\psi : G \rightarrow \mathfrak{G}_n$ for some $n \geq 0$. Because \mathfrak{G}_n is virtually a product of free groups, $\psi(G)$ is either virtually abelian or large. In the former case, the image of φ in \mathfrak{G} has to be virtually abelian; and in the latter case, G has to be large. \square

Proposition A.6. *A finitely generated nilpotent group quasi-isometrically embeds into the Grigorchuk group \mathfrak{G} if and only if it is virtually abelian.*

Proof. Because a finitely generated nilpotent group is automatically finitely presented, it follows from Fact A.2 that such a group that quasi-isometrically embeds in the Grigorchuk group must quasi-isometrically embed in a product of finitely many free groups. On the other hand, a finitely generated nilpotent group can be realised as a uniform lattice in a connected simply-connected nilpotent Lie group; and such nilpotent Lie group does not quasi-isometrically embed in a CAT(0) space such as a product of trees according to [Pau01], unless it is abelian. We conclude that a finitely generated nilpotent group that quasi-isometrically embeds in the Grigorchuk group must be virtually abelian. Conversely, because the Grigorchuk is commensurable to its own square, \mathfrak{Z}^n quasi-isometrically embeds into \mathfrak{G} for every $n \geq 1$, so every finitely generated virtually abelian group quasi-isometrically embeds in the Grigorchuk group. \square

Another source of examples is the following. Given a group G , let $\mathcal{H}(G)$ denote the group $\text{Bij}_c(G) \rtimes G \leq \text{Bij}(G)$ where $\text{Bij}_c(G)$ is the group of finitely supported permutations of G . Observe that $\mathcal{H}(\mathbb{Z})$ coincides with the second Houghton group H_2 . Fixing a symmetric generating set $S \subset G$, $\mathcal{H}(G)$ can be alternatively described as the semidirect product $C(\Gamma) \rtimes G$, where $C(\Gamma)$ is the Coxeter group defined by the labelled complete graph Γ on the edge-set $E := \{e_{g,s} := \{g, gs\} \mid g \in G, s \in S\}$ of the Cayley graph $\text{Cayl}(G, S)$ such that the label of $\{e_1, e_2\}$ for $e_1, e_2 \in E$ distinct is 2 if $e_1 \cap e_2 = \emptyset$ and 3 otherwise. Now, given an $n \geq 1$, let Γ_n denote the labelled complete graph Γ on the edge-set $E := \{e_{g,s} := \{g, gs\} \mid g \in G, s \in S\}$ of the Cayley graph $\text{Cayl}(G, S)$ such that the label of $\{e_1, e_2\}$ for $e_1, e_2 \in E$ distinct is 3 if $e_1 \cap e_2 \neq \emptyset$, 2 if $e_1 \cap e_2 = \emptyset$ and $d(e_1, e_2) \leq n$, and 0 otherwise. Then the semidirect product $\mathcal{H}_n(G) := C(\Gamma_n) \rtimes G$ is obtained by truncating a presentation of $\mathcal{H}(G)$, with the key property that the Coxeter group $C(\Gamma_n)$ is now acylindrically hyperbolic (similarly to the graph products in Section 3). Following the proof of Theorem 1.1 leads to a similar statement for $\mathcal{H}(G)$.

As a variation of the previous construction, one can permute the elements of G with braids instead of just permutations. More precisely, one can define the semidirect product $\mathcal{B}(G) := \mathcal{B}_c(G) \rtimes G$ where $\mathcal{B}_c(G)$ denotes the group of finitely supported braids on G . Then, mimicking the previous argument, truncating a well-chosen presentation leads to a semidirect product between an acylindrically hyperbolic Artin-Tits group and G . But proving a statement like Theorem 1.1 is more difficult here because Artin-Tits groups are poorly understood compared to graph products and Coxeter groups.

Of course, the variations go on and on. There are at least as many possible constructions as generalisations of braid groups.

References

- [AM15] Y. Antolín and A. Minasyan. Tits alternatives for graph products. *J. Reine Angew. Math.*, 704:55–83, 2015.
- [Bau93] G. Baumslag. *Topics in combinatorial group theory*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 1993.
- [BGdlH13] M. Benli, R. Grigorchuk, and P. de la Harpe. Amenable groups without finitely presented amenable covers. *Bull. Math. Sci.*, 3(1):73–131, 2013.
- [BH72] R. Burns and V. Hale. A note on group rings of certain torsion-free groups. *Canad. Math. Bull.*, 15:441–445, 1972.
- [CDH10] I. Chatterji, C. Druţu, and F. Haglund. Kazhdan and Haagerup properties from the median viewpoint. *Advances in Mathematics*, 225:882–921, 2010.
- [CFP96] J. Cannon, W. Floyd, and W. Parry. Introductory notes on Richard Thompson’s groups. *Enseign. Math. (2)*, 42(3-4):215–256, 1996.
- [CG05] C. Champetier and V. Guirardel. Limit groups as limits of free groups. *Israel J. Math.*, 146:1–75, 2005.
- [Che00] V. Chepoi. Graphs of some CAT(0) complexes. *Adv. in Appl. Math.*, 24(2):125–179, 2000.
- [CK11] Y. Cornuier and A. Kar. On property (FA) for wreath products. *J. Group Theory*, 14(1):165–174, 2011.
- [Cor06] Y. Cornuier. Finitely presented wreath products and double coset decompositions. *Geom. Dedicata*, 122:89–108, 2006.
- [Cou14] R. Coulon. On the geometry of Burnside quotients of torsion free hyperbolic groups. *Internat. J. Algebra Comput.*, 24(3):251–345, 2014.
- [DGO17] F. Dahmani, V. Guirardel, and D. Osin. Hyperbolically embedded subgroups and rotating families in groups acting on hyperbolic spaces. *Mem. Amer. Math. Soc.*, 245(1156):v+152, 2017.
- [Gar21] G. Gardam. A counterexample to the unit conjecture for group rings. *arXiv:2102.11818*, to appear in *Ann. of Math.*, 2021.

- [Gen17] A. Genevois. Cubical-like geometry of quasi-median graphs and applications to geometric group theory. *PhD thesis, arxiv:1712.01618*, 2017.
- [Gen18] A. Genevois. Automorphisms of graph products of groups and acylindrical hyperbolicity. *arXiv:1807.00622*, 2018.
- [Gen19a] A. Genevois. Acylindrical hyperbolicity from actions on CAT(0) cube complexes: a few criteria. *New York J. Math.*, 25:1214–1239, 2019.
- [Gen19b] A. Genevois. Hyperbolic and cubical rigidities of Thompson’s group V . *J. Group Theory*, 22(2):313–345, 2019.
- [Gen19c] A. Genevois. On the geometry of van Kampen diagrams of graph products of groups. *arXiv:1901.04538*, 2019.
- [GM19] A. Genevois and A. Martin. Automorphisms of graph products of groups from a geometric perspective. *Proceedings of the London Mathematical Society*, 119(6):1745–1779, 2019.
- [Gre90] E. Green. Graph products of groups. *PhD Thesis*, 1990.
- [GT21] A. Genevois and R. Tessera. Asymptotic geometry of lamplighters over one-ended groups. *arxiv:2105.04878*, 2021.
- [Hig51] G. Higman. A finitely generated infinite simple group. *J. London Math. Soc.*, 26:61–64, 1951.
- [Kat19] M. Kato. On groups whose actions on finite-dimensional CAT(0) spaces have global fixed points. *J. Group Theory*, 22(6):1089–1099, 2019.
- [KK51] M. Krasner and L. Kaloujnine. Produit complet des groupes de permutations et problème d’extension de groupes. III. *Acta Sci. Math. (Szeged)*, 14:69–82, 1951.
- [Lys85] I. Lysënok. A set of defining relations for the Grigorchuk group. *Mat. Zametki*, 38(4):503–516, 634, 1985.
- [Lyu84] A. Lyul’ko. Normal subgroups of Abels groups. *Mat. Zametki*, 36(3):289–294, 1984.
- [MO15] A. Minasyan and D. Osin. Acylindrical hyperbolicity of groups acting on trees. *Mathematische Annalen*, 362(3):1055–1105, 2015.
- [Osa18] D. Osajda. Group cubization. *Duke Math. J.*, 167(6):1049–1055, 2018. With an appendix by M. Pichot.
- [Pas85] D. Passman. *The algebraic structure of group rings*. Robert E. Krieger Publishing Co., Inc., Melbourne, FL, 1985. Reprint of the 1977 original.
- [Pau01] S. Pauls. The large scale geometry of nilpotent Lie groups. *Comm. Anal. Geom.*, 9(5):951–982, 2001.
- [Rol98] M. Roller. Pocsets, median algebras and group actions; an extended study of dunwoody’s construction and sageev’s theorem. *dissertation, arXiv:1607.07747*, 1998.
- [Sag95] Michah Sageev. Ends of group pairs and non-positively curved cube complexes. *Proc. London Math. Soc. (3)*, 71(3):585–617, 1995.

INSTITUT MONTPELLIERAIN ALEXANDER GROTHENDIECK, 499-554 RUE DU TRUEL, 34090 MONTPELLIER, FRANCE.

E-mail address: `anthony.genevois@umontpellier.fr`

INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, PLACE AURÉLIE NEMOURS, 75013 PARIS, FRANCE.

E-mail address: `romain.tessera@imj-prg.fr`